



Australian Government

Department of Health



Schedule of Pharmaceutical Benefits

Section 100 – Items Available under Special Arrangement - Volume 2

Effective 1 October 2019

This Schedule is also available at www.pbs.gov.au



This work is copyright. Apart from any use as permitted under the *Copyright Act 1968*, no part may be reproduced by any process without prior written permission from the Commonwealth. Requests and inquiries concerning reproduction and rights should be addressed to the Commonwealth Copyright Administration, Attorney General's Department, Robert Garran Offices, National Circuit, Barton ACT 2600 or posted at <http://www.ag.gov.au>

This Schedule provides information on the arrangements for the prescribing and supply of pharmaceutical benefits. These arrangements operate under the *National Health Act 1953*. However, at the time of printing, the relevant legislation giving authority for the changes included in this issue of the Schedule may still be subject to the usual Parliamentary scrutiny. This book is not a legal document, and, in cases of discrepancy, the legislation will be the source document for payment for the supply of pharmaceutical benefits. The legislation is available from the Federal Register of Legislation website at www.legislation.gov.au.

The information is not intended to give or replace any legal, medical, dental or optometrical advice. This document is not a legal document and does not constitute legal advice. Neither the information nor this document can be relied upon without first seeking and obtaining independent legal, medical, dental or optometrical advice beforehand. To the extent permitted by law, the Commonwealth of Australia will not be held responsible, nor accept any liability (whether arising out of negligence or otherwise), for any injury, damages, costs, expenses and losses suffered or incurred by a person where such a person has relied on this document or used the information in it as legal, medical, dental or optometrical advice.

Contents

Fees, Patient Contributions and Safety Net Thresholds	1
Summary of Changes	2
About the Schedule	22
Symbols and Abbreviations Used in the Schedule	22
Restricted Benefits	23
Guidelines and General Statements	24
General Statement for Drugs for the Treatment of Hepatitis C.....	24
Pharmaceutical Benefits Schedules	27
Highly Specialised Drugs Program (Private Hospital)	28
Highly Specialised Drugs Program (Public Hospital).....	310
Highly Specialised Drugs Program (Community Access).....	592
Botulinum Toxin Program.....	613
Growth Hormone Program	622
IVF Treatment Program.....	893
Opiate Dependence Treatment Program	899
Extemporaneously Prepared Benefits	902
Drug Tariff	903
Container Prices.....	906
Standard Formula Preparations	907
Codes, Maximum Quantities, and Number of Repeats for Extemporaneously Prepared Benefits	908
Index of Manufacturers' Code.....	909
Generic/Proprietary Index	912

Fees, Patient Contributions and Safety Net Thresholds

The following fees, patient contributions and safety net thresholds apply as at 1 October 2019 and are included, where applicable, in prices published in the Schedule —

Dispensing Fees:	Ready-prepared	\$7.39
	Dangerous drug fee	\$3.11
	Extemporaneously-prepared	\$9.43
	Allowable additional patient charge*	\$4.53
Additional Fees (for safety net prices):	Ready-prepared	\$1.25
	Extemporaneously-prepared	\$1.61
Patient Co-payments:	General	\$40.30
	Concessional	\$6.50
Safety Net Thresholds:	General	\$1550.70
	Concessional	\$390.00
Safety Net Card Issue Fee:		\$10.10

* The allowable additional patient charge is a discretionary charge to general patients if a pharmaceutical item has a dispensed price for maximum quantity less than the general patient co-payment. The pharmacist may charge general patients the allowable additional fee but the fee cannot take the cost of the prescription above the general patient co-payment for the medicine. This fee does not count towards the Safety Net threshold.

Summary of Changes

These changes to the Schedule of Pharmaceutical Benefits are effective from 1 October 2019. The Schedule is updated on the first day of each month and is available on the internet at www.pbs.gov.au.

Prescriber Bag

Alterations

Alteration – Manufacturer Code

		From	To
3485K	<i>Cilicaine</i> – PROCAINE BENZYL PENICILLIN (PROCAINE PENICILLIN) , procaine benzylpenicillin (procaine penicillin) 1.5 g/3.4 mL injection, 5 x 3.4 mL syringes	QA	AF

Advance Notices

1 November 2019

Deletion – Brand

3466K	<i>Frusamide Sandoz, SZ</i> – FUROSEMIDE (FRUSEMIDE) , furosemide (frusemide) 20 mg/2 mL injection, 5 x 2 mL ampoules

General Pharmaceutical Benefits

Additions

Addition – Item

2518M	APREPITANT , aprepitant 165 mg capsule, 1 (<i>Aprepitant APOTEX</i>)
11815W	INSULIN GLARGINE , insulin glargine 100 units/mL injection, 5 x 3 mL pen devices (<i>Semglee</i>)
11798Y	OCRIPLASMIN , ocriplasmin 375 microgram/0.3 mL intraocular injection, 0.3 mL vial (<i>Jetrea RTU</i>)

Addition – Brand

8717T	<i>Aripiprazole generichealth, HQ</i> – ARIPIRAZOLE , aripiprazole 10 mg tablet, 30
8718W	<i>Aripiprazole generichealth, HQ</i> – ARIPIRAZOLE , aripiprazole 15 mg tablet, 30
8719X	<i>Aripiprazole generichealth, HQ</i> – ARIPIRAZOLE , aripiprazole 20 mg tablet, 30
8720Y	<i>Aripiprazole generichealth, HQ</i> – ARIPIRAZOLE , aripiprazole 30 mg tablet, 30
8358X	<i>Clopidogrel Sandoz Pharma, HX</i> – CLOPIDOGREL , clopidogrel 75 mg tablet, 28
9317J	<i>Clopidogrel Sandoz Pharma, HX</i> – CLOPIDOGREL , clopidogrel 75 mg tablet, 28
8245Y	<i>Letrozole GH, HQ</i> – LETROZOLE , letrozole 2.5 mg tablet, 30
8887R	<i>MELOBIC, RF</i> – MELOXICAM , meloxicam 7.5 mg capsule, 30
8888T	<i>MELOBIC, RF</i> – MELOXICAM , meloxicam 15 mg capsule, 30
8627C	<i>MONTELAIR 4, RF</i> – MONTELUKAST , montelukast 4 mg chewable tablet, 28
8628D	<i>MONTELAIR 5, RF</i> – MONTELUKAST , montelukast 5 mg chewable tablet, 28
8228C	<i>APO-Nicorandil, TX</i> – NICORANDIL , nicorandil 10 mg tablet, 60
8229D	<i>APO-Nicorandil, TX</i> – NICORANDIL , nicorandil 20 mg tablet, 60
3010K	<i>APO-Norfloxacin, TX</i> – NORFLOXACIN , norfloxacin 400 mg tablet, 14

Addition – Equivalence Indicator

9039R *Lantus SoloStar, AV* – **INSULIN GLARGINE**, insulin glargine 100 units/mL injection, 5 x 3 mL cartridges

Addition – Note

9039R **INSULIN GLARGINE**, insulin glargine 100 units/mL injection, 5 x 3 mL cartridges (*Lantus SoloStar, Lantus*)

Addition – Restriction

9039R **INSULIN GLARGINE**, insulin glargine 100 units/mL injection, 5 x 3 mL cartridges (*Lantus SoloStar, Lantus*)

Deletions**Deletion – Item**

2544X **BIPERIDEN**, biperiden hydrochloride 2 mg tablet, 100 (*Akineton*)
2018F **SODIUM AUROTHIOMALATE**, sodium aurothiomalate 50 mg/0.5 mL injection, 10 x 0.5 mL ampoules (*Myocrisin*)
10721G **TRASTUZUMAB**, trastuzumab 600 mg/5 mL injection, 5 mL vial (*Herceptin SC*)
10825R **TRASTUZUMAB**, trastuzumab 600 mg/5 mL injection, 5 mL vial (*Herceptin SC*)

Deletion – Brand

8315P *DBL Cefepime, PF* – **CEFEPIME**, cefepime 1 g injection, 1 vial
8316Q *DBL Cefepime, PF* – **CEFEPIME**, cefepime 2 g injection, 1 vial
8246B *Irprestan 75, ZP* – **IRBESARTAN**, irbesartan 75 mg tablet, 30
8247C *Irprestan 150, ZP* – **IRBESARTAN**, irbesartan 150 mg tablet, 30
8248D *Irprestan 300, ZP* – **IRBESARTAN**, irbesartan 300 mg tablet, 30
1596B *Onsetron, ZP* – **ONDANSETRON**, ondansetron 4 mg/2 mL injection, 2 mL ampoule
8226Y *Onsetron, ZP* – **ONDANSETRON**, ondansetron 4 mg/2 mL injection, 2 mL ampoule
1597C *Onsetron, ZP* – **ONDANSETRON**, ondansetron 8 mg/4 mL injection, 4 mL ampoule
8227B *Onsetron, ZP* – **ONDANSETRON**, ondansetron 8 mg/4 mL injection, 4 mL ampoule
1594X *Onsetron 4, ZP* – **ONDANSETRON**, ondansetron 4 mg tablet, 10
8224W *Onsetron 4, ZP* – **ONDANSETRON**, ondansetron 4 mg tablet, 4
1595Y *Onsetron 8, ZP* – **ONDANSETRON**, ondansetron 8 mg tablet, 10
8225X *Onsetron 8, ZP* – **ONDANSETRON**, ondansetron 8 mg tablet, 4
8507R *Parzol 10, ZP* – **RABEPRAZOLE**, rabeprazole sodium 10 mg enteric tablet, 28
11670F *Parzol 20, ZP* – **RABEPRAZOLE**, rabeprazole sodium 20 mg enteric tablet, 30
8508T *Parzol 20, ZP* – **RABEPRAZOLE**, rabeprazole sodium 20 mg enteric tablet, 30
8509W *Parzol 20, ZP* – **RABEPRAZOLE**, rabeprazole sodium 20 mg enteric tablet, 30
2477J *Rivastigmine Patch 5, AF* – **RIVASTIGMINE**, rivastigmine 4.6 mg/24 hours patch, 30
9161E *Rivastigmine Patch 5, AF* – **RIVASTIGMINE**, rivastigmine 4.6 mg/24 hours patch, 30
2551G *Rivastigmine Patch 10, AF* – **RIVASTIGMINE**, rivastigmine 9.5 mg/24 hours patch, 30
9162F *Rivastigmine Patch 10, AF* – **RIVASTIGMINE**, rivastigmine 9.5 mg/24 hours patch, 30
10538P *Rivastigmine Patch 15, AF* – **RIVASTIGMINE**, rivastigmine 13.3 mg/24 hours patch, 30
10541T *Rivastigmine Patch 15, AF* – **RIVASTIGMINE**, rivastigmine 13.3 mg/24 hours patch, 30
5299X *Prograf XL, LL* – **TACROLIMUS**, tacrolimus 500 microgram modified release capsule, 30
5300Y *Prograf XL, LL* – **TACROLIMUS**, tacrolimus 1 mg modified release capsule, 60
5451X *Prograf XL, LL* – **TACROLIMUS**, tacrolimus 5 mg modified release capsule, 30
8523N *Lodam SR 100, ZP* – **TRAMADOL**, tramadol hydrochloride 100 mg modified release tablet, 20
8524P *Lodam SR 150, ZP* – **TRAMADOL**, tramadol hydrochloride 150 mg modified release tablet, 20

Deletion – Equivalence Indicator

1596B *Ondansetron Alphapharm, AF* – **ONDANSETRON**, ondansetron 4 mg/2 mL injection, 2 mL ampoule
8226Y *Ondansetron Alphapharm, AF* – **ONDANSETRON**, ondansetron 4 mg/2 mL injection, 2 mL ampoule
1597C *Ondansetron Alphapharm, AF* – **ONDANSETRON**, ondansetron 8 mg/4 mL injection, 4 mL ampoule

8227B	<i>Ondansetron Alphapharm, AF</i> – ONDANSETRON , ondansetron 8 mg/4 mL injection, 4 mL ampoule
2477J	<i>Exelon Patch 5, NV</i> – RIVASTIGMINE , rivastigmine 4.6 mg/24 hours patch, 30
9161E	<i>Exelon Patch 5, NV</i> – RIVASTIGMINE , rivastigmine 4.6 mg/24 hours patch, 30
2551G	<i>Exelon Patch 10, NV</i> – RIVASTIGMINE , rivastigmine 9.5 mg/24 hours patch, 30
9162F	<i>Exelon Patch 10, NV</i> – RIVASTIGMINE , rivastigmine 9.5 mg/24 hours patch, 30
10538P	<i>Exelon Patch 15, NV</i> – RIVASTIGMINE , rivastigmine 13.3 mg/24 hours patch, 30
10541T	<i>Exelon Patch 15, NV</i> – RIVASTIGMINE , rivastigmine 13.3 mg/24 hours patch, 30
5299X	<i>ADVAGRAF XL, LQ</i> – TACROLIMUS , tacrolimus 500 microgram modified release capsule, 30
5300Y	<i>ADVAGRAF XL, LQ</i> – TACROLIMUS , tacrolimus 1 mg modified release capsule, 60
5451X	<i>ADVAGRAF XL, LQ</i> – TACROLIMUS , tacrolimus 5 mg modified release capsule, 30

Deletion – Note

11416W	PEGINTERFERON ALFA-2A , peginterferon alfa-2a 135 microgram/0.5 mL injection, 4 x 0.5 mL syringes (<i>Pegasys</i>)
11037X	PEGINTERFERON ALFA-2A , peginterferon alfa-2a 180 microgram/0.5 mL injection, 4 x 0.5 mL syringes (<i>Pegasys</i>)

Alterations

Alteration – Note

5281Y	ADALIMUMAB , adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL syringes (<i>Humira</i>)
5282B	ADALIMUMAB , adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL pen devices (<i>Humira</i>)
5283C	ADALIMUMAB , adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL syringes (<i>Humira</i>)
5284D	ADALIMUMAB , adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL pen devices (<i>Humira</i>)
9077R	ADALIMUMAB , adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL syringes (<i>Humira</i>)
9078T	ADALIMUMAB , adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL syringes (<i>Humira</i>)
9103D	ADALIMUMAB , adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL pen devices (<i>Humira</i>)
9104E	ADALIMUMAB , adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL pen devices (<i>Humira</i>)
10137M	CERTOLIZUMAB PEGOL , certolizumab pegol 200 mg/mL injection, 2 x 1 mL syringes (<i>Cimzia</i>)
10897M	CERTOLIZUMAB PEGOL , certolizumab pegol 200 mg/mL injection, 2 x 1 mL syringes (<i>Cimzia</i>)
10904X	CERTOLIZUMAB PEGOL , certolizumab pegol 200 mg/mL injection, 2 x 1 mL syringes (<i>Cimzia</i>)
11318Q	CERTOLIZUMAB PEGOL , certolizumab pegol 200 mg/mL injection, 2 x 1 mL pen devices (<i>Cimzia</i>)
11319R	CERTOLIZUMAB PEGOL , certolizumab pegol 200 mg/mL injection, 2 x 1 mL pen devices (<i>Cimzia</i>)
11320T	CERTOLIZUMAB PEGOL , certolizumab pegol 200 mg/mL injection, 2 x 1 mL pen devices (<i>Cimzia</i>)
9125G	DASATINIB , dasatinib 20 mg tablet, 60 (<i>Sprycel</i>)
9126H	DASATINIB , dasatinib 50 mg tablet, 60 (<i>Sprycel</i>)
9127J	DASATINIB , dasatinib 70 mg tablet, 60 (<i>Sprycel</i>)
9343R	DASATINIB , dasatinib 100 mg tablet, 30 (<i>Sprycel</i>)
11204Q	ETANERCEPT , etanercept 25 mg injection [4 vials] (& inert substance diluent [4 x 1 mL syringes], 1 pack (<i>Enbrel</i>)
3445H	ETANERCEPT , etanercept 25 mg injection [4 vials] (& inert substance diluent [4 x 1 mL syringes], 1 pack (<i>Enbrel</i>)
3448L	ETANERCEPT , etanercept 25 mg injection [4 vials] (& inert substance diluent [4 x 1 mL syringes], 1 pack (<i>Enbrel</i>)
8778B	ETANERCEPT , etanercept 25 mg injection [4 vials] (& inert substance diluent [4 x 1 mL syringes], 1 pack (<i>Enbrel</i>)
8779C	ETANERCEPT , etanercept 25 mg injection [4 vials] (& inert substance diluent [4 x 1 mL syringes], 1 pack (<i>Enbrel</i>)
11201M	ETANERCEPT , etanercept 50 mg/mL injection, 4 x 1 mL pen devices (<i>Brenzys, Enbrel</i>)
11215G	ETANERCEPT , etanercept 50 mg/mL injection, 4 x 1 mL pen devices (<i>Brenzys</i>)
3447K	ETANERCEPT , etanercept 50 mg/mL injection, 4 x 1 mL pen devices (<i>Enbrel</i>)
3450N	ETANERCEPT , etanercept 50 mg/mL injection, 4 x 1 mL pen devices (<i>Enbrel</i>)
9455P	ETANERCEPT , etanercept 50 mg/mL injection, 4 x 1 mL pen devices (<i>Brenzys, Enbrel</i>)

9456Q	ETANERCEPT , etanercept 50 mg/mL injection, 4 x 1 mL pen devices (<i>Brenzys, Enbrel</i>)
11196G	ETANERCEPT , etanercept 50 mg/mL injection, 4 x 1 mL syringes (<i>Brenzys, Enbrel</i>)
11217J	ETANERCEPT , etanercept 50 mg/mL injection, 4 x 1 mL syringes (<i>Brenzys</i>)
3446J	ETANERCEPT , etanercept 50 mg/mL injection, 4 x 1 mL syringes (<i>Enbrel</i>)
3449M	ETANERCEPT , etanercept 50 mg/mL injection, 4 x 1 mL syringes (<i>Enbrel</i>)
9085E	ETANERCEPT , etanercept 50 mg/mL injection, 4 x 1 mL syringes (<i>Brenzys, Enbrel</i>)
9086F	ETANERCEPT , etanercept 50 mg/mL injection, 4 x 1 mL syringes (<i>Brenzys, Enbrel</i>)
11361Y	GOLIMUMAB , golimumab 50 mg/0.5 mL injection, 0.5 mL pen device (<i>Simponi</i>)
11376R	GOLIMUMAB , golimumab 50 mg/0.5 mL injection, 0.5 mL pen device (<i>Simponi</i>)
3434R	GOLIMUMAB , golimumab 50 mg/0.5 mL injection, 0.5 mL syringe (<i>Simponi</i>)
3436W	GOLIMUMAB , golimumab 50 mg/0.5 mL injection, 0.5 mL syringe (<i>Simponi</i>)
11765F	IMATINIB , imatinib 400 mg tablet, 30 (<i>Gilmat, Glivec, IMATINIB RBX, Imatinib-Teva</i>)
11251E	LAPATINIB , lapatinib 250 mg tablet, 70 (<i>Tykerb</i>)
9148L	LAPATINIB , lapatinib 250 mg tablet, 70 (<i>Tykerb</i>)
10890E	SECUKINUMAB , secukinumab 150 mg/mL injection, 1 mL pen device (<i>Cosentyx</i>)
10893H	SECUKINUMAB , secukinumab 150 mg/mL injection, 1 mL pen device (<i>Cosentyx</i>)
10906B	SECUKINUMAB , secukinumab 150 mg/mL injection, 1 mL pen device (<i>Cosentyx</i>)
11720W	TOCILIZUMAB , tocilizumab 162 mg/0.9 mL injection, 4 x 0.9 mL syringes (<i>Actemra Subcutaneous Injection</i>)
11725D	TOCILIZUMAB , tocilizumab 162 mg/0.9 mL injection, 4 x 0.9 mL pen devices (<i>Actemra ACTPen</i>)
11730J	TOCILIZUMAB , tocilizumab 162 mg/0.9 mL injection, 4 x 0.9 mL syringes (<i>Actemra Subcutaneous Injection</i>)
11734N	TOCILIZUMAB , tocilizumab 162 mg/0.9 mL injection, 4 x 0.9 mL pen devices (<i>Actemra ACTPen</i>)
11741Y	TOCILIZUMAB , tocilizumab 162 mg/0.9 mL injection, 4 x 0.9 mL syringes (<i>Actemra Subcutaneous Injection</i>)
11742B	TOCILIZUMAB , tocilizumab 162 mg/0.9 mL injection, 4 x 0.9 mL pen devices (<i>Actemra ACTPen</i>)
11748H	TOCILIZUMAB , tocilizumab 162 mg/0.9 mL injection, 4 x 0.9 mL syringes (<i>Actemra Subcutaneous Injection</i>)
11750K	TOCILIZUMAB , tocilizumab 162 mg/0.9 mL injection, 4 x 0.9 mL pen devices (<i>Actemra ACTPen</i>)
10682F	TRASTUZUMAB , trastuzumab 600 mg/5 mL injection, 5 mL vial (<i>Herceptin SC</i>)
10798H	TRASTUZUMAB , trastuzumab 600 mg/5 mL injection, 5 mL vial (<i>Herceptin SC</i>)
10803N	TRASTUZUMAB , trastuzumab 600 mg/5 mL injection, 5 mL vial (<i>Herceptin SC</i>)

Alteration – Restriction

5281Y	ADALIMUMAB , adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL syringes (<i>Humira</i>)
5282B	ADALIMUMAB , adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL pen devices (<i>Humira</i>)
5283C	ADALIMUMAB , adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL syringes (<i>Humira</i>)
5284D	ADALIMUMAB , adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL pen devices (<i>Humira</i>)
9077R	ADALIMUMAB , adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL syringes (<i>Humira</i>)
9078T	ADALIMUMAB , adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL syringes (<i>Humira</i>)
9103D	ADALIMUMAB , adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL pen devices (<i>Humira</i>)
9104E	ADALIMUMAB , adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL pen devices (<i>Humira</i>)
10137M	CERTOLIZUMAB PEGOL , certolizumab pegol 200 mg/mL injection, 2 x 1 mL syringes (<i>Cimzia</i>)
10897M	CERTOLIZUMAB PEGOL , certolizumab pegol 200 mg/mL injection, 2 x 1 mL syringes (<i>Cimzia</i>)
10904X	CERTOLIZUMAB PEGOL , certolizumab pegol 200 mg/mL injection, 2 x 1 mL syringes (<i>Cimzia</i>)
11318Q	CERTOLIZUMAB PEGOL , certolizumab pegol 200 mg/mL injection, 2 x 1 mL pen devices (<i>Cimzia</i>)
11319R	CERTOLIZUMAB PEGOL , certolizumab pegol 200 mg/mL injection, 2 x 1 mL pen devices (<i>Cimzia</i>)
11320T	CERTOLIZUMAB PEGOL , certolizumab pegol 200 mg/mL injection, 2 x 1 mL pen devices (<i>Cimzia</i>)
9125G	DASATINIB , dasatinib 20 mg tablet, 60 (<i>Sprycel</i>)

9126H	DASATINIB , dasatinib 50 mg tablet, 60 (<i>Sprycel</i>)
9127J	DASATINIB , dasatinib 70 mg tablet, 60 (<i>Sprycel</i>)
9343R	DASATINIB , dasatinib 100 mg tablet, 30 (<i>Sprycel</i>)
11204Q	ETANERCEPT , etanercept 25 mg injection [4 vials] (& inert substance diluent [4 x 1 mL syringes], 1 pack (<i>Enbrel</i>)
3445H	ETANERCEPT , etanercept 25 mg injection [4 vials] (& inert substance diluent [4 x 1 mL syringes], 1 pack (<i>Enbrel</i>)
3448L	ETANERCEPT , etanercept 25 mg injection [4 vials] (& inert substance diluent [4 x 1 mL syringes], 1 pack (<i>Enbrel</i>)
8778B	ETANERCEPT , etanercept 25 mg injection [4 vials] (& inert substance diluent [4 x 1 mL syringes], 1 pack (<i>Enbrel</i>)
8779C	ETANERCEPT , etanercept 25 mg injection [4 vials] (& inert substance diluent [4 x 1 mL syringes], 1 pack (<i>Enbrel</i>)
11201M	ETANERCEPT , etanercept 50 mg/mL injection, 4 x 1 mL pen devices (<i>Brenzys</i> , <i>Enbrel</i>)
11215G	ETANERCEPT , etanercept 50 mg/mL injection, 4 x 1 mL pen devices (<i>Brenzys</i>)
3447K	ETANERCEPT , etanercept 50 mg/mL injection, 4 x 1 mL pen devices (<i>Enbrel</i>)
3450N	ETANERCEPT , etanercept 50 mg/mL injection, 4 x 1 mL pen devices (<i>Enbrel</i>)
9455P	ETANERCEPT , etanercept 50 mg/mL injection, 4 x 1 mL pen devices (<i>Brenzys</i> , <i>Enbrel</i>)
9456Q	ETANERCEPT , etanercept 50 mg/mL injection, 4 x 1 mL pen devices (<i>Brenzys</i> , <i>Enbrel</i>)
11196G	ETANERCEPT , etanercept 50 mg/mL injection, 4 x 1 mL syringes (<i>Brenzys</i> , <i>Enbrel</i>)
11217J	ETANERCEPT , etanercept 50 mg/mL injection, 4 x 1 mL syringes (<i>Brenzys</i>)
3446J	ETANERCEPT , etanercept 50 mg/mL injection, 4 x 1 mL syringes (<i>Enbrel</i>)
3449M	ETANERCEPT , etanercept 50 mg/mL injection, 4 x 1 mL syringes (<i>Enbrel</i>)
9085E	ETANERCEPT , etanercept 50 mg/mL injection, 4 x 1 mL syringes (<i>Brenzys</i> , <i>Enbrel</i>)
9086F	ETANERCEPT , etanercept 50 mg/mL injection, 4 x 1 mL syringes (<i>Brenzys</i> , <i>Enbrel</i>)
11361Y	GOLIMUMAB , golimumab 50 mg/0.5 mL injection, 0.5 mL pen device (<i>Simponi</i>)
11376R	GOLIMUMAB , golimumab 50 mg/0.5 mL injection, 0.5 mL pen device (<i>Simponi</i>)
3434R	GOLIMUMAB , golimumab 50 mg/0.5 mL injection, 0.5 mL syringe (<i>Simponi</i>)
3436W	GOLIMUMAB , golimumab 50 mg/0.5 mL injection, 0.5 mL syringe (<i>Simponi</i>)
11251E	LAPATINIB , lapatinib 250 mg tablet, 70 (<i>Tykerb</i>)
9148L	LAPATINIB , lapatinib 250 mg tablet, 70 (<i>Tykerb</i>)
8598M	MESALAZINE , mesalazine 500 mg modified release granules, 100 sachets (<i>Salofalk</i>)
2234N	MESALAZINE , mesalazine 1 g modified release granules, 120 sachets (<i>Pentasa</i>)
8599N	MESALAZINE , mesalazine 1 g modified release granules, 100 sachets (<i>Salofalk</i>)
9206M	MESALAZINE , mesalazine 1.5 g modified release granules, 60 sachets (<i>Salofalk</i>)
2287J	MESALAZINE , mesalazine 2 g modified release granules, 60 sachets (<i>Pentasa</i>)
10257W	MESALAZINE , mesalazine 3 g modified release granules, 30 sachets (<i>Salofalk</i>)
10254Q	MESALAZINE , mesalazine 4 g modified release granules, 30 sachets (<i>Pentasa</i>)
1611T	MESALAZINE , mesalazine 250 mg enteric tablet, 100 (<i>Mesasal</i>)
2214M	MESALAZINE , mesalazine 500 mg modified release tablet, 100 (<i>Pentasa</i>)
8731M	MESALAZINE , mesalazine 500 mg enteric tablet, 100 (<i>Salofalk</i>)
11210B	MESALAZINE , mesalazine 800 mg enteric tablet, 90 (<i>Asacol</i>)
11554D	MESALAZINE , mesalazine 1 g enteric tablet, 60 (<i>Salofalk</i>)
3413P	MESALAZINE , mesalazine 1 g modified release tablet, 60 (<i>Pentasa</i>)
9353G	MESALAZINE , mesalazine 1.2 g modified release tablet, 60 (<i>Mezavant</i>)
10947E	OCRIPLASMIN , ocriplasmin 500 microgram/0.2 mL injection, 0.2 mL vial (<i>Jetrea</i>)
11454W	PONATINIB , ponatinib 15 mg tablet, 60 (<i>Iclusig</i>)
11453T	PONATINIB , ponatinib 45 mg tablet, 30 (<i>Iclusig</i>)
10890E	SECUKINUMAB , secukinumab 150 mg/mL injection, 1 mL pen device (<i>Cosentyx</i>)

10893H	SECUKINUMAB , secukinumab 150 mg/mL injection, 1 mL pen device (<i>Cosentyx</i>)
10906B	SECUKINUMAB , secukinumab 150 mg/mL injection, 1 mL pen device (<i>Cosentyx</i>)
11720W	TOCILIZUMAB , tocilizumab 162 mg/0.9 mL injection, 4 x 0.9 mL syringes (<i>Actemra Subcutaneous Injection</i>)
11725D	TOCILIZUMAB , tocilizumab 162 mg/0.9 mL injection, 4 x 0.9 mL pen devices (<i>Actemra ACTPen</i>)
11730J	TOCILIZUMAB , tocilizumab 162 mg/0.9 mL injection, 4 x 0.9 mL syringes (<i>Actemra Subcutaneous Injection</i>)
11734N	TOCILIZUMAB , tocilizumab 162 mg/0.9 mL injection, 4 x 0.9 mL pen devices (<i>Actemra ACTPen</i>)
11741Y	TOCILIZUMAB , tocilizumab 162 mg/0.9 mL injection, 4 x 0.9 mL syringes (<i>Actemra Subcutaneous Injection</i>)
11742B	TOCILIZUMAB , tocilizumab 162 mg/0.9 mL injection, 4 x 0.9 mL pen devices (<i>Actemra ACTPen</i>)
11748H	TOCILIZUMAB , tocilizumab 162 mg/0.9 mL injection, 4 x 0.9 mL syringes (<i>Actemra Subcutaneous Injection</i>)
11750K	TOCILIZUMAB , tocilizumab 162 mg/0.9 mL injection, 4 x 0.9 mL pen devices (<i>Actemra ACTPen</i>)
10682F	TRASTUZUMAB , trastuzumab 600 mg/5 mL injection, 5 mL vial (<i>Herceptin SC</i>)
10798H	TRASTUZUMAB , trastuzumab 600 mg/5 mL injection, 5 mL vial (<i>Herceptin SC</i>)
10803N	TRASTUZUMAB , trastuzumab 600 mg/5 mL injection, 5 mL vial (<i>Herceptin SC</i>)

Alteration – Restriction Level

		<i>From</i>	<i>To</i>
8598M	MESALAZINE , mesalazine 500 mg modified release granules, 100 sachets (<i>Salofalk</i>)	streamlined	restricted
2234N	MESALAZINE , mesalazine 1 g modified release granules, 120 sachets (<i>Pentasa</i>)	streamlined	restricted
8599N	MESALAZINE , mesalazine 1 g modified release granules, 100 sachets (<i>Salofalk</i>)	streamlined	restricted
9206M	MESALAZINE , mesalazine 1.5 g modified release granules, 60 sachets (<i>Salofalk</i>)	streamlined	restricted
2287J	MESALAZINE , mesalazine 2 g modified release granules, 60 sachets (<i>Pentasa</i>)	streamlined	restricted
10257W	MESALAZINE , mesalazine 3 g modified release granules, 30 sachets (<i>Salofalk</i>)	streamlined	restricted
10254Q	MESALAZINE , mesalazine 4 g modified release granules, 30 sachets (<i>Pentasa</i>)	streamlined	restricted
1611T	MESALAZINE , mesalazine 250 mg enteric tablet, 100 (<i>Mesasal</i>)	streamlined	restricted
2214M	MESALAZINE , mesalazine 500 mg modified release tablet, 100 (<i>Pentasa</i>)	streamlined	restricted
8731M	MESALAZINE , mesalazine 500 mg enteric tablet, 100 (<i>Salofalk</i>)	streamlined	restricted
11210B	MESALAZINE , mesalazine 800 mg enteric tablet, 90 (<i>Asacol</i>)	streamlined	authority-required
11554D	MESALAZINE , mesalazine 1 g enteric tablet, 60 (<i>Salofalk</i>)	streamlined	restricted
3413P	MESALAZINE , mesalazine 1 g modified release tablet, 60 (<i>Pentasa</i>)	streamlined	restricted
9353G	MESALAZINE , mesalazine 1.2 g modified release tablet, 60 (<i>Mezavant</i>)	streamlined	restricted
10682F	TRASTUZUMAB , trastuzumab 600 mg/5 mL injection, 5 mL vial (<i>Herceptin SC</i>)	authority-required	streamlined
10798H	TRASTUZUMAB , trastuzumab 600 mg/5 mL injection, 5 mL vial (<i>Herceptin SC</i>)	authority-required	streamlined
10803N	TRASTUZUMAB , trastuzumab 600 mg/5 mL injection, 5 mL vial (<i>Herceptin SC</i>)	authority-required	streamlined

Alteration – Manufacturer Code

		<i>From</i>	<i>To</i>
1884E	<i>Cilamox</i> – AMOXICILLIN , amoxicillin 250 mg capsule, 20	QA	AL
3301R	<i>Cilamox</i> – AMOXICILLIN , amoxicillin 250 mg capsule, 20	QA	AL
1889K	<i>Cilamox</i> – AMOXICILLIN , amoxicillin 500 mg capsule, 20	QA	AL
3300Q	<i>Cilamox</i> – AMOXICILLIN , amoxicillin 500 mg capsule, 20	QA	AL

2460L	<i>Aclor 125</i> – CEFACLOR , cefaclor 125 mg/5 mL powder for oral liquid, 100 mL	QA	MQ
2460L	<i>Ceclo</i> r – CEFACLOR , cefaclor 125 mg/5 mL powder for oral liquid, 100 mL	AS	AL
5046N	<i>Aclor 125</i> – CEFACLOR , cefaclor 125 mg/5 mL powder for oral liquid, 100 mL	QA	MQ
5046N	<i>Ceclo</i> r – CEFACLOR , cefaclor 125 mg/5 mL powder for oral liquid, 100 mL	AS	AL
2461M	<i>Aclor 250</i> – CEFACLOR , cefaclor 250 mg/5 mL powder for oral liquid, 75 mL	QA	MQ
2461M	<i>Ceclo</i> r – CEFACLOR , cefaclor 250 mg/5 mL powder for oral liquid, 75 mL	AS	AL
5047P	<i>Aclor 250</i> – CEFACLOR , cefaclor 250 mg/5 mL powder for oral liquid, 75 mL	QA	MQ
5047P	<i>Ceclo</i> r – CEFACLOR , cefaclor 250 mg/5 mL powder for oral liquid, 75 mL	AS	AL
1169M	<i>Ceclo</i> r CD – CEFACLOR , cefaclor 375 mg modified release tablet, 10	AS	AL
1169M	<i>Karlo</i> r CD – CEFACLOR , cefaclor 375 mg modified release tablet, 10	LN	MQ
5045M	<i>Ceclo</i> r CD – CEFACLOR , cefaclor 375 mg modified release tablet, 10	AS	AL
5045M	<i>Karlo</i> r CD – CEFACLOR , cefaclor 375 mg modified release tablet, 10	LN	MQ
2967E	<i>Questran Lite</i> – COLESTYRAMINE , colestyramine 4 g powder for oral liquid, 50 sachets	QA	GO
9249T	<i>Questran Lite</i> – COLESTYRAMINE , colestyramine 4 g powder for oral liquid, 50 sachets	QA	GO
1526H	<i>Flo</i> p <i>en</i> – FLUCLOXACILLIN , flucloxacillin 250 mg capsule, 24	AS	AL
5090X	<i>Flo</i> p <i>en</i> – FLUCLOXACILLIN , flucloxacillin 250 mg capsule, 24	AS	AL
10788T	<i>Flo</i> p <i>en</i> – FLUCLOXACILLIN , flucloxacillin 500 mg capsule, 24	AS	AL
1527J	<i>Flo</i> p <i>en</i> – FLUCLOXACILLIN , flucloxacillin 500 mg capsule, 24	AS	AL
5091Y	<i>Flo</i> p <i>en</i> – FLUCLOXACILLIN , flucloxacillin 500 mg capsule, 24	AS	AL
2546B	<i>Aldi</i> q – IMIQUIMOD , imiquimod 5% cream, 12 x 250 mg sachets	QA	AF
1611T	<i>Mesasa</i> l – MESALAZINE , mesalazine 250 mg enteric tablet, 100	AS	GO
10833E	<i>Advanta</i> n – METHYLPREDNISOLONE , methylprednisolone aceponate 0.1% cream, 15 g	BN	LO
10835G	<i>Advanta</i> n – METHYLPREDNISOLONE , methylprednisolone aceponate 0.1% cream, 15 g	BN	LO
10842P	<i>Advanta</i> n – METHYLPREDNISOLONE , methylprednisolone aceponate 0.1% cream, 15 g	BN	LO
10851D	<i>Advanta</i> n – METHYLPREDNISOLONE , methylprednisolone aceponate 0.1% cream, 15 g	BN	LO
10855H	<i>Advanta</i> n – METHYLPREDNISOLONE , methylprednisolone aceponate 0.1% cream, 15 g	BN	LO
8054X	<i>Advanta</i> n – METHYLPREDNISOLONE , methylprednisolone aceponate 0.1% cream, 15 g	BN	LO
10830B	<i>Advanta</i> n – METHYLPREDNISOLONE , methylprednisolone aceponate 0.1% lotion, 20 g	BN	LO
10838K	<i>Advanta</i> n – METHYLPREDNISOLONE , methylprednisolone aceponate 0.1% lotion, 20 g	BN	LO
10852E	<i>Advanta</i> n – METHYLPREDNISOLONE , methylprednisolone aceponate 0.1% lotion, 20 g	BN	LO
10856J	<i>Advanta</i> n – METHYLPREDNISOLONE , methylprednisolone aceponate 0.1% lotion, 20 g	BN	LO
8618N	<i>Advanta</i> n – METHYLPREDNISOLONE , methylprednisolone aceponate 0.1% lotion, 20 g	BN	LO
10839L	<i>Advanta</i> n – METHYLPREDNISOLONE , methylprednisolone aceponate 0.1% ointment: fatty, 15 g	BN	LO

10840M	<i>Advantan</i> – METHYLPREDNISOLONE , methylprednisolone aceponate 0.1% ointment: fatty, 15 g	BN	LO
10843Q	<i>Advantan</i> – METHYLPREDNISOLONE , methylprednisolone aceponate 0.1% ointment: fatty, 15 g	BN	LO
10844R	<i>Advantan</i> – METHYLPREDNISOLONE , methylprednisolone aceponate 0.1% ointment: fatty, 15 g	BN	LO
10848Y	<i>Advantan</i> – METHYLPREDNISOLONE , methylprednisolone aceponate 0.1% ointment: fatty, 15 g	BN	LO
8128T	<i>Advantan</i> – METHYLPREDNISOLONE , methylprednisolone aceponate 0.1% ointment: fatty, 15 g	BN	LO
10834F	<i>Advantan</i> – METHYLPREDNISOLONE , methylprednisolone aceponate 0.1% ointment, 15 g	BN	LO
10836H	<i>Advantan</i> – METHYLPREDNISOLONE , methylprednisolone aceponate 0.1% ointment, 15 g	BN	LO
10845T	<i>Advantan</i> – METHYLPREDNISOLONE , methylprednisolone aceponate 0.1% ointment, 15 g	BN	LO
10846W	<i>Advantan</i> – METHYLPREDNISOLONE , methylprednisolone aceponate 0.1% ointment, 15 g	BN	LO
10853F	<i>Advantan</i> – METHYLPREDNISOLONE , methylprednisolone aceponate 0.1% ointment, 15 g	BN	LO
8055Y	<i>Advantan</i> – METHYLPREDNISOLONE , methylprednisolone aceponate 0.1% ointment, 15 g	BN	LO
2622B	<i>Endone</i> – OXYCODONE , oxycodone hydrochloride 5 mg tablet, 20	QA	AF
2622B	<i>Oxycodone Aspen</i> – OXYCODONE , oxycodone hydrochloride 5 mg tablet, 20	FM	AL
5195K	<i>Endone</i> – OXYCODONE , oxycodone hydrochloride 5 mg tablet, 20	QA	AF
5195K	<i>Oxycodone Aspen</i> – OXYCODONE , oxycodone hydrochloride 5 mg tablet, 20	FM	AL
1215Y	<i>Codalgin Forte</i> – PARACETAMOL + CODEINE , paracetamol 500 mg + codeine phosphate hemihydrate 30 mg tablet, 20	FM	AF
3316M	<i>Codalgin Forte</i> – PARACETAMOL + CODEINE , paracetamol 500 mg + codeine phosphate hemihydrate 30 mg tablet, 20	FM	AF
8785J	<i>Codalgin Forte</i> – PARACETAMOL + CODEINE , paracetamol 500 mg + codeine phosphate hemihydrate 30 mg tablet, 20	FM	AF
1705R	<i>Cilicaine VK</i> – PHENOXYMETHYLPENICILLIN , phenoxymethylpenicillin 250 mg capsule, 50	FM	AF
1789E	<i>Cilicaine VK</i> – PHENOXYMETHYLPENICILLIN , phenoxymethylpenicillin 250 mg capsule, 50	FM	AF
3363B	<i>Cilicaine VK</i> – PHENOXYMETHYLPENICILLIN , phenoxymethylpenicillin 250 mg capsule, 50	FM	AF
2965C	<i>Cilicaine VK</i> – PHENOXYMETHYLPENICILLIN , phenoxymethylpenicillin 500 mg capsule, 50	FM	AF
3364C	<i>Cilicaine VK</i> – PHENOXYMETHYLPENICILLIN , phenoxymethylpenicillin 500 mg capsule, 50	FM	AF
1703P	<i>Aspecillin VK</i> – PHENOXYMETHYLPENICILLIN , phenoxymethylpenicillin 250 mg tablet, 25	QA	AF
1787C	<i>Aspecillin VK</i> – PHENOXYMETHYLPENICILLIN , phenoxymethylpenicillin 250 mg tablet, 25	QA	AF
3360W	<i>Aspecillin VK</i> – PHENOXYMETHYLPENICILLIN , phenoxymethylpenicillin 250 mg tablet, 25	QA	AF
3028J	<i>Aspecillin VK</i> – PHENOXYMETHYLPENICILLIN , phenoxymethylpenicillin 500 mg tablet, 25	QA	AF
3361X	<i>Aspecillin VK</i> – PHENOXYMETHYLPENICILLIN , phenoxymethylpenicillin 500 mg tablet, 25	QA	AF

1794K	<i>Cilicaine</i> – PROCAINE BENZYL PENICILLIN (PROCAINE PENICILLIN) , procaine benzylpenicillin (procaine penicillin) 1.5 g/3.4 mL injection, 5 x 3.4 mL syringes	QA	AF
3371K	<i>Cilicaine</i> – PROCAINE BENZYL PENICILLIN (PROCAINE PENICILLIN) , procaine benzylpenicillin (procaine penicillin) 1.5 g/3.4 mL injection, 5 x 3.4 mL syringes	QA	AF
2843P	<i>Coumadin</i> – WARFARIN , warfarin sodium 1 mg tablet, 50	QA	GO
2843P	<i>Marevan</i> – WARFARIN , warfarin sodium 1 mg tablet, 50	FM	GT
2209G	<i>Coumadin</i> – WARFARIN , warfarin sodium 2 mg tablet, 50	QA	GO
2844Q	<i>Marevan</i> – WARFARIN , warfarin sodium 3 mg tablet, 50	FM	GT
2211J	<i>Coumadin</i> – WARFARIN , warfarin sodium 5 mg tablet, 50	QA	GO
2211J	<i>Marevan</i> – WARFARIN , warfarin sodium 5 mg tablet, 50	FM	GT

Advance Notices

1 November 2019

Deletion – Brand

3116B	<i>Cal-500, PP</i> – CALCIUM , CALCIUM Tablet (chewable) 500 mg (as carbonate), 60
2413B	<i>Frusemide Sandoz, SZ</i> – FUROSEMIDE (FRUSEMIDE) , furosemide (frusemide) 20 mg/2 mL injection, 5 x 2 mL ampoules
10915L	<i>Glivanib, JU</i> – IMATINIB , imatinib 100 mg capsule, 60
10916M	<i>Glivanib, JU</i> – IMATINIB , imatinib 400 mg capsule, 30
10917N	<i>Glivanib, JU</i> – IMATINIB , imatinib 400 mg capsule, 30
10918P	<i>Glivanib, JU</i> – IMATINIB , imatinib 100 mg capsule, 60
10920R	<i>Glivanib, JU</i> – IMATINIB , imatinib 100 mg capsule, 60
10921T	<i>Glivanib, JU</i> – IMATINIB , imatinib 400 mg capsule, 30
10924Y	<i>Glivanib, JU</i> – IMATINIB , imatinib 100 mg capsule, 60
10925B	<i>Glivanib, JU</i> – IMATINIB , imatinib 400 mg capsule, 30
10933K	<i>Glivanib, JU</i> – IMATINIB , imatinib 400 mg capsule, 30
10935M	<i>Glivanib, JU</i> – IMATINIB , imatinib 400 mg capsule, 30
10939R	<i>Glivanib, JU</i> – IMATINIB , imatinib 400 mg capsule, 30
10940T	<i>Glivanib, JU</i> – IMATINIB , imatinib 100 mg capsule, 60
10941W	<i>Glivanib, JU</i> – IMATINIB , imatinib 100 mg capsule, 60
10942X	<i>Glivanib, JU</i> – IMATINIB , imatinib 100 mg capsule, 60
11756R	<i>Glivanib, JU</i> – IMATINIB , imatinib 400 mg capsule, 30
11757T	<i>Glivanib, JU</i> – IMATINIB , imatinib 100 mg capsule, 60
11763D	<i>Glivanib, JU</i> – IMATINIB , imatinib 400 mg capsule, 30
11764E	<i>Glivanib, JU</i> – IMATINIB , imatinib 400 mg capsule, 30
11770L	<i>Glivanib, JU</i> – IMATINIB , imatinib 100 mg capsule, 60
11771M	<i>Glivanib, JU</i> – IMATINIB , imatinib 400 mg capsule, 30
11772N	<i>Glivanib, JU</i> – IMATINIB , imatinib 400 mg capsule, 30
11776T	<i>Glivanib, JU</i> – IMATINIB , imatinib 100 mg capsule, 60
11777W	<i>Glivanib, JU</i> – IMATINIB , imatinib 100 mg capsule, 60
11779Y	<i>Glivanib, JU</i> – IMATINIB , imatinib 400 mg capsule, 30
11782D	<i>Glivanib, JU</i> – IMATINIB , imatinib 100 mg capsule, 60
11783E	<i>Glivanib, JU</i> – IMATINIB , imatinib 100 mg capsule, 60
5553G	<i>Latanoprost/timolol AN 50/5, JO</i> – LATANOPROST + TIMOLOL , latanoprost 0.005% + timolol 0.5% eye drops, 2.5 mL
5553G	<i>Lantim, JU</i> – LATANOPROST + TIMOLOL , latanoprost 0.005% + timolol 0.5% eye drops, 2.5 mL

8895E	<i>Latanoprost/timolol AN 50/5, JO</i> – LATANOPROST + TIMOLOL , latanoprost 0.005% + timolol 0.5% eye drops, 2.5 mL
8895E	<i>Lantim, JU</i> – LATANOPROST + TIMOLOL , latanoprost 0.005% + timolol 0.5% eye drops, 2.5 mL
11438B	<i>Carbzero, VF</i> – LONG CHAIN TRIGLYCERIDES , long chain triglycerides oral liquid, 15 x 225 mL bottles
11683X	<i>Meprazol, SZ</i> – OMEPRAZOLE , omeprazole 20 mg enteric tablet, 30
8331L	<i>Meprazol, SZ</i> – OMEPRAZOLE , omeprazole 20 mg enteric tablet, 30
8333N	<i>Meprazol, SZ</i> – OMEPRAZOLE , omeprazole 20 mg enteric tablet, 30
8378Y	<i>Temolide, JU</i> – TEMOZOLOMIDE , temozolomide 5 mg capsule, 5
8819E	<i>Temolide, JU</i> – TEMOZOLOMIDE , temozolomide 5 mg capsule, 5
9361Q	<i>Temolide, JU</i> – TEMOZOLOMIDE , temozolomide 140 mg capsule, 5
9362R	<i>Temolide, JU</i> – TEMOZOLOMIDE , temozolomide 140 mg capsule, 5
1925H	<i>Timoptol XE, MF</i> – TIMOLOL , timolol 0.25% eye drops, 2.5 mL
5549C	<i>Timoptol XE, MF</i> – TIMOLOL , timolol 0.25% eye drops, 2.5 mL
1465D	<i>Fasigyn, PF</i> – TINIDAZOLE , tinidazole 500 mg tablet, 4

1 December 2019

Deletion – Brand

1002R	<i>Zovirax, GK</i> – ACICLOVIR , aciclovir 3% eye ointment, 4.5 g
5501M	<i>Zovirax, GK</i> – ACICLOVIR , aciclovir 3% eye ointment, 4.5 g
10784N	<i>Bactrim DS, RO</i> – TRIMETHOPRIM + SULFAMETHOXAZOLE , trimethoprim 160 mg + sulfamethoxazole 800 mg tablet, 10
2951H	<i>Bactrim DS, RO</i> – TRIMETHOPRIM + SULFAMETHOXAZOLE , trimethoprim 160 mg + sulfamethoxazole 800 mg tablet, 10
3390K	<i>Bactrim DS, RO</i> – TRIMETHOPRIM + SULFAMETHOXAZOLE , trimethoprim 160 mg + sulfamethoxazole 800 mg tablet, 10

1 January 2020

Deletion – Brand

8362D	<i>Capecitabine Apotex, TX</i> – CAPECITABINE , capecitabine 500 mg tablet, 120

1 April 2020

Deletion – Brand

11210B	<i>Asacol, EU</i> – MESALAZINE , mesalazine 800 mg enteric tablet, 90

Highly Specialised Drugs Program (Private Hospital)

Additions

Addition – Item

11796W	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflixtra, Renflexis</i>)
11797X	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflixtra, Remicade, Renflexis</i>)
11810N	RITUXIMAB , rituximab 100 mg/10 mL injection, 2 x 10 mL vials (<i>Riximyo</i>)
11790M	RITUXIMAB , rituximab 500 mg/50 mL injection, 50 mL vial (<i>Riximyo</i>)
11804G	RITUXIMAB , rituximab 500 mg/50 mL injection, 50 mL vial (<i>Riximyo</i>)
11795T	TEDUGLUTIDE , teduglutide 5 mg injection [28 vials] (& inert substance diluent [28 x 0.5 mL syringes], 1 pack (<i>Revestive</i>))
11806J	TEDUGLUTIDE , teduglutide 5 mg injection [28 vials] (& inert substance diluent [28 x 0.5 mL syringes], 1 pack (<i>Revestive</i>))
11812Q	TEDUGLUTIDE , teduglutide 5 mg injection [28 vials] (& inert substance diluent [28 x 0.5 mL syringes], 1 pack (<i>Revestive</i>))

Addition – Brand

6291D	<i>Nivestim, PF</i> – FILGRASTIM , filgrastim 300 microgram/0.5 mL injection, 10 x 0.5 mL syringes
6292E	<i>Nivestim, PF</i> – FILGRASTIM , filgrastim 480 microgram/0.5 mL injection, 10 x 0.5 mL syringes

10583B *Riximyo, SZ – RITUXIMAB*, rituximab 100 mg/10 mL injection, 2 x 10 mL vials
 10576P *Riximyo, SZ – RITUXIMAB*, rituximab 500 mg/50 mL injection, 50 mL vial
 9611W *Riximyo, SZ – RITUXIMAB*, rituximab 500 mg/50 mL injection, 50 mL vial

Addition – Equivalence Indicator

2747N *Zarzio, SZ – FILGRASTIM*, filgrastim 300 microgram/0.5 mL injection, 5 x 0.5 mL syringes
 6291D *Neupogen, AN – FILGRASTIM*, filgrastim 300 microgram/0.5 mL injection, 10 x 0.5 mL syringes
 6126K *Neupogen, AN – FILGRASTIM*, filgrastim 300 microgram/mL injection, 10 x 1 mL vials
 2733W *Zarzio, SZ – FILGRASTIM*, filgrastim 480 microgram/0.5 mL injection, 5 x 0.5 mL syringes
 6292E *Neupogen, AN – FILGRASTIM*, filgrastim 480 microgram/0.5 mL injection, 10 x 0.5 mL syringes
 6127L *Neupogen, AN – FILGRASTIM*, filgrastim 480 microgram/1.6 mL injection, 10 x 1.6 mL vials
 10583B *Mabthera, RO – RITUXIMAB*, rituximab 100 mg/10 mL injection, 2 x 10 mL vials
 10576P *Mabthera, RO – RITUXIMAB*, rituximab 500 mg/50 mL injection, 50 mL vial
 9611W *Mabthera, RO – RITUXIMAB*, rituximab 500 mg/50 mL injection, 50 mL vial

Addition – Note

2747N **FILGRASTIM**, filgrastim 300 microgram/0.5 mL injection, 5 x 0.5 mL syringes (*Zarzio*)
 6291D **FILGRASTIM**, filgrastim 300 microgram/0.5 mL injection, 10 x 0.5 mL syringes (*Neupogen, Nivestim*)
 6126K **FILGRASTIM**, filgrastim 300 microgram/mL injection, 10 x 1 mL vials (*Neupogen*)
 2733W **FILGRASTIM**, filgrastim 480 microgram/0.5 mL injection, 5 x 0.5 mL syringes (*Zarzio*)
 6292E **FILGRASTIM**, filgrastim 480 microgram/0.5 mL injection, 10 x 0.5 mL syringes (*Neupogen, Nivestim*)
 6127L **FILGRASTIM**, filgrastim 480 microgram/1.6 mL injection, 10 x 1.6 mL vials (*Neupogen*)
 9611W **RITUXIMAB**, rituximab 500 mg/50 mL injection, 50 mL vial (*Mabthera, Riximyo*)

Deletions

Deletion – Item

9693E **FILGRASTIM**, filgrastim 300 microgram/0.5 mL injection, 10 x 0.5 mL syringes (*Nivestim*)
 9695G **FILGRASTIM**, filgrastim 480 microgram/0.5 mL injection, 10 x 0.5 mL syringes (*Nivestim*)
 11405G **ZOLEDRONIC ACID**, zoledronic acid 4 mg/5 mL injection, 5 x 5 mL vials (*Claris Lifesciences Zoledronic Acid*)

Deletion – Brand

9681M *Prograf XL, LL – TACROLIMUS*, tacrolimus 500 microgram modified release capsule, 30
 9682N *Prograf XL, LL – TACROLIMUS*, tacrolimus 1 mg modified release capsule, 60
 9683P *Prograf XL, LL – TACROLIMUS*, tacrolimus 5 mg modified release capsule, 30

Deletion – Equivalence Indicator

9681M *ADVAGRAF XL, LQ – TACROLIMUS*, tacrolimus 500 microgram modified release capsule, 30
 9682N *ADVAGRAF XL, LQ – TACROLIMUS*, tacrolimus 1 mg modified release capsule, 60
 9683P *ADVAGRAF XL, LQ – TACROLIMUS*, tacrolimus 5 mg modified release capsule, 30

Deletion – Note

6439X **PEGINTERFERON ALFA-2A**, peginterferon alfa-2a 135 microgram/0.5 mL injection, 4 x 0.5 mL syringes (*Pegasys*)
 11044G **PEGINTERFERON ALFA-2A**, peginterferon alfa-2a 180 microgram/0.5 mL injection, 4 x 0.5 mL syringes (*Pegasys*)
 6449K **PEGINTERFERON ALFA-2A**, peginterferon alfa-2a 180 microgram/0.5 mL injection, 4 x 0.5 mL syringes (*Pegasys*)

Deletion – Restriction

10184B **INFLIXIMAB**, infliximab 100 mg injection, 1 vial (*Inflixtra, Remicade, Renflexis*)

Alterations

Alteration – Note

9678J	ADALIMUMAB , adalimumab 20 mg/0.4 mL injection, 2 x 0.4 mL syringes (<i>Humira</i>)
9679K	ADALIMUMAB , adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL syringes (<i>Humira</i>)
9680L	ADALIMUMAB , adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL pen devices (<i>Humira</i>)
6367D	ETANERCEPT , etanercept 25 mg injection [4 vials] (&) inert substance diluent [4 x 1 mL syringes], 1 pack (<i>Enbrel</i>)
9641K	ETANERCEPT , etanercept 50 mg/mL injection, 4 x 1 mL pen devices (<i>Enbrel</i>)
9615C	ETANERCEPT , etanercept 50 mg/mL injection, 4 x 1 mL syringes (<i>Enbrel</i>)
10057H	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflixtra</i> , <i>Remicade</i> , <i>Renflexis</i>)
11396T	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflixtra</i> , <i>Renflexis</i>)
11432Q	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflixtra</i> , <i>Renflexis</i>)
11450P	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflixtra</i> , <i>Renflexis</i>)
11488P	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflixtra</i> , <i>Renflexis</i>)
11489Q	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Remicade</i>)
11515C	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflixtra</i> , <i>Renflexis</i>)
11595G	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflixtra</i> , <i>Renflexis</i>)
6448J	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflixtra</i> , <i>Remicade</i> , <i>Renflexis</i>)
10583B	RITUXIMAB , rituximab 100 mg/10 mL injection, 2 x 10 mL vials (<i>Mabthera</i> , <i>Riximyo</i>)
10576P	RITUXIMAB , rituximab 500 mg/50 mL injection, 50 mL vial (<i>Mabthera</i> , <i>Riximyo</i>)
10068X	TOCILIZUMAB , tocilizumab 80 mg/4 mL injection, 4 mL vial (<i>Actemra</i>)
10073E	TOCILIZUMAB , tocilizumab 80 mg/4 mL injection, 4 mL vial (<i>Actemra</i>)
1419Q	TOCILIZUMAB , tocilizumab 80 mg/4 mL injection, 4 mL vial (<i>Actemra</i>)
10071C	TOCILIZUMAB , tocilizumab 200 mg/10 mL injection, 10 mL vial (<i>Actemra</i>)
10079L	TOCILIZUMAB , tocilizumab 200 mg/10 mL injection, 10 mL vial (<i>Actemra</i>)
1423X	TOCILIZUMAB , tocilizumab 200 mg/10 mL injection, 10 mL vial (<i>Actemra</i>)
10060L	TOCILIZUMAB , tocilizumab 400 mg/20 mL injection, 20 mL vial (<i>Actemra</i>)
10078K	TOCILIZUMAB , tocilizumab 400 mg/20 mL injection, 20 mL vial (<i>Actemra</i>)
1464C	TOCILIZUMAB , tocilizumab 400 mg/20 mL injection, 20 mL vial (<i>Actemra</i>)

Alteration – Restriction

9678J	ADALIMUMAB , adalimumab 20 mg/0.4 mL injection, 2 x 0.4 mL syringes (<i>Humira</i>)
9679K	ADALIMUMAB , adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL syringes (<i>Humira</i>)
9680L	ADALIMUMAB , adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL pen devices (<i>Humira</i>)
10243D	ALEMTUZUMAB , alemtuzumab 12 mg/1.2 mL injection, 1.2 mL vial (<i>Lemtrada</i>)
10246G	ALEMTUZUMAB , alemtuzumab 12 mg/1.2 mL injection, 1.2 mL vial (<i>Lemtrada</i>)
9607P	APOMORPHINE , apomorphine hydrochloride hemihydrate 20 mg/2 mL injection, 5 x 2 mL ampoules (<i>Movapo</i>)
9640J	APOMORPHINE , apomorphine hydrochloride hemihydrate 50 mg/5 mL injection, 5 x 5 mL ampoules (<i>Movapo</i>)
11083H	APOMORPHINE , apomorphine hydrochloride hemihydrate 100 mg/20 mL injection, 5 x 20 mL vials (<i>Apomine Solution for Infusion</i>)
11475Y	APOMORPHINE , apomorphine hydrochloride hemihydrate 30 mg/3 mL injection, 5 x 3 mL pen devices (<i>Movapo Pen</i>)
11688E	APOMORPHINE , apomorphine hydrochloride hemihydrate 30 mg/3 mL injection, 5 x 3 mL cartridges (<i>Apomine Intermittent</i>)
10971K	APOMORPHINE , apomorphine hydrochloride hemihydrate 50 mg/10 mL injection, 5 x 10 mL syringes (<i>Movapo PFS</i>)
6221K	AZITHROMYCIN , azithromycin 600 mg tablet, 8 (<i>Zithromax</i>)
11128Q	BACLOFEN , baclofen 10 mg/5 mL intrathecal injection, 10 x 5 mL ampoules (<i>Sintetica Baclofen Intrathecal</i>)

11420C	BACLOFEN , baclofen 10 mg/5 mL intrathecal injection, 5 x 5 mL ampoules (<i>Bacthecal</i>)
6284R	BACLOFEN , baclofen 10 mg/5 mL intrathecal injection, 5 mL ampoule (<i>Bacthecal, Lioresal Intrathecal</i>)
11194E	BACLOFEN , baclofen 40 mg/20 mL intrathecal injection, 20 mL ampoule (<i>Sintetica Baclofen Intrathecal</i>)
11415T	CLOZAPINE , clozapine 50 mg/mL oral liquid, 100 mL (<i>Versacloz</i>)
9632Y	CLOZAPINE , clozapine 50 mg/mL oral liquid, 100 mL (<i>Clopine Suspension</i>)
6101D	CLOZAPINE , clozapine 25 mg tablet, 100 (<i>Clopine 25, Clozaril 25</i>)
6417R	CLOZAPINE , clozapine 50 mg tablet, 100 (<i>Clopine 50</i>)
6102E	CLOZAPINE , clozapine 100 mg tablet, 100 (<i>Clopine 100, Clozaril 100</i>)
6418T	CLOZAPINE , clozapine 200 mg tablet, 100 (<i>Clopine 200</i>)
11724C	DEFERIPRONE , deferiprone 1 g tablet, 50 (<i>Ferriprox</i>)
6120D	DORNASE ALFA , dornase alfa 2.5 mg/2.5 mL inhalation solution, 30 x 2.5 mL ampoules (<i>Pulmozyme</i>)
6367D	ETANERCEPT , etanercept 25 mg injection [4 vials] (&) inert substance diluent [4 x 1 mL syringes], 1 pack (<i>Enbrel</i>)
9641K	ETANERCEPT , etanercept 50 mg/mL injection, 4 x 1 mL pen devices (<i>Enbrel</i>)
9615C	ETANERCEPT , etanercept 50 mg/mL injection, 4 x 1 mL syringes (<i>Enbrel</i>)
6136Y	GANCICLOVIR , ganciclovir 500 mg injection, 5 vials (<i>Cymevene, GANCICLOVIR SXP</i>)
10057H	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflixtra, Remicade, Renflexis</i>)
11396T	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflixtra, Renflexis</i>)
11432Q	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflixtra, Renflexis</i>)
11450P	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflixtra, Renflexis</i>)
11483J	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflixtra, Renflexis</i>)
11488P	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflixtra, Renflexis</i>)
11489Q	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Remicade</i>)
11515C	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflixtra, Renflexis</i>)
11595G	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflixtra, Renflexis</i>)
6448J	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflixtra, Remicade, Renflexis</i>)
6148N	INTERFERON GAMMA-1B , interferon gamma-1b 2 million units (100 microgram)/0.5 mL injection, 6 x 0.5 mL vials (<i>Imukin</i>)
9744W	LEVODOPA + CARBIDOPA , levodopa 20 mg/mL + carbidopa monohydrate 5 mg/mL intestinal gel, 7 x 100 mL (<i>Duodopa</i>)
2008Q	MANNITOL , mannitol 40 mg powder for inhalation, 280 capsules (<i>bronchitol</i>)
9624M	NATALIZUMAB , natalizumab 300 mg/15 mL injection, 15 mL vial (<i>Tysabri</i>)
11237K	OCRELIZUMAB , ocrelizumab 300 mg/10 mL injection, 10 mL vial (<i>Ocrevus</i>)
6439X	PEGINTERFERON ALFA-2A , peginterferon alfa-2a 135 microgram/0.5 mL injection, 4 x 0.5 mL syringes (<i>Pegasys</i>)
6449K	PEGINTERFERON ALFA-2A , peginterferon alfa-2a 180 microgram/0.5 mL injection, 4 x 0.5 mL syringes (<i>Pegasys</i>)
6195C	RIFABUTIN , rifabutin 150 mg capsule, 30 (<i>Mycobutin</i>)
10583B	RITUXIMAB , rituximab 100 mg/10 mL injection, 2 x 10 mL vials (<i>Mabthera, Riximyo</i>)
10576P	RITUXIMAB , rituximab 500 mg/50 mL injection, 50 mL vial (<i>Mabthera, Riximyo</i>)
9611W	RITUXIMAB , rituximab 500 mg/50 mL injection, 50 mL vial (<i>Mabthera, Riximyo</i>)
10068X	TOCILIZUMAB , tocilizumab 80 mg/4 mL injection, 4 mL vial (<i>Actemra</i>)
10073E	TOCILIZUMAB , tocilizumab 80 mg/4 mL injection, 4 mL vial (<i>Actemra</i>)
1419Q	TOCILIZUMAB , tocilizumab 80 mg/4 mL injection, 4 mL vial (<i>Actemra</i>)
10071C	TOCILIZUMAB , tocilizumab 200 mg/10 mL injection, 10 mL vial (<i>Actemra</i>)
10079L	TOCILIZUMAB , tocilizumab 200 mg/10 mL injection, 10 mL vial (<i>Actemra</i>)
1423X	TOCILIZUMAB , tocilizumab 200 mg/10 mL injection, 10 mL vial (<i>Actemra</i>)

10060L	TOCILIZUMAB , tocilizumab 400 mg/20 mL injection, 20 mL vial (<i>Actemra</i>)
10078K	TOCILIZUMAB , tocilizumab 400 mg/20 mL injection, 20 mL vial (<i>Actemra</i>)
1464C	TOCILIZUMAB , tocilizumab 400 mg/20 mL injection, 20 mL vial (<i>Actemra</i>)

Alteration – Restriction Level

		<i>From</i>	<i>To</i>
10243D	ALEMTUZUMAB , alemtuzumab 12 mg/1.2 mL injection, 1.2 mL vial (<i>Lemtrada</i>)	authority-required	streamlined
10246G	ALEMTUZUMAB , alemtuzumab 12 mg/1.2 mL injection, 1.2 mL vial (<i>Lemtrada</i>)	authority-required	streamlined
9607P	APOMORPHINE , apomorphine hydrochloride hemihydrate 20 mg/2 mL injection, 5 x 2 mL ampoules (<i>Movapo</i>)	authority-required	streamlined
9640J	APOMORPHINE , apomorphine hydrochloride hemihydrate 50 mg/5 mL injection, 5 x 5 mL ampoules (<i>Movapo</i>)	authority-required	streamlined
11083H	APOMORPHINE , apomorphine hydrochloride hemihydrate 100 mg/20 mL injection, 5 x 20 mL vials (<i>Apomine Solution for Infusion</i>)	authority-required	streamlined
11475Y	APOMORPHINE , apomorphine hydrochloride hemihydrate 30 mg/3 mL injection, 5 x 3 mL pen devices (<i>Movapo Pen</i>)	authority-required	streamlined
11688E	APOMORPHINE , apomorphine hydrochloride hemihydrate 30 mg/3 mL injection, 5 x 3 mL cartridges (<i>Apomine Intermittent</i>)	authority-required	streamlined
10971K	APOMORPHINE , apomorphine hydrochloride hemihydrate 50 mg/10 mL injection, 5 x 10 mL syringes (<i>Movapo PFS</i>)	authority-required	streamlined
6221K	AZITHROMYCIN , azithromycin 600 mg tablet, 8 (<i>Zithromax</i>)	authority-required	streamlined
11128Q	BACLOFEN , baclofen 10 mg/5 mL intrathecal injection, 10 x 5 mL ampoules (<i>Sintetica Baclofen Intrathecal</i>)	authority-required	streamlined
11420C	BACLOFEN , baclofen 10 mg/5 mL intrathecal injection, 5 x 5 mL ampoules (<i>Bacethecal</i>)	authority-required	streamlined
6284R	BACLOFEN , baclofen 10 mg/5 mL intrathecal injection, 5 mL ampoule (<i>Bacethecal, Lioresal Intrathecal</i>)	authority-required	streamlined
11194E	BACLOFEN , baclofen 40 mg/20 mL intrathecal injection, 20 mL ampoule (<i>Sintetica Baclofen Intrathecal</i>)	authority-required	streamlined
11415T	CLOZAPINE , clozapine 50 mg/mL oral liquid, 100 mL (<i>Versacloz</i>)	authority-required	streamlined
9632Y	CLOZAPINE , clozapine 50 mg/mL oral liquid, 100 mL (<i>Clopine Suspension</i>)	authority-required	streamlined
6101D	CLOZAPINE , clozapine 25 mg tablet, 100 (<i>Clopine 25, Clozaril 25</i>)	authority-required	streamlined
6417R	CLOZAPINE , clozapine 50 mg tablet, 100 (<i>Clopine 50</i>)	authority-required	streamlined
6102E	CLOZAPINE , clozapine 100 mg tablet, 100 (<i>Clopine 100, Clozaril 100</i>)	authority-required	streamlined
6418T	CLOZAPINE , clozapine 200 mg tablet, 100 (<i>Clopine 200</i>)	authority-required	streamlined
11724C	DEFERIPRONE , deferiprone 1 g tablet, 50 (<i>Ferriprox</i>)	authority-required	streamlined
6120D	DORNASE ALFA , dornase alfa 2.5 mg/2.5 mL inhalation solution, 30 x 2.5 mL ampoules (<i>Pulmozyme</i>)	authority-required	streamlined
6136Y	GANCICLOVIR , ganciclovir 500 mg injection, 5 vials (<i>Cymevene, GANCICLOVIR SXP</i>)	authority-required	streamlined
10057H	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflectra, Remicade, Renflexis</i>)	authority-required	streamlined
11396T	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflectra, Renflexis</i>)	authority-required	streamlined
11432Q	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflectra, Renflexis</i>)	authority-required	streamlined
11450P	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflectra, Renflexis</i>)	authority-required	streamlined
11483J	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflectra, Renflexis</i>)	authority-required	streamlined
11488P	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflectra, Renflexis</i>)	authority-required	streamlined
11515C	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflectra, Renflexis</i>)	authority-required	streamlined

11595G	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflectra, Renflexis</i>)	authority-required	streamlined
6148N	INTERFERON GAMMA-1B , interferon gamma-1b 2 million units (100 microgram)/0.5 mL injection, 6 x 0.5 mL vials (<i>Imukin</i>)	authority-required	streamlined
9744W	LEVODOPA + CARBIDOPA , levodopa 20 mg/mL + carbidopa monohydrate 5 mg/mL intestinal gel, 7 x 100 mL (<i>Duodopa</i>)	authority-required	streamlined
2008Q	MANNITOL , mannitol 40 mg powder for inhalation, 280 capsules (<i>bronchitol</i>)	authority-required	streamlined
9624M	NATALIZUMAB , natalizumab 300 mg/15 mL injection, 15 mL vial (<i>Tysabri</i>)	authority-required	streamlined
11237K	OCRELIZUMAB , ocrelizumab 300 mg/10 mL injection, 10 mL vial (<i>Ocrevus</i>)	authority-required	streamlined
6439X	PEGINTERFERON ALFA-2A , peginterferon alfa-2a 135 microgram/0.5 mL injection, 4 x 0.5 mL syringes (<i>Pegasys</i>)	authority-required	streamlined
6449K	PEGINTERFERON ALFA-2A , peginterferon alfa-2a 180 microgram/0.5 mL injection, 4 x 0.5 mL syringes (<i>Pegasys</i>)	authority-required	streamlined
6195C	RIFABUTIN , rifabutin 150 mg capsule, 30 (<i>Mycobutin</i>)	authority-required	streamlined

Alteration – Maximum Quantity

		From	To
10057H	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflectra, Remicade, Renflexis</i>)	1	5
11396T	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflectra, Renflexis</i>)	1	5
11432Q	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflectra, Renflexis</i>)	1	5
11450P	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflectra, Renflexis</i>)	1	5
11488P	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflectra, Renflexis</i>)	1	5
11515C	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflectra, Renflexis</i>)	1	5
11595G	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflectra, Renflexis</i>)	1	5

Alteration – Number of Repeats

		From	To
11396T	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflectra, Renflexis</i>)	0	2
11432Q	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflectra, Renflexis</i>)	0	2
11450P	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflectra, Renflexis</i>)	0	2
11488P	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflectra, Renflexis</i>)	0	3
11515C	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflectra, Renflexis</i>)	0	2
11595G	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflectra, Renflexis</i>)	0	2

Advance Notices

1 November 2019

Deletion – Brand

6357N	<i>Valganciclovir AN, JO</i> – VALGANCICLOVIR , valganciclovir 450 mg tablet, 60
6357N	<i>Valganciclovir Juno, JU</i> – VALGANCICLOVIR , valganciclovir 450 mg tablet, 60

Highly Specialised Drugs Program (Public Hospital)

Additions

Addition – Item

11813R	RITUXIMAB , rituximab 100 mg/10 mL injection, 2 x 10 mL vials (<i>Riximyo</i>)
11800C	RITUXIMAB , rituximab 500 mg/50 mL injection, 50 mL vial (<i>Riximyo</i>)
11805H	RITUXIMAB , rituximab 500 mg/50 mL injection, 50 mL vial (<i>Riximyo</i>)
11793Q	TEDUGLUTIDE , teduglutide 5 mg injection [28 vials] (&) inert substance diluent [28 x 0.5 mL syringes], 1 pack (<i>Revestive</i>)
11794R	TEDUGLUTIDE , teduglutide 5 mg injection [28 vials] (&) inert substance diluent [28 x 0.5 mL syringes], 1 pack (<i>Revestive</i>)

11808L **TEDUGLUTIDE**, teduglutide 5 mg injection [28 vials] (&) inert substance diluent [28 x 0.5 mL syringes], 1 pack (*Revestive*)

Addition – Brand

5742F *Nivestim, PF* – **FILGRASTIM**, filgrastim 300 microgram/0.5 mL injection, 10 x 0.5 mL syringes
5744H *Nivestim, PF* – **FILGRASTIM**, filgrastim 480 microgram/0.5 mL injection, 10 x 0.5 mL syringes
11459D *Inflectra, PF* – **INFLIXIMAB**, infliximab 100 mg injection, 1 vial
11459D *Renflexis, MK* – **INFLIXIMAB**, infliximab 100 mg injection, 1 vial
10591K *Riximyo, SZ* – **RITUXIMAB**, rituximab 100 mg/10 mL injection, 2 x 10 mL vials
10593M *Riximyo, SZ* – **RITUXIMAB**, rituximab 500 mg/50 mL injection, 50 mL vial
9544H *Riximyo, SZ* – **RITUXIMAB**, rituximab 500 mg/50 mL injection, 50 mL vial

Addition – Equivalence Indicator

2758E *Zarzio, SZ* – **FILGRASTIM**, filgrastim 300 microgram/0.5 mL injection, 5 x 0.5 mL syringes
5742F *Neupogen, AN* – **FILGRASTIM**, filgrastim 300 microgram/0.5 mL injection, 10 x 0.5 mL syringes
5741E *Neupogen, AN* – **FILGRASTIM**, filgrastim 300 microgram/mL injection, 10 x 1 mL vials
2783L *Zarzio, SZ* – **FILGRASTIM**, filgrastim 480 microgram/0.5 mL injection, 5 x 0.5 mL syringes
5744H *Neupogen, AN* – **FILGRASTIM**, filgrastim 480 microgram/0.5 mL injection, 10 x 0.5 mL syringes
5743G *Neupogen, AN* – **FILGRASTIM**, filgrastim 480 microgram/1.6 mL injection, 10 x 1.6 mL vials
11459D *Remicade, JC* – **INFLIXIMAB**, infliximab 100 mg injection, 1 vial
10591K *Mabthera, RO* – **RITUXIMAB**, rituximab 100 mg/10 mL injection, 2 x 10 mL vials
10593M *Mabthera, RO* – **RITUXIMAB**, rituximab 500 mg/50 mL injection, 50 mL vial
9544H *Mabthera, RO* – **RITUXIMAB**, rituximab 500 mg/50 mL injection, 50 mL vial

Addition – Note

2758E **FILGRASTIM**, filgrastim 300 microgram/0.5 mL injection, 5 x 0.5 mL syringes (*Zarzio*)
5742F **FILGRASTIM**, filgrastim 300 microgram/0.5 mL injection, 10 x 0.5 mL syringes (*Neupogen, Nivestim*)
5741E **FILGRASTIM**, filgrastim 300 microgram/mL injection, 10 x 1 mL vials (*Neupogen*)
2783L **FILGRASTIM**, filgrastim 480 microgram/0.5 mL injection, 5 x 0.5 mL syringes (*Zarzio*)
5744H **FILGRASTIM**, filgrastim 480 microgram/0.5 mL injection, 10 x 0.5 mL syringes (*Neupogen, Nivestim*)
5743G **FILGRASTIM**, filgrastim 480 microgram/1.6 mL injection, 10 x 1.6 mL vials (*Neupogen*)
9544H **RITUXIMAB**, rituximab 500 mg/50 mL injection, 50 mL vial (*Mabthera, Riximyo*)

Deletions

Deletion – Item

9692D **FILGRASTIM**, filgrastim 300 microgram/0.5 mL injection, 10 x 0.5 mL syringes (*Nivestim*)
9694F **FILGRASTIM**, filgrastim 480 microgram/0.5 mL injection, 10 x 0.5 mL syringes (*Nivestim*)
11388J **ZOLEDRONIC ACID**, zoledronic acid 4 mg/5 mL injection, 5 x 5 mL vials (*Clarix Lifesciences Zoledronic Acid*)

Deletion – Brand

9664P *Prograf XL, LL* – **TACROLIMUS**, tacrolimus 500 microgram modified release capsule, 30
9665Q *Prograf XL, LL* – **TACROLIMUS**, tacrolimus 1 mg modified release capsule, 60
9666R *Prograf XL, LL* – **TACROLIMUS**, tacrolimus 5 mg modified release capsule, 30

Deletion – Equivalence Indicator

9664P *ADVAGRAF XL, LQ* – **TACROLIMUS**, tacrolimus 500 microgram modified release capsule, 30
9665Q *ADVAGRAF XL, LQ* – **TACROLIMUS**, tacrolimus 1 mg modified release capsule, 60
9666R *ADVAGRAF XL, LQ* – **TACROLIMUS**, tacrolimus 5 mg modified release capsule, 30

Deletion – Note

9515T	PEGINTERFERON ALFA-2A , peginterferon alfa-2a 135 microgram/0.5 mL injection, 4 x 0.5 mL syringes (<i>Pegasys</i>)
11026H	PEGINTERFERON ALFA-2A , peginterferon alfa-2a 180 microgram/0.5 mL injection, 4 x 0.5 mL syringes (<i>Pegasys</i>)
9516W	PEGINTERFERON ALFA-2A , peginterferon alfa-2a 180 microgram/0.5 mL injection, 4 x 0.5 mL syringes (<i>Pegasys</i>)

Alterations**Alteration – Note**

9661L	ADALIMUMAB , adalimumab 20 mg/0.4 mL injection, 2 x 0.4 mL syringes (<i>Humira</i>)
9662M	ADALIMUMAB , adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL syringes (<i>Humira</i>)
9663N	ADALIMUMAB , adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL pen devices (<i>Humira</i>)
5734T	ETANERCEPT , etanercept 25 mg injection [4 vials] (&) inert substance diluent [4 x 1 mL syringes], 1 pack (<i>Enbrel</i>)
5735W	ETANERCEPT , etanercept 50 mg/mL injection, 4 x 1 mL pen devices (<i>Enbrel</i>)
5733R	ETANERCEPT , etanercept 50 mg/mL injection, 4 x 1 mL syringes (<i>Enbrel</i>)
11482H	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Remicade</i>)
11486M	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflixtra</i> , <i>Renflexis</i>)
5753T	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflixtra</i> , <i>Remicade</i> , <i>Renflexis</i>)
10591K	RITUXIMAB , rituximab 100 mg/10 mL injection, 2 x 10 mL vials (<i>Mabthera</i> , <i>Riximyo</i>)
10593M	RITUXIMAB , rituximab 500 mg/50 mL injection, 50 mL vial (<i>Mabthera</i> , <i>Riximyo</i>)
10077J	TOCILIZUMAB , tocilizumab 80 mg/4 mL injection, 4 mL vial (<i>Actemra</i>)
10081N	TOCILIZUMAB , tocilizumab 80 mg/4 mL injection, 4 mL vial (<i>Actemra</i>)
1476Q	TOCILIZUMAB , tocilizumab 80 mg/4 mL injection, 4 mL vial (<i>Actemra</i>)
10056G	TOCILIZUMAB , tocilizumab 200 mg/10 mL injection, 10 mL vial (<i>Actemra</i>)
10058J	TOCILIZUMAB , tocilizumab 200 mg/10 mL injection, 10 mL vial (<i>Actemra</i>)
1481Y	TOCILIZUMAB , tocilizumab 200 mg/10 mL injection, 10 mL vial (<i>Actemra</i>)
10064Q	TOCILIZUMAB , tocilizumab 400 mg/20 mL injection, 20 mL vial (<i>Actemra</i>)
10072D	TOCILIZUMAB , tocilizumab 400 mg/20 mL injection, 20 mL vial (<i>Actemra</i>)
1482B	TOCILIZUMAB , tocilizumab 400 mg/20 mL injection, 20 mL vial (<i>Actemra</i>)

Alteration – Restriction

9661L	ADALIMUMAB , adalimumab 20 mg/0.4 mL injection, 2 x 0.4 mL syringes (<i>Humira</i>)
9662M	ADALIMUMAB , adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL syringes (<i>Humira</i>)
9663N	ADALIMUMAB , adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL pen devices (<i>Humira</i>)
5734T	ETANERCEPT , etanercept 25 mg injection [4 vials] (&) inert substance diluent [4 x 1 mL syringes], 1 pack (<i>Enbrel</i>)
5735W	ETANERCEPT , etanercept 50 mg/mL injection, 4 x 1 mL pen devices (<i>Enbrel</i>)
5733R	ETANERCEPT , etanercept 50 mg/mL injection, 4 x 1 mL syringes (<i>Enbrel</i>)
11482H	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Remicade</i>)
11486M	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflixtra</i> , <i>Renflexis</i>)
5753T	INFLIXIMAB , infliximab 100 mg injection, 1 vial (<i>Inflixtra</i> , <i>Remicade</i> , <i>Renflexis</i>)
10591K	RITUXIMAB , rituximab 100 mg/10 mL injection, 2 x 10 mL vials (<i>Mabthera</i> , <i>Riximyo</i>)
10593M	RITUXIMAB , rituximab 500 mg/50 mL injection, 50 mL vial (<i>Mabthera</i> , <i>Riximyo</i>)
9544H	RITUXIMAB , rituximab 500 mg/50 mL injection, 50 mL vial (<i>Mabthera</i> , <i>Riximyo</i>)
10077J	TOCILIZUMAB , tocilizumab 80 mg/4 mL injection, 4 mL vial (<i>Actemra</i>)
10081N	TOCILIZUMAB , tocilizumab 80 mg/4 mL injection, 4 mL vial (<i>Actemra</i>)
1476Q	TOCILIZUMAB , tocilizumab 80 mg/4 mL injection, 4 mL vial (<i>Actemra</i>)
10056G	TOCILIZUMAB , tocilizumab 200 mg/10 mL injection, 10 mL vial (<i>Actemra</i>)

10058J	TOCILIZUMAB , tocilizumab 200 mg/10 mL injection, 10 mL vial (<i>Actemra</i>)
1481Y	TOCILIZUMAB , tocilizumab 200 mg/10 mL injection, 10 mL vial (<i>Actemra</i>)
10064Q	TOCILIZUMAB , tocilizumab 400 mg/20 mL injection, 20 mL vial (<i>Actemra</i>)
10072D	TOCILIZUMAB , tocilizumab 400 mg/20 mL injection, 20 mL vial (<i>Actemra</i>)
1482B	TOCILIZUMAB , tocilizumab 400 mg/20 mL injection, 20 mL vial (<i>Actemra</i>)

Advance Notices

1 November 2019

Deletion – Brand

9569P	<i>Valganciclovir AN, JO</i> – VALGANCICLOVIR , valganciclovir 450 mg tablet, 60
9569P	<i>Valganciclovir Juno, JU</i> – VALGANCICLOVIR , valganciclovir 450 mg tablet, 60

Highly Specialised Drugs Program (Community Access)

Advance Notices

1 November 2019

Deletion – Brand

10306K	<i>Valganciclovir AN, JO</i> – VALGANCICLOVIR , valganciclovir 450 mg tablet, 60
10306K	<i>Valganciclovir Juno, JU</i> – VALGANCICLOVIR , valganciclovir 450 mg tablet, 60

1 January 2020

Deletion – Brand

10357D	<i>Abacavir/Lamivudine 600/300 APOTEX, TX</i> – ABACAVIR + LAMIVUDINE , abacavir 600 mg + lamivudine 300 mg tablet, 30
10304H	<i>Viramune, BY</i> – NEVIRAPINE , nevirapine 200 mg tablet, 60

1 March 2020

Deletion – Brand

10307L	<i>Stribild, GI</i> – TENOFOVIR + EMTRICITABINE + ELVITEGRAVIR + COBICISTAT , tenofovir disoproxil fumarate 300 mg + emtricitabine 200 mg + elvitegravir 150 mg + cobicistat 150 mg tablet, 30
10314W	<i>Eviplera, GI</i> – TENOFOVIR + EMTRICITABINE + RILPIVIRINE , tenofovir disoproxil fumarate 300 mg + emtricitabine 200 mg + rilpivirine 25 mg tablet, 30

Botulinum Toxin Program

Additions

Addition – Note

10982B	CLOSTRIDIUM BOTULINUM TYPE A TOXIN-HAEMAGGLUTININ COMPLEX , clostridium botulinum type A toxin-haemagglutinin complex 300 units injection, 1 vial (<i>Dysport</i>)
10988H	CLOSTRIDIUM BOTULINUM TYPE A TOXIN-HAEMAGGLUTININ COMPLEX , clostridium botulinum type A toxin-haemagglutinin complex 500 units injection, 1 vial (<i>Dysport</i>)

Alterations

Alteration – Restriction

10982B	CLOSTRIDIUM BOTULINUM TYPE A TOXIN-HAEMAGGLUTININ COMPLEX , clostridium botulinum type A toxin-haemagglutinin complex 300 units injection, 1 vial (<i>Dysport</i>)
10988H	CLOSTRIDIUM BOTULINUM TYPE A TOXIN-HAEMAGGLUTININ COMPLEX , clostridium botulinum type A toxin-haemagglutinin complex 500 units injection, 1 vial (<i>Dysport</i>)

Growth Hormone Program

Deletions

Deletion – Item

10440L	SOMATROPIN , somatropin 10 mg injection [1 vial] (&) inert substance diluent [1 mL syringe], 1 pack (<i>Zomacton</i>)
10455G	SOMATROPIN , somatropin 10 mg injection [1 vial] (&) inert substance diluent [1 mL syringe], 1 pack (<i>Zomacton</i>)
6310D	SOMATROPIN , somatropin 10 mg injection [1 vial] (&) inert substance diluent [1 mL syringe], 1 pack (<i>Zomacton</i>)

Opiate Dependence Treatment Program

Alterations

Alteration – Note

11759X	BUPRENORPHINE , buprenorphine 8 mg/0.16 mL modified release injection, 0.16 mL syringe (<i>Buvidal Weekly</i>)
11774Q	BUPRENORPHINE , buprenorphine 16 mg/0.32 mL modified release injection, 0.32 mL syringe (<i>Buvidal Weekly</i>)
11773P	BUPRENORPHINE , buprenorphine 24 mg/0.48 mL modified release injection, 0.48 mL syringe (<i>Buvidal Weekly</i>)
11766G	BUPRENORPHINE , buprenorphine 32 mg/0.64 mL modified release injection, 0.64 mL syringe (<i>Buvidal Weekly</i>)
11754P	BUPRENORPHINE , buprenorphine 64 mg/0.18 mL modified release injection, 0.18 mL syringe (<i>Buvidal Monthly</i>)
11767H	BUPRENORPHINE , buprenorphine 96 mg/0.27 mL modified release injection, 0.27 mL syringe (<i>Buvidal Monthly</i>)
11768J	BUPRENORPHINE , buprenorphine 128 mg/0.36 mL modified release injection, 0.36 mL syringe (<i>Buvidal Monthly</i>)

Repatriation Pharmaceutical Benefits

Deletions

Deletion – Brand

4592Q	<i>Gabapentin 300, CR</i> – GABAPENTIN , gabapentin 300 mg capsule, 100
4593R	<i>Gabapentin 400, CR</i> – GABAPENTIN , gabapentin 400 mg capsule, 100
4042R	<i>Calmurid, OL</i> – UREA , urea 10% cream, 100 g

Alterations

Alteration – Manufacturer Code

		<i>From</i>	<i>To</i>
4286N	<i>Aspalgin 40</i> – ASPIRIN + CODEINE , aspirin 300 mg + codeine phosphate hemihydrate 8 mg dispersible tablet, 40	QA	GO
4134N	<i>Aldiq</i> – IMIQUIMOD , imiquimod 5% cream, 12 x 250 mg sachets	QA	AF
4559Y	<i>Aldiq</i> – IMIQUIMOD , imiquimod 5% cream, 12 x 250 mg sachets	QA	AF

About the Schedule

The Schedule of Pharmaceutical Benefits lists all of the ready-prepared items subsidised under the Pharmaceutical Benefits Scheme (PBS).

The Schedule is published and is effective on the first day of each month.

For detailed information about the prescribing and supply of pharmaceutical benefits go to www.pbs.gov.au

For information about the operational aspects of the PBS, such as, PBS claiming, authority applications and stationery supplies contact the Department of Human Services at www.humanservices.gov.au

The Repatriation Schedule of Pharmaceutical Benefits provides information about pharmaceutical benefits available under the Repatriation Pharmaceutical Benefits Scheme (RPBS). These may only be prescribed to Department of Veterans' Affairs (DVA) beneficiaries holding a valid repatriation health card. Queries relating to the RPBS can be made to the DVA or go to www.dva.gov.au

Symbols and Abbreviations Used in the Schedule

*	An asterisk in the dispensed price column indicates that the manufacturer's pack does not coincide with the maximum quantity
‡	A double dagger in the maximum quantity column indicates where the maximum quantity has been determined to match the manufacturer's pack. These packs cannot be broken and the maximum quantity should be supplied and claimed
#	A gauge in the dispensed price column indicates that the product is not preconstituted and that the dispensed price therefore included a dispensing fee and where appropriate, an amount for purified water
a or b	Located immediately before brand names of an item indicates that the brands are equivalent for the purposes of substitution. These brands may be interchanged without differences in clinical effect
B	Located immediately before an amount in the premium column indicates a brand premium which applies to that particular brand of the item
T	Located immediately before an amount in the premium column indicates a therapeutic group premium which applies to that particular item
S	Located immediately before an amount in the premium column indicates a special patient contribution which applies to that particular item
DPMQ \$	Dispensed price for maximum quantity
MRVSN \$	Maximum recordable value for safety net
NP	Indicates that the item can be prescribed by an authorised nurse practitioner
MW	Indicates that the item can be prescribed by an authorised midwife
OP	Indicates that the item can be prescribed by an authorised optometrist
DP	Indicates that the item can be prescribed by an authorised dental practitioner

Restricted Benefits

All restricted items have separate headings for authority and non-authority items. In each case these items may be prescribed as pharmaceutical benefits only for use for one of the specified indications. Where more than one indication is specified for an Authority required or Restricted pharmaceutical benefit, each indication is separated from the preceding indication by a semi-colon and commences on the next line. In the case of Authority required (STREAMLINED) items, each indication will also include a four digit streamlined authority code. The drug may be prescribed as a pharmaceutical benefit for a patient who qualifies under any of the specified indications.

Restricted benefits - above an item indicates where an item can only be prescribed for specific therapeutic uses.

Authority required benefits – above an item indicates that a prescriber must seek approval from Department of Human Services or the Department of Veterans' Affairs. The prescriber must declare the specific conditions and circumstances that justify the use of these medicines. This is usually done by phone or in writing

Authority required (STREAMLINED) – authority can be sought electronically.

Guidelines and General Statements

General Statement for Drugs for the Treatment of Hepatitis C

Use the following criteria to determine patient eligibility for subsidisation under the PBS for hepatitis C treating agents.

By writing a PBS prescription, the prescriber is certifying the patient satisfies the qualifying criteria set out below and the use in accordance with the registered indications which differ between agents in this class – refer to the current Product Information for details.

Population criteria:

Patient must be aged 18 years or older.

Treatment criteria:

Must be treated by a medical practitioner or an authorised nurse practitioner¹ experienced in the treatment of chronic hepatitis C infection; or in consultation with a gastroenterologist, hepatologist or infectious diseases physician experienced in the treatment of chronic hepatitis C infection.

The following information must be provided at the time of application:

- the hepatitis C virus genotype; and
- the patient's cirrhotic status (non-cirrhotic or cirrhotic)

The following information must be documented in the patient's medical records:

- evidence of chronic hepatitis C infection (repeatedly antibody to hepatitis C virus (anti-HCV) positive and hepatitis C virus ribonucleic acid (HCV RNA) positive); and
- evidence of the hepatitis C virus genotype

The following matrices identify the regimens which are available for PBS prescription for eligible patients, based on the hepatitis C virus genotype and treatment history.

HEPATITIS C - NON-CIRRHOTIC PATIENTS

	TREATMENT NAÏVE	TREATMENT EXPERIENCED
Genotype 1	LEDIPASVIR + SOFOSBUVIR [8 or 12 weeks] ² OR DACLATASVIR and SOFOSBUVIR [12 weeks] OR SOFOSBUVIR and PEG-IFN and RBV [12 weeks] OR GRAZOPREVIR + ELBASVIR [12 weeks] OR SOFOSBUVIR + VELPATASVIR [12 weeks] OR GLECAPREVIR with PIBRENTASVIR [8 weeks]	LEDIPASVIR + SOFOSBUVIR [12 weeks] OR DACLATASVIR and SOFOSBUVIR [12 or 24 weeks] OR SOFOSBUVIR and PEG-IFN and RBV [12 weeks] OR GRAZOPREVIR + ELBASVIR [12 weeks] OR GRAZOPREVIR + ELBASVIR and RBV [16 weeks] ³ OR SOFOSBUVIR + VELPATASVIR [12 weeks] OR GLECAPREVIR + PIBRENTASVIR [8, 12 and 16 weeks] ⁴ OR SOFOSBUVIR + VELPATASVIR + VOXILAPREVIR [12 weeks] ⁵
Genotype 2	SOFOSBUVIR and RBV [12 weeks] OR SOFOSBUVIR + VELPATASVIR [12 weeks] OR GLECAPREVIR + PIBRENTASVIR [8 weeks]	SOFOSBUVIR and RBV [12 weeks] OR SOFOSBUVIR + VELPATASVIR [12 weeks] OR GLECAPREVIR + PIBRENTASVIR [8 weeks] OR SOFOSBUVIR + VELPATASVIR + VOXILAPREVIR [12 weeks] ⁵
Genotype 3	DACLATASVIR and SOFOSBUVIR [12 weeks]	DACLATASVIR and SOFOSBUVIR [12 weeks]

	OR SOFOSBUVIR and RBV [24 weeks] OR SOFOSBUVIR and PEG-IFN and RBV [12 weeks] OR SOFOSBUVIR + VELPATASVIR [12 weeks] OR GLECAPREVIR + PIBRENTASVIR [8 weeks]	OR SOFOSBUVIR and RBV [24 weeks] OR SOFOSBUVIR and PEG-IFN and RBV [12 weeks] OR SOFOSBUVIR + VELPATASVIR [12 weeks] OR GLECAPREVIR + PIBRENTASVIR [16 weeks] OR SOFOSBUVIR + VELPATASVIR + VOXILAPREVIR [12 weeks] ⁵
Genotype 4	SOFOSBUVIR and PEG-IFN and RBV [12 weeks] OR GRAZOPREVIR + ELBASVIR [12 weeks] OR SOFOSBUVIR + VELPATASVIR [12 weeks] OR GLECAPREVIR + PIBRENTASVIR [8 weeks]	SOFOSBUVIR and PEG-IFN and RBV [12 weeks] OR GRAZOPREVIR + ELBASVIR [12 weeks] OR GRAZOPREVIR + ELBASVIR and RBV [16 weeks] ³ OR SOFOSBUVIR + VELPATASVIR [12 weeks] OR GLECAPREVIR with PIBRENTASVIR [8 weeks] OR SOFOSBUVIR + VELPATASVIR + VOXILAPREVIR [12 weeks] ⁵
Genotype 5 & 6	SOFOSBUVIR and PEG-IFN and RBV [12 weeks] OR SOFOSBUVIR + VELPATASVIR [12 weeks] OR GLECAPREVIR + PIBRENTASVIR [8 weeks]	SOFOSBUVIR and PEG-IFN and RBV [12 weeks] OR SOFOSBUVIR + VELPATASVIR [12 weeks] OR GLECAPREVIR + PIBRENTASVIR [8 weeks] SOFOSBUVIR + VELPATASVIR + VOXILAPREVIR [12 weeks] ⁵

KEY

PEG-IFN - peginterferon alfa-2a

RBV - ribavirin

HEPATITIS C – CIRRHOTIC PATIENTS

	TREATMENT NAÏVE	TREATMENT EXPERIENCED
Genotype 1	LEDIPASVIR + SOFOSBUVIR [12 weeks] OR DACLATASVIR and SOFOSBUVIR and RBV [12 weeks] OR DACLATASVIR and SOFOSBUVIR [24 weeks] OR SOFOSBUVIR and PEG-IFN and RBV [12 weeks] OR GRAZOPREVIR + ELBASVIR [12 weeks] OR SOFOSBUVIR + VELPATASVIR [12 weeks] ⁶ OR GLECAPREVIR + PIBRENTASVIR [12 weeks]	LEDIPASVIR + SOFOSBUVIR [24 weeks] OR DACLATASVIR and SOFOSBUVIR [24 weeks] OR DACLATASVIR and SOFOSBUVIR and RBV [12 weeks] OR SOFOSBUVIR and PEG-IFN and RBV [12 weeks] OR GRAZOPREVIR + ELBASVIR [12 weeks] OR GRAZOPREVIR + ELBASVIR and RBV [16 weeks] ³ OR SOFOSBUVIR + VELPATASVIR [12 weeks] ⁶ OR GLECAPREVIR + PIBRENTASVIR [12 and 16 weeks] ⁷ OR SOFOSBUVIR + VELPATASVIR + VOXILAPREVIR [12 weeks] ⁵
Genotype 2	SOFOSBUVIR and RBV [12 weeks] OR SOFOSBUVIR + VELPATASVIR [12 weeks] ⁶ OR GLECAPREVIR with PIBRENTASVIR [12 weeks]	SOFOSBUVIR and RBV [12 weeks] OR SOFOSBUVIR + VELPATASVIR [12 weeks] ⁶ OR GLECAPREVIR + PIBRENTASVIR [12 weeks] OR SOFOSBUVIR + VELPATASVIR + VOXILAPREVIR [12 weeks] ⁵
Genotype 3	SOFOSBUVIR and RBV [24 weeks] OR DACLATASVIR and SOFOSBUVIR [24 weeks] OR SOFOSBUVIR and PEG-IFN and RBV [12 weeks]	DACLATASVIR and SOFOSBUVIR [24 weeks] OR SOFOSBUVIR and RBV [24 weeks] OR SOFOSBUVIR and PEG-IFN and RBV [12 weeks] OR

	OR DACLATASVIR and SOFOSBUVIR and RBV [12 or 24 weeks] ⁸ OR SOFOSBUVIR + VELPATASVIR [12 weeks] ^{6, 9} OR GLECAPREVIR + PIBRENTASVIR [12 weeks]	DACLATASVIR and SOFOSBUVIR and RBV [12 or 24 weeks] ⁸ OR SOFOSBUVIR + VELPATASVIR [12 weeks] ^{6, 9} OR GLECAPREVIR + PIBRENTASVIR [16 weeks] OR SOFOSBUVIR + VELPATASVIR + VOXILAPREVIR [12 weeks] ⁵
Genotype 4	SOFOSBUVIR and PEG-IFN and RBV [12 weeks] OR GRAZOPREVIR + ELBASVIR [12 weeks] OR SOFOSBUVIR + VELPATASVIR [12 weeks] ⁶ OR GLECAPREVIR + PIBRENTASVIR [12 weeks]	SOFOSBUVIR and PEG-IFN and RBV [12 weeks] OR GRAZOPREVIR + ELBASVIR [12 weeks] OR GRAZOPREVIR + ELBASVIR and RBV [16 weeks] ³ OR SOFOSBUVIR + VELPATASVIR [12 weeks] ⁶ OR GLECAPREVIR with PIBRENTASVIR [12 weeks] OR SOFOSBUVIR + VELPATASVIR + VOXILAPREVIR [12 weeks] ⁵
Genotype 5 & 6	SOFOSBUVIR and PEG-IFN and RBV [12 weeks] OR SOFOSBUVIR + VELPATASVIR [12 weeks] ⁶ OR GLECAPREVIR + PIBRENTASVIR [12 weeks]	SOFOSBUVIR and PEG-IFN and RBV [12 weeks] OR SOFOSBUVIR + VELPATASVIR [12 weeks] ⁶ OR GLECAPREVIR with PIBRENTASVIR [12 weeks] OR SOFOSBUVIR + VELPATASVIR + VOXILAPREVIR [12 weeks] ⁵

KEY

PEG-IFN - peginterferon alfa-2a
 RBV – ribavirin

¹. Medicines for the treatment of hepatitis C are listed for prescribing by authorised nurse practitioners under the General Schedule only. Medicines for the treatment of hepatitis C are not listed for prescribing by authorised nurse practitioners under the S100 Highly Specialised Drugs Program.

². LEDIPASVIR + SOFOSBUVIR [8 or 12 weeks] for treatment-naïve, non-cirrhotic patients:

- consider treatment for 8 weeks where pre-treatment HCV RNA is less than 6 million IU/mL;
- otherwise treatment for 12 weeks where pre-treatment HCV RNA is 6 million IU/mL or greater.

³. GRAZOPREVIR + ELBASVIR and RBV [16 weeks] for treatment-experienced, non-cirrhotic and cirrhotic patients, treatment for 16 weeks in patients with genotype 1a or 4 HCV who have experienced on-treatment virologic failure to prior treatment.

⁴. GLECAPREVIR + PIBRENTASVIR [8 or 12 or 16 weeks]:

- treatment for 8 weeks for treatment-experienced patients with genotype 1 who have failed regimens containing peginterferon, ribavirin, and/or sofosbuvir but no prior treatment experience with an HCV NS3/4A PI or NS5A inhibitor;
- treatment for 12 weeks for treatment-experienced patients with genotype 1 who have failed regimens containing an NS3/4A PI;
- treatment for 16 weeks for treatment-experienced patients with genotype 1 who have failed regimens containing an NS5A inhibitor.

⁵. SOFOSBUVIR + VELPATASVIR + VOXILAPREVIR [12 weeks] only for patients who have failed an NS5A inhibitor.

⁶. SOFOSBUVIR + VELPATASVIR [12 weeks] for patients with decompensated cirrhosis. Use in combination with ribavirin.

⁷. GLECAPREVIR + PIBRENTASVIR [12 or 16 weeks]:

- treatment for 12 weeks for treatment-experienced patients with genotype 1 who have failed regimens containing an NS3/4A PI;
- treatment for 16 weeks for treatment-experienced patients with genotype 1 who have failed regimens containing an NS5A inhibitor.

⁸. DACLATASVIR and SOFOSBUVIR and RBV [12 or 24 weeks] for cirrhotic patients. Consider a 24 week regimen of where clinically appropriate.

⁹. SOFOSBUVIR + VELPATASVIR [12 weeks] for patients with genotype 3 infection with compensated cirrhosis. Consider addition of ribavirin.

Pharmaceutical Benefits Schedules

Highly Specialised Drugs Program (Private Hospital)

ALIMENTARY TRACT AND METABOLISM.....	30
OTHER ALIMENTARY TRACT AND METABOLISM PRODUCTS.....	30
OTHER ALIMENTARY TRACT AND METABOLISM PRODUCTS	30
BLOOD AND BLOOD FORMING ORGANS	33
ANTIHEMORRHAGICS.....	33
VITAMIN K AND OTHER HEMOSTATICS	33
ANTIANEMIC PREPARATIONS	39
OTHER ANTIANEMIC PREPARATIONS	39
CARDIOVASCULAR SYSTEM.....	42
ANTIHYPERTENSIVES	42
OTHER ANTIHYPERTENSIVES	42
SYSTEMIC HORMONAL PREPARATIONS, EXCL. SEX HORMONES AND INSULINS.....	90
PITUITARY AND HYPOTHALAMIC HORMONES AND ANALOGUES.....	90
ANTERIOR PITUITARY LOBE HORMONES AND ANALOGUES	90
HYPOTHALAMIC HORMONES.....	92
ANTIINFECTIVES FOR SYSTEMIC USE	96
ANTIBACTERIALS FOR SYSTEMIC USE.....	96
MACROLIDES, LINCOSAMIDES AND STREPTOGRAMINS.....	96
ANTIMYCOBACTERIALS	96
DRUGS FOR TREATMENT OF TUBERCULOSIS.....	96
ANTIVIRALS FOR SYSTEMIC USE	97
DIRECT ACTING ANTIVIRALS	97
ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS	102
ANTINEOPLASTIC AGENTS.....	102
ANTIMETABOLITES.....	102
CYTOTOXIC ANTIBIOTICS AND RELATED SUBSTANCES	103
OTHER ANTINEOPLASTIC AGENTS.....	104
IMMUNOSTIMULANTS	118
IMMUNOSTIMULANTS	118
IMMUNOSUPPRESSANTS.....	125
IMMUNOSUPPRESSANTS	125

MUSCULO-SKELETAL SYSTEM.....	275
MUSCLE RELAXANTS	275
MUSCLE RELAXANTS, CENTRALLY ACTING AGENTS	275
DRUGS FOR TREATMENT OF BONE DISEASES	277
DRUGS AFFECTING BONE STRUCTURE AND MINERALIZATION.....	277
OTHER DRUGS FOR DISORDERS OF THE MUSCULO-SKELETAL SYSTEM	278
OTHER DRUGS FOR DISORDERS OF THE MUSCULO-SKELETAL SYSTEM	278
NERVOUS SYSTEM.....	281
ANTI-PARKINSON DRUGS	281
DOPAMINERGIC AGENTS	281
PSYCHOLEPTICS.....	282
ANTIPSYCHOTICS.....	282
RESPIRATORY SYSTEM.....	283
DRUGS FOR OBSTRUCTIVE AIRWAY DISEASES	283
OTHER SYSTEMIC DRUGS FOR OBSTRUCTIVE AIRWAY DISEASES	283
COUGH AND COLD PREPARATIONS.....	300
EXPECTORANTS, EXCL. COMBINATIONS WITH COUGH SUPPRESSANTS	300
OTHER RESPIRATORY SYSTEM PRODUCTS	301
OTHER RESPIRATORY SYSTEM PRODUCTS	301
VARIOUS	305
ALL OTHER THERAPEUTIC PRODUCTS	305
ALL OTHER THERAPEUTIC PRODUCTS.....	305

ALIMENTARY TRACT AND METABOLISM

OTHER ALIMENTARY TRACT AND METABOLISM PRODUCTS

OTHER ALIMENTARY TRACT AND METABOLISM PRODUCTS

Various alimentary tract and metabolism products

■ TEDUGLUTIDE

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note A patient may only qualify for PBS-subsidised treatment under this restriction once in a lifetime.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Type III Short bowel syndrome with intestinal failure

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a gastroenterologist; OR
- Must be treated by a specialist within a multidisciplinary intestinal rehabilitation unit.

Clinical criteria:

- Patient must have short bowel syndrome with intestinal failure following major surgery, **AND**
- Patient must have a history of dependence on parenteral support for at least 12 months, **AND**
- Patient must have received a stable parenteral support regimen for at least 3 days per week in the previous 4 weeks, **AND**
- Patient must not have active gastrointestinal malignancy or history of gastrointestinal malignancy within the last 5 years, **AND**
- The treatment must not exceed 12 months under this restriction, **AND**
- Patient must not have previously received PBS-subsidised treatment with this drug for this condition.

Baseline is the mean number of days of parenteral support per week over the four weeks immediately prior to initiating treatment with teduglutide under the PBS initial treatment restriction or four weeks immediately prior to initiating treatment with non-PBS subsidised teduglutide for grandfathered patients.

A stable parenteral support regimen is defined as a minimum of 3 days of parenteral support (parenteral nutrition with or without IV fluids) per week for 4 consecutive weeks to meet caloric, fluid or electrolyte needs.

Baseline number of days of parenteral support should be documented in the patient's medical records.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Short bowel syndrome with intestinal failure form; and
- (3) details of baseline mean number of days on parenteral support per week for 4 consecutive weeks immediately preceding this application; and
- (4) documented duration in months of prior dependence on parenteral support.

teduglutide 5 mg injection [28 vials] (&) inert substance diluent [28 x 0.5 mL syringes], 1 pack

11795T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	11	..	21887.39	Revestive [ZI]

■ TEDUGLUTIDE

Note Special Pricing Arrangements apply.

Authority required

Type III Short bowel syndrome with intestinal failure

Treatment Phase: Initial treatment - Grandfathered patients

Treatment criteria:

- Must be treated by a gastroenterologist; OR
- Must be treated by a specialist within a multidisciplinary intestinal rehabilitation unit.

Clinical criteria:

- Patient must have previously received non-PBS subsidised treatment with this drug for this condition prior to 1 October 2019, **AND**
- Patient must have short bowel syndrome with intestinal failure following major surgery, **AND**
- Patient must have had a history of dependence on parenteral support for at least 12 months prior to initiating non-PBS subsidised treatment with this drug for this condition, **AND**

- Patient must have received a stable parenteral support regimen for at least 3 days per week in the 4 weeks prior to initiating non-PBS subsidised treatment with this drug for this condition, **AND**
- Patient must not have active gastrointestinal malignancy or history of gastrointestinal malignancy within the last 5 years, **AND**
- Patient must have achieved a treatment response if the patient has been on non-PBS subsidised therapy with this drug for more than 12 months.

Baseline is the mean number of days of parenteral support per week over the 4 weeks immediately prior to initiating treatment with non-PBS subsidised teduglutide for grandfathered patients.

A stable parenteral support regimen is defined as a minimum of 3 days of parenteral support (parenteral nutrition with or without IV fluids) per week for 4 consecutive weeks to meet caloric, fluid or electrolyte needs.

A patient has met the criteria for treatment response when there is a reduction in the mean number of days of parenteral support of at least 1 day per week since initiating non-PBS subsidised treatment, or where a patient has completely ceased treatment with parenteral support for a period of at least 4 consecutive weeks prior to application for PBS-subsidised treatment.

The number of days of parenteral support is calculated as the mean number of days in which any parenteral support is required (parenteral nutrition with or without IV fluids) per week to meet caloric, fluid or electrolyte needs between commencement of non-PBS subsidised teduglutide and application for PBS-subsidised treatment.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Short bowel syndrome with intestinal failure Grandfather PBS Authority Application - Supporting Information Form; and
- (3) details of non-PBS subsidised teduglutide treatment start date; and
- (4) details of the number of days on parenteral support per week for 4 consecutive weeks prior to initiating non-PBS subsidised therapy; and
- (5) documented duration in months of dependence on parenteral support prior to initiating non-PBS subsidised treatment; and
- (6) details of response to teduglutide treatment if patient has received 12 or more months of non-PBS subsidised treatment.

A patient may qualify for PBS-subsidised treatment under this restriction once only.

For patients who have been on this drug for less than 12 months, the maximum number of repeats that will be approved will be for an amount equivalent to an initial 12 month supply of PBS and non-PBS subsidised treatment.

For patients who have been on this drug for more than 12 months, a maximum of 5 repeats will be approved.

For continuing PBS-subsidised treatment, a Grandfathered patient must qualify under the First continuing treatment criteria.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Type III Short bowel syndrome with intestinal failure

Treatment Phase: Initial treatment or initial grandfather treatment - balance of supply

Treatment criteria:

- Must be treated by a gastroenterologist; OR
- Must be treated by a specialist within a multidisciplinary intestinal rehabilitation unit.

Clinical criteria:

- Patient must have previously received PBS-subsidised initial treatment with this drug for this condition; OR
- Patient must have received PBS-subsidised treatment with this drug for this condition as a grandfathered patient, **AND**
- Patient must have received insufficient therapy with this drug under the initial or grandfather treatment restriction to complete the maximum duration of 12 months of initial treatment, **AND**
- The treatment must provide no more than the balance of up to 12 months of treatment.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

teduglutide 5 mg injection [28 vials] (&) inert substance diluent [28 x 0.5 mL syringes], 1 pack

11812Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	21887.39	Revestive [ZI]

▪ TEDUGLUTIDE

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Type III Short bowel syndrome with intestinal failure

Treatment Phase: First continuing treatment

Treatment criteria:

- Must be treated by a gastroenterologist; OR
- Must be treated by a specialist within a multidisciplinary intestinal rehabilitation unit.

Clinical criteria:

- Patient must have previously received PBS-subsidised initial treatment with this drug for this condition; OR
- Patient must have received PBS-subsidised treatment with this drug for this condition as a grandfathered patient, **AND**
- Patient must have a reduction in parenteral support frequency of at least one day per week compared to the mean number of days per week at baseline.

Baseline is the mean number of days of parenteral support per week over the four weeks immediately prior to initiating treatment with teduglutide under the PBS initial treatment restriction or four weeks immediately prior to initiating treatment with non-PBS subsidised teduglutide for grandfathered patients.

The current mean number of days of parenteral support is calculated as the mean number of days in which any parenteral support is required (parenteral nutrition with or without IV fluids) per week to meet caloric, fluid or electrolyte needs over the immediately preceding 4 week treatment period

Treatment failure

For applications for first continuing treatment, failure of treatment is defined as no change compared to baseline in the mean number of days per week in parenteral support (parenteral nutrition with or without IV fluids) to meet caloric, fluid or electrolyte needs.

Patients who experience failure of treatment must permanently discontinue treatment.

Current mean number of days of parenteral support should be documented in the patient's medical records.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Short bowel syndrome with intestinal failure Form; and
- (3) details of the baseline mean number of days reduction of parenteral support (parenteral nutrition with or without IV fluids) per week to meet caloric, fluid or electrolyte needs; and
- (4) the current mean number of days per week of parenteral support over the preceding 4 week period.

Note A patient may only qualify for PBS-subsidised treatment under this restriction once in a lifetime.

Authority required

Type III Short bowel syndrome with intestinal failure

Treatment Phase: Subsequent continuing treatment

Treatment criteria:

- Must be treated by a gastroenterologist; OR
- Must be treated by a specialist within a multidisciplinary intestinal rehabilitation unit.

Clinical criteria:

- Patient must have received PBS-subsidised first-continuing treatment with this drug for this condition; OR
- Patient must have received PBS-subsidised recommencement of treatment with this drug for this condition following a trial cessation period, **AND**
- Patient must have achieved a treatment response in the preceding treatment period; OR
- Patient must not have previously experienced a failure to respond to treatment with this drug for this condition.

Treatment response

For applications for subsequent continuing treatment, treatment response is when there was a reduction in the mean number of days of parenteral support of at least 1 day per week since the last assessment for PBS-subsidised treatment, OR where a patient has completely ceased treatment with parenteral support for a period of at least 4 consecutive weeks.

The current mean number of days of parenteral support is calculated as the mean number of days in which any parenteral support is required (parenteral nutrition with or without IV fluids) per week to meet caloric, fluid or electrolyte needs over the immediately preceding 4 week treatment period

Treatment failure

For applications for subsequent continuing treatment, failure of treatment is defined as an increase in the mean number of days per week of parenteral support requirements of at least 1 day per week over the preceding 4 week period compared to the last assessment for PBS-subsidised treatment of parenteral support (parenteral nutrition with or without IV fluids) to meet caloric, fluid or electrolyte needs.

Patients who experience failure of treatment must permanently discontinue treatment.

Treatment stability

Patients who neither demonstrate a treatment response nor a treatment failure since the last assessment for PBS-subsidised treatment are considered to have a stable parenteral support regimen, defined as the same mean number of days of parenteral support (parenteral nutrition with or without IV fluids) per week to meet caloric, fluid or electrolyte needs over the 4 weeks preceding treatment period, where the number of days is greater than zero and the mean number of days of parenteral support is less than baseline. Patients with a stable parenteral support regimen over 6 months must undertake a trial cessation period. Patients who have re-commenced after a trial cessation period are exempt from further trial cessation.

Trial cessation period

Patients who demonstrate a stable frequency of mean days per week of parenteral support in a 6-month period commencing after the initial 12 months of treatment with this drug for this condition are required to undertake a trial of treatment cessation. Patients who have re-commenced after a trial cessation period are exempt from further trial cessation.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Short bowel syndrome with intestinal failure Form; and
- (3) details of the mean number of days reduction of parenteral support (parenteral nutrition with or without IV fluids) per week to meet caloric, fluid or electrolyte needs over the preceding treatment period **or** confirmation the patient has had 4 consecutive weeks without parenteral support; and
- (4) details of completion of a trial cessation period (if applicable); and
- (5) the current mean number of days per week of parenteral support over the preceding 4 week period.

Authority required

Type III Short bowel syndrome with intestinal failure

Treatment Phase: Recommencement of treatment

Treatment criteria:

- Must be treated by a gastroenterologist; **OR**
- Must be treated by a specialist within a multidisciplinary intestinal rehabilitation unit.

Clinical criteria:

- Patient must have received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have undertaken a trial cessation period due to experiencing a stable parenteral support regimen in the first continuing or subsequent continuing treatment phase, and not due to a treatment failure, **AND**
- Patient must have experienced deterioration during a trial cessation period.

Trial cessation period

Patients who demonstrate a stable frequency of mean days per week of parenteral support in a 6-month period commencing after the initial 12 months of treatment with this drug for this condition are required to undertake a trial of treatment cessation. Patients who have re-commenced after a trial cessation period are exempt from further trial cessation.

Deterioration during the trial cessation period includes an increase in parenteral support frequency of more than or equal to one day per week from the pre-cessation level, or other clinical parameters suggestive of deterioration including changes in renal function or urinary sodium levels or changes in body weight.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Short bowel syndrome with intestinal failure Form; and
- (3) details of the reason for recommencement after trial cessation; and
- (4) the current mean number of days per week of parenteral support over the preceding 4 week period

teduglutide 5 mg injection [28 vials] (&) inert substance diluent [28 x 0.5 mL syringes], 1 pack

11806J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	21887.39	Revestive [ZI]

■ BLOOD AND BLOOD FORMING ORGANS

■ ANTIHEMORRHAGICS

VITAMIN K AND OTHER HEMOSTATICS

Other systemic hemostatics

■ ELTROMBOPAG

Note Special Pricing Arrangements apply.

Note No applications for increased repeats will be authorised.

Authority required

Severe thrombocytopenia

Treatment Phase: Initial treatment 1 - New patient

Clinical criteria:

- The condition must be severe chronic immune (idiopathic) thrombocytopenic purpura (ITP), **AND**
- Patient must have had a splenectomy, **AND**
- Patient must have failed to achieve an adequate response to, or be intolerant to, corticosteroid therapy following the splenectomy, **AND**
- Patient must have failed to achieve an adequate response to, or be intolerant to, immunoglobulin therapy following the splenectomy, **AND**
- The treatment must be the sole PBS-subsidised thrombopoietin receptor agonist (TRA) for this condition.

Population criteria:

- Patient must be an adult.

The following criteria indicate failure to achieve an adequate response and must be demonstrated at the time of initial application;

- (a) a platelet count of less than or equal to 20,000 million per L; **OR**
- (b) a platelet count of 20,000 million to 30,000 million per L, where the patient is experiencing significant bleeding or has a history of significant bleeding in this platelet range.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form,
- (2) a signed patient acknowledgement,
- (3) a completed Idiopathic Thrombocytopenic Purpura Initial PBS Authority Application - Supporting Information Form,
- (4) a copy of a full blood count pathology report supporting the diagnosis of ITP, and
- (5) where the application is sought on the basis of a medical contraindication to surgery, a signed and dated letter from the clinician making this assessment which includes the date upon which the patient was assessed for surgery and the clinical grounds upon which surgery is contraindicated.

The full blood count must be no more than 1 month old at the time of application.

A maximum of 24 weeks of treatment with this drug will be authorised under this criterion.

Note Eltrombopag is not PBS-subsidised as an alternative to splenectomy.

Note Patients will be able to trial either eltrombopag or romiplostim within the initial 24 weeks treatment period. Patients who fail to demonstrate a response to treatment with eltrombopag and/or romiplostim under the initial restriction will not be eligible to receive further PBS-subsidised treatment with either of these drugs.

Note Authority approval for sufficient therapy to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written application for authority approval for sufficient therapy to complete a maximum of 24 weeks of treatment should be forwarded to:

Department of Human Services
Prior Written Approval of Complex Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001

Authority required

Severe thrombocytopenia

Treatment Phase: Initial treatment 2 - New patient

Clinical criteria:

- The condition must be severe chronic immune (idiopathic) thrombocytopenic purpura (ITP), **AND**
- Patient must not have had a splenectomy, **AND**
- Patient must have failed to achieve an adequate response to, or be intolerant to, corticosteroid therapy at a dose equivalent to 0.5-2 mg/kg/day of prednisone for at least 4-6 weeks, **AND**
- Patient must have failed to achieve an adequate response to, or be intolerant to, immunoglobulin therapy, **AND**
- Patient must be unsuitable for splenectomy due to medical reasons, **AND**
- The treatment must be the sole PBS-subsidised thrombopoietin receptor agonist (TRA) for this condition.

Population criteria:

- Patient must be an adult.

The following criteria indicate failure to achieve an adequate response and must be demonstrated at the time of initial application;

- (a) a platelet count of less than or equal to 20,000 million per L; OR
- (b) a platelet count of 20,000 million to 30,000 million per L, where the patient is experiencing significant bleeding or has a history of significant bleeding in this platelet range.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form,
- (2) a signed patient acknowledgement,
- (3) a completed Idiopathic Thrombocytopenic Purpura Initial PBS Authority Application - Supporting Information Form,
- (4) a copy of a full blood count pathology report supporting the diagnosis of ITP, and
- (5) where the application is sought on the basis of a medical contraindication to surgery, a signed and dated letter from the clinician making this assessment which includes the date upon which the patient was assessed for surgery and the clinical grounds upon which surgery is contraindicated.

The full blood count must be no more than 1 month old at the time of application.

A maximum of 24 weeks of treatment with this drug will be authorised under this criterion.

Note Eltrombopag is not PBS-subsidised as an alternative to splenectomy.

Note Patients will be able to trial either eltrombopag or romiplostim within the initial 24 weeks treatment period. Patients who fail to demonstrate a response to treatment with eltrombopag and/or romiplostim under the initial restriction will not be eligible to receive further PBS-subsidised treatment with either of these drugs.

Note Authority approval for sufficient therapy to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written application for authority approval for sufficient therapy to complete a maximum of 24 weeks of treatment should be forwarded to:

Department of Human Services
Prior Written Approval of Complex Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001

Authority required

Severe thrombocytopenia

Treatment Phase: First Continuing treatment or Re-initiation of interrupted treatment

Clinical criteria:

- The condition must be severe chronic immune (idiopathic) thrombocytopenic purpura (ITP), **AND**
- Patient must have previously received PBS-subsidised initial treatment with this drug for this condition, **AND**
- Patient must have demonstrated a sustained platelet response to PBS-subsidised treatment with this drug for this condition under the Initial treatment restriction, **AND**
- The treatment must be the sole PBS-subsidised thrombopoietin receptor agonist (TRA) for this condition.

Population criteria:

- Patient must be an adult.

For the purposes of this restriction, a sustained platelet response is defined as:

(a) use of rescue medication (corticosteroids or immunoglobulins) on no more than one occasion during the initial period of PBS-subsidised treatment with this drug,

AND either of the following:

(b) a platelet count greater than or equal to 50,000 million per L on at least four (4) occasions, each at least one week apart;
OR

(c) a platelet count greater than 30,000 million per L and which is double the baseline (pre-treatment) platelet count on at least four (4) occasions, each at least one week apart.

Applications for the First continuing PBS-subsidised treatment or Re-initiation of interrupted PBS-subsidised treatment must be made in writing and must include:

- (1) a completed authority prescription form, and
- (2) a completed Idiopathic Thrombocytopenic Purpura Continuing PBS Authority Application - Supporting Information Form , and

(3) copies of the platelet count pathology reports (unless previously provided for patients re-initiating therapy).

The platelet count must be no more than one month old at the time of application.

A maximum of 24 weeks of treatment with this drug will be authorised under this criterion.

Note Eltrombopag is not PBS-subsidised as an alternative to splenectomy.

Note Patients will be able to trial either eltrombopag or romiplostim within the initial 24 weeks treatment period. Patients who fail to demonstrate a response to treatment with eltrombopag and/or romiplostim under the initial restriction will not be eligible to receive further PBS-subsidised treatment with either of these drugs.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe thrombocytopenia

Treatment Phase: Second or subsequent Continuing treatment

Clinical criteria:

- The condition must be severe chronic immune (idiopathic) thrombocytopenic purpura (ITP), **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have demonstrated a continuing response to treatment with this drug, **AND**
- The treatment must be the sole PBS-subsidised thrombopoietin receptor agonist (TRA) for this condition.

Population criteria:

- Patient must be an adult.

For the purpose of this restriction, a continuing response to treatment with drug is defined as:

(a) use of rescue medication (corticosteroids or immunoglobulins) on no more than one occasion during the most recent 24 week period of PBS-subsidised treatment with this drug

AND either of the following:

(b) a platelet count greater than or equal to 50,000 million per L

OR

(c) a platelet count greater than 30,000 million per L and which is double the baseline platelet count.

The platelet count must be no more than one month old at the time of application.

Authority applications for second and subsequent periods of continuing therapy may be made by telephone

Note Eltrombopag is not PBS-subsidised as an alternative to splenectomy.

Note Authority applications for second and subsequent continuing treatment may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note Patients will be able to trial either eltrombopag or romiplostim within the initial 24 weeks treatment period. Patients who fail to demonstrate a response to treatment with eltrombopag and/or romiplostim under the initial restriction will not be eligible to receive further PBS-subsidised treatment with either of these drugs.

Authority required

Severe thrombocytopenia

Treatment Phase: Initial 1, Initial 2, First Continuing treatment or Re-initiation of interrupted treatment, and Second and Subsequent Continuing treatment - balance of supply

Clinical criteria:

- The condition must be severe chronic immune (idiopathic) thrombocytopenic purpura (ITP), **AND**
- The treatment must be the sole PBS-subsidised thrombopoietin receptor agonist (TRA) for this condition, **AND**
- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the First Continuing treatment or Re-initiation of interrupted treatment restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Second and subsequent Continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restriction.

Population criteria:

- Patient must be an adult.

Note Authority approval for sufficient therapy to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

eltrombopag 50 mg tablet, 28

5828R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	2920.19	Revolade [NV]

eltrombopag 25 mg tablet, 28

5827Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	1483.79	Revolade [NV]

■ ROMIPILOSTIM

Authority required

Severe thrombocytopenia

Treatment Phase: Initial treatment 1 - New patient

Clinical criteria:

- The condition must be severe chronic immune (idiopathic) thrombocytopenic purpura (ITP), **AND**
- Patient must have had a splenectomy, **AND**
- Patient must have failed to achieve an adequate response to, or be intolerant to, corticosteroid therapy following the splenectomy, **AND**
- Patient must have failed to achieve an adequate response to, or be intolerant to, immunoglobulin therapy following the splenectomy, **AND**
- The treatment must be the sole PBS-subsidised thrombopoietin receptor agonist (TRA) for this condition.

Population criteria:

- Patient must be an adult.

The following criteria indicate failure to achieve an adequate response and must be demonstrated at the time of initial application;

(a) a platelet count of less than or equal to 20,000 million per L; OR

(b) a platelet count of 20,000 million to 30,000 million per L, where the patient is experiencing significant bleeding or has a history of significant bleeding in this platelet range.

At the time of the written authority application, medical practitioners should request the appropriate quantity of vials of appropriate strength to provide sufficient drug for a single treatment at a dose of 1 microgram/kg. Up to 1 repeat may be requested with the initial written application.

Subsequently during the initial period of dose titration, authority applications for a single dose and up to 1 repeat may be requested by telephone. The dose (microgram/kg/week) must be provided at the time of application.

Once a patient's dose has been stable for a period of 4 weeks, authority approvals for sufficient vials of appropriate strength based on the weight of the patient and dose (microgram/kg/week) for up to 4 weeks of treatment and up to 4 repeats may be granted, as long as the total period of treatment authorised under this restriction does not exceed 24 weeks.

Authority approval will not be given for doses higher than 10 micrograms/kg/week

The authority application must be made in writing and must include:

- (1) a completed authority prescription form,
- (2) a signed patient acknowledgement,
- (3) a completed Idiopathic Thrombocytopenic Purpura Initial PBS Authority Application - Supporting Information Form,
- (4) a copy of a full blood count pathology report supporting the diagnosis of ITP, and
- (5) where the application is sought on the basis of a medical contraindication to surgery, a signed and dated letter from the clinician making this assessment which includes the date upon which the patient was assessed for surgery and the clinical grounds upon which surgery is contraindicated.

The full blood count must be no more than 1 month old at the time of application.

Note Romiplostim is not PBS-subsidised as an alternative to splenectomy.

Note Patients will be able to trial either eltrombopag or romiplostim within the initial 24 weeks treatment period. Patients who fail to demonstrate a response to treatment with eltrombopag and/or romiplostim under the initial restriction will not be eligible to receive further PBS-subsidised treatment with either of these drugs.

Note Any queries concerning the arrangements to prescribe this drug may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au
Written applications for authority to prescribe this drug should be forwarded to:
Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe thrombocytopenia

Treatment Phase: Initial treatment 2 - New patient

Clinical criteria:

- The condition must be severe chronic immune (idiopathic) thrombocytopenic purpura (ITP), **AND**
- Patient must not have had a splenectomy, **AND**
- Patient must have failed to achieve an adequate response to, or be intolerant to, corticosteroid therapy at a dose equivalent to 0.5-2 mg/kg/day of prednisone for at least 4-6 weeks, **AND**
- Patient must have failed to achieve an adequate response to, or be intolerant to, immunoglobulin therapy, **AND**
- Patient must be unsuitable for splenectomy due to medical reasons, **AND**
- The treatment must be the sole PBS-subsidised thrombopoietin receptor agonist (TRA) for this condition.

Population criteria:

- Patient must be an adult.

The following criteria indicate failure to achieve an adequate response and must be demonstrated at the time of initial application;

(a) a platelet count of less than or equal to 20,000 million per L; OR

(b) a platelet count of 20,000 million to 30,000 million per L, where the patient is experiencing significant bleeding or has a history of significant bleeding in this platelet range.

At the time of the written authority application, medical practitioners should request the appropriate quantity of vials of appropriate strength to provide sufficient drug for a single treatment at a dose of 1 microgram/kg. Up to 1 repeat may be requested with the initial written application.

Subsequently during the initial period of dose titration, authority applications for a single dose and up to 1 repeat may be requested by telephone. The dose (microgram/kg/week) must be provided at the time of application.

Once a patient's dose has been stable for a period of 4 weeks, authority approvals for sufficient vials of appropriate strength based on the weight of the patient and dose (microgram/kg/week) for up to 4 weeks of treatment and up to 4 repeats may be granted, as long as the total period of treatment authorised under this restriction does not exceed 24 weeks.

Authority approval will not be given for doses higher than 10 micrograms/kg/week

The authority application must be made in writing and must include:

- (1) a completed authority prescription form,
- (2) a signed patient acknowledgement,
- (3) a completed Idiopathic Thrombocytopenic Purpura Initial PBS Authority Application - Supporting Information Form,
- (4) a copy of a full blood count pathology report supporting the diagnosis of ITP, and
- (5) where the application is sought on the basis of a medical contraindication to surgery, a signed and dated letter from the clinician making this assessment which includes the date upon which the patient was assessed for surgery and the clinical grounds upon which surgery is contraindicated.

The full blood count must be no more than 1 month old at the time of application.

Note Romiplostim is not PBS-subsidised as an alternative to splenectomy.

Note Patients will be able to trial either eltrombopag or romiplostim within the initial 24 weeks treatment period. Patients who fail to demonstrate a response to treatment with eltrombopag and/or romiplostim under the initial restriction will not be eligible to receive further PBS-subsidised treatment with either of these drugs.

Note Any queries concerning the arrangements to prescribe this drug may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au
Written applications for authority to prescribe this drug should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe thrombocytopenia

Treatment Phase: First Continuing treatment or Re-initiation of interrupted treatment

Clinical criteria:

- The condition must be severe chronic immune (idiopathic) thrombocytopenic purpura (ITP), **AND**
- Patient must have previously received PBS-subsidised initial treatment with this drug for this condition, **AND**
- Patient must have demonstrated a sustained platelet response to PBS-subsidised treatment with this drug for this condition under the Initial treatment restriction, **AND**
- The treatment must be the sole PBS-subsidised thrombopoietin receptor agonist (TRA) for this condition.

Population criteria:

- Patient must be an adult.

For the purposes of this restriction, a sustained platelet response is defined as:

(a) use of rescue medication (corticosteroids or immunoglobulins) on no more than one occasion during the initial period of PBS-subsidised treatment with this drug,

AND either of the following:

(b) a platelet count greater than or equal to 50,000 million per L on at least four (4) occasions, each at least one week apart;
OR

(c) a platelet count greater than 30,000 million per L and which is double the baseline (pre-treatment) platelet count on at least four (4) occasions, each at least one week apart.

The medical practitioner should request sufficient number of vials of appropriate strength based on the weight of the patient and dose (microgram/kg/week) to provide 4 weeks of treatment. Up to a maximum of 5 repeats may be authorised.

Authority approval will not be given for doses higher than 10 micrograms/kg/week

Applications for the First continuing PBS-subsidised treatment or Re-initiation of interrupted PBS-subsidised treatment must be made in writing and must include:

(1) a completed authority prescription form, and

(2) a completed Idiopathic Thrombocytopenic Purpura Continuing PBS Authority Application - Supporting Information Form , and

(3) copies of the platelet count pathology reports (unless previously provided for patients re-initiating therapy).

The platelet count must be no more than one month old at the time of application.

Note Romiplostim is not PBS-subsidised as an alternative to splenectomy.

Note Patients will be able to trial either eltrombopag or romiplostim within the initial 24 weeks treatment period. Patients who fail to demonstrate a response to treatment with eltrombopag and/or romiplostim under the initial restriction will not be eligible to receive further PBS-subsidised treatment with either of these drugs.

Note Any queries concerning the arrangements to prescribe this drug may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Written applications for authority to prescribe this drug should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe thrombocytopenia

Treatment Phase: Second or Subsequent Continuing treatment

Clinical criteria:

- The condition must be severe chronic immune (idiopathic) thrombocytopenic purpura (ITP), **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have demonstrated a continuing response to treatment with this drug, **AND**
- The treatment must be the sole PBS-subsidised thrombopoietin receptor agonist (TRA) for this condition.

Population criteria:

- Patient must be an adult.

For the purpose of this restriction, a continuing response to treatment with drug is defined as:

(a) use of rescue medication (corticosteroids or immunoglobulins) on no more than one occasion during the most recent 24 week period of PBS-subsidised treatment with this drug

AND either of the following:

(b) a platelet count greater than or equal to 50,000 million per L

OR

(c) a platelet count greater than 30,000 million per L and which is double the baseline platelet count.

The platelet count must be no more than one month old at the time of application.

The medical practitioner should request sufficient number of vials of appropriate strength based on the weight of the patient and dose (microgram/kg/week) to provide 4 weeks of treatment. Up to a maximum of 5 repeats may be authorised.

Authority approval will not be given for doses higher than 10 micrograms/kg/week

Authority applications for second and subsequent periods of continuing therapy may be made by telephone

Note Romiplostim is not PBS-subsidised as an alternative to splenectomy.

Note Authority applications for second and subsequent continuing treatment may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note Patients will be able to trial either eltrombopag or romiplostim within the initial 24 weeks treatment period. Patients who fail to demonstrate a response to treatment with eltrombopag and/or romiplostim under the initial restriction will not be eligible to receive further PBS-subsidised treatment with either of these drugs.

Note Any queries concerning the arrangements to prescribe this drug may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Written applications for authority to prescribe this drug should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe thrombocytopenia

Treatment Phase: Initial 1, Initial 2, First Continuing treatment or Re-initiation of interrupted treatment, and Second and Subsequent Continuing treatment - balance of supply

Clinical criteria:

- The condition must be severe chronic immune (idiopathic) thrombocytopenic purpura (ITP), **AND**
- The treatment must be the sole PBS-subsidised thrombopoietin receptor agonist (TRA) for this condition, **AND**
- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the First Continuing treatment or Re-initiation of interrupted treatment restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Second and subsequent Continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restriction.

Population criteria:

- Patient must be an adult.

Note Authority approval for sufficient therapy to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note No applications for increased repeats will be authorised.

romiplostim 500 microgram injection, 1 vial

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
9699L	1	..	..	1145.43	Nplate [AN]

romiplostim 250 microgram injection, 1 vial

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
9697J	1	..	..	578.37	Nplate [AN]

■ ANTIANEMIC PREPARATIONS

OTHER ANTIANEMIC PREPARATIONS

Other antianemic preparations

■ DARBEPOETIN ALFA

Authority required

Anaemia associated with intrinsic renal disease

Clinical criteria:

- Patient must require transfusion, **AND**
- Patient must have a haemoglobin level of less than 100 g per L, **AND**
- Patient must have intrinsic renal disease, as assessed by a nephrologist.

darbepoetin alfa 40 microgram/0.4 mL injection, 4 x 0.4 mL syringes

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
6323T	2	5	..	*898.61	Aranesp [AN]

darbepoetin alfa 60 microgram/0.3 mL injection, 0.3 mL syringe

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
6490N	8	5	..	*1291.47	Aranesp SureClick [AN]

darbepoetin alfa 40 microgram/0.4 mL injection, 0.4 mL syringe

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
6489M	8	5	..	*898.51	Aranesp SureClick [AN]

darbepoetin alfa 150 microgram/0.3 mL injection, 4 x 0.3 mL syringes

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
6365B	2	5	..	*3052.01	Aranesp [AN]

darbepoetin alfa 150 microgram/0.3 mL injection, 0.3 mL syringe

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
6493R	8	5	..	*3052.03	Aranesp SureClick [AN]

darbepoetin alfa 100 microgram/0.5 mL injection, 4 x 0.5 mL syringes

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
6326Y	2	5	..	*2063.95	Aranesp [AN]

BLOOD AND BLOOD FORMING ORGANS

darbepoetin alfa 30 microgram/0.3 mL injection, 4 x 0.3 mL syringes

6322R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*741.63	Aranesp [AN]

darbepoetin alfa 100 microgram/0.5 mL injection, 0.5 mL syringe

6492Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	8	5	..	*2063.87	Aranesp SureClick [AN]

darbepoetin alfa 60 microgram/0.3 mL injection, 4 x 0.3 mL syringes

6325X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*1291.45	Aranesp [AN]

darbepoetin alfa 50 microgram/0.5 mL injection, 4 x 0.5 mL syringes

6324W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*1106.85	Aranesp [AN]

darbepoetin alfa 20 microgram/0.5 mL injection, 0.5 mL syringe

6488L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	8	5	..	*544.11	Aranesp SureClick [AN]

darbepoetin alfa 20 microgram/0.5 mL injection, 4 x 0.5 mL syringes

6321Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*544.07	Aranesp [AN]

darbepoetin alfa 10 microgram/0.4 mL injection, 4 x 0.4 mL syringes

6320P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*292.37	Aranesp [AN]

darbepoetin alfa 80 microgram/0.4 mL injection, 4 x 0.4 mL syringes

6438W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*1684.93	Aranesp [AN]

darbepoetin alfa 80 microgram/0.4 mL injection, 0.4 mL syringe

6491P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	8	5	..	*1684.99	Aranesp SureClick [AN]

■ EPOETIN ALFA

Authority required

Anaemia associated with intrinsic renal disease

Clinical criteria:

- Patient must require transfusion, **AND**
- Patient must have a haemoglobin level of less than 100 g per L, **AND**
- Patient must have intrinsic renal disease, as assessed by a nephrologist.

epoetin alfa 3000 units/0.3 mL injection, 6 x 0.3 mL syringes

6205N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*541.11	Eprex 3000 [JC]

epoetin alfa 6000 units/0.6 mL injection, 6 x 0.6 mL syringes

6303R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*1011.89	Eprex 6000 [JC]

epoetin alfa 8000 units/0.8 mL injection, 6 x 0.8 mL syringes

6305W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*1300.11	Eprex 8000 [JC]

epoetin alfa 10 000 units/mL injection, 6 x 1 mL syringes

6207Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*1563.57	Eprex 10000 [JC]

epoetin alfa 1000 units/0.5 mL injection, 6 x 0.5 mL syringes

6251B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*230.91	Eprex 1000 [JC]

epoetin alfa 4000 units/0.4 mL injection, 6 x 0.4 mL syringes

6206P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*687.09	Eprex 4000 [JC]

epoetin alfa 20 000 units/0.5 mL injection, 6 x 0.5 mL syringes

6434P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*3030.07	Eprex 20,000 [JC]

epoetin alfa 2000 units/0.5 mL injection, 6 x 0.5 mL syringes

6204M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*420.99	Eprex 2000 [JC]

epoetin alfa 40 000 units/mL injection, 1 mL syringe

6339P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*1010.97	Eprex 40,000 [JC]

epoetin alfa 5000 units/0.5 mL injection, 6 x 0.5 mL syringes

6302Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*853.59	Eprex 5000 [JC]

■ EPOETIN BETA**Authority required**

Anaemia associated with intrinsic renal disease

Clinical criteria:

- Patient must require transfusion, **AND**
- Patient must have a haemoglobin level of less than 100 g per L, **AND**
- Patient must have intrinsic renal disease, as assessed by a nephrologist.

epoetin beta 2000 units/0.3 mL injection, 6 x 0.3 mL syringes

6480C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*442.75	NeoRecormon [RO]

epoetin beta 6000 units/0.3 mL injection, 6 x 0.3 mL syringes

6484G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*1064.09	NeoRecormon [RO]

epoetin beta 5000 units/0.3 mL injection, 6 x 0.3 mL syringes

6483F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*898.15	NeoRecormon [RO]

epoetin beta 3000 units/0.3 mL injection, 6 x 0.3 mL syringes

6481D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*569.21	NeoRecormon [RO]

epoetin beta 10 000 units/0.6 mL injection, 6 x 0.6 mL syringes

6485H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*1643.37	NeoRecormon [RO]

epoetin beta 4000 units/0.3 mL injection, 6 x 0.3 mL syringes

6482E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*722.87	NeoRecormon [RO]

■ EPOETIN LAMBDA**Authority required**

Anaemia associated with intrinsic renal disease

Clinical criteria:

- Patient must require transfusion, **AND**
- Patient must have a haemoglobin level of less than 100 g per L, **AND**
- Patient must have intrinsic renal disease, as assessed by a nephrologist.

epoetin lambda 3000 units/0.3 mL injection, 6 x 0.3 mL syringes

9687W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*631.61	Novicrit [SZ]

epoetin lambda 5000 units/0.5 mL injection, 6 x 0.5 mL syringes

9588P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*997.07	Novicrit [SZ]

epoetin lambda 6000 units/0.6 mL injection, 6 x 0.6 mL syringes

9590R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*1177.01	Novicrit [SZ]

CARDIOVASCULAR SYSTEM

epoetin lambda 1000 units/0.5 mL injection, 6 x 0.5 mL syringes

9685R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*268.83	Novicrit [SZ]

epoetin lambda 8000 units/0.8 mL injection, 6 x 0.8 mL syringes

9593X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*1512.51	Novicrit [SZ]

epoetin lambda 2000 units/mL injection, 6 x 1 mL syringes

9686T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*491.11	Novicrit [SZ]

epoetin lambda 10 000 units/mL injection, 6 x 1 mL syringes

9595B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*1820.67	Novicrit [SZ]

epoetin lambda 4000 units/0.4 mL injection, 6 x 0.4 mL syringes

9688X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*802.35	Novicrit [SZ]

■ METHOXY POLYETHYLENE GLYCOL-EPOETIN BETA

Authority required

Anaemia associated with intrinsic renal disease

Clinical criteria:

- Patient must require transfusion, **AND**
- Patient must have a haemoglobin level of less than 100 g per L, **AND**
- Patient must have intrinsic renal disease, as assessed by a nephrologist.

methoxy polyethylene glycol-epoetin beta 30 microgram/0.3 mL injection, 0.3 mL syringe

9574X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*352.97	Mircera [RO]

methoxy polyethylene glycol-epoetin beta 100 microgram/0.3 mL injection, 0.3 mL syringe

9577C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*1090.37	Mircera [RO]

methoxy polyethylene glycol-epoetin beta 75 microgram/0.3 mL injection, 0.3 mL syringe

9576B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*846.09	Mircera [RO]

methoxy polyethylene glycol-epoetin beta 50 microgram/0.3 mL injection, 0.3 mL syringe

9575Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*583.35	Mircera [RO]

methoxy polyethylene glycol-epoetin beta 360 microgram/0.6 mL injection, 0.6 mL syringe

9580F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*3041.37	Mircera [RO]

methoxy polyethylene glycol-epoetin beta 120 microgram/0.3 mL injection, 0.3 mL syringe

9578D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*1254.91	Mircera [RO]

methoxy polyethylene glycol-epoetin beta 200 microgram/0.3 mL injection, 0.3 mL syringe

9579E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*1779.31	Mircera [RO]

■ CARDIOVASCULAR SYSTEM

■ ANTIHYPERTENSIVES

OTHER ANTIHYPERTENSIVES

Antihypertensives for pulmonary arterial hypertension

■ AMBRISENTAN

Caution This is a category X drug and must not be given to pregnant women. Pregnancy must be avoided during treatment and for at least 3 months following cessation of therapy.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 1 (new patients)

Clinical criteria:

- Patient must not have received prior PBS-subsidised treatment with a pulmonary arterial hypertension (PAH) agent, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**
- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH; OR
- Patient must have WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease, **AND**
- Patient must have a mean right atrial pressure of 8 mmHg or less as measured by right heart catheterisation (RHC); OR
- Patient must have right ventricular function assessed by echocardiography (ECHO) where a RHC cannot be performed on clinical grounds, **AND**
- Patient must have failed to respond to 6 or more weeks of appropriate vasodilator treatment unless intolerance or a contraindication to such treatment exists, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6 Minute Walk Test (6MWT); and
- (3) a signed patient acknowledgement.

Idiopathic pulmonary arterial hypertension, anorexigen-induced pulmonary arterial hypertension, hereditary pulmonary arterial hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) are defined as follows:

- (i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary artery wedge pressure (PAWP) less than 15 mmHg; or
- (ii) where a right heart catheter (RHC) cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.

Test requirements to establish baseline for initiation of treatment are as follows:

The first written application for PBS-subsidised treatment with the first PAH agent should be accompanied by the results of a right heart catheter (RHC) composite assessment plus an echocardiograph (ECHO) composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.

Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments;
- (2) RHC composite assessment plus 6MWT;
- (3) RHC composite assessment only.

In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted for consideration based on the results of the following test combinations, which are listed in descending order of preference:

- (1) ECHO composite assessment plus 6MWT;
- (2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

Details of prior vasodilator treatment, including the dose and duration of treatment, must be provided at the time of application. Where the patient has an adverse event to a vasodilator or where vasodilator treatment is contraindicated, details of the nature of the adverse event or contraindication according to the Therapeutic Goods Administration (TGA) approved Product Information must also be provided with the application.

Response to prior vasodilator treatment is defined as follows:

For patients with 2 or more baseline tests, response to treatment is defined as 2 or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information.

A maximum of 5 repeats may be requested.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Prior Written Approval of Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 2 (new patients)

Clinical criteria:

- Patient must not have received prior PBS-subsidised treatment with a pulmonary arterial hypertension (PAH) agent, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**
- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH), or anorexigen-induced PAH or hereditary PAH, and a mean right atrial pressure of greater than 8 mmHg, as measured by right heart catheterisation (RHC); OR
- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH), or anorexigen-induced PAH or hereditary PAH, with right ventricular function assessed by echocardiography (ECHO) where a RHC cannot be performed on clinical grounds; OR
- Patient must have WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease and a mean right atrial pressure greater than 8 mmHg, as measured by RHC; OR
- Patient must have WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease with right ventricular function assessed by ECHO where a RHC cannot be performed on clinical grounds; OR
- Patient must have WHO Functional Class IV idiopathic pulmonary arterial hypertension (iPAH), or anorexigen-induced PAH or hereditary PAH; OR
- Patient must have WHO Functional Class IV pulmonary arterial hypertension secondary to connective tissue disease, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6 Minute Walk Test (6MWT); and
- (3) a signed patient acknowledgement.

Idiopathic pulmonary arterial hypertension, anorexigen-induced pulmonary arterial hypertension, hereditary pulmonary arterial hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) are defined as follows:

- (i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary artery wedge pressure (PAWP) less than 15 mmHg; or
- (ii) where a right heart catheter (RHC) cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.

Test requirements to establish baseline for initiation of treatment are as follows:

The first written application for PBS-subsidised treatment with the first PAH agent should be accompanied by the results of a right heart catheter (RHC) composite assessment plus an echocardiograph (ECHO) composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.

Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments;
- (2) RHC composite assessment plus 6MWT;
- (3) RHC composite assessment only.

In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted for consideration based on the results of the following test combinations, which are listed in descending order of preference:

- (1) ECHO composite assessment plus 6MWT;
- (2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats may be requested.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 3 (change or re-commencement of therapy for all patients)

Clinical criteria:

- Patient must have idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH or PAH secondary to connective tissue disease and must wish to re-commence PBS-subsidised therapy with this agent after a break in therapy and must have demonstrated a response to their most recent course of PBS-subsidised treatment with this agent; OR
- Patient must have idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH or PAH secondary to connective tissue disease and whose most recent course of PBS-subsidised treatment was with a PAH agent other than this agent, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form; and
- (3) the results of the patient's response to treatment with their last course of PBS-subsidised PAH agent.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

Response to a PAH agent is defined as follows:

For patients with two or more baseline tests, response to treatment is defined as two or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats may be requested.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Swapping between PAH agents: Patients can access PAH agents through the PBS according to the relevant restrictions.

Once these patients are approved initial treatment with 1 of these 8 drugs, they may swap between PAH agents at any time without having to re-qualify for treatment with the alternate agent. This means that patients may commence treatment with the alternate agent, subject to that agent's restriction, irrespective of the severity of their disease at the time the application to swap therapy is submitted. It also means that no new baseline measurements will be necessary. New baselines may be submitted where the patient has failed to respond to their current treatment. Eligible patients may only swap between PAH agents if they have not failed prior PBS-subsidised treatment with that agent. For eligible patients, applications to swap between the 8 PAH agents must be made under the relevant initial treatment restriction. Patients should be assessed for response to the treatment they are ceasing at the time the application to swap therapy is being made. Patients who fail to demonstrate a response or for whom no assessment results are submitted with the application to swap therapy may not re-commence PBS-subsidised treatment with the drug they are ceasing.

Note Applications for patients who wish to swap to an alternate PAH agent should be accompanied by the previously approved authority prescription, or remaining repeats, for the treatment the patient is ceasing.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au
Applications for authority to prescribe should be forwarded to:
Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 1 or Initial 2 (new patients) or Initial 3 (change or re-commencement of therapy for all patients) or First Continuing treatment - Balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this agent under the Initial 1 (new patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the Initial 2 (new patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the Initial 3 (change or re-commencement of therapy for all patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the First Continuing treatment restriction to complete a maximum of six months of treatment, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition, **AND**
- The treatment must provide no more than the balance of up to six months treatment available under one of the above restrictions.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authorisation under this criterion should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: First Continuing treatment

Clinical criteria:

- Patient must have received a PBS-subsidised initial course of treatment with this agent for this condition, **AND**
- Patient must have been assessed by a physician from a designated hospital to have achieved a response to the PBS-subsidised initial course of treatment, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

(1) a completed authority prescription form; and

(2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:

(i) RHC composite assessment; and

(ii) ECHO composite assessment; and

(iii) 6 Minute Walk Test (6MWT).

Test requirements to establish response to treatment for continuation of treatment are as follows:

The following list outlines the preferred test combination, in descending order, for the purposes of continuation of PBS-subsidised treatment:

(1) RHC plus ECHO composite assessments plus 6MWT;

(2) RHC plus ECHO composite assessments;

(3) RHC composite assessment plus 6MWT;

(4) ECHO composite assessment plus 6MWT;

(5) RHC composite assessment only;

(6) ECHO composite assessment only.

The results of the same tests as conducted at baseline should be provided with the written First Continuing treatment application, except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a patient specific reason why the test(s) could not be conducted must be provided with the application.

The test results provided with the application for continuing treatment must be no more than 2 months old at the time of application.

Response to a PAH agent is defined as follows:

For patients with two or more baseline tests, response to treatment is defined as two or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats will be authorised.

An application for First Continuing treatment with a PAH agent should be made prior to the completion of the Initial 6 month treatment course to ensure continuity for those patients who respond to treatment, as assessed by the treating physician.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Subsequent Continuing treatment

Clinical criteria:

- Patient must have received a PBS-subsidised treatment under First Continuing treatment with this agent for this condition; OR
- Patient must have previously received PBS-subsidised treatment under this criteria with this agent for this condition, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats will be authorised.

An application for Subsequent Continuing treatment with a PAH agents should be made prior to the completion of the First Continuing treatment course to ensure continuity of treatment.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authorisation under this criterion should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

ambrisentan 10 mg tablet, 30

9649W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	2780.04	Volibris [GK]

ambrisentan 5 mg tablet, 30

9648T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	2780.04	Volibris [GK]

■ BOSENTAN

Caution This is a category X drug and must not be given to pregnant women. Pregnancy must be avoided during treatment and for at least 3 months following cessation of therapy.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 1 (new patients)

Clinical criteria:

- Patient must not have received prior PBS-subsidised treatment with a pulmonary arterial hypertension (PAH) agent, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**
- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH; OR
- Patient must have WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease, **AND**
- Patient must have a mean right atrial pressure of 8 mmHg or less as measured by right heart catheterisation (RHC); OR
- Patient must have right ventricular function assessed by echocardiography (ECHO) where a RHC cannot be performed on clinical grounds, **AND**
- Patient must have failed to respond to 6 or more weeks of appropriate vasodilator treatment unless intolerance or a contraindication to such treatment exists, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

- (1) two completed authority prescription forms; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6 Minute Walk Test (6MWT); and
- (3) a signed patient acknowledgement.

Idiopathic pulmonary arterial hypertension, anorexigen-induced pulmonary arterial hypertension, hereditary pulmonary arterial hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) are defined as follows:

- (i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary artery wedge pressure (PAWP) less than 15 mmHg; or
- (ii) where a right heart catheter (RHC) cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.

Test requirements to establish baseline for initiation of treatment are as follows:

The first written application for PBS-subsidised treatment with the first PAH agent should be accompanied by the results of a right heart catheter (RHC) composite assessment plus an echocardiograph (ECHO) composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.

Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments;
- (2) RHC composite assessment plus 6MWT;
- (3) RHC composite assessment only.

In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted for consideration based on the results of the following test combinations, which are listed in descending order of preference:

- (1) ECHO composite assessment plus 6MWT;
- (2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

Details of prior vasodilator treatment, including the dose and duration of treatment, must be provided at the time of application. Where the patient has an adverse event to a vasodilator or where vasodilator treatment is contraindicated, details of the nature of the adverse event or contraindication according to the Therapeutic Goods Administration (TGA) approved Product Information must also be provided with the application.

Response to prior vasodilator treatment is defined as follows:

For patients with 2 or more baseline tests, response to treatment is defined as 2 or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

Approvals for the first authority prescription will be limited to 1 month of therapy with the 62.5 mg strength tablet, with the quantity approved based on the dosage recommendations in the TGA-approved Product Information. No repeats will be authorised for this prescription.

The second authority prescription may be written for either the 62.5 mg tablet or the 125 mg tablet strengths. Approvals for the second authority prescription will be limited to 1 month of treatment, with the quantity approved based on the dosage recommendations in the TGA-approved Product Information, and a maximum of 4 repeats.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Where the 62.5 mg tablet strength is required for the second authority prescription, please contact the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) for further advice. The approved second authority prescription will be returned to the prescriber by the Department of Human Services two weeks after the date of the approval of the first authority prescription, to allow for the uninterrupted completion of the six months initial treatment course. The Department of Human Services will contact prescribers prior to dispatch of the second authority prescription to confirm the tablet strength required for the patient.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au. Applications for authority to prescribe should be forwarded to:
Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 2 (new patients)

Clinical criteria:

- Patient must not have received prior PBS-subsidised treatment with a pulmonary arterial hypertension (PAH) agent, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**
- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH), or anorexigen-induced PAH or hereditary PAH, and a mean right atrial pressure of greater than 8 mmHg, as measured by right heart catheterisation (RHC); OR
- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH), or anorexigen-induced PAH or hereditary PAH, with right ventricular function assessed by echocardiography (ECHO) where a RHC cannot be performed on clinical grounds; OR
- Patient must have WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease and a mean right atrial pressure greater than 8 mmHg, as measured by RHC; OR
- Patient must have WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease with right ventricular function assessed by ECHO where a RHC cannot be performed on clinical grounds; OR
- Patient must have WHO Functional Class IV idiopathic pulmonary arterial hypertension (iPAH), or anorexigen-induced PAH or hereditary PAH; OR
- Patient must have WHO Functional Class IV pulmonary arterial hypertension secondary to connective tissue disease; OR
- Patient must have WHO Functional Class III or IV pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology), **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

(1) two completed authority prescription forms; and

(2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:

(i) RHC composite assessment; and

(ii) ECHO composite assessment; and

(iii) 6 Minute Walk Test (6MWT); and

(3) a signed patient acknowledgement.

Idiopathic pulmonary arterial hypertension, anorexigen-induced pulmonary arterial hypertension, hereditary pulmonary arterial hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) are defined as follows:

(i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary artery wedge pressure (PAWP) less than 15 mmHg; or

(ii) where a right heart catheter (RHC) cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.

Test requirements to establish baseline for initiation of treatment are as follows:

The first written application for PBS-subsidised treatment with the first PAH agent should be accompanied by the results of a right heart catheter (RHC) composite assessment plus an echocardiograph (ECHO) composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.

Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:

(1) RHC plus ECHO composite assessments;

(2) RHC composite assessment plus 6MWT;

(3) RHC composite assessment only.

In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted for consideration based on the results of the following test combinations, which are listed in descending order of preference:

(1) ECHO composite assessment plus 6MWT;

(2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

Approvals for the first authority prescription will be limited to 1 month of therapy with the 62.5 mg strength tablet, with the quantity approved based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information. No repeats will be authorised for this prescription.

The second authority prescription may be written for either the 62.5 mg tablet or the 125 mg tablet strengths. Approvals for the second authority prescription will be limited to 1 month of treatment, with the quantity approved based on the dosage recommendations in the TGA-approved Product Information, and a maximum of 4 repeats.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Where the 62.5 mg tablet strength is required for the second authority prescription, please contact the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) for further advice.

The approved second authority prescription will be returned to the prescriber by the Department of Human Services two weeks after the date of the approval of the first authority prescription, to allow for the uninterrupted completion of the six months initial treatment course. The Department of Human Services will contact prescribers prior to dispatch of the second authority prescription to confirm the tablet strength required for the patient.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 3 (change or re-commencement of therapy for all patients)

Clinical criteria:

- Patient must have idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH or PAH secondary to connective tissue disease or PAH associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) and must wish to re-commence PBS-subsidised therapy with this agent after a break in therapy and must have demonstrated a response to their most recent course of PBS-subsidised treatment with this agent; OR
- Patient must have idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH or PAH secondary to connective tissue disease or PAH associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) and whose most recent course of PBS-subsidised treatment was with a PAH agent other than this agent, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

(1) two completed authority prescription forms; and

(2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form; and

(3) the results of the patient's response to treatment with their last course of PBS-subsidised PAH agent.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

Response to a PAH agent is defined as follows:

For patients with two or more baseline tests, response to treatment is defined as two or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

Approvals for the first authority prescription will be limited to 1 month of therapy with the 62.5 mg strength tablet, with the quantity approved based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information. No repeats will be authorised for this prescription.

The second authority prescription may be written for either the 62.5 mg tablet or the 125 mg tablet strengths. Approvals for the second authority prescription will be limited to 1 month of treatment, with the quantity approved based on the dosage recommendations in the TGA-approved Product Information, and a maximum of 4 repeats.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Swapping between PAH agents: Patients can access PAH agents through the PBS according to the relevant restrictions. Once these patients are approved initial treatment with 1 of these 8 drugs, they may swap between PAH agents at any time without having to re-qualify for treatment with the alternate agent. This means that patients may commence treatment with the alternate agent, subject to that agent's restriction, irrespective of the severity of their disease at the time the application to swap therapy is submitted. It also means that no new baseline measurements will be necessary. New baselines may be submitted where the patient has failed to respond to their current treatment. Eligible patients may only swap between PAH agents if they have not failed prior PBS-subsidised treatment with that agent. For eligible patients, applications to swap between the 8 PAH agents must be made under the relevant initial treatment restriction. Patients should be assessed for response to the treatment they are ceasing at the time the application to swap therapy is being made. Patients who fail to demonstrate a response or for whom no assessment results are submitted with the application to swap therapy may not re-commence PBS-subsidised treatment with the drug they are ceasing.

Note Where the 62.5 mg tablet strength is required for the second authority prescription, please contact the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) for further advice. The approved second authority prescription will be returned to the prescriber by the Department of Human Services two weeks after the date of the approval of the first authority prescription, to allow for the uninterrupted completion of the six months initial treatment course. The Department of Human Services will contact prescribers prior to dispatch of the second authority prescription to confirm the tablet strength required for the patient.

Note Applications for patients who wish to swap to an alternate PAH agent should be accompanied by the previously approved authority prescription, or remaining repeats, for the treatment the patient is ceasing.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 1 or Initial 2 (new patients) or Initial 3 (change or re-commencement of therapy for all patients) or First Continuing treatment - Balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this agent under the Initial 1 (new patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the Initial 2 (new patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the Initial 3 (change or re-commencement of therapy for all patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the First Continuing treatment restriction to complete a maximum of six months of treatment, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition, **AND**
- The treatment must provide no more than the balance of up to six months treatment available under one of the above restrictions.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authorisation under this criterion should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: First Continuing treatment

Clinical criteria:

- Patient must have received a PBS-subsidised initial course of treatment with this agent for this condition, **AND**
- Patient must have been assessed by a physician from a designated hospital to have achieved a response to the PBS-subsidised initial course of treatment, **AND**

- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6 Minute Walk Test (6MWT).

Test requirements to establish response to treatment for continuation of treatment are as follows:

The following list outlines the preferred test combination, in descending order, for the purposes of continuation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments plus 6MWT;
- (2) RHC plus ECHO composite assessments;
- (3) RHC composite assessment plus 6MWT;
- (4) ECHO composite assessment plus 6MWT;
- (5) RHC composite assessment only;
- (6) ECHO composite assessment only.

The results of the same tests as conducted at baseline should be provided with the written First Continuing treatment application, except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a patient specific reason why the test(s) could not be conducted must be provided with the application.

The test results provided with the application for continuing treatment must be no more than 2 months old at the time of application.

Response to a PAH agent is defined as follows:

For patients with two or more baseline tests, response to treatment is defined as two or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats will be authorised.

An application for First Continuing treatment with a PAH agent should be made prior to the completion of the Initial 6 month treatment course to ensure continuity for those patients who respond to treatment, as assessed by the treating physician.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Subsequent Continuing treatment

Clinical criteria:

- Patient must have received a PBS-subsidised treatment under First Continuing treatment with this agent for this condition; OR
- Patient must have previously received PBS-subsidised treatment under this criteria with this agent for this condition, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats will be authorised.

An application for Subsequent Continuing treatment with a PAH agents should be made prior to the completion of the First Continuing treatment course to ensure continuity of treatment.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authorisation under this criterion should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

bosentan 125 mg tablet, 60

6430K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	1160.47	^a Bosentan APO [GX] ^a BOSENTAN DR. REDDY'S [RI] ^a Bosentan Mylan [AF] ^a Bosentan Sandoz [SZ] ^a Tracleer [AT]	^a BOSENTAN-DRLA [RZ] ^a Bosentan GH [GQ] ^a Bosentan RBX [RA] ^a BOSLEER [RW]

■ BOSENTAN

Caution This is a category X drug and must not be given to pregnant women. Pregnancy must be avoided during treatment and for at least 3 months following cessation of therapy.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 1 (new patients)

Clinical criteria:

- Patient must not have received prior PBS-subsidised treatment with a pulmonary arterial hypertension (PAH) agent, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**
- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH; OR
- Patient must have WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease, **AND**

- Patient must have a mean right atrial pressure of 8 mmHg or less as measured by right heart catheterisation (RHC); OR
- Patient must have right ventricular function assessed by echocardiography (ECHO) where a RHC cannot be performed on clinical grounds, **AND**
- Patient must have failed to respond to 6 or more weeks of appropriate vasodilator treatment unless intolerance or a contraindication to such treatment exists, **AND**

- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

(1) two completed authority prescription forms; and

(2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:

- RHC composite assessment; and
 - ECHO composite assessment; and
 - 6 Minute Walk Test (6MWT); and
- (3) a signed patient acknowledgement.

Idiopathic pulmonary arterial hypertension, anorexigen-induced pulmonary arterial hypertension, hereditary pulmonary arterial hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) are defined as follows:

(i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary artery wedge pressure (PAWP) less than 15 mmHg; or

(ii) where a right heart catheter (RHC) cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.

Test requirements to establish baseline for initiation of treatment are as follows:

The first written application for PBS-subsidised treatment with the first PAH agent should be accompanied by the results of a right heart catheter (RHC) composite assessment plus an echocardiograph (ECHO) composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.

Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments;
- (2) RHC composite assessment plus 6MWT;
- (3) RHC composite assessment only.

In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted for consideration based on the results of the following test combinations, which are listed in descending order of preference:

- (1) ECHO composite assessment plus 6MWT;
- (2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

Details of prior vasodilator treatment, including the dose and duration of treatment, must be provided at the time of application. Where the patient has an adverse event to a vasodilator or where vasodilator treatment is contraindicated, details of the nature of the adverse event or contraindication according to the Therapeutic Goods Administration (TGA) approved Product Information must also be provided with the application.

Response to prior vasodilator treatment is defined as follows:

For patients with 2 or more baseline tests, response to treatment is defined as 2 or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

Approvals for the first authority prescription will be limited to 1 month of therapy with the 62.5 mg strength tablet, with the quantity approved based on the dosage recommendations in the TGA-approved Product Information. No repeats will be authorised for this prescription.

The second authority prescription may be written for either the 62.5 mg tablet or the 125 mg tablet strengths. Approvals for the second authority prescription will be limited to 1 month of treatment, with the quantity approved based on the dosage recommendations in the TGA-approved Product Information, and a maximum of 4 repeats.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Where the 62.5 mg tablet strength is required for the second authority prescription, please contact the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) for further advice.

The approved second authority prescription will be returned to the prescriber by the Department of Human Services two weeks after the date of the approval of the first authority prescription, to allow for the uninterrupted completion of the six months initial treatment course. The Department of Human Services will contact prescribers prior to dispatch of the second authority prescription to confirm the tablet strength required for the patient.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 2 (new patients)

Clinical criteria:

- Patient must not have received prior PBS-subsidised treatment with a pulmonary arterial hypertension (PAH) agent, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**
- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH), or anorexigen-induced PAH or hereditary PAH, and a mean right atrial pressure of greater than 8 mmHg, as measured by right heart catheterisation (RHC); OR
- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH), or anorexigen-induced PAH or hereditary PAH, with right ventricular function assessed by echocardiography (ECHO) where a RHC cannot be performed on clinical grounds; OR
- Patient must have WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease and a mean right atrial pressure greater than 8 mmHg, as measured by RHC; OR
- Patient must have WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease with right ventricular function assessed by ECHO where a RHC cannot be performed on clinical grounds; OR
- Patient must have WHO Functional Class IV idiopathic pulmonary arterial hypertension (iPAH), or anorexigen-induced PAH or hereditary PAH; OR
- Patient must have WHO Functional Class IV pulmonary arterial hypertension secondary to connective tissue disease; OR
- Patient must have WHO Functional Class III or IV pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology), **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

(1) two completed authority prescription forms; and

(2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:

- (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6 Minute Walk Test (6MWT); and
- (3) a signed patient acknowledgement.

Idiopathic pulmonary arterial hypertension, anorexigen-induced pulmonary arterial hypertension, hereditary pulmonary arterial hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) are defined as follows:

- (i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary artery wedge pressure (PAWP) less than 15 mmHg; or
- (ii) where a right heart catheter (RHC) cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.

Test requirements to establish baseline for initiation of treatment are as follows:

The first written application for PBS-subsidised treatment with the first PAH agent should be accompanied by the results of a right heart catheter (RHC) composite assessment plus an echocardiograph (ECHO) composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.

Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments;
- (2) RHC composite assessment plus 6MWT;
- (3) RHC composite assessment only.

In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted for consideration based on the results of the following test combinations, which are listed in descending order of preference:

- (1) ECHO composite assessment plus 6MWT;
- (2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

Approvals for the first authority prescription will be limited to 1 month of therapy with the 62.5 mg strength tablet, with the quantity approved based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information. No repeats will be authorised for this prescription.

The second authority prescription may be written for either the 62.5 mg tablet or the 125 mg tablet strengths. Approvals for the second authority prescription will be limited to 1 month of treatment, with the quantity approved based on the dosage recommendations in the TGA-approved Product Information, and a maximum of 4 repeats.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Where the 62.5 mg tablet strength is required for the second authority prescription, please contact the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) for further advice.

The approved second authority prescription will be returned to the prescriber by the Department of Human Services two weeks after the date of the approval of the first authority prescription, to allow for the uninterrupted completion of the six months initial treatment course. The Department of Human Services will contact prescribers prior to dispatch of the second authority prescription to confirm the tablet strength required for the patient.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 3 (change or re-commencement of therapy for all patients)

Clinical criteria:

- Patient must have idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH or PAH secondary to connective tissue disease or PAH associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) and must wish to re-commence PBS-subsidised therapy with this agent after a break in therapy and must have demonstrated a response to their most recent course of PBS-subsidised treatment with this agent;
OR

- Patient must have idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH or PAH secondary to connective tissue disease or PAH associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) and whose most recent course of PBS-subsidised treatment was with a PAH agent other than this agent, **AND**

- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

- (1) two completed authority prescription forms; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form; and
- (3) the results of the patient's response to treatment with their last course of PBS-subsidised PAH agent.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

Response to a PAH agent is defined as follows:

For patients with two or more baseline tests, response to treatment is defined as two or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

Approvals for the first authority prescription will be limited to 1 month of therapy with the 62.5 mg strength tablet, with the quantity approved based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information. No repeats will be authorised for this prescription.

The second authority prescription may be written for either the 62.5 mg tablet or the 125 mg tablet strengths. Approvals for the second authority prescription will be limited to 1 month of treatment, with the quantity approved based on the dosage recommendations in the TGA-approved Product Information, and a maximum of 4 repeats.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Swapping between PAH agents: Patients can access PAH agents through the PBS according to the relevant restrictions. Once these patients are approved initial treatment with 1 of these 8 drugs, they may swap between PAH agents at any time without having to re-qualify for treatment with the alternate agent. This means that patients may commence treatment with the alternate agent, subject to that agent's restriction, irrespective of the severity of their disease at the time the application to swap therapy is submitted. It also means that no new baseline measurements will be necessary. New baselines may be submitted where the patient has failed to respond to their current treatment. Eligible patients may only swap between PAH agents if they have not failed prior PBS-subsidised treatment with that agent. For eligible patients, applications to swap between the 8 PAH agents must be made under the relevant initial treatment restriction. Patients should be assessed for response to the treatment they are ceasing at the time the application to swap therapy is being made. Patients who fail to demonstrate a response or for whom no assessment results are submitted with the application to swap therapy may not re-commence PBS-subsidised treatment with the drug they are ceasing.

Note Where the 62.5 mg tablet strength is required for the second authority prescription, please contact the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) for further advice. The approved second authority prescription will be returned to the prescriber by the Department of Human Services two weeks after the date of the approval of the first authority prescription, to allow for the uninterrupted completion of the six months initial treatment course. The Department of Human Services will contact prescribers prior to dispatch of the second authority prescription to confirm the tablet strength required for the patient.

Note Applications for patients who wish to swap to an alternate PAH agent should be accompanied by the previously approved authority prescription, or remaining repeats, for the treatment the patient is ceasing.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 1 or Initial 2 (new patients) or Initial 3 (change or re-commencement of therapy for all patients) or First Continuing treatment - Balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this agent under the Initial 1 (new patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the Initial 2 (new patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the Initial 3 (change or re-commencement of therapy for all patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the First Continuing treatment restriction to complete a maximum of six months of treatment, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition, **AND**
- The treatment must provide no more than the balance of up to six months treatment available under one of the above restrictions.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Written applications for authorisation under this criterion should be forwarded to:
Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: First Continuing treatment

Clinical criteria:

- Patient must have received a PBS-subsidised initial course of treatment with this agent for this condition, **AND**
- Patient must have been assessed by a physician from a designated hospital to have achieved a response to the PBS-subsidised initial course of treatment, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

(1) a completed authority prescription form; and

(2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:

(i) RHC composite assessment; and

(ii) ECHO composite assessment; and

(iii) 6 Minute Walk Test (6MWT).

Test requirements to establish response to treatment for continuation of treatment are as follows:

The following list outlines the preferred test combination, in descending order, for the purposes of continuation of PBS-subsidised treatment:

(1) RHC plus ECHO composite assessments plus 6MWT;

(2) RHC plus ECHO composite assessments;

(3) RHC composite assessment plus 6MWT;

(4) ECHO composite assessment plus 6MWT;

(5) RHC composite assessment only;

(6) ECHO composite assessment only.

The results of the same tests as conducted at baseline should be provided with the written First Continuing treatment application, except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a patient specific reason why the test(s) could not be conducted must be provided with the application.

The test results provided with the application for continuing treatment must be no more than 2 months old at the time of application.

Response to a PAH agent is defined as follows:

For patients with two or more baseline tests, response to treatment is defined as two or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats will be authorised.

An application for First Continuing treatment with a PAH agent should be made prior to the completion of the Initial 6 month treatment course to ensure continuity for those patients who respond to treatment, as assessed by the treating physician.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Subsequent Continuing treatment

Clinical criteria:

- Patient must have received a PBS-subsidised treatment under First Continuing treatment with this agent for this condition; OR
- Patient must have previously received PBS-subsidised treatment under this criteria with this agent for this condition, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats will be authorised.

An application for Subsequent Continuing treatment with a PAH agents should be made prior to the completion of the First Continuing treatment course to ensure continuity of treatment.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authorisation under this criterion should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Cessation of treatment (all patients)

Clinical criteria:

- Patient must have received approval for initial PBS-subsidised treatment with this agent, **AND**
 - Patient must have not responded to prior PBS-subsidised therapy with this agent, **AND**
 - The treatment must be for the purpose of gradual dose reduction prior to ceasing therapy, **AND**
 - The treatment must be the sole PBS-subsidised PAH agent for this condition.
- The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment. Treatment beyond 1 month will not be approved.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authorisation under this criterion should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

bosentan 62.5 mg tablet, 60

6429J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	1160.47	^a Bosentan APO [GX] ^a BOSENTAN DR. REDDY'S [RI] ^a Bosentan RBX [RA] ^a BOSLEER [RW]	^a BOSENTAN-DRLA [RZ] ^a Bosentan Mylan [AF] ^a Bosentan Sandoz [SZ] ^a Tracleer [AT]

■ EPOPROSTENOL

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 1 (new patients)

Clinical criteria:

- Patient must not have received prior PBS-subsidised treatment with a pulmonary arterial hypertension (PAH) agent, **AND**

- Patient must have been assessed by a physician at a designated hospital, **AND**
- Patient must have WHO Functional Class IV idiopathic pulmonary arterial hypertension (iPAH), or anorexigen-induced PAH or hereditary PAH; OR
- Patient must have WHO Functional Class IV pulmonary arterial hypertension secondary to connective tissue disease, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6 Minute Walk Test (6MWT); and
- (3) a signed patient acknowledgement.

Idiopathic pulmonary arterial hypertension, anorexigen-induced pulmonary arterial hypertension, hereditary pulmonary arterial hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) are defined as follows:

- (i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary artery wedge pressure (PAWP) less than 15 mmHg; or
- (ii) where a right heart catheter (RHC) cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.

Test requirements to establish baseline for initiation of treatment are as follows:

The first written application for PBS-subsidised treatment with the first PAH agent should be accompanied by the results of a right heart catheter (RHC) composite assessment plus an echocardiograph (ECHO) composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.

Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments;
- (2) RHC composite assessment plus 6MWT;
- (3) RHC composite assessment only.

In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted for consideration based on the results of the following test combinations, which are listed in descending order of preference:

- (1) ECHO composite assessment plus 6MWT;
- (2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats may be requested.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 2 (change or re-commencement of therapy for all patients)

Clinical criteria:

- Patient must have idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH or PAH secondary to connective tissue disease and must wish to re-commence PBS-subsidised therapy with this agent after a break in therapy and must have demonstrated a response to their most recent course of PBS-subsidised treatment with this agent; OR

- Patient must have WHO Functional Class IV idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH or PAH secondary to connective tissue disease and must have received prior treatment with a PBS-subsidised PAH agent other than this agent; OR
- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH or PAH secondary to connective tissue disease and must have failed to respond to a prior PBS-subsidised PAH agent, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form; and
- (3) the results of the patient's response to treatment with their last course of PBS-subsidised PAH agent; and
- (4) for WHO Functional Class III patients, where this is the first application for this agent, assessment details of the PBS-subsidised PAH agent they have failed to respond to.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

Response to a PAH agent is defined as follows:

For patients with two or more baseline tests, response to treatment is defined as two or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats may be requested.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Swapping between PAH agents: Patients can access PAH agents through the PBS according to the relevant restrictions. Once these patients are approved initial treatment with 1 of these 8 drugs, they may swap between PAH agents at any time without having to re-qualify for treatment with the alternate agent. This means that patients may commence treatment with the alternate agent, subject to that agent's restriction, irrespective of the severity of their disease at the time the application to swap therapy is submitted. It also means that no new baseline measurements will be necessary. New baselines may be submitted where the patient has failed to respond to their current treatment. Eligible patients may only swap between PAH agents if they have not failed prior PBS-subsidised treatment with that agent. For eligible patients, applications to swap between the 8 PAH agents must be made under the relevant initial treatment restriction. Patients should be assessed for response to the treatment they are ceasing at the time the application to swap therapy is being made. Patients who fail to demonstrate a response or for whom no assessment results are submitted with the application to swap therapy may not re-commence PBS-subsidised treatment with the drug they are ceasing.

Note Applications for patients who wish to swap to an alternate PAH agent should be accompanied by the previously approved authority prescription, or remaining repeats, for the treatment the patient is ceasing.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Caution This is a category X drug and must not be given to pregnant women. Pregnancy must be avoided during treatment and for at least 3 months following cessation of therapy.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 1 (new patients) or Initial 2 (change or re-commencement of therapy for all patients) or First Continuing treatment - Balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this agent under the Initial 1 (new patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the Initial 2 (change or re-commencement of therapy for all patients) restriction to complete a maximum of six months of treatment; OR

- Patient must have received insufficient therapy with this agent under the First Continuing treatment restriction to complete a maximum of six months of treatment, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition, **AND**
- The treatment must provide no more than the balance of up to six months treatment available under one of the above restrictions.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authorisation under this criterion should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: First Continuing treatment

Clinical criteria:

- Patient must have received a PBS-subsidised initial course of treatment with this agent for this condition, **AND**
- Patient must have been assessed by a physician from a designated hospital to have achieved a response to the PBS-subsidised initial course of treatment, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

(1) a completed authority prescription form; and

(2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:

(i) RHC composite assessment; and

(ii) ECHO composite assessment; and

(iii) 6 Minute Walk Test (6MWT).

Test requirements to establish response to treatment for continuation of treatment are as follows:

The following list outlines the preferred test combination, in descending order, for the purposes of continuation of PBS-subsidised treatment:

(1) RHC plus ECHO composite assessments plus 6MWT;

(2) RHC plus ECHO composite assessments;

(3) RHC composite assessment plus 6MWT;

(4) ECHO composite assessment plus 6MWT;

(5) RHC composite assessment only;

(6) ECHO composite assessment only.

The results of the same tests as conducted at baseline should be provided with the written First Continuing treatment application, except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a patient specific reason why the test(s) could not be conducted must be provided with the application.

The test results provided with the application for continuing treatment must be no more than 2 months old at the time of application.

Response to a PAH agent is defined as follows:

For patients with two or more baseline tests, response to treatment is defined as two or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats will be authorised.

An application for First Continuing treatment with a PAH agent should be made prior to the completion of the Initial 6 month treatment course to ensure continuity for those patients who respond to treatment, as assessed by the treating physician.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Subsequent Continuing treatment

Clinical criteria:

- Patient must have received a PBS-subsidised treatment under First Continuing treatment with this agent for this condition; OR
- Patient must have previously received PBS-subsidised treatment under this criteria with this agent for this condition, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats will be authorised.

An application for Subsequent Continuing treatment with a PAH agents should be made prior to the completion of the First Continuing treatment course to ensure continuity of treatment.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authorisation under this criterion should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

epoprostenol 500 microgram injection, 1 vial

10111E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	44.00	Velettri [AT]

epoprostenol 1.5 mg injection [1 vial] (&) inert substance diluent [2 x 50 mL vials], 1 pack

11082G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	70.72	Flolan [GK]

epoprostenol 1.5 mg injection, 1 vial

10129D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	70.72	Velettri [AT]

epoprostenol 500 microgram injection [1 vial] (&) inert substance diluent [2 x 50 mL vials], 1 pack

11069N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	40.03	Flolan [GK]

■ ILOPROST

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 1 (new patients)

Clinical criteria:

- Patient must not have received prior PBS-subsidised treatment with this agent, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**
- Patient must have WHO Functional Class III drug-induced PAH, **AND**
- Patient must have a mean right atrial pressure of 8 mmHg or less as measured by right heart catheterisation (RHC); OR
- Patient must have right ventricular function assessed by echocardiography (ECHO) where a RHC cannot be performed on clinical grounds, **AND**
- Patient must have failed to respond to 6 or more weeks of appropriate vasodilator treatment unless intolerance or a contraindication to such treatment exists, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

(1) a completed authority prescription form; and

(2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:

(i) RHC composite assessment; and

(ii) ECHO composite assessment; and

- (iii) 6 Minute Walk Test (6MWT); and
- (3) a signed patient acknowledgement.

Idiopathic pulmonary arterial hypertension, anorexigen-induced pulmonary arterial hypertension, hereditary pulmonary arterial hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) are defined as follows:

- (i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary artery wedge pressure (PAWP) less than 15 mmHg; or
- (ii) where a right heart catheter (RHC) cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.

Test requirements to establish baseline for initiation of treatment are as follows:

The first written application for PBS-subsidised treatment with the first PAH agent should be accompanied by the results of a right heart catheter (RHC) composite assessment plus an echocardiograph (ECHO) composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.

Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments;
- (2) RHC composite assessment plus 6MWT;
- (3) RHC composite assessment only.

In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted for consideration based on the results of the following test combinations, which are listed in descending order of preference:

- (1) ECHO composite assessment plus 6MWT;
- (2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

Details of prior vasodilator treatment, including the dose and duration of treatment, must be provided at the time of application. Where the patient has an adverse event to a vasodilator or where vasodilator treatment is contraindicated, details of the nature of the adverse event or contraindication according to the Therapeutic Goods Administration (TGA) approved Product Information must also be provided with the application.

Response to prior vasodilator treatment is defined as follows:

For patients with 2 or more baseline tests, response to treatment is defined as 2 or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information.

A maximum of 5 repeats may be requested.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Note Special Pricing Arrangements apply.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 2 (new patients)

Clinical criteria:

- Patient must not have received prior PBS-subsidised treatment with a pulmonary arterial hypertension (PAH) agent, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**

- Patient must have WHO Functional Class III drug-induced PAH and a mean right atrial pressure greater than 8 mmHg, as measured by right heart catheterisation (RHC); OR
- Patient must have WHO Functional Class III drug-induced PAH with right ventricular function assessed by echocardiography (ECHO) where a RHC cannot be performed on clinical grounds; OR
- Patient must have WHO Functional Class IV idiopathic pulmonary arterial hypertension (iPAH), or anorexigen-induced PAH or hereditary PAH; OR
- Patient must have WHO Functional Class IV pulmonary arterial hypertension secondary to connective tissue disease; OR
- Patient must have WHO Functional Class IV drug-induced PAH, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6 Minute Walk Test (6MWT); and
- (3) a signed patient acknowledgement.

Idiopathic pulmonary arterial hypertension, anorexigen-induced pulmonary arterial hypertension, hereditary pulmonary arterial hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) are defined as follows:

- (i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary artery wedge pressure (PAWP) less than 15 mmHg; or
- (ii) where a right heart catheter (RHC) cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.

Test requirements to establish baseline for initiation of treatment are as follows:

The first written application for PBS-subsidised treatment with the first PAH agent should be accompanied by the results of a right heart catheter (RHC) composite assessment plus an echocardiograph (ECHO) composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.

Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments;
- (2) RHC composite assessment plus 6MWT;
- (3) RHC composite assessment only.

In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted for consideration based on the results of the following test combinations, which are listed in descending order of preference:

- (1) ECHO composite assessment plus 6MWT;
- (2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats may be requested.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Note Special Pricing Arrangements apply.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 3 (change or re-commencement of therapy for all patients)

Clinical criteria:

- Patient must have idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH or PAH secondary to connective tissue disease or drug-induced PAH and must wish to re-commence PBS-subsidised therapy with this agent after a break in therapy and must have demonstrated a response to their most recent course of PBS-subsidised treatment with this agent; OR
- Patient must have WHO Functional Class IV idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH or PAH secondary to connective tissue disease and must have received prior treatment with a PBS-subsidised PAH agent other than this agent; OR
- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH or PAH secondary to connective tissue disease and must have failed to respond to a prior PBS-subsidised PAH agent, **AND**

- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form; and
- (3) the results of the patient's response to treatment with their last course of PBS-subsidised PAH agent; and
- (4) for WHO Functional Class III patients, where this is the first application for this agent, assessment details of the PBS-subsidised PAH agent they have failed to respond to.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

Response to a PAH agent is defined as follows:

For patients with two or more baseline tests, response to treatment is defined as two or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats may be requested.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Swapping between PAH agents: Patients can access PAH agents through the PBS according to the relevant restrictions.

Once these patients are approved initial treatment with 1 of these 8 drugs, they may swap between PAH agents at any time without having to re-qualify for treatment with the alternate agent. This means that patients may commence treatment with the alternate agent, subject to that agent's restriction, irrespective of the severity of their disease at the time the application to swap therapy is submitted. It also means that no new baseline measurements will be necessary. New baselines may be submitted where the patient has failed to respond to their current treatment. Eligible patients may only swap between PAH agents if they have not failed prior PBS-subsidised treatment with that agent. For eligible patients, applications to swap between the 8 PAH agents must be made under the relevant initial treatment restriction. Patients should be assessed for response to the treatment they are ceasing at the time the application to swap therapy is being made. Patients who fail to demonstrate a response or for whom no assessment results are submitted with the application to swap therapy may not re-commence PBS-subsidised treatment with the drug they are ceasing.

Note Applications for patients who wish to swap to an alternate PAH agent should be accompanied by the previously approved authority prescription, or remaining repeats, for the treatment the patient is ceasing.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Note Special Pricing Arrangements apply.

Caution This is a category X drug and must not be given to pregnant women. Pregnancy must be avoided during treatment and for at least 3 months following cessation of therapy.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 1 or Initial 2 (new patients) or Initial 3 (change or re-commencement of therapy for all patients) or First Continuing treatment - Balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this agent under the Initial 1 (new patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the Initial 2 (new patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the Initial 3 (change or re-commencement of therapy for all patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the First Continuing treatment restriction to complete a maximum of six months of treatment, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition, **AND**
- The treatment must provide no more than the balance of up to six months treatment available under one of the above restrictions.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authorisation under this criterion should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: First Continuing treatment

Clinical criteria:

- Patient must have received a PBS-subsidised initial course of treatment with this agent for this condition, **AND**
- Patient must have been assessed by a physician from a designated hospital to have achieved a response to the PBS-subsidised initial course of treatment, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

(1) a completed authority prescription form; and

(2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:

(i) RHC composite assessment; and

(ii) ECHO composite assessment; and

(iii) 6 Minute Walk Test (6MWT).

Test requirements to establish response to treatment for continuation of treatment are as follows:

The following list outlines the preferred test combination, in descending order, for the purposes of continuation of PBS-subsidised treatment:

(1) RHC plus ECHO composite assessments plus 6MWT;

(2) RHC plus ECHO composite assessments;

(3) RHC composite assessment plus 6MWT;

(4) ECHO composite assessment plus 6MWT;

(5) RHC composite assessment only;

(6) ECHO composite assessment only.

The results of the same tests as conducted at baseline should be provided with the written First Continuing treatment application, except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a patient specific reason why the test(s) could not be conducted must be provided with the application.

The test results provided with the application for continuing treatment must be no more than 2 months old at the time of application.

Response to a PAH agent is defined as follows:

For patients with two or more baseline tests, response to treatment is defined as two or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats will be authorised.

An application for First Continuing treatment with a PAH agent should be made prior to the completion of the Initial 6 month treatment course to ensure continuity for those patients who respond to treatment, as assessed by the treating physician.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Subsequent Continuing treatment

Clinical criteria:

- Patient must have received a PBS-subsidised treatment under First Continuing treatment with this agent for this condition; OR
- Patient must have previously received PBS-subsidised treatment under this criteria with this agent for this condition, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats will be authorised.

An application for Subsequent Continuing treatment with a PAH agents should be made prior to the completion of the First Continuing treatment course to ensure continuity of treatment.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authorisation under this criterion should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

iloprost 20 microgram/2 mL inhalation solution, 30 x 2 mL ampoules

6456T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	390.10	Ventavis [BN]

▪ **MACITENTAN**

Caution This is a category X drug and must not be given to pregnant women. Pregnancy must be avoided during treatment and for at least 3 months following cessation of therapy.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 1 (new patients)

Clinical criteria:

- Patient must not have received prior PBS-subsidised treatment with a pulmonary arterial hypertension (PAH) agent, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**
- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH; OR
- Patient must have WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease, **AND**
- Patient must have a mean right atrial pressure of 8 mmHg or less as measured by right heart catheterisation (RHC); OR
- Patient must have right ventricular function assessed by echocardiography (ECHO) where a RHC cannot be performed on clinical grounds, **AND**
- Patient must have failed to respond to 6 or more weeks of appropriate vasodilator treatment unless intolerance or a contraindication to such treatment exists, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

(1) a completed authority prescription form; and

(2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:

(i) RHC composite assessment; and

(ii) ECHO composite assessment; and

- (iii) 6 Minute Walk Test (6MWT); and
- (3) a signed patient acknowledgement.

Idiopathic pulmonary arterial hypertension, anorexigen-induced pulmonary arterial hypertension, hereditary pulmonary arterial hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) are defined as follows:

(i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary artery wedge pressure (PAWP) less than 15 mmHg; or

(ii) where a right heart catheter (RHC) cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.

Test requirements to establish baseline for initiation of treatment are as follows:

The first written application for PBS-subsidised treatment with the first PAH agent should be accompanied by the results of a right heart catheter (RHC) composite assessment plus an echocardiograph (ECHO) composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.

Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments;
- (2) RHC composite assessment plus 6MWT;
- (3) RHC composite assessment only.

In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted for consideration based on the results of the following test combinations, which are listed in descending order of preference:

- (1) ECHO composite assessment plus 6MWT;
- (2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

Details of prior vasodilator treatment, including the dose and duration of treatment, must be provided at the time of application. Where the patient has an adverse event to a vasodilator or where vasodilator treatment is contraindicated, details of the nature of the adverse event or contraindication according to the Therapeutic Goods Administration (TGA) approved Product Information must also be provided with the application.

Response to prior vasodilator treatment is defined as follows:

For patients with 2 or more baseline tests, response to treatment is defined as 2 or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information.

A maximum of 5 repeats may be requested.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Prior Written Approval of Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 2 (new patients)

Clinical criteria:

- Patient must not have received prior PBS-subsidised treatment with a pulmonary arterial hypertension (PAH) agent, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**
- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH), or anorexigen-induced PAH or hereditary PAH, and a mean right atrial pressure of greater than 8 mmHg, as measured by right heart catheterisation (RHC); OR

- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH), or anorexigen-induced PAH or hereditary PAH, with right ventricular function assessed by echocardiography (ECHO) where a RHC cannot be performed on clinical grounds; OR
- Patient must have WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease and a mean right atrial pressure greater than 8 mmHg, as measured by RHC; OR
- Patient must have WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease with right ventricular function assessed by ECHO where a RHC cannot be performed on clinical grounds; OR
- Patient must have WHO Functional Class IV idiopathic pulmonary arterial hypertension (iPAH), or anorexigen-induced PAH or hereditary PAH; OR
- Patient must have WHO Functional Class IV pulmonary arterial hypertension secondary to connective tissue disease; OR
- Patient must have WHO Functional Class III or IV pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology), **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6 Minute Walk Test (6MWT); and
- (3) a signed patient acknowledgement.

Idiopathic pulmonary arterial hypertension, anorexigen-induced pulmonary arterial hypertension, hereditary pulmonary arterial hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) are defined as follows:

- (i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary artery wedge pressure (PAWP) less than 15 mmHg; or
- (ii) where a right heart catheter (RHC) cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.

Test requirements to establish baseline for initiation of treatment are as follows:

The first written application for PBS-subsidised treatment with the first PAH agent should be accompanied by the results of a right heart catheter (RHC) composite assessment plus an echocardiograph (ECHO) composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.

Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments;
- (2) RHC composite assessment plus 6MWT;
- (3) RHC composite assessment only.

In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted for consideration based on the results of the following test combinations, which are listed in descending order of preference:

- (1) ECHO composite assessment plus 6MWT;
- (2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats may be requested.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Prior Written Approval of Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 3 (change or re-commencement of therapy for all patients)

Clinical criteria:

- Patient must have idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH or PAH secondary to connective tissue disease or PAH associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) and must wish to re-commence PBS-subsidised therapy with this agent after a break in therapy and must have demonstrated a response to their most recent course of PBS-subsidised treatment with this agent; OR
- Patient must have idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH or PAH secondary to connective tissue disease or PAH associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) and whose most recent course of PBS-subsidised treatment was with a PAH agent other than this agent, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form; and
- (3) the results of the patient's response to treatment with their last course of PBS-subsidised PAH agent.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

Response to a PAH agent is defined as follows:

For patients with two or more baseline tests, response to treatment is defined as two or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

A maximum of 5 repeats will be authorised.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Swapping between PAH agents: Patients can access PAH agents through the PBS according to the relevant restrictions. Once these patients are approved initial treatment with 1 of these 8 drugs, they may swap between PAH agents at any time without having to re-qualify for treatment with the alternate agent. This means that patients may commence treatment with the alternate agent, subject to that agent's restriction, irrespective of the severity of their disease at the time the application to swap therapy is submitted. It also means that no new baseline measurements will be necessary. New baselines may be submitted where the patient has failed to respond to their current treatment. Eligible patients may only swap between PAH agents if they have not failed prior PBS-subsidised treatment with that agent. For eligible patients, applications to swap between the 8 PAH agents must be made under the relevant initial treatment restriction. Patients should be assessed for response to the treatment they are ceasing at the time the application to swap therapy is being made. Patients who fail to demonstrate a response or for whom no assessment results are submitted with the application to swap therapy may not re-commence PBS-subsidised treatment with the drug they are ceasing.

Note Applications for patients who wish to swap to an alternate PAH agent should be accompanied by the previously approved authority prescription, or remaining repeats, for the treatment the patient is ceasing.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Prior Written Approval of Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 1 or Initial 2 (new patients) or Initial 3 (change or re-commencement of therapy for all patients) or

First Continuing treatment - Balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this agent under the Initial 1 (new patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the Initial 2 (new patients) restriction to complete a maximum of six months of treatment; OR

- Patient must have received insufficient therapy with this agent under the Initial 3 (change or re-commencement of therapy for all patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the First Continuing treatment restriction to complete a maximum of six months of treatment, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition, **AND**
- The treatment must provide no more than the balance of up to six months treatment available under one of the above restrictions.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authorisation under this criterion should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: First Continuing treatment

Clinical criteria:

- Patient must have received a PBS-subsidised initial course of treatment with this agent for this condition, **AND**
- Patient must have been assessed by a physician from a designated hospital to have achieved a response to the PBS-subsidised initial course of treatment, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

(1) a completed authority prescription form; and

(2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:

(i) RHC composite assessment; and

(ii) ECHO composite assessment; and

(iii) 6 Minute Walk Test (6MWT).

Test requirements to establish response to treatment for continuation of treatment are as follows:

The following list outlines the preferred test combination, in descending order, for the purposes of continuation of PBS-subsidised treatment:

(1) RHC plus ECHO composite assessments plus 6MWT;

(2) RHC plus ECHO composite assessments;

(3) RHC composite assessment plus 6MWT;

(4) ECHO composite assessment plus 6MWT;

(5) RHC composite assessment only;

(6) ECHO composite assessment only.

The results of the same tests as conducted at baseline should be provided with the written First Continuing treatment application, except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a patient specific reason why the test(s) could not be conducted must be provided with the application.

The test results provided with the application for continuing treatment must be no more than 2 months old at the time of application.

Response to a PAH agent is defined as follows:

For patients with two or more baseline tests, response to treatment is defined as two or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats will be authorised.

An application for First Continuing treatment with a PAH agent should be made prior to the completion of the Initial 6 month treatment course to ensure continuity for those patients who respond to treatment, as assessed by the treating physician.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available

on the Department of Human Services website at www.humanservices.gov.au
Applications for authority to prescribe should be forwarded to:
Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)
Treatment Phase: Subsequent Continuing treatment

Clinical criteria:

- Patient must have received a PBS-subsidised treatment under First Continuing treatment with this agent for this condition; OR
- Patient must have previously received PBS-subsidised treatment under this criteria with this agent for this condition, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats will be authorised.

An application for Subsequent Continuing treatment with a PAH agents should be made prior to the completion of the First Continuing treatment course to ensure continuity of treatment.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authorisation under this criterion should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

macitentan 10 mg tablet, 30

10134J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	2923.86	Opsumit [AT]

▪ **RIOCIGUAT**

Caution This is a category X drug and must not be given to pregnant women. Pregnancy must be avoided during treatment and for at least 1 month following cessation of therapy, as recommended by the TGA-approved Product Information.

Note Special Pricing Arrangements apply.

Authority required

Chronic thromboembolic pulmonary hypertension (CTEPH)
Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have WHO Functional Class II, III or IV CTEPH, **AND**
- The condition must be inoperable by pulmonary endarterectomy; OR
- The condition must be recurrent or persistent following pulmonary endarterectomy, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Treatment criteria:

- Must be treated in a centre with expertise in the management of CTEPH.

Population criteria:

- Patient must be aged 18 years or older.

CTEPH that is inoperable by pulmonary endarterectomy is defined as follows:

- Right heart catheterisation (RHC) demonstrating pulmonary vascular resistance (PVR) of greater than 300 dyn*sec*cm⁻⁵ measured at least 90 days after start of full anticoagulation; and
- A mean pulmonary artery pressure (PAPmean) of greater than 25 mmHg at least 90 days after start of full anticoagulation.

CTEPH that is recurrent or persistent subsequent to pulmonary endarterectomy is defined as follows:

- RHC demonstrating a PVR of greater than 300 dyn*sec*cm⁻⁵ measured at least 180 days following pulmonary endarterectomy.

Where a RHC cannot be performed due to right ventricular dysfunction, an echocardiogram demonstrating the dysfunction must be provided at the time of application.

Applications for authorisation must be in writing and must include:(1) completed authority prescription forms sufficient for dose titration; and(2) a completed CTEPH PBS Initial Authority Application - Supporting Information form which includes results from the 3 tests below, to establish baseline measurements, where available:(i) RHC composite assessment, and(ii) ECHO composite assessment, and(iii) 6 Minute Walk Test (6MWT); and(3) a signed patient acknowledgment form; and(4)

confirmation of evidence of inoperable CTEPH including results of a pulmonary vascular resistance (PVR), a mean pulmonary artery pressure (PAPmean) and the starting date of full anticoagulation; or(5) confirmation of evidence of recurrent or persistent CTEPH including result of PVR and the date that pulmonary endarterectomy was performed; or(6) confirmation of an echocardiogram demonstrating right ventricular dysfunction.

Where it is not possible to perform all 3 tests above on clinical grounds, applications may be submitted for consideration based on the results of the following test combinations, which are listed in descending order of preference:(1) RHC plus ECHO composite assessments;(2) RHC composite assessment plus 6MWT;(3) RHC composite assessment only.

In circumstance where a RHC cannot be performed on clinical grounds, applications may be submitted for consideration based on the results of the following test combinations, which are listed in descending order of preference:(1) ECHO composite assessment plus 6MWT;(2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

Prescriptions for dose titration must provide sufficient quantity for dose titrations by 0.5 mg increments at 2-week intervals to achieve up to a maximum of 2.5 mg three times daily based on the dosage recommendations for initiation of treatment in the TGA-approved Product Information. No repeats will be authorised for these prescriptions.

Approvals for subsequent authority prescription will be limited to 1 month of treatment. The quantity approved must be based on the dosage recommendations in the TGA-approved Product Information, and a maximum of 3 repeats.

The assessment of the patient's response to the initial 20-week course of treatment should be made following the preceding 16 weeks of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Chronic thromboembolic pulmonary hypertension (CTEPH)

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must demonstrate stable or responding disease, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Treatment criteria:

- Must be treated in a centre with expertise in the management of CTEPH.

Population criteria:

- Patient must be aged 18 years or older.

Applications for authorisation must be in writing and must include:(1) a completed authority prescription form; and(2) a completed CTEPH PBS Continuing Authority Application - Supporting Information form which includes results from the three tests below, where available:(i) RHC composite assessment; and(ii) ECHO composite assessment; and(iii) 6 Minute Walk Test (6MWT).

Test requirements to establish response to treatment for continuation of treatment are as follows:

The following list outlines the preferred test combination, in descending order, for the purposes of continuation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments plus 6MWT;
- (2) RHC plus ECHO composite assessments;
- (3) RHC composite assessment plus 6MWT;
- (4) ECHO composite assessment plus 6MWT;
- (5) RHC composite assessment only;
- (6) ECHO composite assessment only.

The results of the same tests as conducted at baseline should be provided with each written continuing treatment application (i.e., every 6 months), except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a patient specific reason why the test(s) could not be conducted must be provided with the application.

The test results provided with the application for continuing treatment must be no more than 2 months old at the time of application.

Response to this drug is defined as follows:

For patients with two or more baseline tests, response to treatment is defined as two or more tests demonstrating stability or improvement of disease.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease.

The assessment of the patient's response to the continuing 6 month courses of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

The maximum quantity per prescription must be based on the dosage recommendations in the TGA-approved Product Information and be limited to provide sufficient supply for 1 month of treatment.

A maximum of 5 repeats will be authorised.

Applications for continuing treatment with this drug should be made two weeks prior to the completion of the 6-month treatment course to ensure continuity for those patients who respond to treatment, as assessed by the treating physician.

Patients who fail to demonstrate disease stability or improvement to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Chronic thromboembolic pulmonary hypertension (CTEPH)

Treatment Phase: Balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the Initial treatment restriction to complete a maximum of 20 weeks of treatment; OR
- Patient must have received insufficient therapy with this drug under the Continuing treatment restriction to complete a maximum of 24 weeks of treatment, **AND**
- The treatment must provide no more than the balance of up to 20 or 24 weeks of treatment available under the above respective restriction, **AND**
- The treatment must be the sole PBS-subsidised agent for this condition.

Treatment criteria:

- Must be treated in a centre with expertise in the management of CTEPH.

Population criteria:

- Patient must be aged 18 years or older.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

riociguat 2.5 mg tablet, 42

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
10985E	1	..	..	1765.10	Adempas [BN]

riociguat 1 mg tablet, 84

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
11010L	1	..	..	3482.81	Adempas [BN]

riociguat 1 mg tablet, 42

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
10990K	1	..	..	1765.10	Adempas [BN]

riociguat 500 microgram tablet, 84

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
11008J	1	..	..	3482.81	Adempas [BN]

riociguat 2 mg tablet, 84

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
11017W	1	..	..	3482.81	Adempas [BN]

riociguat 2 mg tablet, 42

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
11012N	1	..	..	1765.10	Adempas [BN]

riociguat 500 microgram tablet, 42

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
11009K	1	..	..	1765.10	Adempas [BN]

riociguat 1.5 mg tablet, 84

10975P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	3482.81	Adempas [BN]

riociguat 2.5 mg tablet, 84

11018X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	3482.81	Adempas [BN]

riociguat 1.5 mg tablet, 42

10974N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	1765.10	Adempas [BN]

■ RIOCIQUAT

Caution This is a category X drug and must not be given to pregnant women. Pregnancy must be avoided during treatment and for at least 1 month following cessation of therapy, as recommended by the TGA-approved Product Information.

Note Special Pricing Arrangements apply.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 1 (new patients)

Clinical criteria:

- Patient must not have received prior PBS-subsidised treatment with a pulmonary arterial hypertension (PAH) agent, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**
- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH; OR
- Patient must have WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease, **AND**
- Patient must have a mean right atrial pressure of 8 mmHg or less as measured by right heart catheterisation (RHC); OR
- Patient must have right ventricular function assessed by echocardiography (ECHO) where a RHC cannot be performed on clinical grounds, **AND**
- Patient must have failed to respond to 6 or more weeks of appropriate vasodilator treatment unless intolerance or a contraindication to such treatment exists, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include: (1) completed authority prescription forms sufficient for dose titration; and (2) a completed Pulmonary Arterial Hypertension Initial PBS Authority Application - Supporting Information form which includes results from the three tests below, where available: (i) RHC composite assessment; and (ii) ECHO composite assessment; and (iii) 6 Minute Walk Test (6MWT); and (3) a signed patient acknowledgement. Idiopathic pulmonary arterial hypertension, anorexigen-induced pulmonary arterial hypertension, hereditary pulmonary arterial hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) are defined as follows: (i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary artery wedge pressure (PAWP) less than 15 mmHg; or (ii) where a right heart catheter (RHC) cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.

Test requirements to establish baseline for initiation of treatment are as follows:

The first written application for PBS-subsidised treatment with the first PAH agent should be accompanied by the results of a right heart catheter (RHC) composite assessment plus an echocardiograph (ECHO) composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.

Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments;
- (2) RHC composite assessment plus 6MWT;
- (3) RHC composite assessment only.

In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted for consideration based on the results of the following test combinations, which are listed in descending order of preference:

- (1) ECHO composite assessment plus 6MWT;
- (2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

Details of prior vasodilator treatment, including the dose and duration of treatment, must be provided at the time of application. Where the patient has an adverse event to a vasodilator or where vasodilator treatment is contraindicated, details of the nature of the adverse event or contraindication according to the Therapeutic Goods Administration (TGA) approved Product Information must also be provided with the application.

Response to prior vasodilator treatment is defined as follows:

For patients with 2 or more baseline tests, response to treatment is defined as 2 or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

Approvals for prescriptions for dose titration will provide sufficient quantity for dose titrations by 0.5 mg increments at 2-week intervals to achieve up to a maximum of 2.5 mg three times daily based on the dosage recommendations for initiation of treatment in the TGA-approved Product Information. No repeats will be authorised for these prescriptions.

Approvals for subsequent authority prescription will be limited to 1 month of treatment, with the quantity approved based on the dosage recommendations in the TGA-approved Product Information, and a maximum of 4 repeats.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 2 (new patients)

Clinical criteria:

- Patient must not have received prior PBS-subsidised treatment with a pulmonary arterial hypertension (PAH) agent, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**
- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH), or anorexigen-induced PAH or hereditary PAH, and a mean right atrial pressure of greater than 8 mmHg, as measured by right heart catheterisation (RHC); OR
- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH), or anorexigen-induced PAH or hereditary PAH, with right ventricular function assessed by echocardiography (ECHO) where a RHC cannot be performed on clinical grounds; OR
- Patient must have WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease and a mean right atrial pressure greater than 8 mmHg, as measured by RHC; OR
- Patient must have WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease with right ventricular function assessed by ECHO where a RHC cannot be performed on clinical grounds; OR
- Patient must have WHO Functional Class IV idiopathic pulmonary arterial hypertension (iPAH), or anorexigen-induced PAH or hereditary PAH; OR
- Patient must have WHO Functional Class IV pulmonary arterial hypertension secondary to connective tissue disease; OR
- Patient must have WHO Functional Class III or IV pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology), **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include: (1) completed authority prescription forms sufficient for dose titration; and (2) a completed Pulmonary Arterial Hypertension Initial PBS Authority Application - Supporting Information form which includes results from the three tests below, where available: (i) RHC composite assessment; and (ii) ECHO composite assessment; and (iii) 6 Minute Walk Test (6MWT); and (3) a signed patient acknowledgement. Idiopathic pulmonary arterial hypertension, anorexigen-induced pulmonary arterial hypertension, hereditary pulmonary arterial hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) are defined as follows: (i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary artery wedge pressure (PAWP) less than 15 mmHg; or (ii) where a right heart catheter (RHC) cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.

Test requirements to establish baseline for initiation of treatment are as follows:

The first written application for PBS-subsidised treatment with the first PAH agent should be accompanied by the results of a right heart catheter (RHC) composite assessment plus an echocardiograph (ECHO) composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.

Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments;
- (2) RHC composite assessment plus 6MWT;
- (3) RHC composite assessment only.

In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted for consideration based on the results of the following test combinations, which are listed in descending order of preference:

- (1) ECHO composite assessment plus 6MWT;
- (2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

Approvals for prescriptions for dose titration will provide sufficient quantity for dose titrations by 0.5 mg increments at 2-week intervals to achieve up to a maximum of 2.5 mg three times daily based on the dosage recommendations for initiation of treatment in the TGA-approved Product Information. No repeats will be authorised for these prescriptions.

Approvals for subsequent authority prescription will be limited to 1 month of treatment, with the quantity approved based on the dosage recommendations in the TGA-approved Product Information, and a maximum of 4 repeats.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 3 (change or re-commencement of therapy for all patients)

Clinical criteria:

- Patient must have idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH or PAH secondary to connective tissue disease or PAH associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) and must wish to re-commence PBS-subsidised therapy with this agent after a break in therapy and must have demonstrated a response to their most recent course of PBS-subsidised treatment with this agent; OR
- Patient must have idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH or PAH secondary to connective tissue disease or PAH associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) and whose most recent course of PBS-subsidised treatment was with a PAH agent other than this agent, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include: (1) completed authority prescription forms sufficient for dose titration; and (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form; and (3) the results of the patient's response to treatment with their last course of PBS-subsidised PAH agent. Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application. The test results provided must not be more than 2 months old at the time of application.

Response to a PAH agent is defined as follows:

For patients with two or more baseline tests, response to treatment is defined as two or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

Approvals for prescriptions for dose titration will provide sufficient quantity for dose titrations by 0.5 mg increments at 2-week intervals to achieve up to a maximum of 2.5 mg three times daily based on the dosage recommendations for initiation of treatment in the TGA-approved Product Information. No repeats will be authorised for these prescriptions.

Approvals for subsequent authority prescription will be limited to 1 month of treatment, with the quantity approved based on the dosage recommendations in the TGA-approved Product Information, and a maximum of 4 repeats.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Swapping between PAH agents: Patients can access PAH agents through the PBS according to the relevant restrictions. Once these patients are approved initial treatment with 1 of these 8 drugs, they may swap between PAH agents at any time without having to re-qualify for treatment with the alternate agent. This means that patients may commence treatment with the alternate agent, subject to that agent's restriction, irrespective of the severity of their disease at the time the application to swap therapy is submitted. It also means that no new baseline measurements will be necessary. New baselines may be submitted where the patient has failed to respond to their current treatment. Eligible patients may only swap between PAH agents if they have not failed prior PBS-subsidised treatment with that agent. For eligible patients, applications to swap between the 8 PAH agents must be made under the relevant initial treatment restriction. Patients should be assessed for response to the treatment they are ceasing at the time the application to swap therapy is being made. Patients who fail to demonstrate a response or for whom no assessment results are submitted with the application to swap therapy may not re-commence PBS-subsidised treatment with the drug they are ceasing.

Note Applications for patients who wish to swap to an alternate PAH agent should be accompanied by the previously approved authority prescription, or remaining repeats, for the treatment the patient is ceasing.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 1 or Initial 2 (new patients) or Initial 3 (change or re-commencement of therapy for all patients) or First Continuing treatment - Balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this agent under the Initial 1 (new patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the Initial 2 (new patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the Initial 3 (change or re-commencement of therapy for all patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the First Continuing treatment restriction to complete a maximum of six months of treatment, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition, **AND**
- The treatment must provide no more than the balance of up to six months treatment available under one of the above restrictions.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: First Continuing treatment

Clinical criteria:

- Patient must have received a PBS-subsidised initial course of treatment with this agent for this condition, **AND**
- Patient must have been assessed by a physician from a designated hospital to have achieved a response to the PBS-subsidised initial course of treatment, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:

- (i) RHC composite assessment; and
- (ii) ECHO composite assessment; and
- (iii) 6 Minute Walk Test (6MWT).

Test requirements to establish response to treatment for continuation of treatment are as follows:

The following list outlines the preferred test combination, in descending order, for the purposes of continuation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments plus 6MWT;
- (2) RHC plus ECHO composite assessments;
- (3) RHC composite assessment plus 6MWT;
- (4) ECHO composite assessment plus 6MWT;
- (5) RHC composite assessment only;
- (6) ECHO composite assessment only.

The results of the same tests as conducted at baseline should be provided with the written First Continuing treatment application, except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a patient specific reason why the test(s) could not be conducted must be provided with the application.

The test results provided with the application for continuing treatment must be no more than 2 months old at the time of application.

Response to a PAH agent is defined as follows:

For patients with two or more baseline tests, response to treatment is defined as two or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats will be authorised.

An application for First Continuing treatment with a PAH agent should be made two weeks prior to the completion of the Initial 6 month treatment course to ensure continuity for those patients who respond to treatment, as assessed by the treating physician.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Subsequent Continuing treatment

Clinical criteria:

- Patient must have received a PBS-subsidised treatment under First Continuing treatment with this agent for this condition; OR
- Patient must have previously received PBS-subsidised treatment under this criteria with this agent for this condition, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats will be authorised.

An application for Subsequent Continuing treatment with a PAH agents should be made prior to the completion of the First Continuing treatment course to ensure continuity of treatment.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

riociguat 2.5 mg tablet, 42

11052Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	1765.10	Adempas [BN]

riociguat 1 mg tablet, 84

11060D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	3482.81	Adempas [BN]

riociguat 1 mg tablet, 42

11028K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	1765.10	Adempas [BN]

riociguat 500 microgram tablet, 84

11058B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	3482.81	Adempas [BN]

riociguat 2 mg tablet, 84

11030M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	3482.81	Adempas [BN]

riociguat 2 mg tablet, 42

11045H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	1765.10	Adempas [BN]

riociguat 500 microgram tablet, 42

11031N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	1765.10	Adempas [BN]

riociguat 1.5 mg tablet, 84

11061E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	3482.81	Adempas [BN]

riociguat 2.5 mg tablet, 84

11035T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	3482.81	Adempas [BN]

riociguat 1.5 mg tablet, 42

11046J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	1765.10	Adempas [BN]

▪ **SILDENAFIL**

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 1 (new patients)

Clinical criteria:

- Patient must not have received prior PBS-subsidised treatment with a pulmonary arterial hypertension (PAH) agent, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**
- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH; OR
- Patient must have WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease, **AND**
- Patient must have a mean right atrial pressure of 8 mmHg or less as measured by right heart catheterisation (RHC); OR
- Patient must have right ventricular function assessed by echocardiography (ECHO) where a RHC cannot be performed on clinical grounds, **AND**
- Patient must have failed to respond to 6 or more weeks of appropriate vasodilator treatment unless intolerance or a contraindication to such treatment exists, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

(1) a completed authority prescription form; and

(2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:

(i) RHC composite assessment; and

(ii) ECHO composite assessment; and

(iii) 6 Minute Walk Test (6MWT); and

(3) a signed patient acknowledgement.

Idiopathic pulmonary arterial hypertension, anorexigen-induced pulmonary arterial hypertension, hereditary pulmonary arterial hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) are defined as follows:

(i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary artery wedge pressure (PAWP) less than 15 mmHg; or

(ii) where a right heart catheter (RHC) cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.

Test requirements to establish baseline for initiation of treatment are as follows:

The first written application for PBS-subsidised treatment with the first PAH agent should be accompanied by the results of a right heart catheter (RHC) composite assessment plus an echocardiograph (ECHO) composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.

Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments;
- (2) RHC composite assessment plus 6MWT;
- (3) RHC composite assessment only.

In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted for consideration based on the results of the following test combinations, which are listed in descending order of preference:

- (1) ECHO composite assessment plus 6MWT;
- (2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

Details of prior vasodilator treatment, including the dose and duration of treatment, must be provided at the time of application. Where the patient has an adverse event to a vasodilator or where vasodilator treatment is contraindicated, details of the nature of the adverse event or contraindication according to the Therapeutic Goods Administration (TGA) approved Product Information must also be provided with the application.

Response to prior vasodilator treatment is defined as follows:

For patients with 2 or more baseline tests, response to treatment is defined as 2 or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information.

A maximum of 5 repeats may be requested.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 2 (new patients)

Clinical criteria:

- Patient must not have received prior PBS-subsidised treatment with a pulmonary arterial hypertension (PAH) agent, **AND**
 - Patient must have been assessed by a physician at a designated hospital, **AND**
 - Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH), or anorexigen-induced PAH or hereditary PAH, and a mean right atrial pressure of greater than 8 mmHg, as measured by right heart catheterisation (RHC); OR
 - Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH), or anorexigen-induced PAH or hereditary PAH, with right ventricular function assessed by echocardiography (ECHO) where a RHC cannot be performed on clinical grounds; OR
 - Patient must have WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease and a mean right atrial pressure greater than 8 mmHg, as measured by RHC; OR
 - Patient must have WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease with right ventricular function assessed by ECHO where a RHC cannot be performed on clinical grounds, **AND**
 - The treatment must be the sole PBS-subsidised PAH agent for this condition.
- Applications for authorisation must be in writing and must include:
- (1) a completed authority prescription form; and

(2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:

- (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6 Minute Walk Test (6MWT); and
- (3) a signed patient acknowledgement.

Idiopathic pulmonary arterial hypertension, anorexigen-induced pulmonary arterial hypertension, hereditary pulmonary arterial hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) are defined as follows:

(i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary artery wedge pressure (PAWP) less than 15 mmHg; or

(ii) where a right heart catheter (RHC) cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.

Test requirements to establish baseline for initiation of treatment are as follows:

The first written application for PBS-subsidised treatment with the first PAH agent should be accompanied by the results of a right heart catheter (RHC) composite assessment plus an echocardiograph (ECHO) composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.

Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments;
- (2) RHC composite assessment plus 6MWT;
- (3) RHC composite assessment only.

In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted for consideration based on the results of the following test combinations, which are listed in descending order of preference:

- (1) ECHO composite assessment plus 6MWT;
- (2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats may be requested.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 3 (change or re-commencement of therapy for all patients)

Clinical criteria:

- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH or PAH secondary to connective tissue disease and must wish to re-commence PBS-subsidised therapy with this agent after a break in therapy and must have demonstrated a response to their most recent course of PBS-subsidised treatment with this agent; OR
- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH or PAH secondary to connective tissue disease and whose most recent course of PBS-subsidised treatment was with a PAH agent other than this agent, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form; and
- (3) the results of the patient's response to treatment with their last course of PBS-subsidised PAH agent.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

Response to a PAH agent is defined as follows:

For patients with two or more baseline tests, response to treatment is defined as two or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats may be requested.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Swapping between PAH agents: Patients can access PAH agents through the PBS according to the relevant restrictions.

Once these patients are approved initial treatment with 1 of these 8 drugs, they may swap between PAH agents at any time without having to re-qualify for treatment with the alternate agent. This means that patients may commence treatment with the alternate agent, subject to that agent's restriction, irrespective of the severity of their disease at the time the application to swap therapy is submitted. It also means that no new baseline measurements will be necessary. New baselines may be submitted where the patient has failed to respond to their current treatment. Eligible patients may only swap between PAH agents if they have not failed prior PBS-subsidised treatment with that agent. For eligible patients, applications to swap between the 8 PAH agents must be made under the relevant initial treatment restriction. Patients should be assessed for response to the treatment they are ceasing at the time the application to swap therapy is being made. Patients who fail to demonstrate a response or for whom no assessment results are submitted with the application to swap therapy may not re-commence PBS-subsidised treatment with the drug they are ceasing.

Note Applications for patients who wish to swap to an alternate PAH agent should be accompanied by the previously approved authority prescription, or remaining repeats, for the treatment the patient is ceasing.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Caution This is a category X drug and must not be given to pregnant women. Pregnancy must be avoided during treatment and for at least 3 months following cessation of therapy.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 1 or Initial 2 (new patients) or Initial 3 (change or re-commencement of therapy for all patients) or

First Continuing treatment - Balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this agent under the Initial 1 (new patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the Initial 2 (new patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the Initial 3 (change or re-commencement of therapy for all patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the First Continuing treatment restriction to complete a maximum of six months of treatment, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition, **AND**
- The treatment must provide no more than the balance of up to six months treatment available under one of the above restrictions.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authorisation under this criterion should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: First Continuing treatment

Clinical criteria:

- Patient must have received a PBS-subsidised initial course of treatment with this agent for this condition, **AND**
- Patient must have been assessed by a physician from a designated hospital to have achieved a response to the PBS-subsidised initial course of treatment, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6 Minute Walk Test (6MWT).

Test requirements to establish response to treatment for continuation of treatment are as follows:

The following list outlines the preferred test combination, in descending order, for the purposes of continuation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments plus 6MWT;
- (2) RHC plus ECHO composite assessments;
- (3) RHC composite assessment plus 6MWT;
- (4) ECHO composite assessment plus 6MWT;
- (5) RHC composite assessment only;
- (6) ECHO composite assessment only.

The results of the same tests as conducted at baseline should be provided with the written First Continuing treatment application, except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a patient specific reason why the test(s) could not be conducted must be provided with the application.

The test results provided with the application for continuing treatment must be no more than 2 months old at the time of application.

Response to a PAH agent is defined as follows:

For patients with two or more baseline tests, response to treatment is defined as two or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats will be authorised.

An application for First Continuing treatment with a PAH agent should be made prior to the completion of the Initial 6 month treatment course to ensure continuity for those patients who respond to treatment, as assessed by the treating physician.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Subsequent Continuing treatment

Clinical criteria:

- Patient must have received a PBS-subsidised treatment under First Continuing treatment with this agent for this condition; **OR**
- Patient must have previously received PBS-subsidised treatment under this criteria with this agent for this condition, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**

- The treatment must be the sole PBS-subsidised PAH agent for this condition. The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats will be authorised.

An application for Subsequent Continuing treatment with a PAH agents should be made prior to the completion of the First Continuing treatment course to ensure continuity of treatment.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authorisation under this criterion should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

sildenafil 20 mg tablet, 90

9605M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	271.87	^a APO-Sildenafil PHT [TX] ^a Sildenafil AN PHT 20 [EA] ^a Sildenafil Sandoz PHT 20 [SZ]	^a Revatio [PF] ^a SILDENAFIL-DRx [RZ]

■ TADALAFIL

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 1 (new patients)

Clinical criteria:

- Patient must not have received prior PBS-subsidised treatment with a pulmonary arterial hypertension (PAH) agent, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**
- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH; OR
- Patient must have WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease, **AND**
- Patient must have a mean right atrial pressure of 8 mmHg or less as measured by right heart catheterisation (RHC); OR
- Patient must have right ventricular function assessed by echocardiography (ECHO) where a RHC cannot be performed on clinical grounds, **AND**
- Patient must have failed to respond to 6 or more weeks of appropriate vasodilator treatment unless intolerance or a contraindication to such treatment exists, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:

- (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6 Minute Walk Test (6MWT); and
- (3) a signed patient acknowledgement.

Idiopathic pulmonary arterial hypertension, anorexigen-induced pulmonary arterial hypertension, hereditary pulmonary arterial hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) are defined as follows:

(i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary artery wedge pressure (PAWP) less than 15 mmHg; or

(ii) where a right heart catheter (RHC) cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.

Test requirements to establish baseline for initiation of treatment are as follows:

The first written application for PBS-subsidised treatment with the first PAH agent should be accompanied by the results of a right heart catheter (RHC) composite assessment plus an echocardiograph (ECHO) composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.

Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments;
- (2) RHC composite assessment plus 6MWT;
- (3) RHC composite assessment only.

In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted for consideration based on the results of the following test combinations, which are listed in descending order of preference:

- (1) ECHO composite assessment plus 6MWT;
- (2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

Details of prior vasodilator treatment, including the dose and duration of treatment, must be provided at the time of application. Where the patient has an adverse event to a vasodilator or where vasodilator treatment is contraindicated, details of the nature of the adverse event or contraindication according to the Therapeutic Goods Administration (TGA) approved Product Information must also be provided with the application.

Response to prior vasodilator treatment is defined as follows:

For patients with 2 or more baseline tests, response to treatment is defined as 2 or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information.

A maximum of 5 repeats may be requested.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 2 (new patients)

Clinical criteria:

- Patient must not have received prior PBS-subsidised treatment with a pulmonary arterial hypertension (PAH) agent, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**
- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH), or anorexigen-induced PAH or hereditary PAH, and a mean right atrial pressure of greater than 8 mmHg, as measured by right heart catheterisation (RHC); OR
- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH), or anorexigen-induced PAH or hereditary PAH, with right ventricular function assessed by echocardiography (ECHO) where a RHC cannot be performed on clinical grounds; OR
- Patient must have WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease and a mean right atrial pressure greater than 8 mmHg, as measured by RHC; OR
- Patient must have WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease with right ventricular function assessed by ECHO where a RHC cannot be performed on clinical grounds, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6 Minute Walk Test (6MWT); and
- (3) a signed patient acknowledgement.

Idiopathic pulmonary arterial hypertension, anorexigen-induced pulmonary arterial hypertension, hereditary pulmonary arterial hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) are defined as follows:

(i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary artery wedge pressure (PAWP) less than 15 mmHg; or

(ii) where a right heart catheter (RHC) cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.

Test requirements to establish baseline for initiation of treatment are as follows:

The first written application for PBS-subsidised treatment with the first PAH agent should be accompanied by the results of a right heart catheter (RHC) composite assessment plus an echocardiograph (ECHO) composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.

Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments;
- (2) RHC composite assessment plus 6MWT;
- (3) RHC composite assessment only.

In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted for consideration based on the results of the following test combinations, which are listed in descending order of preference:

- (1) ECHO composite assessment plus 6MWT;
- (2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats may be requested.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 3 (change or re-commencement of therapy for all patients)

Clinical criteria:

- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH or PAH secondary to connective tissue disease and must wish to re-commence PBS-subsidised therapy with this agent after a break in therapy and must have demonstrated a response to their most recent course of PBS-subsidised treatment with this agent; OR
- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH or PAH secondary to connective tissue disease and whose most recent course of PBS-subsidised treatment was with a PAH agent other than this agent, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form; and
- (3) the results of the patient's response to treatment with their last course of PBS-subsidised PAH agent.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

Response to a PAH agent is defined as follows:

For patients with two or more baseline tests, response to treatment is defined as two or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats may be requested.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Swapping between PAH agents: Patients can access PAH agents through the PBS according to the relevant restrictions. Once these patients are approved initial treatment with 1 of these 8 drugs, they may swap between PAH agents at any time without having to re-qualify for treatment with the alternate agent. This means that patients may commence treatment with the alternate agent, subject to that agent's restriction, irrespective of the severity of their disease at the time the application to swap therapy is submitted. It also means that no new baseline measurements will be necessary. New baselines may be submitted where the patient has failed to respond to their current treatment. Eligible patients may only swap between PAH agents if they have not failed prior PBS-subsidised treatment with that agent. For eligible patients, applications to swap between the 8 PAH agents must be made under the relevant initial treatment restriction. Patients should be assessed for response to the treatment they are ceasing at the time the application to swap therapy is being made. Patients who fail to demonstrate a response or for whom no assessment results are submitted with the application to swap therapy may not re-commence PBS-subsidised treatment with the drug they are ceasing.

Note Applications for patients who wish to swap to an alternate PAH agent should be accompanied by the previously approved authority prescription, or remaining repeats, for the treatment the patient is ceasing.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Caution This is a category X drug and must not be given to pregnant women. Pregnancy must be avoided during treatment and for at least 3 months following cessation of therapy.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 1 or Initial 2 (new patients) or Initial 3 (change or re-commencement of therapy for all patients) or

First Continuing treatment - Balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this agent under the Initial 1 (new patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the Initial 2 (new patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the Initial 3 (change or re-commencement of therapy for all patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the First Continuing treatment restriction to complete a maximum of six months of treatment, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition, **AND**
- The treatment must provide no more than the balance of up to six months treatment available under one of the above restrictions.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authorisation under this criterion should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: First Continuing treatment

Clinical criteria:

- Patient must have received a PBS-subsidised initial course of treatment with this agent for this condition, **AND**
- Patient must have been assessed by a physician from a designated hospital to have achieved a response to the PBS-subsidised initial course of treatment, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and

(2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:

- (i) RHC composite assessment; and
- (ii) ECHO composite assessment; and
- (iii) 6 Minute Walk Test (6MWT).

Test requirements to establish response to treatment for continuation of treatment are as follows:

The following list outlines the preferred test combination, in descending order, for the purposes of continuation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments plus 6MWT;
- (2) RHC plus ECHO composite assessments;
- (3) RHC composite assessment plus 6MWT;
- (4) ECHO composite assessment plus 6MWT;
- (5) RHC composite assessment only;
- (6) ECHO composite assessment only.

The results of the same tests as conducted at baseline should be provided with the written First Continuing treatment application, except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a patient specific reason why the test(s) could not be conducted must be provided with the application.

The test results provided with the application for continuing treatment must be no more than 2 months old at the time of application.

Response to a PAH agent is defined as follows:

For patients with two or more baseline tests, response to treatment is defined as two or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats will be authorised.

An application for First Continuing treatment with a PAH agent should be made prior to the completion of the Initial 6 month treatment course to ensure continuity for those patients who respond to treatment, as assessed by the treating physician.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Subsequent Continuing treatment

Clinical criteria:

- Patient must have received a PBS-subsidised treatment under First Continuing treatment with this agent for this condition; OR
- Patient must have previously received PBS-subsidised treatment under this criteria with this agent for this condition, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats will be authorised.

An application for Subsequent Continuing treatment with a PAH agents should be made prior to the completion of the First Continuing treatment course to ensure continuity of treatment.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

SYSTEMIC HORMONAL PREPARATIONS, EXCL. SEX HORMONES AND INSULINS

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authorisation under this criterion should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

tadalafil 20 mg tablet, 56

1304P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	835.85	Adcirca [LY]

SYSTEMIC HORMONAL PREPARATIONS, EXCL. SEX HORMONES AND INSULINS

PITUITARY AND HYPOTHALAMIC HORMONES AND ANALOGUES

ANTERIOR PITUITARY LOBE HORMONES AND ANALOGUES

Other anterior pituitary lobe hormones and analogues

PEGVISOMANT

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs Programs

Reply Paid 9826

HOBART TAS 7001

Authority required

Acromegaly

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must not have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have an age- and sex-adjusted insulin-like growth factor 1 (IGF-1) concentration greater than the upper limit of normal (ULN), **AND**
- The treatment must be after failure to achieve biochemical control with a maximum indicated dose of either 30 mg octreotide LAR or 120 mg lanreotide ATG every 28 days for 24 weeks; unless contraindicated or not tolerated according to the TGA approved Product Information, **AND**
- The treatment must not be given concomitantly with a PBS-subsidised somatostatin analogue.

Somatostatin analogues include octreotide, lanreotide and pasireotide

Failure to achieve biochemical control after completion of a prior therapy with either octreotide or lanreotide is defined as:

1) Growth hormone level greater than 1 mcg/L or 3 mIU/L; OR

2) IGF-1 level is greater than the age- and sex-adjusted ULN.

If treatment with either octreotide or lanreotide is contraindicated according to the relevant TGA-approved Product Information, the application must provide details of contraindication.

If intolerance to either octreotide or lanreotide treatment developed during the relevant period of use which is of a severity to necessitate withdrawal of the treatment, the application must provide details of the nature and severity of this intolerance.

In a patient treated with radiotherapy, pegvisomant should be withdrawn every 2 years in the 10 years after completion of radiotherapy for assessment of remission. Pegvisomant should be withdrawn at least 8 weeks prior to the assessment of remission.

Biochemical evidence of remission is defined as normalisation of sex- and age- adjusted insulin-like growth factor 1 (IGF-1).

Two completed authority prescriptions should be submitted with the initial application for this drug. One prescription should be for the loading dose of 80 mg for a quantity of 4 vials of 20 mg with no repeats. The second prescription should be for subsequent doses, starting from 10 mg daily, and allowing dose adjustments in increments of 5 mg based on serum IGF-1 levels measured every 4 to 6 weeks in order to maintain the serum IGF-1 level within the age-adjusted normal range based on the dosage recommendations in the TGA-approved Product Information.

The authority application must be made in writing and must include:

- a) two completed authority prescription forms ; and
- b) a completed Acromegaly Pegvisomant initial PBS Authority Application - Supporting Information Form; and
- c) in a patient who has been previously treated with radiotherapy for this condition, the date of completion of radiotherapy, the date and result of IGF-1 levels taken at the most recent two yearly assessment in the 10 years after completion of radiotherapy; and
- d) a recent result of the IGF-1 level and the date of assessment ; and

e) demonstration of failure to achieve biochemical control after completion of a prior therapy with either octreotide or lanreotide

No increase in the maximum quantity or number of units may be authorised for the loading dose.

pegvisomant 20 mg injection [1 vial] (&) inert substance diluent [1 syringe], 1 pack

11166Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	4	..	..	*586.83	Somavert [PF]

■ PEGVISOMANT

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Acromegaly

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must not have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have an age- and sex-adjusted insulin-like growth factor 1 (IGF-1) concentration greater than the upper limit of normal (ULN), **AND**
- The treatment must be after failure to achieve biochemical control with a maximum indicated dose of either 30 mg octreotide LAR or 120 mg lanreotide ATG every 28 days for 24 weeks; unless contraindicated or not tolerated according to the TGA approved Product Information, **AND**
- The treatment must not be given concomitantly with a PBS-subsidised somatostatin analogue.

Somatostatin analogues include octreotide, lanreotide and pasireotide

Failure to achieve biochemical control after completion of a prior therapy with either octreotide or lanreotide is defined as:

1) Growth hormone level greater than 1 mcg/L or 3 mIU/L; OR

2) IGF-1 level is greater than the age- and sex-adjusted ULN.

If treatment with either octreotide or lanreotide is contraindicated according to the relevant TGA-approved Product Information, the application must provide details of contraindication.

If intolerance to either octreotide or lanreotide treatment developed during the relevant period of use which is of a severity to necessitate withdrawal of the treatment, the application must provide details of the nature and severity of this intolerance.

In a patient treated with radiotherapy, pegvisomant should be withdrawn every 2 years in the 10 years after completion of radiotherapy for assessment of remission. Pegvisomant should be withdrawn at least 8 weeks prior to the assessment of remission.

Biochemical evidence of remission is defined as normalisation of sex- and age- adjusted insulin-like growth factor 1 (IGF-1). Two completed authority prescriptions should be submitted with the initial application for this drug. One prescription should be for the loading dose of 80 mg for a quantity of 4 vials of 20 mg with no repeats. The second prescription should be for subsequent doses, starting from 10 mg daily, and allowing dose adjustments in increments of 5 mg based on serum IGF-1 levels measured every 4 to 6 weeks in order to maintain the serum IGF-1 level within the age-adjusted normal range based on the dosage recommendations in the TGA-approved Product Information.

The authority application must be made in writing and must include:

- a) two completed authority prescription forms ; and
- b) a completed Acromegaly Pegvisomant initial PBS Authority Application - Supporting Information Form; and
- c) in a patient who has been previously treated with radiotherapy for this condition, the date of completion of radiotherapy, the date and result of IGF-1 levels taken at the most recent two yearly assessment in the 10 years after completion of radiotherapy; and
- d) a recent result of the IGF-1 level and the date of assessment ; and
- e) demonstration of failure to achieve biochemical control after completion of a prior therapy with either octreotide or lanreotide

No increase in the maximum quantity or number of units may be authorised for the loading dose.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs Programs

Reply Paid 9826

HOBART TAS 7001

Authority required

Acromegaly

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
 - The treatment must not be given concomitantly with a PBS-subsidised somatostatin analogue, **AND**
 - The treatment must cease if IGF-1 is not lower after 3 months of pegvisomant treatment at the maximum tolerated dose.
- Somatostatin analogues include octreotide, lanreotide and pasireotide

SYSTEMIC HORMONAL PREPARATIONS, EXCL. SEX HORMONES AND INSULINS

In a patient treated with radiotherapy, pegvisomant should be withdrawn every 2 years in the 10 years after completion of radiotherapy for assessment of remission. Pegvisomant should be withdrawn at least 8 weeks prior to the assessment of remission.

Biochemical evidence of remission is defined as normalisation of sex- and age- adjusted insulin-like growth factor 1 (IGF-1). In a patient who has been previously treated with radiotherapy for this condition, the date of completion of radiotherapy must be provided; and a copy of IGF-1 level taken at the most recent two yearly assessment in the 10 years after completion of radiotherapy must be provided at the time of application.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

pegvisomant 10 mg injection [30 vials] (&) inert substance diluent [30 syringes], 1 pack

11167R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	4225.96	Somavert [PF]

pegvisomant 15 mg injection [30 vials] (&) inert substance diluent [30 syringes], 1 pack

11172B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	4225.96	Somavert [PF]

pegvisomant 20 mg injection [30 vials] (&) inert substance diluent [30 syringes], 1 pack

11174D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	4225.96	Somavert [PF]

HYPOTHALAMIC HORMONES

Somatostatin and analogues

■ LANREOTIDE

Authority required (STREAMLINED)

9225

Acromegaly

Clinical criteria:

- The condition must be active, **AND**
- Patient must have persistent elevation of mean growth hormone levels of greater than 2.5 micrograms per litre, **AND**
- The treatment must be after failure of other therapy including dopamine agonists; OR
- The treatment must be as interim treatment while awaiting the effects of radiotherapy and where treatment with dopamine agonists has failed; OR
- The treatment must be in a patient who is unfit for or unwilling to undergo surgery and where radiotherapy is contraindicated, **AND**
- The treatment must cease in a patient treated with radiotherapy if there is biochemical evidence of remission (normal IGF1) after lanreotide has been withdrawn for at least 4 weeks (6 weeks after the last dose), **AND**
- The treatment must cease if IGF1 is not lower after 3 months of treatment, **AND**
- The treatment must not be given concomitantly with PBS-subsidised pegvisomant.

In a patient treated with radiotherapy, lanreotide should be withdrawn every 2 years in the 10 years after radiotherapy for assessment of remission.

lanreotide 30 mg modified release injection [1 vial] (&) inert substance diluent [2 mL ampoule], 1 pack

6332G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	11	..	*1265.77	Somatuline LA [IS]

■ LANREOTIDE

Authority required (STREAMLINED)

9261

Acromegaly

Clinical criteria:

- The condition must be active, **AND**
- Patient must have persistent elevation of mean growth hormone levels of greater than 2.5 micrograms per litre, **AND**
- The treatment must be after failure of other therapy including dopamine agonists; OR
- The treatment must be as interim treatment while awaiting the effects of radiotherapy and where treatment with dopamine agonists has failed; OR
- The treatment must be in a patient who is unfit for or unwilling to undergo surgery and where radiotherapy is contraindicated, **AND**
- The treatment must cease in a patient treated with radiotherapy if there is biochemical evidence of remission (normal IGF1) after lanreotide has been withdrawn for at least 4 weeks (8 weeks after the last dose), **AND**
- The treatment must cease if IGF1 is not lower after 3 months of treatment, **AND**
- The treatment must not be given concomitantly with PBS-subsidised pegvisomant.

In a patient treated with radiotherapy, lanreotide should be withdrawn every 2 years in the 10 years after radiotherapy for assessment of remission.

Authority required (STREAMLINED)

9260

Functional carcinoid tumour

Clinical criteria:

- The condition must be causing intractable symptoms, **AND**
- Patient must have experienced on average over 1 week, 3 or more episodes per day of diarrhoea and/or flushing, which persisted despite the use of anti-histamines, anti-serotonin agents and anti-diarrhoea agents, **AND**
- Patient must be one in whom surgery or antineoplastic therapy has failed or is inappropriate, **AND**
- The treatment must cease if there is failure to produce a clinically significant reduction in the frequency and severity of symptoms after 3 months' therapy at a dose of 120 mg every 28 days.

Dosage and tolerance to the drug should be assessed regularly and the dosage should be titrated slowly downwards to determine the minimum effective dose.

lanreotide 120 mg/0.5 mL injection, 0.5 mL syringe

6425E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*3686.27	Somatuline Autogel [IS]

lanreotide 90 mg/0.5 mL injection, 0.5 mL syringe

6424D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*2955.25	Somatuline Autogel [IS]

lanreotide 60 mg/0.5 mL injection, 0.5 mL syringe

6423C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*2232.35	Somatuline Autogel [IS]

■ **LANREOTIDE**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

9323

Non-functional gastroenteropancreatic neuroendocrine tumour (GEP-NET)

Clinical criteria:

- The condition must be unresectable locally advanced disease or metastatic disease, **AND**
- The condition must be World Health Organisation (WHO) grade 1 or 2, **AND**
- The treatment must be as monotherapy.

Population criteria:

- Patient must be aged 18 years or older.

WHO grade 1 of GEP-NET is defined as a mitotic count (10HPF) of less than 2 and Ki-67 index (%) of less than or equal to 2.

WHO grade 2 of GEP-NET is defined as a mitotic count (10HPF) of 2-20 and Ki-67 index (%) of 3-20.

Lanreotide is not PBS-subsidised for use in combination with everolimus or sunitinib for this condition.

lanreotide 120 mg/0.5 mL injection, 0.5 mL syringe

11527Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*3686.27	Somatuline Autogel [IS]

■ **OCTREOTIDE**

Authority required (STREAMLINED)

9262

Acromegaly

Clinical criteria:

- The condition must be controlled with octreotide immediate release injections, **AND**
- The treatment must cease in a patient treated with radiotherapy if there is biochemical evidence of remission (normal IGF1) after octreotide has been withdrawn for at least 4 weeks (8 weeks after the last dose), **AND**
- The treatment must cease if IGF1 is not lower after 3 months of treatment, **AND**
- The treatment must not be given concomitantly with PBS-subsidised lanreotide or pegvisomant for this condition.

In a patient treated with radiotherapy, octreotide should be withdrawn every 2 years in the 10 years after radiotherapy for assessment of remission

Authority required (STREAMLINED)

9313

Functional carcinoid tumour

Clinical criteria:

- Patient must have achieved symptom control on octreotide immediate release injections, **AND**
- The treatment must cease if there is failure to produce a clinically significant reduction in the frequency and severity of symptoms after 3 months therapy at a dose of 30 mg every 28 days and having allowed adequate rescue therapy with octreotide immediate release injections.

Dosage and tolerance to the drug should be assessed regularly and the dosage should be titrated slowly downwards to determine the minimum effective dose.

Authority required (STREAMLINED)**9288**

Vasoactive intestinal peptide secreting tumour (VIPoma)

Clinical criteria:

- Patient must have achieved symptom control on octreotide immediate release injections, **AND**
- The treatment must cease if there is failure to produce a clinically significant reduction in the frequency and severity of symptoms after 3 months therapy at a dose of 30 mg every 28 days and having allowed adequate rescue therapy with octreotide immediate release injections.

Dosage and tolerance to the drug should be assessed regularly and the dosage should be titrated slowly downwards to determine the minimum effective dose.

octreotide 20 mg modified release injection [1 vial] (&) inert substance diluent [2 mL syringe], 1 pack

10549F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*3527.01	Sandostatin LAR [NV]

octreotide 10 mg modified release injection [1 vial] (&) inert substance diluent [2 mL syringe], 1 pack

10566D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*2661.11	Sandostatin LAR [NV]

octreotide 30 mg modified release injection [1 vial] (&) inert substance diluent [2 mL syringe], 1 pack

10558Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*4402.31	Sandostatin LAR [NV]

■ OCTREOTIDE**Authority required (STREAMLINED)****9233**

Acromegaly

Clinical criteria:

- The condition must be active, **AND**
 - Patient must have persistent elevation of mean growth hormone levels of greater than 2.5 micrograms per litre, **AND**
 - The treatment must be after failure of other therapy including dopamine agonists; OR
 - The treatment must be as interim treatment while awaiting the effects of radiotherapy and where treatment with dopamine agonists has failed; OR
 - The treatment must be in a patient who is unfit for or unwilling to undergo surgery and where radiotherapy is contraindicated, **AND**
 - The treatment must cease in a patient treated with radiotherapy if there is biochemical evidence of remission (normal IGF1) after octreotide has been withdrawn for at least 4 weeks, **AND**
 - The treatment must cease if IGF1 is not lower after 3 months of treatment at a dose of 100 micrograms 3 time daily, **AND**
 - The treatment must not be given concomitantly with PBS-subsidised lanreotide or pegvisomant for this condition.
- In a patient treated with radiotherapy, octreotide should be withdrawn every 2 years in the 10 years after radiotherapy for assessment of remission

Authority required (STREAMLINED)**9289**

Functional carcinoid tumour

Clinical criteria:

- The condition must be causing intractable symptoms, **AND**
- Patient must have experienced on average over 1 week, 3 or more episodes per day of diarrhoea and/or flushing, which persisted despite the use of anti-histamines, anti-serotonin agents and anti-diarrhoea agents, **AND**
- Patient must be one in whom surgery or antineoplastic therapy has failed or is inappropriate, **AND**
- The treatment must cease if there is failure to produce a clinically significant reduction in the frequency and severity of symptoms after 2 months' therapy.

Dosage and tolerance to the drug should be assessed regularly and the dosage should be titrated slowly downwards to determine the minimum effective dose.

Authority required (STREAMLINED)**9232**

Vasoactive intestinal peptide secreting tumour (VIPoma)

Clinical criteria:

- The condition must be causing intractable symptoms, **AND**
- Patient must have experienced on average over 1 week, 3 or more episodes per day of diarrhoea and/or flushing, which persisted despite the use of anti-histamines, anti-serotonin agents and anti-diarrhoea agents, **AND**
- Patient must be one in whom surgery or antineoplastic therapy has failed or is inappropriate, **AND**
- The treatment must cease if there is failure to produce a clinically significant reduction in the frequency and severity of symptoms after 2 months' therapy.

Dosage and tolerance to the drug should be assessed regularly and the dosage should be titrated slowly downwards to determine the minimum effective dose.

octreotide 500 microgram/mL injection, 5 x 1 mL ampoules

6229W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	18	11	..	*6241.87	^a Hospira Pty Limited [PF] ^a Octreotide (SUN) [RA]	^a Octreotide MaxRx [GQ] ^a Sandostatin 0.5 [NV]

octreotide 50 microgram/mL injection, 5 x 1 mL ampoules

6227R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	18	11	..	*651.25	^a Hospira Pty Limited [PF] ^a Octreotide (SUN) [RA]	^a Octreotide MaxRx [GQ] ^a Sandostatin 0.05 [NV]

octreotide 100 microgram/mL injection, 5 x 1 mL ampoules

6228T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	18	11	..	*1283.77	^a Hospira Pty Limited [PF] ^a Octreotide (SUN) [RA]	^a Octreotide MaxRx [GQ] ^a Sandostatin 0.1 [NV]

■ PASIREOTIDE

Caution Careful monitoring of patients is mandatory due to high risk of developing hyperglycaemia

Note Special Pricing Arrangements apply.

Authority required

Acromegaly

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must not have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have a mean growth hormone (GH) level greater than 1 microgram per litre or 3 mIU/L; OR
- Patient must have an age- and sex-adjusted insulin-like growth factor 1 (IGF-1) concentration greater than the upper limit of normal (ULN), **AND**
- The treatment must be after failure to achieve biochemical control with a maximum indicated dose of either 30 mg octreotide LAR or 120 mg lanreotide ATG every 28 days for 24 weeks; unless contraindicated or not tolerated according to the TGA approved Product Information, **AND**
- The treatment must not be given concomitantly with PBS-subsidised pegvisomant.

Population criteria:

- Patient must be aged 18 years or older.

If treatment with either octreotide or lanreotide is contraindicated according to the relevant TGA-approved Product Information, the application must provide details of contraindication.

If intolerance to either octreotide or lanreotide treatment developed during the relevant period of use which is of a severity to necessitate withdrawal of the treatment, the application must provide details of the nature and severity of this intolerance.

Failure to achieve biochemical control after completion of a prior therapy with either octreotide or lanreotide is defined as:

- 1) Growth hormone level greater than 1 mcg/L or 3 mIU/L; OR
- 2) IGF-1 level is greater than the age- and sex-adjusted ULN.

In a patient treated with radiotherapy, pasireotide should be withdrawn every 2 years in the 10 years after completion of radiotherapy for assessment of remission. Pasireotide should be withdrawn at least 8 weeks prior to the assessment of remission.

Biochemical evidence of remission is defined as:

- 1) Growth hormone (GH) levels of less than 1 mcg/L or 3 mIU/L; OR
- 2) normalisation of sex- and age- adjusted insulin-like growth factor 1 (IGF-1)

The authority application must be made in writing and must include:

- a) a completed authority prescription form; and
- b) a completed Acromegaly PBS Authority Application - Supporting Information Form; and
- c) in a patient who has been previously treated with radiotherapy for this condition, the date of completion of radiotherapy must be provided; the date and result of GH or IGF-1 levels taken at the most recent two yearly assessment in the 10 years after completion of radiotherapy must be provided; and
- d) a recent result of GH or IGF-1 levels must be provided.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Acromegaly

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must not be given concomitantly with PBS-subsidised pegvisomant.

Population criteria:

- Patient must be aged 18 years or older.

In a patient treated with radiotherapy, pasireotide should be withdrawn every 2 years in the 10 years after completion of radiotherapy for assessment of remission. Pasireotide should be withdrawn at least 8 weeks prior to the assessment of remission.

Biochemical evidence of remission is defined as:

- 1) Growth hormone (GH) levels of less than 1 mcg/L or 3 mIU/L; OR
- 2) normalisation of sex- and age- adjusted insulin-like growth factor 1 (IGF-1)

In a patient who has been previously treated with radiotherapy for this condition, the date of completion of radiotherapy and the GH and IGF-1 levels taken at the most recent two yearly assessment in the 10 years after completion of radiotherapy must be provided at the time of approval.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authorisation under this criterion should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

pasireotide 60 mg modified release injection [1 vial] (&) inert substance diluent [2 mL syringe], 1 pack

10887B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*7847.39	Signifor LAR [NV]

pasireotide 40 mg modified release injection [1 vial] (&) inert substance diluent [2 mL syringe], 1 pack

10884W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*7847.39	Signifor LAR [NV]

pasireotide 20 mg modified release injection [1 vial] (&) inert substance diluent [2 mL syringe], 1 pack

10880P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*7847.39	Signifor LAR [NV]

ANTIINFECTIVES FOR SYSTEMIC USE

ANTIBACTERIALS FOR SYSTEMIC USE

MACROLIDES, LINCOSAMIDES AND STREPTOGRAMINS

Macrolides

■ AZITHROMYCIN

Authority required (STREAMLINED)

9604

Mycobacterium avium complex infection

Clinical criteria:

- The treatment must be for prophylaxis, **AND**
- Patient must be human immunodeficiency virus (HIV) positive, **AND**
- Patient must have CD4 cell counts of less than 75 per cubic millimetre.

azithromycin 600 mg tablet, 8

6221K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*122.59	Zithromax [PF]

■ ANTIMYCOBACTERIALS

DRUGS FOR TREATMENT OF TUBERCULOSIS

Antibiotics

■ RIFABUTIN

Authority required (STREAMLINED)

9560

Mycobacterium avium complex infection

Clinical criteria:

- Patient must be human immunodeficiency virus (HIV) positive.

Authority required (STREAMLINED)

9622

Mycobacterium avium complex infection

Clinical criteria:

- The treatment must be for prophylaxis, **AND**
- Patient must be human immunodeficiency virus (HIV) positive, **AND**
- Patient must have CD4 cell counts of less than 75 per cubic millimetre.

rifabutin 150 mg capsule, 30

6195C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	4	5	..	*554.27	Mycobutin [PF]

■ ANTIVIRALS FOR SYSTEMIC USE

DIRECT ACTING ANTIVIRALS

Nucleosides and nucleotides excl. reverse transcriptase inhibitors

■ GANCICLOVIR

Authority required (STREAMLINED)

9404

Cytomegalovirus disease

Treatment Phase: Prophylaxis

Clinical criteria:

- Patient must be a bone marrow transplant recipient at risk of cytomegalovirus disease.

Authority required (STREAMLINED)

9526

Cytomegalovirus disease

Treatment Phase: Prophylaxis

Clinical criteria:

- Patient must be a solid organ transplant recipient at risk of cytomegalovirus disease.

ganciclovir 500 mg injection, 5 vials

6136Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	1	..	*424.79	^a Cymevene [PB]	^a GANCICLOVIR SXP [HN]

■ VALACICLOVIR

Authority required (STREAMLINED)

9267

Cytomegalovirus infection and disease

Treatment Phase: Prophylaxis

Clinical criteria:

- Patient must have undergone a renal transplant, **AND**
- Patient must be at risk of cytomegalovirus disease.

valaciclovir 500 mg tablet, 100

6280M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	5	2	..	*237.24	^a APO-Valaciclovir [TX] ^a Valaciclovir RBX [RA]	^a Valaciclovir APOTEX [GX]
			^B 2.30	*239.54	^a Valtrex [RW]	

■ VALGANCICLOVIR

Authority required (STREAMLINED)

9316

Cytomegalovirus infection and disease

Treatment Phase: Prophylaxis

Clinical criteria:

- Patient must be a solid organ transplant recipient at risk of cytomegalovirus disease.

valganciclovir 450 mg tablet, 60

6357N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	5	..	*1851.93	^a Valcyte [RO] ^a Valganciclovir Juno [JU] ^a Valganciclovir Sandoz [SZ]	^a Valganciclovir AN [JO] ^a Valganciclovir Mylan [AF]

valganciclovir 50 mg/mL powder for oral liquid, 100 mL

9675F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	11	5	..	*#4396.36	Valcyte [RO]

Antivirals for treatment of HCV infections

■ DACLATASVIR

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Chronic hepatitis C infection

Clinical criteria:

- Patient must meet the criteria set out in the General Statement for Drugs for the Treatment of Hepatitis C, **AND**
- Patient must be taking this drug as part of a regimen set out in the matrix in the General Statement for Drugs for the Treatment of Hepatitis C, based on the hepatitis C virus genotype, patient treatment history and cirrhotic status, **AND**
- The treatment must be limited to a maximum duration of 24 weeks.

daclatasvir 60 mg tablet, 28

10631M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	7714.06	Daklinza [BQ]

daclatasvir 30 mg tablet, 28

10630L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	7714.06	Daklinza [BQ]

■ DACLATASVIR

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Chronic hepatitis C infection

Clinical criteria:

- Patient must meet the criteria set out in the General Statement for Drugs for the Treatment of Hepatitis C, **AND**
- Patient must be taking this drug as part of a regimen set out in the matrix in the General Statement for Drugs for the Treatment of Hepatitis C, based on the hepatitis C virus genotype, patient treatment history and cirrhotic status, **AND**
- The treatment must be limited to a maximum duration of 12 weeks.

daclatasvir 60 mg tablet, 28

10644F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	2	..	7714.06	Daklinza [BQ]

daclatasvir 30 mg tablet, 28

10643E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	2	..	7714.06	Daklinza [BQ]

■ ELBASVIR + GRAZOPREVR

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Chronic hepatitis C infection

Clinical criteria:

- Patient must meet the criteria set out in the General Statement for Drugs for the Treatment of Hepatitis C, **AND**
- Patient must be taking this drug as part of a regimen set out in the matrix in the General Statement for Drugs for the Treatment of Hepatitis C, based on the hepatitis C virus genotype, patient treatment history and cirrhotic status, **AND**
- The treatment must be limited to a maximum duration of 12 weeks.

elbasvir 50 mg + grazoprevir 100 mg tablet, 28

10979W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	2	..	8447.39	Zepatier [MK]

■ ELBASVIR + GRAZOPREVR

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Chronic hepatitis C infection

Clinical criteria:

- Patient must meet the criteria set out in the General Statement for Drugs for the Treatment of Hepatitis C, **AND**
- Patient must be taking this drug as part of a regimen set out in the matrix in the General Statement for Drugs for the Treatment of Hepatitis C, based on the hepatitis C virus genotype, patient treatment history and cirrhotic status, **AND**
- The treatment must be limited to a maximum duration of 16 weeks.

elbasvir 50 mg + grazoprevir 100 mg tablet, 28

10991L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	3	..	8447.39	Zepatier [MK]

■ GLECAPREVR + PIBRENTASVIR

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Chronic hepatitis C infection

Clinical criteria:

- Patient must meet the criteria set out in the General Statement for Drugs for the Treatment of Hepatitis C, **AND**
- Patient must be taking this drug as part of a regimen set out in the matrix in the General Statement for Drugs for the Treatment of Hepatitis C, based on the hepatitis C virus genotype, patient treatment history and cirrhotic status, **AND**
- The treatment must be limited to a maximum duration of 16 weeks.

glecaprevir 100 mg + pibrentasvir 40 mg film-coated tablet, 84

11337Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	3	..	18714.06	Maviret [VE]

▪ **GLECAPREVIR + PIBRENTASVIR**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Chronic hepatitis C infection

Clinical criteria:

- Patient must meet the criteria set out in the General Statement for Drugs for the Treatment of Hepatitis C, **AND**
- Patient must be taking this drug as part of a regimen set out in the matrix in the General Statement for Drugs for the Treatment of Hepatitis C, based on the hepatitis C virus genotype, patient treatment history and cirrhotic status, **AND**
- The treatment must be limited to a maximum duration of 12 weeks.

glecaprevir 100 mg + pibrentasvir 40 mg film-coated tablet, 84

11346E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	2	..	18714.06	Maviret [VE]

▪ **GLECAPREVIR + PIBRENTASVIR**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Chronic hepatitis C infection

Clinical criteria:

- Patient must meet the criteria set out in the General Statement for Drugs for the Treatment of Hepatitis C, **AND**
- Patient must be taking this drug as part of a regimen set out in the matrix in the General Statement for Drugs for the Treatment of Hepatitis C, based on the hepatitis C virus genotype, patient treatment history and cirrhotic status, **AND**
- The treatment must be limited to a maximum duration of 8 weeks.

glecaprevir 100 mg + pibrentasvir 40 mg film-coated tablet, 84

11355P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	1	..	18714.06	Maviret [VE]

▪ **LEDIPASVIR + SOFOSBUVIR**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Chronic hepatitis C infection

Clinical criteria:

- Patient must meet the criteria set out in the General Statement for Drugs for the Treatment of Hepatitis C, **AND**
- Patient must be taking this drug as part of a regimen set out in the matrix in the General Statement for Drugs for the Treatment of Hepatitis C, based on the hepatitis C virus genotype, patient treatment history and cirrhotic status, **AND**
- The treatment must be limited to a maximum duration of 8 weeks.

ledipasvir 90 mg + sofosbuvir 400 mg tablet, 28

10653Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	1	..	12547.39	Harvoni [GI]

▪ **LEDIPASVIR + SOFOSBUVIR**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Chronic hepatitis C infection

Clinical criteria:

- Patient must meet the criteria set out in the General Statement for Drugs for the Treatment of Hepatitis C, **AND**
- Patient must be taking this drug as part of a regimen set out in the matrix in the General Statement for Drugs for the Treatment of Hepatitis C, based on the hepatitis C virus genotype, patient treatment history and cirrhotic status, **AND**
- The treatment must be limited to a maximum duration of 12 weeks.

ledipasvir 90 mg + sofosbuvir 400 mg tablet, 28

10672Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	2	..	12547.39	Harvoni [GI]

▪ LEDIPASVIR + SOFOSBUVIR**Note** No increase in the maximum quantity or number of units may be authorised.**Note** No increase in the maximum number of repeats may be authorised.**Note** Special Pricing Arrangements apply.**Authority required**

Chronic hepatitis C infection

Clinical criteria:

- Patient must meet the criteria set out in the General Statement for Drugs for the Treatment of Hepatitis C, **AND**
- Patient must be taking this drug as part of a regimen set out in the matrix in the General Statement for Drugs for the Treatment of Hepatitis C, based on the hepatitis C virus genotype, patient treatment history and cirrhotic status, **AND**
- The treatment must be limited to a maximum duration of 24 weeks.

ledipasvir 90 mg + sofosbuvir 400 mg tablet, 28

10679C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	12547.39	Harvoni [GI]

▪ RIBAVIRIN**Caution** Ribavirin is a category X drug and must not be given to pregnant women. Pregnancy in female patients or in the partners of male patients must be avoided during treatment and during the 6 months period after cessation of treatment.**Note** No increase in the maximum number of repeats may be authorised.**Authority required**

Chronic hepatitis C infection

Clinical criteria:

- Patient must meet the criteria set out in the General Statement for Drugs for the Treatment of Hepatitis C, **AND**
- Patient must be taking this drug as part of a regimen set out in the matrix in the General Statement for Drugs for the Treatment of Hepatitis C, based on the hepatitis C virus genotype, patient treatment history and cirrhotic status, **AND**
- The treatment must be limited to a maximum duration of 12 weeks.

Population criteria:

- Patient must not be pregnant or breastfeeding. Female partners of male patients must not be pregnant. Patients and their partners must each be using an effective form of contraception if of child-bearing age.

ribavirin 600 mg tablet, 28

10675W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	2	..	444.19	Ibavir [IX]

ribavirin 400 mg tablet, 28

10623D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	2	..	298.59	Ibavir [IX]

▪ RIBAVIRIN**Caution** Ribavirin is a category X drug and must not be given to pregnant women. Pregnancy in female patients or in the partners of male patients must be avoided during treatment and during the 6 months period after cessation of treatment.**Note** No increase in the maximum number of repeats may be authorised.**Authority required**

Chronic hepatitis C infection

Clinical criteria:

- Patient must meet the criteria set out in the General Statement for Drugs for the Treatment of Hepatitis C, **AND**
- Patient must be taking this drug as part of a regimen set out in the matrix in the General Statement for Drugs for the Treatment of Hepatitis C, based on the hepatitis C virus genotype, patient treatment history and cirrhotic status, **AND**
- The treatment must be limited to a maximum duration of 24 weeks.

Population criteria:

- Patient must not be pregnant or breastfeeding. Female partners of male patients must not be pregnant. Patients and their partners must each be using an effective form of contraception if of child-bearing age.

ribavirin 600 mg tablet, 28

10637W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	444.19	Ibavyr [IX]

ribavirin 400 mg tablet, 28

10635R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	298.59	Ibavyr [IX]

▪ SOFOSBUVIR

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Chronic hepatitis C infection

Clinical criteria:

- Patient must meet the criteria set out in the General Statement for Drugs for the Treatment of Hepatitis C, **AND**
- Patient must be taking this drug as part of a regimen set out in the matrix in the General Statement for Drugs for the Treatment of Hepatitis C, based on the hepatitis C virus genotype, patient treatment history and cirrhotic status, **AND**
- The treatment must be limited to a maximum duration of 12 weeks.

sofosbuvir 400 mg tablet, 28

10654R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	2	..	12547.39	Sovaldi [GI]

▪ SOFOSBUVIR

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Chronic hepatitis C infection

Clinical criteria:

- Patient must meet the criteria set out in the General Statement for Drugs for the Treatment of Hepatitis C, **AND**
- Patient must be taking this drug as part of a regimen set out in the matrix in the General Statement for Drugs for the Treatment of Hepatitis C, based on the hepatitis C virus genotype, patient treatment history and cirrhotic status, **AND**
- The treatment must be limited to a maximum duration of 24 weeks.

sofosbuvir 400 mg tablet, 28

10676X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	12547.39	Sovaldi [GI]

▪ SOFOSBUVIR + VELPATASVIR

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Chronic hepatitis C infection

Clinical criteria:

- Patient must meet the criteria set out in the General Statement for Drugs for the Treatment of Hepatitis C, **AND**
- Patient must be taking this drug as part of a regimen set out in the matrix in the General Statement for Drugs for the Treatment of Hepatitis C, based on the hepatitis C virus genotype, patient treatment history and cirrhotic status, **AND**
- The treatment must be limited to a maximum duration of 12 weeks.

sofosbuvir 400 mg + velpatasvir 100 mg tablet, 28

11144M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	2	..	12547.39	Eplusa [GI]

▪ SOFOSBUVIR + VELPATASVIR + VOXILAPREVIR

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Chronic hepatitis C infection

Clinical criteria:

- Patient must meet the criteria set out in the General Statement for Drugs for the Treatment of Hepatitis C, **AND**
- Patient must be taking this drug as part of a regimen set out in the matrix in the General Statement for Drugs for the Treatment of Hepatitis C, based on the hepatitis C virus genotype, patient treatment history and cirrhotic status, **AND**

- The treatment must be limited to a maximum duration of 12 weeks.

sofosbuvir 400 mg + velpatasvir 100 mg + voxilaprevir 100 mg tablet, 28

11659P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	2	..	12547.39	Vosevi [GI]

■ ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS

■ ANTINEOPLASTIC AGENTS

ANTIMETABOLITES

Pyrimidine analogues

■ AZACITIDINE

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Myelodysplastic syndrome

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be classified as Intermediate-2 according to the International Prognostic Scoring System (IPSS); OR
 - The condition must be classified as high risk according to the International Prognostic Scoring System (IPSS).
- Classification of the condition as Intermediate-2 requires a score of 1.5 to 2.0 on the IPSS, achieved with the possible combinations:

- 11% to 30% marrow blasts with good karyotypic status (normal, -Y alone, del(5q) alone, del(20q) alone), and 0 to 1 cytopenias; OR
- 11% to 20% marrow blasts with intermediate karyotypic status (other abnormalities), and 0 to 1 cytopenias; OR
- 11% to 20% marrow blasts with good karyotypic status (normal, -Y alone, del(5q) alone, del(20q) alone), and 2 to 3 cytopenias; OR
- 5% to 10% marrow blasts with poor karyotypic status (3 or more abnormalities or chromosome 7 anomalies), regardless of cytopenias; OR
- 5% to 10% marrow blasts with intermediate karyotypic status (other abnormalities), and 2 to 3 cytopenias; OR
- Less than 5% marrow blasts with poor karyotypic status (3 or more abnormalities or chromosome 7 anomalies), and 2 to 3 cytopenias.

Classification of the condition as high risk requires a score of 2.5 or more on the IPSS, achieved with the possible combinations:

- 21% to 30% marrow blasts with good karyotypic status (normal, -Y alone, del(5q) alone, del(20q) alone), and 2 to 3 cytopenias; OR
- 21% to 30% marrow blasts with intermediate (other abnormalities) or poor karyotypic status (3 or more abnormalities or chromosome 7 anomalies), regardless of cytopenias; OR
- 11% to 20% marrow blasts with poor karyotypic status (3 or more abnormalities or chromosome 7 anomalies), regardless of cytopenias; OR
- 11% to 20% marrow blasts with intermediate karyotypic status (other abnormalities), and 2 to 3 cytopenias.

The first authority application must be made in writing and must include:

- a completed authority prescription form; and
- a completed Azacitidine PBS Authority Application - Supporting Information Form; and
- a copy of the bone marrow biopsy report demonstrating that the patient has myelodysplastic syndrome; and
- a copy of the full blood examination report; and
- a copy of the pathology report detailing the cytogenetics demonstrating intermediate-2 or high risk disease according to the International Prognostic Scoring System (IPSS); and
- a signed patient acknowledgment form.

No more than 3 cycles will be authorised.

Authority required

Chronic Myelomonocytic Leukaemia

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must have 10% to 29% marrow blasts without Myeloproliferative Disorder.

The first authority application must be made in writing and must include:

- a completed authority prescription form; and
- a completed Azacitidine PBS Authority Application - Supporting Information Form; and
- a copy of the bone marrow biopsy report demonstrating that the patient has chronic myelomonocytic leukaemia ; and

- (d) a copy of the full blood examination report; and
(e) a signed patient acknowledgement.
No more than 3 cycles will be authorised.

Authority required

Acute Myeloid Leukaemia

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must have 20% to 30% marrow blasts and multi-lineage dysplasia, according to World Health Organisation (WHO) Classification.

The first authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
 - (b) a completed Azacitidine PBS Authority Application - Supporting Information Form; and
 - (c) a copy of the bone marrow biopsy report demonstrating that the patient has acute myeloid leukaemia; and
 - (d) a copy of the full blood examination report; and
 - (e) a signed patient acknowledgement.
- No more than 3 cycles will be authorised.

azacitidine 100 mg injection, 1 vial

6100C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	14	2	..	*3094.39	^a Azacitidine Accord [OC] ^a Azacitidine Juno [JO] ^a Celazadine [JU]	^a AZACITIDINE DR.REDDY'S [RI] ^a Azadine [RZ] ^a Vidaza [CJ]

■ AZACITIDINE

Note Authority applications for continuing treatment may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Myelodysplastic syndrome

Treatment Phase: Continuing treatment

Clinical criteria:

- The condition must be classified as Intermediate-2 according to the International Prognostic Scoring System (IPSS); OR
- The condition must be classified as high risk according to the International Prognostic Scoring System (IPSS), **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have progressive disease.

Applications for continuing therapy may be made by telephone.

Up to 6 cycles will be authorised.

Authority required

Chronic Myelomonocytic Leukaemia

Treatment Phase: Continuing treatment

Clinical criteria:

- The condition must have 10% to 29% marrow blasts without Myeloproliferative Disorder, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have progressive disease.

Applications for continuing therapy may be made by telephone.

Up to 6 cycles will be authorised.

Authority required

Acute Myeloid Leukaemia

Treatment Phase: Continuing treatment

Clinical criteria:

- The condition must have 20% to 30% marrow blasts and multi-lineage dysplasia, according to World Health Organisation (WHO) Classification, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have progressive disease.

Applications for continuing therapy may be made by telephone.

Up to 6 cycles will be authorised.

azacitidine 100 mg injection, 1 vial

6138C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	14	5	..	*3094.39	^a Azacitidine Accord [OC] ^a Azacitidine Juno [JO] ^a Celazadine [JU]	^a AZACITIDINE DR.REDDY'S [RI] ^a Azadine [RZ] ^a Vidaza [CJ]

CYTOTOXIC ANTIBIOTICS AND RELATED SUBSTANCES

Anthracyclines and related substances

■ DOXORUBICIN HYDROCHLORIDE (AS PEGYLATED LIPOSOMAL)

Authority required (STREAMLINED)

9287

Kaposi sarcoma

Clinical criteria:

- The condition must be AIDS-related, **AND**
- Patient must have a CD4 cell count of less than 200 per cubic millimetre, **AND**
- The condition must include extensive mucocutaneous involvement.

Authority required (STREAMLINED)

9223

Kaposi sarcoma

Clinical criteria:

- The condition must be AIDS-related, **AND**
- Patient must have a CD4 cell count of less than 200 per cubic millimetre, **AND**
- The condition must include extensive visceral involvement.

doxorubicin hydrochloride (as pegylated liposomal) 20 mg/10 mL injection, 10 mL vial

6249X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	4	5	..	*936.91	^a Caelyx [JC]	^a Liposomal Doxorubicin SUN [RA]

OTHER ANTINEOPLASTIC AGENTS

Monoclonal antibodies

■ RITUXIMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for adults with severe active rheumatoid arthritis. Where the term biological medicine appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alfa antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab), the T-cell co-stimulation modulator (abatacept) and the Janus kinase (JAK) inhibitors (baricitinib, tofacitinib). A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised biological medicine therapy may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once, and
- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised biological medicines for the treatment of rheumatoid arthritis.

Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Stevens Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

For a patient who has failed PBS-subsidised treatment with 2 or 3 TNF-alfa antagonists prior to 1 August 2010 please contact the Department of Human Services on 1800 700 270. A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction.

A patient who has failed fewer than 5 biological medicines and who has a break in treatment of less than 24 months may commence a further course of treatment with a biological medicine under Initial 2 treatment restriction. A patient who has failed fewer than 5 biological medicines and who has had a break in therapy of longer than 24 months may commence a further course of treatment with a biological medicine under the Initial 3 treatment restriction. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine treatment is stopped to the date of the new application for treatment with a biological medicine.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 April 2019.

(a) Initial treatment.

Applications for initial treatment should be made where:

- a patient has received no prior PBS-subsidised biological medicine treatment and wishes to commence such therapy, excluding rituximab (Initial 1 - new patient); or
 - a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months) [further details are under 'Swapping therapy' below]; or
 - a patient wishes to re-commence treatment with a specific biological medicine following a break of less than 24 months in PBS-subsidised therapy with that agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months).
 - a patient wishes to re-commence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 24 months (Initial 3 - re-commencement of treatment after a break in biological medicine of more than 24 months)
- Initial applications for a new patient (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6-month intensive DMARD trial, but prior to ceasing DMARD therapy.
- Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, baricitinib, etanercept, golimumab, tocilizumab and tofacitinib, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.
- Rituximab patients should be assessed following a minimum of 12 weeks after the first infusion, and the assessment should be submitted to the Department of Human Services within 4 weeks to ensure continuity of treatment for those who meet the

continuing restriction for PBS subsidised treatment with this drug for this condition.

Where a response assessment is not provided the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment. For second and subsequent courses of PBS-subsidised biological medicine (excluding rituximab) treatment it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to the Department of Human Services within sufficient timeframe to allow for authority assessment.

Abatacept patients:

A patient is eligible to receive one I.V. loading dose when commencing treatment with the subcutaneous formulation. Two prescriptions are required, the first prescription for the I.V. loading dose for sufficient vials for one dose based on the patient's weight with no repeats. The second prescription for the subcutaneous formulation, with a maximum quantity of 4 and up to 3 repeats, must be submitted with the initial application.

Rituximab patients:

A further application may be submitted to the Department of Human Services within sufficient timeframe to allow authority assessment. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

Infliximab and etanercept patients:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction. Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

(2) Swapping therapy

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non- biological medicine therapy requirements except if the patient has had a break in therapy of more than 24 months who would need to requalify with respect to the indices of disease severity. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each biological medicine trialled.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent, unless they have a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug.

Abatacept:

A patient swapping from I.V. abatacept to subcutaneous abatacept will not be eligible for an I.V. loading dose when commencing treatment with the subcutaneous formulation.

Rituximab:

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF-alfa antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological medicine during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological medicine therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate biological medicine may do so without having to have any treatment-free period.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and the Department of Human Services will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints.

Applications under the Initial 1 treatment restriction for a new patient must include a joint count and ESR and/or CRP

measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. Applications under the Initial 3 treatment restriction for re-commencement of treatment after a break in biological medicine of more than 24 months must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required (STREAMLINED)

9611

Severe active rheumatoid arthritis

Treatment Phase: Subsequent continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be given concomitantly with methotrexate at a dose of at least 7.5 mg weekly.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The measurement of response to the prior course of therapy must be documented in the patient's medical notes.

If a patient has either failed or ceased to respond to a PBS-subsidised biological medicine for this condition 5 times, they will not be eligible to receive further PBS-subsidised treatment with a biological medicine for this condition.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

rituximab 500 mg/50 mL injection, 50 mL vial

11790M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	1	..	*2446.03	Riximyo [SZ]

■ RITUXIMAB

Note Authority applications for increased quantities/ repeats (where relevant) may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note Risk of end-organ damage or mortality includes a minimum of one of the following: Glomerulonephritis with risk of progression

- Risk to sight including scleritis/episcleritis, sudden visual loss, uveitis, retinal changes (vasculitis/thrombosis/exudates/haemorrhage)
- Bronchial/subglottic obstruction
- Pulmonary haemorrhage
- Parenchymal lung disease
- Sensory neural hearing loss
- Recurrent sinonasal disease requiring recurrent surgical interventions
- Meningitis, organic confusion, seizures, stroke, cord lesion, cranial nerve palsy, sensory peripheral neuropathy, motor mononeuritis multiplex

Note Patients could be considered contraindicated, refractory, or unable to tolerate cyclophosphamide for one of the following reasons: Cyclophosphamide is contraindicated as per the TGA approved Product Information;

- Cyclophosphamide is not recommended due to the need to preserve gonad function;
- Patient experiences severe toxicity to cyclophosphamide that warrants cessation of treatment;
- Patient has life- or organ-threatening deterioration at any time during treatment with cyclophosphamide, where the deterioration is thought to be due to severe uncontrolled active vasculitis;
- Commencing a further treatment cycle with cyclophosphamide would exceed the maximum cumulative dose of cyclophosphamide of 25g; or

- Patient's condition with this indication is persistent despite at least 3 months therapy with cyclophosphamide.

Authority required (STREAMLINED)

9640

Severe active granulomatosis with polyangiitis (Wegeners granulomatosis)

Treatment Phase: Re-induction of remission

Clinical criteria:

- The treatment must be for the re-induction of remission, **AND**
- Patient must have previously received and responded to this drug for this condition, **AND**
- The treatment must in combination with glucocorticoids, **AND**
- Patient must be at risk of end-organ damage or mortality, **AND**
- Patient must be contraindicated, refractory or unable to tolerate cyclophosphamide.

Diagnosis should be made according to the Chapel Hill Consensus Conference Nomenclature of the Vasculitides with anti-neutrophil cytoplasmic antibody (ANCA) positive serology.

This drug is not PBS-subsidised for maintenance of remission

Authority required (STREAMLINED)

9641

Severe active microscopic polyangiitis

Treatment Phase: Re-induction of remission

Clinical criteria:

- The treatment must be for the re-induction of remission, **AND**
- Patient must have previously received and responded to this drug for this condition, **AND**
- The treatment must in combination with glucocorticoids, **AND**
- Patient must be at risk of end-organ damage or mortality, **AND**
- Patient must be contraindicated, refractory or unable to tolerate cyclophosphamide.

Diagnosis should be made according to the Chapel Hill Consensus Conference Nomenclature of the Vasculitides with anti-neutrophil cytoplasmic antibody (ANCA) positive serology.

This drug is not PBS-subsidised for maintenance therapy.

rituximab 500 mg/50 mL injection, 50 mL vial

11804G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	4	..	..	*4844.67	Riximyo [SZ]

rituximab 100 mg/10 mL injection, 2 x 10 mL vials

11810N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	3	..	..	*1486.57	Riximyo [SZ]

▪ RITUXIMAB

Note Risk of end-organ damage or mortality includes a minimum of one of the following: Glomerulonephritis with risk of progression

- Risk to sight including scleritis/episcleritis, sudden visual loss, uveitis, retinal changes (vasculitis/thrombosis/exudates/haemorrhage)
- Bronchial/subglottic obstruction
- Pulmonary haemorrhage
- Parenchymal lung disease
- Sensory neural hearing loss
- Recurrent sinonasal disease requiring recurrent surgical interventions
- Meningitis, organic confusion, seizures, stroke, cord lesion, cranial nerve palsy, sensory peripheral neuropathy, motor mononeuritis multiplex

Note Patients could be considered contraindicated, refractory, or unable to tolerate cyclophosphamide for one of the following reasons: Cyclophosphamide is contraindicated as per the TGA approved Product Information;

- Cyclophosphamide is not recommended due to the need to preserve gonad function;
- Patient experiences severe toxicity to cyclophosphamide that warrants cessation of treatment;
- Patient has life- or organ-threatening deterioration at any time during treatment with cyclophosphamide, where the deterioration is thought to be due to severe uncontrolled active vasculitis;
- Commencing a further treatment cycle with cyclophosphamide would exceed the maximum cumulative dose of cyclophosphamide of 25g; or
- Patient's condition with this indication is persistent despite at least 3 months therapy with cyclophosphamide.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs

Reply Paid 9826
HOBART TAS 7001

Note At the time of authority application, medical practitioners must request the appropriate number of vials to provide sufficient drug for four weeks of treatment.

Authority required

Severe active granulomatosis with polyangiitis (Wegeners granulomatosis)

Treatment Phase: Induction of remission

Clinical criteria:

- The treatment must be for the induction of remission, **AND**
- Patient must not have previously received this drug for this condition, **AND**
- The treatment must in combination with glucocorticoids, **AND**
- Patient must be at risk of end-organ damage or mortality, **AND**
- Patient must be contraindicated, refractory or unable to tolerate cyclophosphamide.

Diagnosis should be made according to the Chapel Hill Consensus Conference Nomenclature of the Vasculitides with anti-neutrophil cytoplasmic antibody (ANCA) positive serology.

This drug is not PBS-subsidised for maintenance of remission

The authority application must be made in writing

Note Biosimilar prescribing policy

Prescribing of the biosimilar brand, Riximyo, is encouraged for treatment naive patients.

Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Authority required

Severe active microscopic polyangiitis

Treatment Phase: Induction of remission

Clinical criteria:

- The treatment must be for the induction of remission, **AND**
- Patient must not have previously received this drug for this condition, **AND**
- The treatment must in combination with glucocorticoids, **AND**
- Patient must be at risk of end-organ damage or mortality, **AND**
- Patient must be contraindicated, refractory or unable to tolerate cyclophosphamide.

Diagnosis should be made according to the Chapel Hill Consensus Conference Nomenclature of the Vasculitides with anti-neutrophil cytoplasmic antibody (ANCA) positive serology.

This drug is not PBS-subsidised for maintenance therapy.

The authority application must be made in writing

Note Biosimilar prescribing policy

Prescribing of the biosimilar brand, Riximyo, is encouraged for treatment naive patients.

Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Authority required

Severe active granulomatosis with polyangiitis (Wegeners granulomatosis)

Treatment Phase: Re-induction of remission

Clinical criteria:

- The treatment must be for the re-induction of remission, **AND**
- Patient must have previously received and responded to this drug for this condition, **AND**
- The treatment must in combination with glucocorticoids, **AND**
- Patient must be at risk of end-organ damage or mortality, **AND**
- Patient must be contraindicated, refractory or unable to tolerate cyclophosphamide.

Diagnosis should be made according to the Chapel Hill Consensus Conference Nomenclature of the Vasculitides with anti-neutrophil cytoplasmic antibody (ANCA) positive serology.

This drug is not PBS-subsidised for maintenance of remission

The authority application must be made in writing

Authority required

Severe active microscopic polyangiitis

Treatment Phase: Re-induction of remission

Clinical criteria:

- The treatment must be for the re-induction of remission, **AND**
- Patient must have previously received and responded to this drug for this condition, **AND**
- The treatment must in combination with glucocorticoids, **AND**
- Patient must be at risk of end-organ damage or mortality, **AND**
- Patient must be contraindicated, refractory or unable to tolerate cyclophosphamide.

Diagnosis should be made according to the Chapel Hill Consensus Conference Nomenclature of the Vasculitides with anti-neutrophil cytoplasmic antibody (ANCA) positive serology.

This drug is not PBS-subsidised for maintenance therapy.

The authority application must be made in writing

rituximab 500 mg/50 mL injection, 50 mL vial

10576P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	1246.71	^a Mabthera [RO]	^a Riximyo [SZ]

rituximab 100 mg/10 mL injection, 2 x 10 mL vials

10583B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	506.31	^a Mabthera [RO]	^a Riximyo [SZ]

■ RITUXIMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for adults with severe active rheumatoid arthritis. Where the term biological medicine appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alfa antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab), the T-cell co-stimulation modulator (abatacept) and the Janus kinase (JAK) inhibitors (baricitinib, tofacitinib). A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised biological medicine therapy may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once, and
- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised biological medicines for the treatment of rheumatoid arthritis.

Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Stevens Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

For a patient who has failed PBS-subsidised treatment with 2 or 3 TNF-alfa antagonists prior to 1 August 2010 please contact the Department of Human Services on 1800 700 270. A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction.

A patient who has failed fewer than 5 biological medicines and who has a break in treatment of less than 24 months may commence a further course of treatment with a biological medicine under Initial 2 treatment restriction. A patient who has failed fewer than 5 biological medicines and who has had a break in therapy of longer than 24 months may commence a further course of treatment with a biological medicine under the Initial 3 treatment restriction. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine treatment is stopped to the date of the new application for treatment with a biological medicine.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 April 2019.

(a) Initial treatment.

Applications for initial treatment should be made where:

- a patient has received no prior PBS-subsidised biological medicine treatment and wishes to commence such therapy, excluding rituximab (Initial 1 - new patient); or
 - a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months) [further details are under 'Swapping therapy' below]; or
 - a patient wishes to re-commence treatment with a specific biological medicine following a break of less than 24 months in PBS-subsidised therapy with that agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months).
 - a patient wishes to re-commence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 24 months (Initial 3 - re-commencement of treatment after a break in biological medicine of more than 24 months)
- Initial applications for a new patient (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6-month intensive DMARD trial, but prior to ceasing DMARD therapy.

Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, baricitinib, etanercept, golimumab, tocilizumab and tofacitinib, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

Rituximab patients should be assessed following a minimum of 12 weeks after the first infusion, and the assessment should be submitted to the Department of Human Services within 4 weeks to ensure continuity of treatment for those who meet the continuing restriction for PBS subsidised treatment with this drug for this condition.

Where a response assessment is not provided the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment. For second and subsequent courses of PBS-subsidised biological medicine (excluding rituximab) treatment it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to the Department of Human Services within sufficient timeframe to allow for authority assessment.

Abatacept patients:

A patient is eligible to receive one I.V. loading dose when commencing treatment with the subcutaneous formulation. Two prescriptions are required, the first prescription for the I.V. loading dose for sufficient vials for one dose based on the patient's weight with no repeats. The second prescription for the subcutaneous formulation, with a maximum quantity of 4

and up to 3 repeats, must be submitted with the initial application.

Rituximab patients:

A further application may be submitted to the Department of Human Services within sufficient timeframe to allow authority assessment. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

Infliximab and etanercept patients:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction. Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

(2) Swapping therapy

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non- biological medicine therapy requirements except if the patient has had a break in therapy of more than 24 months who would need to requalify with respect to the indices of disease severity. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each biological medicine trialled.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent, unless they have a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug.

Abatacept:

A patient swapping from I.V. abatacept to subcutaneous abatacept will not be eligible for an I.V. loading dose when commencing treatment with the subcutaneous formulation.

Rituximab:

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF-alfa antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological medicine during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological medicine therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate biological medicine may do so without having to have any treatment-free period.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and the Department of Human Services will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints.

Applications under the Initial 1 treatment restriction for a new patient must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy.

Applications under the Initial 3 treatment restriction for re-commencement of treatment after a break in biological medicine of more than 24 months must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have failed to respond to at least 1 PBS-subsidised tumour necrosis factor (TNF) alfa antagonist for this condition, **AND**
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with disease modifying anti-rheumatic drugs (DMARDs) which must include at least 3 months continuous treatment with each of at least 2 DMARDs, one of which must be methotrexate at a dose of at least 20 mg weekly and one of which must be: (i) hydroxychloroquine at a dose of at least 200 mg daily; or (ii) leflunomide at a dose of at least 10 mg daily; or (iii) sulfasalazine at a dose of at least 2 g daily; OR
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs which, if methotrexate is contraindicated according to the Therapeutic Goods Administration (TGA)-approved Product Information or cannot be tolerated at a 20 mg weekly dose, must include at least 3 months continuous treatment with each of at least 2 of the following DMARDs: (i) hydroxychloroquine at a dose of at least 200 mg daily; and/or (ii) leflunomide at a dose of at least 10 mg daily; and/or (iii) sulfasalazine at a dose of at least 2 g daily; OR
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs which, if 3 or more of methotrexate, hydroxychloroquine, leflunomide and sulfasalazine are contraindicated according to the relevant TGA-approved Product Information or cannot be tolerated at the doses specified above, must include at least 3 months continuous treatment with each of at least 2 DMARDs, with one or more of the following DMARDs being used in place of the DMARDs which are contraindicated or not tolerated: (i) azathioprine at a dose of at least 1 mg/kg per day; and/or (ii) cyclosporin at a dose of at least 2 mg/kg/day; and/or (iii) sodium aurothiomalate at a dose of 50 mg weekly, **AND**
- Patient must not receive more than 2 infusions of this drug under this restriction, **AND**
- The treatment must be given concomitantly with methotrexate at a dose of at least 7.5 mg weekly.

Population criteria:

- Patient must be aged 18 years or older.

If methotrexate is contraindicated according to the TGA-approved product information or cannot be tolerated at a 20 mg weekly dose, the application must include details of the contraindication or intolerance including severity to methotrexate. The maximum tolerated dose of methotrexate must be documented in the application, if applicable.

The application must include details of the DMARDs trialed, their doses and duration of treatment, and all relevant contraindications and/or intolerances including severity.

The requirement to trial at least 2 DMARDs for periods of at least 3 months each can be met using single agents sequentially or by using one or more combinations of DMARDs.

If the requirement to trial 6 months of intensive DMARD therapy with at least 2 DMARDs cannot be met because of contraindications and/or intolerances of a severity necessitating permanent treatment withdrawal to all of the DMARDs specified above, details of the contraindication or intolerance including severity and dose for each DMARD must be provided in the authority application.

The following criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; **AND** either

- (a) a total active joint count of at least 20 active (swollen and tender) joints; or
- (b) at least 4 active joints from the following list of major joints:
 - (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The joint count and ESR and/or CRP must be determined at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. All measures must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the

date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient whose most recent course of PBS-subsidised therapy was with this drug and whose response to this treatment is demonstrated at 12 weeks, may apply for a further course of this drug under the First continuing treatment restriction.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

If a patient fails to demonstrate a response to this drug and who qualifies to trial an alternate biological medicine according to the interchangeability arrangements for biological medicines for the treatment of severe rheumatoid arthritis, may do so without having to have a 22 week treatment-free period.

Note Biosimilar prescribing policy

Prescribing of the biosimilar brand, Riximyo, is encouraged for treatment naive patients.

Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note The Department of Human Services website (www.humanservices.gov.au) has details of the toxicities, including severity, which will be accepted for the following purposes:

- (a) exempting a patient from the requirement to undertake a minimum 3 month trial of methotrexate at a 20 mg weekly dose;
- (b) substituting azathioprine, cyclosporin or sodium aurothiomalate for another DMARD as part of the 6 month intensive DMARD trial;
- (c) exempting a patient from the requirement for a 6 month trial of intensive DMARD therapy.

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 24 months)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to respond to at least 1 PBS-subsidised tumour necrosis factor (TNF) alfa antagonist for this condition, **AND**
- Patient must not have failed to respond to previous PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised biological medicine treatment for this condition 5 times, **AND**
- Patient must not receive more than 2 infusions of this drug under this restriction, **AND**
- The treatment must be given concomitantly with methotrexate at a dose of at least 7.5 mg weekly.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either of the Initial 1, Initial 2, Initial 3, first or subsequent continuing treatment restrictions, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

A patient may qualify to receive a further course of treatment (every 24 weeks) with this drug provided they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with this drug. The demonstration of response must be submitted within 4 weeks of assessment.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient whose most recent course of PBS-subsidised therapy was with this drug and whose response to this treatment is demonstrated at 12 weeks, may apply for a further course of this drug under the First continuing treatment restriction.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

If a patient fails to demonstrate a response to this drug and who qualifies to trial an alternate biological medicine according to the interchangeability arrangements for biological medicines for the treatment of severe rheumatoid arthritis, may do so without having to have a 22 week treatment-free period.

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 3 (recommencement of treatment after a break in biological medicine of more than 24 months)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 24 months or more from the most recent PBS-subsidised biological medicine for this condition, **AND**
- Patient must have failed to respond to at least 1 PBS-subsidised tumour necrosis factor (TNF) alfa antagonist for this condition, **AND**
- Patient must not have failed to respond to previous PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have already failed , or ceased to respond to, PBS-subsidised biological medicine treatment for this condition 5 times, **AND**
- The condition must have an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; OR
- The condition must have a C-reactive protein (CRP) level greater than 15 mg per L, **AND**
- The condition must have either (a) a total active joint count of at least 20 active (swollen and tender) joints; or (b) at least 4 active major joints, **AND**
- Patient must not receive more than 2 infusions of this drug under this restriction, **AND**
- The treatment must be given concomitantly with methotrexate at a dose of at least 7.5 mg weekly.

Population criteria:

- Patient must be aged 18 years or older.

Major joints are defined as (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

All measures of joint count and ESR and/or CRP must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient whose most recent course of PBS-subsidised therapy was with this drug and whose response to this treatment is demonstrated at 12 weeks, may apply for a further course of this drug under the First continuing treatment restriction.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

If a patient fails to demonstrate a response to this drug and who qualifies to trial an alternate biological medicine according to the interchangeability arrangements for biological medicines for the treatment of severe rheumatoid arthritis, may do so without having to have a 22 week treatment-free period.

Note Biosimilar prescribing policy

Prescribing of the biosimilar brand, Riximyo, is encouraged for treatment naive patients. Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars). Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Authority required

Severe active rheumatoid arthritis

Treatment Phase: First continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 2 infusions of this drug under this restriction, **AND**
- The treatment must be given concomitantly with methotrexate at a dose of at least 7.5 mg weekly.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

A patient may qualify to receive a further course of treatment (every 24 weeks) with this drug provided they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with this drug. The demonstration of response must be submitted within 4 weeks of assessment.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient has either failed or ceased to respond to a PBS-subsidised biological medicine for this condition 5 times, they will not be eligible to receive further PBS-subsidised treatment with a biological medicine for this condition.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Subsequent continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be given concomitantly with methotrexate at a dose of at least 7.5 mg weekly.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

- (a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or
- (b) a reduction in the number of the following active joints, from at least 4, by at least 50%:
- elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

- a completed authority prescription form(s); and
- a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

A patient may qualify to receive a further course of treatment (every 24 weeks) with this drug provided they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with this drug. The demonstration of response must be submitted within 4 weeks of assessment.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient has either failed or ceased to respond to a PBS-subsidised biological medicine for this condition 5 times, they will not be eligible to receive further PBS-subsidised treatment with a biological medicine for this condition.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

rituximab 500 mg/50 mL injection, 50 mL vial

9611W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	1246.71	^a Mabthera [RO]	^a Riximyo [SZ]

HSD (Private)

Protein kinase inhibitors

■ MIDOSTAURIN

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Applications for authority to prescribe may be made by phone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday)

Authority required

Acute Myeloid Leukaemia

Treatment Phase: Induction / Consolidation therapy

Clinical criteria:

- Patient must not have received prior chemotherapy as induction therapy for this condition; OR
- The treatment must be for consolidation treatment following induction treatment with midostaurin in combination with chemotherapy, **AND**
- The condition must be internal tandem duplication (ITD) or tyrosine kinase domain (TKD) FMS tyrosine kinase 3 (FLT3) mutation positive before initiating this drug for this condition, **AND**
- The condition must not be acute promyelocytic leukaemia, **AND**
- The treatment must be in combination with standard intensive remission induction or consolidation chemotherapy for this condition.

A maximum of 6 cycles will be authorised under this restriction in a lifetime.

Standard intensive remission induction combination chemotherapy must include cytarabine and an anthracycline.

The FLT3 ITD or TKD mutation test result and date of testing must be provided at the time of application.

This drug is not PBS-subsidised if it is prescribed to an in-patient in a public hospital setting.

Progressive disease monitoring via a complete blood count must be taken at the end of each cycle.

If abnormal blood counts suggest the potential for relapsed AML, a bone marrow biopsy must be performed to confirm the absence of progressive disease for the patient to be eligible for further cycles.

Progressive disease is defined as the presence of any of the following:

- Leukaemic cells in the CSF;
- Re-appearance of circulating blast cells in the peripheral blood, not attributable to overshoot following recovery from myeloablative therapy;
- Greater than 5 % blasts in the marrow not attributable to bone marrow regeneration or another cause;
- Extramedullary leukaemia.

A patient who has progressive disease when treated with this drug is no longer eligible for PBS-subsidised treatment with this drug.

midostaurin 25 mg capsule, 56

11506N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	2	..	10253.89	Rydapt [NV]

■ MIDOSTAURIN

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Applications for authority to prescribe may be made by phone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday)

Authority required

Acute Myeloid Leukaemia

Treatment Phase: Maintenance therapy - Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the initial maintenance or the initial maintenance grandfathering treatment restriction, **AND**
- Patient must not have developed disease progression while receiving PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not be undergoing or have undergone a stem cell transplant.

A maximum of 9 cycles will be authorised under this restriction in a lifetime.

Progressive disease monitoring via a complete blood count must be taken at the end of each cycle.

If abnormal blood counts suggest the potential for relapsed AML, a bone marrow biopsy must be performed to confirm the absence of progressive disease for the patient to be eligible for further cycles.

Progressive disease is defined as the presence of any of the following:

- Leukaemic cells in the CSF;
- Re-appearance of circulating blast cells in the peripheral blood, not attributable to overshoot following recovery from myeloablative therapy;
- Greater than 5 % blasts in the marrow not attributable to bone marrow regeneration or another cause;
- Extramedullary leukaemia.

A patient who has progressive disease when treated with this drug is no longer eligible for PBS-subsidised treatment with this drug.

midostaurin 25 mg capsule, 112

11518F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	2	..	20460.39	Rydapt [NV]

■ MIDOSTAURIN

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Acute Myeloid Leukaemia

Treatment Phase: Maintenance therapy - Initial treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while receiving PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have demonstrated complete remission after induction and consolidation chemotherapy in combination with midostaurin, **AND**
- Patient must not be undergoing or have undergone a stem cell transplant, **AND**
- The condition must have been internal tandem duplication (ITD) or tyrosine kinase domain (TKD) FMS tyrosine kinase 3 (FLT3) mutation positive before initiating this drug for this condition.

A maximum of 3 cycles will be authorised under this restriction in a lifetime.

Progressive disease monitoring via a complete blood count must be taken at the end of each cycle.

If abnormal blood counts suggest the potential for relapsed AML, a bone marrow biopsy must be performed to confirm the absence of progressive disease for the patient to be eligible for further cycles.

Progressive disease is defined as the presence of any of the following:

- Leukaemic cells in the CSF;

- Re-appearance of circulating blast cells in the peripheral blood, not attributable to overshoot following recovery from myeloablative therapy;
- Greater than 5 % blasts in the marrow not attributable to bone marrow regeneration or another cause;
- Extramedullary leukaemia.

A patient who has progressive disease when treated with this drug is no longer eligible for PBS-subsidised treatment with this drug.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form;
- (2) a completed Acute myeloid leukaemia PBS Authority Application - Supporting Information Form; and
- (3) confirmation that the patient is not undergoing or has not undergone a stem cell transplant; and
- (4) confirmation that the patient does not have progressive disease; and
- (5) a copy of a recent bone marrow biopsy report demonstrating that the patient is in complete remission; and
- (6) a copy of the pathology test demonstrating that the condition was FMS tyrosine kinase 3 (FLT3) (ITD or TKD) mutation positive prior to commencing midostaurin.

midostaurin 25 mg capsule, 112

11531X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	2	..	20460.39	Rydapt [NV]

■ MIDOSTAURIN

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Acute Myeloid Leukaemia

Treatment Phase: Maintenance therapy - Grandfathered treatment

Clinical criteria:

- Patient must have received non-PBS subsidised treatment with this drug for this condition prior to 1 December 2018, **AND**
- Patient must be receiving treatment with this drug for this condition at the time of application, **AND**
- Patient must not have developed disease progression while receiving treatment with this drug for this condition, **AND**
- Patient must have demonstrated complete remission after induction and consolidation chemotherapy in combination with midostaurin, **AND**
- Patient must not be undergoing or have undergone a stem cell transplant, **AND**
- The condition must have been internal tandem duplication (ITD) or tyrosine kinase domain (TKD) FMS tyrosine kinase 3 (FLT3) mutation positive before initiating this drug for this condition.

A maximum of 2 cycles will be authorised under this restriction in a lifetime.

A patient may qualify for PBS-subsidised treatment under this restriction once only.

For continuing PBS-subsidised treatment, a Grandfathered patient must qualify under the maintenance therapy continuing treatment criteria.

Progressive disease monitoring via a complete blood count must be taken at the end of each cycle.

If abnormal blood counts suggest the potential for relapsed AML, a bone marrow biopsy must be performed to confirm the absence of progressive disease for the patient to be eligible for further cycles.

Progressive disease is defined as the presence of any of the following:

- Leukaemic cells in the CSF;
- Re-appearance of circulating blast cells in the peripheral blood, not attributable to overshoot following recovery from myeloablative therapy;
- Greater than 5 % blasts in the marrow not attributable to bone marrow regeneration or another cause;
- Extramedullary leukaemia.

A patient who has progressive disease when treated with this drug is no longer eligible for PBS-subsidised treatment with this drug.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form;
- (2) a completed Acute myeloid leukaemia PBS Authority Application - Supporting Information Form; and
- (3) confirmation that the patient is not undergoing or has not undergone a stem cell transplant; and
- (4) confirmation that the patient does not have progressive disease; and
- (5) a copy of a recent bone marrow biopsy report demonstrating that the patient is in complete remission; and
- (6) a copy of the pathology test demonstrating that the condition was FMS tyrosine kinase 3 (FLT3) (ITD or TKD) mutation positive prior to commencing midostaurin.

midostaurin 25 mg capsule, 112

11541K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	1	..	20460.39	Rydapt [NV]

■ IMMUNOSTIMULANTS
IMMUNOSTIMULANTS
Colony stimulating factors
■ FILGRASTIM
Authority required (STREAMLINED)
8674

Chemotherapy-induced neutropenia

Clinical criteria:

- Patient must be receiving chemotherapy with the intention of achieving a cure or a substantial remission, **AND**
- Patient must be at greater than 20% risk of developing febrile neutropenia; OR
- Patient must be at substantial risk (greater than 20%) of prolonged severe neutropenia for more than or equal to seven days.

Authority required (STREAMLINED)
8667

Chemotherapy-induced neutropenia

Clinical criteria:

- Patient must be receiving chemotherapy with the intention of achieving a cure or a substantial remission, **AND**
- Patient must have had a prior episode of febrile neutropenia; OR
- Patient must have had a prior episode of prolonged severe neutropenia for more than or equal to seven days.

Authority required (STREAMLINED)
8672

Mobilisation of peripheral blood progenitor cells

Clinical criteria:

- The treatment must be to facilitate harvest of peripheral blood progenitor cells for autologous transplantation into a patient with a non-myeloid malignancy who has had myeloablative or myelosuppressive therapy.

Authority required (STREAMLINED)
8668

Mobilisation of peripheral blood progenitor cells

Clinical criteria:

- The treatment must be in a normal volunteer for use in allogeneic transplantation.

Authority required (STREAMLINED)
8671

Assisting bone marrow transplantation

Clinical criteria:

- Patient must be receiving marrow-ablative chemotherapy prior to the transplantation.

Authority required (STREAMLINED)
8696

Assisting autologous peripheral blood progenitor cell transplantation

Clinical criteria:

- The treatment must be following marrow-ablative chemotherapy for non-myeloid malignancy prior to the transplantation.

Authority required (STREAMLINED)
8669

Severe congenital neutropenia

Clinical criteria:

- Patient must have an absolute neutrophil count of less than 100 million cells per litre measured on 3 occasions, with readings at least 2 weeks apart, **AND**
- Patient must have had a bone marrow examination that has shown evidence of maturational arrest of the neutrophil lineage.

Authority required (STREAMLINED)
8670

Severe chronic neutropenia

Clinical criteria:

- Patient must have an absolute neutrophil count of less than 1,000 million cells per litre measured on 3 occasions, with readings at least 2 weeks apart; OR
- Patient must have neutrophil dysfunction, **AND**
- Patient must have experienced a life-threatening infectious episode requiring hospitalisation and treatment with intravenous antibiotics in the previous 12 months; OR
- Patient must have had at least 3 recurrent clinically significant infections in the previous 12 months.

Authority required (STREAMLINED)
8673

Chronic cyclical neutropenia

Clinical criteria:

- Patient must have an absolute neutrophil count of less than 500 million cells per litre lasting for 3 days per cycle, measured over 3 separate cycles, **AND**
- Patient must have experienced a life-threatening infectious episode requiring hospitalisation and treatment with intravenous antibiotics; OR
- Patient must have had at least 3 recurrent clinically significant infections in the previous 12 months.

filgrastim 120 microgram/0.2 mL injection, 10 x 0.2 mL syringes

5830W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	11	..	*264.97	Nivestim [PF]

■ FILGRASTIM

Note Pharmaceutical benefits that have the forms filgrastim 480 microgram/0.5 mL injection, 10 x 0.5 mL syringes and filgrastim 480 microgram/1.6 mL injection, 10 x 1.6 mL vials and filgrastim 480 microgram/0.5 mL injection, 5 x 0.5 mL syringes are equivalent for the purposes of substitution.

Authority required (STREAMLINED)
8674

Chemotherapy-induced neutropenia

Clinical criteria:

- Patient must be receiving chemotherapy with the intention of achieving a cure or a substantial remission, **AND**
- Patient must be at greater than 20% risk of developing febrile neutropenia; OR
- Patient must be at substantial risk (greater than 20%) of prolonged severe neutropenia for more than or equal to seven days.

Authority required (STREAMLINED)
8667

Chemotherapy-induced neutropenia

Clinical criteria:

- Patient must be receiving chemotherapy with the intention of achieving a cure or a substantial remission, **AND**
- Patient must have had a prior episode of febrile neutropenia; OR
- Patient must have had a prior episode of prolonged severe neutropenia for more than or equal to seven days.

Authority required (STREAMLINED)
8672

Mobilisation of peripheral blood progenitor cells

Clinical criteria:

- The treatment must be to facilitate harvest of peripheral blood progenitor cells for autologous transplantation into a patient with a non-myeloid malignancy who has had myeloablative or myelosuppressive therapy.

Authority required (STREAMLINED)
8668

Mobilisation of peripheral blood progenitor cells

Clinical criteria:

- The treatment must be in a normal volunteer for use in allogeneic transplantation.

Authority required (STREAMLINED)
8671

Assisting bone marrow transplantation

Clinical criteria:

- Patient must be receiving marrow-ablative chemotherapy prior to the transplantation.

Authority required (STREAMLINED)
8696

Assisting autologous peripheral blood progenitor cell transplantation

Clinical criteria:

- The treatment must be following marrow-ablative chemotherapy for non-myeloid malignancy prior to the transplantation.

Authority required (STREAMLINED)
8669

Severe congenital neutropenia

Clinical criteria:

- Patient must have an absolute neutrophil count of less than 100 million cells per litre measured on 3 occasions, with readings at least 2 weeks apart, **AND**
- Patient must have had a bone marrow examination that has shown evidence of maturational arrest of the neutrophil lineage.

Authority required (STREAMLINED)
8670

Severe chronic neutropenia

Clinical criteria:

- Patient must have an absolute neutrophil count of less than 1,000 million cells per litre measured on 3 occasions, with readings at least 2 weeks apart; OR
- Patient must have neutrophil dysfunction, **AND**
- Patient must have experienced a life-threatening infectious episode requiring hospitalisation and treatment with intravenous antibiotics in the previous 12 months; OR

- Patient must have had at least 3 recurrent clinically significant infections in the previous 12 months.

Authority required (STREAMLINED)
8673

Chronic cyclical neutropenia

Clinical criteria:

- Patient must have an absolute neutrophil count of less than 500 million cells per litre lasting for 3 days per cycle, measured over 3 separate cycles, **AND**
- Patient must have experienced a life-threatening infectious episode requiring hospitalisation and treatment with intravenous antibiotics; OR
- Patient must have had at least 3 recurrent clinically significant infections in the previous 12 months.

filgrastim 480 microgram/1.6 mL injection, 10 x 1.6 mL vials

6127L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	11	..	*1039.71	^a Neupogen [AN]

filgrastim 480 microgram/0.5 mL injection, 5 x 0.5 mL syringes

2733W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	4	11	..	*1039.71	^a Zarzio [SZ]

filgrastim 480 microgram/0.5 mL injection, 10 x 0.5 mL syringes

6292E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	11	..	*1039.69	^a Neupogen [AN]	^a Nivestim [PF]

■ FILGRASTIM

Note Pharmaceutical benefits that have the forms filgrastim 300 microgram/0.5 mL injection, 10 x 0.5 mL syringes and filgrastim 300 microgram/0.5 mL injection, 5 x 0.5 mL syringes and filgrastim 300 microgram/mL injection, 10 x 1 mL vials are equivalent for the purposes of substitution.

Authority required (STREAMLINED)
8674

Chemotherapy-induced neutropenia

Clinical criteria:

- Patient must be receiving chemotherapy with the intention of achieving a cure or a substantial remission, **AND**
- Patient must be at greater than 20% risk of developing febrile neutropenia; OR
- Patient must be at substantial risk (greater than 20%) of prolonged severe neutropenia for more than or equal to seven days.

Authority required (STREAMLINED)
8667

Chemotherapy-induced neutropenia

Clinical criteria:

- Patient must be receiving chemotherapy with the intention of achieving a cure or a substantial remission, **AND**
- Patient must have had a prior episode of febrile neutropenia; OR
- Patient must have had a prior episode of prolonged severe neutropenia for more than or equal to seven days.

Authority required (STREAMLINED)
8672

Mobilisation of peripheral blood progenitor cells

Clinical criteria:

- The treatment must be to facilitate harvest of peripheral blood progenitor cells for autologous transplantation into a patient with a non-myeloid malignancy who has had myeloablative or myelosuppressive therapy.

Authority required (STREAMLINED)
8668

Mobilisation of peripheral blood progenitor cells

Clinical criteria:

- The treatment must be in a normal volunteer for use in allogeneic transplantation.

Authority required (STREAMLINED)
8671

Assisting bone marrow transplantation

Clinical criteria:

- Patient must be receiving marrow-ablative chemotherapy prior to the transplantation.

Authority required (STREAMLINED)
8696

Assisting autologous peripheral blood progenitor cell transplantation

Clinical criteria:

- The treatment must be following marrow-ablative chemotherapy for non-myeloid malignancy prior to the transplantation.

Authority required (STREAMLINED)
8669

Severe congenital neutropenia

Clinical criteria:

- Patient must have an absolute neutrophil count of less than 100 million cells per litre measured on 3 occasions, with readings at least 2 weeks apart, **AND**
- Patient must have had a bone marrow examination that has shown evidence of maturational arrest of the neutrophil lineage.

Authority required (STREAMLINED)

8670

Severe chronic neutropenia

Clinical criteria:

- Patient must have an absolute neutrophil count of less than 1,000 million cells per litre measured on 3 occasions, with readings at least 2 weeks apart; OR
- Patient must have neutrophil dysfunction, **AND**
- Patient must have experienced a life-threatening infectious episode requiring hospitalisation and treatment with intravenous antibiotics in the previous 12 months; OR
- Patient must have had at least 3 recurrent clinically significant infections in the previous 12 months.

Authority required (STREAMLINED)

8673

Chronic cyclical neutropenia

Clinical criteria:

- Patient must have an absolute neutrophil count of less than 500 million cells per litre lasting for 3 days per cycle, measured over 3 separate cycles, **AND**
- Patient must have experienced a life-threatening infectious episode requiring hospitalisation and treatment with intravenous antibiotics; OR
- Patient must have had at least 3 recurrent clinically significant infections in the previous 12 months.

filgrastim 300 microgram/0.5 mL injection, 5 x 0.5 mL syringes

2747N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	4	11	..	*651.35	^a Zarzio [SZ]

filgrastim 300 microgram/mL injection, 10 x 1 mL vials

6126K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	11	..	*651.35	^a Neupogen [AN]

filgrastim 300 microgram/0.5 mL injection, 10 x 0.5 mL syringes

6291D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	11	..	*651.35	^a Neupogen [AN]	^a Nivestim [PF]

▪ **LENOGRASTIM**

Authority required (STREAMLINED)

9265

Chemotherapy-induced neutropenia

Clinical criteria:

- Patient must be receiving standard dose adjuvant chemotherapy for breast cancer, **AND**
- Patient must have had a prior episode of febrile neutropenia; OR
- Patient must have had a prior episode of prolonged severe neutropenia (neutrophil count of less than 1,000 million cells per litre), **AND**
- The treatment must be used in a patient for whom there is a clinical justification for wishing to continue chemotherapy with the same drug combination, dosage and treatment schedule, **AND**
- Patient must be anticipated to have a good response to treatment providing chemotherapy can be delivered as planned.

Authority required (STREAMLINED)

9326

Chemotherapy-induced neutropenia

Clinical criteria:

- Patient must be receiving first-line chemotherapy for Hodgkin disease, **AND**
- Patient must have had a prior episode of febrile neutropenia; OR
- Patient must have had a prior episode of prolonged severe neutropenia (neutrophil count of less than 1,000 million cells per litre), **AND**
- The treatment must be used in a patient for whom there is a clinical justification for wishing to continue chemotherapy with the same drug combination, dosage and treatment schedule, **AND**
- Patient must be anticipated to have a good response to treatment providing chemotherapy can be delivered as planned.

Authority required (STREAMLINED)

9324

Chemotherapy-induced neutropenia

Clinical criteria:

- Patient must be receiving treatment with aggressive chemotherapy with the intention of achieving a cure or substantial remission in acute lymphoblastic leukaemia.

Authority required (STREAMLINED)

9263

Chemotherapy-induced neutropenia

Clinical criteria:

- Patient must be receiving treatment with aggressive chemotherapy with the intention of achieving a cure or substantial remission in germ cell tumours.

Authority required (STREAMLINED)
9229

Chemotherapy-induced neutropenia

Clinical criteria:

- Patient must be receiving treatment with aggressive chemotherapy with the intention of achieving a cure or substantial remission in relapsed Hodgkin disease.

Authority required (STREAMLINED)
9230

Chemotherapy-induced neutropenia

Clinical criteria:

- Patient must be receiving treatment with aggressive chemotherapy with the intention of achieving a cure or substantial remission in an infant or child with central nervous system tumours.

Authority required (STREAMLINED)
9266

Chemotherapy-induced neutropenia

Clinical criteria:

- Patient must be receiving treatment with aggressive chemotherapy with the intention of achieving a cure or substantial remission in neuroblastoma.

Authority required (STREAMLINED)
9327

Chemotherapy-induced neutropenia

Clinical criteria:

- Patient must be receiving treatment with aggressive chemotherapy with the intention of achieving a cure or substantial remission in Ewing's sarcoma.

Authority required (STREAMLINED)
9264

Chemotherapy-induced neutropenia

Clinical criteria:

- Patient must be receiving treatment with aggressive chemotherapy with the intention of achieving a cure or substantial remission in non-Hodgkin's lymphoma (intermediate or high grade).

Authority required (STREAMLINED)
9226

Chemotherapy-induced neutropenia

Clinical criteria:

- Patient must be receiving treatment with aggressive chemotherapy with the intention of achieving a cure or substantial remission in osteosarcoma.

Authority required (STREAMLINED)
9325

Chemotherapy-induced neutropenia

Clinical criteria:

- Patient must be receiving treatment with aggressive chemotherapy with the intention of achieving a cure or substantial remission in rhabdomyosarcoma.

Authority required (STREAMLINED)
9231

Mobilisation of peripheral blood progenitor cells

Clinical criteria:

- The treatment must be to facilitate harvest of peripheral blood progenitor cells for autologous transplantation into a patient with a non-myeloid malignancy who has had myeloablative or myelosuppressive therapy.

Authority required (STREAMLINED)
9314

Mobilisation of peripheral blood progenitor cells

Clinical criteria:

- The treatment must be in a normal volunteer for use in allogeneic transplantation.

Authority required (STREAMLINED)
9227

Assisting peripheral blood progenitor cell or bone marrow transplantation

Clinical criteria:

- The treatment must be following marrow-ablative chemotherapy for non-myeloid malignancy prior to the transplantation.

lenograstim 33.6 million units (263 microgram) injection, 1 vial

11530W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	20	11	..	*2132.59	Granocyte 34 [PF]

lenograstim 13.4 million units (105 microgram) injection, 1 vial

11517E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	20	11	..	*873.39	Granocyte 13 [PF]

■ LIPEGFILGRASTIM
Authority required (STREAMLINED)
9224

Chemotherapy-induced neutropenia

Clinical criteria:

- Patient must be receiving chemotherapy with the intention of achieving a cure or a substantial remission, **AND**
- Patient must be at greater than 20% risk of developing febrile neutropenia; OR
- Patient must be at substantial risk (greater than 20%) of prolonged severe neutropenia for more than or equal to seven days.

Authority required (STREAMLINED)
9322

Chemotherapy-induced neutropenia

Clinical criteria:

- Patient must be receiving chemotherapy with the intention of achieving a cure or a substantial remission, **AND**
- Patient must have had a prior episode of febrile neutropenia; OR
- Patient must have had a prior episode of prolonged severe neutropenia for more than or equal to seven days.

lipegfilgrastim 6 mg/0.6 mL injection, 0.6 mL syringe

10931H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	11	..	1222.39	Lonquex [TB]

■ PEGFILGRASTIM
Authority required (STREAMLINED)
9235

Chemotherapy-induced neutropenia

Clinical criteria:

- Patient must be receiving chemotherapy with the intention of achieving a cure or a substantial remission, **AND**
- Patient must be at greater than 20% risk of developing febrile neutropenia; OR
- Patient must be at substantial risk (greater than 20%) of prolonged severe neutropenia for more than or equal to seven days.

Authority required (STREAMLINED)
9303

Chemotherapy-induced neutropenia

Clinical criteria:

- Patient must be receiving chemotherapy with the intention of achieving a cure or a substantial remission, **AND**
- Patient must have had a prior episode of febrile neutropenia; OR
- Patient must have had a prior episode of prolonged severe neutropenia for more than or equal to seven days.

pegfilgrastim 6 mg/0.6 mL injection, 0.6 mL syringe

6363X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	11	..	1222.39	^a Neulasta [JU] ^a Tezmota [JX]	^a Ristempa [JO]

Interferons
■ INTERFERON ALFA-2A

Caution Treatment with interferon alfa has been associated with depression and suicide in some patients. Patients with a history of suicidal ideation or depressive illness should be warned of the risks. Psychiatric status during therapy should be monitored.

Authority required (STREAMLINED)
9259

Chronic Myeloid Leukaemia (CML)

Clinical criteria:

- The condition must be Philadelphia chromosome positive.

interferon alfa-2a 3 million units/0.5 mL injection, 0.5 mL syringe

6210W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	30	5	..	*762.79	Roferon-A [RO]

interferon alfa-2a 9 million units/0.5 mL injection, 0.5 mL syringe

6213B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	30	5	..	*2225.29	Roferon-A [RO]

■ INTERFERON GAMMA-1B
Authority required (STREAMLINED)
9639

Chronic granulomatous disease

Clinical criteria:

- Patient must have frequent and severe infections despite adequate prophylaxis with antimicrobial agents.

interferon gamma-1b 2 million units (100 microgram)/0.5 mL injection, 6 x 0.5 mL vials

6148N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	11	..	*2258.19	Imukin [EU]

■ **PEGINTERFERON ALFA-2A**

Caution Treatment with peginterferon alfa has been associated with depression and suicide in some patients. Patients with a history of suicidal ideation or depressive illness should be warned of the risks. Psychiatric status during therapy should be monitored.

Note Treatment centres are required to have access to the following appropriate specialist facilities for the provision of clinical support services for hepatitis C:

- (a) a nurse educator/counsellor for patients; and
- (b) 24-hour access by patients to medical advice; and
- (c) an established liver clinic.

Authority required (STREAMLINED)

9603

Chronic hepatitis C infection

Treatment criteria:

- Must be treated in an accredited treatment centre.

Population criteria:

- Patient must be aged 18 years or older, **AND**
- Patient must not be pregnant or breastfeeding, and must be using an effective form of contraception if female and of child-bearing age.

Clinical criteria:

- Patient must have compensated liver disease, **AND**
- Patient must not have received prior interferon alfa or peginterferon alfa treatment for hepatitis C, **AND**
- Patient must have a contraindication to ribavirin, **AND**
- The treatment must cease unless the results of an HCV RNA quantitative assay at week 12 (performed at the same laboratory using the same test) show that plasma HCV RNA has become undetectable or the viral load has decreased by at least a 2 log drop, **AND**
- The treatment must be limited to a maximum duration of 48 weeks.

Evidence of chronic hepatitis C infection (repeatedly anti-HCV positive and HCV RNA positive) must be documented in the patient's medical records.

peginterferon alfa-2a 135 microgram/0.5 mL injection, 4 x 0.5 mL syringes

6439X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*1155.01	Pegasys [RO]

peginterferon alfa-2a 180 microgram/0.5 mL injection, 4 x 0.5 mL syringes

6449K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*1330.11	Pegasys [RO]

■ **PEGINTERFERON ALFA-2A**

Caution Treatment with peginterferon alfa has been associated with depression and suicide in some patients. Patients with a history of suicidal ideation or depressive illness should be warned of the risks. Psychiatric status during therapy should be monitored.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Chronic hepatitis C infection

Clinical criteria:

- Patient must meet the criteria set out in the General Statement for Drugs for the Treatment of Hepatitis C, **AND**
- Patient must be taking this drug as part of a regimen set out in the matrix in the General Statement for Drugs for the Treatment of Hepatitis C, based on the hepatitis C virus genotype, patient treatment history and cirrhotic status, **AND**
- The treatment must be limited to a maximum duration of 12 weeks.

peginterferon alfa-2a 180 microgram/0.5 mL injection, 4 x 0.5 mL syringes

11044G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	2	..	674.40	Pegasys [RO]

Other immunostimulants

■ **PLERIXAFOR**

Note Special Pricing Arrangements apply.

Note Applications for increased maximum quantities will only be authorised for patients with body weight greater than 100 kg.

Authority required (STREAMLINED)

9329

Mobilisation of haematopoietic stem cells

Clinical criteria:

- The treatment must be in combination with granulocyte-colony stimulating factor (G-CSF), **AND**
- Patient must have lymphoma; OR
- Patient must have multiple myeloma, **AND**
- Patient must require autologous stem cell transplantation, **AND**
- Patient must have failed previous stem cell collection; OR
- Patient must be undergoing chemotherapy plus G-CSF mobilisation and their peripheral blood CD34+ count is less than 10,000 per millilitre or less than 10 million per litre on the day of planned collection; OR
- Patient must be undergoing chemotherapy plus G-CSF mobilisation and the first apheresis has yielded less than 1 million CD34+ cells/kg.

Evidence that the patient meets the PBS restriction criteria must be recorded in the patient's medical records.

plerixafor 24 mg/1.2 mL injection, 1.2 mL vial

10084R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	1	..	7038.39	Mozobil [GZ]

IMMUNOSUPPRESSANTS

IMMUNOSUPPRESSANTS

Selective immunosuppressants

ABATACEPT

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for adults with severe active rheumatoid arthritis. Where the term biological medicine appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alfa antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab), the T-cell co-stimulation modulator (abatacept) and the Janus kinase (JAK) inhibitors (baricitinib, tofacitinib). A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised biological medicine therapy may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once, and
- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised biological medicines for the treatment of rheumatoid arthritis.

Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Stevens Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

For a patient who has failed PBS-subsidised treatment with 2 or 3 TNF-alfa antagonists prior to 1 August 2010 please contact the Department of Human Services on 1800 700 270. A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction.

A patient who has failed fewer than 5 biological medicines and who has a break in treatment of less than 24 months may commence a further course of treatment with a biological medicine under Initial 2 treatment restriction. A patient who has failed fewer than 5 biological medicines and who has had a break in therapy of longer than 24 months may commence a further course of treatment with a biological medicine under the Initial 3 treatment restriction. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine treatment is stopped to the date of the new application for treatment with a biological medicine.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 April 2019.

(a) Initial treatment.

Applications for initial treatment should be made where:

- a patient has received no prior PBS-subsidised biological medicine treatment and wishes to commence such therapy, excluding rituximab (Initial 1 - new patient); or
 - a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months) [further details are under 'Swapping therapy' below]; or
 - a patient wishes to re-commence treatment with a specific biological medicine following a break of less than 24 months in PBS-subsidised therapy with that agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months).
 - a patient wishes to re-commence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 24 months (Initial 3 - re-commencement of treatment after a break in biological medicine of more than 24 months)
- Initial applications for a new patient (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6-month intensive DMARD trial, but prior to ceasing DMARD therapy.
- Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, baricitinib, etanercept, golimumab, tocilizumab and tofacitinib, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

Rituximab patients should be assessed following a minimum of 12 weeks after the first infusion, and the assessment should be submitted to the Department of Human Services within 4 weeks to ensure continuity of treatment for those who meet the continuing restriction for PBS subsidised treatment with this drug for this condition.

Where a response assessment is not provided the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment. For second and subsequent courses of PBS-subsidised biological medicine (excluding rituximab) treatment it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to the Department of Human Services within sufficient timeframe to allow for authority assessment.

Abatacept patients:

A patient is eligible to receive one I.V. loading dose when commencing treatment with the subcutaneous formulation. Two prescriptions are required, the first prescription for the I.V. loading dose for sufficient vials for one dose based on the patient's weight with no repeats. The second prescription for the subcutaneous formulation, with a maximum quantity of 4 and up to 3 repeats, must be submitted with the initial application.

Rituximab patients:

A further application may be submitted to the Department of Human Services within sufficient timeframe to allow authority assessment. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

Infliximab and etanercept patients:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction. Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

(2) Swapping therapy

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non- biological medicine therapy requirements except if the patient has had a break in therapy of more than 24 months who would need to requalify with respect to the indices of disease severity. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each biological medicine trialled.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent, unless they have a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug.

Abatacept:

A patient swapping from I.V. abatacept to subcutaneous abatacept will not be eligible for an I.V. loading dose when commencing treatment with the subcutaneous formulation.

Rituximab:

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF-alfa antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological medicine during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological medicine therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate biological medicine may do so without having to have any treatment-free period.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and the Department of Human Services will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be

demonstrated on the total number of major joints.

Applications under the Initial 1 treatment restriction for a new patient must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy.

Applications under the Initial 3 treatment restriction for re-commencement of treatment after a break in biological medicine of more than 24 months must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with disease modifying anti-rheumatic drugs (DMARDs) which must include at least 3 months continuous treatment with each of at least 2 DMARDs, one of which must be methotrexate at a dose of at least 20 mg weekly and one of which must be: (i) hydroxychloroquine at a dose of at least 200 mg daily; or (ii) leflunomide at a dose of at least 10 mg daily; or (iii) sulfasalazine at a dose of at least 2 g daily; OR
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs which, if methotrexate is contraindicated according to the Therapeutic Goods Administration (TGA)-approved Product Information or cannot be tolerated at a 20 mg weekly dose, must include at least 3 months continuous treatment with each of at least 2 of the following DMARDs: (i) hydroxychloroquine at a dose of at least 200 mg daily; and/or (ii) leflunomide at a dose of at least 10 mg daily; and/or (iii) sulfasalazine at a dose of at least 2 g daily; OR
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs which, if 3 or more of methotrexate, hydroxychloroquine, leflunomide and sulfasalazine are contraindicated according to the relevant TGA-approved Product Information or cannot be tolerated at the doses specified above, must include at least 3 months continuous treatment with each of at least 2 DMARDs, with one or more of the following DMARDs being used in place of the DMARDs which are contraindicated or not tolerated: (i) azathioprine at a dose of at least 1 mg/kg per day; and/or (ii) cyclosporin at a dose of at least 2 mg/kg/day; and/or (iii) sodium aurothiomalate at a dose of 50 mg weekly, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction, **AND**
- The treatment must be given concomitantly with methotrexate at a dose of at least 7.5 mg weekly.

Population criteria:

- Patient must be aged 18 years or older.

If methotrexate is contraindicated according to the TGA-approved product information or cannot be tolerated at a 20 mg weekly dose, the application must include details of the contraindication or intolerance including severity to methotrexate. The maximum tolerated dose of methotrexate must be documented in the application, if applicable.

The application must include details of the DMARDs trialled, their doses and duration of treatment, and all relevant contraindications and/or intolerances including severity.

The requirement to trial at least 2 DMARDs for periods of at least 3 months each can be met using single agents sequentially or by using one or more combinations of DMARDs.

If the requirement to trial 6 months of intensive DMARD therapy with at least 2 DMARDs cannot be met because of contraindications and/or intolerances of a severity necessitating permanent treatment withdrawal to all of the DMARDs specified above, details of the contraindication or intolerance including severity and dose for each DMARD must be provided in the authority application.

The following criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; **AND** either

- (a) a total active joint count of at least 20 active (swollen and tender) joints; or
- (b) at least 4 active joints from the following list of major joints:
 - (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The joint count and ESR and/or CRP must be determined at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. All measures must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

At the time of authority application, medical practitioners should request the appropriate number of vials to provide sufficient drug, based on the weight of the patient, for a single infusion.

Up to a maximum of 4 repeats will be authorised.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and

(2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note The Department of Human Services website (www.humanservices.gov.au) has details of the toxicities, including severity, which will be accepted for the following purposes:

- (a) exempting a patient from the requirement to undertake a minimum 3 month trial of methotrexate at a 20 mg weekly dose;
- (b) substituting azathioprine, cyclosporin or sodium aurothiomalate for another DMARD as part of the 6 month intensive DMARD trial;
- (c) exempting a patient from the requirement for a 6 month trial of intensive DMARD therapy.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 2 (change or re-commencement of treatment after a break in biological medicine of less than 24 months).

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must not have failed to respond to previous PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised biological medicine treatment for this condition 5 times, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction, **AND**
- The treatment must be given concomitantly with methotrexate at a dose of at least 7.5 mg weekly.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either of the Initial 1, Initial 2, Initial 3, or continuing treatment restrictions, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

At the time of authority application, medical practitioners should request the appropriate number of vials to provide sufficient drug, based on the weight of the patient, for a single infusion.

Up to a maximum of 4 repeats will be authorised.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

A patient who has demonstrated a response to a course of rituximab must have a PBS-subsidised biological therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 3 (re-commencement of treatment after a break in biological medicine of more than 24 months)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 24 months or more from the most recent PBS-subsidised biological medicine for this condition, **AND**
- Patient must not have failed to respond to previous PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised biological medicine treatment for this condition 5 times, **AND**
- The condition must have an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; OR
- The condition must have a C-reactive protein (CRP) level greater than 15 mg per L, **AND**
- The condition must have either (a) a total active joint count of at least 20 active (swollen and tender) joints; or (b) at least 4 active major joints, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction, **AND**
- The treatment must be given concomitantly with methotrexate at a dose of at least 7.5 mg weekly.

Population criteria:

- Patient must be aged 18 years or older.

Major joints are defined as (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

All measures of joint count and ESR and/or CRP must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au
Applications for authority to prescribe should be forwarded to:
Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial 1 (new patient) or Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 24 months) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 24 months) - balance of supply

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 24 months) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 24 months) to complete 16 weeks of treatment, **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction, **AND**
- The treatment must be given concomitantly with methotrexate at a dose of at least 7.5 mg weekly.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient has either failed or ceased to respond to a PBS-subsidised biological medicine for this condition 5 times, they will not be eligible to receive further PBS-subsidised treatment with a biological medicine for this condition.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au
Applications for authority to prescribe should be forwarded to:
Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Continuing Treatment - balance of supply.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restriction.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

abatacept 250 mg injection, 1 vial

9621J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	284.38	Orencia [BQ]

■ ALEMTUZUMAB

Note Neurologists prescribing PBS-subsidised alemtuzumab must be registered with the Lemtrada monitoring program.

Note Special Pricing Arrangements apply.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

9589

Multiple sclerosis

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not show continuing progression of disability while on treatment with this drug, **AND**
- Patient must not receive more than one PBS-subsidised treatment per year, **AND**
- The treatment must be a sole PBS-subsidised disease modifying therapy for this condition, **AND**
- Patient must have demonstrated compliance with, and an ability to tolerate this therapy.

Treatment criteria:

- Must be treated by a neurologist.

alemtuzumab 12 mg/1.2 mL injection, 1.2 mL vial

10246G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	3	..	..	*34229.38	Lemtrada [GZ]

■ ALEMTUZUMAB

Note Neurologists prescribing PBS-subsidised alemtuzumab must be registered with the Lemtrada monitoring program.

Note Special Pricing Arrangements apply.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

9636

Multiple sclerosis

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be diagnosed as clinically definite relapsing-remitting multiple sclerosis by magnetic resonance imaging of the brain and/or spinal cord; OR
- The condition must be diagnosed as clinically definite relapsing-remitting multiple sclerosis by accompanying written certification provided by a radiologist that a magnetic resonance imaging scan is contraindicated because of the risk of physical (not psychological) injury to the patient, **AND**
- The treatment must be a sole PBS-subsidised disease modifying therapy for this condition, **AND**
- Patient must have experienced at least 2 documented attacks of neurological dysfunction, believed to be due to multiple sclerosis, in the preceding 2 years of commencing a PBS-subsidised disease modifying therapy for this condition, **AND**
- Patient must be ambulatory (without assistance or support).

Treatment criteria:

- Must be treated by a neurologist.

Where applicable, the date of the magnetic resonance imaging scan must be recorded in the patient's medical records.

alemtuzumab 12 mg/1.2 mL injection, 1.2 mL vial

10243D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	5	..	..	*57017.39	Lemtrada [GZ]

■ ECULIZUMAB

Note At the time of authority application, medical practitioners must request the appropriate number of vials to provide sufficient drug for four weeks of treatment, according to the specified dosage in the approved Product Information (PI).

Applications for treatment with this drug where the dose and dosing frequency exceeds that specified in the approved PI will not be approved.

Note WARNING: Eculizumab increases the risk of meningococcal infections (septicaemia and/or meningitis).

Please consult the approved PI for information about vaccination against meningococcal infection.

Note Eculizumab is not PBS subsidised to treat TMA caused by conditions other than aHUS, such as TMA occurring in the setting of, but not limited to:

- Active malignancy;
- Active HIV infection;
- Hematopoietic stem cell transplants;
- Various drugs including quinine, high-dose calcineurin inhibitors, antiplatelet agents;
- Certain chemotherapy drugs or immunosuppressant drugs associated with microangiopathic haemolytic anaemia/TMA;
- Active autoimmune diseases;

In cases where alternative causes of TMA have not been adequately excluded, additional information may be required from the prescriber to clarify the diagnosis before approval of the application.

Note The Authority application should be accompanied by a cover letter from the prescriber, providing complete details on:

- Presenting clinical features, including history, acute treatment and medications;
- Results of testing for genetic mutations (if available);
- Family history of aHUS, especially in first-degree relatives;
- Patient's prior history of episodes of active and progressing TMA caused by aHUS;
- Exclusion of alternative causes of TMA;
- History of renal or other organ transplant (if any);
- Any other matters considered relevant by the prescriber.

In cases where there are discordant results (for example, an equivocal biopsy result in the absence of objective evidence of haemolysis) the cover letter should articulate the prescriber's interpretation of the clinical data and how a diagnosis of aHUS is supported by the available evidence.

Note The Authority application should include the results of screening for genetic mutations known to confer a high risk of developing aHUS. The results of genetic screening should be provided whether or not a high-risk mutation has been identified.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Written applications for authority to prescribe must be submitted to Department of Human Services. Human Services will then contact the prescriber by telephone.

Authority required

Atypical haemolytic uraemic syndrome (aHUS)

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have active and progressing thrombotic microangiopathy (TMA) caused by aHUS, **AND**
- Patient must have ADAMTS-13 activity of greater than or equal to 10% on a blood sample taken prior to plasma exchange or infusion; or, if ADAMTS-13 activity was not collected prior to plasma exchange or infusion, patient must have platelet counts of greater than $30 \times 10^9/L$ and a serum creatinine of greater than $150 \mu\text{mol/L}$, **AND**
- Patient must have a confirmed negative STEC (Shiga toxin-producing E.Coli) result if the patient has had diarrhoea in the preceding 14 days, **AND**
- Patient must have clinical features of active organ damage or impairment, **AND**
- Patient must not receive more than 4 weeks of treatment under this restriction.

Treatment criteria:

- Must be treated by a paediatric nephrologist, a nephrologist, a paediatric haematologist or a haematologist, or, must be in consultation with a paediatric nephrologist, a nephrologist, a paediatric haematologist or a haematologist.

Evidence of active and progressing TMA is defined by the following:

(1) a platelet count of less than $150 \times 10^9/L$; and evidence of two of the following:

- presence of schistocytes on blood film;
- low or absent haptoglobin;
- lactate dehydrogenase (LDH) above normal range;

OR

(2) in recipients of a kidney transplant for end-stage kidney disease due to aHUS, a kidney biopsy confirming TMA; **AND**

(3) evidence of at least one of the following clinical features of active TMA-related organ damage or impairment is defined as below:

(a) kidney impairment as demonstrated by one of the following:

- (i) a decline in estimated Glomerular Filtration Rate (eGFR) of greater than 20% in a patient who has pre-existing kidney impairment; and/or
- (ii) a serum creatinine (sCr) of greater than the upper limit of normal (ULN) in a patient who has no history of pre-existing kidney impairment; or
- (iii) a sCr of greater than the age-appropriate ULN in paediatric patients; or
- (iv) a renal biopsy consistent with aHUS;
- (b) onset of TMA-related neurological impairment;
- (c) onset of TMA-related cardiac impairment;
- (d) onset of TMA-related gastrointestinal impairment;
- (e) onset of TMA-related pulmonary impairment.

Claims of non-renal TMA-related organ damage should be made at the point of application for initial PBS-subsidised eculizumab (where possible), and should be supported by objective clinical measures. The prescriber's cover letter should establish that the observed organ damage is directly linked to active and progressing TMA, particularly when indirect causes such as severe thrombocytopenia, hypertension and acute renal failure are present at the time of the initial organ impairment.

Serial haematological results (every 3 months while the patient is receiving treatment) must be provided with every subsequent application for treatment.

The authority application must be in writing and must include:

- (1) A completed authority prescription form; and
- (2) A completed aHUS eculizumab Authority Application Supporting Information Form - Initial PBS-subsidised eculizumab treatment; and
- (3) A signed patient acknowledgement or an acknowledgement signed by a parent or authorised guardian, if applicable; and
- (4) A detailed cover letter from the prescriber; and
- (5) A copy of a current Certificate of vaccination or a statement that vaccination has or will be administered and appropriate antibiotic prophylaxis has been prescribed; and
- (6) A measurement of body weight at the time of application; and
- (7) The result of ADAMTS-13 activity on a blood sample taken prior to plasma exchange or infusion; the date and time that the sample for the ADAMTS-13 assay was collected, and the dates and times of any plasma exchanges or infusions that were undertaken in the two weeks prior to collection of the ADAMTS-13 assay; and
- (8) In the case that a sample for ADAMTS-13 assay was not collected prior to plasma exchange or infusion, measurement of ADAMTS-13 activity must be taken 1-2 weeks following the last plasma exchange or infusion. The ADAMTS-13 result must be submitted to the Department of Human Services within 27 days of commencement of eculizumab treatment in order for the patient to be considered as eligible for further PBS-subsidised eculizumab treatment, under **Initial treatment 1-balance of supply**; and
- (9) A confirmed negative STEC result if the patient has had diarrhoea in the preceding 14 days; and
- (10) Evidence of active and progressing TMA, including pathology results where relevant. Evidence of the onset of TMA-related neurological, cardiac, gastrointestinal or pulmonary impairment requires a supporting statement with clinical evidence in patient records. All tests must have been performed within one month of application; and
- (11) For all patients, a recent measurement of eGFR, platelets and two of either LDH, haptoglobin or schistocytes of no more than 1 week old at the time of application.

eculizumab 300 mg/30 mL injection, 30 mL vial

10182X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	5984.89	Soliris [XI]

■ ECULIZUMAB

Note At the time of authority application, medical practitioners must request the appropriate number of vials to provide sufficient drug for 4 weeks and up to 4 repeats, according to the specified dosage in the approved Product Information (PI). Applications for treatment with this drug where the dose and dosing frequency exceeds that specified in the approved PI will not be approved.

Note WARNING: Eculizumab increases the risk of meningococcal infections (septicaemia and/or meningitis). Please consult the approved PI for information about vaccination against meningococcal infection.

Note Eculizumab is not PBS subsidised to treat TMA caused by conditions other than aHUS, such as TMA occurring in the setting of, but not limited to:

- a) Active malignancy;
- b) Active HIV infection;
- c) Hematopoietic stem cell transplants;
- d) Various drugs including quinine, high-dose calcineurin inhibitors, antiplatelet agents;
- e) Certain chemotherapy drugs or immunosuppressant drugs associated with microangiopathic haemolytic anaemia/TMA;
- f) Active autoimmune diseases;

In cases where alternative causes of TMA have not been adequately excluded, additional information may be required from the prescriber to clarify the diagnosis before approval of the application.

Note The Authority application should be accompanied by a cover letter from the prescriber, providing complete details on:

- a) Presenting clinical features, including history, acute treatment and medications;
- b) Results of testing for genetic mutations (if available);
- c) Family history of aHUS, especially in first-degree relatives;
- d) Patient's prior history of episodes of active and progressing TMA caused by aHUS;
- e) Exclusion of alternative causes of TMA;
- f) History of renal or other organ transplant (if any);
- g) Any other matters considered relevant by the prescriber.

In cases where there are discordant results (for example, an equivocal biopsy result in the absence of objective evidence of haemolysis) the cover letter should articulate the prescriber's interpretation of the clinical data and how a diagnosis of aHUS is supported by the available evidence.

Note The Authority application should include the results of screening for genetic mutations known to confer a high risk of developing aHUS. The results of genetic screening should be provided whether or not a high-risk mutation has been identified.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Written applications for authority to prescribe must be submitted to Department of Human Services. Human Services will then contact the prescriber by telephone.

Authority required

Atypical haemolytic uraemic syndrome (aHUS)

Treatment Phase: Initial treatment - Balance of Supply

Treatment criteria:

- Must be treated by a paediatric nephrologist, a nephrologist, a paediatric haematologist or a haematologist, or, must be in consultation with a paediatric nephrologist, a nephrologist, a paediatric haematologist or a haematologist.

Clinical criteria:

- Patient must have received PBS-subsidised initial supply of eculizumab for this condition, **AND**
- Patient must have ADAMTS-13 activity of greater than or equal to 10% on a blood sample, **AND**
- Patient must not receive more than 20 weeks supply under this restriction.

ADAMTS-13 activity result must have been submitted to the Department of Human Services. In the case that a sample for ADAMTS-13 activity taken prior to plasma exchange or infusion was not available at the time of application for **Initial Treatment**, ADAMTS-13 activity must have been measured 1-2 weeks following the last plasma exchange or infusion, and must have been submitted to the Department of Human Services within 27 days of commencement of eculizumab. The date and time that the sample for the ADAMTS-13 assay was collected, and the dates and times of the last, if any, plasma exchange or infusion that was undertaken in the two weeks prior to collection of the ADAMTS-13 assay must also have been provided to Department of Human Services.

Serial haematological results (every 3 months while the patient is receiving treatment) must be provided with every subsequent application for treatment.

eculizumab 300 mg/30 mL injection, 30 mL vial

10192K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	4	..	5984.89	Soliris [XI]

■ ECULIZUMAB

Note At the time of authority application, medical practitioners must request the appropriate number of vials to provide sufficient drug for 4 weeks and up to 6 repeats, according to the specified dosage in the approved Product Information (PI). Applications for treatment with this drug where the dose and dosing frequency exceeds that specified in the approved PI will not be approved.

Note For patients who have received continuing treatment with PBS-subsidised eculizumab prior to 1 January 2016, this restriction is limited to 28 weeks of therapy.

Note WARNING: Eculizumab increases the risk of meningococcal infections (septicaemia and/or meningitis). Please consult the approved PI for information about vaccination against meningococcal infection.

Note Eculizumab is not PBS subsidised to treat TMA caused by conditions other than aHUS, such as TMA occurring in the setting of, but not limited to:

- a) Active malignancy;
- b) Active HIV infection;
- c) Hematopoietic stem cell transplants;
- d) Various drugs including quinine, high-dose calcineurin inhibitors, antiplatelet agents;
- e) Certain chemotherapy drugs or immunosuppressant drugs associated with microangiopathic haemolytic anaemia/TMA;
- f) Active autoimmune diseases;

In cases where alternative causes of TMA have not been adequately excluded, additional information may be required from the prescriber to clarify the diagnosis before approval of the application.

Note The Authority application should be accompanied by a cover letter from the prescriber, providing complete details on:

- a) Presenting clinical features, including history, acute treatment and medications;
- b) Results of testing for genetic mutations (if available);
- c) Family history of aHUS, especially in first-degree relatives;
- d) Patient's prior history of episodes of active and progressing TMA caused by aHUS;
- e) Exclusion of alternative causes of TMA;
- f) History of renal or other organ transplant (if any);
- g) Any other matters considered relevant by the prescriber.

In cases where there are discordant results (for example, an equivocal biopsy result in the absence of objective evidence of haemolysis) the cover letter should articulate the prescriber's interpretation of the clinical data and how a diagnosis of aHUS is supported by the available evidence.

Note The Authority application should include the results of screening for genetic mutations known to confer a high risk of developing aHUS. The results of genetic screening should be provided whether or not a high-risk mutation has been identified.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au. Written applications for authority to prescribe must be submitted to Department of Human Services. Human Services will then contact the prescriber by telephone.

Authority required

Atypical haemolytic uraemic syndrome (aHUS)

Treatment Phase: Extended initial treatment - Assessment phase

Clinical criteria:

- Patient must have received treatment under the initial restriction with PBS subsidised eculizumab for this condition, **AND**
- Patient must have demonstrated on-going treatment response of PBS-subsidised eculizumab treatment for this condition, **AND**
- Patient must not have experienced treatment failure with eculizumab including PBS-subsidised eculizumab for this condition, **AND**
- Patient must not receive more than 56 weeks of treatment under this restriction.

Treatment criteria:

- Must be treated by a paediatric nephrologist, a nephrologist, a paediatric haematologist or a haematologist, or, must be in consultation with a paediatric nephrologist, a nephrologist, a paediatric haematologist or a haematologist.

A treatment response is defined as:

(1) Normalisation of haematology as demonstrated by at least 2 of the following: platelet count, haptoglobin, and LDH; **AND**

(2) One of the following:

a) An increase in eGFR of > 25% from baseline, where the baseline is the eGFR measurement immediately prior to commencing treatment with eculizumab or

b) an eGFR within +/- 25% from baseline; or

c) an avoidance of dialysis-dependence but worsening of kidney function with a reduction in eGFR 25% from baseline.

PBS-subsidised treatment with eculizumab will not be permitted if a patient has experienced treatment failure.

A treatment failure is defined as a patient who is:

(1) dialysis-dependent at the time of application and has failed to demonstrate significant resolution of extra-renal complications if originally presented; or

(2) on dialysis and has been on dialysis for 4 months of the previous 6 months while receiving PBS-subsidised eculizumab and has failed to demonstrate significant resolution of extra-renal complications if originally presented.

A maximum of up to 56 weeks of treatment is allowed under this restriction, however an application must be submitted at 6 months, 12 months, 18 months and 24 months following commencing PBS-subsidised eculizumab.

The authority application must include the following measures of response to the prior course of treatment, including serial haematological results (every 3 months while the patient is receiving treatment).

The authority application must be in writing and must include:

- (1) A completed authority prescription form; and
- (2) A completed aHUS eculizumab Authority Application Supporting Information Form for Extended Initial treatment; and
- (3) A detailed cover letter from the prescriber; and
- (4) A copy of a current Certificate of vaccination or a statement that vaccination has or will be administered and appropriate antibiotic prophylaxis has been prescribed; and
- (5) A measurement of body weight at the time of application; and
- (6) An identified genetic mutation, if applicable; and
- (7) A family history of aHUS, if applicable; and
- (8) A history of multiple episodes of aHUS before commencing eculizumab treatment, if applicable; and
- (9) A history of kidney transplant, if applicable, (especially if required due to aHUS); and
- (10) An inclusion of the individual consequences of recurrent disease, if applicable; and
- (11) Evidence that the patient has had a treatment response including haematological results of no more than 1 week old at the time of application (platelet count, haptoglobin and LDH); and an eGFR level of no more than 1 week old at the time of application; and
- (12) Evidence that the patient has not experienced treatment failure, including a supporting statement with clinical evidence that the patient does not require dialysis, unless the indication for continuing eculizumab is severe extra-renal complications that have significantly improved; and
- (13) If the indication for continuing eculizumab is severe extra-renal complications, then a supporting statement with clinical evidence that any initial extra-renal complications of TMA have significantly improved is required.

This assessment must be submitted no later than 4 weeks from the cessation of the prior treatment. Where a response assessment is not undertaken and submitted within these timeframes, the patient will be deemed to have failed to respond to treatment with eculizumab.

eculizumab 300 mg/30 mL injection, 30 mL vial

10521R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	6	..	5984.89	Soliris [XI]

■ ECULIZUMAB

Note At the time of authority application, medical practitioners must request the appropriate number of vials to provide sufficient drug for 4 weeks and up to 5 repeats, according to the specified dosage in the approved Product Information (PI).

Applications for treatment with this drug where the dose and dosing frequency exceeds that specified in the approved PI will not be approved.

Note WARNING: Eculizumab increases the risk of meningococcal infections (septicaemia and/or meningitis).

Please consult the approved PI for information about vaccination against meningococcal infection.

Note Eculizumab is not PBS subsidised to treat TMA caused by conditions other than aHUS, such as TMA occurring in the setting of, but not limited to:

- a) Active malignancy;
- b) Active HIV infection;
- c) Hematopoietic stem cell transplants;
- d) Various drugs including quinine, high-dose calcineurin inhibitors, antiplatelet agents;
- e) Certain chemotherapy drugs or immunosuppressant drugs associated with microangiopathic haemolytic anaemia/TMA;
- f) Active autoimmune diseases;

In cases where alternative causes of TMA have not been adequately excluded, additional information may be required from the prescriber to clarify the diagnosis before approval of the application.

Note The Authority application should be accompanied by a cover letter from the prescriber, providing complete details on:

- a) Presenting clinical features, including history, acute treatment and medications;
- b) Results of testing for genetic mutations (if available);
- c) Family history of aHUS, especially in first-degree relatives;
- d) Patient's prior history of episodes of active and progressing TMA caused by aHUS;
- e) Exclusion of alternative causes of TMA;
- f) History of renal or other organ transplant (if any);
- g) Any other matters considered relevant by the prescriber.

In cases where there are discordant results (for example, an equivocal biopsy result in the absence of objective evidence of haemolysis) the cover letter should articulate the prescriber's interpretation of the clinical data and how a diagnosis of aHUS is supported by the available evidence.

Note The Authority application should include the results of screening for genetic mutations known to confer a high risk of developing aHUS. The results of genetic screening should be provided whether or not a high-risk mutation has been identified.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Written applications for authority to prescribe must be submitted to Department of Human Services. Human Services will then contact the prescriber by telephone.

Authority required

Atypical haemolytic uraemic syndrome (aHUS)

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have received treatment under Extended Initial restriction with PBS subsidised eculizumab for this condition, **AND**
- Patient must have demonstrated on-going treatment response of PBS-subsidised eculizumab treatment for this condition, **AND**
- Patient must not have experienced treatment failure with eculizumab including PBS-subsidised eculizumab for this condition, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Treatment criteria:

- Must be treated by a paediatric nephrologist, a nephrologist, a paediatric haematologist or a haematologist, or, must be in consultation with a paediatric nephrologist, a nephrologist, a paediatric haematologist or a haematologist.

A treatment response is defined as:

- (1) Normalisation of haematology as demonstrated by at least 2 of the following: platelet count, haptoglobin, and LDH; **AND**
- (2) One of the following:

a) An increase in eGFR of > 25% from baseline, where the baseline is the eGFR measurement immediately prior to commencing treatment with eculizumab or

b) an eGFR within +/- 25% from baseline; or

c) an avoidance of dialysis-dependence but worsening of kidney function with a reduction in eGFR 25% from baseline.

PBS-subsidised treatment with eculizumab will not be permitted if a patient has experienced treatment failure.

A treatment failure is defined as a patient who is:

- (1) dialysis-dependent at the time of application and has failed to demonstrate significant resolution of extra-renal complications if originally presented; or
- (2) on dialysis and has been on dialysis for 4 months of the previous 6 months while receiving PBS-subsidised eculizumab and has failed to demonstrate significant resolution of extra-renal complications if originally presented.

The authority application must include the following measures of response to the prior course of treatment, including serial haematological results (every 3 months while the patient is receiving treatment).

The authority application must be in writing and must include:

- (1) A completed authority prescription form; and
- (2) A completed aHUS eculizumab Authority Application Supporting Information Form for Continuing treatment; and
- (3) A detailed cover letter from the prescriber; and
- (4) A copy of a current Certificate of vaccination or a statement that vaccination has or will be administered and appropriate antibiotic prophylaxis has been prescribed; and
- (5) A measurement of body weight at the time of application; and
- (6) An identified genetic mutation, if applicable; and

- (7) A family history of aHUS, if applicable; and
- (8) A history of multiple episodes of aHUS before recommencing eculizumab treatment, if applicable; and
- (9) A history of kidney transplant if applicable (especially if required due to aHUS); and
- (10) An inclusion of the individual consequences of recurrent disease, if applicable; and
- (11) Evidence that the patient has had a treatment response including haematological results of no more than 1 week old at the time of application (platelet count, haptoglobin and LDH); and an eGFR level of no more than 1 week old at the time of application; and
- (12) Evidence that the patient has not experienced treatment failure, including a supporting statement with clinical evidence that the patient does not require dialysis, unless the indication for continuing eculizumab is severe extra-renal complications that have significantly improved; and
- (13) If the indication for continuing eculizumab is severe extra-renal complications, then a supporting statement with clinical evidence that any initial extra-renal complications of TMA have significantly improved is required.

This assessment must be submitted no later than 4 weeks from the cessation of the prior treatment. Where a response assessment is not undertaken and submitted within these timeframes, the patient will be deemed to have failed to respond to treatment with eculizumab.

Authority required

Atypical haemolytic uraemic syndrome (aHUS)

Treatment Phase: Extended Continuing treatment

Clinical criteria:

- Patient must have received treatment under the Continuing treatment with PBS-subsidised eculizumab for this condition, **AND**
- Patient must have demonstrated on-going treatment response with PBS-subsidised eculizumab for this condition, **AND**
- Patient must not have ever experienced treatment failure with eculizumab including PBS-subsidised eculizumab for this condition, **AND**
- Patient must have a TMA-related cardiomyopathy as evidenced by left ventricular ejection fraction < 40% on current objective measurement; OR
- Patient must have severe TMA-related neurological impairment; OR
- Patient must have severe TMA-related gastrointestinal impairment; OR
- Patient must have severe TMA-related pulmonary impairment on current objective measurement; OR
- Patient must have grade 4 or 5 chronic kidney disease (eGFR of less than 30 mL/min); OR
- Patient must have a high risk of aHUS recurrence in the short term in the absence of continued treatment with eculizumab, **AND**
- Patient must not receive more than 24 weeks of treatment per continuing treatment course authorised under this restriction.

Treatment criteria:

- Must be treated by a paediatric nephrologist, a nephrologist, a paediatric haematologist or a haematologist, or, must be in consultation with a paediatric nephrologist, a nephrologist, a paediatric haematologist or a haematologist.

A treatment response is defined as:

- (1) Normalisation of haematology as demonstrated by at least 2 of the following: platelet count, haptoglobin, and LDH; AND
- (2) One of the following:
 - a) An increase in eGFR of > 25% from baseline, where the baseline is the eGFR measurement immediately prior to commencing treatment with eculizumab or
 - b) an eGFR within +/- 25% from baseline; or
 - c) an avoidance of dialysis-dependence but worsening of kidney function with a reduction in eGFR 25% from baseline.

PBS-subsidised treatment with eculizumab will not be permitted if a patient has experienced treatment failure. A treatment failure is defined as a patient who is:

- (1) dialysis-dependent at the time of application and has failed to demonstrate significant resolution of extra-renal complications if originally presented; or
- (2) on dialysis and has been on dialysis for 4 months of the previous 6 months while receiving PBS-subsidised eculizumab and has failed to demonstrate significant resolution of extra-renal complications if originally presented.

The authority application must include the following measures of response to the prior course of treatment, including serial haematological results (every 3 months while the patient is receiving treatment).

The authority application must be in writing and must include:

- (1) A completed authority prescription form; and
- (2) A completed aHUS eculizumab Authority Application Supporting Information Form for Continuing treatment; and
- (3) A detailed cover letter from the prescriber; and
- (4) A copy of a current Certificate of vaccination or a statement that vaccination has or will be administered and appropriate antibiotic prophylaxis has been prescribed; and
- (5) A measurement of body weight at the time of application; and
- (6) An identified genetic mutation, if applicable; and
- (7) A family history of aHUS, if applicable; and
- (8) A history of multiple episodes of aHUS before commencing eculizumab treatment, if applicable; and
- (9) A history of kidney transplant, if applicable (especially if required due to aHUS); and
- (10) An inclusion of the individual consequences of recurrent disease; and
- (11) A supporting statement with clinical evidence of severe TMA-related cardiomyopathy (including current LVEF result), neurological impairment, gastrointestinal impairment or pulmonary impairment; and

(12) Evidence that the patient has had a treatment response including haematological results of no more than 1 month old at the time of application (platelet count, haptoglobin and LDH); and an eGFR level of no more than 1 month old at the time of application; and

(13) Evidence that the patient has not experienced treatment failure, including a supporting statement with clinical evidence that the patient does not require dialysis, unless the indication for continuing eculizumab is severe extra-renal complications that have significantly improved; and

(14) If the indication for continuing eculizumab is severe extra-renal complications, then a supporting statement with clinical evidence that any initial extra-renal complications of TMA have significantly improved is required.

This assessment must be submitted no later than 4 weeks from the cessation of the prior treatment. Where a response assessment is not undertaken and submitted within these timeframes, the patient will be deemed to have failed to respond to treatment with eculizumab.

Note All applications should be accompanied by a detailed letter that outlines the objective evidence of high risk of critical organ damage if aHUS recurs. The following evidence may be submitted to establish the patient's level of risk of aHUS recurrence in the short term in the absence of continued treatment with eculizumab:

- Evidence of a mutation known to confer a high risk of aHUS recurrence;
- Past history of recurrent episodes of active and progressive TMA due to aHUS, prior to the episode that led to current use of eculizumab;
- Past family history of aHUS recurrence, especially in first-degree relatives;
- Past history of recurrent aHUS following renal transplant for end-stage renal failure due to aHUS.

Authority required

Atypical haemolytic uraemic syndrome (aHUS)

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have demonstrated treatment response to previous treatment with PBS-subsidised eculizumab for this condition, **AND**
- Patient must not have ever experienced treatment failure with eculizumab including PBS-subsidised eculizumab for this condition, **AND**
- Patient must have the following clinical conditions:(i) either significant haemolysis as measured by low/absent haptoglobin; or presence of schistocytes on the blood film; or lactate dehydrogenase (LDH) above normal;AND(ii) either platelet consumption as measured by either 25% decline from patient baseline or thrombocytopenia (platelet count $<150 \times 10^9/L$);OR(iii) TMA-related organ impairment including on recent biopsy, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Treatment criteria:

- Must be treated by a paediatric nephrologist, a nephrologist, a paediatric haematologist or a haematologist, or, must be in consultation with a paediatric nephrologist, a nephrologist, a paediatric haematologist or a haematologist.

A treatment response is defined as:

(1) Normalisation of haematology as demonstrated by at least 2 of the following: platelet count, haptoglobin, and LDH; AND

(2) One of the following:

a) An increase in eGFR of $> 25\%$ from baseline, where the baseline is the eGFR measurement immediately prior to commencing treatment with eculizumab or

b) an eGFR within $\pm 25\%$ from baseline; or

c) an avoidance of dialysis-dependence but worsening of kidney function with a reduction in eGFR 25% from baseline.

PBS-subsidised treatment with eculizumab will not be permitted if a patient has experienced treatment failure. A treatment failure is defined as a patient who is:

(1) dialysis-dependent at the time of application and has failed to demonstrate significant resolution of extra-renal complications if originally presented; or

(2) on dialysis and has been on dialysis for 4 months of the previous 6 months while receiving PBS-subsidised eculizumab and has failed to demonstrate significant resolution of extra-renal complications if originally presented.

The authority application must include the following measures of response to the prior course of treatment, including serial haematological results (every 3 months while the patient is receiving treatment).

The authority application must be in writing and must include:

(1) A completed authority prescription form(s); and

(2) A completed aHUS eculizumab Authority Application Supporting Information Form for Recommencement of treatment; and

(3) A signed patient acknowledgement or an acknowledgement signed by a parent or authorised guardian, if applicable; and

(4) A detailed cover letter from the prescriber; and

(5) A copy of a current Certificate of vaccination or a statement that vaccination has or will be administered and appropriate antibiotic prophylaxis has been prescribed; and

(6) A measurement of body weight at the time of application, and

(7) An identified genetic mutation, if applicable; and

(8) A family history of aHUS if applicable; and

(9) A history of multiple episodes of aHUS following the treatment break, if applicable; and

(10) A history of kidney transplant if applicable (especially if required due to aHUS); and

(11) An inclusion of the individual consequences of recurrent disease; and

(12) A supporting statement with clinical evidence of TMA-related organ damage including current (within one week of application) haematological results (platelet count, haptoglobin and LDH), eGFR level, and, if applicable, on recent biopsy;

(13) Evidence that the patient has had a treatment response to their previous treatment with eculizumab; and

(14) Evidence that the patient has not experienced treatment failure, including a supporting statement with clinical evidence that the patient does not require dialysis, unless the indication for continuing eculizumab is severe extra-renal complications that have significantly improved; and

(15) If the indication for continuing eculizumab is severe extra-renal complications, then a supporting statement with clinical evidence that any initial extra-renal complications of TMA have significantly improved is required.

This assessment must be submitted no later than 4 weeks from the cessation of the prior treatment. Where a response assessment is not undertaken and submitted within these timeframes, the patient will be deemed to have failed to respond to treatment with eculizumab.

Note A raise in LDH alone is not a sufficient reason to re-commence eculizumab, but thrombocytopenia with one marker of haemolysis (such as raised LDH, presence of schistocytes, or low/absence of haptoglobin) is an accepted reason to consider re-commencement of eculizumab treatment.

Note Kidney transplantation/dialysis is not a contraindication to recommencement of eculizumab treatment.

Authority required

Atypical haemolytic uraemic syndrome (aHUS)

Treatment Phase: Continuing recommencement of treatment

Clinical criteria:

- Patient must have received treatment under Recommencement of treatment restriction with PBS-subsidised eculizumab for this condition, **AND**
- Patient must have demonstrated ongoing treatment response to the previous 24 weeks of PBS-subsidised eculizumab for this condition, **AND**
- Patient must not have experienced treatment failure with eculizumab including PBS-subsidised eculizumab for this condition, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Treatment criteria:

- Must be treated by a paediatric nephrologist, a nephrologist, a paediatric haematologist or a haematologist, or, must be in consultation with a paediatric nephrologist, a nephrologist, a paediatric haematologist or a haematologist.

A treatment response is defined as:

- (1) Normalisation of haematology as demonstrated by at least 2 of the following: platelet count, haptoglobin, and LDH; **AND**
- (2) One of the following:

a) An increase in eGFR of > 25% from baseline, where the baseline is the eGFR measurement immediately prior to commencing treatment with eculizumab or

b) an eGFR within +/- 25% from baseline; or

c) an avoidance of dialysis-dependence but worsening of kidney function with a reduction in eGFR 25% from baseline.

PBS-subsidised treatment with eculizumab will not be permitted if a patient has experienced treatment failure. A treatment failure is defined as a patient who is:

(1) dialysis-dependent at the time of application and has failed to demonstrate significant resolution of extra-renal complications if originally presented; or

(2) on dialysis and has been on dialysis for 4 months of the previous 6 months while receiving PBS-subsidised eculizumab and has failed to demonstrate significant resolution of extra-renal complications if originally presented.

The authority application must include the following measures of response to the prior course of treatment, including serial haematological results (every 3 months while the patient is receiving treatment).

The authority application must be in writing and must include:

- (1) A completed authority prescription form; and
 - (2) A completed aHUS eculizumab Authority Application Supporting Information Form for Continuing treatment; and
 - (3) A detailed cover letter from the prescriber; and
 - (4) A copy of a current Certificate of vaccination or a statement that vaccination has or will be administered and appropriate antibiotic prophylaxis has been prescribed; and
 - (5) A measurement of body weight at the time of application; and
 - (6) An identified genetic mutation, if applicable; and
 - (7) A family history of aHUS, if applicable; and
 - (8) A history of multiple episodes of aHUS before recommencing eculizumab treatment, if applicable; and
 - (9) A history of kidney transplant if applicable (especially if required due to aHUS); and
 - (10) An inclusion of the individual consequences of recurrent disease, if applicable; and
 - (11) Evidence that the patient has had a treatment response including haematological results of no more than 1 week old at the time of application (platelet count, haptoglobin and LDH); and an eGFR level of no more than 1 week old at the time of application; and
 - (12) Evidence that the patient has not experienced treatment failure, including a supporting statement with clinical evidence that the patient does not require dialysis, unless the indication for continuing eculizumab is severe extra-renal complications that have significantly improved; and
 - (13) If the indication for continuing eculizumab is severe extra-renal complications, then a supporting statement with clinical evidence that any initial extra-renal complications of TMA have significantly improved is required.
- This assessment must be submitted no later than 4 weeks from the cessation of the prior treatment. Where a response assessment is not undertaken and submitted within these timeframes, the patient will be deemed to have failed to respond to treatment with eculizumab.

eculizumab 300 mg/30 mL injection, 30 mL vial

10194M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	5984.89	Soliris [XI]

EVEROLIMUS

Caution Careful monitoring of patients is mandatory.

Authority required

Management of renal allograft rejection

Treatment Phase: Management (initiation, stabilisation and review of therapy)

Clinical criteria:

- Patient must be receiving this drug for prophylaxis of renal allograft rejection, **AND**
- The treatment must be under the supervision and direction of a transplant unit.

Authority required

Management of cardiac allograft rejection

Treatment Phase: Management (initiation, stabilisation and review of therapy)

Clinical criteria:

- Patient must be receiving this drug for prophylaxis of cardiac allograft rejection, **AND**
- The treatment must be under the supervision and direction of a transplant unit.

everolimus 750 microgram tablet, 60

Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
4	5	..	*2348.51	Certican [NV]

everolimus 500 microgram tablet, 60

Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
2	5	..	*805.11	Certican [NV]

everolimus 1 mg tablet, 60

Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
4	5	..	*3115.55	Certican [NV]

everolimus 250 microgram tablet, 60

Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
2	5	..	*406.25	Certican [NV]

MYCOPHENOLATE

Caution Careful monitoring of patients is mandatory.

Authority required

Management of renal allograft rejection

Treatment Phase: Management (initiation, stabilisation and review of therapy)

Clinical criteria:

- Patient must be receiving this drug for prophylaxis of renal allograft rejection, **AND**
- The treatment must be under the supervision and direction of a transplant unit.

Authority required

Management of cardiac allograft rejection

Treatment Phase: Management (initiation, stabilisation and review of therapy)

Clinical criteria:

- Patient must be receiving this drug for prophylaxis of cardiac allograft rejection, **AND**
- The treatment must be under the supervision and direction of a transplant unit.

mycophenolate mofetil 1 g/5 mL powder for oral liquid, 165 mL

Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
2	5	..	*#518.80	CellCept [RO]

mycophenolate mofetil 500 mg tablet, 50

Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
6	5	..	*281.71	^a APO-Mycophenolate [TX]	^a CellCept [RO]
				^a Ceptolate [AF]	^a MycoCept [RF]
				^a Mycophenolate AN [EA]	^a Mycophenolate Sandoz [SZ]
				^a Pharmacor Mycophenolate 500 [CR]	

MYCOPHENOLATE

Caution Careful monitoring of patients is mandatory.

Note Management includes initiation, stabilisation and review of therapy as required.

Authority required

Prophylaxis of renal allograft rejection

Treatment Phase: Management

Clinical criteria:

- The treatment must be under the supervision and direction of a transplant unit.

Authority required

WHO Class III, IV or V lupus nephritis

Treatment Phase: Management

Clinical criteria:

- The condition must be proven by biopsy.

Treatment criteria:

- Must be treated by a nephrologist or in consultation with a nephrologist.

The name of the consulting nephrologist must be included in the patient medical records.

mycophenolate 360 mg enteric tablet, 120

6370G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*364.23	Myfortic [NV]

mycophenolate 180 mg enteric tablet, 120

6369F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*185.83	Myfortic [NV]

■ **MYCOPHENOLATE**

Caution Careful monitoring of patients is mandatory.

Note For item codes 6208R and 1837Q, pharmaceutical benefits that have the form capsule 250 mg are equivalent for the purposes of substitution.

Authority required

Management of renal allograft rejection

Treatment Phase: Management (initiation, stabilisation and review of therapy)

Clinical criteria:

- Patient must be receiving this drug for prophylaxis of renal allograft rejection, **AND**
- The treatment must be under the supervision and direction of a transplant unit.

Authority required

Management of cardiac allograft rejection

Treatment Phase: Management (initiation, stabilisation and review of therapy)

Clinical criteria:

- Patient must be receiving this drug for prophylaxis of cardiac allograft rejection, **AND**
- The treatment must be under the supervision and direction of a transplant unit.

mycophenolate mofetil 250 mg capsule, 100

6208R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	6	5	..	*281.83	^a APO-Mycophenolate [TX] ^a Mycophenolate Sandoz [SZ]	^a CellCept [RO] ^a Pharmacor Mycophenolate 250 [CR]

mycophenolate mofetil 250 mg capsule, 50

1837Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	12	5	..	*281.83	^a Ceptolate [AF]

■ **NATALIZUMAB**

Caution Progressive multifocal leukoencephalopathy has been reported with this drug.

Authority required (STREAMLINED)

9406

Clinically definite relapsing-remitting multiple sclerosis

Treatment criteria:

- Must be treated by a neurologist.

Clinical criteria:

- The treatment must be a sole PBS-subsidised disease modifying therapy for this condition, **AND**
- Patient must be ambulatory (without assistance or support), **AND**
- Patient must have experienced at least 2 documented attacks of neurological dysfunction, believed to be due to multiple sclerosis, in the preceding 2 years of commencing a PBS-subsidised disease modifying therapy for this condition, **AND**
- The condition must be confirmed by magnetic resonance imaging of the brain and/or spinal cord; OR
- Patient must be deemed unsuitable for magnetic resonance imaging due to the risk of physical (not psychological) injury to the patient.

Population criteria:

- Patient must be aged 18 years or older.

The date of the magnetic resonance imaging scan must be included in the patient's medical notes, unless written certification is provided, in the patient's medical notes, by a radiologist that an MRI scan is contraindicated because of the risk of physical (not psychological) injury to the patient.

Treatment with this drug must cease if there is continuing progression of disability whilst the patient is being treated with this drug.

For continued treatment the patient must demonstrate compliance with, and an ability to tolerate, this drug.

Neurologists prescribing natalizumab under the PBS listing must be registered with the Tysabri Australian Prescribing Program.

natalizumab 300 mg/15 mL injection, 15 mL vial

9624M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	1388.07	Tysabri [BD]

OCRELIZUMAB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)
9523

Multiple sclerosis

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be diagnosed as clinically definite relapsing-remitting multiple sclerosis by magnetic resonance imaging of the brain and/or spinal cord; OR
- The condition must be diagnosed as clinically definite relapsing-remitting multiple sclerosis by accompanying written certification provided by a radiologist that a magnetic resonance imaging scan is contraindicated because of the risk of physical (not psychological) injury to the patient, **AND**
- The treatment must be a sole PBS-subsidised disease modifying therapy for this condition, **AND**
- Patient must have experienced at least 2 documented attacks of neurological dysfunction, believed to be due to multiple sclerosis, in the preceding 2 years of commencing a PBS-subsidised disease modifying therapy for this condition, **AND**
- Patient must be ambulatory (without assistance or support).

Treatment criteria:

- Must be treated by a neurologist.

Where applicable, the date of the magnetic resonance imaging scan must be recorded in the patient's medical records.

Authority required (STREAMLINED)
9635

Multiple sclerosis

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not show continuing progression of disability while on treatment with this drug, **AND**
- The treatment must be a sole PBS-subsidised disease modifying therapy for this condition, **AND**
- Patient must have demonstrated compliance with, and an ability to tolerate this therapy.

Treatment criteria:

- Must be treated by a neurologist.

ocrelizumab 300 mg/10 mL injection, 10 mL vial

11237K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	..	..	*17580.39	Ocrevus [RO]

SIROLIMUS

Caution Careful monitoring of patients is mandatory.

Authority required

Management of renal allograft rejection

Treatment Phase: Management (initiation, stabilisation and review of therapy)

Clinical criteria:

- Patient must be receiving this drug for prophylaxis of renal allograft rejection, **AND**
- The treatment must be under the supervision and direction of a transplant unit.

sirolimus 2 mg tablet, 100

6457W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*2397.51	Rapamune [PF]

sirolimus 1 mg tablet, 100

6436R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*1222.43	Rapamune [PF]

sirolimus 1 mg/mL oral liquid, 60 mL

6437T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*968.37	Rapamune [PF]

sirolimus 500 microgram tablet, 100

9748C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*618.43	Rapamune [PF]

■ VEDOLIZUMAB

Note TREATMENT OF ADULT PATIENTS WITH MODERATE TO SEVERE ULCERATIVE COLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, golimumab, infliximab and vedolizumab for adult patients with ulcerative colitis. Patients are eligible for PBS-subsidised treatment with either adalimumab, golimumab, infliximab or vedolizumab at any one time.

Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, golimumab, infliximab and vedolizumab only.

From 1 June 2018, under the PBS, all adult patients will be able to commence a treatment cycle where they may trial PBS-subsidised adalimumab, golimumab, infliximab or vedolizumab without having to experience a disease flare when swapping to one of the alternate agents. Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with adalimumab, golimumab, infliximab or vedolizumab while they continue to show a response to therapy.

A patient who received PBS-subsidised adalimumab, infliximab, vedolizumab treatment prior to 1 June 2018 is considered to start their first cycle as of 1 June 2018. Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised adalimumab, golimumab, infliximab or vedolizumab more than once. Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised therapy before they are eligible to commence the next cycle.

The 5-year break is measured from the date of the last approval for PBS-subsidised adalimumab, golimumab, infliximab or vedolizumab treatment in the most recent cycle to the date of the first application for initial treatment with adalimumab, golimumab, infliximab or vedolizumab under the new treatment cycle.

A patient who has failed fewer than 3 trials of either adalimumab, golimumab, infliximab or vedolizumab in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

(1) How to prescribe PBS-subsidised treatment with adalimumab, golimumab, infliximab and vedolizumab after 1 June 2018.

(a) Initial treatment. Applications for initial treatment should be made where:

(i) an adult patient has received no prior PBS-subsidised treatment with adalimumab, golimumab, infliximab or vedolizumab in this treatment cycle and wishes to commence such therapy (Initial 1 - New patient or recommencement of treatment after more than 5 years break in therapy); or

(ii) an adult patient has received prior PBS-subsidised (initial or continuing) adalimumab, golimumab, infliximab or vedolizumab therapy and wishes to trial an alternate agent (Initial 2 - Change or Re-commencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) an adult patient wishes to re-commence treatment with adalimumab, golimumab, infliximab or vedolizumab following a break in PBS-subsidised therapy with the same agent (Initial 2 - Change or Re-commencement of treatment after a break in therapy of less than 5 years).

Treatment authorisations under Initial 1 and Initial 2 will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, 14 weeks of therapy for golimumab, infliximab and vedolizumab.

A patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of treatment for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for golimumab, infliximab and vedolizumab. For second and subsequent courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment.

(b) Continuing treatment.

Following the completion of an initial treatment course with adalimumab, golimumab, infliximab or vedolizumab a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed in the month prior to completing their current course of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised treatment is approved, a patient may swap if eligible to the alternate adalimumab, golimumab, infliximab or vedolizumab treatment within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Mayo clinic score or partial Mayo clinic score), or the prior corticosteroid therapy and immunosuppressive therapy. A patient may trial an alternate treatment at any time, regardless of whether they are receiving therapy (initial or continuing) with adalimumab, golimumab, infliximab or vedolizumab at the time of the application. However, they cannot swap to a particular therapy if they have failed to respond to prior treatment with that drug once within the same treatment cycle. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(3) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised adalimumab, golimumab, infliximab or vedolizumab therapy of at least 5 years, must requalify for initial 1 treatment with respect to the scores of disease severity. A patient must have received treatment with a 5-aminosalicylate oral preparation in a standard dose for induction of remission for a minimum of 3 consecutive months, and, either azathioprine or 6-mercaptopurine for a minimum of 3 consecutive months or a tapered course of oral steroids over a 6 week period followed by an appropriately dosed thiopurine agent for a minimum of 3 consecutive months (unless intolerance develops necessitating permanent treatment withdrawal to these agents). These above prior treatments must have been received immediately prior to the time the scores of disease severity being used to trial a second or subsequent course are measured.

(4) Patients 'grandfathered' onto PBS-subsidised treatment with golimumab.

A patient who commenced treatment with golimumab for moderate to severe ulcerative colitis prior to 1 June 2018 and who continues to receive treatment at the time of application, may qualify for treatment under the initial 3 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

Note Special Pricing Arrangements apply.

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Initial treatment (new patient or Recommencement of treatment after more than 5 years break in therapy - Initial 1)

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have failed to achieve an adequate response to a 5-aminosalicylate oral preparation in a standard dose for induction of remission for 3 or more consecutive months or have intolerance necessitating permanent treatment withdrawal, **AND**
- Patient must have failed to achieve an adequate response to azathioprine at a dose of at least 2 mg per kg daily for 3 or more consecutive months or have intolerance necessitating permanent treatment withdrawal; OR
- Patient must have failed to achieve an adequate response to 6-mercaptopurine at a dose of at least 1 mg per kg daily for 3 or more consecutive months or have intolerance necessitating permanent treatment withdrawal; OR
- Patient must have failed to achieve an adequate response to a tapered course of oral steroids, starting at a dose of at least 40 mg prednisolone (or equivalent), over a 6 week period or have intolerance necessitating permanent treatment withdrawal, and followed by a failure to achieve an adequate response to 3 or more consecutive months of treatment of an appropriately dosed thiopurine agent, **AND**
- Patient must have a Mayo clinic score greater than or equal to 6; OR
- Patient must have a partial Mayo clinic score greater than or equal to 6, provided the rectal bleeding and stool frequency subscores are both greater than or equal to 2 (endoscopy subscore is not required for a partial Mayo clinic score), **AND**
- Patient must be appropriately assessed for the risk of developing progressive multifocal leukoencephalopathy whilst on this treatment.

Population criteria:

- Patient must be aged 18 years or older.

Applications for authorisation of initial treatment must be in writing and must include:

(a) a completed authority prescription form; and

(b) a completed Ulcerative Colitis PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed current Mayo clinic or partial Mayo clinic calculation sheet including the date of assessment of the patient's condition; and

(ii) details of prior systemic drug therapy [dosage, date of commencement and duration of therapy]; and

(iii) the signed patient acknowledgement.

A maximum quantity and number of repeats to provide for an initial course of this drug consisting of one vial of 300 mg per dose, with one dose to be administered at weeks 0, 2 and 6, will be authorised.

All tests and assessments should be performed preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior conventional treatment.

The most recent Mayo clinic or partial Mayo clinic score must be no more than 1 month old at the time of application.

Patients who fail to achieve a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1 or have failed to maintain a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1 with continuing treatment with this drug, will not be eligible to receive further PBS-subsidised treatment with this drug.

A partial Mayo clinic assessment of the patient's response to this initial course of treatment must be made up to 12 weeks after the first dose for patients administered doses at weeks 0, 2 and 6 (6 weeks following the third dose) so that there is adequate time for a response to be demonstrated.

Patients must have signed a patient acknowledgement indicating they understand and acknowledge that the PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment.

If treatment with any of the above-mentioned drugs is contraindicated according to the relevant TGA-approved Product Information, details must be provided at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, details of this toxicity must be provided at the time of application.

Note Details of accepted toxicities including severity can be found on the Department of Human Services website at www.humanservices.gov.au.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR

- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have previously been issued with an authority prescription for this drug for this condition, **AND**
- Patient must have demonstrated or sustained an adequate response to treatment by having a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1 while receiving treatment with this drug, **AND**
- Patient must be appropriately assessed for the risk of developing progressive multifocal leukoencephalopathy whilst on this treatment.

Patients who have failed to maintain a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1 with continuing treatment with this drug, will not be eligible to receive further PBS-subsidised treatment with this drug.

Patients are eligible to receive continuing treatment with this drug in courses of up to 24 weeks providing they continue to sustain the response.

At the time of the authority application, medical practitioners should request the appropriate number of vials, to provide for a single infusion of 300 mg per dose.

Up to a maximum of 2 repeats will be authorised.

Note No applications for increased repeats will be authorised.

Note Authority approval for sufficient therapy to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Change or Re-commencement of treatment after a break in therapy of less than 5 years (Initial 2)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with adalimumab, golimumab, infliximab or vedolizumab for this condition in this treatment cycle, **AND**
- Patient must not have failed PBS-subsidised therapy with vedolizumab for this condition in the current treatment cycle, **AND**
- Patient must be appropriately assessed for the risk of developing progressive multifocal leukoencephalopathy whilst on this treatment.

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Population criteria:

- Patient must be aged 18 years or older.

To demonstrate a response to treatment the application must be accompanied by the results of the most recent course of this drug within the timelines specified in the relevant restriction. If the response assessment to the previous course of this drug is not submitted as detailed in the relevant restriction, the patient will be deemed to have failed therapy with this drug. Applications for authorisation of change or recommencement treatment must be in writing and must include:

(a) a completed authority prescription form; and

(b) a completed Ulcerative Colitis PBS Authority Application - Supporting Information Form which includes the following:

(i) Mayo clinical assessment (to demonstrate response to prior treatment).

A maximum quantity and number of repeats to provide for an initial course of this drug consisting of one vial of 300 mg per dose, with one dose to be administered at weeks 0, 2 and 6, will be authorised.

At the time of the authority application, medical practitioners should request the appropriate number of vials, to provide for a single infusion of 300 mg per dose.

Up to a maximum of 2 repeats will be authorised.

Authority approval for sufficient therapy to complete a maximum of 3 initial doses of treatment may be requested by telephone by contacting the Department of Human Services.

Note No applications for increased repeats will be authorised.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Balance of supply

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the Initial 1 (new patient) restriction to complete the 3 doses (i.e. the initial infusion regimen at 0, 2 and 6 weeks); OR

- Patient must have received insufficient therapy with this drug under the Initial 2 (Change or Recommencement of treatment after a break in therapy) restriction to complete the 3 doses (i.e. the initial infusion regimen at 0, 2 and 6 weeks); OR
- Patient must have received insufficient therapy with this drug under the Continuing treatment restriction to complete 24 weeks of treatment, **AND**
- The treatment must provide no more than the balance of up to 3 doses (Initial 1 and Initial 2 restrictions) or 2 repeats (Continuing restriction), **AND**
- Patient must be appropriately assessed for the risk of developing progressive multifocal leukoencephalopathy whilst on this treatment.

Population criteria:

- Patient must be aged 18 years or older.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Initial PBS-subsidised treatment (Grandfather patient)

Clinical criteria:

- Patient must have previously received non-PBS-subsidised therapy with this drug for this condition prior to 1 August 2015, **AND**
- Patient must have had a Mayo clinic score greater than or equal to 6 prior to commencing treatment with this drug; OR
- Patient must have had a partial Mayo clinic score greater than or equal to 6, provided the rectal bleeding and stool frequency subscores were both greater than or equal to 2 (endoscopy subscore is not required for a partial Mayo score) prior to commencing treatment with this drug; OR
- Patient must have a documented history of moderate to severe refractory ulcerative colitis prior to having commenced treatment with this drug where a Mayo clinic, partial Mayo clinic baseline assessment is not available, **AND**
- Patient must have demonstrated or sustained an adequate response to treatment by having a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1 while receiving treatment with this drug, **AND**
- Patient must be appropriately assessed for the risk of developing progressive multifocal leukoencephalopathy whilst on this treatment.

Population criteria:

- Patient must be 18 years of age or older.

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Applications for authorisation of initial treatment must be in writing and must include:

- a completed authority prescription form; and
- a completed Ulcerative Colitis PBS Authority Application - Supporting Information Form which includes the following:
 - the completed current and baseline Mayo clinic or partial Mayo clinic calculation sheet including the date of assessment of the patient's condition; and
 - the date of commencement of this drug; and
 - the signed patient acknowledgement.

The current Mayo clinic or partial Mayo clinic assessment must be no more than 1 month old at the time of application. The baseline assessment must be from immediately prior to commencing treatment with this drug. Where a baseline assessment is not available the prescriber must contact the Department of Human Services to discuss.

Patients are eligible to receive continuing treatment with this drug in courses of up to 24 weeks providing they continue to sustain the response.

At the time of the authority application, medical practitioners should request the appropriate number of vials, to provide for a single infusion of 300 mg per dose.

Up to a maximum of 2 repeats will be authorised.

A patient may qualify for PBS-subsidised treatment under this restriction once only.

For continuing PBS-subsidised treatment, a Grandfathered patient must qualify under the Continuing treatment criteria.

Note The patient must have signed a patient acknowledgement indicating they understand and acknowledge that the PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment.

Note No applications for increased repeats will be authorised.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

vedolizumab 300 mg injection, 1 vial

10398G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	3152.58	Entyvio [TK]

VEDOLIZUMAB
Note TREATMENT OF ADULT PATIENTS WITH SEVERE CROHN DISEASE

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for adult patients with severe Crohn disease. Where the term biological medicine appears in the following NOTES and restrictions, it refers to the tumour necrosis factor (TNF) alpha-antagonists (adalimumab and infliximab), the alpha-4 beta-7 integrin inhibitor (vedolizumab) and the human IgG1kappa monoclonal antibody (ustekinumab).

Patients are eligible for PBS-subsidised treatment with only 1 of the above PBS-subsidised biological medicines at any one time.

From 1 September 2017, under the PBS, all patients will be able to commence a treatment cycle where they may trial PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab without having to experience a disease flare when swapping to the alternate agent. Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with adalimumab, infliximab, vedolizumab or ustekinumab while they continue to show a response to therapy.

A patient who received PBS-subsidised adalimumab, infliximab, or vedolizumab treatment prior to 1 September 2017 is considered to have started their treatment cycle as of 1 September 2017.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab more than once.

Once a patient has either failed or ceased to respond to treatment for this condition 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy for this condition before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab treatment in the most recent cycle to the date of the first application for initial treatment with adalimumab, infliximab, vedolizumab or ustekinumab under the new treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of less than 5 years, may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab therapy after 1 September 2017.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised therapy with adalimumab, infliximab, vedolizumab or ustekinumab in this treatment cycle and wishes to commence such therapy (Initial treatment (new patient or Recommencement of treatment after more than 5 years break in therapy - Initial 1)); or

(ii) a patient has received prior PBS-subsidised (initial or continuing) adalimumab, infliximab, vedolizumab or ustekinumab and wishes to trial an alternate agent (Initial 2 - Change or recommencement) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to re-commence treatment with adalimumab, infliximab, vedolizumab or ustekinumab following a break in PBS-subsidised therapy with that agent (Change or Re-commencement of treatment after a break in therapy of less than 5 years (Initial 2)).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, 14 weeks of therapy for infliximab, 14 weeks of therapy for vedolizumab and 16 weeks for ustekinumab.

From 1 September 2017, a patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy for adalimumab or ustekinumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab or vedolizumab, and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

For subsequent courses of PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

Adalimumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats and the second prescription should be written for 2 doses of 40 mg and 2 repeats for patients weighing 40 kg or greater. For patients weighing less than 40 kg, one prescription should be written for 2 doses of 40 mg with no repeats and the second prescription should be written for 2 doses of 20 mg with 3 repeats.

Ustekinumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be written under S100 (Highly Specialised Drugs) for a weight-based loading dose, containing a quantity of up to 4 vials of 130 mg and no repeats. The second prescription should be written under S85 (General) for the subsequent first dose, containing a quantity of 2 vials of 45 mg and no repeats.

(b) Continuing treatment.

Following the completion of an initial treatment course with adalimumab, infliximab, vedolizumab or ustekinumab, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted supply of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine therapy is approved, a patient may swap if eligible to the alternate adalimumab, infliximab, vedolizumab or ustekinumab within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Crohn Disease Activity Index (CDAI) Score, confirmation of Crohn disease), or the prior conventional therapies of corticosteroid therapy and immunosuppressive therapy.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with adalimumab, infliximab, vedolizumab or ustekinumab at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug once within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment, within the timeframes specified in the relevant restriction.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the CDAI or evidence of intestinal inflammation submitted with the first authority application for adalimumab, infliximab, vedolizumab or ustekinumab. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to assess response to all subsequent treatments.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to recommence treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity. Patients must have received treatment with a corticosteroid and at least 1 immunosuppressive agent, at an adequate dose, for a minimum of 3 consecutive months immediately prior to the time the CDAI score or the indices of intestinal inflammation are measured.

(5) Patients 'grandfathered' onto PBS-subsidised treatment with vedolizumab.

A patient who commenced treatment with vedolizumab for severe Crohn disease prior to 1 August 2015 and who continues to receive treatment at the time of application may qualify for treatment under the initial 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment will be assessed under the continuing treatment restriction of the relevant drug.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must requalify for continuing treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' above for further details.

(6) Patients 'grandfathered' onto PBS-subsidised treatment with ustekinumab.

A patient who commenced treatment with ustekinumab for severe Crohn disease prior to 1 September 2017 and who continues to receive treatment at the time of application may qualify for treatment under the initial 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment will be assessed under the continuing treatment restriction of the relevant drug.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must requalify for continuing treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' above for further details.

Note No applications for increased maximum quantities will be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Special Pricing Arrangements apply.

Authority required

Severe Crohn disease

Treatment Phase: Initial treatment (new patient - initial 1)

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have confirmed severe Crohn disease, defined by standard clinical, endoscopic and/or imaging features, including histological evidence, with the diagnosis confirmed by a gastroenterologist or a consultant physician, **AND**
- Patient must have failed to achieve an adequate response to prior systemic therapy with a tapered course of steroids, starting at a dose of at least 40 mg prednisolone (or equivalent), over a 6 week period, **AND**
- Patient must have failed to achieve adequate response to prior systemic immunosuppressive therapy with azathioprine at a dose of at least 2 mg per kg daily for 3 or more months; OR

- Patient must have failed to achieve adequate response to prior systemic immunosuppressive therapy with 6-mercaptopurine at a dose of at least 1 mg per kg daily for 3 or more months; OR
- Patient must have failed to achieve adequate response to prior systemic immunosuppressive therapy with methotrexate at a dose of at least 15 mg weekly for 3 or more months, **AND**
- Patient must be appropriately assessed for the risk of developing progressive multifocal leukoencephalopathy whilst on this treatment.

Population criteria:

- Patient must be aged 18 years or older.

Clinical criteria:

- Patient must have a Crohn Disease Activity Index (CDAI) Score greater than or equal to 300 as evidence of failure to achieve an adequate response to prior systemic therapy; OR
- Patient must have short gut syndrome with diagnostic imaging or surgical evidence, or have had an ileostomy or colostomy; and must have evidence of intestinal inflammation; and must have evidence of failure to achieve an adequate response to prior systemic therapy as specified below; OR
- Patient must have extensive intestinal inflammation affecting more than 50 cm of the small intestine as evidenced by radiological imaging; and must have a Crohn Disease Activity Index (CDAI) Score greater than or equal to 220; and must have evidence of failure to achieve an adequate response to prior systemic therapy as specified below.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current Crohn Disease Activity Index (CDAI) calculation sheet including the date of assessment of the patient's condition if relevant; and
 - (ii) details of prior systemic drug therapy [dosage, date of commencement and duration of therapy]; and
 - (iii) the reports and dates of the pathology or diagnostic imaging test(s) nominated as the response criterion, if relevant; and
 - (iv) the date of the most recent clinical assessment; and
 - (v) the signed patient acknowledgement indicating they understand and acknowledge that the PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment.

Evidence of failure to achieve an adequate response to prior therapy must include at least one of the following: (a) patient must have evidence of intestinal inflammation; (b) patient must be assessed clinically as being in a high faecal output state; (c) patient must be assessed clinically as requiring surgery or total parenteral nutrition (TPN) as the next therapeutic option, in the absence of this drug, if affected by short gut syndrome, extensive small intestine disease or is an ostomy patient. Evidence of intestinal inflammation includes: (i) blood: higher than normal platelet count, or, an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour, or, a C-reactive protein (CRP) level greater than 15 mg per L; or (ii) faeces: higher than normal lactoferrin or calprotectin level; or (iii) diagnostic imaging: demonstration of increased uptake of intravenous contrast with thickening of the bowel wall or mesenteric lymphadenopathy or fat streaking in the mesentery; All assessments, pathology tests and diagnostic imaging studies must be made within 1 month of the date of application and should be performed preferably whilst still on conventional treatment, but no longer than 1 month following cessation of the most recent prior treatment

If treatment with any of the specified prior conventional drugs is contraindicated according to the relevant TGA-approved Product Information, please provide details at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, details of this toxicity must be provided at the time of application.

Details of the accepted toxicities including severity can be found on the Department of Human Services website.

Any one of the baseline criteria may be used to determine response to an initial course of treatment and eligibility for continued therapy, according to the criteria included in the continuing treatment restriction. However, the same criterion must be used for any subsequent determination of response to treatment, for the purpose of eligibility for continuing PBS-subsidised therapy.

A maximum quantity and number of repeats to provide for an initial course of this drug consisting of one vial of 300 mg per dose, with one dose to be administered at weeks 0, 2 and 6, will be authorised.

If fewer than the maximum stated repeats in the relevant treatment phase are requested at the time of the application, authority approvals for sufficient repeats to complete the balance of the stated repeats in the relevant treatment phase may be requested by telephone by contacting the Department of Human Services and applying through the Balance of Supply restriction. Under no circumstances will telephone approvals be granted for treatment that would otherwise extend the relevant treatment phase.

The assessment of the patient's response to this initial course of treatment must be made up to 12 weeks after the first dose (6 weeks following the third dose) so that there is adequate time for a response to be demonstrated.

Note This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to the Department of Human Services no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with this drug.

Note It is recommended that an application for continuing treatment is submitted to the Department of Human Services at the time of the 12 week assessment, to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised treatment with this drug.

Authority required

Severe Crohn disease

Treatment Phase: Change or Re-commencement of treatment (initial 2)

Clinical criteria:

- Patient must have a documented history of severe Crohn disease, **AND**

- Patient must have received prior PBS-subsidised treatment with a biological disease modifying drug for this condition in this treatment cycle, **AND**
- Patient must not have failed PBS-subsidised therapy with this drug for this condition in the current treatment cycle, **AND**
- Patient must be appropriately assessed for the risk of developing progressive multifocal leukoencephalopathy whilst on this treatment.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form, which includes the following:
 - (i) the completed Crohn Disease Activity Index (CDAI) Score calculation sheet including the date of the assessment of the patient's condition, if relevant; or
 - (ii) the reports and dates of the pathology or diagnostic imaging test(s) used to assess response to therapy for patients with short gut syndrome, extensive small intestine disease or an ostomy, if relevant; and
 - (iii) the date of clinical assessment; and
 - (iv) the details of prior biological disease modifying drug treatment including the details of date and duration of treatment.

To demonstrate a response to treatment the application must be accompanied by the results of the most recent course of biological disease modifying drug (bDMD) therapy within the timeframes specified in the relevant restriction.

Where the most recent course of PBS-subsidised bDMD treatment was approved under an initial treatment restriction, the patient must have been assessed for response to that course following a minimum of 12 weeks of therapy for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab and vedolizumab and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

If the response assessment to the previous course of bDMD treatment is not submitted as detailed above, the patient will be deemed to have failed therapy with that particular course of bDMD.

A maximum quantity and number of repeats to provide for an initial course of this drug consisting of one vial of 300 mg per dose, with one dose to be administered at weeks 0, 2 and 6, will be authorised.

If fewer than the maximum stated repeats in the relevant treatment phase are requested at the time of the application, authority approvals for sufficient repeats to complete the balance of the stated repeats in the relevant treatment phase may be requested by telephone by contacting the Department of Human Services and applying through the Balance of Supply restriction. Under no circumstances will telephone approvals be granted for treatment that would otherwise extend the relevant treatment phase.

The assessment of the patient's response to this initial course of treatment must be made up to 12 weeks after the first dose (6 weeks following the third dose) so that there is adequate time for a response to be demonstrated.

This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to the Department of Human Services no later than 1 month from the date of completion of this initial course of treatment.

Where a response assessment is not undertaken and submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with this drug.

Note It is recommended that an application for continuing treatment is submitted to the Department of Human Services at the time of the 12 week assessment, to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised treatment with this drug.

Authority required

Severe Crohn disease

Treatment Phase: Initial PBS-subsidised treatment (Grandfather)

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have a documented history of severe Crohn disease, **AND**
- Patient must have previously received non-PBS-subsidised therapy with this drug for this condition prior to 1 August 2015.

Population criteria:

- Patient must be aged 18 years or older.

Clinical criteria:

- Patient must have had a Crohn Disease Activity Index (CDAI) Score of greater than or equal to 300 prior to commencing treatment with this drug; OR
- Patient must have a documented history of intestinal inflammation and have diagnostic imaging or surgical evidence of short gut syndrome if affected by the syndrome or has an ileostomy or colostomy; OR
- Patient must have a documented history and radiological evidence of intestinal inflammation if the patient has extensive small intestinal disease affecting more than 50 cm of the small intestine, **AND**
- Patient must have an adequate response to this drug defined as a reduction in Crohn Disease Activity Index (CDAI) Score to a level no greater than 150 if assessed by CDAI or if affected by extensive small intestine disease; OR
- Patient must have an adequate response to this drug defined as (a) an improvement of intestinal inflammation as demonstrated by: (i) blood: normalisation of the platelet count, or an erythrocyte sedimentation rate (ESR) level no greater than 25 mm per hour, or a C-reactive protein (CRP) level no greater than 15 mg per L; or (ii) faeces:

normalisation of lactoferrin or calprotectin level; or (iii) evidence of mucosal healing, as demonstrated by diagnostic imaging findings, compared to the baseline assessment; or (b) reversal of high faecal output state; or (c) avoidance of the need for surgery or total parenteral nutrition (TPN), if affected by short gut syndrome, extensive small intestine or is an ostomy patient, **AND**

- Patient must be appropriately assessed for the risk of developing progressive multifocal leukoencephalopathy whilst on this treatment.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current Crohn Disease Activity Index (CDAI) calculation sheet including the date of assessment of the patient's condition if relevant; and
 - (ii) details of prior systemic drug therapy [dosage, date of commencement and duration of therapy]; and
 - (iii) the reports and dates of the pathology or diagnostic imaging test(s) nominated as the response criterion, if relevant; and
 - (iv) the date of the most recent clinical assessment; and
 - (v) the signed patient acknowledgement indicating they understand and acknowledge that the PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment.

The assessment of the patient's response to a continuing course of therapy must be made within the 4 weeks prior to completion of that course and posted to the Department of Human Services no less than 2 weeks prior to the date the next dose is scheduled, in order to ensure continuity of treatment for those patients who meet the continuation criterion.

Where an assessment is not submitted to the Department of Human Services within these timeframes, patients will be deemed to have failed to respond, or to have failed to sustain a response, to treatment with this drug.

Patients are eligible to receive continuing treatment with this drug in courses of up to 24 weeks providing they continue to sustain the response.

At the time of the authority application, medical practitioners should request the appropriate number of vials, to provide sufficient for a single infusion of 300 mg vedolizumab per dose. Up to a maximum of 2 repeats will be authorised.

If fewer than the maximum stated repeats in the relevant treatment phase are requested at the time of the application, authority approvals for sufficient repeats to complete the balance of the stated repeats in the relevant treatment phase may be requested by telephone by contacting the Department of Human Services and applying through the Balance of Supply restriction. Under no circumstances will telephone approvals be granted for treatment that would otherwise extend the relevant treatment phase.

A patient may qualify for PBS-subsidised treatment under this restriction once only.

Authority required

Severe Crohn disease

Treatment Phase: Balance of supply

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the Initial 1 (new patient) restriction to complete the 3 doses (i.e. the initial infusion regimen at 0, 2 and 6 weeks); OR
- Patient must have received insufficient therapy with this drug under the Continuing treatment restriction to complete 24 weeks of treatment; OR
- Patient must have received insufficient therapy with this drug to complete 24 weeks of treatment under the Initial PBS-subsidised treatment restriction for patients who had previously received non-PBS subsidised treatment (Grandfathered patient), **AND**
- The treatment must provide no more than the balance of up to 3 doses (new patients) or 2 repeats (Continuing or Grandfathered patients), **AND**
- Patient must be appropriately assessed for the risk of developing progressive multifocal leukoencephalopathy whilst on this treatment.

Population criteria:

- Patient must be aged 18 years or older.

Note Authority approval for sufficient therapy to complete a maximum of 3 initial doses or 2 repeats may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Severe Crohn disease

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have a documented history of severe Crohn disease, **AND**
- Patient must have previously been issued with an authority prescription for this drug for this condition, **AND**
- Patient must have demonstrated or sustained an adequate response to treatment with this drug.

Population criteria:

- Patient must be aged 18 years or older.

Clinical criteria:

- Patient must be appropriately assessed for the risk of developing progressive multifocal leukoencephalopathy whilst on this treatment, **AND**
- Patient must have an adequate response to this drug defined as a reduction in Crohn Disease Activity Index (CDAI) Score to a level no greater than 150 if assessed by CDAI or if affected by extensive small intestine disease; OR
- Patient must have an adequate response to this drug defined as (a) an improvement of intestinal inflammation as demonstrated by: (i) blood: normalisation of the platelet count, or an erythrocyte sedimentation rate (ESR) level no greater than 25 mm per hour, or a C-reactive protein (CRP) level no greater than 15 mg per L; or (ii) faeces: normalisation of lactoferrin or calprotectin level; or (iii) evidence of mucosal healing, as demonstrated by diagnostic imaging findings, compared to the baseline assessment; or (b) reversal of high faecal output state; or (c) avoidance of the need for surgery or total parenteral nutrition (TPN), if affected by short gut syndrome, extensive small intestine or is an ostomy patient.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed Crohn Disease Activity Index (CDAI) Score calculation sheet including the date of the assessment of the patient's condition, if relevant; or
 - (ii) the reports and dates of the pathology test or diagnostic imaging test(s) used to assess response to therapy for patients with short gut syndrome, extensive small intestine disease or an ostomy, if relevant; and
 - (iii) the date of clinical assessment.

All assessments, pathology tests and diagnostic imaging studies, must be made within 1 month of the date of application.

If the application is the first application for continuing treatment with this drug, an assessment of the patient's response to the initial course of treatment must be made up to 12 weeks after the first dose so that there is adequate time for a response to be demonstrated.

The assessment of the patient's response to a continuing course of therapy must be made within the 4 weeks prior to completion of that course and posted to the Department of Human Services no less than 2 weeks prior to the date the next dose is scheduled, in order to ensure continuity of treatment for those patients who meet the continuation criterion.

Where an assessment is not submitted to the Department of Human Services within these timeframes, patients will be deemed to have failed to respond, or to have failed to sustain a response, to treatment with this drug.

Patients are eligible to receive continuing treatment with this drug in courses of up to 24 weeks providing they continue to sustain the response.

At the time of the authority application, medical practitioners should request the appropriate number of vials, to provide sufficient for a single infusion of 300 mg vedolizumab per dose. Up to a maximum of 2 repeats will be authorised.

If fewer than the maximum stated repeats in the relevant treatment phase are requested at the time of the application, authority approvals for sufficient repeats to complete the balance of the stated repeats in the relevant treatment phase may be requested by telephone by contacting the Department of Human Services and applying through the Balance of Supply restriction. Under no circumstances will telephone approvals be granted for treatment that would otherwise extend the relevant treatment phase.

vedolizumab 300 mg injection, 1 vial

10415E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	3152.58	Entyvio [TK]

Tumor necrosis factor alpha (TNF-) inhibitors
■ ADALIMUMAB
Note TREATMENT OF PATIENTS WITH SEVERE ACTIVE JUVENILE IDIOPATHIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, etanercept and tocilizumab for a patient who has severe active juvenile idiopathic arthritis. Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, etanercept and tocilizumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 3 biological medicines at any one time.

From 1 April 2014, a patient receiving PBS-subsidised biological medicine is considered to be in a treatment cycle where they may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements, within a single treatment cycle, a patient may:

- (i) continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy; and

- (ii) fail to respond or to sustain a response to each PBS-subsidised biological medicine once only. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single treatment cycle and they must have, at a minimum, a 12 month break in PBS-subsidised biological medicine therapy before they are eligible to commence another cycle.

The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine was approved to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who received PBS-subsidised biological medicine immediately prior to 1 April 2014 is considered to be in their first cycle as of 1 April 2014. A patient who has had a break in biological medicine treatment of at least 12 months immediately prior to making a new application, on or after 1 April 2014, will commence a new treatment cycle under the Initial 3 treatment

restriction.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of less than 12 months may commence a further course of treatment within the same treatment cycle under the Initial 2 treatment restriction.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of more than 12 months must commence a new treatment cycle under the Initial 3 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment therapy after 1 April 2014.

(1) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy

(Initial 1 - New patient); or

(ii) a patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an alternate medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine therapy of less than 12 months) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 12 months with the same medicine (initial 2 - Change or Recommencement of treatment after a break in biological medicine therapy of less than 12 months).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 12 months (initial 3 - Recommencement of treatment after a break in biological medicine of more than 12 months) Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy.

A patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not conducted within these timeframes, the patient will be deemed to have failed this course of treatment.

For the second and subsequent cycles of PBS-subsidised biological medicine, it is recommended that a patient is reviewed in the 4 weeks prior to completing their current course of treatment and that an application is posted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services where applicable. Where a response assessment is not submitted, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (joint count) or the prior non-biological medicine therapy requirements, except if the patient has had a break in therapy of more than 12 months who would then need to requalify under the initial 3 restrictions with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application.

However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, an application for a patient who wishes to swap to an alternate biological medicine should be accompanied by the approved authority prescription or remaining repeats for the biological medicine the patient is ceasing.

(4) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count submitted with the first authority application for a biological medicine. However, prescribers may provide a new baseline measurement any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to the revised baseline measurement.

(5) Recommencement of treatment after a 12 months break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological medicine therapy of at least 12 months, must qualify under the initial 3 restriction and meet the relevant criteria and index of disease severity.

(6) Withdrawal of treatment after sustained remission.

Withdrawal of treatment with biological medicine should be considered in a patient who has achieved and sustained complete remission of disease for 12 months. A demonstration of response to the current treatment should be submitted to the Department of Human Services at the time treatment is ceased.

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Treatment criteria:

- Must be treated by a paediatric rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Clinical criteria:

- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have demonstrated severe intolerance of, or toxicity due to, methotrexate; OR
- Patient must have demonstrated failure to achieve an adequate response to 1 or more of the following treatment regimens: (i) oral or parenteral methotrexate at a dose of at least 20 mg per square metre weekly, alone or in combination with oral or intra-articular corticosteroids, for a minimum of 3 months; or (ii) oral methotrexate at a dose of at least 10 mg per square metre weekly together with at least 1 other disease modifying anti-rheumatic drug (DMARD), alone or in combination with corticosteroids, for a minimum of 3 months, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be under 18 years of age.

Severe intolerance to methotrexate is defined as intractable nausea and vomiting and general malaise unresponsive to manoeuvres, including reducing or omitting concomitant non-steroidal anti-inflammatory drugs (NSAIDs) on the day of methotrexate administration, use of folic acid supplementation, or administering the dose of methotrexate in 2 divided doses over 24 hours.

Toxicity due to methotrexate is defined as evidence of hepatotoxicity with repeated elevations of transaminases, bone marrow suppression temporally related to methotrexate use, pneumonitis, or serious sepsis.

If treatment with methotrexate alone or in combination with another DMARD is contraindicated according to the relevant TGA-approved Product Information, details must be provided at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, details of this toxicity must be provided at the time of application.

The following criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

- (a) an active joint count of at least 20 active (swollen and tender) joints; OR
- (b) at least 4 active joints from the following list:
 - (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - (ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The joint count assessment must be performed preferably whilst still on DMARD treatment, but no longer than 4 weeks following cessation of the most recent prior treatment.

The authority application must be made in writing and must include:

- (1) completed authority prescription form(s); and
- (2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

At the time of authority application, medical practitioners must request the appropriate number of injections of appropriate strength, based on the weight of the patient, to provide sufficient for two doses. Up to a maximum of 3 repeats will be authorised.

An assessment of a patient's response to an initial course of treatment must be conducted following a minimum of 12 weeks of therapy. An application for the continuing treatment must be accompanied with the assessment of response and submitted to the Department of Human Services no later than 4 weeks from the date of completion of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Use of alternative DMARDs in children is dependent on approval by the Therapeutic Goods Administration as age restrictions may apply.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 12 months)

Treatment criteria:

- Must be treated by a paediatric rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be under 18 years of age.

An adequate response to treatment is defined as:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

At the time of authority application, medical practitioners must request the appropriate number of injections of appropriate strength, based on the weight of the patient, to provide sufficient for two doses. Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

(1) completed authority prescription form(s); and

(2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to change or recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient who fails to demonstrate a response to treatment with this drug under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug in this treatment cycle. A patient may re-trial this drug after a minimum of 12 months have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the initial 3 treatment restriction.

If a patient fails to respond to PBS-subsidised biological medicine treatment 3 times (once with each agent) they will not be eligible to receive further PBS-subsidised biological medicine therapy in this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 3 (recommencement of treatment after a break in biological medicine of more than 12 months)

Treatment criteria:

- Must be treated by a paediatric rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have had a break in treatment of 12 months or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must have either (a) a total active joint count of at least 20 active (swollen and tender) joints; or (b) at least 4 active major joints, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be under 18 years of age.

Active joints are defined as:

- (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
(ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

All measures of joint count must be no more than 4 weeks old at the time of this application.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of active joints, the response must be demonstrated on the total number of active joints.

At the time of authority application, medical practitioners must request the appropriate number of injections of appropriate strength, based on the weight of the patient, to provide sufficient for two doses. Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

- (1) completed authority prescription form(s); and
- (2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient) or Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 12 months) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 12 months) - balance of supply

Treatment criteria:

- Must be treated by a paediatric rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 12 months) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 12 months) restriction to complete 16 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**

- Patient must not receive more than 24 weeks of treatment per continuing treatment course authorised under this restriction.

An adequate response to treatment is defined as:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

Determination of whether a response has been demonstrated to initial and subsequent courses of treatment will be based on the baseline measurement of joint count submitted with the initial treatment application.

The authority application must be made in writing and must include:

(1) completed authority prescription form(s); and

(2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

At the time of authority application, medical practitioners must request the appropriate number of injections of appropriate strength, based on the weight of the patient, to provide sufficient for two doses. Up to a maximum of 5 repeats will be authorised.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either Initial 1, Initial 2, or Initial 3 treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 12 months have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

If a patient fails to respond to PBS-subsidised biological medicine treatment 3 times (once with each agent) they will not be eligible to receive further PBS-subsidised biological medicine therapy in this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Continuing treatment - balance of supply

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restriction.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL pen devices

9680L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	1199.26	Humira [VE]

adalimumab 20 mg/0.4 mL injection, 2 x 0.4 mL syringes

9678J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	1199.26	Humira [VE]

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL syringes

9679K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	1199.26	Humira [VE]

■ ETANERCEPT

Note TREATMENT OF PATIENTS WITH SEVERE ACTIVE JUVENILE IDIOPATHIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, etanercept and tocilizumab for a patient who has severe active juvenile idiopathic arthritis. Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, etanercept and tocilizumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 3 biological medicines at any one time.

From 1 April 2014, a patient receiving PBS-subsidised biological medicine is considered to be in a treatment cycle where they may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements, within a single treatment cycle, a patient may:

(i) continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy; and

(ii) fail to respond or to sustain a response to each PBS-subsidised biological medicine once only. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single treatment cycle and they must have, at a minimum, a 12 month break in PBS-subsidised biological medicine therapy before they are eligible to commence another cycle.

The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine was approved to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who received PBS-subsidised biological medicine immediately prior to 1 April 2014 is considered to be in their first cycle as of 1 April 2014. A patient who has had a break in biological medicine treatment of at least 12 months immediately prior to making a new application, on or after 1 April 2014, will commence a new treatment cycle under the Initial 3 treatment restriction.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of less than 12 months may commence a further course of treatment within the same treatment cycle under the Initial 2 treatment restriction.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of more than 12 months must commence a new treatment cycle under the Initial 3 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment therapy after 1 April 2014.

(1) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an alternate medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine therapy of less than 12 months) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 12 months with the same medicine (initial 2 - Change or Recommencement of treatment after a break in biological medicine therapy of less than 12 months).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 12 months (initial 3 - Recommencement of treatment after a break in biological medicine of more than 12 months) Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy.

A patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not conducted within these timeframes, the patient will be deemed to have failed this course of treatment.

For the second and subsequent cycles of PBS-subsidised biological medicine, it is recommended that a patient is reviewed in the 4 weeks prior to completing their current course of treatment and that an application is posted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services where applicable. Where a response assessment is not submitted, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (joint count) or the prior non-biological medicine therapy requirements, except if the patient has had a break in therapy of more than 12 months who would then need to requalify under the initial 3 restrictions with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application.

However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, an application for a patient who wishes to swap to an alternate biological medicine should be accompanied by the approved authority prescription or remaining repeats for the biological medicine the patient is ceasing.

(4) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count submitted with the first authority application for a biological medicine. However, prescribers may provide a new baseline measurement any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to the revised baseline measurement.

(5) Recommencement of treatment after a 12 months break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological medicine therapy of at least 12 months, must qualify under the initial 3 restriction and meet the relevant criteria and index of disease severity.

(6) Withdrawal of treatment after sustained remission.

Withdrawal of treatment with biological medicine should be considered in a patient who has achieved and sustained complete remission of disease for 12 months. A demonstration of response to the current treatment should be submitted to the Department of Human Services at the time treatment is ceased.

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Treatment criteria:

- Must be treated by a paediatric rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Clinical criteria:

- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have demonstrated severe intolerance of, or toxicity due to, methotrexate; OR
- Patient must have demonstrated failure to achieve an adequate response to 1 or more of the following treatment regimens: (i) oral or parenteral methotrexate at a dose of at least 20 mg per square metre weekly, alone or in combination with oral or intra-articular corticosteroids, for a minimum of 3 months; or (ii) oral methotrexate at a dose of at least 10 mg per square metre weekly together with at least 1 other disease modifying anti-rheumatic drug (DMARD), alone or in combination with corticosteroids, for a minimum of 3 months, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be under 18 years of age.

Severe intolerance to methotrexate is defined as intractable nausea and vomiting and general malaise unresponsive to manoeuvres, including reducing or omitting concomitant non-steroidal anti-inflammatory drugs (NSAIDs) on the day of methotrexate administration, use of folic acid supplementation, or administering the dose of methotrexate in 2 divided doses over 24 hours.

Toxicity due to methotrexate is defined as evidence of hepatotoxicity with repeated elevations of transaminases, bone marrow suppression temporally related to methotrexate use, pneumonitis, or serious sepsis.

If treatment with methotrexate alone or in combination with another DMARD is contraindicated according to the relevant TGA-approved Product Information, details must be provided at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, details of this toxicity must be provided at the time of application.

The following criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

- (a) an active joint count of at least 20 active (swollen and tender) joints; OR
- (b) at least 4 active joints from the following list:
 - (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - (ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The joint count assessment must be performed preferably whilst still on DMARD treatment, but no longer than 4 weeks following cessation of the most recent prior treatment.

The authority application must be made in writing and must include:

- (1) completed authority prescription form(s); and
- (2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

At the time of authority application, medical practitioners must request the appropriate number of injections to provide sufficient for four weeks of treatment. Up to a maximum of 3 repeats will be authorised.

An assessment of a patient's response to an initial course of treatment must be conducted following a minimum of 12 weeks of therapy. An application for the continuing treatment must be accompanied with the assessment of response and submitted to the Department of Human Services no later than 4 weeks from the date of completion of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Use of alternative DMARDs in children is dependent on approval by the Therapeutic Goods Administration as age restrictions may apply.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 12 months)

Treatment criteria:

- Must be treated by a paediatric rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be under 18 years of age.

An adequate response to treatment is defined as:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

At the time of authority application, medical practitioners must request the appropriate number of injections to provide sufficient for four weeks of treatment. Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

(1) completed authority prescription form(s); and

(2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to change or recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient who fails to demonstrate a response to treatment with this drug under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug in this treatment cycle. A patient may re-trial this drug after a minimum of 12 months have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the initial 3 treatment restriction.

If a patient fails to respond to PBS-subsidised biological medicine treatment 3 times (once with each agent) they will not be eligible to receive further PBS-subsidised biological medicine therapy in this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 3 (recommencement of treatment after a break in biological medicine of more than 12 months)

Treatment criteria:

- Must be treated by a paediatric rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have had a break in treatment of 12 months or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must have either (a) a total active joint count of at least 20 active (swollen and tender) joints; or (b) at least 4 active major joints, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be under 18 years of age.

Active joints are defined as:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

All measures of joint count must be no more than 4 weeks old at the time of this application.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of active joints, the response must be demonstrated on the total number of active joints.

At the time of authority application, medical practitioners must request the appropriate number of injections to provide sufficient for four weeks of treatment. Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

- (1) completed authority prescription form(s); and
- (2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient) or Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 12 months) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 12 months) - balance of supply

Treatment criteria:

- Must be treated by a paediatric rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 16 weeks treatment; OR

- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 12 months) restriction to complete 16 weeks treatment; OR
 - Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 12 months) restriction to complete 16 weeks treatment, **AND**
 - The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.
- Note** Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment per continuing treatment course authorised under this restriction.

An adequate response to treatment is defined as:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

Determination of whether a response has been demonstrated to initial and subsequent courses of treatment will be based on the baseline measurement of joint count submitted with the initial treatment application.

The authority application must be made in writing and must include:

(1) completed authority prescription form(s); and

(2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

At the time of authority application, medical practitioners must request the appropriate number of injections to provide sufficient for four weeks of treatment. Up to a maximum of 5 repeats will be authorised.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either Initial 1, Initial 2, or Initial 3 treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 12 months have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

If a patient fails to respond to PBS-subsidised biological medicine treatment 3 times (once with each agent) they will not be eligible to receive further PBS-subsidised biological medicine therapy in this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Continuing treatment - balance of supply

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restriction.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

etanercept 50 mg/mL injection, 4 x 1 mL syringes

9615C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	984.21	Enbrel [PF]

etanercept 50 mg/mL injection, 4 x 1 mL pen devices

9641K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	984.21	Enbrel [PF]

etanercept 25 mg injection [4 vials] (&) inert substance diluent [4 x 1 mL syringes], 1 pack

6367D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	495.81	Enbrel [PF]

■ INFLIXIMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CROHN DISEASE

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for adult patients with severe Crohn disease. Where the term biological medicine appears in the following NOTES and restrictions, it refers to the tumour necrosis factor (TNF) alpha-antagonists (adalimumab and infliximab), the alpha-4 beta-7 integrin inhibitor (vedolizumab) and the human IgG1kappa monoclonal antibody (ustekinumab).

Patients are eligible for PBS-subsidised treatment with only 1 of the above PBS-subsidised biological medicines at any one time.

From 1 September 2017, under the PBS, all patients will be able to commence a treatment cycle where they may trial PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab without having to experience a disease flare when swapping to the alternate agent. Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with adalimumab, infliximab, vedolizumab or ustekinumab while they continue to show a response to therapy.

A patient who received PBS-subsidised adalimumab, infliximab, or vedolizumab treatment prior to 1 September 2017 is considered to have started their treatment cycle as of 1 September 2017.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab more than once.

Once a patient has either failed or ceased to respond to treatment for this condition 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy for this condition before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab treatment in the most recent cycle to the date of the first application for initial treatment with adalimumab, infliximab, vedolizumab or ustekinumab under the new treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of less than 5 years, may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab therapy after 1 September 2017.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised therapy with adalimumab, infliximab, vedolizumab or ustekinumab in this treatment cycle and wishes to commence such therapy (Initial treatment (new patient or Recommencement of treatment after more than 5 years break in therapy - Initial 1)); or

(ii) a patient has received prior PBS-subsidised (initial or continuing) adalimumab, infliximab, vedolizumab or ustekinumab and wishes to trial an alternate agent (Initial 2 - Change or recommencement) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to re-commence treatment with adalimumab, infliximab, vedolizumab or ustekinumab following a break in PBS-subsidised therapy with that agent (Change or Re-commencement of treatment after a break in therapy of less than 5 years (Initial 2)).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, 14 weeks of therapy for infliximab, 14 weeks of therapy for vedolizumab and 16 weeks for ustekinumab.

From 1 September 2017, a patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy for adalimumab or ustekinumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab or vedolizumab, and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

For subsequent courses of PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

Adalimumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats and the second

prescription should be written for 2 doses of 40 mg and 2 repeats for patients weighing 40 kg or greater. For patients weighing less than 40 kg, one prescription should be written for 2 doses of 40 mg with no repeats and the second prescription should be written for 2 doses of 20 mg with 3 repeats.

Ustekinumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be written under S100 (Highly Specialised Drugs) for a weight-based loading dose, containing a quantity of up to 4 vials of 130 mg and no repeats. The second prescription should be written under S85 (General) for the subsequent first dose, containing a quantity of 2 vials of 45 mg and no repeats.

(b) Continuing treatment.

Following the completion of an initial treatment course with adalimumab, infliximab, vedolizumab or ustekinumab, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted supply of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine therapy is approved, a patient may swap if eligible to the alternate adalimumab, infliximab, vedolizumab or ustekinumab within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Crohn Disease Activity Index (CDAI) Score, confirmation of Crohn disease), or the prior conventional therapies of corticosteroid therapy and immunosuppressive therapy.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with adalimumab, infliximab, vedolizumab or ustekinumab at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug once within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment, within the timeframes specified in the relevant restriction.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the CDAI or evidence of intestinal inflammation submitted with the first authority application for adalimumab, infliximab, vedolizumab or ustekinumab. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to assess response to all subsequent treatments.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to recommence treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity. Patients must have received treatment with a corticosteroid and at least 1 immunosuppressive agent, at an adequate dose, for a minimum of 3 consecutive months immediately prior to the time the CDAI score or the indices of intestinal inflammation are measured.

(5) Patients 'grandfathered' onto PBS-subsidised treatment with vedolizumab.

A patient who commenced treatment with vedolizumab for severe Crohn disease prior to 1 August 2015 and who continues to receive treatment at the time of application may qualify for treatment under the initial 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment will be assessed under the continuing treatment restriction of the relevant drug.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must requalify for continuing treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' above for further details.

(6) Patients 'grandfathered' onto PBS-subsidised treatment with ustekinumab.

A patient who commenced treatment with ustekinumab for severe Crohn disease prior to 1 September 2017 and who continues to receive treatment at the time of application may qualify for treatment under the initial 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment will be assessed under the continuing treatment restriction of the relevant drug.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must requalify for continuing treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' above for further details.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe Crohn disease

Treatment Phase: Subsequent continuing treatment

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must have an adequate response to this drug defined as a reduction in Crohn Disease Activity Index (CDAI) Score to a level no greater than 150 if assessed by CDAI or if affected by extensive small intestine disease; OR
- Patient must have an adequate response to this drug defined as (a) an improvement of intestinal inflammation as demonstrated by: (i) blood: normalisation of the platelet count, or an erythrocyte sedimentation rate (ESR) level no greater than 25 mm per hour, or a C-reactive protein (CRP) level no greater than 15 mg per L; or (ii) faeces: normalisation of lactoferrin or calprotectin level; or (iii) evidence of mucosal healing, as demonstrated by diagnostic imaging findings, compared to the baseline assessment; or (b) reversal of high faecal output state; or (c) avoidance of the need for surgery or total parenteral nutrition (TPN), if affected by short gut syndrome, extensive small intestine or is an ostomy patient.

Population criteria:

- Patient must be aged 18 years or older.

Applications for authorisation must be made in writing and must include: (a) a completed authority prescription form; and (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form which includes the following: (i) the completed Crohn Disease Activity Index (CDAI) Score; or (ii) the reports and dates of the pathology test or diagnostic imaging test(s) used to assess response to therapy for patients with short gut syndrome, extensive small intestine disease or an ostomy, if relevant; and (iii) the date of the most recent clinical assessment.

Each application for subsequent continuing treatment with this drug must include an assessment of the patient's response to the prior course of therapy. If the response assessment is not provided at the time of application the patient will be deemed to have failed this course of treatment.

Patients are eligible to receive continuing treatment with this drug in courses of up to 24 weeks providing they continue to sustain the response.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg eight weekly. Up to a maximum of 2 repeats will be authorised.

infliximab 100 mg injection, 1 vial

11399Y	Max. Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	340.93	Remicade [JC]

■ INFLIXIMAB
Note TREATMENT OF COMPLEX REFRACTORY FISTULISING CROHN DISEASE

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for patients with complex refractory fistulising Crohn disease. Where the term "biological medicine" appears in the following NOTES and restrictions, it refers to adalimumab and infliximab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the PBS- subsidised biological medicines for this condition at any one time.

From 1 April 2011, under the PBS, all patients will be able to commence a treatment cycle where they may trial PBS-subsidised adalimumab or infliximab without having to experience a disease flare when swapping to the alternate agent. Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with adalimumab or infliximab while they continue to show a response to therapy.

A patient who received PBS-subsidised adalimumab or infliximab treatment prior to 1 April 2011 is considered to have started their treatment cycle as of 1 April 2011.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised adalimumab or infliximab more than twice.

Once a patient has either failed or ceased to respond to treatment for this condition 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy for this condition before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised adalimumab or infliximab treatment in the most recent cycle to the date of the first application for initial treatment with adalimumab or infliximab under the new treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of less than 5 years, may commence a further course of treatment within the same treatment cycle.

A patient who has failed 3 trials or fewer of biological medicine therapy in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised adalimumab or infliximab therapy after 1 April 2011.

(a) Initial treatment. Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised adalimumab or infliximab therapy in this treatment cycle and wishes to commence such therapy (Initial treatment (new patient or Recommencement of treatment after more than 5 years break in therapy - Initial 1); or

(ii) a patient has received prior PBS-subsidised (initial or continuing) adalimumab or infliximab therapy and wishes to trial an alternate agent (Initial 2 - Change or recommencement) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to re-commence treatment with adalimumab or infliximab following a break in PBS-subsidised therapy with that agent (Change or Re-commencement of treatment after a break in therapy of less than 5 years (Initial 2)

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy for adalimumab and 14 weeks of therapy for infliximab.

From 1 April 2011, a patient must be assessed for response to any course of initial PBS-subsidised treatment following a

minimum of 12 weeks of therapy for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab, and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

For subsequent courses of PBS-subsidised adalimumab or infliximab treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

Adalimumab only: Two completed authority prescriptions must be submitted with every initial application for adalimumab. One prescription must be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats. The second prescription must be written for 2 doses of 40 mg and 2 repeats.

(b) Continuing treatment.

Adalimumab patients:

Following the completion of an initial treatment course with adalimumab, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

Infliximab patients:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment under the Initial 1 or Initial 2 treatment restrictions.

For the second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted supply of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap if eligible to the alternate biological medicine within the same treatment cycle.

A patient may trial the alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with adalimumab or infliximab at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug two times within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment within the timeframes specified in the relevant restriction.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements submitted with the first authority application for adalimumab or infliximab. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to these revised baseline measurements.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to recommence treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Complex refractory Fistulising Crohn disease

Treatment Phase: Subsequent continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug.

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

An adequate response is defined as:

- (a) a decrease from baseline in the number of open draining fistulae of greater than or equal to 50%; and/or
- (b) a marked reduction in drainage of all fistula(e) from baseline, together with less pain and induration as reported by the patient.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Fistulising Crohn Disease PBS Authority Application - Supporting Information Form which includes a completed Fistula Assessment form including the date of the assessment of the patient's condition.

The most recent fistula assessment must be no more than 1 month old at the time of application.

Each application for subsequent continuing treatment with this drug must include an assessment of the patient's response to the prior course of therapy. If the response assessment is not provided at the time of application the patient will be deemed to have failed this course of treatment.

Patients are eligible to receive continuing treatment with this drug in courses of up to 24 weeks providing they continue to sustain the response.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg eight weekly. Up to a maximum of 2 repeats will be authorised.

infliximab 100 mg injection, 1 vial

11412P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	340.93	Remicade [JC]

■ INFLIXIMAB

Note TREATMENT OF PAEDIATRIC PATIENTS WITH REFRACTORY CROHN DISEASE

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) for paediatric patients with adalimumab for severe refractory Crohn disease and infliximab for moderate to severe refractory Crohn disease. Where the term "biological medicines" appears in the following NOTES and restrictions, it refers to adalimumab and infliximab only. A patient is eligible for PBS-subsidised treatment with only one PBS-subsidised biological medicine at any one time. For paediatric patients with Crohn disease, infliximab is PBS-subsidised for moderate to severe disease while adalimumab is PBS-subsidised for severe disease.

From 1 August 2015, under the PBS, patients commencing on adalimumab will be able to commence a treatment cycle where they may trial each PBS-subsidised biological medicine without having to experience a disease flare when swapping to infliximab. Patients on infliximab will be able to commence a treatment cycle where they may trial each PBS-subsidised biological medicine but will need to meet a PCDAI score of greater than or equal to 40 when swapping to adalimumab. Under these arrangements, within a single treatment cycle and depending on the disease severity, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy.

A patient who received PBS-subsidised biological medicine treatment prior to 1 August 2015 is considered to have started their treatment cycle as of 1 August 2015.

Within the same treatment cycle, a paediatric patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than twice.

Once a patient has either failed, or ceased to respond to treatment for this condition 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy for this condition before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle.

A patient who has failed 3 trials or fewer of biological medicine therapy in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 August 2015.

(a) Initial treatment.

Applications for initial treatment should be made where:

- i) a patient has received no prior PBS-subsidised biological medicine therapy in this treatment cycle and wishes to commence such therapy - Initial 1 treatment (new patient or Re-commencement of treatment after more than 5 years break in therapy); or
- (ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent - Initial 2 treatment (Change or Re-commencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or
- (iii) a patient wishes to re-commence treatment with a specific biological medicine following a break in PBS-subsidised therapy with that agent - Initial 2 treatment (Change or Re-commencement of treatment after a break in therapy of less than 5 years).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy for adalimumab and 14 weeks of therapy for infliximab.

From 1 August 2015, a patient must be assessed for response to any course of initial PBS-subsidised biological therapy following a minimum of 12 weeks of therapy for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab, and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

For first and subsequent continuing courses of PBS-subsidised biological medicine therapy, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

Adalimumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats and the second prescription should be written for 2 doses of 40 mg and 2 repeats for patients weighing 40 kg or greater. For patients weighing less than 40 kg, one prescription should be written for 2 doses of 40 mg with no repeats and the second prescription should be written for 2 doses of 20 mg with 3 repeats.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient remains eligible to receive up to 24 weeks per course of continuing treatment under the First continuing treatment and Subsequent Continuing

treatment restrictions with that drug providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted supply of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine therapy is approved, a patient with severe disease may swap if eligible to the alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Paediatric Crohn Disease Activity Index (PCDAI) Score, confirmation of Crohn disease), or the prior conventional therapies of corticosteroid therapy, immunosuppressive therapy or enteral nutrition. Patients on infliximab may swap to adalimumab within the same treatment cycle provided that their disease severity has progressed to severe disease (i.e. they have a current PCDAI score of 40 or more).

A patient cannot swap to a biological medicine if they have failed to respond to prior treatment with that drug twice within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment within the timeframes specified in the relevant restriction.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the PCDAI submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to assess response to all subsequent treatments.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to recommence treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity.

Patients must have failed to achieve an adequate response to 2 of the following 3 conventional prior therapies including: (i) a tapered course of steroids, starting at a dose of at least 1 mg per kg or 40 mg (whichever is the lesser) prednisolone (or equivalent), over a 6 week period; (ii) an 8 week course of enteral nutrition; or (iii) immunosuppressive therapy including azathioprine at a dose of at least 2 mg per kg daily for 3 or more months, or, 6-mercaptopurine at a dose of at least 1 mg per kg daily for 3 or more months, or, methotrexate at a dose of at least 10 mg per square metre weekly for 3 or more months immediately prior to the time the PCDAI score is measured.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Moderate to severe Crohn disease

Treatment Phase: Subsequent continuing treatment

Treatment criteria:

- Must be treated by a gastroenterologist (code 87) or a consultant physician [internal medicine specialising in gastroenterology (code 81)] or a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician or a specialist paediatric gastroenterologist.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First Continuing treatment restriction; AND
- Patient must have a reduction in PCDAI Score by at least 15 points from baseline value; AND
- Patient must have a total PCDAI score of 30 points or less.

Population criteria:

- Patient must be aged 6 to 17 years inclusive.

Application for authorisation must be made in writing and must include:

(a) a completed authority prescription form; and

(b) a completed Paediatric Crohn Disease PBS Authority Application - Supporting Information Form, which includes the completed Paediatric Crohn Disease Activity Index (PCDAI) calculation sheet along with the date of the assessment of the patient's condition.

Patients are only eligible to receive subsequent continuing PBS-subsidised treatment with this drug in courses of up to 24 weeks at a dose of 5 mg per kg per dose providing they continue to sustain the response.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg eight weekly. Up to a maximum of 2 repeats will be authorised.

infliximab 100 mg injection, 1 vial

11445J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	340.93	Remicade [JC]

■ INFLIXIMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for adults with severe active rheumatoid arthritis. Where the term biological medicine appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alpha antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab), the T-cell co-stimulation modulator (abatacept) and the Janus kinase (JAK) inhibitors (baricitinib, tofacitinib). A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised biological medicine therapy may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once, and
- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised biological medicines for the treatment of rheumatoid arthritis.

Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Stevens Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

For a patient who has failed PBS-subsidised treatment with 2 or 3 TNF-alfa antagonists prior to 1 August 2010 please contact the Department of Human Services on 1800 700 270. A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction.

A patient who has failed fewer than 5 biological medicines and who has a break in treatment of less than 24 months may commence a further course of treatment with a biological medicine under Initial 2 treatment restriction. A patient who has failed fewer than 5 biological medicines and who has had a break in therapy of longer than 24 months may commence a further course of treatment with a biological medicine under the Initial 3 treatment restriction. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine treatment is stopped to the date of the new application for treatment with a biological medicine.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 April 2019.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised biological medicine treatment and wishes to commence such therapy, excluding rituximab (Initial 1 - new patient); or
 - (ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months) [further details are under 'Swapping therapy' below]; or
 - (iii) a patient wishes to re-commence treatment with a specific biological medicine following a break of less than 24 months in PBS-subsidised therapy with that agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months).
 - (iv) a patient wishes to re-commence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 24 months (Initial 3 - re-commencement of treatment after a break in biological medicine of more than 24 months)
- Initial applications for a new patient (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6-month intensive DMARD trial, but prior to ceasing DMARD therapy.

Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, baricitinib, etanercept, golimumab, tocilizumab and tofacitinib, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

Rituximab patients should be assessed following a minimum of 12 weeks after the first infusion, and the assessment should be submitted to the Department of Human Services within 4 weeks to ensure continuity of treatment for those who meet the continuing restriction for PBS subsidised treatment with this drug for this condition.

Where a response assessment is not provided the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment. For second and subsequent courses of PBS-subsidised biological medicine (excluding rituximab) treatment it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to the Department of Human Services within sufficient timeframe to allow for authority assessment.

Abatacept patients:

A patient is eligible to receive one I.V. loading dose when commencing treatment with the subcutaneous formulation. Two prescriptions are required, the first prescription for the I.V. loading dose for sufficient vials for one dose based on the patient's weight with no repeats. The second prescription for the subcutaneous formulation, with a maximum quantity of 4 and up to 3 repeats, must be submitted with the initial application.

Rituximab patients:

A further application may be submitted to the Department of Human Services within sufficient timeframe to allow authority assessment. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

Infliximab and etanercept patients:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient

is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions. For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction. Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

(2) Swapping therapy

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non- biological medicine therapy requirements except if the patient has had a break in therapy of more than 24 months who would need to requalify with respect to the indices of disease severity. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each biological medicine trialled.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent, unless they have a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug.

Abatacept:

A patient swapping from I.V. abatacept to subcutaneous abatacept will not be eligible for an I.V. loading dose when commencing treatment with the subcutaneous formulation.

Rituximab:

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF-alfa antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological medicine during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological medicine therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate biological medicine may do so without having to have any treatment-free period.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and the Department of Human Services will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints.

Applications under the Initial 1 treatment restriction for a new patient must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy.

Applications under the Initial 3 treatment restriction for re-commencement of treatment after a break in biological medicine of more than 24 months must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note Authority applications for continuing treatment may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required (STREAMLINED)

9615

Severe active rheumatoid arthritis

Treatment Phase: Subsequent continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must not receive more than 24 weeks of treatment per subsequent continuing treatment course authorised under this restriction, **AND**
- The treatment must be given concomitantly with methotrexate at a dose of at least 7.5 mg weekly.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials to provide sufficient drug, based on the weight of the patient, for a single infusion at a dose of 3 mg per kg.

Up to a maximum of 2 repeats will be authorised.

Each application for subsequent continuing treatment with this drug must include an assessment of the patient's response to the prior course of therapy. If the response assessment is not provided at the time of application the patient will be deemed to have failed this course of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

If a patient has either failed or ceased to respond to a PBS-subsidised biological medicine for this condition 5 times, they will not be eligible to receive further PBS-subsidised treatment with a biological medicine for this condition.

infliximab 100 mg injection, 1 vial

11483J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	340.93	^a Inflectra [PF]	^a Renflexis [MK]

■ INFLIXIMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for adults with severe active rheumatoid arthritis. Where the term biological medicine appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alfa antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab), the T-cell co-stimulation modulator (abatacept) and the Janus kinase (JAK) inhibitors (baricitinib, tofacitinib). A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised biological medicine therapy may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once, and
- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised biological medicines for the treatment of rheumatoid arthritis.

Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Stevens Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

For a patient who has failed PBS-subsidised treatment with 2 or 3 TNF-alfa antagonists prior to 1 August 2010 please contact the Department of Human Services on 1800 700 270. A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction.

A patient who has failed fewer than 5 biological medicines and who has a break in treatment of less than 24 months may commence a further course of treatment with a biological medicine under Initial 2 treatment restriction. A patient who has failed fewer than 5 biological medicines and who has had a break in therapy of longer than 24 months may commence a further course of treatment with a biological medicine under the Initial 3 treatment restriction. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine treatment is stopped to the date of the new application for treatment with a biological medicine.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 April 2019.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised biological medicine treatment and wishes to commence such therapy, excluding rituximab (Initial 1 - new patient); or
- (ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months) [further details are under 'Swapping therapy' below]; or
- (iii) a patient wishes to re-commence treatment with a specific biological medicine following a break of less than 24 months in PBS-subsidised therapy with that agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months).

(iv) a patient wishes to re-commence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 24 months (Initial 3 - re-commencement of treatment after a break in biological medicine of more than 24 months) Initial applications for a new patient (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6-month intensive DMARD trial, but prior to ceasing DMARD therapy.

Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, baricitinib, etanercept, golimumab, tocilizumab and tofacitinib, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

Rituximab patients should be assessed following a minimum of 12 weeks after the first infusion, and the assessment should be submitted to the Department of Human Services within 4 weeks to ensure continuity of treatment for those who meet the continuing restriction for PBS subsidised treatment with this drug for this condition.

Where a response assessment is not provided the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment. For second and subsequent courses of PBS-subsidised biological medicine (excluding rituximab) treatment it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to the Department of Human Services within sufficient timeframe to allow for authority assessment.

Abatacept patients:

A patient is eligible to receive one I.V. loading dose when commencing treatment with the subcutaneous formulation. Two prescriptions are required, the first prescription for the I.V. loading dose for sufficient vials for one dose based on the patient's weight with no repeats. The second prescription for the subcutaneous formulation, with a maximum quantity of 4 and up to 3 repeats, must be submitted with the initial application.

Rituximab patients:

A further application may be submitted to the Department of Human Services within sufficient timeframe to allow authority assessment. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

Infliximab and etanercept patients:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions. For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction. Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

(2) Swapping therapy

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non- biological medicine therapy requirements except if the patient has had a break in therapy of more than 24 months who would need to requalify with respect to the indices of disease severity. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each biological medicine trialled.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent, unless they have a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug.

Abatacept:

A patient swapping from I.V. abatacept to subcutaneous abatacept will not be eligible for an I.V. loading dose when commencing treatment with the subcutaneous formulation.

Rituximab:

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF-alfa antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological medicine during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological medicine therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate biological medicine may do so without having to have any treatment-free period.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial treatment authority

application is submitted and the Department of Human Services will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints.

Applications under the Initial 1 treatment restriction for a new patient must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy.

Applications under the Initial 3 treatment restriction for re-commencement of treatment after a break in biological medicine of more than 24 months must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Subsequent continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment per subsequent continuing treatment course authorised under this restriction, **AND**
- The treatment must be given concomitantly with methotrexate at a dose of at least 7.5 mg weekly.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials to provide sufficient drug, based on the weight of the patient, for a single infusion at a dose of 3 mg per kg.

Up to a maximum of 2 repeats will be authorised.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient has either failed or ceased to respond to a PBS-subsidised biological medicine for this condition 5 times, they will not be eligible to receive further PBS-subsidised treatment with a biological medicine for this condition.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

infliximab 100 mg injection, 1 vial

11487N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	340.93	Remicade [JC]

■ INFLIXIMAB

Note Authority applications for increased quantities/ repeats (where relevant) may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)
9632

Acute severe ulcerative colitis

Treatment criteria:

- Must be treated by a gastroenterologist; OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology, or general medicine specialising in gastroenterology].

Clinical criteria:

- Patient must have received an infusion of infliximab for the treatment of this condition as a hospital inpatient no more than two weeks prior to the date of the authority application, **AND**
- Patient must be an adult aged 18 years or older, and prior to initiation of infliximab treatment in hospital must have been experiencing six or more bloody stools per day, plus at least one of the following: (i) Temperature greater than 37.8 degrees Celsius; (ii) Pulse rate greater than 90 beats per minute; (iii) Haemoglobin less than 105 g/L; (iv) Erythrocyte sedimentation rate greater than 30 mm/h; OR
- Patient must be a child aged 6 to 17 years inclusive, and prior to initiation of infliximab treatment in hospital must have had a Paediatric Ulcerative Colitis Activity Index (PUCAI) greater than or equal to 65, with the diagnosis confirmed by a gastroenterologist, or a consultant physician as specified below, **AND**
- Patient must have failed to achieve an adequate response to at least 72 hours treatment with intravenous corticosteroids prior to initiation of infliximab treatment in hospital.

Population criteria:

- Patient must be 6 years of age or older.

For adults aged 18 years or older, failure to achieve an adequate response to intravenous corticosteroid treatment is defined by the Oxford criteria where:

(i) If assessed on day 3, patients pass 8 or more stools per day or 3 or more stools per day with a C-reactive protein (CRP) greater than 45 mg/L

(ii) If assessed on day 7, patients pass 3 or more stools per day with visible blood.

For children aged 6 to 17 years, failure to achieve an adequate response to intravenous corticosteroids means a PUCAI score greater than 45 at 72 hours.

At the time of authority application, prescribers should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg.

Before administering infliximab to a child aged 6 to 17 years, the treating clinician must have consulted with a paediatric gastroenterologist or with an institution experienced in performance of paediatric colectomy. The name of the specialist or institution must be included in the patient's medical records.

Evidence that the patient meets the PBS restriction criteria must be recorded in the patient's medical records.

infliximab 100 mg injection, 1 vial

10057H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	5	1	..	*1650.94	^a Inflectra [PF]	^a Remicade [JC]
					^a Renflexis [MK]	

■ INFLIXIMAB
Note TREATMENT OF ADULT PATIENTS WITH ANKYLOSING SPONDYLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, certolizumab pegol, etanercept, golimumab, infliximab and secukinumab for adult patients with ankylosing spondylitis. Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab and secukinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 6 biological medicines at any 1 time.

Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last prescription for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised biological medicine treatment with adalimumab, certolizumab pegol, etanercept,

golimumab, infliximab and secukinumab.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised biological medicine treatment in this treatment cycle and wishes to commence such therapy (Initial 1 - New patient)
 - (ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or
 - (iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years); or
 - (iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).
- A patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy.

(b) Continuing treatment.

For the first continuing treatment course of PBS-subsidised biological medicine, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1, Initial 2 or Initial 3 treatment restrictions. For second and subsequent continuing courses of PBS-subsidised biological medicine it is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply. Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be conducted within 1 month of the last dose. Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI), or the prior NSAID therapy and exercise program requirements.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the BASDAI, ESR and/or CRP submitted with the first authority application for a biological medicine.

However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to these revised baseline measurements.

For a new patient, the BASDAI used to determine the baseline must be measured while the patient is receiving NSAID therapy and completing their exercise program.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response.

(4) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must qualify under the Initial 3 treatment restriction. The same clinical criteria and indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI) as for the Initial 1 (New patient) restriction will need to be met, but a re-trial of NSAID therapy and exercise therapy is not required.

Note Authority applications for increased quantities/ repeats (where relevant) may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required (STREAMLINED)

9621

Ankylosing spondylitis

Treatment Phase: Subsequent continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**

- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response is defined as an improvement from baseline of at least 2 of the BASDAI and 1 of the following:

- (a) an ESR measurement no greater than 25 mm per hour; or
- (b) a CRP measurement no greater than 10 mg per L; or
- (c) an ESR or CRP measurement reduced by at least 20% from baseline.

Where only 1 acute phase reactant measurement is supplied in the first application for PBS-subsidised treatment, that same marker must be used to determine response for all subsequent continuing treatments.

The measurement of response to the prior course of therapy must be documented in the patient's medical notes.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

infliximab 100 mg injection, 1 vial

11488P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	5	3	..	*1650.94	^a Inflectra [PF]	^a Renflexis [MK]

■ INFLIXIMAB
Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE PSORIATIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab for adult patients with severe active psoriatic arthritis. Therefore, where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient receiving PBS-subsidised treatment for psoriatic arthritis is able to commence a treatment cycle where they may trial biological medicines without having to experience a disease flare when swapping to the alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

A patient who received PBS-subsidised adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab or ustekinumab treatment prior to 1 May 2019 is considered to start their first cycle as of 1 May 2019.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence another cycle [further details are under '(5) Recommencement of treatment after a 5-year break in PBS-subsidised therapy' below].

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe active psoriatic arthritis.

(1) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an alternate medicine (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years) or An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, golimumab secukinumab and tofacitinib, 18 to 20 weeks of therapy for certolizumab pegol (depending upon the dosing regimen), 20 weeks of therapy for ixekizumab, 22 weeks of therapy for infliximab, and 28 weeks of therapy for ustekinumab. It is recommended that a patient be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of PBS-subsidised initial treatment following a minimum of 12 weeks of therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

Grandfather patients (ixekizumab only).

A patient who commenced treatment with ixekizumab for severe psoriatic arthritis prior to 1 March 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

Grandfather patients (tofacitinib only).

A patient who commenced treatment with Tofacitinib for severe psoriatic arthritis prior to 1 May 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment under the Initial 1 or Initial 2 treatment restrictions.

For the second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to re-qualify with respect to either the indices of disease severity (i.e. erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP) level, and active joint count) or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify under the Initial 3 treatment restriction with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application.

However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

Within a treatment cycle a patient may alternate between therapy with any biological medicine of their choice (1 at a time) providing:

(i) they have not received PBS-subsidised treatment with that particular biological medicine previously; or
(ii) they have demonstrated an adequate response to that particular biological medicine if they have previously trialled it on the PBS; and

(iii) they have not previously failed to respond to treatment 3 times in this treatment cycle with that biological medicine.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(4) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the indices of disease severity submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and these revised baseline measurements will be used to assess response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to determine response for all subsequent continuing treatments. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. 20 or more active joints), response will be determined according to a reduction in the total number of active joints.

(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment restriction according to the criteria of the relevant restriction

and index of disease severity. The application must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Subsequent continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment per subsequent continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an erythrocyte sedimentation rate (ESR) no greater than 25 mm per hour or a C-reactive protein (CRP) level no greater than 15 mg per L or either marker reduced by at least 20% from baseline; and

either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following major active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to determine response for all subsequent continuing treatments.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg eight weekly.

Up to a maximum of 2 repeats will be authorised.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Severe Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

Where the most recent course of PBS-subsidised treatment with this drug was approved under the first continuing treatment restriction, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

infliximab 100 mg injection, 1 vial

11498E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	340.93	Remicade [JC]

■ INFLIXIMAB

Note TREATMENT OF ADULT PATIENTS WITH MODERATE TO SEVERE ULCERATIVE COLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, golimumab, infliximab and vedolizumab for adult patients with ulcerative colitis. Patients are eligible for PBS-subsidised treatment with either adalimumab, golimumab, infliximab or vedolizumab at any one time.

Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, golimumab, infliximab and vedolizumab only.

From 1 June 2018, under the PBS, all adult patients will be able to commence a treatment cycle where they may trial PBS-subsidised adalimumab, golimumab, infliximab or vedolizumab without having to experience a disease flare when swapping to one of the alternate agents. Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with adalimumab, golimumab, infliximab or vedolizumab while they continue to show a response to therapy.

A patient who received PBS-subsidised adalimumab, infliximab, vedolizumab treatment prior to 1 June 2018 is considered to start their first cycle as of 1 June 2018. Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised adalimumab, golimumab, infliximab or vedolizumab more than once. Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised therapy before they are eligible to commence the next cycle.

The 5-year break is measured from the date of the last approval for PBS-subsidised adalimumab, golimumab, infliximab or vedolizumab treatment in the most recent cycle to the date of the first application for initial treatment with adalimumab, golimumab, infliximab or vedolizumab under the new treatment cycle.

A patient who has failed fewer than 3 trials of either adalimumab, golimumab, infliximab or vedolizumab in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

(1) How to prescribe PBS-subsidised treatment with adalimumab, golimumab, infliximab and vedolizumab after 1 June 2018.

(a) Initial treatment. Applications for initial treatment should be made where:

(i) an adult patient has received no prior PBS-subsidised treatment with adalimumab, golimumab, infliximab or vedolizumab in this treatment cycle and wishes to commence such therapy (Initial 1 - New patient or recommencement of treatment after more than 5 years break in therapy); or

(ii) an adult patient has received prior PBS-subsidised (initial or continuing) adalimumab, golimumab, infliximab or vedolizumab therapy and wishes to trial an alternate agent (Initial 2 - Change or Re-commencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) an adult patient wishes to re-commence treatment with adalimumab, golimumab, infliximab or vedolizumab following a break in PBS-subsidised therapy with the same agent (Initial 2 - Change or Re-commencement of treatment after a break in therapy of less than 5 years).

Treatment authorisations under Initial 1 and Initial 2 will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, 14 weeks of therapy for golimumab, infliximab and vedolizumab.

A patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of treatment for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for golimumab, infliximab and vedolizumab. For second and subsequent courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment.

(b) Continuing treatment.

Following the completion of an initial treatment course with adalimumab, golimumab, infliximab or vedolizumab a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed in the month prior to completing their current course of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised treatment is approved, a patient may swap if eligible to the alternate adalimumab, golimumab, infliximab or vedolizumab treatment within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Mayo clinic score or partial Mayo clinic score), or the prior corticosteroid therapy and immunosuppressive therapy. A patient may trial an alternate treatment at any time, regardless of whether they are receiving therapy (initial or continuing) with adalimumab, golimumab, infliximab or vedolizumab at the time of the application. However, they cannot swap to a particular therapy if they have failed to respond to prior treatment with that drug once within the same treatment cycle. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(3) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised adalimumab, golimumab, infliximab or vedolizumab therapy of at least 5 years, must requalify for initial 1 treatment with respect to the scores of disease severity. A patient must have received treatment with a 5-aminosalicylate oral preparation in a standard dose for induction of remission for a minimum of 3 consecutive months, and, either azathioprine or 6-mercaptopurine for a minimum of 3 consecutive months or a tapered course of oral steroids over a 6 week period followed by an appropriately dosed thiopurine agent for a minimum of 3 consecutive months (unless intolerance develops necessitating permanent treatment withdrawal to these agents). These above prior treatments must have been received immediately prior to the time the scores of disease severity being used to trial a second or subsequent course are measured.

(4) Patients 'grandfathered' onto PBS-subsidised treatment with golimumab.

A patient who commenced treatment with golimumab for moderate to severe ulcerative colitis prior to 1 June 2018 and who continues to receive treatment at the time of application, may qualify for treatment under the initial 3 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

Note TREATMENT OF PAEDIATRIC PATIENTS WITH MODERATE TO SEVERE ULCERATIVE COLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) for paediatric patients with infliximab or adalimumab for moderate to severe ulcerative colitis; and infliximab for acute severe ulcerative colitis.

Where the term 'biological medicine' appears in the following NOTES and restrictions, it refers to infliximab and adalimumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 2 biological medicines at any one time. Infliximab and

adalimumab are PBS-subsidised for moderate to severe disease while only infliximab is PBS-subsidised for acute severe disease.

From 1 June 2017, under the PBS, all paediatric patients will be able to commence a treatment cycle where they may trial each PBS-subsidised biological medicine without having to experience a disease flare when swapping to the alternate agent. Under these arrangements, within a single treatment cycle and depending on the disease severity, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy. A patient who received PBS-subsidised biological medicine treatment prior to 1 June 2017 is considered to have started their treatment cycle as of 1 June 2017. Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than twice. Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle. A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle. There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 June 2017.

(a) Initial treatment. Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised biological medicine treatment in this treatment cycle and wishes to commence such therapy - Initial 1 treatment (new patient or Re-commencement of treatment after more than 5 years break in therapy); or
- (ii) a patient has received prior PBS-subsidised (initial or continuing) treatment with a biological medicine and wishes to trial an alternate agent - Initial 2 treatment (Change or Re-commencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping treatment' below]; or
- (iii) a patient wishes to re-commence treatment with a specific biological medicine following a break in PBS-subsidised therapy with that agent - Initial 2 treatment (Change or Re-commencement of treatment after a break in therapy of less than 5 years).

Treatment authorisations under Initial 1 and Initial 2 treatment will be limited to provide for a maximum of 16 weeks of treatment for adalimumab and 14 weeks of treatment for infliximab. From 1 June 2017, a patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of treatment for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab. For second and subsequent courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment.

Adalimumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. For patients weighing 40 kg or greater, one prescription should be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats and the second prescription should be written for 2 doses of 40 mg and 2 repeats. For patients weighing less than 40 kg, one prescription should be written for 2 doses of 40 mg with no repeats and the second prescription should be written for 2 doses of 20 mg with 3 repeats.

(b) Continuing treatment. Following the completion of an initial treatment course with a specific biological medicine, a patient remains eligible to receive up to 24 weeks per course of continuing treatment with that drug under the continuing treatment restriction providing they continue to sustain the response. It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure PBS subsidy criteria are met.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap if eligible to the alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Paediatric Ulcerative Colitis Activity Index (PUCAI) Score, confirmation of ulcerative colitis disease), or the prior conventional therapies of corticosteroids or immunosuppressives. A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving treatment (initial or continuing) with infliximab or adalimumab at the time of the application. However, a patient cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug twice within the same treatment cycle. To ensure a patient receives the maximum treatment opportunities allowed under these swapping arrangements, it is important that they are assessed for response to every course of treatment.

(3) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to recommence treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity. A patient must have received treatment with a 5-aminosalicylate oral preparation in a standard dose for induction of remission for a minimum of 3 consecutive months, and, either azathioprine or 6-mercaptopurine for a minimum of 3 consecutive months or a tapered course of oral steroids over a 6 week period followed by an appropriately dosed thiopurine agent for a minimum of 3 consecutive months (unless intolerance develops necessitating permanent treatment withdrawal to these agents) immediately prior to the time the PUCAI score is measured.

Note Authority applications for continuing treatment may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician; OR
- Must be treated by a specialist paediatric gastroenterologist.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have demonstrated or sustained an adequate response to treatment by having a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1 while receiving treatment with this drug; OR
- Patient must have demonstrated or sustained an adequate response to treatment by having a Paediatric Ulcerative Colitis Activity Index (PUCAI) score of less than 10 while receiving treatment with this drug, if aged 6 to 17 years.

Population criteria:

- Patient must be 6 years of age or older.

Patients who have failed to maintain a partial Mayo clinic score of less than or equal to 2, with no subscore greater than 1, or, patients who have failed to maintain a Paediatric Ulcerative Colitis Activity Index (PUCAI) score of less than 10 (if aged 6 to 17 years) with continuing treatment with this drug, will not be eligible to receive further PBS-subsidised treatment with this drug.

Patients are only eligible to receive continuing PBS-subsidised treatment with this drug in courses of up to 24 weeks at a dose of 5 mg per kg per dose providing they continue to sustain the response.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg eight weekly. Up to a maximum of 2 repeats will be authorised.

infliximab 100 mg injection, 1 vial

11797X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	340.93	^a Inflectra [PF] ^a Renflexis [MK]	^a Remicade [JC]

■ INFLIXIMAB
Note TREATMENT OF ADULT PATIENTS WITH SEVERE CROHN DISEASE

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for adult patients with severe Crohn disease. Where the term biological medicine appears in the following NOTES and restrictions, it refers to the tumour necrosis factor (TNF) alpha-antagonists (adalimumab and infliximab), the alpha-4 beta-7 integrin inhibitor (vedolizumab) and the human IgG1kappa monoclonal antibody (ustekinumab).

Patients are eligible for PBS-subsidised treatment with only 1 of the above PBS-subsidised biological medicines at any one time.

From 1 September 2017, under the PBS, all patients will be able to commence a treatment cycle where they may trial PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab without having to experience a disease flare when swapping to the alternate agent. Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with adalimumab, infliximab, vedolizumab or ustekinumab while they continue to show a response to therapy.

A patient who received PBS-subsidised adalimumab, infliximab, or vedolizumab treatment prior to 1 September 2017 is considered to have started their treatment cycle as of 1 September 2017.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab more than once.

Once a patient has either failed or ceased to respond to treatment for this condition 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy for this condition before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab treatment in the most recent cycle to the date of the first application for initial treatment with adalimumab, infliximab, vedolizumab or ustekinumab under the new treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of less than 5 years, may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab therapy after 1 September 2017.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised therapy with adalimumab, infliximab, vedolizumab or ustekinumab in this treatment cycle and wishes to commence such therapy (Initial treatment (new patient or Recommencement of treatment after more than 5 years break in therapy - Initial 1)); or

(ii) a patient has received prior PBS-subsidised (initial or continuing) adalimumab, infliximab, vedolizumab or ustekinumab and wishes to trial an alternate agent (Initial 2 - Change or recommencement) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to re-commence treatment with adalimumab, infliximab, vedolizumab or ustekinumab following a break in PBS-subsidised therapy with that agent (Change or Re-commencement of treatment after a break in therapy of less than 5 years (Initial 2)).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, 14 weeks of therapy for infliximab, 14 weeks of therapy for vedolizumab and 16 weeks for ustekinumab.

From 1 September 2017, a patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy for adalimumab or ustekinumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab or vedolizumab, and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

For subsequent courses of PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that where

required an application is submitted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

Adalimumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats and the second prescription should be written for 2 doses of 40 mg and 2 repeats for patients weighing 40 kg or greater. For patients weighing less than 40 kg, one prescription should be written for 2 doses of 40 mg with no repeats and the second prescription should be written for 2 doses of 20 mg with 3 repeats.

Ustekinumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be written under S100 (Highly Specialised Drugs) for a weight-based loading dose, containing a quantity of up to 4 vials of 130 mg and no repeats. The second prescription should be written under S85 (General) for the subsequent first dose, containing a quantity of 2 vials of 45 mg and no repeats.

(b) Continuing treatment.

Following the completion of an initial treatment course with adalimumab, infliximab, vedolizumab or ustekinumab, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted supply of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine therapy is approved, a patient may swap if eligible to the alternate adalimumab, infliximab, vedolizumab or ustekinumab within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Crohn Disease Activity Index (CDAI) Score, confirmation of Crohn disease), or the prior conventional therapies of corticosteroid therapy and immunosuppressive therapy.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with adalimumab, infliximab, vedolizumab or ustekinumab at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug once within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment, within the timeframes specified in the relevant restriction.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the CDAI or evidence of intestinal inflammation submitted with the first authority application for adalimumab, infliximab, vedolizumab or ustekinumab. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to assess response to all subsequent treatments.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to recommence treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity. Patients must have received treatment with a corticosteroid and at least 1 immunosuppressive agent, at an adequate dose, for a minimum of 3 consecutive months immediately prior to the time the CDAI score or the indices of intestinal inflammation are measured.

(5) Patients 'grandfathered' onto PBS-subsidised treatment with vedolizumab.

A patient who commenced treatment with vedolizumab for severe Crohn disease prior to 1 August 2015 and who continues to receive treatment at the time of application may qualify for treatment under the initial 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment will be assessed under the continuing treatment restriction of the relevant drug.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must requalify for continuing treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' above for further details.

(6) Patients 'grandfathered' onto PBS-subsidised treatment with ustekinumab.

A patient who commenced treatment with ustekinumab for severe Crohn disease prior to 1 September 2017 and who continues to receive treatment at the time of application may qualify for treatment under the initial 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment will be assessed under the continuing treatment restriction of the relevant drug.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must requalify for continuing treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' above for further details.

Note Authority applications for increased quantities/ repeats (where relevant) may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

9552

Severe Crohn disease

Treatment Phase: Subsequent continuing treatment

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR

- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must have an adequate response to this drug defined as a reduction in Crohn Disease Activity Index (CDAI) Score to a level no greater than 150 if assessed by CDAI or if affected by extensive small intestine disease; OR
- Patient must have an adequate response to this drug defined as (a) an improvement of intestinal inflammation as demonstrated by: (i) blood: normalisation of the platelet count, or an erythrocyte sedimentation rate (ESR) level no greater than 25 mm per hour, or a C-reactive protein (CRP) level no greater than 15 mg per L; or (ii) faeces: normalisation of lactoferrin or calprotectin level; or (iii) evidence of mucosal healing, as demonstrated by diagnostic imaging findings, compared to the baseline assessment; or (b) reversal of high faecal output state; or (c) avoidance of the need for surgery or total parenteral nutrition (TPN), if affected by short gut syndrome, extensive small intestine or is an ostomy patient.

Population criteria:

- Patient must be aged 18 years or older.

The measurement of response to the prior course of therapy must be documented in the patient's medical notes.

Patients are eligible to receive subsequent continuing treatment with this drug in courses of up to 24 weeks at a dose of 5 mg per kg per dose providing they continue to sustain the response.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

infliximab 100 mg injection, 1 vial

11396T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	5	2	..	*1650.94	^a Inflectra [PF]	^a Renflexis [MK]

■ INFLIXIMAB
Note TREATMENT OF COMPLEX REFRACTORY FISTULISING CROHN DISEASE

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for patients with complex refractory fistulising Crohn disease. Where the term "biological medicine" appears in the following NOTES and restrictions, it refers to adalimumab and infliximab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the PBS- subsidised biological medicines for this condition at any one time.

From 1 April 2011, under the PBS, all patients will be able to commence a treatment cycle where they may trial PBS-subsidised adalimumab or infliximab without having to experience a disease flare when swapping to the alternate agent. Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with adalimumab or infliximab while they continue to show a response to therapy.

A patient who received PBS-subsidised adalimumab or infliximab treatment prior to 1 April 2011 is considered to have started their treatment cycle as of 1 April 2011.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised adalimumab or infliximab more than twice.

Once a patient has either failed or ceased to respond to treatment for this condition 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy for this condition before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised adalimumab or infliximab treatment in the most recent cycle to the date of the first application for initial treatment with adalimumab or infliximab under the new treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of less than 5 years, may commence a further course of treatment within the same treatment cycle.

A patient who has failed 3 trials or fewer of biological medicine therapy in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised adalimumab or infliximab therapy after 1 April 2011.

(a) Initial treatment. Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised adalimumab or infliximab therapy in this treatment cycle and wishes to commence such therapy (Initial treatment (new patient or Recommencement of treatment after more than 5 years break in therapy - Initial 1); or

(ii) a patient has received prior PBS-subsidised (initial or continuing) adalimumab or infliximab therapy and wishes to trial an alternate agent (Initial 2 - Change or recommencement) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to re-commence treatment with adalimumab or infliximab following a break in PBS-subsidised therapy with that agent (Change or Re-commencement of treatment after a break in therapy of less than 5 years (Initial 2)

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy for adalimumab and 14 weeks of therapy for infliximab.

From 1 April 2011, a patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab, and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

For subsequent courses of PBS-subsidised adalimumab or infliximab treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

Adalimumab only: Two completed authority prescriptions must be submitted with every initial application for adalimumab. One prescription must be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats. The second prescription must be written for 2 doses of 40 mg and 2 repeats.

(b) Continuing treatment.

Adalimumab patients:

Following the completion of an initial treatment course with adalimumab, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

Infliximab patients:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment under the Initial 1 or Initial 2 treatment restrictions.

For the second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted supply of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap if eligible to the alternate biological medicine within the same treatment cycle.

A patient may trial the alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with adalimumab or infliximab at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug two times within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment within the timeframes specified in the relevant restriction.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements submitted with the first authority application for adalimumab or infliximab. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to these revised baseline measurements.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to recommence treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity.

Note Authority applications for increased quantities/ repeats (where relevant) may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

9397

Complex refractory Fistulising Crohn disease

Treatment Phase: Subsequent continuing treatment

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological agent treatment for this condition in this treatment cycle, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug.

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

An adequate response is defined as:

- (a) a decrease from baseline in the number of open draining fistulae of greater than or equal to 50%; and/or
- (b) a marked reduction in drainage of all fistula(e) from baseline, together with less pain and induration as reported by the patient.

The measurement of response to the prior course of therapy must be documented in the patient's medical notes.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Patients are eligible to receive subsequent continuing treatment with this drug in courses of up to 24 weeks at a dose of 5 mg per kg per dose providing they continue to sustain the response.

infliximab 100 mg injection, 1 vial

11432Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	5	2	..	*1650.94	^a Inflectra [PF]	^a Renflexis [MK]

■ INFLIXIMAB

Note TREATMENT OF PAEDIATRIC PATIENTS WITH REFRACTORY CROHN DISEASE

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) for paediatric patients with adalimumab for severe refractory Crohn disease and infliximab for moderate to severe refractory Crohn disease. Where the term "biological medicines" appears in the following NOTES and restrictions, it refers to adalimumab and infliximab only. A patient is eligible for PBS-subsidised treatment with only one PBS-subsidised biological medicine at any one time. For paediatric patients with Crohn disease, infliximab is PBS-subsidised for moderate to severe disease while adalimumab is PBS-subsidised for severe disease.

From 1 August 2015, under the PBS, patients commencing on adalimumab will be able to commence a treatment cycle where they may trial each PBS-subsidised biological medicine without having to experience a disease flare when swapping to infliximab. Patients on infliximab will be able to commence a treatment cycle where they may trial each PBS-subsidised biological medicine but will need to meet a PCDAI score of greater than or equal to 40 when swapping to adalimumab. Under these arrangements, within a single treatment cycle and depending on the disease severity, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy.

A patient who received PBS-subsidised biological medicine treatment prior to 1 August 2015 is considered to have started their treatment cycle as of 1 August 2015.

Within the same treatment cycle, a paediatric patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than twice.

Once a patient has either failed, or ceased to respond to treatment for this condition 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy for this condition before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle.

A patient who has failed 3 trials or fewer of biological medicine therapy in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 August 2015.

(a) Initial treatment.

Applications for initial treatment should be made where:

i) a patient has received no prior PBS-subsidised biological medicine therapy in this treatment cycle and wishes to commence such therapy - Initial 1 treatment (new patient or Re-commencement of treatment after more than 5 years break in therapy); or

(ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent - Initial 2 treatment (Change or Re-commencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to re-commence treatment with a specific biological medicine following a break in PBS-subsidised therapy with that agent - Initial 2 treatment (Change or Re-commencement of treatment after a break in therapy of less than 5 years).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy for adalimumab and 14 weeks of therapy for infliximab.

From 1 August 2015, a patient must be assessed for response to any course of initial PBS-subsidised biological therapy following a minimum of 12 weeks of therapy for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab, and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

For first and subsequent continuing courses of PBS-subsidised biological medicine therapy, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

Adalimumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats and the second prescription should be written for 2 doses of 40 mg and 2 repeats for patients weighing 40 kg or greater. For patients weighing less than 40 kg, one prescription should be written for 2 doses of 40 mg with no repeats and the second prescription should be written for 2 doses of 20 mg with 3 repeats.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient remains eligible to receive up to 24 weeks per course of continuing treatment under the First continuing treatment and Subsequent Continuing treatment restrictions with that drug providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted supply of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine therapy is approved, a patient with severe disease may swap if eligible to the alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Paediatric Crohn Disease Activity Index (PCDAI) Score, confirmation of Crohn disease), or the prior conventional therapies of corticosteroid therapy, immunosuppressive therapy or enteral nutrition.

Patients on infliximab may swap to adalimumab within the same treatment cycle provided that their disease severity has progressed to severe disease (i.e. they have a current PCDAI score of 40 or more).

A patient cannot swap to a biological medicine if they have failed to respond to prior treatment with that drug twice within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment within the timeframes specified in the relevant restriction.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the PCDAI submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to assess response to all subsequent

treatments.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to recommence treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity.

Patients must have failed to achieve an adequate response to 2 of the following 3 conventional prior therapies including: (i) a tapered course of steroids, starting at a dose of at least 1 mg per kg or 40 mg (whichever is the lesser) prednisolone (or equivalent), over a 6 week period; (ii) an 8 week course of enteral nutrition; or (iii) immunosuppressive therapy including azathioprine at a dose of at least 2 mg per kg daily for 3 or more months, or, 6-mercaptopurine at a dose of at least 1 mg per kg daily for 3 or more months, or, methotrexate at a dose of at least 10 mg per square metre weekly for 3 or more months immediately prior to the time the PCDAI score is measured.

Note Authority applications for increased quantities/ repeats (where relevant) may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

9557

Moderate to severe Crohn disease

Treatment Phase: Subsequent continuing treatment

Treatment criteria:

- Must be treated by a gastroenterologist (code 87) or a consultant physician [internal medicine specialising in gastroenterology (code 81)] or a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician or a specialist paediatric gastroenterologist.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First Continuing treatment restriction; AND
- Patient must have a reduction in PCDAI Score by at least 15 points from baseline value; AND
- Patient must have a total PCDAI score of 30 points or less.

Population criteria:

- Patient must be aged 6 to 17 years inclusive.

The PCDAI assessment must be no more than 1 month old at the time of prescribing.

The PCDAI score must be documented in the patient's medical notes as the measurement of response to the prior course of therapy.

Patients are only eligible to receive subsequent continuing PBS-subsidised treatment with this drug in courses of up to 24 weeks at a dose of 5 mg per kg per dose providing they continue to sustain the response.

infliximab 100 mg injection, 1 vial

11450P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	5	2	..	*1650.94	^a Inflectra [PF]	^a Renflexis [MK]

■ INFLIXIMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE PSORIATIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab for adult patients with severe active psoriatic arthritis. Therefore, where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient receiving PBS-subsidised treatment for psoriatic arthritis is able to commence a treatment cycle where they may trial biological medicines without having to experience a disease flare when swapping to the alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

A patient who received PBS-subsidised adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab or ustekinumab treatment prior to 1 May 2019 is considered to start their first cycle as of 1 May 2019.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence another cycle [further details are under '(5) Recommencement of treatment after a 5-year break in PBS-subsidised therapy' below].

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe active psoriatic arthritis.

(1) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy (Initial 1 - New patient); or
- (ii) a patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an alternate medicine (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or
- (iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).
- (iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years) or An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, golimumab secukinumab and tofacitinib, 18 to 20 weeks of therapy for certolizumab pegol (depending upon the dosing regimen), 20 weeks of therapy for ixekizumab, 22 weeks of therapy for infliximab, and 28 weeks of therapy for ustekinumab. It is recommended that a patient be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of PBS-subsidised initial treatment following a minimum of 12 weeks of therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

Grandfather patients (ixekizumab only).

A patient who commenced treatment with ixekizumab for severe psoriatic arthritis prior to 1 March 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

Grandfather patients (tofacitinib only).

A patient who commenced treatment with Tofacitinib for severe psoriatic arthritis prior to 1 May 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment under the Initial 1 or Initial 2 treatment restrictions.

For the second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to re-qualify with respect to either the indices of disease severity (i.e. erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP) level, and active joint count) or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify under the Initial 3 treatment restriction with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application.

However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

Within a treatment cycle a patient may alternate between therapy with any biological medicine of their choice (1 at a time) providing:

- (i) they have not received PBS-subsidised treatment with that particular biological medicine previously; or
- (ii) they have demonstrated an adequate response to that particular biological medicine if they have previously trialled it on the PBS; and
- (iii) they have not previously failed to respond to treatment 3 times in this treatment cycle with that biological medicine.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(4) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the indices of disease severity submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and these revised baseline measurements will be used to assess response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to determine response for all subsequent continuing treatments. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. 20 or more active joints), response will be determined according to a reduction in the total number of active joints.

(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment restriction according to the criteria of the relevant restriction and index of disease severity. The application must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note Authority applications for increased quantities/ repeats (where relevant) may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

9472

Severe psoriatic arthritis

Treatment Phase: Subsequent continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment per subsequent continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an erythrocyte sedimentation rate (ESR) no greater than 25 mm per hour or a C-reactive protein (CRP) level no greater than 15 mg per L or either marker reduced by at least 20% from baseline; and

either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following major active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to determine response for all subsequent continuing treatments.

The measurement of response to the prior course of therapy must have been conducted following a minimum of 12 weeks of therapy with this drug and must be documented in the patient's medical records.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

infliximab 100 mg injection, 1 vial

11515C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	5	2	..	*1650.94	^a Inflectra [PF]	^a Renflexis [MK]

■ INFLIXIMAB

Note TREATMENT OF ADULT PATIENTS WITH ANKYLOSING SPONDYLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, certolizumab pegol, etanercept, golimumab, infliximab and secukinumab for adult patients with ankylosing spondylitis.

Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab and secukinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 6 biological medicines at any 1 time.

Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last prescription for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised biological medicine treatment with adalimumab, certolizumab pegol, etanercept, golimumab, infliximab and secukinumab.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised biological medicine treatment in this treatment cycle and wishes to commence such therapy (Initial 1 - New patient)

(ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years); or

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

A patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy.

(b) Continuing treatment.

For the first continuing treatment course of PBS-subsidised biological medicine, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1, Initial 2 or Initial 3 treatment restrictions. For second and subsequent continuing courses of PBS-subsidised biological medicine it is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply. Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be conducted within 1 month of the last dose. Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI), or the prior NSAID therapy and exercise program requirements.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the BASDAI, ESR and/or CRP submitted with the first authority application for a biological medicine.

However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to these revised baseline measurements.

For a new patient, the BASDAI used to determine the baseline must be measured while the patient is receiving NSAID therapy and completing their exercise program.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response.

(4) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must qualify under the Initial 3 treatment restriction. The same clinical criteria and indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI) as for the Initial 1 (New patient) restriction will need to be met, but a re-trial of NSAID therapy and exercise therapy is not required.

Authority required

Ankylosing spondylitis

Treatment Phase: Subsequent continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Ankylosing Spondylitis PBS Authority Application - Supporting Information Form.

An adequate response is defined as an improvement from baseline of at least 2 of the BASDAI and 1 of the following:

- (a) an ESR measurement no greater than 25 mm per hour; or
- (b) a CRP measurement no greater than 10 mg per L; or
- (c) an ESR or CRP measurement reduced by at least 20% from baseline.

Where only 1 acute phase reactant measurement is supplied in the first application for PBS-subsidised treatment, that same marker must be measured and supplied in all subsequent continuing treatment applications.

All measurements provided must be no more than 1 month old at the time of application.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg.

Up to a maximum of 3 repeats will be authorised.

Each application for subsequent continuing treatment with this drug must include an assessment of the patient's response to the prior course of therapy. If the response assessment is not provided at the time of application the patient will be deemed to have failed this course of treatment, unless the patient has experienced serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Ankylosing spondylitis

Treatment Phase: Continuing treatment - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the first continuing treatment restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the subsequent continuing Authority Required (in writing) treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

infliximab 100 mg injection, 1 vial

11489Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	340.93	Remicade [JC]

■ **INFLIXIMAB**

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, etanercept, guselkumab, infliximab, ixekizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, etanercept, guselkumab, infliximab, ixekizumab, secukinumab, tildrakizumab and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, etanercept, guselkumab, infliximab, ixekizumab, secukinumab, tildrakizumab and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February

2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years); or

(iii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or

(iv) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab and 28 weeks of therapy for tildrakizumab and ustekinumab.

It is recommended that a patient be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

Grandfather patients (guselkumab only).

A patient who commenced treatment with guselkumab for chronic plaque psoriasis prior to (1 February 2019) and who continues to receive treatment at the time of application, may qualify for treatment under the Initial treatment - Grandfather treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

Grandfather patients (tildrakizumab only).

A patient who commenced treatment with tildrakizumab for chronic plaque psoriasis prior to (1 February 2019) and who continues to receive treatment at the time of application, may qualify for treatment under the Initial treatment Grandfather treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment. This assessment must be conducted within 1 month of the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24

weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions. For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 the PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 1 month at the time of application.

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Subsequent continuing treatment, Whole body

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per subsequent continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg eight weekly. Up to a maximum of 2 repeats will be authorised.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet including the date of the assessment of the patient's condition.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Subsequent continuing treatment, Face, hand, foot

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per subsequent continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or
- (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg eight weekly. Up to a maximum of 2 repeats will be authorised.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet and face, hand, foot area diagrams including the date of the assessment of the patient's condition.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

infliximab 100 mg injection, 1 vial

11590B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	340.93	Remicade [JC]

■ INFLIXIMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, etanercept, guselkumab, infliximab, ixekizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, etanercept, guselkumab, infliximab, ixekizumab, secukinumab, tildrakizumab and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, etanercept, guselkumab, infliximab, ixekizumab, secukinumab, tildrakizumab and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patients receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years); or

(iii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or

(iv) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab and 28 weeks of therapy for tildrakizumab and ustekinumab.

It is recommended that a patient be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

Grandfather patients (guselkumab only).

A patient who commenced treatment with guselkumab for chronic plaque psoriasis prior to (1 February 2019) and who continues to receive treatment at the time of application, may qualify for treatment under the Initial treatment - Grandfather treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

Grandfather patients (tildrakizumab only).

A patient who commenced treatment with tildrakizumab for chronic plaque psoriasis prior to (1 February 2019) and who continues to receive treatment at the time of application, may qualify for treatment under the Initial treatment Grandfather treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment. This assessment must be conducted within 1 month of the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 the PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 1 month at the time of application.

Note Authority applications for increased quantities/ repeats (where relevant) may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

9602

Severe chronic plaque psoriasis

Treatment Phase: Subsequent continuing treatment, Whole body

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per subsequent continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

The measurement of response to the prior course of therapy must be documented in the patient's medical notes.

Determination of response must be based on the PASI assessment of response to the most recent course of treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Authority required (STREAMLINED)

9584

Severe chronic plaque psoriasis

Treatment Phase: Subsequent continuing treatment, Face, hand, foot

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per subsequent continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or
- (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

The measurement of response to the prior course of therapy must be documented in the patient's medical notes.

Determination of response must be based on the PASI assessment of response to the most recent course of treatment with this drug.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

infliximab 100 mg injection, 1 vial

11595G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	5	2	..	*1650.94	^a Inflectra [PF]	^a Renflexis [MK]

INFLIXIMAB

Note TREATMENT OF ADULT PATIENTS WITH MODERATE TO SEVERE ULCERATIVE COLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, golimumab, infliximab and vedolizumab for adult patients with ulcerative colitis. Patients are eligible for PBS-subsidised treatment with either adalimumab, golimumab, infliximab or vedolizumab at any one time.

Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, golimumab, infliximab and vedolizumab only.

From 1 June 2018, under the PBS, all adult patients will be able to commence a treatment cycle where they may trial PBS-subsidised adalimumab, golimumab, infliximab or vedolizumab without having to experience a disease flare when swapping to one of the alternate agents. Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with adalimumab, golimumab, infliximab or vedolizumab while they continue to show a response to therapy.

A patient who received PBS-subsidised adalimumab, infliximab, vedolizumab treatment prior to 1 June 2018 is considered to start their first cycle as of 1 June 2018. Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised adalimumab, golimumab, infliximab or vedolizumab more than once. Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised therapy before they are eligible to commence the next cycle.

The 5-year break is measured from the date of the last approval for PBS-subsidised adalimumab, golimumab, infliximab or vedolizumab treatment in the most recent cycle to the date of the first application for initial treatment with adalimumab, golimumab, infliximab or vedolizumab under the new treatment cycle.

A patient who has failed fewer than 3 trials of either adalimumab, golimumab, infliximab or vedolizumab in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

(1) How to prescribe PBS-subsidised treatment with adalimumab, golimumab, infliximab and vedolizumab after 1 June 2018.

(a) Initial treatment. Applications for initial treatment should be made where:

(i) an adult patient has received no prior PBS-subsidised treatment with adalimumab, golimumab, infliximab or vedolizumab in this treatment cycle and wishes to commence such therapy (Initial 1 - New patient or recommencement of treatment after more than 5 years break in therapy); or

(ii) an adult patient has received prior PBS-subsidised (initial or continuing) adalimumab, golimumab, infliximab or vedolizumab therapy and wishes to trial an alternate agent (Initial 2 - Change or Re-commencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) an adult patient wishes to re-commence treatment with adalimumab, golimumab, infliximab or vedolizumab following a break in PBS-subsidised therapy with the same agent (Initial 2 - Change or Re-commencement of treatment after a break in therapy of less than 5 years).

Treatment authorisations under Initial 1 and Initial 2 will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, 14 weeks of therapy for golimumab, infliximab and vedolizumab.

A patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of treatment for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for golimumab,

infliximab and vedolizumab. For second and subsequent courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment.

(b) Continuing treatment.

Following the completion of an initial treatment course with adalimumab, golimumab, infliximab or vedolizumab a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed in the month prior to completing their current course of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised treatment is approved, a patient may swap if eligible to the alternate adalimumab, golimumab, infliximab or vedolizumab treatment within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Mayo clinic score or partial Mayo clinic score), or the prior corticosteroid therapy and immunosuppressive therapy. A patient may trial an alternate treatment at any time, regardless of whether they are receiving therapy (initial or continuing) with adalimumab, golimumab, infliximab or vedolizumab at the time of the application. However, they cannot swap to a particular therapy if they have failed to respond to prior treatment with that drug once within the same treatment cycle. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(3) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised adalimumab, golimumab, infliximab or vedolizumab therapy of at least 5 years, must requalify for initial 1 treatment with respect to the scores of disease severity. A patient must have received treatment with a 5-aminosalicylate oral preparation in a standard dose for induction of remission for a minimum of 3 consecutive months, and, either azathioprine or 6-mercaptopurine for a minimum of 3 consecutive months or a tapered course of oral steroids over a 6 week period followed by an appropriately dosed thiopurine agent for a minimum of 3 consecutive months (unless intolerance develops necessitating permanent treatment withdrawal to these agents). These above prior treatments must have been received immediately prior to the time the scores of disease severity being used to trial a second or subsequent course are measured.

(4) Patients 'grandfathered' onto PBS-subsidised treatment with golimumab.

A patient who commenced treatment with golimumab for moderate to severe ulcerative colitis prior to 1 June 2018 and who continues to receive treatment at the time of application, may qualify for treatment under the initial 3 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

Note TREATMENT OF PAEDIATRIC PATIENTS WITH MODERATE TO SEVERE ULCERATIVE COLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) for paediatric patients with infliximab or adalimumab for moderate to severe ulcerative colitis; and infliximab for acute severe ulcerative colitis. Where the term 'biological medicine' appears in the following NOTES and restrictions, it refers to infliximab and adalimumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 2 biological medicines at any one time. Infliximab and adalimumab are PBS-subsidised for moderate to severe disease while only infliximab is PBS-subsidised for acute severe disease.

From 1 June 2017, under the PBS, all paediatric patients will be able to commence a treatment cycle where they may trial each PBS-subsidised biological medicine without having to experience a disease flare when swapping to the alternate agent. Under these arrangements, within a single treatment cycle and depending on the disease severity, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy. A patient who received PBS-subsidised biological medicine treatment prior to 1 June 2017 is considered to have started their treatment cycle as of 1 June 2017. Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than twice. Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle. A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle. There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 June 2017.

(a) Initial treatment. Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised biological medicine treatment in this treatment cycle and wishes to commence such therapy - Initial 1 treatment (new patient or Re-commencement of treatment after more than 5 years break in therapy); or
- (ii) a patient has received prior PBS-subsidised (initial or continuing) treatment with a biological medicine and wishes to trial an alternate agent - Initial 2 treatment (Change or Re-commencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping treatment' below]; or
- (iii) a patient wishes to re-commence treatment with a specific biological medicine following a break in PBS-subsidised therapy with that agent - Initial 2 treatment (Change or Re-commencement of treatment after a break in therapy of less than 5 years).

Treatment authorisations under Initial 1 and Initial 2 treatment will be limited to provide for a maximum of 16 weeks of treatment for adalimumab and 14 weeks of treatment for infliximab. From 1 June 2017, a patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of treatment for adalimumab and

up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab. For second and subsequent courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment.

Adalimumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. For patients weighing 40 kg or greater, one prescription should be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats and the second prescription should be written for 2 doses of 40 mg and 2 repeats. For patients weighing less than 40 kg, one prescription should be written for 2 doses of 40 mg with no repeats and the second prescription should be written for 2 doses of 20 mg with 3 repeats.

(b) Continuing treatment. Following the completion of an initial treatment course with a specific biological medicine, a patient remains eligible to receive up to 24 weeks per course of continuing treatment with that drug under the continuing treatment restriction providing they continue to sustain the response. It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure PBS subsidy criteria are met.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap if eligible to the alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Paediatric Ulcerative Colitis Activity Index (PUCAI) Score, confirmation of ulcerative colitis disease), or the prior conventional therapies of corticosteroids or immunosuppressives. A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving treatment (initial or continuing) with infliximab or adalimumab at the time of the application. However, a patient cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug twice within the same treatment cycle. To ensure a patient receives the maximum treatment opportunities allowed under these swapping arrangements, it is important that they are assessed for response to every course of treatment.

(3) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to recommence treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity. A patient must have received treatment with a 5-aminosalicylate oral preparation in a standard dose for induction of remission for a minimum of 3 consecutive months, and, either azathioprine or 6-mercaptopurine for a minimum of 3 consecutive months or a tapered course of oral steroids over a 6 week period followed by an appropriately dosed thiopurine agent for a minimum of 3 consecutive months (unless intolerance develops necessitating permanent treatment withdrawal to these agents) immediately prior to the time the PUCAI score is measured.

Note Authority applications for increased quantities/ repeats (where relevant) may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

9499

Moderate to severe ulcerative colitis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician; OR
- Must be treated by a specialist paediatric gastroenterologist.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have demonstrated or sustained an adequate response to treatment by having a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1 while receiving treatment with this drug; OR
- Patient must have demonstrated or sustained an adequate response to treatment by having a Paediatric Ulcerative Colitis Activity Index (PUCAI) score of less than 10 while receiving treatment with this drug, if aged 6 to 17 years.

Population criteria:

- Patient must be 6 years of age or older.

Patients who have failed to maintain a partial Mayo clinic score of less than or equal to 2, with no subscore greater than 1, or, patients who have failed to maintain a Paediatric Ulcerative Colitis Activity Index (PUCAI) score of less than 10 (if aged 6 to 17 years) with continuing treatment with this drug, will not be eligible to receive further PBS-subsidised treatment with this drug.

Patients are only eligible to receive continuing PBS-subsidised treatment with this drug in courses of up to 24 weeks at a dose of 5 mg per kg per dose providing they continue to sustain the response.

The measurement of response to the prior course of therapy must be documented in the patient's medical notes.

infliximab 100 mg injection, 1 vial

11796W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	5	2	..	*1650.94	^a Inflectra [PF]	^a Renflexis [MK]

■ INFLIXIMAB

Note TREATMENT OF ADULT PATIENTS WITH MODERATE TO SEVERE ULCERATIVE COLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, golimumab, infliximab and vedolizumab for adult patients with ulcerative colitis. Patients are eligible for PBS-subsidised treatment with either adalimumab, golimumab, infliximab or vedolizumab at any one time.

Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, golimumab, infliximab and vedolizumab only.

From 1 June 2018, under the PBS, all adult patients will be able to commence a treatment cycle where they may trial PBS-

subsidised adalimumab, golimumab, infliximab or vedolizumab without having to experience a disease flare when swapping to one of the alternate agents. Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with adalimumab, golimumab, infliximab or vedolizumab while they continue to show a response to therapy.

A patient who received PBS-subsidised adalimumab, infliximab, vedolizumab treatment prior to 1 June 2018 is considered to start their first cycle as of 1 June 2018. Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised adalimumab, golimumab, infliximab or vedolizumab more than once. Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised adalimumab, golimumab, infliximab or vedolizumab treatment in the most recent cycle to the date of the first application for initial treatment with adalimumab, golimumab, infliximab or vedolizumab under the new treatment cycle.

A patient who has failed fewer than 3 trials of either adalimumab, golimumab, infliximab or vedolizumab in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

(1) How to prescribe PBS-subsidised treatment with adalimumab, golimumab, infliximab and vedolizumab after 1 June 2018.

(a) Initial treatment. Applications for initial treatment should be made where:

(i) an adult patient has received no prior PBS-subsidised treatment with adalimumab, golimumab, infliximab or vedolizumab in this treatment cycle and wishes to commence such therapy (Initial 1 - New patient or recommencement of treatment after more than 5 years break in therapy); or

(ii) an adult patient has received prior PBS-subsidised (initial or continuing) adalimumab, golimumab, infliximab or vedolizumab therapy and wishes to trial an alternate agent (Initial 2 - Change or Re-commencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) an adult patient wishes to re-commence treatment with adalimumab, golimumab, infliximab or vedolizumab following a break in PBS-subsidised therapy with the same agent (Initial 2 - Change or Re-commencement of treatment after a break in therapy of less than 5 years).

Treatment authorisations under Initial 1 and Initial 2 will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, 14 weeks of therapy for golimumab, infliximab and vedolizumab.

A patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of treatment for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for golimumab, infliximab and vedolizumab. For second and subsequent courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment.

(b) Continuing treatment.

Following the completion of an initial treatment course with adalimumab, golimumab, infliximab or vedolizumab a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed in the month prior to completing their current course of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised treatment is approved, a patient may swap if eligible to the alternate adalimumab, golimumab, infliximab or vedolizumab treatment within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Mayo clinic score or partial Mayo clinic score), or the prior corticosteroid therapy and immunosuppressive therapy. A patient may trial an alternate treatment at any time, regardless of whether they are receiving therapy (initial or continuing) with adalimumab, golimumab, infliximab or vedolizumab at the time of the application. However, they cannot swap to a particular therapy if they have failed to respond to prior treatment with that drug once within the same treatment cycle. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(3) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised adalimumab, golimumab, infliximab or vedolizumab therapy of at least 5 years, must requalify for initial 1 treatment with respect to the scores of disease severity. A patient must have received treatment with a 5-aminosalicylate oral preparation in a standard dose for induction of remission for a minimum of 3 consecutive months, and, either azathioprine or 6-mercaptopurine for a minimum of 3 consecutive months or a tapered course of oral steroids over a 6 week period followed by an appropriately dosed thiopurine agent for a minimum of 3 consecutive months (unless intolerance develops necessitating permanent treatment withdrawal to these agents). These above prior treatments must have been received immediately prior to the time the scores of disease severity being used to trial a second or subsequent course are measured.

(4) Patients 'grandfathered' onto PBS-subsidised treatment with golimumab.

A patient who commenced treatment with golimumab for moderate to severe ulcerative colitis prior to 1 June 2018 and who continues to receive treatment at the time of application, may qualify for treatment under the initial 3 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

Note TREATMENT OF PAEDIATRIC PATIENTS WITH MODERATE TO SEVERE ULCERATIVE COLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) for paediatric patients with infliximab or adalimumab for moderate to severe ulcerative colitis; and infliximab for acute severe ulcerative colitis.

Where the term 'biological medicine' appears in the following NOTES and restrictions, it refers to infliximab and adalimumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 2 biological medicines at any one time. Infliximab and adalimumab are PBS-subsidised for moderate to severe disease while only infliximab is PBS-subsidised for acute severe disease.

From 1 June 2017, under the PBS, all paediatric patients will be able to commence a treatment cycle where they may trial

each PBS-subsidised biological medicine without having to experience a disease flare when swapping to the alternate agent. Under these arrangements, within a single treatment cycle and depending on the disease severity, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy. A patient who received PBS-subsidised biological medicine treatment prior to 1 June 2017 is considered to have started their treatment cycle as of 1 June 2017. Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than twice. Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle. A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle. There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 June 2017.

(a) Initial treatment. Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised biological medicine treatment in this treatment cycle and wishes to commence such therapy - Initial 1 treatment (new patient or Recommencement of treatment after more than 5 years break in therapy); or
- (ii) a patient has received prior PBS-subsidised (initial or continuing) treatment with a biological medicine and wishes to trial an alternate agent - Initial 2 treatment (Change or Re-commencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping treatment' below]; or
- (iii) a patient wishes to re-commence treatment with a specific biological medicine following a break in PBS-subsidised therapy with that agent - Initial 2 treatment (Change or Re-commencement of treatment after a break in therapy of less than 5 years).

Treatment authorisations under Initial 1 and Initial 2 treatment will be limited to provide for a maximum of 16 weeks of treatment for adalimumab and 14 weeks of treatment for infliximab. From 1 June 2017, a patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of treatment for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab. For second and subsequent courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment.

Adalimumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. For patients weighing 40 kg or greater, one prescription should be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats and the second prescription should be written for 2 doses of 40 mg and 2 repeats. For patients weighing less than 40 kg, one prescription should be written for 2 doses of 40 mg with no repeats and the second prescription should be written for 2 doses of 20 mg with 3 repeats.

(b) Continuing treatment. Following the completion of an initial treatment course with a specific biological medicine, a patient remains eligible to receive up to 24 weeks per course of continuing treatment with that drug under the continuing treatment restriction providing they continue to sustain the response. It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure PBS subsidy criteria are met.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap if eligible to the alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Paediatric Ulcerative Colitis Activity Index (PUCAI) Score, confirmation of ulcerative colitis disease), or the prior conventional therapies of corticosteroids or immunosuppressives. A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving treatment (initial or continuing) with infliximab or adalimumab at the time of the application. However, a patient cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug twice within the same treatment cycle. To ensure a patient receives the maximum treatment opportunities allowed under these swapping arrangements, it is important that they are assessed for response to every course of treatment.

(3) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to recommence treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity. A patient must have received treatment with a 5-aminosalicylate oral preparation in a standard dose for induction of remission for a minimum of 3 consecutive months, and, either azathioprine or 6-mercaptopurine for a minimum of 3 consecutive months or a tapered course of oral steroids over a 6 week period followed by an appropriately dosed thiopurine agent for a minimum of 3 consecutive months (unless intolerance develops necessitating permanent treatment withdrawal to these agents) immediately prior to the time the PUCAI score is measured.

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Initial 1 - New patient or recommencement of treatment after more than 5 years break in therapy

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician; OR
- Must be treated by a specialist paediatric gastroenterologist.

Clinical criteria:

- Patient must have failed to achieve an adequate response to a 5-aminosalicylate oral preparation in a standard dose for induction of remission for 3 or more consecutive months or have intolerance necessitating permanent treatment withdrawal, **AND**

- Patient must have failed to achieve an adequate response to azathioprine at a dose of at least 2 mg per kg daily for 3 or more consecutive months or have intolerance necessitating permanent treatment withdrawal; OR
- Patient must have failed to achieve an adequate response to 6-mercaptopurine at a dose of at least 1 mg per kg daily for 3 or more consecutive months or have intolerance necessitating permanent treatment withdrawal; OR
- Patient must have failed to achieve an adequate response to a tapered course of oral steroids, starting at a dose of at least 40 mg (for a child, 1 to 2 mg/kg up to 40 mg) prednisolone (or equivalent), over a 6 week period or have intolerance necessitating permanent treatment withdrawal, and followed by a failure to achieve an adequate response to 3 or more consecutive months of treatment of an appropriately dosed thiopurine agent, **AND**
- Patient must have a Mayo clinic score greater than or equal to 6 if an adult patient; OR
- Patient must have a partial Mayo clinic score greater than or equal to 6, provided the rectal bleeding and stool frequency subscores are both greater than or equal to 2 (endoscopy subscore is not required for a partial Mayo clinic score); OR
- Patient must have a Paediatric Ulcerative Colitis Activity Index (PUCAI) Score greater than or equal to 30 if aged 6 to 17 years; OR
- Patient must have previously received induction therapy with this drug for an acute severe episode of ulcerative colitis in the last 4 months and demonstrated an adequate response to induction therapy by achieving and maintaining a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1, or a PUCAI score less than 10 (if aged 6 to 17 years).

Population criteria:

- Patient must be 6 years of age or older.

Application for authorisation must be made in writing and must include:

(a) a completed authority prescription form; and

(b) a completed Ulcerative Colitis PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed current Mayo clinic or partial Mayo clinic or Paediatric Ulcerative Colitis Activity Index (PUCAI) calculation sheet including the date of assessment of the patient's condition; and

(ii) details of prior systemic drug therapy [dosage, date of commencement and duration of therapy].

A maximum quantity and number of repeats to provide for an initial course of this drug consisting of 3 doses at 5 mg per kg body weight per dose to be administered at weeks 0, 2 and 6, or to be administered at 8-weekly intervals for patients who have received prior treatment for an acute severe episode, will be authorised.

All tests and assessments should be performed preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior conventional treatment.

The most recent Mayo clinic, partial Mayo clinic or Paediatric Ulcerative Colitis Activity Index (PUCAI) score must be no more than 1 month old at the time of application.

Where treatment for an acute severe episode has occurred, an adequate response to induction therapy needs to be demonstrated by achieving and maintaining a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1, or a Paediatric Ulcerative Colitis Activity Index (PUCAI) score less than 10 (if aged 6 to 17 years), within the first 12 weeks of receiving this drug for acute severe ulcerative colitis.

Patients who fail to achieve a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1, or a Paediatric Ulcerative Colitis Activity Index (PUCAI) score less than 10 within the first 12 weeks of receiving this drug for ulcerative colitis, or have failed to maintain a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1, or have failed to maintain a PUCAI score less than 10 (if aged 6 to 17 years) with continuing treatment with this drug, will not be eligible to receive further PBS-subsidised treatment with this drug.

A partial Mayo clinic or Paediatric Ulcerative Colitis Activity Index (PUCAI) assessment of the patient's response to this initial course of treatment must be made up to 12 weeks after the first dose for patients administered doses at weeks 0, 2 and 6 (6 weeks following the third dose) so that there is adequate time for a response to be demonstrated.

The measurement of response to the prior course of therapy must be documented in the patient's medical notes.

If treatment with any of the above-mentioned drugs is contraindicated according to the relevant TGA-approved Product Information, details must be provided at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, details of this toxicity must be provided at the time of application.

Details of the accepted toxicities including severity can be found on the Department of Human Services website.

Note Biosimilar preferred prescribing policy

Prescribing of the biosimilar brand Inflectra or Renflexis is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Initial 2 - Change or Re-commencement of treatment after a break in therapy of less than 5 years

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR

- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician; OR
- Must be treated by a specialist paediatric gastroenterologist.

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle; OR
- Patient must have previously received PBS-subsidised treatment with a biological medicine (adalimumab or infliximab) for this condition in this treatment cycle if aged 6 to 17 years, **AND**
- Patient must not have failed PBS-subsidised treatment with this drug for this condition in the current treatment cycle; OR
- Patient must not have failed PBS-subsidised treatment with this drug for this condition in the current treatment cycle more than once if aged 6 to 17 years.

Population criteria:

- Patient must be 6 years of age or older.

Application for authorisation must be made in writing and must include:

(a) a completed authority prescription form; and

(b) a completed Ulcerative Colitis PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed current Mayo clinic or partial Mayo clinic or Paediatric Ulcerative Colitis Activity Index (PUCAI) calculation sheet including the date of assessment of the patient's condition; and

(ii) the number of total failures to courses of biological medicine therapy in the current treatment cycle.

To demonstrate a response to treatment the application must be accompanied by the results of the most recent course of biological medicine therapy within the timeframes specified in the relevant restriction.

If the response assessment to the previous course of biological medicine therapy is not met, the patient will be deemed to have failed therapy with that particular course of biological medicine.

A maximum quantity and number of repeats to provide for an initial course of this drug consisting of 3 doses at 5 mg per kg body weight per dose to be administered at weeks 0, 2 and 6, will be authorised.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg eight weekly.

Up to a maximum of 2 repeats will be authorised.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Balance of supply

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician; OR
- Must be treated by a specialist paediatric gastroenterologist.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 treatment (New patient or Re commencement of treatment after more than 5 years break in therapy) restriction to complete the 3 doses (the initial infusion regimen at 0, 2 and 6 weeks); OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 treatment (Change or Re commencement of treatment after a break in therapy of less than 5 years) restriction to complete the 3 doses (the initial infusion regimen at 0, 2 and 6 weeks), **AND**
- The treatment must provide no more than the balance of up to 3 doses (Initial 1 or Initial 2 treatment).

Population criteria:

- Patient must be 6 years of age or older.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

infliximab 100 mg injection, 1 vial

10184B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	340.93	^a Inflectra [PF]	^a Remicade [JC]
					^a Renflexis [MK]	

■ INFLIXIMAB
Note TREATMENT OF PAEDIATRIC PATIENTS WITH REFRACTORY CROHN DISEASE

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) for paediatric patients with adalimumab for severe refractory Crohn disease and infliximab for moderate to severe refractory Crohn disease. Where the term "biological medicines" appears in the following NOTES and restrictions, it refers to adalimumab and infliximab only.

A patient is eligible for PBS-subsidised treatment with only one PBS-subsidised biological medicine at any one time. For paediatric patients with Crohn disease, infliximab is PBS-subsidised for moderate to severe disease while adalimumab is PBS-subsidised for severe disease.

From 1 August 2015, under the PBS, patients commencing on adalimumab will be able to commence a treatment cycle where they may trial each PBS-subsidised biological medicine without having to experience a disease flare when swapping to infliximab. Patients on infliximab will be able to commence a treatment cycle where they may trial each PBS-subsidised biological medicine but will need to meet a PCDAI score of greater than or equal to 40 when swapping to adalimumab.

Under these arrangements, within a single treatment cycle and depending on the disease severity, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy.

A patient who received PBS-subsidised biological medicine treatment prior to 1 August 2015 is considered to have started their treatment cycle as of 1 August 2015.

Within the same treatment cycle, a paediatric patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than twice.

Once a patient has either failed, or ceased to respond to treatment for this condition 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy for this condition before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle.

A patient who has failed 3 trials or fewer of biological medicine therapy in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 August 2015.

(a) Initial treatment.

Applications for initial treatment should be made where:

- i) a patient has received no prior PBS-subsidised biological medicine therapy in this treatment cycle and wishes to commence such therapy - Initial 1 treatment (new patient or Recommencement of treatment after more than 5 years break in therapy); or
- (ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent - Initial 2 treatment (Change or Re-commencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or
- (iii) a patient wishes to re-commence treatment with a specific biological medicine following a break in PBS-subsidised therapy with that agent - Initial 2 treatment (Change or Re-commencement of treatment after a break in therapy of less than 5 years).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy for adalimumab and 14 weeks of therapy for infliximab.

From 1 August 2015, a patient must be assessed for response to any course of initial PBS-subsidised biological therapy following a minimum of 12 weeks of therapy for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab, and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

For first and subsequent continuing courses of PBS-subsidised biological medicine therapy, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

Adalimumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats and the second prescription should be written for 2 doses of 40 mg and 2 repeats for patients weighing 40 kg or greater. For patients weighing less than 40 kg, one prescription should be written for 2 doses of 40 mg with no repeats and the second prescription should be written for 2 doses of 20 mg with 3 repeats.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient remains eligible to receive up to 24 weeks per course of continuing treatment under the First continuing treatment and Subsequent Continuing treatment restrictions with that drug providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted supply of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine therapy is approved, a patient with severe disease may swap if eligible to the alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Paediatric Crohn Disease Activity Index (PCDAI) Score, confirmation of Crohn disease), or the prior conventional therapies of corticosteroid therapy, immunosuppressive therapy or enteral nutrition. Patients on infliximab may swap to adalimumab within the same treatment cycle provided that their disease severity has progressed to severe disease (i.e. they have a current PCDAI score of 40 or more).

A patient cannot swap to a biological medicine if they have failed to respond to prior treatment with that drug twice within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment within the timeframes specified in the relevant restriction.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the PCDAI submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to these revised baseline

measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to assess response to all subsequent treatments.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to recommence treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity.

Patients must have failed to achieve an adequate response to 2 of the following 3 conventional prior therapies including: (i) a tapered course of steroids, starting at a dose of at least 1 mg per kg or 40 mg (whichever is the lesser) prednisolone (or equivalent), over a 6 week period; (ii) an 8 week course of enteral nutrition; or (iii) immunosuppressive therapy including azathioprine at a dose of at least 2 mg per kg daily for 3 or more months, or, 6-mercaptopurine at a dose of at least 1 mg per kg daily for 3 or more months, or, methotrexate at a dose of at least 10 mg per square metre weekly for 3 or more months immediately prior to the time the PCDAI score is measured.

Authority required

Moderate to severe Crohn disease

Treatment Phase: Initial 1 - New patient or recommencement of treatment after more than 5 years break in therapy

Treatment criteria:

- Must be treated by a gastroenterologist (code 87) or a consultant physician [internal medicine specialising in gastroenterology (code 81)] or a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician or a specialist paediatric gastroenterologist.

Clinical criteria:

- Patient must have confirmed diagnosis of Crohn disease, defined by standard clinical, endoscopic and/or imaging features including histological evidence, **AND**
- Patient must have failed to achieve an adequate response to 2 of the following 3 conventional prior therapies including: (i) a tapered course of steroids, starting at a dose of at least 1 mg per kg or 40 mg (whichever is the lesser) prednisolone (or equivalent), over a 6 week period; (ii) an 8 week course of enteral nutrition; or (iii) immunosuppressive therapy including azathioprine at a dose of at least 2 mg per kg daily for 3 or more months, or, 6-mercaptopurine at a dose of at least 1 mg per kg daily for 3 or more months, or, methotrexate at a dose of at least 10 mg per square metre weekly for 3 or more months; OR
- Patient must have a documented intolerance of a severity necessitating permanent treatment withdrawal or a contra-indication to each of prednisolone (or equivalent), azathioprine, 6-mercaptopurine and methotrexate, **AND**
- Patient must have a Paediatric Crohn Disease Activity Index (PCDAI) Score greater than or equal to 30 preferably whilst still on treatment.

Population criteria:

- Patient must be aged 6 to 17 years inclusive.

Application for authorisation must be made in writing and must include:

(a) a completed authority prescription form; and

(b) a completed Paediatric Crohn Disease PBS Authority Application -Supporting Information Form which includes the following:

- (i) the completed current Paediatric Crohn Disease Activity Index (PCDAI) calculation sheet including the date of assessment of the patient's condition which must be no more than one month old at the time of application; and
- (ii) details of previous systemic drug therapy [dosage, date of commencement and duration of therapy] or dates of enteral nutrition.

The PCDAI score should preferably be obtained whilst on conventional treatment but must be obtained within one month of the last conventional treatment dose.

If treatment with any of the specified prior conventional drugs is contraindicated according to the relevant TGA-approved Product Information, please provide details at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, details of this toxicity must be provided at the time of application.

Details of the accepted toxicities including severity can be found on the Department of Human Services website.

A maximum quantity and number of repeats to provide for an initial course of this drug consisting of 3 doses at 5 mg per kg body weight per dose to be administered at weeks 0, 2 and 6, will be authorised.

A PCDAI assessment of the patient's response to this initial course of treatment must be made up to 12 weeks after the first dose (6 weeks following the third dose) so that there is adequate time for a response to be demonstrated.

This assessment, which will be used to determine eligibility for the first continuing treatment, must be submitted to the Department of Human Services no later than 1 month from the date of completion of this initial course of treatment.

Where a response assessment is not undertaken and submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with this drug.

Note It is recommended that an application for the first continuing treatment is submitted to the Department of Human Services at the time of the 12 week assessment, to ensure continuity of treatment for those patients who meet the first continuing treatment criteria for PBS-subsidised treatment with this drug.

Note Biosimilar preferred prescribing policy

Prescribing of the biosimilar brand Inflectra or Renflexis is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:
Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Moderate to severe Crohn disease

Treatment Phase: Initial 2 - Change or Re-commencement of treatment after a break in therapy of less than 5 years

Treatment criteria:

- Must be treated by a gastroenterologist (code 87) or a consultant physician [internal medicine specialising in gastroenterology (code 81)] or a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician or a specialist paediatric gastroenterologist.

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have failed PBS-subsidised therapy with this drug for this condition more than once in the current treatment cycle.

Population criteria:

- Patient must be aged 6 to 17 years inclusive.

Application for authorisation must be made in writing and must include:

(a) a completed authority prescription form; and

(b) a completed Paediatric Crohn Disease PBS Authority Application -Supporting Information Form which includes the following:

- (i) the completed current Paediatric Crohn Disease Activity Index (PCDAI) Score calculation sheet; and
- (ii) details of prior biological medicine treatment including details of date and duration of treatment.

To demonstrate a response to treatment the application must be accompanied by the results of the most recent course of biological medicine therapy within the timeframes specified in the relevant restriction.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under an initial treatment restriction, the patient must have been assessed for response to that course following a minimum of 12 weeks therapy for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

If the response assessment to the previous course of biological medicine treatment is not submitted as detailed above, the patient will be deemed to have failed therapy with that particular course of biological medicine.

A maximum quantity and number of repeats to provide for an initial course of this drug consisting of 3 doses at 5 mg per kg body weight per dose to be administered at weeks 0, 2 and 6, will be authorised.

A PCDAI assessment of the patient's response to this initial course of treatment must be made up to 12 weeks after the first dose (6 weeks following the third dose) so that there is adequate time for a response to be demonstrated.

This assessment, which will be used to determine eligibility for the first continuing treatment, must be submitted to the Department of Human Services no later than 1 month from the date of completion of this initial course of treatment.

Where a response assessment is not undertaken and submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with this drug.

Note It is recommended that an application for the first continuing treatment is submitted to the Department of Human Services at the time of the 12 week assessment, to ensure continuity of treatment for those patients who meet the first continuing treatment criteria for PBS-subsidised treatment with this drug.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Moderate to severe Crohn disease

Treatment Phase: Balance of supply

Treatment criteria:

- Must be treated by a gastroenterologist (code 87) or a consultant physician [internal medicine specialising in gastroenterology (code 81)] or a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician or a specialist paediatric gastroenterologist.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 treatment (New patient or Re-commencement of treatment after more than 5 years break in therapy) restriction to complete the 3 doses (the initial infusion regimen at 0, 2 and 6 weeks); OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 treatment (Change or Re-commencement of treatment after a break in therapy of less than 5 years) restriction to complete the 3 doses the initial infusion regimen at 0, 2 and 6 weeks); OR
- Patient must have received insufficient therapy with this drug for this condition under the first continuing treatment or subsequent continuing treatment restrictions to complete 24 weeks of treatment, **AND**

- The treatment must provide no more than the balance of up to 3 doses (Initial 1 or Initial 2 treatment) or 2 repeats (first Continuing or Subsequent Continuing treatment).

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Moderate to severe Crohn disease

Treatment Phase: First continuing treatment

Treatment criteria:

- Must be treated by a gastroenterologist (code 87) or a consultant physician [internal medicine specialising in gastroenterology (code 81)] or a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician or a specialist paediatric gastroenterologist.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have a reduction in PCDAI Score by at least 15 points from baseline value; **AND**
- Patient must have a total PCDAI score of 30 points or less.

Population criteria:

- Patient must be aged 6 to 17 years inclusive.

Application for authorisation must be made in writing and must include:

(a) a completed authority prescription form; and

(b) a completed Paediatric Crohn Disease PBS Authority Application - Supporting Information Form, which includes the completed Paediatric Crohn Disease Activity Index (PCDAI) calculation sheet along with the date of the assessment of the patient's condition.

The PCDAI assessment must be no more than 1 month old at the time of application.

The application for first continuing treatment with this drug must include a PCDAI assessment of the patient's response to the initial course of treatment. The assessment must be made up to 12 weeks after the first dose so that there is adequate time for a response to be demonstrated. This assessment must be submitted no later than 4 weeks from the cessation of that treatment course.

Where a response assessment is not undertaken and submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with this drug.

A maximum of 24 weeks of treatment with this drug will be authorised under this restriction.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg eight weekly.

Up to a maximum of 2 repeats will be authorised.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

infliximab 100 mg injection, 1 vial

9612X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	340.93	^a Inflectra [PF] ^a Renflexis [MK]	^a Remicade [JC]

■ INFLIXIMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CROHN DISEASE

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for adult patients with severe Crohn disease. Where the term biological medicine appears in the following NOTES and restrictions, it refers to the tumour necrosis factor (TNF) alfa-antagonists (adalimumab and infliximab), the alpha-4 beta-7 integrin inhibitor (vedolizumab) and the human IgG1kappa monoclonal antibody (ustekinumab).

Patients are eligible for PBS-subsidised treatment with only 1 of the above PBS-subsidised biological medicines at any one time.

From 1 September 2017, under the PBS, all patients will be able to commence a treatment cycle where they may trial PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab without having to experience a disease flare when swapping to the alternate agent. Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with adalimumab, infliximab, vedolizumab or ustekinumab while they continue to show a response to therapy.

A patient who received PBS-subsidised adalimumab, infliximab, or vedolizumab treatment prior to 1 September 2017 is considered to have started their treatment cycle as of 1 September 2017.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab more than once.

Once a patient has either failed or ceased to respond to treatment for this condition 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy for this condition before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab treatment in the most recent cycle to the date of the first application for initial treatment with adalimumab, infliximab, vedolizumab or ustekinumab under the new treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of less than 5 years, may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab therapy after 1 September 2017.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised therapy with adalimumab, infliximab, vedolizumab or ustekinumab in this treatment cycle and wishes to commence such therapy (Initial treatment (new patient or Recommencement of treatment after more than 5 years break in therapy - Initial 1)); or

(ii) a patient has received prior PBS-subsidised (initial or continuing) adalimumab, infliximab, vedolizumab or ustekinumab and wishes to trial an alternate agent (Initial 2 - Change or recommencement) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to re-commence treatment with adalimumab, infliximab, vedolizumab or ustekinumab following a break in PBS-subsidised therapy with that agent (Change or Re-commencement of treatment after a break in therapy of less than 5 years (Initial 2)).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, 14 weeks of therapy for infliximab, 14 weeks of therapy for vedolizumab and 16 weeks for ustekinumab.

From 1 September 2017, a patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy for adalimumab or ustekinumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab or vedolizumab, and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

For subsequent courses of PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

Adalimumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats and the second prescription should be written for 2 doses of 40 mg and 2 repeats for patients weighing 40 kg or greater. For patients weighing less than 40 kg, one prescription should be written for 2 doses of 40 mg with no repeats and the second prescription should be written for 2 doses of 20 mg with 3 repeats.

Ustekinumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be written under S100 (Highly Specialised Drugs) for a weight-based loading dose, containing a quantity of up to 4 vials of 130 mg and no repeats. The second prescription should be written under S85 (General) for the subsequent first dose, containing a quantity of 2 vials of 45 mg and no repeats.

(b) Continuing treatment.

Following the completion of an initial treatment course with adalimumab, infliximab, vedolizumab or ustekinumab, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted supply of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine therapy is approved, a patient may swap if eligible to the alternate adalimumab, infliximab, vedolizumab or ustekinumab within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Crohn Disease Activity Index (CDAI) Score, confirmation of Crohn disease), or the prior conventional therapies of corticosteroid therapy and immunosuppressive therapy.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with adalimumab, infliximab, vedolizumab or ustekinumab at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug once within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment, within the timeframes specified in the relevant restriction.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the CDAI or evidence of intestinal inflammation submitted with the first authority application for adalimumab, infliximab, vedolizumab or ustekinumab. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to assess response to all subsequent treatments.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to recommence treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity. Patients must have received treatment with a corticosteroid and at least 1 immunosuppressive agent, at an adequate dose, for a minimum of 3 consecutive months immediately prior to the time the CDAI score or the indices of intestinal inflammation are measured.

(5) Patients 'grandfathered' onto PBS-subsidised treatment with vedolizumab.

A patient who commenced treatment with vedolizumab for severe Crohn disease prior to 1 August 2015 and who continues to receive treatment at the time of application may qualify for treatment under the initial 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment will

be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment will be assessed under the continuing treatment restriction of the relevant drug.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must requalify for continuing treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' above for further details.

(6) Patients 'grandfathered' onto PBS-subsidised treatment with ustekinumab.

A patient who commenced treatment with ustekinumab for severe Crohn disease prior to 1 September 2017 and who continues to receive treatment at the time of application may qualify for treatment under the initial 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment will be assessed under the continuing treatment restriction of the relevant drug.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must requalify for continuing treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' above for further details.

Authority required

Severe Crohn disease

Treatment Phase: Initial treatment (new patient or Recommencement of treatment after more than 5 years break in therapy - Initial 1)

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have confirmed severe Crohn disease, defined by standard clinical, endoscopic and/or imaging features, including histological evidence, with the diagnosis confirmed by a gastroenterologist or a consultant physician, **AND**
- Patient must have failed to achieve an adequate response to prior systemic therapy with a tapered course of steroids, starting at a dose of at least 40 mg prednisolone (or equivalent), over a 6 week period, **AND**
- Patient must have failed to achieve adequate response to prior systemic immunosuppressive therapy with azathioprine at a dose of at least 2 mg per kg daily for 3 or more consecutive months; OR
- Patient must have failed to achieve adequate response to prior systemic immunosuppressive therapy with 6-mercaptopurine at a dose of at least 1 mg per kg daily for 3 or more consecutive months; OR
- Patient must have failed to achieve adequate response to prior systemic immunosuppressive therapy with methotrexate at a dose of at least 15 mg weekly for 3 or more consecutive months, **AND**
- Patient must have a Crohn Disease Activity Index (CDAI) Score greater than or equal to 300 as evidence of failure to achieve an adequate response to prior systemic therapy; OR
- Patient must have short gut syndrome with diagnostic imaging or surgical evidence, or have had an ileostomy or colostomy; and must have evidence of intestinal inflammation; and must have evidence of failure to achieve an adequate response to prior systemic therapy as specified below; OR
- Patient must have extensive intestinal inflammation affecting more than 50 cm of the small intestine as evidenced by radiological imaging; and must have a Crohn Disease Activity Index (CDAI) Score greater than or equal to 220; and must have evidence of failure to achieve an adequate response to prior systemic therapy as specified below.

Population criteria:

- Patient must be aged 18 years or older.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current Crohn Disease Activity Index (CDAI) calculation sheet including the date of assessment of the patient's condition if relevant; and
 - (ii) details of prior systemic drug therapy [dosage, date of commencement and duration of therapy]; and
 - (iii) the reports and dates of the pathology or diagnostic imaging test(s) nominated as the response criterion, if relevant; and
 - (iv) the date of the most recent clinical assessment; and
 - (v) the signed patient acknowledgement indicating they understand and acknowledge that the PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment.

Evidence of failure to achieve an adequate response to prior therapy must include at least one of the following: (a) patient must have evidence of intestinal inflammation; (b) patient must be assessed clinically as being in a high faecal output state; (c) patient must be assessed clinically as requiring surgery or total parenteral nutrition (TPN) as the next therapeutic option, in the absence of this drug, if affected by short gut syndrome, extensive small intestine disease or is an ostomy patient. Evidence of intestinal inflammation includes: (i) blood: higher than normal platelet count, or, an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour, or, a C-reactive protein (CRP) level greater than 15 mg per L; or (ii) faeces: higher than normal lactoferrin or calprotectin level; or (iii) diagnostic imaging: demonstration of increased uptake of intravenous contrast with thickening of the bowel wall or mesenteric lymphadenopathy or fat streaking in the mesentery; All assessments, pathology tests, and diagnostic imaging studies must be made within 1 month of the date of application and preferably should be performed whilst still on treatment with the most recent course of prior therapies.

If treatment with any of the specified prior conventional drugs is contraindicated according to the relevant TGA-approved Product Information, please provide details at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, details of this toxicity must be provided at the time of application.

Details of the accepted toxicities including severity can be found on the Department of Human Services website.

Any one of the baseline criteria may be used to determine response to an initial course of treatment and eligibility for continued therapy, according to the criteria included in the first or subsequent continuing treatment restrictions. However, the same criterion must be used for any subsequent determination of response to treatment, for the purpose of eligibility for continuing PBS-subsidised therapy.

A maximum quantity and number of repeats to provide for an initial course of this drug consisting of 3 doses at 5 mg per kg body weight per dose to be administered at weeks 0, 2 and 6, will be authorised.

The assessment of the patient's response to this initial course of treatment must be made up to 12 weeks after the first dose (6 weeks following the third dose) so that there is adequate time for a response to be demonstrated.

This assessment, which will be used to determine eligibility for the first continuing treatment, must be submitted to the Department of Human Services no later than 1 month from the date of completion of this initial course of treatment.

Where a response assessment is not undertaken and submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with this drug.

Note It is recommended that an application for the first continuing treatment is submitted to the Department of Human Services at the time of the 12 week assessment, to ensure continuity of treatment for those patients who meet the first continuing treatment criterion for PBS-subsidised treatment with this drug.

Note Biosimilar preferred prescribing policy

Prescribing of the biosimilar brand Inflectra or Renflexis is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe Crohn disease

Treatment Phase: Change or Re-commencement of treatment after a break in therapy of less than 5 years (Initial 2)

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have failed PBS-subsidised therapy with this drug for this condition in the current treatment cycle.

Population criteria:

- Patient must be aged 18 years or older.

Applications for authorisation must be made in writing and must include: (a) a completed authority prescription form; and (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form, which includes the following: (i) the completed current Crohn Disease Activity Index (CDAI) Score calculation sheet including the date of assessment of the patient's condition if relevant; or (ii) the reports and dates of the pathology or diagnostic imaging test(s) used to assess response to therapy for patients with short gut syndrome, extensive small intestine disease or an ostomy, if relevant; and (iii) the date of clinical assessment; and (iv) the details of prior biological medicine treatment including the details of date and duration of treatment.

To demonstrate a response to treatment the application must be accompanied by the results of the most recent course of biological medicine therapy within the timeframes specified in the relevant restriction.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under an initial treatment restriction, the patient must have been assessed for response to that course following a minimum of 12 weeks of therapy for adalimumab or ustekinumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab and vedolizumab and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

If the response assessment to the previous course of biological medicine treatment is not submitted as detailed above, the patient will be deemed to have failed therapy with that particular course of biological medicine.

A maximum quantity and number of repeats to provide for an initial course of this drug consisting of 3 doses at 5 mg per kg body weight per dose to be administered at weeks 0, 2 and 6, will be authorised.

The assessment of the patient's response to this initial course of treatment must be made up to 12 weeks after the first dose (6 weeks following the third dose) so that there is adequate time for a response to be demonstrated.

This assessment, which will be used to determine eligibility for the first continuing treatment, must be submitted to the Department of Human Services no later than 1 month from the date of completion of this initial course of treatment.

Where a response assessment is not undertaken and submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with this drug.

Note It is recommended that an application for the first continuing treatment is submitted to the Department of Human Services at the time of the 12 week assessment, to ensure continuity of treatment for those patients who meet the first continuing treatment criterion for PBS-subsidised treatment with this drug.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe Crohn disease

Treatment Phase: Balance of supply

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial treatment (new patient or Recommencement of treatment after more than 5 years break in therapy - Initial 1) restriction to complete the 3 doses (the initial infusion regimen at 0, 2 and 6 weeks); OR
- Patient must have received insufficient therapy with this drug for this condition under the Change or Re-commencement of treatment after a break in therapy of less than 5 years (Initial 2) restriction to complete the 3 doses (the initial infusion regimen at 0, 2 and 6 weeks); OR
- Patient must have received insufficient therapy with this drug for this condition under the first continuing treatment or subsequent continuing treatment restrictions to complete 24 weeks of treatment, **AND**
- The treatment must provide no more than the balance of up to 3 doses (Initial 1 or Initial 2 treatment) or 2 repeats (first Continuing or Subsequent Continuing treatment).

Population criteria:

- Patient must be aged 18 years or older.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Severe Crohn disease

Treatment Phase: First continuing treatment

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must have an adequate response to this drug defined as a reduction in Crohn Disease Activity Index (CDAI) Score to a level no greater than 150 if assessed by CDAI or if affected by extensive small intestine disease; OR
- Patient must have an adequate response to this drug defined as (a) an improvement of intestinal inflammation as demonstrated by: (i) blood: normalisation of the platelet count, or an erythrocyte sedimentation rate (ESR) level no greater than 25 mm per hour, or a C-reactive protein (CRP) level no greater than 15 mg per L; or (ii) faeces: normalisation of lactoferrin or calprotectin level; or (iii) evidence of mucosal healing, as demonstrated by diagnostic imaging findings, compared to the baseline assessment; or (b) reversal of high faecal output state; or (c) avoidance of the need for surgery or total parenteral nutrition (TPN), if affected by short gut syndrome, extensive small intestine or is an ostomy patient.

Population criteria:

- Patient must be aged 18 years or older.

Applications for authorisation must be made in writing and must include:

(a) a completed authority prescription form; and

(b) a completed Crohn Disease PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed Crohn Disease Activity Index (CDAI) Score calculation sheet including the date of the assessment of the patient's condition, if relevant; or

(ii) the reports and dates of the pathology test or diagnostic imaging test(s) used to assess response to therapy for patients with short gut syndrome, extensive small intestine disease or an ostomy, if relevant; and

(iii) the date of clinical assessment.

All assessments, pathology tests, and diagnostic imaging studies must be made within 1 month of the date of application.

The application for first continuing treatment with this drug must include an assessment of the patient's response to the initial course of treatment. The assessment must be made up to 12 weeks after the first dose so that there is adequate time for a response to be demonstrated. This assessment must be submitted no later than 4 weeks from the cessation of that treatment course.

Where a response assessment is not undertaken and submitted within these timeframes, the patient will be deemed to have failed to respond to treatment with this drug.

A maximum of 24 weeks of treatment with this drug will be authorised under this criterion.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg eight weekly. Up to a maximum of 2 repeats will be authorised.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

infliximab 100 mg injection, 1 vial

9613Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	340.93	^a Inflectra [PF] ^a Renflexis [MK]	^a Remicade [JC]

■ INFLIXIMAB

Note TREATMENT OF COMPLEX REFRACTORY FISTULISING CROHN DISEASE

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for patients with complex refractory fistulising Crohn disease. Where the term "biological medicine" appears in the following NOTES and restrictions, it refers to adalimumab and infliximab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the PBS- subsidised biological medicines for this condition at any one time.

From 1 April 2011, under the PBS, all patients will be able to commence a treatment cycle where they may trial PBS-subsidised adalimumab or infliximab without having to experience a disease flare when swapping to the alternate agent.

Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with adalimumab or infliximab while they continue to show a response to therapy.

A patient who received PBS-subsidised adalimumab or infliximab treatment prior to 1 April 2011 is considered to have started their treatment cycle as of 1 April 2011.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised adalimumab or infliximab more than twice.

Once a patient has either failed or ceased to respond to treatment for this condition 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy for this condition before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised adalimumab or infliximab treatment in the most recent cycle to the date of the first application for initial treatment with adalimumab or infliximab under the new treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of less than 5 years, may commence a further course of treatment within the same treatment cycle.

A patient who has failed 3 trials or fewer of biological medicine therapy in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised adalimumab or infliximab therapy after 1 April 2011.

(a) Initial treatment. Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised adalimumab or infliximab therapy in this treatment cycle and wishes to commence such therapy (Initial treatment (new patient or Recommencement of treatment after more than 5 years break in therapy - Initial 1); or

(ii) a patient has received prior PBS-subsidised (initial or continuing) adalimumab or infliximab therapy and wishes to trial an alternate agent (Initial 2 - Change or recommencement) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to re-commence treatment with adalimumab or infliximab following a break in PBS-subsidised therapy with that agent (Change or Re-commencement of treatment after a break in therapy of less than 5 years (Initial 2)

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy for adalimumab and 14 weeks of therapy for infliximab.

From 1 April 2011, a patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab, and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

For subsequent courses of PBS-subsidised adalimumab or infliximab treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

Adalimumab only: Two completed authority prescriptions must be submitted with every initial application for adalimumab.

One prescription must be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats. The second prescription must be written for 2 doses of 40 mg and 2 repeats.

(b) Continuing treatment.

Adalimumab patients:

Following the completion of an initial treatment course with adalimumab, a patient may qualify to receive up to 24 weeks of

continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

Infliximab patients:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment under the Initial 1 or Initial 2 treatment restrictions.

For the second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted supply of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap if eligible to the alternate biological medicine within the same treatment cycle.

A patient may trial the alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with adalimumab or infliximab at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug two times within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment within the timeframes specified in the relevant restriction.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements submitted with the first authority application for adalimumab or infliximab. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to these revised baseline measurements.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to recommence treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity.

Authority required

Complex refractory Fistulising Crohn disease

Treatment Phase: Initial treatment (new patient or Re commencement of treatment after more than 5 years break in therapy - Initial 1)

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have confirmed Crohn disease, defined by standard clinical, endoscopic and/or imaging features, including histological evidence, with the diagnosis confirmed by a gastroenterologist or a consultant physician, **AND**
- Patient must have an externally draining enterocutaneous or rectovaginal fistula.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Fistulising Crohn Disease PBS Authority Application - Supporting Information Form which includes the following:
 - (i) a completed current Fistula Assessment Form including the date of assessment of the patient's condition; and
 - (ii) a signed patient acknowledgement.

The most recent fistula assessment must be no more than 1 month old at the time of application.

A maximum quantity and number of repeats to provide for an initial course of this drug consisting of 3 doses at 5 mg per kg body weight per dose to be administered at weeks 0, 2 and 6, will be authorised.

An assessment of the patient's response to this initial course of treatment must be made up to 12 weeks after the first dose (up to 6 weeks following the third dose) so that there is adequate time for a response to be demonstrated.

This assessment, which will be used to determine eligibility for the first continuing treatment, must be submitted to the Department of Human Services no later than 1 month from the date of completion of this initial course of treatment.

Where a response assessment is not undertaken and submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with this drug.

Note Biosimilar preferred prescribing policy

Prescribing of the biosimilar brand Inflectra or Renflexis is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note It is recommended that an application for the first continuing treatment is submitted to the Department of Human Services at the time of the 12 week assessment, to ensure continuity of treatment for those patients who meet the first continuing treatment criteria for PBS-subsidised treatment with this drug.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826
HOBART TAS 7001

Authority required

Complex refractory Fistulising Crohn disease

Treatment Phase: Change or Re-commencement of treatment after a break in therapy of less than 5 years (Initial 2)

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have failed PBS-subsidised therapy with this drug for this condition more than once in the current treatment cycle.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Fistulising Crohn Disease PBS Authority Application - Supporting Information Form which includes the following:
 - (i) a completed current Fistula Assessment Form including the date of assessment of the patient's condition; and
 - (ii) details of prior biological medicine treatment including details of date and duration of treatment.

The most recent fistula assessment must be no more than 1 month old at the time of application.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under an initial treatment restriction, the patient must have been assessed for response to that course following a minimum of 12 weeks therapy for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

To demonstrate a response to treatment the application must be accompanied by the results of the most recent course of biological medicine therapy within the timeframes specified in the relevant restriction.

If the response assessment to the previous course of biological medicine treatment is not submitted as detailed above, the patient will be deemed to have failed therapy with that particular course of biological medicine.

A maximum quantity and number of repeats to provide for an initial course of this drug consisting of 3 doses at 5 mg per kg body weight per dose to be administered at weeks 0, 2 and 6, will be authorised.

An assessment of the patient's response to this initial course of treatment must be made up to 12 weeks after the first dose (up to 6 weeks following the third dose) so that there is adequate time for a response to be demonstrated.

This assessment, which will be used to determine eligibility for the first continuing treatment, must be submitted to the Department of Human Services no later than 1 month from the date of completion of this initial course of treatment.

Where a response assessment is not undertaken and submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with this drug.

Note It is recommended that an application for the first continuing treatment is submitted to the Department of Human Services at the time of the 12 week assessment, to ensure continuity of treatment for those patients who meet the first continuing treatment criteria for PBS-subsidised treatment with this drug.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Complex refractory Fistulising Crohn disease

Treatment Phase: Balance of supply

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial treatment (new patient or Re-commencement of treatment after more than 5 years break in therapy - Initial 1) restriction to complete the 3 doses (the initial infusion regimen at 0, 2 and 6 weeks); OR
- Patient must have received insufficient therapy with this drug for this condition under the Change or Re-commencement of treatment after a break in therapy of less than 5 years (Initial 2) restriction to complete the 3 doses (the initial infusion regimen at 0, 2 and 6 weeks); OR
- Patient must have received insufficient therapy with this drug for this condition under the first continuing treatment or subsequent continuing treatment restrictions to complete 24 weeks of treatment, **AND**
- The treatment must provide no more than the balance of up to 3 doses (Initial 1 or Initial 2 treatment) or 2 repeats (first Continuing or Subsequent Continuing treatment).

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Complex refractory Fistulising Crohn disease

Treatment Phase: First continuing treatment

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug.

An adequate response is defined as:

- (a) a decrease from baseline in the number of open draining fistulae of greater than or equal to 50%; and/or
- (b) a marked reduction in drainage of all fistula(e) from baseline, together with less pain and induration as reported by the patient.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Fistulising Crohn Disease PBS Authority Application - Supporting Information Form which includes a completed Fistula Assessment form including the date of the assessment of the patient's condition.

The most recent fistula assessment must be no more than 1 month old at the time of application.

The application for first continuing treatment with this drug must include an assessment of the patient's response to the initial course of treatment. The assessment must be made up to 12 weeks after the first dose so that there is adequate time for a response to be demonstrated. This assessment must be submitted no later than 4 weeks from the cessation of that treatment course.

Where a response assessment is not undertaken and submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with this drug.

A maximum of 24 weeks of treatment with this drug will be authorised under this restriction.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg eight weekly.

Up to a maximum of 2 repeats will be authorised.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

infliximab 100 mg injection, 1 vial

9674E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	340.93	^a Inflectra [PF]	^a Remicade [JC]
					^a Renflexis [MK]	

INFLIXIMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for adults with severe active rheumatoid arthritis. Where the term biological medicine appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alfa antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab), the T-cell co-stimulation modulator (abatacept) and the Janus kinase (JAK) inhibitors (baricitinib, tofacitinib). A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised biological medicine therapy may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once, and
- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised biological medicines for the treatment of rheumatoid arthritis.

Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Stevens Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

For a patient who has failed PBS-subsidised treatment with 2 or 3 TNF-alfa antagonists prior to 1 August 2010 please contact the Department of Human Services on 1800 700 270. A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction.

A patient who has failed fewer than 5 biological medicines and who has a break in treatment of less than 24 months may commence a further course of treatment with a biological medicine under Initial 2 treatment restriction. A patient who has

failed fewer than 5 biological medicines and who has had a break in therapy of longer than 24 months may commence a further course of treatment with a biological medicine under the Initial 3 treatment restriction. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine treatment is stopped to the date of the new application for treatment with a biological medicine.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 April 2019.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised biological medicine treatment and wishes to commence such therapy, excluding rituximab (Initial 1 - new patient); or

(ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to re-commence treatment with a specific biological medicine following a break of less than 24 months in PBS-subsidised therapy with that agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months).

(iv) a patient wishes to re-commence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 24 months (Initial 3 - re-commencement of treatment after a break in biological medicine of more than 24 months) Initial applications for a new patient (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6-month intensive DMARD trial, but prior to ceasing DMARD therapy.

Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, baricitinib, etanercept, golimumab, tocilizumab and tofacitinib, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

Rituximab patients should be assessed following a minimum of 12 weeks after the first infusion, and the assessment should be submitted to the Department of Human Services within 4 weeks to ensure continuity of treatment for those who meet the continuing restriction for PBS subsidised treatment with this drug for this condition.

Where a response assessment is not provided the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment. For second and subsequent courses of PBS-subsidised biological medicine (excluding rituximab) treatment it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to the Department of Human Services within sufficient timeframe to allow for authority assessment.

Abatacept patients:

A patient is eligible to receive one I.V. loading dose when commencing treatment with the subcutaneous formulation. Two prescriptions are required, the first prescription for the I.V. loading dose for sufficient vials for one dose based on the patient's weight with no repeats. The second prescription for the subcutaneous formulation, with a maximum quantity of 4 and up to 3 repeats, must be submitted with the initial application.

Rituximab patients:

A further application may be submitted to the Department of Human Services within sufficient timeframe to allow authority assessment. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

Infliximab and etanercept patients:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction. Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

(2) Swapping therapy

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non- biological medicine therapy requirements except if the patient has had a break in therapy of more than 24 months who would need to requalify with respect to the indices of disease severity. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each biological medicine trialled.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent, unless they have a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug.

Abatacept:

A patient swapping from I.V. abatacept to subcutaneous abatacept will not be eligible for an I.V. loading dose when commencing treatment with the subcutaneous formulation.

Rituximab:

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF- α antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological medicine during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological medicine therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate biological medicine may do so without having to have any treatment-free period.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and the Department of Human Services will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints.

Applications under the Initial 1 treatment restriction for a new patient must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy.

Applications under the Initial 3 treatment restriction for re-commencement of treatment after a break in biological medicine of more than 24 months must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial 1 (new patient)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with disease modifying anti-rheumatic drugs (DMARDs) which must include at least 3 months continuous treatment with each of at least 2 DMARDs, one of which must be methotrexate at a dose of at least 20 mg weekly and one of which must be: (i) hydroxychloroquine at a dose of at least 200 mg daily; or (ii) leflunomide at a dose of at least 10 mg daily; or (iii) sulfasalazine at a dose of at least 2 g daily; OR
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs which, if methotrexate is contraindicated according to the Therapeutic Goods Administration (TGA)-approved Product Information or cannot be tolerated at a 20 mg weekly dose, must include at least 3 months continuous treatment with each of at least 2 of the following DMARDs: (i) hydroxychloroquine at a dose of at least 200 mg daily; and/or (ii) leflunomide at a dose of at least 10 mg daily; and/or (iii) sulfasalazine at a dose of at least 2 g daily; OR
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs which, if 3 or more of methotrexate, hydroxychloroquine, leflunomide and sulfasalazine are contraindicated according to the relevant TGA-approved Product Information or cannot be tolerated at the doses specified above, must include at least 3 months continuous treatment with each of at least 2 DMARDs, with one or more of the following DMARDs being used in place of the DMARDs which are contraindicated or not tolerated: (i) azathioprine at a dose of at least 1 mg/kg per day; and/or (ii) cyclosporin at a dose of at least 2 mg/kg/day; and/or (iii) sodium aurothiomalate at a dose of 50 mg weekly, **AND**
- Patient must not receive more than 22 weeks of treatment under this restriction, **AND**
- The treatment must be given concomitantly with methotrexate at a dose of at least 7.5 mg weekly.

Population criteria:

- Patient must be aged 18 years or older.

If methotrexate is contraindicated according to the TGA-approved product information or cannot be tolerated at a 20 mg weekly dose, the application must include details of the contraindication or intolerance including severity to methotrexate. The maximum tolerated dose of methotrexate must be documented in the application, if applicable.

The application must include details of the DMARDs trialled, their doses and duration of treatment, and all relevant contraindications and/or intolerances including severity.

The requirement to trial at least 2 DMARDs for periods of at least 3 months each can be met using single agents sequentially or by using one or more combinations of DMARDs.

If the requirement to trial 6 months of intensive DMARD therapy with at least 2 DMARDs cannot be met because of contraindications and/or intolerances of a severity necessitating permanent treatment withdrawal to all of the DMARDs specified above, details of the contraindication or intolerance including severity and dose for each DMARD must be provided in the authority application.

The following criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; AND either

(a) a total active joint count of at least 20 active (swollen and tender) joints; or

(b) at least 4 active joints from the following list of major joints:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The joint count and ESR and/or CRP must be determined at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. All measures must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials to provide sufficient drug, based on the weight of the patient, for a single infusion at a dose of 3 mg per kg.

Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note Biosimilar preferred prescribing policy

Prescribing of the biosimilar brand Inflectra or Renflexis is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note The Department of Human Services website (www.humanservices.gov.au) has details of the toxicities, including severity, which will be accepted for the following purposes:

(a) exempting a patient from the requirement to undertake a minimum 3 month trial of methotrexate at a 20 mg weekly dose;

(b) substituting azathioprine, cyclosporin or sodium aurothiomalate for another DMARD as part of the 6 month intensive DMARD trial;

(c) exempting a patient from the requirement for a 6 month trial of intensive DMARD therapy.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 2 (change or re-commencement of treatment after a break in biological medicine of less than 24 months)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must not have failed to respond to previous PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised biological medicine treatment for this condition 5 times, **AND**
- Patient must not receive more than 22 weeks of treatment under this restriction, **AND**
- The treatment must be given concomitantly with methotrexate at a dose of at least 7.5 mg weekly.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either of the Initial 1, Initial 2, Initial 3, first or subsequent continuing treatment restrictions, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials to provide sufficient drug, based on the weight of the patient, for a single infusion at a dose of 3 mg per kg.

Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

A patient who has demonstrated a response to a course of rituximab must have a PBS-subsidised biological therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 3 (re-commencement of treatment after a break in biological medicine of more than 24 months)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 24 months or more from the most recent PBS-subsidised biological medicine for this condition, **AND**
- Patient must not have failed to respond to previous PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised biological medicine treatment for this condition 5 times, **AND**
- The condition must have an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; OR
- The condition must have a C-reactive protein (CRP) level greater than 15 mg per L, **AND**
- The condition must have either (a) a total active joint count of at least 20 active (swollen and tender) joints; or (b) at least 4 active major joints, **AND**
- Patient must not receive more than 22 weeks of treatment under this restriction, **AND**
- The treatment must be given concomitantly with methotrexate at a dose of at least 7.5 mg weekly.

Population criteria:

- Patient must be aged 18 years or older.

Major joints are defined as (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

All measures of joint count and ESR and/or CRP must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials to provide sufficient drug, based on the weight of the patient, for a single infusion at a dose of 3 mg per kg.

Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note Biosimilar preferred prescribing policy

Prescribing of the biosimilar brand Inflectra or Renflexis is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial 1 (new patient) or Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 24 months) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 24 months) - balance of supply

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 22 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 24 months) restriction to complete 22 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 24 months) to complete 22 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 22 weeks treatment available under the above restrictions.

Population criteria:

- Patient must be aged 18 years or older.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Severe active rheumatoid arthritis

Treatment Phase: First continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction, **AND**
- The treatment must be given concomitantly with methotrexate at a dose of at least 7.5 mg weekly.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials to provide sufficient drug, based on the weight of the patient, for a single infusion at a dose of 3 mg per kg.

Up to a maximum of 2 repeats will be authorised.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient has either failed or ceased to respond to a PBS-subsidised biological medicine for this condition 5 times, they will not be eligible to receive further PBS-subsidised treatment with a biological medicine for this condition.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Continuing Treatment - balance of supply.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the first continuing treatment restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the subsequent continuing Authority Required (in writing) treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restrictions.

Population criteria:

- Patient must be aged 18 years or older.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

infliximab 100 mg injection, 1 vial

6397Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	340.93	^a Inflectra [PF]	^a Remicade [JC]
					^a Renflexis [MK]	

■ INFLIXIMAB

Note TREATMENT OF ADULT PATIENTS WITH ANKYLOSING SPONDYLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, certolizumab pegol, etanercept, golimumab, infliximab and secukinumab for adult patients with ankylosing spondylitis. Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab and secukinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 6 biological medicines at any 1 time.

Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last prescription for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised biological medicine treatment with adalimumab, certolizumab pegol, etanercept, golimumab, infliximab and secukinumab.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised biological medicine treatment in this treatment cycle and wishes to commence such therapy (Initial 1 - New patient)

(ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years); or

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

A patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy.

(b) Continuing treatment.

For the first continuing treatment course of PBS-subsidised biological medicine, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1, Initial 2 or Initial 3 treatment restrictions. For second and subsequent continuing courses of PBS-subsidised biological medicine it is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply. Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be conducted within 1 month of the last dose. Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI), or the prior NSAID therapy and exercise program requirements.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the BASDAI, ESR and/or CRP submitted with the first authority application for a biological medicine.

However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to these revised baseline measurements.

For a new patient, the BASDAI used to determine the baseline must be measured while the patient is receiving NSAID therapy and completing their exercise program.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at

baseline, an ESR or CRP level respectively must be used to determine response.

(4) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must qualify under the Initial 3 treatment restriction. The same clinical criteria and indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI) as for the Initial 1 (New patient) restriction will need to be met, but a re-trial of NSAID therapy and exercise therapy is not required.

Authority required

Ankylosing spondylitis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Clinical criteria:

- The condition must be radiographically (plain X-ray) confirmed Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have at least 2 of the following: (i) low back pain and stiffness for 3 or more months that is relieved by exercise but not by rest; or (ii) limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by a score of at least 1 on each of the lumbar flexion and lumbar side flexion measurements of the Bath Ankylosing Spondylitis Metrology Index (BASMI); or (iii) limitation of chest expansion relative to normal values for age and gender, **AND**
- Patient must have failed to achieve an adequate response following treatment with at least 2 non-steroidal anti-inflammatory drugs (NSAIDs), whilst completing an appropriate exercise program, for a total period of 3 months, **AND**
- Patient must not receive more than 18 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

The application must include details of the NSAIDs trialled, their doses and duration of treatment.

If the NSAID dose is less than the maximum recommended dose in the relevant TGA-approved Product Information, the application must include the reason a higher dose cannot be used.

If treatment with NSAIDs is contraindicated according to the relevant TGA-approved Product Information, the application must provide details of the contraindication.

If intolerance to NSAID treatment develops during the relevant period of use which is of a severity to necessitate permanent treatment withdrawal, the application must provide details of the nature and severity of this intolerance.

The following criteria indicate failure to achieve an adequate response and must be demonstrated at the time of the initial application:

- (a) a Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 4 on a 0-10 scale; AND
- (b) an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 10 mg per L.

The BASDAI must be determined at the completion of the 3 month NSAID and exercise trial, but prior to ceasing NSAID treatment. The BASDAI must be no more than 1 month old at the time of initial application.

Both ESR and CRP measures should be provided with the initial treatment application and both must be no more than 1 month old. If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reason this criterion cannot be satisfied.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Ankylosing Spondylitis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) a copy of the radiological report confirming Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis; and
 - (ii) a completed BASDAI Assessment Form; and
 - (iii) a completed Exercise Program Self Certification Form included in the supporting information form.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg.

A maximum quantity and number of repeats to provide for an initial course of this drug consisting of 3 doses at 5 mg per kg body weight per dose to be administered at weeks 0, 2 and 6, will be authorised.

Up to a maximum of 3 repeats will be authorised.

An assessment of a patient's response to an initial course of treatment must be conducted following a minimum of 12 weeks of therapy. An application for the continuing treatment must be accompanied with the assessment of response and submitted to the Department of Human Services no later than 4 weeks from the date of completion of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Details of the toxicities, including severity, which will be accepted for the purposes of administering this restriction can be found on the Department of Human Services website at www.humanservices.gov.au

Note For details on the appropriate minimum exercise program that will be accepted for the purposes of administering this restriction, please refer to the Department of Human Services website at www.humanservices.gov.au

Note Biosimilar preferred prescribing policy

Prescribing of the biosimilar brand Inflectra or Renflexis is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Ankylosing spondylitis

Treatment Phase: Initial treatment - Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- Patient must not receive more than 18 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

The authority application must be made in writing and must include:

- a completed authority prescription form; and
- a completed Ankylosing Spondylitis PBS Authority Application - Supporting Information Form.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg.

Up to a maximum of 3 repeats will be authorised.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to change or recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

An adequate response is defined as an improvement from baseline of at least 2 of the BASDAI and 1 of the following:

- an ESR measurement no greater than 25 mm per hour; or
- a CRP measurement no greater than 10 mg per L; or
- an ESR or CRP measurement reduced by at least 20% from baseline.

Where only 1 acute phase reactant measurement is supplied in the first application for PBS-subsidised treatment, that same marker must be measured and supplied in all subsequent continuing treatment applications.

All measurements provided must be no more than 1 month old at the time of application.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826
HOBART TAS 7001

Authority required

Ankylosing spondylitis

Treatment Phase: Initial treatment - Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must be radiographically (plain X-ray) confirmed Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis, **AND**
- Patient must have at least 2 of the following: (i) low back pain and stiffness for 3 or more months that is relieved by exercise but not by rest; or (ii) limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by a score of at least 1 on each of the lumbar flexion and lumbar side flexion measurements of the Bath Ankylosing Spondylitis Metrology Index (BASMI); or (iii) limitation of chest expansion relative to normal values for age and gender,

AND

- Patient must have a Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 4 on a 0-10 scale that is no more than 4 weeks old at the time of application, **AND**
- Patient must have an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour that is no more than 4 weeks old at the time of application; OR
- Patient must have a C-reactive protein (CRP) level greater than 10 mg per L that is no more than 4 weeks old at the time of application; OR
- Patient must have a clinical reason as to why demonstration of an elevated ESR or CRP cannot be met and the application must state the reason, **AND**
- Patient must not receive more than 18 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Ankylosing Spondylitis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) a copy of the radiological report confirming Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis; and
 - (ii) a completed BASDAI Assessment Form.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg.

A maximum quantity and number of repeats to provide for an initial course of this drug consisting of 3 doses at 5 mg per kg body weight per dose to be administered at weeks 0, 2 and 6, will be authorised.

Up to a maximum of 3 repeats will be authorised.

An assessment of a patient's response to an initial course of treatment must be conducted following a minimum of 12 weeks of therapy. An application for the continuing treatment must be accompanied with the assessment of response and submitted to the Department of Human Services no later than 4 weeks from the date of completion of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Biosimilar preferred prescribing policy

Prescribing of the biosimilar brand Inflectra or Renflexis is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Ankylosing spondylitis

Treatment Phase: Initial treatment - Initial 1 (new patient), Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years) - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the Initial 1 (new patient) restriction to complete 18 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 18 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 18 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 18 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Ankylosing spondylitis

Treatment Phase: First continuing treatment

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

The authority application must be made in writing and must include:

- a completed authority prescription form; and
- a completed Ankylosing Spondylitis PBS Authority Application - Supporting Information Form.

An adequate response is defined as an improvement from baseline of at least 2 of the BASDAI and 1 of the following:

- an ESR measurement no greater than 25 mm per hour; or
- a CRP measurement no greater than 10 mg per L; or
- an ESR or CRP measurement reduced by at least 20% from baseline.

Where only 1 acute phase reactant measurement is supplied in the first application for PBS-subsidised treatment, that same marker must be measured and supplied in all subsequent continuing treatment applications.

All measurements provided must be no more than 1 month old at the time of application.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg.

Up to a maximum of 3 repeats will be authorised.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Ankylosing spondylitis

Treatment Phase: Continuing treatment - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the first continuing treatment restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the subsequent continuing Authority Required (in writing) treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

infliximab 100 mg injection, 1 vial

6448J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	340.93	^a Inflectra [PF] ^a Renflexis [MK]	^a Remicade [JC]

■ INFLIXIMAB
Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE PSORIATIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab for adult patients with severe active psoriatic arthritis. Therefore, where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient receiving PBS-subsidised treatment for psoriatic arthritis is able to commence a treatment cycle where they may trial biological medicines without having to experience a disease flare when swapping to the alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

A patient who received PBS-subsidised adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab or ustekinumab treatment prior to 1 May 2019 is considered to start their first cycle as of 1 May 2019.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence another cycle [further details are under '(5) Recommencement of treatment after a 5-year break in PBS-subsidised therapy' below].

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction. There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe active psoriatic arthritis.

(1) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an alternate medicine (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years) or An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, golimumab secukinumab and tofacitinib, 18 to 20 weeks of therapy for certolizumab pegol (depending upon the dosing regimen), 20 weeks of therapy for ixekizumab, 22 weeks of therapy for infliximab, and 28 weeks of therapy for ustekinumab. It is recommended that a patient be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of PBS-subsidised initial treatment following a minimum of 12 weeks of therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

Grandfather patients (ixekizumab only).

A patient who commenced treatment with ixekizumab for severe psoriatic arthritis prior to 1 March 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

Grandfather patients (tofacitinib only).

A patient who commenced treatment with Tofacitinib for severe psoriatic arthritis prior to 1 May 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment under the Initial 1 or Initial 2 treatment restrictions.

For the second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to re-qualify with respect to either the indices of disease severity (i.e. erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP) level, and active joint count) or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify under the Initial 3 treatment restriction with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application.

However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

Within a treatment cycle a patient may alternate between therapy with any biological medicine of their choice (1 at a time) providing:

(i) they have not received PBS-subsidised treatment with that particular biological medicine previously; or
(ii) they have demonstrated an adequate response to that particular biological medicine if they have previously trialled it on the PBS; and

(iii) they have not previously failed to respond to treatment 3 times in this treatment cycle with that biological medicine.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(4) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the indices of disease severity submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial or change or commencement treatment application is submitted within a treatment cycle and these revised baseline measurements will be used to assess response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to determine response for all subsequent continuing treatments. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. 20 or more active joints), response will be determined according to a reduction in the total number of active joints.

(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment restriction according to the criteria of the relevant restriction and index of disease severity. The application must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response to methotrexate at a dose of at least 20 mg weekly for a minimum period of 3 months, **AND**
- Patient must have failed to achieve an adequate response to sulfasalazine at a dose of at least 2 g per day for a minimum period of 3 months; OR
- Patient must have failed to achieve an adequate response to leflunomide at a dose of up to 20 mg daily for a minimum period of 3 months, **AND**
- Patient must not receive more than 22 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Where treatment with methotrexate, sulfasalazine or leflunomide is contraindicated according to the relevant TGA-approved Product Information, details must be provided at the time of application.

Where intolerance to treatment with methotrexate, sulfasalazine or leflunomide developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

The following initiation criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; and

either

- (a) an active joint count of at least 20 active (swollen and tender) joints; or
 - (b) at least 4 active joints from the following list of major joints:
 - (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).
- If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg.

Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Severe Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

An assessment of a patient's response to an initial course of treatment must be conducted following a minimum of 12 weeks of therapy. An application for the continuing treatment must be accompanied with the assessment of response and submitted to the Department of Human Services no later than 4 weeks from the date of completion of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Biosimilar preferred prescribing policy

Prescribing of the biosimilar brand Inflectra or Renflexis is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 3 months treatment with methotrexate and 3 months treatment with sulfasalazine or leflunomide can be found on the Department of Human Services website (www.humanservices.gov.au)

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial treatment - Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- Patient must not receive more than 22 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an erythrocyte sedimentation rate (ESR) no greater than 25 mm per hour or a C-reactive protein (CRP) level no greater than 15 mg per L or either marker reduced by at least 20% from baseline; and

either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following major active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg.

Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Severe Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to change or recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3, first or subsequent continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Biosimilar preferred prescribing policy

Prescribing of the biosimilar brand Inflectra or Renflexis is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial treatment - Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must have an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; OR
- The condition must have a C-reactive protein (CRP) level greater than 15 mg per L, **AND**
- The condition must have either (a) a total active joint count of at least 20 active (swollen and tender) joints; or (b) at least 4 active major joints, **AND**
- Patient must not receive more than 22 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Major joints are defined as (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

All measures of joint count and ESR and/or CRP must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg.

Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Severe Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3, first or subsequent continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Biosimilar preferred prescribing policy

Prescribing of the biosimilar brand Inflectra or Renflexis is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial 1 (new patient) or Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years) - balance of supply

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 22 weeks treatment; OR

- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 22 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 22 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 22 weeks treatment available under the above restrictions.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Severe psoriatic arthritis

Treatment Phase: First continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an erythrocyte sedimentation rate (ESR) no greater than 25 mm per hour or a C-reactive protein (CRP) level no greater than 15 mg per L or either marker reduced by at least 20% from baseline; and

either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following major active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to determine response for all subsequent continuing treatments.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg eight weekly.

Up to a maximum of 2 repeats will be authorised.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Severe Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either Initial 1, Initial 2, or Initial 3 treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Continuing treatment - balance of supply

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the first continuing treatment restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the subsequent continuing Authority Required (in writing) treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restrictions.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

infliximab 100 mg injection, 1 vial

6496X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	340.93	^a Inflectra [PF] ^a Renflexis [MK]	^a Remicade [JC]

■ INFLIXIMAB
Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, etanercept, guselkumab, infliximab, ixekizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, etanercept, guselkumab, infliximab, ixekizumab, secukinumab, tildrakizumab and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, etanercept, guselkumab, infliximab, ixekizumab, secukinumab, tildrakizumab and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

- a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or
- a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years); or
- a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or
- a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab and 28 weeks of therapy for tildrakizumab and ustekinumab.

It is recommended that a patient be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

Grandfather patients (guselkumab only).

A patient who commenced treatment with guselkumab for chronic plaque psoriasis prior to (1 February 2019) and who continues to receive treatment at the time of application, may qualify for treatment under the Initial treatment - Grandfather treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

Grandfather patients (tildrakizumab only).

A patient who commenced treatment with tildrakizumab for chronic plaque psoriasis prior to (1 February 2019) and who continues to receive treatment at the time of application, may qualify for treatment under the Initial treatment Grandfather treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment. This assessment must be conducted within 1 month of the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or commencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 the PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 1 month at the time of application.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial 1 - Whole body (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis where lesions have been present for at least 6 months from the time of initial diagnosis, **AND**

- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 4 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; and/or (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; and/or (iii) cyclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; and/or (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 22 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, cyclosporin or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application.

Where intolerance to treatment with phototherapy, methotrexate, cyclosporin or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, cyclosporin, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

- (a) A current Psoriasis Area and Severity Index (PASI) score of greater than 15, as assessed, preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior treatment.
- (b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 1 month following cessation of each course of treatment.
- (c) The most recent PASI assessment must be no more than 1 month old at the time of application.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg. Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and
 - (ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition. Demonstration of response should be provided within this timeframe.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Biosimilar preferred prescribing policy

Prescribing of the biosimilar brand Inflectra or Renflexis is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, cyclosporin or acitretin can be found on the Department of Human Services website (www.humanservices.gov.au)

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Whole body (change or recommencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised therapy with this drug for this condition in the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 22 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either of the Initial 1, Initial 2, Initial 3, first or subsequent continuing treatment restrictions, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition. Demonstration of response should be provided within this timeframe.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg. Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

- a completed authority prescription form(s); and
- a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:

- the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and

- details of prior biological treatment, including dosage, date and duration of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Biosimilar preferred prescribing policy

Prescribing of the biosimilar brand Inflectra or Renflexis is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 3, Whole body (re-commencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**

- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must have a current Psoriasis Area and Severity Index (PASI) score of greater than 15, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 22 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

The most recent PASI assessment must be no more than 1 month old at the time of application.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg. Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

- a completed authority prescription form(s); and
- a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition.

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition. Demonstration of response should be provided within this timeframe.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Biosimilar preferred prescribing policy

Prescribing of the biosimilar brand Inflectra or Renflexis is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Face, hand, foot (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 4 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; and/or (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; and/or (iii) cyclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; and/or (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 22 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, cyclosporin or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application.

Where intolerance to treatment with phototherapy, methotrexate, cyclosporin or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, cyclosporin, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met. The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

(a) Chronic plaque psoriasis classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:

(i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe, as assessed, preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior treatment; or

(ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed, preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior treatment;

(b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 1 month following cessation of each course of treatment.

(c) The most recent PASI assessment must be no more than 1 month old at the time of application.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg. Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition; and

(ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition. Demonstration of response should be provided within this timeframe.

The PASI assessment for first continuing or subsequent continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Biosimilar preferred prescribing policy

Prescribing of the biosimilar brand Inflectra or Renflexis is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, cyclosporin or acitretin can be found on the Department of Human Services website (www.humanservices.gov.au)

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Face, hand, foot (change or recommencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised therapy with this drug for this condition in the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**

- Patient must not receive more than 22 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or
- a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either of the Initial 1, Initial 2, Initial 3, first or subsequent continuing treatment restrictions, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition. Demonstration of response should be provided within this timeframe. The PASI assessment for first continuing or subsequent continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg. Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

- a completed authority prescription form(s); and
- a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
 - the completed current Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition; and
 - details of prior biological treatment, including dosage, date and duration of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Biosimilar preferred prescribing policy

Prescribing of the biosimilar brand Inflectra or Renflexis is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 3, Face, hand, foot (re-commencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must be classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:
 - at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or
 - the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**

- Patient must not receive more than 22 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

The most recent PASI assessment must be no more than 1 month old at the time of application.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg. Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition.

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition. Demonstration of response should be provided within this timeframe.

The PASI assessment for first continuing or subsequent continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Biosimilar preferred prescribing policy

Prescribing of the biosimilar brand Inflectra or Renflexis is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial 1, Whole body or Face, hand, foot (new patient) or Initial 2, Whole body or Face, hand, foot (change or recommencement of treatment after a break in biological medicine of less than 5 years) or Initial 3, Whole body or Face, hand, foot (re-commencement of treatment after a break in biological medicine of more than 5 years) - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1, Whole body (new patient) restriction to complete 22 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2, Whole body (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 22 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3, Whole body (re-commencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 22 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 1, Face, hand, foot (new patient) restriction to complete 22 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2, Face, hand, foot (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 22 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3, Face, hand, foot (re-commencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 22 weeks treatment, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- The treatment must provide no more than the balance of up to 22 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a dermatologist.

Note Authority approval for sufficient therapy to complete a maximum of 22 weeks of treatment may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Severe chronic plaque psoriasis

Treatment Phase: First continuing treatment, Whole body

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg eight weekly. Up to a maximum of 2 repeats will be authorised.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: First continuing treatment, Face, hand, foot

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or
- (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg eight weekly. Up to a maximum of 2 repeats will be authorised.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet and face, hand, foot area diagrams including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

The PASI assessment for first continuing or subsequent continuing treatment must be performed on the same affected area assessed at baseline.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Whole body or Continuing treatment, Face, hand, foot - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the first continuing treatment, Whole body restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the first continuing treatment, Face, hand, foot restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the subsequent continuing treatment Authority Required (in writing), Whole body restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the subsequent continuing treatment Authority Required (in writing), Face, hand, foot restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restrictions, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate).

Treatment criteria:

- Must be treated by a dermatologist.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

infliximab 100 mg injection, 1 vial

9617E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	340.93	^a Inflectra [PF] ^a Renflexis [MK]	^a Remicade [JC]

Interleukin inhibitors

■ ANAKINRA

Note This drug is not PBS-subsidised for conditions other than CAPS.

Authority required (STREAMLINED)

5450

Moderate to severe cryopyrin associated periodic syndromes (CAPS)

Treatment criteria:

- Must be treated by a rheumatologist or in consultation with a rheumatologist; OR
 - Must be treated by a clinical immunologist or in consultation with a clinical immunologist.
- A diagnosis of CAPS must be documented in the patient's medical records.

anakinra 100 mg/0.67 mL injection, 28 x 0.67 mL syringes

10263E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	1532.39	Kineret [FK]

■ TOCILIZUMAB
Note TREATMENT OF PATIENTS WITH SEVERE ACTIVE SYSTEMIC JUVENILE IDIOPATHIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of tocilizumab for a patient who has severe active systemic juvenile idiopathic arthritis (sJIA).

From 1 May 2012, a patient receiving PBS-subsidised tocilizumab therapy is considered to be in a treatment cycle. Under these arrangements, within a single treatment cycle, a patient may:

(i) continue to receive long-term treatment with PBS-subsidised tocilizumab while they continue to show a response to therapy, and

(ii) fail to respond, or to sustain a response, to PBS-subsidised tocilizumab twice.

Once a patient has either failed or ceased to sustain a response to 2 courses of treatment, they are deemed to have completed a single treatment cycle and they must have, at a minimum, a 12 month break in PBS-subsidised tocilizumab therapy before they are eligible to commence another cycle. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised tocilizumab treatment was approved to the date of the first application for initial treatment with tocilizumab under the new treatment cycle.

Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient who received PBS-subsidised tocilizumab treatment immediately prior to 1 May 2012 is considered to be in their first cycle as of 1 May 2012. A patient who has had a break in tocilizumab treatment of at least 12 months immediately prior to making a new application, on or after 1 May 2012, will commence a new treatment cycle. A patient who has failed their first course of tocilizumab in a treatment cycle and who has a break in therapy of less than 12 months may commence a second course of treatment within the same treatment cycle under the Initial 2 treatment restriction.

A patient who has failed their first course of tocilizumab in a treatment cycle and who has a break in therapy of more than 12 months must commence a new treatment cycle under the Initial 3 treatment restriction.

How to prescribe PBS-subsidised tocilizumab treatment therapy after 1 May 2012.

(1) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised tocilizumab treatment and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient has received the first course of PBS-subsidised (initial or continuing) tocilizumab therapy in a treatment cycle and is deemed to have failed to respond or sustain a response and the treating physician wishes to trial a second course, provided any break in therapy is less than 12 months (Initial 2 - retreat or recommencement of treatment after a break of less than 12 months); or

(iii) a patient wishes to recommence treatment with tocilizumab following a break in PBS-subsidised therapy of less than 12 months (Initial 2 - retreat or recommencement of treatment after a break of less than 12 months).

(iv) a patient wishes to recommence treatment following a break in PBS-subsidised tocilizumab therapy of more than 12 months (Initial 3 - recommencement of treatment after a break of more than 12 months).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy.

A patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not conducted within these timeframes, the patient will be deemed to have failed this course of treatment.

For the second and subsequent courses of PBS-subsidised tocilizumab, it is recommended that a patient is reviewed in the 4 weeks prior to completing their current course of treatment and that an application is posted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

(2) Continuing treatment.

Following the completion of an initial treatment course with tocilizumab, a patient may qualify to receive up to 24 weeks of continuing treatment with this drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing tocilizumab treatment in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted tocilizumab supply.

Assessments of response to a course of PBS-subsidised therapy must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased. Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with tocilizumab.

(2) Treatment cycle.

Once initial treatment with PBS-subsidised tocilizumab is approved, a patient deemed to have failed to respond to the first course of treatment may have a second course without having to requalify with respect to the indices of disease severity (joint count, fever and/or CRP level and platelet count), except if the patient has had a break in therapy of more than 12 months.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the relevant baseline measurements of the joint count, fever and/or CRP level and platelet count submitted with the first authority application for tocilizumab.

Where a patient is deemed to have failed to respond or to sustain a response to the first course of therapy in a treatment cycle, prescribers may provide new baseline measurements for the second course of treatment within that cycle. The

Department of Human Services will assess response according to these revised baseline measurements. If new baseline measurements are not submitted with the initial application for the second course of treatment, then those submitted with the first course will be used by the Department of Human Services to assess response to the second course.

(4) Recommencement of treatment after a 12 month break in PBS-subsidised therapy.

A patient who wishes to start a second or subsequent treatment cycle following a break in PBS-subsidised tocilizumab therapy of at least 12 months, must requalify for treatment under the Initial 3 treatment restriction.

(5) Withdrawal of treatment after sustained remission.

Withdrawal of treatment with tocilizumab should be considered in a patient who has achieved and sustained complete remission of disease for 12 months. A demonstration of response to the current treatment should be submitted to the Department of Human Services at the time treatment is ceased.

Authority required

Systemic juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Clinical criteria:

- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have polyarticular course disease which has failed to respond adequately to oral or parenteral methotrexate at a dose of at least 15 mg per square metre weekly, alone or in combination with oral or intra-articular corticosteroids, for a minimum of 3 months; OR
- Patient must have polyarticular course disease and have demonstrated severe intolerance of, or toxicity due to, methotrexate; OR
- Patient must have refractory systemic symptoms, demonstrated by an inability to decrease and maintain the dose of prednisolone (or equivalent) below 0.5 mg per kg per day following a minimum of 2 months of therapy, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be under 18 years of age.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

The following criteria indicate failure to achieve an adequate response to prior methotrexate therapy in a patient with polyarticular course disease and must be demonstrated in the patient at the time of the initial application:

(a) an active joint count of at least 20 active (swollen and tender) joints; or

(b) at least 4 active joints from the following list of major joints:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The following criteria indicate failure to achieve an adequate response to prior therapy in a patient with refractory systemic symptoms and must be demonstrated in the patient at the time of the initial application:

(a) an active joint count of at least 2 active joints; and

(b) persistent fever greater than 38 degrees Celsius for at least 5 out of 14 consecutive days; and/or

(c) a C-reactive protein (CRP) level and platelet count above the upper limits of normal (ULN).

The baseline measurements of joint count, fever and/or CRP level and platelet count must be performed preferably whilst on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment.

The same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to determine response for all subsequent continuing treatments.

Severe intolerance to methotrexate is defined as intractable nausea and vomiting and general malaise unresponsive to manoeuvres, including reducing or omitting concomitant non-steroidal anti-inflammatory drugs (NSAIDs) on the day of methotrexate administration, use of folic acid supplementation, or administering the dose of methotrexate in 2 divided doses over 24 hours.

Toxicity due to methotrexate is defined as evidence of hepatotoxicity with repeated elevations of transaminases, bone marrow suppression temporally related to methotrexate use, pneumonitis, or serious sepsis.

If treatment with methotrexate alone or in combination with other treatments is contraindicated according to the relevant TGA-approved Product Information, details must be provided at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, details of this toxicity must be provided at the time of application.

The authority application must be made in writing and must include:

(1) completed authority prescription form(s); and

(2) a completed Systemic Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form which includes the following:

(i) the date of assessment of severe active systemic juvenile idiopathic arthritis;

(ii) details of prior treatment including dose and duration of treatment;

(iii) pathology reports detailing CRP and platelet count where appropriate.

At the time of authority application, the medical practitioner must request the appropriate number of vials of appropriate strength to provide sufficient drug, based on the weight of the patient, for two infusions (one month's supply). A separate authority prescription form must be completed for each strength requested. Up to a maximum of 3 repeats will be authorised.

An assessment of a patient's response to an initial course of treatment must be conducted following a minimum of 12 weeks of therapy. An application for the continuing treatment must be accompanied with the assessment of response and submitted to the Department of Human Services no later than 4 weeks from the date of completion of the most recent

course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Systemic juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 2 (retial or recommencement of treatment after a break of less than 12 months)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with this drug for this condition in the previous 12 months, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug more than once during the current treatment cycle, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be under 18 years of age.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

An adequate response to treatment is defined as:

(a) in a patient with polyarticular course disease:

(i) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(ii) a reduction in the number of the following major active joints, from at least 4, by at least 50%:

- elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

- shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

(b) in a patient with refractory systemic symptoms:

(i) absence of fever greater than 38 degrees Celsius in the preceding seven days; and/or

(ii) a reduction in the C-reactive protein (CRP) level and platelet count by at least 30% from baseline; and/or

(iii) a reduction in the dose of corticosteroid by at least 30% from baseline.

At the time of authority application, the medical practitioner must request the appropriate number of vials of appropriate strength to provide sufficient drug, based on the weight of the patient, for two infusions (one month's supply). A separate authority prescription form must be completed for each strength requested. Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

(1) completed authority prescription form(s); and

(2) a completed Systemic Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form which includes pathology reports detailing C-reactive protein (CRP) level and platelet count where appropriate.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to retial or recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to 2 courses of treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 12 months have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au. Applications for authority to prescribe should be forwarded to:
Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Systemic juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 3 (recommencement of treatment after a break of more than 12 months)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have had a break in treatment of 12 months or more from this drug for this condition, **AND**
- Patient must have polyarticular course disease and the condition must have (a) an active joint count of at least 20 active (swollen and tender) joints; or (b) at least 4 active joints from the following list of major joints: i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or (ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth); OR
- Patient must have refractory systemic symptoms and the condition must have (a) an active joint count of at least 2 active joints; and (b) persistent fever greater than 38 degrees Celsius for at least 5 out of 14 consecutive days; and/or (c) a C-reactive protein (CRP) level and platelet count above the upper limits of normal (ULN), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Population criteria:

- Patient must be under 18 years of age.

The authority application must be made in writing and must include:

- (1) completed authority prescription form(s); and
- (2) a completed Systemic Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form which includes the following:

- (i) the date of assessment of severe active systemic juvenile idiopathic arthritis;
- (ii) pathology reports detailing C-reactive protein (CRP) level and platelet count where appropriate.

The most recent systemic juvenile idiopathic arthritis assessment must be no more than 4 weeks old at the time of application.

At the time of authority application, the medical practitioner must request the appropriate number of vials of appropriate strength to provide sufficient drug, based on the weight of the patient, for two infusions (one month's supply). A separate authority prescription form must be completed for each strength requested. Up to a maximum of 3 repeats will be authorised.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to 2 courses of treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au. Applications for authority to prescribe should be forwarded to:
Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Systemic juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient) or Initial 2 (retrial or recommencement of treatment after a break of less than 12 months) or Initial 3 (recommencement of treatment after a break of more than 12 months) - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (retrial or recommencement of treatment after a break of less than 12 months) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under Initial 3 (recommencement of treatment after a break of more than 12 months) restriction to complete 16 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Systemic juvenile idiopathic arthritis

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

An adequate response to treatment is defined as:

(a) in a patient with polyarticular course disease:

(i) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(ii) a reduction in the number of the following major active joints, from at least 4, by at least 50%:

- elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

- shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

(b) in a patient with refractory systemic symptoms:

(i) absence of fever greater than 38 degrees Celsius in the preceding seven days; and/or

(ii) a reduction in the C-reactive protein (CRP) level and platelet count by at least 30% from baseline; and/or

(iii) a reduction in the dose of corticosteroid by at least 30% from baseline.

Determination of whether a response has been demonstrated to initial and subsequent courses of treatment will be based on the baseline measurements of disease severity submitted with the initial treatment application.

The authority application must be made in writing and must include:

(1) completed authority prescription form(s); and

(2) a completed Systemic Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form which includes baseline and current pathology reports detailing CRP and platelet count where appropriate.

The most recent systemic juvenile idiopathic arthritis assessment must be no more than 1 month old at the time of application.

At the time of authority application, the medical practitioner must request the appropriate number of vials of appropriate strength to provide sufficient drug, based on the weight of the patient, for two infusions (one month's supply). A separate authority prescription form must be completed for each strength requested. Up to a maximum of 5 repeats will be authorised.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either Initial 1, Initial 2, or Initial 3 treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to 2 courses of treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 12 months have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Systemic juvenile idiopathic arthritis

Treatment Phase: Continuing treatment - balance of supply

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restriction.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

tocilizumab 200 mg/10 mL injection, 10 mL vial

1423X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	219.27	Actemra [RO]

tocilizumab 80 mg/4 mL injection, 4 mL vial

1419Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	93.58	Actemra [RO]

tocilizumab 400 mg/20 mL injection, 20 mL vial

1464C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	429.00	Actemra [RO]

■ TOCILIZUMAB

Note TREATMENT OF PATIENTS WITH SEVERE ACTIVE JUVENILE IDIOPATHIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, etanercept and tocilizumab for a patient who has severe active juvenile idiopathic arthritis. Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, etanercept and tocilizumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 3 biological medicines at any one time.

From 1 April 2014, a patient receiving PBS-subsidised biological medicine is considered to be in a treatment cycle where they may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements, within a single treatment cycle, a patient may:

(i) continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy; and

(ii) fail to respond or to sustain a response to each PBS-subsidised biological medicine once only. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single treatment cycle and they must have, at a minimum, a 12 month break in PBS-subsidised biological medicine therapy before they are eligible to commence another cycle.

The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine was approved to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who received PBS-subsidised biological medicine immediately prior to 1 April 2014 is considered to be in their first cycle as of 1 April 2014. A patient who has had a break in biological medicine treatment of at least 12 months immediately prior to making a new application, on or after 1 April 2014, will commence a new treatment cycle under the Initial 3 treatment restriction.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of less than 12 months may commence a further course of treatment within the same treatment cycle under the Initial 2 treatment restriction.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of more than 12 months must commence a new treatment cycle under the Initial 3 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment therapy after 1 April 2014.

(1) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an alternate medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine therapy of less than 12 months) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised

therapy of less than 12 months with the same medicine (initial 2 - Change or Recommencement of treatment after a break in biological medicine therapy of less than 12 months).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 12 months (initial 3 - Recommencement of treatment after a break in biological medicine of more than 12 months) Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy.

A patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not conducted within these timeframes, the patient will be deemed to have failed this course of treatment.

For the second and subsequent cycles of PBS-subsidised biological medicine, it is recommended that a patient is reviewed in the 4 weeks prior to completing their current course of treatment and that an application is posted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services where applicable. Where a response assessment is not submitted, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (joint count) or the prior non-biological medicine therapy requirements, except if the patient has had a break in therapy of more than 12 months who would then need to requalify under the initial 3 restrictions with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application.

However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, an application for a patient who wishes to swap to an alternate biological medicine should be accompanied by the approved authority prescription or remaining repeats for the biological medicine the patient is ceasing.

(4) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count submitted with the first authority application for a biological medicine. However, prescribers may provide a new baseline measurement any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to the revised baseline measurement.

(5) Recommencement of treatment after a 12 months break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological medicine therapy of at least 12 months, must qualify under the initial 3 restriction and meet the relevant criteria and index of disease severity.

(6) Withdrawal of treatment after sustained remission.

Withdrawal of treatment with biological medicine should be considered in a patient who has achieved and sustained complete remission of disease for 12 months. A demonstration of response to the current treatment should be submitted to the Department of Human Services at the time treatment is ceased.

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Treatment criteria:

- Must be treated by a paediatric rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Clinical criteria:

- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have demonstrated severe intolerance of, or toxicity due to, methotrexate; OR
- Patient must have demonstrated failure to achieve an adequate response to 1 or more of the following treatment regimens: (i) oral or parenteral methotrexate at a dose of at least 20 mg per square metre weekly, alone or in combination with oral or intra-articular corticosteroids, for a minimum of 3 months; or (ii) oral methotrexate at a dose of at least 10 mg per square metre weekly together with at least 1 other disease modifying anti-rheumatic drug (DMARD), alone or in combination with corticosteroids, for a minimum of 3 months, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be under 18 years of age.

Severe intolerance to methotrexate is defined as intractable nausea and vomiting and general malaise unresponsive to manoeuvres, including reducing or omitting concomitant non-steroidal anti-inflammatory drugs (NSAIDs) on the day of

methotrexate administration, use of folic acid supplementation, or administering the dose of methotrexate in 2 divided doses over 24 hours.

Toxicity due to methotrexate is defined as evidence of hepatotoxicity with repeated elevations of transaminases, bone marrow suppression temporally related to methotrexate use, pneumonitis, or serious sepsis.

If treatment with methotrexate alone or in combination with another DMARD is contraindicated according to the relevant TGA-approved Product Information, details must be provided at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, details of this toxicity must be provided at the time of application.

The following criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

(a) an active joint count of at least 20 active (swollen and tender) joints; OR

(b) at least 4 active joints from the following list:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The joint count assessment must be performed preferably whilst still on DMARD treatment, but no longer than 4 weeks following cessation of the most recent prior treatment.

The authority application must be made in writing and must include:

(1) completed authority prescription form(s); and

(2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

At the time of authority application, medical practitioners must request the appropriate number of vials of appropriate strength to provide sufficient drug, based on the weight of the patient, for one infusion. A separate authority prescription form must be completed for each strength requested. Up to a maximum of 3 repeats will be authorised.

An assessment of a patient's response to an initial course of treatment must be conducted following a minimum of 12 weeks of therapy. An application for the continuing treatment must be accompanied with the assessment of response and submitted to the Department of Human Services no later than 4 weeks from the date of completion of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Use of alternative DMARDs in children is dependent on approval by the Therapeutic Goods Administration as age restrictions may apply.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 12 months)

Treatment criteria:

- Must be treated by a paediatric rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be under 18 years of age.

An adequate response to treatment is defined as:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

At the time of authority application, medical practitioners must request the appropriate number of vials of appropriate strength to provide sufficient drug, based on the weight of the patient, for one infusion. A separate authority prescription form must be completed for each strength requested. Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

- (1) completed authority prescription form(s); and
- (2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to change or recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient who fails to demonstrate a response to treatment with this drug under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug in this treatment cycle. A patient may re-trial this drug after a minimum of 12 months have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the initial 3 treatment restriction.

If a patient fails to respond to PBS-subsidised biological medicine treatment 3 times (once with each agent) they will not be eligible to receive further PBS-subsidised biological medicine therapy in this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 3 (recommencement of treatment after a break in biological medicine of more than 12 months)

Treatment criteria:

- Must be treated by a paediatric rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have had a break in treatment of 12 months or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must have either (a) a total active joint count of at least 20 active (swollen and tender) joints; or (b) at least 4 active major joints, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be under 18 years of age.

Active joints are defined as:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

All measures of joint count must be no more than 4 weeks old at the time of this application.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of active joints, the response must be demonstrated on the total number of active joints.

At the time of authority application, medical practitioners must request the appropriate number of vials of appropriate strength to provide sufficient drug, based on the weight of the patient, for one infusion. A separate authority prescription form must be completed for each strength requested. Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

- (1) completed authority prescription form(s); and
- (2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient) or Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 12 months) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 12 months) - balance of supply

Treatment criteria:

- Must be treated by a paediatric rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 12 months) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 12 months) restriction to complete 16 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment per continuing treatment course authorised under this restriction.

An adequate response to treatment is defined as:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

Determination of whether a response has been demonstrated to initial and subsequent courses of treatment will be based on the baseline measurement of joint count submitted with the initial treatment application.

The authority application must be made in writing and must include:

- (1) completed authority prescription form(s); and
- (2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

At the time of authority application, medical practitioners must request the appropriate number of vials of appropriate strength to provide sufficient drug, based on the weight of the patient, for one infusion. A separate authority prescription form must be completed for each strength requested. Up to a maximum of 5 repeats will be authorised.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either Initial 1, Initial 2, or Initial 3 treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 12 months have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

If a patient fails to respond to PBS-subsidised biological medicine treatment 3 times (once with each agent) they will not be eligible to receive further PBS-subsidised biological medicine therapy in this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Continuing treatment - balance of supply

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restriction.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

tocilizumab 200 mg/10 mL injection, 10 mL vial

10079L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	219.27	Actemra [RO]

tocilizumab 80 mg/4 mL injection, 4 mL vial

10068X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	93.58	Actemra [RO]

tocilizumab 400 mg/20 mL injection, 20 mL vial

10060L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	429.00	Actemra [RO]

■ TOCILIZUMAB

Note TREATMENT OF ADULT PATIENTS WITH A HISTORY OF JUVENILE IDIOPATHIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, etanercept and tocilizumab for a patient over 18 years who has a history of juvenile idiopathic arthritis with onset prior to the age of 18 years. Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, etanercept and tocilizumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 3 biological medicines at any one time.

From 1 April 2014, a patient receiving PBS-subsidised biological medicine therapy is considered to be in a treatment cycle where they may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements, within a single treatment cycle, a patient may:

(i) continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy; and

(ii) fail to respond or to sustain a response to each PBS-subsidised biological medicine once only. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment.

Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single treatment cycle and they must have, at a minimum, a 5 year break in PBS-subsidised biological medicine therapy before they are eligible to receive further PBS-subsidised biological medicine therapy.

The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine was approved to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who received PBS-subsidised biological medicine treatment immediately prior to 1 April 2014 is considered to be in their first cycle as of 1 April 2014. A patient who has had a break in biological medicine treatment of at least 24 months immediately prior to making a new application, on or after 1 April 2014, will commence a new treatment cycle under the Initial 3 treatment restriction.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of less than 24 months may commence a further course of treatment within the same treatment cycle under the Initial 2 treatment restriction.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of more than 24 months must commence a new treatment cycle under the Initial 3 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine therapy after 1 April 2014.

(1) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy (Initial 1 - New patient); or
- (ii) a patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an alternate medicine (Initial 2 - Change or recommencement of treatment after a break in biological medicine of less than 24 months) [further details are under 'Swapping therapy' below]; or
- (iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 24 months with the same medicine (Initial 2 - Change or recommencement of treatment after a break in biological medicine therapy of less than 24 months); or
- (iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 24 months (Initial 3 - recommencement of treatment after a break in biological medicine of more than 24 months).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy.

A patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not conducted within these timeframes, the patient will be deemed to have failed this course of treatment.

For second and subsequent cycles of PBS-subsidised biological medicine, it is recommended that a patient is reviewed in the 4 weeks prior to completing their current course of treatment and that an application is posted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services where applicable. Where a response assessment is not submitted, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (joint count and ESR/CRP) or the prior non- biological medicine therapy requirements, except if the patient has had a break in therapy of more than 24 months who would then need to requalify under the Initial 3 restrictions with respect to the indices of disease severity.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, an application for a patient who wishes to swap to an alternate biological medicine should be accompanied by the approved authority prescription or remaining repeats for the biological medicine the patient is ceasing.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count submitted with the first authority application for a biological medicine. However, prescribers may provide a new baseline measurement any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to the revised baseline measurement.

(4) Recommencement of treatment after a 24 months break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological medicine therapy of at least 24 months, must qualify under the Initial 3 restriction and meet the relevant criteria and index of disease severity.

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have a documented history of severe active juvenile idiopathic arthritis with onset prior to the age of 18 years, **AND**
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with disease modifying anti-rheumatic drugs (DMARDs) which must include at least 3 months continuous treatment with each of at least 2 DMARDs, one of which must be methotrexate at a dose of at least 20 mg weekly and one of which must be: (i) hydroxychloroquine at a dose of at least 200 mg daily; or (ii) leflunomide at a dose of at least 10 mg daily; or (iii) sulfasalazine at a dose of at least 2 g daily; OR
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs which, if methotrexate is contraindicated according to the Therapeutic Goods Administration (TGA)-approved Product Information or cannot be tolerated at a 20 mg weekly dose, must include at least 3 months continuous treatment with each of at least 2 of the following DMARDs: (i) hydroxychloroquine at a dose of at least 200 mg daily; and/or (ii) leflunomide at a dose of at least 10 mg daily; and/or (iii) sulfasalazine at a dose of at least 2 g daily; OR
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs which, if 3 or more of methotrexate, hydroxychloroquine, leflunomide and sulfasalazine are contraindicated according to the relevant TGA-approved Product Information or cannot be tolerated at the doses specified above, must include at least 3 months continuous treatment with each of at least 2 DMARDs, with one or more of the following DMARDs being used in place of the DMARDs which are contraindicated or not tolerated: (i) azathioprine at a dose of at least 1 mg/kg per day; and/or (ii) cyclosporin at a dose of at least 2 mg/kg/day; and/or (iii) sodium aurothiomalate at a dose of 50 mg weekly, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

If methotrexate is contraindicated according to the TGA-approved Product Information or cannot be tolerated at a 20 mg weekly dose, the application must include details of the contraindication or intolerance to methotrexate. The maximum tolerated dose of methotrexate must be documented in the application, if applicable.

The application must include details of the DMARDs trialed, their doses and duration of treatment, and all relevant contraindications and/or intolerances.

The requirement to trial at least 2 DMARDs for periods of at least 3 months each can be met using single agents sequentially or by using one or more combinations of DMARDs.

If the requirement to trial 6 months of intensive DMARD therapy with at least 2 DMARDs cannot be met because of contraindications and/or intolerances of a severity necessitating permanent treatment withdrawal to all of the DMARDs specified above, details of the contraindication or intolerance and dose for each DMARD must be provided in the authority application.

The following criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; **AND** either

(a) an active joint count of at least 20 active (swollen and tender) joints; or

(b) at least 4 active joints from the following list:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The joint count and ESR and/or CRP must be determined at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. All measures must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

The authority application must be made in writing and must include:

(1) completed authority prescription form(s); and

(2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

At the time of authority application, medical practitioners must request the appropriate number of vials of appropriate strength to provide sufficient drug, based on the weight of the patient, for one infusion. A separate authority prescription form must be completed for each strength requested. Up to a maximum of 3 repeats will be authorised.

An assessment of a patient's response to an initial course of treatment must be conducted following a minimum of 12 weeks of therapy. An application for the continuing treatment must be accompanied with the assessment of response and submitted to the Department of Human Services no later than 4 weeks from the date of completion of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

- Note** The Department of Human Services website (www.humanservices.gov.au) has details of the toxicities, including severity, which will be accepted for the following purposes:
- (a) exempting a patient from the requirement to undertake a minimum 3 month trial of methotrexate at a 20 mg weekly dose;
 - (b) substituting azathioprine, cyclosporin or sodium aurothiomalate for another DMARD as part of the 6 month intensive DMARD trial;
 - (c) exempting a patient from the requirement for a 6 month trial of intensive DMARD therapy.
- Note** Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au. Applications for authority to prescribe should be forwarded to:
- Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 24 months)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have a documented history of severe active juvenile idiopathic arthritis with onset prior to the age of 18 years, **AND**
- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

- (a) an active joint count of fewer than 10 active (swollen and tender) joints; or
- (b) a reduction in the active (swollen and tender) joint count by at least 50% from baseline; or
- (c) a reduction in the number of the following active joints, from at least 4, by at least 50%:
 - (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - (ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

At the time of authority application, medical practitioners must request the appropriate number of vials of appropriate strength to provide sufficient drug, based on the weight of the patient, for one infusion. A separate authority prescription form must be completed for each strength requested. Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

- (1) completed authority prescription form(s); and
- (2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to change or recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient who fails to demonstrate a response to treatment with this drug under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug in this treatment cycle. A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the initial 3 treatment restriction.

If a patient fails to respond to PBS-subsidised biological medicine treatment 3 times (once with each agent) they will not be eligible to receive further PBS-subsidised biological medicine therapy in this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 3 (recommencement of treatment after a break in biological medicine of more than 24 months)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 24 months or more from the most recently approved PBS-subsidised biological medicine for this condition; OR
- Patient must not have received PBS-subsidised biological medicine for at least 5 years if they failed or ceased to respond to PBS-subsidised biological medicine treatment 3 times in their last treatment cycle, **AND**
- The condition must have an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; OR
- The condition must have a C-reactive protein (CRP) level greater than 15 mg per L, **AND**
- The condition must have either (a) a total active joint count of at least 20 active (swollen and tender) joints; or (b) at least 4 active major joints, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Active joints are defined as:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

All measures of joint count must be no more than 4 weeks old at the time of this application.

At the time of authority application, medical practitioners must request the appropriate number of vials of appropriate strength to provide sufficient drug, based on the weight of the patient, for one infusion. A separate authority prescription form must be completed for each strength requested. Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

(1) completed authority prescription form(s); and

(2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient) or Initial 2 (change or recommencement of treatment after break of less than 24 months) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 24 months) - balance of supply

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 24 months) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 24 months) to complete 16 weeks of treatment, **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment per continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

- (a) an active joint count of fewer than 10 active (swollen and tender) joints; or
- (b) a reduction in the active (swollen and tender) joint count by at least 50% from baseline; or
- (c) a reduction in the number of the following active joints, from at least 4, by at least 50%:
 - (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - (ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

- (1) completed authority prescription form(s); and
- (2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

At the time of authority application, medical practitioners must request the appropriate number of vials of appropriate strength to provide sufficient drug, based on the weight of the patient, for one infusion. A separate authority prescription form must be completed for each strength requested. Up to a maximum of 5 repeats will be authorised.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either Initial 1, Initial 2, or Initial 3 treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

If a patient fails to respond to PBS-subsidised biological medicine treatment 3 times (once with each agent) they will not be eligible to receive further PBS-subsidised biological medicine therapy in this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Continuing Treatment - balance of supply

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restriction.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

tocilizumab 200 mg/10 mL injection, 10 mL vial

10071C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	219.27	Actemra [RO]

tocilizumab 80 mg/4 mL injection, 4 mL vial

10073E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	93.58	Actemra [RO]

tocilizumab 400 mg/20 mL injection, 20 mL vial

10078K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	429.00	Actemra [RO]

■ TOCILIZUMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for adults with severe active rheumatoid arthritis. Where the term biological medicine appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alfa antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab), the T-cell co-stimulation modulator (abatacept) and the Janus kinase (JAK) inhibitors (baricitinib, tofacitinib).

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised biological medicine therapy may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once, and
- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised biological medicines for the treatment of rheumatoid arthritis.

Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Stevens Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

For a patient who has failed PBS-subsidised treatment with 2 or 3 TNF-alfa antagonists prior to 1 August 2010 please contact the Department of Human Services on 1800 700 270. A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction.

A patient who has failed fewer than 5 biological medicines and who has a break in treatment of less than 24 months may commence a further course of treatment with a biological medicine under Initial 2 treatment restriction. A patient who has failed fewer than 5 biological medicines and who has had a break in therapy of longer than 24 months may commence a further course of treatment with a biological medicine under the Initial 3 treatment restriction. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine treatment is stopped to the date of the new application for treatment with a biological medicine.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 April 2019.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised biological medicine treatment and wishes to commence such therapy, excluding rituximab (Initial 1 - new patient); or
- (ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months) [further details are under 'Swapping therapy' below]; or
- (iii) a patient wishes to re-commence treatment with a specific biological medicine following a break of less than 24 months in PBS-subsidised therapy with that agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months).
- (iv) a patient wishes to re-commence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 24 months (Initial 3 - re-commencement of treatment after a break in biological medicine of more than 24 months) Initial applications for a new patient (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6-month intensive DMARD trial, but prior to ceasing DMARD therapy.

Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, baricitinib, etanercept, golimumab, tocilizumab and tofacitinib, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

Rituximab patients should be assessed following a minimum of 12 weeks after the first infusion, and the assessment should be submitted to the Department of Human Services within 4 weeks to ensure continuity of treatment for those who meet the continuing restriction for PBS subsidised treatment with this drug for this condition.

Where a response assessment is not provided the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment. For second and subsequent courses of PBS-subsidised biological medicine (excluding rituximab) treatment it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to the Department of Human Services within sufficient timeframe to allow for authority assessment.

Abatacept patients:

A patient is eligible to receive one I.V. loading dose when commencing treatment with the subcutaneous formulation. Two prescriptions are required, the first prescription for the I.V. loading dose for sufficient vials for one dose based on the patient's weight with no repeats. The second prescription for the subcutaneous formulation, with a maximum quantity of 4 and up to 3 repeats, must be submitted with the initial application.

Rituximab patients:

A further application may be submitted to the Department of Human Services within sufficient timeframe to allow authority assessment. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

Infliximab and etanercept patients:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions. For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction. Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

(2) Swapping therapy

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non- biological medicine therapy requirements except if the patient has had a break in therapy of more than 24 months who would need to requalify with respect to the indices of disease severity. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each biological medicine trialled.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent, unless they have a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug.

Abatacept:

A patient swapping from I.V. abatacept to subcutaneous abatacept will not be eligible for an I.V. loading dose when commencing treatment with the subcutaneous formulation.

Rituximab:

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF-alfa antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological medicine during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological medicine therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate biological medicine may do so without having to have any treatment-free period.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and the Department of Human Services will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints.

Applications under the Initial 1 treatment restriction for a new patient must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy.

Applications under the Initial 3 treatment restriction for re-commencement of treatment after a break in biological medicine of more than 24 months must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with disease modifying anti-rheumatic drugs (DMARDs) which must include at least 3 months continuous treatment with each of at least 2 DMARDs, one of which must be methotrexate at a dose of at least 20 mg weekly and one of which must be: (i) hydroxychloroquine at a dose of at least 200 mg daily; or (ii) leflunomide at a dose of at least 10 mg daily; or (iii) sulfasalazine at a dose of at least 2 g daily; OR
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs which, if methotrexate is contraindicated according to the Therapeutic Goods Administration (TGA)-approved Product Information or cannot be tolerated at a 20 mg weekly dose, must include at least 3 months continuous treatment with each of at least 2 of the following DMARDs: (i) hydroxychloroquine at a dose of at least 200 mg daily; and/or (ii) leflunomide at a dose of at least 10 mg daily; and/or (iii) sulfasalazine at a dose of at least 2 g daily; OR
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs which, if 3 or more of methotrexate, hydroxychloroquine, leflunomide and sulfasalazine are contraindicated according to the relevant TGA-approved Product Information or cannot be tolerated at the doses specified above, must include at least 3 months continuous treatment with each of at least 2 DMARDs, with one or more of the following DMARDs being used in place of the DMARDs which are contraindicated or not tolerated: (i) azathioprine at a dose of at least 1 mg/kg per day; and/or (ii) cyclosporin at a dose of at least 2 mg/kg/day; and/or (iii) sodium aurothiomalate at a dose of 50 mg weekly, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

If methotrexate is contraindicated according to the TGA-approved product information or cannot be tolerated at a 20 mg weekly dose, the application must include details of the contraindication or intolerance including severity to methotrexate. The maximum tolerated dose of methotrexate must be documented in the application, if applicable.

The application must include details of the DMARDs trialled, their doses and duration of treatment, and all relevant contraindications and/or intolerances including severity.

The requirement to trial at least 2 DMARDs for periods of at least 3 months each can be met using single agents sequentially or by using one or more combinations of DMARDs.

If the requirement to trial 6 months of intensive DMARD therapy with at least 2 DMARDs cannot be met because of contraindications and/or intolerances of a severity necessitating permanent treatment withdrawal to all of the DMARDs specified above, details of the contraindication or intolerance including severity and dose for each DMARD must be provided in the authority application.

The following criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; **AND** either

- (a) a total active joint count of at least 20 active (swollen and tender) joints; or
- (b) at least 4 active joints from the following list of major joints:
 - (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth). The joint count and ESR and/or CRP must be determined at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. All measures must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

At the time of the authority application, medical practitioners should request the appropriate number of vials of appropriate strength to provide sufficient drug, based on the weight of the patient, for a single infusion at a dose of 8 mg per kg. A separate authority prescription form must be completed for each strength requested.

Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note The Department of Human Services website (www.humanservices.gov.au) has details of the toxicities, including severity, which will be accepted for the following purposes:

- (a) exempting a patient from the requirement to undertake a minimum 3 month trial of methotrexate at a 20 mg weekly dose;
- (b) substituting azathioprine, cyclosporin or sodium aurothiomalate for another DMARD as part of the 6 month intensive DMARD trial;
- (c) exempting a patient from the requirement for a 6 month trial of intensive DMARD therapy.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 2 (change or re-commencement of treatment after a break in biological medicine of less than 24 months)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must not have failed to respond to previous PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised biological medicine treatment for this condition 5 times, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either of the Initial 1, Initial 2, Initial 3, or continuing treatment restrictions, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

At the time of the authority application, medical practitioners should request the appropriate number of vials of appropriate strength to provide sufficient drug, based on the weight of the patient, for a single infusion at a dose of 8 mg per kg. A separate authority prescription form must be completed for each strength requested.

Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

A patient who has demonstrated a response to a course of rituximab must have a PBS-subsidised biological therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 3 (re-commencement of treatment after a break in biological medicine of more than 24 months)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 24 months or more from the most recent PBS-subsidised biological medicine for this condition, **AND**
- Patient must not have failed to respond to previous PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised biological medicine treatment for this condition 5 times, **AND**
- The condition must have an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; OR
- The condition must have a C-reactive protein (CRP) level greater than 15 mg per L, **AND**
- The condition must have either (a) a total active joint count of at least 20 active (swollen and tender) joints; or (b) at least 4 active major joints, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Major joints are defined as (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

All measures of joint count and ESR and/or CRP must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

At the time of the authority application, medical practitioners should request the appropriate number of vials of appropriate strength to provide sufficient drug, based on the weight of the patient, for a single infusion at a dose of 8 mg per kg. A separate authority prescription form must be completed for each strength requested.

Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial 1 (new patient) or Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 24 months) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 24 months) - balance of supply

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 24 months) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 24 months) to complete 16 weeks of treatment, **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment per continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number

of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

At the time of the authority application, medical practitioners should request the appropriate number of vials of appropriate strength to provide sufficient drug, based on the weight of the patient, for a single infusion at a dose of 8 mg per kg. A separate authority prescription form must be completed for each strength requested.

Up to a maximum of 5 repeats will be authorised.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient has either failed or ceased to respond to a PBS-subsidised biological medicine for this condition 5 times, they will not be eligible to receive further PBS-subsidised treatment with a biological medicine for this condition.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Continuing Treatment - balance of supply.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restriction.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

tocilizumab 200 mg/10 mL injection, 10 mL vial

9672C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	219.27	Actemra [RO]

tocilizumab 80 mg/4 mL injection, 4 mL vial

9671B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	93.58	Actemra [RO]

tocilizumab 400 mg/20 mL injection, 20 mL vial

9673D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	429.00	Actemra [RO]

■ USTEKINUMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CROHN DISEASE

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for adult patients with severe Crohn disease. Where the term biological medicine appears in the following NOTES and restrictions, it refers to the tumour necrosis factor (TNF) alfa-antagonists (adalimumab and infliximab), the alpha-4 beta-7 integrin inhibitor (vedolizumab) and the human IgG1kappa monoclonal antibody (ustekinumab).

Patients are eligible for PBS-subsidised treatment with only 1 of the above PBS-subsidised biological medicines at any one time.

From 1 September 2017, under the PBS, all patients will be able to commence a treatment cycle where they may trial PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab without having to experience a disease flare when swapping to the alternate agent. Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with adalimumab, infliximab, vedolizumab or ustekinumab while they continue to show a response to therapy.

A patient who received PBS-subsidised adalimumab, infliximab, or vedolizumab treatment prior to 1 September 2017 is considered to have started their treatment cycle as of 1 September 2017.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab more than once.

Once a patient has either failed or ceased to respond to treatment for this condition 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy for this condition before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab treatment in the most recent cycle to the date of the first application for initial treatment with adalimumab, infliximab, vedolizumab or ustekinumab under the new treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of less than 5 years, may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab therapy after 1 September 2017. (a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised therapy with adalimumab, infliximab, vedolizumab or ustekinumab in this treatment cycle and wishes to commence such therapy (Initial treatment (new patient or Recommencement of treatment after more than 5 years break in therapy - Initial 1)); or

(ii) a patient has received prior PBS-subsidised (initial or continuing) adalimumab, infliximab, vedolizumab or ustekinumab and wishes to trial an alternate agent (Initial 2 - Change or recommencement) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to re-commence treatment with adalimumab, infliximab, vedolizumab or ustekinumab following a break in PBS-subsidised therapy with that agent (Change or Re-commencement of treatment after a break in therapy of less than 5 years (Initial 2)).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, 14 weeks of therapy for infliximab, 14 weeks of therapy for vedolizumab and 16 weeks for ustekinumab.

From 1 September 2017, a patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy for adalimumab or ustekinumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab or vedolizumab, and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

For subsequent courses of PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

Adalimumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats and the second prescription should be written for 2 doses of 40 mg and 2 repeats for patients weighing 40 kg or greater. For patients weighing less than 40 kg, one prescription should be written for 2 doses of 40 mg with no repeats and the second prescription should be written for 2 doses of 20 mg with 3 repeats.

Ustekinumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be written under S100 (Highly Specialised Drugs) for a weight-based loading dose, containing a quantity of up to 4 vials of 130 mg and no repeats. The second prescription should be written under S85 (General) for the subsequent first dose, containing a quantity of 2 vials of 45 mg and no repeats.

(b) Continuing treatment.

Following the completion of an initial treatment course with adalimumab, infliximab, vedolizumab or ustekinumab, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted supply of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine therapy is approved, a patient may swap if eligible to the alternate adalimumab, infliximab, vedolizumab or ustekinumab within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Crohn Disease Activity Index (CDAI) Score, confirmation of Crohn disease), or the prior conventional therapies of corticosteroid therapy and immunosuppressive therapy.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with adalimumab, infliximab, vedolizumab or ustekinumab at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug once within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment, within the timeframes specified in the relevant restriction.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the CDAI or evidence of intestinal inflammation submitted with the first authority application for adalimumab, infliximab, vedolizumab or ustekinumab. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to assess response to all subsequent treatments.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to recommence treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity. Patients must have received

treatment with a corticosteroid and at least 1 immunosuppressive agent, at an adequate dose, for a minimum of 3 consecutive months immediately prior to the time the CDAI score or the indices of intestinal inflammation are measured.

(5) Patients 'grandfathered' onto PBS-subsidised treatment with vedolizumab.

A patient who commenced treatment with vedolizumab for severe Crohn disease prior to 1 August 2015 and who continues to receive treatment at the time of application may qualify for treatment under the initial 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment will be assessed under the continuing treatment restriction of the relevant drug.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must requalify for continuing treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' above for further details.

(6) Patients 'grandfathered' onto PBS-subsidised treatment with ustekinumab.

A patient who commenced treatment with ustekinumab for severe Crohn disease prior to 1 September 2017 and who continues to receive treatment at the time of application may qualify for treatment under the initial 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment will be assessed under the continuing treatment restriction of the relevant drug.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must requalify for continuing treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' above for further details.

Note It is recommended that an application for continuing treatment is submitted to the Department of Human Services at the time of the 12 week assessment, to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised treatment with this drug.

Note No increase in the maximum number of repeats may be authorised.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs Programs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe Crohn disease

Treatment Phase: Initial treatment (new patient - initial 1)

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have confirmed severe Crohn disease, defined by standard clinical, endoscopic and/or imaging features, including histological evidence, with the diagnosis confirmed by a gastroenterologist or a consultant physician, **AND**
- Patient must have failed to achieve an adequate response to prior systemic therapy with a tapered course of steroids, starting at a dose of at least 40 mg prednisolone (or equivalent), over a 6 week period, **AND**
- Patient must have failed to achieve adequate response to prior systemic immunosuppressive therapy with azathioprine at a dose of at least 2 mg per kg daily for 3 or more months; OR
- Patient must have failed to achieve adequate response to prior systemic immunosuppressive therapy with 6-mercaptopurine at a dose of at least 1 mg per kg daily for 3 or more months; OR
- Patient must have failed to achieve adequate response to prior systemic immunosuppressive therapy with methotrexate at a dose of at least 15 mg weekly for 3 or more months, **AND**
- Patient must have a Crohn Disease Activity Index (CDAI) Score greater than or equal to 300 as evidence of failure to achieve an adequate response to prior systemic therapy; OR
- Patient must have short gut syndrome with diagnostic imaging or surgical evidence, or have had an ileostomy or colostomy; and must have evidence of intestinal inflammation; and must have evidence of failure to achieve an adequate response to prior systemic therapy as specified below; OR
- Patient must have extensive intestinal inflammation affecting more than 50 cm of the small intestine as evidenced by radiological imaging; and must have a Crohn Disease Activity Index (CDAI) Score greater than or equal to 220; and must have evidence of failure to achieve an adequate response to prior systemic therapy as specified below.

Population criteria:

- Patient must be aged 18 years or older.

Applications for authorisation must be made in writing and must include:

(a) two completed authority prescription forms; and

(b) a completed Crohn Disease PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed current Crohn Disease Activity Index (CDAI) calculation sheet including the date of assessment of the patient's condition if relevant; and

(ii) details of prior systemic drug therapy [dosage, date of commencement and duration of therapy]; and

(iii) the reports and dates of the pathology or diagnostic imaging test(s) nominated as the response criterion, if relevant; and

(iv) the date of the most recent clinical assessment; and

(v) the signed patient acknowledgement indicating they understand and acknowledge that the PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment

Evidence of failure to achieve an adequate response to prior therapy must include at least one of the following: (a) patient must have evidence of intestinal inflammation; (b) patient must be assessed clinically as being in a high faecal output state; (c) patient must be assessed clinically as requiring surgery or total parenteral nutrition (TPN) as the next therapeutic option, in the absence of this drug, if affected by short gut syndrome, extensive small intestine disease or is an ostomy patient. Evidence of intestinal inflammation includes: (i) blood: higher than normal platelet count, or, an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour, or, a C-reactive protein (CRP) level greater than 15 mg per L; or (ii) faeces: higher than normal lactoferrin or calprotectin level; or (iii) diagnostic imaging: demonstration of increased uptake of intravenous contrast with thickening of the bowel wall or mesenteric lymphadenopathy or fat streaking in the mesentery;

Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be written under S100 (Highly Specialised Drugs) for a weight-based loading dose, containing a quantity of up to 4 vials of 130 mg and no repeats. The second prescription should be written under S85 (General) for the subsequent first dose, containing a quantity of 2 vials of 45 mg and no repeats.

Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period.

All assessments, pathology tests and diagnostic imaging studies must be made within 1 month of the date of application. If treatment with any of the specified prior conventional drugs is contraindicated according to the relevant TGA-approved Product Information, please provide details at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, details of this toxicity must be provided at the time of application.

Details of the accepted toxicities including severity can be found on the Department of Human Services website.

Any one of the baseline criteria may be used to determine response to an initial course of treatment and eligibility for continued therapy, according to the criteria included in the continuing treatment restriction. However, the same criterion must be used for any subsequent determination of response to treatment, for the purpose of eligibility for continuing PBS-subsidised therapy.

A maximum quantity of a weight based loading dose is up to 4 vials with no repeats and the subsequent dose of 90 mg (2 vials of 45 mg) with no repeats provide for an initial 16 week course of this drug will be authorised.

The assessment of the patient's response to this initial course of treatment must be made following a minimum of 12 weeks of therapy so that there is adequate time for a response to be demonstrated.

This assessment, which will be used to determine eligibility for further continuing treatment, must be submitted to the Department of Human Services no later than 1 month from the date of completion of this course of treatment.

Where a response assessment is not undertaken and submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with this drug.

Note Increase in the maximum quantity or number of units up to 4 may be authorised for the purpose of weight-based loading dose.

Authority required

Severe Crohn disease

Treatment Phase: Change or Re-commencement of treatment (initial 2)

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have a documented history of severe Crohn disease, **AND**
- Patient must have received prior PBS-subsidised treatment with a biological disease modifying drug for this condition in this treatment cycle, **AND**
- Patient must not have failed PBS-subsidised therapy with this drug for this condition in the current treatment cycle.

Population criteria:

- Patient must be aged 18 years or older.

Applications for authorisation must be made in writing and must include:

(a) two completed authority prescription forms; and

(b) a completed Crohn Disease PBS Authority Application - Supporting Information Form, which includes the following:

(i) the completed Crohn Disease Activity Index (CDAI) Score calculation sheet including the date of the assessment of the patient's condition, if relevant; or

(ii) the reports and dates of the pathology or diagnostic imaging test(s) used to assess response to therapy for patients with short gut syndrome, extensive small intestine disease or an ostomy, if relevant; and

(iii) the date of clinical assessment; and

(iv) the details of prior biological disease modifying drug treatment including the details of date and duration of treatment.

Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be written under S100 (Highly Specialised Drugs) for a weight-based loading dose, containing a quantity of up to 4 vials of 130 mg and no repeats. The second prescription should be written under S85 (General) for 2 vials of 45 mg and no repeats.

To demonstrate a response to treatment the application must be accompanied by the results of the most recent course of biological disease modifying drug (bDMD) therapy within the timeframes specified in the relevant restriction.

Where the most recent course of PBS-subsidised bDMD treatment was approved under an initial treatment restriction, the patient must have been assessed for response to that course following a minimum of 12 weeks of therapy for adalimumab

and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab and vedolizumab and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

If the response assessment to the previous course of bDMD treatment is not submitted as detailed above, the patient will be deemed to have failed therapy with that particular course of bDMD.

A maximum quantity of a weight based loading dose is up to 4 vials with no repeats and the subsequent first dose of 90 mg (2 vials of 45 mg) with no repeats provide for an initial 16 week course of this drug will be authorised.

The assessment of the patient's response to this initial course of treatment must be made following a minimum of 12 weeks of therapy so that there is adequate time for a response to be demonstrated.

This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to the Department of Human Services no later than 1 month from the date of completion of this initial course of treatment.

Where a response assessment is not undertaken and submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with this drug.

Note Increase in the maximum quantity or number of units up to 4 may be authorised for the purpose of weight-based loading dose.

ustekinumab 130 mg/26 mL injection, 26 mL vial

11164N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	4	..	..	*16976.63	Stelara [JC]

Calcineurin inhibitors

■ CICLOSPORIN

Caution Careful monitoring of patients is mandatory.

Authority required

Management of transplant rejection

Clinical criteria:

- The treatment must be used by organ or tissue transplant recipients.

ciclosporin 50 mg/mL injection, 10 x 1 mL ampoules

6109M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	65.49	Sandimmun [NV]

■ CICLOSPORIN

Caution Careful monitoring of patients is mandatory.

Authority required

Management of transplant rejection

Treatment Phase: Management (initiation, stabilisation and review of therapy)

Clinical criteria:

- Patient must have had an organ or tissue transplantation, **AND**
- The treatment must be under the supervision and direction of a transplant unit.

Authority required

Severe atopic dermatitis

Treatment Phase: Management (initiation, stabilisation and review of therapy)

Treatment criteria:

- Must be treated by a dermatologist; OR
- Must be treated by a clinical immunologists.

Clinical criteria:

- The condition must be ineffective to other systemic therapies; OR
- The condition must be inappropriate for other systemic therapies.

Authority required

Severe psoriasis

Treatment Phase: Management (initiation, stabilisation and review of therapy)

Clinical criteria:

- The condition must be ineffective to other systemic therapies; OR
- The condition must be inappropriate for other systemic therapies, **AND**
- The condition must have caused significant interference with quality of life.

Treatment criteria:

- Must be treated by a dermatologist.

Authority required

Nephrotic syndrome

Treatment Phase: Management (initiation, stabilisation and review of therapy)

Clinical criteria:

- Patient must have failed prior treatment with steroids and cytostatic drugs; OR
- Patient must be intolerant to treatment with steroids and cytostatic drugs; OR
- The condition must be considered inappropriate for treatment with steroids and cytostatic drugs, **AND**
- Patient must not have renal impairment.

Treatment criteria:

- Must be treated by a nephrologist.

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Management (initiation, stabilisation and review of therapy)

Clinical criteria:

- The condition must have been ineffective to prior treatment with classical slow-acting anti-rheumatic agents (including methotrexate); OR
- The condition must be considered inappropriate for treatment with slow-acting anti-rheumatic agents (including methotrexate).

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist.

ciclosporin 25 mg capsule, 30

6352H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	4	5	..	*141.43	^a Cyclosporin Sandoz [SZ]	^a Neoral 25 [NV]

ciclosporin 10 mg capsule, 60

6232B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*85.79	Neoral 10 [NV]

ciclosporin 100 mg/mL oral liquid, 50 mL

6125J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	4	5	..	*1310.55	Neoral [NV]

ciclosporin 50 mg capsule, 30

6353J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	4	5	..	*286.27	^a Cyclosporin Sandoz [SZ]	^a Neoral 50 [NV]

ciclosporin 100 mg capsule, 30

6354K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	4	5	..	*575.63	^a Cyclosporin Sandoz [SZ]	^a Neoral 100 [NV]

■ TACROLIMUS

Caution Careful monitoring of patients is mandatory.

Authority required

Management of rejection in patients following organ or tissue transplantation

Clinical criteria:

- The treatment must be under the supervision and direction of a transplant unit, **AND**
- The treatment must include initiation, stabilisation, and review of therapy as required.

tacrolimus 1 mg modified release capsule, 60

9682N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*281.21	ADVAGRAF XL [LQ]

tacrolimus 2 mg capsule, 100

10879N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*1047.39	Tacrolimus Sandoz [SZ]

tacrolimus 5 mg capsule, 50

6217F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	5	..	*1143.83	^a Pacrolim [AF] ^a Prograf [LL] ^a TACROLIMUS APOTEX [TX]	^a Pharmacor Tacrolimus 5 [CR] ^a Tacrograf [RW] ^a Tacrolimus Sandoz [SZ]

tacrolimus 750 microgram capsule, 100

10875J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*397.39	Tacrolimus Sandoz [SZ]

tacrolimus 5 mg modified release capsule, 30

9683P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*787.85	ADVAGRAF XL [LQ]

tacrolimus 500 microgram modified release capsule, 30

9681M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*86.43	ADVAGRAF XL [LQ]

tacrolimus 500 microgram capsule, 100

6328C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	5	..	*235.57	^a Pacrolim [AF] ^a Prograf [LL] ^a TACROLIMUS APOTEX [TX]	^a Pharmacor Tacrolimus 0.5 [CR] ^a Tacrograf [RW] ^a Tacrolimus Sandoz [SZ]

tacrolimus 1 mg capsule, 100

6216E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	5	..	*463.75	^a Pacrolim [AF] ^a Prograf [LL] ^a TACROLIMUS APOTEX [TX]	^a Pharmacor Tacrolimus 1 [CR] ^a Tacrograf [RW] ^a Tacrolimus Sandoz [SZ]

Other immunosuppressants
■ LENALIDOMIDE

Note Special Pricing Arrangements apply.

Authority required

Myelodysplastic syndrome

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be limited to a maximum duration of 16 weeks, **AND**
- Patient must be classified as Low risk or Intermediate-1 according to the International Prognostic Scoring System (IPSS), **AND**
- Patient must have a deletion 5q cytogenetic abnormality with or without additional cytogenetic abnormalities, **AND**
- Patient must be red blood cell transfusion dependent.

Classification of a patient as Low risk requires a score of 0 on the IPSS, achieved with the following combination: less than 5% marrow blasts with good karyotypic status (normal, -Y alone, -5q alone, -20q alone), and 0/1 cytopenias.

Classification of a patient as Intermediate-1 requires a score of 0.5 to 1 on the IPSS, achieved with the following possible combinations:

1. 5%-10% marrow blasts with good karyotypic status (normal, -Y alone, -5q alone, -20q alone), and 0/1 cytopenias; OR
2. less than 5% marrow blasts with intermediate karyotypic status (other abnormalities), and 0/1 cytopenias; OR
3. less than 5% marrow blasts with good karyotypic status (normal, -Y alone, -5q alone, -20q alone), and 2/3 cytopenias; OR
4. less than 5% marrow blasts with intermediate karyotypic status (other abnormalities), and 2/3 cytopenias; OR
5. 5%-10% marrow blasts with intermediate karyotypic status (other abnormalities), and 0/1 cytopenias; OR
6. 5%-10% marrow blasts with good karyotypic status (normal, -Y alone, -5q alone, -20q alone), and 2/3 cytopenias; OR
7. less than 5% marrow blasts with poor karyotypic status (complex, greater than 3 abnormalities), and 0/1 cytopenias.

Classification of a patient as red blood cell transfusion dependent requires that:

- (i) the patient has been transfused within the last 8 weeks; and
- (ii) the patient has received at least 8 units of red blood cell in the last 6 months prior to commencing PBS-subsidised therapy with lenalidomide; and would be expected to continue this requirement without lenalidomide treatment.

Patients receiving lenalidomide under the PBS listing must be registered in the i-access risk management program.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Myelodysplastic Syndrome Lenalidomide Authority Application - Supporting Information Form; and
- (c) a copy of the bone marrow biopsy report demonstrating that the patient has myelodysplastic syndrome; and
- (d) a copy of the full blood examination report; and
- (e) a copy of the pathology report detailing the cytogenetics demonstrating Low risk or Intermediate-1 disease according to the IPSS (note: using Fluorescence in Situ Hybridization (FISH) to demonstrate MDS -5q is acceptable); and
- (f) details of transfusion requirements including: (i) the date of most recent transfusion and the number of red blood cell units transfused; and (ii) the total number of red cell units transfused in the 4 and 6 months preceding the date of this application; and
- (g) a signed patient acknowledgement form.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Myelodysplastic syndrome

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must be classified as Low risk or Intermediate-1 according to the International Prognostic Scoring System (IPSS), **AND**
- Patient must have a deletion 5q cytogenetic abnormality with or without additional cytogenetic abnormalities, **AND**

- Patient must have received PBS-subsidised initial therapy with lenalidomide for myelodysplastic syndrome, **AND**
- Patient must have achieved and maintained transfusion independence; or least a 50% reduction in red blood cell unit transfusion requirements compared with the four month period prior to commencing initial PBS-subsidised therapy with lenalidomide, **AND**
- Patient must not have progressive disease.

Patients receiving lenalidomide under the PBS listing must be registered in the i-access risk management program.

The first authority application for continuing supply must be made in writing. Subsequent authority applications for continuing supply may be made by telephone.

The following evidence of response must be provided at each application:

- a haemoglobin level taken within the last 4 weeks; and
- the date of the last transfusion; and
- a statement of the number of units of red cells transfused in the 4 months immediately preceding this application; and
- a statement confirming that the patient has not progressed to acute myeloid leukaemia.

Note Written applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Subsequent authority applications for continuing treatment may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

lenalidomide 5 mg capsule, 21

2798G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	3	..	5170.15	Revlimid [CJ]

lenalidomide 10 mg capsule, 21

2796E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	3	..	5408.55	Revlimid [CJ]

■ LENALIDOMIDE

Note Special Pricing Arrangements apply.

Authority required

Multiple myeloma

Treatment Phase: Initial PBS-subsidised treatment

Clinical criteria:

- The condition must be confirmed by a histological diagnosis, **AND**
 - The treatment must be as monotherapy; OR
 - The treatment must be in combination with dexamethasone, **AND**
 - Patient must have progressive disease after at least one prior therapy, **AND**
 - Patient must have undergone or be ineligible for a primary stem cell transplant, **AND**
 - Patient must not be receiving concomitant PBS-subsidised bortezomib, carfilzomib or thalidomide or its analogues.
- Progressive disease is defined as at least 1 of the following:
- at least a 25% increase and an absolute increase of at least 5 g per L in serum M protein (monoclonal protein); or
 - at least a 25% increase in 24-hour urinary light chain M protein excretion, and an absolute increase of at least 200 mg per 24 hours; or
 - in oligo-secretory and non-secretory myeloma patients only, at least a 50% increase of the difference between involved free light chain and uninvolved free light chain; or
 - at least a 25% relative increase and at least a 10% absolute increase in plasma cells in a bone marrow aspirate or on biopsy; or
 - an increase in the size or number of lytic bone lesions (not including compression fractures); or
 - at least a 25% increase in the size of an existing or the development of a new soft tissue plasmacytoma (determined by clinical examination or diagnostic imaging); or
 - development of hypercalcaemia (corrected serum calcium greater than 2.65 mmol per L not attributable to any other cause).

Oligo-secretory and non-secretory patients are defined as having active disease with less than 10 g per L serum M protein.

The authority application must be made in writing and must include:

- a completed authority prescription form; and
- a completed Multiple Myeloma lenalidomide Authority Application - Supporting Information Form, which includes details of the histological diagnosis of multiple myeloma, prior treatments including name(s) of drug(s) and date of most recent treatment cycle and record of prior stem cell transplant or ineligibility for prior stem cell transplant; details of the basis of the diagnosis of progressive disease or failure to respond; and nomination of which disease activity parameters will be used to assess response; and
- a signed patient acknowledgment.

To enable confirmation of eligibility for treatment, current diagnostic reports of at least one of the following must be provided:

- the level of serum monoclonal protein; or
- Bence-Jones proteinuria - the results of 24-hour urinary light chain M protein excretion; or
- the serum level of free kappa and lambda light chains; or

- (d) bone marrow aspirate or trephine; or
(e) if present, the size and location of lytic bone lesions (not including compression fractures); or
(f) if present, the size and location of all soft tissue plasmacytomas by clinical or radiographic examination i.e. MRI or CT-scan; or
(g) if present, the level of hypercalcaemia, corrected for albumin concentration.

As these parameters will be used to determine response, results for either (a) or (b) or (c) should be provided for all patients. Where the patient has oligo-secretory or non-secretory multiple myeloma, either (c) or (d) or if relevant (e), (f) or (g) should be provided. Where the prescriber plans to assess response in patients with oligo-secretory or non-secretory multiple myeloma with free light chain assays, evidence of the oligo-secretory or non-secretory nature of the multiple myeloma (current serum M protein less than 10 g per L) must be provided.

Patients receiving lenalidomide under the PBS listing must be registered in the i-access risk management program.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Multiple myeloma

Treatment Phase: Continuing PBS-subsidised treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for relapsed or refractory multiple myeloma, **AND**
- The treatment must be as monotherapy; **OR**
- The treatment must be in combination with dexamethasone, **AND**
- Patient must not be receiving concomitant PBS-subsidised bortezomib, carfilzomib or thalidomide or its analogues.

Patients receiving lenalidomide under the PBS listing must be registered in the i-access risk management program.

Note Authority applications for continuing treatment may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note Written applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

lenalidomide 5 mg capsule, 21

9642L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	5170.15	Revlimid [CJ]

lenalidomide 15 mg capsule, 21

9644N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	6299.92	Revlimid [CJ]

lenalidomide 10 mg capsule, 21

9643M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	5408.55	Revlimid [CJ]

lenalidomide 25 mg capsule, 21

9645P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	6634.88	Revlimid [CJ]

■ LENALIDOMIDE

Caution This drug is a category X drug and must not be given to pregnant women. If lenalidomide is taken during pregnancy, a teratogenic effect of lenalidomide in humans cannot be ruled out.

Note Special Pricing Arrangements apply.

Authority required

Multiple myeloma

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be newly diagnosed, **AND**
- The condition must be confirmed by a histological diagnosis, **AND**
- Patient must be ineligible for a primary stem cell transplantation, **AND**
- Patient must not be receiving concomitant PBS-subsidised bortezomib, thalidomide or its analogues, **AND**
- The treatment must be in combination with dexamethasone.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Multiple Myeloma lenalidomide Authority Application - Supporting Information Form, which includes details of the histological diagnosis of multiple myeloma, and ineligibility for prior stem cell transplant; and nomination of which disease activity parameters will be used to assess response; and
- (3) a signed patient acknowledgement.

To enable confirmation of eligibility for treatment, current diagnostic reports of at least one of the following must be provided:

- (a) the level of serum monoclonal protein; or
- (b) Bence-Jones proteinuria - the results of 24-hour urinary light chain M protein excretion; or
- (c) the serum level of free kappa and lambda light chains; or
- (d) bone marrow aspirate or trephine; or
- (e) if present, the size and location of lytic bone lesions (not including compression fractures); or
- (f) if present, the size and location of all soft tissue plasmacytomas by clinical or radiographic examination i.e. MRI or CT-scan; or
- (g) if present, the level of hypercalcaemia, corrected for albumin concentration.

As these parameters will be used to determine response, results for either (a) or (b) or (c) should be provided for all patients. Where the patient has oligo-secretory or non-secretory multiple myeloma, either (c) or (d) or if relevant (e), (f) or (g) should be provided. Where the prescriber plans to assess response in patients with oligo-secretory or non-secretory multiple myeloma with free light chain assays, evidence of the oligo-secretory or non-secretory nature of the multiple myeloma (current serum M protein less than 10 g per L) must be provided.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Patient must be registered in the i-access risk management program.

Authority required

Multiple myeloma

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously been authorised with a PBS prescription with this drug for the condition, **AND**
- Patient must not have demonstrated progressive disease, **AND**
- Patient must not be receiving concomitant PBS-subsidised bortezomib, thalidomide or its analogues, **AND**
- The treatment must be in combination with dexamethasone.

Progressive disease is defined as at least 1 of the following:

- (a) at least a 25% increase and an absolute increase of at least 5 g per L in serum M protein (monoclonal protein); or
- (b) at least a 25% increase in 24-hour urinary light chain M protein excretion, and an absolute increase of at least 200 mg per 24 hours; or
- (c) in oligo-secretory and non-secretory myeloma patients only, at least a 50% increase of the difference between involved free light chain and uninvolved free light chain; or
- (d) at least a 25% relative increase and at least a 10% absolute increase in plasma cells in a bone marrow aspirate or on biopsy; or
- (e) an increase in the size or number of lytic bone lesions (not including compression fractures); or
- (f) at least a 25% increase in the size of an existing or the development of a new soft tissue plasmacytoma (determined by clinical examination or diagnostic imaging); or
- (g) development of hypercalcaemia (corrected serum calcium greater than 2.65 mmol per L not attributable to any other cause).

Oligo-secretory and non-secretory patients are defined as having active disease with less than 10 g per L serum M protein.

Patients receiving this drug under the PBS listing must be registered in the i-access risk management program.

Note Authority applications for continuing treatment may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note Written applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

lenalidomide 5 mg capsule, 21

11036W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	5170.15	Revlimid [CJ]

lenalidomide 15 mg capsule, 21

11042E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	6299.92	Revlimid [CJ]

lenalidomide 10 mg capsule, 21

11063G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	5408.55	Revlimid [CJ]

lenalidomide 25 mg capsule, 21

11055W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	6634.88	Revlimid [CJ]

■ POMALIDOMIDE

Caution This drug is a category X drug and must not be given to pregnant women. Pregnancy in female patients or in the partners of male patients must be avoided during treatment and for 1 month after cessation of treatment.

Note Special Pricing Arrangements apply.

Authority required

Multiple myeloma

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be in combination with dexamethasone, **AND**
- Patient must have undergone or be ineligible for a primary stem cell transplant, **AND**
- Patient must have experienced treatment failure with lenalidomide, unless contraindicated or not tolerated according to the Therapeutic Goods Administration (TGA) approved Product Information, **AND**
- Patient must have experienced treatment failure with bortezomib, unless contraindicated or not tolerated according to the Therapeutic Goods Administration (TGA) approved Product Information, **AND**
- Patient must not be receiving concomitant PBS-subsidised bortezomib, carfilzomib or thalidomide or its analogues. Bortezomib treatment failure is the absence of achieving at least a partial response or as progressive disease during treatment or within 6 months of discontinuing treatment with bortezomib. Lenalidomide treatment failure is progressive disease during treatment or within 6 months of discontinuing treatment with lenalidomide.

If treatment with either bortezomib or lenalidomide is contraindicated according to the relevant TGA-approved Product Information, the application must provide details of the contraindication.

If intolerance to either bortezomib or lenalidomide treatment develops during the relevant period of use which is of a severity to necessitate withdrawal of the treatment, the application must provide details of the nature and severity of this intolerance. Progressive disease is defined as at least 1 of the following:

- at least a 25% increase and an absolute increase of at least 5 g per L in serum M protein (monoclonal protein); or
- at least a 25% increase in 24-hour urinary light chain M protein excretion, and an absolute increase of at least 200 mg per 24 hours; or
- in oligo-secretory and non-secretory myeloma patients only, at least a 50% increase of the difference between involved free light chain and uninvolved free light chain; or
- at least a 25% relative increase and at least a 10% absolute increase in plasma cells in a bone marrow aspirate or on biopsy; or
- an increase in the size or number of lytic bone lesions (not including compression fractures); or
- at least a 25% increase in the size of an existing or the development of a new soft tissue plasmacytoma (determined by clinical examination or diagnostic imaging); or
- development of hypercalcaemia (corrected serum calcium greater than 2.65 mmol per L not attributable to any other cause).

Oligo-secretory and non-secretory patients are defined as having active disease with less than 10 g per L serum M protein.

The authority application must be made in writing and must include:

- a completed authority prescription form; and
- a completed Multiple Myeloma pomalidomide Authority Application Supporting Information form; and
- reports demonstrating the patient has failed treatment with, providing details of the contraindication to or details of the nature and severity of the intolerance to lenalidomide; and
- reports demonstrating the patient has failed treatment with, providing details of the contraindication to or details of the nature and severity of the intolerance to bortezomib.

Patients receiving this drug under the PBS listing must be registered in the i-access risk management program.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Prior Written Approval of Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Multiple myeloma

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously been issued with an authority prescription for this drug, **AND**
- Patient must not have progressive disease, **AND**

- The treatment must be in combination with dexamethasone, **AND**
- Patient must not be receiving concomitant PBS-subsidised bortezomib, carfilzomib or thalidomide or its analogues. Progressive disease is defined as at least 1 of the following:
 - (a) at least a 25% increase and an absolute increase of at least 5 g per L in serum M protein (monoclonal protein); or
 - (b) at least a 25% increase in 24-hour urinary light chain M protein excretion, and an absolute increase of at least 200 mg per 24 hours; or
 - (c) in oligo-secretory and non-secretory myeloma patients only, at least a 50% increase of the difference between involved free light chain and uninvolved free light chain; or
 - (d) at least a 25% relative increase and at least a 10% absolute increase in plasma cells in a bone marrow aspirate or on biopsy; or
 - (e) an increase in the size or number of lytic bone lesions (not including compression fractures); or
 - (f) at least a 25% increase in the size of an existing or the development of a new soft tissue plasmacytoma (determined by clinical examination or diagnostic imaging); or
 - (g) development of hypercalcaemia (corrected serum calcium greater than 2.65 mmol per L not attributable to any other cause).

Patients receiving this drug under the PBS listing must be registered in the i-access risk management program.

Note Authority applications for continuing treatment may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note Written applications for authority to prescribe should be forwarded to:

Department of Human Services
Prior Written Approval of Complex Drugs
Reply Paid 9826
HOBART TAS 7001

pomalidomide 3 mg capsule, 21

10417G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	10547.39	Pomalyst [CJ]

pomalidomide 4 mg capsule, 21

10386P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	10547.39	Pomalyst [CJ]

■ THALIDOMIDE

Caution Thalidomide is a category X drug and must not be given to pregnant women. Pregnancy in female patients or in the partners of male patients must be avoided during treatment and for 1 month after cessation of treatment.

Note Patients receiving thalidomide under the PBS listing must be registered in the i-access risk management program.

Authority required (STREAMLINED)

9290

Multiple myeloma

thalidomide 50 mg capsule, 28

6469L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	4	..	..	*1483.79	Thalomid [CJ]

thalidomide 100 mg capsule, 28

9684Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	..	..	*1483.79	Thalomid [CJ]

■ MUSCULO-SKELETAL SYSTEM

■ MUSCLE RELAXANTS

MUSCLE RELAXANTS, CENTRALLY ACTING AGENTS

Other centrally acting agents

■ BACLOFEN

Authority required (STREAMLINED)

9562

Severe chronic spasticity

Clinical criteria:

- Patient must have failed to respond to treatment with oral antispastic agents; OR
- Patient must have had unacceptable side effects to treatment with oral antispastic agents, **AND**
- Patient must have chronic spasticity of cerebral origin.

Authority required (STREAMLINED)

9525

Severe chronic spasticity

Clinical criteria:

- Patient must have failed to respond to treatment with oral antispastic agents; OR

- Patient must have had unacceptable side effects to treatment with oral antispastic agents, **AND**
- Patient must have chronic spasticity due to multiple sclerosis.

Authority required (STREAMLINED)
9638

Severe chronic spasticity

Clinical criteria:

- Patient must have failed to respond to treatment with oral antispastic agents; OR
- Patient must have had unacceptable side effects to treatment with oral antispastic agents, **AND**
- Patient must have chronic spasticity due to spinal cord injury.

Authority required (STREAMLINED)
9606

Severe chronic spasticity

Clinical criteria:

- Patient must have failed to respond to treatment with oral antispastic agents; OR
- Patient must have had unacceptable side effects to treatment with oral antispastic agents, **AND**
- Patient must have chronic spasticity due to spinal cord disease.

baclofen 40 mg/20 mL intrathecal injection, 20 mL ampoule

11194E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	..	..	*804.27	Sintetica Baclofen Intrathecal [BZ]

■ BACLOFEN

Note Pharmaceutical benefits that have the forms baclofen 10 mg/5 mL intrathecal injection, 5 mL ampoule, baclofen 10 mg/5 mL intrathecal injection, 5 x 5 mL ampoules and baclofen 10 mg/5 mL intrathecal injection, 10 x 5 mL ampoules are equivalent for the purposes of substitution.

Authority required (STREAMLINED)
9488

Severe chronic spasticity

Clinical criteria:

- Patient must have failed to respond to treatment with oral antispastic agents; OR
- Patient must have had unacceptable side effects to treatment with oral antispastic agents, **AND**
- Patient must have chronic spasticity of cerebral origin.

Authority required (STREAMLINED)
9637

Severe chronic spasticity

Clinical criteria:

- Patient must have failed to respond to treatment with oral antispastic agents; OR
- Patient must have had unacceptable side effects to treatment with oral antispastic agents, **AND**
- Patient must have chronic spasticity due to multiple sclerosis.

Authority required (STREAMLINED)
9489

Severe chronic spasticity

Clinical criteria:

- Patient must have failed to respond to treatment with oral antispastic agents; OR
- Patient must have had unacceptable side effects to treatment with oral antispastic agents, **AND**
- Patient must have chronic spasticity due to spinal cord injury.

Authority required (STREAMLINED)
9524

Severe chronic spasticity

Clinical criteria:

- Patient must have failed to respond to treatment with oral antispastic agents; OR
- Patient must have had unacceptable side effects to treatment with oral antispastic agents, **AND**
- Patient must have chronic spasticity due to spinal cord disease.

baclofen 10 mg/5 mL intrathecal injection, 5 x 5 mL ampoules

11420C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	..	..	*1003.71	^a Bacthecal [DZ]

baclofen 10 mg/5 mL intrathecal injection, 5 mL ampoule

6284R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	10	..	..	*1003.69	^a Bacthecal [DZ]	^a Lioresal Intrathecal [NV]

baclofen 10 mg/5 mL intrathecal injection, 10 x 5 mL ampoules

11128Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	1003.71	^a Sintetica Baclofen Intrathecal [BZ]

DRUGS FOR TREATMENT OF BONE DISEASES

DRUGS AFFECTING BONE STRUCTURE AND MINERALIZATION

Bisphosphonates

IBANDRONATE

Authority required (STREAMLINED)

9333

Bone metastases

Clinical criteria:

- The condition must be due to breast cancer.

ibandronate 6 mg/6 mL injection, 6 mL vial

9619G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	11	..	310.92	Bondronat [IX]

PAMIDRONATE DISODIUM

Authority required (STREAMLINED)

9234

Hypercalcaemia of malignancy

Clinical criteria:

- Patient must have a malignancy refractory to anti-neoplastic therapy.

pamidronate disodium 15 mg/5 mL injection, 5 mL vial

6286W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	4	2	..	*66.63	Pamisol [PF]

pamidronate disodium 60 mg/10 mL injection, 10 mL vial

6288Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	2	..	66.65	Pamisol [PF]

pamidronate disodium 30 mg/10 mL injection, 10 mL vial

6287X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	2	..	*66.65	Pamisol [PF]

PAMIDRONATE DISODIUM

Authority required (STREAMLINED)

9234

Hypercalcaemia of malignancy

Clinical criteria:

- Patient must have a malignancy refractory to anti-neoplastic therapy.

Authority required (STREAMLINED)

9335

Multiple myeloma

Authority required (STREAMLINED)

9315

Bone metastases

Clinical criteria:

- The condition must be due to breast cancer.

pamidronate disodium 90 mg/10 mL injection, 10 mL vial

6289B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	11	..	94.28	Pamisol [PF]

ZOLEDRONIC ACID

Note Pharmaceutical benefits that have the form zoledronic acid 4 mg/100 mL injection and pharmaceutical benefits that have the form zoledronic acid 4 mg/5 mL injection are equivalent for the purposes of substitution.

Authority required (STREAMLINED)

9268

Multiple myeloma

Authority required (STREAMLINED)

9328

Bone metastases

Clinical criteria:

- The condition must be due to breast cancer.

Authority required (STREAMLINED)

9304

Bone metastases

Clinical criteria:

- The condition must be due to castration-resistant prostate cancer.

Authority required (STREAMLINED)

9317

Hypercalcaemia of malignancy

Clinical criteria:

- Patient must have a malignancy refractory to anti-neoplastic therapy.

zoledronic acid 4 mg/5 mL injection, 5 mL vial

6371H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	11	..	145.05	^a APO-Zoledronic Acid [TX] ^a DEZTRON [DZ]	^a DBL Zoledronic Acid [PF] ^a Zometa [NV]

zoledronic acid 4 mg/100 mL injection, 100 mL bag

10542W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	11	..	145.05	^a DBL Zoledronic Acid [PF]

■ **OTHER DRUGS FOR DISORDERS OF THE MUSCULO-SKELETAL SYSTEM**

OTHER DRUGS FOR DISORDERS OF THE MUSCULO-SKELETAL SYSTEM

Other drugs for disorders of the musculo-skeletal system

■ **NUSINERSEN**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Spinal muscular atrophy (SMA)

Treatment Phase: Initial treatment - Loading doses

Treatment criteria:

- Must be treated by a specialist medical practitioner experienced in the diagnosis and management of SMA associated with a neuromuscular clinic of a recognised hospital in the management of SMA; or in consultation with a specialist medical practitioner experienced in the diagnosis and management of SMA associated with a neuromuscular clinic of a recognised hospital in the management of SMA.

Clinical criteria:

- The condition must 5q homozygous deletion, mutation of, or compound heterozygous mutation in the SMN1 gene of type I, II or IIIa, **AND**
- Patient must have experienced at least two of the defined signs and symptoms of SMA type I, II or IIIa prior to 3 years of age, **AND**
- The treatment must be given concomitantly with standard of care for this condition, **AND**
- The treatment must not exceed four loading doses (at days 0, 14, 28 and 63) under this restriction.

Population criteria:

- Patient must be 18 years of age or under.

Defined signs and symptoms of type I SMA are:

- Onset before 6 months of age; and
- Failure to meet or regression in ability to perform age-appropriate motor milestones; or
- Proximal weakness; or
- Hypotonia; or
- Absence of deep tendon reflexes; or
- Failure to gain weight appropriate for age; or
- Any active chronic neurogenic changes; or
- A compound muscle action potential below normative values for an age-matched child.

Defined signs and symptoms of type II SMA are:

- Onset between 6 and 18 months; and
- Failure to meet or regression in ability to perform age-appropriate motor milestones; or
- Proximal weakness; or
- Weakness in trunk righting/derotation; or
- Hypotonia; or

- vi) Absence of deep tendon reflexes; or
 - vii) Failure to gain weight appropriate for age; or
 - viii) Any active chronic neurogenic changes; or
 - ix) A compound muscle action potential below normative values for an age-matched child.
- Defined signs and symptoms of type IIIa SMA are:

- i) Onset between 18 months and 3 years of age; and
- ii) Failure to meet or regression in ability to perform age-appropriate motor milestones; or
- iii) Proximal weakness; or
- iv) Hypotonia; or
- v) Absence of deep tendon reflexes; or
- vi) Failure to gain weight appropriate for age; or
- vii) Any active chronic neurogenic changes; or
- viii) A compound muscle action potential below normative values for an age-matched child.

Recognised hospitals in the management of SMA are Lady Cilento Children's Hospital (Brisbane), Royal Children's Hospital Melbourne, Monash Children's Hospital (Melbourne), John Hunter Hospital (Newcastle), Sydney Children's Hospital Randwick, Children's Hospital at Westmead, Adelaide Women and Children's Hospital and Perth Children's Hospital.

Application for authorisation of initial treatment must be in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Spinal muscular atrophy PBS Authority Application - Supporting Information Form which includes the following:
 - i) specification of SMA type (I, II or IIIa); and
 - ii) sign(s) and symptom(s) that the patient has experienced; and
 - iii) patient's age at the onset of sign(s) and symptom(s).

nusinersen 12 mg/5 mL injection, 5 mL vial

11472T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	3	..	110047.39	Spinraza [BD]

■ NUSINERSEN

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Authority applications for continuing treatment may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Spinal muscular atrophy (SMA)

Treatment Phase: Continuing treatment - Maintenance

Treatment criteria:

- Must be treated by a specialist medical practitioner experienced in the diagnosis and management of SMA associated with a neuromuscular clinic of a recognised hospital in the management of SMA; or in consultation with a specialist medical practitioner experienced in the diagnosis and management of SMA associated with a neuromuscular clinic of a recognised hospital in the management of SMA.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be given concomitantly with standard of care for this condition, **AND**
- The treatment must be ceased when invasive permanent assisted ventilation is required in the absence of a potentially reversible cause while being treated with this drug.

Recognised hospitals in the management of SMA are Lady Cilento Children's Hospital (Brisbane), Royal Children's Hospital Melbourne, Monash Children's Hospital (Melbourne), John Hunter Hospital (Newcastle), Sydney Children's Hospital Randwick, Children's Hospital at Westmead, Adelaide Women and Children's Hospital and Perth Children's Hospital.

Invasive permanent assisted ventilation means ventilation via tracheostomy tube for greater than or equal to 16 hours per day.

nusinersen 12 mg/5 mL injection, 5 mL vial

11476B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	110047.39	Spinraza [BD]

■ NUSINERSEN

Note Special Pricing Arrangements apply.

Note No increase in the maximum quantity or number of units may be authorised.

Note A maximum number of repeats of up to 2 may be authorised for patients requiring loading doses for days 14, 28 and 63.

Note A maximum number of repeats of up to 1 may be authorised for patients requiring loading doses for days 28 and 63.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Spinal muscular atrophy (SMA)

Treatment Phase: Grandfather patients

Treatment criteria:

- Must be treated by a specialist medical practitioner experienced in the diagnosis and management of SMA associated with a neuromuscular clinic of a recognised hospital in the management of SMA; or in consultation with a specialist medical practitioner experienced in the diagnosis and management of SMA associated with a neuromuscular clinic of a recognised hospital in the management of SMA.

Clinical criteria:

- Patient must have previously received non-PBS-subsidised treatment for this condition with this drug prior to 1 June 2018, **AND**

- The condition must 5q homozygous deletion, mutation of, or compound heterozygous mutation in the SMN1 gene of type I, II or IIIa, **AND**
- Patient must have had experienced at least two of the defined signs and symptoms of SMA type I, II or IIIa prior to 3 years of age, **AND**
- Patient must have previously received at least one of the four loading doses at days 0, 14, 28 and 63, **AND**
- The treatment must be given concomitantly with standard of care for this condition, **AND**
- The treatment must be ceased when invasive permanent assisted ventilation is required in the absence of a potentially reversible cause while being treated with this drug.

Population criteria:

- Patient must have been 18 years of age or under at the time treatment with this drug was initiated for this condition; OR
- Patient must have previously received treatment with this drug for this condition under the care of clinicians with the authorised prescriber number of AP17/83146.

Defined signs and symptoms of type I SMA are:

- i) Onset before 6 months of age; and
- ii) Failure to meet or regression in ability to perform age-appropriate motor milestones; or
- iii) Proximal weakness; or
- iv) Hypotonia; or
- v) Absence of deep tendon reflexes; or
- vi) Failure to gain weight appropriate for age; or
- vii) Any active chronic neurogenic changes; or
- viii) A compound muscle action potential below normative values for an age-matched child.

Defined signs and symptoms of type II SMA are:

- i) Onset between 6 and 18 months; and
- ii) Failure to meet or regression in ability to perform age-appropriate motor milestones; or
- iii) Proximal weakness; or
- iv) Weakness in trunk righting/derotation; or
- v) Hypotonia; or
- vi) Absence of deep tendon reflexes; or
- vii) Failure to gain weight appropriate for age; or
- viii) Any active chronic neurogenic changes; or
- ix) A compound muscle action potential below normative values for an age-matched child.

Defined signs and symptoms of type IIIa SMA are:

- i) Onset between 18 months and 3 years of age; and
- ii) Failure to meet or regression in ability to perform age-appropriate motor milestones; or
- iii) Proximal weakness; or
- iv) Hypotonia; or
- v) Absence of deep tendon reflexes; or
- vi) Failure to gain weight appropriate for age; or
- vii) Any active chronic neurogenic changes; or
- viii) A compound muscle action potential below normative values for an age-matched child.

Invasive permanent assisted ventilation means ventilation via tracheostomy tube for greater than or equal to 16 hours per day.

Recognised hospitals in the management of SMA are Lady Cilento Children's Hospital (Brisbane), Royal Children's Hospital Melbourne, Monash Children's Hospital (Melbourne), John Hunter Hospital (Newcastle), Sydney Children's Hospital Randwick, Children's Hospital at Westmead, Adelaide Women and Children's Hospital and Perth Children's Hospital.

Application for authorisation of grandfathering treatment must be in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Spinal muscular atrophy PBS Authority Application for Grandfather patients - Supporting Information Form which includes the following:
 - (i) specification of SMA type (I, II or IIIa); and
 - (ii) sign(s) and symptom(s) that the patient has experienced; and
 - (iii) patient's age at the onset of sign(s) and symptom(s); and

(iv) if relevant, a copy of a TGA-approval letter to clinician with the authorised prescriber number of AP17/83146.

A patient may qualify for PBS-subsidised treatment under this restriction once only.

For continuing PBS-subsidised treatment, a Grandfathered patient must qualify under the Continuing treatment criteria.

nusinersen 12 mg/5 mL injection, 5 mL vial

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
11470Q	1	..	..	110047.39	Spinraza [BD]

■ NERVOUS SYSTEM

■ ANTI-PARKINSON DRUGS

DOPAMINERGIC AGENTS

Dopa and dopa derivatives

■ LEVODOPA + CARBIDOPA

Note Special Pricing Arrangements apply.

Note Patients should have adequate cognitive function to manage administration with a portable continuous infusion pump.

Authority required (STREAMLINED)

9405

Advanced Parkinson disease

Clinical criteria:

- Patient must have severe disabling motor fluctuations not adequately controlled by oral therapy, **AND**
- The treatment must be commenced in a hospital-based movement disorder clinic.

levodopa 20 mg/mL + carbidopa monohydrate 5 mg/mL intestinal gel, 7 x 100 mL

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
9744W	8	5	..	*11583.39	Duodopa [VE]

Dopamine agonists

■ APOMORPHINE

Authority required (STREAMLINED)

9561

Parkinson disease

Clinical criteria:

- Patient must have experienced severely disabling motor fluctuations which have not responded to other therapy.

apomorphine hydrochloride hemihydrate 20 mg/2 mL injection, 5 x 2 mL ampoules

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
9607P	72	5	..	*6576.67	Movapo [TD]

apomorphine hydrochloride hemihydrate 50 mg/5 mL injection, 5 x 5 mL ampoules

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
9640J	36	5	..	*8215.39	Movapo [TD]

apomorphine hydrochloride hemihydrate 100 mg/20 mL injection, 5 x 20 mL vials

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
11083H	18	5	..	*7600.15	Apomine Solution for Infusion [PF]

apomorphine hydrochloride hemihydrate 50 mg/10 mL injection, 5 x 10 mL syringes

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
10971K	36	5	..	*8215.39	Movapo PFS [TD]

■ APOMORPHINE

Note No increase in the maximum quantity or number of units may be authorised.

Note Pharmaceutical benefits that have the form apomorphine injection 30 mg/3 mL pen device and pharmaceutical benefits that have the form apomorphine injection 30 mg/3 mL cartridge are equivalent for the purposes of substitution.

Authority required (STREAMLINED)

9561

Parkinson disease

Clinical criteria:

- Patient must have experienced severely disabling motor fluctuations which have not responded to other therapy.

apomorphine hydrochloride hemihydrate 30 mg/3 mL injection, 5 x 3 mL cartridges

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
11688E	20	5	..	*2767.79	^a Apomine Intermittent [PF]

apomorphine hydrochloride hemihydrate 30 mg/3 mL injection, 5 x 3 mL pen devices

11475Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	20	5	..	*2767.79	^a Movapo Pen [TD]

■ **PSYCHOLEPTICS****ANTIPSYCHOTICS***Diazepines, oxazepines, thiazepines and oxepines*■ **CLOZAPINE**

Note Patients receiving clozapine under the PBS listing must be registered in the clozapine patient monitoring program relevant for the brand of clozapine being prescribed and dispensed: Pfizer ClopineCentral.

Authority required (STREAMLINED)**9490**

Schizophrenia

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a psychiatrist or in consultation with the psychiatrist affiliated with the hospital or specialised unit managing the patient.

Clinical criteria:

- Patient must be non-responsive to other neuroleptic agents; OR
- Patient must be intolerant of other neuroleptic agents.

Patients must complete at least 18 weeks of initial treatment under this restriction before being able to qualify for treatment under the continuing restriction.

The name of the consulting psychiatrist should be included in the patient's medical records.

A medical practitioner should request a quantity sufficient for up to one month's supply. Up to 5 repeats will be authorised.

clozapine 50 mg/mL oral liquid, 100 mL

11415T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	±1	..	..	147.79	Versacloz [PF]

■ **CLOZAPINE**

Note Patients receiving clozapine under the PBS listing must be registered in the clozapine patient monitoring program relevant for the brand of clozapine being prescribed and dispensed: Novartis Clozaril Patient Monitoring System (eCPMS) or Hospira Clopineconnect.

Authority required (STREAMLINED)**9490**

Schizophrenia

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a psychiatrist or in consultation with the psychiatrist affiliated with the hospital or specialised unit managing the patient.

Clinical criteria:

- Patient must be non-responsive to other neuroleptic agents; OR
- Patient must be intolerant of other neuroleptic agents.

Patients must complete at least 18 weeks of initial treatment under this restriction before being able to qualify for treatment under the continuing restriction.

The name of the consulting psychiatrist should be included in the patient's medical records.

A medical practitioner should request a quantity sufficient for up to one month's supply. Up to 5 repeats will be authorised.

clozapine 25 mg tablet, 100

6101D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	..	..	*76.03	Clopine 25 [PF]	Clozaril 25 [GO]

clozapine 100 mg tablet, 100

6102E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	..	..	*259.47	Clopine 100 [PF]	Clozaril 100 [GO]

clozapine 50 mg tablet, 100

6417R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	..	..	*141.85	Clopine 50 [PF]

clozapine 200 mg tablet, 100

6418T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	..	..	*511.55	Clopine 200 [PF]

clozapine 50 mg/mL oral liquid, 100 mL

9632Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	147.79	Clopine Suspension [PF]

■ **RESPIRATORY SYSTEM**■ **DRUGS FOR OBSTRUCTIVE AIRWAY DISEASES****OTHER SYSTEMIC DRUGS FOR OBSTRUCTIVE AIRWAY DISEASES***Other systemic drugs for obstructive airway diseases*■ **BENRALIZUMAB****Note TREATMENT OF ADULT AND ADOLESCENT PATIENTS WITH UNCONTROLLED SEVERE EOSINOPHILIC ASTHMA**

Patients are eligible to commence a 'benralizumab treatment cycle' (initial treatment course with or without continuing treatment course/s) if they satisfy the eligibility criteria as detailed under the initial treatment restriction.

Once a patient has either failed to achieve or maintain a response to benralizumab, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 6 month break in PBS-subsidised benralizumab therapy before they are eligible to commence the next 'benralizumab treatment cycle', or if eligible, a 'mepolizumab treatment cycle' or an 'omalizumab treatment cycle'. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised benralizumab is stopped to the date of the first application for initial treatment with benralizumab, mepolizumab or omalizumab under the new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised benralizumab therapy:

(a) Initial treatment:

Applications for initial treatment should be made where:

- i) A patient has received no prior PBS-subsidised benralizumab treatment and wishes to commence such therapy; or
- ii) A patient wishes to recommence treatment with benralizumab following a break in PBS-subsidised therapy of more than 6 months; or
- iii) A patient has received prior PBS-subsidised mepolizumab or omalizumab and wishes to commence treatment with benralizumab after a treatment break of 6 months.

All applications for initial treatment for non-grandfather patients will be limited to provide for a maximum of up to 32 weeks of therapy for benralizumab.

(b) Grandfather patients:

For patients who commenced treatment with non-PBS subsidised benralizumab for uncontrolled severe eosinophilic asthma prior to 1 December 2018 and who continue to receive treatment at the time of application, may qualify for treatment under the initial 'grandfather' treatment restriction. A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment with benralizumab will be authorised under this criterion. Approval will be based on the criteria included in the relevant restriction. Following completion of the Initial PBS-subsidised course, further applications for treatment with benralizumab will be assessed under the continuing treatment restriction.

'Grandfather' arrangements will only apply for the first treatment cycle (initial treatment course with or without continuing treatment course/s). If a 'Grandfathered' patient recommences on second and subsequent cycles after a treatment break, the 'Grandfathered' patient must re-qualify for Initial treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 6 month break in PBS-subsidised therapy' below for further details.

(c) Continuing treatment:

Following the completion of the initial treatment course with benralizumab, a patient may qualify to receive up to a further 24 weeks of continuing treatment with benralizumab providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing benralizumab treatment in courses of up to 24 weeks providing they continue to sustain the response.

(2) Baseline measurements to determine response:

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the Asthma Control Questionnaire (ACQ; 5 item version) or oral corticosteroid dose, submitted with the Initial authority application for benralizumab. However, prescribers may provide new baseline measurements when a new Initial treatment authority application is submitted and the Department of Human Services will assess response according to these revised baseline measurements.

(3) Re-commencement of treatment after a 6 month break in PBS-subsidised therapy:

A patient who wishes to trial a second or subsequent benralizumab treatment cycle, or an initial mepolizumab or omalizumab treatment cycle, following a break in PBS-subsidised therapy of at least 6 months, must re-qualify for initial treatment with respect to the indices of disease severity (oral corticosteroid dose, Asthma Control Questionnaire (ACQ-5) score, and relevant exacerbation history). Patients must have received optimised standard therapy, at adequate doses and for the minimum period specified, immediately prior to the time the new baseline assessments are performed.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Uncontrolled severe eosinophilic asthma

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a respiratory physician, clinical immunologist, allergist or general physician experienced in the management of patients with severe asthma.

Clinical criteria:

- Patient must be under the care of the same physician for at least 6 months; OR

- Patient must have been diagnosed by a multidisciplinary severe asthma clinic team, **AND**
- Patient must have a diagnosis of asthma confirmed and documented by a respiratory physician, clinical immunologist, allergist or general physician experienced in the management of patients with severe asthma, defined by the following standard clinical features: (i) forced expiratory volume (FEV1) reversibility greater than or equal to 12% and greater than or equal to 200 mL at baseline within 30 minutes after administration of salbutamol (200 to 400 micrograms), or (ii) airway hyperresponsiveness defined as a greater than 20% decline in FEV1 during a direct bronchial provocation test or greater than 15% decline during an indirect bronchial provocation test, or (iii) peak expiratory flow (PEF) variability of greater than 15% between the two highest and two lowest peak expiratory flow rates during 14 days, **AND**
- Patient must have a duration of asthma of at least 1 year, **AND**
- Patient must have forced expiratory volume (FEV1) less than or equal to 80% predicted, documented on 1 or more occasions in the previous 12 months, **AND**
- Patient must have blood eosinophil count greater than or equal to 300 cells per microlitre in the last 12 months, **AND**
- Patient must have failed to achieve adequate control with optimised asthma therapy, despite formal assessment of and adherence to correct inhaler technique, which has been documented, **AND**
- The treatment must not be used in combination with, or within 6 months of treatment with, PBS-subsidised omalizumab or mepolizumab.

Population criteria:

- Patient must be aged 12 years or older.

Optimised asthma therapy includes:

- (i) Adherence to maximal inhaled therapy, including high dose inhaled corticosteroid (ICS) plus long-acting beta-2 agonist (LABA) therapy for at least 12 months, unless contraindicated or not tolerated; **AND**
- (ii) treatment with oral corticosteroids, either daily oral corticosteroids for at least 6 weeks, OR a cumulative dose of oral corticosteroids of at least 500 mg prednisolone equivalent in the previous 12 months, unless contraindicated or not tolerated.

If the requirement for treatment with optimised asthma therapy cannot be met because of contraindications according to the relevant TGA-approved Product Information and/or intolerances of a severity necessitating permanent treatment withdrawal, details of the contraindication and/or intolerance must be provided in the Authority application.

The following initiation criteria indicate failure to achieve adequate control and must be demonstrated in all patients at the time of the application:

- (a) an Asthma Control Questionnaire (ACQ-5) score of at least 2.0, as assessed in the previous month, **AND**
- (b) while receiving optimised asthma therapy in the past 12 months, experienced at least 1 admission to hospital for a severe asthma exacerbation, OR 1 severe asthma exacerbation, requiring documented use of systemic corticosteroids (oral corticosteroids initiated or increased for at least 3 days, or parenteral corticosteroids) prescribed/supervised by a physician.

The Asthma Control Questionnaire (5 item version) assessment of the patient must be made at time of application for treatment (to establish baseline score) and again around 20 weeks after the first PBS-subsidised dose of this drug under this restriction so that there is adequate time for a response to be demonstrated and for the application for the first continuing therapy to be processed.

This assessment at around 24 weeks, which will be used to determine eligibility for the first continuing treatment, must be submitted no later than 2 weeks prior to the patient completing their current treatment course, to avoid an interruption to supply. Where a response assessment is not undertaken and submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

A patient who fails to respond to a course of PBS-subsidised benralizumab for the treatment of uncontrolled severe eosinophilic asthma will not be eligible to receive further PBS-subsidised treatment with benralizumab, mepolizumab or omalizumab within 6 months of the date on which treatment was ceased.

A multidisciplinary severe asthma clinic team comprises of:

- A respiratory physician; and
- A pharmacist, nurse or asthma educator.

At the time of the authority application, medical practitioners should request up to 4 repeats to provide for an initial course of benralizumab sufficient for up to 32 weeks of therapy, at a dose of 30 mg every 4 weeks for the first three doses (weeks 0, 4, and 8) then 30 mg every eight weeks thereafter.

Benralizumab must not be used concurrently with omalizumab or mepolizumab or within 6 months of each other. A patient is required to have ceased treatment with omalizumab or mepolizumab for 6 months prior to initiating treatment with benralizumab.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Severe Eosinophilic Asthma Initial PBS Authority Application - Supporting Information Form, which includes the following:
 - (i) details of prior optimised asthma drug therapy (date of commencement and duration of therapy); and
 - (ii) details of severe exacerbation/s experienced in the past 12 months while receiving optimised asthma therapy (date and treatment); and
- (c) a copy of the eosinophil pathology report; and
- (d) a completed Asthma Control Questionnaire (ACQ-5) calculation sheet including the date of assessment of the patient's symptoms.

Note The Department of Human Services website (www.humanservices.gov.au) has details of the accepted toxicities, including severity, which will be accepted for the purposes of exempting a patient from the requirement of treatment with optimised asthma therapy.

Note For copies of the ACQ, please contact AstraZeneca Medical Information on 1800 805 342.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available

on the Department of Human Services website at www.humanservices.gov.au
Applications for authority to prescribe should be forwarded to:
Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Formal assessment and correction of inhaler technique should be performed in accordance with the National Asthma Council (NAC) Information Paper for Health Professionals on Inhaler Technique (available at www.humanservices.gov.au or www.nationalasthma.org.au); the assessment and adherence to correct technique should be documented in the patient's medical records. Patients can obtain support with inhaler technique through their local Asthma Foundation (1800 645 130).

Authority required

Uncontrolled severe eosinophilic asthma

Treatment Phase: Initial treatment - balance of supply

Treatment criteria:

- Must be treated by a respiratory physician, clinical immunologist, allergist or general physician experienced in the management of patients with severe asthma.

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the Initial treatment restriction to complete 32 weeks treatment; AND
- The treatment must provide no more than the balance of up to 32 weeks treatment available under the Initial restriction.

Population criteria:

- Patient must be aged 12 years or older.

Note Authority approval for sufficient therapy to complete a maximum of 32 weeks of treatment under the initial restriction may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

benralizumab 30 mg/mL injection, 1 mL syringe

11523L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	4	..	3358.39	Fasenra [AP]

■ BENRALIZUMAB

Note TREATMENT OF ADULT AND ADOLESCENT PATIENTS WITH UNCONTROLLED SEVERE EOSINOPHILIC ASTHMA

Patients are eligible to commence a 'benralizumab treatment cycle' (initial treatment course with or without continuing treatment course/s) if they satisfy the eligibility criteria as detailed under the initial treatment restriction.

Once a patient has either failed to achieve or maintain a response to benralizumab, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 6 month break in PBS-subsidised benralizumab therapy before they are eligible to commence the next 'benralizumab treatment cycle', or if eligible, a 'mepolizumab treatment cycle' or an 'omalizumab treatment cycle'. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised benralizumab is stopped to the date of the first application for initial treatment with benralizumab, mepolizumab or omalizumab under the new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised benralizumab therapy:

(a) Initial treatment:

Applications for initial treatment should be made where:

- A patient has received no prior PBS-subsidised benralizumab treatment and wishes to commence such therapy; or
- A patient wishes to recommence treatment with benralizumab following a break in PBS-subsidised therapy of more than 6 months; or
- A patient has received prior PBS-subsidised mepolizumab or omalizumab and wishes to commence treatment with benralizumab after a treatment break of 6 months.

All applications for initial treatment for non-grandfather patients will be limited to provide for a maximum of up to 32 weeks of therapy for benralizumab.

(b) Grandfather patients:

For patients who commenced treatment with non-PBS subsidised benralizumab for uncontrolled severe eosinophilic asthma prior to 1 December 2018 and who continue to receive treatment at the time of application, may qualify for treatment under the initial 'grandfather' treatment restriction. A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment with benralizumab will be authorised under this criterion. Approval will be based on the criteria included in the relevant restriction. Following completion of the Initial PBS-subsidised course, further applications for treatment with benralizumab will be assessed under the continuing treatment restriction.

'Grandfather' arrangements will only apply for the first treatment cycle (initial treatment course with or without continuing treatment course/s). If a 'Grandfathered' patient recommences on second and subsequent cycles after a treatment break, the 'Grandfathered' patient must re-qualify for Initial treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 6 month break in PBS-subsidised therapy' below for further details.

(c) Continuing treatment:

Following the completion of the initial treatment course with benralizumab, a patient may qualify to receive up to a further 24 weeks of continuing treatment with benralizumab providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing benralizumab treatment in courses of up to 24 weeks providing they continue to sustain the response.

(2) Baseline measurements to determine response:

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the Asthma Control Questionnaire (ACQ; 5 item version) or oral corticosteroid dose, submitted with the Initial authority application for benralizumab. However, prescribers may provide new baseline measurements when a new Initial treatment authority application is submitted and the Department of Human Services will assess response

according to these revised baseline measurements.

(3) Re-commencement of treatment after a 6 month break in PBS-subsidised therapy:

A patient who wishes to trial a second or subsequent benralizumab treatment cycle, or an initial mepolizumab or omalizumab treatment cycle, following a break in PBS-subsidised therapy of at least 6 months, must re-qualify for initial treatment with respect to the indices of disease severity (oral corticosteroid dose, Asthma Control Questionnaire (ACQ-5) score, and relevant exacerbation history). Patients must have received optimised standard therapy, at adequate doses and for the minimum period specified, immediately prior to the time the new baseline assessments are performed.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Uncontrolled severe eosinophilic asthma

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a respiratory physician, clinical immunologist, allergist or general physician experienced in the management of patients with severe asthma.

Clinical criteria:

- Patient must have demonstrated or sustained an adequate response to PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must not be used in combination with, or within 6 months of treatment with, PBS-subsidised omalizumab or mepolizumab.

Population criteria:

- Patient must be aged 12 years or older.

An adequate response to benralizumab treatment is defined as:

(a) a reduction in the Asthma Control Questionnaire (ACQ-5) score of at least 0.5 from baseline; OR

(b) maintenance oral corticosteroid dose reduced by at least 25% from baseline, and no deterioration in ACQ-5 score from baseline.

All applications for second and subsequent continuing treatments with this drug must include a measurement of response to the prior course of therapy. The Asthma Control Questionnaire (5 item version) assessment of the patient's response to the prior course of treatment, or the assessment of oral corticosteroid dose, should be made at around 16 weeks after the first PBS-subsidised dose of this drug under this restriction so that there is adequate time for a response to be demonstrated and for the application for continuing therapy to be processed.

The assessment should, where possible, be completed by the same physician who initiated treatment with this drug. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted no later than 2 weeks prior to the patient completing their current treatment course, to avoid an interruption to supply. Where a response assessment is not undertaken and submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

A patient who fails to respond to a course of PBS-subsidised benralizumab for the treatment of uncontrolled severe eosinophilic asthma will not be eligible to receive further PBS-subsidised treatment with benralizumab for this condition within 6 months of the date on which treatment was ceased.

At the time of the authority application, medical practitioners should request the appropriate number of repeats to provide for a continuing course of this drug sufficient for up to 24 weeks of therapy.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Severe Eosinophilic Asthma Continuing PBS Authority Application - Supporting Information Form which includes details of maintenance oral corticosteroid dose; or a completed Asthma Control Questionnaire (ACQ-5) calculation sheet including the date of assessment of the patient's symptoms.

Note If the same physician cannot assess the patient please call the Department of Human Services on 1800 700 270.

Note For copies of the ACQ, please contact AstraZeneca Medical Information on 1800 805 342.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Uncontrolled severe eosinophilic asthma

Treatment Phase: Continuing treatment or Grandfathered treatment - balance of supply

Treatment criteria:

- Must be treated by a respiratory physician, clinical immunologist, allergist or general physician experienced in the management of patients with severe asthma.

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the continuing treatment restriction or the grandfather restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the continuing treatment restriction or the grandfather restriction.

Population criteria:

- Patient must be aged 12 years or older.

Note Authority approval for sufficient therapy to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Uncontrolled severe eosinophilic asthma

Treatment Phase: Grandfathered treatment

Treatment criteria:

- Must be treated by a respiratory physician, clinical immunologist, allergist or general physician experienced in the management of patients with severe asthma.

Clinical criteria:

- Patient must have received non-PBS subsidised treatment with this drug for this condition prior to 1 December 2018, **AND**
- Patient must be receiving treatment with this drug for this condition at the time of application, **AND**
- Patient must be under the care of the same physician for at least 6 months; **OR**
- Patient must have been diagnosed by a multidisciplinary severe asthma clinic team, **AND**
- Patient must have had, prior to commencement of non-PBS subsidised treatment with this drug, a diagnosis of asthma confirmed and documented by a respiratory physician, clinical immunologist, allergist or general physician experienced in the management of patients with severe asthma, defined by the following standard clinical features: (i) forced expiratory volume (FEV1) reversibility greater than or equal to 12% and greater than or equal to 200 mL at baseline within 30 minutes after administration of salbutamol (200 to 400 micrograms), or (ii) airway hyperresponsiveness defined as a greater than 20% decline in FEV1 during a direct bronchial provocation test or greater than 15% decline during an indirect bronchial provocation test, or (iii) peak expiratory flow (PEF) variability of greater than 15% between the two highest and two lowest peak expiratory flow rates during 14 days, **AND**
- Patient must have had blood eosinophil count greater than or equal to 300 cells per microlitre prior to commencement of non-PBS subsidised treatment with this drug, **AND**
- Patient must have had a duration of asthma of at least 1 year prior to commencement of non-PBS subsidised treatment with this drug, **AND**
- Patient must have failed to achieve adequate control with optimised asthma therapy prior to non-PBS subsidised treatment with this drug despite formal assessment of and adherence to correct inhaler technique, which has been documented, **AND**
- Patient must have demonstrated an adequate response if the patient has received at least 24 weeks of treatment of non-PBS subsidised benralizumab for this condition, **AND**
- The treatment must not be used in combination with, or within 6 months of treatment with, PBS-subsidised omalizumab or mepolizumab.

Population criteria:

- Patient must be aged 12 years or older.

Optimised asthma therapy includes:

(i) Adherence to maximal inhaled therapy, including high dose inhaled corticosteroid (ICS) plus long-acting beta-2 agonist (LABA) therapy for at least 12 months, unless contraindicated or not tolerated; **AND**

(ii) treatment with oral corticosteroids, either daily oral corticosteroids for at least 6 weeks, **OR** a cumulative dose of oral corticosteroids of at least 500 mg prednisolone equivalent in the previous 12 months, prior to commencing non-PBS subsidised treatment with this drug, unless contraindicated or not tolerated.

If the requirement for treatment with optimised asthma therapy cannot be met because of contraindications according to the relevant TGA-approved Product Information and/or intolerances of a severity necessitating permanent treatment withdrawal, details of the contraindication and/or intolerance must be provided in the Authority application.

The following initiation criteria indicate failure to achieve adequate control and must be demonstrated in all patients at the time of the application:

(a) an Asthma Control Questionnaire (ACQ-5) score of at least 2.0, as assessed in the previous month, **AND**

(b) while receiving optimised asthma therapy in the past 12 months, experienced at least 1 admission to hospital for a severe asthma exacerbation, **OR** 1 severe asthma exacerbation, requiring documented use of systemic corticosteroids (oral corticosteroids initiated or increased for at least 3 days, or parenteral corticosteroids) prescribed/supervised by a physician.

The Asthma Control Questionnaire (5 item version) assessment of the patient must be made at time of application for treatment (to establish baseline score) and again around 20 weeks after the first PBS-subsidised dose of this drug under this restriction so that there is adequate time for a response to be demonstrated and for the application for the first continuing therapy to be processed.

This assessment at around 24 weeks, which will be used to determine eligibility for the first continuing treatment, must be submitted no later than 2 weeks prior to the patient completing their current treatment course, to avoid an interruption to supply. Where a response assessment is not undertaken and submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

An adequate response to benralizumab treatment is defined as:

(a) a reduction in the Asthma Control Questionnaire (ACQ 5) score of at least 0.5 from baseline; **OR**

(b) maintenance oral corticosteroid dose reduced by at least 25% from baseline, and no deterioration in ACQ 5 score from baseline.

A multidisciplinary severe asthma clinic team comprises of:

- A respiratory physician; and
- A pharmacist, nurse or asthma educator.

A review of the patient's records should be conducted to extract pre- and post-benralizumab data on symptoms, quality of life, medication doses, exacerbations and hospitalisations. Parameters to establish response are:

- (i) a reduction in Asthma Control Questionnaire (ACQ-5) score of at least 0.5; and/or
- (ii) maintenance oral corticosteroid dose reduced by at least 25% from baseline and no deterioration in ACQ 5 score from baseline.

The assessment of the patient's response to the initial PBS-subsidised course of treatment under this restriction must be made at around 16 weeks after the first dose of PBS-subsidised treatment with this drug under this restriction so that there is adequate time for a response to be demonstrated and for the application for continuing therapy to be processed. The same parameters used to establish response to non-PBS subsidised therapy with this drug should be used for the assessment.

This assessment, which will be used to determine eligibility for continuing treatment, must be submitted no later than 2 weeks prior to the patient completing their current treatment course, to avoid an interruption to supply. Where a response assessment is not undertaken and submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

Patients will be eligible to receive continuing courses of treatment with this drug of up to 24 weeks providing they continue to demonstrate an adequate response to treatment.

A patient may qualify for PBS-subsidised treatment under this restriction once only.

A patient who fails to respond to a course of PBS-subsidised benralizumab for the treatment of uncontrolled severe eosinophilic asthma will not be eligible to receive further PBS-subsidised treatment with benralizumab, omalizumab or mepolizumab within 6 months of the date on which treatment was ceased.

At the time of the authority application, medical practitioners should request the appropriate maximum quantity and number of repeats to provide for a continuing course of benralizumab sufficient for up to 24 weeks of therapy.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Severe Eosinophilic Asthma Grandfather PBS Authority Application - Supporting Information Form, which includes the following:
 - (i) details of prior optimised asthma drug therapy (date of commencement and duration of therapy); and
 - (ii) details of pre- and post-benralizumab data on symptoms, quality of life, medication doses, severe exacerbation/s and hospitalisations, and
 - (c) a copy of the pre-benralizumab eosinophil pathology report; and
 - (d) a completed Asthma Control Questionnaire (ACQ 5) calculation sheet including the date of assessment of the patient's symptoms; or details of maintenance oral corticosteroid dose.

Note The Department of Human Services website (www.humanservices.gov.au) has details of the accepted toxicities, including severity, which will be accepted for the purposes of exempting a patient from the requirement of treatment with optimised asthma therapy.

Note For copies of the ACQ, please contact AstraZeneca Medical Information on 1800 805 342.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Formal assessment and correction of inhaler technique should be performed in accordance with the National Asthma Council (NAC) Information Paper for Health Professionals on Inhaler Technique (available at www.humanservices.gov.au or www.nationalasthma.org.au); the assessment and adherence to correct technique should be documented in the patient's medical records. Patients can obtain support with inhaler technique through their local Asthma Foundation (1800 645 130).

benralizumab 30 mg/mL injection, 1 mL syringe

11504L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	2	..	3358.39	Fasenra [AP]

■ MEPOLIZUMAB

Note TREATMENT OF ADULT AND ADOLESCENT PATIENTS WITH UNCONTROLLED SEVERE EOSINOPHILIC ASTHMA

Patients are eligible to commence a 'mepolizumab treatment cycle' (initial treatment course with or without continuing treatment course/s) if they satisfy the eligibility criteria as detailed under the initial treatment restriction.

Once a patient has either failed to achieve or maintain a response to mepolizumab, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 6 month break in PBS-subsidised mepolizumab therapy before they are eligible to commence the next mepolizumab treatment cycle, or if eligible, a 'benralizumab treatment cycle' or an 'omalizumab treatment cycle'. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised mepolizumab is stopped to the date of the first application for initial treatment with benralizumab, mepolizumab or omalizumab under the new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised mepolizumab therapy:

(a) Initial treatment:

Applications for initial treatment should be made where:

- i) A patient has received no prior PBS-subsidised mepolizumab treatment and wishes to commence such therapy; or
- ii) A patient wishes to recommence treatment with mepolizumab following a break in PBS-subsidised therapy of more than 6 months; or

iii) A patient has received prior PBS-subsidised benralizumab or omalizumab and wishes to commence treatment with mepolizumab after a treatment break of 6 months.

All applications for initial treatment will be limited to provide for a maximum of up to 32 weeks of therapy for mepolizumab.

(b) Continuing treatment:

Following the completion of the initial treatment course with mepolizumab, a patient may qualify to receive up to a further 24 weeks of continuing treatment with mepolizumab providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing mepolizumab treatment in courses of up to 24 weeks providing they continue to sustain the response.

(2) Baseline measurements to determine response:

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the Asthma Control Questionnaire (ACQ; 5 item version) or oral corticosteroid dose, submitted with the Initial authority application for mepolizumab. However, prescribers may provide new baseline measurements when a new Initial treatment authority application is submitted and the Department of Human Services will assess response according to these revised baseline measurements.

(3) Re-commencement of treatment after a 6 month break in PBS-subsidised therapy:

A patient who wishes to trial a second or subsequent mepolizumab treatment cycle, or an initial benralizumab or omalizumab treatment cycle, following a break in PBS-subsidised therapy of at least 6 months, must re-qualify for initial treatment with respect to the indices of disease severity (oral corticosteroid dose, Asthma Control Questionnaire (ACQ-5) score, and relevant exacerbation history). Patients must have received optimised standard therapy, at adequate doses and for the minimum period specified, immediately prior to the time the new baseline assessments are performed.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note If the same physician cannot assess the patient please call the Department of Human Services on 1800 700 270.

Note For copies of the ACQ, please contact GlaxoSmithKline Medical Information on 1800 033 109.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Uncontrolled severe eosinophilic asthma

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a respiratory physician, clinical immunologist, allergist or general physician experienced in the management of patients with severe asthma.

Clinical criteria:

- Patient must have demonstrated or sustained an adequate response to PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must not be used in combination with, or within 6 months of treatment with, PBS-subsidised benralizumab or omalizumab.

Population criteria:

- Patient must be aged 12 years or older.

An adequate response to mepolizumab treatment is defined as:

(a) a reduction in the Asthma Control Questionnaire (ACQ-5) score of at least 0.5 from baseline,

OR

(b) maintenance oral corticosteroid dose reduced by at least 25% from baseline, and no deterioration in ACQ-5 score from baseline.

All applications for second and subsequent continuing treatments with this drug must include a measurement of response to the prior course of therapy. The Asthma Control Questionnaire (5 item version) assessment of the patient's response to the prior course of treatment or the assessment of oral corticosteroid dose, should be made at around 18 to 22 weeks after the first dose of PBS-subsidised dose of this drug under this restriction so that there is adequate time for a response to be demonstrated and for the application for continuing therapy to be processed.

The assessment should, where possible, be completed by the same physician who initiated treatment with this drug. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted within 4 weeks of the date of assessment, and no later than 2 weeks prior to the patient completing their current treatment course, to avoid an interruption to supply. Where a response assessment is not undertaken and submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

A patient who fails to respond to a course of PBS-subsidised mepolizumab for the treatment of uncontrolled severe eosinophilic asthma will not be eligible to receive further PBS subsidised treatment with mepolizumab for this condition within 6 months of the date on which treatment was ceased.

At the time of the authority application, medical practitioners should request the appropriate number of repeats to provide for a continuing course of this drug sufficient for up to 24 weeks of therapy.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and

(b) a completed Severe Eosinophilic Asthma Continuing PBS Authority Application - Supporting Information Form which includes details of maintenance oral corticosteroid dose; or a completed Asthma Control Questionnaire (ACQ-5) calculation sheet including the date of assessment of the patient's symptoms.

mepolizumab 100 mg injection, 1 vial

11014Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	1685.39	Nucala [GK]

■ MEPOLIZUMAB

Note TREATMENT OF ADULT AND ADOLESCENT PATIENTS WITH UNCONTROLLED SEVERE EOSINOPHILIC ASTHMA

Patients are eligible to commence a 'mepolizumab treatment cycle' (initial treatment course with or without continuing treatment course/s) if they satisfy the eligibility criteria as detailed under the initial treatment restriction.

Once a patient has either failed to achieve or maintain a response to mepolizumab, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 6 month break in PBS-subsidised mepolizumab therapy before they are eligible to commence the next mepolizumab treatment cycle, or if eligible, a 'benralizumab treatment cycle' or an 'omalizumab treatment cycle'. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised mepolizumab is stopped to the date of the first application for initial treatment with benralizumab, mepolizumab or omalizumab under the new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised mepolizumab therapy:

(a) Initial treatment:

Applications for initial treatment should be made where:

- i) A patient has received no prior PBS-subsidised mepolizumab treatment and wishes to commence such therapy; or
- ii) A patient wishes to recommence treatment with mepolizumab following a break in PBS-subsidised therapy of more than 6 months; or
- iii) A patient has received prior PBS-subsidised benralizumab or omalizumab and wishes to commence treatment with mepolizumab after a treatment break of 6 months.

All applications for initial treatment will be limited to provide for a maximum of up to 32 weeks of therapy for mepolizumab.

(b) Continuing treatment:

Following the completion of the initial treatment course with mepolizumab, a patient may qualify to receive up to a further 24 weeks of continuing treatment with mepolizumab providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing mepolizumab treatment in courses of up to 24 weeks providing they continue to sustain the response.

(2) Baseline measurements to determine response:

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the Asthma Control Questionnaire (ACQ; 5 item version) or oral corticosteroid dose, submitted with the Initial authority application for mepolizumab. However, prescribers may provide new baseline measurements when a new Initial treatment authority application is submitted and the Department of Human Services will assess response according to these revised baseline measurements.

(3) Re-commencement of treatment after a 6 month break in PBS-subsidised therapy:

A patient who wishes to trial a second or subsequent mepolizumab treatment cycle, or an initial benralizumab or omalizumab treatment cycle, following a break in PBS-subsidised therapy of at least 6 months, must re-qualify for initial treatment with respect to the indices of disease severity (oral corticosteroid dose, Asthma Control Questionnaire (ACQ-5) score, and relevant exacerbation history). Patients must have received optimised standard therapy, at adequate doses and for the minimum period specified, immediately prior to the time the new baseline assessments are performed.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note The Department of Human Services website (www.humanservices.gov.au) has details of the accepted toxicities, including severity, which will be accepted for the purposes of exempting a patient from the requirement of treatment with optimised asthma therapy.

Note For copies of the ACQ, please contact GlaxoSmithKline Medical Information on 1800 033 109.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application Forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Formal assessment and correction of inhaler technique should be performed in accordance with the National Asthma Council (NAC) Information Paper for Health Professionals on Inhaler Technique (available at www.humanservices.gov.au or www.nationalasthma.org.au); the assessment and adherence to correct technique should be documented in the patient's medical records. Patients can obtain support with inhaler technique through their local Asthma Foundation (1800 645 130).

Authority required

Uncontrolled severe eosinophilic asthma

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a respiratory physician, clinical immunologist, allergist or general physician experienced in the management of patients with severe asthma.

Clinical criteria:

- Patient must be under the care of the same physician for at least 6 months; OR
- Patient must have been diagnosed by a multidisciplinary severe asthma clinic team, **AND**
- Patient must have a diagnosis of asthma confirmed and documented by a respiratory physician, clinical immunologist, allergist or general physician experienced in the management of patients with severe asthma, defined by the following standard clinical features: (i) forced expiratory volume (FEV1) reversibility greater than or equal to 12% and greater than or equal to 200 mL at baseline within 30 minutes after administration of salbutamol (200 to 400 micrograms), or (ii) airway hyperresponsiveness defined as a greater than 20% decline in FEV1 during a direct bronchial provocation test or greater than 15% decline during an indirect bronchial provocation test, or (iii) peak expiratory flow (PEF) variability of greater than 15% between the two highest and two lowest peak expiratory flow rates during 14 days, **AND**
- Patient must have a duration of asthma of at least 1 year, **AND**
- Patient must have forced expiratory volume (FEV1) less than or equal to 80% predicted, documented on 1 or more occasions in the previous 12 months, **AND**
- Patient must have blood eosinophil count greater than or equal to 300 cells per microlitre in the last 12 months, **AND**
- Patient must have failed to achieve adequate control with optimised asthma therapy, despite formal assessment of and adherence to correct inhaler technique, which has been documented, **AND**
- The treatment must not be used in combination with, or within 6 months of treatment with, PBS-subsidised benralizumab or omalizumab.

Population criteria:

- Patient must be aged 12 years or older.

Optimised asthma therapy includes:

- (i) Adherence to maximal inhaled therapy, including high dose inhaled corticosteroid (ICS) plus long-acting beta-2 agonist (LABA) therapy for at least 12 months, unless contraindicated or not tolerated; **AND**
- (ii) treatment with oral corticosteroids, either daily oral corticosteroids for at least 6 weeks, OR a cumulative dose of oral corticosteroids of at least 500 mg prednisolone equivalent in the previous 12 months, unless contraindicated or not tolerated.

If the requirement for treatment with optimised asthma therapy cannot be met because of contraindications according to the relevant TGA-approved Product Information and/or intolerances of a severity necessitating permanent treatment withdrawal, details of the contraindication and/or intolerance must be provided in the Authority application.

The following initiation criteria indicate failure to achieve adequate control and must be demonstrated in all patients at the time of the application:

- (a) an Asthma Control Questionnaire (ACQ-5) score of at least 2.0, as assessed in the previous month, **AND**
 - (b) while receiving optimised asthma therapy in the past 12 months, experienced at least 1 admission to hospital for a severe asthma exacerbation, OR 1 severe asthma exacerbation, requiring documented use of systemic corticosteroids (oral corticosteroids initiated or increased for at least 3 days, or parenteral corticosteroids) prescribed/supervised by a physician.
- The Asthma Control Questionnaire (5 item version) assessment of the patient must be made at time of application for treatment (to establish baseline score) and again around 26 to 30 weeks after the first PBS-subsidised dose of this drug under this restriction so that there is adequate time for a response to be demonstrated and for the application for the first continuing therapy to be processed.

This assessment at around 26 to 30 weeks, which will be used to determine eligibility for the first continuing treatment, must be submitted within 4 weeks of the date of assessment, and no later than 2 weeks prior to the patient completing their current treatment course, to avoid an interruption to supply. Where a response assessment is not undertaken and submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

A patient who fails to respond to a course of PBS-subsidised mepolizumab for the treatment of uncontrolled severe eosinophilic asthma will not be eligible to receive further PBS-subsidised treatment with benralizumab, mepolizumab or omalizumab within 6 months of the date on which treatment was ceased.

At the time of the authority application, medical practitioners should request up to 7 repeats to provide for an initial course of mepolizumab sufficient for up to 32 weeks of therapy.

A multidisciplinary severe asthma clinic team comprises of:

- A respiratory physician; and
- A pharmacist, nurse or asthma educator.

Mepolizumab must not be used concurrently with benralizumab or omalizumab, or within 6 months of each other. A patient is required to have ceased treatment with benralizumab or omalizumab for 6 months prior to initiating treatment with mepolizumab.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Severe Eosinophilic Asthma Initial PBS Authority Application - Supporting Information Form, which includes the following:
 - (i) details of prior optimised asthma drug therapy (date of commencement and duration of therapy); and
 - (ii) details of severe exacerbation/s experienced in the past 12 months while receiving optimised asthma therapy (date and treatment); and
- (c) a copy of the eosinophil pathology report; and
- (d) a completed Asthma Control Questionnaire (ACQ-5) calculation sheet including the date of assessment of the patient's symptoms.

mepolizumab 100 mg injection, 1 vial

11003D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	7	..	1685.39	Nucala [GK]

■ OMALIZUMAB

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au. Applications for authority to prescribe should be forwarded to:
Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic spontaneous urticaria

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a clinical immunologist; OR
- Must be treated by an allergist; OR
- Must be treated by a dermatologist; OR
- Must be treated by a general physician with expertise in the management of chronic spontaneous urticaria (CSU).

Clinical criteria:

- The condition must be based on both physical examination and patient history (to exclude any factors that may be triggering the urticaria), **AND**
- Patient must have experienced itch and hives that persist on a daily basis for at least 6 weeks despite treatment with H1 antihistamines, **AND**
- Patient must have failed to achieve an adequate response after a minimum of 2 weeks treatment with a standard therapy, **AND**
- Patient must not receive more than 12 weeks of treatment under this restriction.

A standard therapy is defined as a combination of therapies that includes H1 antihistamines at maximally tolerated doses in accordance with clinical guidelines, and one of the following:

- 1) a H2 receptor antagonist (150 mg twice per day); or
- 2) a leukotriene receptor antagonist (LTRA) (10 mg per day); or
- 3) doxepin (up to 25 mg three times a day)

If the requirement for treatment with H1 antihistamines and a H2 receptor antagonist, or a leukotriene receptor antagonist or doxepin cannot be met because of contraindications according to the relevant TGA-approved Product Information and/or intolerances of a severity necessitating permanent treatment withdrawal, details of the contraindication and/or intolerance must be provided in the authority application.

A failure to achieve an adequate response to standard therapy is defined as a current Urticaria Activity Score 7 (UAS7) score of equal to or greater than 28 with an itch score of greater than 8, as assessed while still on standard therapy.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Chronic Spontaneous Urticaria Omalizumab Initial PBS Authority Application - Supporting Information Form which must include:
 - (i) demonstration of failure to achieve an adequate response to standard therapy; and
 - (ii) drug names and doses of standard therapies that the patient has failed; and
 - (iii) a signed patient acknowledgment that cessation of therapy should be considered after the patient has demonstrated clinical benefit with omalizumab to re-evaluate the need for continued therapy. Any patient who ceases therapy and whose CSU relapses will need to re-initiate PBS-subsidised omalizumab as a new patient.

omalizumab 150 mg/mL injection, 1 mL syringe

11175E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	2	..	*860.19	Xolair [NV]

■ OMALIZUMAB

Authority required

Severe chronic spontaneous urticaria

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a clinical immunologist; OR
- Must be treated by an allergist; OR
- Must be treated by a dermatologist; OR
- Must be treated by a general physician with expertise in the management of chronic spontaneous urticaria (CSU).

Clinical criteria:

- Patient must have demonstrated a response to the most recent PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not receive more than 24 weeks per authorised course of treatment under this restriction.

Note A proportion of patients respond to 150 mg 4-weekly so where a substantial improvement has been obtained with a 300 mg dose it is reasonable to back-titrate dose after initial treatment.

Note Cessation of therapy should be considered after the patient has demonstrated clinical benefit with omalizumab to re-evaluate the need for continued therapy. Any patient who ceases therapy and whose CSU relapses will need to re-initiate PBS-subsidised omalizumab as a new patient.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Severe chronic spontaneous urticaria

Treatment Phase: Grandfathering treatment

Clinical criteria:

- Patient must have received non-PBS subsidised treatment with this drug for this condition prior to 1 September 2017, **AND**
- Patient must have documented history of itch and hives that persisted on a daily basis for at least 6 weeks despite treatment with H1 antihistamines prior to commencing non-PBS subsidised treatment with this drug for this condition, **AND**
- Patient must have documented history of failure to achieve an adequate response after a minimum of 2 weeks treatment with a standard therapy prior to commencing non-PBS subsidised treatment with this drug for this condition, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Treatment criteria:

- Must be treated by a clinical immunologist; OR
- Must be treated by an allergist; OR
- Must be treated by a dermatologist; OR
- Must be treated by a general physician with expertise in the management of chronic spontaneous urticaria (CSU). A standard therapy is defined as a combination of therapies that includes H1 antihistamines at maximally tolerated doses in accordance with clinical guidelines, and one of the following:
 - 1) a H2 receptor antagonist (150 mg twice per day); or
 - 2) a leukotriene receptor antagonist (LTRA) (10 mg per day); or
 - 3) doxepin (up to 25 mg three times a day)

If the requirement for treatment with H1 antihistamines and a H2 receptor antagonist, or a leukotriene receptor antagonist or doxepin cannot be met because of contraindications according to the relevant TGA-approved Product Information and/or intolerances of a severity necessitating permanent treatment withdrawal, details of the contraindication and/or intolerance must be provided in the authority application.

A failure to achieve an adequate response to standard therapy is defined as a current Urticaria Activity Score 7 (UAS7) score of equal to or greater than 28 with an itch score of greater than 8, as assessed while still on standard therapy.

A patient may qualify for PBS-subsidised treatment under this restriction once only. For continuing PBS-subsidised treatment, a Grandfathered patient must qualify under the Continuing treatment criteria.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Chronic Spontaneous Urticaria Omalizumab Initial Grandfather PBS Authority Application - Supporting Information Form which must include:
 - (i) demonstration of failure to achieve an adequate response to standard therapy; and
 - (ii) drug names and doses of standard therapies that the patient has failed; and
 - (iii) a signed patient acknowledgment that cessation of therapy should be considered after the patient has demonstrated clinical benefit with omalizumab to re-evaluate the need for continued therapy. Any patient who ceases therapy and whose CSU relapses will need to re-initiate PBS-subsidised omalizumab as a new patient.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

omalizumab 150 mg/mL injection, 1 mL syringe

11163M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*860.19	Xolair [NV]

■ OMALIZUMAB

Note TREATMENT OF ADULT AND ADOLESCENT PATIENTS WITH UNCONTROLLED SEVERE ALLERGIC ASTHMA

Patients are eligible to commence an 'omalizumab treatment cycle' (initial treatment course with or without continuing treatment course/s) if they satisfy the eligibility criteria as detailed under the initial treatment restriction.

Once a patient has either failed to achieve or maintain a response to omalizumab, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 6 month break in PBS-subsidised omalizumab therapy before they are eligible to commence the next omalizumab treatment cycle or, if eligible, a 'benralizumab treatment cycle' or a 'mepolizumab treatment cycle'. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised omalizumab is stopped to the date of the first application for initial treatment with benralizumab, omalizumab or mepolizumab under the new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised omalizumab therapy:

(a) Initial treatment:

Applications for initial treatment should be made where:

- i) A patient has received no prior PBS-subsidised omalizumab treatment and wishes to commence such therapy; or
- ii) A patient wishes to recommence treatment with omalizumab following a break in PBS-subsidised therapy of at least 6 months; or
- iii) A patient has received prior PBS-subsidised benralizumab or mepolizumab and wishes to commence treatment with omalizumab after a treatment break of at least 6 months.

All applications for initial treatment will be limited to provide for a maximum of up to 28 weeks of therapy of omalizumab.

(b) Continuing treatment:

Following the completion of the initial treatment course with omalizumab, a patient may qualify to receive up to a further 24 weeks of continuing treatment with omalizumab providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing omalizumab treatment in courses of up to 24 weeks providing they continue to sustain the response.

(2) Baseline measurements to determine response:

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the Asthma Control Questionnaire (ACQ; 5 item version) or oral corticosteroid dose submitted with the Initial authority application for omalizumab. For patients transitioned from the paediatric to the adolescent/adult restriction, the exacerbation history may also be used to determine response. However, prescribers may provide new baseline measurements when a new Initial treatment authority application is submitted and the Department of Human Services will assess response according to these revised baseline measurements.

(3) Re-commencement of treatment after a 6 month break in PBS-subsidised therapy:

A patient who wishes to trial a second or subsequent omalizumab treatment cycle, or an initial benralizumab or mepolizumab treatment cycle, following a break in PBS-subsidised therapy of at least 6 months, must re-qualify for initial treatment with respect to the indices of disease severity (oral corticosteroid dose, Asthma Control Questionnaire (ACQ-5) score, and relevant exacerbation history). Patients must have received optimised standard therapy, at adequate doses and for the minimum period specified, immediately prior to the time the new baseline assessments are performed.

(4) Monitoring of patients:

Anaphylaxis and anaphylactoid reactions have been reported following first or subsequent administration of omalizumab (see Product Information). Patients should be monitored post-injection, and medications for the treatment of anaphylactic reactions should be available for immediate use following administration of omalizumab. Patients should be informed that such reactions are possible and prompt medical attention should be sought if allergic reactions occur.

Note Special Pricing Arrangements apply.

Authority required

Uncontrolled severe allergic asthma

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a respiratory physician, clinical immunologist, allergist or general physician experienced in the management of patients with severe asthma.

Clinical criteria:

- Patient must be under the care of the same physician for at least 6 months; OR
- Patient must have been diagnosed by a multidisciplinary severe asthma clinic team, **AND**
- Patient must have a diagnosis of asthma confirmed and documented by a respiratory physician, clinical immunologist, allergist or general physician experienced in the management of patients with severe asthma, defined by the following standard clinical features: (i) forced expiratory volume (FEV1) reversibility greater than or equal to 12% and greater than or equal to 200 mL at baseline within 30 minutes after administration of salbutamol (200 to 400 micrograms), or (ii) airway hyperresponsiveness defined as a greater than 20% decline in FEV1 during a direct bronchial provocation test or greater than 15% decline during an indirect bronchial provocation test, or (iii) peak expiratory flow (PEF) variability of greater than 15% between the two highest and two lowest peak expiratory flow rates during 14 days, **AND**
- Patient must have a duration of asthma of at least 1 year, **AND**
- Patient must have forced expiratory volume (FEV1) less than or equal to 80% predicted, documented on 1 or more occasions in the previous 12 months, **AND**
- Patient must have past or current evidence of atopy, documented by skin prick testing or RAST, **AND**
- Patient must have total serum human immunoglobulin E greater than or equal to 30 IU/mL, **AND**
- Patient must have failed to achieve adequate control with optimised asthma therapy, despite formal assessment of and adherence to correct inhaler technique, which has been documented, **AND**
- Patient must not receive more than 28 weeks of treatment under this restriction, **AND**
- The treatment must not be used in combination with, or within 6 months of treatment with, PBS-subsidised benralizumab or mepolizumab.

Population criteria:

- Patient must be aged 12 years or older.

Optimised asthma therapy includes:

(i) Adherence to maximal inhaled therapy, including high dose inhaled corticosteroid (ICS) plus long-acting beta-2 agonist (LABA) therapy for at least 12 months, unless contraindicated or not tolerated; **AND**

(ii) treatment with oral corticosteroids, either daily oral corticosteroids for at least 6 weeks, OR a cumulative dose of oral corticosteroids of at least 500 mg prednisolone equivalent in the previous 12 months, unless contraindicated or not tolerated.

If the requirement for treatment with optimised asthma therapy cannot be met because of contraindications according to the relevant TGA-approved Product Information and/or intolerances of a severity necessitating permanent treatment withdrawal, details of the contraindication and/or intolerance must be provided in the Authority application.

The initial IgE assessment must be no more than 12 months old at the time of application.

The following initiation criteria indicate failure to achieve adequate control and must be demonstrated in all patients at the time of the application:

- (a) an Asthma Control Questionnaire (ACQ-5) score of at least 2.0, as assessed in the previous month, AND
 - (b) while receiving optimised asthma therapy in the past 12 months, experienced at least 1 admission to hospital for a severe asthma exacerbation, OR 1 severe asthma exacerbation, requiring documented use of systemic corticosteroids (oral corticosteroids initiated or increased for at least 3 days, or parenteral corticosteroids) prescribed/supervised by a physician.
- The Asthma Control Questionnaire (5 item version) assessment of the patient's response to this initial course of treatment, and the assessment of oral corticosteroid dose, must be made at around 22 to 26 weeks after the first PBS-subsidised dose of this drug under this restriction so that there is adequate time for a response to be demonstrated and for the application for the first continuing therapy to be processed.

This assessment, which will be used to determine eligibility for the first continuing treatment, must be submitted within 4 weeks of the date of assessment, and no later than 2 weeks prior to the patient completing their current treatment course, to avoid an interruption to supply. Where a response assessment is not undertaken and submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

A patient who fails to respond to a course of PBS-subsidised omalizumab for the treatment of uncontrolled severe allergic asthma will not be eligible to receive further PBS-subsidised treatment with benralizumab, omalizumab or mepolizumab for this condition within 6 months of the date on which treatment was ceased.

At the time of the authority application, medical practitioners should request the appropriate maximum quantity and number of repeats to provide for an initial course of omalizumab consisting of the recommended number of doses for the baseline IgE level and body weight of the patient (refer to the TGA-approved Product Information) to be administered every 2 or 4 weeks.

A multidisciplinary severe asthma clinic team comprises of:

- A respiratory physician; and
- A pharmacist, nurse or asthma educator.

Omalizumab must not be used concurrently with benralizumab or mepolizumab, or within 6 months of each other. A patient is required to have ceased treatment with benralizumab or mepolizumab for 6 months prior to initiating treatment with omalizumab.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Severe Allergic Asthma PBS Authority Application - Supporting Information Form, which includes the following:
 - (i) details of prior optimised asthma drug therapy (date of commencement and duration of therapy); and
 - (ii) details of severe exacerbation/s experienced in the past 12 months while receiving optimised asthma therapy (date and treatment); and
- (c) the IgE pathology report; and
- (d) a completed Asthma Control Questionnaire (ACQ-5) calculation sheet including the date of assessment of the patient's symptoms.

Note The Department of Human Services website (www.humanservices.gov.au) has details of the accepted toxicities, including severity, which will be accepted for the purposes of exempting a patient from the requirement of treatment with optimised asthma therapy.

Note For copies of the ACQ and the calculation sheets please contact Novartis Medical Information on 1800 671 203 or medinfo.phauno@novartis.com

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Formal assessment and correction of inhaler technique should be performed in accordance with the National Asthma Council (NAC) Information Paper for Health Professionals on Inhaler Technique (available at www.humanservices.gov.au or www.nationalasthma.org.au); the assessment and adherence to correct technique should be documented in the patient's medical records. Patients can obtain support with inhaler technique through their local Asthma Foundation (1800 645 130).

Authority required

Uncontrolled severe allergic asthma

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have a documented history of severe allergic asthma, **AND**
- Patient must have demonstrated or sustained an adequate response to PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction, **AND**
- The treatment must not be used in combination with, or within 6 months of treatment with, PBS-subsidised benralizumab or mepolizumab.

Treatment criteria:

- Must be treated by a respiratory physician, clinical immunologist, allergist or general physician experienced in the management of patients with severe asthma.

Population criteria:

- Patient must be aged 12 years or older.

An adequate response to omalizumab treatment is defined as:

- (a) a reduction in the Asthma Control Questionnaire (ACQ-5) score of at least 0.5 from baseline, OR
- (b) maintenance oral corticosteroid dose reduced by at least 25% from baseline, and no deterioration in ACQ-5 score from baseline, OR
- (c) a reduction in the time-adjusted exacerbation rates compared to the 12 months prior to baseline (this criterion is only applicable for patients transitioned from the paediatric to the adolescent/adult restriction).

All applications for second and subsequent continuing treatments with this drug must include a measurement of response to the prior course of therapy. The Asthma Control Questionnaire (5 item version) assessment of the patient's response to the prior course of treatment, the assessment of oral corticosteroid dose or the assessment of time adjusted exacerbation rate must be made at around 18 to 22 weeks after the first PBS-subsidised dose of this drug under this restriction so that there is adequate time for a response to be demonstrated and for the application for continuing therapy to be processed.

The assessment should, where possible, be completed by the same physician who initiated treatment with this drug. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted within 4 weeks of the date of assessment, and no later than 2 weeks prior to the patient completing their current treatment course, to avoid an interruption to supply. Where a response assessment is not undertaken and submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

A patient who fails to respond to a course of PBS-subsidised omalizumab for the treatment of uncontrolled severe allergic asthma will not be eligible to receive further PBS-subsidised treatment with omalizumab for this condition within 6 months of the date on which treatment was ceased.

At the time of the authority application, medical practitioners should request the appropriate quantity and number of repeats to provide for a continuing course of omalizumab consisting of the recommended number of doses for the baseline IgE level and body weight of the patient (refer to the TGA-approved Product Information), sufficient for up to 24 weeks of therapy.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Allergic Asthma PBS Authority Application and Supporting Information Form which includes details of maintenance oral corticosteroid dose; or
- (c) a completed Asthma Control Questionnaire (ACQ-5) calculation sheet including the date of assessment of the patient's symptoms and is endorsed with the signature of the prescriber; for patients transitioned from the paediatric to the adolescent/adult restrictions an exacerbation calculation sheet may be submitted.

Note If the same physician cannot assess the patient please call the Department of Human Services on 1800 700 270.

Note For copies of the ACQ and the calculation sheets please contact Novartis Medical Information on 1800 671 203 or medinfo.phauno@novartis.com

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Formal assessment and correction of inhaler technique should be performed in accordance with the National Asthma Council (NAC) Information Paper for Health Professionals on Inhaler Technique (available at www.humanservices.gov.au or www.nationalasthma.org.au); the assessment and adherence to correct technique should be documented in the patient's medical records. Patients can obtain support with inhaler technique through their local Asthma Foundation (1800 645 130).

Authority required

Uncontrolled severe allergic asthma

Treatment Phase: Initial and continuing treatment - balance of supply

Treatment criteria:

- Must be treated by a respiratory physician, clinical immunologist, allergist or general physician experienced in the management of patients with severe asthma.

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the Initial treatment restriction to complete 28 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the Continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 28 weeks treatment available under the Initial restriction or up to 24 weeks treatment available under the Continuing restriction.

Note Authority approval for sufficient therapy to complete a maximum of 28 weeks of treatment under the initial restriction or 24 weeks of treatment under the continuing restriction may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

omalizumab 150 mg/mL injection, 1 mL syringe

10122R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	433.79	Xolair [NV]

omalizumab 75 mg/0.5 mL injection, 0.5 mL syringe

10110D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	220.59	Xolair [NV]

■ OMALIZUMAB**Note TREATMENT OF PAEDIATRIC PATIENTS WITH UNCONTROLLED SEVERE ALLERGIC ASTHMA**

Patients are eligible to commence an 'omalizumab treatment cycle' (initial treatment course with or without continuing treatment course/s) if they satisfy the eligibility criteria as detailed under the initial treatment restriction. Once a patient has either failed to achieve or maintain a response to omalizumab, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 6 month break in PBS-subsidised omalizumab therapy before they are eligible to commence the next cycle. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised omalizumab treatment is stopped to the date of the first application for initial treatment with omalizumab under the new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised omalizumab therapy.

(a) Initial treatment:

Applications for initial treatment should be made where a patient has received no prior PBS-subsidised omalizumab treatment in this treatment cycle and wishes to commence such therapy.

All applications for initial treatment will be limited to provide for a maximum of 28 weeks of therapy for omalizumab.

(b) Continuing treatment:

Following the completion of the initial treatment course with omalizumab, a patient may qualify to receive up to a further 24 weeks of continuing treatment with omalizumab providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing omalizumab treatment in courses of up to 24 weeks providing they continue to sustain the response.

(2) Baseline measurements to determine response:

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the Asthma Control Questionnaire (ACQ; 5 item version) or ACQ-IA, systemic corticosteroid dose and time-adjusted exacerbation rate, submitted with the Initial authority application for omalizumab. However, prescribers may provide new baseline measurements when a new Initial treatment authority application is submitted and The Department of Human Services will assess response according to these revised baseline measurements.

(3) Re-commencement of treatment after a 6 month break in PBS-subsidised therapy:

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised omalizumab therapy of at least 6 months, must re-qualify for initial treatment with respect to the indices of disease severity (systemic corticosteroid dose, Asthma Control Questionnaire (ACQ-5) score or ACQ-IA, and relevant exacerbation history). Patients must have received optimised standard therapy, at adequate doses and for the minimum period specified, immediately prior to the time the new baseline assessments are performed.

(4) Monitoring of patients:

Anaphylaxis and anaphylactoid reactions have been reported following first or subsequent administration of omalizumab (see Product Information). Patients should be monitored post-injection, and medications for the treatment of anaphylactic reactions should be available for immediate use following administration of omalizumab. Patients should be informed that such reactions are possible and prompt medical attention should be sought if allergic reactions occur.

Note Special Pricing Arrangements apply.

Authority required

Uncontrolled severe allergic asthma

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have a diagnosis of asthma confirmed and documented by a paediatric respiratory physician, clinical immunologist, or allergist; or paediatrician or general physician experienced in the management of patients with severe asthma in consultation with a respiratory physician, defined by the following standard clinical features: forced expiratory volume (FEV1) reversibility or airway hyperresponsiveness or peak expiratory flow (PEF) variability, **AND**
- Patient must have a duration of asthma of at least 1 year, **AND**
- Patient must have past or current evidence of atopy, documented by skin prick testing or an in vitro measure of specific IgE, **AND**
- Patient must have total serum human immunoglobulin E greater than or equal to 30 IU/mL, **AND**
- Patient must have failed to achieve adequate control with optimised asthma therapy, despite formal assessment of and adherence to correct inhaler technique, which has been documented, **AND**
- Patient must not receive more than 28 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 6 to less than 12 years.

Treatment criteria:

- Must be treated by a paediatric respiratory physician, clinical immunologist, allergist; or paediatrician or general physician experienced in the management of patients with severe asthma, in consultation with a respiratory physician.

Clinical criteria:

- Patient must be under the care of the same physician for at least 6 months.

Optimised asthma therapy includes:

(i) Adherence to optimal inhaled therapy, including high dose inhaled corticosteroid (ICS) and long-acting beta-2 agonist (LABA) therapy for at least six months. If LABA therapy is contraindicated, not tolerated or not effective, montelukast, cromoglycate or nedocromil may be used as an alternative; **AND**

(ii) treatment with at least 2 courses of oral or IV corticosteroids (daily or alternate day maintenance treatment courses, or 3-5 day exacerbation treatment courses), in the previous 12 months, unless contraindicated or not tolerated.

If the requirement for treatment with optimised asthma therapy cannot be met because of contraindications (including those specified in the relevant TGA-approved Product Information) and/or intolerances of a severity necessitating permanent treatment withdrawal, details of the contraindication and/or intolerance must be provided in the Authority application.

The initial IgE assessment must be no more than 12 months old at the time of application.

The following initiation criteria indicate failure to achieve adequate control and must be demonstrated in all patients at the time of the application:

- (a) An Asthma Control Questionnaire (ACQ-5) score of at least 2.0, as assessed in the previous month (for children aged 6 to 10 years it is recommended that the Interviewer Administered version - the ACQ-IA be used), AND
 - (b) while receiving optimised asthma therapy in the previous 12 months, experienced at least 1 admission to hospital for a severe asthma exacerbation, OR 1 severe asthma exacerbation, requiring documented use of systemic corticosteroids (oral corticosteroids initiated or increased for at least 3 days, or parenteral corticosteroids) prescribed/supervised by a physician.
- The Asthma Control Questionnaire (5 item version) or ACQ-IA assessment of the patient's response to this initial course of treatment, the assessment of oral corticosteroid dose, and the assessment of exacerbation rate must be made at around 22 to 26 weeks after the first dose so that there is adequate time for a response to be demonstrated and for the application for continuing therapy to be processed.

This assessment, which will be used to determine eligibility for continuing treatment, must be submitted within 4 weeks of the date of assessment, and no later than 2 weeks prior to the patient completing their current treatment course, to avoid an interruption to supply. Where a response assessment is not undertaken and submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with omalizumab.

A patient who fails to respond to a course of PBS-subsidised omalizumab for the treatment of uncontrolled severe allergic asthma will not be eligible to receive further PBS-subsidised treatment with omalizumab for this condition within 6 months of the date on which treatment was ceased.

At the time of the authority application, medical practitioners should request the appropriate maximum quantity and number of repeats to provide for an initial course of omalizumab of up to 28 weeks, consisting of the recommended number of doses for the baseline IgE level and body weight of the patient (refer to the TGA-approved Product Information) to be administered every 2 or 4 weeks.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Paediatric Severe Allergic Asthma Initial PBS Authority Application - Supporting Information form, which includes the following:
 - (i) details of prior optimised asthma drug therapy (dosage, date of commencement and duration of therapy); and
 - (ii) details of severe exacerbation/s experienced in the past 12 months while receiving optimised asthma therapy (date and treatment); and
 - (iii) acknowledgement signed by a parent or authorised guardian; and
- (c) a copy of the IgE pathology report; and
- (d) a completed Asthma Control Questionnaire (ACQ-5) or the Asthma Control Questionnaire interviewer administered version (ACQ-IA) calculation sheet including the date of assessment of the patient's symptoms and is endorsed with the prescriber's signature.

Note The Department of Human Services website (www.humanservices.gov.au) has details of the accepted toxicities, including severity, which will be accepted for the purposes of exempting a patient from the requirement of treatment with optimised asthma therapy.

Note For copies of the ACQ please contact Novartis Medical Information on 1800 671 203 or medinfo.phauno@novartis.com

Note It is recommended that an application for continuing treatment is submitted at the time of the 22 to 26 week assessment, to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised omalizumab treatment.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Formal assessment and correction of inhaler technique should be performed in accordance with the National Asthma Council (NAC) Information Paper for Health Professionals on Inhaler Technique (available at www.humanservices.gov.au or www.nationalasthma.org.au); the assessment and adherence to correct technique should be documented in the patient's medical records. Patients can obtain support with inhaler technique through their local Asthma Foundation (1800 645 130).

Authority required

Uncontrolled severe allergic asthma

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have a documented history of severe allergic asthma, **AND**
- Patient must have demonstrated or sustained an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Treatment criteria:

- Must be treated by a paediatric respiratory physician, clinical immunologist, allergist; or paediatrician or general physician experienced in the management of patients with severe asthma, in consultation with a respiratory physician.

An adequate response to omalizumab treatment is defined as:

- (a) a reduction in the Asthma Control Questionnaire (ACQ-5) or ACQ-IA score of at least 0.5 from baseline, OR
 (b) maintenance oral corticosteroid dose reduced by at least 25% from baseline, and no deterioration in ACQ-5 or ACQ-IA score from baseline, OR
 (c) a reduction in the time-adjusted exacerbation rates compared to the 12 months prior to baseline.

All applications for continuing treatment with omalizumab must include a measurement of response to the prior course of therapy. The Asthma Control Questionnaire (5 item version) or Asthma Control Questionnaire interviewer administered version (ACQ-IA) assessment of the patient's response to the prior course of treatment, the assessment of systemic corticosteroid dose, and the assessment of time-adjusted exacerbation rate must be made at around 18 to 22 weeks after the first dose of PBS-subsidised omalizumab so that there is adequate time for a response to be demonstrated and for the application for continuing therapy to be processed.

The first assessment should, where possible, be completed by the same physician who initiated treatment with omalizumab. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted within 4 weeks of the date of assessment, and no later than 2 weeks prior to the patient completing their current treatment course, to avoid an interruption to supply. Where a response assessment is not undertaken and submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with omalizumab.

A patient who fails to respond to a course of PBS-subsidised omalizumab for the treatment of uncontrolled severe allergic asthma will not be eligible to receive further PBS-subsidised treatment with omalizumab for this condition within 6 months of the date on which treatment was ceased.

At the time of the authority application, medical practitioners should request the appropriate quantity and number of repeats to provide for a continuing course of omalizumab consisting of the recommended number of doses for the baseline IgE level and body weight of the patient (refer to the TGA-approved Product Information), sufficient for 24 weeks of therapy.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Paediatric Severe Allergic Asthma Continuing PBS Authority Application - Supporting Information form which includes details of maintenance oral corticosteroid dose; and
- (c) a completed Asthma Control Questionnaire (ACQ-5) or the Asthma Control Questionnaire interviewer administered version (ACQ-IA) calculation sheet including the date of assessment of the patient's symptoms and is endorsed with the signature of the prescriber.

Note If the same physician cannot assess the patient please call the Department of Human Services on 1800 700 270.

Note For copies of the ACQ please contact Novartis Medical Information on 1800 671 203 or medinfo.phauno@novartis.com

Note It is recommended that an application for continuing treatment is submitted at the time of the 18 to 22 week assessment, to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised omalizumab treatment.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
 Complex Drugs
 Reply Paid 9826
 HOBART TAS 7001

Note Formal assessment and correction of inhaler technique should be performed in accordance with the National Asthma Council (NAC) Information Paper for Health Professionals on Inhaler Technique (available at www.humanservices.gov.au or www.nationalasthma.org.au); the assessment and adherence to correct technique should be documented in the patient's medical records. Patients can obtain support with inhaler technique through their local Asthma Foundation (1800 645 130).

Authority required

Uncontrolled severe allergic asthma

Treatment Phase: Initial and continuing treatment - balance of supply

Treatment criteria:

- Must be treated by a paediatric respiratory physician, clinical immunologist, allergist; or paediatrician or general physician experienced in the management of patients with severe asthma, in consultation with a respiratory physician.

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the Initial treatment restriction to complete 28 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the Continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 28 weeks treatment available under the Initial restriction or up to 24 weeks treatment available under the Continuing restriction.

Note Authority approval for sufficient therapy to complete a maximum of 28 weeks of treatment under the initial restriction or 24 weeks of treatment under the continuing restriction may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

omalizumab 150 mg/mL injection, 1 mL syringe

10968G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	433.79	Xolair [NV]

omalizumab 75 mg/0.5 mL injection, 0.5 mL syringe

10956P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	220.59	Xolair [NV]

■ **COUGH AND COLD PREPARATIONS****EXPECTORANTS, EXCL. COMBINATIONS WITH COUGH SUPPRESSANTS***Mucolytics*■ **DORNASE ALFA**

Note It is highly desirable that all patients be included in the national cystic fibrosis patient database.

Authority required (STREAMLINED)**9624**

Cystic fibrosis

Population criteria:

- Patient must be 5 years of age or older.

Patient must be assessed at a cystic fibrosis clinic/centre which is under the control of specialist respiratory physicians with experience and expertise in the management of cystic fibrosis or by a specialist physician or paediatrician in consultation with such a unit.

Prior to therapy with this drug, a baseline measurement of forced expiratory volume in 1 second (FEV1) must be undertaken during a stable period of the disease.

Initial therapy is limited to 3 months treatment with dornase alfa at a dose of 2.5 mg daily.

To be eligible for continued PBS-subsidised treatment with this drug following 3 months of initial treatment:

- (1) the patient must demonstrate no deterioration in FEV1 compared to baseline; AND
- (2) the patient or the patient's family (in the case of paediatric patients) and the treating physician(s) must report a benefit in the clinical status of the patient.

Further reassessments must be undertaken and documented at six-monthly intervals. Therapy with this drug should cease if there is not general agreement of benefit as there is always the possibility of harm from unnecessary use.

Authority required (STREAMLINED)**9591**

Cystic fibrosis

Clinical criteria:

- Patient must have a severe clinical course with frequent respiratory exacerbations or chronic respiratory symptoms (including chronic or recurrent cough, wheeze or tachypnoea) requiring hospital admissions more frequently than 3 times per year; OR
- Patient must have significant bronchiectasis on chest high resolution computed tomography scan; OR
- Patient must have severe cystic fibrosis bronchiolitis with persistent wheeze non-responsive to conventional medicines; OR
- Patient must have severe physiological deficit measure by forced oscillation technique or multiple breath nitrogen washout and failure to respond to conventional therapy.

Population criteria:

- Patient must be less than 5 years of age.

Patient must be assessed at a cystic fibrosis clinic/centre which is under the control of specialist respiratory physicians with experience and expertise in the management of cystic fibrosis or by a specialist physician or paediatrician in consultation with such a unit.

Following an initial 6 months therapy, a comprehensive assessment must be undertaken and documented. Treatment with this drug should cease if there is not agreement of benefit, as there is always the possibility of harm from unnecessary use. Further reassessments must be undertaken and documented at six-monthly intervals.

Authority required (STREAMLINED)**9592**

Cystic fibrosis

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have initiated treatment with dornase alfa at an age of less than 5 years, **AND**
- Patient must have undergone a comprehensive assessment which documents agreement that dornase alfa treatment is continuing to produce worthwhile benefit.

Population criteria:

- Patient must be 5 years of age or older.

Further reassessments must be undertaken and documented at six-monthly intervals. Treatment with this drug should cease if there is not agreement of benefit as there is always the possibility of harm from unnecessary use.

dornase alfa 2.5 mg/2.5 mL inhalation solution, 30 x 2.5 mL ampoules

6120D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*1964.31	Pulmozyme [RO]

■ **MANNITOL**

Note It is highly desirable that all patients be included in the national cystic fibrosis patient database.

Authority required (STREAMLINED)

9527

Cystic fibrosis

Clinical criteria:

- The treatment must be as monotherapy, **AND**
- Patient must be intolerant or inadequately responsive to dornase alfa.

Population criteria:

- Patient must be 6 years of age or older.

Patient must have been assessed for bronchial hyperresponsiveness as per the TGA approved Product Information initiation dose assessment for this drug, prior to therapy with this drug, with a negative result.

Patient must be assessed at a cystic fibrosis clinic/centre which is under the control of specialist respiratory physicians with experience and expertise in the management of cystic fibrosis or by a specialist physician or paediatrician in consultation with such a unit.

Prior to therapy with this drug, a baseline measurement of forced expiratory volume in 1 second (FEV1) must be undertaken during a stable period of the disease.

Initial therapy is limited to 3 months treatment with mannitol at a dose of 400 mg twice daily.

To be eligible for continued PBS-subsidised treatment with this drug following 3 months of initial treatment:

- (1) the patient must demonstrate no deterioration in FEV1 compared to baseline; **AND**
- (2) the patient or the patient's family (in the case of paediatric patients) and the treating physician(s) must report a benefit in the clinical status of the patient.

Further reassessments must be undertaken and documented at six-monthly intervals. Therapy with this drug should cease if there is not general agreement of benefit as there is always the possibility of harm from unnecessary use.

Authority required (STREAMLINED)**9593**

Cystic fibrosis

Clinical criteria:

- The treatment must be in combination with dornase alfa, **AND**
- Patient must be inadequately responsive to dornase alfa, **AND**
- Patient must have trialled hypertonic saline for this condition.

Population criteria:

- Patient must be 6 years of age or older.

Patient must have been assessed for bronchial hyperresponsiveness as per the TGA approved Product Information initiation dose assessment for this drug, prior to therapy with this drug, with a negative result.

Patient must be assessed at a cystic fibrosis clinic/centre which is under the control of specialist respiratory physicians with experience and expertise in the management of cystic fibrosis or by a specialist physician or paediatrician in consultation with such a unit.

Prior to therapy with this drug, a baseline measurement of forced expiratory volume in 1 second (FEV1) must be undertaken during a stable period of the disease.

Initial therapy is limited to 3 months treatment with mannitol at a dose of 400 mg twice daily.

To be eligible for continued PBS-subsidised treatment with this drug following 3 months of initial treatment:

- (1) the patient must demonstrate no deterioration in FEV1 compared to baseline; **AND**
- (2) the patient or the patient's family (in the case of paediatric patients) and the treating physician(s) must report a benefit in the clinical status of the patient.

Further reassessments must be undertaken and documented at six-monthly intervals. Therapy with this drug should cease if there is not general agreement of benefit as there is always the possibility of harm from unnecessary use.

mannitol 40 mg powder for inhalation, 280 capsules

2008Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	4	5	..	*1836.59	bronchitol [XA]

OTHER RESPIRATORY SYSTEM PRODUCTS**OTHER RESPIRATORY SYSTEM PRODUCTS***Other respiratory system products***■ IVACAFTOR**

Note No increase in the maximum number of repeats may be authorised.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Cystic fibrosis

Treatment Phase: Initial treatment - New patients

Clinical criteria:

- Patient must be assessed through a cystic fibrosis clinic/centre which is under the control of specialist respiratory physicians with experience and expertise in the management of cystic fibrosis. If attendance at such a unit is not possible because of geographical isolation, management (including prescribing) may be in consultation with such a unit, **AND**
- Patient must have G551D mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene on at least 1 allele; OR
- Patient must have other gating (class III) mutation in the CFTR gene on at least 1 allele, **AND**
- Patient must have a sweat chloride value of at least 60 mmol/L by quantitative pilocarpine iontophoresis, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction, **AND**
- The treatment must be given concomitantly with standard therapy for this condition.

Population criteria:

- Patient must be aged 12 months or older.

Patients receiving PBS-subsidised ivacaftor must be registered in the Australian Cystic Fibrosis Database Registry.

Treatment must not be given to a patient who has an acute upper or lower respiratory infection, pulmonary exacerbation, or changes in therapy (including antibiotics) for pulmonary disease in the last 4 weeks prior to commencing this drug.

Dosage of ivacaftor must not exceed the dose of one tablet (150 mg) or one sachet twice a week, if the patient is concomitantly receiving one of the following strong CYP3A4 drugs inhibitors: boceprevir, clarithromycin, conivaptan, indinavir, itraconazole, ketoconazole, lopinavir/ritonavir, mibefradil, nefazodone, nelfinavir, posaconazole, ritonavir, saquinavir, telaprevir, telithromycin, voriconazole. Where a patient is concomitantly receiving a strong CYP3A4 inhibitor, a single supply of 56 tablets or sachets of ivacaftor will last for 28 weeks.

Dosage of ivacaftor must not exceed the dose of one tablet (150 mg) or one sachet once daily, if the patient is concomitantly receiving one of the following moderate CYP3A4 inhibitors: amprenavir, aprepitant, atazanavir, darunavir/ritonavir, diltiazem, erythromycin, fluconazole, fosamprenavir, imatinib, verapamil. Where a patient is concomitantly receiving a moderate CYP3A4 inhibitor, a single supply of 56 tablets or sachets of ivacaftor will last for 8 weeks.

Ivacaftor is not PBS-subsidised for this condition as a sole therapy.

Ivacaftor is not PBS-subsidised for this condition in a patient who is currently receiving one of the following CYP3A4 inducers:

Strong CYP3A4 inducers: avasimibe, carbamazepine, phenobarbital, phenytoin, rifabutin, rifampicin, St. John's wort

Moderate CYP3A4 inducers: bosentan, efavirenz, etravirine, modafinil, nafcillin

Weak CYP3A4 inducers: armodafinil, echinacea, pioglitazone, rufinamide.

The authority application must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Cystic Fibrosis Ivacaftor Authority Application Supporting Information Form; and
- (3) a copy of the pathology report detailing the molecular testing for G551D mutation or other gating (class III) mutation on the CFTR gene; and
- (4) the result of a FEV1 measurement performed within a month prior to the date of application, if aged from 6 years or older. Note: FEV1, must be measured in an accredited pulmonary function laboratory, with documented no acute infective exacerbation at the time FEV1 is measured; and
- (5) a copy of a current medication history, including any CYP3A4 inhibitors and/or CYP3A4 inducers; and
- (6) a copy of a sweat chloride result; and
- (7) height and weight measurements at the time of application; and
- (8) a baseline measurement of the number of days of CF-related hospitalisation (including hospital-in-the home) in the previous 12 months.

Authority required

Cystic fibrosis

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must be assessed through a cystic fibrosis clinic/centre which is under the control of specialist respiratory physicians with experience and expertise in the management of cystic fibrosis. If attendance at such a unit is not possible because of geographical isolation, management (including prescribing) may be in consultation with such a unit, **AND**
- Patient must have received PBS-subsidised initial therapy with ivacaftor, given concomitantly with standard therapy, for this condition, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction, **AND**
- The treatment must be given concomitantly with standard therapy for this condition.

Population criteria:

- Patient must be aged 12 months or older.

Patients receiving PBS-subsidised ivacaftor must be registered in the Australian Cystic Fibrosis Database Registry.

Treatment must not be given to a patient who has an acute upper or lower respiratory infection, pulmonary exacerbation, or changes in therapy (including antibiotics) for pulmonary disease in the last 4 weeks prior to commencing this drug.

Patients who have an acute infective exacerbation at the time of assessment for continuing therapy may receive an additional one month's supply in order to enable the assessment to be repeated following resolution of the exacerbation.

Dosage of ivacaftor must not exceed the dose of one tablet (150 mg) or one sachet twice a week, if the patient is concomitantly receiving one of the following strong CYP3A4 drugs inhibitors: boceprevir, clarithromycin, conivaptan, indinavir, itraconazole, ketoconazole, lopinavir/ritonavir, mibefradil, nefazodone, nelfinavir, posaconazole, ritonavir, saquinavir, telaprevir, telithromycin, voriconazole. Where a patient is concomitantly receiving a strong CYP3A4 inhibitor, a single supply of 56 tablets or sachets of ivacaftor will last for 28 weeks.

Dosage of ivacaftor must not exceed the dose of one tablet (150 mg) or one sachet once daily, if the patient is concomitantly receiving one of the following moderate CYP3A4 inhibitors: amprenavir, aprepitant, atazanavir, darunavir/ritonavir, diltiazem,

erythromycin, fluconazole, fosamprenavir, imatinib, verapamil. Where a patient is concomitantly receiving a moderate CYP3A4 inhibitor, a single supply of 56 tablets or sachets of ivacaftor will last for 8 weeks.

Ivacaftor is not PBS-subsidised for this condition as a sole therapy.

Ivacaftor is not PBS-subsidised for this condition in a patient who is currently receiving one of the following CYP3A4 inducers:

Strong CYP3A4 inducers: avasimibe, carbamazepine, phenobarbital, phenytoin, rifabutin, rifampicin, St. John's wort

Moderate CYP3A4 inducers: bosentan, efavirenz, etravirine, modafinil, nafcillin

Weak CYP3A4 inducers: armodafinil, echinacea, pioglitazone, rufinamide.

The authority application must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Cystic Fibrosis Ivacaftor Authority Continuing Application Supporting Information Form; and
- (3) the result of a FEV1 measurement performed within one month prior to the date of application, if aged 6 years or older. Note: FEV1, must be measured in an accredited pulmonary function laboratory, with documented no acute infective exacerbation at the time FEV1 is measured; and
- (4) a copy of a current medication history, including any CYP3A4 inhibitors and/or CYP3A4 inducers; and
- (5) height and weight measurements at the time of application; and
- (6) a measurement of number of days of CF-related hospitalisation (including hospital in the home) in the previous 6 months.

ivacaftor 75 mg granules, 4 x 14 sachets

11109Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	22547.39	Kalydeco [VR]

ivacaftor 150 mg tablet, 56

10175M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	22547.39	Kalydeco [VR]

ivacaftor 50 mg granules, 4 x 14 sachets

11097C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	22547.39	Kalydeco [VR]

■ LUMACAFITOR + IVACAFITOR

Note Managed Access Program:

This medicine has been listed on the PBS via a Managed Access Program (MAP). The Pharmaceutical Benefits Advisory Committee (PBAC) made its recommendation on the basis of 24 weeks of data in children aged 6 - 11 years and 96 weeks of data in patients aged 12 years and over. Information about the long term benefits of this medicine will be collected and analysed under this MAP.

For more information on Managed Access Programs, please visit <http://www.pbs.gov.au/info/industry/listing/elements/pbac-meetings/pbac-outcomes/2015-03/march-2015-other-matters-managed-access-programme-framework>.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Cystic fibrosis

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a specialist respiratory physician with expertise in cystic fibrosis or in consultation with a specialist respiratory physician with expertise in cystic fibrosis if attendance is not possible due to geographic isolation, **AND**
- Must be treated in a centre with expertise in cystic fibrosis or in consultation with a centre with expertise in cystic fibrosis if attendance is not possible due to geographic isolation.

Clinical criteria:

- Patient must be homozygous for the F508del mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene, **AND**
- The treatment must be given concomitantly with standard therapy for this condition.

Population criteria:

- Patient must be 12 years of age or older.

The patient must be registered in the Australian Cystic Fibrosis Database Registry.

Treatment must not be given to a patient who has an acute upper or lower respiratory infection, pulmonary exacerbation, or changes in therapy (including antibiotics) for pulmonary disease in the last 4 weeks prior to commencing this drug.

The authority application must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Cystic Fibrosis Lumacaftor with Ivacaftor Authority Application Supporting Information Form; and

- (3) a copy of the pathology report detailing the molecular testing for the patient being homozygous for the F508del mutation on the CFTR gene; and
- (4) the result of a FEV1 measurement performed within a month prior to the date of application. Note: FEV1 must be measured in an accredited pulmonary function laboratory, with documented no acute infective exacerbation at the time FEV1 is measured; and
- (5) confirmation that the patient has either chronic sinopulmonary disease or gastrointestinal and nutritional abnormalities; and
- (6) a copy of a current medication history, including any CPY3A inhibitors and/or inducers; and
- (7) height and weight measurements at the time of application; and
- (8) a baseline measurement of the number of days of CF-related hospitalisation (including hospital-in-the home) in the previous 12 months.

Authority required

Cystic fibrosis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a specialist respiratory physician with expertise in cystic fibrosis or in consultation with a specialist respiratory physician with expertise in cystic fibrosis if attendance is not possible due to geographic isolation, **AND**
- Must be treated in a centre with expertise in cystic fibrosis or in consultation with a centre with expertise in cystic fibrosis if attendance is not possible due to geographic isolation.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be given concomitantly with standard therapy for this condition.

Population criteria:

- Patient must be 12 years of age or older.

Treatment must not be given to a patient who has an acute upper or lower respiratory infection, pulmonary exacerbation, or changes in therapy (including antibiotics) for pulmonary disease in the last 4 weeks prior to commencing this drug.

Patients who have an acute infective exacerbation at the time of assessment for continuing therapy may receive an additional one month's supply in order to enable the assessment to be repeated following resolution of the exacerbation.

The authority application must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Cystic Fibrosis Lumacaftor with Ivacaftor Continuing Authority Application Supporting Information Form; and
- (3) the result of a FEV1 measurement performed within a month prior to the date of application. Note: FEV1, must be measured in an accredited pulmonary function laboratory, with documented no acute infective exacerbation at the time FEV1 is measured; and
- (4) a copy of a current medication history, including any CYP3A inhibitors and/or inducers; and
- (5) height and weight measurements at the time of application; and
- (6) the number of days of CF-related hospitalisation (including hospital-in-the home) in the previous 6 months.

lumacaftor 200 mg + ivacaftor 125 mg tablet, 112

11463H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	18797.39	Orkambi [VR]

■ LUMACFTOR + IVACFTOR**Note Managed Access Program:**

This medicine has been listed on the PBS via a Managed Access Program (MAP). The Pharmaceutical Benefits Advisory Committee (PBAC) made its recommendation on the basis of 24 weeks of data in children aged 6 - 11 years and 96 weeks of data in patients aged 12 years and over. Information about the long term benefits of this medicine will be collected and analysed under this MAP.

For more information on Managed Access Programs, please visit <http://www.pbs.gov.au/info/industry/listing/elements/pbac-meetings/pbac-outcomes/2015-03/march-2015-other-matters-managed-access-programme-framework>.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Cystic fibrosis

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a specialist respiratory physician with expertise in cystic fibrosis or in consultation with a specialist respiratory physician with expertise in cystic fibrosis if attendance is not possible due to geographic isolation, **AND**
- Must be treated in a centre with expertise in cystic fibrosis or in consultation with a centre with expertise in cystic fibrosis if attendance is not possible due to geographic isolation.

Clinical criteria:

- Patient must be homozygous for the F508del mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene, **AND**
- The treatment must be given concomitantly with standard therapy for this condition.

Population criteria:

- Patient must be aged between 6 and 11 years inclusive.

The patient must be registered in the Australian Cystic Fibrosis Database Registry.

Treatment must not be given to a patient who has an acute upper or lower respiratory infection, pulmonary exacerbation, or changes in therapy (including antibiotics) for pulmonary disease in the last 4 weeks prior to commencing this drug.

The authority application must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Cystic Fibrosis Lumacaftor with Ivacaftor Authority Application Supporting Information Form; and
- (3) a copy of the pathology report detailing the molecular testing for the patient being homozygous for the F508del mutation on the CFTR gene; and
- (4) the result of a FEV1 measurement performed within a month prior to the date of application. Note: FEV1 must be measured in an accredited pulmonary function laboratory, with documented no acute infective exacerbation at the time FEV1 is measured; and
- (5) confirmation that the patient has either chronic sinopulmonary disease or gastrointestinal and nutritional abnormalities; and
- (6) a copy of a current medication history, including any CPY3A inhibitors and/or inducers; and
- (7) height and weight measurements at the time of application; and
- (8) a baseline measurement of the number of days of CF-related hospitalisation (including hospital-in-the home) in the previous 12 months.

Authority required

Cystic fibrosis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a specialist respiratory physician with expertise in cystic fibrosis or in consultation with a specialist respiratory physician with expertise in cystic fibrosis if attendance is not possible due to geographic isolation, **AND**
- Must be treated in a centre with expertise in cystic fibrosis or in consultation with a centre with expertise in cystic fibrosis if attendance is not possible due to geographic isolation.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be given concomitantly with standard therapy for this condition.

Population criteria:

- Patient must be aged between 6 and 11 years inclusive.

Treatment must not be given to a patient who has an acute upper or lower respiratory infection, pulmonary exacerbation, or changes in therapy (including antibiotics) for pulmonary disease in the last 4 weeks prior to commencing this drug.

Patients who have an acute infective exacerbation at the time of assessment for continuing therapy may receive an additional one month's supply in order to enable the assessment to be repeated following resolution of the exacerbation.

The authority application must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Cystic Fibrosis Lumacaftor with Ivacaftor Continuing Authority Application Supporting Information Form; and
- (3) the result of a FEV1 measurement performed within a month prior to the date of application. Note: FEV1, must be measured in an accredited pulmonary function laboratory, with documented no acute infective exacerbation at the time FEV1 is measured; and
- (4) a copy of a current medication history, including any CYP3A inhibitors and/or inducers; and
- (5) height and weight measurements at the time of application; and
- (6) the number of days of CF-related hospitalisation (including hospital-in-the home) in the previous 6 months.

lumacaftor 100 mg + ivacaftor 125 mg tablet, 112

11464J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	18797.39	Orkambi [VR]

VARIOUS

ALL OTHER THERAPEUTIC PRODUCTS

ALL OTHER THERAPEUTIC PRODUCTS

Iron chelating agents

DEFERASIROX

Note Special Pricing Arrangements apply.

Authority required

Chronic iron overload

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must be transfusion dependent, **AND**

- Patient must not have a malignant disorder of erythropoiesis.

Authority required

Chronic iron overload

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must not be transfusion dependent, **AND**
- The condition must be thalassaemia.

deferasirox 180 mg tablet, 30

11546Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	6	5	..	*2615.05	Jadenu [NM]

deferasirox 360 mg tablet, 30

11496C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	6	5	..	*5182.75	Jadenu [NM]

deferasirox 90 mg tablet, 30

11545P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	6	5	..	*1331.29	Jadenu [NM]

▪ **DEFERASIROX****Note** Special Pricing Arrangements apply.**Authority required (STREAMLINED)****9302**

Chronic iron overload

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must be transfusion dependent, **AND**
- Patient must not have a malignant disorder of erythropoiesis, **AND**
- Patient must have previously received PBS-subsidised therapy with deferasirox for this condition.

Authority required (STREAMLINED)**9222**

Chronic iron overload

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must not be transfusion dependent, **AND**
- The condition must be thalassaemia, **AND**
- Patient must have previously received PBS-subsidised therapy with deferasirox for this condition.

Authority required (STREAMLINED)**9258**

Chronic iron overload

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must be red blood cell transfusion dependent, **AND**
- Patient must have a malignant disorder of haemopoiesis, **AND**
- Patient must have previously received PBS-subsidised therapy with deferasirox for this condition.

Note Interruption of treatment should be considered if serum ferritin levels fall consistently below 500 microgram/mL.**deferasirox 180 mg tablet, 30**

11510T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	6	2	..	*2615.05	Jadenu [NM]

deferasirox 360 mg tablet, 30

11547R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	6	2	..	*5182.75	Jadenu [NM]

deferasirox 90 mg tablet, 30

11548T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	6	2	..	*1331.29	Jadenu [NM]

▪ **DEFERASIROX****Note** Special Pricing Arrangements apply.**Note** A patient's median life expectancy is determined by the severity of their underlying disease.**Note** Patients with underlying myelodysplastic syndrome are considered to have a median life expectancy exceeding five years if they are classified as:

- low risk according to the International Prognostic Scoring System (IPSS); or

- very low and low risk according to the Revised International Prognostic Scoring System (IPSS-R); or
- very low and low risk according to the WHO classification based Prognostic Scoring System (WPSS).

Note Patients with underlying myelofibrosis have a median life expectancy exceeding five years if they are classified as:

- low or intermediate risk according to the International Prognostic Scoring System (IPSS); or
- low or intermediate-1 risk according to Dynamic International Prognostic Scoring System (DIPSS).

Authority required

Chronic iron overload

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must be red blood cell transfusion dependent, **AND**
- Patient must have a serum ferritin level of greater than 1000 microgram/L, **AND**
- Patient must have a malignant disorder of haemopoiesis, **AND**
- Patient must have a median life expectancy exceeding five years.

deferasirox 180 mg tablet, 30

11557G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	6	2	..	*2615.05	Jadenu [NM]

deferasirox 360 mg tablet, 30

11511W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	6	2	..	*5182.75	Jadenu [NM]

deferasirox 90 mg tablet, 30

11558H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	6	2	..	*1331.29	Jadenu [NM]

▪ DEFERIPRONE

Authority required (STREAMLINED)

9286

Iron overload

Clinical criteria:

- Patient must have thalassaemia major, **AND**
- Patient must be unable to take desferrioxamine therapy.

Authority required (STREAMLINED)

9228

Iron overload

Clinical criteria:

- Patient must have thalassaemia major, **AND**
- Patient must be one in whom desferrioxamine therapy has proven ineffective.

deferiprone 500 mg tablet, 100

6416Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	3	5	..	*1203.07	Ferriprox [TX]

deferiprone 100 mg/mL oral liquid, 250 mL

9638G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	5	5	..	*1008.99	Ferriprox [TX]

▪ DEFERIPRONE

Authority required (STREAMLINED)

9623

Iron overload

Clinical criteria:

- Patient must have thalassaemia major, **AND**
- Patient must be unable to take desferrioxamine therapy.

Authority required (STREAMLINED)

9590

Iron overload

Clinical criteria:

- Patient must have thalassaemia major, **AND**
- Patient must be one in whom desferrioxamine therapy has proven ineffective.

deferiprone 1 g tablet, 50

11724C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	6	5	..	*2358.79	Ferriprox [TX]

■ DESFERRIOXAMINE

Authority required

Disorders of erythropoiesis

Clinical criteria:

- The condition must be associated with treatment-related chronic iron overload.

desferrioxamine mesilate 2 g injection, 1 vial

6270B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	60	5	..	*1915.99	Hospira Pty Limited [PF]

desferrioxamine mesilate 500 mg injection, 10 vials

6113R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	40	5	..	*5602.59	Hospira Pty Limited [PF]

Drugs for treatment of hyperkalemia and hyperphosphatemia

■ LANTHANUM

Authority required

Hyperphosphataemia

Treatment Phase: Initiation and stabilisation

Clinical criteria:

- The condition must not be adequately controlled by calcium, **AND**
- Patient must have a serum phosphate of greater than 1.6 mmol per L at the commencement of therapy; OR
- The condition must be where a serum calcium times phosphate product is greater than 4 at the commencement of therapy, **AND**
- The treatment must not be used in combination with any other non-calcium phosphate binding agents.

Treatment criteria:

- Patient must be undergoing dialysis for chronic kidney disease.

lanthanum 500 mg chewable tablet, 2 x 45

9635D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*524.65	Fosrenol [ZI]

lanthanum 1 g chewable tablet, 6 x 15

9637F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*886.73	Fosrenol [ZI]

lanthanum 750 mg chewable tablet, 6 x 15

9636E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*788.47	Fosrenol [ZI]

■ SEVELAMER

Authority required

Hyperphosphataemia

Treatment Phase: Initiation and stabilisation

Clinical criteria:

- The condition must not be adequately controlled by calcium, **AND**
- Patient must have a serum phosphate of greater than 1.6 mmol per L at the commencement of therapy; OR
- The condition must be where a serum calcium times phosphate product is greater than 4 at the commencement of therapy, **AND**
- The treatment must not be used in combination with any other non-calcium phosphate binding agents.

Treatment criteria:

- Patient must be undergoing dialysis for chronic kidney disease.

sevelamer hydrochloride 800 mg tablet, 180

9620H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*558.69	Renagel [GZ]

■ SUCROFERRIC OXYHYDROXIDE

Authority required

Hyperphosphataemia

Treatment Phase: Initiation and stabilisation

Clinical criteria:

- The condition must not be adequately controlled by calcium, **AND**
- Patient must have a serum phosphate of greater than 1.6 mmol per L at the commencement of therapy; OR
- The condition must be where a serum calcium times phosphate product is greater than 4 at the commencement of therapy, **AND**
- The treatment must not be used in combination with any other non-calcium phosphate binding agents.

Treatment criteria:

- Patient must be undergoing dialysis for chronic kidney disease.

sucroferric oxyhydroxide 2.5 g (iron 500 mg) chewable tablet, 90

10230K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*790.99	Velphoro [VL]

Highly Specialised Drugs Program (Public Hospital)

ALIMENTARY TRACT AND METABOLISM.....	312
OTHER ALIMENTARY TRACT AND METABOLISM PRODUCTS.....	312
OTHER ALIMENTARY TRACT AND METABOLISM PRODUCTS	312
BLOOD AND BLOOD FORMING ORGANS	315
ANTIHEMORRHAGICS.....	315
VITAMIN K AND OTHER HEMOSTATICS	315
ANTIANEMIC PREPARATIONS	321
OTHER ANTIANEMIC PREPARATIONS	321
CARDIOVASCULAR SYSTEM.....	325
ANTIHYPERTENSIVES	325
OTHER ANTIHYPERTENSIVES	325
SYSTEMIC HORMONAL PREPARATIONS, EXCL. SEX HORMONES AND INSULINS.....	372
PITUITARY AND HYPOTHALAMIC HORMONES AND ANALOGUES	372
ANTERIOR PITUITARY LOBE HORMONES AND ANALOGUES	372
HYPOTHALAMIC HORMONES.....	374
ANTIINFECTIVES FOR SYSTEMIC USE	378
ANTIBACTERIALS FOR SYSTEMIC USE.....	378
MACROLIDES, LINCOSAMIDES AND STREPTOGRAMINS.....	378
ANTIMYCOBACTERIALS	378
DRUGS FOR TREATMENT OF TUBERCULOSIS.....	378
ANTIVIRALS FOR SYSTEMIC USE	379
DIRECT ACTING ANTIVIRALS	379
ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS	384
ANTINEOPLASTIC AGENTS.....	384
ANTIMETABOLITES.....	384
CYTOTOXIC ANTIBIOTICS AND RELATED SUBSTANCES	386
OTHER ANTINEOPLASTIC AGENTS.....	386
IMMUNOSTIMULANTS	400
IMMUNOSTIMULANTS	400
IMMUNOSUPPRESSANTS.....	407
IMMUNOSUPPRESSANTS	407

MUSCULO-SKELETAL SYSTEM.....	558
MUSCLE RELAXANTS	558
MUSCLE RELAXANTS, CENTRALLY ACTING AGENTS	558
DRUGS FOR TREATMENT OF BONE DISEASES	559
DRUGS AFFECTING BONE STRUCTURE AND MINERALIZATION.....	559
OTHER DRUGS FOR DISORDERS OF THE MUSCULO-SKELETAL SYSTEM.....	560
OTHER DRUGS FOR DISORDERS OF THE MUSCULO-SKELETAL SYSTEM	560
NERVOUS SYSTEM.....	563
ANTI-PARKINSON DRUGS	563
DOPAMINERGIC AGENTS	563
PSYCHOLEPTICS.....	564
ANTIPSYCHOTICS.....	564
RESPIRATORY SYSTEM.....	565
DRUGS FOR OBSTRUCTIVE AIRWAY DISEASES	565
OTHER SYSTEMIC DRUGS FOR OBSTRUCTIVE AIRWAY DISEASES	565
COUGH AND COLD PREPARATIONS.....	582
EXPECTORANTS, EXCL. COMBINATIONS WITH COUGH SUPPRESSANTS	582
OTHER RESPIRATORY SYSTEM PRODUCTS	584
OTHER RESPIRATORY SYSTEM PRODUCTS	584
VARIOUS	588
ALL OTHER THERAPEUTIC PRODUCTS	588
ALL OTHER THERAPEUTIC PRODUCTS.....	588

ALIMENTARY TRACT AND METABOLISM

OTHER ALIMENTARY TRACT AND METABOLISM PRODUCTS

OTHER ALIMENTARY TRACT AND METABOLISM PRODUCTS

Various alimentary tract and metabolism products

■ TEDUGLUTIDE

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note A patient may only qualify for PBS-subsidised treatment under this restriction once in a lifetime.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Type III Short bowel syndrome with intestinal failure

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a gastroenterologist; OR
- Must be treated by a specialist within a multidisciplinary intestinal rehabilitation unit.

Clinical criteria:

- Patient must have short bowel syndrome with intestinal failure following major surgery, **AND**
- Patient must have a history of dependence on parenteral support for at least 12 months, **AND**
- Patient must have received a stable parenteral support regimen for at least 3 days per week in the previous 4 weeks, **AND**
- Patient must not have active gastrointestinal malignancy or history of gastrointestinal malignancy within the last 5 years, **AND**
- The treatment must not exceed 12 months under this restriction, **AND**
- Patient must not have previously received PBS-subsidised treatment with this drug for this condition.

Baseline is the mean number of days of parenteral support per week over the four weeks immediately prior to initiating treatment with teduglutide under the PBS initial treatment restriction or four weeks immediately prior to initiating treatment with non-PBS subsidised teduglutide for grandfathered patients.

A stable parenteral support regimen is defined as a minimum of 3 days of parenteral support (parenteral nutrition with or without IV fluids) per week for 4 consecutive weeks to meet caloric, fluid or electrolyte needs.

Baseline number of days of parenteral support should be documented in the patient's medical records.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Short bowel syndrome with intestinal failure form; and
- (3) details of baseline mean number of days on parenteral support per week for 4 consecutive weeks immediately preceding this application; and
- (4) documented duration in months of prior dependence on parenteral support.

teduglutide 5 mg injection [28 vials] (&) inert substance diluent [28 x 0.5 mL syringes], 1 pack

11793Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	11	..	21840.00	Revestive [ZI]

■ TEDUGLUTIDE

Note Special Pricing Arrangements apply.

Authority required

Type III Short bowel syndrome with intestinal failure

Treatment Phase: Initial treatment - Grandfathered patients

Treatment criteria:

- Must be treated by a gastroenterologist; OR
- Must be treated by a specialist within a multidisciplinary intestinal rehabilitation unit.

Clinical criteria:

- Patient must have previously received non-PBS subsidised treatment with this drug for this condition prior to 1 October 2019, **AND**
- Patient must have short bowel syndrome with intestinal failure following major surgery, **AND**
- Patient must have had a history of dependence on parenteral support for at least 12 months prior to initiating non-PBS subsidised treatment with this drug for this condition, **AND**

- Patient must have received a stable parenteral support regimen for at least 3 days per week in the 4 weeks prior to initiating non-PBS subsidised treatment with this drug for this condition, **AND**
- Patient must not have active gastrointestinal malignancy or history of gastrointestinal malignancy within the last 5 years, **AND**
- Patient must have achieved a treatment response if the patient has been on non-PBS subsidised therapy with this drug for more than 12 months.

Baseline is the mean number of days of parenteral support per week over the 4 weeks immediately prior to initiating treatment with non-PBS subsidised teduglutide for grandfathered patients.

A stable parenteral support regimen is defined as a minimum of 3 days of parenteral support (parenteral nutrition with or without IV fluids) per week for 4 consecutive weeks to meet caloric, fluid or electrolyte needs.

A patient has met the criteria for treatment response when there is a reduction in the mean number of days of parenteral support of at least 1 day per week since initiating non-PBS subsidised treatment, or where a patient has completely ceased treatment with parenteral support for a period of at least 4 consecutive weeks prior to application for PBS-subsidised treatment.

The number of days of parenteral support is calculated as the mean number of days in which any parenteral support is required (parenteral nutrition with or without IV fluids) per week to meet caloric, fluid or electrolyte needs between commencement of non-PBS subsidised teduglutide and application for PBS-subsidised treatment.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Short bowel syndrome with intestinal failure Grandfather PBS Authority Application - Supporting Information Form; and
- (3) details of non-PBS subsidised teduglutide treatment start date; and
- (4) details of the number of days on parenteral support per week for 4 consecutive weeks prior to initiating non-PBS subsidised therapy; and
- (5) documented duration in months of dependence on parenteral support prior to initiating non-PBS subsidised treatment; and
- (6) details of response to teduglutide treatment if patient has received 12 or more months of non-PBS subsidised treatment.

A patient may qualify for PBS-subsidised treatment under this restriction once only.

For patients who have been on this drug for less than 12 months, the maximum number of repeats that will be approved will be for an amount equivalent to an initial 12 month supply of PBS and non-PBS subsidised treatment.

For patients who have been on this drug for more than 12 months, a maximum of 5 repeats will be approved.

For continuing PBS-subsidised treatment, a Grandfathered patient must qualify under the First continuing treatment criteria.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Type III Short bowel syndrome with intestinal failure

Treatment Phase: Initial treatment or initial grandfather treatment - balance of supply

Treatment criteria:

- Must be treated by a gastroenterologist; OR
- Must be treated by a specialist within a multidisciplinary intestinal rehabilitation unit.

Clinical criteria:

- Patient must have previously received PBS-subsidised initial treatment with this drug for this condition; OR
- Patient must have received PBS-subsidised treatment with this drug for this condition as a grandfathered patient, **AND**
- Patient must have received insufficient therapy with this drug under the initial or grandfather treatment restriction to complete the maximum duration of 12 months of initial treatment, **AND**
- The treatment must provide no more than the balance of up to 12 months of treatment.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

teduglutide 5 mg injection [28 vials] (&) inert substance diluent [28 x 0.5 mL syringes], 1 pack

11808L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	21840.00	Revestive [ZI]

▪ TEDUGLUTIDE

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Type III Short bowel syndrome with intestinal failure

Treatment Phase: First continuing treatment

Treatment criteria:

- Must be treated by a gastroenterologist; OR
- Must be treated by a specialist within a multidisciplinary intestinal rehabilitation unit.

Clinical criteria:

- Patient must have previously received PBS-subsidised initial treatment with this drug for this condition; OR
- Patient must have received PBS-subsidised treatment with this drug for this condition as a grandfathered patient, **AND**
- Patient must have a reduction in parenteral support frequency of at least one day per week compared to the mean number of days per week at baseline.

Baseline is the mean number of days of parenteral support per week over the four weeks immediately prior to initiating treatment with teduglutide under the PBS initial treatment restriction or four weeks immediately prior to initiating treatment with non-PBS subsidised teduglutide for grandfathered patients.

The current mean number of days of parenteral support is calculated as the mean number of days in which any parenteral support is required (parenteral nutrition with or without IV fluids) per week to meet caloric, fluid or electrolyte needs over the immediately preceding 4 week treatment period

Treatment failure

For applications for first continuing treatment, failure of treatment is defined as no change compared to baseline in the mean number of days per week in parenteral support (parenteral nutrition with or without IV fluids) to meet caloric, fluid or electrolyte needs.

Patients who experience failure of treatment must permanently discontinue treatment.

Current mean number of days of parenteral support should be documented in the patient's medical records.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Short bowel syndrome with intestinal failure Form; and
- (3) details of the baseline mean number of days reduction of parenteral support (parenteral nutrition with or without IV fluids) per week to meet caloric, fluid or electrolyte needs; and
- (4) the current mean number of days per week of parenteral support over the preceding 4 week period.

Note A patient may only qualify for PBS-subsidised treatment under this restriction once in a lifetime.

Authority required

Type III Short bowel syndrome with intestinal failure

Treatment Phase: Subsequent continuing treatment

Treatment criteria:

- Must be treated by a gastroenterologist; OR
- Must be treated by a specialist within a multidisciplinary intestinal rehabilitation unit.

Clinical criteria:

- Patient must have received PBS-subsidised first-continuing treatment with this drug for this condition; OR
- Patient must have received PBS-subsidised recommencement of treatment with this drug for this condition following a trial cessation period, **AND**
- Patient must have achieved a treatment response in the preceding treatment period; OR
- Patient must not have previously experienced a failure to respond to treatment with this drug for this condition.

Treatment response

For applications for subsequent continuing treatment, treatment response is when there was a reduction in the mean number of days of parenteral support of at least 1 day per week since the last assessment for PBS-subsidised treatment, OR where a patient has completely ceased treatment with parenteral support for a period of at least 4 consecutive weeks.

The current mean number of days of parenteral support is calculated as the mean number of days in which any parenteral support is required (parenteral nutrition with or without IV fluids) per week to meet caloric, fluid or electrolyte needs over the immediately preceding 4 week treatment period

Treatment failure

For applications for subsequent continuing treatment, failure of treatment is defined as an increase in the mean number of days per week of parenteral support requirements of at least 1 day per week over the preceding 4 week period compared to the last assessment for PBS-subsidised treatment of parenteral support (parenteral nutrition with or without IV fluids) to meet caloric, fluid or electrolyte needs.

Patients who experience failure of treatment must permanently discontinue treatment.

Treatment stability

Patients who neither demonstrate a treatment response nor a treatment failure since the last assessment for PBS-subsidised treatment are considered to have a stable parenteral support regimen, defined as the same mean number of days of parenteral support (parenteral nutrition with or without IV fluids) per week to meet caloric, fluid or electrolyte needs over the 4 weeks preceding treatment period, where the number of days is greater than zero and the mean number of days of parenteral support is less than baseline. Patients with a stable parenteral support regimen over 6 months must undertake a trial cessation period. Patients who have re-commenced after a trial cessation period are exempt from further trial cessation.

Trial cessation period

Patients who demonstrate a stable frequency of mean days per week of parenteral support in a 6-month period commencing after the initial 12 months of treatment with this drug for this condition are required to undertake a trial of treatment cessation. Patients who have re-commenced after a trial cessation period are exempt from further trial cessation.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Short bowel syndrome with intestinal failure Form; and
- (3) details of the mean number of days reduction of parenteral support (parenteral nutrition with or without IV fluids) per week to meet caloric, fluid or electrolyte needs over the preceding treatment period **or** confirmation the patient has had 4 consecutive weeks without parenteral support; and
- (4) details of completion of a trial cessation period (if applicable); and
- (5) the current mean number of days per week of parenteral support over the preceding 4 week period.

Authority required

Type III Short bowel syndrome with intestinal failure

Treatment Phase: Recommencement of treatment

Treatment criteria:

- Must be treated by a gastroenterologist; **OR**
- Must be treated by a specialist within a multidisciplinary intestinal rehabilitation unit.

Clinical criteria:

- Patient must have received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have undertaken a trial cessation period due to experiencing a stable parenteral support regimen in the first continuing or subsequent continuing treatment phase, and not due to a treatment failure, **AND**
- Patient must have experienced deterioration during a trial cessation period.

Trial cessation period

Patients who demonstrate a stable frequency of mean days per week of parenteral support in a 6-month period commencing after the initial 12 months of treatment with this drug for this condition are required to undertake a trial of treatment cessation. Patients who have re-commenced after a trial cessation period are exempt from further trial cessation.

Deterioration during the trial cessation period includes an increase in parenteral support frequency of more than or equal to one day per week from the pre-cessation level, or other clinical parameters suggestive of deterioration including changes in renal function or urinary sodium levels or changes in body weight.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Short bowel syndrome with intestinal failure Form; and
- (3) details of the reason for recommencement after trial cessation; and
- (4) the current mean number of days per week of parenteral support over the preceding 4 week period

teduglutide 5 mg injection [28 vials] (&) inert substance diluent [28 x 0.5 mL syringes], 1 pack

11794R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	21840.00	Revestive [ZI]

■ BLOOD AND BLOOD FORMING ORGANS

■ ANTIHEMORRHAGICS

VITAMIN K AND OTHER HEMOSTATICS

Other systemic hemostatics

■ ELTROMBOPAG

Note Special Pricing Arrangements apply.

Note No applications for increased repeats will be authorised.

Authority required

Severe thrombocytopenia

Treatment Phase: Initial treatment 1 - New patient

Clinical criteria:

- The condition must be severe chronic immune (idiopathic) thrombocytopenic purpura (ITP), **AND**
- Patient must have had a splenectomy, **AND**
- Patient must have failed to achieve an adequate response to, or be intolerant to, corticosteroid therapy following the splenectomy, **AND**
- Patient must have failed to achieve an adequate response to, or be intolerant to, immunoglobulin therapy following the splenectomy, **AND**
- The treatment must be the sole PBS-subsidised thrombopoietin receptor agonist (TRA) for this condition.

Population criteria:

- Patient must be an adult.

The following criteria indicate failure to achieve an adequate response and must be demonstrated at the time of initial application;

- (a) a platelet count of less than or equal to 20,000 million per L; **OR**
- (b) a platelet count of 20,000 million to 30,000 million per L, where the patient is experiencing significant bleeding or has a history of significant bleeding in this platelet range.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form,
- (2) a signed patient acknowledgement,
- (3) a completed Idiopathic Thrombocytopenic Purpura Initial PBS Authority Application - Supporting Information Form,
- (4) a copy of a full blood count pathology report supporting the diagnosis of ITP, and
- (5) where the application is sought on the basis of a medical contraindication to surgery, a signed and dated letter from the clinician making this assessment which includes the date upon which the patient was assessed for surgery and the clinical grounds upon which surgery is contraindicated.

The full blood count must be no more than 1 month old at the time of application.

A maximum of 24 weeks of treatment with this drug will be authorised under this criterion.

Note Eltrombopag is not PBS-subsidised as an alternative to splenectomy.

Note Patients will be able to trial either eltrombopag or romiplostim within the initial 24 weeks treatment period. Patients who fail to demonstrate a response to treatment with eltrombopag and/or romiplostim under the initial restriction will not be eligible to receive further PBS-subsidised treatment with either of these drugs.

Note Authority approval for sufficient therapy to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written application for authority approval for sufficient therapy to complete a maximum of 24 weeks of treatment should be forwarded to:

Department of Human Services
Prior Written Approval of Complex Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001

Authority required

Severe thrombocytopenia

Treatment Phase: Initial treatment 2 - New patient

Clinical criteria:

- The condition must be severe chronic immune (idiopathic) thrombocytopenic purpura (ITP), **AND**
- Patient must not have had a splenectomy, **AND**
- Patient must have failed to achieve an adequate response to, or be intolerant to, corticosteroid therapy at a dose equivalent to 0.5-2 mg/kg/day of prednisone for at least 4-6 weeks, **AND**
- Patient must have failed to achieve an adequate response to, or be intolerant to, immunoglobulin therapy, **AND**
- Patient must be unsuitable for splenectomy due to medical reasons, **AND**
- The treatment must be the sole PBS-subsidised thrombopoietin receptor agonist (TRA) for this condition.

Population criteria:

- Patient must be an adult.

The following criteria indicate failure to achieve an adequate response and must be demonstrated at the time of initial application;

- (a) a platelet count of less than or equal to 20,000 million per L; OR
- (b) a platelet count of 20,000 million to 30,000 million per L, where the patient is experiencing significant bleeding or has a history of significant bleeding in this platelet range.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form,
- (2) a signed patient acknowledgement,
- (3) a completed Idiopathic Thrombocytopenic Purpura Initial PBS Authority Application - Supporting Information Form,
- (4) a copy of a full blood count pathology report supporting the diagnosis of ITP, and
- (5) where the application is sought on the basis of a medical contraindication to surgery, a signed and dated letter from the clinician making this assessment which includes the date upon which the patient was assessed for surgery and the clinical grounds upon which surgery is contraindicated.

The full blood count must be no more than 1 month old at the time of application.

A maximum of 24 weeks of treatment with this drug will be authorised under this criterion.

Note Eltrombopag is not PBS-subsidised as an alternative to splenectomy.

Note Patients will be able to trial either eltrombopag or romiplostim within the initial 24 weeks treatment period. Patients who fail to demonstrate a response to treatment with eltrombopag and/or romiplostim under the initial restriction will not be eligible to receive further PBS-subsidised treatment with either of these drugs.

Note Authority approval for sufficient therapy to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written application for authority approval for sufficient therapy to complete a maximum of 24 weeks of treatment should be forwarded to:

Department of Human Services
Prior Written Approval of Complex Drugs
Reply Paid 9826
GPO Box 9826
HOBART TAS 7001

Authority required

Severe thrombocytopenia

Treatment Phase: First Continuing treatment or Re-initiation of interrupted treatment

Clinical criteria:

- The condition must be severe chronic immune (idiopathic) thrombocytopenic purpura (ITP), **AND**
- Patient must have previously received PBS-subsidised initial treatment with this drug for this condition, **AND**
- Patient must have demonstrated a sustained platelet response to PBS-subsidised treatment with this drug for this condition under the Initial treatment restriction, **AND**
- The treatment must be the sole PBS-subsidised thrombopoietin receptor agonist (TRA) for this condition.

Population criteria:

- Patient must be an adult.

For the purposes of this restriction, a sustained platelet response is defined as:

(a) use of rescue medication (corticosteroids or immunoglobulins) on no more than one occasion during the initial period of PBS-subsidised treatment with this drug,

AND either of the following:

(b) a platelet count greater than or equal to 50,000 million per L on at least four (4) occasions, each at least one week apart;
OR

(c) a platelet count greater than 30,000 million per L and which is double the baseline (pre-treatment) platelet count on at least four (4) occasions, each at least one week apart.

Applications for the First continuing PBS-subsidised treatment or Re-initiation of interrupted PBS-subsidised treatment must be made in writing and must include:

- (1) a completed authority prescription form, and
- (2) a completed Idiopathic Thrombocytopenic Purpura Continuing PBS Authority Application - Supporting Information Form , and

(3) copies of the platelet count pathology reports (unless previously provided for patients re-initiating therapy).

The platelet count must be no more than one month old at the time of application.

A maximum of 24 weeks of treatment with this drug will be authorised under this criterion.

Note Eltrombopag is not PBS-subsidised as an alternative to splenectomy.

Note Patients will be able to trial either eltrombopag or romiplostim within the initial 24 weeks treatment period. Patients who fail to demonstrate a response to treatment with eltrombopag and/or romiplostim under the initial restriction will not be eligible to receive further PBS-subsidised treatment with either of these drugs.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe thrombocytopenia

Treatment Phase: Second or subsequent Continuing treatment

Clinical criteria:

- The condition must be severe chronic immune (idiopathic) thrombocytopenic purpura (ITP), **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have demonstrated a continuing response to treatment with this drug, **AND**
- The treatment must be the sole PBS-subsidised thrombopoietin receptor agonist (TRA) for this condition.

Population criteria:

- Patient must be an adult.

For the purpose of this restriction, a continuing response to treatment with drug is defined as:

(a) use of rescue medication (corticosteroids or immunoglobulins) on no more than one occasion during the most recent 24 week period of PBS-subsidised treatment with this drug

AND either of the following:

(b) a platelet count greater than or equal to 50,000 million per L

OR

(c) a platelet count greater than 30,000 million per L and which is double the baseline platelet count.

The platelet count must be no more than one month old at the time of application.

Authority applications for second and subsequent periods of continuing therapy may be made by telephone

Note Eltrombopag is not PBS-subsidised as an alternative to splenectomy.

Note Authority applications for second and subsequent continuing treatment may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note Patients will be able to trial either eltrombopag or romiplostim within the initial 24 weeks treatment period. Patients who fail to demonstrate a response to treatment with eltrombopag and/or romiplostim under the initial restriction will not be eligible to receive further PBS-subsidised treatment with either of these drugs.

Authority required

Severe thrombocytopenia

Treatment Phase: Initial 1, Initial 2, First Continuing treatment or Re-initiation of interrupted treatment, and Second and Subsequent Continuing treatment - balance of supply

Clinical criteria:

- The condition must be severe chronic immune (idiopathic) thrombocytopenic purpura (ITP), **AND**
- The treatment must be the sole PBS-subsidised thrombopoietin receptor agonist (TRA) for this condition, **AND**
- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the First Continuing treatment or Re-initiation of interrupted treatment restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Second and subsequent Continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restriction.

Population criteria:

- Patient must be an adult.

Note Authority approval for sufficient therapy to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

eltrombopag 50 mg tablet, 28

5826P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	2872.80	Revolade [NV]

eltrombopag 25 mg tablet, 28

5825N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	1436.40	Revolade [NV]

▪ ROMIPILOSTIM

Authority required

Severe thrombocytopenia

Treatment Phase: Initial treatment 1 - New patient

Clinical criteria:

- The condition must be severe chronic immune (idiopathic) thrombocytopenic purpura (ITP), **AND**
- Patient must have had a splenectomy, **AND**
- Patient must have failed to achieve an adequate response to, or be intolerant to, corticosteroid therapy following the splenectomy, **AND**
- Patient must have failed to achieve an adequate response to, or be intolerant to, immunoglobulin therapy following the splenectomy, **AND**
- The treatment must be the sole PBS-subsidised thrombopoietin receptor agonist (TRA) for this condition.

Population criteria:

- Patient must be an adult.

The following criteria indicate failure to achieve an adequate response and must be demonstrated at the time of initial application;

(a) a platelet count of less than or equal to 20,000 million per L; OR

(b) a platelet count of 20,000 million to 30,000 million per L, where the patient is experiencing significant bleeding or has a history of significant bleeding in this platelet range.

At the time of the written authority application, medical practitioners should request the appropriate quantity of vials of appropriate strength to provide sufficient drug for a single treatment at a dose of 1 microgram/kg. Up to 1 repeat may be requested with the initial written application.

Subsequently during the initial period of dose titration, authority applications for a single dose and up to 1 repeat may be requested by telephone. The dose (microgram/kg/week) must be provided at the time of application.

Once a patient's dose has been stable for a period of 4 weeks, authority approvals for sufficient vials of appropriate strength based on the weight of the patient and dose (microgram/kg/week) for up to 4 weeks of treatment and up to 4 repeats may be granted, as long as the total period of treatment authorised under this restriction does not exceed 24 weeks.

Authority approval will not be given for doses higher than 10 micrograms/kg/week

The authority application must be made in writing and must include:

- (1) a completed authority prescription form,
- (2) a signed patient acknowledgement,
- (3) a completed Idiopathic Thrombocytopenic Purpura Initial PBS Authority Application - Supporting Information Form,
- (4) a copy of a full blood count pathology report supporting the diagnosis of ITP, and
- (5) where the application is sought on the basis of a medical contraindication to surgery, a signed and dated letter from the clinician making this assessment which includes the date upon which the patient was assessed for surgery and the clinical grounds upon which surgery is contraindicated.

The full blood count must be no more than 1 month old at the time of application.

Note Romiplostim is not PBS-subsidised as an alternative to splenectomy.

Note Patients will be able to trial either eltrombopag or romiplostim within the initial 24 weeks treatment period. Patients who fail to demonstrate a response to treatment with eltrombopag and/or romiplostim under the initial restriction will not be eligible to receive further PBS-subsidised treatment with either of these drugs.

Note Any queries concerning the arrangements to prescribe this drug may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au
Written applications for authority to prescribe this drug should be forwarded to:
Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe thrombocytopenia

Treatment Phase: Initial treatment 2 - New patient

Clinical criteria:

- The condition must be severe chronic immune (idiopathic) thrombocytopenic purpura (ITP), **AND**
- Patient must not have had a splenectomy, **AND**
- Patient must have failed to achieve an adequate response to, or be intolerant to, corticosteroid therapy at a dose equivalent to 0.5-2 mg/kg/day of prednisone for at least 4-6 weeks, **AND**
- Patient must have failed to achieve an adequate response to, or be intolerant to, immunoglobulin therapy, **AND**
- Patient must be unsuitable for splenectomy due to medical reasons, **AND**
- The treatment must be the sole PBS-subsidised thrombopoietin receptor agonist (TRA) for this condition.

Population criteria:

- Patient must be an adult.

The following criteria indicate failure to achieve an adequate response and must be demonstrated at the time of initial application;

(a) a platelet count of less than or equal to 20,000 million per L; OR

(b) a platelet count of 20,000 million to 30,000 million per L, where the patient is experiencing significant bleeding or has a history of significant bleeding in this platelet range.

At the time of the written authority application, medical practitioners should request the appropriate quantity of vials of appropriate strength to provide sufficient drug for a single treatment at a dose of 1 microgram/kg. Up to 1 repeat may be requested with the initial written application.

Subsequently during the initial period of dose titration, authority applications for a single dose and up to 1 repeat may be requested by telephone. The dose (microgram/kg/week) must be provided at the time of application.

Once a patient's dose has been stable for a period of 4 weeks, authority approvals for sufficient vials of appropriate strength based on the weight of the patient and dose (microgram/kg/week) for up to 4 weeks of treatment and up to 4 repeats may be granted, as long as the total period of treatment authorised under this restriction does not exceed 24 weeks.

Authority approval will not be given for doses higher than 10 micrograms/kg/week

The authority application must be made in writing and must include:

- (1) a completed authority prescription form,
- (2) a signed patient acknowledgement,
- (3) a completed Idiopathic Thrombocytopenic Purpura Initial PBS Authority Application - Supporting Information Form,
- (4) a copy of a full blood count pathology report supporting the diagnosis of ITP, and
- (5) where the application is sought on the basis of a medical contraindication to surgery, a signed and dated letter from the clinician making this assessment which includes the date upon which the patient was assessed for surgery and the clinical grounds upon which surgery is contraindicated.

The full blood count must be no more than 1 month old at the time of application.

Note Romiplostim is not PBS-subsidised as an alternative to splenectomy.

Note Patients will be able to trial either eltrombopag or romiplostim within the initial 24 weeks treatment period. Patients who fail to demonstrate a response to treatment with eltrombopag and/or romiplostim under the initial restriction will not be eligible to receive further PBS-subsidised treatment with either of these drugs.

Note Any queries concerning the arrangements to prescribe this drug may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au
Written applications for authority to prescribe this drug should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe thrombocytopenia

Treatment Phase: First Continuing treatment or Re-initiation of interrupted treatment

Clinical criteria:

- The condition must be severe chronic immune (idiopathic) thrombocytopenic purpura (ITP), **AND**
- Patient must have previously received PBS-subsidised initial treatment with this drug for this condition, **AND**
- Patient must have demonstrated a sustained platelet response to PBS-subsidised treatment with this drug for this condition under the Initial treatment restriction, **AND**
- The treatment must be the sole PBS-subsidised thrombopoietin receptor agonist (TRA) for this condition.

Population criteria:

- Patient must be an adult.

For the purposes of this restriction, a sustained platelet response is defined as:

(a) use of rescue medication (corticosteroids or immunoglobulins) on no more than one occasion during the initial period of PBS-subsidised treatment with this drug,

AND either of the following:

(b) a platelet count greater than or equal to 50,000 million per L on at least four (4) occasions, each at least one week apart;
OR

(c) a platelet count greater than 30,000 million per L and which is double the baseline (pre-treatment) platelet count on at least four (4) occasions, each at least one week apart.

The medical practitioner should request sufficient number of vials of appropriate strength based on the weight of the patient and dose (microgram/kg/week) to provide 4 weeks of treatment. Up to a maximum of 5 repeats may be authorised.

Authority approval will not be given for doses higher than 10 micrograms/kg/week

Applications for the First continuing PBS-subsidised treatment or Re-initiation of interrupted PBS-subsidised treatment must be made in writing and must include:

(1) a completed authority prescription form, and

(2) a completed Idiopathic Thrombocytopenic Purpura Continuing PBS Authority Application - Supporting Information Form , and

(3) copies of the platelet count pathology reports (unless previously provided for patients re-initiating therapy).

The platelet count must be no more than one month old at the time of application.

Note Romiplostim is not PBS-subsidised as an alternative to splenectomy.

Note Patients will be able to trial either eltrombopag or romiplostim within the initial 24 weeks treatment period. Patients who fail to demonstrate a response to treatment with eltrombopag and/or romiplostim under the initial restriction will not be eligible to receive further PBS-subsidised treatment with either of these drugs.

Note Any queries concerning the arrangements to prescribe this drug may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Written applications for authority to prescribe this drug should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe thrombocytopenia

Treatment Phase: Second or Subsequent Continuing treatment

Clinical criteria:

- The condition must be severe chronic immune (idiopathic) thrombocytopenic purpura (ITP), **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have demonstrated a continuing response to treatment with this drug, **AND**
- The treatment must be the sole PBS-subsidised thrombopoietin receptor agonist (TRA) for this condition.

Population criteria:

- Patient must be an adult.

For the purpose of this restriction, a continuing response to treatment with drug is defined as:

(a) use of rescue medication (corticosteroids or immunoglobulins) on no more than one occasion during the most recent 24 week period of PBS-subsidised treatment with this drug

AND either of the following:

(b) a platelet count greater than or equal to 50,000 million per L

OR

(c) a platelet count greater than 30,000 million per L and which is double the baseline platelet count.

The platelet count must be no more than one month old at the time of application.

The medical practitioner should request sufficient number of vials of appropriate strength based on the weight of the patient and dose (microgram/kg/week) to provide 4 weeks of treatment. Up to a maximum of 5 repeats may be authorised.

Authority approval will not be given for doses higher than 10 micrograms/kg/week

Authority applications for second and subsequent periods of continuing therapy may be made by telephone

Note Romiplostim is not PBS-subsidised as an alternative to splenectomy.

Note Authority applications for second and subsequent continuing treatment may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note Patients will be able to trial either eltrombopag or romiplostim within the initial 24 weeks treatment period. Patients who fail to demonstrate a response to treatment with eltrombopag and/or romiplostim under the initial restriction will not be eligible to receive further PBS-subsidised treatment with either of these drugs.

Note Any queries concerning the arrangements to prescribe this drug may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Written applications for authority to prescribe this drug should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe thrombocytopenia

Treatment Phase: Initial 1, Initial 2, First Continuing treatment or Re-initiation of interrupted treatment, and Second and Subsequent Continuing treatment - balance of supply

Clinical criteria:

- The condition must be severe chronic immune (idiopathic) thrombocytopenic purpura (ITP), **AND**
- The treatment must be the sole PBS-subsidised thrombopoietin receptor agonist (TRA) for this condition, **AND**
- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the First Continuing treatment or Re-initiation of interrupted treatment restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Second and subsequent Continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restriction.

Population criteria:

- Patient must be an adult.

Note Authority approval for sufficient therapy to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note No applications for increased repeats will be authorised.

romiplostim 500 microgram injection, 1 vial

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
9698K	1	..	..	1098.04	Nplate [AN]

romiplostim 250 microgram injection, 1 vial

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
9696H	1	..	..	549.02	Nplate [AN]

■ ANTIANEMIC PREPARATIONS

OTHER ANTIANEMIC PREPARATIONS

Other antianemic preparations

■ DARBEPOETIN ALFA

Authority required (STREAMLINED)

6294

Anaemia associated with intrinsic renal disease

Clinical criteria:

- Patient must require transfusion, **AND**
- Patient must have a haemoglobin level of less than 100 g per L, **AND**
- Patient must have intrinsic renal disease, as assessed by a nephrologist.

darbepoetin alfa 40 microgram/0.4 mL injection, 4 x 0.4 mL syringes

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
5640W	2	5	..	*856.94	Aranesp [AN]

darbepoetin alfa 60 microgram/0.3 mL injection, 0.3 mL syringe

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
5647F	8	5	..	*1244.08	Aranesp SureClick [AN]

darbepoetin alfa 40 microgram/0.4 mL injection, 0.4 mL syringe

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
5646E	8	5	..	*856.88	Aranesp SureClick [AN]

darbepoetin alfa 150 microgram/0.3 mL injection, 4 x 0.3 mL syringes

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
5643B	2	5	..	*3004.62	Aranesp [AN]

darbepoetin alfa 150 microgram/0.3 mL injection, 0.3 mL syringe

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
5650J	8	5	..	*3004.64	Aranesp SureClick [AN]

darbepoetin alfa 100 microgram/0.5 mL injection, 4 x 0.5 mL syringes

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
5651K	2	5	..	*2016.56	Aranesp [AN]

darbepoetin alfa 30 microgram/0.3 mL injection, 4 x 0.3 mL syringes

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
5639T	2	5	..	*706.00	Aranesp [AN]

darbepoetin alfa 100 microgram/0.5 mL injection, 0.5 mL syringe

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
5649H	8	5	..	*2016.48	Aranesp SureClick [AN]

darbepoetin alfa 60 microgram/0.3 mL injection, 4 x 0.3 mL syringes

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
5642Y	2	5	..	*1244.06	Aranesp [AN]

darbepoetin alfa 50 microgram/0.5 mL injection, 4 x 0.5 mL syringes

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
5641X	2	5	..	*1059.46	Aranesp [AN]

darbepoetin alfa 20 microgram/0.5 mL injection, 0.5 mL syringe

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
5645D	8	5	..	*516.08	Aranesp SureClick [AN]

darbepoetin alfa 20 microgram/0.5 mL injection, 4 x 0.5 mL syringes

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
5638R	2	5	..	*516.04	Aranesp [AN]

darbepoetin alfa 10 microgram/0.4 mL injection, 4 x 0.4 mL syringes

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
5637Q	2	5	..	*274.02	Aranesp [AN]

darbepoetin alfa 80 microgram/0.4 mL injection, 4 x 0.4 mL syringes

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
5644C	2	5	..	*1637.54	Aranesp [AN]

darbepoetin alfa 80 microgram/0.4 mL injection, 0.4 mL syringe

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
5648G	8	5	..	*1637.60	Aranesp SureClick [AN]

■ **EPOETIN ALFA**

Authority required (STREAMLINED)

6294

Anaemia associated with intrinsic renal disease

Clinical criteria:

- Patient must require transfusion, **AND**
- Patient must have a haemoglobin level of less than 100 g per L, **AND**
- Patient must have intrinsic renal disease, as assessed by a nephrologist.

epoetin alfa 3000 units/0.3 mL injection, 6 x 0.3 mL syringes

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
5720C	2	5	..	*513.20	Eprex 3000 [JC]

epoetin alfa 6000 units/0.6 mL injection, 6 x 0.6 mL syringes

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
5716W	2	5	..	*965.86	Eprex 6000 [JC]

epoetin alfa 8000 units/0.8 mL injection, 6 x 0.8 mL syringes

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
5717X	2	5	..	*1252.72	Eprex 8000 [JC]

epoetin alfa 10 000 units/mL injection, 6 x 1 mL syringes

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
5722E	2	5	..	*1516.18	Eprex 10000 [JC]

epoetin alfa 1000 units/0.5 mL injection, 6 x 0.5 mL syringes

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
5714R	2	5	..	*214.92	Eprex 1000 [JC]

epoetin alfa 4000 units/0.4 mL injection, 6 x 0.4 mL syringes

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
5721D	2	5	..	*653.56	Eprex 4000 [JC]

epoetin alfa 20 000 units/0.5 mL injection, 6 x 0.5 mL syringes

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
5713Q	2	5	..	*2982.68	Eprex 20,000 [JC]

epoetin alfa 2000 units/0.5 mL injection, 6 x 0.5 mL syringes

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
5719B	2	5	..	*397.70	Eprex 2000 [JC]

epoetin alfa 40 000 units/mL injection, 1 mL syringe

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
5718Y	2	5	..	*964.98	Eprex 40,000 [JC]

epoetin alfa 5000 units/0.5 mL injection, 6 x 0.5 mL syringes

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
5715T	2	5	..	*813.66	Eprex 5000 [JC]

■ EPOETIN BETA**Authority required (STREAMLINED)****6294**

Anaemia associated with intrinsic renal disease

Clinical criteria:

- Patient must require transfusion, **AND**
- Patient must have a haemoglobin level of less than 100 g per L, **AND**
- Patient must have intrinsic renal disease, as assessed by a nephrologist.

epoetin beta 2000 units/0.3 mL injection, 6 x 0.3 mL syringes

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
5724G	2	5	..	*418.62	NeoRecormon [RO]

epoetin beta 6000 units/0.3 mL injection, 6 x 0.3 mL syringes

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
5728L	2	5	..	*1016.70	NeoRecormon [RO]

epoetin beta 5000 units/0.3 mL injection, 6 x 0.3 mL syringes

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
5727K	2	5	..	*856.50	NeoRecormon [RO]

epoetin beta 3000 units/0.3 mL injection, 6 x 0.3 mL syringes

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
5725H	2	5	..	*540.22	NeoRecormon [RO]

epoetin beta 10 000 units/0.6 mL injection, 6 x 0.6 mL syringes

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
5729M	2	5	..	*1595.98	NeoRecormon [RO]

epoetin beta 4000 units/0.3 mL injection, 6 x 0.3 mL syringes

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
5726J	2	5	..	*687.96	NeoRecormon [RO]

■ EPOETIN LAMBDA**Authority required (STREAMLINED)****6294**

Anaemia associated with intrinsic renal disease

Clinical criteria:

- Patient must require transfusion, **AND**
- Patient must have a haemoglobin level of less than 100 g per L, **AND**
- Patient must have intrinsic renal disease, as assessed by a nephrologist.

epoetin lambda 3000 units/0.3 mL injection, 6 x 0.3 mL syringes

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
9670Y	2	5	..	*600.22	Novicrit [SZ]

BLOOD AND BLOOD FORMING ORGANS

epoetin lambda 5000 units/0.5 mL injection, 6 x 0.5 mL syringes

9589Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*951.62	Novicrit [SZ]

epoetin lambda 6000 units/0.6 mL injection, 6 x 0.6 mL syringes

9591T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*1129.62	Novicrit [SZ]

epoetin lambda 1000 units/0.5 mL injection, 6 x 0.5 mL syringes

9668W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*251.38	Novicrit [SZ]

epoetin lambda 8000 units/0.8 mL injection, 6 x 0.8 mL syringes

9594Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*1465.12	Novicrit [SZ]

epoetin lambda 2000 units/mL injection, 6 x 1 mL syringes

9669X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*465.12	Novicrit [SZ]

epoetin lambda 10 000 units/mL injection, 6 x 1 mL syringes

9596C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*1773.28	Novicrit [SZ]

epoetin lambda 4000 units/0.4 mL injection, 6 x 0.4 mL syringes

9587N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*764.38	Novicrit [SZ]

■ METHOXY POLYETHYLENE GLYCOL-EPOETIN BETA

Authority required (STREAMLINED)

6294

Anaemia associated with intrinsic renal disease

Clinical criteria:

- Patient must require transfusion, **AND**
- Patient must have a haemoglobin level of less than 100 g per L, **AND**
- Patient must have intrinsic renal disease, as assessed by a nephrologist.

methoxy polyethylene glycol-epoetin beta 30 microgram/0.3 mL injection, 0.3 mL syringe

5794Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*332.28	Mircera [RO]

methoxy polyethylene glycol-epoetin beta 100 microgram/0.3 mL injection, 0.3 mL syringe

5797D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*1042.98	Mircera [RO]

methoxy polyethylene glycol-epoetin beta 75 microgram/0.3 mL injection, 0.3 mL syringe

5796C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*806.44	Mircera [RO]

methoxy polyethylene glycol-epoetin beta 50 microgram/0.3 mL injection, 0.3 mL syringe

5795B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*553.80	Mircera [RO]

methoxy polyethylene glycol-epoetin beta 360 microgram/0.6 mL injection, 0.6 mL syringe

5800G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*2993.98	Mircera [RO]

methoxy polyethylene glycol-epoetin beta 120 microgram/0.3 mL injection, 0.3 mL syringe

5798E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*1207.52	Mircera [RO]

methoxy polyethylene glycol-epoetin beta 200 microgram/0.3 mL injection, 0.3 mL syringe

5799F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*1731.92	Mircera [RO]

■ CARDIOVASCULAR SYSTEM

■ ANTIHYPERTENSIVES

OTHER ANTIHYPERTENSIVES

Antihypertensives for pulmonary arterial hypertension

■ AMBRISENTAN

Caution This is a category X drug and must not be given to pregnant women. Pregnancy must be avoided during treatment and for at least 3 months following cessation of therapy.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 1 (new patients)

Clinical criteria:

- Patient must not have received prior PBS-subsidised treatment with a pulmonary arterial hypertension (PAH) agent, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**
- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH; OR
- Patient must have WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease, **AND**
- Patient must have a mean right atrial pressure of 8 mmHg or less as measured by right heart catheterisation (RHC); OR
- Patient must have right ventricular function assessed by echocardiography (ECHO) where a RHC cannot be performed on clinical grounds, **AND**
- Patient must have failed to respond to 6 or more weeks of appropriate vasodilator treatment unless intolerance or a contraindication to such treatment exists, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6 Minute Walk Test (6MWT); and
- (3) a signed patient acknowledgement.

Idiopathic pulmonary arterial hypertension, anorexigen-induced pulmonary arterial hypertension, hereditary pulmonary arterial hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) are defined as follows:

- (i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary artery wedge pressure (PAWP) less than 15 mmHg; or
- (ii) where a right heart catheter (RHC) cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.

Test requirements to establish baseline for initiation of treatment are as follows:

The first written application for PBS-subsidised treatment with the first PAH agent should be accompanied by the results of a right heart catheter (RHC) composite assessment plus an echocardiograph (ECHO) composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.

Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments;
- (2) RHC composite assessment plus 6MWT;
- (3) RHC composite assessment only.

In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted for consideration based on the results of the following test combinations, which are listed in descending order of preference:

- (1) ECHO composite assessment plus 6MWT;
- (2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

Details of prior vasodilator treatment, including the dose and duration of treatment, must be provided at the time of application. Where the patient has an adverse event to a vasodilator or where vasodilator treatment is contraindicated, details of the nature of the adverse event or contraindication according to the Therapeutic Goods Administration (TGA) approved Product Information must also be provided with the application.

Response to prior vasodilator treatment is defined as follows:

For patients with 2 or more baseline tests, response to treatment is defined as 2 or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information.

A maximum of 5 repeats may be requested.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Prior Written Approval of Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 2 (new patients)

Clinical criteria:

- Patient must not have received prior PBS-subsidised treatment with a pulmonary arterial hypertension (PAH) agent, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**
- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH), or anorexigen-induced PAH or hereditary PAH, and a mean right atrial pressure of greater than 8 mmHg, as measured by right heart catheterisation (RHC); OR
- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH), or anorexigen-induced PAH or hereditary PAH, with right ventricular function assessed by echocardiography (ECHO) where a RHC cannot be performed on clinical grounds; OR
- Patient must have WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease and a mean right atrial pressure greater than 8 mmHg, as measured by RHC; OR
- Patient must have WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease with right ventricular function assessed by ECHO where a RHC cannot be performed on clinical grounds; OR
- Patient must have WHO Functional Class IV idiopathic pulmonary arterial hypertension (iPAH), or anorexigen-induced PAH or hereditary PAH; OR
- Patient must have WHO Functional Class IV pulmonary arterial hypertension secondary to connective tissue disease, **AND**

- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6 Minute Walk Test (6MWT); and
- (3) a signed patient acknowledgement.

Idiopathic pulmonary arterial hypertension, anorexigen-induced pulmonary arterial hypertension, hereditary pulmonary arterial hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) are defined as follows:

- (i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary artery wedge pressure (PAWP) less than 15 mmHg; or
- (ii) where a right heart catheter (RHC) cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.

Test requirements to establish baseline for initiation of treatment are as follows:

The first written application for PBS-subsidised treatment with the first PAH agent should be accompanied by the results of a right heart catheter (RHC) composite assessment plus an echocardiograph (ECHO) composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.

Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments;

- (2) RHC composite assessment plus 6MWT;
- (3) RHC composite assessment only.

In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted for consideration based on the results of the following test combinations, which are listed in descending order of preference:

- (1) ECHO composite assessment plus 6MWT;
- (2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats may be requested.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 3 (change or re-commencement of therapy for all patients)

Clinical criteria:

- Patient must have idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH or PAH secondary to connective tissue disease and must wish to re-commence PBS-subsidised therapy with this agent after a break in therapy and must have demonstrated a response to their most recent course of PBS-subsidised treatment with this agent; OR
- Patient must have idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH or PAH secondary to connective tissue disease and whose most recent course of PBS-subsidised treatment was with a PAH agent other than this agent, **AND**

- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form; and
- (3) the results of the patient's response to treatment with their last course of PBS-subsidised PAH agent.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

Response to a PAH agent is defined as follows:

For patients with two or more baseline tests, response to treatment is defined as two or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats may be requested.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Swapping between PAH agents: Patients can access PAH agents through the PBS according to the relevant restrictions. Once these patients are approved initial treatment with 1 of these 8 drugs, they may swap between PAH agents at any time without having to re-qualify for treatment with the alternate agent. This means that patients may commence treatment with the alternate agent, subject to that agent's restriction, irrespective of the severity of their disease at the time the application to swap therapy is submitted. It also means that no new baseline measurements will be necessary. New baselines may be submitted where the patient has failed to respond to their current treatment. Eligible patients may only swap between PAH agents if they have not failed prior PBS-subsidised treatment with that agent. For eligible patients, applications to swap between the 8 PAH agents must be made under the relevant initial treatment restriction. Patients should be assessed for response to the treatment they are ceasing at the time the application to swap therapy is being made. Patients who fail to demonstrate a response or for whom no assessment results are submitted with the application to swap therapy may not re-commence PBS-subsidised treatment with the drug they are ceasing.

Note Applications for patients who wish to swap to an alternate PAH agent should be accompanied by the previously approved authority prescription, or remaining repeats, for the treatment the patient is ceasing.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 1 or Initial 2 (new patients) or Initial 3 (change or re-commencement of therapy for all patients) or First Continuing treatment - Balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this agent under the Initial 1 (new patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the Initial 2 (new patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the Initial 3 (change or re-commencement of therapy for all patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the First Continuing treatment restriction to complete a maximum of six months of treatment, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition, **AND**
- The treatment must provide no more than the balance of up to six months treatment available under one of the above restrictions.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authorisation under this criterion should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: First Continuing treatment

Clinical criteria:

- Patient must have received a PBS-subsidised initial course of treatment with this agent for this condition, **AND**
- Patient must have been assessed by a physician from a designated hospital to have achieved a response to the PBS-subsidised initial course of treatment, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

(1) a completed authority prescription form; and

(2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:

(i) RHC composite assessment; and

(ii) ECHO composite assessment; and

(iii) 6 Minute Walk Test (6MWT).

Test requirements to establish response to treatment for continuation of treatment are as follows:

The following list outlines the preferred test combination, in descending order, for the purposes of continuation of PBS-subsidised treatment:

(1) RHC plus ECHO composite assessments plus 6MWT;

(2) RHC plus ECHO composite assessments;

(3) RHC composite assessment plus 6MWT;

(4) ECHO composite assessment plus 6MWT;

(5) RHC composite assessment only;

(6) ECHO composite assessment only.

The results of the same tests as conducted at baseline should be provided with the written First Continuing treatment application, except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a patient specific reason why the test(s) could not be conducted must be provided with the application.

The test results provided with the application for continuing treatment must be no more than 2 months old at the time of application.

Response to a PAH agent is defined as follows:

For patients with two or more baseline tests, response to treatment is defined as two or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats will be authorised.

An application for First Continuing treatment with a PAH agent should be made prior to the completion of the Initial 6 month treatment course to ensure continuity for those patients who respond to treatment, as assessed by the treating physician.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Subsequent Continuing treatment

Clinical criteria:

- Patient must have received a PBS-subsidised treatment under First Continuing treatment with this agent for this condition; OR
- Patient must have previously received PBS-subsidised treatment under this criteria with this agent for this condition, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats will be authorised.

An application for Subsequent Continuing treatment with a PAH agents should be made prior to the completion of the First Continuing treatment course to ensure continuity of treatment.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authorisation under this criterion should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

ambrisentan 10 mg tablet, 30

5608E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	2732.65	Volibris [GK]

ambrisentan 5 mg tablet, 30

5607D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	2732.65	Volibris [GK]

■ **BOSENTAN**

Caution This is a category X drug and must not be given to pregnant women. Pregnancy must be avoided during treatment and for at least 3 months following cessation of therapy.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 1 (new patients)

Clinical criteria:

- Patient must not have received prior PBS-subsidised treatment with a pulmonary arterial hypertension (PAH) agent, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**
- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH; OR
- Patient must have WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease, **AND**
- Patient must have a mean right atrial pressure of 8 mmHg or less as measured by right heart catheterisation (RHC); OR
- Patient must have right ventricular function assessed by echocardiography (ECHO) where a RHC cannot be performed on clinical grounds, **AND**
- Patient must have failed to respond to 6 or more weeks of appropriate vasodilator treatment unless intolerance or a contraindication to such treatment exists, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

- (1) two completed authority prescription forms; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6 Minute Walk Test (6MWT); and
- (3) a signed patient acknowledgement.

Idiopathic pulmonary arterial hypertension, anorexigen-induced pulmonary arterial hypertension, hereditary pulmonary arterial hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) are defined as follows:

- (i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary artery wedge pressure (PAWP) less than 15 mmHg; or
- (ii) where a right heart catheter (RHC) cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.

Test requirements to establish baseline for initiation of treatment are as follows:

The first written application for PBS-subsidised treatment with the first PAH agent should be accompanied by the results of a right heart catheter (RHC) composite assessment plus an echocardiograph (ECHO) composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.

Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments;
- (2) RHC composite assessment plus 6MWT;
- (3) RHC composite assessment only.

In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted for consideration based on the results of the following test combinations, which are listed in descending order of preference:

- (1) ECHO composite assessment plus 6MWT;
- (2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

Details of prior vasodilator treatment, including the dose and duration of treatment, must be provided at the time of application. Where the patient has an adverse event to a vasodilator or where vasodilator treatment is contraindicated, details of the nature of the adverse event or contraindication according to the Therapeutic Goods Administration (TGA) approved Product Information must also be provided with the application.

Response to prior vasodilator treatment is defined as follows:

For patients with 2 or more baseline tests, response to treatment is defined as 2 or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

Approvals for the first authority prescription will be limited to 1 month of therapy with the 62.5 mg strength tablet, with the quantity approved based on the dosage recommendations in the TGA-approved Product Information. No repeats will be authorised for this prescription.

The second authority prescription may be written for either the 62.5 mg tablet or the 125 mg tablet strengths. Approvals for the second authority prescription will be limited to 1 month of treatment, with the quantity approved based on the dosage recommendations in the TGA-approved Product Information, and a maximum of 4 repeats.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Where the 62.5 mg tablet strength is required for the second authority prescription, please contact the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) for further advice. The approved second authority prescription will be returned to the prescriber by the Department of Human Services two weeks after the date of the approval of the first authority prescription, to allow for the uninterrupted completion of the six months initial treatment course. The Department of Human Services will contact prescribers prior to dispatch of the second authority prescription to confirm the tablet strength required for the patient.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 2 (new patients)

Clinical criteria:

- Patient must not have received prior PBS-subsidised treatment with a pulmonary arterial hypertension (PAH) agent, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**
- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH), or anorexigen-induced PAH or hereditary PAH, and a mean right atrial pressure of greater than 8 mmHg, as measured by right heart catheterisation (RHC); OR
- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH), or anorexigen-induced PAH or hereditary PAH, with right ventricular function assessed by echocardiography (ECHO) where a RHC cannot be performed on clinical grounds; OR
- Patient must have WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease and a mean right atrial pressure greater than 8 mmHg, as measured by RHC; OR
- Patient must have WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease with right ventricular function assessed by ECHO where a RHC cannot be performed on clinical grounds; OR
- Patient must have WHO Functional Class IV idiopathic pulmonary arterial hypertension (iPAH), or anorexigen-induced PAH or hereditary PAH; OR
- Patient must have WHO Functional Class IV pulmonary arterial hypertension secondary to connective tissue disease; OR
- Patient must have WHO Functional Class III or IV pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology), **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

- (1) two completed authority prescription forms; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6 Minute Walk Test (6MWT); and
- (3) a signed patient acknowledgement.

Idiopathic pulmonary arterial hypertension, anorexigen-induced pulmonary arterial hypertension, hereditary pulmonary arterial hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) are defined as follows:

(i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary artery wedge pressure (PAWP) less than 15 mmHg; or

(ii) where a right heart catheter (RHC) cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.

Test requirements to establish baseline for initiation of treatment are as follows:

The first written application for PBS-subsidised treatment with the first PAH agent should be accompanied by the results of a right heart catheter (RHC) composite assessment plus an echocardiograph (ECHO) composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.

Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments;
- (2) RHC composite assessment plus 6MWT;
- (3) RHC composite assessment only.

In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted for consideration based on the results of the following test combinations, which are listed in descending order of preference:

- (1) ECHO composite assessment plus 6MWT;
- (2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

Approvals for the first authority prescription will be limited to 1 month of therapy with the 62.5 mg strength tablet, with the quantity approved based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information. No repeats will be authorised for this prescription.

The second authority prescription may be written for either the 62.5 mg tablet or the 125 mg tablet strengths. Approvals for the second authority prescription will be limited to 1 month of treatment, with the quantity approved based on the dosage recommendations in the TGA-approved Product Information, and a maximum of 4 repeats.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Where the 62.5 mg tablet strength is required for the second authority prescription, please contact the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) for further advice.

The approved second authority prescription will be returned to the prescriber by the Department of Human Services two weeks after the date of the approval of the first authority prescription, to allow for the uninterrupted completion of the six months initial treatment course. The Department of Human Services will contact prescribers prior to dispatch of the second authority prescription to confirm the tablet strength required for the patient.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 3 (change or re-commencement of therapy for all patients)

Clinical criteria:

- Patient must have idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH or PAH secondary to connective tissue disease or PAH associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) and must wish to re-commence PBS-subsidised therapy with this agent after a break in therapy and must have demonstrated a response to their most recent course of PBS-subsidised treatment with this agent; OR
- Patient must have idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH or PAH secondary to connective tissue disease or PAH associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) and whose most recent course of PBS-subsidised treatment was with a PAH agent other than this agent, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

- (1) two completed authority prescription forms; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form; and
- (3) the results of the patient's response to treatment with their last course of PBS-subsidised PAH agent.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

Response to a PAH agent is defined as follows:

For patients with two or more baseline tests, response to treatment is defined as two or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

Approvals for the first authority prescription will be limited to 1 month of therapy with the 62.5 mg strength tablet, with the quantity approved based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information. No repeats will be authorised for this prescription.

The second authority prescription may be written for either the 62.5 mg tablet or the 125 mg tablet strengths. Approvals for the second authority prescription will be limited to 1 month of treatment, with the quantity approved based on the dosage recommendations in the TGA-approved Product Information, and a maximum of 4 repeats.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Swapping between PAH agents: Patients can access PAH agents through the PBS according to the relevant restrictions. Once these patients are approved initial treatment with 1 of these 8 drugs, they may swap between PAH agents at any time without having to re-qualify for treatment with the alternate agent. This means that patients may commence treatment with the alternate agent, subject to that agent's restriction, irrespective of the severity of their disease at the time the application to swap therapy is submitted. It also means that no new baseline measurements will be necessary. New baselines may be submitted where the patient has failed to respond to their current treatment. Eligible patients may only swap between PAH agents if they have not failed prior PBS-subsidised treatment with that agent. For eligible patients, applications to swap between the 8 PAH agents must be made under the relevant initial treatment restriction. Patients should be assessed for response to the treatment they are ceasing at the time the application to swap therapy is being made. Patients who fail to demonstrate a response or for whom no assessment results are submitted with the application to swap therapy may not re-commence PBS-subsidised treatment with the drug they are ceasing.

Note Where the 62.5 mg tablet strength is required for the second authority prescription, please contact the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) for further advice. The approved second authority prescription will be returned to the prescriber by the Department of Human Services two weeks after the date of the approval of the first authority prescription, to allow for the uninterrupted completion of the six months initial treatment course. The Department of Human Services will contact prescribers prior to dispatch of the second authority prescription to confirm the tablet strength required for the patient.

Note Applications for patients who wish to swap to an alternate PAH agent should be accompanied by the previously approved authority prescription, or remaining repeats, for the treatment the patient is ceasing.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 1 or Initial 2 (new patients) or Initial 3 (change or re-commencement of therapy for all patients) or First Continuing treatment - Balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this agent under the Initial 1 (new patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the Initial 2 (new patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the Initial 3 (change or re-commencement of therapy for all patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the First Continuing treatment restriction to complete a maximum of six months of treatment, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition, **AND**
- The treatment must provide no more than the balance of up to six months treatment available under one of the above restrictions.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authorisation under this criterion should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: First Continuing treatment

Clinical criteria:

- Patient must have received a PBS-subsidised initial course of treatment with this agent for this condition, **AND**
- Patient must have been assessed by a physician from a designated hospital to have achieved a response to the PBS-subsidised initial course of treatment, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

(1) a completed authority prescription form; and

(2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:

(i) RHC composite assessment; and

(ii) ECHO composite assessment; and

(iii) 6 Minute Walk Test (6MWT).

Test requirements to establish response to treatment for continuation of treatment are as follows:

The following list outlines the preferred test combination, in descending order, for the purposes of continuation of PBS-subsidised treatment:

(1) RHC plus ECHO composite assessments plus 6MWT;

(2) RHC plus ECHO composite assessments;

(3) RHC composite assessment plus 6MWT;

(4) ECHO composite assessment plus 6MWT;

(5) RHC composite assessment only;

(6) ECHO composite assessment only.

The results of the same tests as conducted at baseline should be provided with the written First Continuing treatment application, except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a patient specific reason why the test(s) could not be conducted must be provided with the application.

The test results provided with the application for continuing treatment must be no more than 2 months old at the time of application.

Response to a PAH agent is defined as follows:

For patients with two or more baseline tests, response to treatment is defined as two or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats will be authorised.

An application for First Continuing treatment with a PAH agent should be made prior to the completion of the Initial 6 month treatment course to ensure continuity for those patients who respond to treatment, as assessed by the treating physician.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Subsequent Continuing treatment

Clinical criteria:

- Patient must have received a PBS-subsidised treatment under First Continuing treatment with this agent for this condition; OR
- Patient must have previously received PBS-subsidised treatment under this criteria with this agent for this condition, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats will be authorised.

An application for Subsequent Continuing treatment with a PAH agents should be made prior to the completion of the First Continuing treatment course to ensure continuity of treatment.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authorisation under this criterion should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

bosentan 125 mg tablet, 60

5619R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	1113.08	^a Bosentan APO [GX] ^a BOSENTAN DR. REDDY'S [RI] ^a Bosentan Mylan [AF] ^a Bosentan Sandoz [SZ] ^a Tracleer [AT]	^a BOSENTAN-DRLA [RZ] ^a Bosentan GH [GQ] ^a Bosentan RBX [RA] ^a BOSLEER [RW]

■ BOSENTAN

Caution This is a category X drug and must not be given to pregnant women. Pregnancy must be avoided during treatment and for at least 3 months following cessation of therapy.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 1 (new patients)

Clinical criteria:

- Patient must not have received prior PBS-subsidised treatment with a pulmonary arterial hypertension (PAH) agent, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**
- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH; OR
- Patient must have WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease, **AND**
- Patient must have a mean right atrial pressure of 8 mmHg or less as measured by right heart catheterisation (RHC); OR
- Patient must have right ventricular function assessed by echocardiography (ECHO) where a RHC cannot be performed on clinical grounds, **AND**
- Patient must have failed to respond to 6 or more weeks of appropriate vasodilator treatment unless intolerance or a contraindication to such treatment exists, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

(1) two completed authority prescription forms; and

(2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:

(i) RHC composite assessment; and

(ii) ECHO composite assessment; and

(iii) 6 Minute Walk Test (6MWT); and

(3) a signed patient acknowledgement.

Idiopathic pulmonary arterial hypertension, anorexigen-induced pulmonary arterial hypertension, hereditary pulmonary arterial hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) are defined as follows:

(i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary artery wedge pressure (PAWP) less than 15 mmHg; or

(ii) where a right heart catheter (RHC) cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.

Test requirements to establish baseline for initiation of treatment are as follows:

The first written application for PBS-subsidised treatment with the first PAH agent should be accompanied by the results of a right heart catheter (RHC) composite assessment plus an echocardiograph (ECHO) composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.

Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments;
- (2) RHC composite assessment plus 6MWT;
- (3) RHC composite assessment only.

In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted for consideration based on the results of the following test combinations, which are listed in descending order of preference:

- (1) ECHO composite assessment plus 6MWT;
- (2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

Details of prior vasodilator treatment, including the dose and duration of treatment, must be provided at the time of application. Where the patient has an adverse event to a vasodilator or where vasodilator treatment is contraindicated, details of the nature of the adverse event or contraindication according to the Therapeutic Goods Administration (TGA) approved Product Information must also be provided with the application.

Response to prior vasodilator treatment is defined as follows:

For patients with 2 or more baseline tests, response to treatment is defined as 2 or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

Approvals for the first authority prescription will be limited to 1 month of therapy with the 62.5 mg strength tablet, with the quantity approved based on the dosage recommendations in the TGA-approved Product Information. No repeats will be authorised for this prescription.

The second authority prescription may be written for either the 62.5 mg tablet or the 125 mg tablet strengths. Approvals for the second authority prescription will be limited to 1 month of treatment, with the quantity approved based on the dosage recommendations in the TGA-approved Product Information, and a maximum of 4 repeats.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Where the 62.5 mg tablet strength is required for the second authority prescription, please contact the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) for further advice.

The approved second authority prescription will be returned to the prescriber by the Department of Human Services two weeks after the date of the approval of the first authority prescription, to allow for the uninterrupted completion of the six months initial treatment course. The Department of Human Services will contact prescribers prior to dispatch of the second authority prescription to confirm the tablet strength required for the patient.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 2 (new patients)

Clinical criteria:

- Patient must not have received prior PBS-subsidised treatment with a pulmonary arterial hypertension (PAH) agent, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**
- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH), or anorexigen-induced PAH or hereditary PAH, and a mean right atrial pressure of greater than 8 mmHg, as measured by right heart catheterisation (RHC); OR

- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH), or anorexigen-induced PAH or hereditary PAH, with right ventricular function assessed by echocardiography (ECHO) where a RHC cannot be performed on clinical grounds; OR
- Patient must have WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease and a mean right atrial pressure greater than 8 mmHg, as measured by RHC; OR
- Patient must have WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease with right ventricular function assessed by ECHO where a RHC cannot be performed on clinical grounds; OR
- Patient must have WHO Functional Class IV idiopathic pulmonary arterial hypertension (iPAH), or anorexigen-induced PAH or hereditary PAH; OR
- Patient must have WHO Functional Class IV pulmonary arterial hypertension secondary to connective tissue disease; OR
- Patient must have WHO Functional Class III or IV pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology), **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

(1) two completed authority prescription forms; and

(2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:

(i) RHC composite assessment; and

(ii) ECHO composite assessment; and

(iii) 6 Minute Walk Test (6MWT); and

(3) a signed patient acknowledgement.

Idiopathic pulmonary arterial hypertension, anorexigen-induced pulmonary arterial hypertension, hereditary pulmonary arterial hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) are defined as follows:

(i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary artery wedge pressure (PAWP) less than 15 mmHg; or

(ii) where a right heart catheter (RHC) cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.

Test requirements to establish baseline for initiation of treatment are as follows:

The first written application for PBS-subsidised treatment with the first PAH agent should be accompanied by the results of a right heart catheter (RHC) composite assessment plus an echocardiograph (ECHO) composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.

Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:

(1) RHC plus ECHO composite assessments;

(2) RHC composite assessment plus 6MWT;

(3) RHC composite assessment only.

In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted for consideration based on the results of the following test combinations, which are listed in descending order of preference:

(1) ECHO composite assessment plus 6MWT;

(2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

Approvals for the first authority prescription will be limited to 1 month of therapy with the 62.5 mg strength tablet, with the quantity approved based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information. No repeats will be authorised for this prescription.

The second authority prescription may be written for either the 62.5 mg tablet or the 125 mg tablet strengths. Approvals for the second authority prescription will be limited to 1 month of treatment, with the quantity approved based on the dosage recommendations in the TGA-approved Product Information, and a maximum of 4 repeats.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Where the 62.5 mg tablet strength is required for the second authority prescription, please contact the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) for further advice.

The approved second authority prescription will be returned to the prescriber by the Department of Human Services two weeks after the date of the approval of the first authority prescription, to allow for the uninterrupted completion of the six months initial treatment course. The Department of Human Services will contact prescribers prior to dispatch of the second authority prescription to confirm the tablet strength required for the patient.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 3 (change or re-commencement of therapy for all patients)

Clinical criteria:

- Patient must have idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH or PAH secondary to connective tissue disease or PAH associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) and must wish to re-commence PBS-subsidised therapy with this agent after a break in therapy and must have demonstrated a response to their most recent course of PBS-subsidised treatment with this agent; OR
- Patient must have idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH or PAH secondary to connective tissue disease or PAH associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) and whose most recent course of PBS-subsidised treatment was with a PAH agent other than this agent, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

- (1) two completed authority prescription forms; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form; and
- (3) the results of the patient's response to treatment with their last course of PBS-subsidised PAH agent.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

Response to a PAH agent is defined as follows:

For patients with two or more baseline tests, response to treatment is defined as two or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

Approvals for the first authority prescription will be limited to 1 month of therapy with the 62.5 mg strength tablet, with the quantity approved based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information. No repeats will be authorised for this prescription.

The second authority prescription may be written for either the 62.5 mg tablet or the 125 mg tablet strengths. Approvals for the second authority prescription will be limited to 1 month of treatment, with the quantity approved based on the dosage recommendations in the TGA-approved Product Information, and a maximum of 4 repeats.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Swapping between PAH agents: Patients can access PAH agents through the PBS according to the relevant restrictions. Once these patients are approved initial treatment with 1 of these 8 drugs, they may swap between PAH agents at any time without having to re-qualify for treatment with the alternate agent. This means that patients may commence treatment with the alternate agent, subject to that agent's restriction, irrespective of the severity of their disease at the time the application to swap therapy is submitted. It also means that no new baseline measurements will be necessary. New baselines may be submitted where the patient has failed to respond to their current treatment. Eligible patients may only swap between PAH agents if they have not failed prior PBS-subsidised treatment with that agent. For eligible patients, applications to swap between the 8 PAH agents must be made under the relevant initial treatment restriction. Patients should be assessed for response to the treatment they are ceasing at the time the application to swap therapy is being made. Patients who fail to demonstrate a response or for whom no assessment results are submitted with the application to swap therapy may not re-commence PBS-subsidised treatment with the drug they are ceasing.

Note Where the 62.5 mg tablet strength is required for the second authority prescription, please contact the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) for further advice. The approved second authority prescription will be returned to the prescriber by the Department of Human Services two weeks after the date of the approval of the first authority prescription, to allow for the uninterrupted completion of the six months initial treatment course. The Department of Human Services will contact prescribers prior to dispatch of the second authority prescription to confirm the tablet strength required for the patient.

Note Applications for patients who wish to swap to an alternate PAH agent should be accompanied by the previously approved authority prescription, or remaining repeats, for the treatment the patient is ceasing.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au
Applications for authority to prescribe should be forwarded to:
Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 1 or Initial 2 (new patients) or Initial 3 (change or re-commencement of therapy for all patients) or First Continuing treatment - Balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this agent under the Initial 1 (new patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the Initial 2 (new patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the Initial 3 (change or re-commencement of therapy for all patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the First Continuing treatment restriction to complete a maximum of six months of treatment, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition, **AND**
- The treatment must provide no more than the balance of up to six months treatment available under one of the above restrictions.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authorisation under this criterion should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: First Continuing treatment

Clinical criteria:

- Patient must have received a PBS-subsidised initial course of treatment with this agent for this condition, **AND**
- Patient must have been assessed by a physician from a designated hospital to have achieved a response to the PBS-subsidised initial course of treatment, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6 Minute Walk Test (6MWT).

Test requirements to establish response to treatment for continuation of treatment are as follows:

The following list outlines the preferred test combination, in descending order, for the purposes of continuation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments plus 6MWT;
- (2) RHC plus ECHO composite assessments;
- (3) RHC composite assessment plus 6MWT;
- (4) ECHO composite assessment plus 6MWT;
- (5) RHC composite assessment only;
- (6) ECHO composite assessment only.

The results of the same tests as conducted at baseline should be provided with the written First Continuing treatment application, except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a patient specific reason why the test(s) could not be conducted must be provided with the application.

The test results provided with the application for continuing treatment must be no more than 2 months old at the time of application.

Response to a PAH agent is defined as follows:

For patients with two or more baseline tests, response to treatment is defined as two or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats will be authorised.

An application for First Continuing treatment with a PAH agent should be made prior to the completion of the Initial 6 month treatment course to ensure continuity for those patients who respond to treatment, as assessed by the treating physician.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Subsequent Continuing treatment

Clinical criteria:

- Patient must have received a PBS-subsidised treatment under First Continuing treatment with this agent for this condition; OR
- Patient must have previously received PBS-subsidised treatment under this criteria with this agent for this condition, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats will be authorised.

An application for Subsequent Continuing treatment with a PAH agents should be made prior to the completion of the First Continuing treatment course to ensure continuity of treatment.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authorisation under this criterion should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Cessation of treatment (all patients)

Clinical criteria:

- Patient must have received approval for initial PBS-subsidised treatment with this agent, **AND**
- Patient must have not responded to prior PBS-subsidised therapy with this agent, **AND**
- The treatment must be for the purpose of gradual dose reduction prior to ceasing therapy, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment. Treatment beyond 1 month will not be approved.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authorisation under this criterion should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

bosentan 62.5 mg tablet, 60

5618Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	1113.08	^a Bosentan APO [GX] ^a BOSENTAN DR. REDDY'S [RI] ^a Bosentan RBX [RA] ^a BOSLEER [RW]	^a BOSENTAN-DRLA [RZ] ^a Bosentan Mylan [AF] ^a Bosentan Sandoz [SZ] ^a Tracleer [AT]

■ EPOPROSTENOL**Authority required**

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 1 (new patients)

Clinical criteria:

- Patient must not have received prior PBS-subsidised treatment with a pulmonary arterial hypertension (PAH) agent, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**
- Patient must have WHO Functional Class IV idiopathic pulmonary arterial hypertension (iPAH), or anorexigen-induced PAH or hereditary PAH; OR
- Patient must have WHO Functional Class IV pulmonary arterial hypertension secondary to connective tissue disease, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6 Minute Walk Test (6MWT); and
- (3) a signed patient acknowledgement.

Idiopathic pulmonary arterial hypertension, anorexigen-induced pulmonary arterial hypertension, hereditary pulmonary arterial hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) are defined as follows:

- (i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary artery wedge pressure (PAWP) less than 15 mmHg; or
- (ii) where a right heart catheter (RHC) cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.

Test requirements to establish baseline for initiation of treatment are as follows:

The first written application for PBS-subsidised treatment with the first PAH agent should be accompanied by the results of a right heart catheter (RHC) composite assessment plus an echocardiograph (ECHO) composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.

Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments;
- (2) RHC composite assessment plus 6MWT;
- (3) RHC composite assessment only.

In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted for consideration based on the results of the following test combinations, which are listed in descending order of preference:

- (1) ECHO composite assessment plus 6MWT;
- (2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats may be requested.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 2 (change or re-commencement of therapy for all patients)

Clinical criteria:

- Patient must have idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH or PAH secondary to connective tissue disease and must wish to re-commence PBS-subsidised therapy with this agent after a break in therapy and must have demonstrated a response to their most recent course of PBS-subsidised treatment with this agent; OR
- Patient must have WHO Functional Class IV idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH or PAH secondary to connective tissue disease and must have received prior treatment with a PBS-subsidised PAH agent other than this agent; OR
- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH or PAH secondary to connective tissue disease and must have failed to respond to a prior PBS-subsidised PAH agent, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form; and
- (3) the results of the patient's response to treatment with their last course of PBS-subsidised PAH agent; and
- (4) for WHO Functional Class III patients, where this is the first application for this agent, assessment details of the PBS-subsidised PAH agent they have failed to respond to.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

Response to a PAH agent is defined as follows:

For patients with two or more baseline tests, response to treatment is defined as two or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats may be requested.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Swapping between PAH agents: Patients can access PAH agents through the PBS according to the relevant restrictions. Once these patients are approved initial treatment with 1 of these 8 drugs, they may swap between PAH agents at any time without having to re-qualify for treatment with the alternate agent. This means that patients may commence treatment with the alternate agent, subject to that agent's restriction, irrespective of the severity of their disease at the time the application to swap therapy is submitted. It also means that no new baseline measurements will be necessary. New baselines may be submitted where the patient has failed to respond to their current treatment. Eligible patients may only swap between PAH agents if they have not failed prior PBS-subsidised treatment with that agent. For eligible patients, applications to swap between the 8 PAH agents must be made under the relevant initial treatment restriction. Patients should be assessed for response to the treatment they are ceasing at the time the application to swap therapy is being made. Patients who fail to demonstrate a response or for whom no assessment results are submitted with the application to swap therapy may not re-commence PBS-subsidised treatment with the drug they are ceasing.

Note Applications for patients who wish to swap to an alternate PAH agent should be accompanied by the previously approved authority prescription, or remaining repeats, for the treatment the patient is ceasing.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Caution This is a category X drug and must not be given to pregnant women. Pregnancy must be avoided during treatment and for at least 3 months following cessation of therapy.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 1 (new patients) or Initial 2 (change or re-commencement of therapy for all patients) or First Continuing treatment - Balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this agent under the Initial 1 (new patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the Initial 2 (change or re-commencement of therapy for all patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the First Continuing treatment restriction to complete a maximum of six months of treatment, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition, **AND**
- The treatment must provide no more than the balance of up to six months treatment available under one of the above restrictions.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authorisation under this criterion should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: First Continuing treatment

Clinical criteria:

- Patient must have received a PBS-subsidised initial course of treatment with this agent for this condition, **AND**
- Patient must have been assessed by a physician from a designated hospital to have achieved a response to the PBS-subsidised initial course of treatment, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

(1) a completed authority prescription form; and

(2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:

(i) RHC composite assessment; and

(ii) ECHO composite assessment; and

(iii) 6 Minute Walk Test (6MWT).

Test requirements to establish response to treatment for continuation of treatment are as follows:

The following list outlines the preferred test combination, in descending order, for the purposes of continuation of PBS-subsidised treatment:

(1) RHC plus ECHO composite assessments plus 6MWT;

(2) RHC plus ECHO composite assessments;

(3) RHC composite assessment plus 6MWT;

(4) ECHO composite assessment plus 6MWT;

(5) RHC composite assessment only;

(6) ECHO composite assessment only.

The results of the same tests as conducted at baseline should be provided with the written First Continuing treatment application, except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a patient specific reason why the test(s) could not be conducted must be provided with the application.

The test results provided with the application for continuing treatment must be no more than 2 months old at the time of application.

Response to a PAH agent is defined as follows:

For patients with two or more baseline tests, response to treatment is defined as two or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats will be authorised.

An application for First Continuing treatment with a PAH agent should be made prior to the completion of the Initial 6 month treatment course to ensure continuity for those patients who respond to treatment, as assessed by the treating physician. Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Subsequent Continuing treatment

Clinical criteria:

- Patient must have received a PBS-subsidised treatment under First Continuing treatment with this agent for this condition; OR
- Patient must have previously received PBS-subsidised treatment under this criteria with this agent for this condition, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats will be authorised.

An application for Subsequent Continuing treatment with a PAH agents should be made prior to the completion of the First Continuing treatment course to ensure continuity of treatment.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authorisation under this criterion should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

epoprostenol 500 microgram injection, 1 vial

10130E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	33.28	Veletri [AT]

epoprostenol 1.5 mg injection [1 vial] (&) inert substance diluent [2 x 50 mL vials], 1 pack

11065J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	59.33	Flolan [GK]

epoprostenol 1.5 mg injection, 1 vial

10117L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	59.33	Veletri [AT]

epoprostenol 500 microgram injection [1 vial] (&) inert substance diluent [2 x 50 mL vials], 1 pack

11090Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	29.67	Flolan [GK]

■ ILOPROST

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 1 (new patients)

Clinical criteria:

- Patient must not have received prior PBS-subsidised treatment with this agent, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**

- Patient must have WHO Functional Class III drug-induced PAH, **AND**
- Patient must have a mean right atrial pressure of 8 mmHg or less as measured by right heart catheterisation (RHC); OR
- Patient must have right ventricular function assessed by echocardiography (ECHO) where a RHC cannot be performed on clinical grounds, **AND**
- Patient must have failed to respond to 6 or more weeks of appropriate vasodilator treatment unless intolerance or a contraindication to such treatment exists, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6 Minute Walk Test (6MWT); and
- (3) a signed patient acknowledgement.

Idiopathic pulmonary arterial hypertension, anorexigen-induced pulmonary arterial hypertension, hereditary pulmonary arterial hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) are defined as follows:

- (i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary artery wedge pressure (PAWP) less than 15 mmHg; or
- (ii) where a right heart catheter (RHC) cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.

Test requirements to establish baseline for initiation of treatment are as follows:

The first written application for PBS-subsidised treatment with the first PAH agent should be accompanied by the results of a right heart catheter (RHC) composite assessment plus an echocardiograph (ECHO) composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.

Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments;
- (2) RHC composite assessment plus 6MWT;
- (3) RHC composite assessment only.

In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted for consideration based on the results of the following test combinations, which are listed in descending order of preference:

- (1) ECHO composite assessment plus 6MWT;
- (2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

Details of prior vasodilator treatment, including the dose and duration of treatment, must be provided at the time of application. Where the patient has an adverse event to a vasodilator or where vasodilator treatment is contraindicated, details of the nature of the adverse event or contraindication according to the Therapeutic Goods Administration (TGA) approved Product Information must also be provided with the application.

Response to prior vasodilator treatment is defined as follows:

For patients with 2 or more baseline tests, response to treatment is defined as 2 or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information.

A maximum of 5 repeats may be requested.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:
Department of Human Services

Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Note Special Pricing Arrangements apply.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 2 (new patients)

Clinical criteria:

- Patient must not have received prior PBS-subsidised treatment with a pulmonary arterial hypertension (PAH) agent, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**
- Patient must have WHO Functional Class III drug-induced PAH and a mean right atrial pressure greater than 8 mmHg, as measured by right heart catheterisation (RHC); OR
- Patient must have WHO Functional Class III drug-induced PAH with right ventricular function assessed by echocardiography (ECHO) where a RHC cannot be performed on clinical grounds; OR
- Patient must have WHO Functional Class IV idiopathic pulmonary arterial hypertension (iPAH), or anorexigen-induced PAH or hereditary PAH; OR
- Patient must have WHO Functional Class IV pulmonary arterial hypertension secondary to connective tissue disease; OR
- Patient must have WHO Functional Class IV drug-induced PAH, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:

- (i) RHC composite assessment; and
- (ii) ECHO composite assessment; and
- (iii) 6 Minute Walk Test (6MWT); and
- (3) a signed patient acknowledgement.

Idiopathic pulmonary arterial hypertension, anorexigen-induced pulmonary arterial hypertension, hereditary pulmonary arterial hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) are defined as follows:

- (i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary artery wedge pressure (PAWP) less than 15 mmHg; or
- (ii) where a right heart catheter (RHC) cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.

Test requirements to establish baseline for initiation of treatment are as follows:

The first written application for PBS-subsidised treatment with the first PAH agent should be accompanied by the results of a right heart catheter (RHC) composite assessment plus an echocardiograph (ECHO) composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.

Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments;
- (2) RHC composite assessment plus 6MWT;
- (3) RHC composite assessment only.

In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted for consideration based on the results of the following test combinations, which are listed in descending order of preference:

- (1) ECHO composite assessment plus 6MWT;
- (2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats may be requested.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Note Special Pricing Arrangements apply.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 3 (change or re-commencement of therapy for all patients)

Clinical criteria:

- Patient must have idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH or PAH secondary to connective tissue disease or drug-induced PAH and must wish to re-commence PBS-subsidised therapy with this agent after a break in therapy and must have demonstrated a response to their most recent course of PBS-subsidised treatment with this agent; OR
- Patient must have WHO Functional Class IV idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH or PAH secondary to connective tissue disease and must have received prior treatment with a PBS-subsidised PAH agent other than this agent; OR
- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH or PAH secondary to connective tissue disease and must have failed to respond to a prior PBS-subsidised PAH agent, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form; and
- (3) the results of the patient's response to treatment with their last course of PBS-subsidised PAH agent; and
- (4) for WHO Functional Class III patients, where this is the first application for this agent, assessment details of the PBS-subsidised PAH agent they have failed to respond to.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

Response to a PAH agent is defined as follows:

For patients with two or more baseline tests, response to treatment is defined as two or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats may be requested.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Swapping between PAH agents: Patients can access PAH agents through the PBS according to the relevant restrictions.

Once these patients are approved initial treatment with 1 of these 8 drugs, they may swap between PAH agents at any time without having to re-qualify for treatment with the alternate agent. This means that patients may commence treatment with the alternate agent, subject to that agent's restriction, irrespective of the severity of their disease at the time the application to swap therapy is submitted. It also means that no new baseline measurements will be necessary. New baselines may be submitted where the patient has failed to respond to their current treatment. Eligible patients may only swap between PAH agents if they have not failed prior PBS-subsidised treatment with that agent. For eligible patients, applications to swap between the 8 PAH agents must be made under the relevant initial treatment restriction. Patients should be assessed for response to the treatment they are ceasing at the time the application to swap therapy is being made. Patients who fail to demonstrate a response or for whom no assessment results are submitted with the application to swap therapy may not re-commence PBS-subsidised treatment with the drug they are ceasing.

Note Applications for patients who wish to swap to an alternate PAH agent should be accompanied by the previously approved authority prescription, or remaining repeats, for the treatment the patient is ceasing.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs

Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Note Special Pricing Arrangements apply.

Caution This is a category X drug and must not be given to pregnant women. Pregnancy must be avoided during treatment and for at least 3 months following cessation of therapy.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 1 or Initial 2 (new patients) or Initial 3 (change or re-commencement of therapy for all patients) or First Continuing treatment - Balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this agent under the Initial 1 (new patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the Initial 2 (new patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the Initial 3 (change or re-commencement of therapy for all patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the First Continuing treatment restriction to complete a maximum of six months of treatment, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition, **AND**
- The treatment must provide no more than the balance of up to six months treatment available under one of the above restrictions.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authorisation under this criterion should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: First Continuing treatment

Clinical criteria:

- Patient must have received a PBS-subsidised initial course of treatment with this agent for this condition, **AND**
- Patient must have been assessed by a physician from a designated hospital to have achieved a response to the PBS-subsidised initial course of treatment, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6 Minute Walk Test (6MWT).

Test requirements to establish response to treatment for continuation of treatment are as follows:

The following list outlines the preferred test combination, in descending order, for the purposes of continuation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments plus 6MWT;
- (2) RHC plus ECHO composite assessments;
- (3) RHC composite assessment plus 6MWT;
- (4) ECHO composite assessment plus 6MWT;
- (5) RHC composite assessment only;
- (6) ECHO composite assessment only.

The results of the same tests as conducted at baseline should be provided with the written First Continuing treatment application, except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a patient specific reason why the test(s) could not be conducted must be provided with the application.

The test results provided with the application for continuing treatment must be no more than 2 months old at the time of application.

Response to a PAH agent is defined as follows:

For patients with two or more baseline tests, response to treatment is defined as two or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats will be authorised.

An application for First Continuing treatment with a PAH agent should be made prior to the completion of the Initial 6 month treatment course to ensure continuity for those patients who respond to treatment, as assessed by the treating physician.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Subsequent Continuing treatment

Clinical criteria:

- Patient must have received a PBS-subsidised treatment under First Continuing treatment with this agent for this condition; OR
- Patient must have previously received PBS-subsidised treatment under this criteria with this agent for this condition, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats will be authorised.

An application for Subsequent Continuing treatment with a PAH agents should be made prior to the completion of the First Continuing treatment course to ensure continuity of treatment.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authorisation under this criterion should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

iloprost 20 microgram/2 mL inhalation solution, 30 x 2 mL ampoules

5751Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	367.99	Ventavis [BN]

■ MACITENTAN

Caution This is a category X drug and must not be given to pregnant women. Pregnancy must be avoided during treatment and for at least 3 months following cessation of therapy.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 1 (new patients)

Clinical criteria:

- Patient must not have received prior PBS-subsidised treatment with a pulmonary arterial hypertension (PAH) agent, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**
- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH; OR
- Patient must have WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease, **AND**
- Patient must have a mean right atrial pressure of 8 mmHg or less as measured by right heart catheterisation (RHC); OR

- Patient must have right ventricular function assessed by echocardiography (ECHO) where a RHC cannot be performed on clinical grounds, **AND**
- Patient must have failed to respond to 6 or more weeks of appropriate vasodilator treatment unless intolerance or a contraindication to such treatment exists, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:

- (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6 Minute Walk Test (6MWT); and
- (3) a signed patient acknowledgement.

Idiopathic pulmonary arterial hypertension, anorexigen-induced pulmonary arterial hypertension, hereditary pulmonary arterial hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) are defined as follows:

- (i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary artery wedge pressure (PAWP) less than 15 mmHg; or
- (ii) where a right heart catheter (RHC) cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.

Test requirements to establish baseline for initiation of treatment are as follows:

The first written application for PBS-subsidised treatment with the first PAH agent should be accompanied by the results of a right heart catheter (RHC) composite assessment plus an echocardiograph (ECHO) composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.

Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments;
- (2) RHC composite assessment plus 6MWT;
- (3) RHC composite assessment only.

In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted for consideration based on the results of the following test combinations, which are listed in descending order of preference:

- (1) ECHO composite assessment plus 6MWT;
- (2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

Details of prior vasodilator treatment, including the dose and duration of treatment, must be provided at the time of application. Where the patient has an adverse event to a vasodilator or where vasodilator treatment is contraindicated, details of the nature of the adverse event or contraindication according to the Therapeutic Goods Administration (TGA) approved Product Information must also be provided with the application.

Response to prior vasodilator treatment is defined as follows:

For patients with 2 or more baseline tests, response to treatment is defined as 2 or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information.

A maximum of 5 repeats may be requested.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Prior Written Approval of Complex Drugs

Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 2 (new patients)

Clinical criteria:

- Patient must not have received prior PBS-subsidised treatment with a pulmonary arterial hypertension (PAH) agent, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**
- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH), or anorexigen-induced PAH or hereditary PAH, and a mean right atrial pressure of greater than 8 mmHg, as measured by right heart catheterisation (RHC); OR
- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH), or anorexigen-induced PAH or hereditary PAH, with right ventricular function assessed by echocardiography (ECHO) where a RHC cannot be performed on clinical grounds; OR
- Patient must have WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease and a mean right atrial pressure greater than 8 mmHg, as measured by RHC; OR
- Patient must have WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease with right ventricular function assessed by ECHO where a RHC cannot be performed on clinical grounds; OR
- Patient must have WHO Functional Class IV idiopathic pulmonary arterial hypertension (iPAH), or anorexigen-induced PAH or hereditary PAH; OR
- Patient must have WHO Functional Class IV pulmonary arterial hypertension secondary to connective tissue disease; OR
- Patient must have WHO Functional Class III or IV pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology), **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6 Minute Walk Test (6MWT); and
- (3) a signed patient acknowledgement.

Idiopathic pulmonary arterial hypertension, anorexigen-induced pulmonary arterial hypertension, hereditary pulmonary arterial hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) are defined as follows:

- (i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary artery wedge pressure (PAWP) less than 15 mmHg; or
- (ii) where a right heart catheter (RHC) cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.

Test requirements to establish baseline for initiation of treatment are as follows:

The first written application for PBS-subsidised treatment with the first PAH agent should be accompanied by the results of a right heart catheter (RHC) composite assessment plus an echocardiograph (ECHO) composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.

Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments;
- (2) RHC composite assessment plus 6MWT;
- (3) RHC composite assessment only.

In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted for consideration based on the results of the following test combinations, which are listed in descending order of preference:

- (1) ECHO composite assessment plus 6MWT;
- (2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats may be requested.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au
Applications for authority to prescribe should be forwarded to:
Department of Human Services
Prior Written Approval of Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 3 (change or re-commencement of therapy for all patients)

Clinical criteria:

- Patient must have idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH or PAH secondary to connective tissue disease or PAH associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) and must wish to re-commence PBS-subsidised therapy with this agent after a break in therapy and must have demonstrated a response to their most recent course of PBS-subsidised treatment with this agent; OR
- Patient must have idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH or PAH secondary to connective tissue disease or PAH associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) and whose most recent course of PBS-subsidised treatment was with a PAH agent other than this agent, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form; and
- (3) the results of the patient's response to treatment with their last course of PBS-subsidised PAH agent.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

Response to a PAH agent is defined as follows:

For patients with two or more baseline tests, response to treatment is defined as two or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

A maximum of 5 repeats will be authorised.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Swapping between PAH agents: Patients can access PAH agents through the PBS according to the relevant restrictions. Once these patients are approved initial treatment with 1 of these 8 drugs, they may swap between PAH agents at any time without having to re-qualify for treatment with the alternate agent. This means that patients may commence treatment with the alternate agent, subject to that agent's restriction, irrespective of the severity of their disease at the time the application to swap therapy is submitted. It also means that no new baseline measurements will be necessary. New baselines may be submitted where the patient has failed to respond to their current treatment. Eligible patients may only swap between PAH agents if they have not failed prior PBS-subsidised treatment with that agent. For eligible patients, applications to swap between the 8 PAH agents must be made under the relevant initial treatment restriction. Patients should be assessed for response to the treatment they are ceasing at the time the application to swap therapy is being made. Patients who fail to demonstrate a response or for whom no assessment results are submitted with the application to swap therapy may not re-commence PBS-subsidised treatment with the drug they are ceasing.

Note Applications for patients who wish to swap to an alternate PAH agent should be accompanied by the previously approved authority prescription, or remaining repeats, for the treatment the patient is ceasing.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au
Applications for authority to prescribe should be forwarded to:
Department of Human Services

Prior Written Approval of Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 1 or Initial 2 (new patients) or Initial 3 (change or re-commencement of therapy for all patients) or First Continuing treatment - Balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this agent under the Initial 1 (new patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the Initial 2 (new patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the Initial 3 (change or re-commencement of therapy for all patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the First Continuing treatment restriction to complete a maximum of six months of treatment, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition, **AND**
- The treatment must provide no more than the balance of up to six months treatment available under one of the above restrictions.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authorisation under this criterion should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: First Continuing treatment

Clinical criteria:

- Patient must have received a PBS-subsidised initial course of treatment with this agent for this condition, **AND**
- Patient must have been assessed by a physician from a designated hospital to have achieved a response to the PBS-subsidised initial course of treatment, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6 Minute Walk Test (6MWT).

Test requirements to establish response to treatment for continuation of treatment are as follows:

The following list outlines the preferred test combination, in descending order, for the purposes of continuation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments plus 6MWT;
- (2) RHC plus ECHO composite assessments;
- (3) RHC composite assessment plus 6MWT;
- (4) ECHO composite assessment plus 6MWT;
- (5) RHC composite assessment only;
- (6) ECHO composite assessment only.

The results of the same tests as conducted at baseline should be provided with the written First Continuing treatment application, except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a patient specific reason why the test(s) could not be conducted must be provided with the application.

The test results provided with the application for continuing treatment must be no more than 2 months old at the time of application.

Response to a PAH agent is defined as follows:

For patients with two or more baseline tests, response to treatment is defined as two or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats will be authorised.

An application for First Continuing treatment with a PAH agent should be made prior to the completion of the Initial 6 month treatment course to ensure continuity for those patients who respond to treatment, as assessed by the treating physician.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Subsequent Continuing treatment

Clinical criteria:

- Patient must have received a PBS-subsidised treatment under First Continuing treatment with this agent for this condition; OR
- Patient must have previously received PBS-subsidised treatment under this criteria with this agent for this condition, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats will be authorised.

An application for Subsequent Continuing treatment with a PAH agents should be made prior to the completion of the First Continuing treatment course to ensure continuity of treatment.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authorisation under this criterion should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

macitentan 10 mg tablet, 30

10136L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	2876.47	Opsumit [AT]

▪ **RIOCIGUAT**

Caution This is a category X drug and must not be given to pregnant women. Pregnancy must be avoided during treatment and for at least 1 month following cessation of therapy, as recommended by the TGA-approved Product Information.

Note Special Pricing Arrangements apply.

Authority required

Chronic thromboembolic pulmonary hypertension (CTEPH)

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have WHO Functional Class II, III or IV CTEPH, **AND**
- The condition must be inoperable by pulmonary endarterectomy; OR
- The condition must be recurrent or persistent following pulmonary endarterectomy, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Treatment criteria:

- Must be treated in a centre with expertise in the management of CTEPH.

Population criteria:

- Patient must be aged 18 years or older.

CTEPH that is inoperable by pulmonary endarterectomy is defined as follows:

- Right heart catheterisation (RHC) demonstrating pulmonary vascular resistance (PVR) of greater than 300 dyn*sec*cm⁻⁵ measured at least 90 days after start of full anticoagulation; and
- A mean pulmonary artery pressure (PAPmean) of greater than 25 mmHg at least 90 days after start of full anticoagulation.

CTEPH that is recurrent or persistent subsequent to pulmonary endarterectomy is defined as follows:

- RHC demonstrating a PVR of greater than 300 dyn*sec*cm⁻⁵ measured at least 180 days following pulmonary endarterectomy.

Where a RHC cannot be performed due to right ventricular dysfunction, an echocardiogram demonstrating the dysfunction must be provided at the time of application.

Applications for authorisation must be in writing and must include: (1) completed authority prescription forms sufficient for dose titration; and (2) a completed CTEPH PBS Initial Authority Application - Supporting Information form which includes results from the 3 tests below, to establish baseline measurements, where available: (i) RHC composite assessment, and (ii) ECHO composite assessment, and (iii) 6 Minute Walk Test (6MWT); and (3) a signed patient acknowledgment form; and (4) confirmation of evidence of inoperable CTEPH including results of a pulmonary vascular resistance (PVR), a mean pulmonary artery pressure (PAPmean) and the starting date of full anticoagulation; or (5) confirmation of evidence of recurrent or persistent CTEPH including result of PVR and the date that pulmonary endarterectomy was performed; or (6) confirmation of an echocardiogram demonstrating right ventricular dysfunction.

Where it is not possible to perform all 3 tests above on clinical grounds, applications may be submitted for consideration based on the results of the following test combinations, which are listed in descending order of preference: (1) RHC plus ECHO composite assessments; (2) RHC composite assessment plus 6MWT; (3) RHC composite assessment only.

In circumstance where a RHC cannot be performed on clinical grounds, applications may be submitted for consideration based on the results of the following test combinations, which are listed in descending order of preference: (1) ECHO composite assessment plus 6MWT; (2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

Prescriptions for dose titration must provide sufficient quantity for dose titrations by 0.5 mg increments at 2-week intervals to achieve up to a maximum of 2.5 mg three times daily based on the dosage recommendations for initiation of treatment in the TGA-approved Product Information. No repeats will be authorised for these prescriptions.

Approvals for subsequent authority prescription will be limited to 1 month of treatment. The quantity approved must be based on the dosage recommendations in the TGA-approved Product Information, and a maximum of 3 repeats.

The assessment of the patient's response to the initial 20-week course of treatment should be made following the preceding 16 weeks of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Chronic thromboembolic pulmonary hypertension (CTEPH)

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must demonstrate stable or responding disease, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Treatment criteria:

- Must be treated in a centre with expertise in the management of CTEPH.

Population criteria:

- Patient must be aged 18 years or older.

Applications for authorisation must be in writing and must include: (1) a completed authority prescription form; and (2) a completed CTEPH PBS Continuing Authority Application - Supporting Information form which includes results from the three tests below, where available: (i) RHC composite assessment; and (ii) ECHO composite assessment; and (iii) 6 Minute Walk Test (6MWT).

Test requirements to establish response to treatment for continuation of treatment are as follows:

The following list outlines the preferred test combination, in descending order, for the purposes of continuation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments plus 6MWT;
- (2) RHC plus ECHO composite assessments;
- (3) RHC composite assessment plus 6MWT;
- (4) ECHO composite assessment plus 6MWT;
- (5) RHC composite assessment only;
- (6) ECHO composite assessment only.

The results of the same tests as conducted at baseline should be provided with each written continuing treatment application (i.e., every 6 months), except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a patient specific reason why the test(s) could not be conducted must be provided with the application. The test results provided with the application for continuing treatment must be no more than 2 months old at the time of application.

Response to this drug is defined as follows:

For patients with two or more baseline tests, response to treatment is defined as two or more tests demonstrating stability or improvement of disease.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease.

The assessment of the patient's response to the continuing 6 month courses of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

The maximum quantity per prescription must be based on the dosage recommendations in the TGA-approved Product Information and be limited to provide sufficient supply for 1 month of treatment.

A maximum of 5 repeats will be authorised.

Applications for continuing treatment with this drug should be made two weeks prior to the completion of the 6-month treatment course to ensure continuity for those patients who respond to treatment, as assessed by the treating physician.

Patients who fail to demonstrate disease stability or improvement to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Chronic thromboembolic pulmonary hypertension (CTEPH)

Treatment Phase: Balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the Initial treatment restriction to complete a maximum of 20 weeks of treatment; OR
- Patient must have received insufficient therapy with this drug under the Continuing treatment restriction to complete a maximum of 24 weeks of treatment, **AND**
- The treatment must provide no more than the balance of up to 20 or 24 weeks of treatment available under the above respective restriction, **AND**
- The treatment must be the sole PBS-subsidised agent for this condition.

Treatment criteria:

- Must be treated in a centre with expertise in the management of CTEPH.

Population criteria:

- Patient must be aged 18 years or older.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

riociguat 2.5 mg tablet, 42

11002C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	1717.71	Adempas [BN]

riociguat 1 mg tablet, 84

11020B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	3435.42	Adempas [BN]

riociguat 1 mg tablet, 42

10976Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	1717.71	Adempas [BN]

riociguat 500 microgram tablet, 84

10995Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	3435.42	Adempas [BN]

riociguat 2 mg tablet, 84

11013P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	3435.42	Adempas [BN]

riociguat 2 mg tablet, 42

10984D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	1717.71	Adempas [BN]

riociguat 500 microgram tablet, 42

11001B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	1717.71	Adempas [BN]

riociguat 1.5 mg tablet, 84

10977R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	3435.42	Adempas [BN]

riociguat 2.5 mg tablet, 84

11019Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	3435.42	Adempas [BN]

riociguat 1.5 mg tablet, 42

10989J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	1717.71	Adempas [BN]

■ RIOCIQUAT

Caution This is a category X drug and must not be given to pregnant women. Pregnancy must be avoided during treatment and for at least 1 month following cessation of therapy, as recommended by the TGA-approved Product Information.

Note Special Pricing Arrangements apply.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 1 (new patients)

Clinical criteria:

- Patient must not have received prior PBS-subsidised treatment with a pulmonary arterial hypertension (PAH) agent, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**
- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH; OR
- Patient must have WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease, **AND**
- Patient must have a mean right atrial pressure of 8 mmHg or less as measured by right heart catheterisation (RHC); OR
- Patient must have right ventricular function assessed by echocardiography (ECHO) where a RHC cannot be performed on clinical grounds, **AND**
- Patient must have failed to respond to 6 or more weeks of appropriate vasodilator treatment unless intolerance or a contraindication to such treatment exists, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include: (1) completed authority prescription forms sufficient for dose titration; and (2) a completed Pulmonary Arterial Hypertension Initial PBS Authority Application - Supporting Information form which includes results from the three tests below, where available: (i) RHC composite assessment; and (ii) ECHO composite assessment; and (iii) 6 Minute Walk Test (6MWT); and (3) a signed patient acknowledgement. Idiopathic pulmonary arterial hypertension, anorexigen-induced pulmonary arterial hypertension, hereditary pulmonary arterial hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) are defined as follows: (i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary artery wedge pressure (PAWP) less than 15 mmHg; or (ii) where a right heart catheter (RHC) cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.

Test requirements to establish baseline for initiation of treatment are as follows:

The first written application for PBS-subsidised treatment with the first PAH agent should be accompanied by the results of a right heart catheter (RHC) composite assessment plus an echocardiograph (ECHO) composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.

Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments;
- (2) RHC composite assessment plus 6MWT;
- (3) RHC composite assessment only.

In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted for consideration based on the results of the following test combinations, which are listed in descending order of preference:

- (1) ECHO composite assessment plus 6MWT;
- (2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

Details of prior vasodilator treatment, including the dose and duration of treatment, must be provided at the time of application. Where the patient has an adverse event to a vasodilator or where vasodilator treatment is contraindicated, details of the nature of the adverse event or contraindication according to the Therapeutic Goods Administration (TGA) approved Product Information must also be provided with the application.

Response to prior vasodilator treatment is defined as follows:

For patients with 2 or more baseline tests, response to treatment is defined as 2 or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

Approvals for prescriptions for dose titration will provide sufficient quantity for dose titrations by 0.5 mg increments at 2-week intervals to achieve up to a maximum of 2.5 mg three times daily based on the dosage recommendations for initiation of treatment in the TGA-approved Product Information. No repeats will be authorised for these prescriptions.

Approvals for subsequent authority prescription will be limited to 1 month of treatment, with the quantity approved based on the dosage recommendations in the TGA-approved Product Information, and a maximum of 4 repeats.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 2 (new patients)

Clinical criteria:

- Patient must not have received prior PBS-subsidised treatment with a pulmonary arterial hypertension (PAH) agent, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**
- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH), or anorexigen-induced PAH or hereditary PAH, and a mean right atrial pressure of greater than 8 mmHg, as measured by right heart catheterisation (RHC); OR
- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH), or anorexigen-induced PAH or hereditary PAH, with right ventricular function assessed by echocardiography (ECHO) where a RHC cannot be performed on clinical grounds; OR
- Patient must have WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease and a mean right atrial pressure greater than 8 mmHg, as measured by RHC; OR
- Patient must have WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease with right ventricular function assessed by ECHO where a RHC cannot be performed on clinical grounds; OR
- Patient must have WHO Functional Class IV idiopathic pulmonary arterial hypertension (iPAH), or anorexigen-induced PAH or hereditary PAH; OR
- Patient must have WHO Functional Class IV pulmonary arterial hypertension secondary to connective tissue disease; OR
- Patient must have WHO Functional Class III or IV pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology), **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include: (1) completed authority prescription forms sufficient for dose titration; and (2) a completed Pulmonary Arterial Hypertension Initial PBS Authority Application - Supporting Information form which includes results from the three tests below, where available: (i) RHC composite assessment; and (ii) ECHO composite assessment; and (iii) 6 Minute Walk Test (6MWT); and (3) a signed patient acknowledgement. Idiopathic pulmonary arterial hypertension, anorexigen-induced pulmonary arterial hypertension, hereditary pulmonary arterial hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) are defined as follows: (i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary artery wedge pressure (PAWP) less than 15 mmHg; or (ii) where a right heart catheter

(RHC) cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.

Test requirements to establish baseline for initiation of treatment are as follows:

The first written application for PBS-subsidised treatment with the first PAH agent should be accompanied by the results of a right heart catheter (RHC) composite assessment plus an echocardiograph (ECHO) composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.

Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments;
- (2) RHC composite assessment plus 6MWT;
- (3) RHC composite assessment only.

In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted for consideration based on the results of the following test combinations, which are listed in descending order of preference:

- (1) ECHO composite assessment plus 6MWT;
- (2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

Approvals for prescriptions for dose titration will provide sufficient quantity for dose titrations by 0.5 mg increments at 2-week intervals to achieve up to a maximum of 2.5 mg three times daily based on the dosage recommendations for initiation of treatment in the TGA-approved Product Information. No repeats will be authorised for these prescriptions.

Approvals for subsequent authority prescription will be limited to 1 month of treatment, with the quantity approved based on the dosage recommendations in the TGA-approved Product Information, and a maximum of 4 repeats.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 3 (change or re-commencement of therapy for all patients)

Clinical criteria:

- Patient must have idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH or PAH secondary to connective tissue disease or PAH associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) and must wish to re-commence PBS-subsidised therapy with this agent after a break in therapy and must have demonstrated a response to their most recent course of PBS-subsidised treatment with this agent; OR
- Patient must have idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH or PAH secondary to connective tissue disease or PAH associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) and whose most recent course of PBS-subsidised treatment was with a PAH agent other than this agent, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include: (1) completed authority prescription forms sufficient for dose titration; and (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form; and (3) the results of the patient's response to treatment with their last course of PBS-subsidised PAH agent. Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application. The test results provided must not be more than 2 months old at the time of application.

Response to a PAH agent is defined as follows:

For patients with two or more baseline tests, response to treatment is defined as two or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

Approvals for prescriptions for dose titration will provide sufficient quantity for dose titrations by 0.5 mg increments at 2-week intervals to achieve up to a maximum of 2.5 mg three times daily based on the dosage recommendations for initiation of treatment in the TGA-approved Product Information. No repeats will be authorised for these prescriptions.

Approvals for subsequent authority prescription will be limited to 1 month of treatment, with the quantity approved based on the dosage recommendations in the TGA-approved Product Information, and a maximum of 4 repeats.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Swapping between PAH agents: Patients can access PAH agents through the PBS according to the relevant restrictions.

Once these patients are approved initial treatment with 1 of these 8 drugs, they may swap between PAH agents at any time without having to re-qualify for treatment with the alternate agent. This means that patients may commence treatment with the alternate agent, subject to that agent's restriction, irrespective of the severity of their disease at the time the application to swap therapy is submitted. It also means that no new baseline measurements will be necessary. New baselines may be submitted where the patient has failed to respond to their current treatment. Eligible patients may only swap between PAH agents if they have not failed prior PBS-subsidised treatment with that agent. For eligible patients, applications to swap between the 8 PAH agents must be made under the relevant initial treatment restriction. Patients should be assessed for response to the treatment they are ceasing at the time the application to swap therapy is being made. Patients who fail to demonstrate a response or for whom no assessment results are submitted with the application to swap therapy may not re-commence PBS-subsidised treatment with the drug they are ceasing.

Note Applications for patients who wish to swap to an alternate PAH agent should be accompanied by the previously approved authority prescription, or remaining repeats, for the treatment the patient is ceasing.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 1 or Initial 2 (new patients) or Initial 3 (change or re-commencement of therapy for all patients) or First Continuing treatment - Balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this agent under the Initial 1 (new patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the Initial 2 (new patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the Initial 3 (change or re-commencement of therapy for all patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the First Continuing treatment restriction to complete a maximum of six months of treatment, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition, **AND**
- The treatment must provide no more than the balance of up to six months treatment available under one of the above restrictions.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: First Continuing treatment

Clinical criteria:

- Patient must have received a PBS-subsidised initial course of treatment with this agent for this condition, **AND**
- Patient must have been assessed by a physician from a designated hospital to have achieved a response to the PBS-subsidised initial course of treatment, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

(1) a completed authority prescription form; and

(2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:

(i) RHC composite assessment; and

- (ii) ECHO composite assessment; and
- (iii) 6 Minute Walk Test (6MWT).

Test requirements to establish response to treatment for continuation of treatment are as follows:

The following list outlines the preferred test combination, in descending order, for the purposes of continuation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments plus 6MWT;
- (2) RHC plus ECHO composite assessments;
- (3) RHC composite assessment plus 6MWT;
- (4) ECHO composite assessment plus 6MWT;
- (5) RHC composite assessment only;
- (6) ECHO composite assessment only.

The results of the same tests as conducted at baseline should be provided with the written First Continuing treatment application, except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a patient specific reason why the test(s) could not be conducted must be provided with the application.

The test results provided with the application for continuing treatment must be no more than 2 months old at the time of application.

Response to a PAH agent is defined as follows:

For patients with two or more baseline tests, response to treatment is defined as two or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats will be authorised.

An application for First Continuing treatment with a PAH agent should be made two weeks prior to the completion of the Initial 6 month treatment course to ensure continuity for those patients who respond to treatment, as assessed by the treating physician.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Subsequent Continuing treatment

Clinical criteria:

- Patient must have received a PBS-subsidised treatment under First Continuing treatment with this agent for this condition; OR
- Patient must have previously received PBS-subsidised treatment under this criteria with this agent for this condition, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats will be authorised.

An application for Subsequent Continuing treatment with a PAH agents should be made prior to the completion of the First Continuing treatment course to ensure continuity of treatment.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

CARDIOVASCULAR SYSTEM

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

riociguat 2.5 mg tablet, 42

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
11057Y	1	..	..	1717.71	Adempas [BN]

riociguat 1 mg tablet, 84

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
11053R	1	..	..	3435.42	Adempas [BN]

riociguat 1 mg tablet, 42

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
11054T	1	..	..	1717.71	Adempas [BN]

riociguat 500 microgram tablet, 84

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
11059C	1	..	..	3435.42	Adempas [BN]

riociguat 2 mg tablet, 84

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
11039B	1	..	..	3435.42	Adempas [BN]

riociguat 2 mg tablet, 42

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
11038Y	1	..	..	1717.71	Adempas [BN]

riociguat 500 microgram tablet, 42

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
11040C	1	..	..	1717.71	Adempas [BN]

riociguat 1.5 mg tablet, 84

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
11048L	1	..	..	3435.42	Adempas [BN]

riociguat 2.5 mg tablet, 84

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
11024F	1	..	..	3435.42	Adempas [BN]

riociguat 1.5 mg tablet, 42

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
11047K	1	..	..	1717.71	Adempas [BN]

■ SILDENAFIL

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 1 (new patients)

Clinical criteria:

- Patient must not have received prior PBS-subsidised treatment with a pulmonary arterial hypertension (PAH) agent, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**
- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH; OR
- Patient must have WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease, **AND**
- Patient must have a mean right atrial pressure of 8 mmHg or less as measured by right heart catheterisation (RHC); OR
- Patient must have right ventricular function assessed by echocardiography (ECHO) where a RHC cannot be performed on clinical grounds, **AND**
- Patient must have failed to respond to 6 or more weeks of appropriate vasodilator treatment unless intolerance or a contraindication to such treatment exists, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

(1) a completed authority prescription form; and

(2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:

(i) RHC composite assessment; and

(ii) ECHO composite assessment; and

- (iii) 6 Minute Walk Test (6MWT); and
- (3) a signed patient acknowledgement.

Idiopathic pulmonary arterial hypertension, anorexigen-induced pulmonary arterial hypertension, hereditary pulmonary arterial hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) are defined as follows:

- (i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary artery wedge pressure (PAWP) less than 15 mmHg; or
- (ii) where a right heart catheter (RHC) cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.

Test requirements to establish baseline for initiation of treatment are as follows:

The first written application for PBS-subsidised treatment with the first PAH agent should be accompanied by the results of a right heart catheter (RHC) composite assessment plus an echocardiograph (ECHO) composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.

Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments;
- (2) RHC composite assessment plus 6MWT;
- (3) RHC composite assessment only.

In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted for consideration based on the results of the following test combinations, which are listed in descending order of preference:

- (1) ECHO composite assessment plus 6MWT;
- (2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

Details of prior vasodilator treatment, including the dose and duration of treatment, must be provided at the time of application. Where the patient has an adverse event to a vasodilator or where vasodilator treatment is contraindicated, details of the nature of the adverse event or contraindication according to the Therapeutic Goods Administration (TGA) approved Product Information must also be provided with the application.

Response to prior vasodilator treatment is defined as follows:

For patients with 2 or more baseline tests, response to treatment is defined as 2 or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information.

A maximum of 5 repeats may be requested.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 2 (new patients)

Clinical criteria:

- Patient must not have received prior PBS-subsidised treatment with a pulmonary arterial hypertension (PAH) agent, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**
- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH), or anorexigen-induced PAH or hereditary PAH, and a mean right atrial pressure of greater than 8 mmHg, as measured by right heart catheterisation (RHC); OR

- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH), or anorexigen-induced PAH or hereditary PAH, with right ventricular function assessed by echocardiography (ECHO) where a RHC cannot be performed on clinical grounds; OR
- Patient must have WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease and a mean right atrial pressure greater than 8 mmHg, as measured by RHC; OR
- Patient must have WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease with right ventricular function assessed by ECHO where a RHC cannot be performed on clinical grounds, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

(1) a completed authority prescription form; and

(2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:

(i) RHC composite assessment; and

(ii) ECHO composite assessment; and

(iii) 6 Minute Walk Test (6MWT); and

(3) a signed patient acknowledgement.

Idiopathic pulmonary arterial hypertension, anorexigen-induced pulmonary arterial hypertension, hereditary pulmonary arterial hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) are defined as follows:

(i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary artery wedge pressure (PAWP) less than 15 mmHg; or

(ii) where a right heart catheter (RHC) cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.

Test requirements to establish baseline for initiation of treatment are as follows:

The first written application for PBS-subsidised treatment with the first PAH agent should be accompanied by the results of a right heart catheter (RHC) composite assessment plus an echocardiograph (ECHO) composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.

Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:

(1) RHC plus ECHO composite assessments;

(2) RHC composite assessment plus 6MWT;

(3) RHC composite assessment only.

In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted for consideration based on the results of the following test combinations, which are listed in descending order of preference:

(1) ECHO composite assessment plus 6MWT;

(2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats may be requested.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 3 (change or re-commencement of therapy for all patients)

Clinical criteria:

- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH or PAH secondary to connective tissue disease and must wish to re-commence PBS-subsidised therapy with this agent after a break in therapy and must have demonstrated a response to their most recent course of PBS-subsidised treatment with this agent; OR

- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH or PAH secondary to connective tissue disease and whose most recent course of PBS-subsidised treatment was with a PAH agent other than this agent, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form; and
- (3) the results of the patient's response to treatment with their last course of PBS-subsidised PAH agent.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

Response to a PAH agent is defined as follows:

For patients with two or more baseline tests, response to treatment is defined as two or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats may be requested.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Swapping between PAH agents: Patients can access PAH agents through the PBS according to the relevant restrictions.

Once these patients are approved initial treatment with 1 of these 8 drugs, they may swap between PAH agents at any time without having to re-qualify for treatment with the alternate agent. This means that patients may commence treatment with the alternate agent, subject to that agent's restriction, irrespective of the severity of their disease at the time the application to swap therapy is submitted. It also means that no new baseline measurements will be necessary. New baselines may be submitted where the patient has failed to respond to their current treatment. Eligible patients may only swap between PAH agents if they have not failed prior PBS-subsidised treatment with that agent. For eligible patients, applications to swap between the 8 PAH agents must be made under the relevant initial treatment restriction. Patients should be assessed for response to the treatment they are ceasing at the time the application to swap therapy is being made. Patients who fail to demonstrate a response or for whom no assessment results are submitted with the application to swap therapy may not re-commence PBS-subsidised treatment with the drug they are ceasing.

Note Applications for patients who wish to swap to an alternate PAH agent should be accompanied by the previously approved authority prescription, or remaining repeats, for the treatment the patient is ceasing.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Caution This is a category X drug and must not be given to pregnant women. Pregnancy must be avoided during treatment and for at least 3 months following cessation of therapy.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 1 or Initial 2 (new patients) or Initial 3 (change or re-commencement of therapy for all patients) or First Continuing treatment - Balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this agent under the Initial 1 (new patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the Initial 2 (new patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the Initial 3 (change or re-commencement of therapy for all patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the First Continuing treatment restriction to complete a maximum of six months of treatment, **AND**

- The treatment must be the sole PBS-subsidised PAH agent for this condition, **AND**
- The treatment must provide no more than the balance of up to six months treatment available under one of the above restrictions.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authorisation under this criterion should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: First Continuing treatment

Clinical criteria:

- Patient must have received a PBS-subsidised initial course of treatment with this agent for this condition, **AND**
- Patient must have been assessed by a physician from a designated hospital to have achieved a response to the PBS-subsidised initial course of treatment, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

(1) a completed authority prescription form; and

(2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:

(i) RHC composite assessment; and

(ii) ECHO composite assessment; and

(iii) 6 Minute Walk Test (6MWT).

Test requirements to establish response to treatment for continuation of treatment are as follows:

The following list outlines the preferred test combination, in descending order, for the purposes of continuation of PBS-subsidised treatment:

(1) RHC plus ECHO composite assessments plus 6MWT;

(2) RHC plus ECHO composite assessments;

(3) RHC composite assessment plus 6MWT;

(4) ECHO composite assessment plus 6MWT;

(5) RHC composite assessment only;

(6) ECHO composite assessment only.

The results of the same tests as conducted at baseline should be provided with the written First Continuing treatment application, except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a patient specific reason why the test(s) could not be conducted must be provided with the application.

The test results provided with the application for continuing treatment must be no more than 2 months old at the time of application.

Response to a PAH agent is defined as follows:

For patients with two or more baseline tests, response to treatment is defined as two or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats will be authorised.

An application for First Continuing treatment with a PAH agent should be made prior to the completion of the Initial 6 month treatment course to ensure continuity for those patients who respond to treatment, as assessed by the treating physician.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Subsequent Continuing treatment

Clinical criteria:

- Patient must have received a PBS-subsidised treatment under First Continuing treatment with this agent for this condition; OR
- Patient must have previously received PBS-subsidised treatment under this criteria with this agent for this condition, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats will be authorised.

An application for Subsequent Continuing treatment with a PAH agents should be made prior to the completion of the First Continuing treatment course to ensure continuity of treatment.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authorisation under this criterion should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

sildenafil 20 mg tablet, 90

9547L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	254.31	^a APO-Sildenafil PHT [TX] ^a Sildenafil AN PHT 20 [EA] ^a Sildenafil Sandoz PHT 20 [SZ]	^a Revatio [PF] ^a SILDENAFIL-DRx [RZ]

▪ **TADALAFIL**

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 1 (new patients)

Clinical criteria:

- Patient must not have received prior PBS-subsidised treatment with a pulmonary arterial hypertension (PAH) agent, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**
- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH; OR
- Patient must have WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease, **AND**
- Patient must have a mean right atrial pressure of 8 mmHg or less as measured by right heart catheterisation (RHC); OR
- Patient must have right ventricular function assessed by echocardiography (ECHO) where a RHC cannot be performed on clinical grounds, **AND**
- Patient must have failed to respond to 6 or more weeks of appropriate vasodilator treatment unless intolerance or a contraindication to such treatment exists, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

(1) a completed authority prescription form; and

(2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:

(i) RHC composite assessment; and

(ii) ECHO composite assessment; and

(iii) 6 Minute Walk Test (6MWT); and

(3) a signed patient acknowledgement.

Idiopathic pulmonary arterial hypertension, anorexigen-induced pulmonary arterial hypertension, hereditary pulmonary arterial hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) are defined as follows:

(i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary artery wedge pressure (PAWP) less than 15 mmHg; or

(ii) where a right heart catheter (RHC) cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.

Test requirements to establish baseline for initiation of treatment are as follows:

The first written application for PBS-subsidised treatment with the first PAH agent should be accompanied by the results of a right heart catheter (RHC) composite assessment plus an echocardiograph (ECHO) composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.

Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments;
- (2) RHC composite assessment plus 6MWT;
- (3) RHC composite assessment only.

In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted for consideration based on the results of the following test combinations, which are listed in descending order of preference:

- (1) ECHO composite assessment plus 6MWT;
- (2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

Details of prior vasodilator treatment, including the dose and duration of treatment, must be provided at the time of application. Where the patient has an adverse event to a vasodilator or where vasodilator treatment is contraindicated, details of the nature of the adverse event or contraindication according to the Therapeutic Goods Administration (TGA) approved Product Information must also be provided with the application.

Response to prior vasodilator treatment is defined as follows:

For patients with 2 or more baseline tests, response to treatment is defined as 2 or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the TGA-approved Product Information.

A maximum of 5 repeats may be requested.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 2 (new patients)

Clinical criteria:

- Patient must not have received prior PBS-subsidised treatment with a pulmonary arterial hypertension (PAH) agent, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**
- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH), or anorexigen-induced PAH or hereditary PAH, and a mean right atrial pressure of greater than 8 mmHg, as measured by right heart catheterisation (RHC); OR
- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH), or anorexigen-induced PAH or hereditary PAH, with right ventricular function assessed by echocardiography (ECHO) where a RHC cannot be performed on clinical grounds; OR
- Patient must have WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease and a mean right atrial pressure greater than 8 mmHg, as measured by RHC; OR
- Patient must have WHO Functional Class III pulmonary arterial hypertension secondary to connective tissue disease with right ventricular function assessed by ECHO where a RHC cannot be performed on clinical grounds, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6 Minute Walk Test (6MWT); and
- (3) a signed patient acknowledgement.

Idiopathic pulmonary arterial hypertension, anorexigen-induced pulmonary arterial hypertension, hereditary pulmonary arterial hypertension, drug-induced pulmonary arterial hypertension, pulmonary arterial hypertension secondary to connective tissue disease including scleroderma, or pulmonary arterial hypertension associated with a congenital systemic-to-pulmonary shunt (including Eisenmenger's physiology) are defined as follows:

- (i) mean pulmonary artery pressure (mPAP) greater than 25 mmHg at rest and pulmonary artery wedge pressure (PAWP) less than 15 mmHg; or
- (ii) where a right heart catheter (RHC) cannot be performed on clinical grounds, right ventricular systolic pressure (RVSP), assessed by echocardiography (ECHO), greater than 40 mmHg, with normal left ventricular function.

Test requirements to establish baseline for initiation of treatment are as follows:

The first written application for PBS-subsidised treatment with the first PAH agent should be accompanied by the results of a right heart catheter (RHC) composite assessment plus an echocardiograph (ECHO) composite assessment, plus a 6 minute walk test (6MWT) to establish the patient's baseline measurements.

Where it is not possible to perform all 3 tests above on clinical grounds, the following list outlines the preferred test combination, in descending order, for the purposes of initiation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments;
- (2) RHC composite assessment plus 6MWT;
- (3) RHC composite assessment only.

In circumstances where a RHC cannot be performed on clinical grounds, applications may be submitted for consideration based on the results of the following test combinations, which are listed in descending order of preference:

- (1) ECHO composite assessment plus 6MWT;
- (2) ECHO composite assessment only.

Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.

The test results provided must not be more than 2 months old at the time of application.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats may be requested.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 3 (change or re-commencement of therapy for all patients)

Clinical criteria:

- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH or PAH secondary to connective tissue disease and must wish to re-commence PBS-subsidised therapy with this agent after a break in therapy and must have demonstrated a response to their most recent course of PBS-subsidised treatment with this agent; OR
- Patient must have WHO Functional Class III idiopathic pulmonary arterial hypertension (iPAH) or anorexigen-induced PAH or hereditary PAH or PAH secondary to connective tissue disease and whose most recent course of PBS-subsidised treatment was with a PAH agent other than this agent, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form; and

(3) the results of the patient's response to treatment with their last course of PBS-subsidised PAH agent.
Where fewer than 3 tests are able to be performed on clinical grounds, a patient specific reason outlining why the particular test(s) could not be conducted must be provided with the authority application.
The test results provided must not be more than 2 months old at the time of application.

Response to a PAH agent is defined as follows:

For patients with two or more baseline tests, response to treatment is defined as two or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats may be requested.

The assessment of the patient's response to the initial 6 month course of treatment should be made following the preceding 5 months of treatment, in order to allow sufficient time for a response to be demonstrated.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Swapping between PAH agents: Patients can access PAH agents through the PBS according to the relevant restrictions. Once these patients are approved initial treatment with 1 of these 8 drugs, they may swap between PAH agents at any time without having to re-qualify for treatment with the alternate agent. This means that patients may commence treatment with the alternate agent, subject to that agent's restriction, irrespective of the severity of their disease at the time the application to swap therapy is submitted. It also means that no new baseline measurements will be necessary. New baselines may be submitted where the patient has failed to respond to their current treatment. Eligible patients may only swap between PAH agents if they have not failed prior PBS-subsidised treatment with that agent. For eligible patients, applications to swap between the 8 PAH agents must be made under the relevant initial treatment restriction. Patients should be assessed for response to the treatment they are ceasing at the time the application to swap therapy is being made. Patients who fail to demonstrate a response or for whom no assessment results are submitted with the application to swap therapy may not re-commence PBS-subsidised treatment with the drug they are ceasing.

Note Applications for patients who wish to swap to an alternate PAH agent should be accompanied by the previously approved authority prescription, or remaining repeats, for the treatment the patient is ceasing.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Caution This is a category X drug and must not be given to pregnant women. Pregnancy must be avoided during treatment and for at least 3 months following cessation of therapy.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Initial 1 or Initial 2 (new patients) or Initial 3 (change or re-commencement of therapy for all patients) or First Continuing treatment - Balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this agent under the Initial 1 (new patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the Initial 2 (new patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the Initial 3 (change or re-commencement of therapy for all patients) restriction to complete a maximum of six months of treatment; OR
- Patient must have received insufficient therapy with this agent under the First Continuing treatment restriction to complete a maximum of six months of treatment, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition, **AND**
- The treatment must provide no more than the balance of up to six months treatment available under one of the above restrictions.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authorisation under this criterion should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826
HOBART TAS 7001

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: First Continuing treatment

Clinical criteria:

- Patient must have received a PBS-subsidised initial course of treatment with this agent for this condition, **AND**
- Patient must have been assessed by a physician from a designated hospital to have achieved a response to the PBS-subsidised initial course of treatment, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Pulmonary Arterial Hypertension PBS Authority Application - Supporting Information form which includes results from the three tests below, where available:
 - (i) RHC composite assessment; and
 - (ii) ECHO composite assessment; and
 - (iii) 6 Minute Walk Test (6MWT).

Test requirements to establish response to treatment for continuation of treatment are as follows:

The following list outlines the preferred test combination, in descending order, for the purposes of continuation of PBS-subsidised treatment:

- (1) RHC plus ECHO composite assessments plus 6MWT;
- (2) RHC plus ECHO composite assessments;
- (3) RHC composite assessment plus 6MWT;
- (4) ECHO composite assessment plus 6MWT;
- (5) RHC composite assessment only;
- (6) ECHO composite assessment only.

The results of the same tests as conducted at baseline should be provided with the written First Continuing treatment application, except for patients who were able to undergo all 3 tests at baseline, and whose subsequent ECHO and 6MWT results demonstrate disease stability or improvement, in which case RHC can be omitted. In all other patients, where the same test(s) conducted at baseline cannot be performed for assessment of response on clinical grounds, a patient specific reason why the test(s) could not be conducted must be provided with the application.

The test results provided with the application for continuing treatment must be no more than 2 months old at the time of application.

Response to a PAH agent is defined as follows:

For patients with two or more baseline tests, response to treatment is defined as two or more tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with a RHC composite assessment alone at baseline, response to treatment is defined as a RHC result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients with an ECHO composite assessment alone at baseline, response to treatment is defined as an ECHO result demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

For patients aged less than 18 years, response to treatment is defined as at least one of the baseline tests demonstrating stability or improvement of disease, as assessed by a physician from a designated hospital.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats will be authorised.

An application for First Continuing treatment with a PAH agent should be made prior to the completion of the Initial 6 month treatment course to ensure continuity for those patients who respond to treatment, as assessed by the treating physician.

Patients who fail to demonstrate a response to PBS-subsidised treatment with this agent at the time where an assessment is required must cease PBS-subsidised therapy with this agent.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

Authority required

Pulmonary arterial hypertension (PAH)

Treatment Phase: Subsequent Continuing treatment

Clinical criteria:

SYSTEMIC HORMONAL PREPARATIONS, EXCL. SEX HORMONES AND INSULINS

- Patient must have received a PBS-subsidised treatment under First Continuing treatment with this agent for this condition; OR
- Patient must have previously received PBS-subsidised treatment under this criteria with this agent for this condition, **AND**
- Patient must have been assessed by a physician at a designated hospital, **AND**
- The treatment must be the sole PBS-subsidised PAH agent for this condition.

The maximum quantity authorised will be limited to provide sufficient supply for 1 month of treatment, based on the dosage recommendations in the Therapeutic Goods Administration (TGA) approved Product Information.

A maximum of 5 repeats will be authorised.

An application for Subsequent Continuing treatment with a PAH agents should be made prior to the completion of the First Continuing treatment course to ensure continuity of treatment.

PAH agents are not PBS-subsidised for patients with pulmonary hypertension secondary to interstitial lung disease associated with connective tissue disease, where the total lung capacity is less than 70% of predicted.

The term 'PAH agents' refers to bosentan monohydrate, iloprost trometamol, epoprostenol sodium, sildenafil citrate, ambrisentan, tadalafil, macitentan, and riociguat.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authorisation under this criterion should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Refer to the Department of Human Services website at www.humanservices.gov.au for a list of designated hospitals.

tadalafil 20 mg tablet, 56

1308W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	796.60	Adcirca [LY]

SYSTEMIC HORMONAL PREPARATIONS, EXCL. SEX HORMONES AND INSULINS

PITUITARY AND HYPOTHALAMIC HORMONES AND ANALOGUES

ANTERIOR PITUITARY LOBE HORMONES AND ANALOGUES

Other anterior pituitary lobe hormones and analogues

PEGVISOMANT

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs Programs
Reply Paid 9826
HOBART TAS 7001

Authority required

Acromegaly

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must not have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have an age- and sex-adjusted insulin-like growth factor 1 (IGF-1) concentration greater than the upper limit of normal (ULN), **AND**
- The treatment must be after failure to achieve biochemical control with a maximum indicated dose of either 30 mg octreotide LAR or 120 mg lanreotide ATG every 28 days for 24 weeks; unless contraindicated or not tolerated according to the TGA approved Product Information, **AND**
- The treatment must not be given concomitantly with a PBS-subsidised somatostatin analogue.

Somatostatin analogues include octreotide, lanreotide and pasireotide

Failure to achieve biochemical control after completion of a prior therapy with either octreotide or lanreotide is defined as:

- 1) Growth hormone level greater than 1 mcg/L or 3 mIU/L; OR
- 2) IGF-1 level is greater than the age- and sex-adjusted ULN.

If treatment with either octreotide or lanreotide is contraindicated according to the relevant TGA-approved Product Information, the application must provide details of contraindication.

If intolerance to either octreotide or lanreotide treatment developed during the relevant period of use which is of a severity to necessitate withdrawal of the treatment, the application must provide details of the nature and severity of this intolerance.

In a patient treated with radiotherapy, pegvisomant should be withdrawn every 2 years in the 10 years after completion of radiotherapy for assessment of remission. Pegvisomant should be withdrawn at least 8 weeks prior to the assessment of remission.

Biochemical evidence of remission is defined as normalisation of sex- and age- adjusted insulin-like growth factor 1 (IGF-1).

Two completed authority prescriptions should be submitted with the initial application for this drug. One prescription should be for the loading dose of 80 mg for a quantity of 4 vials of 20 mg with no repeats. The second prescription should be for subsequent doses, starting from 10 mg daily, and allowing dose adjustments in increments of 5 mg based on serum IGF-1 levels measured every 4 to 6 weeks in order to maintain the serum IGF-1 level within the age-adjusted normal range based on the dosage recommendations in the TGA-approved Product Information.

The authority application must be made in writing and must include:

- a) two completed authority prescription forms ; and
- b) a completed Acromegaly Pegvisomant initial PBS Authority Application - Supporting Information Form; and
- c) in a patient who has been previously treated with radiotherapy for this condition, the date of completion of radiotherapy, the date and result of IGF-1 levels taken at the most recent two yearly assessment in the 10 years after completion of radiotherapy; and
- d) a recent result of the IGF-1 level and the date of assessment ; and
- e) demonstration of failure to achieve biochemical control after completion of a prior therapy with either octreotide or lanreotide

No increase in the maximum quantity or number of units may be authorised for the loading dose.

pegvisomant 20 mg injection [1 vial] (&) inert substance diluent [1 syringe], 1 pack

11177G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	4	..	..	*557.16	Somavert [PF]

■ **PEGVISOMANT**

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Acromegaly

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must not have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have an age- and sex-adjusted insulin-like growth factor 1 (IGF-1) concentration greater than the upper limit of normal (ULN), **AND**
- The treatment must be after failure to achieve biochemical control with a maximum indicated dose of either 30 mg octreotide LAR or 120 mg lanreotide ATG every 28 days for 24 weeks; unless contraindicated or not tolerated according to the TGA approved Product Information, **AND**
- The treatment must not be given concomitantly with a PBS-subsidised somatostatin analogue.

Somatostatin analogues include octreotide, lanreotide and pasireotide

Failure to achieve biochemical control after completion of a prior therapy with either octreotide or lanreotide is defined as:

- 1) Growth hormone level greater than 1 mcg/L or 3 mIU/L; OR
- 2) IGF-1 level is greater than the age- and sex-adjusted ULN.

If treatment with either octreotide or lanreotide is contraindicated according to the relevant TGA-approved Product Information, the application must provide details of contraindication.

If intolerance to either octreotide or lanreotide treatment developed during the relevant period of use which is of a severity to necessitate withdrawal of the treatment, the application must provide details of the nature and severity of this intolerance.

In a patient treated with radiotherapy, pegvisomant should be withdrawn every 2 years in the 10 years after completion of radiotherapy for assessment of remission. Pegvisomant should be withdrawn at least 8 weeks prior to the assessment of remission.

Biochemical evidence of remission is defined as normalisation of sex- and age- adjusted insulin-like growth factor 1 (IGF-1).

Two completed authority prescriptions should be submitted with the initial application for this drug. One prescription should be for the loading dose of 80 mg for a quantity of 4 vials of 20 mg with no repeats. The second prescription should be for subsequent doses, starting from 10 mg daily, and allowing dose adjustments in increments of 5 mg based on serum IGF-1 levels measured every 4 to 6 weeks in order to maintain the serum IGF-1 level within the age-adjusted normal range based on the dosage recommendations in the TGA-approved Product Information.

The authority application must be made in writing and must include:

- a) two completed authority prescription forms ; and
- b) a completed Acromegaly Pegvisomant initial PBS Authority Application - Supporting Information Form; and
- c) in a patient who has been previously treated with radiotherapy for this condition, the date of completion of radiotherapy, the date and result of IGF-1 levels taken at the most recent two yearly assessment in the 10 years after completion of radiotherapy; and
- d) a recent result of the IGF-1 level and the date of assessment ; and
- e) demonstration of failure to achieve biochemical control after completion of a prior therapy with either octreotide or lanreotide

No increase in the maximum quantity or number of units may be authorised for the loading dose.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs Programs

Reply Paid 9826
HOBART TAS 7001

Authority required

Acromegaly

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must not be given concomitantly with a PBS-subsidised somatostatin analogue, **AND**
- The treatment must cease if IGF-1 is not lower after 3 months of pegvisomant treatment at the maximum tolerated dose. Somatostatin analogues include octreotide, lanreotide and pasireotide

In a patient treated with radiotherapy, pegvisomant should be withdrawn every 2 years in the 10 years after completion of radiotherapy for assessment of remission. Pegvisomant should be withdrawn at least 8 weeks prior to the assessment of remission.

Biochemical evidence of remission is defined as normalisation of sex- and age- adjusted insulin-like growth factor 1 (IGF-1). In a patient who has been previously treated with radiotherapy for this condition, the date of completion of radiotherapy must be provided; and a copy of IGF-1 level taken at the most recent two yearly assessment in the 10 years after completion of radiotherapy must be provided at the time of application.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

pegvisomant 10 mg injection [30 vials] (&) inert substance diluent [30 syringes], 1 pack

11179J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	4178.57	Somavert [PF]

pegvisomant 15 mg injection [30 vials] (&) inert substance diluent [30 syringes], 1 pack

11173C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	4178.57	Somavert [PF]

pegvisomant 20 mg injection [30 vials] (&) inert substance diluent [30 syringes], 1 pack

11181L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	4178.57	Somavert [PF]

HYPOTHALAMIC HORMONES*Somatostatin and analogues***■ LANREOTIDE****Authority required (STREAMLINED)**

7042

Acromegaly

Clinical criteria:

- The condition must be active, **AND**
- Patient must have persistent elevation of mean growth hormone levels of greater than 2.5 micrograms per litre, **AND**
- The treatment must be after failure of other therapy including dopamine agonists; OR
- The treatment must be as interim treatment while awaiting the effects of radiotherapy and where treatment with dopamine agonists has failed; OR
- The treatment must be in a patient who is unfit for or unwilling to undergo surgery and where radiotherapy is contraindicated, **AND**
- The treatment must cease in a patient treated with radiotherapy if there is biochemical evidence of remission (normal IGF1) after lanreotide has been withdrawn for at least 4 weeks (6 weeks after the last dose), **AND**
- The treatment must cease if IGF1 is not lower after 3 months of treatment, **AND**
- The treatment must not be given concomitantly with PBS-subsidised pegvisomant.

In a patient treated with radiotherapy, lanreotide should be withdrawn every 2 years in the 10 years after radiotherapy for assessment of remission.

lanreotide 30 mg modified release injection [1 vial] (&) inert substance diluent [2 mL ampoule], 1 pack

5776B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	11	..	*1218.38	Somatuline LA [IS]

■ LANREOTIDE**Authority required (STREAMLINED)**

7025

Acromegaly

Clinical criteria:

- The condition must be active, **AND**
- Patient must have persistent elevation of mean growth hormone levels of greater than 2.5 micrograms per litre, **AND**
- The treatment must be after failure of other therapy including dopamine agonists; OR

- The treatment must be as interim treatment while awaiting the effects of radiotherapy and where treatment with dopamine agonists has failed; OR
- The treatment must be in a patient who is unfit for or unwilling to undergo surgery and where radiotherapy is contraindicated, **AND**
- The treatment must cease in a patient treated with radiotherapy if there is biochemical evidence of remission (normal IGF1) after lanreotide has been withdrawn for at least 4 weeks (8 weeks after the last dose), **AND**
- The treatment must cease if IGF1 is not lower after 3 months of treatment, **AND**
- The treatment must not be given concomitantly with PBS-subsidised pegvisomant.

In a patient treated with radiotherapy, lanreotide should be withdrawn every 2 years in the 10 years after radiotherapy for assessment of remission.

Authority required (STREAMLINED)

4575

Functional carcinoid tumour

Clinical criteria:

- The condition must be causing intractable symptoms, **AND**
- Patient must have experienced on average over 1 week, 3 or more episodes per day of diarrhoea and/or flushing, which persisted despite the use of anti-histamines, anti-serotonin agents and anti-diarrhoea agents, **AND**
- Patient must be one in whom surgery or antineoplastic therapy has failed or is inappropriate, **AND**
- The treatment must cease if there is failure to produce a clinically significant reduction in the frequency and severity of symptoms after 3 months' therapy at a dose of 120 mg every 28 days.

Dosage and tolerance to the drug should be assessed regularly and the dosage should be titrated slowly downwards to determine the minimum effective dose.

lanreotide 120 mg/0.5 mL injection, 0.5 mL syringe

5779E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*3638.88	Somatuline Autogel [IS]

lanreotide 90 mg/0.5 mL injection, 0.5 mL syringe

5778D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*2907.86	Somatuline Autogel [IS]

lanreotide 60 mg/0.5 mL injection, 0.5 mL syringe

5777C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*2184.96	Somatuline Autogel [IS]

■ LANREOTIDE

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

8260

Non-functional gastroenteropancreatic neuroendocrine tumour (GEP-NET)

Clinical criteria:

- The condition must be unresectable locally advanced disease or metastatic disease, **AND**
- The condition must be World Health Organisation (WHO) grade 1 or 2, **AND**
- The treatment must be as monotherapy.

Population criteria:

- Patient must be aged 18 years or older.
- WHO grade 1 of GEP-NET is defined as a mitotic count (10HPF) of less than 2 and Ki-67 index (%) of less than or equal to 2.

WHO grade 2 of GEP-NET is defined as a mitotic count (10HPF) of 2-20 and Ki-67 index (%) of 3-20.

Lanreotide is not PBS-subsidised for use in combination with everolimus or sunitinib for this condition.

lanreotide 120 mg/0.5 mL injection, 0.5 mL syringe

11513Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*3638.88	Somatuline Autogel [IS]

■ OCTREOTIDE

Authority required (STREAMLINED)

8161

Acromegaly

Clinical criteria:

- The condition must be controlled with octreotide immediate release injections, **AND**
- The treatment must cease in a patient treated with radiotherapy if there is biochemical evidence of remission (normal IGF1) after octreotide has been withdrawn for at least 4 weeks (8 weeks after the last dose), **AND**
- The treatment must cease if IGF1 is not lower after 3 months of treatment, **AND**
- The treatment must not be given concomitantly with PBS-subsidised lanreotide or pegvisomant for this condition.

In a patient treated with radiotherapy, octreotide should be withdrawn every 2 years in the 10 years after radiotherapy for assessment of remission

Authority required (STREAMLINED)

5901

Functional carcinoid tumour

Clinical criteria:

- Patient must have achieved symptom control on octreotide immediate release injections, **AND**
- The treatment must cease if there is failure to produce a clinically significant reduction in the frequency and severity of symptoms after 3 months therapy at a dose of 30 mg every 28 days and having allowed adequate rescue therapy with octreotide immediate release injections.

Dosage and tolerance to the drug should be assessed regularly and the dosage should be titrated slowly downwards to determine the minimum effective dose.

Authority required (STREAMLINED)

5906

Vasoactive intestinal peptide secreting tumour (VIPoma)

Clinical criteria:

- Patient must have achieved symptom control on octreotide immediate release injections, **AND**
- The treatment must cease if there is failure to produce a clinically significant reduction in the frequency and severity of symptoms after 3 months therapy at a dose of 30 mg every 28 days and having allowed adequate rescue therapy with octreotide immediate release injections.

Dosage and tolerance to the drug should be assessed regularly and the dosage should be titrated slowly downwards to determine the minimum effective dose.

octreotide 20 mg modified release injection [1 vial] (&) inert substance diluent [2 mL syringe], 1 pack

10533J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*3479.62	Sandostatin LAR [NV]

octreotide 10 mg modified release injection [1 vial] (&) inert substance diluent [2 mL syringe], 1 pack

10543X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*2613.72	Sandostatin LAR [NV]

octreotide 30 mg modified release injection [1 vial] (&) inert substance diluent [2 mL syringe], 1 pack

10550G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*4354.92	Sandostatin LAR [NV]

■ **OCTREOTIDE**

Authority required (STREAMLINED)

8165

Acromegaly

Clinical criteria:

- The condition must be active, **AND**
- Patient must have persistent elevation of mean growth hormone levels of greater than 2.5 micrograms per litre, **AND**
- The treatment must be after failure of other therapy including dopamine agonists; OR
- The treatment must be as interim treatment while awaiting the effects of radiotherapy and where treatment with dopamine agonists has failed; OR
- The treatment must be in a patient who is unfit for or unwilling to undergo surgery and where radiotherapy is contraindicated, **AND**
- The treatment must cease in a patient treated with radiotherapy if there is biochemical evidence of remission (normal IGF1) after octreotide has been withdrawn for at least 4 weeks, **AND**
- The treatment must cease if IGF1 is not lower after 3 months of treatment at a dose of 100 micrograms 3 time daily, **AND**
- The treatment must not be given concomitantly with PBS-subsidised lanreotide or pegvisomant for this condition.

In a patient treated with radiotherapy, octreotide should be withdrawn every 2 years in the 10 years after radiotherapy for assessment of remission

Authority required (STREAMLINED)

6390

Functional carcinoid tumour

Clinical criteria:

- The condition must be causing intractable symptoms, **AND**
- Patient must have experienced on average over 1 week, 3 or more episodes per day of diarrhoea and/or flushing, which persisted despite the use of anti-histamines, anti-serotonin agents and anti-diarrhoea agents, **AND**
- Patient must be one in whom surgery or antineoplastic therapy has failed or is inappropriate, **AND**
- The treatment must cease if there is failure to produce a clinically significant reduction in the frequency and severity of symptoms after 2 months' therapy.

Dosage and tolerance to the drug should be assessed regularly and the dosage should be titrated slowly downwards to determine the minimum effective dose.

Authority required (STREAMLINED)

6369

Vasoactive intestinal peptide secreting tumour (VIPoma)

Clinical criteria:

- The condition must be causing intractable symptoms, **AND**
- Patient must have experienced on average over 1 week, 3 or more episodes per day of diarrhoea and/or flushing, which persisted despite the use of anti-histamines, anti-serotonin agents and anti-diarrhoea agents, **AND**
- Patient must be one in whom surgery or antineoplastic therapy has failed or is inappropriate, **AND**
- The treatment must cease if there is failure to produce a clinically significant reduction in the frequency and severity of symptoms after 2 months' therapy.

Dosage and tolerance to the drug should be assessed regularly and the dosage should be titrated slowly downwards to determine the minimum effective dose.

octreotide 500 microgram/mL injection, 5 x 1 mL ampoules

9510M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	18	11	..	*6194.52	^a Hospira Pty Limited [PF] ^a Octreotide (SUN) [RA]	^a Octreotide MaxRx [GQ] ^a Sandostatin 0.5 [NV]

octreotide 50 microgram/mL injection, 5 x 1 mL ampoules

9508K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	18	11	..	*619.02	^a Hospira Pty Limited [PF] ^a Octreotide (SUN) [RA]	^a Octreotide MaxRx [GQ] ^a Sandostatin 0.05 [NV]

octreotide 100 microgram/mL injection, 5 x 1 mL ampoules

9509L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	18	11	..	*1236.42	^a Hospira Pty Limited [PF] ^a Octreotide (SUN) [RA]	^a Octreotide MaxRx [GQ] ^a Sandostatin 0.1 [NV]

■ PASIREOTIDE

Caution Careful monitoring of patients is mandatory due to high risk of developing hyperglycaemia

Note Special Pricing Arrangements apply.

Authority required

Acromegaly

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must not have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have a mean growth hormone (GH) level greater than 1 microgram per litre or 3 mIU/L; OR
- Patient must have an age- and sex-adjusted insulin-like growth factor 1 (IGF-1) concentration greater than the upper limit of normal (ULN), **AND**
- The treatment must be after failure to achieve biochemical control with a maximum indicated dose of either 30 mg octreotide LAR or 120 mg lanreotide ATG every 28 days for 24 weeks; unless contraindicated or not tolerated according to the TGA approved Product Information, **AND**
- The treatment must not be given concomitantly with PBS-subsidised pegvisomant.

Population criteria:

- Patient must be aged 18 years or older.

If treatment with either octreotide or lanreotide is contraindicated according to the relevant TGA-approved Product Information, the application must provide details of contraindication.

If intolerance to either octreotide or lanreotide treatment developed during the relevant period of use which is of a severity to necessitate withdrawal of the treatment, the application must provide details of the nature and severity of this intolerance.

Failure to achieve biochemical control after completion of a prior therapy with either octreotide or lanreotide is defined as:

- 1) Growth hormone level greater than 1 mcg/L or 3 mIU/L; OR
- 2) IGF-1 level is greater than the age- and sex-adjusted ULN.

In a patient treated with radiotherapy, pasireotide should be withdrawn every 2 years in the 10 years after completion of radiotherapy for assessment of remission. Pasireotide should be withdrawn at least 8 weeks prior to the assessment of remission.

Biochemical evidence of remission is defined as:

- 1) Growth hormone (GH) levels of less than 1 mcg/L or 3 mIU/L; OR
- 2) normalisation of sex- and age- adjusted insulin-like growth factor 1 (IGF-1)

The authority application must be made in writing and must include:

- a) a completed authority prescription form; and
- b) a completed Acromegaly PBS Authority Application - Supporting Information Form; and
- c) in a patient who has been previously treated with radiotherapy for this condition, the date of completion of radiotherapy must be provided; the date and result of GH or IGF-1 levels taken at the most recent two yearly assessment in the 10 years after completion of radiotherapy must be provided; and
- d) a recent result of GH or IGF-1 levels must be provided.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

ANTIINFECTIVES FOR SYSTEMIC USE

Reply Paid 9826
HOBART TAS 7001

Authority required

Acromegaly

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must not be given concomitantly with PBS-subsidised pegvisomant.

Population criteria:

- Patient must be aged 18 years or older.

In a patient treated with radiotherapy, pasireotide should be withdrawn every 2 years in the 10 years after completion of radiotherapy for assessment of remission. Pasireotide should be withdrawn at least 8 weeks prior to the assessment of remission.

Biochemical evidence of remission is defined as:

- 1) Growth hormone (GH) levels of less than 1 mcg/L or 3 mIU/L; OR
- 2) normalisation of sex- and age- adjusted insulin-like growth factor 1 (IGF-1)

In a patient who has been previously treated with radiotherapy for this condition, the date of completion of radiotherapy and the GH and IGF-1 levels taken at the most recent two yearly assessment in the 10 years after completion of radiotherapy must be provided at the time of approval.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written applications for authorisation under this criterion should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

pasireotide 60 mg modified release injection [1 vial] (& inert substance diluent [2 mL syringe], 1 pack

10882R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*7800.00	Signifor LAR [NV]

pasireotide 40 mg modified release injection [1 vial] (& inert substance diluent [2 mL syringe], 1 pack

10883T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*7800.00	Signifor LAR [NV]

pasireotide 20 mg modified release injection [1 vial] (& inert substance diluent [2 mL syringe], 1 pack

10886Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*7800.00	Signifor LAR [NV]

■ ANTIINFECTIVES FOR SYSTEMIC USE

■ ANTIBACTERIALS FOR SYSTEMIC USE

MACROLIDES, LINCOSAMIDES AND STREPTOGRAMINS

Macrolides

■ AZITHROMYCIN

Authority required (STREAMLINED)

6356

Mycobacterium avium complex infection

Clinical criteria:

- The treatment must be for prophylaxis, **AND**
- Patient must be human immunodeficiency virus (HIV) positive, **AND**
- Patient must have CD4 cell counts of less than 75 per cubic millimetre.

azithromycin 600 mg tablet, 8

5616N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*110.76	Zithromax [PF]

■ ANTIMYCOBACTERIALS

DRUGS FOR TREATMENT OF TUBERCULOSIS

Antibiotics

■ RIFABUTIN

Authority required (STREAMLINED)

6350

Mycobacterium avium complex infection

Clinical criteria:

- Patient must be human immunodeficiency virus (HIV) positive.

Authority required (STREAMLINED)**6356**

Mycobacterium avium complex infection

Clinical criteria:

- The treatment must be for prophylaxis, **AND**
- Patient must be human immunodeficiency virus (HIV) positive, **AND**
- Patient must have CD4 cell counts of less than 75 per cubic millimetre.

rifabutin 150 mg capsule, 30

9541E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	4	5	..	*525.84	Mycobutin [PF]

■ ANTIVIRALS FOR SYSTEMIC USE

DIRECT ACTING ANTIVIRALS

Nucleosides and nucleotides excl. reverse transcriptase inhibitors

■ GANCICLOVIR

Authority required (STREAMLINED)**4972**

Cytomegalovirus disease

Treatment Phase: Prophylaxis

Clinical criteria:

- Patient must be a bone marrow transplant recipient at risk of cytomegalovirus disease.

Authority required (STREAMLINED)**4999**

Cytomegalovirus disease

Treatment Phase: Prophylaxis

Clinical criteria:

- Patient must be a solid organ transplant recipient at risk of cytomegalovirus disease.

ganciclovir 500 mg injection, 5 vials

5749N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	1	..	*401.34	^a Cymevene [PB]	^a GANCICLOVIR SXP [HN]

■ VALACICLOVIR

Authority required (STREAMLINED)**5975**

Cytomegalovirus infection and disease

Treatment Phase: Prophylaxis

Clinical criteria:

- Patient must have undergone a renal transplant, **AND**
- Patient must be at risk of cytomegalovirus disease.

valaciclovir 500 mg tablet, 100

9568N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	5	2	..	*221.00	^a APO-Valaciclovir [TX]	^a Valaciclovir APOTEX [GX]
			^b 2.20	*223.20	^a Valaciclovir RBX [RA]	
					^a Valtrex [RW]	

■ VALGANCICLOVIR

Authority required (STREAMLINED)**4989**

Cytomegalovirus infection and disease

Treatment Phase: Prophylaxis

Clinical criteria:

- Patient must be a solid organ transplant recipient at risk of cytomegalovirus disease.

valganciclovir 450 mg tablet, 60

9569P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	5	..	*1804.54	^a Valcyte [RO]	^a Valganciclovir AN [JO]
					^a Valganciclovir Juno [JU]	^a Valganciclovir Mylan [AF]
					^a Valganciclovir Sandoz [SZ]	

valganciclovir 50 mg/mL powder for oral liquid, 100 mL

9655E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	11	5	..	*4346.10	Valcyte [RO]

Antivirals for treatment of HCV infections

▪ **DACLATASVIR**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Chronic hepatitis C infection

Clinical criteria:

- Patient must meet the criteria set out in the General Statement for Drugs for the Treatment of Hepatitis C, **AND**
- Patient must be taking this drug as part of a regimen set out in the matrix in the General Statement for Drugs for the Treatment of Hepatitis C, based on the hepatitis C virus genotype, patient treatment history and cirrhotic status, **AND**
- The treatment must be limited to a maximum duration of 24 weeks.

daclatasvir 60 mg tablet, 28

10641C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	7666.67	Daklinza [BQ]

daclatasvir 30 mg tablet, 28

10629K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	7666.67	Daklinza [BQ]

▪ **DACLATASVIR**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Chronic hepatitis C infection

Clinical criteria:

- Patient must meet the criteria set out in the General Statement for Drugs for the Treatment of Hepatitis C, **AND**
- Patient must be taking this drug as part of a regimen set out in the matrix in the General Statement for Drugs for the Treatment of Hepatitis C, based on the hepatitis C virus genotype, patient treatment history and cirrhotic status, **AND**
- The treatment must be limited to a maximum duration of 12 weeks.

daclatasvir 60 mg tablet, 28

10660C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	2	..	7666.67	Daklinza [BQ]

daclatasvir 30 mg tablet, 28

10651N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	2	..	7666.67	Daklinza [BQ]

▪ **ELBASVIR + GRAZOPREVR**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Chronic hepatitis C infection

Clinical criteria:

- Patient must meet the criteria set out in the General Statement for Drugs for the Treatment of Hepatitis C, **AND**
- Patient must be taking this drug as part of a regimen set out in the matrix in the General Statement for Drugs for the Treatment of Hepatitis C, based on the hepatitis C virus genotype, patient treatment history and cirrhotic status, **AND**
- The treatment must be limited to a maximum duration of 12 weeks.

elbasvir 50 mg + grazoprevir 100 mg tablet, 28

10978T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	2	..	8400.00	Zepatier [MK]

▪ **ELBASVIR + GRAZOPREVR**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Chronic hepatitis C infection

Clinical criteria:

- Patient must meet the criteria set out in the General Statement for Drugs for the Treatment of Hepatitis C, **AND**
- Patient must be taking this drug as part of a regimen set out in the matrix in the General Statement for Drugs for the Treatment of Hepatitis C, based on the hepatitis C virus genotype, patient treatment history and cirrhotic status, **AND**

- The treatment must be limited to a maximum duration of 16 weeks.

elbasvir 50 mg + grazoprevir 100 mg tablet, 28

10986F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	3	..	8400.00	Zepatier [MK]

▪ GLECAPREVIR + PIBRENTASVIR

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Chronic hepatitis C infection

Clinical criteria:

- Patient must meet the criteria set out in the General Statement for Drugs for the Treatment of Hepatitis C, **AND**
- Patient must be taking this drug as part of a regimen set out in the matrix in the General Statement for Drugs for the Treatment of Hepatitis C, based on the hepatitis C virus genotype, patient treatment history and cirrhotic status, **AND**
- The treatment must be limited to a maximum duration of 8 weeks.

glecaprevir 100 mg + pibrentasvir 40 mg film-coated tablet, 84

11332K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	1	..	18666.67	Maviret [VE]

▪ GLECAPREVIR + PIBRENTASVIR

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Chronic hepatitis C infection

Clinical criteria:

- Patient must meet the criteria set out in the General Statement for Drugs for the Treatment of Hepatitis C, **AND**
- Patient must be taking this drug as part of a regimen set out in the matrix in the General Statement for Drugs for the Treatment of Hepatitis C, based on the hepatitis C virus genotype, patient treatment history and cirrhotic status, **AND**
- The treatment must be limited to a maximum duration of 16 weeks.

glecaprevir 100 mg + pibrentasvir 40 mg film-coated tablet, 84

11333L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	3	..	18666.67	Maviret [VE]

▪ GLECAPREVIR + PIBRENTASVIR

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Chronic hepatitis C infection

Clinical criteria:

- Patient must meet the criteria set out in the General Statement for Drugs for the Treatment of Hepatitis C, **AND**
- Patient must be taking this drug as part of a regimen set out in the matrix in the General Statement for Drugs for the Treatment of Hepatitis C, based on the hepatitis C virus genotype, patient treatment history and cirrhotic status, **AND**
- The treatment must be limited to a maximum duration of 12 weeks.

glecaprevir 100 mg + pibrentasvir 40 mg film-coated tablet, 84

11345D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	2	..	18666.67	Maviret [VE]

▪ LEDIPASVIR + SOFOSBUVIR

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Chronic hepatitis C infection

Clinical criteria:

- Patient must meet the criteria set out in the General Statement for Drugs for the Treatment of Hepatitis C, **AND**
- Patient must be taking this drug as part of a regimen set out in the matrix in the General Statement for Drugs for the Treatment of Hepatitis C, based on the hepatitis C virus genotype, patient treatment history and cirrhotic status, **AND**
- The treatment must be limited to a maximum duration of 12 weeks.

ledipasvir 90 mg + sofosbuvir 400 mg tablet, 28

10661D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	2	..	12500.00	Harvoni [GI]

▪ LEDIPASVIR + SOFOSBUVIR

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Chronic hepatitis C infection

Clinical criteria:

- Patient must meet the criteria set out in the General Statement for Drugs for the Treatment of Hepatitis C, **AND**
- Patient must be taking this drug as part of a regimen set out in the matrix in the General Statement for Drugs for the Treatment of Hepatitis C, based on the hepatitis C virus genotype, patient treatment history and cirrhotic status, **AND**
- The treatment must be limited to a maximum duration of 8 weeks.

ledipasvir 90 mg + sofosbuvir 400 mg tablet, 28

10667K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	1	..	12500.00	Harvoni [GI]

▪ LEDIPASVIR + SOFOSBUVIR

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Chronic hepatitis C infection

Clinical criteria:

- Patient must meet the criteria set out in the General Statement for Drugs for the Treatment of Hepatitis C, **AND**
- Patient must be taking this drug as part of a regimen set out in the matrix in the General Statement for Drugs for the Treatment of Hepatitis C, based on the hepatitis C virus genotype, patient treatment history and cirrhotic status, **AND**
- The treatment must be limited to a maximum duration of 24 weeks.

ledipasvir 90 mg + sofosbuvir 400 mg tablet, 28

10669M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	12500.00	Harvoni [GI]

▪ RIBAVIRIN

Caution Ribavirin is a category X drug and must not be given to pregnant women. Pregnancy in female patients or in the partners of male patients must be avoided during treatment and during the 6 months period after cessation of treatment.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Chronic hepatitis C infection

Clinical criteria:

- Patient must meet the criteria set out in the General Statement for Drugs for the Treatment of Hepatitis C, **AND**
- Patient must be taking this drug as part of a regimen set out in the matrix in the General Statement for Drugs for the Treatment of Hepatitis C, based on the hepatitis C virus genotype, patient treatment history and cirrhotic status, **AND**
- The treatment must be limited to a maximum duration of 24 weeks.

Population criteria:

- Patient must not be pregnant or breastfeeding. Female partners of male patients must not be pregnant. Patients and their partners must each be using an effective form of contraception if of child-bearing age.

ribavirin 600 mg tablet, 28

10638X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	420.00	Ibavir [IX]

ribavirin 400 mg tablet, 28

10646H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	280.00	Ibavir [IX]

▪ RIBAVIRIN

Caution Ribavirin is a category X drug and must not be given to pregnant women. Pregnancy in female patients or in the partners of male patients must be avoided during treatment and during the 6 months period after cessation of treatment.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Chronic hepatitis C infection

Clinical criteria:

- Patient must meet the criteria set out in the General Statement for Drugs for the Treatment of Hepatitis C, **AND**
- Patient must be taking this drug as part of a regimen set out in the matrix in the General Statement for Drugs for the Treatment of Hepatitis C, based on the hepatitis C virus genotype, patient treatment history and cirrhotic status, **AND**
- The treatment must be limited to a maximum duration of 12 weeks.

Population criteria:

- Patient must not be pregnant or breastfeeding. Female partners of male patients must not be pregnant. Patients and their partners must each be using an effective form of contraception if of child-bearing age.

ribavirin 600 mg tablet, 28

10663F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	2	..	420.00	Ibavyr [IX]

ribavirin 400 mg tablet, 28

10678B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	2	..	280.00	Ibavyr [IX]

▪ SOFOSBUVIR

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Chronic hepatitis C infection

Clinical criteria:

- Patient must meet the criteria set out in the General Statement for Drugs for the Treatment of Hepatitis C, **AND**
- Patient must be taking this drug as part of a regimen set out in the matrix in the General Statement for Drugs for the Treatment of Hepatitis C, based on the hepatitis C virus genotype, patient treatment history and cirrhotic status, **AND**
- The treatment must be limited to a maximum duration of 12 weeks.

sofosbuvir 400 mg tablet, 28

10625F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	2	..	12500.00	Sovaldi [GI]

▪ SOFOSBUVIR

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Chronic hepatitis C infection

Clinical criteria:

- Patient must meet the criteria set out in the General Statement for Drugs for the Treatment of Hepatitis C, **AND**
- Patient must be taking this drug as part of a regimen set out in the matrix in the General Statement for Drugs for the Treatment of Hepatitis C, based on the hepatitis C virus genotype, patient treatment history and cirrhotic status, **AND**
- The treatment must be limited to a maximum duration of 24 weeks.

sofosbuvir 400 mg tablet, 28

10648K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	12500.00	Sovaldi [GI]

▪ SOFOSBUVIR + VELPATASVIR

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Chronic hepatitis C infection

Clinical criteria:

- Patient must meet the criteria set out in the General Statement for Drugs for the Treatment of Hepatitis C, **AND**
- Patient must be taking this drug as part of a regimen set out in the matrix in the General Statement for Drugs for the Treatment of Hepatitis C, based on the hepatitis C virus genotype, patient treatment history and cirrhotic status, **AND**
- The treatment must be limited to a maximum duration of 12 weeks.

sofosbuvir 400 mg + velpatasvir 100 mg tablet, 28

11145N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	2	..	12500.00	Epclusa [GI]

▪ SOFOSBUVIR + VELPATASVIR + VOXILAPREVIR

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Chronic hepatitis C infection

Clinical criteria:

- Patient must meet the criteria set out in the General Statement for Drugs for the Treatment of Hepatitis C, **AND**
- Patient must be taking this drug as part of a regimen set out in the matrix in the General Statement for Drugs for the Treatment of Hepatitis C, based on the hepatitis C virus genotype, patient treatment history and cirrhotic status, **AND**
- The treatment must be limited to a maximum duration of 12 weeks.

sofosbuvir 400 mg + velpatasvir 100 mg + voxilaprevir 100 mg tablet, 28

11665Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	2	..	12500.00	Vosevi [GI]

■ ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS

■ ANTINEOPLASTIC AGENTS

ANTIMETABOLITES

Pyrimidine analogues

■ AZACITIDINE

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Myelodysplastic syndrome

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be classified as Intermediate-2 according to the International Prognostic Scoring System (IPSS); OR
 - The condition must be classified as high risk according to the International Prognostic Scoring System (IPSS).
- Classification of the condition as Intermediate-2 requires a score of 1.5 to 2.0 on the IPSS, achieved with the possible combinations:
- 11% to 30% marrow blasts with good karyotypic status (normal, -Y alone, del(5q) alone, del(20q) alone), and 0 to 1 cytopenias; OR
 - 11% to 20% marrow blasts with intermediate karyotypic status (other abnormalities), and 0 to 1 cytopenias; OR
 - 11% to 20% marrow blasts with good karyotypic status (normal, -Y alone, del(5q) alone, del(20q) alone), and 2 to 3 cytopenias; OR
 - 5% to 10% marrow blasts with poor karyotypic status (3 or more abnormalities or chromosome 7 anomalies), regardless of cytopenias; OR
 - 5% to 10% marrow blasts with intermediate karyotypic status (other abnormalities), and 2 to 3 cytopenias; OR
 - Less than 5% marrow blasts with poor karyotypic status (3 or more abnormalities or chromosome 7 anomalies), and 2 to 3 cytopenias.

Classification of the condition as high risk requires a score of 2.5 or more on the IPSS, achieved with the possible combinations:

- 21% to 30% marrow blasts with good karyotypic status (normal, -Y alone, del(5q) alone, del(20q) alone), and 2 to 3 cytopenias; OR
- 21% to 30% marrow blasts with intermediate (other abnormalities) or poor karyotypic status (3 or more abnormalities or chromosome 7 anomalies), regardless of cytopenias; OR
- 11% to 20% marrow blasts with poor karyotypic status (3 or more abnormalities or chromosome 7 anomalies), regardless of cytopenias; OR
- 11% to 20% marrow blasts with intermediate karyotypic status (other abnormalities), and 2 to 3 cytopenias.

The first authority application must be made in writing and must include:

- a completed authority prescription form; and
- a completed Azacitidine PBS Authority Application - Supporting Information Form; and
- a copy of the bone marrow biopsy report demonstrating that the patient has myelodysplastic syndrome; and
- a copy of the full blood examination report; and
- a copy of the pathology report detailing the cytogenetics demonstrating intermediate-2 or high risk disease according to the International Prognostic Scoring System (IPSS); and
- a signed patient acknowledgment form.

No more than 3 cycles will be authorised.

Authority required

Chronic Myelomonocytic Leukaemia

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must have 10% to 29% marrow blasts without Myeloproliferative Disorder.

The first authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Azacitidine PBS Authority Application - Supporting Information Form; and
- (c) a copy of the bone marrow biopsy report demonstrating that the patient has chronic myelomonocytic leukaemia ; and
- (d) a copy of the full blood examination report; and
- (e) a signed patient acknowledgement.

No more than 3 cycles will be authorised.

Authority required

Acute Myeloid Leukaemia

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must have 20% to 30% marrow blasts and multi-lineage dysplasia, according to World Health Organisation (WHO) Classification.

The first authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Azacitidine PBS Authority Application - Supporting Information Form; and
- (c) a copy of the bone marrow biopsy report demonstrating that the patient has acute myeloid leukaemia; and
- (d) a copy of the full blood examination report; and
- (e) a signed patient acknowledgement.

No more than 3 cycles will be authorised.

azacitidine 100 mg injection, 1 vial

9597D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	14	2	..	*3046.96	^a Azacitidine Accord [OC] ^a Azacitidine Juno [JO] ^a Celazadine [JU]	^a AZACITIDINE DR.REDDY'S [RI] ^a Azadine [RZ] ^a Vidaza [CJ]

▪ **AZACITIDINE**

Note Authority applications for continuing treatment may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Myelodysplastic syndrome

Treatment Phase: Continuing treatment

Clinical criteria:

- The condition must be classified as Intermediate-2 according to the International Prognostic Scoring System (IPSS); OR
- The condition must be classified as high risk according to the International Prognostic Scoring System (IPSS), **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have progressive disease.

Applications for continuing therapy may be made by telephone.

Up to 6 cycles will be authorised.

Authority required

Chronic Myelomonocytic Leukaemia

Treatment Phase: Continuing treatment

Clinical criteria:

- The condition must have 10% to 29% marrow blasts without Myeloproliferative Disorder, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have progressive disease.

Applications for continuing therapy may be made by telephone.

Up to 6 cycles will be authorised.

Authority required

Acute Myeloid Leukaemia

Treatment Phase: Continuing treatment

Clinical criteria:

- The condition must have 20% to 30% marrow blasts and multi-lineage dysplasia, according to World Health Organisation (WHO) Classification, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have progressive disease.

Applications for continuing therapy may be made by telephone.

Up to 6 cycles will be authorised.

azacitidine 100 mg injection, 1 vial

9598E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	14	5	..	*3046.96	^a Azacitidine Accord [OC] ^a Azacitidine Juno [JO]	^a AZACITIDINE DR.REDDY'S [RI] ^a Azadine [RZ]

CYTOTOXIC ANTIBIOTICS AND RELATED SUBSTANCES

Anthracyclines and related substances

■ DOXORUBICIN HYDROCHLORIDE (AS PEGYLATED LIPOSOMAL)

Authority required (STREAMLINED)

6234

Kaposi sarcoma

Clinical criteria:

- The condition must be AIDS-related, **AND**
- Patient must have a CD4 cell count of less than 200 per cubic millimetre, **AND**
- The condition must include extensive mucocutaneous involvement.

Authority required (STREAMLINED)

6274

Kaposi sarcoma

Clinical criteria:

- The condition must be AIDS-related, **AND**
- Patient must have a CD4 cell count of less than 200 per cubic millimetre, **AND**
- The condition must include extensive visceral involvement.

doxorubicin hydrochloride (as pegylated liposomal) 20 mg/10 mL injection, 10 mL vial

5705G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	4	5	..	*893.76	^a Caelyx [JC]	^a Liposomal Doxorubicin SUN [RA]

OTHER ANTINEOPLASTIC AGENTS

Monoclonal antibodies

■ RITUXIMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for adults with severe active rheumatoid arthritis. Where the term biological medicine appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alfa antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab), the T-cell co-stimulation modulator (abatacept) and the Janus kinase (JAK) inhibitors (baricitinib, tofacitinib). A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time. In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised biological medicine therapy may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once, and
- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised biological medicines for the treatment of rheumatoid arthritis.

Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Stevens Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

For a patient who has failed PBS-subsidised treatment with 2 or 3 TNF-alfa antagonists prior to 1 August 2010 please contact the Department of Human Services on 1800 700 270. A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction.

A patient who has failed fewer than 5 biological medicines and who has a break in treatment of less than 24 months may commence a further course of treatment with a biological medicine under Initial 2 treatment restriction. A patient who has failed fewer than 5 biological medicines and who has had a break in therapy of longer than 24 months may commence a further course of treatment with a biological medicine under the Initial 3 treatment restriction. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine treatment is stopped to the date of the new application for treatment with a biological medicine.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 April 2019.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised biological medicine treatment and wishes to commence such therapy, excluding rituximab (Initial 1 - new patient); or
- (ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months) [further details are under 'Swapping therapy' below]; or
- (iii) a patient wishes to re-commence treatment with a specific biological medicine following a break of less than 24 months in PBS-subsidised therapy with that agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months).
- (iv) a patient wishes to re-commence treatment with a biological medicine following a break in PBS-subsidised therapy of

more than 24 months (Initial 3 - re-commencement of treatment after a break in biological medicine of more than 24 months) Initial applications for a new patient (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6-month intensive DMARD trial, but prior to ceasing DMARD therapy.

Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, baricitinib, etanercept, golimumab, tocilizumab and tofacitinib, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

Rituximab patients should be assessed following a minimum of 12 weeks after the first infusion, and the assessment should be submitted to the Department of Human Services within 4 weeks to ensure continuity of treatment for those who meet the continuing restriction for PBS subsidised treatment with this drug for this condition.

Where a response assessment is not provided the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment. For second and subsequent courses of PBS-subsidised biological medicine (excluding rituximab) treatment it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to the Department of Human Services within sufficient timeframe to allow for authority assessment.

Abatacept patients:

A patient is eligible to receive one I.V. loading dose when commencing treatment with the subcutaneous formulation. Two prescriptions are required, the first prescription for the I.V. loading dose for sufficient vials for one dose based on the patient's weight with no repeats. The second prescription for the subcutaneous formulation, with a maximum quantity of 4 and up to 3 repeats, must be submitted with the initial application.

Rituximab patients:

A further application may be submitted to the Department of Human Services within sufficient timeframe to allow authority assessment. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

Infliximab and etanercept patients:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction. Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

(2) Swapping therapy

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non- biological medicine therapy requirements except if the patient has had a break in therapy of more than 24 months who would need to requalify with respect to the indices of disease severity. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each biological medicine trialled.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent, unless they have a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug.

Abatacept:

A patient swapping from I.V. abatacept to subcutaneous abatacept will not be eligible for an I.V. loading dose when commencing treatment with the subcutaneous formulation.

Rituximab:

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF-alfa antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological medicine during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological medicine therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate biological medicine may do so without having to have any treatment-free period.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and the Department of Human Services will assess response according to these revised baseline

measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints.

Applications under the Initial 1 treatment restriction for a new patient must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy.

Applications under the Initial 3 treatment restriction for re-commencement of treatment after a break in biological medicine of more than 24 months must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required (STREAMLINED)

9446

Severe active rheumatoid arthritis

Treatment Phase: Subsequent continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be given concomitantly with methotrexate at a dose of at least 7.5 mg weekly.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The measurement of response to the prior course of therapy must be documented in the patient's medical notes.

If a patient has either failed or ceased to respond to a PBS-subsidised biological medicine for this condition 5 times, they will not be eligible to receive further PBS-subsidised treatment with a biological medicine for this condition.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

rituximab 500 mg/50 mL injection, 50 mL vial

11805H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	1	..	*2398.64	Riximyo [SZ]

■ RITUXIMAB

Note Authority applications for increased quantities/ repeats (where relevant) may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note Risk of end-organ damage or mortality includes a minimum of one of the following: Glomerulonephritis with risk of progression

- Risk to sight including scleritis/episcleritis, sudden visual loss, uveitis, retinal changes (vasculitis/thrombosis/exudates/haemorrhage)
- Bronchial/subglottic obstruction
- Pulmonary haemorrhage
- Parenchymal lung disease
- Sensory neural hearing loss
- Recurrent sinonasal disease requiring recurrent surgical interventions
- Meningitis, organic confusion, seizures, stroke, cord lesion, cranial nerve palsy, sensory peripheral neuropathy, motor mononeuritis multiplex

Note Patients could be considered contraindicated, refractory, or unable to tolerate cyclophosphamide for one of the following reasons: Cyclophosphamide is contraindicated as per the TGA approved Product Information;

- Cyclophosphamide is not recommended due to the need to preserve gonad function;
- Patient experiences severe toxicity to cyclophosphamide that warrants cessation of treatment;
- Patient has life- or organ-threatening deterioration at any time during treatment with cyclophosphamide, where the deterioration is thought to be due to severe uncontrolled active vasculitis;
- Commencing a further treatment cycle with cyclophosphamide would exceed the maximum cumulative dose of cyclophosphamide of 25g; or
- Patient's condition with this indication is persistent despite at least 3 months therapy with cyclophosphamide.

Authority required (STREAMLINED)

9336

Severe active granulomatosis with polyangiitis (Wegeners granulomatosis)

Treatment Phase: Re-induction of remission

Clinical criteria:

- The treatment must be for the re-induction of remission, **AND**
- Patient must have previously received and responded to this drug for this condition, **AND**
- The treatment must in combination with glucocorticoids, **AND**
- Patient must be at risk of end-organ damage or mortality, **AND**
- Patient must be contraindicated, refractory or unable to tolerate cyclophosphamide.

Diagnosis should be made according to the Chapel Hill Consensus Conference Nomenclature of the Vasculitides with anti-neutrophil cytoplasmic antibody (ANCA) positive serology.

This drug is not PBS-subsidised for maintenance of remission

Authority required (STREAMLINED)

9539

Severe active microscopic polyangiitis

Treatment Phase: Re-induction of remission

Clinical criteria:

- The treatment must be for the re-induction of remission, **AND**
- Patient must have previously received and responded to this drug for this condition, **AND**
- The treatment must in combination with glucocorticoids, **AND**
- Patient must be at risk of end-organ damage or mortality, **AND**
- Patient must be contraindicated, refractory or unable to tolerate cyclophosphamide.

Diagnosis should be made according to the Chapel Hill Consensus Conference Nomenclature of the Vasculitides with anti-neutrophil cytoplasmic antibody (ANCA) positive serology.

This drug is not PBS-subsidised for maintenance therapy.

rituximab 500 mg/50 mL injection, 50 mL vial

11800C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	4	..	..	*4797.28	Riximyo [SZ]

rituximab 100 mg/10 mL injection, 2 x 10 mL vials

11813R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	3	..	..	*1439.19	Riximyo [SZ]

▪ RITUXIMAB

Note Risk of end-organ damage or mortality includes a minimum of one of the following: Glomerulonephritis with risk of progression

- Risk to sight including scleritis/episcleritis, sudden visual loss, uveitis, retinal changes (vasculitis/thrombosis/exudates/haemorrhage)
- Bronchial/subglottic obstruction
- Pulmonary haemorrhage
- Parenchymal lung disease
- Sensory neural hearing loss
- Recurrent sinonasal disease requiring recurrent surgical interventions
- Meningitis, organic confusion, seizures, stroke, cord lesion, cranial nerve palsy, sensory peripheral neuropathy, motor mononeuritis multiplex

Note Patients could be considered contraindicated, refractory, or unable to tolerate cyclophosphamide for one of the following reasons: Cyclophosphamide is contraindicated as per the TGA approved Product Information;

- Cyclophosphamide is not recommended due to the need to preserve gonad function;
- Patient experiences severe toxicity to cyclophosphamide that warrants cessation of treatment;
- Patient has life- or organ-threatening deterioration at any time during treatment with cyclophosphamide, where the deterioration is thought to be due to severe uncontrolled active vasculitis;
- Commencing a further treatment cycle with cyclophosphamide would exceed the maximum cumulative dose of cyclophosphamide of 25g; or
- Patient's condition with this indication is persistent despite at least 3 months therapy with cyclophosphamide.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au
Applications for authority to prescribe should be forwarded to:
Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note At the time of authority application, medical practitioners must request the appropriate number of vials to provide sufficient drug for four weeks of treatment.

Authority required

Severe active granulomatosis with polyangiitis (Wegeners granulomatosis)

Treatment Phase: Induction of remission

Clinical criteria:

- The treatment must be for the induction of remission, **AND**
 - Patient must not have previously received this drug for this condition, **AND**
 - The treatment must in combination with glucocorticoids, **AND**
 - Patient must be at risk of end-organ damage or mortality, **AND**
 - Patient must be contraindicated, refractory or unable to tolerate cyclophosphamide.
- Diagnosis should be made according to the Chapel Hill Consensus Conference Nomenclature of the Vasculitides with anti-neutrophil cytoplasmic antibody (ANCA) positive serology.

This drug is not PBS-subsidised for maintenance of remission

The authority application must be made in writing

Note Biosimilar prescribing policy

Prescribing of the biosimilar brand, Riximyo, is encouraged for treatment naive patients.

Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Authority required

Severe active microscopic polyangiitis

Treatment Phase: Induction of remission

Clinical criteria:

- The treatment must be for the induction of remission, **AND**
 - Patient must not have previously received this drug for this condition, **AND**
 - The treatment must in combination with glucocorticoids, **AND**
 - Patient must be at risk of end-organ damage or mortality, **AND**
 - Patient must be contraindicated, refractory or unable to tolerate cyclophosphamide.
- Diagnosis should be made according to the Chapel Hill Consensus Conference Nomenclature of the Vasculitides with anti-neutrophil cytoplasmic antibody (ANCA) positive serology.

This drug is not PBS-subsidised for maintenance therapy.

The authority application must be made in writing

Note Biosimilar prescribing policy

Prescribing of the biosimilar brand, Riximyo, is encouraged for treatment naive patients.

Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Authority required

Severe active granulomatosis with polyangiitis (Wegeners granulomatosis)

Treatment Phase: Re-induction of remission

Clinical criteria:

- The treatment must be for the re-induction of remission, **AND**
- Patient must have previously received and responded to this drug for this condition, **AND**
- The treatment must in combination with glucocorticoids, **AND**
- Patient must be at risk of end-organ damage or mortality, **AND**
- Patient must be contraindicated, refractory or unable to tolerate cyclophosphamide.

Diagnosis should be made according to the Chapel Hill Consensus Conference Nomenclature of the Vasculitides with anti-neutrophil cytoplasmic antibody (ANCA) positive serology.

This drug is not PBS-subsidised for maintenance of remission

The authority application must be made in writing

Authority required

Severe active microscopic polyangiitis

Treatment Phase: Re-induction of remission

Clinical criteria:

- The treatment must be for the re-induction of remission, **AND**

- Patient must have previously received and responded to this drug for this condition, **AND**
- The treatment must in combination with glucocorticoids, **AND**
- Patient must be at risk of end-organ damage or mortality, **AND**
- Patient must be contraindicated, refractory or unable to tolerate cyclophosphamide.

Diagnosis should be made according to the Chapel Hill Consensus Conference Nomenclature of the Vasculitides with anti-neutrophil cytoplasmic antibody (ANCA) positive serology.

This drug is not PBS-subsidised for maintenance therapy.

The authority application must be made in writing

rituximab 500 mg/50 mL injection, 50 mL vial

10593M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	1199.32	^a Mabthera [RO]	^a Riximyo [SZ]

rituximab 100 mg/10 mL injection, 2 x 10 mL vials

10591K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	479.73	^a Mabthera [RO]	^a Riximyo [SZ]

■ RITUXIMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for adults with severe active rheumatoid arthritis. Where the term biological medicine appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alfa antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab), the T-cell co-stimulation modulator (abatacept) and the Janus kinase (JAK) inhibitors (baricitinib, tofacitinib). A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised biological medicine therapy may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once, and
- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised biological medicines for the treatment of rheumatoid arthritis.

Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Stevens Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

For a patient who has failed PBS-subsidised treatment with 2 or 3 TNF-alfa antagonists prior to 1 August 2010 please contact the Department of Human Services on 1800 700 270. A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction.

A patient who has failed fewer than 5 biological medicines and who has a break in treatment of less than 24 months may commence a further course of treatment with a biological medicine under Initial 2 treatment restriction. A patient who has failed fewer than 5 biological medicines and who has had a break in therapy of longer than 24 months may commence a further course of treatment with a biological medicine under the Initial 3 treatment restriction. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine treatment is stopped to the date of the new application for treatment with a biological medicine.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 April 2019.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised biological medicine treatment and wishes to commence such therapy, excluding rituximab (Initial 1 - new patient); or

(ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to re-commence treatment with a specific biological medicine following a break of less than 24 months in PBS-subsidised therapy with that agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months).

(iv) a patient wishes to re-commence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 24 months (Initial 3 - re-commencement of treatment after a break in biological medicine of more than 24 months) Initial applications for a new patient (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6-month intensive DMARD trial, but prior to ceasing DMARD therapy.

Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, baricitinib, etanercept, golimumab, tocilizumab and tofacitinib, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

Rituximab patients should be assessed following a minimum of 12 weeks after the first infusion, and the assessment should be submitted to the Department of Human Services within 4 weeks to ensure continuity of treatment for those who meet the continuing restriction for PBS subsidised treatment with this drug for this condition.

Where a response assessment is not provided the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment. For second and subsequent courses of PBS-subsidised biological medicine (excluding

rituximab) treatment it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to the Department of Human Services within sufficient timeframe to allow for authority assessment.

Abatacept patients:

A patient is eligible to receive one I.V. loading dose when commencing treatment with the subcutaneous formulation. Two prescriptions are required, the first prescription for the I.V. loading dose for sufficient vials for one dose based on the patient's weight with no repeats. The second prescription for the subcutaneous formulation, with a maximum quantity of 4 and up to 3 repeats, must be submitted with the initial application.

Rituximab patients:

A further application may be submitted to the Department of Human Services within sufficient timeframe to allow authority assessment. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

Infliximab and etanercept patients:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions. For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction. Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

(2) Swapping therapy

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non- biological medicine therapy requirements except if the patient has had a break in therapy of more than 24 months who would need to requalify with respect to the indices of disease severity. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each biological medicine trialled.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent, unless they have a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug.

Abatacept:

A patient swapping from I.V. abatacept to subcutaneous abatacept will not be eligible for an I.V. loading dose when commencing treatment with the subcutaneous formulation.

Rituximab:

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF-alfa antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological medicine during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological medicine therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate biological medicine may do so without having to have any treatment-free period.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and the Department of Human Services will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints.

Applications under the Initial 1 treatment restriction for a new patient must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy.

Applications under the Initial 3 treatment restriction for re-commencement of treatment after a break in biological medicine of more than 24 months must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au
Applications for authority to prescribe should be forwarded to:
Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have failed to respond to at least 1 PBS-subsidised tumour necrosis factor (TNF) alfa antagonist for this condition, **AND**
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with disease modifying anti-rheumatic drugs (DMARDs) which must include at least 3 months continuous treatment with each of at least 2 DMARDs, one of which must be methotrexate at a dose of at least 20 mg weekly and one of which must be: (i) hydroxychloroquine at a dose of at least 200 mg daily; or (ii) leflunomide at a dose of at least 10 mg daily; or (iii) sulfasalazine at a dose of at least 2 g daily; OR
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs which, if methotrexate is contraindicated according to the Therapeutic Goods Administration (TGA)-approved Product Information or cannot be tolerated at a 20 mg weekly dose, must include at least 3 months continuous treatment with each of at least 2 of the following DMARDs: (i) hydroxychloroquine at a dose of at least 200 mg daily; and/or (ii) leflunomide at a dose of at least 10 mg daily; and/or (iii) sulfasalazine at a dose of at least 2 g daily; OR
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs which, if 3 or more of methotrexate, hydroxychloroquine, leflunomide and sulfasalazine are contraindicated according to the relevant TGA-approved Product Information or cannot be tolerated at the doses specified above, must include at least 3 months continuous treatment with each of at least 2 DMARDs, with one or more of the following DMARDs being used in place of the DMARDs which are contraindicated or not tolerated: (i) azathioprine at a dose of at least 1 mg/kg per day; and/or (ii) cyclosporin at a dose of at least 2 mg/kg/day; and/or (iii) sodium aurothiomalate at a dose of 50 mg weekly, **AND**
- Patient must not receive more than 2 infusions of this drug under this restriction, **AND**
- The treatment must be given concomitantly with methotrexate at a dose of at least 7.5 mg weekly.

Population criteria:

- Patient must be aged 18 years or older.

If methotrexate is contraindicated according to the TGA-approved product information or cannot be tolerated at a 20 mg weekly dose, the application must include details of the contraindication or intolerance including severity to methotrexate. The maximum tolerated dose of methotrexate must be documented in the application, if applicable.

The application must include details of the DMARDs trialed, their doses and duration of treatment, and all relevant contraindications and/or intolerances including severity.

The requirement to trial at least 2 DMARDs for periods of at least 3 months each can be met using single agents sequentially or by using one or more combinations of DMARDs.

If the requirement to trial 6 months of intensive DMARD therapy with at least 2 DMARDs cannot be met because of contraindications and/or intolerances of a severity necessitating permanent treatment withdrawal to all of the DMARDs specified above, details of the contraindication or intolerance including severity and dose for each DMARD must be provided in the authority application.

The following criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; **AND** either

(a) a total active joint count of at least 20 active (swollen and tender) joints; or

(b) at least 4 active joints from the following list of major joints:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The joint count and ESR and/or CRP must be determined at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. All measures must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and

(2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient whose most recent course of PBS-subsidised therapy was with this drug and whose response to this treatment is demonstrated at 12 weeks, may apply for a further course of this drug under the First continuing treatment restriction.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

If a patient fails to demonstrate a response to this drug and who qualifies to trial an alternate biological medicine according to the interchangeability arrangements for biological medicines for the treatment of severe rheumatoid arthritis, may do so without having to have a 22 week treatment-free period.

Note Biosimilar prescribing policy

Prescribing of the biosimilar brand, Riximyo, is encouraged for treatment naive patients.

Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note The Department of Human Services website (www.humanservices.gov.au) has details of the toxicities, including severity, which will be accepted for the following purposes:

- (a) exempting a patient from the requirement to undertake a minimum 3 month trial of methotrexate at a 20 mg weekly dose;
- (b) substituting azathioprine, cyclosporin or sodium aurothiomalate for another DMARD as part of the 6 month intensive DMARD trial;
- (c) exempting a patient from the requirement for a 6 month trial of intensive DMARD therapy.

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 24 months)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to respond to at least 1 PBS-subsidised tumour necrosis factor (TNF) alfa antagonist for this condition, **AND**
- Patient must not have failed to respond to previous PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised biological medicine treatment for this condition 5 times, **AND**
- Patient must not receive more than 2 infusions of this drug under this restriction, **AND**
- The treatment must be given concomitantly with methotrexate at a dose of at least 7.5 mg weekly.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either of the Initial 1, Initial 2, Initial 3, first or subsequent continuing treatment restrictions, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the

date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

A patient may qualify to receive a further course of treatment (every 24 weeks) with this drug provided they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with this drug. The demonstration of response must be submitted within 4 weeks of assessment.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient whose most recent course of PBS-subsidised therapy was with this drug and whose response to this treatment is demonstrated at 12 weeks, may apply for a further course of this drug under the First continuing treatment restriction.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

If a patient fails to demonstrate a response to this drug and who qualifies to trial an alternate biological medicine according to the interchangeability arrangements for biological medicines for the treatment of severe rheumatoid arthritis, may do so without having to have a 22 week treatment-free period.

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 3 (recommencement of treatment after a break in biological medicine of more than 24 months)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 24 months or more from the most recent PBS-subsidised biological medicine for this condition, **AND**
- Patient must have failed to respond to at least 1 PBS-subsidised tumour necrosis factor (TNF) alfa antagonist for this condition, **AND**
- Patient must not have failed to respond to previous PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised biological medicine treatment for this condition 5 times, **AND**
- The condition must have an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; OR
- The condition must have a C-reactive protein (CRP) level greater than 15 mg per L, **AND**
- The condition must have either (a) a total active joint count of at least 20 active (swollen and tender) joints; or (b) at least 4 active major joints, **AND**
- Patient must not receive more than 2 infusions of this drug under this restriction, **AND**
- The treatment must be given concomitantly with methotrexate at a dose of at least 7.5 mg weekly.

Population criteria:

- Patient must be aged 18 years or older.

Major joints are defined as (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

All measures of joint count and ESR and/or CRP must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient whose most recent course of PBS-subsidised therapy was with this drug and whose response to this treatment is demonstrated at 12 weeks, may apply for a further course of this drug under the First continuing treatment restriction.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

If a patient fails to demonstrate a response to this drug and who qualifies to trial an alternate biological medicine according to the interchangeability arrangements for biological medicines for the treatment of severe rheumatoid arthritis, may do so without having to have a 22 week treatment-free period.

Note Biosimilar prescribing policy

Prescribing of the biosimilar brand, Riximyo, is encouraged for treatment naive patients.

Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Authority required

Severe active rheumatoid arthritis

Treatment Phase: First continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 2 infusions of this drug under this restriction, **AND**
- The treatment must be given concomitantly with methotrexate at a dose of at least 7.5 mg weekly.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

A patient may qualify to receive a further course of treatment (every 24 weeks) with this drug provided they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with this drug. The demonstration of response must be submitted within 4 weeks of assessment.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient has either failed or ceased to respond to a PBS-subsidised biological medicine for this condition 5 times, they will not be eligible to receive further PBS-subsidised treatment with a biological medicine for this condition.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Subsequent continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be given concomitantly with methotrexate at a dose of at least 7.5 mg weekly.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

A patient may qualify to receive a further course of treatment (every 24 weeks) with this drug provided they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with this drug. The demonstration of response must be submitted within 4 weeks of assessment.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient has either failed or ceased to respond to a PBS-subsidised biological medicine for this condition 5 times, they will not be eligible to receive further PBS-subsidised treatment with a biological medicine for this condition.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

rituximab 500 mg/50 mL injection, 50 mL vial

9544H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	1199.32	^a Mabthera [RO]	^a Riximyo [SZ]

HSD (Public)

Protein kinase inhibitors

▪ MIDOSTAURIN

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Applications for authority to prescribe may be made by phone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday)

Authority required

Acute Myeloid Leukaemia

Treatment Phase: Maintenance therapy - Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the initial maintenance or the initial maintenance grandfathering treatment restriction, **AND**
- Patient must not have developed disease progression while receiving PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not be undergoing or have undergone a stem cell transplant.

A maximum of 9 cycles will be authorised under this restriction in a lifetime.

Progressive disease monitoring via a complete blood count must be taken at the end of each cycle.

If abnormal blood counts suggest the potential for relapsed AML, a bone marrow biopsy must be performed to confirm the absence of progressive disease for the patient to be eligible for further cycles.

Progressive disease is defined as the presence of any of the following:

- Leukaemic cells in the CSF;
- Re-appearance of circulating blast cells in the peripheral blood, not attributable to overshoot following recovery from myeloablative therapy;
- Greater than 5 % blasts in the marrow not attributable to bone marrow regeneration or another cause;
- Extramedullary leukaemia.

A patient who has progressive disease when treated with this drug is no longer eligible for PBS-subsidised treatment with this drug.

midostaurin 25 mg capsule, 112

11505M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	2	..	20413.00	Rydapt [NV]

■ MIDOSTAURIN

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Acute Myeloid Leukaemia

Treatment Phase: Maintenance therapy - Grandfathered treatment

Clinical criteria:

- Patient must have received non-PBS subsidised treatment with this drug for this condition prior to 1 December 2018, **AND**
- Patient must be receiving treatment with this drug for this condition at the time of application, **AND**
- Patient must not have developed disease progression while receiving treatment with this drug for this condition, **AND**
- Patient must have demonstrated complete remission after induction and consolidation chemotherapy in combination with midostaurin, **AND**
- Patient must not be undergoing or have undergone a stem cell transplant, **AND**
- The condition must have been internal tandem duplication (ITD) or tyrosine kinase domain (TKD) FMS tyrosine kinase 3 (FLT3) mutation positive before initiating this drug for this condition.

A maximum of 2 cycles will be authorised under this restriction in a lifetime.

A patient may qualify for PBS-subsidised treatment under this restriction once only.

For continuing PBS-subsidised treatment, a Grandfathered patient must qualify under the maintenance therapy continuing treatment criteria.

Progressive disease monitoring via a complete blood count must be taken at the end of each cycle.

If abnormal blood counts suggest the potential for relapsed AML, a bone marrow biopsy must be performed to confirm the absence of progressive disease for the patient to be eligible for further cycles.

Progressive disease is defined as the presence of any of the following:

- Leukaemic cells in the CSF;
- Re-appearance of circulating blast cells in the peripheral blood, not attributable to overshoot following recovery from myeloablative therapy;
- Greater than 5 % blasts in the marrow not attributable to bone marrow regeneration or another cause;
- Extramedullary leukaemia.

A patient who has progressive disease when treated with this drug is no longer eligible for PBS-subsidised treatment with this drug.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form;
- (2) a completed Acute myeloid leukaemia PBS Authority Application - Supporting Information Form; and
- (3) confirmation that the patient is not undergoing or has not undergone a stem cell transplant; and
- (4) confirmation that the patient does not have progressive disease; and
- (5) a copy of a recent bone marrow biopsy report demonstrating that the patient is in complete remission; and
- (6) a copy of the pathology test demonstrating that the condition was FMS tyrosine kinase 3 (FLT3) (ITD or TKD) mutation positive prior to commencing midostaurin.

midostaurin 25 mg capsule, 112

11542L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	1	..	20413.00	Rydapt [NV]

■ MIDOSTAURIN

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Acute Myeloid Leukaemia

Treatment Phase: Maintenance therapy - Initial treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while receiving PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have demonstrated complete remission after induction and consolidation chemotherapy in combination with midostaurin, **AND**
- Patient must not be undergoing or have undergone a stem cell transplant, **AND**
- The condition must have been internal tandem duplication (ITD) or tyrosine kinase domain (TKD) FMS tyrosine kinase 3 (FLT3) mutation positive before initiating this drug for this condition.

A maximum of 3 cycles will be authorised under this restriction in a lifetime.

Progressive disease monitoring via a complete blood count must be taken at the end of each cycle.

If abnormal blood counts suggest the potential for relapsed AML, a bone marrow biopsy must be performed to confirm the absence of progressive disease for the patient to be eligible for further cycles.

Progressive disease is defined as the presence of any of the following:

- Leukaemic cells in the CSF;
- Re-appearance of circulating blast cells in the peripheral blood, not attributable to overshoot following recovery from myeloablative therapy;
- Greater than 5 % blasts in the marrow not attributable to bone marrow regeneration or another cause;
- Extramedullary leukaemia.

A patient who has progressive disease when treated with this drug is no longer eligible for PBS-subsidised treatment with this drug.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form;
- (2) a completed Acute myeloid leukaemia PBS Authority Application - Supporting Information Form; and
- (3) confirmation that the patient is not undergoing or has not undergone a stem cell transplant; and
- (4) confirmation that the patient does not have progressive disease; and
- (5) a copy of a recent bone marrow biopsy report demonstrating that the patient is in complete remission; and
- (6) a copy of the pathology test demonstrating that the condition was FMS tyrosine kinase 3 (FLT3) (ITD or TKD) mutation positive prior to commencing midostaurin.

midostaurin 25 mg capsule, 112

11552B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	2	..	20413.00	Rydapt [NV]

▪ **MIDOSTAURIN**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Applications for authority to prescribe may be made by phone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday)

Authority required

Acute Myeloid Leukaemia

Treatment Phase: Induction / Consolidation therapy

Clinical criteria:

- Patient must not have received prior chemotherapy as induction therapy for this condition; OR
- The treatment must be for consolidation treatment following induction treatment with midostaurin in combination with chemotherapy, **AND**
- The condition must be internal tandem duplication (ITD) or tyrosine kinase domain (TKD) FMS tyrosine kinase 3 (FLT3) mutation positive before initiating this drug for this condition, **AND**
- The condition must not be acute promyelocytic leukaemia, **AND**
- The treatment must be in combination with standard intensive remission induction or consolidation chemotherapy for this condition.

A maximum of 6 cycles will be authorised under this restriction in a lifetime.

Standard intensive remission induction combination chemotherapy must include cytarabine and an anthracycline.

The FLT3 ITD or TKD mutation test result and date of testing must be provided at the time of application.

This drug is not PBS-subsidised if it is prescribed to an in-patient in a public hospital setting.

Progressive disease monitoring via a complete blood count must be taken at the end of each cycle.

If abnormal blood counts suggest the potential for relapsed AML, a bone marrow biopsy must be performed to confirm the absence of progressive disease for the patient to be eligible for further cycles.

Progressive disease is defined as the presence of any of the following:

- Leukaemic cells in the CSF;

- Re-appearance of circulating blast cells in the peripheral blood, not attributable to overshoot following recovery from myeloablative therapy;
- Greater than 5 % blasts in the marrow not attributable to bone marrow regeneration or another cause;
- Extramedullary leukaemia.

A patient who has progressive disease when treated with this drug is no longer eligible for PBS-subsidised treatment with this drug.

midostaurin 25 mg capsule, 56

11553C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	2	..	10206.50	Rydapt [NV]

IMMUNOSTIMULANTS

IMMUNOSTIMULANTS

Colony stimulating factors

FILGRASTIM

Authority required (STREAMLINED)

7822

Chemotherapy-induced neutropenia

Clinical criteria:

- Patient must be receiving chemotherapy with the intention of achieving a cure or a substantial remission, **AND**
- Patient must be at greater than 20% risk of developing febrile neutropenia; OR
- Patient must be at substantial risk (greater than 20%) of prolonged severe neutropenia for more than or equal to seven days.

Authority required (STREAMLINED)

7843

Chemotherapy-induced neutropenia

Clinical criteria:

- Patient must be receiving chemotherapy with the intention of achieving a cure or a substantial remission, **AND**
- Patient must have had a prior episode of febrile neutropenia; OR
- Patient must have had a prior episode of prolonged severe neutropenia for more than or equal to seven days.

Authority required (STREAMLINED)

6653

Mobilisation of peripheral blood progenitor cells

Clinical criteria:

- The treatment must be to facilitate harvest of peripheral blood progenitor cells for autologous transplantation into a patient with a non-myeloid malignancy who has had myeloablative or myelosuppressive therapy.

Authority required (STREAMLINED)

6654

Mobilisation of peripheral blood progenitor cells

Clinical criteria:

- The treatment must be in a normal volunteer for use in allogeneic transplantation.

Authority required (STREAMLINED)

6679

Assisting bone marrow transplantation

Clinical criteria:

- Patient must be receiving marrow-ablative chemotherapy prior to the transplantation.

Authority required (STREAMLINED)

6655

Assisting autologous peripheral blood progenitor cell transplantation

Clinical criteria:

- The treatment must be following marrow-ablative chemotherapy for non-myeloid malignancy prior to the transplantation.

Authority required (STREAMLINED)

6680

Severe congenital neutropenia

Clinical criteria:

- Patient must have an absolute neutrophil count of less than 100 million cells per litre measured on 3 occasions, with readings at least 2 weeks apart, **AND**
- Patient must have had a bone marrow examination that has shown evidence of maturational arrest of the neutrophil lineage.

Authority required (STREAMLINED)

6621

Severe chronic neutropenia

Clinical criteria:

- Patient must have an absolute neutrophil count of less than 1,000 million cells per litre measured on 3 occasions, with readings at least 2 weeks apart; OR
- Patient must have neutrophil dysfunction, **AND**

- Patient must have experienced a life-threatening infectious episode requiring hospitalisation and treatment with intravenous antibiotics in the previous 12 months; OR
- Patient must have had at least 3 recurrent clinically significant infections in the previous 12 months.

Authority required (STREAMLINED)

6640

Chronic cyclical neutropenia

Clinical criteria:

- Patient must have an absolute neutrophil count of less than 500 million cells per litre lasting for 3 days per cycle, measured over 3 separate cycles, **AND**
- Patient must have experienced a life-threatening infectious episode requiring hospitalisation and treatment with intravenous antibiotics; OR
- Patient must have had at least 3 recurrent clinically significant infections in the previous 12 months.

filgrastim 120 microgram/0.2 mL injection, 10 x 0.2 mL syringes

5829T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	11	..	*247.68	Nivestim [PF]

▪ **FILGRASTIM**

Note Pharmaceutical benefits that have the forms filgrastim 300 microgram/0.5 mL injection, 10 x 0.5 mL syringes and filgrastim 300 microgram/0.5 mL injection, 5 x 0.5 mL syringes and filgrastim 300 microgram/mL injection, 10 x 1 mL vials are equivalent for the purposes of substitution.

Authority required (STREAMLINED)

7822

Chemotherapy-induced neutropenia

Clinical criteria:

- Patient must be receiving chemotherapy with the intention of achieving a cure or a substantial remission, **AND**
- Patient must be at greater than 20% risk of developing febrile neutropenia; OR
- Patient must be at substantial risk (greater than 20%) of prolonged severe neutropenia for more than or equal to seven days.

Authority required (STREAMLINED)

7843

Chemotherapy-induced neutropenia

Clinical criteria:

- Patient must be receiving chemotherapy with the intention of achieving a cure or a substantial remission, **AND**
- Patient must have had a prior episode of febrile neutropenia; OR
- Patient must have had a prior episode of prolonged severe neutropenia for more than or equal to seven days.

Authority required (STREAMLINED)

6653

Mobilisation of peripheral blood progenitor cells

Clinical criteria:

- The treatment must be to facilitate harvest of peripheral blood progenitor cells for autologous transplantation into a patient with a non-myeloid malignancy who has had myeloablative or myelosuppressive therapy.

Authority required (STREAMLINED)

6654

Mobilisation of peripheral blood progenitor cells

Clinical criteria:

- The treatment must be in a normal volunteer for use in allogeneic transplantation.

Authority required (STREAMLINED)

6679

Assisting bone marrow transplantation

Clinical criteria:

- Patient must be receiving marrow-ablative chemotherapy prior to the transplantation.

Authority required (STREAMLINED)

6655

Assisting autologous peripheral blood progenitor cell transplantation

Clinical criteria:

- The treatment must be following marrow-ablative chemotherapy for non-myeloid malignancy prior to the transplantation.

Authority required (STREAMLINED)

6680

Severe congenital neutropenia

Clinical criteria:

- Patient must have an absolute neutrophil count of less than 100 million cells per litre measured on 3 occasions, with readings at least 2 weeks apart, **AND**
- Patient must have had a bone marrow examination that has shown evidence of maturational arrest of the neutrophil lineage.

Authority required (STREAMLINED)

6621

Severe chronic neutropenia

Clinical criteria:

- Patient must have an absolute neutrophil count of less than 1,000 million cells per litre measured on 3 occasions, with readings at least 2 weeks apart; OR
- Patient must have neutrophil dysfunction, **AND**
- Patient must have experienced a life-threatening infectious episode requiring hospitalisation and treatment with intravenous antibiotics in the previous 12 months; OR
- Patient must have had at least 3 recurrent clinically significant infections in the previous 12 months.

Authority required (STREAMLINED)
6640

Chronic cyclical neutropenia

Clinical criteria:

- Patient must have an absolute neutrophil count of less than 500 million cells per litre lasting for 3 days per cycle, measured over 3 separate cycles, **AND**
- Patient must have experienced a life-threatening infectious episode requiring hospitalisation and treatment with intravenous antibiotics; OR
- Patient must have had at least 3 recurrent clinically significant infections in the previous 12 months.

filgrastim 300 microgram/0.5 mL injection, 5 x 0.5 mL syringes

2758E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	4	11	..	*619.20	^a Zarzio [SZ]

filgrastim 300 microgram/mL injection, 10 x 1 mL vials

5741E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	11	..	*619.20	^a Neupogen [AN]

filgrastim 300 microgram/0.5 mL injection, 10 x 0.5 mL syringes

5742F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	11	..	*619.20	^a Neupogen [AN]	^a Nivestim [PF]

■ FILGRASTIM

Note Pharmaceutical benefits that have the forms filgrastim 480 microgram/0.5 mL injection, 10 x 0.5 mL syringes and filgrastim 480 microgram/1.6 mL injection, 10 x 1.6 mL vials and filgrastim 480 microgram/0.5 mL injection, 5 x 0.5 mL syringes are equivalent for the purposes of substitution.

Authority required (STREAMLINED)
7822

Chemotherapy-induced neutropenia

Clinical criteria:

- Patient must be receiving chemotherapy with the intention of achieving a cure or a substantial remission, **AND**
- Patient must be at greater than 20% risk of developing febrile neutropenia; OR
- Patient must be at substantial risk (greater than 20%) of prolonged severe neutropenia for more than or equal to seven days.

Authority required (STREAMLINED)
7843

Chemotherapy-induced neutropenia

Clinical criteria:

- Patient must be receiving chemotherapy with the intention of achieving a cure or a substantial remission, **AND**
- Patient must have had a prior episode of febrile neutropenia; OR
- Patient must have had a prior episode of prolonged severe neutropenia for more than or equal to seven days.

Authority required (STREAMLINED)
6653

Mobilisation of peripheral blood progenitor cells

Clinical criteria:

- The treatment must be to facilitate harvest of peripheral blood progenitor cells for autologous transplantation into a patient with a non-myeloid malignancy who has had myeloablative or myelosuppressive therapy.

Authority required (STREAMLINED)
6654

Mobilisation of peripheral blood progenitor cells

Clinical criteria:

- The treatment must be in a normal volunteer for use in allogeneic transplantation.

Authority required (STREAMLINED)
6679

Assisting bone marrow transplantation

Clinical criteria:

- Patient must be receiving marrow-ablative chemotherapy prior to the transplantation.

Authority required (STREAMLINED)
6655

Assisting autologous peripheral blood progenitor cell transplantation

Clinical criteria:

- The treatment must be following marrow-ablative chemotherapy for non-myeloid malignancy prior to the transplantation.

Authority required (STREAMLINED)

6680

Severe congenital neutropenia

Clinical criteria:

- Patient must have an absolute neutrophil count of less than 100 million cells per litre measured on 3 occasions, with readings at least 2 weeks apart, **AND**
- Patient must have had a bone marrow examination that has shown evidence of maturational arrest of the neutrophil lineage.

Authority required (STREAMLINED)

6621

Severe chronic neutropenia

Clinical criteria:

- Patient must have an absolute neutrophil count of less than 1,000 million cells per litre measured on 3 occasions, with readings at least 2 weeks apart; OR
- Patient must have neutrophil dysfunction, **AND**
- Patient must have experienced a life-threatening infectious episode requiring hospitalisation and treatment with intravenous antibiotics in the previous 12 months; OR
- Patient must have had at least 3 recurrent clinically significant infections in the previous 12 months.

Authority required (STREAMLINED)

6640

Chronic cyclical neutropenia

Clinical criteria:

- Patient must have an absolute neutrophil count of less than 500 million cells per litre lasting for 3 days per cycle, measured over 3 separate cycles, **AND**
- Patient must have experienced a life-threatening infectious episode requiring hospitalisation and treatment with intravenous antibiotics; OR
- Patient must have had at least 3 recurrent clinically significant infections in the previous 12 months.

filgrastim 480 microgram/1.6 mL injection, 10 x 1.6 mL vials

5743G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	11	..	*992.62	^a Neupogen [AN]

filgrastim 480 microgram/0.5 mL injection, 5 x 0.5 mL syringes

2783L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	4	11	..	*992.60	^a Zarzio [SZ]

filgrastim 480 microgram/0.5 mL injection, 10 x 0.5 mL syringes

5744H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	11	..	*992.60	^a Neupogen [AN]	^a Nivestim [PF]

■ LENOGRASTIM

Authority required (STREAMLINED)

6522

Chemotherapy-induced neutropenia

Clinical criteria:

- Patient must be receiving standard dose adjuvant chemotherapy for breast cancer, **AND**
- Patient must have had a prior episode of febrile neutropenia; OR
- Patient must have had a prior episode of prolonged severe neutropenia (neutrophil count of less than 1,000 million cells per litre), **AND**
- The treatment must be used in a patient for whom there is a clinical justification for wishing to continue chemotherapy with the same drug combination, dosage and treatment schedule, **AND**
- Patient must be anticipated to have a good response to treatment providing chemotherapy can be delivered as planned.

Authority required (STREAMLINED)

6532

Chemotherapy-induced neutropenia

Clinical criteria:

- Patient must be receiving first-line chemotherapy for Hodgkin disease, **AND**
- Patient must have had a prior episode of febrile neutropenia; OR
- Patient must have had a prior episode of prolonged severe neutropenia (neutrophil count of less than 1,000 million cells per litre), **AND**
- The treatment must be used in a patient for whom there is a clinical justification for wishing to continue chemotherapy with the same drug combination, dosage and treatment schedule, **AND**
- Patient must be anticipated to have a good response to treatment providing chemotherapy can be delivered as planned.

Authority required (STREAMLINED)

6507

Chemotherapy-induced neutropenia

Clinical criteria:

- Patient must be receiving treatment with aggressive chemotherapy with the intention of achieving a cure or substantial remission in acute lymphoblastic leukaemia.

Authority required (STREAMLINED)

6523

Chemotherapy-induced neutropenia

Clinical criteria:

- Patient must be receiving treatment with aggressive chemotherapy with the intention of achieving a cure or substantial remission in germ cell tumours.

Authority required (STREAMLINED)

6535

Chemotherapy-induced neutropenia

Clinical criteria:

- Patient must be receiving treatment with aggressive chemotherapy with the intention of achieving a cure or substantial remission in relapsed Hodgkin disease.

Authority required (STREAMLINED)

6502

Chemotherapy-induced neutropenia

Clinical criteria:

- Patient must be receiving treatment with aggressive chemotherapy with the intention of achieving a cure or substantial remission in an infant or child with central nervous system tumours.

Authority required (STREAMLINED)

6516

Chemotherapy-induced neutropenia

Clinical criteria:

- Patient must be receiving treatment with aggressive chemotherapy with the intention of achieving a cure or substantial remission in neuroblastoma.

Authority required (STREAMLINED)

6644

Chemotherapy-induced neutropenia

Clinical criteria:

- Patient must be receiving treatment with aggressive chemotherapy with the intention of achieving a cure or substantial remission in Ewing's sarcoma.

Authority required (STREAMLINED)

6673

Chemotherapy-induced neutropenia

Clinical criteria:

- Patient must be receiving treatment with aggressive chemotherapy with the intention of achieving a cure or substantial remission in non-Hodgkin's lymphoma (intermediate or high grade).

Authority required (STREAMLINED)

6634

Chemotherapy-induced neutropenia

Clinical criteria:

- Patient must be receiving treatment with aggressive chemotherapy with the intention of achieving a cure or substantial remission in osteosarcoma.

Authority required (STREAMLINED)

6682

Chemotherapy-induced neutropenia

Clinical criteria:

- Patient must be receiving treatment with aggressive chemotherapy with the intention of achieving a cure or substantial remission in rhabdomyosarcoma.

Authority required (STREAMLINED)

6653

Mobilisation of peripheral blood progenitor cells

Clinical criteria:

- The treatment must be to facilitate harvest of peripheral blood progenitor cells for autologous transplantation into a patient with a non-myeloid malignancy who has had myeloablative or myelosuppressive therapy.

Authority required (STREAMLINED)

6654

Mobilisation of peripheral blood progenitor cells

Clinical criteria:

- The treatment must be in a normal volunteer for use in allogeneic transplantation.

Authority required (STREAMLINED)

6657

Assisting peripheral blood progenitor cell or bone marrow transplantation

Clinical criteria:

- The treatment must be following marrow-ablative chemotherapy for non-myeloid malignancy prior to the transplantation.

lenograstim 33.6 million units (263 microgram) injection, 1 vial

11551Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	20	11	..	*2085.20	Granocyte 34 [PF]

lenograstim 13.4 million units (105 microgram) injection, 1 vial

11550X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	20	11	..	*832.60	Granocyte 13 [PF]

■ LIPEGFILGRASTIM
Authority required (STREAMLINED)
7822

Chemotherapy-induced neutropenia

Clinical criteria:

- Patient must be receiving chemotherapy with the intention of achieving a cure or a substantial remission, **AND**
- Patient must be at greater than 20% risk of developing febrile neutropenia; OR
- Patient must be at substantial risk (greater than 20%) of prolonged severe neutropenia for more than or equal to seven days.

Authority required (STREAMLINED)
7843

Chemotherapy-induced neutropenia

Clinical criteria:

- Patient must be receiving chemotherapy with the intention of achieving a cure or a substantial remission, **AND**
- Patient must have had a prior episode of febrile neutropenia; OR
- Patient must have had a prior episode of prolonged severe neutropenia for more than or equal to seven days.

lipegfilgrastim 6 mg/0.6 mL injection, 0.6 mL syringe

10936N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	11	..	1175.00	Lonquex [TB]

■ PEGFILGRASTIM
Authority required (STREAMLINED)
7822

Chemotherapy-induced neutropenia

Clinical criteria:

- Patient must be receiving chemotherapy with the intention of achieving a cure or a substantial remission, **AND**
- Patient must be at greater than 20% risk of developing febrile neutropenia; OR
- Patient must be at substantial risk (greater than 20%) of prolonged severe neutropenia for more than or equal to seven days.

Authority required (STREAMLINED)
7843

Chemotherapy-induced neutropenia

Clinical criteria:

- Patient must be receiving chemotherapy with the intention of achieving a cure or a substantial remission, **AND**
- Patient must have had a prior episode of febrile neutropenia; OR
- Patient must have had a prior episode of prolonged severe neutropenia for more than or equal to seven days.

pegfilgrastim 6 mg/0.6 mL injection, 0.6 mL syringe

9514R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	11	..	1175.00	^a Neulasta [JU] ^a Tezmota [JX]	^a Ristempa [JO]

Interferons
■ INTERFERON ALFA-2A

Caution Treatment with interferon alfa has been associated with depression and suicide in some patients. Patients with a history of suicidal ideation or depressive illness should be warned of the risks. Psychiatric status during therapy should be monitored.

Authority required (STREAMLINED)
5042

Chronic Myeloid Leukaemia (CML)

Clinical criteria:

- The condition must be Philadelphia chromosome positive.

interferon alfa-2a 3 million units/0.5 mL injection, 0.5 mL syringe

5759D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	30	5	..	*726.30	Roferon-A [RO]

interferon alfa-2a 9 million units/0.5 mL injection, 0.5 mL syringe

5762G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	30	5	..	*2178.00	Roferon-A [RO]

■ INTERFERON GAMMA-1B

Authority required (STREAMLINED)

6222

Chronic granulomatous disease

Clinical criteria:

- Patient must have frequent and severe infections despite adequate prophylaxis with antimicrobial agents.

interferon gamma-1b 2 million units (100 microgram)/0.5 mL injection, 6 x 0.5 mL vials

5769P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	11	..	*2210.80	Imukin [EU]

■ PEGINTERFERON ALFA-2A

Caution Treatment with peginterferon alfa has been associated with depression and suicide in some patients. Patients with a history of suicidal ideation or depressive illness should be warned of the risks. Psychiatric status during therapy should be monitored.

Note Treatment centres are required to have access to the following appropriate specialist facilities for the provision of clinical support services for hepatitis C:

- (a) a nurse educator/counsellor for patients; and
- (b) 24-hour access by patients to medical advice; and
- (c) an established liver clinic.

Authority required (STREAMLINED)

5004

Chronic hepatitis C infection

Treatment criteria:

- Must be treated in an accredited treatment centre.

Population criteria:

- Patient must be aged 18 years or older, **AND**
- Patient must not be pregnant or breastfeeding, and must be using an effective form of contraception if female and of child-bearing age.

Clinical criteria:

- Patient must have compensated liver disease, **AND**
- Patient must not have received prior interferon alfa or peginterferon alfa treatment for hepatitis C, **AND**
- Patient must have a contraindication to ribavirin, **AND**
- The treatment must cease unless the results of an HCV RNA quantitative assay at week 12 (performed at the same laboratory using the same test) show that plasma HCV RNA has become undetectable or the viral load has decreased by at least a 2 log drop, **AND**
- The treatment must be limited to a maximum duration of 48 weeks.

Evidence of chronic hepatitis C infection (repeatedly anti-HCV positive and HCV RNA positive) must be documented in the patient's medical records.

peginterferon alfa-2a 135 microgram/0.5 mL injection, 4 x 0.5 mL syringes

9515T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*1107.62	Pegasys [RO]

peginterferon alfa-2a 180 microgram/0.5 mL injection, 4 x 0.5 mL syringes

9516W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*1282.72	Pegasys [RO]

■ PEGINTERFERON ALFA-2A

Caution Treatment with peginterferon alfa has been associated with depression and suicide in some patients. Patients with a history of suicidal ideation or depressive illness should be warned of the risks. Psychiatric status during therapy should be monitored.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Chronic hepatitis C infection

Clinical criteria:

- Patient must meet the criteria set out in the General Statement for Drugs for the Treatment of Hepatitis C, **AND**
- Patient must be taking this drug as part of a regimen set out in the matrix in the General Statement for Drugs for the Treatment of Hepatitis C, based on the hepatitis C virus genotype, patient treatment history and cirrhotic status, **AND**
- The treatment must be limited to a maximum duration of 12 weeks.

peginterferon alfa-2a 180 microgram/0.5 mL injection, 4 x 0.5 mL syringes

11026H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	2	..	641.36	Pegasys [RO]

Other immunostimulants

■ PLERIXAFOR

Note Special Pricing Arrangements apply.

Note Applications for increased maximum quantities will only be authorised for patients with body weight greater than 100 kg.

Authority required (STREAMLINED)

4549

Mobilisation of haematopoietic stem cells

Clinical criteria:

- The treatment must be in combination with granulocyte-colony stimulating factor (G-CSF), **AND**
- Patient must have lymphoma; OR
- Patient must have multiple myeloma, **AND**
- Patient must require autologous stem cell transplantation, **AND**
- Patient must have failed previous stem cell collection; OR
- Patient must be undergoing chemotherapy plus G-CSF mobilisation and their peripheral blood CD34+ count is less than 10,000 per millilitre or less than 10 million per litre on the day of planned collection; OR
- Patient must be undergoing chemotherapy plus G-CSF mobilisation and the first apheresis has yielded less than 1 million CD34+ cells/kg.

Evidence that the patient meets the PBS restriction criteria must be recorded in the patient's medical records.

plerixafor 24 mg/1.2 mL injection, 1.2 mL vial

10083Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	1	..	6991.00	Mozobil [GZ]

■ IMMUNOSUPPRESSANTS

IMMUNOSUPPRESSANTS

Selective immunosuppressants

■ ABATACEPT

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for adults with severe active rheumatoid arthritis. Where the term biological medicine appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alfa antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab), the T-cell co-stimulation modulator (abatacept) and the Janus kinase (JAK) inhibitors (baricitinib, tofacitinib). A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised biological medicine therapy may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once, and
- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised biological medicines for the treatment of rheumatoid arthritis.

Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Stevens Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

For a patient who has failed PBS-subsidised treatment with 2 or 3 TNF-alfa antagonists prior to 1 August 2010 please contact the Department of Human Services on 1800 700 270. A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction.

A patient who has failed fewer than 5 biological medicines and who has a break in treatment of less than 24 months may commence a further course of treatment with a biological medicine under Initial 2 treatment restriction. A patient who has failed fewer than 5 biological medicines and who has had a break in therapy of longer than 24 months may commence a further course of treatment with a biological medicine under the Initial 3 treatment restriction. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine treatment is stopped to the date of the new application for treatment with a biological medicine.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 April 2019.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised biological medicine treatment and wishes to commence such therapy, excluding rituximab (Initial 1 - new patient); or

(ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to re-commence treatment with a specific biological medicine following a break of less than 24 months in PBS-subsidised therapy with that agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months).

(iv) a patient wishes to re-commence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 24 months (Initial 3 - re-commencement of treatment after a break in biological medicine of more than 24 months)

Initial applications for a new patient (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6-month intensive DMARD trial, but prior to ceasing DMARD therapy.

Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, baricitinib, etanercept, golimumab, tocilizumab and tofacitinib, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

Rituximab patients should be assessed following a minimum of 12 weeks after the first infusion, and the assessment should be submitted to the Department of Human Services within 4 weeks to ensure continuity of treatment for those who meet the continuing restriction for PBS subsidised treatment with this drug for this condition.

Where a response assessment is not provided the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment. For second and subsequent courses of PBS-subsidised biological medicine (excluding rituximab) treatment it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to the Department of Human Services within sufficient timeframe to allow for authority assessment.

Abatacept patients:

A patient is eligible to receive one I.V. loading dose when commencing treatment with the subcutaneous formulation. Two prescriptions are required, the first prescription for the I.V. loading dose for sufficient vials for one dose based on the patient's weight with no repeats. The second prescription for the subcutaneous formulation, with a maximum quantity of 4 and up to 3 repeats, must be submitted with the initial application.

Rituximab patients:

A further application may be submitted to the Department of Human Services within sufficient timeframe to allow authority assessment. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

Infliximab and etanercept patients:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction. Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

(2) Swapping therapy

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non- biological medicine therapy requirements except if the patient has had a break in therapy of more than 24 months who would need to requalify with respect to the indices of disease severity. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each biological medicine trialled.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent, unless they have a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug.

Abatacept:

A patient swapping from I.V. abatacept to subcutaneous abatacept will not be eligible for an I.V. loading dose when commencing treatment with the subcutaneous formulation.

Rituximab:

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF-alfa antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological medicine during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological medicine therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate biological medicine may do so without having to have any treatment-free period.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and the Department of Human Services will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints.

Applications under the Initial 1 treatment restriction for a new patient must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy.

Applications under the Initial 3 treatment restriction for re-commencement of treatment after a break in biological medicine of more than 24 months must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with disease modifying anti-rheumatic drugs (DMARDs) which must include at least 3 months continuous treatment with each of at least 2 DMARDs, one of which must be methotrexate at a dose of at least 20 mg weekly and one of which must be: (i) hydroxychloroquine at a dose of at least 200 mg daily; or (ii) leflunomide at a dose of at least 10 mg daily; or (iii) sulfasalazine at a dose of at least 2 g daily; OR
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs which, if methotrexate is contraindicated according to the Therapeutic Goods Administration (TGA)-approved Product Information or cannot be tolerated at a 20 mg weekly dose, must include at least 3 months continuous treatment with each of at least 2 of the following DMARDs: (i) hydroxychloroquine at a dose of at least 200 mg daily; and/or (ii) leflunomide at a dose of at least 10 mg daily; and/or (iii) sulfasalazine at a dose of at least 2 g daily; OR
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs which, if 3 or more of methotrexate, hydroxychloroquine, leflunomide and sulfasalazine are contraindicated according to the relevant TGA-approved Product Information or cannot be tolerated at the doses specified above, must include at least 3 months continuous treatment with each of at least 2 DMARDs, with one or more of the following DMARDs being used in place of the DMARDs which are contraindicated or not tolerated: (i) azathioprine at a dose of at least 1 mg/kg per day; and/or (ii) cyclosporin at a dose of at least 2 mg/kg/day; and/or (iii) sodium aurothiomalate at a dose of 50 mg weekly, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction, **AND**
- The treatment must be given concomitantly with methotrexate at a dose of at least 7.5 mg weekly.

Population criteria:

- Patient must be aged 18 years or older.

If methotrexate is contraindicated according to the TGA-approved product information or cannot be tolerated at a 20 mg weekly dose, the application must include details of the contraindication or intolerance including severity to methotrexate. The maximum tolerated dose of methotrexate must be documented in the application, if applicable.

The application must include details of the DMARDs trialed, their doses and duration of treatment, and all relevant contraindications and/or intolerances including severity.

The requirement to trial at least 2 DMARDs for periods of at least 3 months each can be met using single agents sequentially or by using one or more combinations of DMARDs.

If the requirement to trial 6 months of intensive DMARD therapy with at least 2 DMARDs cannot be met because of contraindications and/or intolerances of a severity necessitating permanent treatment withdrawal to all of the DMARDs specified above, details of the contraindication or intolerance including severity and dose for each DMARD must be provided in the authority application.

The following criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; **AND** either

- (a) a total active joint count of at least 20 active (swollen and tender) joints; or
- (b) at least 4 active joints from the following list of major joints:
 - (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The joint count and ESR and/or CRP must be determined at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. All measures must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

At the time of authority application, medical practitioners should request the appropriate number of vials to provide sufficient drug, based on the weight of the patient, for a single infusion.

Up to a maximum of 4 repeats will be authorised.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note The Department of Human Services website (www.humanservices.gov.au) has details of the toxicities, including severity, which will be accepted for the following purposes:

- (a) exempting a patient from the requirement to undertake a minimum 3 month trial of methotrexate at a 20 mg weekly dose;
- (b) substituting azathioprine, cyclosporin or sodium aurothiomalate for another DMARD as part of the 6 month intensive DMARD trial;
- (c) exempting a patient from the requirement for a 6 month trial of intensive DMARD therapy.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 2 (change or re-commencement of treatment after a break in biological medicine of less than 24 months).

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must not have failed to respond to previous PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised biological medicine treatment for this condition 5 times, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction, **AND**
- The treatment must be given concomitantly with methotrexate at a dose of at least 7.5 mg weekly.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either of the Initial 1, Initial 2, Initial 3, or continuing treatment restrictions, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the

date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

At the time of authority application, medical practitioners should request the appropriate number of vials to provide sufficient drug, based on the weight of the patient, for a single infusion.

Up to a maximum of 4 repeats will be authorised.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

A patient who has demonstrated a response to a course of rituximab must have a PBS-subsidised biological therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 3 (re-commencement of treatment after a break in biological medicine of more than 24 months)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 24 months or more from the most recent PBS-subsidised biological medicine for this condition, **AND**
- Patient must not have failed to respond to previous PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised biological medicine treatment for this condition 5 times, **AND**
- The condition must have an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; OR
- The condition must have a C-reactive protein (CRP) level greater than 15 mg per L, **AND**
- The condition must have either (a) a total active joint count of at least 20 active (swollen and tender) joints; or (b) at least 4 active major joints, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction, **AND**
- The treatment must be given concomitantly with methotrexate at a dose of at least 7.5 mg weekly.

Population criteria:

- Patient must be aged 18 years or older.

Major joints are defined as (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

All measures of joint count and ESR and/or CRP must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial 1 (new patient) or Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 24 months) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 24 months) - balance of supply

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 24 months) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 24 months) to complete 16 weeks of treatment, **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction, **AND**
- The treatment must be given concomitantly with methotrexate at a dose of at least 7.5 mg weekly.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient has either failed or ceased to respond to a PBS-subsidised biological medicine for this condition 5 times, they will not be eligible to receive further PBS-subsidised treatment with a biological medicine for this condition.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Continuing Treatment - balance of supply.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restriction.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

abatacept 250 mg injection, 1 vial

5605B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	266.34	Orencia [BQ]

■ ALEMTUZUMAB

Note Neurologists prescribing PBS-subsidised alemtuzumab must be registered with the Lemtrada monitoring program.

Note Special Pricing Arrangements apply.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

6847

Multiple sclerosis

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not show continuing progression of disability while on treatment with this drug, **AND**
- Patient must not receive more than one PBS-subsidised treatment per year, **AND**
- The treatment must be a sole PBS-subsidised disease modifying therapy for this condition, **AND**
- Patient must have demonstrated compliance with, and an ability to tolerate this therapy.

Treatment criteria:

- Must be treated by a neurologist.

alemtuzumab 12 mg/1.2 mL injection, 1.2 mL vial

10232M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	3	..	..	*34182.00	Lemtrada [GZ]

■ ALEMTUZUMAB

Note Neurologists prescribing PBS-subsidised alemtuzumab must be registered with the Lemtrada monitoring program.

Note Special Pricing Arrangements apply.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

7714

Multiple sclerosis

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be diagnosed as clinically definite relapsing-remitting multiple sclerosis by magnetic resonance imaging of the brain and/or spinal cord; OR

- The condition must be diagnosed as clinically definite relapsing-remitting multiple sclerosis by accompanying written certification provided by a radiologist that a magnetic resonance imaging scan is contraindicated because of the risk of physical (not psychological) injury to the patient, **AND**
- The treatment must be a sole PBS-subsidised disease modifying therapy for this condition, **AND**
- Patient must have experienced at least 2 documented attacks of neurological dysfunction, believed to be due to multiple sclerosis, in the preceding 2 years of commencing a PBS-subsidised disease modifying therapy for this condition, **AND**
- Patient must be ambulatory (without assistance or support).

Treatment criteria:

- Must be treated by a neurologist.

Where applicable, the date of the magnetic resonance imaging scan must be recorded in the patient's medical records.

alemtuzumab 12 mg/1.2 mL injection, 1.2 mL vial

10228H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	5	..	..	*56970.00	Lemtrada [GZ]

■ ECULIZUMAB

Note At the time of authority application, medical practitioners must request the appropriate number of vials to provide sufficient drug for 4 weeks and up to 4 repeats, according to the specified dosage in the approved Product Information (PI). Applications for treatment with this drug where the dose and dosing frequency exceeds that specified in the approved PI will not be approved.

Note WARNING: Eculizumab increases the risk of meningococcal infections (septicaemia and/or meningitis).

Please consult the approved PI for information about vaccination against meningococcal infection.

Note Eculizumab is not PBS subsidised to treat TMA caused by conditions other than aHUS, such as TMA occurring in the setting of, but not limited to:

- Active malignancy;
- Active HIV infection;
- Hematopoietic stem cell transplants;
- Various drugs including quinine, high-dose calcineurin inhibitors, antiplatelet agents;
- Certain chemotherapy drugs or immunosuppressant drugs associated with microangiopathic haemolytic anaemia/TMA;
- Active autoimmune diseases;

In cases where alternative causes of TMA have not been adequately excluded, additional information may be required from the prescriber to clarify the diagnosis before approval of the application.

Note The Authority application should be accompanied by a cover letter from the prescriber, providing complete details on:

- Presenting clinical features, including history, acute treatment and medications;
- Results of testing for genetic mutations (if available);
- Family history of aHUS, especially in first-degree relatives;
- Patient's prior history of episodes of active and progressing TMA caused by aHUS;
- Exclusion of alternative causes of TMA;
- History of renal or other organ transplant (if any);
- Any other matters considered relevant by the prescriber.

In cases where there are discordant results (for example, an equivocal biopsy result in the absence of objective evidence of haemolysis) the cover letter should articulate the prescriber's interpretation of the clinical data and how a diagnosis of aHUS is supported by the available evidence.

Note The Authority application should include the results of screening for genetic mutations known to confer a high risk of developing aHUS. The results of genetic screening should be provided whether or not a high-risk mutation has been identified.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Written applications for authority to prescribe must be submitted to Department of Human Services. Human Services will then contact the prescriber by telephone.

Authority required

Atypical haemolytic uraemic syndrome (aHUS)

Treatment Phase: Initial treatment - Balance of Supply

Treatment criteria:

- Must be treated by a paediatric nephrologist, a nephrologist, a paediatric haematologist or a haematologist, or, must be in consultation with a paediatric nephrologist, a nephrologist, a paediatric haematologist or a haematologist.

Clinical criteria:

- Patient must have received PBS-subsidised initial supply of eculizumab for this condition, **AND**
- Patient must have ADAMTS-13 activity of greater than or equal to 10% on a blood sample, **AND**
- Patient must not receive more than 20 weeks supply under this restriction.

ADAMTS-13 activity result must have been submitted to the Department of Human Services. In the case that a sample for ADAMTS-13 activity taken prior to plasma exchange or infusion was not available at the time of application for **Initial**

Treatment, ADAMTS-13 activity must have been measured 1-2 weeks following the last plasma exchange or infusion, and must have been submitted to the Department of Human Services within 27 days of commencement of eculizumab. The date and time that the sample for the ADAMTS-13 assay was collected, and the dates and times of the last, if any, plasma exchange or infusion that was undertaken in the two weeks prior to collection of the ADAMTS-13 assay must also have been provided to Department of Human Services.

Serial haematological results (every 3 months while the patient is receiving treatment) must be provided with every subsequent application for treatment.

eculizumab 300 mg/30 mL injection, 30 mL vial

10190H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	4	..	5937.50	Soliris [XI]

■ ECULIZUMAB

Note At the time of authority application, medical practitioners must request the appropriate number of vials to provide sufficient drug for four weeks of treatment, according to the specified dosage in the approved Product Information (PI). Applications for treatment with this drug where the dose and dosing frequency exceeds that specified in the approved PI will not be approved.

Note WARNING: Eculizumab increases the risk of meningococcal infections (septicaemia and/or meningitis). Please consult the approved PI for information about vaccination against meningococcal infection.

Note Eculizumab is not PBS subsidised to treat TMA caused by conditions other than aHUS, such as TMA occurring in the setting of, but not limited to:

- Active malignancy;
- Active HIV infection;
- Hematopoietic stem cell transplants;
- Various drugs including quinine, high-dose calcineurin inhibitors, antiplatelet agents;
- Certain chemotherapy drugs or immunosuppressant drugs associated with microangiopathic haemolytic anaemia/TMA;
- Active autoimmune diseases;

In cases where alternative causes of TMA have not been adequately excluded, additional information may be required from the prescriber to clarify the diagnosis before approval of the application.

Note The Authority application should be accompanied by a cover letter from the prescriber, providing complete details on:

- Presenting clinical features, including history, acute treatment and medications;
- Results of testing for genetic mutations (if available);
- Family history of aHUS, especially in first-degree relatives;
- Patient's prior history of episodes of active and progressing TMA caused by aHUS;
- Exclusion of alternative causes of TMA;
- History of renal or other organ transplant (if any);
- Any other matters considered relevant by the prescriber.

In cases where there are discordant results (for example, an equivocal biopsy result in the absence of objective evidence of haemolysis) the cover letter should articulate the prescriber's interpretation of the clinical data and how a diagnosis of aHUS is supported by the available evidence.

Note The Authority application should include the results of screening for genetic mutations known to confer a high risk of developing aHUS. The results of genetic screening should be provided whether or not a high-risk mutation has been identified.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Written applications for authority to prescribe must be submitted to Department of Human Services. Human Services will then contact the prescriber by telephone.

Authority required

Atypical haemolytic uraemic syndrome (aHUS)

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have active and progressing thrombotic microangiopathy (TMA) caused by aHUS, **AND**
- Patient must have ADAMTS-13 activity of greater than or equal to 10% on a blood sample taken prior to plasma exchange or infusion; or, if ADAMTS-13 activity was not collected prior to plasma exchange or infusion, patient must have platelet counts of greater than $30 \times 10^9/L$ and a serum creatinine of greater than $150 \mu\text{mol/L}$, **AND**
- Patient must have a confirmed negative STEC (Shiga toxin-producing E.Coli) result if the patient has had diarrhoea in the preceding 14 days, **AND**
- Patient must have clinical features of active organ damage or impairment, **AND**
- Patient must not receive more than 4 weeks of treatment under this restriction.

Treatment criteria:

- Must be treated by a paediatric nephrologist, a nephrologist, a paediatric haematologist or a haematologist, or, must be in consultation with a paediatric nephrologist, a nephrologist, a paediatric haematologist or a haematologist.

Evidence of active and progressing TMA is defined by the following:

(1) a platelet count of less than $150 \times 10^9/L$; and evidence of two of the following:

- presence of schistocytes on blood film;
- low or absent haptoglobin;
- lactate dehydrogenase (LDH) above normal range;

OR

(2) in recipients of a kidney transplant for end-stage kidney disease due to aHUS, a kidney biopsy confirming TMA; **AND**

(3) evidence of at least one of the following clinical features of active TMA-related organ damage or impairment is defined as below:

(a) kidney impairment as demonstrated by one of the following:

- a decline in estimated Glomerular Filtration Rate (eGFR) of greater than 20% in a patient who has pre-existing kidney impairment; and/or

- (ii) a serum creatinine (sCr) of greater than the upper limit of normal (ULN) in a patient who has no history of pre-existing kidney impairment; or
- (iii) a sCr of greater than the age-appropriate ULN in paediatric patients; or
- (iv) a renal biopsy consistent with aHUS;
- (b) onset of TMA-related neurological impairment;
- (c) onset of TMA-related cardiac impairment;
- (d) onset of TMA-related gastrointestinal impairment;
- (e) onset of TMA-related pulmonary impairment.

Claims of non-renal TMA-related organ damage should be made at the point of application for initial PBS-subsidised eculizumab (where possible), and should be supported by objective clinical measures. The prescriber's cover letter should establish that the observed organ damage is directly linked to active and progressing TMA, particularly when indirect causes such as severe thrombocytopenia, hypertension and acute renal failure are present at the time of the initial organ impairment.

Serial haematological results (every 3 months while the patient is receiving treatment) must be provided with every subsequent application for treatment.

The authority application must be in writing and must include:

- (1) A completed authority prescription form; and
- (2) A completed aHUS eculizumab Authority Application Supporting Information Form - Initial PBS-subsidised eculizumab treatment; and
- (3) A signed patient acknowledgement or an acknowledgement signed by a parent or authorised guardian, if applicable; and
- (4) A detailed cover letter from the prescriber; and
- (5) A copy of a current Certificate of vaccination or a statement that vaccination has or will be administered and appropriate antibiotic prophylaxis has been prescribed; and
- (6) A measurement of body weight at the time of application; and
- (7) The result of ADAMTS-13 activity on a blood sample taken prior to plasma exchange or infusion; the date and time that the sample for the ADAMTS-13 assay was collected, and the dates and times of any plasma exchanges or infusions that were undertaken in the two weeks prior to collection of the ADAMTS-13 assay; and
- (8) In the case that a sample for ADAMTS-13 assay was not collected prior to plasma exchange or infusion, measurement of ADAMTS-13 activity must be taken 1-2 weeks following the last plasma exchange or infusion. The ADAMTS-13 result must be submitted to the Department of Human Services within 27 days of commencement of eculizumab treatment in order for the patient to be considered as eligible for further PBS-subsidised eculizumab treatment, under **Initial treatment 1-balance of supply**; and
- (9) A confirmed negative STEC result if the patient has had diarrhoea in the preceding 14 days; and
- (10) Evidence of active and progressing TMA, including pathology results where relevant. Evidence of the onset of TMA-related neurological, cardiac, gastrointestinal or pulmonary impairment requires a supporting statement with clinical evidence in patient records. All tests must have been performed within one month of application; and
- (11) For all patients, a recent measurement of eGFR, platelets and two of either LDH, haptoglobin or schistocytes of no more than 1 week old at the time of application.

eculizumab 300 mg/30 mL injection, 30 mL vial

10191J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	5937.50	Soliris [XI]

■ ECULIZUMAB

Note At the time of authority application, medical practitioners must request the appropriate number of vials to provide sufficient drug for 4 weeks and up to 6 repeats, according to the specified dosage in the approved Product Information (PI). Applications for treatment with this drug where the dose and dosing frequency exceeds that specified in the approved PI will not be approved.

Note For patients who have received continuing treatment with PBS-subsidised eculizumab prior to 1 January 2016, this restriction is limited to 28 weeks of therapy.

Note WARNING: Eculizumab increases the risk of meningococcal infections (septicaemia and/or meningitis). Please consult the approved PI for information about vaccination against meningococcal infection.

Note Eculizumab is not PBS subsidised to treat TMA caused by conditions other than aHUS, such as TMA occurring in the setting of, but not limited to:

- a) Active malignancy;
- b) Active HIV infection;
- c) Hematopoietic stem cell transplants;
- d) Various drugs including quinine, high-dose calcineurin inhibitors, antiplatelet agents;
- e) Certain chemotherapy drugs or immunosuppressant drugs associated with microangiopathic haemolytic anaemia/TMA;
- f) Active autoimmune diseases;

In cases where alternative causes of TMA have not been adequately excluded, additional information may be required from the prescriber to clarify the diagnosis before approval of the application.

Note The Authority application should be accompanied by a cover letter from the prescriber, providing complete details on:

- a) Presenting clinical features, including history, acute treatment and medications;
- b) Results of testing for genetic mutations (if available);
- c) Family history of aHUS, especially in first-degree relatives;
- d) Patient's prior history of episodes of active and progressing TMA caused by aHUS;
- e) Exclusion of alternative causes of TMA;
- f) History of renal or other organ transplant (if any);
- g) Any other matters considered relevant by the prescriber.

In cases where there are discordant results (for example, an equivocal biopsy result in the absence of objective evidence of haemolysis) the cover letter should articulate the prescriber's interpretation of the clinical data and how a diagnosis of aHUS is supported by the available evidence.

Note The Authority application should include the results of screening for genetic mutations known to confer a high risk of developing aHUS. The results of genetic screening should be provided whether or not a high-risk mutation has been identified.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Written applications for authority to prescribe must be submitted to Department of Human Services. Human Services will then contact the prescriber by telephone.

Authority required

Atypical haemolytic uraemic syndrome (aHUS)

Treatment Phase: Extended initial treatment - Assessment phase

Clinical criteria:

- Patient must have received treatment under the initial restriction with PBS subsidised eculizumab for this condition, **AND**
- Patient must have demonstrated on-going treatment response of PBS-subsidised eculizumab treatment for this condition, **AND**
- Patient must not have experienced treatment failure with eculizumab including PBS-subsidised eculizumab for this condition, **AND**
- Patient must not receive more than 56 weeks of treatment under this restriction.

Treatment criteria:

- Must be treated by a paediatric nephrologist, a nephrologist, a paediatric haematologist or a haematologist, or, must be in consultation with a paediatric nephrologist, a nephrologist, a paediatric haematologist or a haematologist.

A treatment response is defined as:

(1) Normalisation of haematology as demonstrated by at least 2 of the following: platelet count, haptoglobin, and LDH; **AND**

(2) One of the following:

a) An increase in eGFR of > 25% from baseline, where the baseline is the eGFR measurement immediately prior to commencing treatment with eculizumab or

b) an eGFR within +/- 25% from baseline; or

c) an avoidance of dialysis-dependence but worsening of kidney function with a reduction in eGFR 25% from baseline.

PBS-subsidised treatment with eculizumab will not be permitted if a patient has experienced treatment failure.

A treatment failure is defined as a patient who is:

(1) dialysis-dependent at the time of application and has failed to demonstrate significant resolution of extra-renal complications if originally presented; or

(2) on dialysis and has been on dialysis for 4 months of the previous 6 months while receiving PBS-subsidised eculizumab and has failed to demonstrate significant resolution of extra-renal complications if originally presented.

A maximum of up to 56 weeks of treatment is allowed under this restriction, however an application must be submitted at 6 months, 12 months, 18 months and 24 months following commencing PBS-subsidised eculizumab.

The authority application must include the following measures of response to the prior course of treatment, including serial haematological results (every 3 months while the patient is receiving treatment).

The authority application must be in writing and must include:

- (1) A completed authority prescription form; and
- (2) A completed aHUS eculizumab Authority Application Supporting Information Form for Extended Initial treatment; and
- (3) A detailed cover letter from the prescriber; and
- (4) A copy of a current Certificate of vaccination or a statement that vaccination has or will be administered and appropriate antibiotic prophylaxis has been prescribed; and
- (5) A measurement of body weight at the time of application; and
- (6) An identified genetic mutation, if applicable; and
- (7) A family history of aHUS, if applicable; and
- (8) A history of multiple episodes of aHUS before commencing eculizumab treatment, if applicable; and
- (9) A history of kidney transplant, if applicable, (especially if required due to aHUS); and
- (10) An inclusion of the individual consequences of recurrent disease, if applicable; and
- (11) Evidence that the patient has had a treatment response including haematological results of no more than 1 week old at the time of application (platelet count, haptoglobin and LDH); and an eGFR level of no more than 1 week old at the time of application; and
- (12) Evidence that the patient has not experienced treatment failure, including a supporting statement with clinical evidence that the patient does not require dialysis, unless the indication for continuing eculizumab is severe extra-renal complications that have significantly improved; and
- (13) If the indication for continuing eculizumab is severe extra-renal complications, then a supporting statement with clinical evidence that any initial extra-renal complications of TMA have significantly improved is required.

This assessment must be submitted no later than 4 weeks from the cessation of the prior treatment. Where a response assessment is not undertaken and submitted within these timeframes, the patient will be deemed to have failed to respond to treatment with eculizumab.

eculizumab 300 mg/30 mL injection, 30 mL vial

10525Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	6	..	5937.50	Soliris [XI]

■ ECULIZUMAB

Note At the time of authority application, medical practitioners must request the appropriate number of vials to provide sufficient drug for 4 weeks and up to 5 repeats, according to the specified dosage in the approved Product Information (PI). Applications for treatment with this drug where the dose and dosing frequency exceeds that specified in the approved PI will not be approved.

Note WARNING: Eculizumab increases the risk of meningococcal infections (septicaemia and/or meningitis). Please consult the approved PI for information about vaccination against meningococcal infection.

Note Eculizumab is not PBS subsidised to treat TMA caused by conditions other than aHUS, such as TMA occurring in the setting of, but not limited to:

- Active malignancy;
- Active HIV infection;
- Hematopoietic stem cell transplants;
- Various drugs including quinine, high-dose calcineurin inhibitors, antiplatelet agents;
- Certain chemotherapy drugs or immunosuppressant drugs associated with microangiopathic haemolytic anaemia/TMA;
- Active autoimmune diseases;

In cases where alternative causes of TMA have not been adequately excluded, additional information may be required from the prescriber to clarify the diagnosis before approval of the application.

Note The Authority application should be accompanied by a cover letter from the prescriber, providing complete details on:

- Presenting clinical features, including history, acute treatment and medications;
- Results of testing for genetic mutations (if available);
- Family history of aHUS, especially in first-degree relatives;
- Patient's prior history of episodes of active and progressing TMA caused by aHUS;
- Exclusion of alternative causes of TMA;
- History of renal or other organ transplant (if any);
- Any other matters considered relevant by the prescriber.

In cases where there are discordant results (for example, an equivocal biopsy result in the absence of objective evidence of haemolysis) the cover letter should articulate the prescriber's interpretation of the clinical data and how a diagnosis of aHUS is supported by the available evidence.

Note The Authority application should include the results of screening for genetic mutations known to confer a high risk of developing aHUS. The results of genetic screening should be provided whether or not a high-risk mutation has been identified.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Written applications for authority to prescribe must be submitted to Department of Human Services. Human Services will then contact the prescriber by telephone.

Authority required

Atypical haemolytic uraemic syndrome (aHUS)

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have received treatment under Extended Initial restriction with PBS subsidised eculizumab for this condition, **AND**
- Patient must have demonstrated on-going treatment response of PBS-subsidised eculizumab treatment for this condition, **AND**
- Patient must not have experienced treatment failure with eculizumab including PBS-subsidised eculizumab for this condition, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Treatment criteria:

- Must be treated by a paediatric nephrologist, a nephrologist, a paediatric haematologist or a haematologist, or, must be in consultation with a paediatric nephrologist, a nephrologist, a paediatric haematologist or a haematologist.

A treatment response is defined as:

- Normalisation of haematology as demonstrated by at least 2 of the following: platelet count, haptoglobin, and LDH; **AND**
- One of the following:
 - An increase in eGFR of > 25% from baseline, where the baseline is the eGFR measurement immediately prior to commencing treatment with eculizumab or
 - an eGFR within +/- 25% from baseline; or
 - an avoidance of dialysis-dependence but worsening of kidney function with a reduction in eGFR 25% from baseline.

PBS-subsidised treatment with eculizumab will not be permitted if a patient has experienced treatment failure.

A treatment failure is defined as a patient who is:

- dialysis-dependent at the time of application and has failed to demonstrate significant resolution of extra-renal complications if originally presented; or
- on dialysis and has been on dialysis for 4 months of the previous 6 months while receiving PBS-subsidised eculizumab and has failed to demonstrate significant resolution of extra-renal complications if originally presented.

The authority application must include the following measures of response to the prior course of treatment, including serial haematological results (every 3 months while the patient is receiving treatment).

The authority application must be in writing and must include:

- (1) A completed authority prescription form; and
- (2) A completed aHUS eculizumab Authority Application Supporting Information Form for Continuing treatment; and
- (3) A detailed cover letter from the prescriber; and
- (4) A copy of a current Certificate of vaccination or a statement that vaccination has or will be administered and appropriate antibiotic prophylaxis has been prescribed; and
- (5) A measurement of body weight at the time of application; and
- (6) An identified genetic mutation, if applicable; and
- (7) A family history of aHUS, if applicable; and
- (8) A history of multiple episodes of aHUS before recommencing eculizumab treatment, if applicable; and
- (9) A history of kidney transplant if applicable (especially if required due to aHUS); and
- (10) An inclusion of the individual consequences of recurrent disease, if applicable; and
- (11) Evidence that the patient has had a treatment response including haematological results of no more than 1 week old at the time of application (platelet count, haptoglobin and LDH); and an eGFR level of no more than 1 week old at the time of application; and
- (12) Evidence that the patient has not experienced treatment failure, including a supporting statement with clinical evidence that the patient does not require dialysis, unless the indication for continuing eculizumab is severe extra-renal complications that have significantly improved; and
- (13) If the indication for continuing eculizumab is severe extra-renal complications, then a supporting statement with clinical evidence that any initial extra-renal complications of TMA have significantly improved is required.

This assessment must be submitted no later than 4 weeks from the cessation of the prior treatment. Where a response assessment is not undertaken and submitted within these timeframes, the patient will be deemed to have failed to respond to treatment with eculizumab.

Authority required

Atypical haemolytic uraemic syndrome (aHUS)

Treatment Phase: Extended Continuing treatment

Clinical criteria:

- Patient must have received treatment under the Continuing treatment with PBS-subsidised eculizumab for this condition, **AND**
- Patient must have demonstrated on-going treatment response with PBS-subsidised eculizumab for this condition, **AND**
- Patient must not have ever experienced treatment failure with eculizumab including PBS-subsidised eculizumab for this condition, **AND**
- Patient must have a TMA-related cardiomyopathy as evidenced by left ventricular ejection fraction < 40% on current objective measurement; OR
- Patient must have severe TMA-related neurological impairment; OR
- Patient must have severe TMA-related gastrointestinal impairment; OR
- Patient must have severe TMA-related pulmonary impairment on current objective measurement; OR
- Patient must have grade 4 or 5 chronic kidney disease (eGFR of less than 30 mL/min); OR
- Patient must have a high risk of aHUS recurrence in the short term in the absence of continued treatment with eculizumab, **AND**
- Patient must not receive more than 24 weeks of treatment per continuing treatment course authorised under this restriction.

Treatment criteria:

- Must be treated by a paediatric nephrologist, a nephrologist, a paediatric haematologist or a haematologist, or, must be in consultation with a paediatric nephrologist, a nephrologist, a paediatric haematologist or a haematologist.

A treatment response is defined as:

- (1) Normalisation of haematology as demonstrated by at least 2 of the following: platelet count, haptoglobin, and LDH; AND
- (2) One of the following:
 - a) An increase in eGFR of > 25% from baseline, where the baseline is the eGFR measurement immediately prior to commencing treatment with eculizumab or
 - b) an eGFR within +/- 25% from baseline; or
 - c) an avoidance of dialysis-dependence but worsening of kidney function with a reduction in eGFR 25% from baseline.

PBS-subsidised treatment with eculizumab will not be permitted if a patient has experienced treatment failure. A treatment failure is defined as a patient who is:

- (1) dialysis-dependent at the time of application and has failed to demonstrate significant resolution of extra-renal complications if originally presented; or
- (2) on dialysis and has been on dialysis for 4 months of the previous 6 months while receiving PBS-subsidised eculizumab and has failed to demonstrate significant resolution of extra-renal complications if originally presented.

The authority application must include the following measures of response to the prior course of treatment, including serial haematological results (every 3 months while the patient is receiving treatment).

The authority application must be in writing and must include:

- (1) A completed authority prescription form; and
- (2) A completed aHUS eculizumab Authority Application Supporting Information Form for Continuing treatment; and
- (3) A detailed cover letter from the prescriber; and
- (4) A copy of a current Certificate of vaccination or a statement that vaccination has or will be administered and appropriate antibiotic prophylaxis has been prescribed; and

- (5) A measurement of body weight at the time of application; and
- (6) An identified genetic mutation, if applicable; and
- (7) A family history of aHUS, if applicable; and
- (8) A history of multiple episodes of aHUS before commencing eculizumab treatment, if applicable; and
- (9) A history of kidney transplant, if applicable (especially if required due to aHUS); and
- (10) An inclusion of the individual consequences of recurrent disease; and
- (11) A supporting statement with clinical evidence of severe TMA-related cardiomyopathy (including current LVEF result), neurological impairment, gastrointestinal impairment or pulmonary impairment; and
- (12) Evidence that the patient has had a treatment response including haematological results of no more than 1 month old at the time of application (platelet count, haptoglobin and LDH); and an eGFR level of no more than 1 month old at the time of application; and
- (13) Evidence that the patient has not experienced treatment failure, including a supporting statement with clinical evidence that the patient does not require dialysis, unless the indication for continuing eculizumab is severe extra-renal complications that have significantly improved; and
- (14) If the indication for continuing eculizumab is severe extra-renal complications, then a supporting statement with clinical evidence that any initial extra-renal complications of TMA have significantly improved is required.

This assessment must be submitted no later than 4 weeks from the cessation of the prior treatment. Where a response assessment is not undertaken and submitted within these timeframes, the patient will be deemed to have failed to respond to treatment with eculizumab.

Note All applications should be accompanied by a detailed letter that outlines the objective evidence of high risk of critical organ damage if aHUS recurs. The following evidence may be submitted to establish the patient's level of risk of aHUS recurrence in the short term in the absence of continued treatment with eculizumab:

- a) Evidence of a mutation known to confer a high risk of aHUS recurrence;
- b) Past history of recurrent episodes of active and progressive TMA due to aHUS, prior to the episode that led to current use of eculizumab;
- c) Past family history of aHUS recurrence, especially in first-degree relatives;
- d) Past history of recurrent aHUS following renal transplant for end-stage renal failure due to aHUS.

Authority required

Atypical haemolytic uraemic syndrome (aHUS)

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have demonstrated treatment response to previous treatment with PBS-subsidised eculizumab for this condition, **AND**
- Patient must not have ever experienced treatment failure with eculizumab including PBS-subsidised eculizumab for this condition, **AND**
- Patient must have the following clinical conditions:(i) either significant haemolysis as measured by low/absent haptoglobin; or presence of schistocytes on the blood film; or lactate dehydrogenase (LDH) above normal;AND(ii) either platelet consumption as measured by either 25% decline from patient baseline or thrombocytopenia (platelet count $<150 \times 10^9/L$);OR(iii) TMA-related organ impairment including on recent biopsy, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Treatment criteria:

- Must be treated by a paediatric nephrologist, a nephrologist, a paediatric haematologist or a haematologist, or, must be in consultation with a paediatric nephrologist, a nephrologist, a paediatric haematologist or a haematologist.

A treatment response is defined as:

- (1) Normalisation of haematology as demonstrated by at least 2 of the following: platelet count, haptoglobin, and LDH; AND
- (2) One of the following:

a) An increase in eGFR of $> 25\%$ from baseline, where the baseline is the eGFR measurement immediately prior to commencing treatment with eculizumab or

b) an eGFR within $\pm 25\%$ from baseline; or

c) an avoidance of dialysis-dependence but worsening of kidney function with a reduction in eGFR 25% from baseline.

PBS-subsidised treatment with eculizumab will not be permitted if a patient has experienced treatment failure. A treatment failure is defined as a patient who is:

- (1) dialysis-dependent at the time of application and has failed to demonstrate significant resolution of extra-renal complications if originally presented; or
- (2) on dialysis and has been on dialysis for 4 months of the previous 6 months while receiving PBS-subsidised eculizumab and has failed to demonstrate significant resolution of extra-renal complications if originally presented.

The authority application must include the following measures of response to the prior course of treatment, including serial haematological results (every 3 months while the patient is receiving treatment).

The authority application must be in writing and must include:

- (1) A completed authority prescription form(s); and
- (2) A completed aHUS eculizumab Authority Application Supporting Information Form for Recommencement of treatment; and
- (3) A signed patient acknowledgement or an acknowledgement signed by a parent or authorised guardian, if applicable; and
- (4) A detailed cover letter from the prescriber; and
- (5) A copy of a current Certificate of vaccination or a statement that vaccination has or will be administered and appropriate antibiotic prophylaxis has been prescribed; and
- (6) A measurement of body weight at the time of application, and
- (7) An identified genetic mutation, if applicable; and

- (8) A family history of aHUS if applicable; and
- (9) A history of multiple episodes of aHUS following the treatment break, if applicable; and
- (10) A history of kidney transplant if applicable (especially if required due to aHUS); and
- (11) An inclusion of the individual consequences of recurrent disease; and
- (12) A supporting statement with clinical evidence of TMA-related organ damage including current (within one week of application) haematological results (platelet count, haptoglobin and LDH), eGFR level, and, if applicable, on recent biopsy;
- (13) Evidence that the patient has had a treatment response to their previous treatment with eculizumab; and
- (14) Evidence that the patient has not experienced treatment failure, including a supporting statement with clinical evidence that the patient does not require dialysis, unless the indication for continuing eculizumab is severe extra-renal complications that have significantly improved; and
- (15) If the indication for continuing eculizumab is severe extra-renal complications, then a supporting statement with clinical evidence that any initial extra-renal complications of TMA have significantly improved is required.

This assessment must be submitted no later than 4 weeks from the cessation of the prior treatment. Where a response assessment is not undertaken and submitted within these timeframes, the patient will be deemed to have failed to respond to treatment with eculizumab.

Note A raise in LDH alone is not a sufficient reason to re-commence eculizumab, but thrombocytopenia with one marker of haemolysis (such as raised LDH, presence of schistocytes, or low/absence of haptoglobin) is an accepted reason to consider re-commencement of eculizumab treatment.

Note Kidney transplantation/dialysis is not a contraindication to recommencement of eculizumab treatment.

Authority required

Atypical haemolytic uraemic syndrome (aHUS)

Treatment Phase: Continuing recommencement of treatment

Clinical criteria:

- Patient must have received treatment under Recommencement of treatment restriction with PBS-subsidised eculizumab for this condition, **AND**
- Patient must have demonstrated ongoing treatment response to the previous 24 weeks of PBS-subsidised eculizumab for this condition, **AND**
- Patient must not have experienced treatment failure with eculizumab including PBS-subsidised eculizumab for this condition, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Treatment criteria:

- Must be treated by a paediatric nephrologist, a nephrologist, a paediatric haematologist or a haematologist, or, must be in consultation with a paediatric nephrologist, a nephrologist, a paediatric haematologist or a haematologist.

A treatment response is defined as:

- (1) Normalisation of haematology as demonstrated by at least 2 of the following: platelet count, haptoglobin, and LDH; AND
- (2) One of the following:

- a) An increase in eGFR of > 25% from baseline, where the baseline is the eGFR measurement immediately prior to commencing treatment with eculizumab or
- b) an eGFR within +/- 25% from baseline; or
- c) an avoidance of dialysis-dependence but worsening of kidney function with a reduction in eGFR 25% from baseline.

PBS-subsidised treatment with eculizumab will not be permitted if a patient has experienced treatment failure. A treatment failure is defined as a patient who is:

- (1) dialysis-dependent at the time of application and has failed to demonstrate significant resolution of extra-renal complications if originally presented; or
- (2) on dialysis and has been on dialysis for 4 months of the previous 6 months while receiving PBS-subsidised eculizumab and has failed to demonstrate significant resolution of extra-renal complications if originally presented.

The authority application must include the following measures of response to the prior course of treatment, including serial haematological results (every 3 months while the patient is receiving treatment).

The authority application must be in writing and must include:

- (1) A completed authority prescription form; and
- (2) A completed aHUS eculizumab Authority Application Supporting Information Form for Continuing treatment; and
- (3) A detailed cover letter from the prescriber; and
- (4) A copy of a current Certificate of vaccination or a statement that vaccination has or will be administered and appropriate antibiotic prophylaxis has been prescribed; and
- (5) A measurement of body weight at the time of application; and
- (6) An identified genetic mutation, if applicable; and
- (7) A family history of aHUS, if applicable; and
- (8) A history of multiple episodes of aHUS before recommencing eculizumab treatment, if applicable; and
- (9) A history of kidney transplant if applicable (especially if required due to aHUS); and
- (10) An inclusion of the individual consequences of recurrent disease, if applicable; and
- (11) Evidence that the patient has had a treatment response including haematological results of no more than 1 week old at the time of application (platelet count, haptoglobin and LDH); and an eGFR level of no more than 1 week old at the time of application; and
- (12) Evidence that the patient has not experienced treatment failure, including a supporting statement with clinical evidence that the patient does not require dialysis, unless the indication for continuing eculizumab is severe extra-renal complications that have significantly improved; and
- (13) If the indication for continuing eculizumab is severe extra-renal complications, then a supporting statement with clinical evidence that any initial extra-renal complications of TMA have significantly improved is required.

This assessment must be submitted no later than 4 weeks from the cessation of the prior treatment. Where a response assessment is not undertaken and submitted within these timeframes, the patient will be deemed to have failed to respond to treatment with eculizumab.

eculizumab 300 mg/30 mL injection, 30 mL vial

10183Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	5937.50	Soliris [XI]

■ EVEROLIMUS

Caution Careful monitoring of patients is mandatory.

Authority required (STREAMLINED)

5795

Management of renal allograft rejection

Treatment Phase: Management (initiation, stabilisation and review of therapy)

Clinical criteria:

- Patient must be receiving this drug for prophylaxis of renal allograft rejection, **AND**
- The treatment must be under the supervision and direction of a transplant unit.

Authority required (STREAMLINED)

5554

Management of cardiac allograft rejection

Treatment Phase: Management (initiation, stabilisation and review of therapy)

Clinical criteria:

- Patient must be receiving this drug for prophylaxis of cardiac allograft rejection, **AND**
- The treatment must be under the supervision and direction of a transplant unit.

everolimus 750 microgram tablet, 60

5740D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	4	5	..	*2301.12	Certican [NV]

everolimus 500 microgram tablet, 60

5739C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*767.04	Certican [NV]

everolimus 1 mg tablet, 60

5737Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	4	5	..	*3068.16	Certican [NV]

everolimus 250 microgram tablet, 60

5738B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*383.52	Certican [NV]

■ MYCOPHENOLATE

Caution Careful monitoring of patients is mandatory.

Authority required (STREAMLINED)

5795

Management of renal allograft rejection

Treatment Phase: Management (initiation, stabilisation and review of therapy)

Clinical criteria:

- Patient must be receiving this drug for prophylaxis of renal allograft rejection, **AND**
- The treatment must be under the supervision and direction of a transplant unit.

Authority required (STREAMLINED)

5554

Management of cardiac allograft rejection

Treatment Phase: Management (initiation, stabilisation and review of therapy)

Clinical criteria:

- Patient must be receiving this drug for prophylaxis of cardiac allograft rejection, **AND**
- The treatment must be under the supervision and direction of a transplant unit.

mycophenolate mofetil 1 g/5 mL powder for oral liquid, 165 mL

9500B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*489.02	CellCept [RO]

mycophenolate mofetil 500 mg tablet, 50

9502D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	6	5	..	*263.76	^a APO-Mycophenolate [TX] ^a Ceptolate [AF] ^a Mycophenolate AN [EA] ^a Pharmacor Mycophenolate 500 [CR]	^a CellCept [RO] ^a MycoCept [RF] ^a Mycophenolate Sandoz [SZ]

MYCOPHENOLATE

Caution Careful monitoring of patients is mandatory.

Note Management includes initiation, stabilisation and review of therapy as required.

Authority required (STREAMLINED)

4084

Prophylaxis of renal allograft rejection

Treatment Phase: Management

Clinical criteria:

- The treatment must be under the supervision and direction of a transplant unit.

Authority required (STREAMLINED)

4095

WHO Class III, IV or V lupus nephritis

Treatment Phase: Management

Clinical criteria:

- The condition must be proven by biopsy.

Treatment criteria:

- Must be treated by a nephrologist or in consultation with a nephrologist.
- The name of the consulting nephrologist must be included in the patient medical records.

mycophenolate 360 mg enteric tablet, 120

9504F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*343.12	Myfortic [NV]

mycophenolate 180 mg enteric tablet, 120

9503E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*171.58	Myfortic [NV]

MYCOPHENOLATE

Caution Careful monitoring of patients is mandatory.

Note For item codes 9501C and 1839T, pharmaceutical benefits that have the form capsule 250 mg are equivalent for the purposes of substitution

Authority required (STREAMLINED)

5653

Management of renal allograft rejection

Treatment Phase: Management (initiation, stabilisation and review of therapy)

Clinical criteria:

- Patient must be receiving this drug for prophylaxis of renal allograft rejection, **AND**
- The treatment must be under the supervision and direction of a transplant unit.

Authority required (STREAMLINED)

5600

Management of cardiac allograft rejection

Treatment Phase: Management (initiation, stabilisation and review of therapy)

Clinical criteria:

- Patient must be receiving this drug for prophylaxis of cardiac allograft rejection, **AND**
- The treatment must be under the supervision and direction of a transplant unit.

mycophenolate mofetil 250 mg capsule, 100

9501C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	6	5	..	*263.88	^a APO-Mycophenolate [TX] ^a Mycophenolate Sandoz [SZ]	^a CellCept [RO] ^a Pharmacor Mycophenolate 250 [CR]

mycophenolate mofetil 250 mg capsule, 50

1839T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	12	5	..	*263.88	^a Ceptolate [AF]

NATALIZUMAB

Caution Progressive multifocal leukoencephalopathy has been reported with this drug.

Authority required (STREAMLINED)

7697

Clinically definite relapsing-remitting multiple sclerosis

Treatment criteria:

- Must be treated by a neurologist.

Clinical criteria:

- The treatment must be a sole PBS-subsidised disease modifying therapy for this condition, **AND**
- Patient must be ambulatory (without assistance or support), **AND**
- Patient must have experienced at least 2 documented attacks of neurological dysfunction, believed to be due to multiple sclerosis, in the preceding 2 years of commencing a PBS-subsidised disease modifying therapy for this condition, **AND**

- The condition must be confirmed by magnetic resonance imaging of the brain and/or spinal cord; OR
- Patient must be deemed unsuitable for magnetic resonance imaging due to the risk of physical (not psychological) injury to the patient.

Population criteria:

- Patient must be aged 18 years or older.

The date of the magnetic resonance imaging scan must be included in the patient's medical notes, unless written certification is provided, in the patient's medical notes, by a radiologist that an MRI scan is contraindicated because of the risk of physical (not psychological) injury to the patient.

Treatment with this drug must cease if there is continuing progression of disability whilst the patient is being treated with this drug.

For continued treatment the patient must demonstrate compliance with, and an ability to tolerate, this drug.

Neurologists prescribing natalizumab under the PBS listing must be registered with the Tysabri Australian Prescribing Program.

natalizumab 300 mg/15 mL injection, 15 mL vial

9505G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	1340.68	Tysabri [BD]

■ OCRELIZUMAB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

7699

Multiple sclerosis

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be diagnosed as clinically definite relapsing-remitting multiple sclerosis by magnetic resonance imaging of the brain and/or spinal cord; OR
- The condition must be diagnosed as clinically definite relapsing-remitting multiple sclerosis by accompanying written certification provided by a radiologist that a magnetic resonance imaging scan is contraindicated because of the risk of physical (not psychological) injury to the patient, **AND**
- The treatment must be a sole PBS-subsidised disease modifying therapy for this condition, **AND**
- Patient must have experienced at least 2 documented attacks of neurological dysfunction, believed to be due to multiple sclerosis, in the preceding 2 years of commencing a PBS-subsidised disease modifying therapy for this condition, **AND**
- Patient must be ambulatory (without assistance or support).

Treatment criteria:

- Must be treated by a neurologist.

Where applicable, the date of the magnetic resonance imaging scan must be recorded in the patient's medical records.

Authority required (STREAMLINED)

7386

Multiple sclerosis

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not show continuing progression of disability while on treatment with this drug, **AND**
- The treatment must be a sole PBS-subsidised disease modifying therapy for this condition, **AND**
- Patient must have demonstrated compliance with, and an ability to tolerate this therapy.

Treatment criteria:

- Must be treated by a neurologist.

ocrelizumab 300 mg/10 mL injection, 10 mL vial

11242Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	..	..	*17533.00	Ocrevus [RO]

■ SIROLIMUS

Caution Careful monitoring of patients is mandatory.

Authority required (STREAMLINED)

5795

Management of renal allograft rejection

Treatment Phase: Management (initiation, stabilisation and review of therapy)

Clinical criteria:

- Patient must be receiving this drug for prophylaxis of renal allograft rejection, **AND**
- The treatment must be under the supervision and direction of a transplant unit.

sirolimus 2 mg tablet, 100

9548M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*2350.12	Rapamune [PF]

sirolimus 1 mg tablet, 100

9549N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*1175.04	Rapamune [PF]

sirolimus 1 mg/mL oral liquid, 60 mL

9550P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*924.02	Rapamune [PF]

sirolimus 500 microgram tablet, 100

9747B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*587.54	Rapamune [PF]

VEDOLIZUMAB
Note TREATMENT OF ADULT PATIENTS WITH MODERATE TO SEVERE ULCERATIVE COLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, golimumab, infliximab and vedolizumab for adult patients with ulcerative colitis. Patients are eligible for PBS-subsidised treatment with either adalimumab, golimumab, infliximab or vedolizumab at any one time.

Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, golimumab, infliximab and vedolizumab only.

From 1 June 2018, under the PBS, all adult patients will be able to commence a treatment cycle where they may trial PBS-subsidised adalimumab, golimumab, infliximab or vedolizumab without having to experience a disease flare when swapping to one of the alternate agents. Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with adalimumab, golimumab, infliximab or vedolizumab while they continue to show a response to therapy.

A patient who received PBS-subsidised adalimumab, infliximab, vedolizumab treatment prior to 1 June 2018 is considered to start their first cycle as of 1 June 2018. Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised adalimumab, golimumab, infliximab or vedolizumab more than once. Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised adalimumab, golimumab, infliximab or vedolizumab treatment in the most recent cycle to the date of the first application for initial treatment with adalimumab, golimumab, infliximab or vedolizumab under the new treatment cycle.

A patient who has failed fewer than 3 trials of either adalimumab, golimumab, infliximab or vedolizumab in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

(1) How to prescribe PBS-subsidised treatment with adalimumab, golimumab, infliximab and vedolizumab after 1 June 2018.

(a) Initial treatment. Applications for initial treatment should be made where:

(i) an adult patient has received no prior PBS-subsidised treatment with adalimumab, golimumab, infliximab or vedolizumab in this treatment cycle and wishes to commence such therapy (Initial 1 - New patient or recommencement of treatment after more than 5 years break in therapy); or

(ii) an adult patient has received prior PBS-subsidised (initial or continuing) adalimumab, golimumab, infliximab or vedolizumab therapy and wishes to trial an alternate agent (Initial 2 - Change or Re-commencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) an adult patient wishes to re-commence treatment with adalimumab, golimumab, infliximab or vedolizumab following a break in PBS-subsidised therapy with the same agent (Initial 2 - Change or Re-commencement of treatment after a break in therapy of less than 5 years).

Treatment authorisations under Initial 1 and Initial 2 will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, 14 weeks of therapy for golimumab, infliximab and vedolizumab.

A patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of treatment for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for golimumab, infliximab and vedolizumab. For second and subsequent courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment.

(b) Continuing treatment.

Following the completion of an initial treatment course with adalimumab, golimumab, infliximab or vedolizumab a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed in the month prior to completing their current course of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised treatment is approved, a patient may swap if eligible to the alternate adalimumab, golimumab, infliximab or vedolizumab treatment within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Mayo clinic score or partial Mayo clinic score), or the prior corticosteroid therapy and immunosuppressive therapy. A patient may trial an alternate treatment at any time, regardless of whether they are receiving therapy (initial or continuing) with adalimumab, golimumab, infliximab or vedolizumab at the time of the application. However, they cannot swap to a particular therapy if they have failed to respond to prior treatment with that drug once within the same treatment cycle. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(3) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised adalimumab,

golimumab, infliximab or vedolizumab therapy of at least 5 years, must requalify for initial 1 treatment with respect to the scores of disease severity. A patient must have received treatment with a 5-aminosalicylate oral preparation in a standard dose for induction of remission for a minimum of 3 consecutive months, and, either azathioprine or 6-mercaptopurine for a minimum of 3 consecutive months or a tapered course of oral steroids over a 6 week period followed by an appropriately dosed thiopurine agent for a minimum of 3 consecutive months (unless intolerance develops necessitating permanent treatment withdrawal to these agents). These above prior treatments must have been received immediately prior to the time the scores of disease severity being used to trial a second or subsequent course are measured.

(4) Patients 'grandfathered' onto PBS-subsidised treatment with golimumab.

A patient who commenced treatment with golimumab for moderate to severe ulcerative colitis prior to 1 June 2018 and who continues to receive treatment at the time of application, may qualify for treatment under the initial 3 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

Note Special Pricing Arrangements apply.

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Initial treatment (new patient or Recommencement of treatment after more than 5 years break in therapy - Initial 1)

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have failed to achieve an adequate response to a 5-aminosalicylate oral preparation in a standard dose for induction of remission for 3 or more consecutive months or have intolerance necessitating permanent treatment withdrawal, **AND**
- Patient must have failed to achieve an adequate response to azathioprine at a dose of at least 2 mg per kg daily for 3 or more consecutive months or have intolerance necessitating permanent treatment withdrawal; OR
- Patient must have failed to achieve an adequate response to 6-mercaptopurine at a dose of at least 1 mg per kg daily for 3 or more consecutive months or have intolerance necessitating permanent treatment withdrawal; OR
- Patient must have failed to achieve an adequate response to a tapered course of oral steroids, starting at a dose of at least 40 mg prednisolone (or equivalent), over a 6 week period or have intolerance necessitating permanent treatment withdrawal, and followed by a failure to achieve an adequate response to 3 or more consecutive months of treatment of an appropriately dosed thiopurine agent, **AND**
- Patient must have a Mayo clinic score greater than or equal to 6; OR
- Patient must have a partial Mayo clinic score greater than or equal to 6, provided the rectal bleeding and stool frequency subscores are both greater than or equal to 2 (endoscopy subscore is not required for a partial Mayo clinic score), **AND**
- Patient must be appropriately assessed for the risk of developing progressive multifocal leukoencephalopathy whilst on this treatment.

Population criteria:

- Patient must be aged 18 years or older.

Applications for authorisation of initial treatment must be in writing and must include:

(a) a completed authority prescription form; and

(b) a completed Ulcerative Colitis PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed current Mayo clinic or partial Mayo clinic calculation sheet including the date of assessment of the patient's condition; and

(ii) details of prior systemic drug therapy [dosage, date of commencement and duration of therapy]; and

(iii) the signed patient acknowledgement.

A maximum quantity and number of repeats to provide for an initial course of this drug consisting of one vial of 300 mg per dose, with one dose to be administered at weeks 0, 2 and 6, will be authorised.

All tests and assessments should be performed preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior conventional treatment.

The most recent Mayo clinic or partial Mayo clinic score must be no more than 1 month old at the time of application.

Patients who fail to achieve a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1 or have failed to maintain a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1 with continuing treatment with this drug, will not be eligible to receive further PBS-subsidised treatment with this drug.

A partial Mayo clinic assessment of the patient's response to this initial course of treatment must be made up to 12 weeks after the first dose for patients administered doses at weeks 0, 2 and 6 (6 weeks following the third dose) so that there is adequate time for a response to be demonstrated.

Patients must have signed a patient acknowledgement indicating they understand and acknowledge that the PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment.

If treatment with any of the above-mentioned drugs is contraindicated according to the relevant TGA-approved Product Information, details must be provided at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, details of this toxicity must be provided at the time of application.

Note Details of accepted toxicities including severity can be found on the Department of Human Services website at www.humanservices.gov.au.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au
Applications for authority to prescribe should be forwarded to:
Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have previously been issued with an authority prescription for this drug for this condition, **AND**
- Patient must have demonstrated or sustained an adequate response to treatment by having a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1 while receiving treatment with this drug, **AND**
- Patient must be appropriately assessed for the risk of developing progressive multifocal leukoencephalopathy whilst on this treatment.

Patients who have failed to maintain a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1 with continuing treatment with this drug, will not be eligible to receive further PBS-subsidised treatment with this drug.

Patients are eligible to receive continuing treatment with this drug in courses of up to 24 weeks providing they continue to sustain the response.

At the time of the authority application, medical practitioners should request the appropriate number of vials, to provide for a single infusion of 300 mg per dose.

Up to a maximum of 2 repeats will be authorised.

Note No applications for increased repeats will be authorised.

Note Authority approval for sufficient therapy to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Change or Re-commencement of treatment after a break in therapy of less than 5 years (Initial 2)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with adalimumab, golimumab, infliximab or vedolizumab for this condition in this treatment cycle, **AND**
- Patient must not have failed PBS-subsidised therapy with vedolizumab for this condition in the current treatment cycle, **AND**
- Patient must be appropriately assessed for the risk of developing progressive multifocal leukoencephalopathy whilst on this treatment.

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Population criteria:

- Patient must be aged 18 years or older.

To demonstrate a response to treatment the application must be accompanied by the results of the most recent course of this drug within the timelines specified in the relevant restriction. If the response assessment to the previous course of this drug is not submitted as detailed in the relevant restriction, the patient will be deemed to have failed therapy with this drug. Applications for authorisation of change or recommencement treatment must be in writing and must include:

(a) a completed authority prescription form; and

(b) a completed Ulcerative Colitis PBS Authority Application - Supporting Information Form which includes the following:

(i) Mayo clinical assessment (to demonstrate response to prior treatment).

A maximum quantity and number of repeats to provide for an initial course of this drug consisting of one vial of 300 mg per dose, with one dose to be administered at weeks 0, 2 and 6, will be authorised.

At the time of the authority application, medical practitioners should request the appropriate number of vials, to provide for a single infusion of 300 mg per dose.

Up to a maximum of 2 repeats will be authorised.

Authority approval for sufficient therapy to complete a maximum of 3 initial doses of treatment may be requested by telephone by contacting the Department of Human Services.

Note No applications for increased repeats will be authorised.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au
Applications for authority to prescribe should be forwarded to:
Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Balance of supply

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the Initial 1 (new patient) restriction to complete the 3 doses (i.e. the initial infusion regimen at 0, 2 and 6 weeks); OR
- Patient must have received insufficient therapy with this drug under the Initial 2 (Change or Recommencement of treatment after a break in therapy) restriction to complete the 3 doses (i.e. the initial infusion regimen at 0, 2 and 6 weeks); OR
- Patient must have received insufficient therapy with this drug under the Continuing treatment restriction to complete 24 weeks of treatment, **AND**
- The treatment must provide no more than the balance of up to 3 doses (Initial 1 and Initial 2 restrictions) or 2 repeats (Continuing restriction), **AND**
- Patient must be appropriately assessed for the risk of developing progressive multifocal leukoencephalopathy whilst on this treatment.

Population criteria:

- Patient must be aged 18 years or older.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Initial PBS-subsidised treatment (Grandfather patient)

Clinical criteria:

- Patient must have previously received non-PBS-subsidised therapy with this drug for this condition prior to 1 August 2015, **AND**
- Patient must have had a Mayo clinic score greater than or equal to 6 prior to commencing treatment with this drug; OR
- Patient must have had a partial Mayo clinic score greater than or equal to 6, provided the rectal bleeding and stool frequency subscores were both greater than or equal to 2 (endoscopy subscore is not required for a partial Mayo score) prior to commencing treatment with this drug; OR
- Patient must have a documented history of moderate to severe refractory ulcerative colitis prior to having commenced treatment with this drug where a Mayo clinic, partial Mayo clinic baseline assessment is not available, **AND**
- Patient must have demonstrated or sustained an adequate response to treatment by having a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1 while receiving treatment with this drug, **AND**
- Patient must be appropriately assessed for the risk of developing progressive multifocal leukoencephalopathy whilst on this treatment.

Population criteria:

- Patient must be 18 years of age or older.

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Applications for authorisation of initial treatment must be in writing and must include:

- a completed authority prescription form; and
- a completed Ulcerative Colitis PBS Authority Application - Supporting Information Form which includes the following:
 - the completed current and baseline Mayo clinic or partial Mayo clinic calculation sheet including the date of assessment of the patient's condition; and
 - the date of commencement of this drug; and
 - the signed patient acknowledgement.

The current Mayo clinic or partial Mayo clinic assessment must be no more than 1 month old at the time of application. The baseline assessment must be from immediately prior to commencing treatment with this drug. Where a baseline assessment is not available the prescriber must contact the Department of Human Services to discuss.

Patients are eligible to receive continuing treatment with this drug in courses of up to 24 weeks providing they continue to sustain the response.

At the time of the authority application, medical practitioners should request the appropriate number of vials, to provide for a single infusion of 300 mg per dose.

Up to a maximum of 2 repeats will be authorised.

A patient may qualify for PBS-subsidised treatment under this restriction once only.

For continuing PBS-subsidised treatment, a Grandfathered patient must qualify under the Continuing treatment criteria.

Note The patient must have signed a patient acknowledgement indicating they understand and acknowledge that the PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment.

Note No applications for increased repeats will be authorised.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

vedolizumab 300 mg injection, 1 vial

10384M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	3105.19	Entyvio [TK]

■ VEDOLIZUMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CROHN DISEASE

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for adult patients with severe Crohn disease. Where the term biological medicine appears in the following NOTES and restrictions, it refers to the tumour necrosis factor (TNF) alfa-antagonists (adalimumab and infliximab), the alpha-4 beta-7 integrin inhibitor (vedolizumab) and the human IgG1kappa monoclonal antibody (ustekinumab).

Patients are eligible for PBS-subsidised treatment with only 1 of the above PBS-subsidised biological medicines at any one time.

From 1 September 2017, under the PBS, all patients will be able to commence a treatment cycle where they may trial PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab without having to experience a disease flare when swapping to the alternate agent. Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with adalimumab, infliximab, vedolizumab or ustekinumab while they continue to show a response to therapy.

A patient who received PBS-subsidised adalimumab, infliximab, or vedolizumab treatment prior to 1 September 2017 is considered to have started their treatment cycle as of 1 September 2017.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab more than once.

Once a patient has either failed or ceased to respond to treatment for this condition 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy for this condition before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab treatment in the most recent cycle to the date of the first application for initial treatment with adalimumab, infliximab, vedolizumab or ustekinumab under the new treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of less than 5 years, may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab therapy after 1 September 2017.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised therapy with adalimumab, infliximab, vedolizumab or ustekinumab in this treatment cycle and wishes to commence such therapy (Initial treatment (new patient or Recommencement of treatment after more than 5 years break in therapy - Initial 1)); or

(ii) a patient has received prior PBS-subsidised (initial or continuing) adalimumab, infliximab, vedolizumab or ustekinumab and wishes to trial an alternate agent (Initial 2 - Change or recommencement) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to re-commence treatment with adalimumab, infliximab, vedolizumab or ustekinumab following a break in PBS-subsidised therapy with that agent (Change or Re-commencement of treatment after a break in therapy of less than 5 years (Initial 2)).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, 14 weeks of therapy for infliximab, 14 weeks of therapy for vedolizumab and 16 weeks for ustekinumab.

From 1 September 2017, a patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy for adalimumab or ustekinumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab or vedolizumab, and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

For subsequent courses of PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

Adalimumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats and the second

prescription should be written for 2 doses of 40 mg and 2 repeats for patients weighing 40 kg or greater. For patients weighing less than 40 kg, one prescription should be written for 2 doses of 40 mg with no repeats and the second prescription should be written for 2 doses of 20 mg with 3 repeats.

Ustekinumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be written under S100 (Highly Specialised Drugs) for a weight-based loading dose, containing a quantity of up to 4 vials of 130 mg and no repeats. The second prescription should be written under S85 (General) for the subsequent first dose, containing a quantity of 2 vials of 45 mg and no repeats.

(b) Continuing treatment.

Following the completion of an initial treatment course with adalimumab, infliximab, vedolizumab or ustekinumab, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted supply of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine therapy is approved, a patient may swap if eligible to the alternate adalimumab, infliximab, vedolizumab or ustekinumab within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Crohn Disease Activity Index (CDAI) Score, confirmation of Crohn disease), or the prior conventional therapies of corticosteroid therapy and immunosuppressive therapy.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with adalimumab, infliximab, vedolizumab or ustekinumab at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug once within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment, within the timeframes specified in the relevant restriction.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the CDAI or evidence of intestinal inflammation submitted with the first authority application for adalimumab, infliximab, vedolizumab or ustekinumab. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to assess response to all subsequent treatments.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to recommence treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity. Patients must have received treatment with a corticosteroid and at least 1 immunosuppressive agent, at an adequate dose, for a minimum of 3 consecutive months immediately prior to the time the CDAI score or the indices of intestinal inflammation are measured.

(5) Patients 'grandfathered' onto PBS-subsidised treatment with vedolizumab.

A patient who commenced treatment with vedolizumab for severe Crohn disease prior to 1 August 2015 and who continues to receive treatment at the time of application may qualify for treatment under the initial 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment will be assessed under the continuing treatment restriction of the relevant drug.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must requalify for continuing treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' above for further details.

(6) Patients 'grandfathered' onto PBS-subsidised treatment with ustekinumab.

A patient who commenced treatment with ustekinumab for severe Crohn disease prior to 1 September 2017 and who continues to receive treatment at the time of application may qualify for treatment under the initial 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment will be assessed under the continuing treatment restriction of the relevant drug.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must requalify for continuing treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' above for further details.

Note No applications for increased maximum quantities will be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Special Pricing Arrangements apply.

Authority required

Severe Crohn disease

Treatment Phase: Initial treatment (new patient - initial 1)

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have confirmed severe Crohn disease, defined by standard clinical, endoscopic and/or imaging features, including histological evidence, with the diagnosis confirmed by a gastroenterologist or a consultant physician, **AND**
- Patient must have failed to achieve an adequate response to prior systemic therapy with a tapered course of steroids, starting at a dose of at least 40 mg prednisolone (or equivalent), over a 6 week period, **AND**
- Patient must have failed to achieve adequate response to prior systemic immunosuppressive therapy with azathioprine at a dose of at least 2 mg per kg daily for 3 or more months; OR
- Patient must have failed to achieve adequate response to prior systemic immunosuppressive therapy with 6-mercaptopurine at a dose of at least 1 mg per kg daily for 3 or more months; OR
- Patient must have failed to achieve adequate response to prior systemic immunosuppressive therapy with methotrexate at a dose of at least 15 mg weekly for 3 or more months, **AND**
- Patient must be appropriately assessed for the risk of developing progressive multifocal leukoencephalopathy whilst on this treatment.

Population criteria:

- Patient must be aged 18 years or older.

Clinical criteria:

- Patient must have a Crohn Disease Activity Index (CDAI) Score greater than or equal to 300 as evidence of failure to achieve an adequate response to prior systemic therapy; OR
- Patient must have short gut syndrome with diagnostic imaging or surgical evidence, or have had an ileostomy or colostomy; and must have evidence of intestinal inflammation; and must have evidence of failure to achieve an adequate response to prior systemic therapy as specified below; OR
- Patient must have extensive intestinal inflammation affecting more than 50 cm of the small intestine as evidenced by radiological imaging; and must have a Crohn Disease Activity Index (CDAI) Score greater than or equal to 220; and must have evidence of failure to achieve an adequate response to prior systemic therapy as specified below.

Applications for authorisation must be made in writing and must include:

- a completed authority prescription form; and
- a completed Crohn Disease PBS Authority Application - Supporting Information Form which includes the following:
 - the completed current Crohn Disease Activity Index (CDAI) calculation sheet including the date of assessment of the patient's condition if relevant; and
 - details of prior systemic drug therapy [dosage, date of commencement and duration of therapy]; and
 - the reports and dates of the pathology or diagnostic imaging test(s) nominated as the response criterion, if relevant; and
 - the date of the most recent clinical assessment; and
 - the signed patient acknowledgement indicating they understand and acknowledge that the PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment.

Evidence of failure to achieve an adequate response to prior therapy must include at least one of the following: (a) patient must have evidence of intestinal inflammation; (b) patient must be assessed clinically as being in a high faecal output state; (c) patient must be assessed clinically as requiring surgery or total parenteral nutrition (TPN) as the next therapeutic option, in the absence of this drug, if affected by short gut syndrome, extensive small intestine disease or is an ostomy patient. Evidence of intestinal inflammation includes: (i) blood: higher than normal platelet count, or, an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour, or, a C-reactive protein (CRP) level greater than 15 mg per L; or (ii) faeces: higher than normal lactoferrin or calprotectin level; or (iii) diagnostic imaging: demonstration of increased uptake of intravenous contrast with thickening of the bowel wall or mesenteric lymphadenopathy or fat streaking in the mesentery;

All assessments, pathology tests and diagnostic imaging studies must be made within 1 month of the date of application and should be performed preferably whilst still on conventional treatment, but no longer than 1 month following cessation of the most recent prior treatment

If treatment with any of the specified prior conventional drugs is contraindicated according to the relevant TGA-approved Product Information, please provide details at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, details of this toxicity must be provided at the time of application.

Details of the accepted toxicities including severity can be found on the Department of Human Services website.

Any one of the baseline criteria may be used to determine response to an initial course of treatment and eligibility for continued therapy, according to the criteria included in the continuing treatment restriction. However, the same criterion must be used for any subsequent determination of response to treatment, for the purpose of eligibility for continuing PBS-subsidised therapy.

A maximum quantity and number of repeats to provide for an initial course of this drug consisting of one vial of 300 mg per dose, with one dose to be administered at weeks 0, 2 and 6, will be authorised.

If fewer than the maximum stated repeats in the relevant treatment phase are requested at the time of the application, authority approvals for sufficient repeats to complete the balance of the stated repeats in the relevant treatment phase may be requested by telephone by contacting the Department of Human Services and applying through the Balance of Supply restriction. Under no circumstances will telephone approvals be granted for treatment that would otherwise extend the relevant treatment phase.

The assessment of the patient's response to this initial course of treatment must be made up to 12 weeks after the first dose (6 weeks following the third dose) so that there is adequate time for a response to be demonstrated.

Note This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to the Department of Human Services no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with this drug.

Note It is recommended that an application for continuing treatment is submitted to the Department of Human Services at the time of the 12 week assessment, to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised treatment with this drug.

Authority required

Severe Crohn disease

Treatment Phase: Change or Re-commencement of treatment (initial 2)

Clinical criteria:

- Patient must have a documented history of severe Crohn disease, **AND**
- Patient must have received prior PBS-subsidised treatment with a biological disease modifying drug for this condition in this treatment cycle, **AND**
- Patient must not have failed PBS-subsidised therapy with this drug for this condition in the current treatment cycle, **AND**
- Patient must be appropriately assessed for the risk of developing progressive multifocal leukoencephalopathy whilst on this treatment.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form, which includes the following:
 - (i) the completed Crohn Disease Activity Index (CDAI) Score calculation sheet including the date of the assessment of the patient's condition, if relevant; or
 - (ii) the reports and dates of the pathology or diagnostic imaging test(s) used to assess response to therapy for patients with short gut syndrome, extensive small intestine disease or an ostomy, if relevant; and
 - (iii) the date of clinical assessment; and
 - (iv) the details of prior biological disease modifying drug treatment including the details of date and duration of treatment.

To demonstrate a response to treatment the application must be accompanied by the results of the most recent course of biological disease modifying drug (bDMD) therapy within the timeframes specified in the relevant restriction.

Where the most recent course of PBS-subsidised bDMD treatment was approved under an initial treatment restriction, the patient must have been assessed for response to that course following a minimum of 12 weeks of therapy for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab and vedolizumab and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

If the response assessment to the previous course of bDMD treatment is not submitted as detailed above, the patient will be deemed to have failed therapy with that particular course of bDMD.

A maximum quantity and number of repeats to provide for an initial course of this drug consisting of one vial of 300 mg per dose, with one dose to be administered at weeks 0, 2 and 6, will be authorised.

If fewer than the maximum stated repeats in the relevant treatment phase are requested at the time of the application, authority approvals for sufficient repeats to complete the balance of the stated repeats in the relevant treatment phase may be requested by telephone by contacting the Department of Human Services and applying through the Balance of Supply restriction. Under no circumstances will telephone approvals be granted for treatment that would otherwise extend the relevant treatment phase.

The assessment of the patient's response to this initial course of treatment must be made up to 12 weeks after the first dose (6 weeks following the third dose) so that there is adequate time for a response to be demonstrated.

This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to the Department of Human Services no later than 1 month from the date of completion of this initial course of treatment.

Where a response assessment is not undertaken and submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with this drug.

Note It is recommended that an application for continuing treatment is submitted to the Department of Human Services at the time of the 12 week assessment, to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised treatment with this drug.

Authority required

Severe Crohn disease

Treatment Phase: Initial PBS-subsidised treatment (Grandfather)

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have a documented history of severe Crohn disease, **AND**
- Patient must have previously received non-PBS-subsidised therapy with this drug for this condition prior to 1 August 2015.

Population criteria:

- Patient must be aged 18 years or older.

Clinical criteria:

- Patient must have had a Crohn Disease Activity Index (CDAI) Score of greater than or equal to 300 prior to commencing treatment with this drug; OR
- Patient must have a documented history of intestinal inflammation and have diagnostic imaging or surgical evidence of short gut syndrome if affected by the syndrome or has an ileostomy or colostomy; OR
- Patient must have a documented history and radiological evidence of intestinal inflammation if the patient has extensive small intestinal disease affecting more than 50 cm of the small intestine, **AND**
- Patient must have an adequate response to this drug defined as a reduction in Crohn Disease Activity Index (CDAI) Score to a level no greater than 150 if assessed by CDAI or if affected by extensive small intestine disease; OR
- Patient must have an adequate response to this drug defined as (a) an improvement of intestinal inflammation as demonstrated by: (i) blood: normalisation of the platelet count, or an erythrocyte sedimentation rate (ESR) level no greater than 25 mm per hour, or a C-reactive protein (CRP) level no greater than 15 mg per L; or (ii) faeces: normalisation of lactoferrin or calprotectin level; or (iii) evidence of mucosal healing, as demonstrated by diagnostic imaging findings, compared to the baseline assessment; or (b) reversal of high faecal output state; or (c) avoidance of the need for surgery or total parenteral nutrition (TPN), if affected by short gut syndrome, extensive small intestine or is an ostomy patient, **AND**
- Patient must be appropriately assessed for the risk of developing progressive multifocal leukoencephalopathy whilst on this treatment.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current Crohn Disease Activity Index (CDAI) calculation sheet including the date of assessment of the patient's condition if relevant; and
 - (ii) details of prior systemic drug therapy [dosage, date of commencement and duration of therapy]; and
 - (iii) the reports and dates of the pathology or diagnostic imaging test(s) nominated as the response criterion, if relevant; and
 - (iv) the date of the most recent clinical assessment; and
 - (v) the signed patient acknowledgement indicating they understand and acknowledge that the PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment.

The assessment of the patient's response to a continuing course of therapy must be made within the 4 weeks prior to completion of that course and posted to the Department of Human Services no less than 2 weeks prior to the date the next dose is scheduled, in order to ensure continuity of treatment for those patients who meet the continuation criterion.

Where an assessment is not submitted to the Department of Human Services within these timeframes, patients will be deemed to have failed to respond, or to have failed to sustain a response, to treatment with this drug.

Patients are eligible to receive continuing treatment with this drug in courses of up to 24 weeks providing they continue to sustain the response.

At the time of the authority application, medical practitioners should request the appropriate number of vials, to provide sufficient for a single infusion of 300 mg vedolizumab per dose. Up to a maximum of 2 repeats will be authorised.

If fewer than the maximum stated repeats in the relevant treatment phase are requested at the time of the application, authority approvals for sufficient repeats to complete the balance of the stated repeats in the relevant treatment phase may be requested by telephone by contacting the Department of Human Services and applying through the Balance of Supply restriction. Under no circumstances will telephone approvals be granted for treatment that would otherwise extend the relevant treatment phase.

A patient may qualify for PBS-subsidised treatment under this restriction once only.

Authority required

Severe Crohn disease

Treatment Phase: Balance of supply

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the Initial 1 (new patient) restriction to complete the 3 doses (i.e. the initial infusion regimen at 0, 2 and 6 weeks); OR
- Patient must have received insufficient therapy with this drug under the Continuing treatment restriction to complete 24 weeks of treatment; OR
- Patient must have received insufficient therapy with this drug to complete 24 weeks of treatment under the Initial PBS-subsidised treatment restriction for patients who had previously received non-PBS subsidised treatment (Grandfathered patient), **AND**
- The treatment must provide no more than the balance of up to 3 doses (new patients) or 2 repeats (Continuing or Grandfathered patients), **AND**
- Patient must be appropriately assessed for the risk of developing progressive multifocal leukoencephalopathy whilst on this treatment.

Population criteria:

- Patient must be aged 18 years or older.

Note Authority approval for sufficient therapy to complete a maximum of 3 initial doses or 2 repeats may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Severe Crohn disease

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have a documented history of severe Crohn disease, **AND**
- Patient must have previously been issued with an authority prescription for this drug for this condition, **AND**
- Patient must have demonstrated or sustained an adequate response to treatment with this drug.

Population criteria:

- Patient must be aged 18 years or older.

Clinical criteria:

- Patient must be appropriately assessed for the risk of developing progressive multifocal leukoencephalopathy whilst on this treatment, **AND**
- Patient must have an adequate response to this drug defined as a reduction in Crohn Disease Activity Index (CDAI) Score to a level no greater than 150 if assessed by CDAI or if affected by extensive small intestine disease; OR
- Patient must have an adequate response to this drug defined as (a) an improvement of intestinal inflammation as demonstrated by: (i) blood: normalisation of the platelet count, or an erythrocyte sedimentation rate (ESR) level no greater than 25 mm per hour, or a C-reactive protein (CRP) level no greater than 15 mg per L; or (ii) faeces: normalisation of lactoferrin or calprotectin level; or (iii) evidence of mucosal healing, as demonstrated by diagnostic imaging findings, compared to the baseline assessment; or (b) reversal of high faecal output state; or (c) avoidance of the need for surgery or total parenteral nutrition (TPN), if affected by short gut syndrome, extensive small intestine or is an ostomy patient.

Applications for authorisation must be made in writing and must include:

- a completed authority prescription form; and
- a completed Crohn Disease PBS Authority Application - Supporting Information Form which includes the following:
 - the completed Crohn Disease Activity Index (CDAI) Score calculation sheet including the date of the assessment of the patient's condition, if relevant; or
 - the reports and dates of the pathology test or diagnostic imaging test(s) used to assess response to therapy for patients with short gut syndrome, extensive small intestine disease or an ostomy, if relevant; and
 - the date of clinical assessment.

All assessments, pathology tests and diagnostic imaging studies, must be made within 1 month of the date of application.

If the application is the first application for continuing treatment with this drug, an assessment of the patient's response to the initial course of treatment must be made up to 12 weeks after the first dose so that there is adequate time for a response to be demonstrated.

The assessment of the patient's response to a continuing course of therapy must be made within the 4 weeks prior to completion of that course and posted to the Department of Human Services no less than 2 weeks prior to the date the next dose is scheduled, in order to ensure continuity of treatment for those patients who meet the continuation criterion.

Where an assessment is not submitted to the Department of Human Services within these timeframes, patients will be deemed to have failed to respond, or to have failed to sustain a response, to treatment with this drug.

Patients are eligible to receive continuing treatment with this drug in courses of up to 24 weeks providing they continue to sustain the response.

At the time of the authority application, medical practitioners should request the appropriate number of vials, to provide sufficient for a single infusion of 300 mg vedolizumab per dose. Up to a maximum of 2 repeats will be authorised.

If fewer than the maximum stated repeats in the relevant treatment phase are requested at the time of the application, authority approvals for sufficient repeats to complete the balance of the stated repeats in the relevant treatment phase may be requested by telephone by contacting the Department of Human Services and applying through the Balance of Supply restriction. Under no circumstances will telephone approvals be granted for treatment that would otherwise extend the relevant treatment phase.

vedolizumab 300 mg injection, 1 vial

10390W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	3105.19	Entyvio [TK]

Tumor necrosis factor alpha (TNF-) inhibitors

ADALIMUMAB

Note TREATMENT OF PATIENTS WITH SEVERE ACTIVE JUVENILE IDIOPATHIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, etanercept and tocilizumab for a patient who has severe active juvenile idiopathic arthritis. Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, etanercept and tocilizumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 3 biological medicines at any one time.

From 1 April 2014, a patient receiving PBS-subsidised biological medicine is considered to be in a treatment cycle where they may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements, within a single treatment cycle, a patient may:

- continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy; and
- fail to respond or to sustain a response to each PBS-subsidised biological medicine once only. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological

medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single treatment cycle and they must have, at a minimum, a 12 month break in PBS-subsidised biological medicine therapy before they are eligible to commence another cycle.

The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine was approved to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who received PBS-subsidised biological medicine immediately prior to 1 April 2014 is considered to be in their first cycle as of 1 April 2014. A patient who has had a break in biological medicine treatment of at least 12 months immediately prior to making a new application, on or after 1 April 2014, will commence a new treatment cycle under the Initial 3 treatment restriction.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of less than 12 months may commence a further course of treatment within the same treatment cycle under the Initial 2 treatment restriction.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of more than 12 months must commence a new treatment cycle under the Initial 3 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment therapy after 1 April 2014.

(1) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an alternate medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine therapy of less than 12 months) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 12 months with the same medicine (initial 2 - Change or Recommencement of treatment after a break in biological medicine therapy of less than 12 months).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 12 months (initial 3 - Recommencement of treatment after a break in biological medicine of more than 12 months) Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy.

A patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not conducted within these timeframes, the patient will be deemed to have failed this course of treatment.

For the second and subsequent cycles of PBS-subsidised biological medicine, it is recommended that a patient is reviewed in the 4 weeks prior to completing their current course of treatment and that an application is posted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services where applicable. Where a response assessment is not submitted, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (joint count) or the prior non-biological medicine therapy requirements, except if the patient has had a break in therapy of more than 12 months who would then need to requalify under the initial 3 restrictions with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application.

However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, an application for a patient who wishes to swap to an alternate biological medicine should be accompanied by the approved authority prescription or remaining repeats for the biological medicine the patient is ceasing.

(4) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count submitted with the first authority application for a biological medicine. However, prescribers may provide a new baseline measurement any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to the revised baseline measurement.

(5) Recommencement of treatment after a 12 months break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological medicine therapy of at least 12 months, must qualify under the initial 3 restriction and meet the relevant criteria and index of disease severity.

(6) Withdrawal of treatment after sustained remission.

Withdrawal of treatment with biological medicine should be considered in a patient who has achieved and sustained complete remission of disease for 12 months. A demonstration of response to the current treatment should be submitted to the Department of Human Services at the time treatment is ceased.

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Treatment criteria:

- Must be treated by a paediatric rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Clinical criteria:

- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have demonstrated severe intolerance of, or toxicity due to, methotrexate; OR
- Patient must have demonstrated failure to achieve an adequate response to 1 or more of the following treatment regimens: (i) oral or parenteral methotrexate at a dose of at least 20 mg per square metre weekly, alone or in combination with oral or intra-articular corticosteroids, for a minimum of 3 months; or (ii) oral methotrexate at a dose of at least 10 mg per square metre weekly together with at least 1 other disease modifying anti-rheumatic drug (DMARD), alone or in combination with corticosteroids, for a minimum of 3 months, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be under 18 years of age.

Severe intolerance to methotrexate is defined as intractable nausea and vomiting and general malaise unresponsive to manoeuvres, including reducing or omitting concomitant non-steroidal anti-inflammatory drugs (NSAIDs) on the day of methotrexate administration, use of folic acid supplementation, or administering the dose of methotrexate in 2 divided doses over 24 hours.

Toxicity due to methotrexate is defined as evidence of hepatotoxicity with repeated elevations of transaminases, bone marrow suppression temporally related to methotrexate use, pneumonitis, or serious sepsis.

If treatment with methotrexate alone or in combination with another DMARD is contraindicated according to the relevant TGA-approved Product Information, details must be provided at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, details of this toxicity must be provided at the time of application.

The following criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

- (a) an active joint count of at least 20 active (swollen and tender) joints; OR
- (b) at least 4 active joints from the following list:
 - (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - (ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The joint count assessment must be performed preferably whilst still on DMARD treatment, but no longer than 4 weeks following cessation of the most recent prior treatment.

The authority application must be made in writing and must include:

- (1) completed authority prescription form(s); and
- (2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

At the time of authority application, medical practitioners must request the appropriate number of injections of appropriate strength, based on the weight of the patient, to provide sufficient for two doses. Up to a maximum of 3 repeats will be authorised.

An assessment of a patient's response to an initial course of treatment must be conducted following a minimum of 12 weeks of therapy. An application for the continuing treatment must be accompanied with the assessment of response and submitted to the Department of Human Services no later than 4 weeks from the date of completion of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Use of alternative DMARDs in children is dependent on approval by the Therapeutic Goods Administration as age restrictions may apply.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs

Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 12 months)

Treatment criteria:

- Must be treated by a paediatric rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be under 18 years of age.

An adequate response to treatment is defined as:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

- (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
- (ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

At the time of authority application, medical practitioners must request the appropriate number of injections of appropriate strength, based on the weight of the patient, to provide sufficient for two doses. Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

- (1) completed authority prescription form(s); and
- (2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to change or recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient who fails to demonstrate a response to treatment with this drug under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug in this treatment cycle. A patient may re-trial this drug after a minimum of 12 months have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the initial 3 treatment restriction.

If a patient fails to respond to PBS-subsidised biological medicine treatment 3 times (once with each agent) they will not be eligible to receive further PBS-subsidised biological medicine therapy in this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 3 (recommencement of treatment after a break in biological medicine of more than 12 months)

Treatment criteria:

- Must be treated by a paediatric rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have had a break in treatment of 12 months or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must have either (a) a total active joint count of at least 20 active (swollen and tender) joints; or (b) at least 4 active major joints, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be under 18 years of age.

Active joints are defined as:

- (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
- (ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

All measures of joint count must be no more than 4 weeks old at the time of this application.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of active joints, the response must be demonstrated on the total number of active joints.

At the time of authority application, medical practitioners must request the appropriate number of injections of appropriate strength, based on the weight of the patient, to provide sufficient for two doses. Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

- (1) completed authority prescription form(s); and
- (2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient) or Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 12 months) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 12 months) - balance of supply

Treatment criteria:

- Must be treated by a paediatric rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 12 months) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 12 months) restriction to complete 16 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment per continuing treatment course authorised under this restriction.

An adequate response to treatment is defined as:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

Determination of whether a response has been demonstrated to initial and subsequent courses of treatment will be based on the baseline measurement of joint count submitted with the initial treatment application.

The authority application must be made in writing and must include:

(1) completed authority prescription form(s); and

(2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

At the time of authority application, medical practitioners must request the appropriate number of injections of appropriate strength, based on the weight of the patient, to provide sufficient for two doses. Up to a maximum of 5 repeats will be authorised.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either Initial 1, Initial 2, or Initial 3 treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 12 months have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

If a patient fails to respond to PBS-subsidised biological medicine treatment 3 times (once with each agent) they will not be eligible to receive further PBS-subsidised biological medicine therapy in this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Continuing treatment - balance of supply

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restriction.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL pen devices

9663N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	1151.87	Humira [VE]

adalimumab 20 mg/0.4 mL injection, 2 x 0.4 mL syringes

9661L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	1151.87	Humira [VE]

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL syringes

9662M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	1151.87	Humira [VE]

■ ETANERCEPT
Note TREATMENT OF PATIENTS WITH SEVERE ACTIVE JUVENILE IDIOPATHIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, etanercept and tocilizumab for a patient who has severe active juvenile idiopathic arthritis. Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, etanercept and tocilizumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 3 biological medicines at any one time.

From 1 April 2014, a patient receiving PBS-subsidised biological medicine is considered to be in a treatment cycle where they may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements, within a single treatment cycle, a patient may:

(i) continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy; and

(ii) fail to respond or to sustain a response to each PBS-subsidised biological medicine once only. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single treatment cycle and they must have, at a minimum, a 12 month break in PBS-subsidised biological medicine therapy before they are eligible to commence another cycle.

The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine was approved to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who received PBS-subsidised biological medicine immediately prior to 1 April 2014 is considered to be in their first cycle as of 1 April 2014. A patient who has had a break in biological medicine treatment of at least 12 months immediately prior to making a new application, on or after 1 April 2014, will commence a new treatment cycle under the Initial 3 treatment restriction.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of less than 12 months may commence a further course of treatment within the same treatment cycle under the Initial 2 treatment restriction.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of more than 12 months must commence a new treatment cycle under the Initial 3 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment therapy after 1 April 2014.

(1) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an alternate medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine therapy of less than 12 months) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 12 months with the same medicine (initial 2 - Change or Recommencement of treatment after a break in biological medicine therapy of less than 12 months).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 12 months (initial 3 - Recommencement of treatment after a break in biological medicine of more than 12 months) Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy.

A patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not conducted within these timeframes, the patient will be deemed to have failed this course of treatment.

For the second and subsequent cycles of PBS-subsidised biological medicine, it is recommended that a patient is reviewed in the 4 weeks prior to completing their current course of treatment and that an application is posted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services where applicable. Where a response assessment is not submitted, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (joint count) or the prior non-biological medicine therapy requirements, except if the patient has had a break in therapy of more than 12 months who would then need to requalify under the initial 3 restrictions with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application.

However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, an application for a patient who wishes to swap to an alternate biological medicine should be accompanied by the approved authority prescription or remaining repeats for the biological medicine the patient is ceasing.

(4) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count submitted with the first authority application for a biological medicine. However, prescribers may provide a new baseline measurement any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to the revised baseline measurement.

(5) Recommencement of treatment after a 12 months break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological medicine therapy of at least 12 months, must qualify under the initial 3 restriction and meet the relevant criteria and index of disease severity.

(6) Withdrawal of treatment after sustained remission.

Withdrawal of treatment with biological medicine should be considered in a patient who has achieved and sustained complete remission of disease for 12 months. A demonstration of response to the current treatment should be submitted to the Department of Human Services at the time treatment is ceased.

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Treatment criteria:

- Must be treated by a paediatric rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Clinical criteria:

- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have demonstrated severe intolerance of, or toxicity due to, methotrexate; OR
- Patient must have demonstrated failure to achieve an adequate response to 1 or more of the following treatment regimens: (i) oral or parenteral methotrexate at a dose of at least 20 mg per square metre weekly, alone or in combination with oral or intra-articular corticosteroids, for a minimum of 3 months; or (ii) oral methotrexate at a dose of at least 10 mg per square metre weekly together with at least 1 other disease modifying anti-rheumatic drug (DMARD), alone or in combination with corticosteroids, for a minimum of 3 months, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be under 18 years of age.

Severe intolerance to methotrexate is defined as intractable nausea and vomiting and general malaise unresponsive to manoeuvres, including reducing or omitting concomitant non-steroidal anti-inflammatory drugs (NSAIDs) on the day of methotrexate administration, use of folic acid supplementation, or administering the dose of methotrexate in 2 divided doses over 24 hours.

Toxicity due to methotrexate is defined as evidence of hepatotoxicity with repeated elevations of transaminases, bone marrow suppression temporally related to methotrexate use, pneumonitis, or serious sepsis.

If treatment with methotrexate alone or in combination with another DMARD is contraindicated according to the relevant TGA-approved Product Information, details must be provided at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, details of this toxicity must be provided at the time of application.

The following criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

- (a) an active joint count of at least 20 active (swollen and tender) joints; OR
- (b) at least 4 active joints from the following list:
 - (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - (ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The joint count assessment must be performed preferably whilst still on DMARD treatment, but no longer than 4 weeks following cessation of the most recent prior treatment.

The authority application must be made in writing and must include:

- (1) completed authority prescription form(s); and
- (2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

At the time of authority application, medical practitioners must request the appropriate number of injections to provide sufficient for four weeks of treatment. Up to a maximum of 3 repeats will be authorised.

An assessment of a patient's response to an initial course of treatment must be conducted following a minimum of 12 weeks of therapy. An application for the continuing treatment must be accompanied with the assessment of response and submitted to the Department of Human Services no later than 4 weeks from the date of completion of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Use of alternative DMARDs in children is dependent on approval by the Therapeutic Goods Administration as age restrictions may apply.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 12 months)

Treatment criteria:

- Must be treated by a paediatric rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be under 18 years of age.

An adequate response to treatment is defined as:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

At the time of authority application, medical practitioners must request the appropriate number of injections to provide sufficient for four weeks of treatment. Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

- (1) completed authority prescription form(s); and
- (2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to change or recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient who fails to demonstrate a response to treatment with this drug under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug in this treatment cycle. A patient may re-trial this drug after a minimum of 12 months have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the initial 3 treatment restriction.

If a patient fails to respond to PBS-subsidised biological medicine treatment 3 times (once with each agent) they will not be eligible to receive further PBS-subsidised biological medicine therapy in this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 3 (recommencement of treatment after a break in biological medicine of more than 12 months)

Treatment criteria:

- Must be treated by a paediatric rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have had a break in treatment of 12 months or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must have either (a) a total active joint count of at least 20 active (swollen and tender) joints; or (b) at least 4 active major joints, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be under 18 years of age.

Active joints are defined as:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

All measures of joint count must be no more than 4 weeks old at the time of this application.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of active joints, the response must be demonstrated on the total number of active joints.

At the time of authority application, medical practitioners must request the appropriate number of injections to provide sufficient for four weeks of treatment. Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

(1) completed authority prescription form(s); and

(2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient) or Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 12 months) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 12 months) - balance of supply

Treatment criteria:

- Must be treated by a paediatric rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 12 months) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 12 months) restriction to complete 16 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment per continuing treatment course authorised under this restriction.

An adequate response to treatment is defined as:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

Determination of whether a response has been demonstrated to initial and subsequent courses of treatment will be based on the baseline measurement of joint count submitted with the initial treatment application.

The authority application must be made in writing and must include:

(1) completed authority prescription form(s); and

(2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

At the time of authority application, medical practitioners must request the appropriate number of injections to provide sufficient for four weeks of treatment. Up to a maximum of 5 repeats will be authorised.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either Initial 1, Initial 2, or Initial 3 treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 12 months have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

If a patient fails to respond to PBS-subsidised biological medicine treatment 3 times (once with each agent) they will not be eligible to receive further PBS-subsidised biological medicine therapy in this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au
Applications for authority to prescribe should be forwarded to:
Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis
Treatment Phase: Continuing treatment - balance of supply

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restriction.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

etanercept 50 mg/mL injection, 4 x 1 mL syringes

5733R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	939.25	Enbrel [PF]

etanercept 50 mg/mL injection, 4 x 1 mL pen devices

5735W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	939.25	Enbrel [PF]

etanercept 25 mg injection [4 vials] (&) inert substance diluent [4 x 1 mL syringes], 1 pack

5734T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	469.63	Enbrel [PF]

■ INFLIXIMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CROHN DISEASE

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for adult patients with severe Crohn disease. Where the term biological medicine appears in the following NOTES and restrictions, it refers to the tumour necrosis factor (TNF) alpha-antagonists (adalimumab and infliximab), the alpha-4 beta-7 integrin inhibitor (vedolizumab) and the human IgG1kappa monoclonal antibody (ustekinumab).

Patients are eligible for PBS-subsidised treatment with only 1 of the above PBS-subsidised biological medicines at any one time.

From 1 September 2017, under the PBS, all patients will be able to commence a treatment cycle where they may trial PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab without having to experience a disease flare when swapping to the alternate agent. Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with adalimumab, infliximab, vedolizumab or ustekinumab while they continue to show a response to therapy.

A patient who received PBS-subsidised adalimumab, infliximab, or vedolizumab treatment prior to 1 September 2017 is considered to have started their treatment cycle as of 1 September 2017.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab more than once.

Once a patient has either failed or ceased to respond to treatment for this condition 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy for this condition before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab treatment in the most recent cycle to the date of the first application for initial treatment with adalimumab, infliximab, vedolizumab or ustekinumab under the new treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of less than 5 years, may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab therapy after 1 September 2017.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised therapy with adalimumab, infliximab, vedolizumab or ustekinumab in this treatment cycle and wishes to commence such therapy (Initial treatment (new patient or Recommencement of treatment after more than 5 years break in therapy - Initial 1)); or
- (ii) a patient has received prior PBS-subsidised (initial or continuing) adalimumab, infliximab, vedolizumab or ustekinumab and wishes to trial an alternate agent (Initial 2 - Change or recommencement) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to re-commence treatment with adalimumab, infliximab, vedolizumab or ustekinumab following a break in PBS-subsidised therapy with that agent (Change or Re-commencement of treatment after a break in therapy of less than 5 years (Initial 2)).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, 14 weeks of therapy for infliximab, 14 weeks of therapy for vedolizumab and 16 weeks for ustekinumab.

From 1 September 2017, a patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy for adalimumab or ustekinumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab or vedolizumab, and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

For subsequent courses of PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

Adalimumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats and the second prescription should be written for 2 doses of 40 mg and 2 repeats for patients weighing 40 kg or greater. For patients weighing less than 40 kg, one prescription should be written for 2 doses of 40 mg with no repeats and the second prescription should be written for 2 doses of 20 mg with 3 repeats.

Ustekinumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be written under S100 (Highly Specialised Drugs) for a weight-based loading dose, containing a quantity of up to 4 vials of 130 mg and no repeats. The second prescription should be written under S85 (General) for the subsequent first dose, containing a quantity of 2 vials of 45 mg and no repeats.

(b) Continuing treatment.

Following the completion of an initial treatment course with adalimumab, infliximab, vedolizumab or ustekinumab, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted supply of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine therapy is approved, a patient may swap if eligible to the alternate adalimumab, infliximab, vedolizumab or ustekinumab within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Crohn Disease Activity Index (CDAI) Score, confirmation of Crohn disease), or the prior conventional therapies of corticosteroid therapy and immunosuppressive therapy.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with adalimumab, infliximab, vedolizumab or ustekinumab at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug once within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment, within the timeframes specified in the relevant restriction.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the CDAI or evidence of intestinal inflammation submitted with the first authority application for adalimumab, infliximab, vedolizumab or ustekinumab. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to assess response to all subsequent treatments.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to recommence treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity. Patients must have received treatment with a corticosteroid and at least 1 immunosuppressive agent, at an adequate dose, for a minimum of 3 consecutive months immediately prior to the time the CDAI score or the indices of intestinal inflammation are measured.

(5) Patients 'grandfathered' onto PBS-subsidised treatment with vedolizumab.

A patient who commenced treatment with vedolizumab for severe Crohn disease prior to 1 August 2015 and who continues to receive treatment at the time of application may qualify for treatment under the initial 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment will be assessed under the continuing treatment restriction of the relevant drug.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must requalify for continuing treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' above for further details.

(6) Patients 'grandfathered' onto PBS-subsidised treatment with ustekinumab.

A patient who commenced treatment with ustekinumab for severe Crohn disease prior to 1 September 2017 and who continues to receive treatment at the time of application may qualify for treatment under the initial 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment will be assessed under the continuing treatment restriction of the relevant drug.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must requalify for continuing treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' above for further details.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe Crohn disease

Treatment Phase: Subsequent continuing treatment

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must have an adequate response to this drug defined as a reduction in Crohn Disease Activity Index (CDAI) Score to a level no greater than 150 if assessed by CDAI or if affected by extensive small intestine disease; OR
- Patient must have an adequate response to this drug defined as (a) an improvement of intestinal inflammation as demonstrated by: (i) blood: normalisation of the platelet count, or an erythrocyte sedimentation rate (ESR) level no greater than 25 mm per hour, or a C-reactive protein (CRP) level no greater than 15 mg per L; or (ii) faeces: normalisation of lactoferrin or calprotectin level; or (iii) evidence of mucosal healing, as demonstrated by diagnostic imaging findings, compared to the baseline assessment; or (b) reversal of high faecal output state; or (c) avoidance of the need for surgery or total parenteral nutrition (TPN), if affected by short gut syndrome, extensive small intestine or is an ostomy patient.

Population criteria:

- Patient must be aged 18 years or older.

Applications for authorisation must be made in writing and must include: (a) a completed authority prescription form; and (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form which includes the following: (i) the completed Crohn Disease Activity Index (CDAI) Score; or (ii) the reports and dates of the pathology test or diagnostic imaging test(s) used to assess response to therapy for patients with short gut syndrome, extensive small intestine disease or an ostomy, if relevant; and (iii) the date of the most recent clinical assessment.

Each application for subsequent continuing treatment with this drug must include an assessment of the patient's response to the prior course of therapy. If the response assessment is not provided at the time of application the patient will be deemed to have failed this course of treatment.

Patients are eligible to receive continuing treatment with this drug in courses of up to 24 weeks providing they continue to sustain the response.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg eight weekly. Up to a maximum of 2 repeats will be authorised.

infliximab 100 mg injection, 1 vial

11389K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	320.71	Remicade [JC]

INFLIXIMAB

Note TREATMENT OF COMPLEX REFRACTORY FISTULISING CROHN DISEASE

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for patients with complex refractory fistulising Crohn disease. Where the term "biological medicine" appears in the following NOTES and restrictions, it refers to adalimumab and infliximab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the PBS- subsidised biological medicines for this condition at any one time.

From 1 April 2011, under the PBS, all patients will be able to commence a treatment cycle where they may trial PBS-subsidised adalimumab or infliximab without having to experience a disease flare when swapping to the alternate agent. Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with adalimumab or infliximab while they continue to show a response to therapy.

A patient who received PBS-subsidised adalimumab or infliximab treatment prior to 1 April 2011 is considered to have started their treatment cycle as of 1 April 2011.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised adalimumab or infliximab more than twice.

Once a patient has either failed or ceased to respond to treatment for this condition 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy for this condition before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised adalimumab or infliximab treatment in the most recent cycle to the date of the first application for initial treatment with adalimumab or infliximab under the new treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of less than 5 years, may commence a further course of treatment within the same treatment cycle.
A patient who has failed 3 trials or fewer of biological medicine therapy in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised adalimumab or infliximab therapy after 1 April 2011.

(a) Initial treatment. Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised adalimumab or infliximab therapy in this treatment cycle and wishes to commence such therapy (Initial treatment (new patient or Recommencement of treatment after more than 5 years break in therapy - Initial 1); or
 - (ii) a patient has received prior PBS-subsidised (initial or continuing) adalimumab or infliximab therapy and wishes to trial an alternate agent (Initial 2 - Change or recommencement) [further details are under 'Swapping therapy' below]; or
 - (iii) a patient wishes to re-commence treatment with adalimumab or infliximab following a break in PBS-subsidised therapy with that agent (Change or Re-commencement of treatment after a break in therapy of less than 5 years (Initial 2)
- Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy for adalimumab and 14 weeks of therapy for infliximab.

From 1 April 2011, a patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab, and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

For subsequent courses of PBS-subsidised adalimumab or infliximab treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

Adalimumab only: Two completed authority prescriptions must be submitted with every initial application for adalimumab. One prescription must be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats. The second prescription must be written for 2 doses of 40 mg and 2 repeats.

(b) Continuing treatment.

Adalimumab patients:

Following the completion of an initial treatment course with adalimumab, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

Infliximab patients:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment under the Initial 1 or Initial 2 treatment restrictions.

For the second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted supply of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap if eligible to the alternate biological medicine within the same treatment cycle.

A patient may trial the alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with adalimumab or infliximab at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug two times within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment within the timeframes specified in the relevant restriction.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements submitted with the first authority application for adalimumab or infliximab. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to these revised baseline measurements.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to recommence treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Complex refractory Fistulising Crohn disease

Treatment Phase: Subsequent continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug.

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

An adequate response is defined as:

- (a) a decrease from baseline in the number of open draining fistulae of greater than or equal to 50%; and/or
- (b) a marked reduction in drainage of all fistula(e) from baseline, together with less pain and induration as reported by the patient.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Fistulising Crohn Disease PBS Authority Application - Supporting Information Form which includes a completed Fistula Assessment form including the date of the assessment of the patient's condition.

The most recent fistula assessment must be no more than 1 month old at the time of application.

Each application for subsequent continuing treatment with this drug must include an assessment of the patient's response to the prior course of therapy. If the response assessment is not provided at the time of application the patient will be deemed to have failed this course of treatment.

Patients are eligible to receive continuing treatment with this drug in courses of up to 24 weeks providing they continue to sustain the response.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg eight weekly. Up to a maximum of 2 repeats will be authorised.

infliximab 100 mg injection, 1 vial

11424G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	320.71	Remicade [JC]

■ INFLIXIMAB
Note TREATMENT OF PAEDIATRIC PATIENTS WITH REFRACTORY CROHN DISEASE

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) for paediatric patients with adalimumab for severe refractory Crohn disease and infliximab for moderate to severe refractory Crohn disease. Where the term "biological medicines" appears in the following NOTES and restrictions, it refers to adalimumab and infliximab only. A patient is eligible for PBS-subsidised treatment with only one PBS-subsidised biological medicine at any one time. For paediatric patients with Crohn disease, infliximab is PBS-subsidised for moderate to severe disease while adalimumab is PBS-subsidised for severe disease.

From 1 August 2015, under the PBS, patients commencing on adalimumab will be able to commence a treatment cycle where they may trial each PBS-subsidised biological medicine without having to experience a disease flare when swapping to infliximab. Patients on infliximab will be able to commence a treatment cycle where they may trial each PBS-subsidised biological medicine but will need to meet a PCDAl score of greater than or equal to 40 when swapping to adalimumab. Under these arrangements, within a single treatment cycle and depending on the disease severity, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy.

A patient who received PBS-subsidised biological medicine treatment prior to 1 August 2015 is considered to have started their treatment cycle as of 1 August 2015.

Within the same treatment cycle, a paediatric patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than twice.

Once a patient has either failed, or ceased to respond to treatment for this condition 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy for this condition before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle.

A patient who has failed 3 trials or fewer of biological medicine therapy in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 August 2015.

(a) Initial treatment.

Applications for initial treatment should be made where:

- i) a patient has received no prior PBS-subsidised biological medicine therapy in this treatment cycle and wishes to commence such therapy - Initial 1 treatment (new patient or Recommencement of treatment after more than 5 years break in therapy); or
- (ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent - Initial 2 treatment (Change or Re-commencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or
- (iii) a patient wishes to re-commence treatment with a specific biological medicine following a break in PBS-subsidised therapy with that agent - Initial 2 treatment (Change or Re-commencement of treatment after a break in therapy of less than 5 years).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy for adalimumab and 14 weeks of therapy for infliximab.

From 1 August 2015, a patient must be assessed for response to any course of initial PBS-subsidised biological therapy following a minimum of 12 weeks of therapy for adalimumab and up to 12 weeks after the first dose (6 weeks following the

third dose) for infliximab, and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

For first and subsequent continuing courses of PBS-subsidised biological medicine therapy, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

Adalimumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats and the second prescription should be written for 2 doses of 40 mg and 2 repeats for patients weighing 40 kg or greater. For patients weighing less than 40 kg, one prescription should be written for 2 doses of 40 mg with no repeats and the second prescription should be written for 2 doses of 20 mg with 3 repeats.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient remains eligible to receive up to 24 weeks per course of continuing treatment under the First continuing treatment and Subsequent Continuing treatment restrictions with that drug providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted supply of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine therapy is approved, a patient with severe disease may swap if eligible to the alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Paediatric Crohn Disease Activity Index (PCDAI) Score, confirmation of Crohn disease), or the prior conventional therapies of corticosteroid therapy, immunosuppressive therapy or enteral nutrition. Patients on infliximab may swap to adalimumab within the same treatment cycle provided that their disease severity has progressed to severe disease (i.e. they have a current PCDAI score of 40 or more).

A patient cannot swap to a biological medicine if they have failed to respond to prior treatment with that drug twice within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment within the timeframes specified in the relevant restriction.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the PCDAI submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to assess response to all subsequent treatments.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to recommence treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity.

Patients must have failed to achieve an adequate response to 2 of the following 3 conventional prior therapies including: (i) a tapered course of steroids, starting at a dose of at least 1 mg per kg or 40 mg (whichever is the lesser) prednisolone (or equivalent), over a 6 week period; (ii) an 8 week course of enteral nutrition; or (iii) immunosuppressive therapy including azathioprine at a dose of at least 2 mg per kg daily for 3 or more months, or, 6-mercaptopurine at a dose of at least 1 mg per kg daily for 3 or more months, or, methotrexate at a dose of at least 10 mg per square metre weekly for 3 or more months immediately prior to the time the PCDAI score is measured.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Moderate to severe Crohn disease

Treatment Phase: Subsequent continuing treatment

Treatment criteria:

- Must be treated by a gastroenterologist (code 87) or a consultant physician [internal medicine specialising in gastroenterology (code 81)] or a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician or a specialist paediatric gastroenterologist.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First Continuing treatment restriction; AND
- Patient must have a reduction in PCDAI Score by at least 15 points from baseline value; AND
- Patient must have a total PCDAI score of 30 points or less.

Population criteria:

- Patient must be aged 6 to 17 years inclusive.

Application for authorisation must be made in writing and must include:

(a) a completed authority prescription form; and

(b) a completed Paediatric Crohn Disease PBS Authority Application - Supporting Information Form, which includes the completed Paediatric Crohn Disease Activity Index (PCDAI) calculation sheet along with the date of the assessment of the patient's condition.

Patients are only eligible to receive subsequent continuing PBS-subsidised treatment with this drug in courses of up to 24 weeks at a dose of 5 mg per kg per dose providing they continue to sustain the response.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg eight weekly. Up to a maximum of 2 repeats will be authorised.

infliximab 100 mg injection, 1 vial

11448M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	320.71	Remicade [JC]

■ INFLIXIMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for adults with severe active rheumatoid arthritis. Where the term biological medicine appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alfa antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab), the T-cell co-stimulation modulator (abatacept) and the Janus kinase (JAK) inhibitors (baricitinib, tofacitinib). A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised biological medicine therapy may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once, and
- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised biological medicines for the treatment of rheumatoid arthritis.

Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Stevens Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

For a patient who has failed PBS-subsidised treatment with 2 or 3 TNF-alfa antagonists prior to 1 August 2010 please contact the Department of Human Services on 1800 700 270. A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction.

A patient who has failed fewer than 5 biological medicines and who has a break in treatment of less than 24 months may commence a further course of treatment with a biological medicine under Initial 2 treatment restriction. A patient who has failed fewer than 5 biological medicines and who has had a break in therapy of longer than 24 months may commence a further course of treatment with a biological medicine under the Initial 3 treatment restriction. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine treatment is stopped to the date of the new application for treatment with a biological medicine.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 April 2019.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised biological medicine treatment and wishes to commence such therapy, excluding rituximab (Initial 1 - new patient); or
- (ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months) [further details are under 'Swapping therapy' below]; or
- (iii) a patient wishes to re-commence treatment with a specific biological medicine following a break of less than 24 months in PBS-subsidised therapy with that agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months).
- (iv) a patient wishes to re-commence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 24 months (Initial 3 - re-commencement of treatment after a break in biological medicine of more than 24 months) Initial applications for a new patient (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6-month intensive DMARD trial, but prior to ceasing DMARD therapy.

Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, baricitinib, etanercept, golimumab, tocilizumab and tofacitinib, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

Rituximab patients should be assessed following a minimum of 12 weeks after the first infusion, and the assessment should be submitted to the Department of Human Services within 4 weeks to ensure continuity of treatment for those who meet the continuing restriction for PBS subsidised treatment with this drug for this condition.

Where a response assessment is not provided the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment. For second and subsequent courses of PBS-subsidised biological medicine (excluding rituximab) treatment it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to the Department of Human Services within sufficient timeframe to allow for authority assessment.

Abatacept patients:

A patient is eligible to receive one I.V. loading dose when commencing treatment with the subcutaneous formulation. Two prescriptions are required, the first prescription for the I.V. loading dose for sufficient vials for one dose based on the patient's weight with no repeats. The second prescription for the subcutaneous formulation, with a maximum quantity of 4 and up to 3 repeats, must be submitted with the initial application.

Rituximab patients:

A further application may be submitted to the Department of Human Services within sufficient timeframe to allow authority assessment. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

Infliximab and etanercept patients:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction. Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

(2) Swapping therapy

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non- biological medicine therapy requirements except if the patient has had a break in therapy of more than 24 months who would need to requalify with respect to the indices of disease severity. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each biological medicine trialled.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent, unless they have a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug.

Abatacept:

A patient swapping from I.V. abatacept to subcutaneous abatacept will not be eligible for an I.V. loading dose when commencing treatment with the subcutaneous formulation.

Rituximab:

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF-alfa antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological medicine during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological medicine therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate biological medicine may do so without having to have any treatment-free period.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and the Department of Human Services will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints.

Applications under the Initial 1 treatment restriction for a new patient must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy.

Applications under the Initial 3 treatment restriction for re-commencement of treatment after a break in biological medicine of more than 24 months must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:
Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Subsequent continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment per subsequent continuing treatment course authorised under this restriction, **AND**
- The treatment must be given concomitantly with methotrexate at a dose of at least 7.5 mg weekly.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials to provide sufficient drug, based on the weight of the patient, for a single infusion at a dose of 3 mg per kg.

Up to a maximum of 2 repeats will be authorised.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient has either failed or ceased to respond to a PBS-subsidised biological medicine for this condition 5 times, they will not be eligible to receive further PBS-subsidised treatment with a biological medicine for this condition.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

infliximab 100 mg injection, 1 vial

11481G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	320.71	Remicade [JC]

INFLIXIMAB

Note Authority applications for increased quantities/ repeats (where relevant) may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

4524

Acute severe ulcerative colitis

Treatment criteria:

- Must be treated by a gastroenterologist; OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology, or general medicine specialising in gastroenterology].

Clinical criteria:

- Patient must have received an infusion of infliximab for the treatment of this condition as a hospital inpatient no more than two weeks prior to the date of the authority application, **AND**
- Patient must be an adult aged 18 years or older, and prior to initiation of infliximab treatment in hospital must have been experiencing six or more bloody stools per day, plus at least one of the following: (i) Temperature greater than 37.8 degrees Celsius; (ii) Pulse rate greater than 90 beats per minute; (iii) Haemoglobin less than 105 g/L; (iv) Erythrocyte sedimentation rate greater than 30 mm/h; OR
- Patient must be a child aged 6 to 17 years inclusive, and prior to initiation of infliximab treatment in hospital must have had a Paediatric Ulcerative Colitis Activity Index (PUCAI) greater than or equal to 65, with the diagnosis confirmed by a gastroenterologist, or a consultant physician as specified below, **AND**
- Patient must have failed to achieve an adequate response to at least 72 hours treatment with intravenous corticosteroids prior to initiation of infliximab treatment in hospital.

Population criteria:

- Patient must be 6 years of age or older.

For adults aged 18 years or older, failure to achieve an adequate response to intravenous corticosteroid treatment is defined by the Oxford criteria where:

(i) If assessed on day 3, patients pass 8 or more stools per day or 3 or more stools per day with a C-reactive protein (CRP) greater than 45 mg/L

(ii) If assessed on day 7, patients pass 3 or more stools per day with visible blood.

For children aged 6 to 17 years, failure to achieve an adequate response to intravenous corticosteroids means a PUCAI score greater than 45 at 72 hours.

At the time of authority application, prescribers should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single infusion at a dose of 5 mg per kg.

Before administering infliximab to a child aged 6 to 17 years, the treating clinician must have consulted with a paediatric gastroenterologist or with an institution experienced in performance of paediatric colectomy. The name of the specialist or institution must be included in the patient's medical records.

Evidence that the patient meets the PBS restriction criteria must be recorded in the patient's medical records.

infliximab 100 mg injection, 1 vial

10067W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	5	1	..	*1603.55	^a Inflectra [PF] ^a Renflexis [MK]	^a Remicade [JC]

■ INFLIXIMAB

Note TREATMENT OF ADULT PATIENTS WITH ANKYLOSING SPONDYLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, certolizumab pegol, etanercept, golimumab, infliximab and secukinumab for adult patients with ankylosing spondylitis. Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab and secukinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 6 biological medicines at any 1 time.

Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last prescription for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised biological medicine treatment with adalimumab, certolizumab pegol, etanercept, golimumab, infliximab and secukinumab.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised biological medicine treatment in this treatment cycle and wishes to commence such therapy (Initial 1 - New patient)

(ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years); or

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

A patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy.

(b) Continuing treatment.

For the first continuing treatment course of PBS-subsidised biological medicine, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1, Initial 2 or Initial 3 treatment restrictions. For second and subsequent continuing courses of PBS-subsidised biological medicine it is recommended that a patient be

reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply. Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be conducted within 1 month of the last dose. Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI), or the prior NSAID therapy and exercise program requirements.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the BASDAI, ESR and/or CRP submitted with the first authority application for a biological medicine.

However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to these revised baseline measurements.

For a new patient, the BASDAI used to determine the baseline must be measured while the patient is receiving NSAID therapy and completing their exercise program.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response.

(4) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must qualify under the Initial 3 treatment restriction. The same clinical criteria and indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI) as for the Initial 1 (New patient) restriction will need to be met, but a re-trial of NSAID therapy and exercise therapy is not required.

Note Authority applications for increased quantities/ repeats (where relevant) may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required (STREAMLINED)

9481

Ankylosing spondylitis

Treatment Phase: Subsequent continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response is defined as an improvement from baseline of at least 2 of the BASDAI and 1 of the following:

- (a) an ESR measurement no greater than 25 mm per hour; or
- (b) a CRP measurement no greater than 10 mg per L; or
- (c) an ESR or CRP measurement reduced by at least 20% from baseline.

Where only 1 acute phase reactant measurement is supplied in the first application for PBS-subsidised treatment, that same marker must be used to determine response for all subsequent continuing treatments.

The measurement of response to the prior course of therapy must be documented in the patient's medical notes.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

infliximab 100 mg injection, 1 vial

11486M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	5	3	..	*1603.55	^a Inflectra [PF]	^a Renflexis [MK]

■ INFLIXIMAB
Note TREATMENT OF ADULT PATIENTS WITH MODERATE TO SEVERE ULCERATIVE COLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, golimumab, infliximab and vedolizumab for adult patients with ulcerative colitis. Patients are eligible for PBS-subsidised treatment with either adalimumab, golimumab, infliximab or vedolizumab at any one time.

Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, golimumab, infliximab and vedolizumab only.

From 1 June 2018, under the PBS, all adult patients will be able to commence a treatment cycle where they may trial PBS-subsidised adalimumab, golimumab, infliximab or vedolizumab without having to experience a disease flare when swapping to one of the alternate agents. Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with adalimumab, golimumab, infliximab or vedolizumab while they continue to show a response to therapy.

A patient who received PBS-subsidised adalimumab, infliximab, vedolizumab treatment prior to 1 June 2018 is considered to start their first cycle as of 1 June 2018. Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised adalimumab, golimumab, infliximab or vedolizumab more than once. Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised adalimumab, golimumab, infliximab or vedolizumab treatment in the most recent cycle to the date of the first application for initial treatment with adalimumab, golimumab, infliximab or vedolizumab under the new treatment cycle.

A patient who has failed fewer than 3 trials of either adalimumab, golimumab, infliximab or vedolizumab in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

(1) How to prescribe PBS-subsidised treatment with adalimumab, golimumab, infliximab and vedolizumab after 1 June 2018.

(a) Initial treatment. Applications for initial treatment should be made where:

(i) an adult patient has received no prior PBS-subsidised treatment with adalimumab, golimumab, infliximab or vedolizumab in this treatment cycle and wishes to commence such therapy (Initial 1 - New patient or recommencement of treatment after more than 5 years break in therapy); or

(ii) an adult patient has received prior PBS-subsidised (initial or continuing) adalimumab, golimumab, infliximab or vedolizumab therapy and wishes to trial an alternate agent (Initial 2 - Change or Re-commencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) an adult patient wishes to re-commence treatment with adalimumab, golimumab, infliximab or vedolizumab following a break in PBS-subsidised therapy with the same agent (Initial 2 - Change or Re-commencement of treatment after a break in therapy of less than 5 years).

Treatment authorisations under Initial 1 and Initial 2 will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, 14 weeks of therapy for golimumab, infliximab and vedolizumab.

A patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of treatment for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for golimumab, infliximab and vedolizumab. For second and subsequent courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment.

(b) Continuing treatment.

Following the completion of an initial treatment course with adalimumab, golimumab, infliximab or vedolizumab a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed in the month prior to completing their current course of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised treatment is approved, a patient may swap if eligible to the alternate adalimumab, golimumab, infliximab or vedolizumab treatment within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Mayo clinic score or partial Mayo clinic score), or the prior corticosteroid therapy and immunosuppressive therapy. A patient may trial an alternate treatment at any time, regardless of whether they are receiving therapy (initial or continuing) with adalimumab, golimumab, infliximab or vedolizumab at the time of the application. However, they cannot swap to a particular therapy if they have failed to respond to prior treatment with that drug once within the same treatment cycle. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(3) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised adalimumab, golimumab, infliximab or vedolizumab therapy of at least 5 years, must requalify for initial 1 treatment with respect to the scores of disease severity. A patient must have received treatment with a 5-aminosalicylate oral preparation in a standard dose for induction of remission for a minimum of 3 consecutive months, and, either azathioprine or 6-mercaptopurine for a minimum of 3 consecutive months or a tapered course of oral steroids over a 6 week period followed by an appropriately dosed thiopurine agent for a minimum of 3 consecutive months (unless intolerance develops necessitating permanent treatment withdrawal to these agents). These above prior treatments must have been received immediately prior to the time the scores of disease severity being used to trial a second or subsequent course are measured.

(4) Patients 'grandfathered' onto PBS-subsidised treatment with golimumab.

A patient who commenced treatment with golimumab for moderate to severe ulcerative colitis prior to 1 June 2018 and who continues to receive treatment at the time of application, may qualify for treatment under the initial 3 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment

must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

Note TREATMENT OF PAEDIATRIC PATIENTS WITH MODERATE TO SEVERE ULCERATIVE COLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) for paediatric patients with infliximab or adalimumab for moderate to severe ulcerative colitis; and infliximab for acute severe ulcerative colitis. Where the term 'biological medicine' appears in the following NOTES and restrictions, it refers to infliximab and adalimumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 2 biological medicines at any one time. Infliximab and adalimumab are PBS-subsidised for moderate to severe disease while only infliximab is PBS-subsidised for acute severe disease.

From 1 June 2017, under the PBS, all paediatric patients will be able to commence a treatment cycle where they may trial each PBS-subsidised biological medicine without having to experience a disease flare when swapping to the alternate agent. Under these arrangements, within a single treatment cycle and depending on the disease severity, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy. A patient who received PBS-subsidised biological medicine treatment prior to 1 June 2017 is considered to have started their treatment cycle as of 1 June 2017. Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than twice. Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle. A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle. There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 June 2017.

(a) Initial treatment. Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised biological medicine treatment in this treatment cycle and wishes to commence such therapy - Initial 1 treatment (new patient or Re-commencement of treatment after more than 5 years break in therapy); or
- (ii) a patient has received prior PBS-subsidised (initial or continuing) treatment with a biological medicine and wishes to trial an alternate agent - Initial 2 treatment (Change or Re-commencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping treatment' below]; or
- (iii) a patient wishes to re-commence treatment with a specific biological medicine following a break in PBS-subsidised therapy with that agent - Initial 2 treatment (Change or Re-commencement of treatment after a break in therapy of less than 5 years).

Treatment authorisations under Initial 1 and Initial 2 treatment will be limited to provide for a maximum of 16 weeks of treatment for adalimumab and 14 weeks of treatment for infliximab. From 1 June 2017, a patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of treatment for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab. For second and subsequent courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment.

Adalimumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. For patients weighing 40 kg or greater, one prescription should be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats and the second prescription should be written for 2 doses of 40 mg and 2 repeats. For patients weighing less than 40 kg, one prescription should be written for 2 doses of 40 mg with no repeats and the second prescription should be written for 2 doses of 20 mg with 3 repeats.

(b) Continuing treatment. Following the completion of an initial treatment course with a specific biological medicine, a patient remains eligible to receive up to 24 weeks per course of continuing treatment with that drug under the continuing treatment restriction providing they continue to sustain the response. It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure PBS subsidy criteria are met.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap if eligible to the alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Paediatric Ulcerative Colitis Activity Index (PUCAI) Score, confirmation of ulcerative colitis disease), or the prior conventional therapies of corticosteroids or immunosuppressives. A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving treatment (initial or continuing) with infliximab or adalimumab at the time of the application. However, a patient cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug twice within the same treatment cycle. To ensure a patient receives the maximum treatment opportunities allowed under these swapping arrangements, it is important that they are assessed for response to every course of treatment.

(3) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to recommence treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity. A patient must have received treatment with a 5-aminosalicylate oral preparation in a standard dose for induction of remission for a minimum of 3 consecutive months, and, either azathioprine or 6-mercaptopurine for a minimum of 3 consecutive months or a tapered course of oral steroids over a 6 week period followed by an appropriately dosed thiopurine agent for a minimum of 3 consecutive months (unless intolerance develops necessitating permanent treatment withdrawal to these agents) immediately prior to the time the PUCAI score is measured.

Note Authority applications for continuing treatment may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician; OR
- Must be treated by a specialist paediatric gastroenterologist.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have demonstrated or sustained an adequate response to treatment by having a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1 while receiving treatment with this drug; OR
- Patient must have demonstrated or sustained an adequate response to treatment by having a Paediatric Ulcerative Colitis Activity Index (PUCAI) score of less than 10 while receiving treatment with this drug, if aged 6 to 17 years.

Population criteria:

- Patient must be 6 years of age or older.

Patients who have failed to maintain a partial Mayo clinic score of less than or equal to 2, with no subscore greater than 1, or, patients who have failed to maintain a Paediatric Ulcerative Colitis Activity Index (PUCAI) score of less than 10 (if aged 6 to 17 years) with continuing treatment with this drug, will not be eligible to receive further PBS-subsidised treatment with this drug.

Patients are only eligible to receive continuing PBS-subsidised treatment with this drug in courses of up to 24 weeks at a dose of 5 mg per kg per dose providing they continue to sustain the response.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg eight weekly. Up to a maximum of 2 repeats will be authorised.

infliximab 100 mg injection, 1 vial

11459D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	320.71	^a Inflectra [PF] ^a Renflexis [MK]	^a Remicade [JC]

INFLIXIMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE PSORIATIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab for adult patients with severe active psoriatic arthritis. Therefore, where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient receiving PBS-subsidised treatment for psoriatic arthritis is able to commence a treatment cycle where they may trial biological medicines without having to experience a disease flare when swapping to the alternate biological medicine.

Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

A patient who received PBS-subsidised adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab or ustekinumab treatment prior to 1 May 2019 is considered to start their first cycle as of 1 May 2019.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment.

Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence another cycle [further details are under '(5) Recommencement of treatment after a 5-year break in PBS-subsidised therapy' below].

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe active psoriatic arthritis.

(1) Initial treatment.

Applications for initial treatment should be made where:

- a patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy (Initial 1 - New patient); or
- a patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an alternate medicine (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years)

[further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years) or An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, golimumab secukinumab and tofacitinib, 18 to 20 weeks of therapy for certolizumab pegol (depending upon the dosing regimen), 20 weeks of therapy for ixekizumab, 22 weeks of therapy for infliximab, and 28 weeks of therapy for ustekinumab. It is recommended that a patient be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of PBS-subsidised initial treatment following a minimum of 12 weeks of therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

Grandfather patients (ixekizumab only).

A patient who commenced treatment with ixekizumab for severe psoriatic arthritis prior to 1 March 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

Grandfather patients (tofacitinib only).

A patient who commenced treatment with Tofacitinib for severe psoriatic arthritis prior to 1 May 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment under the Initial 1 or Initial 2 treatment restrictions.

For the second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to re-qualify with respect to either the indices of disease severity (i.e. erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP) level, and active joint count) or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify under the Initial 3 treatment restriction with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application.

However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

Within a treatment cycle a patient may alternate between therapy with any biological medicine of their choice (1 at a time) providing:

- (i) they have not received PBS-subsidised treatment with that particular biological medicine previously; or
- (ii) they have demonstrated an adequate response to that particular biological medicine if they have previously trialled it on the PBS; and
- (iii) they have not previously failed to respond to treatment 3 times in this treatment cycle with that biological medicine.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(4) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the indices of disease severity submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial or change or

recommencement treatment application is submitted within a treatment cycle and these revised baseline measurements will be used to assess response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to determine response for all subsequent continuing treatments. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. 20 or more active joints), response will be determined according to a reduction in the total number of active joints. (5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment restriction according to the criteria of the relevant restriction and index of disease severity. The application must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Subsequent continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment per subsequent continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an erythrocyte sedimentation rate (ESR) no greater than 25 mm per hour or a C-reactive protein (CRP) level no greater than 15 mg per L or either marker reduced by at least 20% from baseline; and

either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following major active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to determine response for all subsequent continuing treatments.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg eight weekly.

Up to a maximum of 2 repeats will be authorised.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Severe Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

Where the most recent course of PBS-subsidised treatment with this drug was approved under the first continuing treatment restriction, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

infliximab 100 mg injection, 1 vial

11497D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	320.71	Remicade [JC]

INFLIXIMAB
Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for adults with severe active rheumatoid arthritis. Where the term biological medicine appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alfa antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab), the T-cell co-stimulation modulator (abatacept) and the Janus kinase (JAK) inhibitors (baricitinib, tofacitinib). A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised biological medicine therapy may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once, and
- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised biological medicines for the treatment of rheumatoid arthritis.

Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Stevens Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

For a patient who has failed PBS-subsidised treatment with 2 or 3 TNF-alfa antagonists prior to 1 August 2010 please contact the Department of Human Services on 1800 700 270. A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction.

A patient who has failed fewer than 5 biological medicines and who has a break in treatment of less than 24 months may commence a further course of treatment with a biological medicine under Initial 2 treatment restriction. A patient who has failed fewer than 5 biological medicines and who has had a break in therapy of longer than 24 months may commence a further course of treatment with a biological medicine under the Initial 3 treatment restriction. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine treatment is stopped to the date of the new application for treatment with a biological medicine.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 April 2019.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised biological medicine treatment and wishes to commence such therapy, excluding rituximab (Initial 1 - new patient); or
 - (ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months) [further details are under 'Swapping therapy' below]; or
 - (iii) a patient wishes to re-commence treatment with a specific biological medicine following a break of less than 24 months in PBS-subsidised therapy with that agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months).
 - (iv) a patient wishes to re-commence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 24 months (Initial 3 - re-commencement of treatment after a break in biological medicine of more than 24 months)
- Initial applications for a new patient (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6-month intensive DMARD trial, but prior to ceasing DMARD therapy.

Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, baricitinib, etanercept, golimumab, tocilizumab and tofacitinib, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

Rituximab patients should be assessed following a minimum of 12 weeks after the first infusion, and the assessment should be submitted to the Department of Human Services within 4 weeks to ensure continuity of treatment for those who meet the continuing restriction for PBS subsidised treatment with this drug for this condition.

Where a response assessment is not provided the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment. For second and subsequent courses of PBS-subsidised biological medicine (excluding rituximab) treatment it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to the Department of Human Services within sufficient timeframe to allow for authority assessment.

Abatacept patients:

A patient is eligible to receive one I.V. loading dose when commencing treatment with the subcutaneous formulation. Two prescriptions are required, the first prescription for the I.V. loading dose for sufficient vials for one dose based on the patient's weight with no repeats. The second prescription for the subcutaneous formulation, with a maximum quantity of 4 and up to 3 repeats, must be submitted with the initial application.

Rituximab patients:

A further application may be submitted to the Department of Human Services within sufficient timeframe to allow authority assessment. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate

response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

Infliximab and etanercept patients:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction. Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

(2) Swapping therapy

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non- biological medicine therapy requirements except if the patient has had a break in therapy of more than 24 months who would need to requalify with respect to the indices of disease severity. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each biological medicine trialled.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent, unless they have a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug.

Abatacept:

A patient swapping from I.V. abatacept to subcutaneous abatacept will not be eligible for an I.V. loading dose when commencing treatment with the subcutaneous formulation.

Rituximab:

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF-alfa antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological medicine during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological medicine therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate biological medicine may do so without having to have any treatment-free period.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and the Department of Human Services will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints.

Applications under the Initial 1 treatment restriction for a new patient must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy.

Applications under the Initial 3 treatment restriction for re-commencement of treatment after a break in biological medicine of more than 24 months must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note Authority applications for increased quantities/ repeats (where relevant) may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

8755

Severe active rheumatoid arthritis

Treatment Phase: Subsequent continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have demonstrated an adequate response to treatment with this drug, **AND**

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must not receive more than 24 weeks of treatment per subsequent continuing treatment course authorised under this restriction, **AND**
- The treatment must be given concomitantly with methotrexate at a dose of at least 7.5 mg weekly.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The measurement of response to the prior course of therapy must be documented in the patient's medical notes.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

If a patient has either failed or ceased to respond to a PBS-subsidised biological medicine for this condition 5 times, they will not be eligible to receive further PBS-subsidised treatment with a biological medicine for this condition.

infliximab 100 mg injection, 1 vial

11490R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	3	2	..	*962.13	^a Inflectra [PF]	^a Renflexis [MK]

■ INFLIXIMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CROHN DISEASE

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for adult patients with severe Crohn disease. Where the term biological medicine appears in the following NOTES and restrictions, it refers to the tumour necrosis factor (TNF) alpha-antagonists (adalimumab and infliximab), the alpha-4 beta-7 integrin inhibitor (vedolizumab) and the human IgG1kappa monoclonal antibody (ustekinumab).

Patients are eligible for PBS-subsidised treatment with only 1 of the above PBS-subsidised biological medicines at any one time.

From 1 September 2017, under the PBS, all patients will be able to commence a treatment cycle where they may trial PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab without having to experience a disease flare when swapping to the alternate agent. Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with adalimumab, infliximab, vedolizumab or ustekinumab while they continue to show a response to therapy.

A patient who received PBS-subsidised adalimumab, infliximab, or vedolizumab treatment prior to 1 September 2017 is considered to have started their treatment cycle as of 1 September 2017.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab more than once.

Once a patient has either failed or ceased to respond to treatment for this condition 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy for this condition before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab treatment in the most recent cycle to the date of the first application for initial treatment with adalimumab, infliximab, vedolizumab or ustekinumab under the new treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of less than 5 years, may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab therapy after 1 September 2017.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised therapy with adalimumab, infliximab, vedolizumab or ustekinumab in this treatment cycle and wishes to commence such therapy (Initial treatment (new patient or Recommencement of treatment after more than 5 years break in therapy - Initial 1)); or

(ii) a patient has received prior PBS-subsidised (initial or continuing) adalimumab, infliximab, vedolizumab or ustekinumab and wishes to trial an alternate agent (Initial 2 - Change or recommencement) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to re-commence treatment with adalimumab, infliximab, vedolizumab or ustekinumab following a break in PBS-subsidised therapy with that agent (Change or Re-commencement of treatment after a break in therapy of less than 5 years (Initial 2)).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, 14 weeks of

therapy for infliximab, 14 weeks of therapy for vedolizumab and 16 weeks for ustekinumab.

From 1 September 2017, a patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy for adalimumab or ustekinumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab or vedolizumab, and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

For subsequent courses of PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

Adalimumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats and the second prescription should be written for 2 doses of 40 mg and 2 repeats for patients weighing 40 kg or greater. For patients weighing less than 40 kg, one prescription should be written for 2 doses of 40 mg with no repeats and the second prescription should be written for 2 doses of 20 mg with 3 repeats.

Ustekinumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be written under S100 (Highly Specialised Drugs) for a weight-based loading dose, containing a quantity of up to 4 vials of 130 mg and no repeats. The second prescription should be written under S85 (General) for the subsequent first dose, containing a quantity of 2 vials of 45 mg and no repeats.

(b) Continuing treatment.

Following the completion of an initial treatment course with adalimumab, infliximab, vedolizumab or ustekinumab, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted supply of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine therapy is approved, a patient may swap if eligible to the alternate adalimumab, infliximab, vedolizumab or ustekinumab within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Crohn Disease Activity Index (CDAI) Score, confirmation of Crohn disease), or the prior conventional therapies of corticosteroid therapy and immunosuppressive therapy.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with adalimumab, infliximab, vedolizumab or ustekinumab at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug once within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment, within the timeframes specified in the relevant restriction.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the CDAI or evidence of intestinal inflammation submitted with the first authority application for adalimumab, infliximab, vedolizumab or ustekinumab. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to assess response to all subsequent treatments.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to recommence treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity. Patients must have received treatment with a corticosteroid and at least 1 immunosuppressive agent, at an adequate dose, for a minimum of 3 consecutive months immediately prior to the time the CDAI score or the indices of intestinal inflammation are measured.

(5) Patients 'grandfathered' onto PBS-subsidised treatment with vedolizumab.

A patient who commenced treatment with vedolizumab for severe Crohn disease prior to 1 August 2015 and who continues to receive treatment at the time of application may qualify for treatment under the initial 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment will be assessed under the continuing treatment restriction of the relevant drug.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must requalify for continuing treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' above for further details.

(6) Patients 'grandfathered' onto PBS-subsidised treatment with ustekinumab.

A patient who commenced treatment with ustekinumab for severe Crohn disease prior to 1 September 2017 and who continues to receive treatment at the time of application may qualify for treatment under the initial 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment will be assessed under the continuing treatment restriction of the relevant drug.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must requalify for continuing treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' above for further details.

Note Authority applications for increased quantities/ repeats (where relevant) may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

7763

Severe Crohn disease

Treatment Phase: Subsequent continuing treatment

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug.

Population criteria:

- Patient must be aged 18 years or older.

Clinical criteria:

- Patient must have an adequate response to this drug defined as a reduction in Crohn Disease Activity Index (CDAI) Score to a level no greater than 150 if assessed by CDAI or if affected by extensive small intestine disease; OR
- Patient must have an adequate response to this drug defined as (a) an improvement of intestinal inflammation as demonstrated by: (i) blood: normalisation of the platelet count, or an erythrocyte sedimentation rate (ESR) level no greater than 25 mm per hour, or a C-reactive protein (CRP) level no greater than 15 mg per L; or (ii) faeces: normalisation of lactoferrin or calprotectin level; or (iii) evidence of mucosal healing, as demonstrated by diagnostic imaging findings, compared to the baseline assessment; or (b) reversal of high faecal output state; or (c) avoidance of the need for surgery or total parenteral nutrition (TPN), if affected by short gut syndrome, extensive small intestine or is an ostomy patient.

The measurement of response to the prior course of therapy must be documented in the patient's medical notes.

Patients are eligible to receive subsequent continuing treatment with this drug in courses of up to 24 weeks at a dose of 5 mg per kg per dose providing they continue to sustain the response.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

infliximab 100 mg injection, 1 vial

11400B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	5	2	..	*1603.55	^a Inflectra [PF]	^a Renflexis [MK]

■ INFLIXIMAB

Note TREATMENT OF COMPLEX REFRACTORY FISTULISING CROHN DISEASE

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for patients with complex refractory fistulising Crohn disease. Where the term "biological medicine" appears in the following NOTES and restrictions, it refers to adalimumab and infliximab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the PBS- subsidised biological medicines for this condition at any one time.

From 1 April 2011, under the PBS, all patients will be able to commence a treatment cycle where they may trial PBS-subsidised adalimumab or infliximab without having to experience a disease flare when swapping to the alternate agent. Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with adalimumab or infliximab while they continue to show a response to therapy.

A patient who received PBS-subsidised adalimumab or infliximab treatment prior to 1 April 2011 is considered to have started their treatment cycle as of 1 April 2011.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised adalimumab or infliximab more than twice.

Once a patient has either failed or ceased to respond to treatment for this condition 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy for this condition before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised adalimumab or infliximab treatment in the most recent cycle to the date of the first application for initial treatment with adalimumab or infliximab under the new treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of less than 5 years, may commence a further course of treatment within the same treatment cycle.

A patient who has failed 3 trials or fewer of biological medicine therapy in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised adalimumab or infliximab therapy after 1 April 2011.

(a) Initial treatment. Applications for initial treatment should be made where:

- a patient has received no prior PBS-subsidised adalimumab or infliximab therapy in this treatment cycle and wishes to commence such therapy (Initial treatment (new patient or Recommencement of treatment after more than 5 years break in therapy - Initial 1); or
 - a patient has received prior PBS-subsidised (initial or continuing) adalimumab or infliximab therapy and wishes to trial an alternate agent (Initial 2 - Change or recommencement) [further details are under 'Swapping therapy' below]; or
 - a patient wishes to re-commence treatment with adalimumab or infliximab following a break in PBS-subsidised therapy with that agent (Change or Re-commencement of treatment after a break in therapy of less than 5 years (Initial 2)
- Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy for adalimumab and 14

weeks of therapy for infliximab.

From 1 April 2011, a patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab, and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

For subsequent courses of PBS-subsidised adalimumab or infliximab treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

Adalimumab only: Two completed authority prescriptions must be submitted with every initial application for adalimumab.

One prescription must be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats. The second prescription must be written for 2 doses of 40 mg and 2 repeats.

(b) Continuing treatment.

Adalimumab patients:

Following the completion of an initial treatment course with adalimumab, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

Infliximab patients:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment under the Initial 1 or Initial 2 treatment restrictions.

For the second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted supply of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap if eligible to the alternate biological medicine within the same treatment cycle.

A patient may trial the alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with adalimumab or infliximab at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug two times within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment within the timeframes specified in the relevant restriction.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements submitted with the first authority application for adalimumab or infliximab. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to these revised baseline measurements.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to recommence treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity.

Note Authority applications for increased quantities/ repeats (where relevant) may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

7847

Complex refractory Fistulising Crohn disease

Treatment Phase: Subsequent continuing treatment

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological agent treatment for this condition in this treatment cycle, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug.

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

An adequate response is defined as:

(a) a decrease from baseline in the number of open draining fistulae of greater than or equal to 50%; and/or

(b) a marked reduction in drainage of all fistula(e) from baseline, together with less pain and induration as reported by the patient.

The measurement of response to the prior course of therapy must be documented in the patient's medical notes.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Patients are eligible to receive subsequent continuing treatment with this drug in courses of up to 24 weeks at a dose of 5 mg per kg per dose providing they continue to sustain the response.

infliximab 100 mg injection, 1 vial

11423F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	5	2	..	*1603.55	^a Inflectra [PF]	^a Renflexis [MK]

■ INFLIXIMAB
Note TREATMENT OF PAEDIATRIC PATIENTS WITH REFRACTORY CROHN DISEASE

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) for paediatric patients with adalimumab for severe refractory Crohn disease and infliximab for moderate to severe refractory Crohn disease. Where the term "biological medicines" appears in the following NOTES and restrictions, it refers to adalimumab and infliximab only. A patient is eligible for PBS-subsidised treatment with only one PBS-subsidised biological medicine at any one time. For paediatric patients with Crohn disease, infliximab is PBS-subsidised for moderate to severe disease while adalimumab is PBS-subsidised for severe disease.

From 1 August 2015, under the PBS, patients commencing on adalimumab will be able to commence a treatment cycle where they may trial each PBS-subsidised biological medicine without having to experience a disease flare when swapping to infliximab. Patients on infliximab will be able to commence a treatment cycle where they may trial each PBS-subsidised biological medicine but will need to meet a PCDAI score of greater than or equal to 40 when swapping to adalimumab. Under these arrangements, within a single treatment cycle and depending on the disease severity, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy.

A patient who received PBS-subsidised biological medicine treatment prior to 1 August 2015 is considered to have started their treatment cycle as of 1 August 2015.

Within the same treatment cycle, a paediatric patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than twice.

Once a patient has either failed, or ceased to respond to treatment for this condition 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy for this condition before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle.

A patient who has failed 3 trials or fewer of biological medicine therapy in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 August 2015.

(a) Initial treatment.

Applications for initial treatment should be made where:

- i) a patient has received no prior PBS-subsidised biological medicine therapy in this treatment cycle and wishes to commence such therapy - Initial 1 treatment (new patient or Re-commencement of treatment after more than 5 years break in therapy); or
- (ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent - Initial 2 treatment (Change or Re-commencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or
- (iii) a patient wishes to re-commence treatment with a specific biological medicine following a break in PBS-subsidised therapy with that agent - Initial 2 treatment (Change or Re-commencement of treatment after a break in therapy of less than 5 years).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy for adalimumab and 14 weeks of therapy for infliximab.

From 1 August 2015, a patient must be assessed for response to any course of initial PBS-subsidised biological therapy following a minimum of 12 weeks of therapy for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab, and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

For first and subsequent continuing courses of PBS-subsidised biological medicine therapy, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

Adalimumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats and the second prescription should be written for 2 doses of 40 mg and 2 repeats for patients weighing 40 kg or greater. For patients weighing less than 40 kg, one prescription should be written for 2 doses of 40 mg with no repeats and the second prescription should be written for 2 doses of 20 mg with 3 repeats.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient remains eligible to receive up to 24 weeks per course of continuing treatment under the First continuing treatment and Subsequent Continuing treatment restrictions with that drug providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted supply of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine therapy is approved, a patient with severe disease may swap if eligible to the alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Paediatric Crohn Disease Activity Index (PCDAI) Score, confirmation of Crohn disease), or the prior conventional therapies of corticosteroid therapy, immunosuppressive therapy or enteral nutrition.

Patients on infliximab may swap to adalimumab within the same treatment cycle provided that their disease severity has

progressed to severe disease (i.e. they have a current PCDAI score of 40 or more).

A patient cannot swap to a biological medicine if they have failed to respond to prior treatment with that drug twice within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment within the timeframes specified in the relevant restriction.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the PCDAI submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to assess response to all subsequent treatments.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to recommence treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity.

Patients must have failed to achieve an adequate response to 2 of the following 3 conventional prior therapies including: (i) a tapered course of steroids, starting at a dose of at least 1 mg per kg or 40 mg (whichever is the lesser) prednisolone (or equivalent), over a 6 week period; (ii) an 8 week course of enteral nutrition; or (iii) immunosuppressive therapy including azathioprine at a dose of at least 2 mg per kg daily for 3 or more months, or, 6-mercaptopurine at a dose of at least 1 mg per kg daily for 3 or more months, or, methotrexate at a dose of at least 10 mg per square metre weekly for 3 or more months immediately prior to the time the PCDAI score is measured.

Note Authority applications for increased quantities/ repeats (where relevant) may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

7931

Moderate to severe Crohn disease

Treatment Phase: Subsequent continuing treatment

Treatment criteria:

- Must be treated by a gastroenterologist (code 87) or a consultant physician [internal medicine specialising in gastroenterology (code 81)] or a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician or a specialist paediatric gastroenterologist.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First Continuing treatment restriction; AND
- Patient must have a reduction in PCDAI Score by at least 15 points from baseline value; AND
- Patient must have a total PCDAI score of 30 points or less.

Population criteria:

- Patient must be aged 6 to 17 years inclusive.

The PCDAI assessment must be no more than 1 month old at the time of prescribing.

The PCDAI score must be documented in the patient's medical notes as the measurement of response to the prior course of therapy.

Patients are only eligible to receive subsequent continuing PBS-subsidised treatment with this drug in courses of up to 24 weeks at a dose of 5 mg per kg per dose providing they continue to sustain the response.

infliximab 100 mg injection, 1 vial

11449N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	5	2	..	*1603.55	^a Inflectra [PF]	^a Renflexis [MK]

■ INFLIXIMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE PSORIATIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab for adult patients with severe active psoriatic arthritis. Therefore, where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient receiving PBS-subsidised treatment for psoriatic arthritis is able to commence a treatment cycle where they may trial biological medicines without having to experience a disease flare when swapping to the alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

A patient who received PBS-subsidised adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab or ustekinumab treatment prior to 1 May 2019 is considered to start their first cycle as of 1 May 2019.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment.

Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence another cycle [further details are under '(5) Recommencement of treatment after a 5-year break in PBS-subsidised therapy' below].

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe active psoriatic arthritis.

(1) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an alternate medicine (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years) or An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, golimumab secukinumab and tofacitinib, 18 to 20 weeks of therapy for certolizumab pegol (depending upon the dosing regimen), 20 weeks of therapy for ixekizumab, 22 weeks of therapy for infliximab, and 28 weeks of therapy for ustekinumab. It is recommended that a patient be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of PBS-subsidised initial treatment following a minimum of 12 weeks of therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

Grandfather patients (ixekizumab only).

A patient who commenced treatment with ixekizumab for severe psoriatic arthritis prior to 1 March 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

Grandfather patients (tofacitinib only).

A patient who commenced treatment with Tofacitinib for severe psoriatic arthritis prior to 1 May 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment under the Initial 1 or Initial 2 treatment restrictions.

For the second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to re-qualify with respect to either the indices of disease severity (i.e. erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP) level, and active joint count) or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify under the Initial 3 treatment restriction with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application.

However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

Within a treatment cycle a patient may alternate between therapy with any biological medicine of their choice (1 at a time) providing:

- (i) they have not received PBS-subsidised treatment with that particular biological medicine previously; or
- (ii) they have demonstrated an adequate response to that particular biological medicine if they have previously trialled it on the PBS; and
- (iii) they have not previously failed to respond to treatment 3 times in this treatment cycle with that biological medicine.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(4) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the indices of disease severity submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and these revised baseline measurements will be used to assess response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to determine response for all subsequent continuing treatments. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. 20 or more active joints), response will be determined according to a reduction in the total number of active joints.

(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment restriction according to the criteria of the relevant restriction and index of disease severity. The application must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note Authority applications for increased quantities/ repeats (where relevant) may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

9188

Severe psoriatic arthritis

Treatment Phase: Subsequent continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment per subsequent continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an erythrocyte sedimentation rate (ESR) no greater than 25 mm per hour or a C-reactive protein (CRP) level no greater than 15 mg per L or either marker reduced by at least 20% from baseline; and

either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following major active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to determine response for all subsequent continuing treatments.

The measurement of response to the prior course of therapy must have been conducted following a minimum of 12 weeks of therapy with this drug and must be documented in the patient's medical records.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

infliximab 100 mg injection, 1 vial

11514B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	5	2	..	*1603.55	^a Inflectra [PF]	^a Renflexis [MK]

■ INFLIXIMAB
Note TREATMENT OF ADULT PATIENTS WITH ANKYLOSING SPONDYLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, certolizumab pegol, etanercept, golimumab, infliximab and secukinumab for adult patients with ankylosing spondylitis. Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab and secukinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 6 biological medicines at any 1 time.

Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last prescription for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised biological medicine treatment with adalimumab, certolizumab pegol, etanercept, golimumab, infliximab and secukinumab.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised biological medicine treatment in this treatment cycle and wishes to commence such therapy (Initial 1 - New patient)

(ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years); or

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

A patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy.

(b) Continuing treatment.

For the first continuing treatment course of PBS-subsidised biological medicine, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1, Initial 2 or Initial 3 treatment restrictions. For second and subsequent continuing courses of PBS-subsidised biological medicine it is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply. Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be conducted within 1 month of the last dose. Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI), or the prior NSAID therapy and exercise program requirements.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the BASDAI, ESR and/or CRP submitted with the first authority application for a biological medicine.

However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to these revised

baseline measurements.

For a new patient, the BASDAI used to determine the baseline must be measured while the patient is receiving NSAID therapy and completing their exercise program.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response.

(4) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must qualify under the Initial 3 treatment restriction. The same clinical criteria and indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI) as for the Initial 1 (New patient) restriction will need to be met, but a re-trial of NSAID therapy and exercise therapy is not required.

Authority required

Ankylosing spondylitis

Treatment Phase: Subsequent continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Ankylosing Spondylitis PBS Authority Application - Supporting Information Form.

An adequate response is defined as an improvement from baseline of at least 2 of the BASDAI and 1 of the following:

- (a) an ESR measurement no greater than 25 mm per hour; or
- (b) a CRP measurement no greater than 10 mg per L; or
- (c) an ESR or CRP measurement reduced by at least 20% from baseline.

Where only 1 acute phase reactant measurement is supplied in the first application for PBS-subsidised treatment, that same marker must be measured and supplied in all subsequent continuing treatment applications.

All measurements provided must be no more than 1 month old at the time of application.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg.

Up to a maximum of 3 repeats will be authorised.

Each application for subsequent continuing treatment with this drug must include an assessment of the patient's response to the prior course of therapy. If the response assessment is not provided at the time of application the patient will be deemed to have failed this course of treatment, unless the patient has experienced serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Ankylosing spondylitis

Treatment Phase: Continuing treatment - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the first continuing treatment restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the subsequent continuing Authority Required (in writing) treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

infliximab 100 mg injection, 1 vial

11482H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	320.71	Remicade [JC]

■ INFLIXIMAB
Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, etanercept, guselkumab, infliximab, ixekizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, etanercept, guselkumab, infliximab, ixekizumab, secukinumab, tildrakizumab and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, etanercept, guselkumab, infliximab, ixekizumab, secukinumab, tildrakizumab and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patients receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction. There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

- (i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or
- (ii) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years); or
- (iii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or
- (iv) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab and 28 weeks of therapy for tildrakizumab and ustekinumab.

It is recommended that a patient be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

Grandfather patients (guselkumab only).

A patient who commenced treatment with guselkumab for chronic plaque psoriasis prior to (1 February 2019) and who continues to receive treatment at the time of application, may qualify for treatment under the Initial treatment - Grandfather treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

Grandfather patients (tildrakizumab only).

A patient who commenced treatment with tildrakizumab for chronic plaque psoriasis prior to (1 February 2019) and who continues to receive treatment at the time of application, may qualify for treatment under the Initial treatment Grandfather treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment. This assessment must be conducted within 1 month of the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 the PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 1 month at the time of application.

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Subsequent continuing treatment, Whole body

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**

- Patient must not receive more than 24 weeks of treatment per subsequent continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg eight weekly. Up to a maximum of 2 repeats will be authorised.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet including the date of the assessment of the patient's condition.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Subsequent continuing treatment, Face, hand, foot

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per subsequent continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

(i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or

(ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg eight weekly. Up to a maximum of 2 repeats will be authorised.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet and face, hand, foot area diagrams including the date of the assessment of the patient's condition.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

infliximab 100 mg injection, 1 vial

11606W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	320.71	Remicade [JC]

■ INFLIXIMAB

Note TREATMENT OF ADULT PATIENTS WITH MODERATE TO SEVERE ULCERATIVE COLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, golimumab, infliximab and vedolizumab for adult patients with ulcerative colitis. Patients are eligible for PBS-subsidised treatment with either adalimumab, golimumab, infliximab or vedolizumab at any one time.

Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, golimumab, infliximab and vedolizumab only.

From 1 June 2018, under the PBS, all adult patients will be able to commence a treatment cycle where they may trial PBS-subsidised adalimumab, golimumab, infliximab or vedolizumab without having to experience a disease flare when swapping to one of the alternate agents. Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with adalimumab, golimumab, infliximab or vedolizumab while they continue to show a response to therapy.

A patient who received PBS-subsidised adalimumab, infliximab, vedolizumab treatment prior to 1 June 2018 is considered to start their first cycle as of 1 June 2018. Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised adalimumab, golimumab, infliximab or vedolizumab more than once. Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised adalimumab, golimumab, infliximab or vedolizumab treatment in the most recent cycle to the date of the first application for initial treatment with adalimumab, golimumab, infliximab or vedolizumab under the new treatment cycle.

A patient who has failed fewer than 3 trials of either adalimumab, golimumab, infliximab or vedolizumab in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

(1) How to prescribe PBS-subsidised treatment with adalimumab, golimumab, infliximab and vedolizumab after 1 June 2018.

(a) Initial treatment. Applications for initial treatment should be made where:

(i) an adult patient has received no prior PBS-subsidised treatment with adalimumab, golimumab, infliximab or vedolizumab in this treatment cycle and wishes to commence such therapy (Initial 1 - New patient or recommencement of treatment after more than 5 years break in therapy); or

(ii) an adult patient has received prior PBS-subsidised (initial or continuing) adalimumab, golimumab, infliximab or vedolizumab therapy and wishes to trial an alternate agent (Initial 2 - Change or Re-commencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) an adult patient wishes to re-commence treatment with adalimumab, golimumab, infliximab or vedolizumab following a break in PBS-subsidised therapy with the same agent (Initial 2 - Change or Re-commencement of treatment after a break in therapy of less than 5 years).

Treatment authorisations under Initial 1 and Initial 2 will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, 14 weeks of therapy for golimumab, infliximab and vedolizumab.

A patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of treatment for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for golimumab, infliximab and vedolizumab. For second and subsequent courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment.

(b) Continuing treatment.

Following the completion of an initial treatment course with adalimumab, golimumab, infliximab or vedolizumab a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed in the month prior to completing their current course of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised treatment is approved, a patient may swap if eligible to the alternate adalimumab, golimumab, infliximab or vedolizumab treatment within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Mayo clinic score or partial Mayo clinic score), or the prior corticosteroid therapy and immunosuppressive therapy. A patient may trial an alternate treatment at any time, regardless of whether they are receiving therapy (initial or continuing) with adalimumab, golimumab, infliximab or vedolizumab at the time of the application. However, they cannot swap to a particular therapy if they have failed to respond to prior treatment with that drug once within the same treatment cycle. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(3) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised adalimumab, golimumab, infliximab or vedolizumab therapy of at least 5 years, must requalify for initial 1 treatment with respect to the scores of disease severity. A patient must have received treatment with a 5-aminosalicylate oral preparation in a standard dose for induction of remission for a minimum of 3 consecutive months, and, either azathioprine or 6-mercaptopurine for a minimum of 3 consecutive months or a tapered course of oral steroids over a 6 week period followed by an appropriately dosed thiopurine agent for a minimum of 3 consecutive months (unless intolerance develops necessitating permanent treatment withdrawal to these agents). These above prior treatments must have been received immediately prior to the time the scores of disease severity being used to trial a second or subsequent course are measured.

(4) Patients 'grandfathered' onto PBS-subsidised treatment with golimumab.

A patient who commenced treatment with golimumab for moderate to severe ulcerative colitis prior to 1 June 2018 and who continues to receive treatment at the time of application, may qualify for treatment under the initial 3 'grandfather' treatment

restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

Note TREATMENT OF PAEDIATRIC PATIENTS WITH MODERATE TO SEVERE ULCERATIVE COLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) for paediatric patients with infliximab or adalimumab for moderate to severe ulcerative colitis; and infliximab for acute severe ulcerative colitis. Where the term 'biological medicine' appears in the following NOTES and restrictions, it refers to infliximab and adalimumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 2 biological medicines at any one time. Infliximab and adalimumab are PBS-subsidised for moderate to severe disease while only infliximab is PBS-subsidised for acute severe disease.

From 1 June 2017, under the PBS, all paediatric patients will be able to commence a treatment cycle where they may trial each PBS-subsidised biological medicine without having to experience a disease flare when swapping to the alternate agent. Under these arrangements, within a single treatment cycle and depending on the disease severity, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy. A patient who received PBS-subsidised biological medicine treatment prior to 1 June 2017 is considered to have started their treatment cycle as of 1 June 2017. Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than twice. Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle. A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle. There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 June 2017.

(a) Initial treatment. Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised biological medicine treatment in this treatment cycle and wishes to commence such therapy - Initial 1 treatment (new patient or Recommencement of treatment after more than 5 years break in therapy); or
- (ii) a patient has received prior PBS-subsidised (initial or continuing) treatment with a biological medicine and wishes to trial an alternate agent - Initial 2 treatment (Change or Re-commencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping treatment' below]; or
- (iii) a patient wishes to re-commence treatment with a specific biological medicine following a break in PBS-subsidised therapy with that agent - Initial 2 treatment (Change or Re-commencement of treatment after a break in therapy of less than 5 years).

Treatment authorisations under Initial 1 and Initial 2 treatment will be limited to provide for a maximum of 16 weeks of treatment for adalimumab and 14 weeks of treatment for infliximab. From 1 June 2017, a patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of treatment for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab. For second and subsequent courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment.

Adalimumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. For patients weighing 40 kg or greater, one prescription should be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats and the second prescription should be written for 2 doses of 40 mg and 2 repeats. For patients weighing less than 40 kg, one prescription should be written for 2 doses of 40 mg with no repeats and the second prescription should be written for 2 doses of 20 mg with 3 repeats.

(b) Continuing treatment. Following the completion of an initial treatment course with a specific biological medicine, a patient remains eligible to receive up to 24 weeks per course of continuing treatment with that drug under the continuing treatment restriction providing they continue to sustain the response. It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure PBS subsidy criteria are met.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap if eligible to the alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Paediatric Ulcerative Colitis Activity Index (PUCAI) Score, confirmation of ulcerative colitis disease), or the prior conventional therapies of corticosteroids or immunosuppressives. A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving treatment (initial or continuing) with infliximab or adalimumab at the time of the application. However, a patient cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug twice within the same treatment cycle. To ensure a patient receives the maximum treatment opportunities allowed under these swapping arrangements, it is important that they are assessed for response to every course of treatment.

(3) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to recommence treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity. A patient must have received treatment with a 5-aminosalicylate oral preparation in a standard dose for induction of remission for a minimum of 3 consecutive months, and, either azathioprine or 6-mercaptopurine for a minimum of 3 consecutive months or a tapered course of oral steroids over a 6 week period followed by an appropriately dosed thiopurine agent for a minimum of 3

consecutive months (unless intolerance develops necessitating permanent treatment withdrawal to these agents) immediately prior to the time the PUCAI score is measured.

Note Authority applications for increased quantities/ repeats (where relevant) may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

7989

Moderate to severe ulcerative colitis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician; OR
- Must be treated by a specialist paediatric gastroenterologist.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have demonstrated or sustained an adequate response to treatment by having a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1 while receiving treatment with this drug; OR
- Patient must have demonstrated or sustained an adequate response to treatment by having a Paediatric Ulcerative Colitis Activity Index (PUCAI) score of less than 10 while receiving treatment with this drug, if aged 6 to 17 years.

Population criteria:

- Patient must be 6 years of age or older.

Patients who have failed to maintain a partial Mayo clinic score of less than or equal to 2, with no subscore greater than 1, or, patients who have failed to maintain a Paediatric Ulcerative Colitis Activity Index (PUCAI) score of less than 10 (if aged 6 to 17 years) with continuing treatment with this drug, will not be eligible to receive further PBS-subsidised treatment with this drug.

Patients are only eligible to receive continuing PBS-subsidised treatment with this drug in courses of up to 24 weeks at a dose of 5 mg per kg per dose providing they continue to sustain the response.

The measurement of response to the prior course of therapy must be documented in the patient's medical notes.

infliximab 100 mg injection, 1 vial

11461F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	5	2	..	*1603.55	^a Inflectra [PF]	^a Renflexis [MK]

■ INFLIXIMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, etanercept, guselkumab, infliximab, ixekizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, etanercept, guselkumab, infliximab, ixekizumab, secukinumab, tildrakizumab and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, etanercept, guselkumab, infliximab, ixekizumab, secukinumab, tildrakizumab and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patients receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

- (i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or
- (ii) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years); or
- (iii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or
- (iv) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab and 28 weeks of therapy for tildrakizumab and ustekinumab.

It is recommended that a patient be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

Grandfather patients (guselkumab only).

A patient who commenced treatment with guselkumab for chronic plaque psoriasis prior to (1 February 2019) and who continues to receive treatment at the time of application, may qualify for treatment under the Initial treatment - Grandfather treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

Grandfather patients (tildrakizumab only).

A patient who commenced treatment with tildrakizumab for chronic plaque psoriasis prior to (1 February 2019) and who continues to receive treatment at the time of application, may qualify for treatment under the Initial treatment Grandfather treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment. This assessment must be conducted within 1 month of the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions. For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle.

To ensure a patients receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the

baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 the PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 1 month at the time of application.

Note No increase in the maximum number of repeats may be authorised.

Note Authority applications for increased quantities/ repeats (where relevant) may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required (STREAMLINED)

8844

Severe chronic plaque psoriasis

Treatment Phase: Subsequent continuing treatment, Whole body

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per subsequent continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

The measurement of response to the prior course of therapy must be documented in the patient's medical notes.

Determination of response must be based on the PASI assessment of response to the most recent course of treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Authority required (STREAMLINED)

8940

Severe chronic plaque psoriasis

Treatment Phase: Subsequent continuing treatment, Face, hand, foot

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per subsequent continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

(i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or

(ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

The measurement of response to the prior course of therapy must be documented in the patient's medical notes.

Determination of response must be based on the PASI assessment of response to the most recent course of treatment with this drug.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

infliximab 100 mg injection, 1 vial

11605T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	5	2	..	*1603.55	^a Inflectra [PF]	^a Renflexis [MK]

■ INFLIXIMAB

Note TREATMENT OF ADULT PATIENTS WITH MODERATE TO SEVERE ULCERATIVE COLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, golimumab, infliximab and vedolizumab for adult patients with ulcerative colitis. Patients are eligible for PBS-subsidised treatment with either adalimumab, golimumab, infliximab or vedolizumab at any one time.

Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, golimumab, infliximab and vedolizumab only.

From 1 June 2018, under the PBS, all adult patients will be able to commence a treatment cycle where they may trial PBS-subsidised adalimumab, golimumab, infliximab or vedolizumab without having to experience a disease flare when swapping to one of the alternate agents. Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with adalimumab, golimumab, infliximab or vedolizumab while they continue to show a response to therapy.

A patient who received PBS-subsidised adalimumab, infliximab, vedolizumab treatment prior to 1 June 2018 is considered to start their first cycle as of 1 June 2018. Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised adalimumab, golimumab, infliximab or vedolizumab more than once. Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised adalimumab, golimumab, infliximab or vedolizumab treatment in the most recent cycle to the date of the first application for initial treatment with adalimumab, golimumab, infliximab or vedolizumab under the new treatment cycle.

A patient who has failed fewer than 3 trials of either adalimumab, golimumab, infliximab or vedolizumab in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

(1) How to prescribe PBS-subsidised treatment with adalimumab, golimumab, infliximab and vedolizumab after 1 June 2018.

(a) Initial treatment. Applications for initial treatment should be made where:

(i) an adult patient has received no prior PBS-subsidised treatment with adalimumab, golimumab, infliximab or vedolizumab in this treatment cycle and wishes to commence such therapy (Initial 1 - New patient or recommencement of treatment after more than 5 years break in therapy); or

(ii) an adult patient has received prior PBS-subsidised (initial or continuing) adalimumab, golimumab, infliximab or vedolizumab therapy and wishes to trial an alternate agent (Initial 2 - Change or Re-commencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) an adult patient wishes to re-commence treatment with adalimumab, golimumab, infliximab or vedolizumab following a break in PBS-subsidised therapy with the same agent (Initial 2 - Change or Re-commencement of treatment after a break in therapy of less than 5 years).

Treatment authorisations under Initial 1 and Initial 2 will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, 14 weeks of therapy for golimumab, infliximab and vedolizumab.

A patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of treatment for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for golimumab, infliximab and vedolizumab. For second and subsequent courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment.

(b) Continuing treatment.

Following the completion of an initial treatment course with adalimumab, golimumab, infliximab or vedolizumab a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed in the month prior to completing their current course of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised treatment is approved, a patient may swap if eligible to the alternate adalimumab, golimumab, infliximab or vedolizumab treatment within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Mayo clinic score or partial Mayo clinic score), or the prior corticosteroid therapy and immunosuppressive therapy. A patient may trial an alternate treatment at any time, regardless of whether they are receiving therapy (initial or continuing) with adalimumab, golimumab, infliximab or vedolizumab at the time of the application. However, they cannot swap to a particular therapy if they have failed to respond to prior treatment with that drug once within the same treatment cycle. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(3) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised adalimumab, golimumab, infliximab or vedolizumab therapy of at least 5 years, must requalify for initial 1 treatment with respect to the scores of disease severity. A patient must have received treatment with a 5-aminosalicylate oral preparation in a standard dose for induction of remission for a minimum of 3 consecutive months, and, either azathioprine or 6-mercaptopurine for a minimum of 3 consecutive months or a tapered course of oral steroids over a 6 week period followed by an appropriately dosed thiopurine agent for a minimum of 3 consecutive months (unless intolerance develops necessitating permanent treatment withdrawal to these agents). These above prior treatments must have been received immediately prior to the time the scores of disease severity being used to trial a second or subsequent course are measured.

(4) Patients 'grandfathered' onto PBS-subsidised treatment with golimumab.

A patient who commenced treatment with golimumab for moderate to severe ulcerative colitis prior to 1 June 2018 and who continues to receive treatment at the time of application, may qualify for treatment under the initial 3 'grandfather' treatment

restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

Note TREATMENT OF PAEDIATRIC PATIENTS WITH MODERATE TO SEVERE ULCERATIVE COLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) for paediatric patients with infliximab or adalimumab for moderate to severe ulcerative colitis; and infliximab for acute severe ulcerative colitis.

Where the term 'biological medicine' appears in the following NOTES and restrictions, it refers to infliximab and adalimumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 2 biological medicines at any one time. Infliximab and adalimumab are PBS-subsidised for moderate to severe disease while only infliximab is PBS-subsidised for acute severe disease.

From 1 June 2017, under the PBS, all paediatric patients will be able to commence a treatment cycle where they may trial each PBS-subsidised biological medicine without having to experience a disease flare when swapping to the alternate agent. Under these arrangements, within a single treatment cycle and depending on the disease severity, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy. A patient who received PBS-subsidised biological medicine treatment prior to 1 June 2017 is considered to have started their treatment cycle as of 1 June 2017. Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than twice. Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle. A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle. There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 June 2017.

(a) Initial treatment. Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised biological medicine treatment in this treatment cycle and wishes to commence such therapy - Initial 1 treatment (new patient or Recommencement of treatment after more than 5 years break in therapy); or
- (ii) a patient has received prior PBS-subsidised (initial or continuing) treatment with a biological medicine and wishes to trial an alternate agent - Initial 2 treatment (Change or Re-commencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping treatment' below]; or
- (iii) a patient wishes to re-commence treatment with a specific biological medicine following a break in PBS-subsidised therapy with that agent - Initial 2 treatment (Change or Re-commencement of treatment after a break in therapy of less than 5 years).

Treatment authorisations under Initial 1 and Initial 2 treatment will be limited to provide for a maximum of 16 weeks of treatment for adalimumab and 14 weeks of treatment for infliximab. From 1 June 2017, a patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of treatment for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab. For second and subsequent courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment.

Adalimumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. For patients weighing 40 kg or greater, one prescription should be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats and the second prescription should be written for 2 doses of 40 mg and 2 repeats. For patients weighing less than 40 kg, one prescription should be written for 2 doses of 40 mg with no repeats and the second prescription should be written for 2 doses of 20 mg with 3 repeats.

(b) Continuing treatment. Following the completion of an initial treatment course with a specific biological medicine, a patient remains eligible to receive up to 24 weeks per course of continuing treatment with that drug under the continuing treatment restriction providing they continue to sustain the response. It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure PBS subsidy criteria are met.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap if eligible to the alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Paediatric Ulcerative Colitis Activity Index (PUCAI) Score, confirmation of ulcerative colitis disease), or the prior conventional therapies of corticosteroids or immunosuppressives. A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving treatment (initial or continuing) with infliximab or adalimumab at the time of the application. However, a patient cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug twice within the same treatment cycle. To ensure a patient receives the maximum treatment opportunities allowed under these swapping arrangements, it is important that they are assessed for response to every course of treatment.

(3) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to recommence treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity. A patient must have received treatment with a 5-aminosalicylate oral preparation in a standard dose for induction of remission for a minimum of 3 consecutive months, and, either azathioprine or 6-mercaptopurine for a minimum of 3 consecutive months or a tapered course of oral steroids over a 6 week period followed by an appropriately dosed thiopurine agent for a minimum of 3

consecutive months (unless intolerance develops necessitating permanent treatment withdrawal to these agents) immediately prior to the time the PUCAI score is measured.

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Initial 1 - New patient or recommencement of treatment after more than 5 years break in therapy

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician; OR
- Must be treated by a specialist paediatric gastroenterologist.

Clinical criteria:

- Patient must have failed to achieve an adequate response to a 5-aminosalicylate oral preparation in a standard dose for induction of remission for 3 or more consecutive months or have intolerance necessitating permanent treatment withdrawal, **AND**
- Patient must have failed to achieve an adequate response to azathioprine at a dose of at least 2 mg per kg daily for 3 or more consecutive months or have intolerance necessitating permanent treatment withdrawal; OR
- Patient must have failed to achieve an adequate response to 6-mercaptopurine at a dose of at least 1 mg per kg daily for 3 or more consecutive months or have intolerance necessitating permanent treatment withdrawal; OR
- Patient must have failed to achieve an adequate response to a tapered course of oral steroids, starting at a dose of at least 40 mg (for a child, 1 to 2 mg/kg up to 40 mg) prednisolone (or equivalent), over a 6 week period or have intolerance necessitating permanent treatment withdrawal, and followed by a failure to achieve an adequate response to 3 or more consecutive months of treatment of an appropriately dosed thiopurine agent, **AND**
- Patient must have a Mayo clinic score greater than or equal to 6 if an adult patient; OR
- Patient must have a partial Mayo clinic score greater than or equal to 6, provided the rectal bleeding and stool frequency subscores are both greater than or equal to 2 (endoscopy subscore is not required for a partial Mayo clinic score); OR
- Patient must have a Paediatric Ulcerative Colitis Activity Index (PUCAI) Score greater than or equal to 30 if aged 6 to 17 years; OR
- Patient must have previously received induction therapy with this drug for an acute severe episode of ulcerative colitis in the last 4 months and demonstrated an adequate response to induction therapy by achieving and maintaining a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1, or a PUCAI score less than 10 (if aged 6 to 17 years).

Population criteria:

- Patient must be 6 years of age or older.

Application for authorisation must be made in writing and must include:

(a) a completed authority prescription form; and

(b) a completed Ulcerative Colitis PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed current Mayo clinic or partial Mayo clinic or Paediatric Ulcerative Colitis Activity Index (PUCAI) calculation sheet including the date of assessment of the patient's condition; and

(ii) details of prior systemic drug therapy [dosage, date of commencement and duration of therapy].

A maximum quantity and number of repeats to provide for an initial course of this drug consisting of 3 doses at 5 mg per kg body weight per dose to be administered at weeks 0, 2 and 6, or to be administered at 8-weekly intervals for patients who have received prior treatment for an acute severe episode, will be authorised.

All tests and assessments should be performed preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior conventional treatment.

The most recent Mayo clinic, partial Mayo clinic or Paediatric Ulcerative Colitis Activity Index (PUCAI) score must be no more than 1 month old at the time of application.

Where treatment for an acute severe episode has occurred, an adequate response to induction therapy needs to be demonstrated by achieving and maintaining a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1, or a Paediatric Ulcerative Colitis Activity Index (PUCAI) score less than 10 (if aged 6 to 17 years), within the first 12 weeks of receiving this drug for acute severe ulcerative colitis.

Patients who fail to achieve a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1, or a Paediatric Ulcerative Colitis Activity Index (PUCAI) score less than 10 within the first 12 weeks of receiving this drug for ulcerative colitis, or have failed to maintain a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1, or have failed to maintain a PUCAI score less than 10 (if aged 6 to 17 years) with continuing treatment with this drug, will not be eligible to receive further PBS-subsidised treatment with this drug.

A partial Mayo clinic or Paediatric Ulcerative Colitis Activity Index (PUCAI) assessment of the patient's response to this initial course of treatment must be made up to 12 weeks after the first dose for patients administered doses at weeks 0, 2 and 6 (6 weeks following the third dose) so that there is adequate time for a response to be demonstrated.

The measurement of response to the prior course of therapy must be documented in the patient's medical notes.

If treatment with any of the above-mentioned drugs is contraindicated according to the relevant TGA-approved Product Information, details must be provided at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, details of this toxicity must be provided at the time of application.

Details of the accepted toxicities including severity can be found on the Department of Human Services website.

Note Biosimilar preferred prescribing policy

Prescribing of the biosimilar brand Inflectra or Renflexis is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Initial 2 - Change or Re-commencement of treatment after a break in therapy of less than 5 years

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician; OR
- Must be treated by a specialist paediatric gastroenterologist.

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle; OR
- Patient must have previously received PBS-subsidised treatment with a biological medicine (adalimumab or infliximab) for this condition in this treatment cycle if aged 6 to 17 years, **AND**
- Patient must not have failed PBS-subsidised treatment with this drug for this condition in the current treatment cycle; OR
- Patient must not have failed PBS-subsidised treatment with this drug for this condition in the current treatment cycle more than once if aged 6 to 17 years.

Population criteria:

- Patient must be 6 years of age or older.

Application for authorisation must be made in writing and must include:

(a) a completed authority prescription form; and

(b) a completed Ulcerative Colitis PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed current Mayo clinic or partial Mayo clinic or Paediatric Ulcerative Colitis Activity Index (PUCAI) calculation sheet including the date of assessment of the patient's condition; and

(ii) the number of total failures to courses of biological medicine therapy in the current treatment cycle.

To demonstrate a response to treatment the application must be accompanied by the results of the most recent course of biological medicine therapy within the timeframes specified in the relevant restriction.

If the response assessment to the previous course of biological medicine therapy is not met, the patient will be deemed to have failed therapy with that particular course of biological medicine.

A maximum quantity and number of repeats to provide for an initial course of this drug consisting of 3 doses at 5 mg per kg body weight per dose to be administered at weeks 0, 2 and 6, will be authorised.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg eight weekly.

Up to a maximum of 2 repeats will be authorised.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Balance of supply

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician; OR
- Must be treated by a specialist paediatric gastroenterologist.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 treatment (New patient or Re commencement of treatment after more than 5 years break in therapy) restriction to complete the 3 doses (the initial infusion regimen at 0, 2 and 6 weeks); OR

- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 treatment (Change or Re-commencement of treatment after a break in therapy of less than 5 years) restriction to complete the 3 doses (the initial infusion regimen at 0, 2 and 6 weeks), **AND**
- The treatment must provide no more than the balance of up to 3 doses (Initial 1 or Initial 2 treatment).

Population criteria:

- Patient must be 6 years of age or older.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

infliximab 100 mg injection, 1 vial

10196P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	320.71	^a Inflectra [PF]	^a Remicade [JC]
					^a Renflexis [MK]	

■ INFLIXIMAB
Note TREATMENT OF ADULT PATIENTS WITH SEVERE CROHN DISEASE

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for adult patients with severe Crohn disease. Where the term biological medicine appears in the following NOTES and restrictions, it refers to the tumour necrosis factor (TNF) alfa-antagonists (adalimumab and infliximab), the alpha-4 beta-7 integrin inhibitor (vedolizumab) and the human IgG1kappa monoclonal antibody (ustekinumab).

Patients are eligible for PBS-subsidised treatment with only 1 of the above PBS-subsidised biological medicines at any one time.

From 1 September 2017, under the PBS, all patients will be able to commence a treatment cycle where they may trial PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab without having to experience a disease flare when swapping to the alternate agent. Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with adalimumab, infliximab, vedolizumab or ustekinumab while they continue to show a response to therapy.

A patient who received PBS-subsidised adalimumab, infliximab, or vedolizumab treatment prior to 1 September 2017 is considered to have started their treatment cycle as of 1 September 2017.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab more than once.

Once a patient has either failed or ceased to respond to treatment for this condition 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy for this condition before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab treatment in the most recent cycle to the date of the first application for initial treatment with adalimumab, infliximab, vedolizumab or ustekinumab under the new treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of less than 5 years, may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab therapy after 1 September 2017.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised therapy with adalimumab, infliximab, vedolizumab or ustekinumab in this treatment cycle and wishes to commence such therapy (Initial treatment (new patient or Re-commencement of treatment after more than 5 years break in therapy - Initial 1)); or

(ii) a patient has received prior PBS-subsidised (initial or continuing) adalimumab, infliximab, vedolizumab or ustekinumab and wishes to trial an alternate agent (Initial 2 - Change or recommencement) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to re-commence treatment with adalimumab, infliximab, vedolizumab or ustekinumab following a break in PBS-subsidised therapy with that agent (Change or Re-commencement of treatment after a break in therapy of less than 5 years (Initial 2)).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, 14 weeks of therapy for infliximab, 14 weeks of therapy for vedolizumab and 16 weeks for ustekinumab.

From 1 September 2017, a patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy for adalimumab or ustekinumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab or vedolizumab, and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

For subsequent courses of PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

Adalimumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats and the second prescription should be written for 2 doses of 40 mg and 2 repeats for patients weighing 40 kg or greater. For patients weighing less than 40 kg, one prescription should be written for 2 doses of 40 mg with no repeats and the second prescription should be written for 2 doses of 20 mg with 3 repeats.

Ustekinumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be written under S100 (Highly Specialised Drugs) for a weight-based loading dose, containing a quantity

of up to 4 vials of 130 mg and no repeats. The second prescription should be written under S85 (General) for the subsequent first dose, containing a quantity of 2 vials of 45 mg and no repeats.

(b) Continuing treatment.

Following the completion of an initial treatment course with adalimumab, infliximab, vedolizumab or ustekinumab, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted supply of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine therapy is approved, a patient may swap if eligible to the alternate adalimumab, infliximab, vedolizumab or ustekinumab within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Crohn Disease Activity Index (CDAI) Score, confirmation of Crohn disease), or the prior conventional therapies of corticosteroid therapy and immunosuppressive therapy.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with adalimumab, infliximab, vedolizumab or ustekinumab at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug once within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment, within the timeframes specified in the relevant restriction.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the CDAI or evidence of intestinal inflammation submitted with the first authority application for adalimumab, infliximab, vedolizumab or ustekinumab. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to assess response to all subsequent treatments.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to recommence treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity. Patients must have received treatment with a corticosteroid and at least 1 immunosuppressive agent, at an adequate dose, for a minimum of 3 consecutive months immediately prior to the time the CDAI score or the indices of intestinal inflammation are measured.

(5) Patients 'grandfathered' onto PBS-subsidised treatment with vedolizumab.

A patient who commenced treatment with vedolizumab for severe Crohn disease prior to 1 August 2015 and who continues to receive treatment at the time of application may qualify for treatment under the initial 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment will be assessed under the continuing treatment restriction of the relevant drug.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must requalify for continuing treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' above for further details.

(6) Patients 'grandfathered' onto PBS-subsidised treatment with ustekinumab.

A patient who commenced treatment with ustekinumab for severe Crohn disease prior to 1 September 2017 and who continues to receive treatment at the time of application may qualify for treatment under the initial 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment will be assessed under the continuing treatment restriction of the relevant drug.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must requalify for continuing treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' above for further details.

Authority required

Severe Crohn disease

Treatment Phase: Initial treatment (new patient or Re-commencement of treatment after more than 5 years break in therapy - Initial 1)

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have confirmed severe Crohn disease, defined by standard clinical, endoscopic and/or imaging features, including histological evidence, with the diagnosis confirmed by a gastroenterologist or a consultant physician, **AND**
- Patient must have failed to achieve an adequate response to prior systemic therapy with a tapered course of steroids, starting at a dose of at least 40 mg prednisolone (or equivalent), over a 6 week period, **AND**
- Patient must have failed to achieve adequate response to prior systemic immunosuppressive therapy with azathioprine at a dose of at least 2 mg per kg daily for 3 or more consecutive months; OR
- Patient must have failed to achieve adequate response to prior systemic immunosuppressive therapy with 6-mercaptopurine at a dose of at least 1 mg per kg daily for 3 or more consecutive months; OR

- Patient must have failed to achieve adequate response to prior systemic immunosuppressive therapy with methotrexate at a dose of at least 15 mg weekly for 3 or more consecutive months, **AND**
- Patient must have a Crohn Disease Activity Index (CDAI) Score greater than or equal to 300 as evidence of failure to achieve an adequate response to prior systemic therapy; OR
- Patient must have short gut syndrome with diagnostic imaging or surgical evidence, or have had an ileostomy or colostomy; and must have evidence of intestinal inflammation; and must have evidence of failure to achieve an adequate response to prior systemic therapy as specified below; OR
- Patient must have extensive intestinal inflammation affecting more than 50 cm of the small intestine as evidenced by radiological imaging; and must have a Crohn Disease Activity Index (CDAI) Score greater than or equal to 220; and must have evidence of failure to achieve an adequate response to prior systemic therapy as specified below.

Population criteria:

- Patient must be aged 18 years or older.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current Crohn Disease Activity Index (CDAI) calculation sheet including the date of assessment of the patient's condition if relevant; and
 - (ii) details of prior systemic drug therapy [dosage, date of commencement and duration of therapy]; and
 - (iii) the reports and dates of the pathology or diagnostic imaging test(s) nominated as the response criterion, if relevant; and
 - (iv) the date of the most recent clinical assessment; and
 - (v) the signed patient acknowledgement indicating they understand and acknowledge that the PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment.

Evidence of failure to achieve an adequate response to prior therapy must include at least one of the following: (a) patient must have evidence of intestinal inflammation; (b) patient must be assessed clinically as being in a high faecal output state; (c) patient must be assessed clinically as requiring surgery or total parenteral nutrition (TPN) as the next therapeutic option, in the absence of this drug, if affected by short gut syndrome, extensive small intestine disease or is an ostomy patient. Evidence of intestinal inflammation includes: (i) blood: higher than normal platelet count, or, an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour, or, a C-reactive protein (CRP) level greater than 15 mg per L; or (ii) faeces: higher than normal lactoferrin or calprotectin level; or (iii) diagnostic imaging: demonstration of increased uptake of intravenous contrast with thickening of the bowel wall or mesenteric lymphadenopathy or fat streaking in the mesentery; All assessments, pathology tests, and diagnostic imaging studies must be made within 1 month of the date of application and preferably should be performed whilst still on treatment with the most recent course of prior therapies.

If treatment with any of the specified prior conventional drugs is contraindicated according to the relevant TGA-approved Product Information, please provide details at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, details of this toxicity must be provided at the time of application.

Details of the accepted toxicities including severity can be found on the Department of Human Services website.

Any one of the baseline criteria may be used to determine response to an initial course of treatment and eligibility for continued therapy, according to the criteria included in the first or subsequent continuing treatment restrictions. However, the same criterion must be used for any subsequent determination of response to treatment, for the purpose of eligibility for continuing PBS-subsidised therapy.

A maximum quantity and number of repeats to provide for an initial course of this drug consisting of 3 doses at 5 mg per kg body weight per dose to be administered at weeks 0, 2 and 6, will be authorised.

The assessment of the patient's response to this initial course of treatment must be made up to 12 weeks after the first dose (6 weeks following the third dose) so that there is adequate time for a response to be demonstrated.

This assessment, which will be used to determine eligibility for the first continuing treatment, must be submitted to the Department of Human Services no later than 1 month from the date of completion of this initial course of treatment.

Where a response assessment is not undertaken and submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with this drug.

Note It is recommended that an application for the first continuing treatment is submitted to the Department of Human Services at the time of the 12 week assessment, to ensure continuity of treatment for those patients who meet the first continuing treatment criterion for PBS-subsidised treatment with this drug.

Note Biosimilar preferred prescribing policy

Prescribing of the biosimilar brand Inflectra or Renflexis is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe Crohn disease

Treatment Phase: Change or Re-commencement of treatment after a break in therapy of less than 5 years (Initial 2)

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have failed PBS-subsidised therapy with this drug for this condition in the current treatment cycle.

Population criteria:

- Patient must be aged 18 years or older.

Applications for authorisation must be made in writing and must include: (a) a completed authority prescription form; and (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form, which includes the following: (i) the completed current Crohn Disease Activity Index (CDAI) Score calculation sheet including the date of assessment of the patient's condition if relevant; or (ii) the reports and dates of the pathology or diagnostic imaging test(s) used to assess response to therapy for patients with short gut syndrome, extensive small intestine disease or an ostomy, if relevant; and (iii) the date of clinical assessment; and (iv) the details of prior biological medicine treatment including the details of date and duration of treatment.

To demonstrate a response to treatment the application must be accompanied by the results of the most recent course of biological medicine therapy within the timeframes specified in the relevant restriction.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under an initial treatment restriction, the patient must have been assessed for response to that course following a minimum of 12 weeks of therapy for adalimumab or ustekinumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab and vedolizumab and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

If the response assessment to the previous course of biological medicine treatment is not submitted as detailed above, the patient will be deemed to have failed therapy with that particular course of biological medicine.

A maximum quantity and number of repeats to provide for an initial course of this drug consisting of 3 doses at 5 mg per kg body weight per dose to be administered at weeks 0, 2 and 6, will be authorised.

The assessment of the patient's response to this initial course of treatment must be made up to 12 weeks after the first dose (6 weeks following the third dose) so that there is adequate time for a response to be demonstrated.

This assessment, which will be used to determine eligibility for the first continuing treatment, must be submitted to the Department of Human Services no later than 1 month from the date of completion of this initial course of treatment.

Where a response assessment is not undertaken and submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with this drug.

Note It is recommended that an application for the first continuing treatment is submitted to the Department of Human Services at the time of the 12 week assessment, to ensure continuity of treatment for those patients who meet the first continuing treatment criterion for PBS-subsidised treatment with this drug.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe Crohn disease

Treatment Phase: Balance of supply

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial treatment (new patient or Recommencement of treatment after more than 5 years break in therapy - Initial 1) restriction to complete the 3 doses (the initial infusion regimen at 0, 2 and 6 weeks); OR
- Patient must have received insufficient therapy with this drug for this condition under the Change or Re-commencement of treatment after a break in therapy of less than 5 years (Initial 2) restriction to complete the 3 doses (the initial infusion regimen at 0, 2 and 6 weeks); OR
- Patient must have received insufficient therapy with this drug for this condition under the first continuing treatment or subsequent continuing treatment restrictions to complete 24 weeks of treatment, **AND**
- The treatment must provide no more than the balance of up to 3 doses (Initial 1 or Initial 2 treatment) or 2 repeats (first Continuing or Subsequent Continuing treatment).

Population criteria:

- Patient must be aged 18 years or older.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Severe Crohn disease

Treatment Phase: First continuing treatment

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must have an adequate response to this drug defined as a reduction in Crohn Disease Activity Index (CDAI) Score to a level no greater than 150 if assessed by CDAI or if affected by extensive small intestine disease; OR
- Patient must have an adequate response to this drug defined as (a) an improvement of intestinal inflammation as demonstrated by: (i) blood: normalisation of the platelet count, or an erythrocyte sedimentation rate (ESR) level no greater than 25 mm per hour, or a C-reactive protein (CRP) level no greater than 15 mg per L; or (ii) faeces: normalisation of lactoferrin or calprotectin level; or (iii) evidence of mucosal healing, as demonstrated by diagnostic imaging findings, compared to the baseline assessment; or (b) reversal of high faecal output state; or (c) avoidance of the need for surgery or total parenteral nutrition (TPN), if affected by short gut syndrome, extensive small intestine or is an ostomy patient.

Population criteria:

- Patient must be aged 18 years or older.

Applications for authorisation must be made in writing and must include:

(a) a completed authority prescription form; and

(b) a completed Crohn Disease PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed Crohn Disease Activity Index (CDAI) Score calculation sheet including the date of the assessment of the patient's condition, if relevant; or

(ii) the reports and dates of the pathology test or diagnostic imaging test(s) used to assess response to therapy for patients with short gut syndrome, extensive small intestine disease or an ostomy, if relevant; and

(iii) the date of clinical assessment.

All assessments, pathology tests, and diagnostic imaging studies must be made within 1 month of the date of application.

The application for first continuing treatment with this drug must include an assessment of the patient's response to the initial course of treatment. The assessment must be made up to 12 weeks after the first dose so that there is adequate time for a response to be demonstrated. This assessment must be submitted no later than 4 weeks from the cessation of that treatment course.

Where a response assessment is not undertaken and submitted within these timeframes, the patient will be deemed to have failed to respond to treatment with this drug.

A maximum of 24 weeks of treatment with this drug will be authorised under this criterion.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg eight weekly. Up to a maximum of 2 repeats will be authorised.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

infliximab 100 mg injection, 1 vial

5754W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	320.71	^a Inflectra [PF]	^a Remicade [JC]
					^a Renflexis [MK]	

■ INFLIXIMAB

Note TREATMENT OF PAEDIATRIC PATIENTS WITH REFRACTORY CROHN DISEASE

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) for paediatric patients with adalimumab for severe refractory Crohn disease and infliximab for moderate to severe refractory Crohn disease. Where the term "biological medicines" appears in the following NOTES and restrictions, it refers to adalimumab and infliximab only. A patient is eligible for PBS-subsidised treatment with only one PBS-subsidised biological medicine at any one time. For paediatric patients with Crohn disease, infliximab is PBS-subsidised for moderate to severe disease while adalimumab is PBS-subsidised for severe disease.

From 1 August 2015, under the PBS, patients commencing on adalimumab will be able to commence a treatment cycle where they may trial each PBS-subsidised biological medicine without having to experience a disease flare when swapping to infliximab. Patients on infliximab will be able to commence a treatment cycle where they may trial each PBS-subsidised biological medicine but will need to meet a PCDAI score of greater than or equal to 40 when swapping to adalimumab.

Under these arrangements, within a single treatment cycle and depending on the disease severity, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy.

A patient who received PBS-subsidised biological medicine treatment prior to 1 August 2015 is considered to have started their treatment cycle as of 1 August 2015.

Within the same treatment cycle, a paediatric patient cannot trial and fail, or cease to respond to, the same PBS-subsidised

biological medicine more than twice.

Once a patient has either failed, or ceased to respond to treatment for this condition 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy for this condition before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle.

A patient who has failed 3 trials or fewer of biological medicine therapy in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 August 2015.

(a) Initial treatment.

Applications for initial treatment should be made where:

- i) a patient has received no prior PBS-subsidised biological medicine therapy in this treatment cycle and wishes to commence such therapy - Initial 1 treatment (new patient or Re-commencement of treatment after more than 5 years break in therapy); or
- (ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent - Initial 2 treatment (Change or Re-commencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or
- (iii) a patient wishes to re-commence treatment with a specific biological medicine following a break in PBS-subsidised therapy with that agent - Initial 2 treatment (Change or Re-commencement of treatment after a break in therapy of less than 5 years).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy for adalimumab and 14 weeks of therapy for infliximab.

From 1 August 2015, a patient must be assessed for response to any course of initial PBS-subsidised biological therapy following a minimum of 12 weeks of therapy for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab, and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

For first and subsequent continuing courses of PBS-subsidised biological medicine therapy, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

Adalimumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats and the second prescription should be written for 2 doses of 40 mg and 2 repeats for patients weighing 40 kg or greater. For patients weighing less than 40 kg, one prescription should be written for 2 doses of 40 mg with no repeats and the second prescription should be written for 2 doses of 20 mg with 3 repeats.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient remains eligible to receive up to 24 weeks per course of continuing treatment under the First continuing treatment and Subsequent Continuing treatment restrictions with that drug providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted supply of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine therapy is approved, a patient with severe disease may swap if eligible to the alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Paediatric Crohn Disease Activity Index (PCDAI) Score, confirmation of Crohn disease), or the prior conventional therapies of corticosteroid therapy, immunosuppressive therapy or enteral nutrition. Patients on infliximab may swap to adalimumab within the same treatment cycle provided that their disease severity has progressed to severe disease (i.e. they have a current PCDAI score of 40 or more).

A patient cannot swap to a biological medicine if they have failed to respond to prior treatment with that drug twice within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment within the timeframes specified in the relevant restriction.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the PCDAI submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to assess response to all subsequent treatments.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to recommence treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity.

Patients must have failed to achieve an adequate response to 2 of the following 3 conventional prior therapies including: (i) a tapered course of steroids, starting at a dose of at least 1 mg per kg or 40 mg (whichever is the lesser) prednisolone (or equivalent), over a 6 week period; (ii) an 8 week course of enteral nutrition; or (iii) immunosuppressive therapy including azathioprine at a dose of at least 2 mg per kg daily for 3 or more months, or, 6-mercaptopurine at a dose of at least 1 mg per

kg daily for 3 or more months, or, methotrexate at a dose of at least 10 mg per square metre weekly for 3 or more months immediately prior to the time the PCDAI score is measured.

Authority required

Moderate to severe Crohn disease

Treatment Phase: Initial 1 - New patient or recommencement of treatment after more than 5 years break in therapy

Treatment criteria:

- Must be treated by a gastroenterologist (code 87) or a consultant physician [internal medicine specialising in gastroenterology (code 81)] or a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician or a specialist paediatric gastroenterologist.

Clinical criteria:

- Patient must have confirmed diagnosis of Crohn disease, defined by standard clinical, endoscopic and/or imaging features including histological evidence, **AND**
- Patient must have failed to achieve an adequate response to 2 of the following 3 conventional prior therapies including: (i) a tapered course of steroids, starting at a dose of at least 1 mg per kg or 40 mg (whichever is the lesser) prednisolone (or equivalent), over a 6 week period; (ii) an 8 week course of enteral nutrition; or (iii) immunosuppressive therapy including azathioprine at a dose of at least 2 mg per kg daily for 3 or more months, or, 6-mercaptopurine at a dose of at least 1 mg per kg daily for 3 or more months, or, methotrexate at a dose of at least 10 mg per square metre weekly for 3 or more months; OR
- Patient must have a documented intolerance of a severity necessitating permanent treatment withdrawal or a contra-indication to each of prednisolone (or equivalent), azathioprine, 6-mercaptopurine and methotrexate, **AND**
- Patient must have a Paediatric Crohn Disease Activity Index (PCDAI) Score greater than or equal to 30 preferably whilst still on treatment.

Population criteria:

- Patient must be aged 6 to 17 years inclusive.

Application for authorisation must be made in writing and must include:

(a) a completed authority prescription form; and

(b) a completed Paediatric Crohn Disease PBS Authority Application -Supporting Information Form which includes the following:

(i) the completed current Paediatric Crohn Disease Activity Index (PCDAI) calculation sheet including the date of assessment of the patient's condition which must be no more than one month old at the time of application; and

(ii) details of previous systemic drug therapy [dosage, date of commencement and duration of therapy] or dates of enteral nutrition.

The PCDAI score should preferably be obtained whilst on conventional treatment but must be obtained within one month of the last conventional treatment dose.

If treatment with any of the specified prior conventional drugs is contraindicated according to the relevant TGA-approved Product Information, please provide details at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, details of this toxicity must be provided at the time of application.

Details of the accepted toxicities including severity can be found on the Department of Human Services website.

A maximum quantity and number of repeats to provide for an initial course of this drug consisting of 3 doses at 5 mg per kg body weight per dose to be administered at weeks 0, 2 and 6, will be authorised.

A PCDAI assessment of the patient's response to this initial course of treatment must be made up to 12 weeks after the first dose (6 weeks following the third dose) so that there is adequate time for a response to be demonstrated.

This assessment, which will be used to determine eligibility for the first continuing treatment, must be submitted to the Department of Human Services no later than 1 month from the date of completion of this initial course of treatment.

Where a response assessment is not undertaken and submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with this drug.

Note It is recommended that an application for the first continuing treatment is submitted to the Department of Human Services at the time of the 12 week assessment, to ensure continuity of treatment for those patients who meet the first continuing treatment criteria for PBS-subsidised treatment with this drug.

Note Biosimilar preferred prescribing policy

Prescribing of the biosimilar brand Inflectra or Renflexis is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Moderate to severe Crohn disease

Treatment Phase: Initial 2 - Change or Re-commencement of treatment after a break in therapy of less than 5 years

Treatment criteria:

- Must be treated by a gastroenterologist (code 87) or a consultant physician [internal medicine specialising in gastroenterology (code 81)] or a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician or a specialist paediatric gastroenterologist.

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have failed PBS-subsidised therapy with this drug for this condition more than once in the current treatment cycle.

Population criteria:

- Patient must be aged 6 to 17 years inclusive.

Application for authorisation must be made in writing and must include:

(a) a completed authority prescription form; and

(b) a completed Paediatric Crohn Disease PBS Authority Application -Supporting Information Form which includes the following:

(i) the completed current Paediatric Crohn Disease Activity Index (PCDAI) Score calculation sheet; and

(ii) details of prior biological medicine treatment including details of date and duration of treatment.

To demonstrate a response to treatment the application must be accompanied by the results of the most recent course of biological medicine therapy within the timeframes specified in the relevant restriction.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under an initial treatment restriction, the patient must have been assessed for response to that course following a minimum of 12 weeks therapy for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

If the response assessment to the previous course of biological medicine treatment is not submitted as detailed above, the patient will be deemed to have failed therapy with that particular course of biological medicine.

A maximum quantity and number of repeats to provide for an initial course of this drug consisting of 3 doses at 5 mg per kg body weight per dose to be administered at weeks 0, 2 and 6, will be authorised.

A PCDAI assessment of the patient's response to this initial course of treatment must be made up to 12 weeks after the first dose (6 weeks following the third dose) so that there is adequate time for a response to be demonstrated.

This assessment, which will be used to determine eligibility for the first continuing treatment, must be submitted to the Department of Human Services no later than 1 month from the date of completion of this initial course of treatment.

Where a response assessment is not undertaken and submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with this drug.

Note It is recommended that an application for the first continuing treatment is submitted to the Department of Human Services at the time of the 12 week assessment, to ensure continuity of treatment for those patients who meet the first continuing treatment criteria for PBS-subsidised treatment with this drug.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Moderate to severe Crohn disease

Treatment Phase: Balance of supply

Treatment criteria:

- Must be treated by a gastroenterologist (code 87) or a consultant physician [internal medicine specialising in gastroenterology (code 81)] or a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician or a specialist paediatric gastroenterologist.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 treatment (New patient or Re-commencement of treatment after more than 5 years break in therapy) restriction to complete the 3 doses (the initial infusion regimen at 0, 2 and 6 weeks); OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 treatment (Change or Re-commencement of treatment after a break in therapy of less than 5 years) restriction to complete the 3 doses the initial infusion regimen at 0, 2 and 6 weeks); OR
- Patient must have received insufficient therapy with this drug for this condition under the first continuing treatment or subsequent continuing treatment restrictions to complete 24 weeks of treatment, **AND**
- The treatment must provide no more than the balance of up to 3 doses (Initial 1 or Initial 2 treatment) or 2 repeats (first Continuing or Subsequent Continuing treatment).

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Moderate to severe Crohn disease

Treatment Phase: First continuing treatment

Treatment criteria:

- Must be treated by a gastroenterologist (code 87) or a consultant physician [internal medicine specialising in gastroenterology (code 81)] or a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician or a specialist paediatric gastroenterologist.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have a reduction in PCDAI Score by at least 15 points from baseline value; **AND**
- Patient must have a total PCDAI score of 30 points or less.

Population criteria:

- Patient must be aged 6 to 17 years inclusive.

Application for authorisation must be made in writing and must include:

(a) a completed authority prescription form; and

(b) a completed Paediatric Crohn Disease PBS Authority Application - Supporting Information Form, which includes the completed Paediatric Crohn Disease Activity Index (PCDAI) calculation sheet along with the date of the assessment of the patient's condition.

The PCDAI assessment must be no more than 1 month old at the time of application.

The application for first continuing treatment with this drug must include a PCDAI assessment of the patient's response to the initial course of treatment. The assessment must be made up to 12 weeks after the first dose so that there is adequate time for a response to be demonstrated. This assessment must be submitted no later than 4 weeks from the cessation of that treatment course.

Where a response assessment is not undertaken and submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with this drug.

A maximum of 24 weeks of treatment with this drug will be authorised under this restriction.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg eight weekly.

Up to a maximum of 2 repeats will be authorised.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

infliximab 100 mg injection, 1 vial

5755X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	320.71	^a Inflectra [PF]	^a Remicade [JC]
					^a Renflexis [MK]	

■ INFLIXIMAB
Note TREATMENT OF COMPLEX REFRACTORY FISTULISING CROHN DISEASE

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for patients with complex refractory fistulising Crohn disease. Where the term "biological medicine" appears in the following NOTES and restrictions, it refers to adalimumab and infliximab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the PBS- subsidised biological medicines for this condition at any one time.

From 1 April 2011, under the PBS, all patients will be able to commence a treatment cycle where they may trial PBS-subsidised adalimumab or infliximab without having to experience a disease flare when swapping to the alternate agent. Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with adalimumab or infliximab while they continue to show a response to therapy.

A patient who received PBS-subsidised adalimumab or infliximab treatment prior to 1 April 2011 is considered to have started their treatment cycle as of 1 April 2011.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised adalimumab or infliximab more than twice.

Once a patient has either failed or ceased to respond to treatment for this condition 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy for this condition before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised adalimumab or infliximab treatment in the most recent cycle to the date of the first application for initial treatment with adalimumab or infliximab under the new treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of less than 5 years, may commence a further course of treatment within the same treatment cycle.

A patient who has failed 3 trials or fewer of biological medicine therapy in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised adalimumab or infliximab therapy after 1 April 2011.

(a) Initial treatment. Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised adalimumab or infliximab therapy in this treatment cycle and wishes to commence such therapy (Initial treatment (new patient or Recommencement of treatment after more than 5 years break in therapy - Initial 1); or

(ii) a patient has received prior PBS-subsidised (initial or continuing) adalimumab or infliximab therapy and wishes to trial an alternate agent (Initial 2 - Change or recommencement) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to re-commence treatment with adalimumab or infliximab following a break in PBS-subsidised therapy with that agent (Change or Re-commencement of treatment after a break in therapy of less than 5 years (Initial 2)) Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy for adalimumab and 14 weeks of therapy for infliximab.

From 1 April 2011, a patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab, and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

For subsequent courses of PBS-subsidised adalimumab or infliximab treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

Adalimumab only: Two completed authority prescriptions must be submitted with every initial application for adalimumab. One prescription must be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats. The second prescription must be written for 2 doses of 40 mg and 2 repeats.

(b) Continuing treatment.

Adalimumab patients:

Following the completion of an initial treatment course with adalimumab, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

Infliximab patients:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment under the Initial 1 or Initial 2 treatment restrictions.

For the second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted supply of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap if eligible to the alternate biological medicine within the same treatment cycle.

A patient may trial the alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with adalimumab or infliximab at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug two times within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment within the timeframes specified in the relevant restriction.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements submitted with the first authority application for adalimumab or infliximab. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to these revised baseline measurements.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to recommence treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity.

Authority required

Complex refractory Fistulising Crohn disease

Treatment Phase: Initial treatment (new patient or Re-commencement of treatment after more than 5 years break in therapy - Initial 1)

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have confirmed Crohn disease, defined by standard clinical, endoscopic and/or imaging features, including histological evidence, with the diagnosis confirmed by a gastroenterologist or a consultant physician, **AND**
- Patient must have an externally draining enterocutaneous or rectovaginal fistula.

Applications for authorisation must be made in writing and must include:

(a) a completed authority prescription form; and

(b) a completed Fistulising Crohn Disease PBS Authority Application - Supporting Information Form which includes the following:

- (i) a completed current Fistula Assessment Form including the date of assessment of the patient's condition; and
- (ii) a signed patient acknowledgement.

The most recent fistula assessment must be no more than 1 month old at the time of application.

A maximum quantity and number of repeats to provide for an initial course of this drug consisting of 3 doses at 5 mg per kg body weight per dose to be administered at weeks 0, 2 and 6, will be authorised.

An assessment of the patient's response to this initial course of treatment must be made up to 12 weeks after the first dose (up to 6 weeks following the third dose) so that there is adequate time for a response to be demonstrated.

This assessment, which will be used to determine eligibility for the first continuing treatment, must be submitted to the Department of Human Services no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with this drug.

Note Biosimilar preferred prescribing policy

Prescribing of the biosimilar brand Inflectra or Renflexis is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note It is recommended that an application for the first continuing treatment is submitted to the Department of Human Services at the time of the 12 week assessment, to ensure continuity of treatment for those patients who meet the first continuing treatment criteria for PBS-subsidised treatment with this drug.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Complex refractory Fistulising Crohn disease

Treatment Phase: Change or Re-commencement of treatment after a break in therapy of less than 5 years (Initial 2)

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have failed PBS-subsidised therapy with this drug for this condition more than once in the current treatment cycle.

Applications for authorisation must be made in writing and must include:

(a) a completed authority prescription form; and

(b) a completed Fistulising Crohn Disease PBS Authority Application - Supporting Information Form which includes the following:

(i) a completed current Fistula Assessment Form including the date of assessment of the patient's condition; and

(ii) details of prior biological medicine treatment including details of date and duration of treatment.

The most recent fistula assessment must be no more than 1 month old at the time of application.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under an initial treatment restriction, the patient must have been assessed for response to that course following a minimum of 12 weeks therapy for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

To demonstrate a response to treatment the application must be accompanied by the results of the most recent course of biological medicine therapy within the timeframes specified in the relevant restriction.

If the response assessment to the previous course of biological medicine treatment is not submitted as detailed above, the patient will be deemed to have failed therapy with that particular course of biological medicine.

A maximum quantity and number of repeats to provide for an initial course of this drug consisting of 3 doses at 5 mg per kg body weight per dose to be administered at weeks 0, 2 and 6, will be authorised.

An assessment of the patient's response to this initial course of treatment must be made up to 12 weeks after the first dose (up to 6 weeks following the third dose) so that there is adequate time for a response to be demonstrated.

This assessment, which will be used to determine eligibility for the first continuing treatment, must be submitted to the Department of Human Services no later than 1 month from the date of completion of this initial course of treatment.

Where a response assessment is not undertaken and submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with this drug.

Note It is recommended that an application for the first continuing treatment is submitted to the Department of Human Services at the time of the 12 week assessment, to ensure continuity of treatment for those patients who meet the first continuing treatment criteria for PBS-subsidised treatment with this drug.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Complex refractory Fistulising Crohn disease

Treatment Phase: Balance of supply

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial treatment (new patient or Recommencement of treatment after more than 5 years break in therapy - Initial 1) restriction to complete the 3 doses (the initial infusion regimen at 0, 2 and 6 weeks); OR
- Patient must have received insufficient therapy with this drug for this condition under the Change or Re-commencement of treatment after a break in therapy of less than 5 years (Initial 2) restriction to complete the 3 doses (the initial infusion regimen at 0, 2 and 6 weeks); OR
- Patient must have received insufficient therapy with this drug for this condition under the first continuing treatment or subsequent continuing treatment restrictions to complete 24 weeks of treatment, **AND**
- The treatment must provide no more than the balance of up to 3 doses (Initial 1 or Initial 2 treatment) or 2 repeats (first Continuing or Subsequent Continuing treatment).

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Complex refractory Fistulising Crohn disease

Treatment Phase: First continuing treatment

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug.

An adequate response is defined as:

- (a) a decrease from baseline in the number of open draining fistulae of greater than or equal to 50%; and/or
- (b) a marked reduction in drainage of all fistula(e) from baseline, together with less pain and induration as reported by the patient.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Fistulising Crohn Disease PBS Authority Application - Supporting Information Form which includes a completed Fistula Assessment form including the date of the assessment of the patient's condition.

The most recent fistula assessment must be no more than 1 month old at the time of application.

The application for first continuing treatment with this drug must include an assessment of the patient's response to the initial course of treatment. The assessment must be made up to 12 weeks after the first dose so that there is adequate time for a response to be demonstrated. This assessment must be submitted no later than 4 weeks from the cessation of that treatment course.

Where a response assessment is not undertaken and submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with this drug.

A maximum of 24 weeks of treatment with this drug will be authorised under this restriction.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg eight weekly.

Up to a maximum of 2 repeats will be authorised.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

infliximab 100 mg injection, 1 vial

9654D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	320.71	^a Inflectra [PF] ^a Renflexis [MK]	^a Remicade [JC]

▪ **INFLIXIMAB**

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for adults with severe active rheumatoid arthritis. Where the term biological medicine appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alfa antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor

(tocilizumab), the T-cell co-stimulation modulator (abatacept) and the Janus kinase (JAK) inhibitors (baricitinib, tofacitinib).

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised biological medicine therapy may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once, and
- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised biological medicines for the treatment of rheumatoid arthritis.

Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Stevens Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

For a patient who has failed PBS-subsidised treatment with 2 or 3 TNF-alfa antagonists prior to 1 August 2010 please contact the Department of Human Services on 1800 700 270. A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction.

A patient who has failed fewer than 5 biological medicines and who has a break in treatment of less than 24 months may commence a further course of treatment with a biological medicine under Initial 2 treatment restriction. A patient who has failed fewer than 5 biological medicines and who has had a break in therapy of longer than 24 months may commence a further course of treatment with a biological medicine under the Initial 3 treatment restriction. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine treatment is stopped to the date of the new application for treatment with a biological medicine.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 April 2019.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised biological medicine treatment and wishes to commence such therapy, excluding rituximab (Initial 1 - new patient); or

(ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to re-commence treatment with a specific biological medicine following a break of less than 24 months in PBS-subsidised therapy with that agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months).

(iv) a patient wishes to re-commence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 24 months (Initial 3 - re-commencement of treatment after a break in biological medicine of more than 24 months) Initial applications for a new patient (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6-month intensive DMARD trial, but prior to ceasing DMARD therapy.

Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, baricitinib, etanercept, golimumab, tocilizumab and tofacitinib, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

Rituximab patients should be assessed following a minimum of 12 weeks after the first infusion, and the assessment should be submitted to the Department of Human Services within 4 weeks to ensure continuity of treatment for those who meet the continuing restriction for PBS subsidised treatment with this drug for this condition.

Where a response assessment is not provided the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment. For second and subsequent courses of PBS-subsidised biological medicine (excluding rituximab) treatment it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to the Department of Human Services within sufficient timeframe to allow for authority assessment.

Abatacept patients:

A patient is eligible to receive one I.V. loading dose when commencing treatment with the subcutaneous formulation. Two prescriptions are required, the first prescription for the I.V. loading dose for sufficient vials for one dose based on the patient's weight with no repeats. The second prescription for the subcutaneous formulation, with a maximum quantity of 4 and up to 3 repeats, must be submitted with the initial application.

Rituximab patients:

A further application may be submitted to the Department of Human Services within sufficient timeframe to allow authority assessment. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

Infliximab and etanercept patients:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most

recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction. Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

(2) Swapping therapy

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non- biological medicine therapy requirements except if the patient has had a break in therapy of more than 24 months who would need to requalify with respect to the indices of disease severity. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each biological medicine trialled.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent, unless they have a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug.

Abatacept:

A patient swapping from I.V. abatacept to subcutaneous abatacept will not be eligible for an I.V. loading dose when commencing treatment with the subcutaneous formulation.

Rituximab:

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF-alfa antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological medicine during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological medicine therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate biological medicine may do so without having to have any treatment-free period.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and the Department of Human Services will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints.

Applications under the Initial 1 treatment restriction for a new patient must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy.

Applications under the Initial 3 treatment restriction for re-commencement of treatment after a break in biological medicine of more than 24 months must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial 1 (new patient)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with disease modifying anti-rheumatic drugs (DMARDs) which must include at least 3 months continuous treatment with each of at least 2 DMARDs, one of which must be methotrexate at a dose of at least 20 mg weekly and one of which must be: (i) hydroxychloroquine at a dose of at least 200 mg daily; or (ii) leflunomide at a dose of at least 10 mg daily; or (iii) sulfasalazine at a dose of at least 2 g daily; OR
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs which, if methotrexate is contraindicated according to the Therapeutic Goods Administration (TGA)-approved Product Information or cannot be tolerated at a 20 mg weekly dose, must include at least 3 months continuous treatment with each of at least 2 of the following DMARDs: (i) hydroxychloroquine at a dose of at least 200 mg daily; and/or (ii) leflunomide at a dose of at least 10 mg daily; and/or (iii) sulfasalazine at a dose of at least 2 g daily; OR
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs which, if 3 or more of methotrexate, hydroxychloroquine, leflunomide and sulfasalazine are contraindicated according to the relevant TGA-approved Product Information or cannot be tolerated at the doses specified above, must include at least 3 months continuous treatment

with each of at least 2 DMARDs, with one or more of the following DMARDs being used in place of the DMARDs which are contraindicated or not tolerated: (i) azathioprine at a dose of at least 1 mg/kg per day; and/or (ii) cyclosporin at a dose of at least 2 mg/kg/day; and/or (iii) sodium aurothiomalate at a dose of 50 mg weekly, **AND**

- Patient must not receive more than 22 weeks of treatment under this restriction, **AND**
- The treatment must be given concomitantly with methotrexate at a dose of at least 7.5 mg weekly.

Population criteria:

- Patient must be aged 18 years or older.

If methotrexate is contraindicated according to the TGA-approved product information or cannot be tolerated at a 20 mg weekly dose, the application must include details of the contraindication or intolerance including severity to methotrexate. The maximum tolerated dose of methotrexate must be documented in the application, if applicable.

The application must include details of the DMARDs trialled, their doses and duration of treatment, and all relevant contraindications and/or intolerances including severity.

The requirement to trial at least 2 DMARDs for periods of at least 3 months each can be met using single agents sequentially or by using one or more combinations of DMARDs.

If the requirement to trial 6 months of intensive DMARD therapy with at least 2 DMARDs cannot be met because of contraindications and/or intolerances of a severity necessitating permanent treatment withdrawal to all of the DMARDs specified above, details of the contraindication or intolerance including severity and dose for each DMARD must be provided in the authority application.

The following criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; **AND** either

(a) a total active joint count of at least 20 active (swollen and tender) joints; or

(b) at least 4 active joints from the following list of major joints:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The joint count and ESR and/or CRP must be determined at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. All measures must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials to provide sufficient drug, based on the weight of the patient, for a single infusion at a dose of 3 mg per kg.

Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note Biosimilar preferred prescribing policy

Prescribing of the biosimilar brand Inflectra or Renflexis is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note The Department of Human Services website (www.humanservices.gov.au) has details of the toxicities, including severity, which will be accepted for the following purposes:

(a) exempting a patient from the requirement to undertake a minimum 3 month trial of methotrexate at a 20 mg weekly dose;

(b) substituting azathioprine, cyclosporin or sodium aurothiomalate for another DMARD as part of the 6 month intensive DMARD trial;

(c) exempting a patient from the requirement for a 6 month trial of intensive DMARD therapy.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 2 (change or re-commencement of treatment after a break in biological medicine of less than 24 months)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must not have failed to respond to previous PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised biological medicine treatment for this condition 5 times, **AND**
- Patient must not receive more than 22 weeks of treatment under this restriction, **AND**
- The treatment must be given concomitantly with methotrexate at a dose of at least 7.5 mg weekly.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either of the Initial 1, Initial 2, Initial 3, first or subsequent continuing treatment restrictions, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials to provide sufficient drug, based on the weight of the patient, for a single infusion at a dose of 3 mg per kg.

Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

A patient who has demonstrated a response to a course of rituximab must have a PBS-subsidised biological therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 3 (re-commencement of treatment after a break in biological medicine of more than 24 months)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 24 months or more from the most recent PBS-subsidised biological medicine for this condition, **AND**
- Patient must not have failed to respond to previous PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised biological medicine treatment for this condition 5 times, **AND**
- The condition must have an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; OR
- The condition must have a C-reactive protein (CRP) level greater than 15 mg per L, **AND**
- The condition must have either (a) a total active joint count of at least 20 active (swollen and tender) joints; or (b) at least 4 active major joints, **AND**
- Patient must not receive more than 22 weeks of treatment under this restriction, **AND**
- The treatment must be given concomitantly with methotrexate at a dose of at least 7.5 mg weekly.

Population criteria:

- Patient must be aged 18 years or older.

Major joints are defined as (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

All measures of joint count and ESR and/or CRP must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials to provide sufficient drug, based on the weight of the patient, for a single infusion at a dose of 3 mg per kg.

Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note Biosimilar preferred prescribing policy

Prescribing of the biosimilar brand Inflectra or Renflexis is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial 1 (new patient) or Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 24 months) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 24 months) - balance of supply

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 22 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 24 months) restriction to complete 22 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 24 months) to complete 22 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 22 weeks treatment available under the above restrictions.

Population criteria:

- Patient must be aged 18 years or older.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Severe active rheumatoid arthritis

Treatment Phase: First continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction, **AND**
- The treatment must be given concomitantly with methotrexate at a dose of at least 7.5 mg weekly.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials to provide sufficient drug, based on the weight of the patient, for a single infusion at a dose of 3 mg per kg.

Up to a maximum of 2 repeats will be authorised.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient has either failed or ceased to respond to a PBS-subsidised biological medicine for this condition 5 times, they will not be eligible to receive further PBS-subsidised treatment with a biological medicine for this condition.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Continuing Treatment - balance of supply.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the first continuing treatment restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the subsequent continuing Authority Required (in writing) treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restrictions.

Population criteria:

- Patient must be aged 18 years or older.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

infliximab 100 mg injection, 1 vial

5757B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	320.71	^a Inflectra [PF] ^a Renflexis [MK]	^a Remicade [JC]

■ INFLIXIMAB
Note TREATMENT OF ADULT PATIENTS WITH ANKYLOSING SPONDYLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, certolizumab pegol, etanercept, golimumab, infliximab and secukinumab for adult patients with ankylosing spondylitis. Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab and secukinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 6 biological medicines at any 1 time.

Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last prescription for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised biological medicine treatment with adalimumab, certolizumab pegol, etanercept, golimumab, infliximab and secukinumab.

(a) Initial treatment.

Applications for initial treatment should be made where:

- a patient has received no prior PBS-subsidised biological medicine treatment in this treatment cycle and wishes to commence such therapy (Initial 1 - New patient)
- a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or
- a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years); or
- a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

A patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy.

(b) Continuing treatment.

For the first continuing treatment course of PBS-subsidised biological medicine, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1, Initial 2 or Initial 3 treatment restrictions. For second and subsequent continuing courses of PBS-subsidised biological medicine it is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply. Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be conducted within 1 month of the last dose. Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI), or the prior NSAID

therapy and exercise program requirements.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the BASDAI, ESR and/or CRP submitted with the first authority application for a biological medicine.

However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to these revised baseline measurements.

For a new patient, the BASDAI used to determine the baseline must be measured while the patient is receiving NSAID therapy and completing their exercise program.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response.

(4) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must qualify under the Initial 3 treatment restriction. The same clinical criteria and indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI) as for the Initial 1 (New patient) restriction will need to be met, but a re-trial of NSAID therapy and exercise therapy is not required.

Authority required

Ankylosing spondylitis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Clinical criteria:

- The condition must be radiographically (plain X-ray) confirmed Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have at least 2 of the following: (i) low back pain and stiffness for 3 or more months that is relieved by exercise but not by rest; or (ii) limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by a score of at least 1 on each of the lumbar flexion and lumbar side flexion measurements of the Bath Ankylosing Spondylitis Metrology Index (BASMI); or (iii) limitation of chest expansion relative to normal values for age and gender,

AND

- Patient must have failed to achieve an adequate response following treatment with at least 2 non-steroidal anti-inflammatory drugs (NSAIDs), whilst completing an appropriate exercise program, for a total period of 3 months, **AND**
- Patient must not receive more than 18 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

The application must include details of the NSAIDs trialled, their doses and duration of treatment.

If the NSAID dose is less than the maximum recommended dose in the relevant TGA-approved Product Information, the application must include the reason a higher dose cannot be used.

If treatment with NSAIDs is contraindicated according to the relevant TGA-approved Product Information, the application must provide details of the contraindication.

If intolerance to NSAID treatment develops during the relevant period of use which is of a severity to necessitate permanent treatment withdrawal, the application must provide details of the nature and severity of this intolerance.

The following criteria indicate failure to achieve an adequate response and must be demonstrated at the time of the initial application:

- (a) a Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 4 on a 0-10 scale; AND
- (b) an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 10 mg per L.

The BASDAI must be determined at the completion of the 3 month NSAID and exercise trial, but prior to ceasing NSAID treatment. The BASDAI must be no more than 1 month old at the time of initial application.

Both ESR and CRP measures should be provided with the initial treatment application and both must be no more than 1 month old. If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reason this criterion cannot be satisfied.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Ankylosing Spondylitis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) a copy of the radiological report confirming Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis; and
 - (ii) a completed BASDAI Assessment Form; and

(iii) a completed Exercise Program Self Certification Form included in the supporting information form.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg.

A maximum quantity and number of repeats to provide for an initial course of this drug consisting of 3 doses at 5 mg per kg body weight per dose to be administered at weeks 0, 2 and 6, will be authorised.

Up to a maximum of 3 repeats will be authorised.

An assessment of a patient's response to an initial course of treatment must be conducted following a minimum of 12 weeks of therapy. An application for the continuing treatment must be accompanied with the assessment of response and submitted to the Department of Human Services no later than 4 weeks from the date of completion of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Details of the toxicities, including severity, which will be accepted for the purposes of administering this restriction can be found on the Department of Human Services website at www.humanservices.gov.au

Note For details on the appropriate minimum exercise program that will be accepted for the purposes of administering this restriction, please refer to the Department of Human Services website at www.humanservices.gov.au

Note Biosimilar preferred prescribing policy

Prescribing of the biosimilar brand Inflectra or Renflexis is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Ankylosing spondylitis

Treatment Phase: Initial treatment - Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- Patient must not receive more than 18 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Ankylosing Spondylitis PBS Authority Application - Supporting Information Form.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg.

Up to a maximum of 3 repeats will be authorised.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to change or recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

An adequate response is defined as an improvement from baseline of at least 2 of the BASDAI and 1 of the following:

- (a) an ESR measurement no greater than 25 mm per hour; or

- (b) a CRP measurement no greater than 10 mg per L; or
(c) an ESR or CRP measurement reduced by at least 20% from baseline.

Where only 1 acute phase reactant measurement is supplied in the first application for PBS-subsidised treatment, that same marker must be measured and supplied in all subsequent continuing treatment applications.

All measurements provided must be no more than 1 month old at the time of application.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Ankylosing spondylitis

Treatment Phase: Initial treatment - Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must be radiographically (plain X-ray) confirmed Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis, **AND**
- Patient must have at least 2 of the following: (i) low back pain and stiffness for 3 or more months that is relieved by exercise but not by rest; or (ii) limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by a score of at least 1 on each of the lumbar flexion and lumbar side flexion measurements of the Bath Ankylosing Spondylitis Metrology Index (BASMI); or (iii) limitation of chest expansion relative to normal values for age and gender,

AND

- Patient must have a Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 4 on a 0-10 scale that is no more than 4 weeks old at the time of application, **AND**
- Patient must have an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour that is no more than 4 weeks old at the time of application; OR
- Patient must have a C-reactive protein (CRP) level greater than 10 mg per L that is no more than 4 weeks old at the time of application; OR
- Patient must have a clinical reason as to why demonstration of an elevated ESR or CRP cannot be met and the application must state the reason, **AND**
- Patient must not receive more than 18 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

The authority application must be made in writing and must include:

(a) a completed authority prescription form; and

(b) a completed Ankylosing Spondylitis PBS Authority Application - Supporting Information Form which includes the following:

- (i) a copy of the radiological report confirming Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis; and
- (ii) a completed BASDAI Assessment Form.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg.

A maximum quantity and number of repeats to provide for an initial course of this drug consisting of 3 doses at 5 mg per kg body weight per dose to be administered at weeks 0, 2 and 6, will be authorised.

Up to a maximum of 3 repeats will be authorised.

An assessment of a patient's response to an initial course of treatment must be conducted following a minimum of 12 weeks of therapy. An application for the continuing treatment must be accompanied with the assessment of response and submitted to the Department of Human Services no later than 4 weeks from the date of completion of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Biosimilar preferred prescribing policy

Prescribing of the biosimilar brand Inflectra or Renflexis is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Ankylosing spondylitis

Treatment Phase: Initial treatment - Initial 1 (new patient), Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years) - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the Initial 1 (new patient) restriction to complete 18 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 18 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 18 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 18 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Ankylosing spondylitis

Treatment Phase: First continuing treatment

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Ankylosing Spondylitis PBS Authority Application - Supporting Information Form.

An adequate response is defined as an improvement from baseline of at least 2 of the BASDAI and 1 of the following:

- (a) an ESR measurement no greater than 25 mm per hour; or
- (b) a CRP measurement no greater than 10 mg per L; or
- (c) an ESR or CRP measurement reduced by at least 20% from baseline.

Where only 1 acute phase reactant measurement is supplied in the first application for PBS-subsidised treatment, that same marker must be measured and supplied in all subsequent continuing treatment applications.

All measurements provided must be no more than 1 month old at the time of application.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg.

Up to a maximum of 3 repeats will be authorised.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Ankylosing spondylitis

Treatment Phase: Continuing treatment - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the first continuing treatment restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the subsequent continuing Authority Required (in writing) treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

infliximab 100 mg injection, 1 vial

5753T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	320.71	^a Inflectra [PF] ^a Renflexis [MK]	^a Remicade [JC]

■ INFLIXIMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE PSORIATIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab for adult patients with severe active psoriatic arthritis. Therefore, where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient receiving PBS-subsidised treatment for psoriatic arthritis is able to commence a treatment cycle where they may trial biological medicines without having to experience a disease flare when swapping to the alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

A patient who received PBS-subsidised adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab or ustekinumab treatment prior to 1 May 2019 is considered to start their first cycle as of 1 May 2019.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence another cycle [further details are under '(5) Recommencement of treatment after a 5-year break in PBS-subsidised therapy' below].

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe active psoriatic arthritis.

(1) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy

(Initial 1 - New patient); or

(ii) a patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an alternate medicine (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years) or An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, golimumab secukinumab and tofacitinib, 18 to 20 weeks of therapy for certolizumab pegol (depending upon the dosing regimen), 20 weeks of therapy for ixekizumab, 22 weeks of therapy for infliximab, and 28 weeks of therapy for ustekinumab. It is recommended that a patient be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of PBS-subsidised initial treatment following a minimum of 12 weeks of therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

Grandfather patients (ixekizumab only).

A patient who commenced treatment with ixekizumab for severe psoriatic arthritis prior to 1 March 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

Grandfather patients (tofacitinib only).

A patient who commenced treatment with Tofacitinib for severe psoriatic arthritis prior to 1 May 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment under the Initial 1 or Initial 2 treatment restrictions.

For the second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to re-qualify with respect to either the indices of disease severity (i.e. erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP) level, and active joint count) or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify under the Initial 3 treatment restriction with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application.

However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

Within a treatment cycle a patient may alternate between therapy with any biological medicine of their choice (1 at a time) providing:

(i) they have not received PBS-subsidised treatment with that particular biological medicine previously; or

(ii) they have demonstrated an adequate response to that particular biological medicine if they have previously trialled it on the PBS; and

(iii) they have not previously failed to respond to treatment 3 times in this treatment cycle with that biological medicine.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(4) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the indices of disease severity submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and these revised baseline measurements will be used to assess response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to determine response for all subsequent continuing treatments. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. 20 or more active joints), response will be determined according to a reduction in the total number of active joints. (5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment restriction according to the criteria of the relevant restriction and index of disease severity. The application must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response to methotrexate at a dose of at least 20 mg weekly for a minimum period of 3 months, **AND**
- Patient must have failed to achieve an adequate response to sulfasalazine at a dose of at least 2 g per day for a minimum period of 3 months; OR
- Patient must have failed to achieve an adequate response to leflunomide at a dose of up to 20 mg daily for a minimum period of 3 months, **AND**
- Patient must not receive more than 22 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Where treatment with methotrexate, sulfasalazine or leflunomide is contraindicated according to the relevant TGA-approved Product Information, details must be provided at the time of application.

Where intolerance to treatment with methotrexate, sulfasalazine or leflunomide developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

The following initiation criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; and

either

- (a) an active joint count of at least 20 active (swollen and tender) joints; or
- (b) at least 4 active joints from the following list of major joints:
 - (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg.

Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Severe Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

An assessment of a patient's response to an initial course of treatment must be conducted following a minimum of 12 weeks of therapy. An application for the continuing treatment must be accompanied with the assessment of response and submitted to the Department of Human Services no later than 4 weeks from the date of completion of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Biosimilar preferred prescribing policy

Prescribing of the biosimilar brand Inflectra or Renflexis is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 3 months treatment with methotrexate and 3 months treatment with sulfasalazine or leflunomide can be found on the Department of Human Services website (www.humanservices.gov.au)

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial treatment - Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- Patient must not receive more than 22 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an erythrocyte sedimentation rate (ESR) no greater than 25 mm per hour or a C-reactive protein (CRP) level no greater than 15 mg per L or either marker reduced by at least 20% from baseline; and

either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following major active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg.

Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Severe Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to change or recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3, first or subsequent continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Biosimilar preferred prescribing policy

Prescribing of the biosimilar brand Inflectra or Renflexis is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial treatment - Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must have an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; OR
- The condition must have a C-reactive protein (CRP) level greater than 15 mg per L, **AND**
- The condition must have either (a) a total active joint count of at least 20 active (swollen and tender) joints; or (b) at least 4 active major joints, **AND**
- Patient must not receive more than 22 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Major joints are defined as (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

All measures of joint count and ESR and/or CRP must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg.

Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Severe Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3, first or subsequent continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Biosimilar preferred prescribing policy

Prescribing of the biosimilar brand Inflectra or Renflexis is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial 1 (new patient) or Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years) - balance of supply

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 22 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 22 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 22 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 22 weeks treatment available under the above restrictions.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Severe psoriatic arthritis

Treatment Phase: First continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an erythrocyte sedimentation rate (ESR) no greater than 25 mm per hour or a C-reactive protein (CRP) level no greater than 15 mg per L or either marker reduced by at least 20% from baseline; and

either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following major active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to determine response for all subsequent continuing treatments.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg eight weekly.

Up to a maximum of 2 repeats will be authorised.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Severe Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either Initial 1, Initial 2, or Initial 3 treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Continuing treatment - balance of supply

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the first continuing treatment restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the subsequent continuing Authority Required (in writing) treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restrictions.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

infliximab 100 mg injection, 1 vial

5756Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	320.71	^a Inflectra [PF] ^a Renflexis [MK]	^a Remicade [JC]

■ INFLIXIMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, etanercept, guselkumab, infliximab, ixekizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, etanercept, guselkumab, infliximab, ixekizumab, secukinumab, tildrakizumab and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, etanercept, guselkumab, infliximab, ixekizumab, secukinumab, tildrakizumab and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patients receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the

new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years); or

(iii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or

(iv) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab and 28 weeks of therapy for tildrakizumab and ustekinumab.

It is recommended that a patient be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

Grandfather patients (guselkumab only).

A patient who commenced treatment with guselkumab for chronic plaque psoriasis prior to (1 February 2019) and who continues to receive treatment at the time of application, may qualify for treatment under the Initial treatment - Grandfather treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

Grandfather patients (tildrakizumab only).

A patient who commenced treatment with tildrakizumab for chronic plaque psoriasis prior to (1 February 2019) and who continues to receive treatment at the time of application, may qualify for treatment under the Initial treatment Grandfather treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment. This assessment must be conducted within 1 month of the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle. To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 the PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 1 month at the time of application.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial 1 - Whole body (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis where lesions have been present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 4 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; and/or (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; and/or (iii) cyclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; and/or (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 22 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, cyclosporin or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application.

Where intolerance to treatment with phototherapy, methotrexate, cyclosporin or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, cyclosporin, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

- (a) A current Psoriasis Area and Severity Index (PASI) score of greater than 15, as assessed, preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior treatment.
- (b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 1 month following cessation of each course of treatment.
- (c) The most recent PASI assessment must be no more than 1 month old at the time of application.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg. Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and
 - (ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition. Demonstration of response should be provided within this timeframe.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Biosimilar preferred prescribing policy

Prescribing of the biosimilar brand Inflectra or Renflexis is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, cyclosporin or acitretin can be found on the Department of Human Services website (www.humanservices.gov.au)

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Whole body (change or recommencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised therapy with this drug for this condition in the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 22 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either of the Initial 1, Initial 2, Initial 3, first or subsequent continuing treatment restrictions, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition. Demonstration of response should be provided within this timeframe.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg. Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and
 - (ii) details of prior biological treatment, including dosage, date and duration of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Biosimilar preferred prescribing policy

Prescribing of the biosimilar brand Inflectra or Renflexis is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 3, Whole body (re-commencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must have a current Psoriasis Area and Severity Index (PASI) score of greater than 15, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 22 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

The most recent PASI assessment must be no more than 1 month old at the time of application.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg. Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition.

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition. Demonstration of response should be provided within this timeframe.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Biosimilar preferred prescribing policy

Prescribing of the biosimilar brand Inflectra or Renflexis is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Face, hand, foot (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 4 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; and/or (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; and/or (iii) cyclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; and/or (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 22 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, cyclosporin or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application.

Where intolerance to treatment with phototherapy, methotrexate, cyclosporin or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, cyclosporin, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

(a) Chronic plaque psoriasis classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:

(i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe, as assessed, preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior treatment; or

(ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed, preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior treatment;

(b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 1 month following cessation of each course of treatment.

(c) The most recent PASI assessment must be no more than 1 month old at the time of application.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg. Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition; and

(ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition. Demonstration of response should be provided within this timeframe.

The PASI assessment for first continuing or subsequent continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Biosimilar preferred prescribing policy

Prescribing of the biosimilar brand Inflectra or Renflexis is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, cyclosporin or acitretin can be found on the Department of Human Services website (www.humanservices.gov.au)

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Face, hand, foot (change or recommencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised therapy with this drug for this condition in the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 22 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or
- (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either of the Initial 1, Initial 2, Initial 3, first or subsequent continuing treatment restrictions, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition. Demonstration of response should be provided within this timeframe. The PASI assessment for first continuing or subsequent continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg. Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition; and
 - (ii) details of prior biological treatment, including dosage, date and duration of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Biosimilar preferred prescribing policy

Prescribing of the biosimilar brand Inflectra or Renflexis is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 3, Face, hand, foot (re-commencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must be classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:
 - (i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or
 - (ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 22 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

The most recent PASI assessment must be no more than 1 month old at the time of application.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg. Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition.

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition. Demonstration of response should be provided within this timeframe.

The PASI assessment for first continuing or subsequent continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Biosimilar preferred prescribing policy

Prescribing of the biosimilar brand Inflectra or Renflexis is encouraged for treatment naive patients.

Note Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial 1, Whole body or Face, hand, foot (new patient) or Initial 2, Whole body or Face, hand, foot (change or recommencement of treatment after a break in biological medicine of less than 5 years) or Initial 3, Whole body or Face, hand, foot (re-commencement of treatment after a break in biological medicine of more than 5 years) - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1, Whole body (new patient) restriction to complete 22 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2, Whole body (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 22 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3, Whole body (re-commencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 22 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 1, Face, hand, foot (new patient) restriction to complete 22 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2, Face, hand, foot (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 22 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3, Face, hand, foot (re-commencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 22 weeks treatment, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- The treatment must provide no more than the balance of up to 22 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a dermatologist.

Note Authority approval for sufficient therapy to complete a maximum of 22 weeks of treatment may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Severe chronic plaque psoriasis

Treatment Phase: First continuing treatment, Whole body

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg eight weekly. Up to a maximum of 2 repeats will be authorised.

The authority application must be made in writing and must include:

- a completed authority prescription form(s); and
- a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available

on the Department of Human Services website at www.humanservices.gov.au
Applications for authority to prescribe should be forwarded to:
Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: First continuing treatment, Face, hand, foot

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or
- (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

At the time of the authority application, medical practitioners should request the appropriate quantity of vials, based on the weight of the patient, to provide for infusions at a dose of 5 mg per kg eight weekly. Up to a maximum of 2 repeats will be authorised.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet and face, hand, foot area diagrams including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

The PASI assessment for first continuing or subsequent continuing treatment must be performed on the same affected area assessed at baseline.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Whole body or Continuing treatment, Face, hand, foot - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the first continuing treatment, Whole body restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the first continuing treatment, Face, hand, foot restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the subsequent continuing treatment Authority Required (in writing), Whole body restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the subsequent continuing treatment Authority Required (in writing), Face, hand, foot restriction to complete 24 weeks treatment, **AND**

- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restrictions, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate).

Treatment criteria:

- Must be treated by a dermatologist.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

infliximab 100 mg injection, 1 vial

5758C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	320.71	^a Inflectra [PF] ^a Renflexis [MK]	^a Remicade [JC]

Interleukin inhibitors

▪ **ANAKINRA**

Note This drug is not PBS-subsidised for conditions other than CAPS.

Authority required (STREAMLINED)

5450

Moderate to severe cryopyrin associated periodic syndromes (CAPS)

Treatment criteria:

- Must be treated by a rheumatologist or in consultation with a rheumatologist; OR
- Must be treated by a clinical immunologist or in consultation with a clinical immunologist.

A diagnosis of CAPS must be documented in the patient's medical records.

anakinra 100 mg/0.67 mL injection, 28 x 0.67 mL syringes

10264F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	1485.00	Kineret [FK]

▪ **TOCILIZUMAB**

Note TREATMENT OF PATIENTS WITH SEVERE ACTIVE SYSTEMIC JUVENILE IDIOPATHIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of tocilizumab for a patient who has severe active systemic juvenile idiopathic arthritis (sJIA).

From 1 May 2012, a patient receiving PBS-subsidised tocilizumab therapy is considered to be in a treatment cycle. Under these arrangements, within a single treatment cycle, a patient may:

- continue to receive long-term treatment with PBS-subsidised tocilizumab while they continue to show a response to therapy, and
- fail to respond, or to sustain a response, to PBS-subsidised tocilizumab twice.

Once a patient has either failed or ceased to sustain a response to 2 courses of treatment, they are deemed to have completed a single treatment cycle and they must have, at a minimum, a 12 month break in PBS-subsidised tocilizumab therapy before they are eligible to commence another cycle. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised tocilizumab treatment was approved to the date of the first application for initial treatment with tocilizumab under the new treatment cycle.

Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient who received PBS-subsidised tocilizumab treatment immediately prior to 1 May 2012 is considered to be in their first cycle as of 1 May 2012. A patient who has had a break in tocilizumab treatment of at least 12 months immediately prior to making a new application, on or after 1 May 2012, will commence a new treatment cycle. A patient who has failed their first course of tocilizumab in a treatment cycle and who has a break in therapy of less than 12 months may commence a second course of treatment within the same treatment cycle under the Initial 2 treatment restriction.

A patient who has failed their first course of tocilizumab in a treatment cycle and who has a break in therapy of more than 12 months must commence a new treatment cycle under the Initial 3 treatment restriction.

How to prescribe PBS-subsidised tocilizumab treatment therapy after 1 May 2012.

(1) Initial treatment.

Applications for initial treatment should be made where:

- a patient has not received prior PBS-subsidised tocilizumab treatment and wishes to commence such therapy (Initial 1 - New patient); or
- a patient has received the first course of PBS-subsidised (initial or continuing) tocilizumab therapy in a treatment cycle and is deemed to have failed to respond or sustain a response and the treating physician wishes to trial a second course, provided any break in therapy is less than 12 months (Initial 2 - retreat or recommencement of treatment after a break of less than 12 months); or
- a patient wishes to recommence treatment with tocilizumab following a break in PBS-subsidised therapy of less than 12 months (Initial 2 - retreat or recommencement of treatment after a break of less than 12 months).
- a patient wishes to recommence treatment following a break in PBS-subsidised tocilizumab therapy of more than 12 months (Initial 3 - recommencement of treatment after a break of more than 12 months).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy.

A patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not conducted within these timeframes, the patient will be deemed to have failed this course of treatment.

For the second and subsequent courses of PBS-subsidised tocilizumab, it is recommended that a patient is reviewed in the 4 weeks prior to completing their current course of treatment and that an application is posted to the Department of Human

Services no later than 2 weeks prior to the patient completing their current treatment course.

(2) Continuing treatment.

Following the completion of an initial treatment course with tocilizumab, a patient may qualify to receive up to 24 weeks of continuing treatment with this drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing tocilizumab treatment in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted tocilizumab supply.

Assessments of response to a course of PBS-subsidised therapy must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased. Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with tocilizumab.

(2) Treatment cycle.

Once initial treatment with PBS-subsidised tocilizumab is approved, a patient deemed to have failed to respond to the first course of treatment may have a second course without having to requalify with respect to the indices of disease severity (joint count, fever and/or CRP level and platelet count), except if the patient has had a break in therapy of more than 12 months.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the relevant baseline measurements of the joint count, fever and/or CRP level and platelet count submitted with the first authority application for tocilizumab.

Where a patient is deemed to have failed to respond or to sustain a response to the first course of therapy in a treatment cycle, prescribers may provide new baseline measurements for the second course of treatment within that cycle. The Department of Human Services will assess response according to these revised baseline measurements. If new baseline measurements are not submitted with the initial application for the second course of treatment, then those submitted with the first course will be used by the Department of Human Services to assess response to the second course.

(4) Recommencement of treatment after a 12 month break in PBS-subsidised therapy.

A patient who wishes to start a second or subsequent treatment cycle following a break in PBS-subsidised tocilizumab therapy of at least 12 months, must requalify for treatment under the Initial 3 treatment restriction.

(5) Withdrawal of treatment after sustained remission.

Withdrawal of treatment with tocilizumab should be considered in a patient who has achieved and sustained complete remission of disease for 12 months. A demonstration of response to the current treatment should be submitted to the Department of Human Services at the time treatment is ceased.

Authority required

Systemic juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Clinical criteria:

- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have polyarticular course disease which has failed to respond adequately to oral or parenteral methotrexate at a dose of at least 15 mg per square metre weekly, alone or in combination with oral or intra-articular corticosteroids, for a minimum of 3 months; OR
- Patient must have polyarticular course disease and have demonstrated severe intolerance of, or toxicity due to, methotrexate; OR
- Patient must have refractory systemic symptoms, demonstrated by an inability to decrease and maintain the dose of prednisolone (or equivalent) below 0.5 mg per kg per day following a minimum of 2 months of therapy, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be under 18 years of age.

Treatment criteria:

- Must be treated by a rheumatologist; OR
 - Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.
- The following criteria indicate failure to achieve an adequate response to prior methotrexate therapy in a patient with polyarticular course disease and must be demonstrated in the patient at the time of the initial application:
- (a) an active joint count of at least 20 active (swollen and tender) joints; or
 - (b) at least 4 active joints from the following list of major joints:
 - (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - (ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The following criteria indicate failure to achieve an adequate response to prior therapy in a patient with refractory systemic symptoms and must be demonstrated in the patient at the time of the initial application:

- (a) an active joint count of at least 2 active joints; and
- (b) persistent fever greater than 38 degrees Celsius for at least 5 out of 14 consecutive days; and/or
- (c) a C-reactive protein (CRP) level and platelet count above the upper limits of normal (ULN).

The baseline measurements of joint count, fever and/or CRP level and platelet count must be performed preferably whilst on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment.

The same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to determine response for all subsequent continuing treatments.

Severe intolerance to methotrexate is defined as intractable nausea and vomiting and general malaise unresponsive to manoeuvres, including reducing or omitting concomitant non-steroidal anti-inflammatory drugs (NSAIDs) on the day of methotrexate administration, use of folic acid supplementation, or administering the dose of methotrexate in 2 divided doses over 24 hours.

Toxicity due to methotrexate is defined as evidence of hepatotoxicity with repeated elevations of transaminases, bone marrow suppression temporally related to methotrexate use, pneumonitis, or serious sepsis.

If treatment with methotrexate alone or in combination with other treatments is contraindicated according to the relevant TGA-approved Product Information, details must be provided at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, details of this toxicity must be provided at the time of application.

The authority application must be made in writing and must include:

- (1) completed authority prescription form(s); and
- (2) a completed Systemic Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form which includes the following:

- (i) the date of assessment of severe active systemic juvenile idiopathic arthritis;
- (ii) details of prior treatment including dose and duration of treatment;
- (iii) pathology reports detailing CRP and platelet count where appropriate.

At the time of authority application, the medical practitioner must request the appropriate number of vials of appropriate strength to provide sufficient drug, based on the weight of the patient, for two infusions (one month's supply). A separate authority prescription form must be completed for each strength requested. Up to a maximum of 3 repeats will be authorised.

An assessment of a patient's response to an initial course of treatment must be conducted following a minimum of 12 weeks of therapy. An application for the continuing treatment must be accompanied with the assessment of response and submitted to the Department of Human Services no later than 4 weeks from the date of completion of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Systemic juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 2 (retial or recommencement of treatment after a break of less than 12 months)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with this drug for this condition in the previous 12 months, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug more than once during the current treatment cycle, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be under 18 years of age.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

An adequate response to treatment is defined as:

(a) in a patient with polyarticular course disease:

(i) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(ii) a reduction in the number of the following major active joints, from at least 4, by at least 50%:

- elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

- shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

(b) in a patient with refractory systemic symptoms:

(i) absence of fever greater than 38 degrees Celsius in the preceding seven days; and/or

(ii) a reduction in the C-reactive protein (CRP) level and platelet count by at least 30% from baseline; and/or

(iii) a reduction in the dose of corticosteroid by at least 30% from baseline.

At the time of authority application, the medical practitioner must request the appropriate number of vials of appropriate strength to provide sufficient drug, based on the weight of the patient, for two infusions (one month's supply). A separate authority prescription form must be completed for each strength requested. Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

(1) completed authority prescription form(s); and

(2) a completed Systemic Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form which includes pathology reports detailing C-reactive protein (CRP) level and platelet count where appropriate.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to re-trial or recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to 2 courses of treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 12 months have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Systemic juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 3 (recommencement of treatment after a break of more than 12 months)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have had a break in treatment of 12 months or more from this drug for this condition, **AND**
- Patient must have polyarticular course disease and the condition must have (a) an active joint count of at least 20 active (swollen and tender) joints; or (b) at least 4 active joints from the following list of major joints: i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or (ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth); OR
- Patient must have refractory systemic symptoms and the condition must have (a) an active joint count of at least 2 active joints; and (b) persistent fever greater than 38 degrees Celsius for at least 5 out of 14 consecutive days; and/or (c) a C-reactive protein (CRP) level and platelet count above the upper limits of normal (ULN), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Population criteria:

- Patient must be under 18 years of age.

The authority application must be made in writing and must include:

(1) completed authority prescription form(s); and

(2) a completed Systemic Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form which includes the following:

(i) the date of assessment of severe active systemic juvenile idiopathic arthritis;

(ii) pathology reports detailing C-reactive protein (CRP) level and platelet count where appropriate.

The most recent systemic juvenile idiopathic arthritis assessment must be no more than 4 weeks old at the time of application.

At the time of authority application, the medical practitioner must request the appropriate number of vials of appropriate strength to provide sufficient drug, based on the weight of the patient, for two infusions (one month's supply). A separate authority prescription form must be completed for each strength requested. Up to a maximum of 3 repeats will be authorised.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a

minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to 2 courses of treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Systemic juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient) or Initial 2 (retiral or recommencement of treatment after a break of less than 12 months) or Initial 3 (recommencement of treatment after a break of more than 12 months) - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (retiral or recommencement of treatment after a break of less than 12 months) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under Initial 3 (recommencement of treatment after a break of more than 12 months) restriction to complete 16 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Systemic juvenile idiopathic arthritis

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

An adequate response to treatment is defined as:

(a) in a patient with polyarticular course disease:

(i) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(ii) a reduction in the number of the following major active joints, from at least 4, by at least 50%:

- elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

- shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

(b) in a patient with refractory systemic symptoms:

(i) absence of fever greater than 38 degrees Celsius in the preceding seven days; and/or

(ii) a reduction in the C-reactive protein (CRP) level and platelet count by at least 30% from baseline; and/or

(iii) a reduction in the dose of corticosteroid by at least 30% from baseline.

Determination of whether a response has been demonstrated to initial and subsequent courses of treatment will be based on the baseline measurements of disease severity submitted with the initial treatment application.

The authority application must be made in writing and must include:

(1) completed authority prescription form(s); and

(2) a completed Systemic Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form which includes baseline and current pathology reports detailing CRP and platelet count where appropriate.

The most recent systemic juvenile idiopathic arthritis assessment must be no more than 1 month old at the time of application.

At the time of authority application, the medical practitioner must request the appropriate number of vials of appropriate strength to provide sufficient drug, based on the weight of the patient, for two infusions (one month's supply). A separate authority prescription form must be completed for each strength requested. Up to a maximum of 5 repeats will be authorised. Where the most recent course of PBS-subsidised treatment with this drug was approved under either Initial 1, Initial 2, or Initial 3 treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to 2 courses of treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 12 months have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Systemic juvenile idiopathic arthritis

Treatment Phase: Continuing treatment - balance of supply

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restriction.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

tocilizumab 200 mg/10 mL injection, 10 mL vial

1481Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	203.73	Actemra [RO]

tocilizumab 80 mg/4 mL injection, 4 mL vial

1476Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	82.19	Actemra [RO]

tocilizumab 400 mg/20 mL injection, 20 mL vial

1482B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	405.39	Actemra [RO]

■ TOCILIZUMAB

Note TREATMENT OF PATIENTS WITH SEVERE ACTIVE JUVENILE IDIOPATHIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, etanercept and tocilizumab for a patient who has severe active juvenile idiopathic arthritis. Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, etanercept and tocilizumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 3 biological medicines at any one time.

From 1 April 2014, a patient receiving PBS-subsidised biological medicine is considered to be in a treatment cycle where they may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements, within a single treatment cycle, a patient may:

(i) continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy; and

(ii) fail to respond or to sustain a response to each PBS-subsidised biological medicine once only. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single treatment cycle and they must have, at a minimum, a 12 month break in PBS-subsidised biological medicine therapy before they are eligible to commence another cycle.

The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine was approved to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who received PBS-subsidised biological medicine immediately prior to 1 April 2014 is considered to be in their first cycle as of 1 April 2014. A patient who has had a break in biological medicine treatment of at least 12 months immediately prior to making a new application, on or after 1 April 2014, will commence a new treatment cycle under the Initial 3 treatment restriction.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of less than 12 months may commence a further course of treatment within the same treatment cycle under the Initial 2 treatment restriction.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of more than 12 months must commence a new treatment cycle under the Initial 3 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment therapy after 1 April 2014.

(1) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an alternate medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine therapy of less than 12 months) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 12 months with the same medicine (initial 2 - Change or Recommencement of treatment after a break in biological medicine therapy of less than 12 months).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 12 months (initial 3 - Recommencement of treatment after a break in biological medicine of more than 12 months). Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy.

A patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not conducted within these timeframes, the patient will be deemed to have failed this course of treatment.

For the second and subsequent cycles of PBS-subsidised biological medicine, it is recommended that a patient is reviewed in the 4 weeks prior to completing their current course of treatment and that an application is posted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services where applicable. Where a response assessment is not submitted, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (joint count) or the prior non-biological medicine therapy requirements, except if the patient has had a break in therapy of more than 12 months who would then need to requalify under the initial 3 restrictions with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application.

However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, an application for a patient who wishes to swap to an alternate biological medicine should be accompanied by the approved authority prescription or remaining repeats for the biological medicine the patient is ceasing.

(4) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count submitted with the first authority application for a biological medicine. However, prescribers may provide a new baseline measurement any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to the revised baseline measurement.

(5) Recommencement of treatment after a 12 months break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological medicine therapy of at least 12 months, must qualify under the initial 3 restriction and meet the relevant criteria and index of disease severity.

(6) Withdrawal of treatment after sustained remission.

Withdrawal of treatment with biological medicine should be considered in a patient who has achieved and sustained complete remission of disease for 12 months. A demonstration of response to the current treatment should be submitted to the Department of Human Services at the time treatment is ceased.

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Treatment criteria:

- Must be treated by a paediatric rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Clinical criteria:

- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have demonstrated severe intolerance of, or toxicity due to, methotrexate; OR
- Patient must have demonstrated failure to achieve an adequate response to 1 or more of the following treatment regimens: (i) oral or parenteral methotrexate at a dose of at least 20 mg per square metre weekly, alone or in combination with oral or intra-articular corticosteroids, for a minimum of 3 months; or (ii) oral methotrexate at a dose of at least 10 mg per square metre weekly together with at least 1 other disease modifying anti-rheumatic drug (DMARD), alone or in combination with corticosteroids, for a minimum of 3 months, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be under 18 years of age.

Severe intolerance to methotrexate is defined as intractable nausea and vomiting and general malaise unresponsive to manoeuvres, including reducing or omitting concomitant non-steroidal anti-inflammatory drugs (NSAIDs) on the day of methotrexate administration, use of folic acid supplementation, or administering the dose of methotrexate in 2 divided doses over 24 hours.

Toxicity due to methotrexate is defined as evidence of hepatotoxicity with repeated elevations of transaminases, bone marrow suppression temporally related to methotrexate use, pneumonitis, or serious sepsis.

If treatment with methotrexate alone or in combination with another DMARD is contraindicated according to the relevant TGA-approved Product Information, details must be provided at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, details of this toxicity must be provided at the time of application.

The following criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

- (a) an active joint count of at least 20 active (swollen and tender) joints; OR
- (b) at least 4 active joints from the following list:
 - (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - (ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The joint count assessment must be performed preferably whilst still on DMARD treatment, but no longer than 4 weeks following cessation of the most recent prior treatment.

The authority application must be made in writing and must include:

- (1) completed authority prescription form(s); and
- (2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

At the time of authority application, medical practitioners must request the appropriate number of vials of appropriate strength to provide sufficient drug, based on the weight of the patient, for one infusion. A separate authority prescription form must be completed for each strength requested. Up to a maximum of 3 repeats will be authorised.

An assessment of a patient's response to an initial course of treatment must be conducted following a minimum of 12 weeks of therapy. An application for the continuing treatment must be accompanied with the assessment of response and submitted to the Department of Human Services no later than 4 weeks from the date of completion of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Use of alternative DMARDs in children is dependent on approval by the Therapeutic Goods Administration as age restrictions may apply.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 12 months)

Treatment criteria:

- Must be treated by a paediatric rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be under 18 years of age.

An adequate response to treatment is defined as:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

At the time of authority application, medical practitioners must request the appropriate number of vials of appropriate strength to provide sufficient drug, based on the weight of the patient, for one infusion. A separate authority prescription form must be completed for each strength requested. Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

(1) completed authority prescription form(s); and

(2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to change or recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient who fails to demonstrate a response to treatment with this drug under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug in this treatment cycle. A patient may re-trial this drug after a minimum of 12 months have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the initial 3 treatment restriction.

If a patient fails to respond to PBS-subsidised biological medicine treatment 3 times (once with each agent) they will not be eligible to receive further PBS-subsidised biological medicine therapy in this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 3 (recommencement of treatment after a break in biological medicine of more than 12 months)

Treatment criteria:

- Must be treated by a paediatric rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**

- Patient must have had a break in treatment of 12 months or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must have either (a) a total active joint count of at least 20 active (swollen and tender) joints; or (b) at least 4 active major joints, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be under 18 years of age.

Active joints are defined as:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

All measures of joint count must be no more than 4 weeks old at the time of this application.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of active joints, the response must be demonstrated on the total number of active joints.

At the time of authority application, medical practitioners must request the appropriate number of vials of appropriate strength to provide sufficient drug, based on the weight of the patient, for one infusion. A separate authority prescription form must be completed for each strength requested. Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

(1) completed authority prescription form(s); and

(2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient) or Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 12 months) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 12 months) - balance of supply

Treatment criteria:

- Must be treated by a paediatric rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 12 months) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 12 months) restriction to complete 16 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment per continuing treatment course authorised under this restriction.

An adequate response to treatment is defined as:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

Determination of whether a response has been demonstrated to initial and subsequent courses of treatment will be based on the baseline measurement of joint count submitted with the initial treatment application.

The authority application must be made in writing and must include:

(1) completed authority prescription form(s); and

(2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

At the time of authority application, medical practitioners must request the appropriate number of vials of appropriate strength to provide sufficient drug, based on the weight of the patient, for one infusion. A separate authority prescription form must be completed for each strength requested. Up to a maximum of 5 repeats will be authorised.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either Initial 1, Initial 2, or Initial 3 treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 12 months have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

If a patient fails to respond to PBS-subsidised biological medicine treatment 3 times (once with each agent) they will not be eligible to receive further PBS-subsidised biological medicine therapy in this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Continuing treatment - balance of supply

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restriction.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

tocilizumab 200 mg/10 mL injection, 10 mL vial

10056G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	203.73	Actemra [RO]

tocilizumab 80 mg/4 mL injection, 4 mL vial

10077J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	82.19	Actemra [RO]

tocilizumab 400 mg/20 mL injection, 20 mL vial

10064Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	405.39	Actemra [RO]

■ TOCILIZUMAB
Note TREATMENT OF ADULT PATIENTS WITH A HISTORY OF JUVENILE IDIOPATHIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, etanercept and tocilizumab for a patient over 18 years who has a history of juvenile idiopathic arthritis with onset prior to the age of 18 years. Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, etanercept and tocilizumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 3 biological medicines at any one time.

From 1 April 2014, a patient receiving PBS-subsidised biological medicine therapy is considered to be in a treatment cycle where they may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements, within a single treatment cycle, a patient may:

- (i) continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy; and
- (ii) fail to respond or to sustain a response to each PBS-subsidised biological medicine once only. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment.

Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single treatment cycle and they must have, at a minimum, a 5 year break in PBS-subsidised biological medicine therapy before they are eligible to receive further PBS-subsidised biological medicine therapy.

The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine was approved to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who received PBS-subsidised biological medicine treatment immediately prior to 1 April 2014 is considered to be in their first cycle as of 1 April 2014. A patient who has had a break in biological medicine treatment of at least 24 months immediately prior to making a new application, on or after 1 April 2014, will commence a new treatment cycle under the Initial 3 treatment restriction.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of less than 24 months may commence a further course of treatment within the same treatment cycle under the Initial 2 treatment restriction.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of more than 24 months must commence a new treatment cycle under the Initial 3 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine therapy after 1 April 2014.

(1) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy (Initial 1 - New patient); or
- (ii) a patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an alternate medicine (Initial 2 - Change or recommencement of treatment after a break in biological medicine of less than 24 months) [further details are under 'Swapping therapy' below]; or
- (iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 24 months with the same medicine (Initial 2 - Change or recommencement of treatment after a break in biological medicine therapy of less than 24 months); or
- (iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 24 months (Initial 3 - recommencement of treatment after a break in biological medicine of more than 24 months).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy. A patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not conducted within these timeframes, the patient will be deemed to have failed this course of treatment.

For second and subsequent cycles of PBS-subsidised biological medicine, it is recommended that a patient is reviewed in the 4 weeks prior to completing their current course of treatment and that an application is posted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services where applicable. Where a response assessment is not submitted, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious

adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (joint count and ESR/CRP) or the prior non- biological medicine therapy requirements, except if the patient has had a break in therapy of more than 24 months who would then need to requalify under the Initial 3 restrictions with respect to the indices of disease severity.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, an application for a patient who wishes to swap to an alternate biological medicine should be accompanied by the approved authority prescription or remaining repeats for the biological medicine the patient is ceasing.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count submitted with the first authority application for a biological medicine. However, prescribers may provide a new baseline measurement any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to the revised baseline measurement.

(4) Recommencement of treatment after a 24 months break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological medicine therapy of at least 24 months, must qualify under the Initial 3 restriction and meet the relevant criteria and index of disease severity.

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have a documented history of severe active juvenile idiopathic arthritis with onset prior to the age of 18 years, **AND**
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with disease modifying anti-rheumatic drugs (DMARDs) which must include at least 3 months continuous treatment with each of at least 2 DMARDs, one of which must be methotrexate at a dose of at least 20 mg weekly and one of which must be: (i) hydroxychloroquine at a dose of at least 200 mg daily; or (ii) leflunomide at a dose of at least 10 mg daily; or (iii) sulfasalazine at a dose of at least 2 g daily; OR
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs which, if methotrexate is contraindicated according to the Therapeutic Goods Administration (TGA)-approved Product Information or cannot be tolerated at a 20 mg weekly dose, must include at least 3 months continuous treatment with each of at least 2 of the following DMARDs: (i) hydroxychloroquine at a dose of at least 200 mg daily; and/or (ii) leflunomide at a dose of at least 10 mg daily; and/or (iii) sulfasalazine at a dose of at least 2 g daily; OR
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs which, if 3 or more of methotrexate, hydroxychloroquine, leflunomide and sulfasalazine are contraindicated according to the relevant TGA-approved Product Information or cannot be tolerated at the doses specified above, must include at least 3 months continuous treatment with each of at least 2 DMARDs, with one or more of the following DMARDs being used in place of the DMARDs which are contraindicated or not tolerated: (i) azathioprine at a dose of at least 1 mg/kg per day; and/or (ii) cyclosporin at a dose of at least 2 mg/kg/day; and/or (iii) sodium aurothiomalate at a dose of 50 mg weekly, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

If methotrexate is contraindicated according to the TGA-approved Product Information or cannot be tolerated at a 20 mg weekly dose, the application must include details of the contraindication or intolerance to methotrexate. The maximum tolerated dose of methotrexate must be documented in the application, if applicable.

The application must include details of the DMARDs trialed, their doses and duration of treatment, and all relevant contraindications and/or intolerances.

The requirement to trial at least 2 DMARDs for periods of at least 3 months each can be met using single agents sequentially or by using one or more combinations of DMARDs.

If the requirement to trial 6 months of intensive DMARD therapy with at least 2 DMARDs cannot be met because of contraindications and/or intolerances of a severity necessitating permanent treatment withdrawal to all of the DMARDs specified above, details of the contraindication or intolerance and dose for each DMARD must be provided in the authority application.

The following criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; **AND** either

- (a) an active joint count of at least 20 active (swollen and tender) joints; or
- (b) at least 4 active joints from the following list:

- (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
(ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The joint count and ESR and/or CRP must be determined at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. All measures must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

The authority application must be made in writing and must include:

- (1) completed authority prescription form(s); and
- (2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

At the time of authority application, medical practitioners must request the appropriate number of vials of appropriate strength to provide sufficient drug, based on the weight of the patient, for one infusion. A separate authority prescription form must be completed for each strength requested. Up to a maximum of 3 repeats will be authorised.

An assessment of a patient's response to an initial course of treatment must be conducted following a minimum of 12 weeks of therapy. An application for the continuing treatment must be accompanied with the assessment of response and submitted to the Department of Human Services no later than 4 weeks from the date of completion of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note The Department of Human Services website (www.humanservices.gov.au) has details of the toxicities, including severity, which will be accepted for the following purposes:

- (a) exempting a patient from the requirement to undertake a minimum 3 month trial of methotrexate at a 20 mg weekly dose;
- (b) substituting azathioprine, cyclosporin or sodium aurothiomalate for another DMARD as part of the 6 month intensive DMARD trial;
- (c) exempting a patient from the requirement for a 6 month trial of intensive DMARD therapy.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 24 months)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have a documented history of severe active juvenile idiopathic arthritis with onset prior to the age of 18 years, **AND**
- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

- (a) an active joint count of fewer than 10 active (swollen and tender) joints; or
- (b) a reduction in the active (swollen and tender) joint count by at least 50% from baseline; or
- (c) a reduction in the number of the following active joints, from at least 4, by at least 50%:
 - (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - (ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

At the time of authority application, medical practitioners must request the appropriate number of vials of appropriate strength to provide sufficient drug, based on the weight of the patient, for one infusion. A separate authority prescription form must be completed for each strength requested. Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

- (1) completed authority prescription form(s); and
- (2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to change or recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient who fails to demonstrate a response to treatment with this drug under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug in this treatment cycle. A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the initial 3 treatment restriction.

If a patient fails to respond to PBS-subsidised biological medicine treatment 3 times (once with each agent) they will not be eligible to receive further PBS-subsidised biological medicine therapy in this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 3 (recommencement of treatment after a break in biological medicine of more than 24 months)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 24 months or more from the most recently approved PBS-subsidised biological medicine for this condition; OR
- Patient must not have received PBS-subsidised biological medicine for at least 5 years if they failed or ceased to respond to PBS-subsidised biological medicine treatment 3 times in their last treatment cycle, **AND**
- The condition must have an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; OR
- The condition must have a C-reactive protein (CRP) level greater than 15 mg per L, **AND**
- The condition must have either (a) a total active joint count of at least 20 active (swollen and tender) joints; or (b) at least 4 active major joints, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Active joints are defined as:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

All measures of joint count must be no more than 4 weeks old at the time of this application.

At the time of authority application, medical practitioners must request the appropriate number of vials of appropriate strength to provide sufficient drug, based on the weight of the patient, for one infusion. A separate authority prescription form must be completed for each strength requested. Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

- (1) completed authority prescription form(s); and
- (2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient) or Initial 2 (change or recommencement of treatment after break of less than 24 months) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 24 months) - balance of supply

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 24 months) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 24 months) to complete 16 weeks of treatment, **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment per continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

- (a) an active joint count of fewer than 10 active (swollen and tender) joints; or
- (b) a reduction in the active (swollen and tender) joint count by at least 50% from baseline; or
- (c) a reduction in the number of the following active joints, from at least 4, by at least 50%:
 - (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - (ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number

of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

(1) completed authority prescription form(s); and

(2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

At the time of authority application, medical practitioners must request the appropriate number of vials of appropriate strength to provide sufficient drug, based on the weight of the patient, for one infusion. A separate authority prescription form must be completed for each strength requested. Up to a maximum of 5 repeats will be authorised.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either Initial 1, Initial 2, or Initial 3 treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

If a patient fails to respond to PBS-subsidised biological medicine treatment 3 times (once with each agent) they will not be eligible to receive further PBS-subsidised biological medicine therapy in this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Continuing Treatment - balance of supply

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restriction.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

tocilizumab 200 mg/10 mL injection, 10 mL vial

10058J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	203.73	Actemra [RO]

tocilizumab 80 mg/4 mL injection, 4 mL vial

10081N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	82.19	Actemra [RO]

tocilizumab 400 mg/20 mL injection, 20 mL vial

10072D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	405.39	Actemra [RO]

■ TOCILIZUMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for adults with severe active rheumatoid arthritis. Where the term biological medicine appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alfa antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab), the T-cell co-stimulation modulator (abatacept) and the Janus kinase (JAK) inhibitors (baricitinib, tofacitinib). A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate

a response to at least 1 course of treatment with a PBS-subsidised TNF- α antagonist.

A patient receiving PBS-subsidised biological medicine therapy may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once, and
- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised biological medicines for the treatment of rheumatoid arthritis.

Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Stevens Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

For a patient who has failed PBS-subsidised treatment with 2 or 3 TNF- α antagonists prior to 1 August 2010 please contact the Department of Human Services on 1800 700 270. A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction.

A patient who has failed fewer than 5 biological medicines and who has a break in treatment of less than 24 months may commence a further course of treatment with a biological medicine under Initial 2 treatment restriction. A patient who has failed fewer than 5 biological medicines and who has had a break in therapy of longer than 24 months may commence a further course of treatment with a biological medicine under the Initial 3 treatment restriction. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine treatment is stopped to the date of the new application for treatment with a biological medicine.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 April 2019.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised biological medicine treatment and wishes to commence such therapy, excluding rituximab (Initial 1 - new patient); or
 - (ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months) [further details are under 'Swapping therapy' below]; or
 - (iii) a patient wishes to re-commence treatment with a specific biological medicine following a break of less than 24 months in PBS-subsidised therapy with that agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months).
 - (iv) a patient wishes to re-commence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 24 months (Initial 3 - re-commencement of treatment after a break in biological medicine of more than 24 months)
- Initial applications for a new patient (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6-month intensive DMARD trial, but prior to ceasing DMARD therapy.

Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, baricitinib, etanercept, golimumab, tocilizumab and tofacitinib, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

Rituximab patients should be assessed following a minimum of 12 weeks after the first infusion, and the assessment should be submitted to the Department of Human Services within 4 weeks to ensure continuity of treatment for those who meet the continuing restriction for PBS subsidised treatment with this drug for this condition.

Where a response assessment is not provided the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment. For second and subsequent courses of PBS-subsidised biological medicine (excluding rituximab) treatment it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to the Department of Human Services within sufficient timeframe to allow for authority assessment.

Abatacept patients:

A patient is eligible to receive one I.V. loading dose when commencing treatment with the subcutaneous formulation. Two prescriptions are required, the first prescription for the I.V. loading dose for sufficient vials for one dose based on the patient's weight with no repeats. The second prescription for the subcutaneous formulation, with a maximum quantity of 4 and up to 3 repeats, must be submitted with the initial application.

Rituximab patients:

A further application may be submitted to the Department of Human Services within sufficient timeframe to allow authority assessment. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

Infliximab and etanercept patients:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions. For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction. Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment

with that biological medicine.

(2) Swapping therapy

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non- biological medicine therapy requirements except if the patient has had a break in therapy of more than 24 months who would need to requalify with respect to the indices of disease severity. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each biological medicine trialled.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent, unless they have a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug.

Abatacept:

A patient swapping from I.V. abatacept to subcutaneous abatacept will not be eligible for an I.V. loading dose when commencing treatment with the subcutaneous formulation.

Rituximab:

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF-alfa antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological medicine during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological medicine therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate biological medicine may do so without having to have any treatment-free period.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and the Department of Human Services will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints.

Applications under the Initial 1 treatment restriction for a new patient must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy.

Applications under the Initial 3 treatment restriction for re-commencement of treatment after a break in biological medicine of more than 24 months must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with disease modifying anti-rheumatic drugs (DMARDs) which must include at least 3 months continuous treatment with each of at least 2 DMARDs, one of which must be methotrexate at a dose of at least 20 mg weekly and one of which must be: (i) hydroxychloroquine at a dose of at least 200 mg daily; or (ii) leflunomide at a dose of at least 10 mg daily; or (iii) sulfasalazine at a dose of at least 2 g daily; OR
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs which, if methotrexate is contraindicated according to the Therapeutic Goods Administration (TGA)-approved Product Information or cannot be tolerated at a 20 mg weekly dose, must include at least 3 months continuous treatment with each of at least 2 of the following DMARDs: (i) hydroxychloroquine at a dose of at least 200 mg daily; and/or (ii) leflunomide at a dose of at least 10 mg daily; and/or (iii) sulfasalazine at a dose of at least 2 g daily; OR
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs which, if 3 or more of methotrexate, hydroxychloroquine, leflunomide and sulfasalazine are contraindicated according to the relevant TGA-approved Product Information or cannot be tolerated at the doses specified above, must include at least 3 months continuous treatment with each of at least 2 DMARDs, with one or more of the following DMARDs being used in place of the DMARDs which are contraindicated or not tolerated: (i) azathioprine at a dose of at least 1 mg/kg per day; and/or (ii) cyclosporin at a dose of at least 2 mg/kg/day; and/or (iii) sodium aurothiomalate at a dose of 50 mg weekly, **AND**

- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

If methotrexate is contraindicated according to the TGA-approved product information or cannot be tolerated at a 20 mg weekly dose, the application must include details of the contraindication or intolerance including severity to methotrexate. The maximum tolerated dose of methotrexate must be documented in the application, if applicable.

The application must include details of the DMARDs trialled, their doses and duration of treatment, and all relevant contraindications and/or intolerances including severity.

The requirement to trial at least 2 DMARDs for periods of at least 3 months each can be met using single agents sequentially or by using one or more combinations of DMARDs.

If the requirement to trial 6 months of intensive DMARD therapy with at least 2 DMARDs cannot be met because of contraindications and/or intolerances of a severity necessitating permanent treatment withdrawal to all of the DMARDs specified above, details of the contraindication or intolerance including severity and dose for each DMARD must be provided in the authority application.

The following criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; AND either

(a) a total active joint count of at least 20 active (swollen and tender) joints; or

(b) at least 4 active joints from the following list of major joints:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The joint count and ESR and/or CRP must be determined at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. All measures must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

At the time of the authority application, medical practitioners should request the appropriate number of vials of appropriate strength to provide sufficient drug, based on the weight of the patient, for a single infusion at a dose of 8 mg per kg. A separate authority prescription form must be completed for each strength requested.

Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note The Department of Human Services website (www.humanservices.gov.au) has details of the toxicities, including severity, which will be accepted for the following purposes:

(a) exempting a patient from the requirement to undertake a minimum 3 month trial of methotrexate at a 20 mg weekly dose;

(b) substituting azathioprine, cyclosporin or sodium aurothiomalate for another DMARD as part of the 6 month intensive DMARD trial;

(c) exempting a patient from the requirement for a 6 month trial of intensive DMARD therapy.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 2 (change or re-commencement of treatment after a break in biological medicine of less than 24 months)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must not have failed to respond to previous PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have already failed , or ceased to respond to, PBS-subsidised biological medicine treatment for this condition 5 times, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either of the Initial 1, Initial 2, Initial 3, or continuing treatment restrictions, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

At the time of the authority application, medical practitioners should request the appropriate number of vials of appropriate strength to provide sufficient drug, based on the weight of the patient, for a single infusion at a dose of 8 mg per kg. A separate authority prescription form must be completed for each strength requested.

Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

A patient who has demonstrated a response to a course of rituximab must have a PBS-subsidised biological therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 3 (re-commencement of treatment after a break in biological medicine of more than 24 months)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 24 months or more from the most recent PBS-subsidised biological medicine for this condition, **AND**
- Patient must not have failed to respond to previous PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have already failed , or ceased to respond to, PBS-subsidised biological medicine treatment for this condition 5 times, **AND**
- The condition must have an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; OR

- The condition must have a C-reactive protein (CRP) level greater than 15 mg per L, **AND**
- The condition must have either (a) a total active joint count of at least 20 active (swollen and tender) joints; or (b) at least 4 active major joints, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Major joints are defined as (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

All measures of joint count and ESR and/or CRP must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

At the time of the authority application, medical practitioners should request the appropriate number of vials of appropriate strength to provide sufficient drug, based on the weight of the patient, for a single infusion at a dose of 8 mg per kg. A separate authority prescription form must be completed for each strength requested.

Up to a maximum of 3 repeats will be authorised.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial 1 (new patient) or Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 24 months) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 24 months) - balance of supply

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 24 months) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 24 months) to complete 16 weeks of treatment, **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment per continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

At the time of the authority application, medical practitioners should request the appropriate number of vials of appropriate strength to provide sufficient drug, based on the weight of the patient, for a single infusion at a dose of 8 mg per kg. A separate authority prescription form must be completed for each strength requested.

Up to a maximum of 5 repeats will be authorised.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient has either failed or ceased to respond to a PBS-subsidised biological medicine for this condition 5 times, they will not be eligible to receive further PBS-subsidised treatment with a biological medicine for this condition.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Continuing Treatment - balance of supply.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restriction.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

tocilizumab 200 mg/10 mL injection, 10 mL vial

9658H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	203.73	Actemra [RO]

tocilizumab 80 mg/4 mL injection, 4 mL vial

9657G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	82.19	Actemra [RO]

tocilizumab 400 mg/20 mL injection, 20 mL vial

9659J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	405.39	Actemra [RO]

■ USTEKINUMAB
Note TREATMENT OF ADULT PATIENTS WITH SEVERE CROHN DISEASE

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for adult patients with severe Crohn disease. Where the term biological medicine appears in the following NOTES and restrictions, it refers to the tumour necrosis factor (TNF) alfa-antagonists (adalimumab and infliximab), the alpha-4 beta-7 integrin inhibitor (vedolizumab) and the human IgG1kappa monoclonal antibody (ustekinumab).

Patients are eligible for PBS-subsidised treatment with only 1 of the above PBS-subsidised biological medicines at any one time.

From 1 September 2017, under the PBS, all patients will be able to commence a treatment cycle where they may trial PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab without having to experience a disease flare when swapping to the alternate agent. Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with adalimumab, infliximab, vedolizumab or ustekinumab while they continue to show a response to therapy.

A patient who received PBS-subsidised adalimumab, infliximab, or vedolizumab treatment prior to 1 September 2017 is considered to have started their treatment cycle as of 1 September 2017.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab more than once.

Once a patient has either failed or ceased to respond to treatment for this condition 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy for this condition before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab treatment in the most recent cycle to the date of the first application for initial treatment with adalimumab, infliximab, vedolizumab or ustekinumab under the new treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of less than 5 years, may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab therapy after 1 September 2017.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised therapy with adalimumab, infliximab, vedolizumab or ustekinumab in this treatment cycle and wishes to commence such therapy (Initial treatment (new patient or Recommencement of treatment after more than 5 years break in therapy - Initial 1)); or

(ii) a patient has received prior PBS-subsidised (initial or continuing) adalimumab, infliximab, vedolizumab or ustekinumab and wishes to trial an alternate agent (Initial 2 - Change or recommencement) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to re-commence treatment with adalimumab, infliximab, vedolizumab or ustekinumab following a break in PBS-subsidised therapy with that agent (Change or Re-commencement of treatment after a break in therapy of less than 5 years (Initial 2)).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, 14 weeks of therapy for infliximab, 14 weeks of therapy for vedolizumab and 16 weeks for ustekinumab.

From 1 September 2017, a patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy for adalimumab or ustekinumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab or vedolizumab, and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

For subsequent courses of PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

Adalimumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats and the second prescription should be written for 2 doses of 40 mg and 2 repeats for patients weighing 40 kg or greater. For patients weighing less than 40 kg, one prescription should be written for 2 doses of 40 mg with no repeats and the second prescription should be written for 2 doses of 20 mg with 3 repeats.

Ustekinumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be written under S100 (Highly Specialised Drugs) for a weight-based loading dose, containing a quantity of up to 4 vials of 130 mg and no repeats. The second prescription should be written under S85 (General) for the subsequent first dose, containing a quantity of 2 vials of 45 mg and no repeats.

(b) Continuing treatment.

Following the completion of an initial treatment course with adalimumab, infliximab, vedolizumab or ustekinumab, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted supply of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine therapy is approved, a patient may swap if eligible to the alternate adalimumab, infliximab, vedolizumab or ustekinumab within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Crohn Disease Activity Index (CDAI) Score, confirmation of Crohn disease), or the prior conventional therapies of corticosteroid therapy and immunosuppressive therapy.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with adalimumab, infliximab, vedolizumab or ustekinumab at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug once within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment, within the timeframes specified in the relevant restriction.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the CDAI or evidence of intestinal inflammation submitted with the first authority application for adalimumab, infliximab, vedolizumab or ustekinumab. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to assess response to all subsequent treatments.

(4) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to recommence treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity. Patients must have received treatment with a corticosteroid and at least 1 immunosuppressive agent, at an adequate dose, for a minimum of 3 consecutive months immediately prior to the time the CDAI score or the indices of intestinal inflammation are measured.

(5) Patients 'grandfathered' onto PBS-subsidised treatment with vedolizumab.

A patient who commenced treatment with vedolizumab for severe Crohn disease prior to 1 August 2015 and who continues to receive treatment at the time of application may qualify for treatment under the initial 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment will be assessed under the continuing treatment restriction of the relevant drug.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must requalify for continuing treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' above for further details.

(6) Patients 'grandfathered' onto PBS-subsidised treatment with ustekinumab.

A patient who commenced treatment with ustekinumab for severe Crohn disease prior to 1 September 2017 and who continues to receive treatment at the time of application may qualify for treatment under the initial 'grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this criterion once. A maximum of 24 weeks of treatment will be authorised under this criterion.

Following completion of the initial PBS-subsidised course, further applications for treatment will be assessed under the continuing treatment restriction of the relevant drug.

'Grandfather' arrangements will only apply for the first treatment cycle. For the second and subsequent cycles, a 'grandfather' patient must requalify for continuing treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 5-year break in PBS-subsidised therapy' above for further details.

Note It is recommended that an application for continuing treatment is submitted to the Department of Human Services at the time of the 12 week assessment, to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised treatment with this drug.

Note No increase in the maximum number of repeats may be authorised.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs Programs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe Crohn disease

Treatment Phase: Initial treatment (new patient - initial 1)

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have confirmed severe Crohn disease, defined by standard clinical, endoscopic and/or imaging features, including histological evidence, with the diagnosis confirmed by a gastroenterologist or a consultant physician, **AND**
- Patient must have failed to achieve an adequate response to prior systemic therapy with a tapered course of steroids, starting at a dose of at least 40 mg prednisolone (or equivalent), over a 6 week period, **AND**
- Patient must have failed to achieve adequate response to prior systemic immunosuppressive therapy with azathioprine at a dose of at least 2 mg per kg daily for 3 or more months; OR

- Patient must have failed to achieve adequate response to prior systemic immunosuppressive therapy with 6-mercaptopurine at a dose of at least 1 mg per kg daily for 3 or more months; OR
- Patient must have failed to achieve adequate response to prior systemic immunosuppressive therapy with methotrexate at a dose of at least 15 mg weekly for 3 or more months, **AND**
- Patient must have a Crohn Disease Activity Index (CDAI) Score greater than or equal to 300 as evidence of failure to achieve an adequate response to prior systemic therapy; OR
- Patient must have short gut syndrome with diagnostic imaging or surgical evidence, or have had an ileostomy or colostomy; and must have evidence of intestinal inflammation; and must have evidence of failure to achieve an adequate response to prior systemic therapy as specified below; OR
- Patient must have extensive intestinal inflammation affecting more than 50 cm of the small intestine as evidenced by radiological imaging; and must have a Crohn Disease Activity Index (CDAI) Score greater than or equal to 220; and must have evidence of failure to achieve an adequate response to prior systemic therapy as specified below.

Population criteria:

- Patient must be aged 18 years or older.

Applications for authorisation must be made in writing and must include:

(a) two completed authority prescription forms; and

(b) a completed Crohn Disease PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed current Crohn Disease Activity Index (CDAI) calculation sheet including the date of assessment of the patient's condition if relevant; and

(ii) details of prior systemic drug therapy [dosage, date of commencement and duration of therapy]; and

(iii) the reports and dates of the pathology or diagnostic imaging test(s) nominated as the response criterion, if relevant; and

(iv) the date of the most recent clinical assessment; and

(v) the signed patient acknowledgement indicating they understand and acknowledge that the PBS-subsidised treatment will cease if they do not meet the predetermined response criterion for ongoing PBS-subsidised treatment, as outlined in the restriction for continuing treatment

Evidence of failure to achieve an adequate response to prior therapy must include at least one of the following: (a) patient must have evidence of intestinal inflammation; (b) patient must be assessed clinically as being in a high faecal output state; (c) patient must be assessed clinically as requiring surgery or total parenteral nutrition (TPN) as the next therapeutic option, in the absence of this drug, if affected by short gut syndrome, extensive small intestine disease or is an ostomy patient. Evidence of intestinal inflammation includes: (i) blood: higher than normal platelet count, or, an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour, or, a C-reactive protein (CRP) level greater than 15 mg per L; or (ii) faeces: higher than normal lactoferrin or calprotectin level; or (iii) diagnostic imaging: demonstration of increased uptake of intravenous contrast with thickening of the bowel wall or mesenteric lymphadenopathy or fat streaking in the mesentery;

Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be written under S100 (Highly Specialised Drugs) for a weight-based loading dose, containing a quantity of up to 4 vials of 130 mg and no repeats. The second prescription should be written under S85 (General) for the subsequent first dose, containing a quantity of 2 vials of 45 mg and no repeats.

Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period.

All assessments, pathology tests and diagnostic imaging studies must be made within 1 month of the date of application.

If treatment with any of the specified prior conventional drugs is contraindicated according to the relevant TGA-approved Product Information, please provide details at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, details of this toxicity must be provided at the time of application.

Details of the accepted toxicities including severity can be found on the Department of Human Services website.

Any one of the baseline criteria may be used to determine response to an initial course of treatment and eligibility for continued therapy, according to the criteria included in the continuing treatment restriction. However, the same criterion must be used for any subsequent determination of response to treatment, for the purpose of eligibility for continuing PBS-subsidised therapy.

A maximum quantity of a weight based loading dose is up to 4 vials with no repeats and the subsequent dose of 90 mg (2 vials of 45 mg) with no repeats provide for an initial 16 week course of this drug will be authorised.

The assessment of the patient's response to this initial course of treatment must be made following a minimum of 12 weeks of therapy so that there is adequate time for a response to be demonstrated.

This assessment, which will be used to determine eligibility for further continuing treatment, must be submitted to the Department of Human Services no later than 1 month from the date of completion of this course of treatment.

Where a response assessment is not undertaken and submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with this drug.

Note Increase in the maximum quantity or number of units up to 4 may be authorised for the purpose of weight-based loading dose.

Authority required

Severe Crohn disease

Treatment Phase: Change or Re-commencement of treatment (initial 2)

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have a documented history of severe Crohn disease, **AND**

- Patient must have received prior PBS-subsidised treatment with a biological disease modifying drug for this condition in this treatment cycle, **AND**
- Patient must not have failed PBS-subsidised therapy with this drug for this condition in the current treatment cycle.

Population criteria:

- Patient must be aged 18 years or older.

Applications for authorisation must be made in writing and must include:

(a) two completed authority prescription forms; and

(b) a completed Crohn Disease PBS Authority Application - Supporting Information Form, which includes the following:

(i) the completed Crohn Disease Activity Index (CDAI) Score calculation sheet including the date of the assessment of the patient's condition, if relevant; or

(ii) the reports and dates of the pathology or diagnostic imaging test(s) used to assess response to therapy for patients with short gut syndrome, extensive small intestine disease or an ostomy, if relevant; and

(iii) the date of clinical assessment; and

(iv) the details of prior biological disease modifying drug treatment including the details of date and duration of treatment.

Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be written under S100 (Highly Specialised Drugs) for a weight-based loading dose, containing a quantity of up to 4 vials of 130 mg and no repeats. The second prescription should be written under S85 (General) for 2 vials of 45 mg and no repeats.

To demonstrate a response to treatment the application must be accompanied by the results of the most recent course of biological disease modifying drug (bDMD) therapy within the timeframes specified in the relevant restriction.

Where the most recent course of PBS-subsidised bDMD treatment was approved under an initial treatment restriction, the patient must have been assessed for response to that course following a minimum of 12 weeks of therapy for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab and vedolizumab and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

If the response assessment to the previous course of bDMD treatment is not submitted as detailed above, the patient will be deemed to have failed therapy with that particular course of bDMD.

A maximum quantity of a weight based loading dose is up to 4 vials with no repeats and the subsequent first dose of 90 mg (2 vials of 45 mg) with no repeats provide for an initial 16 week course of this drug will be authorised.

The assessment of the patient's response to this initial course of treatment must be made following a minimum of 12 weeks of therapy so that there is adequate time for a response to be demonstrated.

This assessment, which will be used to determine eligibility for continuing treatment, must be submitted to the Department of Human Services no later than 1 month from the date of completion of this initial course of treatment.

Where a response assessment is not undertaken and submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with this drug.

Note Increase in the maximum quantity or number of units up to 4 may be authorised for the purpose of weight-based loading dose.

ustekinumab 130 mg/26 mL injection, 26 mL vial

11182M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	4	..	..	*16929.24	Stelara [JC]

Calcineurin inhibitors
■ CICLOSPORIN

Caution Careful monitoring of patients is mandatory.

Authority required (STREAMLINED)

6628

Management of transplant rejection

Clinical criteria:

- The treatment must be used by organ or tissue transplant recipients.

ciclosporin 50 mg/mL injection, 10 x 1 mL ampoules

5631J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	54.10	Sandimmun [NV]

■ CICLOSPORIN

Caution Careful monitoring of patients is mandatory.

Authority required (STREAMLINED)

6643

Management of transplant rejection

Treatment Phase: Management (initiation, stabilisation and review of therapy)

Clinical criteria:

- Patient must have had an organ or tissue transplantation, **AND**
- The treatment must be under the supervision and direction of a transplant unit.

Authority required (STREAMLINED)

6660

Severe atopic dermatitis

Treatment Phase: Management (initiation, stabilisation and review of therapy)

Treatment criteria:

- Must be treated by a dermatologist; OR
- Must be treated by a clinical immunologists.

Clinical criteria:

- The condition must be ineffective to other systemic therapies; OR
- The condition must be inappropriate for other systemic therapies.

Authority required (STREAMLINED)
6676

Severe psoriasis

Treatment Phase: Management (initiation, stabilisation and review of therapy)

Clinical criteria:

- The condition must be ineffective to other systemic therapies; OR
- The condition must be inappropriate for other systemic therapies, **AND**
- The condition must have caused significant interference with quality of life.

Treatment criteria:

- Must be treated by a dermatologist.

Authority required (STREAMLINED)
6631

Nephrotic syndrome

Treatment Phase: Management (initiation, stabilisation and review of therapy)

Clinical criteria:

- Patient must have failed prior treatment with steroids and cytostatic drugs; OR
- Patient must be intolerant to treatment with steroids and cytostatic drugs; OR
- The condition must be considered inappropriate for treatment with steroids and cytostatic drugs, **AND**
- Patient must not have renal impairment.

Treatment criteria:

- Must be treated by a nephrologist.

Authority required (STREAMLINED)
6638

Severe active rheumatoid arthritis

Treatment Phase: Management (initiation, stabilisation and review of therapy)

Clinical criteria:

- The condition must have been ineffective to prior treatment with classical slow-acting anti-rheumatic agents (including methotrexate); OR
- The condition must be considered inappropriate for treatment with slow-acting anti-rheumatic agents (including methotrexate).

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist.

ciclosporin 25 mg capsule, 30

5634M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	4	5	..	*128.88	^a Cyclosporin Sandoz [SZ]	^a Neoral 25 [NV]

ciclosporin 10 mg capsule, 60

5632K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	5	..	*74.40	Neoral 10 [NV]	

ciclosporin 100 mg/mL oral liquid, 50 mL

5633L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	4	5	..	*1263.16	Neoral [NV]	

ciclosporin 50 mg capsule, 30

5635N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	4	5	..	*268.16	^a Cyclosporin Sandoz [SZ]	^a Neoral 50 [NV]

ciclosporin 100 mg capsule, 30

5636P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	4	5	..	*546.40	^a Cyclosporin Sandoz [SZ]	^a Neoral 100 [NV]

■ TACROLIMUS
Caution Careful monitoring of patients is mandatory.

Authority required (STREAMLINED)
5569

Management of rejection in patients following organ or tissue transplantation

Clinical criteria:

- The treatment must be under the supervision and direction of a transplant unit, **AND**

- The treatment must include initiation, stabilisation, and review of therapy as required.

tacrolimus 1 mg modified release capsule, 60

9665Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*263.28	ADVAGRAF XL [LQ]

tacrolimus 2 mg capsule, 100

10860N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*1000.00	Tacrolimus Sandoz [SZ]

tacrolimus 5 mg capsule, 50

9561F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	5	..	*1096.44	^a Pacrolim [AF] ^a Prograf [LL] ^a TACROLIMUS APOTEX [TX]	^a Pharmacor Tacrolimus 5 [CR] ^a Tacrograf [RW] ^a Tacrolimus Sandoz [SZ]

tacrolimus 750 microgram capsule, 100

10859M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*375.00	Tacrolimus Sandoz [SZ]

tacrolimus 5 mg modified release capsule, 30

9666R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*750.44	ADVAGRAF XL [LQ]

tacrolimus 500 microgram modified release capsule, 30

9664P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*75.04	ADVAGRAF XL [LQ]

tacrolimus 500 microgram capsule, 100

9558C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	5	..	*219.40	^a Pacrolim [AF] ^a Prograf [LL] ^a TACROLIMUS APOTEX [TX]	^a Pharmacor Tacrolimus 0.5 [CR] ^a Tacrograf [RW] ^a Tacrolimus Sandoz [SZ]

tacrolimus 1 mg capsule, 100

9560E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	5	..	*438.80	^a Pacrolim [AF] ^a Prograf [LL] ^a TACROLIMUS APOTEX [TX]	^a Pharmacor Tacrolimus 1 [CR] ^a Tacrograf [RW] ^a Tacrolimus Sandoz [SZ]

Other immunosuppressants
■ LENALIDOMIDE

Note Special Pricing Arrangements apply.

Authority required

Myelodysplastic syndrome

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be limited to a maximum duration of 16 weeks, **AND**
- Patient must be classified as Low risk or Intermediate-1 according to the International Prognostic Scoring System (IPSS), **AND**
- Patient must have a deletion 5q cytogenetic abnormality with or without additional cytogenetic abnormalities, **AND**
- Patient must be red blood cell transfusion dependent.

Classification of a patient as Low risk requires a score of 0 on the IPSS, achieved with the following combination: less than 5% marrow blasts with good karyotypic status (normal, -Y alone, -5q alone, -20q alone), and 0/1 cytopenias.

Classification of a patient as Intermediate-1 requires a score of 0.5 to 1 on the IPSS, achieved with the following possible combinations:

- 5%-10% marrow blasts with good karyotypic status (normal, -Y alone, -5q alone, -20q alone), and 0/1 cytopenias; OR
- less than 5% marrow blasts with intermediate karyotypic status (other abnormalities), and 0/1 cytopenias; OR
- less than 5% marrow blasts with good karyotypic status (normal, -Y alone, -5q alone, -20q alone), and 2/3 cytopenias; OR
- less than 5% marrow blasts with intermediate karyotypic status (other abnormalities), and 2/3 cytopenias; OR
- 5%-10% marrow blasts with intermediate karyotypic status (other abnormalities), and 0/1 cytopenias; OR
- 5%-10% marrow blasts with good karyotypic status (normal, -Y alone, -5q alone, -20q alone), and 2/3 cytopenias; OR
- less than 5% marrow blasts with poor karyotypic status (complex, greater than 3 abnormalities), and 0/1 cytopenias.

Classification of a patient as red blood cell transfusion dependent requires that:

- the patient has been transfused within the last 8 weeks; and
- the patient has received at least 8 units of red blood cell in the last 6 months prior to commencing PBS-subsidised therapy with lenalidomide; and would be expected to continue this requirement without lenalidomide treatment.

Patients receiving lenalidomide under the PBS listing must be registered in the i-access risk management program.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Myelodysplastic Syndrome Lenalidomide Authority Application - Supporting Information Form; and
- (c) a copy of the bone marrow biopsy report demonstrating that the patient has myelodysplastic syndrome; and
- (d) a copy of the full blood examination report; and
- (e) a copy of the pathology report detailing the cytogenetics demonstrating Low risk or Intermediate-1 disease according to the IPSS (note: using Fluorescence in Situ Hybridization (FISH) to demonstrate MDS -5q is acceptable); and
- (f) details of transfusion requirements including: (i) the date of most recent transfusion and the number of red blood cell units transfused; and (ii) the total number of red cell units transfused in the 4 and 6 months preceding the date of this application; and
- (g) a signed patient acknowledgement form.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Myelodysplastic syndrome

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must be classified as Low risk or Intermediate-1 according to the International Prognostic Scoring System (IPSS), **AND**
- Patient must have a deletion 5q cytogenetic abnormality with or without additional cytogenetic abnormalities, **AND**
- Patient must have received PBS-subsidised initial therapy with lenalidomide for myelodysplastic syndrome, **AND**
- Patient must have achieved and maintained transfusion independence; or least a 50% reduction in red blood cell unit transfusion requirements compared with the four month period prior to commencing initial PBS-subsidised therapy with lenalidomide, **AND**
- Patient must not have progressive disease.

Patients receiving lenalidomide under the PBS listing must be registered in the i-access risk management program.

The first authority application for continuing supply must be made in writing. Subsequent authority applications for continuing supply may be made by telephone.

The following evidence of response must be provided at each application:

- (i) a haemoglobin level taken within the last 4 weeks; and
- (ii) the date of the last transfusion; and
- (iii) a statement of the number of units of red cells transfused in the 4 months immediately preceding this application; and
- (iv) a statement confirming that the patient has not progressed to acute myeloid leukaemia.

Note Written applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Subsequent authority applications for continuing treatment may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

lenalidomide 5 mg capsule, 21

2799H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	3	..	5122.76	Revlimid [CJ]

lenalidomide 10 mg capsule, 21

2802L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	3	..	5361.16	Revlimid [CJ]

▪ LENALIDOMIDE

Note Special Pricing Arrangements apply.

Authority required

Multiple myeloma

Treatment Phase: Initial PBS-subsidised treatment

Clinical criteria:

- The condition must be confirmed by a histological diagnosis, **AND**
- The treatment must be as monotherapy; OR
- The treatment must be in combination with dexamethasone, **AND**
- Patient must have progressive disease after at least one prior therapy, **AND**
- Patient must have undergone or be ineligible for a primary stem cell transplant, **AND**
- Patient must not be receiving concomitant PBS-subsidised bortezomib, carfilzomib or thalidomide or its analogues.

Progressive disease is defined as at least 1 of the following:

- (a) at least a 25% increase and an absolute increase of at least 5 g per L in serum M protein (monoclonal protein); or
- (b) at least a 25% increase in 24-hour urinary light chain M protein excretion, and an absolute increase of at least 200 mg per 24 hours; or
- (c) in oligo-secretory and non-secretory myeloma patients only, at least a 50% increase of the difference between involved free light chain and uninvolved free light chain; or
- (d) at least a 25% relative increase and at least a 10% absolute increase in plasma cells in a bone marrow aspirate or on biopsy; or
- (e) an increase in the size or number of lytic bone lesions (not including compression fractures); or
- (f) at least a 25% increase in the size of an existing or the development of a new soft tissue plasmacytoma (determined by clinical examination or diagnostic imaging); or
- (g) development of hypercalcaemia (corrected serum calcium greater than 2.65 mmol per L not attributable to any other cause).

Oligo-secretory and non-secretory patients are defined as having active disease with less than 10 g per L serum M protein.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Multiple Myeloma lenalidomide Authority Application - Supporting Information Form, which includes details of the histological diagnosis of multiple myeloma, prior treatments including name(s) of drug(s) and date of most recent treatment cycle and record of prior stem cell transplant or ineligibility for prior stem cell transplant; details of the basis of the diagnosis of progressive disease or failure to respond; and nomination of which disease activity parameters will be used to assess response; and
- (3) a signed patient acknowledgment.

To enable confirmation of eligibility for treatment, current diagnostic reports of at least one of the following must be provided:

- (a) the level of serum monoclonal protein; or
- (b) Bence-Jones proteinuria - the results of 24-hour urinary light chain M protein excretion; or
- (c) the serum level of free kappa and lambda light chains; or
- (d) bone marrow aspirate or trephine; or
- (e) if present, the size and location of lytic bone lesions (not including compression fractures); or
- (f) if present, the size and location of all soft tissue plasmacytomas by clinical or radiographic examination i.e. MRI or CT-scan; or
- (g) if present, the level of hypercalcaemia, corrected for albumin concentration.

As these parameters will be used to determine response, results for either (a) or (b) or (c) should be provided for all patients. Where the patient has oligo-secretory or non-secretory multiple myeloma, either (c) or (d) or if relevant (e), (f) or (g) should be provided. Where the prescriber plans to assess response in patients with oligo-secretory or non-secretory multiple myeloma with free light chain assays, evidence of the oligo-secretory or non-secretory nature of the multiple myeloma (current serum M protein less than 10 g per L) must be provided.

Patients receiving lenalidomide under the PBS listing must be registered in the i-access risk management program.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Multiple myeloma

Treatment Phase: Continuing PBS-subsidised treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for relapsed or refractory multiple myeloma, **AND**
- The treatment must be as monotherapy; OR
- The treatment must be in combination with dexamethasone, **AND**
- Patient must not be receiving concomitant PBS-subsidised bortezomib, carfilzomib or thalidomide or its analogues.

Patients receiving lenalidomide under the PBS listing must be registered in the i-access risk management program.

Note Authority applications for continuing treatment may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note Written applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

lenalidomide 5 mg capsule, 21

5783J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	5122.76	Revlimid [CJ]

lenalidomide 15 mg capsule, 21

5785L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	6252.53	Revlimid [CJ]

lenalidomide 10 mg capsule, 21

5784K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	5361.16	Revlimid [CJ]

lenalidomide 25 mg capsule, 21

5786M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	6587.49	Revlimid [CJ]

▪ **LENALIDOMIDE**

Caution This drug is a category X drug and must not be given to pregnant women. If lenalidomide is taken during pregnancy, a teratogenic effect of lenalidomide in humans cannot be ruled out.

Note Special Pricing Arrangements apply.

Authority required

Multiple myeloma

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be newly diagnosed, **AND**
- The condition must be confirmed by a histological diagnosis, **AND**
- Patient must be ineligible for a primary stem cell transplantation, **AND**
- Patient must not be receiving concomitant PBS-subsidised bortezomib, thalidomide or its analogues, **AND**
- The treatment must be in combination with dexamethasone.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Multiple Myeloma lenalidomide Authority Application - Supporting Information Form, which includes details of the histological diagnosis of multiple myeloma, and ineligibility for prior stem cell transplant; and nomination of which disease activity parameters will be used to assess response; and
- (3) a signed patient acknowledgement.

To enable confirmation of eligibility for treatment, current diagnostic reports of at least one of the following must be provided:

- (a) the level of serum monoclonal protein; or
- (b) Bence-Jones proteinuria - the results of 24-hour urinary light chain M protein excretion; or
- (c) the serum level of free kappa and lambda light chains; or
- (d) bone marrow aspirate or trephine; or
- (e) if present, the size and location of lytic bone lesions (not including compression fractures); or
- (f) if present, the size and location of all soft tissue plasmacytomas by clinical or radiographic examination i.e. MRI or CT-scan; or
- (g) if present, the level of hypercalcaemia, corrected for albumin concentration.

As these parameters will be used to determine response, results for either (a) or (b) or (c) should be provided for all patients. Where the patient has oligo-secretory or non-secretory multiple myeloma, either (c) or (d) or if relevant (e), (f) or (g) should be provided. Where the prescriber plans to assess response in patients with oligo-secretory or non-secretory multiple myeloma with free light chain assays, evidence of the oligo-secretory or non-secretory nature of the multiple myeloma (current serum M protein less than 10 g per L) must be provided.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Patient must be registered in the i-access risk management program.

Authority required

Multiple myeloma

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously been authorised with a PBS prescription with this drug for the condition, **AND**
- Patient must not have demonstrated progressive disease, **AND**
- Patient must not be receiving concomitant PBS-subsidised bortezomib, thalidomide or its analogues, **AND**
- The treatment must be in combination with dexamethasone.

Progressive disease is defined as at least 1 of the following:

- (a) at least a 25% increase and an absolute increase of at least 5 g per L in serum M protein (monoclonal protein); or

- (b) at least a 25% increase in 24-hour urinary light chain M protein excretion, and an absolute increase of at least 200 mg per 24 hours; or
- (c) in oligo-secretory and non-secretory myeloma patients only, at least a 50% increase of the difference between involved free light chain and uninvolved free light chain; or
- (d) at least a 25% relative increase and at least a 10% absolute increase in plasma cells in a bone marrow aspirate or on biopsy; or
- (e) an increase in the size or number of lytic bone lesions (not including compression fractures); or
- (f) at least a 25% increase in the size of an existing or the development of a new soft tissue plasmacytoma (determined by clinical examination or diagnostic imaging); or
- (g) development of hypercalcaemia (corrected serum calcium greater than 2.65 mmol per L not attributable to any other cause).

Oligo-secretory and non-secretory patients are defined as having active disease with less than 10 g per L serum M protein.

Patients receiving this drug under the PBS listing must be registered in the i-access risk management program.

Note Authority applications for continuing treatment may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note Written applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

lenalidomide 5 mg capsule, 21

11029L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	5122.76	Revlimid [CJ]

lenalidomide 15 mg capsule, 21

11062F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	6252.53	Revlimid [CJ]

lenalidomide 10 mg capsule, 21

11064H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	5361.16	Revlimid [CJ]

lenalidomide 25 mg capsule, 21

11041D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	6587.49	Revlimid [CJ]

■ POMALIDOMIDE

Caution This drug is a category X drug and must not be given to pregnant women. Pregnancy in female patients or in the partners of male patients must be avoided during treatment and for 1 month after cessation of treatment.

Note Special Pricing Arrangements apply.

Authority required

Multiple myeloma

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be in combination with dexamethasone, **AND**
- Patient must have undergone or be ineligible for a primary stem cell transplant, **AND**
- Patient must have experienced treatment failure with lenalidomide, unless contraindicated or not tolerated according to the Therapeutic Goods Administration (TGA) approved Product Information, **AND**
- Patient must have experienced treatment failure with bortezomib, unless contraindicated or not tolerated according to the Therapeutic Goods Administration (TGA) approved Product Information, **AND**
- Patient must not be receiving concomitant PBS-subsidised bortezomib, carfilzomib or thalidomide or its analogues. Bortezomib treatment failure is the absence of achieving at least a partial response or as progressive disease during treatment or within 6 months of discontinuing treatment with bortezomib. Lenalidomide treatment failure is progressive disease during treatment or within 6 months of discontinuing treatment with lenalidomide.

If treatment with either bortezomib or lenalidomide is contraindicated according to the relevant TGA-approved Product Information, the application must provide details of the contraindication.

If intolerance to either bortezomib or lenalidomide treatment develops during the relevant period of use which is of a severity to necessitate withdrawal of the treatment, the application must provide details of the nature and severity of this intolerance.

Progressive disease is defined as at least 1 of the following:

- (a) at least a 25% increase and an absolute increase of at least 5 g per L in serum M protein (monoclonal protein); or
- (b) at least a 25% increase in 24-hour urinary light chain M protein excretion, and an absolute increase of at least 200 mg per 24 hours; or
- (c) in oligo-secretory and non-secretory myeloma patients only, at least a 50% increase of the difference between involved free light chain and uninvolved free light chain; or
- (d) at least a 25% relative increase and at least a 10% absolute increase in plasma cells in a bone marrow aspirate or on biopsy; or
- (e) an increase in the size or number of lytic bone lesions (not including compression fractures); or

- (f) at least a 25% increase in the size of an existing or the development of a new soft tissue plasmacytoma (determined by clinical examination or diagnostic imaging); or
(g) development of hypercalcaemia (corrected serum calcium greater than 2.65 mmol per L not attributable to any other cause).

Oligo-secretory and non-secretory patients are defined as having active disease with less than 10 g per L serum M protein. The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Multiple Myeloma pomalidomide Authority Application Supporting Information form; and
- (3) reports demonstrating the patient has failed treatment with, providing details of the contraindication to or details of the nature and severity of the intolerance to lenalidomide; and
- (4) reports demonstrating the patient has failed treatment with, providing details of the contraindication to or details of the nature and severity of the intolerance to bortezomib.

Patients receiving this drug under the PBS listing must be registered in the i-access risk management program.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Prior Written Approval of Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Multiple myeloma

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously been issued with an authority prescription for this drug, **AND**
- Patient must not have progressive disease, **AND**
- The treatment must be in combination with dexamethasone, **AND**
- Patient must not be receiving concomitant PBS-subsidised bortezomib, carfilzomib or thalidomide or its analogues.

Progressive disease is defined as at least 1 of the following:

- (a) at least a 25% increase and an absolute increase of at least 5 g per L in serum M protein (monoclonal protein); or
- (b) at least a 25% increase in 24-hour urinary light chain M protein excretion, and an absolute increase of at least 200 mg per 24 hours; or
- (c) in oligo-secretory and non-secretory myeloma patients only, at least a 50% increase of the difference between involved free light chain and uninvolved free light chain; or
- (d) at least a 25% relative increase and at least a 10% absolute increase in plasma cells in a bone marrow aspirate or on biopsy; or
- (e) an increase in the size or number of lytic bone lesions (not including compression fractures); or
- (f) at least a 25% increase in the size of an existing or the development of a new soft tissue plasmacytoma (determined by clinical examination or diagnostic imaging); or
- (g) development of hypercalcaemia (corrected serum calcium greater than 2.65 mmol per L not attributable to any other cause).

Patients receiving this drug under the PBS listing must be registered in the i-access risk management program.

Note Authority applications for continuing treatment may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note Written applications for authority to prescribe should be forwarded to:

Department of Human Services
Prior Written Approval of Complex Drugs
Reply Paid 9826
HOBART TAS 7001

pomalidomide 3 mg capsule, 21

10406Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	10500.00	Pomalyst [CJ]

pomalidomide 4 mg capsule, 21

10387Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	10500.00	Pomalyst [CJ]

■ THALIDOMIDE

Caution Thalidomide is a category X drug and must not be given to pregnant women. Pregnancy in female patients or in the partners of male patients must be avoided during treatment and for 1 month after cessation of treatment.

Note Patients receiving thalidomide under the PBS listing must be registered in the i-access risk management program.

Authority required (STREAMLINED)

5914

Multiple myeloma

MUSCULO-SKELETAL SYSTEM

thalidomide 50 mg capsule, 28

9566L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	4	..	..	*1436.40	Thalomid [CJ]

thalidomide 100 mg capsule, 28

9667T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	..	..	*1436.40	Thalomid [CJ]

■ MUSCULO-SKELETAL SYSTEM

■ MUSCLE RELAXANTS

MUSCLE RELAXANTS, CENTRALLY ACTING AGENTS

Other centrally acting agents

■ BACLOFEN

Authority required (STREAMLINED)

7152

Severe chronic spasticity

Clinical criteria:

- Patient must have failed to respond to treatment with oral antispastic agents; OR
- Patient must have had unacceptable side effects to treatment with oral antispastic agents, **AND**
- Patient must have chronic spasticity of cerebral origin.

Authority required (STREAMLINED)

7134

Severe chronic spasticity

Clinical criteria:

- Patient must have failed to respond to treatment with oral antispastic agents; OR
- Patient must have had unacceptable side effects to treatment with oral antispastic agents, **AND**
- Patient must have chronic spasticity due to multiple sclerosis.

Authority required (STREAMLINED)

7153

Severe chronic spasticity

Clinical criteria:

- Patient must have failed to respond to treatment with oral antispastic agents; OR
- Patient must have had unacceptable side effects to treatment with oral antispastic agents, **AND**
- Patient must have chronic spasticity due to spinal cord injury.

Authority required (STREAMLINED)

7148

Severe chronic spasticity

Clinical criteria:

- Patient must have failed to respond to treatment with oral antispastic agents; OR
- Patient must have had unacceptable side effects to treatment with oral antispastic agents, **AND**
- Patient must have chronic spasticity due to spinal cord disease.

baclofen 40 mg/20 mL intrathecal injection, 20 mL ampoule

11195F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	..	..	*766.24	Sintetica Baclofen Intrathecal [BZ]

■ BACLOFEN

Note Pharmaceutical benefits that have the forms baclofen 10 mg/5 mL intrathecal injection, 5 mL ampoule, baclofen 10 mg/5 mL intrathecal injection, 5 x 5 mL ampoules and baclofen 10 mg/5 mL intrathecal injection, 10 x 5 mL ampoules are equivalent for the purposes of substitution.

Authority required (STREAMLINED)

6925

Severe chronic spasticity

Clinical criteria:

- Patient must have failed to respond to treatment with oral antispastic agents; OR
- Patient must have had unacceptable side effects to treatment with oral antispastic agents, **AND**
- Patient must have chronic spasticity of cerebral origin.

Authority required (STREAMLINED)

6939

Severe chronic spasticity

Clinical criteria:

- Patient must have failed to respond to treatment with oral antispastic agents; OR
- Patient must have had unacceptable side effects to treatment with oral antispastic agents, **AND**
- Patient must have chronic spasticity due to multiple sclerosis.

Authority required (STREAMLINED)**6940**

Severe chronic spasticity

Clinical criteria:

- Patient must have failed to respond to treatment with oral antispastic agents; OR
- Patient must have had unacceptable side effects to treatment with oral antispastic agents, **AND**
- Patient must have chronic spasticity due to spinal cord injury.

Authority required (STREAMLINED)**6911**

Severe chronic spasticity

Clinical criteria:

- Patient must have failed to respond to treatment with oral antispastic agents; OR
- Patient must have had unacceptable side effects to treatment with oral antispastic agents, **AND**
- Patient must have chronic spasticity due to spinal cord disease.

baclofen 10 mg/5 mL intrathecal injection, 5 x 5 mL ampoules

11421D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	..	..	*958.00	^a Bacthecal [DZ]

baclofen 10 mg/5 mL intrathecal injection, 5 mL ampoule

5617P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	10	..	..	*958.00	^a Bacthecal [DZ]	^a Lioresal Intrathecal [NV]

baclofen 10 mg/5 mL intrathecal injection, 10 x 5 mL ampoules

11126N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	958.00	^a Sintetica Baclofen Intrathecal [BZ]

DRUGS FOR TREATMENT OF BONE DISEASES**DRUGS AFFECTING BONE STRUCTURE AND MINERALIZATION***Bisphosphonates***IBANDRONATE****Authority required (STREAMLINED)****5291**

Bone metastases

Clinical criteria:

- The condition must be due to breast cancer.

ibandronate 6 mg/6 mL injection, 6 mL vial

5750P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	11	..	291.86	Bondronat [IX]

PAMIDRONATE DISODIUM**Authority required (STREAMLINED)****4433**

Hypercalcaemia of malignancy

Clinical criteria:

- Patient must have a malignancy refractory to anti-neoplastic therapy.

pamidronate disodium 15 mg/5 mL injection, 5 mL vial

5667G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	4	2	..	*55.24	Pamisol [PF]

pamidronate disodium 60 mg/10 mL injection, 10 mL vial

5669J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	2	..	55.26	Pamisol [PF]

pamidronate disodium 30 mg/10 mL injection, 10 mL vial

5668H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	2	..	*55.26	Pamisol [PF]

PAMIDRONATE DISODIUM**Authority required (STREAMLINED)****4433**

Hypercalcaemia of malignancy

Clinical criteria:

- Patient must have a malignancy refractory to anti-neoplastic therapy.

Authority required (STREAMLINED)**5218**

Multiple myeloma

Authority required (STREAMLINED)**5291**

Bone metastases

Clinical criteria:

- The condition must be due to breast cancer.

pamidronate disodium 90 mg/10 mL injection, 10 mL vial

5670K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	11	..	82.89	Pamisol [PF]

■ ZOLEDRONIC ACID

Note Pharmaceutical benefits that have the form zoledronic acid 4 mg/100 mL injection and pharmaceutical benefits that have the form zoledronic acid 4 mg/5 mL injection are equivalent for the purposes of substitution.

Authority required (STREAMLINED)**5735**

Multiple myeloma

Authority required (STREAMLINED)**5605**

Bone metastases

Clinical criteria:

- The condition must be due to breast cancer.

Authority required (STREAMLINED)**5703**

Bone metastases

Clinical criteria:

- The condition must be due to castration-resistant prostate cancer.

Authority required (STREAMLINED)**5704**

Hypercalcaemia of malignancy

Clinical criteria:

- Patient must have a malignancy refractory to anti-neoplastic therapy.

zoledronic acid 4 mg/5 mL injection, 5 mL vial

9653C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	11	..	132.37	^a APO-Zoledronic Acid [TX] ^a DEZTRON [DZ]	^a DBL Zoledronic Acid [PF] ^a Zometa [NV]

zoledronic acid 4 mg/100 mL injection, 100 mL bag

10561W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	11	..	132.37	^a DBL Zoledronic Acid [PF]

■ OTHER DRUGS FOR DISORDERS OF THE MUSCULO-SKELETAL SYSTEM**OTHER DRUGS FOR DISORDERS OF THE MUSCULO-SKELETAL SYSTEM***Other drugs for disorders of the musculo-skeletal system***■ NUSINERSEN**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Spinal muscular atrophy (SMA)

Treatment Phase: Initial treatment - Loading doses

Treatment criteria:

- Must be treated by a specialist medical practitioner experienced in the diagnosis and management of SMA associated with a neuromuscular clinic of a recognised hospital in the management of SMA; or in consultation with a specialist

medical practitioner experienced in the diagnosis and management of SMA associated with a neuromuscular clinic of a recognised hospital in the management of SMA.

Clinical criteria:

- The condition must 5q homozygous deletion, mutation of, or compound heterozygous mutation in the SMN1 gene of type I, II or IIIa, **AND**
- Patient must have experienced at least two of the defined signs and symptoms of SMA type I, II or IIIa prior to 3 years of age, **AND**
- The treatment must be given concomitantly with standard of care for this condition, **AND**
- The treatment must not exceed four loading doses (at days 0, 14, 28 and 63) under this restriction.

Population criteria:

- Patient must be 18 years of age or under.

Defined signs and symptoms of type I SMA are:

- Onset before 6 months of age; and
- Failure to meet or regression in ability to perform age-appropriate motor milestones; or
- Proximal weakness; or
- Hypotonia; or
- Absence of deep tendon reflexes; or
- Failure to gain weight appropriate for age; or
- Any active chronic neurogenic changes; or
- A compound muscle action potential below normative values for an age-matched child.

Defined signs and symptoms of type II SMA are:

- Onset between 6 and 18 months; and
- Failure to meet or regression in ability to perform age-appropriate motor milestones; or
- Proximal weakness; or
- Weakness in trunk righting/derotation; or
- Hypotonia; or
- Absence of deep tendon reflexes; or
- Failure to gain weight appropriate for age; or
- Any active chronic neurogenic changes; or
- A compound muscle action potential below normative values for an age-matched child.

Defined signs and symptoms of type IIIa SMA are:

- Onset between 18 months and 3 years of age; and
- Failure to meet or regression in ability to perform age-appropriate motor milestones; or
- Proximal weakness; or
- Hypotonia; or
- Absence of deep tendon reflexes; or
- Failure to gain weight appropriate for age; or
- Any active chronic neurogenic changes; or
- A compound muscle action potential below normative values for an age-matched child.

Recognised hospitals in the management of SMA are Lady Cilento Children's Hospital (Brisbane), Royal Children's Hospital Melbourne, Monash Children's Hospital (Melbourne), John Hunter Hospital (Newcastle), Sydney Children's Hospital Randwick, Children's Hospital at Westmead, Adelaide Women and Children's Hospital and Perth Children's Hospital.

Application for authorisation of initial treatment must be in writing and must include:

- a completed authority prescription form; and
- a completed Spinal muscular atrophy PBS Authority Application - Supporting Information Form which includes the following:
 - specification of SMA type (I, II or IIIa); and
 - sign(s) and symptom(s) that the patient has experienced; and
 - patient's age at the onset of sign(s) and symptom(s).

nusinersen 12 mg/5 mL injection, 5 mL vial

11363C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	3	..	110000.00	Spinraza [BD]

▪ **NUSINERSEN**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Authority applications for continuing treatment may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Spinal muscular atrophy (SMA)

Treatment Phase: Continuing treatment - Maintenance

Treatment criteria:

- Must be treated by a specialist medical practitioner experienced in the diagnosis and management of SMA associated with a neuromuscular clinic of a recognised hospital in the management of SMA; or in consultation with a specialist

medical practitioner experienced in the diagnosis and management of SMA associated with a neuromuscular clinic of a recognised hospital in the management of SMA.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be given concomitantly with standard of care for this condition, **AND**
- The treatment must be ceased when invasive permanent assisted ventilation is required in the absence of a potentially reversible cause while being treated with this drug.

Recognised hospitals in the management of SMA are Lady Cilento Children's Hospital (Brisbane), Royal Children's Hospital Melbourne, Monash Children's Hospital (Melbourne), John Hunter Hospital (Newcastle), Sydney Children's Hospital Randwick, Children's Hospital at Westmead, Adelaide Women and Children's Hospital and Perth Children's Hospital.

Invasive permanent assisted ventilation means ventilation via tracheostomy tube for greater than or equal to 16 hours per day.

nusinersen 12 mg/5 mL injection, 5 mL vial

11378W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	110000.00	Spinraza [BD]

■ NUSINERSEN

Note Special Pricing Arrangements apply.

Note No increase in the maximum quantity or number of units may be authorised.

Note A maximum number of repeats of up to 2 may be authorised for patients requiring loading doses for days 14, 28 and 63.

Note A maximum number of repeats of up to 1 may be authorised for patients requiring loading doses for days 28 and 63.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Spinal muscular atrophy (SMA)

Treatment Phase: Grandfather patients

Treatment criteria:

- Must be treated by a specialist medical practitioner experienced in the diagnosis and management of SMA associated with a neuromuscular clinic of a recognised hospital in the management of SMA; or in consultation with a specialist medical practitioner experienced in the diagnosis and management of SMA associated with a neuromuscular clinic of a recognised hospital in the management of SMA.

Clinical criteria:

- Patient must have previously received non-PBS-subsidised treatment for this condition with this drug prior to 1 June 2018, **AND**
- The condition must 5q homozygous deletion, mutation of, or compound heterozygous mutation in the SMN1 gene of type I, II or IIIa, **AND**
- Patient must have had experienced at least two of the defined signs and symptoms of SMA type I, II or IIIa prior to 3 years of age, **AND**
- Patient must have previously received at least one of the four loading doses at days 0, 14, 28 and 63, **AND**
- The treatment must be given concomitantly with standard of care for this condition, **AND**
- The treatment must be ceased when invasive permanent assisted ventilation is required in the absence of a potentially reversible cause while being treated with this drug.

Population criteria:

- Patient must have been 18 years of age or under at the time treatment with this drug was initiated for this condition; OR
- Patient must have previously received treatment with this drug for this condition under the care of clinicians with the authorised prescriber number of AP17/83146.

Defined signs and symptoms of type I SMA are:

- Onset before 6 months of age; and
- Failure to meet or regression in ability to perform age-appropriate motor milestones; or
- Proximal weakness; or
- Hypotonia; or
- Absence of deep tendon reflexes; or
- Failure to gain weight appropriate for age; or
- Any active chronic neurogenic changes; or
- A compound muscle action potential below normative values for an age-matched child.

Defined signs and symptoms of type II SMA are:

- Onset between 6 and 18 months; and
- Failure to meet or regression in ability to perform age-appropriate motor milestones; or
- Proximal weakness; or
- Weakness in trunk righting/derotation; or

- v) Hypotonia; or
- vi) Absence of deep tendon reflexes; or
- vii) Failure to gain weight appropriate for age; or
- viii) Any active chronic neurogenic changes; or
- ix) A compound muscle action potential below normative values for an age-matched child.

Defined signs and symptoms of type IIIa SMA are:

- i) Onset between 18 months and 3 years of age; and
- ii) Failure to meet or regression in ability to perform age-appropriate motor milestones; or
- iii) Proximal weakness; or
- iv) Hypotonia; or
- v) Absence of deep tendon reflexes; or
- vi) Failure to gain weight appropriate for age; or
- vii) Any active chronic neurogenic changes; or
- viii) A compound muscle action potential below normative values for an age-matched child.

Invasive permanent assisted ventilation means ventilation via tracheostomy tube for greater than or equal to 16 hours per day.

Recognised hospitals in the management of SMA are Lady Cilento Children's Hospital (Brisbane), Royal Children's Hospital Melbourne, Monash Children's Hospital (Melbourne), John Hunter Hospital (Newcastle), Sydney Children's Hospital Randwick, Children's Hospital at Westmead, Adelaide Women and Children's Hospital and Perth Children's Hospital.

Application for authorisation of grandfathering treatment must be in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Spinal muscular atrophy PBS Authority Application for Grandfather patients - Supporting Information Form which includes the following:
 - (i) specification of SMA type (I, II or IIIa); and
 - (ii) sign(s) and symptom(s) that the patient has experienced; and
 - (iii) patient's age at the onset of sign(s) and symptom(s); and
 - (iv) if relevant, a copy of a TGA-approval letter to clinician with the authorised prescriber number of AP17/83146.

A patient may qualify for PBS-subsidised treatment under this restriction once only.

For continuing PBS-subsidised treatment, a Grandfathered patient must qualify under the Continuing treatment criteria.

nusinersen 12 mg/5 mL injection, 5 mL vial

11370K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	110000.00	Spinraza [BD]

■ NERVOUS SYSTEM

■ ANTI-PARKINSON DRUGS

DOPAMINERGIC AGENTS

Dopa and dopa derivatives

■ LEVODOPA + CARBIDOPA

Note Special Pricing Arrangements apply.

Note Patients should have adequate cognitive function to manage administration with a portable continuous infusion pump.

Authority required (STREAMLINED)

6863

Advanced Parkinson disease

Clinical criteria:

- Patient must have severe disabling motor fluctuations not adequately controlled by oral therapy, **AND**
- The treatment must be commenced in a hospital-based movement disorder clinic.

levodopa 20 mg/mL + carbidopa monohydrate 5 mg/mL intestinal gel, 7 x 100 mL

9743T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	8	5	..	*11536.00	Duodopa [VE]

Dopamine agonists

■ APOMORPHINE

Authority required (STREAMLINED)

6813

Parkinson disease

Clinical criteria:

- Patient must have experienced severely disabling motor fluctuations which have not responded to other therapy.

apomorphine hydrochloride hemihydrate 100 mg/20 mL injection, 5 x 20 mL vials

11093W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	18	5	..	*7552.80	Apomine Solution for Infusion [PF]

■ APOMORPHINE

Authority required (STREAMLINED)**4833**

Parkinson disease

Clinical criteria:

- Patient must have experienced severely disabling motor fluctuations which have not responded to other therapy.

apomorphine hydrochloride hemihydrate 20 mg/2 mL injection, 5 x 2 mL ampoules

5609F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	72	5	..	*6528.96	Movapo [TD]

apomorphine hydrochloride hemihydrate 50 mg/5 mL injection, 5 x 5 mL ampoules

5610G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	36	5	..	*8168.04	Movapo [TD]

apomorphine hydrochloride hemihydrate 50 mg/10 mL injection, 5 x 10 mL syringes

10950H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	36	5	..	*8168.04	Movapo PFS [TD]

■ APOMORPHINE

Note No increase in the maximum quantity or number of units may be authorised.

Note Pharmaceutical benefits that have the form apomorphine injection 30 mg/3 mL pen device and pharmaceutical benefits that have the form apomorphine injection 30 mg/3 mL cartridge are equivalent for the purposes of substitution.

Authority required (STREAMLINED)**4833**

Parkinson disease

Clinical criteria:

- Patient must have experienced severely disabling motor fluctuations which have not responded to other therapy.

apomorphine hydrochloride hemihydrate 30 mg/3 mL injection, 5 x 3 mL cartridges

11672H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	20	5	..	*2720.40	^a Apomine Intermittent [PF]

apomorphine hydrochloride hemihydrate 30 mg/3 mL injection, 5 x 3 mL pen devices

11477C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	20	5	..	*2720.40	^a Movapo Pen [TD]

■ PSYCHOLEPTICS

ANTIPSYCHOTICS

Diazepines, oxazepines, thiazepines and oxepines

■ CLOZAPINE

Note Patients receiving clozapine under the PBS listing must be registered in the clozapine patient monitoring program relevant for the brand of clozapine being prescribed and dispensed: Pfizer ClopineCentral.

Authority required (STREAMLINED)**5015**

Schizophrenia

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a psychiatrist or in consultation with the psychiatrist affiliated with the hospital or specialised unit managing the patient.

Clinical criteria:

- Patient must be non-responsive to other neuroleptic agents; OR
- Patient must be intolerant of other neuroleptic agents.

Patients must complete at least 18 weeks of initial treatment under this restriction before being able to qualify for treatment under the continuing restriction.

The name of the consulting psychiatrist should be included in the patient's medical records.

A medical practitioner should request a quantity sufficient for up to one month's supply. Up to 5 repeats will be authorised.

clozapine 50 mg/mL oral liquid, 100 mL

11433R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	±1	..	..	135.00	Versacloz [PF]

■ CLOZAPINE

Note Patients receiving clozapine under the PBS listing must be registered in the clozapine patient monitoring program relevant for the brand of clozapine being prescribed and dispensed: Novartis Clozaril Patient Monitoring System (eCPMS) or Hospira Clopineconnect.

Authority required (STREAMLINED)**5015**

Schizophrenia

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a psychiatrist or in consultation with the psychiatrist affiliated with the hospital or specialised unit managing the patient.

Clinical criteria:

- Patient must be non-responsive to other neuroleptic agents; OR
- Patient must be intolerant of other neuroleptic agents.

Patients must complete at least 18 weeks of initial treatment under this restriction before being able to qualify for treatment under the continuing restriction.

The name of the consulting psychiatrist should be included in the patient's medical records.

A medical practitioner should request a quantity sufficient for up to one month's supply. Up to 5 repeats will be authorised.

clozapine 25 mg tablet, 100

5628F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	..	..	*64.64	Clopine 25 [PF]	Clozaril 25 [GO]

clozapine 100 mg tablet, 100

5629G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	..	..	*242.38	Clopine 100 [PF]	Clozaril 100 [GO]

clozapine 50 mg tablet, 100

5626D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	..	..	*129.28	Clopine 50 [PF]

clozapine 200 mg tablet, 100

5627E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	..	..	*484.76	Clopine 200 [PF]

clozapine 50 mg/mL oral liquid, 100 mL

5630H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	135.00	Clopine Suspension [PF]

■ RESPIRATORY SYSTEM

■ DRUGS FOR OBSTRUCTIVE AIRWAY DISEASES

OTHER SYSTEMIC DRUGS FOR OBSTRUCTIVE AIRWAY DISEASES

Other systemic drugs for obstructive airway diseases

■ BENRALIZUMAB

Note TREATMENT OF ADULT AND ADOLESCENT PATIENTS WITH UNCONTROLLED SEVERE EOSINOPHILIC ASTHMA

Patients are eligible to commence a 'benralizumab treatment cycle' (initial treatment course with or without continuing treatment course/s) if they satisfy the eligibility criteria as detailed under the initial treatment restriction.

Once a patient has either failed to achieve or maintain a response to benralizumab, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 6 month break in PBS-subsidised benralizumab therapy before they are eligible to commence the next 'benralizumab treatment cycle', or if eligible, a 'mepolizumab treatment cycle' or an 'omalizumab treatment cycle'. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised benralizumab is stopped to the date of the first application for initial treatment with benralizumab, mepolizumab or omalizumab under the new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised benralizumab therapy:

(a) Initial treatment:

Applications for initial treatment should be made where:

- A patient has received no prior PBS-subsidised benralizumab treatment and wishes to commence such therapy; or
- A patient wishes to recommence treatment with benralizumab following a break in PBS-subsidised therapy of more than 6 months; or
- A patient has received prior PBS-subsidised mepolizumab or omalizumab and wishes to commence treatment with benralizumab after a treatment break of 6 months.

All applications for initial treatment for non-grandfather patients will be limited to provide for a maximum of up to 32 weeks of therapy for benralizumab.

(b) Grandfather patients:

For patients who commenced treatment with non-PBS subsidised benralizumab for uncontrolled severe eosinophilic asthma prior to 1 December 2018 and who continue to receive treatment at the time of application, may qualify for treatment under the initial 'grandfather' treatment restriction. A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment with benralizumab will be authorised under this criterion. Approval will be based on the criteria included in the relevant restriction. Following completion of the Initial PBS-subsidised course, further applications for treatment with benralizumab will be assessed under the continuing treatment restriction.

'Grandfather' arrangements will only apply for the first treatment cycle (initial treatment course with or without continuing

treatment course(s). If a 'Grandfathered' patient recommences on second and subsequent cycles after a treatment break, the 'Grandfathered' patient must re-qualify for Initial treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 6 month break in PBS-subsidised therapy' below for further details.

(c) Continuing treatment:

Following the completion of the initial treatment course with benralizumab, a patient may qualify to receive up to a further 24 weeks of continuing treatment with benralizumab providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing benralizumab treatment in courses of up to 24 weeks providing they continue to sustain the response.

(2) Baseline measurements to determine response:

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the Asthma Control Questionnaire (ACQ; 5 item version) or oral corticosteroid dose, submitted with the Initial authority application for benralizumab. However, prescribers may provide new baseline measurements when a new Initial treatment authority application is submitted and the Department of Human Services will assess response according to these revised baseline measurements.

(3) Re-commencement of treatment after a 6 month break in PBS-subsidised therapy:

A patient who wishes to trial a second or subsequent benralizumab treatment cycle, or an initial mepolizumab or omalizumab treatment cycle, following a break in PBS-subsidised therapy of at least 6 months, must re-qualify for initial treatment with respect to the indices of disease severity (oral corticosteroid dose, Asthma Control Questionnaire (ACQ-5) score, and relevant exacerbation history). Patients must have received optimised standard therapy, at adequate doses and for the minimum period specified, immediately prior to the time the new baseline assessments are performed.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Uncontrolled severe eosinophilic asthma

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a respiratory physician, clinical immunologist, allergist or general physician experienced in the management of patients with severe asthma.

Clinical criteria:

- Patient must be under the care of the same physician for at least 6 months; OR
- Patient must have been diagnosed by a multidisciplinary severe asthma clinic team, **AND**
- Patient must have a diagnosis of asthma confirmed and documented by a respiratory physician, clinical immunologist, allergist or general physician experienced in the management of patients with severe asthma, defined by the following standard clinical features: (i) forced expiratory volume (FEV1) reversibility greater than or equal to 12% and greater than or equal to 200 mL at baseline within 30 minutes after administration of salbutamol (200 to 400 micrograms), or (ii) airway hyperresponsiveness defined as a greater than 20% decline in FEV1 during a direct bronchial provocation test or greater than 15% decline during an indirect bronchial provocation test, or (iii) peak expiratory flow (PEF) variability of greater than 15% between the two highest and two lowest peak expiratory flow rates during 14 days, **AND**
- Patient must have a duration of asthma of at least 1 year, **AND**
- Patient must have forced expiratory volume (FEV1) less than or equal to 80% predicted, documented on 1 or more occasions in the previous 12 months, **AND**
- Patient must have blood eosinophil count greater than or equal to 300 cells per microlitre in the last 12 months, **AND**
- Patient must have failed to achieve adequate control with optimised asthma therapy, despite formal assessment of and adherence to correct inhaler technique, which has been documented, **AND**
- The treatment must not be used in combination with, or within 6 months of treatment with, PBS-subsidised omalizumab or mepolizumab.

Population criteria:

- Patient must be aged 12 years or older.

Optimised asthma therapy includes:

(i) Adherence to maximal inhaled therapy, including high dose inhaled corticosteroid (ICS) plus long-acting beta-2 agonist (LABA) therapy for at least 12 months, unless contraindicated or not tolerated; **AND**

(ii) treatment with oral corticosteroids, either daily oral corticosteroids for at least 6 weeks, OR a cumulative dose of oral corticosteroids of at least 500 mg prednisolone equivalent in the previous 12 months, unless contraindicated or not tolerated.

If the requirement for treatment with optimised asthma therapy cannot be met because of contraindications according to the relevant TGA-approved Product Information and/or intolerances of a severity necessitating permanent treatment withdrawal, details of the contraindication and/or intolerance must be provided in the Authority application.

The following initiation criteria indicate failure to achieve adequate control and must be demonstrated in all patients at the time of the application:

(a) an Asthma Control Questionnaire (ACQ-5) score of at least 2.0, as assessed in the previous month, **AND**

(b) while receiving optimised asthma therapy in the past 12 months, experienced at least 1 admission to hospital for a severe asthma exacerbation, OR 1 severe asthma exacerbation, requiring documented use of systemic corticosteroids (oral corticosteroids initiated or increased for at least 3 days, or parenteral corticosteroids) prescribed/supervised by a physician.

The Asthma Control Questionnaire (5 item version) assessment of the patient must be made at time of application for treatment (to establish baseline score) and again around 20 weeks after the first PBS-subsidised dose of this drug under this restriction so that there is adequate time for a response to be demonstrated and for the application for the first continuing therapy to be processed.

This assessment at around 24 weeks, which will be used to determine eligibility for the first continuing treatment, must be submitted no later than 2 weeks prior to the patient completing their current treatment course, to avoid an interruption to

supply. Where a response assessment is not undertaken and submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

A patient who fails to respond to a course of PBS-subsidised benralizumab for the treatment of uncontrolled severe eosinophilic asthma will not be eligible to receive further PBS-subsidised treatment with benralizumab, mepolizumab or omalizumab within 6 months of the date on which treatment was ceased.

A multidisciplinary severe asthma clinic team comprises of:

- A respiratory physician; and
- A pharmacist, nurse or asthma educator.

At the time of the authority application, medical practitioners should request up to 4 repeats to provide for an initial course of benralizumab sufficient for up to 32 weeks of therapy, at a dose of 30 mg every 4 weeks for the first three doses (weeks 0, 4, and 8) then 30 mg every eight weeks thereafter.

Benralizumab must not be used concurrently with omalizumab or mepolizumab or within 6 months of each other. A patient is required to have ceased treatment with omalizumab or mepolizumab for 6 months prior to initiating treatment with benralizumab.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Severe Eosinophilic Asthma Initial PBS Authority Application - Supporting Information Form, which includes the following:
 - (i) details of prior optimised asthma drug therapy (date of commencement and duration of therapy); and
 - (ii) details of severe exacerbation/s experienced in the past 12 months while receiving optimised asthma therapy (date and treatment); and
- (c) a copy of the eosinophil pathology report; and
- (d) a completed Asthma Control Questionnaire (ACQ-5) calculation sheet including the date of assessment of the patient's symptoms.

Note The Department of Human Services website (www.humanservices.gov.au) has details of the accepted toxicities, including severity, which will be accepted for the purposes of exempting a patient from the requirement of treatment with optimised asthma therapy.

Note For copies of the ACQ, please contact AstraZeneca Medical Information on 1800 805 342.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Formal assessment and correction of inhaler technique should be performed in accordance with the National Asthma Council (NAC) Information Paper for Health Professionals on Inhaler Technique (available at www.humanservices.gov.au or www.nationalasthma.org.au); the assessment and adherence to correct technique should be documented in the patient's medical records. Patients can obtain support with inhaler technique through their local Asthma Foundation (1800 645 130).

Authority required

Uncontrolled severe eosinophilic asthma

Treatment Phase: Initial treatment - balance of supply

Treatment criteria:

- Must be treated by a respiratory physician, clinical immunologist, allergist or general physician experienced in the management of patients with severe asthma.

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the Initial treatment restriction to complete 32 weeks treatment; AND
- The treatment must provide no more than the balance of up to 32 weeks treatment available under the Initial restriction.

Population criteria:

- Patient must be aged 12 years or older.

Note Authority approval for sufficient therapy to complete a maximum of 32 weeks of treatment under the initial restriction may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

benralizumab 30 mg/mL injection, 1 mL syringe

11549W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	4	..	3311.00	Fasenra [AP]

■ BENRALIZUMAB

Note TREATMENT OF ADULT AND ADOLESCENT PATIENTS WITH UNCONTROLLED SEVERE EOSINOPHILIC ASTHMA
Patients are eligible to commence a 'benralizumab treatment cycle' (initial treatment course with or without continuing treatment course/s) if they satisfy the eligibility criteria as detailed under the initial treatment restriction.

Once a patient has either failed to achieve or maintain a response to benralizumab, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 6 month break in PBS-subsidised benralizumab therapy before they are eligible to commence the next 'benralizumab treatment cycle', or if eligible, a 'mepolizumab treatment cycle' or an 'omalizumab treatment cycle'. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised benralizumab is stopped to the date of the first application for initial treatment with benralizumab,

mepolizumab or omalizumab under the new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised benralizumab therapy:

(a) Initial treatment:

Applications for initial treatment should be made where:

- i) A patient has received no prior PBS-subsidised benralizumab treatment and wishes to commence such therapy; or
- ii) A patient wishes to recommence treatment with benralizumab following a break in PBS-subsidised therapy of more than 6 months; or
- iii) A patient has received prior PBS-subsidised mepolizumab or omalizumab and wishes to commence treatment with benralizumab after a treatment break of 6 months.

All applications for initial treatment for non-grandfather patients will be limited to provide for a maximum of up to 32 weeks of therapy for benralizumab.

(b) Grandfather patients:

For patients who commenced treatment with non-PBS subsidised benralizumab for uncontrolled severe eosinophilic asthma prior to 1 December 2018 and who continue to receive treatment at the time of application, may qualify for treatment under the initial 'grandfather' treatment restriction. A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment with benralizumab will be authorised under this criterion. Approval will be based on the criteria included in the relevant restriction. Following completion of the Initial PBS-subsidised course, further applications for treatment with benralizumab will be assessed under the continuing treatment restriction.

'Grandfather' arrangements will only apply for the first treatment cycle (initial treatment course with or without continuing treatment course/s). If a 'Grandfathered' patient recommences on second and subsequent cycles after a treatment break, the 'Grandfathered' patient must re-qualify for Initial treatment under the criteria that apply to a new patient. See 'Re-commencement of treatment after a 6 month break in PBS-subsidised therapy' below for further details.

(c) Continuing treatment:

Following the completion of the initial treatment course with benralizumab, a patient may qualify to receive up to a further 24 weeks of continuing treatment with benralizumab providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing benralizumab treatment in courses of up to 24 weeks providing they continue to sustain the response.

(2) Baseline measurements to determine response:

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the Asthma Control Questionnaire (ACQ; 5 item version) or oral corticosteroid dose, submitted with the Initial authority application for benralizumab. However, prescribers may provide new baseline measurements when a new Initial treatment authority application is submitted and the Department of Human Services will assess response according to these revised baseline measurements.

(3) Re-commencement of treatment after a 6 month break in PBS-subsidised therapy:

A patient who wishes to trial a second or subsequent benralizumab treatment cycle, or an initial mepolizumab or omalizumab treatment cycle, following a break in PBS-subsidised therapy of at least 6 months, must re-qualify for initial treatment with respect to the indices of disease severity (oral corticosteroid dose, Asthma Control Questionnaire (ACQ-5) score, and relevant exacerbation history). Patients must have received optimised standard therapy, at adequate doses and for the minimum period specified, immediately prior to the time the new baseline assessments are performed.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Uncontrolled severe eosinophilic asthma

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a respiratory physician, clinical immunologist, allergist or general physician experienced in the management of patients with severe asthma.

Clinical criteria:

- Patient must have demonstrated or sustained an adequate response to PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must not be used in combination with, or within 6 months of treatment with, PBS-subsidised omalizumab or mepolizumab.

Population criteria:

- Patient must be aged 12 years or older.

An adequate response to benralizumab treatment is defined as:

- (a) a reduction in the Asthma Control Questionnaire (ACQ-5) score of at least 0.5 from baseline; OR
- (b) maintenance oral corticosteroid dose reduced by at least 25% from baseline, and no deterioration in ACQ-5 score from baseline.

All applications for second and subsequent continuing treatments with this drug must include a measurement of response to the prior course of therapy. The Asthma Control Questionnaire (5 item version) assessment of the patient's response to the prior course of treatment, or the assessment of oral corticosteroid dose, should be made at around 16 weeks after the first PBS-subsidised dose of this drug under this restriction so that there is adequate time for a response to be demonstrated and for the application for continuing therapy to be processed.

The assessment should, where possible, be completed by the same physician who initiated treatment with this drug. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted no later than 2 weeks prior to the patient completing their current treatment course, to avoid an interruption to supply. Where a response assessment is not undertaken and submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

A patient who fails to respond to a course of PBS-subsidised benralizumab for the treatment of uncontrolled severe eosinophilic asthma will not be eligible to receive further PBS-subsidised treatment with benralizumab for this condition within 6 months of the date on which treatment was ceased.

At the time of the authority application, medical practitioners should request the appropriate number of repeats to provide for a continuing course of this drug sufficient for up to 24 weeks of therapy.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Severe Eosinophilic Asthma Continuing PBS Authority Application - Supporting Information Form which includes details of maintenance oral corticosteroid dose; or a completed Asthma Control Questionnaire (ACQ-5) calculation sheet including the date of assessment of the patient's symptoms.

Note If the same physician cannot assess the patient please call the Department of Human Services on 1800 700 270.

Note For copies of the ACQ, please contact AstraZeneca Medical Information on 1800 805 342.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Uncontrolled severe eosinophilic asthma

Treatment Phase: Continuing treatment or Grandfathered treatment - balance of supply

Treatment criteria:

- Must be treated by a respiratory physician, clinical immunologist, allergist or general physician experienced in the management of patients with severe asthma.

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the continuing treatment restriction or the grandfather restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the continuing treatment restriction or the grandfather restriction.

Population criteria:

- Patient must be aged 12 years or older.

Note Authority approval for sufficient therapy to complete a maximum of 24 weeks of treatment may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Uncontrolled severe eosinophilic asthma

Treatment Phase: Grandfathered treatment

Treatment criteria:

- Must be treated by a respiratory physician, clinical immunologist, allergist or general physician experienced in the management of patients with severe asthma.

Clinical criteria:

- Patient must have received non-PBS subsidised treatment with this drug for this condition prior to 1 December 2018, **AND**
- Patient must be receiving treatment with this drug for this condition at the time of application, **AND**
- Patient must be under the care of the same physician for at least 6 months; OR
- Patient must have been diagnosed by a multidisciplinary severe asthma clinic team, **AND**
- Patient must have had, prior to commencement of non-PBS subsidised treatment with this drug, a diagnosis of asthma confirmed and documented by a respiratory physician, clinical immunologist, allergist or general physician experienced in the management of patients with severe asthma, defined by the following standard clinical features: (i) forced expiratory volume (FEV1) reversibility greater than or equal to 12% and greater than or equal to 200 mL at baseline within 30 minutes after administration of salbutamol (200 to 400 micrograms), or (ii) airway hyperresponsiveness defined as a greater than 20% decline in FEV1 during a direct bronchial provocation test or greater than 15% decline during an indirect bronchial provocation test, or (iii) peak expiratory flow (PEF) variability of greater than 15% between the two highest and two lowest peak expiratory flow rates during 14 days, **AND**
- Patient must have had blood eosinophil count greater than or equal to 300 cells per microlitre prior to commencement of non-PBS subsidised treatment with this drug, **AND**
- Patient must have had a duration of asthma of at least 1 year prior to commencement of non-PBS subsidised treatment with this drug, **AND**
- Patient must have failed to achieve adequate control with optimised asthma therapy prior to non-PBS subsidised treatment with this drug despite formal assessment of and adherence to correct inhaler technique, which has been documented, **AND**
- Patient must have demonstrated an adequate response if the patient has received at least 24 weeks of treatment of non-PBS subsidised benralizumab for this condition, **AND**
- The treatment must not be used in combination with, or within 6 months of treatment with, PBS-subsidised omalizumab or mepolizumab.

Population criteria:

- Patient must be aged 12 years or older.

Optimised asthma therapy includes:

(i) Adherence to maximal inhaled therapy, including high dose inhaled corticosteroid (ICS) plus long-acting beta-2 agonist (LABA) therapy for at least 12 months, unless contraindicated or not tolerated; AND

(ii) treatment with oral corticosteroids, either daily oral corticosteroids for at least 6 weeks, OR a cumulative dose of oral corticosteroids of at least 500 mg prednisolone equivalent in the previous 12 months, prior to commencing non-PBS subsidised treatment with this drug, unless contraindicated or not tolerated.

If the requirement for treatment with optimised asthma therapy cannot be met because of contraindications according to the relevant TGA-approved Product Information and/or intolerances of a severity necessitating permanent treatment withdrawal, details of the contraindication and/or intolerance must be provided in the Authority application.

The following initiation criteria indicate failure to achieve adequate control and must be demonstrated in all patients at the time of the application:

(a) an Asthma Control Questionnaire (ACQ-5) score of at least 2.0, as assessed in the previous month, AND

(b) while receiving optimised asthma therapy in the past 12 months, experienced at least 1 admission to hospital for a severe asthma exacerbation, OR 1 severe asthma exacerbation, requiring documented use of systemic corticosteroids (oral corticosteroids initiated or increased for at least 3 days, or parenteral corticosteroids) prescribed/supervised by a physician.

The Asthma Control Questionnaire (5 item version) assessment of the patient must be made at time of application for treatment (to establish baseline score) and again around 20 weeks after the first PBS-subsidised dose of this drug under this restriction so that there is adequate time for a response to be demonstrated and for the application for the first continuing therapy to be processed.

This assessment at around 24 weeks, which will be used to determine eligibility for the first continuing treatment, must be submitted no later than 2 weeks prior to the patient completing their current treatment course, to avoid an interruption to supply. Where a response assessment is not undertaken and submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

An adequate response to benralizumab treatment is defined as:

(a) a reduction in the Asthma Control Questionnaire (ACQ 5) score of at least 0.5 from baseline; OR

(b) maintenance oral corticosteroid dose reduced by at least 25% from baseline, and no deterioration in ACQ 5 score from baseline.

A multidisciplinary severe asthma clinic team comprises of:

- A respiratory physician; and
- A pharmacist, nurse or asthma educator.

A review of the patient's records should be conducted to extract pre- and post-benralizumab data on symptoms, quality of life, medication doses, exacerbations and hospitalisations. Parameters to establish response are:

(i) a reduction in Asthma Control Questionnaire (ACQ-5) score of at least 0.5; and/or

(ii) maintenance oral corticosteroid dose reduced by at least 25% from baseline and no deterioration in ACQ 5 score from baseline.

The assessment of the patient's response to the initial PBS-subsidised course of treatment under this restriction must be made at around 16 weeks after the first dose of PBS-subsidised treatment with this drug under this restriction so that there is adequate time for a response to be demonstrated and for the application for continuing therapy to be processed. The same parameters used to establish response to non-PBS subsidised therapy with this drug should be used for the assessment.

This assessment, which will be used to determine eligibility for continuing treatment, must be submitted no later than 2 weeks prior to the patient completing their current treatment course, to avoid an interruption to supply. Where a response assessment is not undertaken and submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

Patients will be eligible to receive continuing courses of treatment with this drug of up to 24 weeks providing they continue to demonstrate an adequate response to treatment.

A patient may qualify for PBS-subsidised treatment under this restriction once only.

A patient who fails to respond to a course of PBS-subsidised benralizumab for the treatment of uncontrolled severe eosinophilic asthma will not be eligible to receive further PBS-subsidised treatment with benralizumab, omalizumab or mepolizumab within 6 months of the date on which treatment was ceased.

At the time of the authority application, medical practitioners should request the appropriate maximum quantity and number of repeats to provide for a continuing course of benralizumab sufficient for up to 24 weeks of therapy.

The authority application must be made in writing and must include:

(a) a completed authority prescription form; and

(b) a completed Severe Eosinophilic Asthma Grandfather PBS Authority Application - Supporting Information Form, which includes the following:

(i) details of prior optimised asthma drug therapy (date of commencement and duration of therapy); and

(ii) details of pre- and post-benralizumab data on symptoms, quality of life, medication doses, severe exacerbation/s and hospitalisations, and

(c) a copy of the pre-benralizumab eosinophil pathology report; and

(d) a completed Asthma Control Questionnaire (ACQ 5) calculation sheet including the date of assessment of the patient's symptoms; or details of maintenance oral corticosteroid dose.

Note The Department of Human Services website (www.humanservices.gov.au) has details of the accepted toxicities, including severity, which will be accepted for the purposes of exempting a patient from the requirement of treatment with optimised asthma therapy.

Note For copies of the ACQ, please contact AstraZeneca Medical Information on 1800 805 342.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Formal assessment and correction of inhaler technique should be performed in accordance with the National Asthma Council (NAC) Information Paper for Health Professionals on Inhaler Technique (available at www.humanservices.gov.au or www.nationalasthma.org.au); the assessment and adherence to correct technique should be documented in the patient's medical records. Patients can obtain support with inhaler technique through their local Asthma Foundation (1800 645 130).

benralizumab 30 mg/mL injection, 1 mL syringe

11529T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	2	..	3311.00	Fasenra [AP]

■ MEPOLIZUMAB

Note TREATMENT OF ADULT AND ADOLESCENT PATIENTS WITH UNCONTROLLED SEVERE EOSINOPHILIC ASTHMA

Patients are eligible to commence a 'mepolizumab treatment cycle' (initial treatment course with or without continuing treatment course/s) if they satisfy the eligibility criteria as detailed under the initial treatment restriction.

Once a patient has either failed to achieve or maintain a response to mepolizumab, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 6 month break in PBS-subsidised mepolizumab therapy before they are eligible to commence the next mepolizumab treatment cycle, or if eligible, a 'benralizumab treatment cycle' or an 'omalizumab treatment cycle'. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised mepolizumab is stopped to the date of the first application for initial treatment with benralizumab, mepolizumab or omalizumab under the new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised mepolizumab therapy:

(a) Initial treatment:

Applications for initial treatment should be made where:

- i) A patient has received no prior PBS-subsidised mepolizumab treatment and wishes to commence such therapy; or
- ii) A patient wishes to recommence treatment with mepolizumab following a break in PBS-subsidised therapy of more than 6 months; or
- iii) A patient has received prior PBS-subsidised benralizumab or omalizumab and wishes to commence treatment with mepolizumab after a treatment break of 6 months.

All applications for initial treatment will be limited to provide for a maximum of up to 32 weeks of therapy for mepolizumab.

(b) Continuing treatment:

Following the completion of the initial treatment course with mepolizumab, a patient may qualify to receive up to a further 24 weeks of continuing treatment with mepolizumab providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing mepolizumab treatment in courses of up to 24 weeks providing they continue to sustain the response.

(2) Baseline measurements to determine response:

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the Asthma Control Questionnaire (ACQ; 5 item version) or oral corticosteroid dose, submitted with the Initial authority application for mepolizumab. However, prescribers may provide new baseline measurements when a new Initial treatment authority application is submitted and the Department of Human Services will assess response according to these revised baseline measurements.

(3) Re-commencement of treatment after a 6 month break in PBS-subsidised therapy:

A patient who wishes to trial a second or subsequent mepolizumab treatment cycle, or an initial benralizumab or omalizumab treatment cycle, following a break in PBS-subsidised therapy of at least 6 months, must re-qualify for initial treatment with respect to the indices of disease severity (oral corticosteroid dose, Asthma Control Questionnaire (ACQ-5) score, and relevant exacerbation history). Patients must have received optimised standard therapy, at adequate doses and for the minimum period specified, immediately prior to the time the new baseline assessments are performed.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note If the same physician cannot assess the patient please call the Department of Human Services on 1800 700 270.

Note For copies of the ACQ, please contact GlaxoSmithKline Medical Information on 1800 033 109.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Uncontrolled severe eosinophilic asthma

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a respiratory physician, clinical immunologist, allergist or general physician experienced in the management of patients with severe asthma.

Clinical criteria:

- Patient must have demonstrated or sustained an adequate response to PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must not be used in combination with, or within 6 months of treatment with, PBS-subsidised benralizumab or omalizumab.

Population criteria:

- Patient must be aged 12 years or older.

An adequate response to mepolizumab treatment is defined as:

(a) a reduction in the Asthma Control Questionnaire (ACQ-5) score of at least 0.5 from baseline,

OR

(b) maintenance oral corticosteroid dose reduced by at least 25% from baseline, and no deterioration in ACQ-5 score from baseline.

All applications for second and subsequent continuing treatments with this drug must include a measurement of response to the prior course of therapy. The Asthma Control Questionnaire (5 item version) assessment of the patient's response to the prior course of treatment or the assessment of oral corticosteroid dose, should be made at around 18 to 22 weeks after the first dose of PBS-subsidised dose of this drug under this restriction so that there is adequate time for a response to be demonstrated and for the application for continuing therapy to be processed.

The assessment should, where possible, be completed by the same physician who initiated treatment with this drug. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted within 4 weeks of the date of assessment, and no later than 2 weeks prior to the patient completing their current treatment course, to avoid an interruption to supply. Where a response assessment is not undertaken and submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

A patient who fails to respond to a course of PBS-subsidised mepolizumab for the treatment of uncontrolled severe eosinophilic asthma will not be eligible to receive further PBS subsidised treatment with mepolizumab for this condition within 6 months of the date on which treatment was ceased.

At the time of the authority application, medical practitioners should request the appropriate number of repeats to provide for a continuing course of this drug sufficient for up to 24 weeks of therapy.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Severe Eosinophilic Asthma Continuing PBS Authority Application - Supporting Information Form which includes details of maintenance oral corticosteroid dose; or a completed Asthma Control Questionnaire (ACQ-5) calculation sheet including the date of assessment of the patient's symptoms.

mepolizumab 100 mg injection, 1 vial

10980X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	1638.00	Nucala [GK]

■ MEPOLIZUMAB**Note TREATMENT OF ADULT AND ADOLESCENT PATIENTS WITH UNCONTROLLED SEVERE EOSINOPHILIC ASTHMA**

Patients are eligible to commence a 'mepolizumab treatment cycle' (initial treatment course with or without continuing treatment course/s) if they satisfy the eligibility criteria as detailed under the initial treatment restriction.

Once a patient has either failed to achieve or maintain a response to mepolizumab, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 6 month break in PBS-subsidised mepolizumab therapy before they are eligible to commence the next mepolizumab treatment cycle, or if eligible, a 'benralizumab treatment cycle' or an 'omalizumab treatment cycle'. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised mepolizumab is stopped to the date of the first application for initial treatment with benralizumab, mepolizumab or omalizumab under the new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised mepolizumab therapy:

(a) Initial treatment:

Applications for initial treatment should be made where:

- i) A patient has received no prior PBS-subsidised mepolizumab treatment and wishes to commence such therapy; or
- ii) A patient wishes to recommence treatment with mepolizumab following a break in PBS-subsidised therapy of more than 6 months; or
- iii) A patient has received prior PBS-subsidised benralizumab or omalizumab and wishes to commence treatment with mepolizumab after a treatment break of 6 months.

All applications for initial treatment will be limited to provide for a maximum of up to 32 weeks of therapy for mepolizumab.

(b) Continuing treatment:

Following the completion of the initial treatment course with mepolizumab, a patient may qualify to receive up to a further 24 weeks of continuing treatment with mepolizumab providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing mepolizumab treatment in courses of up to 24 weeks providing they continue to sustain the response.

(2) Baseline measurements to determine response:

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the Asthma Control Questionnaire (ACQ; 5 item version) or oral corticosteroid dose, submitted with the Initial authority application for mepolizumab. However, prescribers may provide new baseline measurements when a new Initial treatment authority application is submitted and the Department of Human Services will assess response according to these revised baseline measurements.

(3) Re-commencement of treatment after a 6 month break in PBS-subsidised therapy:

A patient who wishes to trial a second or subsequent mepolizumab treatment cycle, or an initial benralizumab or

omalizumab treatment cycle, following a break in PBS-subsidised therapy of at least 6 months, must re-qualify for initial treatment with respect to the indices of disease severity (oral corticosteroid dose, Asthma Control Questionnaire (ACQ-5) score, and relevant exacerbation history). Patients must have received optimised standard therapy, at adequate doses and for the minimum period specified, immediately prior to the time the new baseline assessments are performed.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note The Department of Human Services website (www.humanservices.gov.au) has details of the accepted toxicities, including severity, which will be accepted for the purposes of exempting a patient from the requirement of treatment with optimised asthma therapy.

Note For copies of the ACQ, please contact GlaxoSmithKline Medical Information on 1800 033 109.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Formal assessment and correction of inhaler technique should be performed in accordance with the National Asthma Council (NAC) Information Paper for Health Professionals on Inhaler Technique (available at www.humanservices.gov.au or www.nationalasthma.org.au); the assessment and adherence to correct technique should be documented in the patient's medical records. Patients can obtain support with inhaler technique through their local Asthma Foundation (1800 645 130).

Authority required

Uncontrolled severe eosinophilic asthma

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a respiratory physician, clinical immunologist, allergist or general physician experienced in the management of patients with severe asthma.

Clinical criteria:

- Patient must be under the care of the same physician for at least 6 months; OR
- Patient must have been diagnosed by a multidisciplinary severe asthma clinic team, **AND**
- Patient must have a diagnosis of asthma confirmed and documented by a respiratory physician, clinical immunologist, allergist or general physician experienced in the management of patients with severe asthma, defined by the following standard clinical features: (i) forced expiratory volume (FEV1) reversibility greater than or equal to 12% and greater than or equal to 200 mL at baseline within 30 minutes after administration of salbutamol (200 to 400 micrograms), or (ii) airway hyperresponsiveness defined as a greater than 20% decline in FEV1 during a direct bronchial provocation test or greater than 15% decline during an indirect bronchial provocation test, or (iii) peak expiratory flow (PEF) variability of greater than 15% between the two highest and two lowest peak expiratory flow rates during 14 days, **AND**
- Patient must have a duration of asthma of at least 1 year, **AND**
- Patient must have forced expiratory volume (FEV1) less than or equal to 80% predicted, documented on 1 or more occasions in the previous 12 months, **AND**
- Patient must have blood eosinophil count greater than or equal to 300 cells per microlitre in the last 12 months, **AND**
- Patient must have failed to achieve adequate control with optimised asthma therapy, despite formal assessment of and adherence to correct inhaler technique, which has been documented, **AND**
- The treatment must not be used in combination with, or within 6 months of treatment with, PBS-subsidised benralizumab or omalizumab.

Population criteria:

- Patient must be aged 12 years or older.

Optimised asthma therapy includes:

(i) Adherence to maximal inhaled therapy, including high dose inhaled corticosteroid (ICS) plus long-acting beta-2 agonist (LABA) therapy for at least 12 months, unless contraindicated or not tolerated; **AND**

(ii) treatment with oral corticosteroids, either daily oral corticosteroids for at least 6 weeks, OR a cumulative dose of oral corticosteroids of at least 500 mg prednisolone equivalent in the previous 12 months, unless contraindicated or not tolerated.

If the requirement for treatment with optimised asthma therapy cannot be met because of contraindications according to the relevant TGA-approved Product Information and/or intolerances of a severity necessitating permanent treatment withdrawal, details of the contraindication and/or intolerance must be provided in the Authority application.

The following initiation criteria indicate failure to achieve adequate control and must be demonstrated in all patients at the time of the application:

(a) an Asthma Control Questionnaire (ACQ-5) score of at least 2.0, as assessed in the previous month, **AND**

(b) while receiving optimised asthma therapy in the past 12 months, experienced at least 1 admission to hospital for a severe asthma exacerbation, OR 1 severe asthma exacerbation, requiring documented use of systemic corticosteroids (oral corticosteroids initiated or increased for at least 3 days, or parenteral corticosteroids) prescribed/supervised by a physician.

The Asthma Control Questionnaire (5 item version) assessment of the patient must be made at time of application for treatment (to establish baseline score) and again around 26 to 30 weeks after the first PBS-subsidised dose of this drug under this restriction so that there is adequate time for a response to be demonstrated and for the application for the first continuing therapy to be processed.

This assessment at around 26 to 30 weeks, which will be used to determine eligibility for the first continuing treatment, must be submitted within 4 weeks of the date of assessment, and no later than 2 weeks prior to the patient completing their current treatment course, to avoid an interruption to supply. Where a response assessment is not undertaken and submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

A patient who fails to respond to a course of PBS-subsidised mepolizumab for the treatment of uncontrolled severe eosinophilic asthma will not be eligible to receive further PBS-subsidised treatment with benralizumab, mepolizumab or omalizumab within 6 months of the date on which treatment was ceased.

At the time of the authority application, medical practitioners should request up to 7 repeats to provide for an initial course of mepolizumab sufficient for up to 32 weeks of therapy.

A multidisciplinary severe asthma clinic team comprises of:

- A respiratory physician; and
- A pharmacist, nurse or asthma educator.

Mepolizumab must not be used concurrently with benralizumab or omalizumab, or within 6 months of each other. A patient is required to have ceased treatment with benralizumab or omalizumab for 6 months prior to initiating treatment with mepolizumab.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Severe Eosinophilic Asthma Initial PBS Authority Application - Supporting Information Form, which includes the following:
 - (i) details of prior optimised asthma drug therapy (date of commencement and duration of therapy); and
 - (ii) details of severe exacerbation/s experienced in the past 12 months while receiving optimised asthma therapy (date and treatment); and
- (c) a copy of the eosinophil pathology report; and
- (d) a completed Asthma Control Questionnaire (ACQ-5) calculation sheet including the date of assessment of the patient's symptoms.

mepolizumab 100 mg injection, 1 vial

Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
1	7	..	1638.00	Nucala [GK]

■ OMALIZUMAB

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic spontaneous urticaria

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a clinical immunologist; OR
- Must be treated by an allergist; OR
- Must be treated by a dermatologist; OR
- Must be treated by a general physician with expertise in the management of chronic spontaneous urticaria (CSU).

Clinical criteria:

- The condition must be based on both physical examination and patient history (to exclude any factors that may be triggering the urticaria), **AND**
- Patient must have experienced itch and hives that persist on a daily basis for at least 6 weeks despite treatment with H1 antihistamines, **AND**
- Patient must have failed to achieve an adequate response after a minimum of 2 weeks treatment with a standard therapy, **AND**
- Patient must not receive more than 12 weeks of treatment under this restriction.

A standard therapy is defined as a combination of therapies that includes H1 antihistamines at maximally tolerated doses in accordance with clinical guidelines, and one of the following:

- 1) a H2 receptor antagonist (150 mg twice per day); or
- 2) a leukotriene receptor antagonist (LTRA) (10 mg per day); or
- 3) doxepin (up to 25 mg three times a day)

If the requirement for treatment with H1 antihistamines and a H2 receptor antagonist, or a leukotriene receptor antagonist or doxepin cannot be met because of contraindications according to the relevant TGA-approved Product Information and/or intolerances of a severity necessitating permanent treatment withdrawal, details of the contraindication and/or intolerance must be provided in the authority application.

A failure to achieve an adequate response to standard therapy is defined as a current Urticaria Activity Score 7 (UAS7) score of equal to or greater than 28 with an itch score of greater than 8, as assessed while still on standard therapy.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and

(b) a completed Chronic Spontaneous Urticaria Omalizumab Initial PBS Authority Application - Supporting Information Form which must include:

- (i) demonstration of failure to achieve an adequate response to standard therapy; and
- (ii) drug names and doses of standard therapies that the patient has failed; and
- (iii) a signed patient acknowledgment that cessation of therapy should be considered after the patient has demonstrated clinical benefit with omalizumab to re-evaluate the need for continued therapy. Any patient who ceases therapy and whose CSU relapses will need to re-initiate PBS-subsidised omalizumab as a new patient.

omalizumab 150 mg/mL injection, 1 mL syringe

11176F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	2	..	*820.00	Xolair [NV]

■ OMALIZUMAB

Authority required

Severe chronic spontaneous urticaria

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a clinical immunologist; OR
- Must be treated by an allergist; OR
- Must be treated by a dermatologist; OR
- Must be treated by a general physician with expertise in the management of chronic spontaneous urticaria (CSU).

Clinical criteria:

- Patient must have demonstrated a response to the most recent PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not receive more than 24 weeks per authorised course of treatment under this restriction.

Note A proportion of patients respond to 150 mg 4-weekly so where a substantial improvement has been obtained with a 300 mg dose it is reasonable to back-titrate dose after initial treatment.

Note Cessation of therapy should be considered after the patient has demonstrated clinical benefit with omalizumab to re-evaluate the need for continued therapy. Any patient who ceases therapy and whose CSU relapses will need to re-initiate PBS-subsidised omalizumab as a new patient.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Severe chronic spontaneous urticaria

Treatment Phase: Grandfathering treatment

Clinical criteria:

- Patient must have received non-PBS subsidised treatment with this drug for this condition prior to 1 September 2017, **AND**
- Patient must have documented history of itch and hives that persisted on a daily basis for at least 6 weeks despite treatment with H1 antihistamines prior to commencing non-PBS subsidised treatment with this drug for this condition, **AND**
- Patient must have documented history of failure to achieve an adequate response after a minimum of 2 weeks treatment with a standard therapy prior to commencing non-PBS subsidised treatment with this drug for this condition, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Treatment criteria:

- Must be treated by a clinical immunologist; OR
- Must be treated by an allergist; OR
- Must be treated by a dermatologist; OR
- Must be treated by a general physician with expertise in the management of chronic spontaneous urticaria (CSU).

A standard therapy is defined as a combination of therapies that includes H1 antihistamines at maximally tolerated doses in accordance with clinical guidelines, and one of the following:

- 1) a H2 receptor antagonist (150 mg twice per day); or
- 2) a leukotriene receptor antagonist (LTRA) (10 mg per day); or
- 3) doxepin (up to 25 mg three times a day)

If the requirement for treatment with H1 antihistamines and a H2 receptor antagonist, or a leukotriene receptor antagonist or doxepin cannot be met because of contraindications according to the relevant TGA-approved Product Information and/or intolerances of a severity necessitating permanent treatment withdrawal, details of the contraindication and/or intolerance must be provided in the authority application.

A failure to achieve an adequate response to standard therapy is defined as a current Urticaria Activity Score 7 (UAS7) score of equal to or greater than 28 with an itch score of greater than 8, as assessed while still on standard therapy.

A patient may qualify for PBS-subsidised treatment under this restriction once only. For continuing PBS-subsidised treatment, a Grandfathered patient must qualify under the Continuing treatment criteria.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Chronic Spontaneous Urticaria Omalizumab Initial Grandfather PBS Authority Application - Supporting Information Form which must include:
 - (i) demonstration of failure to achieve an adequate response to standard therapy; and
 - (ii) drug names and doses of standard therapies that the patient has failed; and

(iii) a signed patient acknowledgment that cessation of therapy should be considered after the patient has demonstrated clinical benefit with omalizumab to re-evaluate the need for continued therapy. Any patient who ceases therapy and whose CSU relapses will need to re-initiate PBS-subsidised omalizumab as a new patient.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

omalizumab 150 mg/mL injection, 1 mL syringe

11168T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*820.00	Xolair [NV]

■ OMALIZUMAB

Note TREATMENT OF ADULT AND ADOLESCENT PATIENTS WITH UNCONTROLLED SEVERE ALLERGIC ASTHMA

Patients are eligible to commence an 'omalizumab treatment cycle' (initial treatment course with or without continuing treatment course/s) if they satisfy the eligibility criteria as detailed under the initial treatment restriction.

Once a patient has either failed to achieve or maintain a response to omalizumab, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 6 month break in PBS-subsidised omalizumab therapy before they are eligible to commence the next omalizumab treatment cycle or, if eligible, a 'benralizumab treatment cycle' or a 'mepolizumab treatment cycle'. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised omalizumab is stopped to the date of the first application for initial treatment with benralizumab, omalizumab or mepolizumab under the new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised omalizumab therapy:

(a) Initial treatment:

Applications for initial treatment should be made where:

i) A patient has received no prior PBS-subsidised omalizumab treatment and wishes to commence such therapy; or

ii) A patient wishes to recommence treatment with omalizumab following a break in PBS-subsidised therapy of at least 6 months; or

iii) A patient has received prior PBS-subsidised benralizumab or mepolizumab and wishes to commence treatment with omalizumab after a treatment break of at least 6 months.

All applications for initial treatment will be limited to provide for a maximum of up to 28 weeks of therapy of omalizumab.

(b) Continuing treatment:

Following the completion of the initial treatment course with omalizumab, a patient may qualify to receive up to a further 24 weeks of continuing treatment with omalizumab providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing omalizumab treatment in courses of up to 24 weeks providing they continue to sustain the response.

(2) Baseline measurements to determine response:

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the Asthma Control Questionnaire (ACQ; 5 item version) or oral corticosteroid dose submitted with the Initial authority application for omalizumab. For patients transitioned from the paediatric to the adolescent/adult restriction, the exacerbation history may also be used to determine response. However, prescribers may provide new baseline measurements when a new Initial treatment authority application is submitted and the Department of Human Services will assess response according to these revised baseline measurements.

(3) Re-commencement of treatment after a 6 month break in PBS-subsidised therapy:

A patient who wishes to trial a second or subsequent omalizumab treatment cycle, or an initial benralizumab or mepolizumab treatment cycle, following a break in PBS-subsidised therapy of at least 6 months, must re-qualify for initial treatment with respect to the indices of disease severity (oral corticosteroid dose, Asthma Control Questionnaire (ACQ-5) score, and relevant exacerbation history). Patients must have received optimised standard therapy, at adequate doses and for the minimum period specified, immediately prior to the time the new baseline assessments are performed.

(4) Monitoring of patients:

Anaphylaxis and anaphylactoid reactions have been reported following first or subsequent administration of omalizumab (see Product Information). Patients should be monitored post-injection, and medications for the treatment of anaphylactic reactions should be available for immediate use following administration of omalizumab. Patients should be informed that such reactions are possible and prompt medical attention should be sought if allergic reactions occur.

Note Special Pricing Arrangements apply.

Authority required

Uncontrolled severe allergic asthma

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a respiratory physician, clinical immunologist, allergist or general physician experienced in the management of patients with severe asthma.

Clinical criteria:

- Patient must be under the care of the same physician for at least 6 months; OR
- Patient must have been diagnosed by a multidisciplinary severe asthma clinic team, **AND**
- Patient must have a diagnosis of asthma confirmed and documented by a respiratory physician, clinical immunologist, allergist or general physician experienced in the management of patients with severe asthma, defined by the following

standard clinical features: (i) forced expiratory volume (FEV1) reversibility greater than or equal to 12% and greater than or equal to 200 mL at baseline within 30 minutes after administration of salbutamol (200 to 400 micrograms), or (ii) airway hyperresponsiveness defined as a greater than 20% decline in FEV1 during a direct bronchial provocation test or greater than 15% decline during an indirect bronchial provocation test, or (iii) peak expiratory flow (PEF) variability of greater than 15% between the two highest and two lowest peak expiratory flow rates during 14 days, **AND**

- Patient must have a duration of asthma of at least 1 year, **AND**
- Patient must have forced expiratory volume (FEV1) less than or equal to 80% predicted, documented on 1 or more occasions in the previous 12 months, **AND**
- Patient must have past or current evidence of atopy, documented by skin prick testing or RAST, **AND**
- Patient must have total serum human immunoglobulin E greater than or equal to 30 IU/mL, **AND**
- Patient must have failed to achieve adequate control with optimised asthma therapy, despite formal assessment of and adherence to correct inhaler technique, which has been documented, **AND**
- Patient must not receive more than 28 weeks of treatment under this restriction, **AND**
- The treatment must not be used in combination with, or within 6 months of treatment with, PBS-subsidised benralizumab or mepolizumab.

Population criteria:

- Patient must be aged 12 years or older.

Optimised asthma therapy includes:

- (i) Adherence to maximal inhaled therapy, including high dose inhaled corticosteroid (ICS) plus long-acting beta-2 agonist (LABA) therapy for at least 12 months, unless contraindicated or not tolerated; **AND**
- (ii) treatment with oral corticosteroids, either daily oral corticosteroids for at least 6 weeks, OR a cumulative dose of oral corticosteroids of at least 500 mg prednisolone equivalent in the previous 12 months, unless contraindicated or not tolerated.

If the requirement for treatment with optimised asthma therapy cannot be met because of contraindications according to the relevant TGA-approved Product Information and/or intolerances of a severity necessitating permanent treatment withdrawal, details of the contraindication and/or intolerance must be provided in the Authority application.

The initial IgE assessment must be no more than 12 months old at the time of application.

The following initiation criteria indicate failure to achieve adequate control and must be demonstrated in all patients at the time of the application:

- (a) an Asthma Control Questionnaire (ACQ-5) score of at least 2.0, as assessed in the previous month, **AND**
 - (b) while receiving optimised asthma therapy in the past 12 months, experienced at least 1 admission to hospital for a severe asthma exacerbation, OR 1 severe asthma exacerbation, requiring documented use of systemic corticosteroids (oral corticosteroids initiated or increased for at least 3 days, or parenteral corticosteroids) prescribed/supervised by a physician.
- The Asthma Control Questionnaire (5 item version) assessment of the patient's response to this initial course of treatment, and the assessment of oral corticosteroid dose, must be made at around 22 to 26 weeks after the first PBS-subsidised dose of this drug under this restriction so that there is adequate time for a response to be demonstrated and for the application for the first continuing therapy to be processed.

This assessment, which will be used to determine eligibility for the first continuing treatment, must be submitted within 4 weeks of the date of assessment, and no later than 2 weeks prior to the patient completing their current treatment course, to avoid an interruption to supply. Where a response assessment is not undertaken and submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

A patient who fails to respond to a course of PBS-subsidised omalizumab for the treatment of uncontrolled severe allergic asthma will not be eligible to receive further PBS-subsidised treatment with benralizumab, omalizumab or mepolizumab for this condition within 6 months of the date on which treatment was ceased.

At the time of the authority application, medical practitioners should request the appropriate maximum quantity and number of repeats to provide for an initial course of omalizumab consisting of the recommended number of doses for the baseline IgE level and body weight of the patient (refer to the TGA-approved Product Information) to be administered every 2 or 4 weeks.

A multidisciplinary severe asthma clinic team comprises of:

- A respiratory physician; and
- A pharmacist, nurse or asthma educator.

Omalizumab must not be used concurrently with benralizumab or mepolizumab, or within 6 months of each other. A patient is required to have ceased treatment with benralizumab or mepolizumab for 6 months prior to initiating treatment with omalizumab.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Severe Allergic Asthma PBS Authority Application - Supporting Information Form, which includes the following:
 - (i) details of prior optimised asthma drug therapy (date of commencement and duration of therapy); and
 - (ii) details of severe exacerbation/s experienced in the past 12 months while receiving optimised asthma therapy (date and treatment); and
- (c) the IgE pathology report; and
- (d) a completed Asthma Control Questionnaire (ACQ-5) calculation sheet including the date of assessment of the patient's symptoms.

Note The Department of Human Services website (www.humanservices.gov.au) has details of the accepted toxicities, including severity, which will be accepted for the purposes of exempting a patient from the requirement of treatment with optimised asthma therapy.

Note For copies of the ACQ and the calculation sheets please contact Novartis Medical Information on 1800 671 203 or medinfo.phauno@novartis.com

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au. Applications for authority to prescribe should be forwarded to:
Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Formal assessment and correction of inhaler technique should be performed in accordance with the National Asthma Council (NAC) Information Paper for Health Professionals on Inhaler Technique (available at www.humanservices.gov.au or www.nationalasthma.org.au); the assessment and adherence to correct technique should be documented in the patient's medical records. Patients can obtain support with inhaler technique through their local Asthma Foundation (1800 645 130).

Authority required

Uncontrolled severe allergic asthma
Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have a documented history of severe allergic asthma, **AND**
- Patient must have demonstrated or sustained an adequate response to PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction, **AND**
- The treatment must not be used in combination with, or within 6 months of treatment with, PBS-subsidised benralizumab or mepolizumab.

Treatment criteria:

- Must be treated by a respiratory physician, clinical immunologist, allergist or general physician experienced in the management of patients with severe asthma.

Population criteria:

- Patient must be aged 12 years or older.

An adequate response to omalizumab treatment is defined as:

- (a) a reduction in the Asthma Control Questionnaire (ACQ-5) score of at least 0.5 from baseline, OR
(b) maintenance oral corticosteroid dose reduced by at least 25% from baseline, and no deterioration in ACQ-5 score from baseline, OR

(c) a reduction in the time-adjusted exacerbation rates compared to the 12 months prior to baseline (this criterion is only applicable for patients transitioned from the paediatric to the adolescent/adult restriction).

All applications for second and subsequent continuing treatments with this drug must include a measurement of response to the prior course of therapy. The Asthma Control Questionnaire (5 item version) assessment of the patient's response to the prior course of treatment, the assessment of oral corticosteroid dose or the assessment of time adjusted exacerbation rate must be made at around 18 to 22 weeks after the first PBS-subsidised dose of this drug under this restriction so that there is adequate time for a response to be demonstrated and for the application for continuing therapy to be processed.

The assessment should, where possible, be completed by the same physician who initiated treatment with this drug. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted within 4 weeks of the date of assessment, and no later than 2 weeks prior to the patient completing their current treatment course, to avoid an interruption to supply. Where a response assessment is not undertaken and submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

A patient who fails to respond to a course of PBS-subsidised omalizumab for the treatment of uncontrolled severe allergic asthma will not be eligible to receive further PBS-subsidised treatment with omalizumab for this condition within 6 months of the date on which treatment was ceased.

At the time of the authority application, medical practitioners should request the appropriate quantity and number of repeats to provide for a continuing course of omalizumab consisting of the recommended number of doses for the baseline IgE level and body weight of the patient (refer to the TGA-approved Product Information), sufficient for up to 24 weeks of therapy.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
(b) a completed Severe Allergic Asthma PBS Authority Application and Supporting Information Form which includes details of maintenance oral corticosteroid dose; or
(c) a completed Asthma Control Questionnaire (ACQ-5) calculation sheet including the date of assessment of the patient's symptoms and is endorsed with the signature of the prescriber; for patients transitioned from the paediatric to the adolescent/adult restrictions an exacerbation calculation sheet may be submitted.

Note If the same physician cannot assess the patient please call the Department of Human Services on 1800 700 270.

Note For copies of the ACQ and the calculation sheets please contact Novartis Medical Information on 1800 671 203 or medinfo.phauno@novartis.com

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au. Applications for authority to prescribe should be forwarded to:
Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Formal assessment and correction of inhaler technique should be performed in accordance with the National Asthma Council (NAC) Information Paper for Health Professionals on Inhaler Technique (available at www.humanservices.gov.au or www.nationalasthma.org.au); the assessment and adherence to correct technique should be documented in the patient's medical records. Patients can obtain support with inhaler technique through their local Asthma Foundation (1800 645 130).

Authority required

Uncontrolled severe allergic asthma

Treatment Phase: Initial and continuing treatment - balance of supply

Treatment criteria:

- Must be treated by a respiratory physician, clinical immunologist, allergist or general physician experienced in the management of patients with severe asthma.

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the Initial treatment restriction to complete 28 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the Continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 28 weeks treatment available under the Initial restriction or up to 24 weeks treatment available under the Continuing restriction.

Note Authority approval for sufficient therapy to complete a maximum of 28 weeks of treatment under the initial restriction or 24 weeks of treatment under the continuing restriction may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

omalizumab 150 mg/mL injection, 1 mL syringe

10109C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	410.00	Xolair [NV]

omalizumab 75 mg/0.5 mL injection, 0.5 mL syringe

10118M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	205.00	Xolair [NV]

▪ **OMALIZUMAB**

Note TREATMENT OF PAEDIATRIC PATIENTS WITH UNCONTROLLED SEVERE ALLERGIC ASTHMA

Patients are eligible to commence an 'omalizumab treatment cycle' (initial treatment course with or without continuing treatment course/s) if they satisfy the eligibility criteria as detailed under the initial treatment restriction. Once a patient has either failed to achieve or maintain a response to omalizumab, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 6 month break in PBS-subsidised omalizumab therapy before they are eligible to commence the next cycle. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised omalizumab treatment is stopped to the date of the first application for initial treatment with omalizumab under the new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised omalizumab therapy.

(a) Initial treatment:

Applications for initial treatment should be made where a patient has received no prior PBS-subsidised omalizumab treatment in this treatment cycle and wishes to commence such therapy.

All applications for initial treatment will be limited to provide for a maximum of 28 weeks of therapy for omalizumab.

(b) Continuing treatment:

Following the completion of the initial treatment course with omalizumab, a patient may qualify to receive up to a further 24 weeks of continuing treatment with omalizumab providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing omalizumab treatment in courses of up to 24 weeks providing they continue to sustain the response.

(2) Baseline measurements to determine response:

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the Asthma Control Questionnaire (ACQ; 5 item version) or ACQ-IA, systemic corticosteroid dose and time-adjusted exacerbation rate, submitted with the Initial authority application for omalizumab. However, prescribers may provide new baseline measurements when a new Initial treatment authority application is submitted and The Department of Human Services will assess response according to these revised baseline measurements.

(3) Re-commencement of treatment after a 6 month break in PBS-subsidised therapy:

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised omalizumab therapy of at least 6 months, must re-qualify for initial treatment with respect to the indices of disease severity (systemic corticosteroid dose, Asthma Control Questionnaire (ACQ-5) score or ACQ-IA, and relevant exacerbation history). Patients must have received optimised standard therapy, at adequate doses and for the minimum period specified, immediately prior to the time the new baseline assessments are performed.

(4) Monitoring of patients:

Anaphylaxis and anaphylactoid reactions have been reported following first or subsequent administration of omalizumab (see Product Information). Patients should be monitored post-injection, and medications for the treatment of anaphylactic reactions should be available for immediate use following administration of omalizumab. Patients should be informed that such reactions are possible and prompt medical attention should be sought if allergic reactions occur.

Note Special Pricing Arrangements apply.

Authority required

Uncontrolled severe allergic asthma

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have a diagnosis of asthma confirmed and documented by a paediatric respiratory physician, clinical immunologist, or allergist; or paediatrician or general physician experienced in the management of patients with severe asthma in consultation with a respiratory physician, defined by the following standard clinical features: forced expiratory volume (FEV1) reversibility or airway hyperresponsiveness or peak expiratory flow (PEF) variability, **AND**
- Patient must have a duration of asthma of at least 1 year, **AND**
- Patient must have past or current evidence of atopy, documented by skin prick testing or an in vitro measure of specific IgE, **AND**
- Patient must have total serum human immunoglobulin E greater than or equal to 30 IU/mL, **AND**
- Patient must have failed to achieve adequate control with optimised asthma therapy, despite formal assessment of and adherence to correct inhaler technique, which has been documented, **AND**
- Patient must not receive more than 28 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 6 to less than 12 years.

Treatment criteria:

- Must be treated by a paediatric respiratory physician, clinical immunologist, allergist; or paediatrician or general physician experienced in the management of patients with severe asthma, in consultation with a respiratory physician.

Clinical criteria:

- Patient must be under the care of the same physician for at least 6 months.

Optimised asthma therapy includes:

(i) Adherence to optimal inhaled therapy, including high dose inhaled corticosteroid (ICS) and long-acting beta-2 agonist (LABA) therapy for at least six months. If LABA therapy is contraindicated, not tolerated or not effective, montelukast, cromoglycate or nedocromil may be used as an alternative; **AND**

(ii) treatment with at least 2 courses of oral or IV corticosteroids (daily or alternate day maintenance treatment courses, or 3-5 day exacerbation treatment courses), in the previous 12 months, unless contraindicated or not tolerated.

If the requirement for treatment with optimised asthma therapy cannot be met because of contraindications (including those specified in the relevant TGA-approved Product Information) and/or intolerances of a severity necessitating permanent treatment withdrawal, details of the contraindication and/or intolerance must be provided in the Authority application.

The initial IgE assessment must be no more than 12 months old at the time of application.

The following initiation criteria indicate failure to achieve adequate control and must be demonstrated in all patients at the time of the application:

(a) An Asthma Control Questionnaire (ACQ-5) score of at least 2.0, as assessed in the previous month (for children aged 6 to 10 years it is recommended that the Interviewer Administered version - the ACQ-IA be used), **AND**

(b) while receiving optimised asthma therapy in the previous 12 months, experienced at least 1 admission to hospital for a severe asthma exacerbation, OR 1 severe asthma exacerbation, requiring documented use of systemic corticosteroids (oral corticosteroids initiated or increased for at least 3 days, or parenteral corticosteroids) prescribed/supervised by a physician. The Asthma Control Questionnaire (5 item version) or ACQ-IA assessment of the patient's response to this initial course of treatment, the assessment of oral corticosteroid dose, and the assessment of exacerbation rate must be made at around 22 to 26 weeks after the first dose so that there is adequate time for a response to be demonstrated and for the application for continuing therapy to be processed.

This assessment, which will be used to determine eligibility for continuing treatment, must be submitted within 4 weeks of the date of assessment, and no later than 2 weeks prior to the patient completing their current treatment course, to avoid an interruption to supply. Where a response assessment is not undertaken and submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with omalizumab.

A patient who fails to respond to a course of PBS-subsidised omalizumab for the treatment of uncontrolled severe allergic asthma will not be eligible to receive further PBS-subsidised treatment with omalizumab for this condition within 6 months of the date on which treatment was ceased.

At the time of the authority application, medical practitioners should request the appropriate maximum quantity and number of repeats to provide for an initial course of omalizumab of up to 28 weeks, consisting of the recommended number of doses for the baseline IgE level and body weight of the patient (refer to the TGA-approved Product Information) to be administered every 2 or 4 weeks.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Paediatric Severe Allergic Asthma Initial PBS Authority Application - Supporting Information form, which includes the following:
 - (i) details of prior optimised asthma drug therapy (dosage, date of commencement and duration of therapy); and
 - (ii) details of severe exacerbation/s experienced in the past 12 months while receiving optimised asthma therapy (date and treatment); and
 - (iii) acknowledgement signed by a parent or authorised guardian; and
- (c) a copy of the IgE pathology report; and
- (d) a completed Asthma Control Questionnaire (ACQ-5) or the Asthma Control Questionnaire interviewer administered version (ACQ-IA) calculation sheet including the date of assessment of the patient's symptoms and is endorsed with the prescriber's signature.

Note The Department of Human Services website (www.humanservices.gov.au) has details of the accepted toxicities, including severity, which will be accepted for the purposes of exempting a patient from the requirement of treatment with optimised asthma therapy.

Note For copies of the ACQ please contact Novartis Medical Information on 1800 671 203 or medinfo.phauno@novartis.com

Note It is recommended that an application for continuing treatment is submitted at the time of the 22 to 26 week assessment, to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised omalizumab treatment.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Formal assessment and correction of inhaler technique should be performed in accordance with the National Asthma Council (NAC) Information Paper for Health Professionals on Inhaler Technique (available at www.humanservices.gov.au or www.nationalasthma.org.au); the assessment and adherence to correct technique should be documented in the patient's medical records. Patients can obtain support with inhaler technique through their local Asthma Foundation (1800 645 130).

Authority required

Uncontrolled severe allergic asthma

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have a documented history of severe allergic asthma, **AND**
- Patient must have demonstrated or sustained an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Treatment criteria:

- Must be treated by a paediatric respiratory physician, clinical immunologist, allergist; or paediatrician or general physician experienced in the management of patients with severe asthma, in consultation with a respiratory physician.

An adequate response to omalizumab treatment is defined as:

(a) a reduction in the Asthma Control Questionnaire (ACQ-5) or ACQ-IA score of at least 0.5 from baseline, OR

(b) maintenance oral corticosteroid dose reduced by at least 25% from baseline, and no deterioration in ACQ-5 or ACQ-IA score from baseline, OR

(c) a reduction in the time-adjusted exacerbation rates compared to the 12 months prior to baseline.

All applications for continuing treatment with omalizumab must include a measurement of response to the prior course of therapy. The Asthma Control Questionnaire (5 item version) or Asthma Control Questionnaire interviewer administered version (ACQ-IA) assessment of the patient's response to the prior course of treatment, the assessment of systemic corticosteroid dose, and the assessment of time-adjusted exacerbation rate must be made at around 18 to 22 weeks after the first dose of PBS-subsidised omalizumab so that there is adequate time for a response to be demonstrated and for the application for continuing therapy to be processed.

The first assessment should, where possible, be completed by the same physician who initiated treatment with omalizumab. This assessment, which will be used to determine eligibility for continuing treatment, must be submitted within 4 weeks of the date of assessment, and no later than 2 weeks prior to the patient completing their current treatment course, to avoid an interruption to supply. Where a response assessment is not undertaken and submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with omalizumab.

A patient who fails to respond to a course of PBS-subsidised omalizumab for the treatment of uncontrolled severe allergic asthma will not be eligible to receive further PBS-subsidised treatment with omalizumab for this condition within 6 months of the date on which treatment was ceased.

At the time of the authority application, medical practitioners should request the appropriate quantity and number of repeats to provide for a continuing course of omalizumab consisting of the recommended number of doses for the baseline IgE level and body weight of the patient (refer to the TGA-approved Product Information), sufficient for 24 weeks of therapy.

The authority application must be made in writing and must include:

(a) a completed authority prescription form; and

(b) a completed Paediatric Severe Allergic Asthma Continuing PBS Authority Application - Supporting Information form which includes details of maintenance oral corticosteroid dose; and

(c) a completed Asthma Control Questionnaire (ACQ-5) or the Asthma Control Questionnaire interviewer administered version (ACQ-IA) calculation sheet including the date of assessment of the patient's symptoms and is endorsed with the signature of the prescriber.

Note If the same physician cannot assess the patient please call the Department of Human Services on 1800 700 270.

Note For copies of the ACQ please contact Novartis Medical Information on 1800 671 203 or medinfo.phauno@novartis.com

Note It is recommended that an application for continuing treatment is submitted at the time of the 18 to 22 week assessment, to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised omalizumab treatment.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Formal assessment and correction of inhaler technique should be performed in accordance with the National Asthma Council (NAC) Information Paper for Health Professionals on Inhaler Technique (available at www.humanservices.gov.au or www.nationalasthma.org.au); the assessment and adherence to correct technique should be documented in the patient's medical records. Patients can obtain support with inhaler technique through their local Asthma Foundation (1800 645 130).

Authority required

Uncontrolled severe allergic asthma

Treatment Phase: Initial and continuing treatment - balance of supply

Treatment criteria:

- Must be treated by a paediatric respiratory physician, clinical immunologist, allergist; or paediatrician or general physician experienced in the management of patients with severe asthma, in consultation with a respiratory physician.

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the Initial treatment restriction to complete 28 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the Continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 28 weeks treatment available under the Initial restriction or up to 24 weeks treatment available under the Continuing restriction.

Note Authority approval for sufficient therapy to complete a maximum of 28 weeks of treatment under the initial restriction or 24 weeks of treatment under the continuing restriction may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

omalizumab 150 mg/mL injection, 1 mL syringe

10973M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	410.00	Xolair [NV]

omalizumab 75 mg/0.5 mL injection, 0.5 mL syringe

10967F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	..	..	205.00	Xolair [NV]

■ **COUGH AND COLD PREPARATIONS**

EXPECTORANTS, EXCL. COMBINATIONS WITH COUGH SUPPRESSANTS

Mucolytics

■ **DORNASE ALFA**

Note It is highly desirable that all patients be included in the national cystic fibrosis patient database.

Authority required (STREAMLINED)

5740

Cystic fibrosis

Population criteria:

- Patient must be 5 years of age or older.

Patient must be assessed at a cystic fibrosis clinic/centre which is under the control of specialist respiratory physicians with experience and expertise in the management of cystic fibrosis or by a specialist physician or paediatrician in consultation with such a unit.

Prior to therapy with this drug, a baseline measurement of forced expiratory volume in 1 second (FEV1) must be undertaken during a stable period of the disease.

Initial therapy is limited to 3 months treatment with dornase alfa at a dose of 2.5 mg daily.

To be eligible for continued PBS-subsidised treatment with this drug following 3 months of initial treatment:

- (1) the patient must demonstrate no deterioration in FEV1 compared to baseline; AND
- (2) the patient or the patient's family (in the case of paediatric patients) and the treating physician(s) must report a benefit in the clinical status of the patient.

Further reassessments must be undertaken and documented at six-monthly intervals. Therapy with this drug should cease if there is not general agreement of benefit as there is always the possibility of harm from unnecessary use.

Authority required (STREAMLINED)

5634

Cystic fibrosis

Clinical criteria:

- Patient must have a severe clinical course with frequent respiratory exacerbations or chronic respiratory symptoms (including chronic or recurrent cough, wheeze or tachypnoea) requiring hospital admissions more frequently than 3 times per year; OR
- Patient must have significant bronchiectasis on chest high resolution computed tomography scan; OR
- Patient must have severe cystic fibrosis bronchiolitis with persistent wheeze non-responsive to conventional medicines; OR
- Patient must have severe physiological deficit measure by forced oscillation technique or multiple breath nitrogen washout and failure to respond to conventional therapy.

Population criteria:

- Patient must be less than 5 years of age.

Patient must be assessed at a cystic fibrosis clinic/centre which is under the control of specialist respiratory physicians with experience and expertise in the management of cystic fibrosis or by a specialist physician or paediatrician in consultation with such a unit.

Following an initial 6 months therapy, a comprehensive assessment must be undertaken and documented. Treatment with this drug should cease if there is not agreement of benefit, as there is always the possibility of harm from unnecessary use. Further reassessments must be undertaken and documented at six-monthly intervals.

Authority required (STREAMLINED)

5635

Cystic fibrosis

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have initiated treatment with dornase alfa at an age of less than 5 years, **AND**
- Patient must have undergone a comprehensive assessment which documents agreement that dornase alfa treatment is continuing to produce worthwhile benefit.

Population criteria:

- Patient must be 5 years of age or older.

Further reassessments must be undertaken and documented at six-monthly intervals. Treatment with this drug should cease if there is not agreement of benefit as there is always the possibility of harm from unnecessary use.

dornase alfa 2.5 mg/2.5 mL inhalation solution, 30 x 2.5 mL ampoules

5704F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*1916.92	Pulmozyme [RO]

■ **MANNITOL**

Note It is highly desirable that all patients be included in the national cystic fibrosis patient database.

Authority required (STREAMLINED)

7362

Cystic fibrosis

Clinical criteria:

- The treatment must be as monotherapy, **AND**
- Patient must be intolerant or inadequately responsive to dornase alfa.

Population criteria:

- Patient must be 6 years of age or older.

Patient must have been assessed for bronchial hyperresponsiveness as per the TGA approved Product Information initiation dose assessment for this drug, prior to therapy with this drug, with a negative result.

Patient must be assessed at a cystic fibrosis clinic/centre which is under the control of specialist respiratory physicians with experience and expertise in the management of cystic fibrosis or by a specialist physician or paediatrician in consultation with such a unit.

Prior to therapy with this drug, a baseline measurement of forced expiratory volume in 1 second (FEV1) must be undertaken during a stable period of the disease.

Initial therapy is limited to 3 months treatment with mannitol at a dose of 400 mg twice daily.

To be eligible for continued PBS-subsidised treatment with this drug following 3 months of initial treatment:

- (1) the patient must demonstrate no deterioration in FEV1 compared to baseline; **AND**
- (2) the patient or the patient's family (in the case of paediatric patients) and the treating physician(s) must report a benefit in the clinical status of the patient.

Further reassessments must be undertaken and documented at six-monthly intervals. Therapy with this drug should cease if there is not general agreement of benefit as there is always the possibility of harm from unnecessary use.

Authority required (STREAMLINED)

7367

Cystic fibrosis

Clinical criteria:

- The treatment must be in combination with dornase alfa, **AND**
- Patient must be inadequately responsive to dornase alfa, **AND**
- Patient must have trialled hypertonic saline for this condition.

Population criteria:

- Patient must be 6 years of age or older.

Patient must have been assessed for bronchial hyperresponsiveness as per the TGA approved Product Information initiation dose assessment for this drug, prior to therapy with this drug, with a negative result.

Patient must be assessed at a cystic fibrosis clinic/centre which is under the control of specialist respiratory physicians with experience and expertise in the management of cystic fibrosis or by a specialist physician or paediatrician in consultation with such a unit.

Prior to therapy with this drug, a baseline measurement of forced expiratory volume in 1 second (FEV1) must be undertaken during a stable period of the disease.

Initial therapy is limited to 3 months treatment with mannitol at a dose of 400 mg twice daily.

To be eligible for continued PBS-subsidised treatment with this drug following 3 months of initial treatment:

- (1) the patient must demonstrate no deterioration in FEV1 compared to baseline; **AND**
- (2) the patient or the patient's family (in the case of paediatric patients) and the treating physician(s) must report a benefit in the clinical status of the patient.

Further reassessments must be undertaken and documented at six-monthly intervals. Therapy with this drug should cease if there is not general agreement of benefit as there is always the possibility of harm from unnecessary use.

mannitol 40 mg powder for inhalation, 280 capsules

2015C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	4	5	..	*1789.20	bronchitol [XA]

OTHER RESPIRATORY SYSTEM PRODUCTS

OTHER RESPIRATORY SYSTEM PRODUCTS

Other respiratory system products

■ IVACAFTOR

Note No increase in the maximum number of repeats may be authorised.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Cystic fibrosis

Treatment Phase: Initial treatment - New patients

Clinical criteria:

- Patient must be assessed through a cystic fibrosis clinic/centre which is under the control of specialist respiratory physicians with experience and expertise in the management of cystic fibrosis. If attendance at such a unit is not possible because of geographical isolation, management (including prescribing) may be in consultation with such a unit, **AND**
- Patient must have G551D mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene on at least 1 allele; OR
- Patient must have other gating (class III) mutation in the CFTR gene on at least 1 allele, **AND**
- Patient must have a sweat chloride value of at least 60 mmol/L by quantitative pilocarpine iontophoresis, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction, **AND**
- The treatment must be given concomitantly with standard therapy for this condition.

Population criteria:

- Patient must be aged 12 months or older.

Patients receiving PBS-subsidised ivacaftor must be registered in the Australian Cystic Fibrosis Database Registry.

Treatment must not be given to a patient who has an acute upper or lower respiratory infection, pulmonary exacerbation, or changes in therapy (including antibiotics) for pulmonary disease in the last 4 weeks prior to commencing this drug.

Dosage of ivacaftor must not exceed the dose of one tablet (150 mg) or one sachet twice a week, if the patient is concomitantly receiving one of the following strong CYP3A4 drugs inhibitors: boceprevir, clarithromycin, conivaptan, indinavir, itraconazole, ketoconazole, lopinavir/ritonavir, mibefradil, nefazodone, nelfinavir, posaconazole, ritonavir, saquinavir, telaprevir, telithromycin, voriconazole. Where a patient is concomitantly receiving a strong CYP3A4 inhibitor, a single supply of 56 tablets or sachets of ivacaftor will last for 28 weeks.

Dosage of ivacaftor must not exceed the dose of one tablet (150 mg) or one sachet once daily, if the patient is concomitantly receiving one of the following moderate CYP3A4 inhibitors: amprenavir, aprepitant, atazanavir, darunavir/ritonavir, diltiazem, erythromycin, fluconazole, fosamprenavir, imatinib, verapamil. Where a patient is concomitantly receiving a moderate CYP3A4 inhibitor, a single supply of 56 tablets or sachets of ivacaftor will last for 8 weeks.

Ivacaftor is not PBS-subsidised for this condition as a sole therapy.

Ivacaftor is not PBS-subsidised for this condition in a patient who is currently receiving one of the following CYP3A4 inducers:

Strong CYP3A4 inducers: avasimibe, carbamazepine, phenobarbital, phenytoin, rifabutin, rifampicin, St. John's wort

Moderate CYP3A4 inducers: bosentan, efavirenz, etravirine, modafinil, nafcillin

Weak CYP3A4 inducers: armodafinil, echinacea, pioglitazone, rufinamide.

The authority application must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Cystic Fibrosis Ivacaftor Authority Application Supporting Information Form; and
- (3) a copy of the pathology report detailing the molecular testing for G551D mutation or other gating (class III) mutation on the CFTR gene; and
- (4) the result of a FEV1 measurement performed within a month prior to the date of application, if aged from 6 years or older. Note: FEV1, must be measured in an accredited pulmonary function laboratory, with documented no acute infective exacerbation at the time FEV1 is measured; and
- (5) a copy of a current medication history, including any CYP3A4 inhibitors and/or CYP3A4 inducers; and
- (6) a copy of a sweat chloride result; and
- (7) height and weight measurements at the time of application; and
- (8) a baseline measurement of the number of days of CF-related hospitalisation (including hospital-in-the home) in the previous 12 months.

Authority required

Cystic fibrosis

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must be assessed through a cystic fibrosis clinic/centre which is under the control of specialist respiratory physicians with experience and expertise in the management of cystic fibrosis. If attendance at such a unit is not possible because of geographical isolation, management (including prescribing) may be in consultation with such a unit, **AND**
- Patient must have received PBS-subsidised initial therapy with ivacaftor, given concomitantly with standard therapy, for this condition, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction, **AND**
- The treatment must be given concomitantly with standard therapy for this condition.

Population criteria:

- Patient must be aged 12 months or older.

Patients receiving PBS-subsidised ivacaftor must be registered in the Australian Cystic Fibrosis Database Registry.

Treatment must not be given to a patient who has an acute upper or lower respiratory infection, pulmonary exacerbation, or changes in therapy (including antibiotics) for pulmonary disease in the last 4 weeks prior to commencing this drug.

Patients who have an acute infective exacerbation at the time of assessment for continuing therapy may receive an additional one month's supply in order to enable the assessment to be repeated following resolution of the exacerbation.

Dosage of ivacaftor must not exceed the dose of one tablet (150 mg) or one sachet twice a week, if the patient is concomitantly receiving one of the following strong CYP3A4 drugs inhibitors: boceprevir, clarithromycin, conivaptan, indinavir, itraconazole, ketoconazole, lopinavir/ritonavir, mibefradil, nefazodone, nelfinavir, posaconazole, ritonavir, saquinavir, telaprevir, telithromycin, voriconazole. Where a patient is concomitantly receiving a strong CYP3A4 inhibitor, a single supply of 56 tablets or sachets of ivacaftor will last for 28 weeks.

Dosage of ivacaftor must not exceed the dose of one tablet (150 mg) or one sachet once daily, if the patient is concomitantly receiving one of the following moderate CYP3A4 inhibitors: amprenavir, aprepitant, atazanavir, darunavir/ritonavir, diltiazem, erythromycin, fluconazole, fosamprenavir, imatinib, verapamil. Where a patient is concomitantly receiving a moderate CYP3A4 inhibitor, a single supply of 56 tablets or sachets of ivacaftor will last for 8 weeks.

Ivacaftor is not PBS-subsidised for this condition as a sole therapy.

Ivacaftor is not PBS-subsidised for this condition in a patient who is currently receiving one of the following CYP3A4 inducers:

Strong CYP3A4 inducers: avasimibe, carbamazepine, phenobarbital, phenytoin, rifabutin, rifampicin, St. John's wort

Moderate CYP3A4 inducers: bosentan, efavirenz, etravirine, modafinil, nafcillin

Weak CYP3A4 inducers: armodafinil, echinacea, pioglitazone, rufinamide.

The authority application must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Cystic Fibrosis Ivacaftor Authority Continuing Application Supporting Information Form; and
- (3) the result of a FEV1 measurement performed within one month prior to the date of application, if aged 6 years or older. Note: FEV1, must be measured in an accredited pulmonary function laboratory, with documented no acute infective exacerbation at the time FEV1 is measured; and
- (4) a copy of a current medication history, including any CYP3A4 inhibitors and/or CYP3A4 inducers; and
- (5) height and weight measurements at the time of application; and
- (6) a measurement of number of days of CF-related hospitalisation (including hospital in the home) in the previous 6 months.

ivacaftor 75 mg granules, 4 x 14 sachets

11098D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	22500.00	Kalydeco [VR]

ivacaftor 150 mg tablet, 56

10170G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	22500.00	Kalydeco [VR]

ivacaftor 50 mg granules, 4 x 14 sachets

11105L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	22500.00	Kalydeco [VR]

■ LUMACAFITOR + IVACAFITOR**Note Managed Access Program:**

This medicine has been listed on the PBS via a Managed Access Program (MAP). The Pharmaceutical Benefits Advisory Committee (PBAC) made its recommendation on the basis of 24 weeks of data in children aged 6 - 11 years and 96 weeks of data in patients aged 12 years and over. Information about the long term benefits of this medicine will be collected and analysed under this MAP.

For more information on Managed Access Programs, please visit <http://www.pbs.gov.au/info/industry/listing/elements/pbac-meetings/pbac-outcomes/2015-03/march-2015-other-matters-managed-access-programme-framework>.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Cystic fibrosis

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a specialist respiratory physician with expertise in cystic fibrosis or in consultation with a specialist respiratory physician with expertise in cystic fibrosis if attendance is not possible due to geographic isolation, **AND**
- Must be treated in a centre with expertise in cystic fibrosis or in consultation with a centre with expertise in cystic fibrosis if attendance is not possible due to geographic isolation.

Clinical criteria:

- Patient must be homozygous for the F508del mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene, **AND**
- The treatment must be given concomitantly with standard therapy for this condition.

Population criteria:

- Patient must be aged between 6 and 11 years inclusive.

The patient must be registered in the Australian Cystic Fibrosis Database Registry.

Treatment must not be given to a patient who has an acute upper or lower respiratory infection, pulmonary exacerbation, or changes in therapy (including antibiotics) for pulmonary disease in the last 4 weeks prior to commencing this drug.

The authority application must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Cystic Fibrosis Lumacaftor with Ivacaftor Authority Application Supporting Information Form; and
- (3) a copy of the pathology report detailing the molecular testing for the patient being homozygous for the F508del mutation on the CFTR gene; and
- (4) the result of a FEV1 measurement performed within a month prior to the date of application. Note: FEV1 must be measured in an accredited pulmonary function laboratory, with documented no acute infective exacerbation at the time FEV1 is measured; and
- (5) confirmation that the patient has either chronic sinopulmonary disease or gastrointestinal and nutritional abnormalities; and
- (6) a copy of a current medication history, including any CPY3A inhibitors and/or inducers; and
- (7) height and weight measurements at the time of application; and
- (8) a baseline measurement of the number of days of CF-related hospitalisation (including hospital-in-the home) in the previous 12 months.

Authority required

Cystic fibrosis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a specialist respiratory physician with expertise in cystic fibrosis or in consultation with a specialist respiratory physician with expertise in cystic fibrosis if attendance is not possible due to geographic isolation, **AND**
- Must be treated in a centre with expertise in cystic fibrosis or in consultation with a centre with expertise in cystic fibrosis if attendance is not possible due to geographic isolation.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be given concomitantly with standard therapy for this condition.

Population criteria:

- Patient must be aged between 6 and 11 years inclusive.

Treatment must not be given to a patient who has an acute upper or lower respiratory infection, pulmonary exacerbation, or changes in therapy (including antibiotics) for pulmonary disease in the last 4 weeks prior to commencing this drug.

Patients who have an acute infective exacerbation at the time of assessment for continuing therapy may receive an additional one month's supply in order to enable the assessment to be repeated following resolution of the exacerbation.

The authority application must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Cystic Fibrosis Lumacaftor with Ivacaftor Continuing Authority Application Supporting Information Form; and
- (3) the result of a FEV1 measurement performed within a month prior to the date of application. Note: FEV1, must be measured in an accredited pulmonary function laboratory, with documented no acute infective exacerbation at the time FEV1 is measured; and
- (4) a copy of a current medication history, including any CYP3A inhibitors and/or inducers; and
- (5) height and weight measurements at the time of application; and
- (6) the number of days of CF-related hospitalisation (including hospital-in-the home) in the previous 6 months.

lumacaftor 100 mg + ivacaftor 125 mg tablet, 112

11465K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	18750.00	Orkambi [VR]

■ LUMACAFITOR + IVACAFITOR

Note Managed Access Program:

This medicine has been listed on the PBS via a Managed Access Program (MAP). The Pharmaceutical Benefits Advisory Committee (PBAC) made its recommendation on the basis of 24 weeks of data in children aged 6 - 11 years and 96 weeks of data in patients aged 12 years and over. Information about the long term benefits of this medicine will be collected and analysed under this MAP.

For more information on Managed Access Programs, please visit <http://www.pbs.gov.au/info/industry/listing/elements/pbac-meetings/pbac-outcomes/2015-03/march-2015-other-matters-managed-access-programme-framework>.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Cystic fibrosis

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a specialist respiratory physician with expertise in cystic fibrosis or in consultation with a specialist respiratory physician with expertise in cystic fibrosis if attendance is not possible due to geographic isolation, **AND**
- Must be treated in a centre with expertise in cystic fibrosis or in consultation with a centre with expertise in cystic fibrosis if attendance is not possible due to geographic isolation.

Clinical criteria:

- Patient must be homozygous for the F508del mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene, **AND**
- The treatment must be given concomitantly with standard therapy for this condition.

Population criteria:

- Patient must be 12 years of age or older.

The patient must be registered in the Australian Cystic Fibrosis Database Registry.

Treatment must not be given to a patient who has an acute upper or lower respiratory infection, pulmonary exacerbation, or changes in therapy (including antibiotics) for pulmonary disease in the last 4 weeks prior to commencing this drug.

The authority application must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Cystic Fibrosis Lumacaftor with Ivacaftor Authority Application Supporting Information Form; and
- (3) a copy of the pathology report detailing the molecular testing for the patient being homozygous for the F508del mutation on the CFTR gene; and
- (4) the result of a FEV1 measurement performed within a month prior to the date of application. Note: FEV1 must be measured in an accredited pulmonary function laboratory, with documented no acute infective exacerbation at the time FEV1 is measured; and
- (5) confirmation that the patient has either chronic sinopulmonary disease or gastrointestinal and nutritional abnormalities; and
- (6) a copy of a current medication history, including any CPY3A inhibitors and/or inducers; and
- (7) height and weight measurements at the time of application; and
- (8) a baseline measurement of the number of days of CF-related hospitalisation (including hospital-in-the home) in the previous 12 months.

Authority required

Cystic fibrosis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a specialist respiratory physician with expertise in cystic fibrosis or in consultation with a specialist respiratory physician with expertise in cystic fibrosis if attendance is not possible due to geographic isolation, **AND**
- Must be treated in a centre with expertise in cystic fibrosis or in consultation with a centre with expertise in cystic fibrosis if attendance is not possible due to geographic isolation.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be given concomitantly with standard therapy for this condition.

Population criteria:

- Patient must be 12 years of age or older.

Treatment must not be given to a patient who has an acute upper or lower respiratory infection, pulmonary exacerbation, or changes in therapy (including antibiotics) for pulmonary disease in the last 4 weeks prior to commencing this drug.

Patients who have an acute infective exacerbation at the time of assessment for continuing therapy may receive an additional one month's supply in order to enable the assessment to be repeated following resolution of the exacerbation.

The authority application must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Cystic Fibrosis Lumacaftor with Ivacaftor Continuing Authority Application Supporting Information Form; and
- (3) the result of a FEV1 measurement performed within a month prior to the date of application. Note: FEV1, must be measured in an accredited pulmonary function laboratory, with documented no acute infective exacerbation at the time FEV1 is measured; and
- (4) a copy of a current medication history, including any CYP3A inhibitors and/or inducers; and
- (5) height and weight measurements at the time of application; and
- (6) the number of days of CF-related hospitalisation (including hospital-in-the home) in the previous 6 months.

lumacaftor 200 mg + ivacaftor 125 mg tablet, 112

11466L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	1	5	..	18750.00	Orkambi [VR]

VARIOUS

ALL OTHER THERAPEUTIC PRODUCTS

ALL OTHER THERAPEUTIC PRODUCTS

Iron chelating agents

DEFERASIROX

Note Special Pricing Arrangements apply.

Authority required

Chronic iron overload

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must be transfusion dependent, **AND**
- Patient must not have a malignant disorder of erythropoiesis.

Authority required

Chronic iron overload

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must not be transfusion dependent, **AND**
- The condition must be thalassaemia.

deferasirox 180 mg tablet, 30

11556F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	6	5	..	*2567.64	Jadenu [NM]

deferasirox 360 mg tablet, 30

11533B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	6	5	..	*5135.34	Jadenu [NM]

deferasirox 90 mg tablet, 30

11499F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	6	5	..	*1283.88	Jadenu [NM]

DEFERASIROX

Note Special Pricing Arrangements apply.

Note A patient's median life expectancy is determined by the severity of their underlying disease.

Note Patients with underlying myelodysplastic syndrome are considered to have a median life expectancy exceeding five years if they are classified as:

- low risk according to the International Prognostic Scoring System (IPSS); or
- very low and low risk according to the Revised International Prognostic Scoring System (IPSS-R); or
- very low and low risk according to the WHO classification based Prognostic Scoring System (WPSS).

Note Patients with underlying myelofibrosis have a median life expectancy exceeding five years if they are classified as:

- low or intermediate risk according to the International Prognostic Scoring System (IPSS); or
- low or intermediate-1 risk according to Dynamic International Prognostic Scoring System (DIPSS).

Authority required

Chronic iron overload

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must be red blood cell transfusion dependent, **AND**
- Patient must have a serum ferritin level of greater than 1000 microgram/L, **AND**
- Patient must have a malignant disorder of haemopoiesis, **AND**
- Patient must have a median life expectancy exceeding five years.

deferasirox 180 mg tablet, 30

11500G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	6	2	..	*2567.64	Jadenu [NM]

deferasirox 360 mg tablet, 30

11536E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	6	2	..	*5135.34	Jadenu [NM]

deferasirox 90 mg tablet, 30

11519G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	6	2	..	*1283.88	Jadenu [NM]

▪ DEFERASIROX

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)**8328**

Chronic iron overload

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must be transfusion dependent, **AND**
- Patient must not have a malignant disorder of erythropoiesis, **AND**
- Patient must have previously received PBS-subsidised therapy with deferasirox for this condition.

Authority required (STREAMLINED)**8329**

Chronic iron overload

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must not be transfusion dependent, **AND**
- The condition must be thalassaemia, **AND**
- Patient must have previously received PBS-subsidised therapy with deferasirox for this condition.

Authority required (STREAMLINED)**8326**

Chronic iron overload

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must be red blood cell transfusion dependent, **AND**
- Patient must have a malignant disorder of haemopoiesis, **AND**
- Patient must have previously received PBS-subsidised therapy with deferasirox for this condition.

Note Interruption of treatment should be considered if serum ferritin levels fall consistently below 500 microgram/mL.

deferasirox 180 mg tablet, 30

11535D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	6	2	..	*2567.64	Jadenu [NM]

deferasirox 360 mg tablet, 30

11555E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	6	2	..	*5135.34	Jadenu [NM]

deferasirox 90 mg tablet, 30

11534C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	6	2	..	*1283.88	Jadenu [NM]

▪ DEFERIPRONE**Authority required (STREAMLINED)****6448**

Iron overload

Clinical criteria:

- Patient must have thalassaemia major, **AND**
- Patient must be unable to take desferrioxamine therapy.

Authority required (STREAMLINED)**6403**

Iron overload

Clinical criteria:

- Patient must have thalassaemia major, **AND**
- Patient must be one in whom desferrioxamine therapy has proven ineffective.

deferiprone 500 mg tablet, 100

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
5657R	3	5	..	*1155.69	Ferriprox [TX]

deferiprone 1 g tablet, 50

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
11747G	6	5	..	*2311.38	Ferriprox [TX]

deferiprone 100 mg/mL oral liquid, 250 mL

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
5658T	5	5	..	*963.10	Ferriprox [TX]

■ **DEFERRIOXAMINE****Authority required (STREAMLINED)****6394**

Disorders of erythropoiesis

Clinical criteria:

- The condition must be associated with treatment-related chronic iron overload.

desferrioxamine mesilate 2 g injection, 1 vial

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
5661Y	60	5	..	*1868.40	Hospira Pty Limited [PF]

desferrioxamine mesilate 500 mg injection, 10 vials

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
5662B	40	5	..	*5555.20	Hospira Pty Limited [PF]

Drugs for treatment of hyperkalemia and hyperphosphatemia■ **LANTHANUM****Authority required (STREAMLINED)****5530**

Hyperphosphataemia

Treatment Phase: Initiation and stabilisation

Clinical criteria:

- The condition must not be adequately controlled by calcium, **AND**
- Patient must have a serum phosphate of greater than 1.6 mmol per L at the commencement of therapy; OR
- The condition must be where a serum calcium times phosphate product is greater than 4 at the commencement of therapy, **AND**
- The treatment must not be used in combination with any other non-calcium phosphate binding agents.

Treatment criteria:

- Patient must be undergoing dialysis for chronic kidney disease.

lanthanum 500 mg chewable tablet, 2 x 45

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
5780F	2	5	..	*497.36	Fosrenol [ZI]

lanthanum 1 g chewable tablet, 6 x 15

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
5782H	2	5	..	*845.52	Fosrenol [ZI]

lanthanum 750 mg chewable tablet, 6 x 15

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
5781G	2	5	..	*751.04	Fosrenol [ZI]

■ **SEVELAMER****Authority required (STREAMLINED)****5530**

Hyperphosphataemia

Treatment Phase: Initiation and stabilisation

Clinical criteria:

- The condition must not be adequately controlled by calcium, **AND**
- Patient must have a serum phosphate of greater than 1.6 mmol per L at the commencement of therapy; OR
- The condition must be where a serum calcium times phosphate product is greater than 4 at the commencement of therapy, **AND**
- The treatment must not be used in combination with any other non-calcium phosphate binding agents.

Treatment criteria:

- Patient must be undergoing dialysis for chronic kidney disease.

sevelamer hydrochloride 800 mg tablet, 180

9546K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*530.10	Renagel [GZ]

■ SUCROFERRIC OXYHYDROXIDE**Authority required (STREAMLINED)****5530**

Hyperphosphataemia

Treatment Phase: Initiation and stabilisation

Clinical criteria:

- The condition must not be adequately controlled by calcium, **AND**
- Patient must have a serum phosphate of greater than 1.6 mmol per L at the commencement of therapy; OR
- The condition must be where a serum calcium times phosphate product is greater than 4 at the commencement of therapy, **AND**
- The treatment must not be used in combination with any other non-calcium phosphate binding agents.

Treatment criteria:

- Patient must be undergoing dialysis for chronic kidney disease.

sucroferric oxyhydroxide 2.5 g (iron 500 mg) chewable tablet, 90

10233N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	Brand Name and Manufacturer
	2	5	..	*753.46	Velphoro [VL]

Highly Specialised Drugs Program (Community Access)

SYSTEMIC HORMONAL PREPARATIONS, EXCL. SEX HORMONES AND INSULINS.....	593
PITUITARY AND HYPOTHALAMIC HORMONES AND ANALOGUES	593
HYPOTHALAMIC HORMONES.....	593
ANTIINFECTIVES FOR SYSTEMIC USE	594
ANTIVIRALS FOR SYSTEMIC USE	594
DIRECT ACTING ANTIVIRALS	594
ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS	611
IMMUNOSTIMULANTS	611
IMMUNOSTIMULANTS	611
NERVOUS SYSTEM.....	611
PSYCHOLEPTICS.....	611
ANTIPSYCHOTICS.....	611

SYSTEMIC HORMONAL PREPARATIONS, EXCL. SEX HORMONES AND INSULINS

PITUITARY AND HYPOTHALAMIC HORMONES AND ANALOGUES

HYPOTHALAMIC HORMONES

Somatostatin and analogues

LANREOTIDE

Authority required (STREAMLINED)

7532

Acromegaly

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The condition must be active, **AND**
- Patient must have persistent elevation of mean growth hormone levels of greater than 2.5 micrograms per litre, **AND**
- The treatment must be after failure of other therapy including dopamine agonists; OR
- The treatment must be as interim treatment while awaiting the effects of radiotherapy and where treatment with dopamine agonists has failed; OR
- The treatment must be in a patient who is unfit for or unwilling to undergo surgery and where radiotherapy is contraindicated, **AND**
- The treatment must cease in a patient treated with radiotherapy if there is biochemical evidence of remission (normal IGF1) after lanreotide has been withdrawn for at least 4 weeks (8 weeks after the last dose), **AND**
- The treatment must cease if IGF1 is not lower after 3 months of treatment, **AND**
- The treatment must not be given concomitantly with PBS-subsidised pegvisomant.

In a patient treated with radiotherapy, lanreotide should be withdrawn every 2 years in the 10 years after radiotherapy for assessment of remission.

Authority required (STREAMLINED)

7509

Functional carcinoid tumour

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The condition must be causing intractable symptoms, **AND**
- Patient must have experienced on average over 1 week, 3 or more episodes per day of diarrhoea and/or flushing, which persisted despite the use of anti-histamines, anti-serotonin agents and anti-diarrhoea agents, **AND**
- Patient must be one in whom surgery or antineoplastic therapy has failed or is inappropriate, **AND**
- The treatment must cease if there is failure to produce a clinically significant reduction in the frequency and severity of symptoms after 3 months' therapy at a dose of 120 mg every 28 days.

Dosage and tolerance to the drug should be assessed regularly and the dosage should be titrated slowly downwards to determine the minimum effective dose.

lanreotide 120 mg/0.5 mL injection, 0.5 mL syringe

11289E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*3686.27	40.30	Somatuline Autogel [IS]

lanreotide 90 mg/0.5 mL injection, 0.5 mL syringe

11316N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*2955.25	40.30	Somatuline Autogel [IS]

lanreotide 60 mg/0.5 mL injection, 0.5 mL syringe

11315M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*2232.35	40.30	Somatuline Autogel [IS]

LANREOTIDE

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

9161

Non-functional gastroenteropancreatic neuroendocrine tumour (GEP-NET)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The condition must be unresectable locally advanced disease or metastatic disease, **AND**
- The condition must be World Health Organisation (WHO) grade 1 or 2, **AND**
- The treatment must be as monotherapy.

Population criteria:

- Patient must be aged 18 years or older.

ANTIINFECTIVES FOR SYSTEMIC USE

WHO grade 1 of GEP-NET is defined as a mitotic count (10HPF) of less than 2 and Ki-67 index (%) of less than or equal to 2.

WHO grade 2 of GEP-NET is defined as a mitotic count (10HPF) of 2-20 and Ki-67 index (%) of 3-20.

Lanreotide is not PBS-subsidised for use in combination with everolimus or sunitinib for this condition.

lanreotide 120 mg/0.5 mL injection, 0.5 mL syringe

11736Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*3686.27	40.30	Somatuline Autogel [IS]

■ OCTREOTIDE

Authority required (STREAMLINED)

8197

Acromegaly

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
 - The condition must be controlled with octreotide immediate release injections, **AND**
 - The treatment must cease in a patient treated with radiotherapy if there is biochemical evidence of remission (normal IGF1) after octreotide has been withdrawn for at least 4 weeks (8 weeks after the last dose), **AND**
 - The treatment must cease if IGF1 is not lower after 3 months of treatment, **AND**
 - The treatment must not be given concomitantly with PBS-subsidised lanreotide or pegvisomant for this condition.
- In a patient treated with radiotherapy, octreotide should be withdrawn every 2 years in the 10 years after radiotherapy for assessment of remission

Authority required (STREAMLINED)

8208

Functional carcinoid tumour

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have achieved symptom control on octreotide immediate release injections, **AND**
- The treatment must cease if there is failure to produce a clinically significant reduction in the frequency and severity of symptoms after 3 months therapy at a dose of 30 mg every 28 days and having allowed adequate rescue therapy with octreotide immediate release injections.

Dosage and tolerance to the drug should be assessed regularly and the dosage should be titrated slowly downwards to determine the minimum effective dose.

Authority required (STREAMLINED)

8198

Vasoactive intestinal peptide secreting tumour (VIPoma)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have achieved symptom control on octreotide immediate release injections, **AND**
- The treatment must cease if there is failure to produce a clinically significant reduction in the frequency and severity of symptoms after 3 months therapy at a dose of 30 mg every 28 days and having allowed adequate rescue therapy with octreotide immediate release injections.

Dosage and tolerance to the drug should be assessed regularly and the dosage should be titrated slowly downwards to determine the minimum effective dose.

octreotide 20 mg modified release injection [1 vial] (&) inert substance diluent [2 mL syringe], 1 pack

11537F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*3527.01	40.30	Sandostatin LAR [NV]

octreotide 10 mg modified release injection [1 vial] (&) inert substance diluent [2 mL syringe], 1 pack

11501H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*2661.11	40.30	Sandostatin LAR [NV]

octreotide 30 mg modified release injection [1 vial] (&) inert substance diluent [2 mL syringe], 1 pack

11512X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*4402.31	40.30	Sandostatin LAR [NV]

■ ANTIINFECTIVES FOR SYSTEMIC USE

■ ANTIVIRALS FOR SYSTEMIC USE

DIRECT ACTING ANTIVIRALS

Nucleosides and nucleotides excl. reverse transcriptase inhibitors

■ GANCICLOVIR

Authority required (STREAMLINED)

5000

Cytomegalovirus retinitis

Clinical criteria:

- Patient must be severely immunocompromised, including due to HIV infection.

ganciclovir 500 mg injection, 5 vials

10328N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	1	..	*424.79	40.30	^a Cymevene [PB]	^a GANCICLOVIR SXP [HN]

■ VALGANCICLOVIR**Authority required (STREAMLINED)****4980**

Cytomegalovirus retinitis

Clinical criteria:

- Patient must have HIV infection.

valganciclovir 450 mg tablet, 60

10306K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	5	..	*1851.93	40.30	^a Valcyte [RO] ^a Valganciclovir Juno [JU] ^a Valganciclovir Sandoz [SZ]	^a Valganciclovir AN [JO] ^a Valganciclovir Mylan [AF]

valganciclovir 50 mg/mL powder for oral liquid, 100 mL

10277X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	11	5	..	*#4396.36	40.30	Valcyte [RO]

Protease inhibitors**■ ATAZANAVIR****Authority required (STREAMLINED)****4512**

HIV infection

Treatment Phase: Initial

Clinical criteria:

- Patient must be antiretroviral treatment naive, **AND**
- The treatment must be in combination with other antiretroviral agents.

Authority required (STREAMLINED)**4454**

HIV infection

Treatment Phase: Continuing

Clinical criteria:

- Patient must have previously received PBS-subsidised therapy for HIV infection, **AND**
- The treatment must be in combination with other antiretroviral agents.

atazanavir 150 mg capsule, 60

10276W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*703.51	40.30	Reyataz [BQ]

atazanavir 200 mg capsule, 60

10349Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	5	..	*935.57	40.30	^a Atazanavir Mylan [AF]	^a Reyataz [BQ]

■ ATAZANAVIR

Note Pharmaceutical benefits that have the form atazanavir 300 mg capsule, 30 and pharmaceutical benefits that have the form atazanavir 300 mg capsule, 60 are equivalent for the purposes of substitution.

Authority required (STREAMLINED)**4512**

HIV infection

Treatment Phase: Initial

Clinical criteria:

- Patient must be antiretroviral treatment naive, **AND**
- The treatment must be in combination with other antiretroviral agents.

Authority required (STREAMLINED)**4454**

HIV infection

Treatment Phase: Continuing

Clinical criteria:

- Patient must have previously received PBS-subsidised therapy for HIV infection, **AND**
- The treatment must be in combination with other antiretroviral agents.

atazanavir 300 mg capsule, 60

11657M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	703.50	40.30	^a Atazanavir Mylan [AF]

atazanavir 300 mg capsule, 30

10321F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*703.51	40.30	^a Reyataz [BQ]

■ **ATAZANAVIR + COBICISTAT**

Authority required (STREAMLINED)

4512

HIV infection

Treatment Phase: Initial

Clinical criteria:

- Patient must be antiretroviral treatment naive, **AND**
- The treatment must be in combination with other antiretroviral agents.

Authority required (STREAMLINED)

4454

HIV infection

Treatment Phase: Continuing

Clinical criteria:

- Patient must have previously received PBS-subsidised therapy for HIV infection, **AND**
- The treatment must be in combination with other antiretroviral agents.

atazanavir 300 mg + cobicistat 150 mg tablet, 30

10692R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*784.41	40.30	Evotaz [BQ]

■ **DARUNAVIR**

Authority required (STREAMLINED)

5094

Human immunodeficiency virus (HIV) infection

Clinical criteria:

- The treatment must be in addition to optimised background therapy, **AND**
- The treatment must be in combination with other antiretroviral agents, **AND**
- The treatment must be co-administered with 100 mg ritonavir twice daily, **AND**
- Patient must have experienced virological failure or clinical failure or genotypic resistance after at least one antiretroviral regimen.

Virological failure is defined as a viral load greater than 400 copies per mL on two consecutive occasions, while clinical failure is linked to emerging signs and symptoms of progressing HIV infection or treatment-limiting toxicity.

darunavir 600 mg tablet, 60

10329P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*1840.67	40.30	Prezista [JC]

darunavir 150 mg tablet, 240

10287K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	939.90	40.30	Prezista [JC]

■ **DARUNAVIR**

Authority required (STREAMLINED)

4313

Human immunodeficiency virus (HIV) infection

Clinical criteria:

- The treatment must be in addition to optimised background therapy, **AND**
- The treatment must be in combination with other antiretroviral agents, **AND**
- The treatment must be co-administered with 100 mg ritonavir, **AND**
- Patient must have experienced virological failure or clinical failure or genotypic resistance after at least one antiretroviral regimen, **AND**
- Patient must not have demonstrated darunavir resistance associated mutations detected on resistance testing.

Virological failure is defined as a viral load greater than 400 copies per mL on two consecutive occasions, while clinical failure is linked to emerging signs and symptoms of progressing HIV infection or treatment-limiting toxicity.

darunavir 800 mg tablet, 30

10367P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*1242.91	40.30	Prezista [JC]

■ FOSAMPRENAVIR

Authority required (STREAMLINED)

4512

HIV infection

Treatment Phase: Initial

Clinical criteria:

- Patient must be antiretroviral treatment naive, **AND**
- The treatment must be in combination with other antiretroviral agents.

Authority required (STREAMLINED)

4454

HIV infection

Treatment Phase: Continuing

Clinical criteria:

- Patient must have previously received PBS-subsidised therapy for HIV infection, **AND**
- The treatment must be in combination with other antiretroviral agents.

fosamprenavir 700 mg tablet, 60

10337C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*681.69	40.30	Telzir [VI]

■ RITONAVIR

Authority required (STREAMLINED)

4512

HIV infection

Treatment Phase: Initial

Clinical criteria:

- Patient must be antiretroviral treatment naive, **AND**
- The treatment must be in combination with other antiretroviral agents.

Authority required (STREAMLINED)

4454

HIV infection

Treatment Phase: Continuing

Clinical criteria:

- Patient must have previously received PBS-subsidised therapy for HIV infection, **AND**
- The treatment must be in combination with other antiretroviral agents.

ritonavir 100 mg tablet, 30

10273Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	24	5	..	*837.55	40.30	Norvir [VE]

■ SAQUINAVIR

Authority required (STREAMLINED)

4512

HIV infection

Treatment Phase: Initial

Clinical criteria:

- Patient must be antiretroviral treatment naive, **AND**
- The treatment must be in combination with other antiretroviral agents.

Authority required (STREAMLINED)

4454

HIV infection

Treatment Phase: Continuing

Clinical criteria:

- Patient must have previously received PBS-subsidised therapy for HIV infection, **AND**
- The treatment must be in combination with other antiretroviral agents.

saquinavir 500 mg tablet, 120

10335Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*861.53	40.30	Invirase [RO]

■ TIPRANAVIR

Authority required (STREAMLINED)

5764

HIV infection

Clinical criteria:

- The treatment must be in addition to optimised background therapy, **AND**
- The treatment must be in combination with other antiretroviral agents, **AND**

- Patient must be antiretroviral experienced, **AND**
 - The treatment must be co-administered with 200 mg ritonavir twice daily, **AND**
 - Patient must have experienced virological failure or clinical failure or genotypic resistance after each of at least 3 different antiretroviral regimens that have included one drug from at least 3 different antiretroviral classes.
- Virological failure is defined as a viral load greater than 400 copies per mL on two consecutive occasions, while clinical failure is linked to emerging signs and symptoms of progressing HIV infection or treatment-limiting toxicity.

tipranavir 250 mg capsule, 120

10344K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*1512.51	40.30	Aptivus [BY]

Nucleoside and nucleotide reverse transcriptase inhibitors

■ ABACAVIR

Authority required (STREAMLINED)

4512

HIV infection

Treatment Phase: Initial

Clinical criteria:

- Patient must be antiretroviral treatment naive, **AND**
- The treatment must be in combination with other antiretroviral agents.

Authority required (STREAMLINED)

4454

HIV infection

Treatment Phase: Continuing

Clinical criteria:

- Patient must have previously received PBS-subsidised therapy for HIV infection, **AND**
- The treatment must be in combination with other antiretroviral agents.

abacavir 300 mg tablet, 60

10294T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*483.81	40.30	Ziagen [VI]

abacavir 20 mg/mL oral liquid, 240 mL

10356C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	8	5	..	*562.51	40.30	Ziagen [VI]

■ ADEFOVIR DIPIVOXIL

Note Patients may receive treatment in combination with lamivudine but not with other PBS-subsidised antihepadnaviral therapy.

Authority required (STREAMLINED)

4490

Chronic hepatitis B infection

Clinical criteria:

- Patient must not have cirrhosis, **AND**
- Patient must have failed antihepadnaviral therapy, **AND**
- Patient must have repeatedly elevated serum ALT levels while on concurrent antihepadnaviral therapy of greater than or equal to 6 months duration, in conjunction with documented chronic hepatitis B infection; OR
- Patient must have repeatedly elevated HBV DNA levels one log greater than the nadir value or failure to achieve a 1 log reduction in HBV DNA within 3 months whilst on previous antihepadnaviral therapy, except in patients with evidence of poor compliance.

Authority required (STREAMLINED)

4510

Chronic hepatitis B infection

Clinical criteria:

- Patient must have cirrhosis, **AND**
- Patient must have failed antihepadnaviral therapy, **AND**
- Patient must have detectable HBV DNA.

Patients with Child's class B or C cirrhosis (ascites, variceal bleeding, encephalopathy, albumin less than 30 g per L, bilirubin greater than 30 micromoles per L) should have their treatment discussed with a transplant unit prior to initiating therapy.

adefovir dipivoxil 10 mg tablet, 30

10290N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	5	..	*898.13	40.30	^a APO-Adefovir [TX]	^a Hepsera [GI]

■ ENTECAVIR

Authority required (STREAMLINED)

4993

Chronic hepatitis B infection

Clinical criteria:

- Patient must not have cirrhosis, **AND**
- Patient must have elevated HBV DNA levels greater than 20,000 IU/mL (100,000 copies/mL) if HBeAg positive, in conjunction with documented hepatitis B infection; OR
- Patient must have elevated HBV DNA levels greater than 2,000 IU/mL (10,000 copies/mL) if HBeAg negative, in conjunction with documented hepatitis B infection, **AND**
- Patient must have evidence of chronic liver injury determined by confirmed elevated serum ALT or liver biopsy.

Authority required (STREAMLINED)**5036**

Chronic hepatitis B infection

Clinical criteria:

- Patient must have cirrhosis, **AND**
 - Patient must have detectable HBV DNA.
- Patients with Child's class B or C cirrhosis (ascites, variceal bleeding, encephalopathy, albumin less than 30 g per L, bilirubin greater than 30 micromoles per L) should have their treatment discussed with a transplant unit prior to initiating therapy.

entecavir 500 microgram tablet, 30

10279B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	5	..	*307.93	40.30	^a Baraclude [BQ]	^a ENTAC [LR]
						^a Entecavir Amneal [EA]	^a ENTECAVIR APO [GX]
						^a Entecavir APOTEX [TX]	^a Entecavir GH [GQ]
						^a Entecavir Mylan [AF]	^a ENTECAVIR RBX [RA]
						^a Entecavir Sandoz [SZ]	^a ENTECLUDE [RW]

■ ENTECAVIR**Note** PBS-subsidised entecavir monohydrate must be used as monotherapy.**Authority required (STREAMLINED)****5044**

Chronic hepatitis B infection

Clinical criteria:

- Patient must not have cirrhosis, **AND**
- Patient must have failed lamivudine, **AND**
- Patient must have repeatedly elevated serum ALT levels while on concurrent antihepadnaviral therapy of greater than or equal to 6 months duration, in conjunction with documented chronic hepatitis B infection; OR
- Patient must have repeatedly elevated HBV DNA levels one log greater than the nadir value or failure to achieve a 1 log reduction in HBV DNA within 3 months whilst on previous antihepadnaviral therapy, except in patients with evidence of poor compliance.

Authority required (STREAMLINED)**5037**

Chronic hepatitis B infection

Clinical criteria:

- Patient must have cirrhosis, **AND**
- Patient must have failed lamivudine, **AND**
- Patient must have detectable HBV DNA.

Patients with Child's class B or C cirrhosis (ascites, variceal bleeding, encephalopathy, albumin less than 30 g per L, bilirubin greater than 30 micromoles per L) should have their treatment discussed with a transplant unit prior to initiating therapy.

entecavir 1 mg tablet, 30

10353X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	5	..	*496.15	40.30	^a Baraclude [BQ]	^a ENTAC [LR]
						^a Entecavir Amneal [EA]	^a ENTECAVIR APO [GX]
						^a Entecavir APOTEX [TX]	^a Entecavir GH [GQ]
						^a Entecavir Mylan [AF]	^a ENTECAVIR RBX [RA]
						^a Entecavir Sandoz [SZ]	^a ENTECLUDE [RW]

■ LAMIVUDINE**Authority required (STREAMLINED)****4512**

HIV infection

Treatment Phase: Initial

Clinical criteria:

- Patient must be antiretroviral treatment naive, **AND**
- The treatment must be in combination with other antiretroviral agents.

Authority required (STREAMLINED)**4454**

HIV infection

Treatment Phase: Continuing

Clinical criteria:

- Patient must have previously received PBS-subsidised therapy for HIV infection, **AND**
- The treatment must be in combination with other antiretroviral agents.

lamivudine 10 mg/mL oral liquid, 240 mL

10320E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	8	5	..	*473.15	40.30	3TC [VI]

lamivudine 300 mg tablet, 30

10311Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	5	..	*143.65	40.30	^a 3TC [VI]	^a Lamivudine Alphapharm [AF]

lamivudine 150 mg tablet, 60

10348P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	5	..	*143.65	40.30	^a 3TC [VI]	^a Lamivudine Alphapharm [AF]

■ LAMIVUDINE

Authority required (STREAMLINED)

4993

Chronic hepatitis B infection

Clinical criteria:

- Patient must not have cirrhosis, **AND**
- Patient must have elevated HBV DNA levels greater than 20,000 IU/mL (100,000 copies/mL) if HBeAg positive, in conjunction with documented hepatitis B infection; OR
- Patient must have elevated HBV DNA levels greater than 2,000 IU/mL (10,000 copies/mL) if HBeAg negative, in conjunction with documented hepatitis B infection, **AND**
- Patient must have evidence of chronic liver injury determined by confirmed elevated serum ALT or liver biopsy.

Authority required (STREAMLINED)

5036

Chronic hepatitis B infection

Clinical criteria:

- Patient must have cirrhosis, **AND**
 - Patient must have detectable HBV DNA.
- Patients with Child's class B or C cirrhosis (ascites, variceal bleeding, encephalopathy, albumin less than 30 g per L, bilirubin greater than 30 micromoles per L) should have their treatment discussed with a transplant unit prior to initiating therapy.

lamivudine 100 mg tablet, 28

10315X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*80.79	40.30	^a Zetlam [AF]
			^b 1.20	*81.99	40.30	^a Zeffix [RW]

■ TENOFOVIR DISOPROXIL

Note Pharmaceutical benefits that have the forms tenofovir disoproxil phosphate 291 mg tablet, tenofovir disoproxil maleate 300 mg tablet, and tenofovir disoproxil fumarate 300 mg tablet are equivalent for the purposes of substitution.

Authority required (STREAMLINED)

6998

HIV infection

Treatment Phase: Initial

Clinical criteria:

- Patient must be antiretroviral treatment naive, **AND**
- The treatment must be in combination with other antiretroviral agents.

Authority required (STREAMLINED)

6982

HIV infection

Treatment Phase: Continuing

Clinical criteria:

- Patient must have previously received PBS-subsidised therapy for HIV infection, **AND**
- The treatment must be in combination with other antiretroviral agents.

Authority required (STREAMLINED)

6980

Chronic hepatitis B infection

Clinical criteria:

- Patient must have cirrhosis, **AND**
- Patient must be nucleoside analogue naive, **AND**
- Patient must have detectable HBV DNA, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Patients with Child's class B or C cirrhosis (ascites, variceal bleeding, encephalopathy, albumin less than 30 g per L, bilirubin greater than 30 micromoles per L) should have their treatment discussed with a transplant unit prior to initiating therapy.

Note Patients may receive treatment in combination with lamivudine but not with other PBS-subsidised antihepadnaviral therapy.

Authority required (STREAMLINED)

6992

Chronic hepatitis B infection

Clinical criteria:

- Patient must not have cirrhosis, **AND**
- Patient must be nucleoside analogue naive, **AND**
- Patient must have elevated HBV DNA levels greater than 20,000 IU/mL (100,000 copies/mL) if HBeAg positive, in conjunction with documented hepatitis B infection; OR
- Patient must have elevated HBV DNA levels greater than 2,000 IU/mL (10,000 copies/mL) if HBeAg negative, in conjunction with documented hepatitis B infection, **AND**
- Patient must have evidence of chronic liver injury determined by: (i) confirmed elevated serum ALT; or (ii) liver biopsy, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Note Patients may receive treatment in combination with lamivudine but not with other PBS-subsidised antihepadnaviral therapy.

Authority required (STREAMLINED)

6983

Chronic hepatitis B infection

Clinical criteria:

- Patient must have cirrhosis, **AND**
- Patient must have failed antihepadnaviral therapy, **AND**
- Patient must have detectable HBV DNA.

Patients with Child's class B or C cirrhosis (ascites, variceal bleeding, encephalopathy, albumin less than 30 g per L, bilirubin greater than 30 micromoles per L) should have their treatment discussed with a transplant unit prior to initiating therapy.

Note Patients may receive treatment in combination with lamivudine but not with other PBS-subsidised antihepadnaviral therapy.

Authority required (STREAMLINED)

6984

Chronic hepatitis B infection

Clinical criteria:

- Patient must not have cirrhosis, **AND**
- Patient must have failed antihepadnaviral therapy, **AND**
- Patient must have repeatedly elevated serum ALT levels while on concurrent antihepadnaviral therapy of greater than or equal to 6 months duration, in conjunction with documented chronic hepatitis B infection; OR
- Patient must have repeatedly elevated HBV DNA levels one log greater than the nadir value or failure to achieve a 1 log reduction in HBV DNA within 3 months whilst on previous antihepadnaviral therapy, except in patients with evidence of poor compliance.

Note Patients may receive treatment in combination with lamivudine but not with other PBS-subsidised antihepadnaviral therapy.

tenofovir disoproxil fumarate 300 mg tablet, 30

10310P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	5	..	*713.27	40.30	^a Tenofovir APOTEX [TX]	^a Viread [GI]

tenofovir disoproxil maleate 300 mg tablet, 30

11155D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*713.27	40.30	^a Tenofovir Disoproxil Mylan [AF]

tenofovir disoproxil phosphate 291 mg tablet, 30

11142K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*713.27	40.30	^a Tenofovir GH [GQ]

■ **ZIDOVUDINE**

Authority required (STREAMLINED)

4512

HIV infection

Treatment Phase: Initial

Clinical criteria:

- Patient must be antiretroviral treatment naive, **AND**
- The treatment must be in combination with other antiretroviral agents.

Authority required (STREAMLINED)

4454

HIV infection

Treatment Phase: Continuing

Clinical criteria:

- Patient must have previously received PBS-subsidised therapy for HIV infection, **AND**
- The treatment must be in combination with other antiretroviral agents.

zidovudine 100 mg capsule, 100

10266H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	4	5	..	*701.67	40.30	Retrovir [VI]

zidovudine 50 mg/5 mL oral liquid, 200 mL

10361H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	15	5	..	*576.19	40.30	Retrovir [VI]

zidovudine 250 mg capsule, 40

10360G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	6	5	..	*1048.75	40.30	Retrovir [VI]

Non-nucleoside reverse transcriptase inhibitors

■ EFAVIRENZ

Authority required (STREAMLINED)

4512

HIV infection

Treatment Phase: Initial

Clinical criteria:

- Patient must be antiretroviral treatment naive, **AND**
- The treatment must be in combination with other antiretroviral agents.

Authority required (STREAMLINED)

4454

HIV infection

Treatment Phase: Continuing

Clinical criteria:

- Patient must have previously received PBS-subsidised therapy for HIV infection, **AND**
- The treatment must be in combination with other antiretroviral agents.

efavirenz 30 mg/mL oral liquid, 180 mL

10275T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	7	5	..	*489.20	40.30	Stocrin [MK]

efavirenz 200 mg tablet, 90

10336B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*466.21	40.30	Stocrin [MK]

efavirenz 600 mg tablet, 30

10366N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*466.21	40.30	Stocrin [MK]

■ ETRAVIRINE

Authority required (STREAMLINED)

5014

HIV infection

Clinical criteria:

- The treatment must be in addition to optimised background therapy, **AND**
 - The treatment must be in combination with other antiretroviral agents, **AND**
 - Patient must be antiretroviral experienced, **AND**
 - Patient must have experienced virological failure or clinical failure or genotypic resistance after each of at least 3 different antiretroviral regimens that have included one drug from at least 3 different antiretroviral classes.
- Virological failure is defined as a viral load greater than 400 copies per mL on two consecutive occasions, while clinical failure is linked to emerging signs and symptoms of progressing HIV infection or treatment-limiting toxicity.

etravirine 200 mg tablet, 60

10301E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*1218.75	40.30	Intelence [JC]

■ NEVIRAPINE

Authority required (STREAMLINED)

4526

HIV infection

Treatment Phase: Initial

Clinical criteria:

- Patient must have been stabilised on nevirapine immediate release, **AND**
- The treatment must be in combination with other antiretroviral agents.

Authority required (STREAMLINED)

4454

HIV infection

Treatment Phase: Continuing

Clinical criteria:

- Patient must have previously received PBS-subsidised therapy for HIV infection, **AND**
- The treatment must be in combination with other antiretroviral agents.

nevirapine 400 mg modified release tablet, 30

10303G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	5	..	*220.23	40.30	^a Nevirapine XR APOTEX [TX]	^a Viramune XR [BY]

▪ **NEVIRAPINE**

Authority required (STREAMLINED)

4512

HIV infection

Treatment Phase: Initial

Clinical criteria:

- Patient must be antiretroviral treatment naive, **AND**
- The treatment must be in combination with other antiretroviral agents.

Authority required (STREAMLINED)

4454

HIV infection

Treatment Phase: Continuing

Clinical criteria:

- Patient must have previously received PBS-subsidised therapy for HIV infection, **AND**
- The treatment must be in combination with other antiretroviral agents.

nevirapine 10 mg/mL oral liquid, 240 mL

10319D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	10	5	..	*1397.39	40.30	Viramune [BY]

nevirapine 200 mg tablet, 60

10304H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	5	..	*220.23	40.30	^a Nevirapine Alphapharm [AF]	^a Viramune [BY]

▪ **RILPIVIRINE**

Authority required (STREAMLINED)

4512

HIV infection

Treatment Phase: Initial

Clinical criteria:

- Patient must be antiretroviral treatment naive, **AND**
- The treatment must be in combination with other antiretroviral agents.

Authority required (STREAMLINED)

4454

HIV infection

Treatment Phase: Continuing

Clinical criteria:

- Patient must have previously received PBS-subsidised therapy for HIV infection, **AND**
- The treatment must be in combination with other antiretroviral agents.

rilpivirine 25 mg tablet, 30

10298B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*544.03	40.30	Edurant [JC]

Antivirals for treatment of HIV infections, combinations

▪ **ABACAVIR + LAMIVUDINE**

Note Pharmaceutical benefits that have the form tablet containing abacavir 600 mg (as sulfate) with lamivudine 300 mg and pharmaceutical benefits that have the form tablet containing abacavir 600 mg (as hydrochloride) with lamivudine 300 mg are equivalent for the purposes of substitution.

Authority required (STREAMLINED)

4527

HIV infection

Treatment Phase: Initial

Clinical criteria:

- Patient must be antiretroviral treatment naive, **AND**
- The treatment must be in combination with other antiretroviral agents.

Population criteria:

- Patient must be aged 12 years or older, **AND**
- Patient must weigh 40 kg or more.

Authority required (STREAMLINED)

4528

HIV infection

Treatment Phase: Continuing

Clinical criteria:

- Patient must have previously received PBS-subsidised therapy for HIV infection, **AND**
- The treatment must be in combination with other antiretroviral agents.

Population criteria:

- Patient must be aged 12 years or older, **AND**
- Patient must weigh 40 kg or more.

abacavir 600 mg + lamivudine 300 mg tablet, 30

10357D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	5	..	*611.73	40.30	^a Abacavir/Lamivudine 600/300 APOTEX [TX]	^a ABACAVIR/LAMIVUDINE 600/300 SUN [RA]
						^a Abacavir/Lamivudine Mylan [AF]	^a Kivexa [VI]

abacavir 600 mg + lamivudine 300 mg tablet, 30

11246X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*611.73	40.30	^a Abacavir/Lamivudine GH 600/300 [GQ]

■ **ABACAVIR + LAMIVUDINE + ZIDOVUDINE**

Authority required (STREAMLINED)

4495

HIV infection

Treatment Phase: Initial

Clinical criteria:

- Patient must be antiretroviral treatment naive.

Population criteria:

- Patient must be aged 12 years or older, **AND**
- Patient must weigh 40 kg or more.

Authority required (STREAMLINED)

4480

HIV infection

Treatment Phase: Continuing

Clinical criteria:

- Patient must have previously received PBS-subsidised therapy for HIV infection.

Population criteria:

- Patient must be aged 12 years or older, **AND**
- Patient must weigh 40 kg or more.

abacavir 300 mg + lamivudine 150 mg + zidovudine 300 mg tablet, 60

10305J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*678.79	40.30	Trizivir [VI]

■ **BICTEGRAVIR + EMTRICITABINE + TENOFOVIR ALAFENAMIDE**

Authority required (STREAMLINED)

4522

HIV infection

Treatment Phase: Initial

Clinical criteria:

- Patient must be antiretroviral treatment naive.

Authority required (STREAMLINED)

4470

HIV infection

Treatment Phase: Continuing

Clinical criteria:

- Patient must have previously received PBS-subsidised therapy for HIV infection.

bictegravir 50 mg + emtricitabine 200 mg + tenofovir alafenamide 25 mg tablet, 30

11649D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*1848.07	40.30	Biktarvy [GI]

■ DARUNAVIR + COBICISTAT**Authority required (STREAMLINED)****6413**

Human immunodeficiency virus (HIV) infection

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must be antiretroviral treatment naive, **AND**
- The treatment must be in combination with other antiretroviral agents, **AND**
- The treatment must not be in combination with ritonavir.

Note The cobicistat component of the darunavir + cobicistat combination product provides the necessary pharmacokinetic enhancement of darunavir to achieve therapeutic levels of darunavir.

Authority required (STREAMLINED)**6428**

Human immunodeficiency virus (HIV) infection

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised therapy for HIV infection, **AND**
- The treatment must be in combination with other antiretroviral agents, **AND**
- The treatment must not be in combination with ritonavir.

Note The cobicistat component of the darunavir + cobicistat combination product provides the necessary pharmacokinetic enhancement of darunavir to achieve therapeutic levels of darunavir.

Authority required (STREAMLINED)**6377**

Human immunodeficiency virus (HIV) infection

Clinical criteria:

- The treatment must be in addition to optimised background therapy, **AND**
- The treatment must be in combination with other antiretroviral agents, **AND**
- The treatment must not be in combination with ritonavir, **AND**
- Patient must have experienced virological failure or clinical failure or genotypic resistance after at least one antiretroviral regimen.

Virological failure is defined as a viral load greater than 400 copies per mL on two consecutive occasions, while clinical failure is linked to emerging signs and symptoms of progressing HIV infection or treatment-limiting toxicity.

Note The cobicistat component of the darunavir + cobicistat combination product provides the necessary pharmacokinetic enhancement of darunavir to achieve therapeutic levels of darunavir.

darunavir 800 mg + cobicistat 150 mg tablet, 30

10903W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*1259.55	40.30	Prezcobix [JC]

■ DOLUTEGRAVIR + ABACAVIR + LAMIVUDINE**Authority required (STREAMLINED)****4495**

HIV infection

Treatment Phase: Initial

Clinical criteria:

- Patient must be antiretroviral treatment naive.

Population criteria:

- Patient must be aged 12 years or older, **AND**
- Patient must weigh 40 kg or more.

Authority required (STREAMLINED)**4480**

HIV infection

Treatment Phase: Continuing

Clinical criteria:

- Patient must have previously received PBS-subsidised therapy for HIV infection.

Population criteria:

- Patient must be aged 12 years or older, **AND**
- Patient must weigh 40 kg or more.

dolutegravir 50 mg + abacavir 600 mg + lamivudine 300 mg tablet, 30

10345L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*1785.65	40.30	Triumeq [VI]

▪ DOLUTEGRAVIR + RILPIVIRINE

Authority required (STREAMLINED)

8214

HIV infection

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must be virologically suppressed on a stable antiretroviral regimen for at least 6 months, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Authority required (STREAMLINED)

8226

HIV infection

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised therapy with this drug for this condition, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

dolutegravir 50 mg + rilpivirine 25 mg tablet, 30

11540J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*1732.25	40.30	Juluca [VI]

▪ EMTRICITABINE + RILPIVIRINE + TENOFOVIR ALAFENAMIDE

Authority required (STREAMLINED)

4522

HIV infection

Treatment Phase: Initial

Clinical criteria:

- Patient must be antiretroviral treatment naive.

Authority required (STREAMLINED)

4470

HIV infection

Treatment Phase: Continuing

Clinical criteria:

- Patient must have previously received PBS-subsidised therapy for HIV infection.

emtricitabine 200 mg + rilpivirine 25 mg + tenofovir alafenamide 25 mg tablet, 30

11104K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*1963.51	40.30	Odefsey [GI]

▪ EMTRICITABINE + TENOFOVIR ALAFENAMIDE

Authority required (STREAMLINED)

4512

HIV infection

Treatment Phase: Initial

Clinical criteria:

- Patient must be antiretroviral treatment naive, **AND**
- The treatment must be in combination with other antiretroviral agents.

Authority required (STREAMLINED)

4454

HIV infection

Treatment Phase: Continuing

Clinical criteria:

- Patient must have previously received PBS-subsidised therapy for HIV infection, **AND**
- The treatment must be in combination with other antiretroviral agents.

emtricitabine 200 mg + tenofovir alafenamide 10 mg tablet, 30

11099E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*1447.51	40.30	Descovy [GI]

emtricitabine 200 mg + tenofovir alafenamide 25 mg tablet, 30

11113X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*1447.51	40.30	Descovy [GI]

■ LAMIVUDINE + ZIDOVUDINE

Authority required (STREAMLINED)

4512

HIV infection

Treatment Phase: Initial

Clinical criteria:

- Patient must be antiretroviral treatment naive, **AND**
- The treatment must be in combination with other antiretroviral agents.

Authority required (STREAMLINED)

4454

HIV infection

Treatment Phase: Continuing

Clinical criteria:

- Patient must have previously received PBS-subsidised therapy for HIV infection, **AND**
- The treatment must be in combination with other antiretroviral agents.

lamivudine 150 mg + zidovudine 300 mg tablet, 60

10284G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	5	..	*217.49	40.30	^a Combivir [VI]	^a Lamivudine 150 mg + Zidovudine 300 mg Alphapharm [AF]

■ LOPINAVIR + RITONAVIR

Authority required (STREAMLINED)

4512

HIV infection

Treatment Phase: Initial

Clinical criteria:

- Patient must be antiretroviral treatment naive, **AND**
- The treatment must be in combination with other antiretroviral agents.

Authority required (STREAMLINED)

4454

HIV infection

Treatment Phase: Continuing

Clinical criteria:

- Patient must have previously received PBS-subsidised therapy for HIV infection, **AND**
- The treatment must be in combination with other antiretroviral agents.

lopinavir 100 mg + ritonavir 25 mg tablet, 60

10285H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*355.59	40.30	Kaletra [VE]

lopinavir 200 mg + ritonavir 50 mg tablet, 120

10272P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*1386.61	40.30	Kaletra [VE]

lopinavir 400 mg/5 mL + ritonavir 100 mg/5 mL oral liquid, 60 mL

10327M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	10	5	..	*1308.89	40.30	Kaletra [VE]

■ TENOFOVIR + EMTRICITABINE

Note Pharmaceutical benefits that have the forms tenofovir disoproxil phosphate 291 mg with emtricitabine 200 mg tablet, tenofovir disoproxil maleate 300 mg with emtricitabine 200 mg tablet, and tenofovir disoproxil fumarate 300 mg with emtricitabine 200 mg tablet are equivalent for the purposes of substitution.

Authority required (STREAMLINED)

6985

HIV infection

Treatment Phase: Initial

Clinical criteria:

- Patient must be antiretroviral treatment naive, **AND**
- The treatment must be in combination with other antiretroviral agents.

Authority required (STREAMLINED)

6986

HIV infection

Treatment Phase: Continuing

Clinical criteria:

- Patient must have previously received PBS-subsidised therapy for HIV infection, **AND**

- The treatment must be in combination with other antiretroviral agents.

tenofovir disoproxil fumarate 300 mg + emtricitabine 200 mg tablet, 30

10347N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	5	..	*306.11	40.30	^a Tenofovir/Emtricitabine 300/200 APOTEX [TX]	^a Truvada [GI]

tenofovir disoproxil phosphate 291 mg + emtricitabine 200 mg tablet, 30

11146P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*306.11	40.30	^a Tenofovir EMT GH [GQ]

tenofovir disoproxil maleate 300 mg + emtricitabine 200 mg tablet, 30

11149T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*306.11	40.30	^a Tenofovir Disoproxil Emtricitabine Mylan 300/200 [AF]

■ TENOFOVIR + EMTRICITABINE + EFAVIRENZ

Note Pharmaceutical benefits that have the form tenofovir disoproxil fumarate 300 mg with emtricitabine 200 mg and efavirenz 600 mg tablet and pharmaceutical benefits that have the form tenofovir disoproxil maleate 300 mg with emtricitabine 200 mg and efavirenz 600 mg tablet are equivalent for the purposes of substitution.

Authority required (STREAMLINED)

4522

HIV infection

Treatment Phase: Initial

Clinical criteria:

- Patient must be antiretroviral treatment naive.

Authority required (STREAMLINED)

4470

HIV infection

Treatment Phase: Continuing

Clinical criteria:

- Patient must have previously received PBS-subsidised therapy for HIV infection.

tenofovir disoproxil fumarate 300 mg + emtricitabine 200 mg + efavirenz 600 mg tablet, 30

10297Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*650.23	40.30	^a Atripla [GI]

tenofovir disoproxil maleate 300 mg + emtricitabine 200 mg + efavirenz 600 mg tablet, 30

11732L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*650.23	40.30	^a Tenofovir Disoproxil/Emtricitabine/Efavirenz Mylan 300/200/600 [AF]

■ TENOFOVIR + EMTRICITABINE + ELVITEGRAVIR + COBICISTAT

Authority required (STREAMLINED)

4522

HIV infection

Treatment Phase: Initial

Clinical criteria:

- Patient must be antiretroviral treatment naive.

Authority required (STREAMLINED)

4470

HIV infection

Treatment Phase: Continuing

Clinical criteria:

- Patient must have previously received PBS-subsidised therapy for HIV infection.

tenofovir disoproxil fumarate 300 mg + emtricitabine 200 mg + elvitegravir 150 mg + cobicistat 150 mg tablet, 30

10307L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*1784.49	40.30	Stribild [GI]

■ TENOFOVIR + EMTRICITABINE + RILPIVIRINE

Authority required (STREAMLINED)

4522

HIV infection

Treatment Phase: Initial

Clinical criteria:

- Patient must be antiretroviral treatment naive.

Authority required (STREAMLINED)

4470

HIV infection

Treatment Phase: Continuing

Clinical criteria:

- Patient must have previously received PBS-subsidised therapy for HIV infection.

tenofovir disoproxil fumarate 300 mg + emtricitabine 200 mg + rilpivirine 25 mg tablet, 30

10314W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*1784.49	40.30	Eviplera [GI]

▪ TENOFOVIR ALAFENAMIDE + EMTRICITABINE + ELVITEGRAVIR + COBICISTAT**Authority required (STREAMLINED)****4522**

HIV infection

Treatment Phase: Initial

Clinical criteria:

- Patient must be antiretroviral treatment naive.

Authority required (STREAMLINED)**4470**

HIV infection

Treatment Phase: Continuing

Clinical criteria:

- Patient must have previously received PBS-subsidised therapy for HIV infection.

tenofovir alafenamide 10 mg + emtricitabine 200 mg + elvitegravir 150 mg + cobicistat 150 mg tablet, 30

11114Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*1963.51	40.30	Genvoya [GI]

HSD
(Community)***Other antiretrovirals*****▪ DOLUTEGRAVIR****Authority required (STREAMLINED)****4512**

HIV infection

Treatment Phase: Initial

Clinical criteria:

- Patient must be antiretroviral treatment naive, **AND**
- The treatment must be in combination with other antiretroviral agents.

Authority required (STREAMLINED)**4454**

HIV infection

Treatment Phase: Continuing

Clinical criteria:

- Patient must have previously received PBS-subsidised therapy for HIV infection, **AND**
- The treatment must be in combination with other antiretroviral agents.

dolutegravir 50 mg tablet, 30

10283F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*1311.93	40.30	Tivicay [VI]

▪ ENFUVIRTIDE**Authority required (STREAMLINED)****5014**

HIV infection

Clinical criteria:

- The treatment must be in addition to optimised background therapy, **AND**
 - The treatment must be in combination with other antiretroviral agents, **AND**
 - Patient must be antiretroviral experienced, **AND**
 - Patient must have experienced virological failure or clinical failure or genotypic resistance after each of at least 3 different antiretroviral regimens that have included one drug from at least 3 different antiretroviral classes.
- Virological failure is defined as a viral load greater than 400 copies per mL on two consecutive occasions, while clinical failure is linked to emerging signs and symptoms of progressing HIV infection or treatment-limiting toxicity.

enfuvirtide 90 mg injection [60 vials] (&) inert substance diluent [60 x 1.1 mL vials], 1 pack

10365M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*3831.63	40.30	Fuzeon [RO]

▪ MARAVIROC**Authority required (STREAMLINED)**

5008

HIV infection

Clinical criteria:

- Patient must be infected with CCR5-tropic HIV-1, **AND**
 - The treatment must be in addition to optimised background therapy, **AND**
 - The treatment must be in combination with other antiretroviral agents, **AND**
 - Patient must have experienced virological failure or clinical failure or genotypic resistance after each of at least 3 different antiretroviral regimens that have included one drug from at least 3 different antiretroviral classes.
- Virological failure is defined as a viral load greater than 400 copies per mL on two consecutive occasions, while clinical failure is linked to emerging signs and symptoms of progressing HIV infection or treatment-limiting toxicity.

A tropism assay to determine CCR5 only strain status must be performed prior to initiation. Individuals with CXCR4 tropism demonstrated at any time point are not eligible.

maraviroc 300 mg tablet, 60

10355B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*1791.03	40.30	Celsentri [VI]

maraviroc 150 mg tablet, 60

10318C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*1791.03	40.30	Celsentri [VI]

■ RALTEGRAVIR

Authority required (STREAMLINED)

4512

HIV infection

Treatment Phase: Initial

Clinical criteria:

- Patient must be antiretroviral treatment naive, **AND**
- The treatment must be in combination with other antiretroviral agents.

Authority required (STREAMLINED)

4454

HIV infection

Treatment Phase: Continuing

Clinical criteria:

- Patient must have previously received PBS-subsidised therapy for HIV infection, **AND**
- The treatment must be in combination with other antiretroviral agents.

raltegravir 600 mg tablet, 60

11248B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*1185.47	40.30	Isentress HD [MK]

raltegravir 400 mg tablet, 60

10286J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*1185.47	40.30	Isentress [MK]

■ RALTEGRAVIR

Authority required (STREAMLINED)

4275

HIV infection

Treatment Phase: Initial

Clinical criteria:

- The treatment must be in combination with other antiretroviral agents, **AND**
- Patient must be antiretroviral experienced with at least 6 months therapy with 2 alternate classes of anti-retroviral therapy, **AND**
- Patient must have a CD4 count of less than 500 per cubic millimetre; OR
- Patient must have symptomatic HIV disease.

Population criteria:

- Patient must be aged 2 years or older.

Authority required (STREAMLINED)

4274

HIV infection

Treatment Phase: Continuing

Clinical criteria:

- The treatment must be in combination with other antiretroviral agents, **AND**
- Patient must be antiretroviral experienced with at least 6 months therapy with 2 alternate classes of anti-retroviral therapy, **AND**
- Patient must have previously received PBS-subsidised therapy for HIV infection.

Population criteria:

- Patient must be aged 2 years or older.

raltegravir 25 mg chewable tablet, 60

10299C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	6	5	..	*457.57	40.30	Isentress [MK]

raltegravir 100 mg chewable tablet, 60

10326L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	6	5	..	*1778.83	40.30	Isentress [MK]

■ ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS

■ IMMUNOSTIMULANTS

IMMUNOSTIMULANTS

Interferons

■ INTERFERON ALFA-2A

Authority required (STREAMLINED)

4993

Chronic hepatitis B infection

Clinical criteria:

- Patient must not have cirrhosis, **AND**
- Patient must have elevated HBV DNA levels greater than 20,000 IU/mL (100,000 copies/mL) if HBeAg positive, in conjunction with documented hepatitis B infection; OR
- Patient must have elevated HBV DNA levels greater than 2,000 IU/mL (10,000 copies/mL) if HBeAg negative, in conjunction with documented hepatitis B infection, **AND**
- Patient must have evidence of chronic liver injury determined by confirmed elevated serum ALT or liver biopsy.

Authority required (STREAMLINED)

5036

Chronic hepatitis B infection

Clinical criteria:

- Patient must have cirrhosis, **AND**
- Patient must have detectable HBV DNA.

Patients with Child's class B or C cirrhosis (ascites, variceal bleeding, encephalopathy, albumin less than 30 g per L, bilirubin greater than 30 micromoles per L) should have their treatment discussed with a transplant unit prior to initiating therapy.

interferon alfa-2a 3 million units/0.5 mL injection, 0.5 mL syringe

10317B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	30	5	..	*762.79	40.30	Roferon-A [RO]

interferon alfa-2a 9 million units/0.5 mL injection, 0.5 mL syringe

10369R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	30	5	..	*2225.29	40.30	Roferon-A [RO]

■ NERVOUS SYSTEM

■ PSYCHOLEPTICS

ANTIPSYCHOTICS

Diazepines, oxazepines, thiazepines and oxepines

■ CLOZAPINE

Note Patients receiving clozapine under the PBS listing must be registered in the clozapine patient monitoring program relevant for the brand of clozapine being prescribed and dispensed: Pfizer ClopineCentral.

Authority required (STREAMLINED)

4998

Schizophrenia

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a psychiatrist; OR
- Must be treated by an authorised medical practitioner, with the agreement of the treating psychiatrist.

Clinical criteria:

- Patient must have previously received PBS-subsidised therapy with this drug for this condition, **AND**
- Patient must have completed at least 18 weeks therapy, **AND**
- Patient must be on a clozapine dosage considered stable by a treating psychiatrist, **AND**

NERVOUS SYSTEM

- The treatment must be under the supervision and direction of a psychiatrist reviewing the patient at regular intervals. A medical practitioner should request a quantity sufficient for up to one month's supply. Up to 5 repeats will be authorised.

clozapine 50 mg/mL oral liquid, 100 mL

11422E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	147.79	40.30	Versacloz [PF]

■ CLOZAPINE

Note Patients receiving clozapine under the PBS listing must be registered in the clozapine patient monitoring program relevant for the brand of clozapine being prescribed and dispensed: Novartis Clozaril Patient Monitoring System (eCPMS) or Hospira Clopineconnect.

Authority required (STREAMLINED)

4998

Schizophrenia

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a psychiatrist; OR
- Must be treated by an authorised medical practitioner, with the agreement of the treating psychiatrist.

Clinical criteria:

- Patient must have previously received PBS-subsidised therapy with this drug for this condition, **AND**
- Patient must have completed at least 18 weeks therapy, **AND**
- Patient must be on a clozapine dosage considered stable by a treating psychiatrist, **AND**
- The treatment must be under the supervision and direction of a psychiatrist reviewing the patient at regular intervals. A medical practitioner should request a quantity sufficient for up to one month's supply. Up to 5 repeats will be authorised.

clozapine 25 mg tablet, 100

10289M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	..	..	*76.03	40.30	Clopine 25 [PF]	Clozaril 25 [GO]

clozapine 100 mg tablet, 100

10358E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	..	..	*259.47	40.30	Clopine 100 [PF]	Clozaril 100 [GO]

clozapine 50 mg tablet, 100

10302F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*141.85	40.30	Clopine 50 [PF]

clozapine 200 mg tablet, 100

10288L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*511.55	40.30	Clopine 200 [PF]

clozapine 50 mg/mL oral liquid, 100 mL

10341G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	147.79	40.30	Clopine Suspension [PF]

Botulinum Toxin Program

MUSCULO-SKELETAL SYSTEM	614
MUSCLE RELAXANTS	614
MUSCLE RELAXANTS, PERIPHERALLY ACTING AGENTS	614

■ MUSCULO-SKELETAL SYSTEM

■ MUSCLE RELAXANTS

MUSCLE RELAXANTS, PERIPHERALLY ACTING AGENTS

Other muscle relaxants, peripherally acting agents

■ BOTULINUM TOXIN TYPE A

Caution Contraindications to treatment include known sensitivity to botulinum toxin.

Note The units used to express the potency of botulinum toxin preparations currently available for PBS subsidy are not equivalent.

Authority required (STREAMLINED)

5221

Blepharospasm or hemifacial spasm

Clinical criteria:

- Patient must have blepharospasm; OR
- Patient must have hemifacial spasm.

Treatment criteria:

- Must be treated by a neurologist; OR
- Must be treated by an ophthalmologist; OR
- Must be treated by an otolaryngology head and neck surgeon; OR
- Must be treated by a plastic surgeon.

Population criteria:

- Patient must be aged 12 years or older.

botulinum toxin type A 100 units injection, 1 vial

10997T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	4	..	..	*1397.35	40.30	Botox [AG]

■ BOTULINUM TOXIN TYPE A

Caution Contraindications to treatment include known sensitivity to botulinum toxin.

Note The units used to express the potency of botulinum toxin preparations currently available for PBS subsidy are not equivalent.

Authority required (STREAMLINED)

5406

Spasmodic torticollis

Clinical criteria:

- Patient must have spasmodic torticollis, **AND**
- The treatment must be as monotherapy; OR
- The treatment must be as adjunctive therapy to current standard care.

Treatment criteria:

- Must be treated by a neurologist; OR
- Must be treated by a plastic surgeon; OR
- Must be treated by a rehabilitation specialist.

botulinum toxin type A 100 units injection, 1 vial

11023E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	4	..	..	*1397.35	40.30	Botox [AG]

■ BOTULINUM TOXIN TYPE A

Caution Contraindications to treatment include known sensitivity to botulinum toxin.

Note The units used to express the potency of botulinum toxin preparations currently available for PBS subsidy are not equivalent.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

5409

Urinary incontinence

Clinical criteria:

- The condition must be due to neurogenic detrusor overactivity, as demonstrated by urodynamic study, **AND**
- The condition must be inadequately controlled by anti-cholinergic therapy, **AND**
- Patient must experience at least 14 episodes of urinary incontinence per week prior to commencement of treatment with Botulinum Toxin Type A Neurotoxin Complex, **AND**
- Patient must be willing and able to self-catheterise, **AND**
- The treatment must not continue if the patient does not achieve a 50% or greater reduction from baseline in urinary incontinence episodes 6-12 weeks after the first treatment, **AND**
- Patient must have multiple sclerosis; OR
- Patient must have a spinal cord injury; OR
- Patient must be aged 18 years or older and have spina bifida.

Treatment criteria:

- Must be treated by a urologist; OR

- Must be treated by a urogynaecologist.

botulinum toxin type A 100 units injection, 1 vial

10993N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	4	..	..	*1397.35	40.30	Botox [AG]

▪ BOTULINUM TOXIN TYPE A

Caution Contraindications to treatment include known sensitivity to botulinum toxin.

Note The units used to express the potency of botulinum toxin preparations currently available for PBS subsidy are not equivalent.

Authority required (STREAMLINED)**5359**

Dynamic equinus foot deformity

Clinical criteria:

- The condition must be due to spasticity, **AND**
- Patient must have cerebral palsy, **AND**
- Patient must be ambulant.

Population criteria:

- Patient must be aged from 2 to 17 years inclusive.

Treatment criteria:

- Must be treated by a neurologist; OR
- Must be treated by an orthopaedic surgeon; OR
- Must be treated by a paediatrician; OR
- Must be treated by a rehabilitation specialist.

Authority required (STREAMLINED)**8822**

Dynamic equinus foot deformity

Clinical criteria:

- The condition must be due to spasticity, **AND**
- Patient must have cerebral palsy, **AND**
- Patient must be ambulant.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a neurologist; OR
- Must be treated by an orthopaedic surgeon; OR
- Must be treated by a paediatrician; OR
- Must be treated by a rehabilitation specialist.

botulinum toxin type A 100 units injection, 1 vial

10998W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	4	..	..	*1397.35	40.30	Botox [AG]

▪ BOTULINUM TOXIN TYPE A

Caution Contraindications to treatment include known sensitivity to botulinum toxin.

Note The units used to express the potency of botulinum toxin preparations currently available for PBS subsidy are not equivalent.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)**5262**

Chronic migraine

Treatment criteria:

- Must be treated by a neurologist.

Clinical criteria:

- Patient must have experienced an average of 15 or more headache days per month, with at least 8 days of migraine, over a period of at least 6 months, prior to commencement of treatment with botulinum toxin type A neurotoxin, **AND**
- Patient must have experienced an inadequate response, intolerance or a contraindication to at least three prophylactic migraine medications prior to commencement of treatment with botulinum toxin type A neurotoxin, **AND**
- Patient must have achieved and maintained a 50% or greater reduction from baseline in the number of headache days per month after two treatment cycles (each of 12 weeks duration) in order to be eligible for continuing PBS-subsidised treatment, **AND**
- Patient must be appropriately managed by his or her practitioner for medication overuse headache, prior to initiation of treatment with botulinum toxin.

Population criteria:

- Patient must be aged 18 years or older.

Prophylactic migraine medications are propranolol, amitriptylin, methsergide, pizotifen, cyproheptadine or topiramate.

botulinum toxin type A 100 units injection, 1 vial

11000Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	4	..	..	*1397.35	40.30	Botox [AG]

▪ BOTULINUM TOXIN TYPE A

Caution Contraindications to treatment include known sensitivity to botulinum toxin.

Note The units used to express the potency of botulinum toxin preparations currently available for PBS subsidy are not equivalent.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)**6953**

Urinary incontinence

Treatment criteria:

- Must be treated by a urologist; OR
- Must be treated by a gynaecologist.

Clinical criteria:

- The condition must be due to idiopathic overactive bladder, **AND**
- The condition must have been inadequately controlled by therapy involving at least two alternative anti-cholinergic agents, **AND**
- Patient must experience at least 14 episodes of urinary incontinence per week prior to commencement of treatment with botulinum toxin type A neurotoxin complex, **AND**
- Patient must be willing and able to self-catheterise, **AND**
- The treatment must not continue if the patient does not achieve a 50% or greater reduction from baseline in urinary incontinence episodes 6-12 weeks after the first treatment.

Population criteria:

- Patient must be aged 18 years or older.

botulinum toxin type A 100 units injection, 1 vial

11004E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	4	..	..	*1397.35	40.30	Botox [AG]

▪ BOTULINUM TOXIN TYPE A

Caution Contraindications to treatment include known sensitivity to botulinum toxin.

Note The units used to express the potency of botulinum toxin preparations currently available for PBS subsidy are not equivalent.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)**5408**

Severe primary axillary hyperhidrosis

Clinical criteria:

- Patient must have previously failed topical aluminium chloride hexahydrate after one to two months of treatment; OR
- Patient must be intolerant to topical aluminium chloride hexahydrate treatment.

Population criteria:

- Patient must be aged 12 years or older.

Treatment criteria:

- Must be treated by a dermatologist; OR
- Must be treated by a neurologist; OR
- Must be treated by a paediatrician.

Maximum number of treatments per year is 3, with no less than 4 months to elapse between treatments.

botulinum toxin type A 100 units injection, 1 vial

11016T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	4	..	..	*1397.35	40.30	Botox [AG]

▪ BOTULINUM TOXIN TYPE A

Caution Contraindications to treatment include known sensitivity to botulinum toxin.

Note The units used to express the potency of botulinum toxin preparations currently available for PBS subsidy are not equivalent.

Authority required (STREAMLINED)**5178**

Moderate to severe spasticity of the upper limb

Clinical criteria:

- Patient must have cerebral palsy.

Population criteria:

- Patient must be aged from 2 to 17 years inclusive.

Treatment criteria:

- Must be treated by a neurologist; OR
- Must be treated by an orthopaedic surgeon; OR
- Must be treated by a paediatrician; OR
- Must be treated by a rehabilitation specialist; OR

- Must be treated by a plastic surgeon.

Authority required (STREAMLINED)**8929**

Moderate to severe spasticity of the upper limb

Clinical criteria:

- Patient must have cerebral palsy.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a neurologist; OR
- Must be treated by an orthopaedic surgeon; OR
- Must be treated by a paediatrician; OR
- Must be treated by a rehabilitation specialist; OR
- Must be treated by a plastic surgeon.

Authority required (STREAMLINED)**5220**

Moderate to severe spasticity of the upper limb following a stroke

Clinical criteria:

- The condition must be moderate to severe spasticity of the upper limb/s following stroke, defined as a Modified Ashworth Scale rating of 3 or more, **AND**
- The treatment must not be initiated until three months post-stroke, **AND**
- The treatment must only be used as second line therapy when standard management has failed; OR
- The treatment must only be used as an adjunct to physical therapy, **AND**
- The treatment must not continue if the patient does not respond (defined as not having had a decrease in spasticity rating greater than 1, using the Modified Ashworth Scale, in at least one joint) after two treatment periods (total Botox, Dysport, and Xeomin), **AND**
- The treatment must not exceed 4 treatment periods (total Botox, Dysport, and Xeomin) per upper limb per lifetime, **AND**
- Patient must not have established severe contracture in the limb to be treated.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a neurologist; OR
- Must be treated by an orthopaedic surgeon; OR
- Must be treated by a rehabilitation specialist; OR
- Must be treated by a plastic surgeon; OR
- Must be treated by a geriatrician.

The date of the stroke must be documented in the patient's medical records when treatment is initiated.

Standard management includes physiotherapy and/or oral spasticity agents.

botulinum toxin type A 100 units injection, 1 vial

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
10999X	4	..	..	*1397.35	40.30	Botox [AG]

■ BOTULINUM TOXIN TYPE A**Caution** Contraindications to treatment include known sensitivity to botulinum toxin.**Note** The units used to express the potency of botulinum toxin preparations currently available for PBS subsidy are not equivalent.**Note** Special Pricing Arrangements apply.**Note** An acute event may be a clinical or external event that leads to upper motor neuron lesions resulting in spasticity for example stroke, traumatic brain injury, spinal cord injury, infection or hypoxia.**Authority required (STREAMLINED)****9334**

Moderate to severe spasticity of the lower limb following an acute event

Treatment criteria:

- Must be treated by a neurologist; OR
- Must be treated by an orthopaedic surgeon; OR
- Must be treated by a rehabilitation specialist; OR
- Must be treated by a plastic surgeon; OR
- Must be treated by a geriatrician.

Clinical criteria:

- The condition must be moderate to severe spasticity of the lower limb/s following stroke or other acute neurological event, defined as a Modified Ashworth Scale rating of 3 or more, **AND**
- The treatment must only be used as second line therapy when standard management has failed; OR
- The treatment must only be used as an adjunct to physical therapy, **AND**
- The treatment must not continue if the patient does not respond (defined as not having had a decrease in spasticity rating of at least 1, using the Modified Ashworth Scale, in at least one joint) after two treatment periods (with any botulinum toxin type A), **AND**
- Patient must not have established severe contracture in the limb to be treated, **AND**

- The treatment must not exceed a maximum of 4 treatment periods (with any botulinum toxin type A) per lower limb in the first year of treatment, and 2 treatment periods (with any botulinum toxin type A) per lower limb each year thereafter.

Population criteria:

- Patient must be aged 18 years or older.

Standard management includes physiotherapy and/or oral spasticity agents.

Authority required (STREAMLINED)

9271

Moderate to severe spasticity of the lower limb following an acute event

Treatment Phase: Grandfathered treatment

Treatment criteria:

- Must be treated by a neurologist; OR
- Must be treated by an orthopaedic surgeon; OR
- Must be treated by a rehabilitation specialist; OR
- Must be treated by a plastic surgeon; OR
- Must be treated by a geriatrician.

Clinical criteria:

- Patient must have previously received non-PBS subsidised treatment with this drug for this condition prior to 1 September 2019, **AND**
- The condition must have been moderate to severe spasticity of the lower limb/s following an acute event, defined as a Modified Ashworth Scale rating of 3 or more prior to commencing non-PBS subsidised treatment, **AND**
- The treatment must only be used as second line therapy when standard management has failed; OR
- The treatment must only be used as an adjunct to physical therapy, **AND**
- The treatment must not continue if the patient does not respond (defined as not having had a decrease in spasticity rating of at least 1, using the Modified Ashworth Scale, in at least one joint) after two treatment periods (with any botulinum toxin type A), **AND**
- Patient must not have established severe contracture in the limb to be treated, **AND**
- The treatment must not exceed a maximum of 4 treatment periods (with any botulinum toxin type A) per lower limb in the first year of treatment, and 2 treatment periods (with any botulinum toxin type A) per lower limb each year thereafter.

Population criteria:

- Patient must be aged 18 years or older.

Standard management includes physiotherapy and/or oral spasticity agents.

botulinum toxin type A 100 units injection, 1 vial

11751L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	4	..	..	*1397.35	40.30	Botox [AG]

■ CLOSTRIDIUM BOTULINUM TYPE A TOXIN-HAEMAGGLUTININ COMPLEX

Caution Contraindications to treatment include known sensitivity to botulinum toxin.

Note The units used to express the potency of botulinum toxin preparations currently available for PBS subsidy are not equivalent.

Authority required (STREAMLINED)

5405

Blepharospasm or hemifacial spasm

Clinical criteria:

- Patient must have blepharospasm; OR
- Patient must have hemifacial spasm.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a neurologist; OR
- Must be treated by an ophthalmologist; OR
- Must be treated by an otolaryngology head and neck surgeon; OR
- Must be treated by a plastic surgeon.

clostridium botulinum type A toxin-haemagglutinin complex 300 units injection, 1 vial

10987G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	4	..	..	*1221.95	40.30	Dysport [IS]

clostridium botulinum type A toxin-haemagglutinin complex 500 units injection, 1 vial

11022D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*1094.89	40.30	Dysport [IS]

■ CLOSTRIDIUM BOTULINUM TYPE A TOXIN-HAEMAGGLUTININ COMPLEX

Caution Contraindications to treatment include known sensitivity to botulinum toxin.

Note The units used to express the potency of botulinum toxin preparations currently available for PBS subsidy are not equivalent.

Authority required (STREAMLINED)

5406

Spasmodic torticollis

Clinical criteria:

- Patient must have spasmodic torticollis, **AND**
- The treatment must be as monotherapy; OR
- The treatment must be as adjunctive therapy to current standard care.

Treatment criteria:

- Must be treated by a neurologist; OR
- Must be treated by a plastic surgeon; OR
- Must be treated by a rehabilitation specialist.

clostridium botulinum type A toxin-haemagglutinin complex 300 units injection, 1 vial

11007H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	4	..	..	*1221.95	40.30	Dysport [IS]

clostridium botulinum type A toxin-haemagglutinin complex 500 units injection, 1 vial

11015R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*1094.89	40.30	Dysport [IS]

■ CLOSTRIDIUM BOTULINUM TYPE A TOXIN-HAEMAGGLUTININ COMPLEX

Caution Contraindications to treatment include known sensitivity to botulinum toxin.

Note The units used to express the potency of botulinum toxin preparations currently available for PBS subsidy are not equivalent.

Authority required (STREAMLINED)**5359**

Dynamic equinus foot deformity

Clinical criteria:

- The condition must be due to spasticity, **AND**
- Patient must have cerebral palsy, **AND**
- Patient must be ambulant.

Population criteria:

- Patient must be aged from 2 to 17 years inclusive.

Treatment criteria:

- Must be treated by a neurologist; OR
- Must be treated by an orthopaedic surgeon; OR
- Must be treated by a paediatrician; OR
- Must be treated by a rehabilitation specialist.

Authority required (STREAMLINED)**8822**

Dynamic equinus foot deformity

Clinical criteria:

- The condition must be due to spasticity, **AND**
- Patient must have cerebral palsy, **AND**
- Patient must be ambulant.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a neurologist; OR
- Must be treated by an orthopaedic surgeon; OR
- Must be treated by a paediatrician; OR
- Must be treated by a rehabilitation specialist.

clostridium botulinum type A toxin-haemagglutinin complex 300 units injection, 1 vial

10981Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	4	..	..	*1221.95	40.30	Dysport [IS]

clostridium botulinum type A toxin-haemagglutinin complex 500 units injection, 1 vial

11006G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*1094.89	40.30	Dysport [IS]

■ CLOSTRIDIUM BOTULINUM TYPE A TOXIN-HAEMAGGLUTININ COMPLEX

Caution Contraindications to treatment include known sensitivity to botulinum toxin.

Note The units used to express the potency of botulinum toxin preparations currently available for PBS subsidy are not equivalent.

Note An acute event may be a clinical or external event that leads to upper motor neuron lesions resulting in spasticity for example stroke, traumatic brain injury, spinal cord injury, infection or hypoxia.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)**9547**

Moderate to severe spasticity of the upper limb following an acute event

Clinical criteria:

- The condition must be moderate to severe spasticity of the upper limb/s following an acute event, defined as a Modified Ashworth Scale rating of 3 or more, **AND**
- The treatment must only be used as second line therapy when standard management has failed; OR
- The treatment must only be used as an adjunct to physical therapy, **AND**
- The treatment must not continue if the patient does not respond (defined as not having had a decrease in spasticity rating greater than 1, using the Modified Ashworth Scale, in at least one joint) after two treatment periods (with any botulinum toxin type A), **AND**
- The treatment must not exceed a maximum of 4 treatment periods (with any botulinum toxin type A) per upper limb in the first year of treatment, and 2 treatment periods (with any botulinum toxin type A) per upper limb each year thereafter,

AND

- Patient must not have established severe contracture in the limb to be treated.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a neurologist; OR
- Must be treated by an orthopaedic surgeon; OR
- Must be treated by a rehabilitation specialist; OR
- Must be treated by a plastic surgeon; OR
- Must be treated by a geriatrician.

Standard management includes physiotherapy and/or oral spasticity agents.

Authority required (STREAMLINED)

9463

Moderate to severe spasticity of the upper limb following an acute event

Treatment Phase: Grandfathered treatment

Clinical criteria:

- Patient must have previously received non-PBS subsidised treatment with this drug for this condition prior to 1 October 2019, **AND**
- The condition must have been moderate to severe spasticity of the upper limb/s following an acute event, defined as a Modified Ashworth Scale rating of 3 or more prior to commencing non-PBS subsidised treatment, **AND**
- The treatment must only be used as second line therapy when standard management has failed; OR
- The treatment must only be used as an adjunct to physical therapy, **AND**
- The treatment must not continue if the patient did not respond (defined as not having had a decrease in spasticity rating greater than 1, using the Modified Ashworth Scale, in at least one joint) after two treatment periods (with any botulinum toxin type A), **AND**
- The treatment must not exceed a maximum of 4 treatment periods (with any botulinum toxin type A) per upper limb in the first year of treatment, and 2 treatment periods (with any botulinum toxin type A) per upper limb each year thereafter,

AND

- Patient must not have established severe contracture in the limb to be treated.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a neurologist; OR
- Must be treated by an orthopaedic surgeon; OR
- Must be treated by a rehabilitation specialist; OR
- Must be treated by a plastic surgeon; OR
- Must be treated by a geriatrician.

Standard management includes physiotherapy and/or oral spasticity agents.

clostridium botulinum type A toxin-haemagglutinin complex 300 units injection, 1 vial

10982B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	4	..	..	*1221.95	40.30	Dysport [IS]

clostridium botulinum type A toxin-haemagglutinin complex 500 units injection, 1 vial

10988H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*1094.89	40.30	Dysport [IS]

▪ **INCOBOTULINUMTOXINA**

Caution Contraindications to treatment include known sensitivity to botulinum toxin.

Note The units used to express the potency of botulinum toxin preparations currently available for PBS subsidy are not equivalent.

Authority required (STREAMLINED)

5220

Moderate to severe spasticity of the upper limb following a stroke

Clinical criteria:

- The condition must be moderate to severe spasticity of the upper limb/s following stroke, defined as a Modified Ashworth Scale rating of 3 or more, **AND**
- The treatment must not be initiated until three months post-stroke, **AND**
- The treatment must only be used as second line therapy when standard management has failed; OR
- The treatment must only be used as an adjunct to physical therapy, **AND**

- The treatment must not continue if the patient does not respond (defined as not having had a decrease in spasticity rating greater than 1, using the Modified Ashworth Scale, in at least one joint) after two treatment periods (total Botox, Dysport, and Xeomin), **AND**
- The treatment must not exceed 4 treatment periods (total Botox, Dysport, and Xeomin) per upper limb per lifetime, **AND**
- Patient must not have established severe contracture in the limb to be treated.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a neurologist; OR
- Must be treated by an orthopaedic surgeon; OR
- Must be treated by a rehabilitation specialist; OR
- Must be treated by a plastic surgeon; OR
- Must be treated by a geriatrician.

The date of the stroke must be documented in the patient's medical records when treatment is initiated.

Standard management includes physiotherapy and/or oral spasticity agents.

incobotulinumtoxinA 100 units injection, 1 vial

10983C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	4	..	..	*1547.39	40.30	Xeomin [EZ]

■ INCOBOTULINUMTOXINA

Caution Contraindications to treatment include known sensitivity to botulinum toxin.

Note The units used to express the potency of botulinum toxin preparations currently available for PBS subsidy are not equivalent.

Authority required (STREAMLINED)**5360**

Blepharospasm

Clinical criteria:

- Patient must have blepharospasm.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a neurologist; OR
- Must be treated by an ophthalmologist; OR
- Must be treated by an otolaryngology head and neck surgeon; OR
- Must be treated by a plastic surgeon.

incobotulinumtoxinA 100 units injection, 1 vial

10994P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	4	..	..	*1547.39	40.30	Xeomin [EZ]

■ INCOBOTULINUMTOXINA

Caution Contraindications to treatment include known sensitivity to botulinum toxin.

Note The units used to express the potency of botulinum toxin preparations currently available for PBS subsidy are not equivalent.

Authority required (STREAMLINED)**5222**

Spasmodic torticollis

Clinical criteria:

- Patient must have spasmodic torticollis, **AND**
- The treatment must be as monotherapy; OR
- The treatment must be as adjunctive therapy to current standard care.

Treatment criteria:

- Must be treated by a neurologist; OR
- Must be treated by a plastic surgeon; OR
- Must be treated by a rehabilitation specialist.

Population criteria:

- Patient must be aged 18 years or older.

incobotulinumtoxinA 100 units injection, 1 vial

11005F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	4	..	..	*1547.39	40.30	Xeomin [EZ]

Growth Hormone Program

SYSTEMIC HORMONAL PREPARATIONS, EXCL. SEX HORMONES AND INSULINS.....	623
PITUITARY AND HYPOTHALAMIC HORMONES AND ANALOGUES	623
ANTERIOR PITUITARY LOBE HORMONES AND ANALOGUES	623

SYSTEMIC HORMONAL PREPARATIONS, EXCL. SEX HORMONES AND INSULINS

PITUITARY AND HYPOTHALAMIC HORMONES AND ANALOGUES

ANTERIOR PITUITARY LOBE HORMONES AND ANALOGUES

Somatropin and somatropin agonists

■ SOMATROPIN

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs Programs

Reply Paid 9826

HOBART TAS 7001

Note No increase in the maximum number of repeats may be authorised.

Authority required

Severe growth hormone deficiency

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by an endocrinologist.

Clinical criteria:

- Patient must have a documented childhood onset growth hormone deficiency due to a congenital, genetic or structural cause; OR
- Patient must have adult onset growth hormone deficiency secondary to organic hypothalamic or pituitary disease, **AND**
- Patient must have an insulin tolerance test with maximum serum growth hormone (GH) less than 2.5 micrograms per litre; OR
- Patient must have an arginine infusion test with maximum serum GH less than 0.4 micrograms per litre; OR
- Patient must have a glucagon provocation test with maximum serum GH less than 3 micrograms per litre.

Population criteria:

- Patient must be aged 18 years or older.

Grandfathered patient who has previously received non-PBS subsidised treatment with this drug for this condition prior to 1 December 2018 must have met all the initial restriction criteria prior to initiating non-PBS subsidised treatment. Additional information of a baseline serum IGF-1 measurement, including the date of testing and laboratory reference range for age and sex, of less than 12 weeks prior to initiating non-PBS subsidised treatment with this drug for this condition must be provided at the time of application. A Grandfathered patient may qualify for PBS-subsidised treatment under this restriction once only. For continuing PBS-subsidised treatment, a Grandfathered patient must qualify under the Continuing treatment criteria.

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Severe Growth Hormone Deficiency supporting information form; AND
3. Confirmation of childhood onset growth hormone deficiency due to a congenital, genetic or structural cause; OR
4. Confirmation of adult onset growth hormone deficiency due to organic hypothalamic or pituitary disease; AND
5. Results of the growth hormone stimulation testing, including the date of testing, the type of test performed, the peak growth hormone concentration, and laboratory reference range for age/gender; AND
6. A baseline serum IGF-1 measurement, including the date of testing and laboratory reference range for age and sex, of less than 12 weeks old at the time of application.

Authority required

Severe growth hormone deficiency

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by an endocrinologist or in consultation with an endocrinologist.

Clinical criteria:

- Patient must have previously received PBS-subsidised therapy with this drug for this condition at the age of 18 years or older, **AND**
- Patient must maintain IGF-1 levels within the normal range for age and sex.

Population criteria:

- Patient must be aged 18 years or older.

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Severe Growth Hormone Deficiency supporting information form; AND
3. A serum IGF-1 measurement, including the date of testing and laboratory reference range for age and sex, of less than 12 weeks old at the time of application.

somatropin 10 mg/2 mL injection, 2 mL cartridge

11650E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	426.41	40.30	NutropinAq [IS]

SOMATROPIN (Recombinant human growth hormone) Powder for injection 12 mg (36 i.u.) with diluent in pre-filled pen (with preservative), 1

11495B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	510.20	40.30	Genotropin GoQuick [PF]

SOMATROPIN (Recombinant human growth hormone) Powder for injection 5 mg (15 i.u.) with diluent in pre-filled pen (with preservative), 1

11493X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	216.90	40.30	Genotropin GoQuick [PF]

■ SOMATROPIN

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Prior Written Approval of Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Short stature and slow growth

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have a current height at or below the 1st percentile for age and sex, **AND**
- Patient must have a growth velocity below the 25th percentile for bone age and sex measured over a 12 month interval (or a 6 month interval for an older child), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must not have a bone age of 2.5 years or less, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7 cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0 cm, **AND**
- Patient must be male and must not have maturational or constitutional delay in combination with an estimated mature height equal to or above 160.1 cm; OR
- Patient must be female and must not have maturational or constitutional delay in combination with an estimated mature height equal to or above 148.0 cm.

Population criteria:

- Patient must be aged 3 years or older.

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; **AND**
3. A minimum of 12 months of recent growth data (height and weight measurements) or a minimum of 6 months of recent growth data for an older child. The most recent data must not be more than three months old at the time of application; **AND**
4. A bone age result performed within the last 12 months; **AND**
5. Confirmation of the patient's maturational or constitutional delay status; **AND**
6. If the patient has maturational or constitutional delay, confirmation that the patient has an estimated mature height below the 1st adult height percentile; **AND**

7. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with biochemical growth hormone deficiency

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must have a current height at or below the 1st percentile for age and sex; OR
- Patient must have a current height above the 1st and at or below the 25th percentiles for age and sex and a growth velocity below the 25th percentile for bone age and sex measured over a 12 month interval (or a 6 month interval for an older child); OR
- Patient must have a current height above the 1st and at or below the 25th percentiles for age and sex and an annual growth velocity of 8 cm per year or less if the patient has a bone age of 2.5 years or less, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Population criteria:

- Patient must be aged 3 years or older.

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; AND
3. (a) A minimum of 12 months of recent growth data (height and weight measurements) or a minimum of 6 months of recent growth data for an older child. The most recent data must not be more than three months old at the time of application; OR
- (b) Height and weight measurements, not more than three months old at the time of application, for a patient whose current height is at or below the 1st percentile for age and sex; AND
4. A bone age result performed within the last 12 months; AND
5. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; AND
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Biochemical growth hormone deficiency should not be secondary to an intracranial lesion or cranial irradiation for applications under this category.

Authority required

Growth retardation secondary to an intracranial lesion, or cranial irradiation

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have had an intracranial lesion and have undergone a 12 month period of observation following completion of treatment for the intracranial lesion (all treatment); OR
- Patient must have had an intracranial lesion, have received medical advice that it is unsafe to treat the intracranial lesion, and have undergone a 12 month period of observation since initial diagnosis of the intracranial lesion; OR
- Patient must have received cranial irradiation without having had an intracranial lesion, and have undergone a 12 month period of observation following completion of treatment for the condition for which cranial irradiation was received, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must have a current height at or below the 1st percentile for age and sex; OR
- Patient must have a current height above the 1st percentile for age and sex and a growth velocity below the 25th percentile for bone age and sex measured over a 12 month interval (or a 6 month interval for an older child); OR
- Patient must have a current height above the 1st percentile for age and sex and an annual growth velocity of 8 cm per year or less if the patient has a bone age of 2.5 years or less, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Population criteria:

- Patient must be aged 3 years or older.

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; **AND**
3. (a) A minimum of 12 months of recent growth data (height and weight measurements) or a minimum of 6 months of recent growth data for an older child. The most recent data must not be more than three months old at the time of application; OR
- (b) Height and weight measurements, not more than three months old at the time of application, for a patient whose current height is at or below the 1st percentile for age and sex; **AND**

4. A bone age result performed within the last 12 months; AND
 5. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; AND
 6. (a) Confirmation that the patient has had an intracranial lesion and has undergone a 12 month period of observation following completion of treatment for the intracranial lesion (all treatment); OR
(b) Confirmation that the patient has had an intracranial lesion, has received medical advice that it is unsafe to treat the intracranial lesion, and has undergone a 12 month period of observation since initial diagnosis of the intracranial lesion; OR
(c) Confirmation that the patient has received cranial irradiation without having had an intracranial lesion, and has undergone a 12 month period of observation following completion of treatment for the condition for which cranial irradiation was received; AND
 7. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).
- Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Biochemical growth hormone deficiency and precocious puberty

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must be male and have commenced puberty (demonstrated by Tanner stage 2 genital or pubic hair development or testicular volumes greater than or equal to 4 mL) before the chronological age of 9 years; OR
- Patient must be female and have commenced puberty (demonstrated by Tanner stage 2 breast or pubic hair development) before the chronological age of 8 years; OR
- Patient must be female and menarche occurred before the chronological age of 10 years, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must be undergoing Gonadotrophin Releasing Hormone agonist therapy for pubertal suppression, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Population criteria:

- Patient must be aged 3 years or older.

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; AND
3. (a) A minimum of 12 months of recent growth data (height and weight) at intervals no greater than six months. The most recent data must not be older than three months; OR

(b) A minimum of 6 months of recent growth data (height and weight) for older children (males chronological age 12 and over or bone age 10 and over, females chronological age 10 and over or bone age 8 and over). The most recent data must not be older than three months; AND

4. A bone age result performed within the last 12 months; AND

5. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; AND

6. Confirmation that the patient has precocious puberty; AND

7. Confirmation that the patient is undergoing Gonadotropin Releasing Hormone agonist therapy, for pubertal suppression; AND

8. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

Clinical criteria:

- Patient must have a structural lesion that is not neoplastic; OR
- Patient must have had a structural lesion that was neoplastic and have undergone a 12 month period of observation following completion of treatment for the structural lesion (all treatment); OR
- Patient must have a structural lesion that is neoplastic, have received medical advice that it is unsafe to treat the structural lesion, and have undergone a 12 month period of observation since initial diagnosis of the structural lesion,

AND

- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must have other hypothalamic/pituitary hormone deficits (includes ACTH, TSH, GnRH and/or vasopressin/ADH deficiencies), **AND**
- Patient must have hypothalamic obesity, **AND**
- Patient must have a growth velocity above the 25th percentile for bone age and sex measured over a 12 month interval (or a 6 month interval for an older child); OR
- Patient must have an annual growth velocity of greater than 8 cm per year if the patient has a bone age of 2.5 years or less, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Population criteria:

- Patient must be aged 3 years or older.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; AND
3. A minimum of 12 months of recent growth data (height and weight measurements) or a minimum of 6 months of recent growth data for an older child. The most recent data must not be more than three months old at the time of application; AND
4. A bone age result performed within the last 12 months; AND
5. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; AND
6. (a) Confirmation that the patient has a structural lesion that is not neoplastic; OR
(b) Confirmation that the patient had a structural lesion that was neoplastic and has undergone a 12 month period of observation following completion of treatment for the structural lesion (all treatment); OR
(c) Confirmation that the patient has a structural lesion that is neoplastic, has received medical advice that it is unsafe to treat the structural lesion, and has undergone a 12 month period of observation since initial diagnosis of the structural lesion; AND
7. Confirmation that the patient has other hypothalamic/pituitary hormone deficits; AND
8. Confirmation that the patient has hypothalamic obesity; AND
9. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Testing for biochemical growth hormone deficiency must have been performed at a time when all other pituitary hormone deficits were being adequately replaced.

Authority required

Short stature associated with Turner syndrome

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

Clinical criteria:

- Patient must have a current height at or below the 95th percentile for age on the Turner syndrome growth curve for girls, **AND**
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as a loss of a whole X chromosome in all cells (45X), and gender of rearing is female; OR
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as a loss of a whole X chromosome in some cells (mosaic 46XX/45X), and gender of rearing is female; OR
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as genetic loss or rearrangement of an X chromosome (such as isochromosome X, ring-chromosome, or partial deletion of an X chromosome), and gender of rearing is female, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must not have a bone age of 2.5 years or less, **AND**
- Patient must not have a height greater than or equal to 155.0 cm, **AND**
- Patient must not have a bone age of 13.5 years or greater.

Population criteria:

- Patient must be aged 3 years or older.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; AND
3. (a) A minimum of 12 months of recent growth data (height and weight) at intervals no greater than six months. The most recent data must not be older than three months; OR
(b) A minimum of 6 months of recent growth data (height and weight) for older children (females chronological age 10 and over or bone age 8 and over). The most recent data must not be older than three months; AND
4. A bone age result performed within the last 12 months; AND
5. Confirmation that the patient has diagnostic results consistent with Turner syndrome; AND

6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature due to short stature homeobox (SHOX) gene disorders

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as a karyotype confirming the presence of a SHOX mutation/deletion without the presence of mixed gonadal dysgenesis; OR
- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as mixed gonadal dysgenesis (45X mosaic karyotype with the presence of any Y chromosome material and/or SRY gene positive by FISH study) and have an appropriate plan of management in place for the patient's increased risk of gonadoblastoma, **AND**
- Patient must have a current height at or below the 1st percentile for age and sex, **AND**
- Patient must have a growth velocity below the 25th percentile for bone age and sex measured over a 12 month interval (or a 6 month interval for an older child); OR
- Patient must have an annual growth velocity of 8 cm per year or less if the patient has a bone or chronological age of 2.5 years or less, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes (excluding gonadoblastoma secondary to mixed gonadal dysgenesis), **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Population criteria:

- Patient must be aged 3 years or older.

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special**

Arrangement 2015 and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; AND
3. A minimum of 12 months of recent growth data (height and weight measurements) or a minimum of 6 months of recent growth data for an older child. The most recent data must not be more than three months old at the time of application; AND
4. A bone age result performed within the last 12 months; AND
5. Confirmation that the patient has diagnostic results consistent with a short stature homeobox (SHOX) gene disorder; AND
6. If the patient's condition is secondary to mixed gonadal dysgenesis, confirmation that an appropriate plan of management for the patient's increased risk of gonadoblastoma is in place; AND
7. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with chronic renal insufficiency

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

Clinical criteria:

- Patient must have an estimated glomerular filtration rate less than 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula, and not have undergone a renal transplant; OR
- Patient must have an estimated glomerular filtration rate less than 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula, have undergone a renal transplant, and have undergone a 12 month period of observation following the transplant, **AND**

- Patient must have a current height at or below the 1st percentile for age and sex; OR
- Patient must have a current height above the 1st and at or below the 25th percentiles for age and sex and a growth velocity less than or equal to the 25th percentile for bone age and sex measured over a 12 month interval (or a 6 month interval for an older child); OR
- Patient must have a current height above the 1st and at or below the 25th percentiles for age and sex and an annual growth velocity of 8 cm per year or less if the patient has a bone age of 2.5 years or less, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7 cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0 cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Population criteria:

- Patient must be aged 3 years or older.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; AND
3. (a) A minimum of 12 months of recent growth data (height and weight measurements) or a minimum of 6 months of recent growth data for an older child. The most recent data must not be more than three months old at the time of application; OR
(b) Height and weight measurements, not more than three months old at the time of application, for a patient whose current height is at or below the 1st percentile for age and sex; AND
4. A bone age result performed within the last 12 months; AND
5. Confirmation that the patient has an estimated glomerular filtration rate less than 30mL/minute/1.73m² ; AND
6. If a renal transplant has taken place, confirmation that the patient has undergone a 12 month period of observation following transplantation; AND
7. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

somatropin 5 mg/1.5 mL injection, 1.5 mL cartridge

10518N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	223.61	40.30	Omnitrope Surepal 5 [SZ]

somatropin 5 mg/1.5 mL injection, 1.5 mL cartridge

6476W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	223.61	40.30	Scitropin A [SA]

somatropin 10 mg/1.5 mL injection, 1.5 mL cartridge

10514J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	439.82	40.30	Omnitrope Surepal 10 [SZ]

somatropin 10 mg/1.5 mL injection, 1.5 mL cartridge

6311E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	439.82	40.30	SciTropin A [SA]

somatropin 15 mg/1.5 mL injection, 1.5 mL cartridge

10446T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	656.04	40.30	Omnitrope Surepal 15 [SZ]

■ SOMATROPIN

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Prior Written Approval of Complex Drugs

Reply Paid 9826
HOBART TAS 7001

Authority required

Short stature and slow growth

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have a current height at or below the 1st percentile for age and sex, **AND**
- Patient must have a growth velocity below the 25th percentile for bone age and sex measured over a 12 month interval (or a 6 month interval for an older child), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must not have a bone age of 2.5 years or less, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7 cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0 cm, **AND**
- Patient must be male and must not have maturational or constitutional delay in combination with an estimated mature height equal to or above 160.1 cm; OR
- Patient must be female and must not have maturational or constitutional delay in combination with an estimated mature height equal to or above 148.0 cm.

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; AND
3. A minimum of 12 months of recent growth data (height and weight measurements) or a minimum of 6 months of recent growth data for an older child. The most recent data must not be more than three months old at the time of application; AND
4. A bone age result performed within the last 12 months; AND
5. Confirmation of the patient's maturational or constitutional delay status; AND
6. If the patient has maturational or constitutional delay, confirmation that the patient has an estimated mature height below the 1st adult height percentile; AND
7. The proprietary name (brand), form and strength of somatotropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with biochemical growth hormone deficiency

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test

(pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR

- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must have a current height at or below the 1st percentile for age and sex; OR
- Patient must have a current height above the 1st and at or below the 25th percentiles for age and sex and a growth velocity below the 25th percentile for bone age and sex measured over a 12 month interval (or a 6 month interval for an older child); OR
- Patient must have a current height above the 1st and at or below the 25th percentiles for age and sex and an annual growth velocity of 14 cm per year or less if the patient has a chronological age of 2 years or less; OR
- Patient must have a current height above the 1st and at or below the 25th percentiles for age and sex and an annual growth velocity of 8 cm per year or less if the patient has a bone or chronological age of 2.5 years or less, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; **AND**
3. (a) A minimum of 12 months of recent growth data (height and weight measurements) or a minimum of 6 months of recent growth data for an older child. The most recent data must not be more than three months old at the time of application; OR
(b) Height and weight measurements, not more than three months old at the time of application, for a patient whose current height is at or below the 1st percentile for age and sex; **AND**
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
5. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; **AND**
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Biochemical growth hormone deficiency should not be secondary to an intracranial lesion or cranial irradiation for applications under this category.

Authority required

Growth retardation secondary to an intracranial lesion, or cranial irradiation

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have had an intracranial lesion and have undergone a 12 month period of observation following completion of treatment for the intracranial lesion (all treatment); OR
- Patient must have had an intracranial lesion, have received medical advice that it is unsafe to treat the intracranial lesion, and have undergone a 12 month period of observation since initial diagnosis of the intracranial lesion; OR
- Patient must have received cranial irradiation without having had an intracranial lesion, and have undergone a 12 month period of observation following completion of treatment for the condition for which cranial irradiation was received, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test

(pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR

- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels, **AND**
- Patient must have a current height at or below the 1st percentile for age and sex; OR
- Patient must have a current height above the 1st percentile for age and sex and a growth velocity below the 25th percentile for bone age and sex measured over a 12 month interval (or a 6 month interval for an older child); OR
- Patient must have a current height above the 1st percentile for age and sex and an annual growth velocity of 14 cm per year or less if the patient has a chronological age of 2 years or less; OR
- Patient must have a current height above the 1st percentile for age and sex and an annual growth velocity of 8 cm per year or less if the patient has a bone or chronological age of 2.5 years or less, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; AND
3. (a) A minimum of 12 months of recent growth data (height and weight measurements) or a minimum of 6 months of recent growth data for an older child. The most recent data must not be more than three months old at the time of application; OR
(b) Height and weight measurements, not more than three months old at the time of application, for a patient whose current height is at or below the 1st percentile for age and sex; AND
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
5. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; AND
6. (a) Confirmation that the patient has had an intracranial lesion and has undergone a 12 month period of observation following completion of treatment for the intracranial lesion (all treatment); OR
(b) Confirmation that the patient has had an intracranial lesion, has received medical advice that it is unsafe to treat the intracranial lesion, and has undergone a 12 month period of observation since initial diagnosis of the intracranial lesion; OR
(c) Confirmation that the patient has received cranial irradiation without having had an intracranial lesion, and has undergone a 12 month period of observation following completion of treatment for the condition for which cranial irradiation was received; AND
7. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

Clinical criteria:

- Patient must have a chronological age of less than 2 years, **AND**
- Patient must have a documented clinical risk of hypoglycaemia, **AND**
- Patient must have documented evidence that the risk of hypoglycaemia is secondary to biochemical growth hormone deficiency, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; **AND**
3. Recent growth data (height and weight, not older than three months); **AND**
4. Confirmation that the patient has a documented clinical risk of hypoglycaemia; **AND**
5. Confirmation that the patient has documented evidence that the risk of hypoglycaemia is secondary to biochemical growth hormone deficiency; **AND**
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Biochemical growth hormone deficiency and precocious puberty

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must be male and have commenced puberty (demonstrated by Tanner stage 2 genital or pubic hair development or testicular volumes greater than or equal to 4 mL) before the chronological age of 9 years; **OR**
- Patient must be female and have commenced puberty (demonstrated by Tanner stage 2 breast or pubic hair development) before the chronological age of 8 years; **OR**
- Patient must be female and menarche occurred before the chronological age of 10 years, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); **OR**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); **OR**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); **OR**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; **OR**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must be undergoing Gonadotrophin Releasing Hormone agonist therapy for pubertal suppression, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; **OR**
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; **OR**
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; AND
3. (a) A minimum of 12 months of recent growth data (height and weight) at intervals no greater than six months. The most recent data must not be older than three months; OR
(b) A minimum of 6 months of recent growth data (height and weight) for older children (males chronological age 12 and over or bone age 10 and over, females chronological age 10 and over or bone age 8 and over). The most recent data must not be older than three months; AND
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
5. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; AND
6. Confirmation that the patient has precocious puberty; AND
7. Confirmation that the patient is undergoing Gonadotropin Releasing Hormone agonist therapy, for pubertal suppression; AND
8. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have a structural lesion that is not neoplastic; OR
- Patient must have had a structural lesion that was neoplastic and have undergone a 12 month period of observation following completion of treatment for the structural lesion (all treatment); OR
- Patient must have a structural lesion that is neoplastic, have received medical advice that it is unsafe to treat the structural lesion, and have undergone a 12 month period of observation since initial diagnosis of the structural lesion, AND
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, AND
- Patient must have other hypothalamic/pituitary hormone deficits (includes ACTH, TSH, GnRH and/or vasopressin/ADH deficiencies), AND
- Patient must have hypothalamic obesity, AND
- Patient must have a growth velocity above the 25th percentile for bone age and sex measured over a 12 month interval (or a 6 month interval for an older child); OR
- Patient must have an annual growth velocity of greater than 14 cm per year if the patient has a chronological age of 2 years or less; OR
- Patient must have an annual growth velocity of greater than 8 cm per year if the patient has a bone or chronological age of 2.5 years or less, AND
- Patient must not have diabetes mellitus, AND
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, AND
- Patient must not have an active tumour or evidence of tumour growth or activity, AND

- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; AND
3. A minimum of 12 months of recent growth data (height and weight measurements) or a minimum of 6 months of recent growth data for an older child. The most recent data must not be more than three months old at the time of application; AND
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
5. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; AND
6. (a) Confirmation that the patient has a structural lesion that is not neoplastic; OR
(b) Confirmation that the patient had a structural lesion that was neoplastic and has undergone a 12 month period of observation following completion of treatment for the structural lesion (all treatment); OR
(c) Confirmation that the patient has a structural lesion that is neoplastic, has received medical advice that it is unsafe to treat the structural lesion, and has undergone a 12 month period of observation since initial diagnosis of the structural lesion; AND
7. Confirmation that the patient has other hypothalamic/pituitary hormone deficits; AND
8. Confirmation that the patient has hypothalamic obesity; AND
9. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Testing for biochemical growth hormone deficiency must have been performed at a time when all other pituitary hormone deficits were being adequately replaced.

Authority required

Short stature associated with Turner syndrome

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

Clinical criteria:

- Patient must have a current height at or below the 95th percentile for age on the Turner syndrome growth curve for girls, **AND**
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as a loss of a whole X chromosome in all cells (45X), and gender of rearing is female; OR
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as a loss of a whole X chromosome in some cells (mosaic 46XX/45X), and gender of rearing is female; OR
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as genetic loss or rearrangement of an X chromosome (such as isochromosome X, ring-chromosome, or partial deletion of an X chromosome), and gender of rearing is female, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must not have a bone age of 2.5 years or less, **AND**
- Patient must not have a height greater than or equal to 155.0cm, **AND**
- Patient must not have a bone age of 13.5 years or greater.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND

2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; AND
 3. (a) A minimum of 12 months of recent growth data (height and weight) at intervals no greater than six months. The most recent data must not be older than three months; OR
 - (b) A minimum of 6 months of recent growth data (height and weight) for older children (females chronological age 10 and over or bone age 8 and over). The most recent data must not be older than three months; AND
 4. A bone age result performed within the last 12 months; AND
 5. Confirmation that the patient has diagnostic results consistent with Turner syndrome; AND
 6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).
- Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature due to short stature homeobox (SHOX) gene disorders

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as a karyotype confirming the presence of a SHOX mutation/deletion without the presence of mixed gonadal dysgenesis; OR
- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as mixed gonadal dysgenesis (45X mosaic karyotype with the presence of any Y chromosome material and/or SRY gene positive by FISH study) and have an appropriate plan of management in place for the patient's increased risk of gonadoblastoma, **AND**
- Patient must have a current height at or below the 1st percentile for age and sex, **AND**
- Patient must have a growth velocity below the 25th percentile for bone age and sex measured over a 12 month interval (or a 6 month interval for an older child); OR
- Patient must have an annual growth velocity of 14 cm per year or less if the patient has a chronological age of 2 years or less; OR
- Patient must have an annual growth velocity of 8 cm per year or less if the patient has a bone or chronological age of 2.5 years or less, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes (excluding gonadoblastoma secondary to mixed gonadal dysgenesis), **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; AND
3. A minimum of 12 months of recent growth data (height and weight measurements) or a minimum of 6 months of recent growth data for an older child. The most recent data must not be more than three months old at the time of application; AND
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
5. Confirmation that the patient has diagnostic results consistent with a short stature homeobox (SHOX) gene disorder; AND
6. If the patient's condition is secondary to mixed gonadal dysgenesis, confirmation that an appropriate plan of management for the patient's increased risk of gonadoblastoma is in place; AND
7. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

somatropin 4 mg injection [1 vial] (&) inert substance diluent [1 vial], 1 pack

6266T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	180.36	40.30	Zomacton [FP]

■ SOMATROPIN

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Prior Written Approval of Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Short stature and slow growth

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have a current height at or below the 1st percentile for age and sex, **AND**
- Patient must have a growth velocity below the 25th percentile for bone age and sex measured over a 12 month interval (or a 6 month interval for an older child), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must not have a bone age of 2.5 years or less, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7 cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0 cm, **AND**
- Patient must be male and must not have maturational or constitutional delay in combination with an estimated mature height equal to or above 160.1 cm; OR
- Patient must be female and must not have maturational or constitutional delay in combination with an estimated mature height equal to or above 148.0 cm.

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special**

Arrangement 2015 and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; AND
3. A minimum of 12 months of recent growth data (height and weight measurements) or a minimum of 6 months of recent growth data for an older child. The most recent data must not be more than three months old at the time of application; AND
4. A bone age result performed within the last 12 months; AND
5. Confirmation of the patient's maturational or constitutional delay status; AND
6. If the patient has maturational or constitutional delay, confirmation that the patient has an estimated mature height below the 1st adult height percentile; AND
7. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with biochemical growth hormone deficiency

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test

(pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR

- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels, **AND**
- Patient must have a current height at or below the 1st percentile for age and sex; OR
- Patient must have a current height above the 1st and at or below the 25th percentiles for age and sex and a growth velocity below the 25th percentile for bone age and sex measured over a 12 month interval (or a 6 month interval for an older child); OR
- Patient must have a current height above the 1st and at or below the 25th percentiles for age and sex and an annual growth velocity of 14 cm per year or less if the patient has a chronological age of 2 years or less; OR
- Patient must have a current height above the 1st and at or below the 25th percentiles for age and sex and an annual growth velocity of 8 cm per year or less if the patient has a bone or chronological age of 2.5 years or less, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special**

Arrangement 2015 and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; **AND**
3. (a) A minimum of 12 months of recent growth data (height and weight measurements) or a minimum of 6 months of recent growth data for an older child. The most recent data must not be more than three months old at the time of application; OR
(b) Height and weight measurements, not more than three months old at the time of application, for a patient whose current height is at or below the 1st percentile for age and sex; **AND**
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
5. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; **AND**
6. The proprietary name (brand), form and strength of somatotropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Biochemical growth hormone deficiency should not be secondary to an intracranial lesion or cranial irradiation for applications under this category.

Authority required

Growth retardation secondary to an intracranial lesion, or cranial irradiation

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have had an intracranial lesion and have undergone a 12 month period of observation following completion of treatment for the intracranial lesion (all treatment); OR
- Patient must have had an intracranial lesion, have received medical advice that it is unsafe to treat the intracranial lesion, and have undergone a 12 month period of observation since initial diagnosis of the intracranial lesion; OR
- Patient must have received cranial irradiation without having had an intracranial lesion, and have undergone a 12 month period of observation following completion of treatment for the condition for which cranial irradiation was received, **AND**

- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must have a current height at or below the 1st percentile for age and sex; OR
- Patient must have a current height above the 1st percentile for age and sex and a growth velocity below the 25th percentile for bone age and sex measured over a 12 month interval (or a 6 month interval for an older child); OR
- Patient must have a current height above the 1st percentile for age and sex and an annual growth velocity of 14 cm per year or less if the patient has a chronological age of 2 years or less; OR
- Patient must have a current height above the 1st percentile for age and sex and an annual growth velocity of 8 cm per year or less if the patient has a bone or chronological age of 2.5 years or less, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; **AND**
3. (a) A minimum of 12 months of recent growth data (height and weight measurements) or a minimum of 6 months of recent growth data for an older child. The most recent data must not be more than three months old at the time of application; OR
(b) Height and weight measurements, not more than three months old at the time of application, for a patient whose current height is at or below the 1st percentile for age and sex; **AND**
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
5. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; **AND**
6. (a) Confirmation that the patient has had an intracranial lesion and has undergone a 12 month period of observation following completion of treatment for the intracranial lesion (all treatment); OR
(b) Confirmation that the patient has had an intracranial lesion, has received medical advice that it is unsafe to treat the intracranial lesion, and has undergone a 12 month period of observation since initial diagnosis of the intracranial lesion; OR
(c) Confirmation that the patient has received cranial irradiation without having had an intracranial lesion, and has undergone a 12 month period of observation following completion of treatment for the condition for which cranial irradiation was received; **AND**
7. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

Clinical criteria:

- Patient must have a chronological age of less than 2 years, **AND**
- Patient must have a documented clinical risk of hypoglycaemia, **AND**
- Patient must have documented evidence that the risk of hypoglycaemia is secondary to biochemical growth hormone deficiency, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; **AND**
3. Recent growth data (height and weight, not older than three months); **AND**
4. Confirmation that the patient has a documented clinical risk of hypoglycaemia; **AND**
5. Confirmation that the patient has documented evidence that the risk of hypoglycaemia is secondary to biochemical growth hormone deficiency; **AND**
6. The proprietary name (brand), form and strength of somatotropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Biochemical growth hormone deficiency and precocious puberty

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must be male and have commenced puberty (demonstrated by Tanner stage 2 genital or pubic hair development or testicular volumes greater than or equal to 4 mL) before the chronological age of 9 years; OR
- Patient must be female and have commenced puberty (demonstrated by Tanner stage 2 breast or pubic hair development) before the chronological age of 8 years; OR
- Patient must be female and menarche occurred before the chronological age of 10 years, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must be undergoing Gonadotrophin Releasing Hormone agonist therapy for pubertal suppression, **AND**

- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; **AND**
3. (a) A minimum of 12 months of recent growth data (height and weight) at intervals no greater than six months. The most recent data must not be older than three months; OR
(b) A minimum of 6 months of recent growth data (height and weight) for older children (males chronological age 12 and over or bone age 10 and over, females chronological age 10 and over or bone age 8 and over). The most recent data must not be older than three months; **AND**
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
5. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; **AND**
6. Confirmation that the patient has precocious puberty; **AND**
7. Confirmation that the patient is undergoing Gonadotropin Releasing Hormone agonist therapy, for pubertal suppression; **AND**
8. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have a structural lesion that is not neoplastic; OR
- Patient must have had a structural lesion that was neoplastic and have undergone a 12 month period of observation following completion of treatment for the structural lesion (all treatment); OR
- Patient must have a structural lesion that is neoplastic, have received medical advice that it is unsafe to treat the structural lesion, and have undergone a 12 month period of observation since initial diagnosis of the structural lesion, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must have other hypothalamic/pituitary hormone deficits (includes ACTH, TSH, GnRH and/or vasopressin/ADH deficiencies), **AND**

- Patient must have hypothalamic obesity, **AND**
- Patient must have a growth velocity above the 25th percentile for bone age and sex measured over a 12 month interval (or a 6 month interval for an older child); OR
- Patient must have an annual growth velocity of greater than 14 cm per year if the patient has a chronological age of 2 years or less; OR
- Patient must have an annual growth velocity of greater than 8 cm per year if the patient has a bone or chronological age of 2.5 years or less, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; AND
3. A minimum of 12 months of recent growth data (height and weight measurements) or a minimum of 6 months of recent growth data for an older child. The most recent data must not be more than three months old at the time of application; AND
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
5. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; AND
6. (a) Confirmation that the patient has a structural lesion that is not neoplastic; OR
(b) Confirmation that the patient had a structural lesion that was neoplastic and has undergone a 12 month period of observation following completion of treatment for the structural lesion (all treatment); OR
(c) Confirmation that the patient has a structural lesion that is neoplastic, has received medical advice that it is unsafe to treat the structural lesion, and has undergone a 12 month period of observation since initial diagnosis of the structural lesion; AND
7. Confirmation that the patient has other hypothalamic/pituitary hormone deficits; AND
8. Confirmation that the patient has hypothalamic obesity; AND
9. The proprietary name (brand), form and strength of somatotropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Testing for biochemical growth hormone deficiency must have been performed at a time when all other pituitary hormone deficits were being adequately replaced.

Authority required

Short stature associated with Turner syndrome

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

Clinical criteria:

- Patient must have a current height at or below the 95th percentile for age on the Turner syndrome growth curve for girls, **AND**
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as a loss of a whole X chromosome in all cells (45X), and gender of rearing is female; OR
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as a loss of a whole X chromosome in some cells (mosaic 46XX/45X), and gender of rearing is female; OR
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as genetic loss or rearrangement of an X chromosome (such as isochromosome X, ring-chromosome, or partial deletion of an X chromosome), and gender of rearing is female, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**

- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must not have a bone age of 2.5 years or less, **AND**
- Patient must not have a height greater than or equal to 155.0cm, **AND**
- Patient must not have a bone age of 13.5 years or greater.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; **AND**
3. (a) A minimum of 12 months of recent growth data (height and weight) at intervals no greater than six months. The most recent data must not be older than three months; **OR**
(b) A minimum of 6 months of recent growth data (height and weight) for older children (females chronological age 10 and over or bone age 8 and over). The most recent data must not be older than three months; **AND**
4. A bone age result performed within the last 12 months; **AND**
5. Confirmation that the patient has diagnostic results consistent with Turner syndrome; **AND**
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature due to short stature homeobox (SHOX) gene disorders

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as a karyotype confirming the presence of a SHOX mutation/deletion without the presence of mixed gonadal dysgenesis; **OR**
- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as mixed gonadal dysgenesis (45X mosaic karyotype with the presence of any Y chromosome material and/or SRY gene positive by FISH study) and have an appropriate plan of management in place for the patient's increased risk of gonadoblastoma, **AND**
- Patient must have a current height at or below the 1st percentile for age and sex, **AND**
- Patient must have a growth velocity below the 25th percentile for bone age and sex measured over a 12 month interval (or a 6 month interval for an older child); **OR**
- Patient must have an annual growth velocity of 14 cm per year or less if the patient has a chronological age of 2 years or less; **OR**
- Patient must have an annual growth velocity of 8 cm per year or less if the patient has a bone or chronological age of 2.5 years or less, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes (excluding gonadoblastoma secondary to mixed gonadal dysgenesis), **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; **OR**
- Patient must be female and must not have a height greater than or equal to 155.0cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; **OR**
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; **OR**
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; **AND**
3. A minimum of 12 months of recent growth data (height and weight measurements) or a minimum of 6 months of recent growth data for an older child. The most recent data must not be more than three months old at the time of application; **AND**
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
5. Confirmation that the patient has diagnostic results consistent with a short stature homeobox (SHOX) gene disorder; **AND**
6. If the patient's condition is secondary to mixed gonadal dysgenesis, confirmation that an appropriate plan of management for the patient's increased risk of gonadoblastoma is in place; **AND**

7. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with chronic renal insufficiency

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

Clinical criteria:

- Patient must have an estimated glomerular filtration rate less than 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula, and not have undergone a renal transplant; OR
- Patient must have an estimated glomerular filtration rate less than 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula, have undergone a renal transplant, and have undergone a 12 month period of observation following the transplant, **AND**
- Patient must have a current height at or below the 1st percentile for age and sex; OR
- Patient must have a current height above the 1st and at or below the 25th percentiles for age and sex and a growth velocity less than or equal to the 25th percentile for bone age and sex measured over a 12 month interval (or a 6 month interval for an older child); OR
- Patient must have a current height above the 1st and at or below the 25th percentiles for age and sex and an annual growth velocity of 8 cm per year or less if the patient has a bone age of 2.5 years or less, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7 cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0 cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Population criteria:

- Patient must be aged 3 years or older.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; **AND**
3. (a) A minimum of 12 months of recent growth data (height and weight measurements) or a minimum of 6 months of recent growth data for an older child. The most recent data must not be more than three months old at the time of application; **OR**
(b) Height and weight measurements, not more than three months old at the time of application, for a patient whose current height is at or below the 1st percentile for age and sex; **AND**
4. A bone age result performed within the last 12 months; **AND**
5. Confirmation that the patient has an estimated glomerular filtration rate less than 30mL/minute/1.73m² ; **AND**
6. If a renal transplant has taken place, confirmation that the patient has undergone a 12 month period of observation following transplantation; **AND**
7. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

somatropin 12 mg/1.5 mL injection, 1.5 mL cartridge

5824M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	526.31	40.30	Saizen [SG]

somatropin 6 mg/1.03 mL injection, 1.03 mL cartridge

5822K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	266.85	40.30	Saizen [SG]

somatropin 20 mg/2.5 mL injection, 2.5 mL cartridge

3388H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	872.25	40.30	Saizen [SG]

■ SOMATROPIN

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Prior Written Approval of Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Short stature and slow growth

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have a current height at or below the 1st percentile for age and sex, **AND**
- Patient must have a growth velocity below the 25th percentile for bone age and sex measured over a 12 month interval (or a 6 month interval for an older child), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must not have a bone age of 2.5 years or less, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7 cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0 cm, **AND**
- Patient must be male and must not have maturational or constitutional delay in combination with an estimated mature height equal to or above 160.1 cm; OR
- Patient must be female and must not have maturational or constitutional delay in combination with an estimated mature height equal to or above 148.0 cm.

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; **AND**
3. A minimum of 12 months of recent growth data (height and weight measurements) or a minimum of 6 months of recent growth data for an older child. The most recent data must not be more than three months old at the time of application; **AND**
4. A bone age result performed within the last 12 months; **AND**
5. Confirmation of the patient's maturational or constitutional delay status; **AND**
6. If the patient has maturational or constitutional delay, confirmation that the patient has an estimated mature height below the 1st adult height percentile; **AND**
7. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with biochemical growth hormone deficiency

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone

stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR

- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must have a current height at or below the 1st percentile for age and sex; OR
- Patient must have a current height above the 1st and at or below the 25th percentiles for age and sex and a growth velocity below the 25th percentile for bone age and sex measured over a 12 month interval (or a 6 month interval for an older child); OR
- Patient must have a current height above the 1st and at or below the 25th percentiles for age and sex and an annual growth velocity of 14 cm per year or less if the patient has a chronological age of 2 years or less; OR
- Patient must have a current height above the 1st and at or below the 25th percentiles for age and sex and an annual growth velocity of 8 cm per year or less if the patient has a bone or chronological age of 2.5 years or less, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; AND
3. (a) A minimum of 12 months of recent growth data (height and weight measurements) or a minimum of 6 months of recent growth data for an older child. The most recent data must not be more than three months old at the time of application; OR
(b) Height and weight measurements, not more than three months old at the time of application, for a patient whose current height is at or below the 1st percentile for age and sex; AND
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
5. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; AND
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Biochemical growth hormone deficiency should not be secondary to an intracranial lesion or cranial irradiation for applications under this category.

Authority required

Growth retardation secondary to an intracranial lesion, or cranial irradiation

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have had an intracranial lesion and have undergone a 12 month period of observation following completion of treatment for the intracranial lesion (all treatment); OR
- Patient must have had an intracranial lesion, have received medical advice that it is unsafe to treat the intracranial lesion, and have undergone a 12 month period of observation since initial diagnosis of the intracranial lesion; OR

- Patient must have received cranial irradiation without having had an intracranial lesion, and have undergone a 12 month period of observation following completion of treatment for the condition for which cranial irradiation was received, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must have a current height at or below the 1st percentile for age and sex; OR
- Patient must have a current height above the 1st percentile for age and sex and a growth velocity below the 25th percentile for bone age and sex measured over a 12 month interval (or a 6 month interval for an older child); OR
- Patient must have a current height above the 1st percentile for age and sex and an annual growth velocity of 14 cm per year or less if the patient has a chronological age of 2 years or less; OR
- Patient must have a current height above the 1st percentile for age and sex and an annual growth velocity of 8 cm per year or less if the patient has a bone or chronological age of 2.5 years or less, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; **AND**
3. (a) A minimum of 12 months of recent growth data (height and weight measurements) or a minimum of 6 months of recent growth data for an older child. The most recent data must not be more than three months old at the time of application; OR
(b) Height and weight measurements, not more than three months old at the time of application, for a patient whose current height is at or below the 1st percentile for age and sex; **AND**
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
5. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; **AND**
6. (a) Confirmation that the patient has had an intracranial lesion and has undergone a 12 month period of observation following completion of treatment for the intracranial lesion (all treatment); OR
(b) Confirmation that the patient has had an intracranial lesion, has received medical advice that it is unsafe to treat the intracranial lesion, and has undergone a 12 month period of observation since initial diagnosis of the intracranial lesion; OR
(c) Confirmation that the patient has received cranial irradiation without having had an intracranial lesion, and has undergone a 12 month period of observation following completion of treatment for the condition for which cranial irradiation was received; **AND**
7. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

Clinical criteria:

- Patient must have a chronological age of less than 2 years, **AND**
- Patient must have a documented clinical risk of hypoglycaemia, **AND**
- Patient must have documented evidence that the risk of hypoglycaemia is secondary to biochemical growth hormone deficiency, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; **AND**
3. Recent growth data (height and weight, not older than three months); **AND**
4. Confirmation that the patient has a documented clinical risk of hypoglycaemia; **AND**
5. Confirmation that the patient has documented evidence that the risk of hypoglycaemia is secondary to biochemical growth hormone deficiency; **AND**
6. The proprietary name (brand), form and strength of somatotropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Biochemical growth hormone deficiency and precocious puberty

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must be male and have commenced puberty (demonstrated by Tanner stage 2 genital or pubic hair development or testicular volumes greater than or equal to 4 mL) before the chronological age of 9 years; OR
- Patient must be female and have commenced puberty (demonstrated by Tanner stage 2 breast or pubic hair development) before the chronological age of 8 years; OR
- Patient must be female and menarche occurred before the chronological age of 10 years, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must be undergoing Gonadotrophin Releasing Hormone agonist therapy for pubertal suppression, **AND**

- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; **AND**
3. (a) A minimum of 12 months of recent growth data (height and weight) at intervals no greater than six months. The most recent data must not be older than three months; OR
(b) A minimum of 6 months of recent growth data (height and weight) for older children (males chronological age 12 and over or bone age 10 and over, females chronological age 10 and over or bone age 8 and over). The most recent data must not be older than three months; **AND**
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
5. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; **AND**
6. Confirmation that the patient has precocious puberty; **AND**
7. Confirmation that the patient is undergoing Gonadotropin Releasing Hormone agonist therapy, for pubertal suppression; **AND**
8. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have a structural lesion that is not neoplastic; OR
- Patient must have had a structural lesion that was neoplastic and have undergone a 12 month period of observation following completion of treatment for the structural lesion (all treatment); OR
- Patient must have a structural lesion that is neoplastic, have received medical advice that it is unsafe to treat the structural lesion, and have undergone a 12 month period of observation since initial diagnosis of the structural lesion, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must have other hypothalamic/pituitary hormone deficits (includes ACTH, TSH, GnRH and/or vasopressin/ADH deficiencies), **AND**

- Patient must have hypothalamic obesity, **AND**
- Patient must have a growth velocity above the 25th percentile for bone age and sex measured over a 12 month interval (or a 6 month interval for an older child); **OR**
- Patient must have an annual growth velocity of greater than 14 cm per year if the patient has a chronological age of 2 years or less; **OR**
- Patient must have an annual growth velocity of greater than 8 cm per year if the patient has a bone or chronological age of 2.5 years or less, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; **OR**
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; **OR**
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; **AND**
3. A minimum of 12 months of recent growth data (height and weight measurements) or a minimum of 6 months of recent growth data for an older child. The most recent data must not be more than three months old at the time of application; **AND**
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
5. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; **AND**
6. (a) Confirmation that the patient has a structural lesion that is not neoplastic; **OR**
(b) Confirmation that the patient had a structural lesion that was neoplastic and has undergone a 12 month period of observation following completion of treatment for the structural lesion (all treatment); **OR**
(c) Confirmation that the patient has a structural lesion that is neoplastic, has received medical advice that it is unsafe to treat the structural lesion, and has undergone a 12 month period of observation since initial diagnosis of the structural lesion; **AND**
7. Confirmation that the patient has other hypothalamic/pituitary hormone deficits; **AND**
8. Confirmation that the patient has hypothalamic obesity; **AND**
9. The proprietary name (brand), form and strength of somatotropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Testing for biochemical growth hormone deficiency must have been performed at a time when all other pituitary hormone deficits were being adequately replaced.

Authority required

Short stature associated with Turner syndrome

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; **OR**
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

Clinical criteria:

- Patient must have a current height at or below the 95th percentile for age on the Turner syndrome growth curve for girls, **AND**
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as a loss of a whole X chromosome in all cells (45X), and gender of rearing is female; **OR**
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as a loss of a whole X chromosome in some cells (mosaic 46XX/45X), and gender of rearing is female; **OR**
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as genetic loss or rearrangement of an X chromosome (such as isochromosome X, ring-chromosome, or partial deletion of an X chromosome), and gender of rearing is female, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**

- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must not have a bone age of 2.5 years or less, **AND**
- Patient must not have a height greater than or equal to 155.0cm, **AND**
- Patient must not have a bone age of 13.5 years or greater.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; **AND**
3. (a) A minimum of 12 months of recent growth data (height and weight) at intervals no greater than six months. The most recent data must not be older than three months; **OR**
(b) A minimum of 6 months of recent growth data (height and weight) for older children (females chronological age 10 and over or bone age 8 and over). The most recent data must not be older than three months; **AND**
4. A bone age result performed within the last 12 months; **AND**
5. Confirmation that the patient has diagnostic results consistent with Turner syndrome; **AND**
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature due to short stature homeobox (SHOX) gene disorders

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as a karyotype confirming the presence of a SHOX mutation/deletion without the presence of mixed gonadal dysgenesis; **OR**
- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as mixed gonadal dysgenesis (45X mosaic karyotype with the presence of any Y chromosome material and/or SRY gene positive by FISH study) and have an appropriate plan of management in place for the patient's increased risk of gonadoblastoma, **AND**
- Patient must have a current height at or below the 1st percentile for age and sex, **AND**
- Patient must have a growth velocity below the 25th percentile for bone age and sex measured over a 12 month interval (or a 6 month interval for an older child); **OR**
- Patient must have an annual growth velocity of 14 cm per year or less if the patient has a chronological age of 2 years or less; **OR**
- Patient must have an annual growth velocity of 8 cm per year or less if the patient has a bone or chronological age of 2.5 years or less, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes (excluding gonadoblastoma secondary to mixed gonadal dysgenesis), **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; **OR**
- Patient must be female and must not have a height greater than or equal to 155.0cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; **OR**
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; **OR**
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; **AND**
3. A minimum of 12 months of recent growth data (height and weight measurements) or a minimum of 6 months of recent growth data for an older child. The most recent data must not be more than three months old at the time of application; **AND**
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
5. Confirmation that the patient has diagnostic results consistent with a short stature homeobox (SHOX) gene disorder; **AND**
6. If the patient's condition is secondary to mixed gonadal dysgenesis, confirmation that an appropriate plan of management for the patient's increased risk of gonadoblastoma is in place; **AND**

7. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with chronic renal insufficiency

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have an estimated glomerular filtration rate less than 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula, and not have undergone a renal transplant; OR
- Patient must have an estimated glomerular filtration rate less than 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula, have undergone a renal transplant, and have undergone a 12 month period of observation following the transplant, **AND**
- Patient must have a current height at or below the 1st percentile for age and sex; OR
- Patient must have a current height above the 1st and at or below the 25th percentiles for age and sex and a growth velocity less than or equal to the 25th percentile for bone age and sex measured over a 12 month interval (or a 6 month interval for an older child); OR
- Patient must have a current height above the 1st and at or below the 25th percentiles for age and sex and an annual growth velocity of 14 cm per year or less if the patient has a chronological age of 2 years or less; OR
- Patient must have a current height above the 1st and at or below the 25th percentiles for age and sex and an annual growth velocity of 8 cm per year or less if the patient has a bone or chronological age of 2.5 years or less, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; AND
3. (a) A minimum of 12 months of recent growth data (height and weight measurements) or a minimum of 6 months of recent growth data for an older child. The most recent data must not be more than three months old at the time of application; OR
(b) Height and weight measurements, not more than three months old at the time of application, for a patient whose current height is at or below the 1st percentile for age and sex; AND
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
5. Confirmation that the patient has an estimated glomerular filtration rate less than 30mL/minute/1.73m² ; AND
6. If a renal transplant has taken place, confirmation that the patient has undergone a 12 month period of observation following transplantation; AND
7. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

somatropin 5 mg/1.5 mL injection, 1.5 mL cartridge

5818F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	223.61	40.30	Norditropin FlexPro [NO]

somatropin 5 mg/1.5 mL injection, 1.5 mL cartridge

6295H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	223.61	40.30	Norditropin SimpleXx [NO]

somatropin 10 mg/1.5 mL injection, 1.5 mL cartridge

5819G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	439.82	40.30	Norditropin FlexPro [NO]

somatropin 10 mg/1.5 mL injection, 1.5 mL cartridge

6296J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	439.82	40.30	Norditropin SimpleXx [NO]

somatropin 10 mg/2 mL injection, 2 mL cartridge

9604L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	426.41	40.30	NutropinAq [IS]

somatropin 15 mg/1.5 mL injection, 1.5 mL cartridge

5820H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	656.04	40.30	Norditropin FlexPro [NO]

somatropin 15 mg/1.5 mL injection, 1.5 mL cartridge

6297K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	656.04	40.30	Norditropin SimpleXx [NO]

■ SOMATROPIN

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Prior Written Approval of Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Short stature and slow growth

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have a current height at or below the 1st percentile for age and sex, **AND**
- Patient must have a growth velocity below the 25th percentile for bone age and sex measured over a 12 month interval (or a 6 month interval for an older child), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must not have a bone age of 2.5 years or less, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7 cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0 cm, **AND**
- Patient must be male and must not have maturational or constitutional delay in combination with an estimated mature height equal to or above 160.1 cm; OR
- Patient must be female and must not have maturational or constitutional delay in combination with an estimated mature height equal to or above 148.0 cm.

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; **AND**
3. A minimum of 12 months of recent growth data (height and weight measurements) or a minimum of 6 months of recent growth data for an older child. The most recent data must not be more than three months old at the time of application; **AND**
4. A bone age result performed within the last 12 months; **AND**
5. Confirmation of the patient's maturational or constitutional delay status; **AND**

6. If the patient has maturational or constitutional delay, confirmation that the patient has an estimated mature height below the 1st adult height percentile; AND
7. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with biochemical growth hormone deficiency

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must have a current height at or below the 1st percentile for age and sex; OR
- Patient must have a current height above the 1st and at or below the 25th percentiles for age and sex and a growth velocity below the 25th percentile for bone age and sex measured over a 12 month interval (or a 6 month interval for an older child); OR
- Patient must have a current height above the 1st and at or below the 25th percentiles for age and sex and an annual growth velocity of 14 cm per year or less if the patient has a chronological age of 2 years or less; OR
- Patient must have a current height above the 1st and at or below the 25th percentiles for age and sex and an annual growth velocity of 8 cm per year or less if the patient has a bone or chronological age of 2.5 years or less, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; AND
3. (a) A minimum of 12 months of recent growth data (height and weight measurements) or a minimum of 6 months of recent growth data for an older child. The most recent data must not be more than three months old at the time of application; OR
(b) Height and weight measurements, not more than three months old at the time of application, for a patient whose current height is at or below the 1st percentile for age and sex; AND
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND

5. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; AND

6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Biochemical growth hormone deficiency should not be secondary to an intracranial lesion or cranial irradiation for applications under this category.

Authority required

Growth retardation secondary to an intracranial lesion, or cranial irradiation

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have had an intracranial lesion and have undergone a 12 month period of observation following completion of treatment for the intracranial lesion (all treatment); OR
- Patient must have had an intracranial lesion, have received medical advice that it is unsafe to treat the intracranial lesion, and have undergone a 12 month period of observation since initial diagnosis of the intracranial lesion; OR
- Patient must have received cranial irradiation without having had an intracranial lesion, and have undergone a 12 month period of observation following completion of treatment for the condition for which cranial irradiation was received, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must have a current height at or below the 1st percentile for age and sex; OR
- Patient must have a current height above the 1st percentile for age and sex and a growth velocity below the 25th percentile for bone age and sex measured over a 12 month interval (or a 6 month interval for an older child); OR
- Patient must have a current height above the 1st percentile for age and sex and an annual growth velocity of 14 cm per year or less if the patient has a chronological age of 2 years or less; OR
- Patient must have a current height above the 1st percentile for age and sex and an annual growth velocity of 8 cm per year or less if the patient has a bone or chronological age of 2.5 years or less, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; AND

3. (a) A minimum of 12 months of recent growth data (height and weight measurements) or a minimum of 6 months of recent growth data for an older child. The most recent data must not be more than three months old at the time of application; OR
- (b) Height and weight measurements, not more than three months old at the time of application, for a patient whose current height is at or below the 1st percentile for age and sex; AND
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
5. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; AND
6. (a) Confirmation that the patient has had an intracranial lesion and has undergone a 12 month period of observation following completion of treatment for the intracranial lesion (all treatment); OR
- (b) Confirmation that the patient has had an intracranial lesion, has received medical advice that it is unsafe to treat the intracranial lesion, and has undergone a 12 month period of observation since initial diagnosis of the intracranial lesion; OR
- (c) Confirmation that the patient has received cranial irradiation without having had an intracranial lesion, and has undergone a 12 month period of observation following completion of treatment for the condition for which cranial irradiation was received; AND
7. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

Clinical criteria:

- Patient must have a chronological age of less than 2 years, **AND**
- Patient must have a documented clinical risk of hypoglycaemia, **AND**
- Patient must have documented evidence that the risk of hypoglycaemia is secondary to biochemical growth hormone deficiency, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; AND
3. Recent growth data (height and weight, not older than three months); AND
4. Confirmation that the patient has a documented clinical risk of hypoglycaemia; AND
5. Confirmation that the patient has documented evidence that the risk of hypoglycaemia is secondary to biochemical growth hormone deficiency; AND
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Biochemical growth hormone deficiency and precocious puberty

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must be male and have commenced puberty (demonstrated by Tanner stage 2 genital or pubic hair development or testicular volumes greater than or equal to 4 mL) before the chronological age of 9 years; OR
- Patient must be female and have commenced puberty (demonstrated by Tanner stage 2 breast or pubic hair development) before the chronological age of 8 years; OR
- Patient must be female and menarche occurred before the chronological age of 10 years, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone

stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR

- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must be undergoing Gonadotrophin Releasing Hormone agonist therapy for pubertal suppression, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; **AND**
3. (a) A minimum of 12 months of recent growth data (height and weight) at intervals no greater than six months. The most recent data must not be older than three months; OR
- (b) A minimum of 6 months of recent growth data (height and weight) for older children (males chronological age 12 and over or bone age 10 and over, females chronological age 10 and over or bone age 8 and over). The most recent data must not be older than three months; **AND**
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
5. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; **AND**
6. Confirmation that the patient has precocious puberty; **AND**
7. Confirmation that the patient is undergoing Gonadotropin Releasing Hormone agonist therapy, for pubertal suppression; **AND**
8. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have a structural lesion that is not neoplastic; OR
- Patient must have had a structural lesion that was neoplastic and have undergone a 12 month period of observation following completion of treatment for the structural lesion (all treatment); OR
- Patient must have a structural lesion that is neoplastic, have received medical advice that it is unsafe to treat the structural lesion, and have undergone a 12 month period of observation since initial diagnosis of the structural lesion, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone

stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR

- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must have other hypothalamic/pituitary hormone deficits (includes ACTH, TSH, GnRH and/or vasopressin/ADH deficiencies), **AND**
- Patient must have hypothalamic obesity, **AND**
- Patient must have a growth velocity above the 25th percentile for bone age and sex measured over a 12 month interval (or a 6 month interval for an older child); OR
- Patient must have an annual growth velocity of greater than 14 cm per year if the patient has a chronological age of 2 years or less; OR
- Patient must have an annual growth velocity of greater than 8 cm per year if the patient has a bone or chronological age of 2.5 years or less, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; **AND**
3. A minimum of 12 months of recent growth data (height and weight measurements) or a minimum of 6 months of recent growth data for an older child. The most recent data must not be more than three months old at the time of application; **AND**
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
5. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; **AND**
6. (a) Confirmation that the patient has a structural lesion that is not neoplastic; OR
- (b) Confirmation that the patient had a structural lesion that was neoplastic and has undergone a 12 month period of observation following completion of treatment for the structural lesion (all treatment); OR
- (c) Confirmation that the patient has a structural lesion that is neoplastic, has received medical advice that it is unsafe to treat the structural lesion, and has undergone a 12 month period of observation since initial diagnosis of the structural lesion; **AND**
7. Confirmation that the patient has other hypothalamic/pituitary hormone deficits; **AND**
8. Confirmation that the patient has hypothalamic obesity; **AND**
9. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Testing for biochemical growth hormone deficiency must have been performed at a time when all other pituitary hormone deficits were being adequately replaced.

Authority required

Short stature associated with Turner syndrome

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

Clinical criteria:

- Patient must have a current height at or below the 95th percentile for age on the Turner syndrome growth curve for girls, **AND**
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as a loss of a whole X chromosome in all cells (45X), and gender of rearing is female; OR
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as a loss of a whole X chromosome in some cells (mosaic 46XX/45X), and gender of rearing is female; OR
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as genetic loss or rearrangement of an X chromosome (such as isochromosome X, ring-chromosome, or partial deletion of an X chromosome), and gender of rearing is female, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must not have a bone age of 2.5 years or less, **AND**
- Patient must not have a height greater than or equal to 155.0cm, **AND**
- Patient must not have a bone age of 13.5 years or greater.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; **AND**
3. (a) A minimum of 12 months of recent growth data (height and weight) at intervals no greater than six months. The most recent data must not be older than three months; **OR**
(b) A minimum of 6 months of recent growth data (height and weight) for older children (females chronological age 10 and over or bone age 8 and over). The most recent data must not be older than three months; **AND**
4. A bone age result performed within the last 12 months; **AND**
5. Confirmation that the patient has diagnostic results consistent with Turner syndrome; **AND**
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature due to short stature homeobox (SHOX) gene disorders

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as a karyotype confirming the presence of a SHOX mutation/deletion without the presence of mixed gonadal dysgenesis; OR
- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as mixed gonadal dysgenesis (45X mosaic karyotype with the presence of any Y chromosome material and/or SRY gene positive by FISH study) and have an appropriate plan of management in place for the patient's increased risk of gonadoblastoma, **AND**
- Patient must have a current height at or below the 1st percentile for age and sex, **AND**
- Patient must have a growth velocity below the 25th percentile for bone age and sex measured over a 12 month interval (or a 6 month interval for an older child); **OR**
- Patient must have an annual growth velocity of 14 cm per year or less if the patient has a chronological age of 2 years or less; **OR**
- Patient must have an annual growth velocity of 8 cm per year or less if the patient has a bone or chronological age of 2.5 years or less, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes (excluding gonadoblastoma secondary to mixed gonadal dysgenesis), **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; **OR**
- Patient must be female and must not have a height greater than or equal to 155.0cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; **OR**
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; AND
3. A minimum of 12 months of recent growth data (height and weight measurements) or a minimum of 6 months of recent growth data for an older child. The most recent data must not be more than three months old at the time of application; AND
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
5. Confirmation that the patient has diagnostic results consistent with a short stature homeobox (SHOX) gene disorder; AND
6. If the patient's condition is secondary to mixed gonadal dysgenesis, confirmation that an appropriate plan of management for the patient's increased risk of gonadoblastoma is in place; AND
7. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with chronic renal insufficiency

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

Clinical criteria:

- Patient must have an estimated glomerular filtration rate less than 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula, and not have undergone a renal transplant; OR
- Patient must have an estimated glomerular filtration rate less than 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula, have undergone a renal transplant, and have undergone a 12 month period of observation following the transplant, **AND**
- Patient must have a current height at or below the 1st percentile for age and sex; OR
- Patient must have a current height above the 1st and at or below the 25th percentiles for age and sex and a growth velocity less than or equal to the 25th percentile for bone age and sex measured over a 12 month interval (or a 6 month interval for an older child); OR
- Patient must have a current height above the 1st and at or below the 25th percentiles for age and sex and an annual growth velocity of 14 cm per year or less if the patient has a chronological age of 2 years or less; OR
- Patient must have a current height above the 1st and at or below the 25th percentiles for age and sex and an annual growth velocity of 8 cm per year or less if the patient has a bone or chronological age of 2.5 years or less, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7 cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0 cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Population criteria:

- Patient must be prepubertal.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; AND
3. (a) A minimum of 12 months of recent growth data (height and weight measurements) or a minimum of 6 months of recent growth data for an older child. The most recent data must not be more than three months old at the time of application; OR

- (b) Height and weight measurements, not more than three months old at the time of application, for a patient whose current height is at or below the 1st percentile for age and sex; AND
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
5. Confirmation that the patient has an estimated glomerular filtration rate less than 30mL/minute/1.73m² ; AND
6. If a renal transplant has taken place, confirmation that the patient has undergone a 12 month period of observation following transplantation; AND
7. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).
- Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

somatropin 24 mg injection [1 cartridge] (&) inert substance diluent [3.15 mL syringe], 1 pack

6345Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	1045.23	40.30	Humatrope [LY]

somatropin 6 mg injection [1 cartridge] (&) inert substance diluent [3.15 mL syringe], 1 pack

6169Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	266.85	40.30	Humatrope [LY]

somatropin 12 mg injection [1 cartridge] (&) inert substance diluent [3.15 mL syringe], 1 pack

6170R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	526.31	40.30	Humatrope [LY]

■ SOMATROPIN

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au Applications for authority to prescribe should be forwarded to:
Department of Human Services
Prior Written Approval of Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Short stature and slow growth

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have a current height at or below the 1st percentile for age and sex, **AND**
- Patient must have a growth velocity below the 25th percentile for bone age and sex measured over a 12 month interval (or a 6 month interval for an older child), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must not have a bone age of 2.5 years or less, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7 cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0 cm, **AND**
- Patient must be male and must not have maturational or constitutional delay in combination with an estimated mature height equal to or above 160.1 cm; OR
- Patient must be female and must not have maturational or constitutional delay in combination with an estimated mature height equal to or above 148.0 cm.

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; AND

3. A minimum of 12 months of recent growth data (height and weight measurements) or a minimum of 6 months of recent growth data for an older child. The most recent data must not be more than three months old at the time of application; AND
4. A bone age result performed within the last 12 months; AND
5. Confirmation of the patient's maturational or constitutional delay status; AND
6. If the patient has maturational or constitutional delay, confirmation that the patient has an estimated mature height below the 1st adult height percentile; AND
7. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with biochemical growth hormone deficiency

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; **AND**
- Patient must have a current height at or below the 1st percentile for age and sex; OR
- Patient must have a current height above the 1st and at or below the 25th percentiles for age and sex and a growth velocity below the 25th percentile for bone age and sex measured over a 12 month interval (or a 6 month interval for an older child); OR
- Patient must have a current height above the 1st and at or below the 25th percentiles for age and sex and an annual growth velocity of 14 cm per year or less if the patient has a chronological age of 2 years or less; OR
- Patient must have a current height above the 1st and at or below the 25th percentiles for age and sex and an annual growth velocity of 8 cm per year or less if the patient has a bone or chronological age of 2.5 years or less; **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; AND
3. (a) A minimum of 12 months of recent growth data (height and weight measurements) or a minimum of 6 months of recent growth data for an older child. The most recent data must not be more than three months old at the time of application; OR

(b) Height and weight measurements, not more than three months old at the time of application, for a patient whose current height is at or below the 1st percentile for age and sex; AND

4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND

5. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; AND

6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Biochemical growth hormone deficiency should not be secondary to an intracranial lesion or cranial irradiation for applications under this category.

Authority required

Growth retardation secondary to an intracranial lesion, or cranial irradiation

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have had an intracranial lesion and have undergone a 12 month period of observation following completion of treatment for the intracranial lesion (all treatment); OR
- Patient must have had an intracranial lesion, have received medical advice that it is unsafe to treat the intracranial lesion, and have undergone a 12 month period of observation since initial diagnosis of the intracranial lesion; OR
- Patient must have received cranial irradiation without having had an intracranial lesion, and have undergone a 12 month period of observation following completion of treatment for the condition for which cranial irradiation was received, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must have a current height at or below the 1st percentile for age and sex; OR
- Patient must have a current height above the 1st percentile for age and sex and a growth velocity below the 25th percentile for bone age and sex measured over a 12 month interval (or a 6 month interval for an older child); OR
- Patient must have a current height above the 1st percentile for age and sex and an annual growth velocity of 14 cm per year or less if the patient has a chronological age of 2 years or less; OR
- Patient must have a current height above the 1st percentile for age and sex and an annual growth velocity of 8 cm per year or less if the patient has a bone or chronological age of 2.5 years or less, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; AND
3. (a) A minimum of 12 months of recent growth data (height and weight measurements) or a minimum of 6 months of recent growth data for an older child. The most recent data must not be more than three months old at the time of application; OR
(b) Height and weight measurements, not more than three months old at the time of application, for a patient whose current height is at or below the 1st percentile for age and sex; AND
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
5. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; AND
6. (a) Confirmation that the patient has had an intracranial lesion and has undergone a 12 month period of observation following completion of treatment for the intracranial lesion (all treatment); OR
(b) Confirmation that the patient has had an intracranial lesion, has received medical advice that it is unsafe to treat the intracranial lesion, and has undergone a 12 month period of observation since initial diagnosis of the intracranial lesion; OR
(c) Confirmation that the patient has received cranial irradiation without having had an intracranial lesion, and has undergone a 12 month period of observation following completion of treatment for the condition for which cranial irradiation was received; AND
7. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

Clinical criteria:

- Patient must have a chronological age of less than 2 years, **AND**
- Patient must have a documented clinical risk of hypoglycaemia, **AND**
- Patient must have documented evidence that the risk of hypoglycaemia is secondary to biochemical growth hormone deficiency, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; AND
3. Recent growth data (height and weight, not older than three months); AND
4. Confirmation that the patient has a documented clinical risk of hypoglycaemia; AND
5. Confirmation that the patient has documented evidence that the risk of hypoglycaemia is secondary to biochemical growth hormone deficiency; AND
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Biochemical growth hormone deficiency and precocious puberty

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must be male and have commenced puberty (demonstrated by Tanner stage 2 genital or pubic hair development or testicular volumes greater than or equal to 4 mL) before the chronological age of 9 years; OR
- Patient must be female and have commenced puberty (demonstrated by Tanner stage 2 breast or pubic hair development) before the chronological age of 8 years; OR
- Patient must be female and menarche occurred before the chronological age of 10 years, **AND**

- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must be undergoing Gonadotrophin Releasing Hormone agonist therapy for pubertal suppression, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; **AND**
3. (a) A minimum of 12 months of recent growth data (height and weight) at intervals no greater than six months. The most recent data must not be older than three months; OR
(b) A minimum of 6 months of recent growth data (height and weight) for older children (males chronological age 12 and over or bone age 10 and over, females chronological age 10 and over or bone age 8 and over). The most recent data must not be older than three months; **AND**
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
5. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; **AND**
6. Confirmation that the patient has precocious puberty; **AND**
7. Confirmation that the patient is undergoing Gonadotropin Releasing Hormone agonist therapy, for pubertal suppression; **AND**
8. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have a structural lesion that is not neoplastic; OR
- Patient must have had a structural lesion that was neoplastic and have undergone a 12 month period of observation following completion of treatment for the structural lesion (all treatment); OR
- Patient must have a structural lesion that is neoplastic, have received medical advice that it is unsafe to treat the structural lesion, and have undergone a 12 month period of observation since initial diagnosis of the structural lesion, **AND**

- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must have other hypothalamic/pituitary hormone deficits (includes ACTH, TSH, GnRH and/or vasopressin/ADH deficiencies), **AND**
- Patient must have hypothalamic obesity, **AND**
- Patient must have a growth velocity above the 25th percentile for bone age and sex measured over a 12 month interval (or a 6 month interval for an older child); OR
- Patient must have an annual growth velocity of greater than 14 cm per year if the patient has a chronological age of 2 years or less; OR
- Patient must have an annual growth velocity of greater than 8 cm per year if the patient has a bone or chronological age of 2.5 years or less, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; **AND**
3. A minimum of 12 months of recent growth data (height and weight measurements) or a minimum of 6 months of recent growth data for an older child. The most recent data must not be more than three months old at the time of application; **AND**
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
5. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; **AND**
6. (a) Confirmation that the patient has a structural lesion that is not neoplastic; OR
(b) Confirmation that the patient had a structural lesion that was neoplastic and has undergone a 12 month period of observation following completion of treatment for the structural lesion (all treatment); OR
(c) Confirmation that the patient has a structural lesion that is neoplastic, has received medical advice that it is unsafe to treat the structural lesion, and has undergone a 12 month period of observation since initial diagnosis of the structural lesion; **AND**
7. Confirmation that the patient has other hypothalamic/pituitary hormone deficits; **AND**
8. Confirmation that the patient has hypothalamic obesity; **AND**
9. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Testing for biochemical growth hormone deficiency must have been performed at a time when all other pituitary hormone deficits were being adequately replaced.

Authority required

Short stature associated with Turner syndrome

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

Clinical criteria:

- Patient must have a current height at or below the 95th percentile for age on the Turner syndrome growth curve for girls, **AND**
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as a loss of a whole X chromosome in all cells (45X), and gender of rearing is female; OR
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as a loss of a whole X chromosome in some cells (mosaic 46XX/45X), and gender of rearing is female; OR
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as genetic loss or rearrangement of an X chromosome (such as isochromosome X, ring-chromosome, or partial deletion of an X chromosome), and gender of rearing is female, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must not have a bone age of 2.5 years or less, **AND**
- Patient must not have a height greater than or equal to 155.0cm, **AND**
- Patient must not have a bone age of 13.5 years or greater.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; **AND**
3. (a) A minimum of 12 months of recent growth data (height and weight) at intervals no greater than six months. The most recent data must not be older than three months; OR
- (b) A minimum of 6 months of recent growth data (height and weight) for older children (females chronological age 10 and over or bone age 8 and over). The most recent data must not be older than three months; **AND**
4. A bone age result performed within the last 12 months; **AND**
5. Confirmation that the patient has diagnostic results consistent with Turner syndrome; **AND**
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature due to short stature homeobox (SHOX) gene disorders

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as a karyotype confirming the presence of a SHOX mutation/deletion without the presence of mixed gonadal dysgenesis; OR
- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as mixed gonadal dysgenesis (45X mosaic karyotype with the presence of any Y chromosome material and/or SRY gene positive by FISH study) and have an appropriate plan of management in place for the patient's increased risk of gonadoblastoma, **AND**
- Patient must have a current height at or below the 1st percentile for age and sex, **AND**
- Patient must have a growth velocity below the 25th percentile for bone age and sex measured over a 12 month interval (or a 6 month interval for an older child); OR
- Patient must have an annual growth velocity of 14 cm per year or less if the patient has a chronological age of 2 years or less; OR
- Patient must have an annual growth velocity of 8 cm per year or less if the patient has a bone or chronological age of 2.5 years or less, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes (excluding gonadoblastoma secondary to mixed gonadal dysgenesis), **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**

- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; **AND**
3. A minimum of 12 months of recent growth data (height and weight measurements) or a minimum of 6 months of recent growth data for an older child. The most recent data must not be more than three months old at the time of application; **AND**
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
5. Confirmation that the patient has diagnostic results consistent with a short stature homeobox (SHOX) gene disorder; **AND**
6. If the patient's condition is secondary to mixed gonadal dysgenesis, confirmation that an appropriate plan of management for the patient's increased risk of gonadoblastoma is in place; **AND**
7. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with chronic renal insufficiency

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have an estimated glomerular filtration rate less than 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula, and not have undergone a renal transplant; OR
- Patient must have an estimated glomerular filtration rate less than 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula, have undergone a renal transplant, and have undergone a 12 month period of observation following the transplant, **AND**
- Patient must have a current height at or below the 1st percentile for age and sex; OR
- Patient must have a current height above the 1st and at or below the 25th percentiles for age and sex and a growth velocity less than or equal to the 25th percentile for bone age and sex measured over a 12 month interval (or a 6 month interval for an older child); OR
- Patient must have a current height above the 1st and at or below the 25th percentiles for age and sex and an annual growth velocity of 14 cm per year or less if the patient has a chronological age of 2 years or less; OR
- Patient must have a current height above the 1st and at or below the 25th percentiles for age and sex and an annual growth velocity of 8 cm per year or less if the patient has a bone or chronological age of 2.5 years or less, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**

2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; AND
 3. (a) A minimum of 12 months of recent growth data (height and weight measurements) or a minimum of 6 months of recent growth data for an older child. The most recent data must not be more than three months old at the time of application; OR
 - (b) Height and weight measurements, not more than three months old at the time of application, for a patient whose current height is at or below the 1st percentile for age and sex; AND
 4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
 5. Confirmation that the patient has an estimated glomerular filtration rate less than 30mL/minute/1.73m² ; AND
 6. If a renal transplant has taken place, confirmation that the patient has undergone a 12 month period of observation following transplantation; AND
 7. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).
- Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature and poor body composition due to Prader-Willi syndrome

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have diagnostic results consistent with Prader-Willi syndrome (the condition must be genetically proven); OR
- Patient must have a clinical diagnosis of Prader-Willi syndrome, confirmed by a clinical geneticist, **AND**
- Patient must have been evaluated via polysomnography for airway obstruction and apnoea within the last 12 months with no sleep disorders identified; OR
- Patient must have been evaluated via polysomnography for airway obstruction and apnoea within the last 12 months with sleep disorders identified which are not of sufficient severity to require treatment; OR
- Patient must have been evaluated via polysomnography for airway obstruction and apnoea within the last 12 months with sleep disorders identified for which the patient is currently receiving ameliorative treatment, **AND**
- Patient must not have uncontrolled morbid obesity, defined as a body weight greater than 200% of ideal body weight for height and sex, with ideal body weight derived by calculating the 50th percentile weight for the patient's current height,

AND

- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have previously received treatment under the PBS S100 Growth Hormone Program, **AND**
- Patient must not have a chronological age of 18 years or greater.

Treatment criteria:

- Must be treated by a specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a specialist or consultant physician in general paediatrics in consultation with a nominated specialist or consultant physician in paediatric endocrinology.

The maximum duration of the initial treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for initial treatment; AND
3. A minimum of 6 months of recent growth data (height, weight and waist circumference). The most recent data must not be older than three months; AND
4. The date at which skeletal maturity was achieved (if applicable) [Note: In patients whose chronological age is greater than 2.5 years, a bone age reading should be performed at least once every 12 months prior to attainment of skeletal maturity]; AND
5. (a) Confirmation that the patient has diagnostic results consistent with Prader-Willi syndrome; OR
- (b) Confirmation that the patient has a clinical diagnosis of Prader-Willi syndrome, confirmed by a clinical geneticist
6. Confirmation that the patient has been evaluated via polysomnography for airway obstruction and apnoea within the last 12 months and any sleep disorders identified via polysomnography that required treatment have been addressed; AND
7. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with 1 repeat allowed)

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

somatropin 2 mg injection [7] (&) inert substance diluent [7 x 0.25 mL syringes], 1 pack

6319N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	612.79	40.30	Genotropin MiniQuick [PF]

somatropin 1.6 mg injection [7] (&) inert substance diluent [7 x 0.25 mL syringes], 1 pack

6317L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	491.72	40.30	Genotropin MiniQuick [PF]

somatropin 1.4 mg injection [7] (&) inert substance diluent [7 x 0.25 mL syringes], 1 pack

6316K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	431.17	40.30	Genotropin MiniQuick [PF]

somatropin 1.8 mg injection [7] (&) inert substance diluent [7 x 0.25 mL syringes], 1 pack

6318M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	552.26	40.30	Genotropin MiniQuick [PF]

SOMATROPIN (Recombinant human growth hormone) Powder for injection 12 mg (36 i.u.) with diluent in pre-filled pen (with preservative), 1

9586M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	510.20	40.30	Genotropin GoQuick [PF]

SOMATROPIN (Recombinant human growth hormone) Powder for injection 5 mg (15 i.u.) with diluent in pre-filled pen (with preservative), 1

9585L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	216.90	40.30	Genotropin GoQuick [PF]

somatropin 800 microgram injection [7] (&) inert substance diluent [7 x 0.25 mL syringes], 1 pack

6313G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	249.55	40.30	Genotropin MiniQuick [PF]

somatropin 1.2 mg injection [7] (&) inert substance diluent [7 x 0.25 mL syringes], 1 pack

6315J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	370.63	40.30	Genotropin MiniQuick [PF]

somatropin 600 microgram injection [7] (&) inert substance diluent [7 x 0.25 mL syringes], 1 pack

9628R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	189.02	40.30	Genotropin MiniQuick [PF]

somatropin 400 microgram injection [7] (&) inert substance diluent [7 x 0.25 mL syringes], 1 pack

10902T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	128.47	40.30	Genotropin MiniQuick [PF]

somatropin 1 mg injection [7] (&) inert substance diluent [7 x 0.25 mL syringes], 1 pack

6314H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	310.09	40.30	Genotropin MiniQuick [PF]

■ SOMATROPIN

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Prior Written Approval of Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Short stature and slow growth

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature and slow growth category, **AND**
- Patient must not have been on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR

- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved and maintained mid parental height standard deviation score while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; AND
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
4. A bone age result performed within the last 12 months; AND
5. The final adult height (in cm) of the patient's mother and father (where available); AND
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with biochemical growth hormone deficiency

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature associated with biochemical growth hormone deficiency category, **AND**
- Patient must not have been on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved and maintained mid parental height standard deviation score while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; AND
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
5. The final adult height (in cm) of the patient's mother and father (where available); AND

6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Growth retardation secondary to an intracranial lesion, or cranial irradiation

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the growth retardation secondary to an intracranial lesion, or cranial irradiation category, **AND**
- Patient must not have been on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved and maintained mid parental height standard deviation score while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; **AND**
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; **AND**
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
5. The final adult height (in cm) of the patient's mother and father (where available); **AND**
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants category, **AND**
- Patient must not have been on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR

- Patient must have achieved and maintained mid parental height standard deviation score while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have a chronological age of 5 years or greater.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; **AND**
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; **AND**
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
5. The final adult height (in cm) of the patient's mother and father (where available); **AND**
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

When a patient receiving treatment under the indication risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants reaches or surpasses 5 years of age (chronological), prescribers should seek reclassification to the indication 'short stature due to biochemical growth hormone deficiency'.

Authority required

Biochemical growth hormone deficiency and precocious puberty

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the biochemical growth hormone deficiency and precocious puberty category, **AND**
- Patient must not have been on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); **OR**
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); **OR**
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); **OR**
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); **OR**
- Patient must have achieved and maintained mid parental height standard deviation score while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; **OR**
- Patient must be female and must not have a bone age of 13.5 years or more.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; **AND**
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; **AND**
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
5. The final adult height (in cm) of the patient's mother and father (where available); **AND**
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth category, **AND**
- Patient must not have been on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved and maintained mid parental height standard deviation score while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; **AND**
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; **AND**
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
5. The final adult height (in cm) of the patient's mother and father (where available); **AND**
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with Turner syndrome

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature associated with Turner syndrome category, **AND**
- Patient must not have been on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an annualised growth velocity for bone age at or above the mean growth velocity for untreated Turner Syndrome girls (using the Turner Syndrome - Ranke growth velocity chart) while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**

- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have a bone age of 13.5 years or greater, **AND**
- Patient must not have a height greater than or equal to 155.0 cm.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; **AND**
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; **AND**
4. A bone age result performed within the last 12 months; **AND**
5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature due to short stature homeobox (SHOX) gene disorders

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature due to short stature homeobox (SHOX) gene disorders category, **AND**
- Patient must not have been on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); **OR**
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); **OR**
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); **OR**
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); **OR**
- Patient must have achieved and maintained mid parental height standard deviation score while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes (excluding gonadoblastoma secondary to mixed gonadal dysgenesis), **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7 cm; **OR**
- Patient must be female and must not have a height greater than or equal to 155.0 cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; **OR**
- Patient must be female and must not have a bone age of 13.5 years or more.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; **AND**
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; **AND**
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
5. The final adult height (in cm) of the patient's mother and father (where available); **AND**
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature and slow growth

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than short stature and slow growth, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have previously received treatment under the indication short stature associated with chronic renal insufficiency, have undergone a renal transplant and a 12 month period of observation following the transplant, and have an estimated glomerular filtration rate of greater than or equal to 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula; OR
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment and a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; **AND**
3. (a) A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment; OR
- (b) Confirmation that the patient has previously received treatment under the indication **short stature associated with chronic renal insufficiency**, has undergone a renal transplant and a 12 month period of observation following the transplant, and has an estimated glomerular filtration rate of greater than or equal to 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula; **AND**
4. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; **AND**
5. A bone age result performed within the last 12 months; **AND**
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with biochemical growth hormone deficiency

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than short stature associated with biochemical growth hormone deficiency, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have previously received treatment under the indication **risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants** and have reached or surpassed 5 years of age (chronological); OR
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment; OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and an annual growth velocity of 14 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and an annual growth velocity of 8 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; AND
3. (a) A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); OR
(b) Height and weight measurements from within three months prior to commencement of treatment for a patient whose height was at or below the 1st percentile for age and sex immediately prior to commencing treatment; OR
(c) Confirmation that the patient has previously received treatment under the indication **risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants** and has reached or surpassed 5 years of age (chronological); AND
4. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; AND
5. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
6. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
7. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Biochemical growth hormone deficiency should not be secondary to an intracranial lesion or cranial irradiation for applications under this category.

Authority required

Growth retardation secondary to an intracranial lesion, or cranial irradiation

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than growth retardation secondary to an intracranial lesion, or cranial irradiation, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have had an intracranial lesion and have undergone a 12 month period of observation following completion of treatment for the intracranial lesion (all treatment); OR
- Patient must have had an intracranial lesion, have received medical advice that it is unsafe to treat the intracranial lesion, and have undergone a 12 month period of observation since initial diagnosis of the intracranial lesion; OR
- Patient must have received cranial irradiation without having had an intracranial lesion, and have undergone a 12 month period of observation following completion of treatment for the condition for which cranial irradiation was received, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR

- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment; OR
- Patient must have had both a height above the 1st percentile for age and sex immediately prior to commencing treatment and a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had both a height above the 1st percentile for age and sex immediately prior to commencing treatment and an annual growth velocity of 14 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had both a height above the 1st percentile for age and sex immediately prior to commencing treatment and an annual growth velocity of 8 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; AND
3. (a) A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); OR
(b) Height and weight measurements from within three months prior to commencement of treatment for a patient whose height was at or below the 1st percentile for age and sex immediately prior to commencing treatment; AND
4. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; AND
5. (a) Confirmation that the patient has had an intracranial lesion and has undergone a 12 month period of observation following completion of treatment for the intracranial lesion (all treatment); OR
(b) Confirmation that the patient has had an intracranial lesion, has received medical advice that it is unsafe to treat the intracranial lesion, and has undergone a 12 month period of observation since initial diagnosis of the intracranial lesion; OR
(c) Confirmation that the patient has received cranial irradiation without having had an intracranial lesion, and has undergone a 12 month period of observation following completion of treatment for the condition for which cranial irradiation was received; AND
6. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
7. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND

8. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

Authority required

Risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under a category other than risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have a chronological age of less than 2 years, **AND**
- Patient must have a documented clinical risk of hypoglycaemia, **AND**
- Patient must have documented evidence that the risk of hypoglycaemia is secondary to biochemical growth hormone deficiency, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; **AND**
3. Confirmation that the patient has a documented clinical risk of hypoglycaemia; **AND**
4. Confirmation that the patient has documented evidence that the risk of hypoglycaemia is secondary to biochemical growth hormone deficiency; **AND**
5. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; **AND**
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Biochemical growth hormone deficiency and precocious puberty

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under a category other than biochemical growth hormone deficiency and precocious puberty, **AND**

- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must be male and have commenced puberty (demonstrated by Tanner stage 2 genital or pubic hair development or testicular volumes greater than or equal to 4 mL) before the chronological age of 9 years; OR
- Patient must be female and have commenced puberty (demonstrated by Tanner stage 2 breast or pubic hair development) before the chronological age of 8 years; OR
- Patient must be female and menarche occurred before the chronological age of 10 years, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must be undergoing Gonadotrophin Releasing Hormone agonist therapy for pubertal suppression, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; **AND**
3. Confirmation that the patient has precocious puberty; **AND**
4. Confirmation that the patient is undergoing Gonadotrophin Releasing Hormone agonist therapy for pubertal suppression; **AND**
5. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; **AND**

6. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
7. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND

8. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have a structural lesion that is not neoplastic; OR
- Patient must have had a structural lesion that was neoplastic and have undergone a 12 month period of observation following completion of treatment for the structural lesion (all treatment); OR
- Patient must have a structural lesion that is neoplastic, have received medical advice that it is unsafe to treat the structural lesion, and have undergone a 12 month period of observation since initial diagnosis of the structural lesion,

AND

- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must have other hypothalamic/pituitary hormone deficits (includes ACTH, TSH, GnRH and/or vasopressin/ADH deficiencies), **AND**
- Patient must have hypothalamic obesity, **AND**
- Patient must have had a growth velocity above the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR

- Patient must have had an annual growth velocity of greater than 14 cm per year in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had an annual growth velocity of greater than 8 cm per year in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; **AND**
3. A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); **AND**
4. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; **AND**
5. (a) Confirmation that the patient has a structural lesion that is not neoplastic; OR
(b) Confirmation that the patient had a structural lesion that was neoplastic and has undergone a 12 month period of observation following completion of treatment for the structural lesion (all treatment); OR
(c) Confirmation that the patient has a structural lesion that is neoplastic, has received medical advice that it is unsafe to treat the structural lesion, and has undergone a 12 month period of observation since initial diagnosis of the structural lesion; **AND**
6. Confirmation that the patient has other hypothalamic/pituitary hormone deficits; **AND**
7. Confirmation that the patient has hypothalamic obesity; **AND**
8. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; **AND**
9. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
10. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with Turner syndrome

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under a category other than short stature associated with Turner syndrome, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies).

continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR

- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have had a height at or below the 95th percentile for age on the Turner syndrome growth curve for girls immediately prior to commencing growth hormone treatment, **AND**
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as a loss of a whole X chromosome in all cells (45X), and gender of rearing is female; OR
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as a loss of a whole X chromosome in some cells (mosaic 46XX/45X), and gender of rearing is female; OR
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as genetic loss or rearrangement of an X chromosome (such as isochromosome X, ring-chromosome, or partial deletion of an X chromosome), and gender of rearing is female, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have a bone age of 2.5 years or less, **AND**
- Patient must not have a bone age of 13.5 years or greater, **AND**
- Patient must not have a height greater than or equal to 155.0 cm.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; **AND**
3. A height measurement from immediately prior to commencement of growth hormone treatment; **AND**
4. Confirmation that the patient has diagnostic results consistent with Turner syndrome; **AND**
5. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; **AND**
6. A bone age result performed within the last 12 months; **AND**
7. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature due to short stature homeobox (SHOX) gene disorders

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than short stature due to short stature homeobox (SHOX) gene disorders, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**

- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as a karyotype confirming the presence of a SHOX mutation/deletion without the presence of mixed gonadal dysgenesis; OR
- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as mixed gonadal dysgenesis (45X mosaic karyotype with the presence of any Y chromosome material and/or SRY gene positive by FISH study) and have an appropriate plan of management in place for the patient's increased risk of gonadoblastoma, **AND**
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment, **AND**
- Patient must have had a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had an annual growth velocity of 14 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had an annual growth velocity of 8 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes (excluding gonadoblastoma secondary to mixed gonadal dysgenesis), **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; AND
3. A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); AND
4. Confirmation that the patient has diagnostic results consistent with a short stature homeobox (SHOX) gene disorder; AND
5. If the patient's condition is secondary to mixed gonadal dysgenesis, confirmation that an appropriate plan of management for the patient's increased risk of gonadoblastoma is in place; AND
6. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
7. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
8. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

somatropin 4 mg injection [1 vial] (& inert substance diluent [1 vial], 1 pack

10452D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	180.36	40.30	Zomacton [FP]

▪ SOMATROPIN

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Prior Written Approval of Complex Drugs

Reply Paid 9826
HOBART TAS 7001

Authority required

Short stature and slow growth

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature and slow growth category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program)**

Special Arrangement 2015 and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; **AND**
3. Recent growth data (height and weight, not older than three months); **AND**
4. A bone age result performed within the last 12 months; **AND**
5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Short stature associated with biochemical growth hormone deficiency

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature associated with biochemical growth hormone deficiency category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR

- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; **AND**
3. Recent growth data (height and weight, not older than three months); **AND**
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Growth retardation secondary to an intracranial lesion, or cranial irradiation

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the growth retardation secondary to an intracranial lesion, or cranial irradiation category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must not have diabetes mellitus, **AND**

- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; AND
3. Recent growth data (height and weight, not older than three months); AND
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have a chronological age of 5 years or greater.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; AND

3. Recent growth data (height and weight, not older than three months); AND
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Biochemical growth hormone deficiency and precocious puberty

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the biochemical growth hormone deficiency and precocious puberty category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must be undergoing Gonadotrophin Releasing Hormone agonist therapy for pubertal suppression, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program)**

Special Arrangement 2015 and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; AND
3. Recent growth data (height and weight, not older than three months); AND
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; **AND**
3. Recent growth data (height and weight, not older than three months); **AND**
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Short stature associated with Turner syndrome

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature associated with Turner syndrome category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR

- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be female and must not have a bone age of 13.5 years or more, **AND**
- Patient must be female and must not have a height greater than or equal to 155.0cm.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; AND
3. Recent growth data (height and weight, not older than three months); AND
4. A bone age result performed within the last 12 months; AND
5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Short stature due to short stature homeobox (SHOX) gene disorders

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature due to short stature homeobox (SHOX) gene disorders category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as a karyotype confirming the presence of a SHOX mutation/deletion without the presence of mixed gonadal dysgenesis; OR
- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as mixed gonadal dysgenesis (45X mosaic karyotype with the presence of any Y chromosome material and/or SRY gene positive by FISH study) and have an appropriate plan of management in place for the patient's increased risk of gonadoblastoma, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes (excluding gonadoblastoma secondary to mixed gonadal dysgenesis), **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more, **AND**

- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program)**

Special Arrangement 2015 and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; AND
3. Recent growth data (height and weight, not older than three months); AND
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
5. The proprietary name (brand), form and strength of somatotropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Short stature and slow growth

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than short stature and slow growth, **AND**
- Patient must have had a lapse in treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have previously received treatment under the indication short stature associated with chronic renal insufficiency, have undergone a renal transplant and a 12 month period of observation following the transplant, and have an estimated glomerular filtration rate of greater than or equal to 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula; OR
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment and a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7 cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0 cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program)**

Special Arrangement 2015 and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; AND
3. (a) A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment; OR
(b) Confirmation that the patient has previously received treatment under the indication short stature associated with chronic renal insufficiency, has undergone a renal transplant and a 12 month period of observation following the transplant, and has an estimated glomerular filtration rate of greater than or equal to 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula; AND
4. Recent growth data (height and weight, not older than three months); AND
5. A bone age result performed within the last 12 months; AND
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with biochemical growth hormone deficiency

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than short stature associated with biochemical growth hormone deficiency, **AND**
- Patient must have had a lapse in treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have previously received treatment under the indication **risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants** and have reached or surpassed 5 years of age (chronological); OR
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment; OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and an annual growth velocity of 14 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and an annual growth velocity of 8 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**

- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; **AND**
3. (a) A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); OR
(b) Height and weight measurements from within three months prior to commencement of treatment for a patient whose height was at or below the 1st percentile for age and sex immediately prior to commencing treatment; OR
(c) Confirmation that the patient has previously received treatment under the indication risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants and has reached or surpassed 5 years of age (chronological); **AND**
4. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; **AND**
5. Recent growth data (height and weight, not older than three months); **AND**
6. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
7. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Biochemical growth hormone deficiency should not be secondary to an intracranial lesion or cranial irradiation for applications under this category.

Authority required

Growth retardation secondary to an intracranial lesion, or cranial irradiation

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than growth retardation secondary to an intracranial lesion, or cranial irradiation, **AND**
- Patient must have had a lapse in treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have had an intracranial lesion and have undergone a 12 month period of observation following completion of treatment for the intracranial lesion (all treatment); OR
- Patient must have had an intracranial lesion, have received medical advice that it is unsafe to treat the intracranial lesion, and have undergone a 12 month period of observation since initial diagnosis of the intracranial lesion; OR
- Patient must have received cranial irradiation without having had an intracranial lesion, and have undergone a 12 month period of observation following completion of treatment for the condition for which cranial irradiation was received, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment; OR
- Patient must have had both a height above the 1st percentile for age and sex immediately prior to commencing treatment and a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had both a height above the 1st percentile for age and sex immediately prior to commencing treatment and an annual growth velocity of 14 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had both a height above the 1st percentile for age and sex immediately prior to commencing treatment and an annual growth velocity of 8 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; AND
3. (a) A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); OR
(b) Height and weight measurements from within three months prior to commencement of treatment for a patient whose height was at or below the 1st percentile for age and sex immediately prior to commencing treatment; AND
4. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; AND
5. (a) Confirmation that the patient has had an intracranial lesion and has undergone a 12 month period of observation following completion of treatment for the intracranial lesion (all treatment); OR
(b) Confirmation that the patient has had an intracranial lesion, has received medical advice that it is unsafe to treat the intracranial lesion, and has undergone a 12 month period of observation since initial diagnosis of the intracranial lesion; OR
(c) Confirmation that the patient has received cranial irradiation without having had an intracranial lesion, and has undergone a 12 month period of observation following completion of treatment for the condition for which cranial irradiation was received; AND
6. Recent growth data (height and weight, not older than three months); AND
7. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
8. The proprietary name (brand), form and strength of somatotropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under a category other than risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have a chronological age of less than 2 years, **AND**
- Patient must have a documented clinical risk of hypoglycaemia, **AND**
- Patient must have documented evidence that the risk of hypoglycaemia is secondary to biochemical growth hormone deficiency, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; AND
3. Confirmation that the patient has a documented clinical risk of hypoglycaemia; AND
4. Confirmation that the patient has documented evidence that the risk of hypoglycaemia is secondary to biochemical growth hormone deficiency; AND
5. Recent growth data (height and weight, not older than three months); AND
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Biochemical growth hormone deficiency and precocious puberty

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under a category other than biochemical growth hormone deficiency and precocious puberty, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must be male and have commenced puberty (demonstrated by Tanner stage 2 genital or pubic hair development or testicular volumes greater than or equal to 4 mL) before the chronological age of 9 years; OR
- Patient must be female and have commenced puberty (demonstrated by Tanner stage 2 breast or pubic hair development) before the chronological age of 8 years; OR
- Patient must be female and menarche occurred before the chronological age of 10 years, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR

- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must be undergoing Gonadotrophin Releasing Hormone agonist therapy for pubertal suppression, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; AND
3. Confirmation that the patient has precocious puberty; AND
4. Confirmation that the patient is undergoing Gonadotrophin Releasing Hormone agonist therapy for pubertal suppression; AND
5. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; AND
6. Recent growth data (height and weight, not older than three months); AND
7. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
8. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth, **AND**
- Patient must have had a lapse in treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR

- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must have a structural lesion that is not neoplastic; OR
- Patient must have had a structural lesion that was neoplastic and have undergone a 12 month period of observation following completion of treatment for the structural lesion (all treatment); OR
- Patient must have a structural lesion that is neoplastic, have received medical advice that it is unsafe to treat the structural lesion, and have undergone a 12 month period of observation since initial diagnosis of the structural lesion,

AND

- Patient must have other hypothalamic/pituitary hormone deficits (includes ACTH, TSH, GnRH and/or vasopressin/ADH deficiencies), **AND**
- Patient must have hypothalamic obesity, **AND**
- Patient must have had a growth velocity above the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had an annual growth velocity of greater than 14 cm per year in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had an annual growth velocity of greater than 8 cm per year in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; **AND**
3. A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); **AND**
4. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; **AND**
5. (a) Confirmation that the patient has a structural lesion that is not neoplastic; OR
(b) Confirmation that the patient had a structural lesion that was neoplastic and has undergone a 12 month period of observation following completion of treatment for the structural lesion (all treatment); OR
(c) Confirmation that the patient has a structural lesion that is neoplastic, has received medical advice that it is unsafe to treat the structural lesion, and has undergone a 12 month period of observation since initial diagnosis of the structural lesion; **AND**
6. Confirmation that the patient has other hypothalamic/pituitary hormone deficits; **AND**
7. Confirmation that the patient has hypothalamic obesity; **AND**

8. Recent growth data (height and weight, not older than three months); AND
9. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
10. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with Turner syndrome

Treatment Phase: Recommencement of treatment as a reclassified patient

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under a category other than short stature associated with Turner syndrome, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have had a height at or below the 95th percentile for age on the Turner syndrome growth curve for girls immediately prior to commencing growth hormone treatment, **AND**
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as a loss of a whole X chromosome in all cells (45X), and gender of rearing is female; OR
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as a loss of a whole X chromosome in some cells (mosaic 46XX/45X), and gender of rearing is female; OR
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as genetic loss or rearrangement of an X chromosome (such as isochromosome X, ring-chromosome, or partial deletion of an X chromosome), and gender of rearing is female, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have a bone age of 2.5 years or less, **AND**
- Patient must not have a height greater than or equal to 155.0 cm, **AND**
- Patient must not have a bone age of 13.5 years or greater.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program)**

Special Arrangement 2015 and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; AND
3. A height measurement from immediately prior to commencement of growth hormone treatment; AND
4. Confirmation that the patient has diagnostic results consistent with Turner syndrome; AND
5. Recent growth data (height and weight, not older than three months); AND
6. A bone age result performed within the last 12 months; AND

The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature due to short stature homeobox (SHOX) gene disorders

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than short stature due to short stature homeobox (SHOX) gene disorders, **AND**
- Patient must have had a lapse in treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as a karyotype confirming the presence of a SHOX mutation/deletion without the presence of mixed gonadal dysgenesis; OR
- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as mixed gonadal dysgenesis (45X mosaic karyotype with the presence of any Y chromosome material and/or SRY gene positive by FISH study) and have an appropriate plan of management in place for the patient's increased risk of gonadoblastoma, **AND**
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment, **AND**
- Patient must have had a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had an annual growth velocity of 14 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had an annual growth velocity of 8 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes (excluding gonadoblastoma secondary to mixed gonadal dysgenesis), **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program)**

Special Arrangement 2015 and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; **AND**
3. A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12

months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); AND

4. Confirmation that the patient has diagnostic results consistent with a short stature homeobox (SHOX) gene disorder; AND

5. If the patient's condition is secondary to mixed gonadal dysgenesis, confirmation that an appropriate plan of management for the patient's increased risk of gonadoblastoma is in place; AND

6. Recent growth data (height and weight, not older than three months); AND

7. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND

8. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

somatropin 4 mg injection [1 vial] (&) inert substance diluent [1 vial], 1 pack

10447W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	180.36	40.30	Zomacton [FP]

■ SOMATROPIN

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Prior Written Approval of Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Short stature and slow growth

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature and slow growth category, **AND**
- Patient must not have been on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved and maintained mid parental height standard deviation score while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm.

Population criteria:

- Patient must be aged 3 years or older.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; AND
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
4. A bone age result performed within the last 12 months; AND

5. The final adult height (in cm) of the patient's mother and father (where available); AND
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with biochemical growth hormone deficiency

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature associated with biochemical growth hormone deficiency category, **AND**
- Patient must not have been on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved and maintained mid parental height standard deviation score while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Population criteria:

- Patient must be aged 3 years or older.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; AND
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
4. A bone age result performed within the last 12 months; AND
5. The final adult height (in cm) of the patient's mother and father (where available); AND
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Growth retardation secondary to an intracranial lesion, or cranial irradiation

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the growth retardation secondary to an intracranial lesion, or cranial irradiation category, **AND**
- Patient must not have been on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR

- Patient must have achieved and maintained mid parental height standard deviation score while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Population criteria:

- Patient must be aged 3 years or older.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; **AND**
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; **AND**
4. A bone age result performed within the last 12 months; **AND**
5. The final adult height (in cm) of the patient's mother and father (where available); **AND**
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants category, **AND**
- Patient must not have been on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved and maintained mid parental height standard deviation score while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have a chronological age of 5 years or greater.

Population criteria:

- Patient must be aged 3 years or older.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; **AND**
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; **AND**
4. A bone age result performed within the last 12 months; **AND**
5. The final adult height (in cm) of the patient's mother and father (where available); **AND**
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

When a patient receiving treatment under the indication risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants reaches or surpasses 5 years of age (chronological), prescribers should seek reclassification to the indication 'short stature due to biochemical growth hormone deficiency'.

Authority required

Biochemical growth hormone deficiency and precocious puberty

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the biochemical growth hormone deficiency and precocious puberty category, **AND**
- Patient must not have been on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved and maintained mid parental height standard deviation score while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Population criteria:

- Patient must be aged 3 years or older.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; AND
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
4. A bone age result performed within the last 12 months; AND
5. The final adult height (in cm) of the patient's mother and father (where available); AND
6. The proprietary name (brand), form and strength of somatotropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth category, **AND**
- Patient must not have been on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR

- Patient must have achieved and maintained mid parental height standard deviation score while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Population criteria:

- Patient must be aged 3 years or older.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; **AND**
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; **AND**
4. A bone age result performed within the last 12 months; **AND**
5. The final adult height (in cm) of the patient's mother and father (where available); **AND**
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with Turner syndrome

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature associated with Turner syndrome category, **AND**
- Patient must not have been on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an annualised growth velocity for bone age at or above the mean growth velocity for untreated Turner Syndrome girls (using the Turner Syndrome - Ranke growth velocity chart) while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have a bone age of 13.5 years or greater, **AND**
- Patient must not have a height greater than or equal to 155.0 cm.

Population criteria:

- Patient must be aged 3 years or older.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; **AND**
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; **AND**
4. A bone age result performed within the last 12 months; **AND**

5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature due to short stature homeobox (SHOX) gene disorders

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature due to short stature homeobox (SHOX) gene disorders category, **AND**
- Patient must not have been on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved and maintained mid parental height standard deviation score while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes (excluding gonadoblastoma secondary to mixed gonadal dysgenesis), **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7 cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0 cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Population criteria:

- Patient must be aged 3 years or older.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; **AND**
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; **AND**
4. A bone age result performed within the last 12 months; **AND**
5. The final adult height (in cm) of the patient's mother and father (where available); **AND**
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with chronic renal insufficiency

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature associated with chronic renal insufficiency category, **AND**
- Patient must not have been on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR

- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved and maintained mid parental height standard deviation score while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have undergone a renal transplant within the 12 month period immediately prior to the date of application, **AND**
- Patient must not have an eGFR equal to or greater than 30mL/min/1.73m², **AND**
- Patient must be male and must not have a height greater than or equal to 167.7 cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0 cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Population criteria:

- Patient must be aged 3 years or older.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; AND
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
4. A bone age result performed within the last 12 months; AND
5. The final adult height (in cm) of the patient's mother and father (where available); AND
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If a patient receiving treatment under the indication short stature due to chronic renal insufficiency undergoes a renal transplant and 12 months post-transplant has an eGFR of equal to or greater than 30mL/min/1.73m² prescribers should seek reclassification to the indication short stature and slow growth.

Authority required

Short stature and slow growth

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than short stature and slow growth, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have previously received treatment under the indication short stature associated with chronic renal insufficiency, have undergone a renal transplant and a 12 month period of observation following the transplant, and have an estimated glomerular filtration rate of greater than or equal to 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula; OR
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment and a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately

prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment), **AND**

- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7 cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0 cm.

Population criteria:

- Patient must be aged 3 years or older.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; **AND**
3. (a) A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment; OR
(b) Confirmation that the patient has previously received treatment under the indication **short stature associated with chronic renal insufficiency**, has undergone a renal transplant and a 12 month period of observation following the transplant, and has an estimated glomerular filtration rate of greater than or equal to 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula; **AND**
4. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; **AND**
5. A bone age result performed within the last 12 months; **AND**
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with biochemical growth hormone deficiency

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than short stature associated with biochemical growth hormone deficiency, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have previously received treatment under the indication **risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants** and have reached or surpassed 5 years of age (chronological); OR

- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment; OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and an annual growth velocity of 14 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and an annual growth velocity of 8 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Population criteria:

- Patient must be aged 3 years or older.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; **AND**
3. (a) A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); OR
(b) Height and weight measurements from within three months prior to commencement of treatment for a patient whose height was at or below the 1st percentile for age and sex immediately prior to commencing treatment; OR
(c) Confirmation that the patient has previously received treatment under the indication **risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants** and has reached or surpassed 5 years of age (chronological); **AND**
4. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; **AND**

5. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
6. A bone age result performed within the last 12 months; AND
7. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Biochemical growth hormone deficiency should not be secondary to an intracranial lesion or cranial irradiation for applications under this category.

Authority required

Growth retardation secondary to an intracranial lesion, or cranial irradiation

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than growth retardation secondary to an intracranial lesion, or cranial irradiation, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have had an intracranial lesion and have undergone a 12 month period of observation following completion of treatment for the intracranial lesion (all treatment); OR
- Patient must have had an intracranial lesion, have received medical advice that it is unsafe to treat the intracranial lesion, and have undergone a 12 month period of observation since initial diagnosis of the intracranial lesion; OR
- Patient must have received cranial irradiation without having had an intracranial lesion, and have undergone a 12 month period of observation following completion of treatment for the condition for which cranial irradiation was received, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-3 levels, **AND**
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment; OR
- Patient must have had both a height above the 1st percentile for age and sex immediately prior to commencing treatment and a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR

- Patient must have had both a height above the 1st percentile for age and sex immediately prior to commencing treatment and an annual growth velocity of 14 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had both a height above the 1st percentile for age and sex immediately prior to commencing treatment and an annual growth velocity of 8 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Population criteria:

- Patient must be aged 3 years or older.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; AND
3. (a) A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); OR
(b) Height and weight measurements from within three months prior to commencement of treatment for a patient whose height was at or below the 1st percentile for age and sex immediately prior to commencing treatment; AND
4. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; AND
5. (a) Confirmation that the patient has had an intracranial lesion and has undergone a 12 month period of observation following completion of treatment for the intracranial lesion (all treatment); OR
(b) Confirmation that the patient has had an intracranial lesion, has received medical advice that it is unsafe to treat the intracranial lesion, and has undergone a 12 month period of observation since initial diagnosis of the intracranial lesion; OR
(c) Confirmation that the patient has received cranial irradiation without having had an intracranial lesion, and has undergone a 12 month period of observation following completion of treatment for the condition for which cranial irradiation was received; AND
6. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
7. A bone age result performed within the last 12 months; AND
8. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Biochemical growth hormone deficiency and precocious puberty

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under a category other than biochemical growth hormone deficiency and precocious puberty, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR

- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must be male and have commenced puberty (demonstrated by Tanner stage 2 genital or pubic hair development or testicular volumes greater than or equal to 4 mL) before the chronological age of 9 years; OR
- Patient must be female and have commenced puberty (demonstrated by Tanner stage 2 breast or pubic hair development) before the chronological age of 8 years; OR
- Patient must be female and menarche occurred before the chronological age of 10 years, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must be undergoing Gonadotrophin Releasing Hormone agonist therapy for pubertal suppression, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Population criteria:

- Patient must be aged 3 years or older.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; **AND**
3. Confirmation that the patient has precocious puberty; **AND**
4. Confirmation that the patient is undergoing Gonadotrophin Releasing Hormone agonist therapy for pubertal suppression; **AND**
5. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; **AND**
6. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; **AND**
7. A bone age result performed within the last 12 months; **AND**
8. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have a structural lesion that is not neoplastic; OR
- Patient must have had a structural lesion that was neoplastic and have undergone a 12 month period of observation following completion of treatment for the structural lesion (all treatment); OR
- Patient must have a structural lesion that is neoplastic, have received medical advice that it is unsafe to treat the structural lesion, and have undergone a 12 month period of observation since initial diagnosis of the structural lesion,

AND

- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must have other hypothalamic/pituitary hormone deficits (includes ACTH, TSH, GnRH and/or vasopressin/ADH deficiencies), **AND**
- Patient must have hypothalamic obesity, **AND**
- Patient must have had a growth velocity above the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had an annual growth velocity of greater than 14 cm per year in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had an annual growth velocity of greater than 8 cm per year in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must not have diabetes mellitus, **AND**

- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Population criteria:

- Patient must be aged 3 years or older.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; AND
3. A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); AND
4. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; AND
5. (a) Confirmation that the patient has a structural lesion that is not neoplastic; OR
(b) Confirmation that the patient had a structural lesion that was neoplastic and has undergone a 12 month period of observation following completion of treatment for the structural lesion (all treatment); OR
(c) Confirmation that the patient has a structural lesion that is neoplastic, has received medical advice that it is unsafe to treat the structural lesion, and has undergone a 12 month period of observation since initial diagnosis of the structural lesion; AND
6. Confirmation that the patient has other hypothalamic/pituitary hormone deficits; AND
7. Confirmation that the patient has hypothalamic obesity; AND
8. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
9. A bone age result performed within the last 12 months; AND
10. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with Turner syndrome

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under a category other than short stature associated with Turner syndrome, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**

- Patient must have had a height at or below the 95th percentile for age on the Turner syndrome growth curve for girls immediately prior to commencing growth hormone treatment, **AND**
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as a loss of a whole X chromosome in all cells (45X), and gender of rearing is female; OR
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as a loss of a whole X chromosome in some cells (mosaic 46XX/45X), and gender of rearing is female; OR
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as genetic loss or rearrangement of an X chromosome (such as isochromosome X, ring-chromosome, or partial deletion of an X chromosome), and gender of rearing is female, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have a bone age of 2.5 years or less, **AND**
- Patient must not have a bone age of 13.5 years or greater, **AND**
- Patient must not have a height greater than or equal to 155.0 cm.

Population criteria:

- Patient must be aged 3 years or older.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; AND
3. A height measurement from immediately prior to commencement of growth hormone treatment; AND
4. Confirmation that the patient has diagnostic results consistent with Turner syndrome; AND
5. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
6. A bone age result performed within the last 12 months; AND
7. The proprietary name (brand), form and strength of somatotropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature due to short stature homeobox (SHOX) gene disorders

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than short stature due to short stature homeobox (SHOX) gene disorders, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or commencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or commencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or commencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or commencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or commencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as a karyotype confirming the presence of a SHOX mutation/deletion without the presence of mixed gonadal dysgenesis; OR

- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as mixed gonadal dysgenesis (45X mosaic karyotype with the presence of any Y chromosome material and/or SRY gene positive by FISH study) and have an appropriate plan of management in place for the patient's increased risk of gonadoblastoma, **AND**
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment, **AND**
- Patient must have had a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had an annual growth velocity of 14 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had an annual growth velocity of 8 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes (excluding gonadoblastoma secondary to mixed gonadal dysgenesis), **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Population criteria:

- Patient must be aged 3 years or older.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; AND
3. A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); AND
4. Confirmation that the patient has diagnostic results consistent with a short stature homeobox (SHOX) gene disorder; AND
5. If the patient's condition is secondary to mixed gonadal dysgenesis, confirmation that an appropriate plan of management for the patient's increased risk of gonadoblastoma is in place; AND
6. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
7. A bone age result performed within the last 12 months; AND
8. The proprietary name (brand), form and strength of somatotropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with chronic renal insufficiency

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than short stature associated with chronic renal insufficiency, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a

continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR

- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment; OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and a growth velocity less than or equal to the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and an annual growth velocity of 14 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and an annual growth velocity of 8 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must have an estimated glomerular filtration rate less than 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula, and not have undergone a renal transplant; OR
- Patient must have an estimated glomerular filtration rate less than 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula, have undergone a renal transplant, and have undergone a 12 month period of observation following the transplant, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Population criteria:

- Patient must be aged 3 years or older.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; **AND**
3. (a) A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); OR
(b) Height and weight measurements from within three months prior to commencement of treatment for a patient whose height was at or below the 1st percentile for age and sex immediately prior to commencing treatment; **AND**
4. Confirmation that the patient has an estimated glomerular filtration rate less than 30mL/minute/1.73m²; **AND**
5. If a renal transplant has taken place, confirmation that the patient has undergone a 12 month period of observation following transplantation; **AND**
6. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; **AND**
7. A bone age result performed within the last 12 months; **AND**

The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

SYSTEMIC HORMONAL PREPARATIONS, EXCL. SEX HORMONES AND INSULINS

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If a patient receiving treatment under the indication short stature due to chronic renal insufficiency undergoes a renal transplant and 12 months post-transplant has an eGFR of equal to or greater than 30mL/min/1.73m² prescribers should seek reclassification to the indication short stature and slow growth.

somatropin 5 mg/1.5 mL injection, 1.5 mL cartridge

10427T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	223.61	40.30	Scitropin A [SA]

somatropin 5 mg/1.5 mL injection, 1.5 mL cartridge

10507B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	223.61	40.30	Omnitrope Surepal 5 [SZ]

somatropin 10 mg/1.5 mL injection, 1.5 mL cartridge

10441M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	439.82	40.30	SciTropin A [SA]

somatropin 10 mg/1.5 mL injection, 1.5 mL cartridge

10506Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	439.82	40.30	Omnitrope Surepal 10 [SZ]

somatropin 15 mg/1.5 mL injection, 1.5 mL cartridge

10490D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	656.04	40.30	Omnitrope Surepal 15 [SZ]

■ SOMATROPIN

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Prior Written Approval of Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Short stature and slow growth

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature and slow growth category, **AND**
- Patient must not have been on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved and maintained mid parental height standard deviation score while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special**

Arrangement 2015 and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; AND
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
4. A bone age result performed within the last 12 months; AND
5. The final adult height (in cm) of the patient's mother and father (where available); AND
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with biochemical growth hormone deficiency

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature associated with biochemical growth hormone deficiency category, **AND**
- Patient must not have been on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved and maintained mid parental height standard deviation score while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; AND
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
5. The final adult height (in cm) of the patient's mother and father (where available); AND
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Growth retardation secondary to an intracranial lesion, or cranial irradiation

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the growth retardation secondary to an intracranial lesion, or cranial irradiation category, **AND**
- Patient must not have been on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR

- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved and maintained mid parental height standard deviation score while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; **AND**
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; **AND**
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
5. The final adult height (in cm) of the patient's mother and father (where available); **AND**
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants category, **AND**
- Patient must not have been on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved and maintained mid parental height standard deviation score while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have a chronological age of 5 years or greater.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; **AND**

3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
5. The final adult height (in cm) of the patient's mother and father (where available); AND
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

When a patient receiving treatment under the indication risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants reaches or surpasses 5 years of age (chronological), prescribers should seek reclassification to the indication 'short stature due to biochemical growth hormone deficiency'.

Authority required

Biochemical growth hormone deficiency and precocious puberty

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the biochemical growth hormone deficiency and precocious puberty category, **AND**
- Patient must not have been on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved and maintained mid parental height standard deviation score while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; AND
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
5. The final adult height (in cm) of the patient's mother and father (where available); AND
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth category, **AND**
- Patient must not have been on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR

- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved and maintained mid parental height standard deviation score while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; **AND**
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; **AND**
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
5. The final adult height (in cm) of the patient's mother and father (where available); **AND**
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with Turner syndrome

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature associated with Turner syndrome category, **AND**
- Patient must not have been on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an annualised growth velocity for bone age at or above the mean growth velocity for untreated Turner Syndrome girls (using the Turner Syndrome - Ranke growth velocity chart) while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have a bone age of 13.5 years or greater, **AND**
- Patient must not have a height greater than or equal to 155.0 cm.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; **AND**
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; **AND**

4. A bone age result performed within the last 12 months; AND

5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature due to short stature homeobox (SHOX) gene disorders

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature due to short stature homeobox (SHOX) gene disorders category, **AND**
- Patient must not have been on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved and maintained mid parental height standard deviation score while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes (excluding gonadoblastoma secondary to mixed gonadal dysgenesis), **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7 cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0 cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; AND
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
5. The final adult height (in cm) of the patient's mother and father (where available); AND
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with chronic renal insufficiency

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature associated with chronic renal insufficiency category, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have been on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR

- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved and maintained mid parental height standard deviation score while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have undergone a renal transplant within the 12 month period immediately prior to the date of application, **AND**
- Patient must not have an eGFR equal to or greater than 30mL/min/1.73m², **AND**
- Patient must be male and must not have a height greater than or equal to 167.7 cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0 cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; **AND**
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; **AND**
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
5. The final adult height (in cm) of the patient's mother and father (where available); **AND**
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If a patient receiving treatment under the indication short stature due to chronic renal insufficiency undergoes a renal transplant and 12 months post-transplant has an eGFR of equal to or greater than 30mL/min/1.73m² prescribers should seek reclassification to the indication short stature and slow growth.

Authority required

Short stature and slow growth

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than short stature and slow growth, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have previously received treatment under the indication short stature associated with chronic renal insufficiency, have undergone a renal transplant and a 12 month period of observation following the transplant, and have an estimated glomerular filtration rate of greater than or equal to 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula; OR
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment and a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment), **AND**
- Patient must not have diabetes mellitus, **AND**

- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; AND
3. (a) A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment; OR
(b) Confirmation that the patient has previously received treatment under the indication **short stature associated with chronic renal insufficiency**, has undergone a renal transplant and a 12 month period of observation following the transplant, and has an estimated glomerular filtration rate of greater than or equal to 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula; AND
4. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
5. A bone age result performed within the last 12 months; AND
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with biochemical growth hormone deficiency

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than short stature associated with biochemical growth hormone deficiency, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have previously received treatment under the indication **risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants** and have reached or surpassed 5 years of age (chronological); OR
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment; OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR

- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and an annual growth velocity of 14 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and an annual growth velocity of 8 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; **AND**
3. (a) A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); OR
(b) Height and weight measurements from within three months prior to commencement of treatment for a patient whose height was at or below the 1st percentile for age and sex immediately prior to commencing treatment; OR
(c) Confirmation that the patient has previously received treatment under the indication **risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants** and has reached or surpassed 5 years of age (chronological); **AND**
4. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; **AND**
5. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; **AND**
6. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
7. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Biochemical growth hormone deficiency should not be secondary to an intracranial lesion or cranial irradiation for applications under this category.

Authority required

Growth retardation secondary to an intracranial lesion, or cranial irradiation

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than growth retardation secondary to an intracranial lesion, or cranial irradiation, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have had an intracranial lesion and have undergone a 12 month period of observation following completion of treatment for the intracranial lesion (all treatment); OR
- Patient must have had an intracranial lesion, have received medical advice that it is unsafe to treat the intracranial lesion, and have undergone a 12 month period of observation since initial diagnosis of the intracranial lesion; OR
- Patient must have received cranial irradiation without having had an intracranial lesion, and have undergone a 12 month period of observation following completion of treatment for the condition for which cranial irradiation was received, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment; OR
- Patient must have had both a height above the 1st percentile for age and sex immediately prior to commencing treatment and a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had both a height above the 1st percentile for age and sex immediately prior to commencing treatment and an annual growth velocity of 14 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had both a height above the 1st percentile for age and sex immediately prior to commencing treatment and an annual growth velocity of 8 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must not have diabetes mellitus, **AND**

- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; AND
3. (a) A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); OR
(b) Height and weight measurements from within three months prior to commencement of treatment for a patient whose height was at or below the 1st percentile for age and sex immediately prior to commencing treatment; AND
4. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; AND
5. (a) Confirmation that the patient has had an intracranial lesion and has undergone a 12 month period of observation following completion of treatment for the intracranial lesion (all treatment); OR
(b) Confirmation that the patient has had an intracranial lesion, has received medical advice that it is unsafe to treat the intracranial lesion, and has undergone a 12 month period of observation since initial diagnosis of the intracranial lesion; OR
(c) Confirmation that the patient has received cranial irradiation without having had an intracranial lesion, and has undergone a 12 month period of observation following completion of treatment for the condition for which cranial irradiation was received; AND
6. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
7. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
8. The proprietary name (brand), form and strength of somatotropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

Authority required

Risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under a category other than risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have a chronological age of less than 2 years, **AND**

- Patient must have a documented clinical risk of hypoglycaemia, **AND**
- Patient must have documented evidence that the risk of hypoglycaemia is secondary to biochemical growth hormone deficiency, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; AND
3. Confirmation that the patient has a documented clinical risk of hypoglycaemia; AND
4. Confirmation that the patient has documented evidence that the risk of hypoglycaemia is secondary to biochemical growth hormone deficiency; AND
5. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Biochemical growth hormone deficiency and precocious puberty

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under a category other than biochemical growth hormone deficiency and precocious puberty, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or commencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or commencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or commencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or commencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or commencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must be male and have commenced puberty (demonstrated by Tanner stage 2 genital or pubic hair development or testicular volumes greater than or equal to 4 mL) before the chronological age of 9 years; OR
- Patient must be female and have commenced puberty (demonstrated by Tanner stage 2 breast or pubic hair development) before the chronological age of 8 years; OR
- Patient must be female and menarche occurred before the chronological age of 10 years, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline

- abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
 - Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
 - Patient must be undergoing Gonadotrophin Releasing Hormone agonist therapy for pubertal suppression, **AND**
 - Patient must not have diabetes mellitus, **AND**
 - Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
 - Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
 - Patient must be male and must not have a bone age of 15.5 years or more; OR
 - Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; AND
3. Confirmation that the patient has precocious puberty; AND
4. Confirmation that the patient is undergoing Gonadotrophin Releasing Hormone agonist therapy for pubertal suppression; AND
5. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; AND
6. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
7. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
8. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or commencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or commencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or commencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or commencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or commencement treatment period and 26 weeks for a continuing treatment period, whichever applies).

continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**

- Patient must have a structural lesion that is not neoplastic; OR
- Patient must have had a structural lesion that was neoplastic and have undergone a 12 month period of observation following completion of treatment for the structural lesion (all treatment); OR
- Patient must have a structural lesion that is neoplastic, have received medical advice that it is unsafe to treat the structural lesion, and have undergone a 12 month period of observation since initial diagnosis of the structural lesion,

AND

- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must have other hypothalamic/pituitary hormone deficits (includes ACTH, TSH, GnRH and/or vasopressin/ADH deficiencies), **AND**
- Patient must have hypothalamic obesity, **AND**
- Patient must have had a growth velocity above the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had an annual growth velocity of greater than 14 cm per year in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had an annual growth velocity of greater than 8 cm per year in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; **AND**
3. A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); **AND**
4. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; **AND**

5. (a) Confirmation that the patient has a structural lesion that is not neoplastic; OR
 - (b) Confirmation that the patient had a structural lesion that was neoplastic and has undergone a 12 month period of observation following completion of treatment for the structural lesion (all treatment); OR
 - (c) Confirmation that the patient has a structural lesion that is neoplastic, has received medical advice that it is unsafe to treat the structural lesion, and has undergone a 12 month period of observation since initial diagnosis of the structural lesion; AND
 6. Confirmation that the patient has other hypothalamic/pituitary hormone deficits; AND
 7. Confirmation that the patient has hypothalamic obesity; AND
 8. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
 9. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
 10. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).
- Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with Turner syndrome

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under a category other than short stature associated with Turner syndrome, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have had a height at or below the 95th percentile for age on the Turner syndrome growth curve for girls immediately prior to commencing growth hormone treatment, **AND**
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as a loss of a whole X chromosome in all cells (45X), and gender of rearing is female; OR
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as a loss of a whole X chromosome in some cells (mosaic 46XX/45X), and gender of rearing is female; OR
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as genetic loss or rearrangement of an X chromosome (such as isochromosome X, ring-chromosome, or partial deletion of an X chromosome), and gender of rearing is female, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have a bone age of 2.5 years or less, **AND**
- Patient must not have a bone age of 13.5 years or greater, **AND**
- Patient must not have a height greater than or equal to 155.0 cm.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND

2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; AND
 3. A height measurement from immediately prior to commencement of growth hormone treatment; AND
 4. Confirmation that the patient has diagnostic results consistent with Turner syndrome; AND
 5. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
 6. A bone age result performed within the last 12 months; AND
 7. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).
- Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature due to short stature homeobox (SHOX) gene disorders

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than short stature due to short stature homeobox (SHOX) gene disorders, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as a karyotype confirming the presence of a SHOX mutation/deletion without the presence of mixed gonadal dysgenesis; OR
- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as mixed gonadal dysgenesis (45X mosaic karyotype with the presence of any Y chromosome material and/or SRY gene positive by FISH study) and have an appropriate plan of management in place for the patient's increased risk of gonadoblastoma, **AND**
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment, **AND**
- Patient must have had a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had an annual growth velocity of 14 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had an annual growth velocity of 8 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes (excluding gonadoblastoma secondary to mixed gonadal dysgenesis), **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special**

Arrangement 2015 and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; AND
3. A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); AND
4. Confirmation that the patient has diagnostic results consistent with a short stature homeobox (SHOX) gene disorder; AND
5. If the patient's condition is secondary to mixed gonadal dysgenesis, confirmation that an appropriate plan of management for the patient's increased risk of gonadoblastoma is in place; AND
6. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
7. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
8. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with chronic renal insufficiency

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than short stature associated with chronic renal insufficiency, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment; OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and a growth velocity less than or equal to the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and an annual growth velocity of 14 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and an annual growth velocity of 8 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must have an estimated glomerular filtration rate less than 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula, and not have undergone a renal transplant; OR
- Patient must have an estimated glomerular filtration rate less than 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula, have undergone a renal transplant, and have undergone a 12 month period of observation following the transplant, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**

- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; AND
3. (a) A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); OR
(b) Height and weight measurements from within three months prior to commencement of treatment for a patient whose height was at or below the 1st percentile for age and sex immediately prior to commencing treatment; AND
4. Confirmation that the patient has an estimated glomerular filtration rate less than 30ml/minute/1.73m²; AND
5. If a renal transplant has taken place, confirmation that the patient has undergone a 12 month period of observation following transplantation; AND
6. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
7. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND

The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If a patient receiving treatment under the indication short stature due to chronic renal insufficiency undergoes a renal transplant and 12 months post-transplant has an eGFR of equal to or greater than 30mL/min/1.73m² prescribers should seek reclassification to the indication short stature and slow growth.

somatropin 5 mg/1.5 mL injection, 1.5 mL cartridge

10432C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	223.61	40.30	Norditropin FlexPro [NO]

somatropin 5 mg/1.5 mL injection, 1.5 mL cartridge

10469B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	223.61	40.30	Norditropin SimpleXx [NO]

somatropin 10 mg/1.5 mL injection, 1.5 mL cartridge

10439K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	439.82	40.30	Norditropin SimpleXx [NO]

somatropin 10 mg/1.5 mL injection, 1.5 mL cartridge

10451C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	439.82	40.30	Norditropin FlexPro [NO]

somatropin 10 mg/2 mL injection, 2 mL cartridge

10478L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	426.41	40.30	NutropinAq [IS]

somatropin 15 mg/1.5 mL injection, 1.5 mL cartridge

10449Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	656.04	40.30	Norditropin FlexPro [NO]

somatropin 15 mg/1.5 mL injection, 1.5 mL cartridge

10468Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	656.04	40.30	Norditropin SimpleXx [NO]

■ SOMATROPIN

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Prior Written Approval of Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Short stature and slow growth

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature and slow growth category, **AND**
- Patient must not have been on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); **OR**
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); **OR**
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); **OR**
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); **OR**
- Patient must have achieved and maintained mid parental height standard deviation score while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; **OR**
- Patient must be female and must not have a bone age of 13.5 years or more, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; **OR**
- Patient must be female and must not have a height greater than or equal to 155.0cm.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; **AND**
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; **AND**
4. A bone age result performed within the last 12 months; **AND**
5. The final adult height (in cm) of the patient's mother and father (where available); **AND**
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with biochemical growth hormone deficiency

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature associated with biochemical growth hormone deficiency category, **AND**
- Patient must not have been on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); **OR**

- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved and maintained mid parental height standard deviation score while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; **AND**
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; **AND**
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
5. The final adult height (in cm) of the patient's mother and father (where available); **AND**
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Growth retardation secondary to an intracranial lesion, or cranial irradiation

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the growth retardation secondary to an intracranial lesion, or cranial irradiation category, **AND**
- Patient must not have been on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved and maintained mid parental height standard deviation score while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; **AND**

3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND

5. The final adult height (in cm) of the patient's mother and father (where available); AND

6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants category, **AND**
- Patient must not have been on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved and maintained mid parental height standard deviation score while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have a chronological age of 5 years or greater.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; AND
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
5. The final adult height (in cm) of the patient's mother and father (where available); AND
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

When a patient receiving treatment under the indication risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants reaches or surpasses 5 years of age (chronological), prescribers should seek reclassification to the indication 'short stature due to biochemical growth hormone deficiency'.

Authority required

Biochemical growth hormone deficiency and precocious puberty

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the biochemical growth hormone deficiency and precocious puberty category, **AND**
- Patient must not have been on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR

- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved and maintained mid parental height standard deviation score while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; **AND**
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; **AND**
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
5. The final adult height (in cm) of the patient's mother and father (where available); **AND**
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth category, **AND**
- Patient must not have been on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved and maintained mid parental height standard deviation score while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; **AND**
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; **AND**

4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
5. The final adult height (in cm) of the patient's mother and father (where available); AND
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with Turner syndrome

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature associated with Turner syndrome category, **AND**
- Patient must not have been on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an annualised growth velocity for bone age at or above the mean growth velocity for untreated Turner Syndrome girls (using the Turner Syndrome - Ranke growth velocity chart) while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have a bone age of 13.5 years or greater, **AND**
- Patient must not have a height greater than or equal to 155.0 cm.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; AND
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
4. A bone age result performed within the last 12 months; AND
5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature due to short stature homeobox (SHOX) gene disorders

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature due to short stature homeobox (SHOX) gene disorders category, **AND**
- Patient must not have been on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR

- Patient must have achieved and maintained mid parental height standard deviation score while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes (excluding gonadoblastoma secondary to mixed gonadal dysgenesis), **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7 cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0 cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; **AND**
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; **AND**
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
5. The final adult height (in cm) of the patient's mother and father (where available); **AND**
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature and slow growth

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than short stature and slow growth, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have previously received treatment under the indication short stature associated with chronic renal insufficiency, have undergone a renal transplant and a 12 month period of observation following the transplant, and have an estimated glomerular filtration rate of greater than or equal to 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula; OR
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment and a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; AND
3. (a) A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment; OR
(b) Confirmation that the patient has previously received treatment under the indication **short stature associated with chronic renal insufficiency**, has undergone a renal transplant and a 12 month period of observation following the transplant, and has an estimated glomerular filtration rate of greater than or equal to 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula; AND
4. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
5. A bone age result performed within the last 12 months; AND
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with biochemical growth hormone deficiency

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than short stature associated with biochemical growth hormone deficiency, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have previously received treatment under the indication **risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants** and have reached or surpassed 5 years of age (chronological); OR
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment; OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and an annual growth velocity of 14 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and an annual growth velocity of 8 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**

- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; **AND**
3. (a) A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); OR
(b) Height and weight measurements from within three months prior to commencement of treatment for a patient whose height was at or below the 1st percentile for age and sex immediately prior to commencing treatment; OR
(c) Confirmation that the patient has previously received treatment under the indication **risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants** and has reached or surpassed 5 years of age (chronological); **AND**
4. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; **AND**
5. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; **AND**
6. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
7. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Biochemical growth hormone deficiency should not be secondary to an intracranial lesion or cranial irradiation for applications under this category.

Authority required

Growth retardation secondary to an intracranial lesion, or cranial irradiation

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than growth retardation secondary to an intracranial lesion, or cranial irradiation, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have had an intracranial lesion and have undergone a 12 month period of observation following completion of treatment for the intracranial lesion (all treatment); OR
- Patient must have had an intracranial lesion, have received medical advice that it is unsafe to treat the intracranial lesion, and have undergone a 12 month period of observation since initial diagnosis of the intracranial lesion; OR
- Patient must have received cranial irradiation without having had an intracranial lesion, and have undergone a 12 month period of observation following completion of treatment for the condition for which cranial irradiation was received, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment; OR
- Patient must have had both a height above the 1st percentile for age and sex immediately prior to commencing treatment and a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had both a height above the 1st percentile for age and sex immediately prior to commencing treatment and an annual growth velocity of 14 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had both a height above the 1st percentile for age and sex immediately prior to commencing treatment and an annual growth velocity of 8 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; AND
3. (a) A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); OR
(b) Height and weight measurements from within three months prior to commencement of treatment for a patient whose height was at or below the 1st percentile for age and sex immediately prior to commencing treatment; AND
4. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; AND
5. (a) Confirmation that the patient has had an intracranial lesion and has undergone a 12 month period of observation following completion of treatment for the intracranial lesion (all treatment); OR
(b) Confirmation that the patient has had an intracranial lesion, has received medical advice that it is unsafe to treat the intracranial lesion, and has undergone a 12 month period of observation since initial diagnosis of the intracranial lesion; OR
(c) Confirmation that the patient has received cranial irradiation without having had an intracranial lesion, and has undergone a 12 month period of observation following completion of treatment for the condition for which cranial irradiation was received; AND
6. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
7. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
8. The proprietary name (brand), form and strength of somatotropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

Authority required

Risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under a category other than risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have a chronological age of less than 2 years, **AND**
- Patient must have a documented clinical risk of hypoglycaemia, **AND**
- Patient must have documented evidence that the risk of hypoglycaemia is secondary to biochemical growth hormone deficiency, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; AND
3. Confirmation that the patient has a documented clinical risk of hypoglycaemia; AND
4. Confirmation that the patient has documented evidence that the risk of hypoglycaemia is secondary to biochemical growth hormone deficiency; AND
5. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Biochemical growth hormone deficiency and precocious puberty

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under a category other than biochemical growth hormone deficiency and precocious puberty, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or commencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or commencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or commencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or commencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or commencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must be male and have commenced puberty (demonstrated by Tanner stage 2 genital or pubic hair development or testicular volumes greater than or equal to 4 mL) before the chronological age of 9 years; OR
- Patient must be female and have commenced puberty (demonstrated by Tanner stage 2 breast or pubic hair development) before the chronological age of 8 years; OR
- Patient must be female and menarche occurred before the chronological age of 10 years, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR

- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must be undergoing Gonadotrophin Releasing Hormone agonist therapy for pubertal suppression, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; **AND**
3. Confirmation that the patient has precocious puberty; **AND**
4. Confirmation that the patient is undergoing Gonadotrophin Releasing Hormone agonist therapy for pubertal suppression; **AND**
5. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; **AND**
6. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; **AND**
7. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
8. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have a structural lesion that is not neoplastic; OR
- Patient must have had a structural lesion that was neoplastic and have undergone a 12 month period of observation following completion of treatment for the structural lesion (all treatment); OR
- Patient must have a structural lesion that is neoplastic, have received medical advice that it is unsafe to treat the structural lesion, and have undergone a 12 month period of observation since initial diagnosis of the structural lesion, **AND**

- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must have other hypothalamic/pituitary hormone deficits (includes ACTH, TSH, GnRH and/or vasopressin/ADH deficiencies), **AND**
- Patient must have hypothalamic obesity, **AND**
- Patient must have had a growth velocity above the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had an annual growth velocity of greater than 14 cm per year in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had an annual growth velocity of greater than 8 cm per year in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; **AND**
3. A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); **AND**
4. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; **AND**
5. (a) Confirmation that the patient has a structural lesion that is not neoplastic; OR
(b) Confirmation that the patient had a structural lesion that was neoplastic and has undergone a 12 month period of observation following completion of treatment for the structural lesion (all treatment); OR
(c) Confirmation that the patient has a structural lesion that is neoplastic, has received medical advice that it is unsafe to treat the structural lesion, and has undergone a 12 month period of observation since initial diagnosis of the structural lesion; **AND**
6. Confirmation that the patient has other hypothalamic/pituitary hormone deficits; **AND**

7. Confirmation that the patient has hypothalamic obesity; AND
 8. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
 9. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
 10. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).
- Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with Turner syndrome

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under a category other than short stature associated with Turner syndrome, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have had a height at or below the 95th percentile for age on the Turner syndrome growth curve for girls immediately prior to commencing growth hormone treatment, **AND**
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as a loss of a whole X chromosome in all cells (45X), and gender of rearing is female; OR
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as a loss of a whole X chromosome in some cells (mosaic 46XX/45X), and gender of rearing is female; OR
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as genetic loss or rearrangement of an X chromosome (such as isochromosome X, ring-chromosome, or partial deletion of an X chromosome), and gender of rearing is female, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have a bone age of 2.5 years or less, **AND**
- Patient must not have a bone age of 13.5 years or greater, **AND**
- Patient must not have a height greater than or equal to 155.0 cm.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; AND
3. A height measurement from immediately prior to commencement of growth hormone treatment; AND
4. Confirmation that the patient has diagnostic results consistent with Turner syndrome; AND
5. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
6. A bone age result performed within the last 12 months; AND

7. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature due to short stature homeobox (SHOX) gene disorders

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than short stature due to short stature homeobox (SHOX) gene disorders, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as a karyotype confirming the presence of a SHOX mutation/deletion without the presence of mixed gonadal dysgenesis; OR
- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as mixed gonadal dysgenesis (45X mosaic karyotype with the presence of any Y chromosome material and/or SRY gene positive by FISH study) and have an appropriate plan of management in place for the patient's increased risk of gonadoblastoma, **AND**
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment, **AND**
- Patient must have had a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had an annual growth velocity of 14 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had an annual growth velocity of 8 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes (excluding gonadoblastoma secondary to mixed gonadal dysgenesis), **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; **AND**
3. A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient

was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); AND

4. Confirmation that the patient has diagnostic results consistent with a short stature homeobox (SHOX) gene disorder; AND

5. If the patient's condition is secondary to mixed gonadal dysgenesis, confirmation that an appropriate plan of management for the patient's increased risk of gonadoblastoma is in place; AND

6. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND

7. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND

8. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with chronic renal insufficiency

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature associated with chronic renal insufficiency category, **AND**
- Patient must not have been on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved and maintained mid parental height standard deviation score while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have undergone a renal transplant within the 12 month period immediately prior to the date of application, **AND**
- Patient must not have an eGFR equal to or greater than 30mL/min/1.73m², **AND**
- Patient must be male and must not have a height greater than or equal to 167.7 cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0 cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Population criteria:

- Patient must be aged 3 years or older.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; AND
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
4. A bone age result performed within the last 12 months; AND
5. The final adult height (in cm) of the patient's mother and father (where available); AND
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If a patient receiving treatment under the indication short stature due to chronic renal insufficiency undergoes a renal transplant and 12 months post-transplant has an eGFR of equal to or greater than 30mL/min/1.73m² prescribers should seek reclassification to the indication short stature and slow growth.

Authority required

Short stature associated with chronic renal insufficiency

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than short stature associated with chronic renal insufficiency, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment; OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and a growth velocity less than or equal to the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and an annual growth velocity of 14 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and an annual growth velocity of 8 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must have an estimated glomerular filtration rate less than 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula, and not have undergone a renal transplant; OR
- Patient must have an estimated glomerular filtration rate less than 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula, have undergone a renal transplant, and have undergone a 12 month period of observation following the transplant, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Population criteria:

- Patient must be aged 3 years or older.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; **AND**
3. (a) A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient

was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); OR

(b) Height and weight measurements from within three months prior to commencement of treatment for a patient whose height was at or below the 1st percentile for age and sex immediately prior to commencing treatment; AND

4. Confirmation that the patient has an estimated glomerular filtration rate less than 30ml/minute/1.73m² ; AND

5. If a renal transplant has taken place, confirmation that the patient has undergone a 12 month period of observation following transplantation; AND

6. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND

7. A bone age result performed within the last 12 months; AND

The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If a patient receiving treatment under the indication short stature due to chronic renal insufficiency undergoes a renal transplant and 12 months post-transplant has an eGFR of equal to or greater than 30mL/min/1.73m² prescribers should seek reclassification to the indication short stature and slow growth.

somatropin 12 mg/1.5 mL injection, 1.5 mL cartridge

10483R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	526.31	40.30	Saizen [SG]

somatropin 6 mg/1.03 mL injection, 1.03 mL cartridge

10462P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	266.85	40.30	Saizen [SG]

somatropin 20 mg/2.5 mL injection, 2.5 mL cartridge

10497L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	872.25	40.30	Saizen [SG]

■ SOMATROPIN

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Prior Written Approval of Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Short stature and slow growth

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature and slow growth category, **AND**
- Patient must not have been on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved and maintained mid parental height standard deviation score while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more, **AND**

- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
 - Patient must be female and must not have a height greater than or equal to 155.0cm.
- The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; AND
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
4. A bone age result performed within the last 12 months; AND
5. The final adult height (in cm) of the patient's mother and father (where available); AND
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with biochemical growth hormone deficiency

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature associated with biochemical growth hormone deficiency category, **AND**
- Patient must not have been on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved and maintained mid parental height standard deviation score while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; AND
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
5. The final adult height (in cm) of the patient's mother and father (where available); AND
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Growth retardation secondary to an intracranial lesion, or cranial irradiation

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the growth retardation secondary to an intracranial lesion, or cranial irradiation category, **AND**

- Patient must not have been on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved and maintained mid parental height standard deviation score while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; **AND**
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; **AND**
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
5. The final adult height (in cm) of the patient's mother and father (where available); **AND**
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants category, **AND**
- Patient must not have been on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved and maintained mid parental height standard deviation score while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have a chronological age of 5 years or greater.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; AND
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
5. The final adult height (in cm) of the patient's mother and father (where available); AND
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

When a patient receiving treatment under the indication risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants reaches or surpasses 5 years of age (chronological), prescribers should seek reclassification to the indication 'short stature due to biochemical growth hormone deficiency'.

Authority required

Biochemical growth hormone deficiency and precocious puberty

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the biochemical growth hormone deficiency and precocious puberty category, **AND**
- Patient must not have been on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved and maintained mid parental height standard deviation score while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; AND
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
5. The final adult height (in cm) of the patient's mother and father (where available); AND
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth category, **AND**
- Patient must not have been on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR

- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved and maintained mid parental height standard deviation score while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; **AND**
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; **AND**
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
5. The final adult height (in cm) of the patient's mother and father (where available); **AND**
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with Turner syndrome

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature associated with Turner syndrome category, **AND**
- Patient must not have been on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an annualised growth velocity for bone age at or above the mean growth velocity for untreated Turner Syndrome girls (using the Turner Syndrome - Ranke growth velocity chart) while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have a bone age of 13.5 years or greater, **AND**
- Patient must not have a height greater than or equal to 155.0 cm.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**

2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; AND
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
4. A bone age result performed within the last 12 months; AND
5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature due to short stature homeobox (SHOX) gene disorders

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature due to short stature homeobox (SHOX) gene disorders category, **AND**
- Patient must not have been on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved and maintained mid parental height standard deviation score while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes (excluding gonadoblastoma secondary to mixed gonadal dysgenesis), **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7 cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0 cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; AND
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
5. The final adult height (in cm) of the patient's mother and father (where available); AND
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with chronic renal insufficiency

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature associated with chronic renal insufficiency category, **AND**
- Patient must not have been on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR

- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved and maintained mid parental height standard deviation score while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have undergone a renal transplant within the 12 month period immediately prior to the date of application, **AND**
- Patient must not have an eGFR equal to or greater than 30mL/min/1.73m², **AND**
- Patient must be male and must not have a height greater than or equal to 167.7 cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0 cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Population criteria:

- Patient must be prepubertal.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; **AND**
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; **AND**
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
5. The final adult height (in cm) of the patient's mother and father (where available); **AND**
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If a patient receiving treatment under the indication short stature due to chronic renal insufficiency undergoes a renal transplant and 12 months post-transplant has an eGFR of equal to or greater than 30mL/min/1.73m² prescribers should seek reclassification to the indication short stature and slow growth.

Authority required

Short stature and slow growth

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than short stature and slow growth, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have previously received treatment under the indication short stature associated with chronic renal insufficiency, have undergone a renal transplant and a 12 month period of observation following the transplant, and have

an estimated glomerular filtration rate of greater than or equal to 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula; OR

- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment and a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; AND
3. (a) A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment; OR
(b) Confirmation that the patient has previously received treatment under the indication **short stature associated with chronic renal insufficiency**, has undergone a renal transplant and a 12 month period of observation following the transplant, and has an estimated glomerular filtration rate of greater than or equal to 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula; AND
4. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
5. A bone age result performed within the last 12 months; AND
6. The proprietary name (brand), form and strength of somatotropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with biochemical growth hormone deficiency

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than short stature associated with biochemical growth hormone deficiency, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**

- Patient must have previously received treatment under the indication **risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants** and have reached or surpassed 5 years of age (chronological); OR
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment; OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and an annual growth velocity of 14 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and an annual growth velocity of 8 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; **AND**
3. (a) A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); OR
(b) Height and weight measurements from within three months prior to commencement of treatment for a patient whose height was at or below the 1st percentile for age and sex immediately prior to commencing treatment; OR
(c) Confirmation that the patient has previously received treatment under the indication **risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants** and has reached or surpassed 5 years of age (chronological); **AND**
4. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; **AND**

5. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
6. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND

7. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Biochemical growth hormone deficiency should not be secondary to an intracranial lesion or cranial irradiation for applications under this category.

Authority required

Growth retardation secondary to an intracranial lesion, or cranial irradiation

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than growth retardation secondary to an intracranial lesion, or cranial irradiation, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have had an intracranial lesion and have undergone a 12 month period of observation following completion of treatment for the intracranial lesion (all treatment); OR
- Patient must have had an intracranial lesion, have received medical advice that it is unsafe to treat the intracranial lesion, and have undergone a 12 month period of observation since initial diagnosis of the intracranial lesion; OR
- Patient must have received cranial irradiation without having had an intracranial lesion, and have undergone a 12 month period of observation following completion of treatment for the condition for which cranial irradiation was received, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment; OR
- Patient must have had both a height above the 1st percentile for age and sex immediately prior to commencing treatment and a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR

- Patient must have had both a height above the 1st percentile for age and sex immediately prior to commencing treatment and an annual growth velocity of 14 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had both a height above the 1st percentile for age and sex immediately prior to commencing treatment and an annual growth velocity of 8 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; **AND**
3. (a) A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); OR
(b) Height and weight measurements from within three months prior to commencement of treatment for a patient whose height was at or below the 1st percentile for age and sex immediately prior to commencing treatment; **AND**
4. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; **AND**
5. (a) Confirmation that the patient has had an intracranial lesion and has undergone a 12 month period of observation following completion of treatment for the intracranial lesion (all treatment); OR
(b) Confirmation that the patient has had an intracranial lesion, has received medical advice that it is unsafe to treat the intracranial lesion, and has undergone a 12 month period of observation since initial diagnosis of the intracranial lesion; OR
(c) Confirmation that the patient has received cranial irradiation without having had an intracranial lesion, and has undergone a 12 month period of observation following completion of treatment for the condition for which cranial irradiation was received; **AND**
6. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; **AND**
7. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
8. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

Authority required

Risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under a category other than risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies).

continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR

- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have a chronological age of less than 2 years, **AND**
- Patient must have a documented clinical risk of hypoglycaemia, **AND**
- Patient must have documented evidence that the risk of hypoglycaemia is secondary to biochemical growth hormone deficiency, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; **AND**
3. Confirmation that the patient has a documented clinical risk of hypoglycaemia; **AND**
4. Confirmation that the patient has documented evidence that the risk of hypoglycaemia is secondary to biochemical growth hormone deficiency; **AND**
5. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; **AND**
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Biochemical growth hormone deficiency and precocious puberty

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under a category other than biochemical growth hormone deficiency and precocious puberty, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must be male and have commenced puberty (demonstrated by Tanner stage 2 genital or pubic hair development or testicular volumes greater than or equal to 4 mL) before the chronological age of 9 years; OR
- Patient must be female and have commenced puberty (demonstrated by Tanner stage 2 breast or pubic hair development) before the chronological age of 8 years; OR
- Patient must be female and menarche occurred before the chronological age of 10 years, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR

- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must be undergoing Gonadotrophin Releasing Hormone agonist therapy for pubertal suppression, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; AND
3. Confirmation that the patient has precocious puberty; AND
4. Confirmation that the patient is undergoing Gonadotrophin Releasing Hormone agonist therapy for pubertal suppression; AND
5. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; AND
6. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
7. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
8. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies).

continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR

- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have a structural lesion that is not neoplastic; OR
- Patient must have had a structural lesion that was neoplastic and have undergone a 12 month period of observation following completion of treatment for the structural lesion (all treatment); OR
- Patient must have a structural lesion that is neoplastic, have received medical advice that it is unsafe to treat the structural lesion, and have undergone a 12 month period of observation since initial diagnosis of the structural lesion,

AND

- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels, **AND**
- Patient must have other hypothalamic/pituitary hormone deficits (includes ACTH, TSH, GnRH and/or vasopressin/ADH deficiencies), **AND**
- Patient must have hypothalamic obesity, **AND**
- Patient must have had a growth velocity above the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had an annual growth velocity of greater than 14 cm per year in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had an annual growth velocity of greater than 8 cm per year in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**

2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; AND
 3. A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); AND
 4. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; AND
 5. (a) Confirmation that the patient has a structural lesion that is not neoplastic; OR
(b) Confirmation that the patient had a structural lesion that was neoplastic and has undergone a 12 month period of observation following completion of treatment for the structural lesion (all treatment); OR
(c) Confirmation that the patient has a structural lesion that is neoplastic, has received medical advice that it is unsafe to treat the structural lesion, and has undergone a 12 month period of observation since initial diagnosis of the structural lesion; AND
 6. Confirmation that the patient has other hypothalamic/pituitary hormone deficits; AND
 7. Confirmation that the patient has hypothalamic obesity; AND
 8. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
 9. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
 10. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).
- Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with Turner syndrome

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under a category other than short stature associated with Turner syndrome, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have had a height at or below the 95th percentile for age on the Turner syndrome growth curve for girls immediately prior to commencing growth hormone treatment, **AND**
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as a loss of a whole X chromosome in all cells (45X), and gender of rearing is female; OR
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as a loss of a whole X chromosome in some cells (mosaic 46XX/45X), and gender of rearing is female; OR
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as genetic loss or rearrangement of an X chromosome (such as isochromosome X, ring-chromosome, or partial deletion of an X chromosome), and gender of rearing is female, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have a bone age of 2.5 years or less, **AND**
- Patient must not have a bone age of 13.5 years or greater, **AND**
- Patient must not have a height greater than or equal to 155.0 cm.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; AND
3. A height measurement from immediately prior to commencement of growth hormone treatment; AND
4. Confirmation that the patient has diagnostic results consistent with Turner syndrome; AND
5. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
6. A bone age result performed within the last 12 months; AND
7. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature due to short stature homeobox (SHOX) gene disorders

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than short stature due to short stature homeobox (SHOX) gene disorders, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as a karyotype confirming the presence of a SHOX mutation/deletion without the presence of mixed gonadal dysgenesis; OR
- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as mixed gonadal dysgenesis (45X mosaic karyotype with the presence of any Y chromosome material and/or SRY gene positive by FISH study) and have an appropriate plan of management in place for the patient's increased risk of gonadoblastoma, **AND**
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment, **AND**
- Patient must have had a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had an annual growth velocity of 14 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had an annual growth velocity of 8 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes (excluding gonadoblastoma secondary to mixed gonadal dysgenesis), **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; AND
3. A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); AND
4. Confirmation that the patient has diagnostic results consistent with a short stature homeobox (SHOX) gene disorder; AND
5. If the patient's condition is secondary to mixed gonadal dysgenesis, confirmation that an appropriate plan of management for the patient's increased risk of gonadoblastoma is in place; AND
6. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
7. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
8. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with chronic renal insufficiency

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than short stature associated with chronic renal insufficiency, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment; OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and a growth velocity less than or equal to the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and an annual growth velocity of 14 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and an annual growth velocity of 8 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**

- Patient must have an estimated glomerular filtration rate less than 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula, and not have undergone a renal transplant; OR
- Patient must have an estimated glomerular filtration rate less than 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula, have undergone a renal transplant, and have undergone a 12 month period of observation following the transplant, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Population criteria:

- Patient must be prepubertal.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; AND
3. (a) A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); OR
(b) Height and weight measurements from within three months prior to commencement of treatment for a patient whose height was at or below the 1st percentile for age and sex immediately prior to commencing treatment; AND
4. Confirmation that the patient has an estimated glomerular filtration rate less than 30mL/minute/1.73m² ; AND
5. If a renal transplant has taken place, confirmation that the patient has undergone a 12 month period of observation following transplantation; AND
6. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
7. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND

The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If a patient receiving treatment under the indication short stature due to chronic renal insufficiency undergoes a renal transplant and 12 months post-transplant has an eGFR of equal to or greater than 30mL/min/1.73m² prescribers should seek reclassification to the indication short stature and slow growth.

somatropin 24 mg injection [1 cartridge] (&) inert substance diluent [3.15 mL syringe], 1 pack

10476J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	1045.23	40.30	Humatrope [LY]

somatropin 6 mg injection [1 cartridge] (&) inert substance diluent [3.15 mL syringe], 1 pack

10482Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	266.85	40.30	Humatrope [LY]

somatropin 12 mg injection [1 cartridge] (&) inert substance diluent [3.15 mL syringe], 1 pack

10487Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	526.31	40.30	Humatrope [LY]

■ SOMATROPIN

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au
Applications for authority to prescribe should be forwarded to:
Department of Human Services
Prior Written Approval of Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Short stature and slow growth

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature and slow growth category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

Population criteria:

- Patient must be aged 3 years or older.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; **AND**
3. Recent growth data (height and weight, not older than three months); **AND**
4. A bone age result performed within the last 12 months; **AND**
5. The proprietary name (brand), form and strength of somatotropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Short stature associated with biochemical growth hormone deficiency

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature associated with biochemical growth hormone deficiency category, **AND**

- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

Population criteria:

- Patient must be aged 3 years or older.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; AND
3. Recent growth data (height and weight, not older than three months); AND
4. A bone age result performed within the last 12 months; AND
5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Growth retardation secondary to an intracranial lesion, or cranial irradiation

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the growth retardation secondary to an intracranial lesion, or cranial irradiation category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR

- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

Population criteria:

- Patient must be aged 3 years or older.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program)**

Special Arrangement 2015 and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; **AND**
3. Recent growth data (height and weight, not older than three months); **AND**
4. A bone age result performed within the last 12 months; **AND**
5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have a chronological age of 5 years or greater.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

Population criteria:

- Patient must be aged 3 years or older.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; AND
3. Recent growth data (height and weight, not older than three months); AND
4. A bone age result performed within the last 12 months; AND
5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Biochemical growth hormone deficiency and precocious puberty

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the biochemical growth hormone deficiency and precocious puberty category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must be undergoing Gonadotrophin Releasing Hormone agonist therapy for pubertal suppression, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

Population criteria:

- Patient must be aged 3 years or older.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; AND
3. Recent growth data (height and weight, not older than three months); AND
4. A bone age result performed within the last 12 months; AND
5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); **OR**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; **OR**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); **OR**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; **OR**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; **OR**
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; **OR**
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

Population criteria:

- Patient must be aged 3 years or older.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program)**

Special Arrangement 2015 and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; **AND**
3. Recent growth data (height and weight, not older than three months); **AND**
4. A bone age result performed within the last 12 months; **AND**
5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Short stature associated with Turner syndrome

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature associated with Turner syndrome category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**

- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be female and must not have a bone age of 13.5 years or more, **AND**
- Patient must be female and must not have a height greater than or equal to 155.0cm.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

Population criteria:

- Patient must be aged 3 years or older.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; **AND**
3. Recent growth data (height and weight, not older than three months); **AND**
4. A bone age result performed within the last 12 months; **AND**
5. The proprietary name (brand), form and strength of somatotropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Short stature due to short stature homeobox (SHOX) gene disorders

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature due to short stature homeobox (SHOX) gene disorders category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR

- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as a karyotype confirming the presence of a SHOX mutation/deletion without the presence of mixed gonadal dysgenesis; OR
- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as mixed gonadal dysgenesis (45X mosaic karyotype with the presence of any Y chromosome material and/or SRY gene positive by FISH study) and have an appropriate plan of management in place for the patient's increased risk of gonadoblastoma, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes (excluding gonadoblastoma secondary to mixed gonadal dysgenesis), **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm.

Population criteria:

- Patient must be aged 3 years or older.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program)**

Special Arrangement 2015 and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; **AND**
3. Recent growth data (height and weight, not older than three months); **AND**
4. A bone age result performed within the last 12 months; **AND**
5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Short stature associated with chronic renal insufficiency

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature associated with chronic renal insufficiency category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**

- Patient must not have undergone a renal transplant within the 12 month period immediately prior to the date of application, **AND**
- Patient must not have an eGFR equal to or greater than 30mL/min/1.73m², **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

Population criteria:

- Patient must be aged 3 years or older.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program)**

Special Arrangement 2015 and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; AND
3. Recent growth data (height and weight, not older than three months); AND
4. A bone age result performed within the last 12 months; AND
5. Confirmation that the patient has an estimated glomerular filtration rate less than 30mL/minute/1.73m²; AND
6. If a renal transplant has taken place, confirmation that the patient has undergone a 12 month period of observation following transplantation; AND
7. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

If a patient receiving treatment under the indication 'short stature associated with chronic renal insufficiency' undergoes a renal transplant and 12 months post-transplant has an eGFR of equal to or greater than 30mL/min/1.73m² prescribers should seek reclassification to the indication short stature and slow growth.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Short stature and slow growth

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than short stature and slow growth, **AND**
- Patient must have had a lapse in treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have previously received treatment under the indication short stature associated with chronic renal insufficiency, have undergone a renal transplant and a 12 month period of observation following the transplant, and have an estimated glomerular filtration rate of greater than or equal to 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula; OR
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment and a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment), **AND**

- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7 cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0 cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Population criteria:

- Patient must be aged 3 years or older.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program)**

Special Arrangement 2015 and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; **AND**
3. (a) A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment; **OR**
(b) Confirmation that the patient has previously received treatment under the indication short stature associated with chronic renal insufficiency, has undergone a renal transplant and a 12 month period of observation following the transplant, and has an estimated glomerular filtration rate of greater than or equal to 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula; **AND**
4. Recent growth data (height and weight, not older than three months); **AND**
5. A bone age result performed within the last 12 months; **AND**
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with biochemical growth hormone deficiency

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than short stature associated with biochemical growth hormone deficiency, **AND**
- Patient must have had a lapse in treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); **OR**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; **OR**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); **OR**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; **OR**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have previously received treatment under the indication **risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants** and have reached or surpassed 5 years of age (chronological); **OR**
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment; **OR**

- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and an annual growth velocity of 14 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and an annual growth velocity of 8 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Population criteria:

- Patient must be aged 3 years or older.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; **AND**
3. (a) A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); OR
(b) Height and weight measurements from within three months prior to commencement of treatment for a patient whose height was at or below the 1st percentile for age and sex immediately prior to commencing treatment; OR
(c) Confirmation that the patient has previously received treatment under the indication risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants and has reached or surpassed 5 years of age (chronological); **AND**
4. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; **AND**
5. Recent growth data (height and weight, not older than three months); **AND**
6. A bone age result performed within the last 12 months; **AND**

7. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Biochemical growth hormone deficiency should not be secondary to an intracranial lesion or cranial irradiation for applications under this category.

Authority required

Growth retardation secondary to an intracranial lesion, or cranial irradiation

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than growth retardation secondary to an intracranial lesion, or cranial irradiation, **AND**
- Patient must have had a lapse in treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have had an intracranial lesion and have undergone a 12 month period of observation following completion of treatment for the intracranial lesion (all treatment); OR
- Patient must have had an intracranial lesion, have received medical advice that it is unsafe to treat the intracranial lesion, and have undergone a 12 month period of observation since initial diagnosis of the intracranial lesion; OR
- Patient must have received cranial irradiation without having had an intracranial lesion, and have undergone a 12 month period of observation following completion of treatment for the condition for which cranial irradiation was received, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment; OR
- Patient must have had both a height above the 1st percentile for age and sex immediately prior to commencing treatment and a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had both a height above the 1st percentile for age and sex immediately prior to commencing treatment and an annual growth velocity of 14 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR

- Patient must have had both a height above the 1st percentile for age and sex immediately prior to commencing treatment and an annual growth velocity of 8 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Population criteria:

- Patient must be aged 3 years or older.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; **AND**
3. (a) A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); OR
(b) Height and weight measurements from within three months prior to commencement of treatment for a patient whose height was at or below the 1st percentile for age and sex immediately prior to commencing treatment; **AND**
4. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; **AND**
5. (a) Confirmation that the patient has had an intracranial lesion and has undergone a 12 month period of observation following completion of treatment for the intracranial lesion (all treatment); OR
(b) Confirmation that the patient has had an intracranial lesion, has received medical advice that it is unsafe to treat the intracranial lesion, and has undergone a 12 month period of observation since initial diagnosis of the intracranial lesion; OR
(c) Confirmation that the patient has received cranial irradiation without having had an intracranial lesion, and has undergone a 12 month period of observation following completion of treatment for the condition for which cranial irradiation was received; **AND**
6. Recent growth data (height and weight, not older than three months); **AND**
7. A bone age result performed within the last 12 months; **AND**
8. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Biochemical growth hormone deficiency and precocious puberty

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under a category other than biochemical growth hormone deficiency and precocious puberty, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a

continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR

- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must be male and have commenced puberty (demonstrated by Tanner stage 2 genital or pubic hair development or testicular volumes greater than or equal to 4 mL) before the chronological age of 9 years; OR
- Patient must be female and have commenced puberty (demonstrated by Tanner stage 2 breast or pubic hair development) before the chronological age of 8 years; OR
- Patient must be female and menarche occurred before the chronological age of 10 years, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must be undergoing Gonadotrophin Releasing Hormone agonist therapy for pubertal suppression, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Population criteria:

- Patient must be aged 3 years or older.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; **AND**
3. Confirmation that the patient has precocious puberty; **AND**
4. Confirmation that the patient is undergoing Gonadotrophin Releasing Hormone agonist therapy for pubertal suppression; **AND**
5. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; **AND**
6. Recent growth data (height and weight, not older than three months); **AND**
7. A bone age result performed within the last 12 months; **AND**
8. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth, **AND**
- Patient must have had a lapse in treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have a structural lesion that is not neoplastic; OR
- Patient must have had a structural lesion that was neoplastic and have undergone a 12 month period of observation following completion of treatment for the structural lesion (all treatment); OR
- Patient must have a structural lesion that is neoplastic, have received medical advice that it is unsafe to treat the structural lesion, and have undergone a 12 month period of observation since initial diagnosis of the structural lesion,

AND

- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must have other hypothalamic/pituitary hormone deficits (includes ACTH, TSH, GnRH and/or vasopressin/ADH deficiencies), **AND**
- Patient must have hypothalamic obesity, **AND**
- Patient must have had a growth velocity above the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had an annual growth velocity of greater than 14 cm per year in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had an annual growth velocity of greater than 8 cm per year in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Population criteria:

- Patient must be aged 3 years or older.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; AND
3. A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); AND
4. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; AND
5. (a) Confirmation that the patient has a structural lesion that is not neoplastic; OR
(b) Confirmation that the patient had a structural lesion that was neoplastic and has undergone a 12 month period of observation following completion of treatment for the structural lesion (all treatment); OR
(c) Confirmation that the patient has a structural lesion that is neoplastic, has received medical advice that it is unsafe to treat the structural lesion, and has undergone a 12 month period of observation since initial diagnosis of the structural lesion; AND
6. Confirmation that the patient has other hypothalamic/pituitary hormone deficits; AND
7. Confirmation that the patient has hypothalamic obesity; AND
8. Recent growth data (height and weight, not older than three months); AND
9. A bone age result performed within the last 12 months; AND
10. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with Turner syndrome

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under a category other than short stature associated with Turner syndrome, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have had a height at or below the 95th percentile for age on the Turner syndrome growth curve for girls immediately prior to commencing growth hormone treatment, **AND**
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as a loss of a whole X chromosome in all cells (45X), and gender of rearing is female; OR
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as a loss of a whole X chromosome in some cells (mosaic 46XX/45X), and gender of rearing is female; OR

- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as genetic loss or rearrangement of an X chromosome (such as isochromosome X, ring-chromosome, or partial deletion of an X chromosome), and gender of rearing is female, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have a bone age of 2.5 years or less, **AND**
- Patient must not have a height greater than or equal to 155.0 cm, **AND**
- Patient must not have a bone age of 13.5 years or greater.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

Population criteria:

- Patient must be aged 3 years or older.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program)**

Special Arrangement 2015 and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; AND
3. A height measurement from immediately prior to commencement of growth hormone treatment; AND
4. Confirmation that the patient has diagnostic results consistent with Turner syndrome; AND
5. Recent growth data (height and weight, not older than three months); AND
6. A bone age result performed within the last 12 months; AND

The proprietary name (brand), form and strength of somatotropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature due to short stature homeobox (SHOX) gene disorders

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than short stature due to short stature homeobox (SHOX) gene disorders, **AND**
- Patient must have had a lapse in treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as a karyotype confirming the presence of a SHOX mutation/deletion without the presence of mixed gonadal dysgenesis; OR
- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as mixed gonadal dysgenesis (45X mosaic karyotype with the presence of any Y chromosome material and/or SRY gene positive by FISH study) and have an appropriate plan of management in place for the patient's increased risk of gonadoblastoma, **AND**
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment, **AND**
- Patient must have had a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR

- Patient must have had an annual growth velocity of 14 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; **OR**
- Patient must have had an annual growth velocity of 8 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes (excluding gonadoblastoma secondary to mixed gonadal dysgenesis), **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; **OR**
- Patient must be female and must not have a height greater than or equal to 155.0cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; **OR**
- Patient must be female and must not have a bone age of 13.5 years or more.

Population criteria:

- Patient must be aged 3 years or older.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; **OR**
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program)**

Special Arrangement 2015 and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; **AND**
3. A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); **AND**
4. Confirmation that the patient has diagnostic results consistent with a short stature homeobox (SHOX) gene disorder; **AND**
5. If the patient's condition is secondary to mixed gonadal dysgenesis, confirmation that an appropriate plan of management for the patient's increased risk of gonadoblastoma is in place; **AND**
6. Recent growth data (height and weight, not older than three months); **AND**
7. A bone age result performed within the last 12 months; **AND**
8. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with chronic renal insufficiency

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than short stature associated with chronic renal insufficiency, **AND**
- Patient must have had a lapse in treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); **OR**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; **OR**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); **OR**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; **OR**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies).

continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**

- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment; **OR**
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and a growth velocity less than or equal to the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); **OR**
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and an annual growth velocity of 14 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; **OR**
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and an annual growth velocity of 8 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must have an estimated glomerular filtration rate less than 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula, and not have undergone a renal transplant; **OR**
- Patient must have an estimated glomerular filtration rate less than 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula, have undergone a renal transplant, and have undergone a 12 month period of observation following the transplant, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; **OR**
- Patient must be female and must not have a height greater than or equal to 155.0cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; **OR**
- Patient must be female and must not have a bone age of 13.5 years or more.

Population criteria:

- Patient must be aged 3 years or older.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; **OR**
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; **AND**
3. (a) A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); **OR**
(b) Height and weight measurements from within three months prior to commencement of treatment for a patient whose height was at or below the 1st percentile for age and sex immediately prior to commencing treatment; **AND**
4. Confirmation that the patient has an estimated glomerular filtration rate less than 30mL/minute/1.73m² ; **AND**
5. If a renal transplant has taken place, confirmation that the patient has undergone a 12 month period of observation following transplantation; **AND**
6. Recent growth data (height and weight, not older than three months); **AND**
7. A bone age result performed within the last 12 months; **AND**
8. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If a patient receiving treatment under the indication short stature due to chronic renal insufficiency undergoes a renal transplant and 12 months post-transplant has an eGFR of equal to or greater than 30mL/min/1.73m² prescribers should seek reclassification to the indication short stature and slow growth.

SYSTEMIC HORMONAL PREPARATIONS, EXCL. SEX HORMONES AND INSULINS

somatropin 5 mg/1.5 mL injection, 1.5 mL cartridge

10484T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	223.61	40.30	Scitropin A [SA]

somatropin 5 mg/1.5 mL injection, 1.5 mL cartridge

10512G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	223.61	40.30	Omnitrope Surepal 5 [SZ]

somatropin 10 mg/1.5 mL injection, 1.5 mL cartridge

10481P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	439.82	40.30	SciTropin A [SA]

somatropin 10 mg/1.5 mL injection, 1.5 mL cartridge

10519P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	439.82	40.30	Omnitrope Surepal 10 [SZ]

somatropin 15 mg/1.5 mL injection, 1.5 mL cartridge

10485W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	656.04	40.30	Omnitrope Surepal 15 [SZ]

■ SOMATROPIN

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Prior Written Approval of Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Short stature and slow growth

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature and slow growth category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; AND
3. Recent growth data (height and weight, not older than three months); AND
4. A bone age result performed within the last 12 months; AND
5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Short stature associated with biochemical growth hormone deficiency

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature associated with biochemical growth hormone deficiency category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; AND
3. Recent growth data (height and weight, not older than three months); AND
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Growth retardation secondary to an intracranial lesion, or cranial irradiation

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the growth retardation secondary to an intracranial lesion, or cranial irradiation category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program)**

Special Arrangement 2015 and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; **AND**
3. Recent growth data (height and weight, not older than three months); **AND**
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR

continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR

- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have a chronological age of 5 years or greater.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program)**

Special Arrangement 2015 and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; **AND**
3. Recent growth data (height and weight, not older than three months); **AND**
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Biochemical growth hormone deficiency and precocious puberty

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the biochemical growth hormone deficiency and precocious puberty category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must be undergoing Gonadotrophin Releasing Hormone agonist therapy for pubertal suppression, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; AND
3. Recent growth data (height and weight, not older than three months); AND
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; AND
3. Recent growth data (height and weight, not older than three months); AND
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND

5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Short stature associated with Turner syndrome

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature associated with Turner syndrome category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be female and must not have a bone age of 13.5 years or more, **AND**
- Patient must be female and must not have a height greater than or equal to 155.0cm.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; AND
3. Recent growth data (height and weight, not older than three months); AND
4. A bone age result performed within the last 12 months; AND
5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Short stature due to short stature homeobox (SHOX) gene disorders

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature due to short stature homeobox (SHOX) gene disorders category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**

- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as a karyotype confirming the presence of a SHOX mutation/deletion without the presence of mixed gonadal dysgenesis; OR
- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as mixed gonadal dysgenesis (45X mosaic karyotype with the presence of any Y chromosome material and/or SRY gene positive by FISH study) and have an appropriate plan of management in place for the patient's increased risk of gonadoblastoma, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes (excluding gonadoblastoma secondary to mixed gonadal dysgenesis), **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; **AND**
3. Recent growth data (height and weight, not older than three months); **AND**
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Short stature associated with chronic renal insufficiency

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature associated with chronic renal insufficiency category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**

continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR

- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have undergone a renal transplant within the 12 month period immediately prior to the date of application, **AND**
- Patient must not have an eGFR equal to or greater than 30mL/min/1.73m², **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

Population criteria:

- Patient must be prepubertal.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program)**

Special Arrangement 2015 and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; AND
3. Recent growth data (height and weight, not older than three months); AND
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
5. Confirmation that the patient has an estimated glomerular filtration rate less than 30mL/minute/1.73m²; AND
6. If a renal transplant has taken place, confirmation that the patient has undergone a 12 month period of observation following transplantation; AND
7. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

If a patient receiving treatment under the indication 'short stature associated with chronic renal insufficiency' undergoes a renal transplant and 12 months post-transplant has an eGFR of equal to or greater than 30mL/min/1.73m² prescribers should seek reclassification to the indication short stature and slow growth.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Short stature and slow growth

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than short stature and slow growth, **AND**
- Patient must have had a lapse in treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR

- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have previously received treatment under the indication short stature associated with chronic renal insufficiency, have undergone a renal transplant and a 12 month period of observation following the transplant, and have an estimated glomerular filtration rate of greater than or equal to 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula; OR
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment and a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7 cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0 cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; **AND**
3. (a) A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment; OR
(b) Confirmation that the patient has previously received treatment under the indication short stature associated with chronic renal insufficiency, has undergone a renal transplant and a 12 month period of observation following the transplant, and has an estimated glomerular filtration rate of greater than or equal to 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula; **AND**
4. Recent growth data (height and weight, not older than three months); **AND**
5. A bone age result performed within the last 12 months; **AND**
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with biochemical growth hormone deficiency

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than short stature associated with biochemical growth hormone deficiency, **AND**
- Patient must have had a lapse in treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR

continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR

- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have previously received treatment under the indication **risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants** and have reached or surpassed 5 years of age (chronological); OR
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment; OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and an annual growth velocity of 14 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and an annual growth velocity of 8 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; **AND**

3. (a) A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); OR
- (b) Height and weight measurements from within three months prior to commencement of treatment for a patient whose height was at or below the 1st percentile for age and sex immediately prior to commencing treatment; OR
- (c) Confirmation that the patient has previously received treatment under the indication risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants and has reached or surpassed 5 years of age (chronological); AND
4. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; AND
5. Recent growth data (height and weight, not older than three months); AND
6. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND

7. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Biochemical growth hormone deficiency should not be secondary to an intracranial lesion or cranial irradiation for applications under this category.

Authority required

Growth retardation secondary to an intracranial lesion, or cranial irradiation

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than growth retardation secondary to an intracranial lesion, or cranial irradiation, **AND**
- Patient must have had a lapse in treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have had an intracranial lesion and have undergone a 12 month period of observation following completion of treatment for the intracranial lesion (all treatment); OR
- Patient must have had an intracranial lesion, have received medical advice that it is unsafe to treat the intracranial lesion, and have undergone a 12 month period of observation since initial diagnosis of the intracranial lesion; OR
- Patient must have received cranial irradiation without having had an intracranial lesion, and have undergone a 12 month period of observation following completion of treatment for the condition for which cranial irradiation was received, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR

- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment; **OR**
- Patient must have had both a height above the 1st percentile for age and sex immediately prior to commencing treatment and a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); **OR**
- Patient must have had both a height above the 1st percentile for age and sex immediately prior to commencing treatment and an annual growth velocity of 14 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; **OR**
- Patient must have had both a height above the 1st percentile for age and sex immediately prior to commencing treatment and an annual growth velocity of 8 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; **OR**
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; **OR**
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; **AND**
3. (a) A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); **OR**
(b) Height and weight measurements from within three months prior to commencement of treatment for a patient whose height was at or below the 1st percentile for age and sex immediately prior to commencing treatment; **AND**
4. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; **AND**
5. (a) Confirmation that the patient has had an intracranial lesion and has undergone a 12 month period of observation following completion of treatment for the intracranial lesion (all treatment); **OR**
(b) Confirmation that the patient has had an intracranial lesion, has received medical advice that it is unsafe to treat the intracranial lesion, and has undergone a 12 month period of observation since initial diagnosis of the intracranial lesion; **OR**
(c) Confirmation that the patient has received cranial irradiation without having had an intracranial lesion, and has undergone a 12 month period of observation following completion of treatment for the condition for which cranial irradiation was received; **AND**
6. Recent growth data (height and weight, not older than three months); **AND**
7. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
8. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under a category other than risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**

- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have a chronological age of less than 2 years, **AND**
- Patient must have a documented clinical risk of hypoglycaemia, **AND**
- Patient must have documented evidence that the risk of hypoglycaemia is secondary to biochemical growth hormone deficiency, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; **AND**
3. Confirmation that the patient has a documented clinical risk of hypoglycaemia; **AND**
4. Confirmation that the patient has documented evidence that the risk of hypoglycaemia is secondary to biochemical growth hormone deficiency; **AND**
5. Recent growth data (height and weight, not older than three months); **AND**
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Biochemical growth hormone deficiency and precocious puberty

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under a category other than biochemical growth hormone deficiency and precocious puberty, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR

- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must be male and have commenced puberty (demonstrated by Tanner stage 2 genital or pubic hair development or testicular volumes greater than or equal to 4 mL) before the chronological age of 9 years; OR
- Patient must be female and have commenced puberty (demonstrated by Tanner stage 2 breast or pubic hair development) before the chronological age of 8 years; OR
- Patient must be female and menarche occurred before the chronological age of 10 years, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must be undergoing Gonadotrophin Releasing Hormone agonist therapy for pubertal suppression, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; AND
3. Confirmation that the patient has precocious puberty; AND
4. Confirmation that the patient is undergoing Gonadotrophin Releasing Hormone agonist therapy for pubertal suppression; AND
5. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; AND
6. Recent growth data (height and weight, not older than three months); AND
7. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
8. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth, **AND**
 - Patient must have had a lapse in treatment, **AND**
 - The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
 - The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
 - The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
 - The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
 - The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
 - Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
 - Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
 - Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
 - Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
 - Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels, **AND**
 - Patient must have a structural lesion that is not neoplastic; OR
 - Patient must have had a structural lesion that was neoplastic and have undergone a 12 month period of observation following completion of treatment for the structural lesion (all treatment); OR
 - Patient must have a structural lesion that is neoplastic, have received medical advice that it is unsafe to treat the structural lesion, and have undergone a 12 month period of observation since initial diagnosis of the structural lesion, **AND**
 - Patient must have other hypothalamic/pituitary hormone deficits (includes ACTH, TSH, GnRH and/or vasopressin/ADH deficiencies), **AND**
 - Patient must have hypothalamic obesity, **AND**
 - Patient must have had a growth velocity above the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
 - Patient must have had an annual growth velocity of greater than 14 cm per year in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
 - Patient must have had an annual growth velocity of greater than 8 cm per year in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
 - Patient must not have diabetes mellitus, **AND**
 - Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
 - Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
 - Patient must be male and must not have a bone age of 15.5 years or more; OR
 - Patient must be female and must not have a bone age of 13.5 years or more.
- Treatment criteria:**

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; AND
3. A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); AND
4. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; AND
5. (a) Confirmation that the patient has a structural lesion that is not neoplastic; OR
(b) Confirmation that the patient had a structural lesion that was neoplastic and has undergone a 12 month period of observation following completion of treatment for the structural lesion (all treatment); OR
(c) Confirmation that the patient has a structural lesion that is neoplastic, has received medical advice that it is unsafe to treat the structural lesion, and has undergone a 12 month period of observation since initial diagnosis of the structural lesion; AND
6. Confirmation that the patient has other hypothalamic/pituitary hormone deficits; AND
7. Confirmation that the patient has hypothalamic obesity; AND
8. Recent growth data (height and weight, not older than three months); AND
9. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
10. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with Turner syndrome

Treatment Phase: Recommencement of treatment as a reclassified patient

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under a category other than short stature associated with Turner syndrome, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have had a height at or below the 95th percentile for age on the Turner syndrome growth curve for girls immediately prior to commencing growth hormone treatment, **AND**

- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as a loss of a whole X chromosome in all cells (45X), and gender of rearing is female; OR
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as a loss of a whole X chromosome in some cells (mosaic 46XX/45X), and gender of rearing is female; OR
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as genetic loss or rearrangement of an X chromosome (such as isochromosome X, ring-chromosome, or partial deletion of an X chromosome), and gender of rearing is female, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have a bone age of 2.5 years or less, **AND**
- Patient must not have a height greater than or equal to 155.0 cm, **AND**
- Patient must not have a bone age of 13.5 years or greater.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; **AND**
3. A height measurement from immediately prior to commencement of growth hormone treatment; **AND**
4. Confirmation that the patient has diagnostic results consistent with Turner syndrome; **AND**
5. Recent growth data (height and weight, not older than three months); **AND**
6. A bone age result performed within the last 12 months; **AND**

The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature due to short stature homeobox (SHOX) gene disorders

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than short stature due to short stature homeobox (SHOX) gene disorders, **AND**
- Patient must have had a lapse in treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as a karyotype confirming the presence of a SHOX mutation/deletion without the presence of mixed gonadal dysgenesis; OR
- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as mixed gonadal dysgenesis (45X mosaic karyotype with the presence of any Y chromosome material and/or SRY gene positive by FISH study) and have an appropriate plan of management in place for the patient's increased risk of gonadoblastoma, **AND**
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment, **AND**
- Patient must have had a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had an annual growth velocity of 14 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR

- Patient must have had an annual growth velocity of 8 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes (excluding gonadoblastoma secondary to mixed gonadal dysgenesis), **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program)**

Special Arrangement 2015 and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; **AND**
3. A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); **AND**
4. Confirmation that the patient has diagnostic results consistent with a short stature homeobox (SHOX) gene disorder; **AND**
5. If the patient's condition is secondary to mixed gonadal dysgenesis, confirmation that an appropriate plan of management for the patient's increased risk of gonadoblastoma is in place; **AND**
6. Recent growth data (height and weight, not older than three months); **AND**
7. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
8. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with chronic renal insufficiency

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than short stature associated with chronic renal insufficiency, **AND**
- Patient must have had a lapse in treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment; OR

- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and a growth velocity less than or equal to the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and an annual growth velocity of 14 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and an annual growth velocity of 8 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must have an estimated glomerular filtration rate less than 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula, and not have undergone a renal transplant; OR
- Patient must have an estimated glomerular filtration rate less than 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula, have undergone a renal transplant, and have undergone a 12 month period of observation following the transplant, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Population criteria:

- Patient must be prepubertal.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program)**

Special Arrangement 2015 and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; **AND**
3. (a) A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); OR
(b) Height and weight measurements from within three months prior to commencement of treatment for a patient whose height was at or below the 1st percentile for age and sex immediately prior to commencing treatment; **AND**
4. Confirmation that the patient has an estimated glomerular filtration rate less than 30mL/minute/1.73m² ; **AND**
5. If a renal transplant has taken place, confirmation that the patient has undergone a 12 month period of observation following transplantation; **AND**
6. Recent growth data (height and weight, not older than three months); **AND**
7. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
8. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If a patient receiving treatment under the indication short stature due to chronic renal insufficiency undergoes a renal transplant and 12 months post-transplant has an eGFR of equal to or greater than 30mL/min/1.73m² prescribers should seek reclassification to the indication short stature and slow growth.

somatropin 24 mg injection [1 cartridge] (&) inert substance diluent [3.15 mL syringe], 1 pack

10502R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	1045.23	40.30	Humatrope [LY]

somatropin 6 mg injection [1 cartridge] (&) inert substance diluent [3.15 mL syringe], 1 pack

10429X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	266.85	40.30	Humatrope [LY]

somatropin 12 mg injection [1 cartridge] (&) inert substance diluent [3.15 mL syringe], 1 pack

10461N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	526.31	40.30	Humatrope [LY]

■ SOMATROPIN

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Prior Written Approval of Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Short stature and slow growth

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature and slow growth category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; **AND**
3. Recent growth data (height and weight, not older than three months); **AND**
4. A bone age result performed within the last 12 months; **AND**
5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Short stature associated with biochemical growth hormone deficiency

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature associated with biochemical growth hormone deficiency category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; **AND**
3. Recent growth data (height and weight, not older than three months); **AND**
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Growth retardation secondary to an intracranial lesion, or cranial irradiation

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the growth retardation secondary to an intracranial lesion, or cranial irradiation category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR

- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; **AND**
3. Recent growth data (height and weight, not older than three months); **AND**
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must not have diabetes mellitus, **AND**

- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have a chronological age of 5 years or greater.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; AND
3. Recent growth data (height and weight, not older than three months); AND
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Biochemical growth hormone deficiency and precocious puberty

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the biochemical growth hormone deficiency and precocious puberty category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must be undergoing Gonadotrophin Releasing Hormone agonist therapy for pubertal suppression, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND

2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; AND
3. Recent growth data (height and weight, not older than three months); AND
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program)**

Special Arrangement 2015 and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; AND
3. Recent growth data (height and weight, not older than three months); AND
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Short stature associated with Turner syndrome

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature associated with Turner syndrome category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be female and must not have a bone age of 13.5 years or more, **AND**
- Patient must be female and must not have a height greater than or equal to 155.0cm.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; AND
3. Recent growth data (height and weight, not older than three months); AND
4. A bone age result performed within the last 12 months; AND
5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Short stature due to short stature homeobox (SHOX) gene disorders

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature due to short stature homeobox (SHOX) gene disorders category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR

- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as a karyotype confirming the presence of a SHOX mutation/deletion without the presence of mixed gonadal dysgenesis; OR
- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as mixed gonadal dysgenesis (45X mosaic karyotype with the presence of any Y chromosome material and/or SRY gene positive by FISH study) and have an appropriate plan of management in place for the patient's increased risk of gonadoblastoma, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes (excluding gonadoblastoma secondary to mixed gonadal dysgenesis), **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; **AND**
3. Recent growth data (height and weight, not older than three months); **AND**
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Short stature associated with chronic renal insufficiency

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature associated with chronic renal insufficiency category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**

- Patient must not have undergone a renal transplant within the 12 month period immediately prior to the date of application, **AND**
- Patient must not have an eGFR equal to or greater than 30mL/min/1.73m², **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; **AND**
3. Recent growth data (height and weight, not older than three months); **AND**
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
5. Confirmation that the patient has an estimated glomerular filtration rate less than 30mL/minute/1.73m²; **AND**
6. If a renal transplant has taken place, confirmation that the patient has undergone a 12 month period of observation following transplantation; **AND**
7. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

If a patient receiving treatment under the indication 'short stature associated with chronic renal insufficiency' undergoes a renal transplant and 12 months post-transplant has an eGFR of equal to or greater than 30mL/min/1.73m² prescribers should seek reclassification to the indication short stature and slow growth.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Short stature and slow growth

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than short stature and slow growth, **AND**
- Patient must have had a lapse in treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have previously received treatment under the indication short stature associated with chronic renal insufficiency, have undergone a renal transplant and a 12 month period of observation following the transplant, and have an estimated glomerular filtration rate of greater than or equal to 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula; OR
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment and a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment), **AND**
- Patient must not have diabetes mellitus, **AND**

- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7 cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0 cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; AND
3. (a) A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment; OR
(b) Confirmation that the patient has previously received treatment under the indication short stature associated with chronic renal insufficiency, has undergone a renal transplant and a 12 month period of observation following the transplant, and has an estimated glomerular filtration rate of greater than or equal to 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula; AND
4. Recent growth data (height and weight, not older than three months); AND
5. A bone age result performed within the last 12 months; AND
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with biochemical growth hormone deficiency

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than short stature associated with biochemical growth hormone deficiency, **AND**
- Patient must have had a lapse in treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have previously received treatment under the indication **risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants** and have reached or surpassed 5 years of age (chronological); OR
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment; OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR

- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and an annual growth velocity of 14 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and an annual growth velocity of 8 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program)**

Special Arrangement 2015 and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; **AND**
3. (a) A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); OR
(b) Height and weight measurements from within three months prior to commencement of treatment for a patient whose height was at or below the 1st percentile for age and sex immediately prior to commencing treatment; OR
(c) Confirmation that the patient has previously received treatment under the indication risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants and has reached or surpassed 5 years of age (chronological); **AND**
4. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; **AND**
5. Recent growth data (height and weight, not older than three months); **AND**
6. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
7. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Biochemical growth hormone deficiency should not be secondary to an intracranial lesion or cranial irradiation for applications under this category.

Authority required

Growth retardation secondary to an intracranial lesion, or cranial irradiation

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than growth retardation secondary to an intracranial lesion, or cranial irradiation, **AND**
- Patient must have had a lapse in treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have had an intracranial lesion and have undergone a 12 month period of observation following completion of treatment for the intracranial lesion (all treatment); OR
- Patient must have had an intracranial lesion, have received medical advice that it is unsafe to treat the intracranial lesion, and have undergone a 12 month period of observation since initial diagnosis of the intracranial lesion; OR
- Patient must have received cranial irradiation without having had an intracranial lesion, and have undergone a 12 month period of observation following completion of treatment for the condition for which cranial irradiation was received, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment; OR
- Patient must have had both a height above the 1st percentile for age and sex immediately prior to commencing treatment and a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had both a height above the 1st percentile for age and sex immediately prior to commencing treatment and an annual growth velocity of 14 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had both a height above the 1st percentile for age and sex immediately prior to commencing treatment and an annual growth velocity of 8 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**

- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; AND
3. (a) A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); OR
(b) Height and weight measurements from within three months prior to commencement of treatment for a patient whose height was at or below the 1st percentile for age and sex immediately prior to commencing treatment; AND
4. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; AND
5. (a) Confirmation that the patient has had an intracranial lesion and has undergone a 12 month period of observation following completion of treatment for the intracranial lesion (all treatment); OR
(b) Confirmation that the patient has had an intracranial lesion, has received medical advice that it is unsafe to treat the intracranial lesion, and has undergone a 12 month period of observation since initial diagnosis of the intracranial lesion; OR
(c) Confirmation that the patient has received cranial irradiation without having had an intracranial lesion, and has undergone a 12 month period of observation following completion of treatment for the condition for which cranial irradiation was received; AND
6. Recent growth data (height and weight, not older than three months); AND
7. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
8. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under a category other than risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have a chronological age of less than 2 years, **AND**
- Patient must have a documented clinical risk of hypoglycaemia, **AND**

- Patient must have documented evidence that the risk of hypoglycaemia is secondary to biochemical growth hormone deficiency, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; AND
3. Confirmation that the patient has a documented clinical risk of hypoglycaemia; AND
4. Confirmation that the patient has documented evidence that the risk of hypoglycaemia is secondary to biochemical growth hormone deficiency; AND
5. Recent growth data (height and weight, not older than three months); AND
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Biochemical growth hormone deficiency and precocious puberty

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under a category other than biochemical growth hormone deficiency and precocious puberty, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must be male and have commenced puberty (demonstrated by Tanner stage 2 genital or pubic hair development or testicular volumes greater than or equal to 4 mL) before the chronological age of 9 years; OR
- Patient must be female and have commenced puberty (demonstrated by Tanner stage 2 breast or pubic hair development) before the chronological age of 8 years; OR
- Patient must be female and menarche occurred before the chronological age of 10 years, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or

absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR

- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must be undergoing Gonadotrophin Releasing Hormone agonist therapy for pubertal suppression, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; **AND**
3. Confirmation that the patient has precocious puberty; **AND**
4. Confirmation that the patient is undergoing Gonadotrophin Releasing Hormone agonist therapy for pubertal suppression; **AND**
5. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; **AND**
6. Recent growth data (height and weight, not older than three months); **AND**
7. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
8. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth, **AND**
- Patient must have had a lapse in treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**

- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must have a structural lesion that is not neoplastic; OR
- Patient must have had a structural lesion that was neoplastic and have undergone a 12 month period of observation following completion of treatment for the structural lesion (all treatment); OR
- Patient must have a structural lesion that is neoplastic, have received medical advice that it is unsafe to treat the structural lesion, and have undergone a 12 month period of observation since initial diagnosis of the structural lesion, **AND**
- Patient must have other hypothalamic/pituitary hormone deficits (includes ACTH, TSH, GnRH and/or vasopressin/ADH deficiencies), **AND**
- Patient must have hypothalamic obesity, **AND**
- Patient must have had a growth velocity above the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had an annual growth velocity of greater than 14 cm per year in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had an annual growth velocity of greater than 8 cm per year in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; **AND**
3. A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); **AND**
4. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; **AND**
5. (a) Confirmation that the patient has a structural lesion that is not neoplastic; OR

- (b) Confirmation that the patient had a structural lesion that was neoplastic and has undergone a 12 month period of observation following completion of treatment for the structural lesion (all treatment); OR
- (c) Confirmation that the patient has a structural lesion that is neoplastic, has received medical advice that it is unsafe to treat the structural lesion, and has undergone a 12 month period of observation since initial diagnosis of the structural lesion; AND
- 6. Confirmation that the patient has other hypothalamic/pituitary hormone deficits; AND
- 7. Confirmation that the patient has hypothalamic obesity; AND
- 8. Recent growth data (height and weight, not older than three months); AND
- 9. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
- 10. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with Turner syndrome

Treatment Phase: Recommencement of treatment as a reclassified patient

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under a category other than short stature associated with Turner syndrome, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have had a height at or below the 95th percentile for age on the Turner syndrome growth curve for girls immediately prior to commencing growth hormone treatment, **AND**
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as a loss of a whole X chromosome in all cells (45X), and gender of rearing is female; OR
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as a loss of a whole X chromosome in some cells (mosaic 46XX/45X), and gender of rearing is female; OR
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as genetic loss or rearrangement of an X chromosome (such as isochromosome X, ring-chromosome, or partial deletion of an X chromosome), and gender of rearing is female, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have a bone age of 2.5 years or less, **AND**
- Patient must not have a height greater than or equal to 155.0 cm, **AND**
- Patient must not have a bone age of 13.5 years or greater.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program)**

Special Arrangement 2015 and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; AND

3. A height measurement from immediately prior to commencement of growth hormone treatment; AND
4. Confirmation that the patient has diagnostic results consistent with Turner syndrome; AND
5. Recent growth data (height and weight, not older than three months); AND
6. A bone age result performed within the last 12 months; AND

The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature due to short stature homeobox (SHOX) gene disorders

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than short stature due to short stature homeobox (SHOX) gene disorders, **AND**
- Patient must have had a lapse in treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as a karyotype confirming the presence of a SHOX mutation/deletion without the presence of mixed gonadal dysgenesis; OR
- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as mixed gonadal dysgenesis (45X mosaic karyotype with the presence of any Y chromosome material and/or SRY gene positive by FISH study) and have an appropriate plan of management in place for the patient's increased risk of gonadoblastoma, **AND**
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment, **AND**
- Patient must have had a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had an annual growth velocity of 14 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had an annual growth velocity of 8 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes (excluding gonadoblastoma secondary to mixed gonadal dysgenesis), **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program)**

Special Arrangement 2015 and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; AND
3. A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); AND
4. Confirmation that the patient has diagnostic results consistent with a short stature homeobox (SHOX) gene disorder; AND
5. If the patient's condition is secondary to mixed gonadal dysgenesis, confirmation that an appropriate plan of management for the patient's increased risk of gonadoblastoma is in place; AND
6. Recent growth data (height and weight, not older than three months); AND
7. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
8. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with chronic renal insufficiency

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than short stature associated with chronic renal insufficiency, **AND**
- Patient must have had a lapse in treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment; OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and a growth velocity less than or equal to the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and an annual growth velocity of 14 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and an annual growth velocity of 8 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must have an estimated glomerular filtration rate less than 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula, and not have undergone a renal transplant; OR
- Patient must have an estimated glomerular filtration rate less than 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula, have undergone a renal transplant, and have undergone a 12 month period of observation following the transplant, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR

- Patient must be female and must not have a height greater than or equal to 155.0cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; AND
3. (a) A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); OR
(b) Height and weight measurements from within three months prior to commencement of treatment for a patient whose height was at or below the 1st percentile for age and sex immediately prior to commencing treatment; AND
4. Confirmation that the patient has an estimated glomerular filtration rate less than 30mL/minute/1.73m²; AND
5. If a renal transplant has taken place, confirmation that the patient has undergone a 12 month period of observation following transplantation; AND
6. Recent growth data (height and weight, not older than three months); AND
7. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
8. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If a patient receiving treatment under the indication short stature due to chronic renal insufficiency undergoes a renal transplant and 12 months post-transplant has an eGFR of equal to or greater than 30mL/min/1.73m² prescribers should seek reclassification to the indication short stature and slow growth.

somatropin 5 mg/1.5 mL injection, 1.5 mL cartridge

10437H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	223.61	40.30	Norditropin SimpleXx [NO]

somatropin 5 mg/1.5 mL injection, 1.5 mL cartridge

10467X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	223.61	40.30	Norditropin FlexPro [NO]

somatropin 10 mg/1.5 mL injection, 1.5 mL cartridge

10448X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	439.82	40.30	Norditropin SimpleXx [NO]

somatropin 10 mg/1.5 mL injection, 1.5 mL cartridge

10496K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	439.82	40.30	Norditropin FlexPro [NO]

somatropin 10 mg/2 mL injection, 2 mL cartridge

10438J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	426.41	40.30	NutropinAq [IS]

somatropin 15 mg/1.5 mL injection, 1.5 mL cartridge

10470C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	656.04	40.30	Norditropin SimpleXx [NO]

somatropin 15 mg/1.5 mL injection, 1.5 mL cartridge

10489C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	656.04	40.30	Norditropin FlexPro [NO]

■ SOMATROPIN

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Prior Written Approval of Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Short stature and slow growth

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature and slow growth category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; **AND**
3. Recent growth data (height and weight, not older than three months); **AND**
4. A bone age result performed within the last 12 months; **AND**
5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Short stature associated with biochemical growth hormone deficiency

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature associated with biochemical growth hormone deficiency category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; AND
3. Recent growth data (height and weight, not older than three months); AND
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Growth retardation secondary to an intracranial lesion, or cranial irradiation

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the growth retardation secondary to an intracranial lesion, or cranial irradiation category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR

- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; **AND**
3. Recent growth data (height and weight, not older than three months); **AND**
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have a chronological age of 5 years or greater.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; AND
3. Recent growth data (height and weight, not older than three months); AND
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Biochemical growth hormone deficiency and precocious puberty

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the biochemical growth hormone deficiency and precocious puberty category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must be undergoing Gonadotrophin Releasing Hormone agonist therapy for pubertal suppression, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; AND
3. Recent growth data (height and weight, not older than three months); AND
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program)**

Special Arrangement 2015 and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; **AND**
3. Recent growth data (height and weight, not older than three months); **AND**
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Short stature associated with Turner syndrome

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature associated with Turner syndrome category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR

- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be female and must not have a bone age of 13.5 years or more, **AND**
- Patient must be female and must not have a height greater than or equal to 155.0cm.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; **AND**
3. Recent growth data (height and weight, not older than three months); **AND**
4. A bone age result performed within the last 12 months; **AND**
5. The proprietary name (brand), form and strength of somatotropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Short stature due to short stature homeobox (SHOX) gene disorders

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature due to short stature homeobox (SHOX) gene disorders category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as a karyotype confirming the presence of a SHOX mutation/deletion without the presence of mixed gonadal dysgenesis; OR
- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as mixed gonadal dysgenesis (45X mosaic karyotype with the presence of any Y chromosome material and/or SRY gene positive by FISH study) and have an appropriate plan of management in place for the patient's increased risk of gonadoblastoma, **AND**

- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes (excluding gonadoblastoma secondary to mixed gonadal dysgenesis), **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; **AND**
3. Recent growth data (height and weight, not older than three months); **AND**
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Short stature and slow growth

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than short stature and slow growth, **AND**
- Patient must have had a lapse in treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have previously received treatment under the indication short stature associated with chronic renal insufficiency, have undergone a renal transplant and a 12 month period of observation following the transplant, and have an estimated glomerular filtration rate of greater than or equal to 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula; OR
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment and a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7 cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0 cm, **AND**

- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; AND
3. (a) A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment; OR
(b) Confirmation that the patient has previously received treatment under the indication short stature associated with chronic renal insufficiency, has undergone a renal transplant and a 12 month period of observation following the transplant, and has an estimated glomerular filtration rate of greater than or equal to 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula; AND
4. Recent growth data (height and weight, not older than three months); AND
5. A bone age result performed within the last 12 months; AND
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with biochemical growth hormone deficiency

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than short stature associated with biochemical growth hormone deficiency, **AND**
- Patient must have had a lapse in treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have previously received treatment under the indication **risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants** and have reached or surpassed 5 years of age (chronological); OR
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment; OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and an annual growth velocity of 14 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR

- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and an annual growth velocity of 8 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); **OR**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); **OR**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); **OR**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; **OR**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; **OR**
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; **OR**
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program)**

Special Arrangement 2015 and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; **AND**
3. (a) A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); **OR**
(b) Height and weight measurements from within three months prior to commencement of treatment for a patient whose height was at or below the 1st percentile for age and sex immediately prior to commencing treatment; **OR**
(c) Confirmation that the patient has previously received treatment under the indication risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants and has reached or surpassed 5 years of age (chronological); **AND**
4. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; **AND**
5. Recent growth data (height and weight, not older than three months); **AND**
6. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
7. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Biochemical growth hormone deficiency should not be secondary to an intracranial lesion or cranial irradiation for applications under this category.

Authority required

Growth retardation secondary to an intracranial lesion, or cranial irradiation

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than growth retardation secondary to an intracranial lesion, or cranial irradiation, **AND**
- Patient must have had a lapse in treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have had an intracranial lesion and have undergone a 12 month period of observation following completion of treatment for the intracranial lesion (all treatment); OR
- Patient must have had an intracranial lesion, have received medical advice that it is unsafe to treat the intracranial lesion, and have undergone a 12 month period of observation since initial diagnosis of the intracranial lesion; OR
- Patient must have received cranial irradiation without having had an intracranial lesion, and have undergone a 12 month period of observation following completion of treatment for the condition for which cranial irradiation was received, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment; OR
- Patient must have had both a height above the 1st percentile for age and sex immediately prior to commencing treatment and a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had both a height above the 1st percentile for age and sex immediately prior to commencing treatment and an annual growth velocity of 14 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had both a height above the 1st percentile for age and sex immediately prior to commencing treatment and an annual growth velocity of 8 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program)**

Special Arrangement 2015 and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; AND
3. (a) A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); OR
(b) Height and weight measurements from within three months prior to commencement of treatment for a patient whose height was at or below the 1st percentile for age and sex immediately prior to commencing treatment; AND
4. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; AND
5. (a) Confirmation that the patient has had an intracranial lesion and has undergone a 12 month period of observation following completion of treatment for the intracranial lesion (all treatment); OR
(b) Confirmation that the patient has had an intracranial lesion, has received medical advice that it is unsafe to treat the intracranial lesion, and has undergone a 12 month period of observation since initial diagnosis of the intracranial lesion; OR
(c) Confirmation that the patient has received cranial irradiation without having had an intracranial lesion, and has undergone a 12 month period of observation following completion of treatment for the condition for which cranial irradiation was received; AND
6. Recent growth data (height and weight, not older than three months); AND
7. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
8. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under a category other than risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have a chronological age of less than 2 years, **AND**
- Patient must have a documented clinical risk of hypoglycaemia, **AND**
- Patient must have documented evidence that the risk of hypoglycaemia is secondary to biochemical growth hormone deficiency, **AND**
- Patient must not have diabetes mellitus, **AND**

- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; AND
3. Confirmation that the patient has a documented clinical risk of hypoglycaemia; AND
4. Confirmation that the patient has documented evidence that the risk of hypoglycaemia is secondary to biochemical growth hormone deficiency; AND
5. Recent growth data (height and weight, not older than three months); AND
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Biochemical growth hormone deficiency and precocious puberty

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under a category other than biochemical growth hormone deficiency and precocious puberty, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must be male and have commenced puberty (demonstrated by Tanner stage 2 genital or pubic hair development or testicular volumes greater than or equal to 4 mL) before the chronological age of 9 years; OR
- Patient must be female and have commenced puberty (demonstrated by Tanner stage 2 breast or pubic hair development) before the chronological age of 8 years; OR
- Patient must be female and menarche occurred before the chronological age of 10 years, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR

- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; **AND**
- Patient must be undergoing Gonadotrophin Releasing Hormone agonist therapy for pubertal suppression, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; **AND**
3. Confirmation that the patient has precocious puberty; **AND**
4. Confirmation that the patient is undergoing Gonadotrophin Releasing Hormone agonist therapy for pubertal suppression; **AND**
5. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; **AND**
6. Recent growth data (height and weight, not older than three months); **AND**
7. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
8. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth, **AND**
- Patient must have had a lapse in treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR

- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must have a structural lesion that is not neoplastic; OR
- Patient must have had a structural lesion that was neoplastic and have undergone a 12 month period of observation following completion of treatment for the structural lesion (all treatment); OR
- Patient must have a structural lesion that is neoplastic, have received medical advice that it is unsafe to treat the structural lesion, and have undergone a 12 month period of observation since initial diagnosis of the structural lesion, **AND**
- Patient must have other hypothalamic/pituitary hormone deficits (includes ACTH, TSH, GnRH and/or vasopressin/ADH deficiencies), **AND**
- Patient must have hypothalamic obesity, **AND**
- Patient must have had a growth velocity above the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had an annual growth velocity of greater than 14 cm per year in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had an annual growth velocity of greater than 8 cm per year in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; **AND**
3. A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); **AND**
4. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; **AND**
5. (a) Confirmation that the patient has a structural lesion that is not neoplastic; OR
(b) Confirmation that the patient had a structural lesion that was neoplastic and has undergone a 12 month period of observation following completion of treatment for the structural lesion (all treatment); OR

(c) Confirmation that the patient has a structural lesion that is neoplastic, has received medical advice that it is unsafe to treat the structural lesion, and has undergone a 12 month period of observation since initial diagnosis of the structural lesion; AND

6. Confirmation that the patient has other hypothalamic/pituitary hormone deficits; AND

7. Confirmation that the patient has hypothalamic obesity; AND

8. Recent growth data (height and weight, not older than three months); AND

9. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND

10. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with Turner syndrome

Treatment Phase: Recommencement of treatment as a reclassified patient

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under a category other than short stature associated with Turner syndrome, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have had a height at or below the 95th percentile for age on the Turner syndrome growth curve for girls immediately prior to commencing growth hormone treatment, **AND**
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as a loss of a whole X chromosome in all cells (45X), and gender of rearing is female; OR
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as a loss of a whole X chromosome in some cells (mosaic 46XX/45X), and gender of rearing is female; OR
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as genetic loss or rearrangement of an X chromosome (such as isochromosome X, ring-chromosome, or partial deletion of an X chromosome), and gender of rearing is female, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have a bone age of 2.5 years or less, **AND**
- Patient must not have a height greater than or equal to 155.0 cm, **AND**
- Patient must not have a bone age of 13.5 years or greater.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program)**

Special Arrangement 2015 and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; AND
3. A height measurement from immediately prior to commencement of growth hormone treatment; AND
4. Confirmation that the patient has diagnostic results consistent with Turner syndrome; AND

5. Recent growth data (height and weight, not older than three months); AND

6. A bone age result performed within the last 12 months; AND

The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature due to short stature homeobox (SHOX) gene disorders

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than short stature due to short stature homeobox (SHOX) gene disorders, **AND**
- Patient must have had a lapse in treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as a karyotype confirming the presence of a SHOX mutation/deletion without the presence of mixed gonadal dysgenesis; OR
- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as mixed gonadal dysgenesis (45X mosaic karyotype with the presence of any Y chromosome material and/or SRY gene positive by FISH study) and have an appropriate plan of management in place for the patient's increased risk of gonadoblastoma, **AND**
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment, **AND**
- Patient must have had a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had an annual growth velocity of 14 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had an annual growth velocity of 8 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes (excluding gonadoblastoma secondary to mixed gonadal dysgenesis), **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND

2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; AND
 3. A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); AND
 4. Confirmation that the patient has diagnostic results consistent with a short stature homeobox (SHOX) gene disorder; AND
 5. If the patient's condition is secondary to mixed gonadal dysgenesis, confirmation that an appropriate plan of management for the patient's increased risk of gonadoblastoma is in place; AND
 6. Recent growth data (height and weight, not older than three months); AND
 7. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
 8. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).
- Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with chronic renal insufficiency

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature associated with chronic renal insufficiency category, AND
- Patient must have had a lapse in growth hormone treatment, AND
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, AND
- Patient must not have diabetes mellitus, AND
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, AND
- Patient must not have an active tumour or evidence of tumour growth or activity, AND
- Patient must not have undergone a renal transplant within the 12 month period immediately prior to the date of application, AND
- Patient must not have an eGFR equal to or greater than 30mL/min/1.73m², AND
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more, AND
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

Population criteria:

- Patient must be aged 3 years or older.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program)**

Special Arrangement 2015 and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; AND
3. Recent growth data (height and weight, not older than three months); AND

4. A bone age result performed within the last 12 months; AND
5. Confirmation that the patient has an estimated glomerular filtration rate less than 30mL/minute/1.73m² ; AND
6. If a renal transplant has taken place, confirmation that the patient has undergone a 12 month period of observation following transplantation; AND
7. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

If a patient receiving treatment under the indication 'short stature associated with chronic renal insufficiency' undergoes a renal transplant and 12 months post-transplant has an eGFR of equal to or greater than 30mL/min/1.73m² prescribers should seek reclassification to the indication short stature and slow growth.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Short stature associated with chronic renal insufficiency

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than short stature associated with chronic renal insufficiency, **AND**
- Patient must have had a lapse in treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment; OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and a growth velocity less than or equal to the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and an annual growth velocity of 14 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and an annual growth velocity of 8 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must have an estimated glomerular filtration rate less than 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula, and not have undergone a renal transplant; OR
- Patient must have an estimated glomerular filtration rate less than 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula, have undergone a renal transplant, and have undergone a 12 month period of observation following the transplant, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Population criteria:

- Patient must be aged 3 years or older.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; AND
3. (a) A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); OR
(b) Height and weight measurements from within three months prior to commencement of treatment for a patient whose height was at or below the 1st percentile for age and sex immediately prior to commencing treatment; AND
4. Confirmation that the patient has an estimated glomerular filtration rate less than 30mL/minute/1.73m²; AND
5. If a renal transplant has taken place, confirmation that the patient has undergone a 12 month period of observation following transplantation; AND
6. Recent growth data (height and weight, not older than three months); AND
7. A bone age result performed within the last 12 months; AND
8. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If a patient receiving treatment under the indication short stature due to chronic renal insufficiency undergoes a renal transplant and 12 months post-transplant has an eGFR of equal to or greater than 30mL/min/1.73m² prescribers should seek reclassification to the indication short stature and slow growth.

somatropin 12 mg/1.5 mL injection, 1.5 mL cartridge

10495J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	526.31	40.30	Saizen [SG]

somatropin 6 mg/1.03 mL injection, 1.03 mL cartridge

10458K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	266.85	40.30	Saizen [SG]

somatropin 20 mg/2.5 mL injection, 2.5 mL cartridge

10442N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	872.25	40.30	Saizen [SG]

■ SOMATROPIN

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Prior Written Approval of Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Short stature and slow growth

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature and slow growth category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR

- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; **AND**
3. Recent growth data (height and weight, not older than three months); **AND**
4. A bone age result performed within the last 12 months; **AND**
5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Short stature associated with biochemical growth hormone deficiency

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature associated with biochemical growth hormone deficiency category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**

- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; AND
3. Recent growth data (height and weight, not older than three months); AND
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Growth retardation secondary to an intracranial lesion, or cranial irradiation

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the growth retardation secondary to an intracranial lesion, or cranial irradiation category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; AND
3. Recent growth data (height and weight, not older than three months); AND
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have a chronological age of 5 years or greater.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program)**

Special Arrangement 2015 and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; AND
3. Recent growth data (height and weight, not older than three months); AND
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Biochemical growth hormone deficiency and precocious puberty

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the biochemical growth hormone deficiency and precocious puberty category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must be undergoing Gonadotrophin Releasing Hormone agonist therapy for pubertal suppression, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; **AND**
3. Recent growth data (height and weight, not older than three months); **AND**
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**

continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR

- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; **AND**
3. Recent growth data (height and weight, not older than three months); **AND**
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Short stature associated with Turner syndrome

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature associated with Turner syndrome category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be female and must not have a bone age of 13.5 years or more, **AND**
- Patient must be female and must not have a height greater than or equal to 155.0cm.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; AND
3. Recent growth data (height and weight, not older than three months); AND
4. A bone age result performed within the last 12 months; AND
5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Short stature due to short stature homeobox (SHOX) gene disorders

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature due to short stature homeobox (SHOX) gene disorders category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as a karyotype confirming the presence of a SHOX mutation/deletion without the presence of mixed gonadal dysgenesis; OR
- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as mixed gonadal dysgenesis (45X mosaic karyotype with the presence of any Y chromosome material and/or SRY gene positive by FISH study) and have an appropriate plan of management in place for the patient's increased risk of gonadoblastoma, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes (excluding gonadoblastoma secondary to mixed gonadal dysgenesis), **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; AND
3. Recent growth data (height and weight, not older than three months); AND

4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Short stature associated with chronic renal insufficiency

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature associated with chronic renal insufficiency category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have undergone a renal transplant within the 12 month period immediately prior to the date of application, **AND**
- Patient must not have an eGFR equal to or greater than 30mL/min/1.73m², **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; AND
3. Recent growth data (height and weight, not older than three months); AND
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
5. Confirmation that the patient has an estimated glomerular filtration rate less than 30mL/minute/1.73m²; AND
6. If a renal transplant has taken place, confirmation that the patient has undergone a 12 month period of observation following transplantation; AND
7. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

If a patient receiving treatment under the indication 'short stature associated with chronic renal insufficiency' undergoes a renal transplant and 12 months post-transplant has an eGFR of equal to or greater than 30mL/min/1.73m² prescribers should seek reclassification to the indication short stature and slow growth.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Short stature and poor body composition due to Prader-Willi syndrome

Treatment Phase: Recommencement of treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature and poor body composition due to Prader Willi syndrome category, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- Patient must have had a bone age below skeletal maturity (15.5 years for males and 13.5 years for females) (except where the patient had a chronological age of 2.5 years or less) at the last application and treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have had a bone age below skeletal maturity (15.5 years for males and 13.5 years for females) (except where the patient had a chronological age of 2.5 years or less) at the last application and treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- Patient must have had a bone age below skeletal maturity (15.5 years for males and 13.5 years for females) (except where the patient had a chronological age of 2.5 years or less) at the last application and treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- Patient must have had a bone age below skeletal maturity (15.5 years for males and 13.5 years for females) (except where the patient had a chronological age of 2.5 years or less) at the last application and treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- Patient must have had a bone age below skeletal maturity (15.5 years for males and 13.5 years for females) (except where the patient had a chronological age of 2.5 years or less) at the last application and treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems; OR
- Patient must have had a bone age at or above skeletal maturity (15.5 years for males and 13.5 years for females) at the last application and treatment must not have lapsed due to failure to respond to growth hormone at a dose of 0.04mg/kg/wk or greater for the most recent treatment period (32 weeks for the initial treatment period or 26 weeks for subsequent treatment periods, whichever applies); OR
- Patient must have had a bone age at or above skeletal maturity (15.5 years for males and 13.5 years for females) at the last application and treatment must not have lapsed due to failure to respond to growth hormone at a dose of 0.04mg/kg/wk or greater for the most recent treatment period (32 weeks for the initial treatment period or 26 weeks for subsequent treatment periods, whichever applies), unless response was affected by a significant medical illness; OR
- Patient must have had a bone age at or above skeletal maturity (15.5 years for males and 13.5 years for females) at the last application and treatment must not have lapsed due to failure to respond to growth hormone at a dose of 0.04mg/kg/wk or greater for the most recent treatment period (32 weeks for the initial treatment period or 26 weeks for subsequent treatment periods, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- Patient must have had a bone age at or above skeletal maturity (15.5 years for males and 13.5 years for females) at the last application and treatment must not have lapsed due to failure to respond to growth hormone at a dose of 0.04mg/kg/wk or greater for the most recent treatment period (32 weeks for the initial treatment period or 26 weeks for subsequent treatment periods, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- Patient must have had a bone age at or above skeletal maturity (15.5 years for males and 13.5 years for females) at the last application and treatment must not have lapsed due to failure to respond to growth hormone at a dose of 0.04mg/kg/wk or greater for the most recent treatment period (32 weeks for the initial treatment period or 26 weeks for subsequent treatment periods, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have been re-evaluated via polysomnography for airway obstruction and apnoea during the initial 32 week treatment period and any sleep disorders identified that required treatment must have been addressed, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have developed uncontrolled morbid obesity, defined as a body weight greater than 200% of ideal body weight for height and sex, with ideal body weight derived by calculating the 50th percentile weight for the patient's current height.

Population criteria:

- Patient must not have a chronological age of equal to or greater than 18 years.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment; AND
3. Recent growth data (height, weight, and waist circumference, not older than three months); AND
4. The date at which skeletal maturity was achieved (if applicable) [Note: In patients whose chronological age is greater than 2.5 years, a bone age reading should be performed at least once every 12 months prior to attainment of skeletal maturity.]; AND
5. Confirmation that during the initial 32 week treatment period, the patient was re-evaluated via polysomnography for airway obstruction and apnoea, and any sleep disorders that were identified have been addressed; AND
6. The proprietary name (brand), form and strength of somatotropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If recommencement of treatment is sought under a different indication than that under which the patient was previously receiving treatment an application for **recommencement of treatment as a reclassified patient** should be submitted.

Authority required

Short stature and slow growth

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than short stature and slow growth, **AND**
- Patient must have had a lapse in treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have previously received treatment under the indication short stature associated with chronic renal insufficiency, have undergone a renal transplant and a 12 month period of observation following the transplant, and have an estimated glomerular filtration rate of greater than or equal to 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula; OR
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment and a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7 cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0 cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program)**

Special Arrangement 2015 and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; AND
3. (a) A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment; OR
(b) Confirmation that the patient has previously received treatment under the indication short stature associated with chronic renal insufficiency, has undergone a renal transplant and a 12 month period of observation following the transplant, and has an estimated glomerular filtration rate of greater than or equal to 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula; AND
4. Recent growth data (height and weight, not older than three months); AND
5. A bone age result performed within the last 12 months; AND
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with biochemical growth hormone deficiency

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than short stature associated with biochemical growth hormone deficiency, **AND**
- Patient must have had a lapse in treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have previously received treatment under the indication **risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants** and have reached or surpassed 5 years of age (chronological); OR
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment; OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and an annual growth velocity of 14 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and an annual growth velocity of 8 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**

- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; **AND**
3. (a) A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); OR
(b) Height and weight measurements from within three months prior to commencement of treatment for a patient whose height was at or below the 1st percentile for age and sex immediately prior to commencing treatment; OR
(c) Confirmation that the patient has previously received treatment under the indication risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants and has reached or surpassed 5 years of age (chronological); **AND**
4. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; **AND**
5. Recent growth data (height and weight, not older than three months); **AND**
6. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
7. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Biochemical growth hormone deficiency should not be secondary to an intracranial lesion or cranial irradiation for applications under this category.

Authority required

Growth retardation secondary to an intracranial lesion, or cranial irradiation

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than growth retardation secondary to an intracranial lesion, or cranial irradiation, **AND**
- Patient must have had a lapse in treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have had an intracranial lesion and have undergone a 12 month period of observation following completion of treatment for the intracranial lesion (all treatment); OR
- Patient must have had an intracranial lesion, have received medical advice that it is unsafe to treat the intracranial lesion, and have undergone a 12 month period of observation since initial diagnosis of the intracranial lesion; OR
- Patient must have received cranial irradiation without having had an intracranial lesion, and have undergone a 12 month period of observation following completion of treatment for the condition for which cranial irradiation was received, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment; OR
- Patient must have had both a height above the 1st percentile for age and sex immediately prior to commencing treatment and a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had both a height above the 1st percentile for age and sex immediately prior to commencing treatment and an annual growth velocity of 14 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had both a height above the 1st percentile for age and sex immediately prior to commencing treatment and an annual growth velocity of 8 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; AND
3. (a) A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); OR
(b) Height and weight measurements from within three months prior to commencement of treatment for a patient whose height was at or below the 1st percentile for age and sex immediately prior to commencing treatment; AND
4. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; AND
5. (a) Confirmation that the patient has had an intracranial lesion and has undergone a 12 month period of observation following completion of treatment for the intracranial lesion (all treatment); OR
(b) Confirmation that the patient has had an intracranial lesion, has received medical advice that it is unsafe to treat the intracranial lesion, and has undergone a 12 month period of observation since initial diagnosis of the intracranial lesion; OR
(c) Confirmation that the patient has received cranial irradiation without having had an intracranial lesion, and has undergone a 12 month period of observation following completion of treatment for the condition for which cranial irradiation was received; AND
6. Recent growth data (height and weight, not older than three months); AND
7. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
8. The proprietary name (brand), form and strength of somatotropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under a category other than risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have a chronological age of less than 2 years, **AND**
- Patient must have a documented clinical risk of hypoglycaemia, **AND**
- Patient must have documented evidence that the risk of hypoglycaemia is secondary to biochemical growth hormone deficiency, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; AND
3. Confirmation that the patient has a documented clinical risk of hypoglycaemia; AND
4. Confirmation that the patient has documented evidence that the risk of hypoglycaemia is secondary to biochemical growth hormone deficiency; AND
5. Recent growth data (height and weight, not older than three months); AND
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Biochemical growth hormone deficiency and precocious puberty

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under a category other than biochemical growth hormone deficiency and precocious puberty, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must be male and have commenced puberty (demonstrated by Tanner stage 2 genital or pubic hair development or testicular volumes greater than or equal to 4 mL) before the chronological age of 9 years; OR
- Patient must be female and have commenced puberty (demonstrated by Tanner stage 2 breast or pubic hair development) before the chronological age of 8 years; OR
- Patient must be female and menarche occurred before the chronological age of 10 years, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR

- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must be undergoing Gonadotrophin Releasing Hormone agonist therapy for pubertal suppression, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; AND
3. Confirmation that the patient has precocious puberty; AND
4. Confirmation that the patient is undergoing Gonadotrophin Releasing Hormone agonist therapy for pubertal suppression; AND
5. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; AND
6. Recent growth data (height and weight, not older than three months); AND
7. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
8. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth, **AND**
- Patient must have had a lapse in treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR

- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels; **AND**
- Patient must have a structural lesion that is not neoplastic; OR
- Patient must have had a structural lesion that was neoplastic and have undergone a 12 month period of observation following completion of treatment for the structural lesion (all treatment); OR
- Patient must have a structural lesion that is neoplastic, have received medical advice that it is unsafe to treat the structural lesion, and have undergone a 12 month period of observation since initial diagnosis of the structural lesion;

AND

- Patient must have other hypothalamic/pituitary hormone deficits (includes ACTH, TSH, GnRH and/or vasopressin/ADH deficiencies); **AND**
- Patient must have hypothalamic obesity; **AND**
- Patient must have had a growth velocity above the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had an annual growth velocity of greater than 14 cm per year in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had an annual growth velocity of greater than 8 cm per year in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment; **AND**
- Patient must not have diabetes mellitus; **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes; **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity; **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; **AND**
3. A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); **AND**
4. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; **AND**
5. (a) Confirmation that the patient has a structural lesion that is not neoplastic; OR
(b) Confirmation that the patient had a structural lesion that was neoplastic and has undergone a 12 month period of observation following completion of treatment for the structural lesion (all treatment); OR
(c) Confirmation that the patient has a structural lesion that is neoplastic, has received medical advice that it is unsafe to treat the structural lesion, and has undergone a 12 month period of observation since initial diagnosis of the structural lesion; **AND**
6. Confirmation that the patient has other hypothalamic/pituitary hormone deficits; **AND**
7. Confirmation that the patient has hypothalamic obesity; **AND**

8. Recent growth data (height and weight, not older than three months); AND
9. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
10. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with Turner syndrome

Treatment Phase: Recommencement of treatment as a reclassified patient

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under a category other than short stature associated with Turner syndrome, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have had a height at or below the 95th percentile for age on the Turner syndrome growth curve for girls immediately prior to commencing growth hormone treatment, **AND**
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as a loss of a whole X chromosome in all cells (45X), and gender of rearing is female; OR
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as a loss of a whole X chromosome in some cells (mosaic 46XX/45X), and gender of rearing is female; OR
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as genetic loss or rearrangement of an X chromosome (such as isochromosome X, ring-chromosome, or partial deletion of an X chromosome), and gender of rearing is female, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have a bone age of 2.5 years or less, **AND**
- Patient must not have a height greater than or equal to 155.0 cm, **AND**
- Patient must not have a bone age of 13.5 years or greater.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program)**

Special Arrangement 2015 and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; AND
3. A height measurement from immediately prior to commencement of growth hormone treatment; AND
4. Confirmation that the patient has diagnostic results consistent with Turner syndrome; AND
5. Recent growth data (height and weight, not older than three months); AND
6. A bone age result performed within the last 12 months; AND

The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature due to short stature homeobox (SHOX) gene disorders

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than short stature due to short stature homeobox (SHOX) gene disorders, **AND**
- Patient must have had a lapse in treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as a karyotype confirming the presence of a SHOX mutation/deletion without the presence of mixed gonadal dysgenesis; OR
- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as mixed gonadal dysgenesis (45X mosaic karyotype with the presence of any Y chromosome material and/or SRY gene positive by FISH study) and have an appropriate plan of management in place for the patient's increased risk of gonadoblastoma, **AND**
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment, **AND**
- Patient must have had a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had an annual growth velocity of 14 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had an annual growth velocity of 8 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes (excluding gonadoblastoma secondary to mixed gonadal dysgenesis), **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program)**

Special Arrangement 2015 and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; **AND**
3. A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12

months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); AND

4. Confirmation that the patient has diagnostic results consistent with a short stature homeobox (SHOX) gene disorder; AND

5. If the patient's condition is secondary to mixed gonadal dysgenesis, confirmation that an appropriate plan of management for the patient's increased risk of gonadoblastoma is in place; AND

6. Recent growth data (height and weight, not older than three months); AND

7. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND

8. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Authority required

Short stature associated with chronic renal insufficiency

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than short stature associated with chronic renal insufficiency, **AND**
- Patient must have had a lapse in treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment; OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and a growth velocity less than or equal to the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and an annual growth velocity of 14 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and an annual growth velocity of 8 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must have an estimated glomerular filtration rate less than 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula, and not have undergone a renal transplant; OR
- Patient must have an estimated glomerular filtration rate less than 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula, have undergone a renal transplant, and have undergone a 12 month period of observation following the transplant, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; AND
3. (a) A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); OR
(b) Height and weight measurements from within three months prior to commencement of treatment for a patient whose height was at or below the 1st percentile for age and sex immediately prior to commencing treatment; AND
4. Confirmation that the patient has an estimated glomerular filtration rate less than 30mL/minute/1.73m²; AND
5. If a renal transplant has taken place, confirmation that the patient has undergone a 12 month period of observation following transplantation; AND
6. Recent growth data (height and weight, not older than three months); AND
7. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
8. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If a patient receiving treatment under the indication short stature due to chronic renal insufficiency undergoes a renal transplant and 12 months post-transplant has an eGFR of equal to or greater than 30mL/min/1.73m² prescribers should seek reclassification to the indication short stature and slow growth.

Authority required

Short stature and poor body composition due to Prader-Willi syndrome

Treatment Phase: Recommencement of treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under a category other than short stature and poor body composition due to Prader-Willi syndrome, **AND**
- Patient must have had a lapse in growth hormone treatment, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have diagnostic results consistent with Prader-Willi syndrome (the condition must be genetically proven); OR
- Patient must have a clinical diagnosis of Prader-Willi syndrome, confirmed by a clinical geneticist, **AND**
- Patient must have been evaluated via polysomnography for airway obstruction and apnoea whilst on growth hormone treatment and any sleep disorders identified that required treatment must have been addressed; OR
- Patient must have been evaluated via polysomnography for airway obstruction and apnoea within the last 12 months with no sleep disorders identified; OR
- Patient must have been evaluated via polysomnography for airway obstruction and apnoea within the last 12 months with sleep disorders identified which are not of sufficient severity to require treatment; OR
- Patient must have been evaluated via polysomnography for airway obstruction and apnoea within the last 12 months with sleep disorders identified for which the patient is currently receiving ameliorative treatment, **AND**

- Patient must not have uncontrolled morbid obesity, defined as a body weight greater than 200% of ideal body weight for height and sex, with ideal body weight derived by calculating the 50th percentile weight for the patient's current height, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have a chronological age of 18 years or greater.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each recommencement treatment phase is 32 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for recommencement of treatment as a reclassified patient; AND
3. (a) Confirmation that the patient has diagnostic results consistent with Prader-Willi syndrome, OR
(b) Confirmation that the patient has a clinical diagnosis of Prader-Willi syndrome, confirmed by a clinical geneticist; AND
4. Confirmation that the patient has been evaluated via polysomnography for airway obstruction and apnoea whilst on growth hormone treatment or within the last 12 months, and any sleep disorders identified via the polysomnography that required treatment have been addressed; AND
5. Recent growth data (height and weight, not older than three months); AND
6. The date at which skeletal maturity was achieved (if applicable) [Note: In patients whose chronological age is greater than 2.5 years, a bone age reading should be performed at least once every 12 months prior to attainment of skeletal maturity]; AND
7. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 16 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

somatropin 2 mg injection [7] (&) inert substance diluent [7 x 0.25 mL syringes], 1 pack

10472E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	612.79	40.30	Genotropin MiniQuick [PF]

somatropin 1.6 mg injection [7] (&) inert substance diluent [7 x 0.25 mL syringes], 1 pack

10498M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	491.72	40.30	Genotropin MiniQuick [PF]

somatropin 1.4 mg injection [7] (&) inert substance diluent [7 x 0.25 mL syringes], 1 pack

10434E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	431.17	40.30	Genotropin MiniQuick [PF]

somatropin 1.8 mg injection [7] (&) inert substance diluent [7 x 0.25 mL syringes], 1 pack

10501Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	552.26	40.30	Genotropin MiniQuick [PF]

SOMATROPIN (Recombinant human growth hormone) Powder for injection 12 mg (36 i.u.) with diluent in pre-filled pen (with preservative), 1

10426R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	510.20	40.30	Genotropin GoQuick [PF]

SOMATROPIN (Recombinant human growth hormone) Powder for injection 5 mg (15 i.u.) with diluent in pre-filled pen (with preservative), 1

10435F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	216.90	40.30	Genotropin GoQuick [PF]

somatropin 800 microgram injection [7] (&) inert substance diluent [7 x 0.25 mL syringes], 1 pack

10463Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	249.55	40.30	Genotropin MiniQuick [PF]

somatropin 1.2 mg injection [7] (&) inert substance diluent [7 x 0.25 mL syringes], 1 pack

10457J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	370.63	40.30	Genotropin MiniQuick [PF]

somatropin 600 microgram injection [7] (&) inert substance diluent [7 x 0.25 mL syringes], 1 pack

10477K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	189.02	40.30	Genotropin MiniQuick [PF]

somatropin 400 microgram injection [7] (&) inert substance diluent [7 x 0.25 mL syringes], 1 pack

10908D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	128.47	40.30	Genotropin MiniQuick [PF]

somatropin 1 mg injection [7] (&) inert substance diluent [7 x 0.25 mL syringes], 1 pack

10430Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	310.09	40.30	Genotropin MiniQuick [PF]

■ SOMATROPIN**Authority required**

Short stature and slow growth

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature and slow growth category, **AND**
- Patient must not have been on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved and maintained mid parental height standard deviation score while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; AND
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
4. A bone age result performed within the last 12 months; AND
5. The final adult height (in cm) of the patient's mother and father (where available); AND
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Prior Written Approval of Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Short stature associated with biochemical growth hormone deficiency

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature associated with biochemical growth hormone deficiency category, **AND**
- Patient must not have been on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved and maintained mid parental height standard deviation score while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; **AND**
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; **AND**
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
5. The final adult height (in cm) of the patient's mother and father (where available); **AND**
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Prior Written Approval of Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Growth retardation secondary to an intracranial lesion, or cranial irradiation

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the growth retardation secondary to an intracranial lesion, or cranial irradiation category, **AND**
- Patient must not have been on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR

- Patient must have achieved and maintained mid parental height standard deviation score while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; **AND**
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; **AND**
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
5. The final adult height (in cm) of the patient's mother and father (where available); **AND**
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Prior Written Approval of Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants category, **AND**
- Patient must not have been on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved and maintained mid parental height standard deviation score while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have a chronological age of 5 years or greater.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; **AND**

3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND

5. The final adult height (in cm) of the patient's mother and father (where available); AND

6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

When a patient receiving treatment under the indication risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants reaches or surpasses 5 years of age (chronological), prescribers should seek reclassification to the indication 'short stature due to biochemical growth hormone deficiency'.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Prior Written Approval of Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Biochemical growth hormone deficiency and precocious puberty

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the biochemical growth hormone deficiency and precocious puberty category, **AND**
- Patient must not have been on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved and maintained mid parental height standard deviation score while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; AND
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
5. The final adult height (in cm) of the patient's mother and father (where available); AND
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:
Department of Human Services
Prior Written Approval of Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth category, **AND**
- Patient must not have been on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved and maintained mid parental height standard deviation score while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; **AND**
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; **AND**
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
5. The final adult height (in cm) of the patient's mother and father (where available); **AND**
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Prior Written Approval of Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Short stature associated with Turner syndrome

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature associated with Turner syndrome category, **AND**
- Patient must not have been on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR

- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an annualised growth velocity for bone age at or above the mean growth velocity for untreated Turner Syndrome girls (using the Turner Syndrome - Ranke growth velocity chart) while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have a bone age of 13.5 years or greater, **AND**
- Patient must not have a height greater than or equal to 155.0 cm.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; **AND**
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; **AND**
4. A bone age result performed within the last 12 months; **AND**
5. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Prior Written Approval of Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Short stature due to short stature homeobox (SHOX) gene disorders

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature due to short stature homeobox (SHOX) gene disorders category, **AND**
- Patient must not have been on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved and maintained mid parental height standard deviation score while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes (excluding gonadoblastoma secondary to mixed gonadal dysgenesis), **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7 cm; OR

- Patient must be female and must not have a height greater than or equal to 155.0 cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; **AND**
3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; **AND**
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**
5. The final adult height (in cm) of the patient's mother and father (where available); **AND**
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Prior Written Approval of Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Short stature associated with chronic renal insufficiency

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature associated with chronic renal insufficiency category, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have been on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved the 50th percentile growth velocity for bone age and sex while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved an increase in height standard deviation score for chronological age and sex while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved a minimum growth velocity of 4cm/year while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have achieved and maintained mid parental height standard deviation score while on the maximum dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have undergone a renal transplant within the 12 month period immediately prior to the date of application, **AND**
- Patient must not have an eGFR equal to or greater than 30mL/min/1.73m², **AND**
- Patient must be male and must not have a height greater than or equal to 167.7 cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0 cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; **AND**

3. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
4. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
5. The final adult height (in cm) of the patient's mother and father (where available); AND
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

- Note** If a patient receiving treatment under the indication short stature due to chronic renal insufficiency undergoes a renal transplant and 12 months post-transplant has an eGFR of equal to or greater than 30mL/min/1.73m² prescribers should seek reclassification to the indication short stature and slow growth.
- Note** Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au. Applications for authority to prescribe should be forwarded to:
Department of Human Services
Prior Written Approval of Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Short stature and poor body composition due to Prader-Willi syndrome

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under the short stature and poor body composition due to Prader-Willi syndrome category, **AND**
- Patient must have been re-evaluated via polysomnography for airway obstruction and apnoea during the initial 32 week treatment period and any sleep disorders identified that required treatment must have been addressed, **AND**
- Patient must have had a bone age below skeletal maturity (15.5 years for males and 13.5 years for females) (except where the patient had a chronological age of 2.5 years or less) at the last application and must not have been on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or commencement treatment period and 26 weeks for a continuing treatment period, whichever applies; OR
- Patient must have had a bone age below skeletal maturity (15.5 years for males and 13.5 years for females) (except where the patient had a chronological age of 2.5 years or less) at the last application and must have maintained or improved height percentile for age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or commencement treatment period and 26 weeks for a continuing treatment period, whichever applies; OR
- Patient must have had a bone age below skeletal maturity (15.5 years for males and 13.5 years for females) (except where the patient had a chronological age of 2.5 years or less) at the last application and must have maintained or improved body mass index SDS for age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or commencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have had a bone age below skeletal maturity (15.5 years for males and 13.5 years for females) (except where the patient had a chronological age of 2.5 years or less) at the last application and must have maintained or improved waist circumference while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or commencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have had a bone age below skeletal maturity (15.5 years for males and 13.5 years for females) (except where the patient had a chronological age of 2.5 years or less) at the last application and must have maintained or improved waist/height ratio (waist circumference in centimetres divided by height in centimetres) while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or commencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have had a bone age below skeletal maturity (15.5 years for males and 13.5 years for females) (except where the patient had a chronological age of 2.5 years or less) at the last application and must have achieved an increase in height percentile with reference to the untreated Prader-Willi syndrome standards for age and sex while on the maximum dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or commencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have had a bone age at or above skeletal maturity (15.5 years for males and 13.5 years for females) at the last application and must not have been on the maximum dose of 0.04mg/kg/week or greater for the most recent treatment period (32 weeks for an initial or commencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have had a bone age at or above skeletal maturity (15.5 years for males and 13.5 years for females) at the last application and must have maintained or improved body mass index while on the maximum dose of 0.04mg/kg/week or greater for the most recent treatment period (32 weeks for an initial or commencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have had a bone age at or above skeletal maturity (15.5 years for males and 13.5 years for females) at the last application and must have maintained or improved body mass index SDS for age and sex while on the maximum dose of 0.04mg/kg/week or greater for the most recent treatment period (32 weeks for an initial or commencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR

- Patient must have had a bone age at or above skeletal maturity (15.5 years for males and 13.5 years for females) at the last application and must have maintained or improved waist circumference while on the maximum dose of 0.04mg/kg/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have had a bone age at or above skeletal maturity (15.5 years for males and 13.5 years for females) at the last application and must have maintained or improved waist/height ratio (waist circumference in centimetres divided by height in centimetres) while on the maximum dose of 0.04mg/kg/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- Patient must have had a bone age at or above skeletal maturity (15.5 years for males and 13.5 years for females) at the last application and must have maintained or improved weight SDS for age and sex while on the maximum dose of 0.04mg/kg/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have developed uncontrolled morbid obesity, defined as a body weight greater than 200% of ideal body weight for height and sex, with ideal body weight derived by calculating the 50th percentile weight for the patient's current height.

Population criteria:

- Patient must not have a chronological age of equal to or greater than 18 years.
- The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment; **AND**
3. Growth data (height, weight and waist circumference) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; **AND**
4. The date at which skeletal maturity was achieved (if applicable) [Note: In patients whose chronological age is greater than 2.5 years, a bone age reading should be performed at least once every 12 months prior to attainment of skeletal maturity]; **AND**
5. Confirmation that during the initial 32 week treatment period, the patient was re-evaluated via polysomnography for airway obstruction and apnoea, and any sleep disorders that were identified have been addressed; **AND**
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Maintenance is defined as a value within a 5% tolerance (this allows for seasonal and other measurement variations).

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Prior Written Approval of Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Short stature and slow growth

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than short stature and slow growth, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies).

continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR

- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have previously received treatment under the indication short stature associated with chronic renal insufficiency, have undergone a renal transplant and a 12 month period of observation following the transplant, and have an estimated glomerular filtration rate of greater than or equal to 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula; OR
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment and a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment), **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; **AND**
3. (a) A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment; OR
(b) Confirmation that the patient has previously received treatment under the indication **short stature associated with chronic renal insufficiency**, has undergone a renal transplant and a 12 month period of observation following the transplant, and has an estimated glomerular filtration rate of greater than or equal to 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula; **AND**
4. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; **AND**
5. A bone age result performed within the last 12 months; **AND**
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Prior Written Approval of Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Short stature associated with biochemical growth hormone deficiency

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than short stature associated with biochemical growth hormone deficiency, **AND**

- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have previously received treatment under the indication **risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants** and have reached or surpassed 5 years of age (chronological); OR
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment; OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and an annual growth velocity of 14 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and an annual growth velocity of 8 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special**

Arrangement 2015 and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; AND
3. (a) A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); OR
(b) Height and weight measurements from within three months prior to commencement of treatment for a patient whose height was at or below the 1st percentile for age and sex immediately prior to commencing treatment; OR
(c) Confirmation that the patient has previously received treatment under the indication **risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants** and has reached or surpassed 5 years of age (chronological); AND
4. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; AND
5. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
6. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
7. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Biochemical growth hormone deficiency should not be secondary to an intracranial lesion or cranial irradiation for applications under this category.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Prior Written Approval of Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Growth retardation secondary to an intracranial lesion, or cranial irradiation

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than growth retardation secondary to an intracranial lesion, or cranial irradiation, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have had an intracranial lesion and have undergone a 12 month period of observation following completion of treatment for the intracranial lesion (all treatment); OR
- Patient must have had an intracranial lesion, have received medical advice that it is unsafe to treat the intracranial lesion, and have undergone a 12 month period of observation since initial diagnosis of the intracranial lesion; OR
- Patient must have received cranial irradiation without having had an intracranial lesion, and have undergone a 12 month period of observation following completion of treatment for the condition for which cranial irradiation was received, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR

- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment; OR
- Patient must have had both a height above the 1st percentile for age and sex immediately prior to commencing treatment and a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had both a height above the 1st percentile for age and sex immediately prior to commencing treatment and an annual growth velocity of 14 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had both a height above the 1st percentile for age and sex immediately prior to commencing treatment and an annual growth velocity of 8 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; **AND**
3. (a) A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); OR
(b) Height and weight measurements from within three months prior to commencement of treatment for a patient whose height was at or below the 1st percentile for age and sex immediately prior to commencing treatment; **AND**
4. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; **AND**
5. (a) Confirmation that the patient has had an intracranial lesion and has undergone a 12 month period of observation following completion of treatment for the intracranial lesion (all treatment); OR
(b) Confirmation that the patient has had an intracranial lesion, has received medical advice that it is unsafe to treat the intracranial lesion, and has undergone a 12 month period of observation since initial diagnosis of the intracranial lesion; OR
(c) Confirmation that the patient has received cranial irradiation without having had an intracranial lesion, and has undergone a 12 month period of observation following completion of treatment for the condition for which cranial irradiation was received; **AND**
6. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; **AND**
7. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**

8. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Prior Written Approval of Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under a category other than risk of hypoglycaemia secondary to growth hormone deficiency in neonates/infants, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have a chronological age of less than 2 years, **AND**
- Patient must have a documented clinical risk of hypoglycaemia, **AND**
- Patient must have documented evidence that the risk of hypoglycaemia is secondary to biochemical growth hormone deficiency, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; AND
3. Confirmation that the patient has a documented clinical risk of hypoglycaemia; AND
4. Confirmation that the patient has documented evidence that the risk of hypoglycaemia is secondary to biochemical growth hormone deficiency; AND
5. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
6. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Prior Written Approval of Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Biochemical growth hormone deficiency and precocious puberty

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under a category other than biochemical growth hormone deficiency and precocious puberty, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or commencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or commencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or commencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or commencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or commencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must be male and have commenced puberty (demonstrated by Tanner stage 2 genital or pubic hair development or testicular volumes greater than or equal to 4 mL) before the chronological age of 9 years; OR
- Patient must be female and have commenced puberty (demonstrated by Tanner stage 2 breast or pubic hair development) before the chronological age of 8 years; OR
- Patient must be female and menarche occurred before the chronological age of 10 years, **AND**
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGFBP-3 levels, **AND**
- Patient must be undergoing Gonadotrophin Releasing Hormone agonist therapy for pubertal suppression, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; AND
3. Confirmation that the patient has precocious puberty; AND
4. Confirmation that the patient is undergoing Gonadotrophin Releasing Hormone agonist therapy for pubertal suppression; AND
5. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; AND
6. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
7. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
8. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Prior Written Approval of Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than hypothalamic-pituitary disease secondary to a structural lesion, with hypothalamic obesity driven growth, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have a structural lesion that is not neoplastic; OR
- Patient must have had a structural lesion that was neoplastic and have undergone a 12 month period of observation following completion of treatment for the structural lesion (all treatment); OR
- Patient must have a structural lesion that is neoplastic, have received medical advice that it is unsafe to treat the structural lesion, and have undergone a 12 month period of observation since initial diagnosis of the structural lesion, **AND**

AND

- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 2 pharmacological growth hormone stimulation tests (e.g. arginine, clonidine, glucagon, insulin); OR

- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 pharmacological growth hormone stimulation test (e.g. arginine, clonidine, glucagon, insulin) and 1 physiological growth hormone stimulation test (e.g. sleep, exercise); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) with other evidence of growth hormone deficiency, including septo-optic dysplasia (absent corpus callosum and/or septum pellucidum), midline abnormality including optic nerve hypoplasia, cleft lip and palate, midfacial hypoplasia and central incisor, ectopic and/or absent posterior pituitary bright spot, absent empty sella syndrome, hypoplastic anterior pituitary gland and/or pituitary stalk/infundibulum, and genetically proven biochemical growth hormone deficiency either isolated or as part of hypopituitarism in association with pituitary deficits (ACTH, TSH, GnRH or vasopressin/ADH deficiency); OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; OR
- Patient must have evidence of biochemical growth hormone deficiency, with a peak serum growth hormone concentration less than 10 mU/L or less than or equal to 3.3 micrograms per litre in response to 1 growth hormone stimulation test (pharmacological or physiological e.g. arginine, clonidine, glucagon, insulin, sleep, exercise) and low plasma IGF-1 levels; AND
- Patient must have other hypothalamic/pituitary hormone deficits (includes ACTH, TSH, GnRH and/or vasopressin/ADH deficiencies), AND
- Patient must have hypothalamic obesity, AND
- Patient must have had a growth velocity above the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had an annual growth velocity of greater than 14 cm per year in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had an annual growth velocity of greater than 8 cm per year in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, AND
- Patient must not have diabetes mellitus, AND
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, AND
- Patient must not have an active tumour or evidence of tumour growth or activity, AND
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; AND
3. A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); AND
4. Evidence of biochemical growth hormone deficiency, including the type of tests performed and peak growth hormone concentrations; AND
5. (a) Confirmation that the patient has a structural lesion that is not neoplastic; OR
(b) Confirmation that the patient had a structural lesion that was neoplastic and has undergone a 12 month period of observation following completion of treatment for the structural lesion (all treatment); OR
(c) Confirmation that the patient has a structural lesion that is neoplastic, has received medical advice that it is unsafe to treat the structural lesion, and has undergone a 12 month period of observation since initial diagnosis of the structural lesion; AND
6. Confirmation that the patient has other hypothalamic/pituitary hormone deficits; AND
7. Confirmation that the patient has hypothalamic obesity; AND
8. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND

9. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
10. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).
- Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:
 Department of Human Services
 Prior Written Approval of Complex Drugs
 Reply Paid 9826
 HOBART TAS 7001

Authority required

Short stature associated with Turner syndrome

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under a category other than short stature associated with Turner syndrome, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have had a height at or below the 95th percentile for age on the Turner syndrome growth curve for girls immediately prior to commencing growth hormone treatment, **AND**
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as a loss of a whole X chromosome in all cells (45X), and gender of rearing is female; OR
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as a loss of a whole X chromosome in some cells (mosaic 46XX/45X), and gender of rearing is female; OR
- Patient must have diagnostic results consistent with Turner syndrome (the condition must be genetically proven), defined as genetic loss or rearrangement of an X chromosome (such as isochromosome X, ring-chromosome, or partial deletion of an X chromosome), and gender of rearing is female, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have a bone age of 2.5 years or less, **AND**
- Patient must not have a bone age of 13.5 years or greater, **AND**
- Patient must not have a height greater than or equal to 155.0 cm.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; AND
3. A height measurement from immediately prior to commencement of growth hormone treatment; AND

4. Confirmation that the patient has diagnostic results consistent with Turner syndrome; AND
 5. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
 6. A bone age result performed within the last 12 months; AND
 7. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).
- Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Prior Written Approval of Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Short stature due to short stature homeobox (SHOX) gene disorders

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than short stature due to short stature homeobox (SHOX) gene disorders, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as a karyotype confirming the presence of a SHOX mutation/deletion without the presence of mixed gonadal dysgenesis; OR
- Patient must have diagnostic results consistent with a SHOX mutation/deletion, defined as mixed gonadal dysgenesis (45X mosaic karyotype with the presence of any Y chromosome material and/or SRY gene positive by FISH study) and have an appropriate plan of management in place for the patient's increased risk of gonadoblastoma, **AND**
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment, **AND**
- Patient must have had a growth velocity below the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had an annual growth velocity of 14 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR
- Patient must have had an annual growth velocity of 8 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes (excluding gonadoblastoma secondary to mixed gonadal dysgenesis), **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; AND
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; AND
3. A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); AND
4. Confirmation that the patient has diagnostic results consistent with a short stature homeobox (SHOX) gene disorder; AND
5. If the patient's condition is secondary to mixed gonadal dysgenesis, confirmation that an appropriate plan of management for the patient's increased risk of gonadoblastoma is in place; AND
6. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; AND
7. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); AND
8. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Prior Written Approval of Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Short stature associated with chronic renal insufficiency

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program (treatment) under a category other than short stature associated with chronic renal insufficiency, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 9.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have had a height at or below the 1st percentile for age and sex immediately prior to commencing treatment; OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and a growth velocity less than or equal to the 25th percentile for bone age and sex measured over the 12 month interval immediately prior to commencement of treatment (or the 6 month interval immediately prior to commencement of treatment if the patient was an older child at commencement of treatment); OR
- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and an annual growth velocity of 14 cm per year or less in the 12 month period immediately

prior to commencement of treatment, if the patient had a chronological age of 2 years or less at commencement of treatment; OR

- Patient must have had both a height above the 1st and at or below the 25th percentiles for age and sex immediately prior to commencing treatment and an annual growth velocity of 8 cm per year or less in the 12 month period immediately prior to commencement of treatment, if the patient had a bone or chronological age of 2.5 years or less at commencement of treatment, **AND**
- Patient must have an estimated glomerular filtration rate less than 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula, and not have undergone a renal transplant; OR
- Patient must have an estimated glomerular filtration rate less than 30mL/minute/1.73m² measured by creatinine clearance, excretion of radionuclides such as DTPA, or by the height/creatinine formula, have undergone a renal transplant, and have undergone a 12 month period of observation following the transplant, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must be male and must not have a height greater than or equal to 167.7cm; OR
- Patient must be female and must not have a height greater than or equal to 155.0cm, **AND**
- Patient must be male and must not have a bone age of 15.5 years or more; OR
- Patient must be female and must not have a bone age of 13.5 years or more.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

An older child is defined as a male with a chronological age of at least 12 years or a bone age of at least 10 years, or a female with a chronological age of at least 10 years or a bone age of at least 8 years.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; **AND**
3. (a) A minimum of 12 months of growth data (height and weight measurements) from immediately prior to commencement of treatment, or a minimum of 6 months of growth data from immediately prior to commencement of treatment if the patient was an older child at commencement of treatment; and the result of a bone age assessment performed within the 12 months immediately prior to commencement of treatment (except for a patient whose chronological age was 2.5 years or less at commencement of treatment); **OR**
(b) Height and weight measurements from within three months prior to commencement of treatment for a patient whose height was at or below the 1st percentile for age and sex immediately prior to commencing treatment; **AND**
4. Confirmation that the patient has an estimated glomerular filtration rate less than 30mL/minute/1.73m² ; **AND**
5. If a renal transplant has taken place, confirmation that the patient has undergone a 12 month period of observation following transplantation; **AND**
6. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; **AND**
7. A bone age result performed within the last 12 months (except for a patient whose chronological age is 2.5 years or less); **AND**

The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note If a patient receiving treatment under the indication short stature due to chronic renal insufficiency undergoes a renal transplant and 12 months post-transplant has an eGFR of equal to or greater than 30mL/min/1.73m² prescribers should seek reclassification to the indication short stature and slow growth.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Prior Written Approval of Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Short stature and poor body composition due to Prader-Willi syndrome

Treatment Phase: Continuing treatment as a reclassified patient

Clinical criteria:

- Patient must have previously received treatment under the PBS S100 Growth Hormone Program under a category other than short stature and poor body composition due to Prader-Willi syndrome, **AND**
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by a significant medical illness; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by major surgery (e.g. renal transplant); OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by an adverse reaction to growth hormone; OR
- The treatment must not have lapsed due to failure to respond to growth hormone at a dose of 7.5mg/m²/week or greater for the most recent treatment period (32 weeks for an initial or recommencement treatment period and 26 weeks for a continuing treatment period, whichever applies), unless response was affected by non-compliance due to social/family problems, **AND**
- Patient must have diagnostic results consistent with Prader-Willi syndrome (the condition must be genetically proven); OR
- Patient must have a clinical diagnosis of Prader-Willi syndrome, confirmed by a clinical geneticist, **AND**
- Patient must have been evaluated via polysomnography for airway obstruction and apnoea whilst on growth hormone treatment and any sleep disorders identified that required treatment must have been addressed, **AND**
- Patient must not have uncontrolled morbid obesity, defined as a body weight greater than 200% of ideal body weight for height and sex, with ideal body weight derived by calculating the 50th percentile weight for the patient's current height, **AND**
- Patient must not have diabetes mellitus, **AND**
- Patient must not have a condition with a known risk of malignancy including chromosomal abnormalities such as Down and Bloom syndromes, **AND**
- Patient must not have an active tumour or evidence of tumour growth or activity, **AND**
- Patient must not have a chronological age of 18 years or greater.

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in paediatric endocrinology; OR
- Must be treated by a medical practitioner in consultation with a nominated specialist or consultant physician in general paediatrics.

The maximum duration of each continuing treatment phase is 26 weeks. Prescribers must determine an appropriate weekly dose in accordance with the dosing arrangements detailed in the **National Health (Growth Hormone Program) Special Arrangement 2015** and request the appropriate number of vials/cartridges required to provide sufficient drug for 13 weeks' worth of treatment (with up to 1 repeat allowed).

The authority application must be in writing and must include:

1. A completed authority prescription form; **AND**
2. A completed Growth Hormone Authority Application Supporting Information Form for continuing treatment as a reclassified patient; **AND**
3. (a) Confirmation that the patient has diagnostic results consistent with Prader-Willi syndrome, OR
(b) Confirmation that the patient has a clinical diagnosis of Prader-Willi syndrome, confirmed by a clinical geneticist; **AND**
4. Confirmation that the patient has been evaluated via polysomnography for airway obstruction and apnoea whilst on growth hormone treatment, and any sleep disorders identified via the polysomnography that required treatment have been addressed; **AND**
5. Growth data (height and weight) for the most recent 6 month treatment period, including data at both the start and end of the treatment period. The most recent data must not be older than three months; **AND**
6. The date at which skeletal maturity was achieved (if applicable) [Note: In patients whose chronological age is greater than 2.5 years, a bone age reading should be performed at least once every 12 months prior to attainment of skeletal maturity]; **AND**
7. The proprietary name (brand), form and strength of somatropin requested, and the number of vials/cartridges required to provide sufficient drug for 13 weeks worth of treatment (with up to 1 repeat allowed).

Prescribers must keep a copy of any clinical records relating to the prescription, including such records required to demonstrate that the prescription was written in compliance with any relevant circumstances and/or purposes. These records must be kept for 2 years after the date the prescription to which the records relate is written.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

SYSTEMIC HORMONAL PREPARATIONS, EXCL. SEX HORMONES AND INSULINS

somatropin 2 mg injection [7] (&) inert substance diluent [7 x 0.25 mL syringes], 1 pack

10428W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	612.79	40.30	Genotropin MiniQuick [PF]

somatropin 1.6 mg injection [7] (&) inert substance diluent [7 x 0.25 mL syringes], 1 pack

10454F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	491.72	40.30	Genotropin MiniQuick [PF]

somatropin 1.4 mg injection [7] (&) inert substance diluent [7 x 0.25 mL syringes], 1 pack

10488B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	431.17	40.30	Genotropin MiniQuick [PF]

somatropin 1.8 mg injection [7] (&) inert substance diluent [7 x 0.25 mL syringes], 1 pack

10500P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	552.26	40.30	Genotropin MiniQuick [PF]

SOMATROPIN (Recombinant human growth hormone) Powder for injection 12 mg (36 i.u.) with diluent in pre-filled pen (with preservative), 1

10431B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	510.20	40.30	Genotropin GoQuick [PF]

SOMATROPIN (Recombinant human growth hormone) Powder for injection 5 mg (15 i.u.) with diluent in pre-filled pen (with preservative), 1

10443P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	216.90	40.30	Genotropin GoQuick [PF]

somatropin 800 microgram injection [7] (&) inert substance diluent [7 x 0.25 mL syringes], 1 pack

10479M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	249.55	40.30	Genotropin MiniQuick [PF]

somatropin 1.2 mg injection [7] (&) inert substance diluent [7 x 0.25 mL syringes], 1 pack

10453E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	370.63	40.30	Genotropin MiniQuick [PF]

somatropin 600 microgram injection [7] (&) inert substance diluent [7 x 0.25 mL syringes], 1 pack

10456H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	189.02	40.30	Genotropin MiniQuick [PF]

somatropin 400 microgram injection [7] (&) inert substance diluent [7 x 0.25 mL syringes], 1 pack

10891F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	128.47	40.30	Genotropin MiniQuick [PF]

somatropin 1 mg injection [7] (&) inert substance diluent [7 x 0.25 mL syringes], 1 pack

10480N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	310.09	40.30	Genotropin MiniQuick [PF]

IVF Treatment Program

GENITO URINARY SYSTEM AND SEX HORMONES	894
SEX HORMONES AND MODULATORS OF THE GENITAL SYSTEM	894
PROGESTOGENS.....	894
GONADOTROPINS AND OTHER OVULATION STIMULANTS	894
SYSTEMIC HORMONAL PREPARATIONS, EXCL. SEX HORMONES AND INSULINS.....	897
PITUITARY AND HYPOTHALAMIC HORMONES AND ANALOGUES	897
HYPOTHALAMIC HORMONES.....	897
ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS	898
ENDOCRINE THERAPY	898
HORMONES AND RELATED AGENTS	898

GENITO URINARY SYSTEM AND SEX HORMONES

SEX HORMONES AND MODULATORS OF THE GENITAL SYSTEM

PROGESTOGENS

Pregnen (4) derivatives

PROGESTERONE

Authority required (STREAMLINED)

4997

Assisted Reproductive Technology

Clinical criteria:

- The treatment must be for luteal phase support as part of an assisted reproductive technology (ART) treatment cycle for infertile women, **AND**
- Patient must be receiving medical services as described in items 13200 or 13201 of the Medicare Benefits Schedule. The luteal phase is defined as the time span from embryo transfer until implantation confirmed by positive B-hCG measurement.

progesterone 200 mg pessary, 15

9609R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	..	..	*172.18	40.30	Oripro [ON]

progesterone 100 mg pessary, 21

10116K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*91.63	40.30	Endometrin [FP]

progesterone 200 mg capsule, 42

10930G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	86.92	40.30	Utrogestan [HB]

progesterone 100 mg pessary, 15

9608Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	..	..	*156.79	40.30	Oripro [ON]

PROGESTERONE

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

5045

Assisted Reproductive Technology

Clinical criteria:

- The treatment must be for luteal phase support as part of an assisted reproductive technology (ART) treatment cycle for infertile women, **AND**
- Patient must be receiving medical services as described in items 13200 or 13201 of the Medicare Benefits Schedule. The luteal phase is defined as the time span from embryo transfer until implantation confirmed by positive B-hCG measurement.

progesterone 8% vaginal gel, 15 applications

6366C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*258.27	40.30	Crinone 8% [SG]

GONADOTROPINS AND OTHER OVULATION STIMULANTS

Gonadotropins

CHORIOGONADOTROPIN ALFA

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

5019

Assisted Reproductive Technology

Clinical criteria:

- Patient must be receiving medical services as described in items 13200, 13201, 13202 or 13203 of the Medicare Benefits Schedule.

choriogonadotropin alfa 250 microgram/0.5 mL injection, 0.5 mL pen device

6182J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	58.24	40.30	Ovidrel [SG]

CORIFOLLITROPIN ALFA

Authority required (STREAMLINED)

5009

Assisted Reproductive Technology

Clinical criteria:

- The treatment must be for controlled ovarian stimulation, **AND**
- Patient must have an antral follicle count of 20 or less, **AND**
- Patient must be receiving medical services as described in items 13200, 13201, or 13202 of the Medicare Benefits Schedule, **AND**
- Patient must be undergoing a gonadotrophin releasing antagonist cycle.

corifollitropin alfa 100 microgram/0.5 mL injection, 0.5 mL syringe

5816D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	412.61	40.30	Elonva [MK]

corifollitropin alfa 150 microgram/0.5 mL injection, 0.5 mL syringe

5817E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	672.81	40.30	Elonva [MK]

■ FOLLITROPIN ALFA

Note Biosimilar prescribing policy Prescribing of the biosimilar brand, Bemfola, is encouraged for treatment naive patients. Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Authority required (STREAMLINED)**5027**

Assisted Reproductive Technology

Clinical criteria:

- Patient must be receiving medical services as described in items 13200, 13201, 13202 or 13203 of the Medicare Benefits Schedule.

follitropin alfa 75 units (5.5 microgram)/0.125 mL injection, 5 x 0.125 mL pen devices

10861P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	..	..	*433.12	40.30	Bemfola [FX]

follitropin alfa 900 units (65.52 microgram)/1.5 mL injection, 1.5 mL pen device

6433N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	5	..	..	*1684.79	40.30	Gonal-f Pen [SG]

follitropin alfa 300 units (21.84 microgram)/0.5 mL injection, 0.5 mL pen device

6431L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*234.45	40.30	Gonal-f Pen [SG]

follitropin alfa 300 units (22 microgram)/0.5 mL injection, 5 x 0.5 mL pen devices

10866X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	..	..	*1684.78	40.30	Bemfola [FX]

follitropin alfa 225 units (16.5 microgram)/0.375 mL injection, 5 x 0.375 mL pen devices

10872F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	..	..	*1275.43	40.30	Bemfola [FX]

follitropin alfa 150 units (11 microgram)/0.25 mL injection, 5 x 0.25 mL pen devices

10873G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	..	..	*858.85	40.30	Bemfola [FX]

follitropin alfa 450 units (33 microgram)/0.75 mL injection, 5 x 0.75 mL pen devices

10867Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	..	..	*2503.48	40.30	Bemfola [FX]

follitropin alfa 450 units (32.76 microgram)/0.75 mL injection, 0.75 mL pen device

6432M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*347.97	40.30	Gonal-f Pen [SG]

■ FOLLITROPIN ALFA + LUTROPIN ALFA**Authority required (STREAMLINED)****5250**

Stimulation of follicular development

Clinical criteria:

- Patient must have severe LH deficiency, **AND**

- Patient must be considered appropriate for treatment with the combination product after titration of FSH and LH after at least one cycle of treatment, **AND**
- Patient must be receiving medical treatment as described in items 13200, 13201, 13202 or 13203 of the Medicare Benefits Schedule.

follitropin alfa 150 units (10.92 microgram) + lutropin alfa 75 units injection [1 vial] (&) inert substance diluent [1 mL vial], 1 pack

10491E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	14	..	..	*2187.19	40.30	Pergoveris [SG]

follitropin alfa 900 units (65.52 microgram)/1.44 mL + lutropin alfa 450 units/1.44 mL injection, 1.44 mL pen device

11667C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*1881.47	40.30	Pergoveris [SG]

■ FOLLITROPIN BETA
Authority required (STREAMLINED)
5027

Assisted Reproductive Technology

Clinical criteria:

- Patient must be receiving medical services as described in items 13200, 13201, 13202 or 13203 of the Medicare Benefits Schedule.

follitropin beta 900 units/1.08 mL injection, 1.08 mL cartridge

6464F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	5	..	..	*1814.64	40.30	Puregon 900 IU/1.08 mL [MK]

follitropin beta 300 units/0.36 mL injection, 0.36 mL cartridge

6335K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*260.81	40.30	Puregon 300 IU/0.36 mL [MK]

follitropin beta 600 units/0.72 mL injection, 0.72 mL cartridge

6336L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	4	..	..	*996.47	40.30	Puregon 600 IU/0.72 mL [MK]

■ FOLLITROPIN DELTA
Authority required (STREAMLINED)
5027

Assisted Reproductive Technology

Clinical criteria:

- Patient must be receiving medical services as described in items 13200, 13201, 13202 or 13203 of the Medicare Benefits Schedule.

follitropin delta 36 microgram/1.08 mL injection, 1.08 mL pen device

11431P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	5	..	..	*1374.09	40.30	Rekovele [FP]

follitropin delta 12 microgram/0.36 mL injection, 0.36 mL pen device

11430N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	5	..	..	*467.34	40.30	Rekovele [FP]

follitropin delta 72 microgram/2.16 mL injection, 2.16 mL pen device

11414R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	4	..	..	*2170.15	40.30	Rekovele [FP]

■ HUMAN CHORIONIC GONADOTROPHIN
Authority required (STREAMLINED)
6991

Assisted Reproductive Technology

Clinical criteria:

- Patient must be receiving medical services as described in items 13200, 13201, 13202 or 13203 of the Medicare Benefits Schedule.

human chorionic gonadotrophin 5000 units injection [1 vial] (&) inert substance diluent [1 mL vial], 1 pack

11156E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*27.57	28.82	Pregnyl [MK]

SYSTEMIC HORMONAL PREPARATIONS, EXCL. SEX HORMONES AND INSULINS

human chorionic gonadotrophin 1500 units injection [3 vials] (&) inert substance diluent [3 x 1 mL vials], 1 pack

11154C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	42.13	40.30	Pregnyl [MK]

■ HUMAN MENOPAUSAL GONADOTROPHIN

Authority required (STREAMLINED)

5027

Assisted Reproductive Technology

Clinical criteria:

- Patient must be receiving medical services as described in items 13200, 13201, 13202 or 13203 of the Medicare Benefits Schedule.

human menopausal gonadotrophin 1200 units injection [1 vial] (&) inert substance diluent [2 x 1 mL syringes], 1 pack

2038G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	4	..	..	*2065.87	40.30	Menopur 1200 [FP]

human menopausal gonadotrophin 600 units injection [1 vial] (&) inert substance diluent [1 mL syringe], 1 pack

2036E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	..	..	*794.59	40.30	Menopur 600 [FP]

■ LUTROPIN ALFA

Authority required (STREAMLINED)

5251

Stimulation of follicular development

Clinical criteria:

- Patient must have severe LH deficiency, **AND**
- Patient must be receiving medical treatment as described in items 13200, 13201, 13202 or 13203 of the Medicare Benefits Schedule.

lutropin alfa 75 units injection [1 vial] (&) inert substance diluent [1 mL vial], 1 pack

10465T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	14	..	..	*1422.93	40.30	Luveris [SG]

■ SYSTEMIC HORMONAL PREPARATIONS, EXCL. SEX HORMONES AND INSULINS

■ PITUITARY AND HYPOTHALAMIC HORMONES AND ANALOGUES

HYPOTHALAMIC HORMONES

Gonadotropin-releasing hormones

■ NAFARELIN

Authority required (STREAMLINED)

5046

Assisted Reproductive Technology

Clinical criteria:

- The treatment must be for prevention of premature luteinisation and ovulation, **AND**
- Patient must be undergoing controlled ovarian stimulation, **AND**
- Patient must be receiving medical services as described in items 13200, 13201, 13202 or 13203 of the Medicare Benefits Schedule.

nafarelin 200 microgram/actuation nasal spray, 60 actuations

5815C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*220.79	40.30	Synarel [PF]

Anti-gonadotropin-releasing hormones

■ CETRORELIX

Authority required (STREAMLINED)

5046

Assisted Reproductive Technology

Clinical criteria:

- The treatment must be for prevention of premature luteinisation and ovulation, **AND**
- Patient must be undergoing controlled ovarian stimulation, **AND**
- Patient must be receiving medical services as described in items 13200, 13201, 13202 or 13203 of the Medicare Benefits Schedule.

ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS

cetrorelix 250 microgram injection [1 vial] (&) inert substance diluent [1 mL syringe], 1 pack

9599F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	10	..	..	*462.69	40.30	Cetrotide [SG]

■ GANIRELIX

Authority required (STREAMLINED)

5046

Assisted Reproductive Technology

Clinical criteria:

- The treatment must be for prevention of premature luteinisation and ovulation, **AND**
- Patient must be undergoing controlled ovarian stimulation, **AND**
- Patient must be receiving medical services as described in items 13200, 13201, 13202 or 13203 of the Medicare Benefits Schedule.

ganirelix 250 microgram/0.5 mL injection, 0.5 mL syringe

9583J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	10	..	..	*462.69	40.30	Orgalutran [MK]

ganirelix 250 microgram/0.5 mL injection, 5 x 0.5 mL syringes

9584K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*462.71	40.30	Orgalutran [MK]

■ ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS

■ ENDOCRINE THERAPY

HORMONES AND RELATED AGENTS

Gonadotropin releasing hormone analogues

■ TRIPTORELIN

Authority required (STREAMLINED)

5046

Assisted Reproductive Technology

Clinical criteria:

- The treatment must be for prevention of premature luteinisation and ovulation, **AND**
- Patient must be undergoing controlled ovarian stimulation, **AND**
- Patient must be receiving medical services as described in items 13200, 13201, 13202 or 13203 of the Medicare Benefits Schedule.

triptorelin acetate 100 microgram/mL injection, 7 x 1 mL syringes

10907C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	4	..	..	*195.91	40.30	Decapeptyl [FP]

Opiate Dependence Treatment Program

NERVOUS SYSTEM.....	900
OTHER NERVOUS SYSTEM DRUGS.....	900
DRUGS USED IN ADDICTIVE DISORDERS	900

NERVOUS SYSTEM

OTHER NERVOUS SYSTEM DRUGS

DRUGS USED IN ADDICTIVE DISORDERS

Drugs used in opioid dependence

■ BUPRENORPHINE

Note Care must be taken to comply with the provisions of State/Territory law when prescribing this drug.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Opiate dependence

Treatment Phase: Maintenance and detoxification (withdrawal)

Clinical criteria:

- The treatment must be within a framework of medical, social and psychological treatment.

buprenorphine 400 microgram sublingual tablet, 7

6307Y	Max.Qty Packs	Price ex manufacturer \$	Brand Name and Manufacturer
NP	1	5.85	Subutex [IR]

buprenorphine 2 mg sublingual tablet, 7

6308B	Max.Qty Packs	Price ex manufacturer \$	Brand Name and Manufacturer
NP	1	9.98	Subutex [IR]

buprenorphine 8 mg sublingual tablet, 7

6309C	Max.Qty Packs	Price ex manufacturer \$	Brand Name and Manufacturer
NP	1	28.60	Subutex [IR]

■ BUPRENORPHINE

Note In the first 6 months of supply, buprenorphine subcutaneous injections will only be available when prescribed within hospital and specialist drug rehabilitation clinics by physicians who have demonstrated that they have reviewed the relevant education materials.

Note Care must be taken to comply with the provisions of State/Territory law when prescribing this drug.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Opiate dependence

Treatment criteria:

- Must be treated by a health care professional.

Clinical criteria:

- The treatment must be within a framework of medical, social and psychological treatment, **AND**
- Patient must be stabilised on sublingual buprenorphine or buprenorphine/naloxone prior to commencing treatment with this drug for this condition.

buprenorphine 32 mg/0.64 mL modified release injection, 0.64 mL syringe

11766G	Max.Qty Packs	Price ex manufacturer \$	Brand Name and Manufacturer
NP	1	92.40	Buvidal Weekly [UR]

buprenorphine 64 mg/0.18 mL modified release injection, 0.18 mL syringe

11754P	Max.Qty Packs	Price ex manufacturer \$	Brand Name and Manufacturer
NP	1	369.60	Buvidal Monthly [UR]

buprenorphine 24 mg/0.48 mL modified release injection, 0.48 mL syringe

11773P	Max.Qty Packs	Price ex manufacturer \$	Brand Name and Manufacturer
NP	1	92.40	Buvidal Weekly [UR]

buprenorphine 8 mg/0.16 mL modified release injection, 0.16 mL syringe

11759X	Max.Qty Packs	Price ex manufacturer \$	Brand Name and Manufacturer
NP	1	92.40	Buvidal Weekly [UR]

buprenorphine 16 mg/0.32 mL modified release injection, 0.32 mL syringe

11774Q	Max.Qty Packs	Price ex manufacturer \$	Brand Name and Manufacturer
NP	1	92.40	Buvidal Weekly [UR]

buprenorphine 128 mg/0.36 mL modified release injection, 0.36 mL syringe

11768J	Max.Qty Packs	Price ex manufacturer \$	Brand Name and Manufacturer
NP	1	369.60	Buvidal Monthly [UR]

buprenorphine 96 mg/0.27 mL modified release injection, 0.27 mL syringe

11767H	Max.Qty Packs	Price ex manufacturer \$	Brand Name and Manufacturer
NP	1	369.60	Buvidal Monthly [UR]

■ BUPRENORPHINE + NALOXONE

Note Buprenorphine with naloxone soluble film and buprenorphine with naloxone sublingual tablet do not meet all the criteria for bioequivalence. Patients being switched between sublingual tablets and soluble films may therefore require a dosage adjustment.

Note Care must be taken to comply with the provisions of State/Territory law when prescribing this drug.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Opiate dependence

Clinical criteria:

- The treatment must be within a framework of medical, social and psychological treatment.

buprenorphine 2 mg + naloxone 500 microgram sublingual film, 28

9749D	Max.Qty Packs	Price ex manufacturer \$	Brand Name and Manufacturer
NP	1	46.20	Suboxone Film 2/0.5 [IR]

buprenorphine 8 mg + naloxone 2 mg sublingual film, 28

9750E	Max.Qty Packs	Price ex manufacturer \$	Brand Name and Manufacturer
NP	1	132.44	Suboxone Film 8/2 [IR]

■ METHADONE

Caution The risk of drug dependence is high.

Note Care must be taken to comply with the provisions of State/Territory law when prescribing this drug.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Opiate dependence

methadone hydrochloride 5 mg/mL oral liquid, 1 L

6172W	Max.Qty Packs	Price ex manufacturer \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	33.20	^a Aspen Methadone Syrup [QA]	^a Biodone Forte [MW]

methadone hydrochloride 5 mg/mL oral liquid, 200 mL

6171T	Max.Qty Packs	Price ex manufacturer \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	7.91	^a Aspen Methadone Syrup [QA]	^a Biodone Forte [MW]

Extemporaneously Prepared Benefits

Drug Tariff

Drug	Standard	Recovery Prices			
		0.1 g/mL \$	1 g/mL \$	10 g/mL \$	100 g/mL \$
Acacia Mucilage (by weight)	APF 15	0.01	0.11	0.88	7.79
Acacia, powdered	BP	0.03	0.20	1.58	14.02
Acetic Acid (33 per cent)	BP	0.01	0.06	0.47	4.14
Acetic Acid (6 per cent)	BP	0.01	0.02	0.16	1.40
Acetic Acid Glacial BP	BP	0.02	0.14	1.09	9.72
Acetone (use as additive only)	BP	0.02	0.19	1.49	13.22
Alum	BP	0.01	0.07	0.59	5.28
Aluminium Acetate Solution	BP	0.02	0.17	1.38	12.27
Anise Oil BP	BP	0.18	1.44	11.52	102.44
Anise Water Concentrated 1 in 40	BP	0.01	0.07	0.59	5.27
Aqueous Cream (for use only as a base combined with active ingredients)	APF	0.01	0.05	0.39	3.44
Ascorbic Acid (for use only as an ingredient of ferrous sulfate mixtures)	BP	0.38	3.00	24.00	213.30
Aspirin	BP	0.14	1.11	8.90	79.09
Belladonna Tincture	BP	0.18	1.45	11.59	103.06
Benzocaine	BP	0.13	1.06	8.50	75.59
Benzoic Acid	BP	0.07	0.52	4.13	36.69
Benzoic Acid Compound Ointment	APF	0.02	0.17	1.37	12.16
Benzoic Acid Solution	BP	0.02	0.14	1.09	9.69
Benzoin Compound Tincture	BP	0.06	0.51	4.04	35.92
Boric Acid (use as additive only)	BP	0.02	0.17	1.38	12.26
Boric Acid, Olive Oil and Zinc Oxide Ointment	QHF	0.02	0.14	1.13	10.06
Calcium Hydroxide	BP	0.11	0.86	6.85	60.85
Calcium Hydroxide Solution	BP	0.01	0.02	0.19	1.65
Castor Oil (use as additive only)	BP	0.02	0.17	1.37	12.14
Cetomacrogol Aqueous Cream (for use only as a base combined with active ingredients)	APF	0.01	0.03	0.26	2.31
Cetrimide Aqueous Cream (for use only as a base combined with active ingredients)	APF	0.02	0.17	1.37	12.21
Chlorhexidine Acetate (use as additive only)	BP	0.63	5.03	40.21	357.42
Chlorhexidine Aqueous Cream (for use only as a base combined with active ingredients)	APF	0.03	0.24	1.95	17.36
Chloroform (use as additive only)	BP	0.09	0.72	5.77	51.32
Chloroform Spirit	BP	0.01	0.09	0.70	6.19
Chloroform Water Concentrated 1 in 40	APF 15	0.01	0.11	0.86	7.65
Citric Acid Monohydrate	BP	0.04	0.29	2.28	20.28
Coal Tar	BP	0.31	2.51	20.09	178.55
Coal Tar Solution	BP	0.02	0.19	1.48	13.17
Cocaine Hydrochloride	BP	6.03	48.22	385.72	3428.65
Coconut Oil	BP	0.01	0.06	0.48	4.24
Codeine Linctus	APF	0.02	0.13	1.00	8.88
Codeine Phosphate (may only be prescribed in linctuses, mixtures or mixtures for children)	BP	2.68	21.45	171.56	1524.99
Collodion Flexible	BP	0.19	1.55	12.37	109.96
Dithranol	BP	4.29	34.31	274.51	2440.09
Emulsifying Ointment (for use only as a base combined with active ingredients)	BP	0.01	0.07	0.57	5.11
Ephedrine Hydrochloride (may only be prescribed in nasal instillations)	BP	2.38	19.03	152.26	1353.39
Ethanol (90 per cent) (use as additive only)	BP	0.01	0.04	0.31	2.72
Ethanol (96 per cent) (use as additive only)	BP	0.01	0.05	0.39	3.47
Ether Solvent (use as additive only)	BP	0.27	2.15	17.18	152.70

Drug	Standard	Recovery Prices			
		0.1 g/mL \$	1 g/mL \$	10 g/mL \$	100 g/mL \$
Eucalyptus Oil (use as additive only)	BP	0.03	0.20	1.62	14.42
Ferrous Sulfate	BP	0.05	0.37	2.94	26.12
Formaldehyde Solution	BP	0.07	0.53	4.26	37.86
Gentian Alkaline Mixture	APF	0.01	0.09	0.68	6.00
Glycerol	BP	0.02	0.12	0.99	8.78
Honey Purified (use as additive only)	BP 1993	0.01	0.03	0.27	2.44
Hydroxybenzoate Compound Solution	APF	0.09	0.68	5.40	47.98
Iodine	BP	0.33	2.61	20.86	185.45
Iodine Alcoholic Solution	BP	0.04	0.31	2.46	21.89
Iodine Aqueous Oral Solution	BP	0.04	0.30	2.41	21.45
Kaolin Mixture	BPC 1968	0.03	0.23	1.84	16.32
Kaolin and Opium Mixture	APF 14	0.01	0.10	0.82	7.27
Lactic Acid	BP	0.35	2.76	22.04	195.91
Lavender Spike Oil	BPC 1968	0.13	1.04	8.32	73.91
Liquorice Liquid Extract	BP	0.03	0.23	1.84	16.37
Magnesium Carbonate Light	BP	0.05	0.37	2.92	25.99
Magnesium Sulfate (may only be prescribed for other than oral use)	BP	0.01	0.03	0.27	2.37
Magnesium Trisilicate	BP	0.05	0.41	3.24	28.76
Menthol, Racemic or Levomenthol	BP	0.28	2.25	17.96	159.62
Methyl Hydroxybenzoate	BP	0.46	3.71	29.64	263.43
Methyl Hydroxybenzoate Solution	APF	0.05	0.38	3.00	26.70
Methylated Industrial Spirit (use as additive only)	BP	0.01	0.01	0.11	1.00
Olive Oil (use as additive only)	BP	0.02	0.13	1.01	9.00
Paraffin Hard	BP	0.06	0.45	3.57	31.74
Paraffin Light Liquid	BP	0.02	0.17	1.39	12.32
Paraffin Liquid (use as additive only)	BP	0.01	0.07	0.57	5.08
Paraffin Soft White	BP	0.01	0.06	0.47	4.18
Paraffin Soft Yellow	BP	0.01	0.05	0.42	3.75
Peppermint Oil (use as additive only)	BP	0.08	0.64	5.09	45.29
Peppermint Water Concentrated 1 in 40 (use as additive only)	APF 16	0.04	0.28	2.26	20.12
Phenobarbitone Sodium (may only be prescribed for the treatment of epilepsy)	BP	9.37	74.92	599.36	5327.61
Phenol Liquefied (not available for ear drops)	BP	0.24	1.90	15.17	134.83
Podophyllum Resin	BP	3.78	30.26	242.09	2151.88
Potassium Citrate	BP	0.03	0.21	1.66	14.78
Potassium Iodide	BP	0.16	1.24	9.91	88.09
Potassium Permanganate	BP	0.05	0.37	2.99	26.57
Propyl Hydroxybenzoate	BP	0.42	3.36	26.84	238.54
Propylene Glycol	BP	0.02	0.12	0.92	8.17
Red Syrup	APF 15	0.02	0.13	1.04	9.23
Resorcinol	BP	0.42	3.33	26.60	236.41
Salicylic Acid	BP	0.03	0.25	1.98	17.60
Salicylic Acid Ointment	APF	0.02	0.17	1.38	12.23
Salicylic Acid Ointment	BP	0.02	0.17	1.38	12.23
Simple Ointment (white) (for use only as a base combined with active ingredients)	BP	0.02	0.14	1.11	9.88
Simple Ointment (yellow) (for use only as a base combined with active ingredients)	BP	0.02	0.14	1.11	9.88
Sodium Bicarbonate	BP	0.02	0.19	1.49	13.25
Sodium Chloride	BP	0.02	0.18	1.46	13.02
Sodium Chloride Solution	BP	0.01	0.01	0.11	0.95
Sodium Citrate	BP	0.07	0.54	4.31	38.34
Sodium Thiosulfate (use as additive only)	BP	0.05	0.37	2.96	26.34
Starch	BP	0.02	0.12	0.95	8.45
Sulfur Ointment (for use only as a base combined with active ingredients)	BP 1980	0.02	0.15	1.22	10.89
Sulfur Precipitated	BP 1980	0.03	0.25	1.99	17.68
Syrup	BP	0.01	0.05	0.43	3.85
Talc Purified, sterilised	BP	0.06	0.48	3.81	33.83
Thymol	BP	0.51	4.11	32.84	291.88
Thymol Compound Mouth Wash	APF 15	0.02	0.13	1.00	8.89
Tragacanth Compound Powder	BP 1980	0.07	0.57	4.58	40.73
Tragacanth Mucilage	APF 13	0.01	0.08	0.66	5.87

Drug	Standard	Recovery Prices			
		0.1 g/mL \$	1 g/mL \$	10 g/mL \$	100 g/mL \$
Tragacanth Mucilage	BPC 1973	0.01	0.07	0.59	5.20
Tragacanth, powdered	BP	0.41	3.27	26.15	232.44
Trichloroacetic Acid	BP 1980	0.38	3.04	24.29	215.91
Triethanolamine	BP	0.15	1.19	9.52	84.66
Water For Injections, sterilised (b) (extemporaneously prepared eye drops and eye lotions)	BP	0.00	0.00	11.95	11.95
Water Purified	BP	0.01	0.01	0.09	0.79
Wool Alcohols Ointment (white) (for use only as a base combined with active ingredients)	BP	0.02	0.19	1.51	13.42
Wool Alcohols Ointment (yellow) (for use only as a base combined with active ingredients)	BP	0.02	0.19	1.48	13.18
Wool Fat	BP	0.02	0.18	1.44	12.79
Wool Fat Hydrous	BP	0.04	0.28	2.27	20.18
Zinc Compound Paste	BP	0.05	0.40	3.17	28.17
Zinc Cream (for use only as a base combined with active ingredients)	BP	0.01	0.09	0.73	6.49
Zinc Oxide	BP	0.03	0.21	1.71	15.22
Zinc Sulfate	BP	0.04	0.31	2.49	22.13
Zinc and Salicylic Acid Paste	BP	0.05	0.37	2.95	26.22

Container Prices

Type	Container	Price \$
Dispensing Bottles	25mL	0.66
Dispensing Bottles	50mL	0.70
Dispensing Bottles	100mL	0.82
Dispensing Bottles	200mL	1.00
Dispensing Bottles	500mL	1.28
Poison Bottles	25mL	0.72
Poison Bottles	50mL	0.73
Poison Bottles	100mL	0.79
Poison Bottles	200mL	1.08
Poison Bottles	500mL	1.97
Dropper Containers (Glass)	15mL	1.15
Dropper Containers (Polythene)	15mL	0.98
	150ml	1.00
Screw Cap Jars	25g	0.95
Screw Cap Jars	50g	1.01
Screw Cap Jars	100g	1.39
Screw Cap Jars	200g	0.77
Screw Cap Jars	500g	2.19
	25ml	0.51

Standard Formula Preparations

Code	Item	Reference	DPMQ \$	MRVSN \$
	Creams			
7502W	(Maximum Quantity 100 g and 1 Repeat) Salicylic Acid and Sulfur Aqueous	APF	15.12	16.73
	Dusting Powders			
7458M	(Maximum Quantity 100 g and 1 Repeat) Zinc, Starch and Talc	APF 15 & BPC 1973	35.89	37.50
	Ear Drops			
7643G	(Maximum Quantity 15 ml and 2 Repeats) Aluminium Acetate	BP	12.48	14.09
7642F	Aluminium Acetate	APF	11.72	13.33
7314Y	Sodium Bicarbonate	APF & BP	11.20	12.81
7313X	Spirit	APF	10.71	12.32
	Inhalations			
	(Maximum Quantity 50 ml and 1 Repeat)			
7484X	Benzoin and Menthol	APF	32.62	34.23
7308P	Menthol	APF	13.94	15.55
7310R	Menthol and Eucalyptus	BP1980	15.10	16.71
	Linctuses containing Codeine Phosphate			
	(Maximum Quantity 100 ml and 0 Repeat)			
7530H	Codeine	APF	19.13	20.74
	Lotions			
	(Maximum Quantity 200 ml and 2 Repeats)			
7709R	Aluminium Acetate Aqueous	APF	13.40	15.01
	Mixtures, Other			
	(Maximum Quantity 200 ml and 4 Repeats)			
7604F	Gentian Alkaline	APF	22.43	24.04
7348R	Kaolin	BPC 1968	43.07	40.30
7301G	Kaolin and Opium	APF 14	24.97	26.58
7342K	Magnesium Trisilicate	BPC 1968	21.04	22.65
7343L	Magnesium Trisilicate and Belladonna	BPC 1968	32.48	34.09
	Mouth Washes			
	(Maximum Quantity 200 ml and 1 Repeat)			
7457L	Thymol Compound	APF 15	28.29	29.90
	Ointments, Waxes			
	(Maximum Quantity 100 g and 1 Repeat)			
7914M	Benzoic Acid Compound	APF & BP	22.98	24.59
7902X	Boric Acid, Olive Oil and Zinc Oxide	QHF	20.88	22.49
7926E	Salicylic Acid	APF	23.05	24.66
7928G	Salicylic Acid (extemporaneous formula)	BP	23.05	24.66
	Paints			
	(Maximum Quantity 25 ml and 1 Repeat)			
7567G	Podophyllin Compound	APF 16 & BP	133.73	40.30
7568H	Salicylic Acid	APF	41.69	40.30
	Pastes, Other			
	(Maximum Quantity 100 g and 1 Repeat)			
7558T	Zinc	APF & BP	38.99	40.30
	Powders for Internal Use			
	(Maximum Quantity 100 g and 2 Repeats)			
7545D	Magnesium Trisilicate	BP	38.96	40.30

Codes, Maximum Quantities, and Number of Repeats for Extemporaneously Prepared Benefits

Code	Preparation	Maximum Quantity	Number of Repeats
13Q	Creams	100 g	1
48M	Dusting Powders	100 g	1
15T	Ear Drops	15 ml	2
19B	Eye Drops containing Cocaine Hydrochloride	15 ml	..
22E	Eye Drops, Other	15 ml	5
23F	Eye Lotions	200 ml	2
29M	Inhalations	50 ml	1
64J	Linctuses containing Codeine Phosphate	100 ml	..
34T	Linctuses, Other	100 ml	2
39C	Lotions	200 ml	2
65K	Mixtures containing Codeine Phosphate	200 ml	..
66L	Mixtures for Children containing Codeine Phosphate	100 ml	..
41E	Mixtures for Children, Other	100 ml	4
40D	Mixtures, Other	200 ml	4
30N	Mouth Washes	200 ml	1
42F	Nasal Instillations	15 ml	2
43G	Ointments, Waxes	100 g	1
44H	Paints	25 ml	1
63H	Pastes containing Cocaine Hydrochloride	25 g	..
45J	Pastes, Other	100 g	1
49N	Powders for Internal Use	100 g	2
52R	Solutions	200 ml	2

Index of Manufacturers' Code

Code	Manufacturer	Code	Manufacturer
AB	Abbott Australasia Pty Ltd	GX	Apotex Pty Ltd
AE	AFT Pharmaceuticals (AU) Pty Ltd	GZ	sanofi-aventis Australia Pty Ltd
AF	Alphapharm Pty Ltd	HB	Besins Healthcare Australia Pty Ltd
AG	Allergan Australia Pty Limited	HN	Horizon Hospital Healthcare Pty Ltd
AL	Alphapharm Pty Ltd	HQ	Generic Health Pty Ltd
AN	Amgen Australia Pty Limited	HR	Paul Hartmann Pty Ltd
AP	AstraZeneca Pty Ltd	HX	Sandoz Pty Ltd
AQ	Alcon Laboratories (Australia) Pty Ltd	IB	Apotex Pty Ltd
AS	Aspen Pharmacare Australia Pty Limited	IG	Sigma Company Limited
AT	Actelion Pharmaceuticals Australia Pty Ltd	IJ	I-Care Pharma Distributors Pty Ltd
AV	sanofi-aventis Australia Pty Ltd	IL	iNova Pharmaceuticals (Australia) Pty Limited
BB	Blackmores Limited	IM	iNova Pharmaceuticals (Australia) Pty Limited
BD	Biogen Australia Pty Ltd	IO	BioMarin Pharmaceutical Australia Pty Ltd
BE	Beiersdorf Australia Ltd	IQ	Alcon Laboratories (Australia) Pty Ltd
BG	Sandoz Pty Ltd	IR	Indivior Pty Ltd
BI	Biotech Pharmaceuticals Pty Ltd	IS	Ipsen Pty Ltd
BN	Bayer Australia Ltd	IX	Clinect Pty Ltd
BQ	Bristol-Myers Squibb Australia Pty Ltd	IY	Clinect Pty Ltd
BR	B. Braun Australia Pty Ltd	JB	Apotex Pty Ltd
BV	BSN medical (Aust.) Pty Ltd	JC	Janssen-Cilag Pty Ltd
BX	Baxter Healthcare Pty Limited	JJ	Johnson & Johnson Medical Pty Ltd
BY	Boehringer Ingelheim Pty Ltd	JO	Juno Pharmaceuticals Pty Ltd
BZ	Boucher & Muir Pty Ltd	JT	Johnson & Johnson Pacific Pty Limited
CC	ConvaTec A Division of Bristol-Myers Squibb Australia Pty Ltd	JU	Juno Pharmaceuticals Pty Ltd
CF	CNS Pharma Pty Ltd	JX	Juno Pharmaceuticals Pty Ltd
CH	Apotex Pty Ltd	KE	Kendall Australasia Pty Ltd
CJ	Celgene Pty Limited	KI	KCI Medical Australia Pty Ltd
CR	Pharmacor Pty Limited	KP	Eli Lilly Australia Pty Ltd
CS	Seqirus (Australia) Pty Ltd	KY	Key Pharmaceuticals Pty Ltd
CT	Coloplast Pty Ltd	LC	Lohmann & Rauscher Pty Ltd
CU	Care Pharmaceuticals Pty Limited	LI	Luminarie Pty Ltd
DE	Stallergenes Australia Pty Ltd	LL	Astellas Pharma Australia Pty Ltd
DJ	De Fries Industries Pty Ltd	LM	Link Medical Products Pty Ltd
DO	Fair-Med Healthcare (Australia) Pty Ltd	LN	Aspen Pharmacare Australia Pty Limited
DQ	Church & Dwight (Australia) Pty Ltd	LO	Leo Pharma Pty Ltd
DV	Medical Developments International Limited	LQ	Astellas Pharma Australia Pty Ltd
DX	Ascensia Diabetes Care Australia Pty Ltd	LR	Cipla Australia Pty Ltd
DZ	Medsurge Healthcare Pty Ltd	LS	Astellas Pharma Australia Pty Ltd
EA	Amneal Pharmaceuticals Pty Ltd	LU	Lundbeck Australia Pty Ltd
ED	Amneal Pharmaceuticals Pty Ltd	LV	Luminarie Pty Ltd
EF	Amneal Pharmaceuticals Pty Ltd	LX	Lawley Pharmaceuticals Pty Ltd
EI	Eisai Australia Pty Ltd	LY	Eli Lilly Australia Pty Ltd
EL	Eli Lilly Australia Pty Ltd	MF	Mundipharma Pty Limited
EO	Ego Pharmaceuticals Proprietary Limited	MH	Molnlycke Health Care Pty Ltd
ER	Eris Pharmaceuticals (Australia) Pty Ltd	MK	Merck Sharp & Dohme (Australia) Pty Ltd
EU	Emerge Health Pty Ltd	MM	3M Pharmaceuticals Australia Pty Ltd
EV	Teva Pharma Australia Pty Limited	MQ	Alphapharm Pty Ltd
EZ	Merz Australia Pty Ltd	MT	Mentholatum Australasia Pty Ltd
FB	Pierre Fabre Australia Pty Ltd	MW	Biomed Aust Pty Limited
FF	Phebra Pty Ltd	NE	Norgine Pty Limited
FG	Phebra Pty Ltd	NF	Novo Nordisk Pharmaceuticals Pty Limited
FI	Boehringer Ingelheim Pty Ltd	NI	Novo Nordisk Pharmaceuticals Pty Limited
FK	A. Menarini Australia Pty Limited	NM	Novartis Pharmaceuticals Australia Pty Limited
FM	Fawns and McAllan Proprietary Limited	NO	Novo Nordisk Pharmaceuticals Pty Limited
FO	For Benefit Medicines Pty Ltd	NQ	Takeda Pharmaceuticals Australia Pty Ltd
FP	Ferring Pharmaceuticals Pty Limited	NT	Nestle Australia Ltd
FR	Merck Sharp & Dohme (Australia) Pty Ltd	NU	Nutricia Australia Pty Limited
FX	Gedeon Richter Australia Pty Ltd	NV	Novartis Pharmaceuticals Australia Pty Limited
FZ	Pfizer Australia Pty Ltd	OA	Orphan Australia Pty Ltd
GA	Galderma Australia Pty Ltd	OB	Oral B Laboratories Pty Ltd
GC	GlaxoSmithKline Australia Pty Ltd	OC	Accord Healthcare Pty Ltd
GH	Amdipharm Mercury (Australia) Pty Limited	OD	Accord Healthcare Pty Ltd
GI	Gilead Sciences Pty Limited	OE	Omegapharm Pty Ltd
GK	GlaxoSmithKline Australia Pty Ltd	OH	Orpharma Pty Ltd
GN	Actavis Pty Ltd	OM	Colgate Oral Care
GO	Mylan Health Pty Ltd	ON	Orion Laboratories Pty Ltd
GQ	Generic Health Pty Ltd	OS	Otsuka Australia Pharmaceutical Pty Ltd
GT	Mylan Health Pty Ltd	OU	Oraderm Pharmaceuticals Pty Ltd
		OW	Arrow Pharma Pty Ltd

Code	Manufacturer
PB	Pharmaco (Australia) Limited
PE	Allergan Australia Pty Limited
PF	Pfizer Australia Pty Ltd
PK	Fresenius Kabi Australia Pty Limited
PP	Petrus Pharmaceuticals Pty Ltd
PY	Procter & Gamble Pharmaceuticals Australia Pty Ltd
QA	Aspen Pharma Pty Ltd
QH	Cortex Health Pty Ltd
RA	Sun Pharma ANZ Pty Ltd
RB	Bio Revive Pty Ltd
RC	Reckitt Benckiser (Australia) Pty Limited
RF	Arrow Pharma Pty Ltd
RI	Dr Reddy's Laboratories (Australia) Pty Ltd
RJ	Recordati Rare Diseases Australia Pty. Ltd.
RN	Sun Pharma ANZ Pty Ltd
RO	Roche Products Pty Ltd
RW	Arrow Pharma Pty Ltd
RX	Servier Laboratories (Aust.) Pty Ltd
RZ	Dr Reddy's Laboratories (Australia) Pty Ltd
SA	SciGen (Australia) Pty Limited
SB	Nutricia Australia Pty Limited
SE	Servier Laboratories (Aust.) Pty Ltd
SG	Merck Serono Australia Pty Ltd
SI	Sigma Company Limited
SN	Smith & Nephew Pty Limited
SS	SSL Australia Pty Ltd
SW	sanofi-aventis Australia Pty Ltd
SY	Bayer Australia Ltd
SZ	Sandoz Pty Ltd
TB	Teva Pharma Australia Pty Limited
TD	STADA Pharmaceuticals Australia Pty Limited
TF	Te Arai BioFarma Limited
TK	Takeda Pharmaceuticals Australia Pty Ltd
TM	Technipro Marketing Pty Ltd
TS	Specialised Therapeutics Australia Pty Ltd
TT	Theramex Australia Pty Ltd
TW	Apotex Pty Ltd
TX	Apotex Pty Ltd
TY	Apotex Pty Ltd
UC	UCB Australia Proprietary Limited
UL	Bausch & Lomb (Australia) Pty Ltd
UM	Unomedical Pty Ltd
UN	Unilever Australia Limited
UO	Bausch & Lomb (Australia) Pty Ltd
UR	Camurus Pty Ltd
VE	AbbVie Pty Ltd
VF	VitaFlo Australia Pty Limited
VI	ViiV Healthcare Pty Ltd
VL	Vifor Pharma Pty Limited
VO	Avallon Pharmaceuticals Pty Limited
VR	Vertex Pharmaceuticals (Australia) Pty Ltd
VZ	Sanofi-aventis Healthcare Pty Ltd
WA	sanofi-aventis Australia Pty Ltd
WC	Wockhardt Bio Pty Ltd
XA	Pharmaxis Ltd
XC	Southern Cross Pharma Pty Ltd
XH	MS Health Pty Ltd
XI	Alexion Pharmaceuticals Australasia Pty Ltd
XM	The Medicines Company (Australia) Pty Limited
YB	Bayport Brands Pty Ltd
YC	Cipla Australia Pty Ltd
YN	Mayne Pharma International Pty Ltd
YT	Mayne Products Pty Ltd
ZA	AstraZeneca Pty Ltd
ZI	Shire Australia Pty Limited
ZP	Medis Pharma Pty Ltd
ZX	Zenex Pharmaceuticals Pty Ltd

Generic/Proprietary Index

3TC(VI)	604
ABACAVIR	602
ABACAVIR + LAMIVUDINE	608
ABACAVIR + LAMIVUDINE + ZIDOVUDINE	609
Abacavir/Lamivudine 600/300 APOTEX(TX)	608
ABACAVIR/LAMIVUDINE 600/300 SUN (RA)	608
Abacavir/Lamivudine GH 600/300(GQ)	609
Abacavir/Lamivudine Mylan(AF)	608
ABATACEPT	126, 410
Actemra(RO).....	249, 254, 260, 266, 532, 533, 538, 544, 550
ADALIMUMAB	153, 438
Adcirca(LY)	90, 374
ADEFOVIR DIPIVOXIL	602
Adempas(BN)	74, 75, 80, 358, 359, 364
ADVAGRAF XL(LQ)	271, 555
ALEMTUZUMAB	132, 416
AMBRISENTAN	43, 327
ANAKINRA	243, 527
APO-Adefovir(TX)	603
Apomine Intermittent(PF)	284, 568
Apomine Solution for Infusion(PF)	283, 567
APOMORPHINE	283, 567
APO-Mycophenolate(TX)	141, 142, 426
APO-Sildenafil PHT(TX)	85, 369
APO-Valaciclovir(TX)	98, 382
APO-Zoledronic Acid(TX)	280, 564
Aptivus(BY)	602
Aranesp SureClick(AN)	39, 40, 323, 324
Aranesp(AN)	39, 40, 323, 324
Aspen Methadone Syrup(QA) .Opiate Dependence Treatment Program	908
ATAZANAVIR	599
ATAZANAVIR + COBICISTAT	600
Atazanavir Mylan(AF)	599, 600
Atripla(GI)	613
AZACITIDINE	103, 104, 387, 388
Azacitidine Accord(OC)	104, 388
AZACITIDINE DR.REDDY'S (RI)	104, 388
Azacitidine Juno(JO)	104, 388
Azadine (RZ)	104, 388
AZITHROMYCIN .Highly Specialised Drugs Program (Private Hospital) .	97
.Highly Specialised Drugs Program (Public Hospital) .	381
BACLOFEN	277, 278, 561, 562
Bacthecal(DZ)	278, 562
Baraclude(BQ)	603, 604
Bemfola(FX)	902
BENRALIZUMAB	285, 287, 569, 571
BICTEGRAVIR + EMTRICITABINE + TENOFOVIR ALAFENAMIDE	609
Biktarvy(GI)	609
Biodone Forte (MW)	908
Bondronat(IX)	279, 563
BOSENTAN	48, 53, 332, 337
Bosentan APO(GX)	53, 59, 337, 343
BOSENTAN DR. REDDY'S(RI)	53, 59, 337, 343
Bosentan GH (GQ)	53, 337
Bosentan Mylan (AF)	59, 343
Bosentan Mylan(AF)	53, 337
Bosentan RBX (RA)	53, 337
Bosentan RBX(RA)	59, 343
Bosentan Sandoz (SZ)	59, 343
Bosentan Sandoz(SZ)	53, 337
BOSENTAN-DRLA (RZ)	53, 59, 337, 343
BOSLEER (RW)	53, 337
BOSLEER(RW)	59, 343
Botox(AG)	619, 620, 621, 622, 623
BOTULINUM TOXIN TYPE A	619, 620, 621, 622
bronchitol(XA)	303, 587

BUPRENORPHINE .Opiate Dependence Treatment Program	907
BUPRENORPHINE + NALOXONE	908
Buvidal Monthly(UR)	907, 908
Buvidal Weekly(UR)	907, 908
Caelyx(JC)	105, 389
Celazadine(JU)	104, 388
CellCept (RO)	141, 142, 426
CellCept(RO)	141, 425
Celsenti(VI)	615
Ceptolate(AF)	142, 426, 427
Certican(NV)	141, 425
CETRORELIX	905
Cetrotide(SG)	905
CHORIOGONADOTROPIN ALFA	901
CICLOSPORIN	270, 553, 554
Clopine 100(PF)	284, 568, 617
Clopine 200(PF)	285, 569, 617
Clopine 25(PF)	284, 568, 617
Clopine 50(PF)	285, 568, 617
Clopine Suspension(PF)	285, 569, 617
CLOSTRIDIUM BOTULINUM TYPE A TOXIN- HAEMAGGLUTININ COMPLEX	623, 624, 625
CLOZAPINE	284, 568, 616, 617
Clozaril 100 (GO)	284, 568, 617
Clozaril 25 (GO)	284, 568, 617
Combivir(VI)	612
CORIFOLLITROPIN ALFA	902
Crinone 8%(SG)	901
Cyclosporin Sandoz(SZ)	271, 554, 555
Cymevene(PB)	98, 382, 599
DACLATASVIR	98, 99, 382, 383
Daklinza(BQ)	98, 99, 382, 383
DARBEPOETIN ALFA	39, 323
DARUNAVIR	600
DARUNAVIR + COBICISTAT	609
DBL Zoledronic Acid (PF)	280, 564
DBL Zoledronic Acid(PF)	280, 564
Decapeptyl(FP)	905
DEFERASIROX	308, 309, 592, 593
DEFERIPRONE	309, 593
Descovy(GI)	611
DEFERRIOXAMINE	310, 594
DEZTRON(DZ)	280, 564
DOLUTEGRAVIR	614
DOLUTEGRAVIR + ABACAVIR + LAMIVUDINE	610
DOLUTEGRAVIR + RILPIVIRINE	610
DORNASE ALFA	302, 586
DOXORUBICIN HYDROCHLORIDE (AS PEGYLATED LIPOSOMAL)	105, 389
Duodopa(VE)	283, 567
Dysport(IS)	623, 624, 625
ECULIZUMAB	133, 134, 135, 137, 417, 418, 419, 421
Edurant(JC)	608
EFAVIRENZ	606
ELBASVIR + GRAZOPREVR	99, 383
Elonva(MK)	902
ELTROMBOPAG	33, 317
EMTRICITABINE + RILPIVIRINE + TENOFOVIR ALAFENAMIDE	611
EMTRICITABINE + TENOFOVIR ALAFENAMIDE	611
Enbrel(PF)	164, 448
Endometrin(FP)	901
ENFUVIRTIDE	614
ENTAC (LR)	603, 604
ENTECAVIR	603
Entecavir Amneal(EA)	603, 604
ENTECAVIR APO (GX)	603, 604
Entecavir APOTEX(TX)	603, 604

<i>Entecavir GH (GQ)</i>	603, 604	<i>INFLIXIMAB</i> ..	164, 166, 168, 170, 172, 175, 176, 177, 180, 182, 184, 186, 188, 190, 192, 195, 198, 200, 204, 208, 212, 216, 222, 227, 234, 448, 450, 452, 454, 457, 459, 461, 464, 466, 468, 470, 472, 474, 476, 479, 481, 484, 488, 493, 496, 500, 506, 511, 517
<i>Entecavir Mylan(AF)</i>	603, 604	<i>Intelence(JC)</i>	607
<i>ENTECAVIR RBX (RA)</i>	603, 604	<i>INTERFERON ALFA-2A</i>	124, 408, 616
<i>Entecavir Sandoz(SZ)</i>	603, 604	<i>INTERFERON GAMMA-1B</i>	125, 409
<i>ENTECLUDE (RW)</i>	603, 604	<i>Invirase(RO)</i>	602
<i>Entyvio(TK)</i>	148, 153, 432, 438	<i>Isentress HD(MK)</i>	615
<i>Epclusa(GI)</i>	102, 386	<i>Isentress(MK)</i>	615, 616
<i>EPOETIN ALFA</i>	40, 324	<i>IVACAFTOR</i>	303, 587
<i>EPOETIN BETA</i>	41, 325	<i>Jadenu(NM)</i>	308, 309, 592, 593
<i>EPOETIN LAMBDA</i>	41, 325	<i>Juluca(VI)</i>	610
<i>EPOPROSTENOL</i>	59, 343	<i>Kaletra(VE)</i>	612
<i>Eprex 1000(JC)</i>	41, 325	<i>Kalydeco(VR)</i>	305, 589
<i>Eprex 10000(JC)</i>	40, 324	<i>Kineret(FK)</i>	243, 527
<i>Eprex 20,000(JC)</i>	41, 325	<i>Kivexa (VI)</i>	608
<i>Eprex 2000(JC)</i>	41, 325	<i>LAMIVUDINE</i>	604
<i>Eprex 3000(JC)</i>	40, 324	<i>LAMIVUDINE + ZIDOVUDINE</i>	611
<i>Eprex 40,000(JC)</i>	41, 325	<i>Lamivudine 150 mg + Zidovudine 300 mg Alphapharm (AF)</i>	612
<i>Eprex 4000(JC)</i>	41, 325	<i>Lamivudine Alphapharm (AF)</i>	604
<i>Eprex 5000(JC)</i>	41, 325	<i>LANREOTIDE</i>	92, 93, 376, 377, 597
<i>Eprex 6000(JC)</i>	40, 324	<i>LANTHANUM</i>	310, 594
<i>Eprex 8000(JC)</i>	40, 324	<i>LEDIPASVIR + SOFOSBUVIR</i>	100, 101, 384, 385
<i>ETANERCEPT</i>	159, 443	<i>Lemtrada(GZ)</i>	132, 133, 416, 417
<i>ETRAVIRINE</i>	607	<i>LENALIDOMIDE</i>	272, 273, 274, 556, 557, 558
<i>EVEROLIMUS</i>		<i>LENOGRASTIM</i>	122, 406
.Highly Specialised Drugs Program (Private Hospital) 141		<i>LEVODOPA + CARBIDOPA</i>	283, 567
.Highly Specialised Drugs Program (Public Hospital) 425		<i>Lioresal Intrathecal (NV)</i>	278, 562
<i>Eviplera(GI)</i>	614	<i>LIPEGILGRASTIM</i>	124, 408
<i>Evotaz(BQ)</i>	600	<i>Liposomal Doxorubicin SUN (RA)</i>	105, 389
<i>Fasenra(AP)</i>	287, 290, 571, 574	<i>Lonquex(TB)</i>	124, 408
<i>Ferriprox(TX)</i>	309, 310, 593	<i>LOPINAVIR + RITONAVIR</i>	612
<i>FILGRASTIM</i>	119, 120, 121, 403, 404, 405	<i>LUMACAFTOR + IVACAFTOR</i>	305, 306, 589, 590
<i>Filolan(GK)</i>	62, 63, 346, 347	<i>LUTROPIN ALFA</i>	904
<i>FOLLITROPIN ALFA</i>	902	<i>Luveris(SG)</i>	904
<i>FOLLITROPIN ALFA + LUTROPIN ALFA</i>	903	<i>Mabthera(RO)</i>	110, 116, 394, 400
<i>FOLLITROPIN BETA</i>	903	<i>MACITENTAN</i>	68, 352
<i>FOLLITROPIN DELTA</i>	903	<i>MANNITOL</i>	303, 587
<i>FOSAMPRENAVIR</i>	601	<i>MARAVIROC</i>	614
<i>Fosrenol(ZI)</i>	310, 594	<i>Maviret(VE)</i>	100, 384
<i>Fuzeon(RO)</i>	614	<i>Menopur 1200(FP)</i>	904
<i>GANCICLOVIR</i>	97, 381, 598	<i>Menopur 600(FP)</i>	904
<i>GANCICLOVIR SXP (HN)</i>	98, 382, 599	<i>MEPOLIZUMAB</i>	290, 292, 574, 576
<i>GANIRELIX</i>	905	<i>METHADONE</i>	
<i>Genotropin GoQuick(PF)</i>	630, 678, 876, 899	.Opiate Dependence Treatment Program	908
<i>Genotropin MiniQuick(PF)</i>	678, 679, 876, 877, 899	<i>METHOXY POLYETHYLENE GLYCOL-EPOETIN BETA42,</i>	326
<i>Genvoya(GI)</i>	614	<i>MIDOSTAURIN</i>	116, 117, 118, 400, 401, 402
<i>GLECAPREVIR + PIBRENTASVIR</i>	99, 100, 383, 384	<i>Mircera(RO)</i>	42, 43, 326, 327
<i>Gonal-f Pen(SG)</i>	902	<i>Movapo Pen(TD)</i>	284, 568
<i>Granocyte 13(PF)</i>	124, 408	<i>Movapo PFS(TD)</i>	283, 567
<i>Granocyte 34(PF)</i>	124, 408	<i>Movapo(TD)</i>	283, 567
<i>Harvoni(GI)</i>	100, 101, 384, 385	<i>Mozobil(GZ)</i>	126, 410
<i>Hepsera (GI)</i>	603	<i>Mycobutin(PF)</i>	97, 381
<i>Hospira Pty Limited(PF)</i>		<i>MycoCept (RF)</i>	142, 426
.Highly Specialised Drugs Program (Private Hospital) 95, 310		<i>MYCOPHENOLATE</i>	141, 142, 425, 426
.Highly Specialised Drugs Program (Public Hospital) 379, 594		<i>Mycophenolate AN(EA)</i>	142, 426
<i>HUMAN CHORIONIC GONADOTROPHIN</i>	903	<i>Mycophenolate Sandoz (SZ)</i>	142, 426
<i>HUMAN MENOPAUSAL GONADOTROPHIN</i>	904	<i>Mycophenolate Sandoz(SZ)</i>	142, 426
<i>Humatrope(LY)</i>	669, 780, 817	<i>Myfortic(NV)</i>	142, 426
<i>Humira(VE)</i>	159, 443	<i>NAFARELIN</i>	904
<i>IBANDRONATE</i>	279, 563	<i>NATALIZUMAB</i>	142, 427
<i>Ibavyr(IX)</i>	101, 385	<i>Neoral 10(NV)</i>	271, 554
<i>ILOPROST</i>	63, 347	<i>Neoral 100 (NV)</i>	271, 555
<i>Imukin(EU)</i>	125, 409	<i>Neoral 25 (NV)</i>	271, 554
<i>INCOBOTULINUMTOXINA</i>	626	<i>Neoral 50 (NV)</i>	271, 555
<i>Inflectra(PF)</i> ...	172, 176, 177, 182, 184, 186, 187, 190, 198, 200, 204, 208, 212, 216, 222, 227, 233, 243, 457, 459, 461, 466, 468, 470, 471, 474, 481, 484, 488, 493, 496, 500, 506, 511, 517, 527	<i>Neoral(NV)</i>	271, 555

NeoRecormon(RO) 41, 325
Neulasta(JU) 124, 408
Neupogen(AN) 121, 122, 405, 406
NEVIRAPINE 607
Nevirapine Alphapharm(AF) 608
Nevirapine XR APOTEX(TX) 607
Nivestim (PF) 121, 122, 405, 406
Nivestim(PF) 120, 404
Norditropin FlexPro(NO) 661, 745, 836
Norditropin SimpleXx(NO) 661, 745, 836
Norvir(VE) 601
Novicrit(SZ) 42, 326
Nplate(AN) 39, 323
Nucala(GK) 292, 294, 576, 578
NUSINERSEN 280, 281, 564, 565
NutropinAq(IS) 630, 661, 745, 836
OCRELIZUMAB 143, 427
Ocrevus(RO) 143, 428
OCTREOTIDE 94, 378, 598
Octreotide (SUN)(RA) 95, 379
Octreotide MaxRx (GQ)
 .Highly Specialised Drugs Program (Private Hospital) . 95
 .Highly Specialised Drugs Program (Public Hospital) . 379
Odefsey(GI) 611
OMALIZUMAB 294, 295, 299, 578, 579, 583
Omnitrope Surepal 10(SZ) 637, 727, 798
Omnitrope Surepal 15(SZ) 637, 727, 798
Omnitrope Surepal 5(SZ) 637, 727, 798
Opsumit(AT) 72, 356
Orencia(BQ) 132, 416
Orgalutran(MK) 905
Oriprio(ON) 901
Orkambi(VR) 306, 307, 590, 591
Ovidrel(SG) 901
Pacrolim(AF) 271, 272, 555
PAMIDRONATE DISODIUM 279, 563
Pamisol(PF) 279, 563
PASIREOTIDE 95, 379
Pegasys(RO) 125, 126, 409, 410
PEGFILGRASTIM 124, 408
PEGINTERFERON ALFA-2A 125, 409
PEGVISOMANT 90, 91, 374, 375
Pergoveris(SG) 903
Pharmacor Mycophenolate 250 (CR) 142, 426
Pharmacor Mycophenolate 500(CR) 142, 426
Pharmacor Tacrolimus 0.5 (CR) 271, 555
Pharmacor Tacrolimus 1 (CR) 272, 555
Pharmacor Tacrolimus 5 (CR) 271, 555
PLERIXAFOR 126, 410
POMALIDOMIDE 276, 560
Pomalyst(CJ) 277, 561
Pregnyl(MK) 904
Prezcobix(JC) 610
Prezista(JC) 600, 601
PROGESTERONE 901
Prograf(LL) 271, 272, 555
Pulmozyme(RO) 303, 587
Puregon 300 IU/0.36 mL(MK) 903
Puregon 600 IU/0.72 mL(MK) 903
Puregon 900 IU/1.08 mL(MK) 903
RALTEGRAVIR 615
Rapamune(PF) 144, 428
Rekovele(FP) 903
Remicade (JC) 176, 182, 204, 208, 212, 216, 222, 227, 233,
 243, 457, 461, 488, 493, 496, 500, 506, 511, 517, 527
Remicade(JC) 166, 168, 170, 175, 180, 192, 195, 450, 452,
 454, 457, 464, 476, 479
Renagel(GZ) 311, 594
Renflexis (MK) 172, 177, 184, 186, 187, 190, 198, 200, 459,
 466, 468, 470, 471, 474, 481, 484

Renflexis(MK) 176, 182, 204, 208, 212, 216, 222, 227, 233,
 243, 457, 461, 488, 493, 496, 500, 506, 511, 517, 527
Retrovir(VI) 606
Revatio (PF) 85, 369
Revestive(ZI) 30, 31, 33, 314, 315, 317
Revlimid(CJ) 273, 274, 275, 276, 557, 558, 559
Revolade(NV) 36, 320
Reyataz (BQ) 599
Reyataz(BQ) 599, 600
RIBAVIRIN 101, 385
RIFABUTIN 97, 381
RILPIVIRINE 608
RIOCIGUAT 72, 75, 356, 359
Ristempa (JO) 124, 408
RITONAVIR 601
RITUXIMAB 105, 107, 108, 110, 389, 391, 392, 394
Riximyo (SZ) 110, 116, 394, 400
Riximyo(SZ) 107, 108, 391, 392
Roferon-A(RO) 124, 125, 408, 409, 616
ROMIPLOSTIM 36, 320
Rydapt(NV) 117, 118, 119, 400, 401, 402, 403
Saizen(SG) 653, 762, 855
Sandimmun(NV) 270, 554
Sandostatin 0.05 (NV) 95, 379
Sandostatin 0.1 (NV) 95, 379
Sandostatin 0.5 (NV) 95, 379
Sandostatin LAR(NV) 94, 378, 598
SAQUINAVIR 601
Scitropin A(SA) 637, 727, 798
SciTropin A(SA) 637, 727, 798
SEVELAMER 310, 594
Signifor LAR(NV) 96, 97, 380, 381
SILDENAFIL
 .Highly Specialised Drugs Program (Private Hospital) ..80
 .Highly Specialised Drugs Program (Public Hospital) .364
Sildenafil AN PHT 20(EA) 85, 369
Sildenafil Sandoz PHT 20(SZ) 85, 369
SILDENAFIL-DRx (RZ) 85, 369
Sintetica Baclofen Intrathecal(BZ) 278, 279, 562
SIROLIMUS 143, 428
SOFOSBUVIR 102, 386
SOFOSBUVIR + VELPATASVIR 102, 386
SOFOSBUVIR + VELPATASVIR + VOXILAPREVIR 102,
 386
Soliris(XI) 134, 135, 137, 141, 418, 419, 421, 425
SOMATROPIN 629, 630, 637, 645, 653, 661, 669, 679, 694,
 710, 727, 745, 763, 780, 798, 817, 836, 855, 877
Somatuline Autogel(IS) 93, 94, 377, 378, 597, 598
Somatuline LA(IS) 93, 377
Somavert(PF) 91, 92, 375, 376
Sovaldi(GI) 102, 386
Spinraza(BD) 281, 283, 565, 567
Stelara(JC) 270, 553
Stocrin(MK) 607
Stribild(GI) 613
Suboxone Film 2/0.5(IR) 908
Suboxone Film 8/2(IR) 908
Subutex(IR) 907
SUCROFERRIC OXYHYDROXIDE 311, 594
Synarel(PF) 904
Tacrograf (RW) 271, 272, 555
TACROLIMUS 271, 555
TACROLIMUS APOTEX(TX) 271, 272, 555, 556
Tacrolimus Sandoz (SZ) 271, 272, 555, 556
Tacrolimus Sandoz(SZ) 271, 555
TADALAFIL
 .Highly Specialised Drugs Program (Private Hospital) ..85
 .Highly Specialised Drugs Program (Public Hospital) .369
TEDUGLUTIDE 30, 31, 314, 315
Telzir(VI) 601

TENOFOVIR + EMTRICITABINE	612	Valcyte(RO)	98, 382, 599
TENOFOVIR + EMTRICITABINE + EFAVIRENZ	613	VALGANCICLOVIR	98, 382, 599
TENOFOVIR + EMTRICITABINE + ELVITEGRAVIR + COBICISTAT	613	Valganciclovir AN (JO)	98, 382, 599
TENOFOVIR + EMTRICITABINE + RILPIVIRINE	613	Valganciclovir Juno(JU)	98, 382, 599
TENOFOVIR ALAFENAMIDE + EMTRICITABINE + ELVITEGRAVIR + COBICISTAT	614	Valganciclovir Mylan (AF)	98, 382, 599
Tenofovir APOTEX(TX)	606	Valganciclovir Sandoz(SZ)	98, 382, 599
TENOFOVIR DISOPROXIL	605	Valtrex(RW)	98, 382
Tenofovir Disoproxil Emtricitabine Mylan 300/200(AF) ..	613	VEDOLIZUMAB	144, 148, 428, 432
Tenofovir Disoproxil Mylan(AF)	606	Veletri(AT)	62, 346
Tenofovir Disoproxil/Emtricitabine/Efavirenz Mylan 300/200/600(AF)	613	Velphoro(VL)	311, 595
Tenofovir EMT GH(GQ)	612	Ventavis(BN)	67, 351
Tenofovir GH(GQ)	606	Versacloz(PF)	284, 568, 617
Tenofovir/Emtricitabine 300/200 APOTEX(TX)	612	Vidaza (CJ)	104, 388
Tezmota(JX)	124, 408	Viramune (BY)	608
THALIDOMIDE	277, 561	Viramune XR (BY)	607
Thalomid(CJ)	277, 561	Viramune(BY)	608
TIPRANAVIR	602	Viread (GI)	606
Tivicay(VI)	614	Volibris(GK)	48, 332
TOCILIZUMAB	243, 249, 254, 260, 527, 533, 538, 544	Vosevi(GI)	102, 386
Tracleer (AT)	59, 343	Xeomin(EZ)	626, 627
Tracleer(AT)	53, 337	Xolair(NV)	294, 295, 299, 302, 578, 579, 583, 586
TRIPTORELIN	905	Zarzio(SZ)	121, 122, 405, 406
Triumeq(VI)	610	Zeffix(RW)	605
Trizivir(VI)	609	Zepatier(MK)	99, 383
Truvada (GI)	612	Zetlam(AF)	604
Tysabri(BD)	143, 427	Ziagen(VI)	602
USTEKINUMAB	266, 550	ZIDOVDINE	606
Utrogestan(HB)	901	Zithromax(PF) .Highly Specialised Drugs Program (Private Hospital) ..	97
VALACICLOVIR	98, 382	.Highly Specialised Drugs Program (Public Hospital) ..	381
Valaciclovir APOTEX (GX)	98, 382	ZOLEDRONIC ACID	279, 563
Valaciclovir RBX(RA)	98, 382	Zomacton(FP)	645, 694, 710
		Zometa (NV)	280, 564