



Australian Government

Department of Health



Schedule of Pharmaceutical Benefits

General Pharmaceutical Schedule - Volume 1

Effective 1 April 2021

This Schedule is also available at www.pbs.gov.au



This work is copyright. Apart from any use as permitted under the *Copyright Act 1968*, no part may be reproduced by any process without prior written permission from the Commonwealth. Requests and inquiries concerning reproduction and rights should be addressed to the Commonwealth Copyright Administration, Attorney General's Department, Robert Garran Offices, National Circuit, Barton ACT 2600 or posted at <http://www.ag.gov.au>

This Schedule provides information on the arrangements for the prescribing and supply of pharmaceutical benefits. These arrangements operate under the *National Health Act 1953*. However, at the time of printing, the relevant legislation giving authority for the changes included in this issue of the Schedule may still be subject to the usual Parliamentary scrutiny. This book is not a legal document, and, in cases of discrepancy, the legislation will be the source document for payment for the supply of pharmaceutical benefits. The legislation is available from the Federal Register of Legislation website at www.legislation.gov.au.

The information is not intended to give or replace any legal, medical, dental or optometrical advice. This document is not a legal document and does not constitute legal advice. Neither the information nor this document can be relied upon without first seeking and obtaining independent legal, medical, dental or optometrical advice beforehand. To the extent permitted by law, the Commonwealth of Australia will not be held responsible, nor accept any liability (whether arising out of negligence or otherwise), for any injury, damages, costs, expenses and losses suffered or incurred by a person where such a person has relied on this document or used the information in it as legal, medical, dental or optometrical advice.

Contents

Fees, Patient Contributions and Safety Net Thresholds	1
Summary of Changes	2
About the Schedule	13
Symbols and Abbreviations Used in the Schedule	13
Restricted Benefits	14
Guidelines and General Statements	15
General Statement for Drugs for the Treatment of Hepatitis C.....	15
Pharmaceutical Benefits Schedules	17
Prescriber Bag	18
General Pharmaceutical Benefits	23
Palliative Care	993
Repatriation Pharmaceutical Benefits Scheme	1006
Extemporaneously Prepared Benefits	1059
Drug Tariff	1060
Container Prices.....	1063
Standard Formula Preparations	1064
Codes, Maximum Quantities, and Number of Repeats for Extemporaneously Prepared Benefits	1065
Index of Manufacturers' Code.....	1066
Generic/Proprietary Index	1069

Fees, Patient Contributions and Safety Net Thresholds

The following fees, patient contributions and safety net thresholds apply as at 1 April 2021 and are included, where applicable, in prices published in the Schedule —

Dispensing Fees:	Ready-prepared	\$7.74
	Dangerous drug fee	\$4.80
	Extemporaneously-prepared	\$9.78
	Allowable additional patient charge*	\$4.42
Additional Fees (for safety net prices):	Ready-prepared	\$1.29
	Extemporaneously-prepared	\$1.66
Patient Co-payments:	General	\$41.30
	Concessional	\$6.60
Safety Net Thresholds:	General	\$1497.20
	Concessional	\$316.80
Safety Net Card Issue Fee:		\$10.34

* The allowable additional patient charge is a discretionary charge to general patients if a pharmaceutical item has a dispensed price for maximum quantity less than the general patient co-payment. The pharmacist may charge general patients the allowable additional fee but the fee cannot take the cost of the prescription above the general patient co-payment for the medicine. This fee does not count towards the Safety Net threshold.

Summary of Changes

These changes to the Schedule of Pharmaceutical Benefits are effective from 1 April 2021. The Schedule is updated on the first day of each month and is available on the internet at www.pbs.gov.au.

Prescriber Bag

Additions

Addition – Item

12222G **FUROSEMIDE (FRUSEMIDE)**, furosemide (frusemide) 20 mg tablet, 50 (*Frusemix-M, Lasix-M, Urex-M*)

General Pharmaceutical Benefits

Additions

Addition – Item

12320K **APOMORPHINE**, apomorphine hydrochloride hemihydrate 20 mg/2 mL injection, 5 x 2 mL ampoules (*Movapo*)

12306Q **APOMORPHINE**, apomorphine hydrochloride hemihydrate 50 mg/5 mL injection, 5 x 5 mL ampoules (*Movapo*)

12319J **APOMORPHINE**, apomorphine hydrochloride hemihydrate 50 mg/10 mL injection, 5 x 10 mL syringes (*Movapo PFS*)

12300J **FULVESTRANT**, fulvestrant 250 mg/5 mL injection, 2 x 5 mL syringes (*Fulvestrant Sandoz*)

12295D **INDACATEROL + GLYCOPYRRONIUM + MOMETASONE**, indacaterol 114 microgram + glycopyrronium 46 microgram + mometasone furoate 136 microgram powder for inhalation, 30 capsules (*Enerzair Breezhaler*)

12298G **INDACATEROL + GLYCOPYRRONIUM + MOMETASONE**, indacaterol 114 microgram + glycopyrronium 46 microgram + mometasone furoate 68 microgram powder for inhalation, 30 capsules (*Enerzair Breezhaler*)

12311Y **OXYCODONE**, oxycodone hydrochloride 5 mg capsule, 10 (*OxyNorm*)

12314D **OXYCODONE**, oxycodone hydrochloride 5 mg capsule, 10 (*OxyNorm*)

12301K **ROMOSOZUMAB**, romosozumab 105 mg/1.17 mL injection, 2 x 1.17 mL syringes (*Evenity*)

12297F **SECUKINUMAB**, secukinumab 150 mg/mL injection, 1 mL pen device (*Cosentyx*)

12307R **SECUKINUMAB**, secukinumab 150 mg/mL injection, 1 mL pen device (*Cosentyx*)

12321L **SECUKINUMAB**, secukinumab 150 mg/mL injection, 1 mL pen device (*Cosentyx*)

12299H **SERTRALINE**, sertraline 50 mg tablet, 30 (*Sertraline tablets USP (Medsurge)*)

12305P **SERTRALINE**, sertraline 50 mg tablet, 30 (*Sertraline tablets USP (Medsurge)*)

12296E **SERTRALINE**, sertraline 100 mg tablet, 30 (*Sertraline tablets USP (Medsurge)*)

12317G **SERTRALINE**, sertraline 100 mg tablet, 30 (*Sertraline tablets USP (Medsurge)*)

Addition – Brand

10756D *Buprenorphine Sandoz, SZ* – **BUPRENORPHINE**, buprenorphine 25 microgram/hour patch, 2

10755C *Buprenorphine Sandoz, SZ* – **BUPRENORPHINE**, buprenorphine 30 microgram/hour patch, 2

10746N *Buprenorphine Sandoz, SZ* – **BUPRENORPHINE**, buprenorphine 40 microgram/hour patch, 2

9157Y *Cinacalcet Mylan, AF* – **CINACALCET**, cinacalcet 30 mg tablet, 28

9158B *Cinacalcet Mylan, AF* – **CINACALCET**, cinacalcet 60 mg tablet, 28

9159C *Cinacalcet Mylan, AF* – **CINACALCET**, cinacalcet 90 mg tablet, 28

10022L	<i>Erlotinib Sandoz, SZ – ERLOTINIB, erlotinib 25 mg tablet, 30</i>
10028T	<i>Erlotinib Sandoz, SZ – ERLOTINIB, erlotinib 25 mg tablet, 30</i>
11263T	<i>Erlotinib Sandoz, SZ – ERLOTINIB, erlotinib 25 mg tablet, 30</i>
10019H	<i>Erlotinib Sandoz, SZ – ERLOTINIB, erlotinib 100 mg tablet, 30</i>
10020J	<i>Erlotinib Sandoz, SZ – ERLOTINIB, erlotinib 100 mg tablet, 30</i>
11260P	<i>Erlotinib Sandoz, SZ – ERLOTINIB, erlotinib 100 mg tablet, 30</i>
10014C	<i>Erlotinib Sandoz, SZ – ERLOTINIB, erlotinib 150 mg tablet, 30</i>
10025P	<i>Erlotinib Sandoz, SZ – ERLOTINIB, erlotinib 150 mg tablet, 30</i>
11259N	<i>Erlotinib Sandoz, SZ – ERLOTINIB, erlotinib 150 mg tablet, 30</i>
11692J	<i>APO-Esomeprazole, TY – ESOMEPRAZOLE, esomeprazole 20 mg enteric tablet, 30</i>
11692J	<i>Esopreze, BG – ESOMEPRAZOLE, esomeprazole 20 mg enteric tablet, 30</i>
12287Q	<i>APO-Esomeprazole, TY – ESOMEPRAZOLE, esomeprazole 20 mg enteric tablet, 30</i>
12287Q	<i>Esopreze, BG – ESOMEPRAZOLE, esomeprazole 20 mg enteric tablet, 30</i>
8600P	<i>APO-Esomeprazole, TY – ESOMEPRAZOLE, esomeprazole 20 mg enteric tablet, 30</i>
8600P	<i>Esopreze, BG – ESOMEPRAZOLE, esomeprazole 20 mg enteric tablet, 30</i>
8886Q	<i>APO-Esomeprazole, TY – ESOMEPRAZOLE, esomeprazole 20 mg enteric tablet, 30</i>
8886Q	<i>Esopreze, BG – ESOMEPRAZOLE, esomeprazole 20 mg enteric tablet, 30</i>
12283L	<i>APO-Esomeprazole, TY – ESOMEPRAZOLE, esomeprazole 40 mg enteric tablet, 30</i>
12283L	<i>Esopreze, BG – ESOMEPRAZOLE, esomeprazole 40 mg enteric tablet, 30</i>
3401B	<i>APO-Esomeprazole, TY – ESOMEPRAZOLE, esomeprazole 40 mg enteric tablet, 30</i>
3401B	<i>Esopreze, BG – ESOMEPRAZOLE, esomeprazole 40 mg enteric tablet, 30</i>
8601Q	<i>APO-Esomeprazole, TY – ESOMEPRAZOLE, esomeprazole 40 mg enteric tablet, 30</i>
8601Q	<i>Esopreze, BG – ESOMEPRAZOLE, esomeprazole 40 mg enteric tablet, 30</i>
8516F	<i>Axotide Junior, GC – FLUTICASONE PROPIONATE, fluticasone propionate 50 microgram/actuation inhalation, 120 actuations</i>
8147T	<i>Axotide Junior Accuhaler, GC – FLUTICASONE PROPIONATE, fluticasone propionate 100 microgram/actuation powder for inhalation, 60 actuations</i>
8148W	<i>Axotide Accuhaler, GC – FLUTICASONE PROPIONATE, fluticasone propionate 250 microgram/actuation powder for inhalation, 60 actuations</i>
8517G	<i>PAVTIDE MDI 50/25, GC – FLUTICASONE PROPIONATE + SALMETEROL, fluticasone propionate 50 microgram/actuation + salmeterol 25 microgram/actuation inhalation, 120 actuations</i>
8430Q	<i>PAVTIDE ACCUHALER 100/50, GC – FLUTICASONE PROPIONATE + SALMETEROL, fluticasone propionate 100 microgram/actuation + salmeterol 50 microgram/actuation powder for inhalation, 60 actuations</i>
8431R	<i>PAVTIDE ACCUHALER 250/50, GC – FLUTICASONE PROPIONATE + SALMETEROL, fluticasone propionate 250 microgram/actuation + salmeterol 50 microgram/actuation powder for inhalation, 60 actuations</i>
8432T	<i>PAVTIDE ACCUHALER 500/50, GC – FLUTICASONE PROPIONATE + SALMETEROL, fluticasone propionate 500 microgram/actuation + salmeterol 50 microgram/actuation powder for inhalation, 60 actuations</i>
8505P	<i>APX-Gabapentin, GX – GABAPENTIN, gabapentin 100 mg capsule, 100</i>
1834M	<i>APX-Gabapentin, GX – GABAPENTIN, gabapentin 300 mg capsule, 100</i>
1835N	<i>APX-Gabapentin, GX – GABAPENTIN, gabapentin 400 mg capsule, 100</i>
9316H	<i>Nepiten, AF – NEBIVOLOL, nebivolol 1.25 mg tablet, 28</i>
9311C	<i>Nepiten, AF – NEBIVOLOL, nebivolol 5 mg tablet, 28</i>
9312D	<i>Nepiten, AF – NEBIVOLOL, nebivolol 10 mg tablet, 28</i>
11683X	<i>Maxor EC Tabs, AF – OMEPRAZOLE, omeprazole 20 mg enteric tablet, 30</i>
12272X	<i>Maxor EC Tabs, AF – OMEPRAZOLE, omeprazole 20 mg enteric tablet, 30</i>
8331L	<i>Maxor EC Tabs, AF – OMEPRAZOLE, omeprazole 20 mg enteric tablet, 30</i>
8333N	<i>Maxor EC Tabs, AF – OMEPRAZOLE, omeprazole 20 mg enteric tablet, 30</i>

10460M	<i>Posaconazole ARX, XT</i> – POSACONAZOLE , posaconazole 100 mg modified release tablet, 24
2590H	<i>Rosuvastatin Lupin, GQ</i> – ROSUVASTATIN , rosuvastatin 5 mg tablet, 30
2606E	<i>Rosuvastatin Lupin, GQ</i> – ROSUVASTATIN , rosuvastatin 5 mg tablet, 30
2584B	<i>Rosuvastatin Lupin, GQ</i> – ROSUVASTATIN , rosuvastatin 10 mg tablet, 30
2628H	<i>Rosuvastatin Lupin, GQ</i> – ROSUVASTATIN , rosuvastatin 10 mg tablet, 30
2574L	<i>Rosuvastatin Lupin, GQ</i> – ROSUVASTATIN , rosuvastatin 20 mg tablet, 30
2609H	<i>Rosuvastatin Lupin, GQ</i> – ROSUVASTATIN , rosuvastatin 20 mg tablet, 30
2594M	<i>Rosuvastatin Lupin, GQ</i> – ROSUVASTATIN , rosuvastatin 40 mg tablet, 30
2636R	<i>Rosuvastatin Lupin, GQ</i> – ROSUVASTATIN , rosuvastatin 40 mg tablet, 30
2898M	<i>Terimide, AF</i> – TERIFLUNOMIDE , teriflunomide 14 mg tablet, 28
9288W	<i>Zoledronic Acid SUN, RA</i> – ZOLEDRONIC ACID , zoledronic acid 5 mg/100 mL injection, 100 mL vial
9350D	<i>Zoledronic Acid SUN, RA</i> – ZOLEDRONIC ACID , zoledronic acid 5 mg/100 mL injection, 100 mL vial

Addition – Equivalence Indicator

9157Y	<i>Pharmacor Cinacalcet, CR</i> – CINACALCET , cinacalcet 30 mg tablet, 28
9158B	<i>Pharmacor Cinacalcet, CR</i> – CINACALCET , cinacalcet 60 mg tablet, 28
9159C	<i>Pharmacor Cinacalcet, CR</i> – CINACALCET , cinacalcet 90 mg tablet, 28
8516F	<i>Flixotide Junior, GK</i> – FLUTICASONE PROPIONATE , fluticasone propionate 50 microgram/actuation inhalation, 120 actuations
8147T	<i>Flixotide Junior Accuhaler, GK</i> – FLUTICASONE PROPIONATE , fluticasone propionate 100 microgram/actuation powder for inhalation, 60 actuations
8148W	<i>Flixotide Accuhaler, GK</i> – FLUTICASONE PROPIONATE , fluticasone propionate 250 microgram/actuation powder for inhalation, 60 actuations
8517G	<i>Seretide MDI 50/25, GK</i> – FLUTICASONE PROPIONATE + SALMETEROL , fluticasone propionate 50 microgram/actuation + salmeterol 25 microgram/actuation inhalation, 120 actuations
8430Q	<i>Seretide Accuhaler 100/50, GK</i> – FLUTICASONE PROPIONATE + SALMETEROL , fluticasone propionate 100 microgram/actuation + salmeterol 50 microgram/actuation powder for inhalation, 60 actuations
8431R	<i>Seretide Accuhaler 250/50, GK</i> – FLUTICASONE PROPIONATE + SALMETEROL , fluticasone propionate 250 microgram/actuation + salmeterol 50 microgram/actuation powder for inhalation, 60 actuations
8432T	<i>Seretide Accuhaler 500/50, GK</i> – FLUTICASONE PROPIONATE + SALMETEROL , fluticasone propionate 500 microgram/actuation + salmeterol 50 microgram/actuation powder for inhalation, 60 actuations

Addition – Note

11361Y	GOLIMUMAB , golimumab 50 mg/0.5 mL injection, 0.5 mL pen device (<i>Simponi</i>)
11365E	GOLIMUMAB , golimumab 50 mg/0.5 mL injection, 0.5 mL pen device (<i>Simponi</i>)
11372M	GOLIMUMAB , golimumab 50 mg/0.5 mL injection, 0.5 mL pen device (<i>Simponi</i>)
11373N	GOLIMUMAB , golimumab 50 mg/0.5 mL injection, 0.5 mL pen device (<i>Simponi</i>)
11375Q	GOLIMUMAB , golimumab 50 mg/0.5 mL injection, 0.5 mL pen device (<i>Simponi</i>)
11376R	GOLIMUMAB , golimumab 50 mg/0.5 mL injection, 0.5 mL pen device (<i>Simponi</i>)
3426H	GOLIMUMAB , golimumab 50 mg/0.5 mL injection, 0.5 mL syringe (<i>Simponi</i>)
3428K	GOLIMUMAB , golimumab 50 mg/0.5 mL injection, 0.5 mL syringe (<i>Simponi</i>)
3430M	GOLIMUMAB , golimumab 50 mg/0.5 mL injection, 0.5 mL syringe (<i>Simponi</i>)
3432P	GOLIMUMAB , golimumab 50 mg/0.5 mL injection, 0.5 mL syringe (<i>Simponi</i>)
3434R	GOLIMUMAB , golimumab 50 mg/0.5 mL injection, 0.5 mL syringe (<i>Simponi</i>)
3436W	GOLIMUMAB , golimumab 50 mg/0.5 mL injection, 0.5 mL syringe (<i>Simponi</i>)
11381B	GOLIMUMAB , golimumab 100 mg/mL injection, 1 mL pen device (<i>Simponi</i>)
11382C	GOLIMUMAB , golimumab 100 mg/mL injection, 1 mL pen device (<i>Simponi</i>)
11502J	GOLIMUMAB , golimumab 100 mg/mL injection, 1 mL pen device (<i>Simponi</i>)

11385F	RIBOCICLIB , ribociclib 200 mg tablet, 21 (<i>Kisqali</i>)
11386G	RIBOCICLIB , ribociclib 200 mg tablet, 63 (<i>Kisqali</i>)
11397W	RIBOCICLIB , ribociclib 200 mg tablet, 42 (<i>Kisqali</i>)
10703H	RITUXIMAB , rituximab 1.4 g/11.7 mL injection, 11.7 mL vial (<i>Mabthera SC</i>)
10709P	RITUXIMAB , rituximab 1.4 g/11.7 mL injection, 11.7 mL vial (<i>Mabthera SC</i>)
10719E	RITUXIMAB , rituximab 1.4 g/11.7 mL injection, 11.7 mL vial (<i>Mabthera SC</i>)
10742J	RITUXIMAB , rituximab 1.4 g/11.7 mL injection, 11.7 mL vial (<i>Mabthera SC</i>)
11988Y	RITUXIMAB , rituximab 1.4 g/11.7 mL injection, 11.7 mL vial (<i>Mabthera SC</i>)
10425Q	SECUKINUMAB , secukinumab 150 mg/mL injection, 2 x 1 mL pen devices (<i>Cosentyx</i>)
10494H	SECUKINUMAB , secukinumab 150 mg/mL injection, 2 x 1 mL pen devices (<i>Cosentyx</i>)
10890E	SECUKINUMAB , secukinumab 150 mg/mL injection, 1 mL pen device (<i>Cosentyx</i>)
10893H	SECUKINUMAB , secukinumab 150 mg/mL injection, 1 mL pen device (<i>Cosentyx</i>)
10894J	SECUKINUMAB , secukinumab 150 mg/mL injection, 2 x 1 mL pen devices (<i>Cosentyx</i>)
10895K	SECUKINUMAB , secukinumab 150 mg/mL injection, 1 mL pen device (<i>Cosentyx</i>)
10898N	SECUKINUMAB , secukinumab 150 mg/mL injection, 1 mL pen device (<i>Cosentyx</i>)
10899P	SECUKINUMAB , secukinumab 150 mg/mL injection, 2 x 1 mL pen devices (<i>Cosentyx</i>)
10900Q	SECUKINUMAB , secukinumab 150 mg/mL injection, 1 mL pen device (<i>Cosentyx</i>)
10901R	SECUKINUMAB , secukinumab 150 mg/mL injection, 2 x 1 mL pen devices (<i>Cosentyx</i>)
10906B	SECUKINUMAB , secukinumab 150 mg/mL injection, 1 mL pen device (<i>Cosentyx</i>)
10910F	SECUKINUMAB , secukinumab 150 mg/mL injection, 2 x 1 mL pen devices (<i>Cosentyx</i>)
2236Q	SERTRALINE , sertraline 50 mg tablet, 30 (<i>APO-Sertraline, Auro-Sertraline 50, Eleva 50, NOUMED SERTRALINE, Sertra 50, Sertraline AN, Sertraline Sandoz, Sertraline generichealth, Setrona, Zolof</i>)
8836C	SERTRALINE , sertraline 50 mg tablet, 30 (<i>Auro-Sertraline 50, Eleva 50, NOUMED SERTRALINE, Sertraline AN, Zolof</i>)
2237R	SERTRALINE , sertraline 100 mg tablet, 30 (<i>APO-Sertraline, Auro-Sertraline 100, Eleva 100, NOUMED SERTRALINE, Sertra 100, Sertraline AN, Sertraline Sandoz, Sertraline generichealth, Setrona, Zolof</i>)
8837D	SERTRALINE , sertraline 100 mg tablet, 30 (<i>Auro-Sertraline 100, Eleva 100, NOUMED SERTRALINE, Sertraline AN, Zolof</i>)

Addition – Restriction

9438R	AMINO ACID FORMULA WITH VITAMINS AND MINERALS WITHOUT LYSINE AND LOW IN TRYPTOPHAN , amino acid formula with vitamins and minerals without lysine and low in tryptophan powder for oral liquid, 30 x 24 g sachets (<i>GA gel</i>)
5484P	AMINO ACID FORMULA WITH VITAMINS AND MINERALS WITHOUT LYSINE AND LOW IN TRYPTOPHAN , amino acid formula with vitamins and minerals without lysine and low in tryptophan powder for oral liquid, 30 x 25 g sachets (<i>GA express 15</i>)

Deletions

Deletion – Item

2002J	CEFUROXIME , cefuroxime 125 mg/5 mL powder for oral liquid, 70 mL (<i>Zinnat</i>)
5499K	CEFUROXIME , cefuroxime 125 mg/5 mL powder for oral liquid, 70 mL (<i>Zinnat</i>)
12025X	OXYCODONE , oxycodone hydrochloride 5 mg capsule, 20 (<i>OxyNorm, Oxycodone BNM</i>)
12044X	OXYCODONE , oxycodone hydrochloride 5 mg capsule, 20 (<i>OxyNorm, Oxycodone BNM</i>)

Deletion – Brand

1148K	<i>Zedace, AF</i> – CAPTOPRIL , captopril 25 mg tablet, 90
1149L	<i>Zedace, AF</i> – CAPTOPRIL , captopril 50 mg tablet, 90
10022L	<i>Tarceva, RO</i> – ERLOTINIB , erlotinib 25 mg tablet, 30
10028T	<i>Tarceva, RO</i> – ERLOTINIB , erlotinib 25 mg tablet, 30
11263T	<i>Tarceva, RO</i> – ERLOTINIB , erlotinib 25 mg tablet, 30

10019H	<i>Tarceva, RO</i> – ERLOTINIB , erlotinib 100 mg tablet, 30
10020J	<i>Tarceva, RO</i> – ERLOTINIB , erlotinib 100 mg tablet, 30
11260P	<i>Tarceva, RO</i> – ERLOTINIB , erlotinib 100 mg tablet, 30
10014C	<i>Tarceva, RO</i> – ERLOTINIB , erlotinib 150 mg tablet, 30
10025P	<i>Tarceva, RO</i> – ERLOTINIB , erlotinib 150 mg tablet, 30
11259N	<i>Tarceva, RO</i> – ERLOTINIB , erlotinib 150 mg tablet, 30

Alterations

Alteration – Brand Name

From

2138M *Fawns and McAllan Proprietary Limited, AS* – **PHENOBARBITAL (PHENOBARBITONE)**, phenobarbital (phenobarbitone) sodium 219 mg/mL injection, 5 x 1 mL ampoules

To

2138M *Phenobarbitone Injection (Aspen Pharmacare Australia Pty Ltd), AS* – **PHENOBARBITAL (PHENOBARBITONE)**, phenobarbital (phenobarbitone) sodium 219 mg/mL injection, 5 x 1 mL ampoules

Alteration – Note

12005W	CERTOLIZUMAB PEGOL , certolizumab pegol 200 mg/mL injection, 2 x 1 mL syringes (<i>Cimzia</i>)
12040Q	CERTOLIZUMAB PEGOL , certolizumab pegol 200 mg/mL injection, 2 x 1 mL syringes (<i>Cimzia</i>)
12063X	CERTOLIZUMAB PEGOL , certolizumab pegol 200 mg/mL injection, 2 x 1 mL syringes (<i>Cimzia</i>)
12013G	CERTOLIZUMAB PEGOL , certolizumab pegol 200 mg/mL injection, 2 x 1 mL pen devices (<i>Cimzia</i>)
12027B	CERTOLIZUMAB PEGOL , certolizumab pegol 200 mg/mL injection, 2 x 1 mL pen devices (<i>Cimzia</i>)
12028C	CERTOLIZUMAB PEGOL , certolizumab pegol 200 mg/mL injection, 2 x 1 mL pen devices (<i>Cimzia</i>)
12291X	DUPILUMAB , dupilumab 200 mg/1.14 mL injection, 2 x 1.14 mL syringes (<i>Dupixent</i>)
12292Y	DUPILUMAB , dupilumab 300 mg/2 mL injection, 2 x 2 mL syringes (<i>Dupixent</i>)
11516D	GOLIMUMAB , golimumab 50 mg/0.5 mL injection, 0.5 mL syringe (<i>Simponi</i>)
11521J	GOLIMUMAB , golimumab 50 mg/0.5 mL injection, 0.5 mL pen device (<i>Simponi</i>)
11538G	GOLIMUMAB , golimumab 50 mg/0.5 mL injection, 0.5 mL pen device (<i>Simponi</i>)
11560K	GOLIMUMAB , golimumab 50 mg/0.5 mL injection, 0.5 mL syringe (<i>Simponi</i>)
9411H	TERIPARATIDE , teriparatide 250 microgram/mL injection, 2.4 mL pen device (<i>Forteo</i>)

Alteration – Restriction

11442F	BARICITINIB , baricitinib 2 mg tablet, 28 (<i>Olumiant</i>)
11443G	BARICITINIB , baricitinib 4 mg tablet, 28 (<i>Olumiant</i>)
12063X	CERTOLIZUMAB PEGOL , certolizumab pegol 200 mg/mL injection, 2 x 1 mL syringes (<i>Cimzia</i>)
12027B	CERTOLIZUMAB PEGOL , certolizumab pegol 200 mg/mL injection, 2 x 1 mL pen devices (<i>Cimzia</i>)
12291X	DUPILUMAB , dupilumab 200 mg/1.14 mL injection, 2 x 1.14 mL syringes (<i>Dupixent</i>)
12292Y	DUPILUMAB , dupilumab 300 mg/2 mL injection, 2 x 2 mL syringes (<i>Dupixent</i>)
11538G	GOLIMUMAB , golimumab 50 mg/0.5 mL injection, 0.5 mL pen device (<i>Simponi</i>)
11560K	GOLIMUMAB , golimumab 50 mg/0.5 mL injection, 0.5 mL syringe (<i>Simponi</i>)
11385F	RIBOCICLIB , ribociclib 200 mg tablet, 21 (<i>Kisqali</i>)
11386G	RIBOCICLIB , ribociclib 200 mg tablet, 63 (<i>Kisqali</i>)
11397W	RIBOCICLIB , ribociclib 200 mg tablet, 42 (<i>Kisqali</i>)
9411H	TERIPARATIDE , teriparatide 250 microgram/mL injection, 2.4 mL pen device (<i>Forteo</i>)

Supply Only

From 1 November 2020 when a product is deleted from the Schedule it may now be available under new Supply Only rules. Supply Only items/brands are available on the Schedule for dispensing but not for prescribing, usually for a period of up to 12 months from when it is deleted.

Substitution of Supply Only items/brands with products flagged as "equivalent for substitution" still apply as specified in the Schedule at the time the script was written. Further information on Supply Only arrangements is available at www.pbs.gov.au

Supply Only commencing April 2021

10687L	<i>Fraxiparine</i> , AS – NADROPARIN , nadroparin calcium 1900 anti-Xa units/0.2 mL injection, 2 x 0.2 mL syringes
10735B	<i>Fraxiparine</i> , AS – NADROPARIN , nadroparin calcium 1900 anti-Xa units/0.2 mL injection, 2 x 0.2 mL syringes
10686K	<i>Fraxiparine</i> , AS – NADROPARIN , nadroparin calcium 2850 anti-Xa units/0.3 mL injection, 2 x 0.3 mL syringes
10701F	<i>Fraxiparine</i> , AS – NADROPARIN , nadroparin calcium 2850 anti-Xa units/0.3 mL injection, 2 x 0.3 mL syringes
10685J	<i>Fraxiparine</i> , AS – NADROPARIN , nadroparin calcium 3800 anti-Xa units/0.4 mL injection, 2 x 0.4 mL syringes
10717C	<i>Fraxiparine</i> , AS – NADROPARIN , nadroparin calcium 3800 anti-Xa units/0.4 mL injection, 2 x 0.4 mL syringes
10716B	<i>Fraxiparine</i> , AS – NADROPARIN , nadroparin calcium 5700 anti-Xa units/0.6 mL injection, 2 x 0.6 mL syringes
10718D	<i>Fraxiparine</i> , AS – NADROPARIN , nadroparin calcium 5700 anti-Xa units/0.6 mL injection, 2 x 0.6 mL syringes
10734Y	<i>Fraxiparine</i> , AS – NADROPARIN , nadroparin calcium 7600 anti-Xa units/0.8 mL injection, 2 x 0.8 mL syringes
10740G	<i>Fraxiparine</i> , AS – NADROPARIN , nadroparin calcium 7600 anti-Xa units/0.8 mL injection, 2 x 0.8 mL syringes
10702G	<i>Fraxiparine</i> , AS – NADROPARIN , nadroparin calcium 9500 anti-Xa units/mL injection, 2 x 1 mL syringes
10733X	<i>Fraxiparine</i> , AS – NADROPARIN , nadroparin calcium 9500 anti-Xa units/mL injection, 2 x 1 mL syringes
10706L	<i>Fraxiparine Forte</i> , AS – NADROPARIN , nadroparin calcium 11 400 anti-Xa units/0.6 mL injection, 2 x 0.6 mL syringes
10725L	<i>Fraxiparine Forte</i> , AS – NADROPARIN , nadroparin calcium 15 200 anti-Xa units/0.8 mL injection, 2 x 0.8 mL syringes
10707M	<i>Fraxiparine Forte</i> , AS – NADROPARIN , nadroparin calcium 19 000 anti-Xa units/mL injection, 2 x 1 mL syringes

Advance Notices

1 May 2021

Deletion – Brand

9155W	<i>Andeptra</i> , EL – DULOXETINE , duloxetine 30 mg enteric capsule, 28
9156X	<i>Andeptra</i> , EL – DULOXETINE , duloxetine 60 mg enteric capsule, 28
3050M	<i>NOUMED PERINDOPRIL</i> , VO – PERINDOPRIL , perindopril erbumine 2 mg tablet, 30
3051N	<i>NOUMED PERINDOPRIL</i> , VO – PERINDOPRIL , perindopril erbumine 4 mg tablet, 30
8704D	<i>NOUMED PERINDOPRIL</i> , VO – PERINDOPRIL , perindopril erbumine 8 mg tablet, 30
2011W	<i>Ransim</i> , RA – SIMVASTATIN , simvastatin 10 mg tablet, 30
8313M	<i>Ransim</i> , RA – SIMVASTATIN , simvastatin 80 mg tablet, 30
9242K	<i>Ransim</i> , RA – SIMVASTATIN , simvastatin 10 mg tablet, 30
9245N	<i>Ransim</i> , RA – SIMVASTATIN , simvastatin 80 mg tablet, 30

1 October 2021

Deletion – Brand

10703H	<i>Mabthera SC</i> , RO – RITUXIMAB , rituximab 1.4 g/11.7 mL injection, 11.7 mL vial
10709P	<i>Mabthera SC</i> , RO – RITUXIMAB , rituximab 1.4 g/11.7 mL injection, 11.7 mL vial
10719E	<i>Mabthera SC</i> , RO – RITUXIMAB , rituximab 1.4 g/11.7 mL injection, 11.7 mL vial
10742J	<i>Mabthera SC</i> , RO – RITUXIMAB , rituximab 1.4 g/11.7 mL injection, 11.7 mL vial
11988Y	<i>Mabthera SC</i> , RO – RITUXIMAB , rituximab 1.4 g/11.7 mL injection, 11.7 mL vial

Palliative Care

Additions

Addition – Brand

- 10964C *Buprenorphine Sandoz, SZ* – **BUPRENORPHINE**, buprenorphine 25 microgram/hour patch, 2
- 10949G *Buprenorphine Sandoz, SZ* – **BUPRENORPHINE**, buprenorphine 30 microgram/hour patch, 2
- 10959T *Buprenorphine Sandoz, SZ* – **BUPRENORPHINE**, buprenorphine 40 microgram/hour patch, 2

Supply Only

From 1 November 2020 when a product is deleted from the Schedule it may now be available under new Supply Only rules. Supply Only items/brands are available on the Schedule for dispensing but not for prescribing, usually for a period of up to 12 months from when it is deleted.

Substitution of Supply Only items/brands with products flagged as "equivalent for substitution" still apply as specified in the Schedule at the time the script was written. Further information on Supply Only arrangements is available at www.pbs.gov.au

Supply Only commencing April 2021

- 5319Y *Panadol, GC* – **PARACETAMOL**, paracetamol 500 mg suppository, 24

Highly Specialised Drugs Program (Private Hospital)

Additions

Addition – Item

- 12313C **DUPILUMAB**, dupilumab 200 mg/1.14 mL injection, 2 x 1.14 mL syringes (*Dupixent*)
- 12316F **DUPILUMAB**, dupilumab 200 mg/1.14 mL injection, 2 x 1.14 mL syringes (*Dupixent*)
- 12294C **DUPILUMAB**, dupilumab 300 mg/2 mL injection, 2 x 2 mL syringes (*Dupixent*)
- 12310X **DUPILUMAB**, dupilumab 300 mg/2 mL injection, 2 x 2 mL syringes (*Dupixent*)

Addition – Brand

- 11891W *Cinacalcet Mylan, AF* – **CINACALCET**, cinacalcet 30 mg tablet, 28
- 9625N *Cinacalcet Mylan, AF* – **CINACALCET**, cinacalcet 30 mg tablet, 28
- 11889R *Cinacalcet Mylan, AF* – **CINACALCET**, cinacalcet 60 mg tablet, 28
- 9626P *Cinacalcet Mylan, AF* – **CINACALCET**, cinacalcet 60 mg tablet, 28
- 11888Q *Cinacalcet Mylan, AF* – **CINACALCET**, cinacalcet 90 mg tablet, 28
- 9627Q *Cinacalcet Mylan, AF* – **CINACALCET**, cinacalcet 90 mg tablet, 28

Addition – Equivalence Indicator

- 11891W *Pharmacor Cinacalcet, CR* – **CINACALCET**, cinacalcet 30 mg tablet, 28
- 9625N *Pharmacor Cinacalcet, CR* – **CINACALCET**, cinacalcet 30 mg tablet, 28
- 11889R *Pharmacor Cinacalcet, CR* – **CINACALCET**, cinacalcet 60 mg tablet, 28
- 9626P *Pharmacor Cinacalcet, CR* – **CINACALCET**, cinacalcet 60 mg tablet, 28
- 11888Q *Pharmacor Cinacalcet, CR* – **CINACALCET**, cinacalcet 90 mg tablet, 28
- 9627Q *Pharmacor Cinacalcet, CR* – **CINACALCET**, cinacalcet 90 mg tablet, 28

Deletions

Deletion – Brand

- 10583B *Mabthera, RO* – **RITUXIMAB**, rituximab 100 mg/10 mL injection, 2 x 10 mL vials
- 10576P *Mabthera, RO* – **RITUXIMAB**, rituximab 500 mg/50 mL injection, 50 mL vial
- 9611W *Mabthera, RO* – **RITUXIMAB**, rituximab 500 mg/50 mL injection, 50 mL vial

Alterations

Alteration – Note

- 11504L **BENRALIZUMAB**, benralizumab 30 mg/mL injection, 1 mL syringe (*Fasenra*)
- 11523L **BENRALIZUMAB**, benralizumab 30 mg/mL injection, 1 mL syringe (*Fasenra*)
- 11847M **BENRALIZUMAB**, benralizumab 30 mg/mL injection, 1 mL syringe (*Fasenra*)
- 11996J **BENRALIZUMAB**, benralizumab 30 mg/mL injection, 1 mL pen device (*Fasenra Pen*)
- 11997K **BENRALIZUMAB**, benralizumab 30 mg/mL injection, 1 mL pen device (*Fasenra Pen*)

11999M	BENRALIZUMAB , benralizumab 30 mg/mL injection, 1 mL pen device (<i>Fasenra Pen</i>)
12030E	MEPOLIZUMAB , mepolizumab 100 mg/mL injection, 1 mL pen device (<i>Nucala</i>)
12043W	MEPOLIZUMAB , mepolizumab 100 mg/mL injection, 1 mL pen device (<i>Nucala</i>)
12051G	MEPOLIZUMAB , mepolizumab 100 mg/mL injection, 1 mL pen device (<i>Nucala</i>)
12052H	MEPOLIZUMAB , mepolizumab 100 mg/mL injection, 1 mL pen device (<i>Nucala</i>)
11003D	MEPOLIZUMAB , mepolizumab 100 mg injection, 1 vial (<i>Nucala</i>)
11014Q	MEPOLIZUMAB , mepolizumab 100 mg injection, 1 vial (<i>Nucala</i>)
11829N	MEPOLIZUMAB , mepolizumab 100 mg injection, 1 vial (<i>Nucala</i>)
10110D	OMALIZUMAB , omalizumab 75 mg/0.5 mL injection, 0.5 mL syringe (<i>Xolair</i>)
11826K	OMALIZUMAB , omalizumab 75 mg/0.5 mL injection, 0.5 mL syringe (<i>Xolair</i>)
11840E	OMALIZUMAB , omalizumab 75 mg/0.5 mL injection, 0.5 mL syringe (<i>Xolair</i>)
10122R	OMALIZUMAB , omalizumab 150 mg/mL injection, 1 mL syringe (<i>Xolair</i>)
11825J	OMALIZUMAB , omalizumab 150 mg/mL injection, 1 mL syringe (<i>Xolair</i>)
11864K	OMALIZUMAB , omalizumab 150 mg/mL injection, 1 mL syringe (<i>Xolair</i>)

Alteration – Restriction

9607P	APOMORPHINE , apomorphine hydrochloride hemihydrate 20 mg/2 mL injection, 5 x 2 mL ampoules (<i>Movapo</i>)
9640J	APOMORPHINE , apomorphine hydrochloride hemihydrate 50 mg/5 mL injection, 5 x 5 mL ampoules (<i>Movapo</i>)
10971K	APOMORPHINE , apomorphine hydrochloride hemihydrate 50 mg/10 mL injection, 5 x 10 mL syringes (<i>Movapo PFS</i>)
11523L	BENRALIZUMAB , benralizumab 30 mg/mL injection, 1 mL syringe (<i>Fasenra</i>)
11997K	BENRALIZUMAB , benralizumab 30 mg/mL injection, 1 mL pen device (<i>Fasenra Pen</i>)
12051G	MEPOLIZUMAB , mepolizumab 100 mg/mL injection, 1 mL pen device (<i>Nucala</i>)
11003D	MEPOLIZUMAB , mepolizumab 100 mg injection, 1 vial (<i>Nucala</i>)
10110D	OMALIZUMAB , omalizumab 75 mg/0.5 mL injection, 0.5 mL syringe (<i>Xolair</i>)
10122R	OMALIZUMAB , omalizumab 150 mg/mL injection, 1 mL syringe (<i>Xolair</i>)

Highly Specialised Drugs Program (Public Hospital)

Additions

Addition – Item

12309W	DUPILUMAB , dupilumab 200 mg/1.14 mL injection, 2 x 1.14 mL syringes (<i>Dupixent</i>)
12318H	DUPILUMAB , dupilumab 200 mg/1.14 mL injection, 2 x 1.14 mL syringes (<i>Dupixent</i>)
12293B	DUPILUMAB , dupilumab 300 mg/2 mL injection, 2 x 2 mL syringes (<i>Dupixent</i>)
12302L	DUPILUMAB , dupilumab 300 mg/2 mL injection, 2 x 2 mL syringes (<i>Dupixent</i>)

Addition – Brand

11887P	<i>Cinacalcet Mylan, AF</i> – CINACALCET , cinacalcet 30 mg tablet, 28
5621W	<i>Cinacalcet Mylan, AF</i> – CINACALCET , cinacalcet 30 mg tablet, 28
11886N	<i>Cinacalcet Mylan, AF</i> – CINACALCET , cinacalcet 60 mg tablet, 28
5622X	<i>Cinacalcet Mylan, AF</i> – CINACALCET , cinacalcet 60 mg tablet, 28
11885M	<i>Cinacalcet Mylan, AF</i> – CINACALCET , cinacalcet 90 mg tablet, 28
5623Y	<i>Cinacalcet Mylan, AF</i> – CINACALCET , cinacalcet 90 mg tablet, 28

Addition – Equivalence Indicator

11887P	<i>Pharmacor Cinacalcet, CR</i> – CINACALCET , cinacalcet 30 mg tablet, 28
5621W	<i>Pharmacor Cinacalcet, CR</i> – CINACALCET , cinacalcet 30 mg tablet, 28
11886N	<i>Pharmacor Cinacalcet, CR</i> – CINACALCET , cinacalcet 60 mg tablet, 28
5622X	<i>Pharmacor Cinacalcet, CR</i> – CINACALCET , cinacalcet 60 mg tablet, 28

11885M	<i>Pharmacor Cinacalcet, CR – CINACALCET</i> , cinacalcet 90 mg tablet, 28
5623Y	<i>Pharmacor Cinacalcet, CR – CINACALCET</i> , cinacalcet 90 mg tablet, 28

Deletions

Deletion – Brand

10591K	<i>Mabthera, RO – RITUXIMAB</i> , rituximab 100 mg/10 mL injection, 2 x 10 mL vials
10593M	<i>Mabthera, RO – RITUXIMAB</i> , rituximab 500 mg/50 mL injection, 50 mL vial
9544H	<i>Mabthera, RO – RITUXIMAB</i> , rituximab 500 mg/50 mL injection, 50 mL vial

Alterations

Alteration – Note

11529T	BENRALIZUMAB , benralizumab 30 mg/mL injection, 1 mL syringe (<i>Fasenra</i>)
11549W	BENRALIZUMAB , benralizumab 30 mg/mL injection, 1 mL syringe (<i>Fasenra</i>)
11830P	BENRALIZUMAB , benralizumab 30 mg/mL injection, 1 mL syringe (<i>Fasenra</i>)
11994G	BENRALIZUMAB , benralizumab 30 mg/mL injection, 1 mL pen device (<i>Fasenra Pen</i>)
11995H	BENRALIZUMAB , benralizumab 30 mg/mL injection, 1 mL pen device (<i>Fasenra Pen</i>)
12000N	BENRALIZUMAB , benralizumab 30 mg/mL injection, 1 mL pen device (<i>Fasenra Pen</i>)
12007Y	MEPOLIZUMAB , mepolizumab 100 mg/mL injection, 1 mL pen device (<i>Nucala</i>)
12021Q	MEPOLIZUMAB , mepolizumab 100 mg/mL injection, 1 mL pen device (<i>Nucala</i>)
12064Y	MEPOLIZUMAB , mepolizumab 100 mg/mL injection, 1 mL pen device (<i>Nucala</i>)
12073K	MEPOLIZUMAB , mepolizumab 100 mg/mL injection, 1 mL pen device (<i>Nucala</i>)
10980X	MEPOLIZUMAB , mepolizumab 100 mg injection, 1 vial (<i>Nucala</i>)
10996R	MEPOLIZUMAB , mepolizumab 100 mg injection, 1 vial (<i>Nucala</i>)
11839D	MEPOLIZUMAB , mepolizumab 100 mg injection, 1 vial (<i>Nucala</i>)
10118M	OMALIZUMAB , omalizumab 75 mg/0.5 mL injection, 0.5 mL syringe (<i>Xolair</i>)
11835X	OMALIZUMAB , omalizumab 75 mg/0.5 mL injection, 0.5 mL syringe (<i>Xolair</i>)
11846L	OMALIZUMAB , omalizumab 75 mg/0.5 mL injection, 0.5 mL syringe (<i>Xolair</i>)
10109C	OMALIZUMAB , omalizumab 150 mg/mL injection, 1 mL syringe (<i>Xolair</i>)
11824H	OMALIZUMAB , omalizumab 150 mg/mL injection, 1 mL syringe (<i>Xolair</i>)
11828M	OMALIZUMAB , omalizumab 150 mg/mL injection, 1 mL syringe (<i>Xolair</i>)

Alteration – Restriction

5609F	APOMORPHINE , apomorphine hydrochloride hemihydrate 20 mg/2 mL injection, 5 x 2 mL ampoules (<i>Movapo</i>)
5610G	APOMORPHINE , apomorphine hydrochloride hemihydrate 50 mg/5 mL injection, 5 x 5 mL ampoules (<i>Movapo</i>)
10950H	APOMORPHINE , apomorphine hydrochloride hemihydrate 50 mg/10 mL injection, 5 x 10 mL syringes (<i>Movapo PFS</i>)
11549W	BENRALIZUMAB , benralizumab 30 mg/mL injection, 1 mL syringe (<i>Fasenra</i>)
11994G	BENRALIZUMAB , benralizumab 30 mg/mL injection, 1 mL pen device (<i>Fasenra Pen</i>)
12007Y	MEPOLIZUMAB , mepolizumab 100 mg/mL injection, 1 mL pen device (<i>Nucala</i>)
10996R	MEPOLIZUMAB , mepolizumab 100 mg injection, 1 vial (<i>Nucala</i>)
10118M	OMALIZUMAB , omalizumab 75 mg/0.5 mL injection, 0.5 mL syringe (<i>Xolair</i>)
10109C	OMALIZUMAB , omalizumab 150 mg/mL injection, 1 mL syringe (<i>Xolair</i>)

Highly Specialised Drugs Program (Community Access)

Additions

Addition – Brand

10310P	<i>Tenofovir Sandoz, SZ – TENOFOVIR DISOPROXIL</i> , tenofovir disoproxil fumarate 300 mg tablet, 30
11992E	<i>Tenofovir Sandoz, SZ – TENOFOVIR DISOPROXIL</i> , tenofovir disoproxil fumarate 300 mg tablet, 30

Growth Hormone Program

Deletions

Deletion – Item

10437H	SOMATROPIN , somatropin 5 mg/1.5 mL injection, 1.5 mL cartridge (<i>Norditropin SimpleXx</i>)
10469B	SOMATROPIN , somatropin 5 mg/1.5 mL injection, 1.5 mL cartridge (<i>Norditropin SimpleXx</i>)
6295H	SOMATROPIN , somatropin 5 mg/1.5 mL injection, 1.5 mL cartridge (<i>Norditropin SimpleXx</i>)
10439K	SOMATROPIN , somatropin 10 mg/1.5 mL injection, 1.5 mL cartridge (<i>Norditropin SimpleXx</i>)
10448X	SOMATROPIN , somatropin 10 mg/1.5 mL injection, 1.5 mL cartridge (<i>Norditropin SimpleXx</i>)
6296J	SOMATROPIN , somatropin 10 mg/1.5 mL injection, 1.5 mL cartridge (<i>Norditropin SimpleXx</i>)
10468Y	SOMATROPIN , somatropin 15 mg/1.5 mL injection, 1.5 mL cartridge (<i>Norditropin SimpleXx</i>)
10470C	SOMATROPIN , somatropin 15 mg/1.5 mL injection, 1.5 mL cartridge (<i>Norditropin SimpleXx</i>)
6297K	SOMATROPIN , somatropin 15 mg/1.5 mL injection, 1.5 mL cartridge (<i>Norditropin SimpleXx</i>)

IVF Program

Additions

Addition – Brand

9583J	<i>GANIRELIX SUN, RA</i> – GANIRELIX , ganirelix 250 microgram/0.5 mL injection, 0.5 mL syringe
9584K	<i>GANIRELIX SUN, RA</i> – GANIRELIX , ganirelix 250 microgram/0.5 mL injection, 5 x 0.5 mL syringes

Addition – Equivalence Indicator

9583J	<i>Orgalutran, OQ</i> – GANIRELIX , ganirelix 250 microgram/0.5 mL injection, 0.5 mL syringe
9584K	<i>Orgalutran, OQ</i> – GANIRELIX , ganirelix 250 microgram/0.5 mL injection, 5 x 0.5 mL syringes

Repatriation Pharmaceutical Benefits

Advance Notices

1 June 2021

Deletion – Brand

4642H	<i>Mepilex Border 295200, MH</i> – DRESSING FOAM WITH SILICONE HEAVY EXUDATE , dressing foam with silicone heavy exudate 7.5 cm x 7.5 cm dressing, 5
4643J	<i>Mepilex Border 295300, MH</i> – DRESSING FOAM WITH SILICONE HEAVY EXUDATE , dressing foam with silicone heavy exudate 10 cm x 10 cm dressing, 5

About the Schedule

The Schedule of Pharmaceutical Benefits lists all of the ready-prepared items subsidised under the Pharmaceutical Benefits Scheme (PBS).

The Schedule is published and is effective on the first day of each month.

For detailed information about the prescribing and supply of pharmaceutical benefits go to www.pbs.gov.au

For information about the operational aspects of the PBS, such as, PBS claiming, authority applications and stationery supplies contact the Department of Human Services at www.humanservices.gov.au

The Repatriation Schedule of Pharmaceutical Benefits provides information about pharmaceutical benefits available under the Repatriation Pharmaceutical Benefits Scheme (RPBS). These may only be prescribed to Department of Veterans' Affairs (DVA) beneficiaries holding a valid repatriation health card. Queries relating to the RPBS can be made to the DVA or go to www.dva.gov.au

Symbols and Abbreviations Used in the Schedule

*	An asterisk in the dispensed price column indicates that the manufacturer's pack does not coincide with the maximum quantity
‡	A double dagger in the maximum quantity column indicates where the maximum quantity has been determined to match the manufacturer's pack. These packs cannot be broken and the maximum quantity should be supplied and claimed
#	A gauge in the dispensed price column indicates that the product is not preconstituted and that the dispensed price therefore included a dispensing fee and where appropriate, an amount for purified water
a or b	Located immediately before brand names of an item indicates that the brands are equivalent for the purposes of substitution. These brands may be interchanged without differences in clinical effect
B	Located immediately before an amount in the premium column indicates a brand premium which applies to that particular brand of the item
T	Located immediately before an amount in the premium column indicates a therapeutic group premium which applies to that particular item
S	Located immediately before an amount in the premium column indicates a special patient contribution which applies to that particular item
DPMQ \$	Dispensed price for maximum quantity
MRVSN \$	Maximum recordable value for safety net
NP	Indicates that the item can be prescribed by an authorised nurse practitioner
MW	Indicates that the item can be prescribed by an authorised midwife
OP	Indicates that the item can be prescribed by an authorised optometrist
DP	Indicates that the item can be prescribed by an authorised dental practitioner

Restricted Benefits

All restricted items have separate headings for authority and non-authority items. In each case these items may be prescribed as pharmaceutical benefits only for use for one of the specified indications. Where more than one indication is specified for an Authority required or Restricted pharmaceutical benefit, each indication is separated from the preceding indication by a semi-colon and commences on the next line. In the case of Authority required (STREAMLINED) items, each indication will also include a four digit streamlined authority code. The drug may be prescribed as a pharmaceutical benefit for a patient who qualifies under any of the specified indications.

Restricted benefits - above an item indicates where an item can only be prescribed for specific therapeutic uses.

Authority required benefits – above an item indicates that a prescriber must seek approval from Department of Human Services or the Department of Veterans' Affairs. The prescriber must declare the specific conditions and circumstances that justify the use of these medicines. This is usually done by phone or in writing

Authority required (STREAMLINED) – authority can be sought electronically.

Guidelines and General Statements

General Statement for Drugs for the Treatment of Hepatitis C

Use the following criteria to determine patient eligibility for subsidisation under the PBS for hepatitis C treating agents.

By writing a PBS prescription, the prescriber is certifying the patient satisfies the qualifying criteria set out below and the use in accordance with the registered indications which differ between agents in this class – refer to the current Product Information for details.

Treatment criteria:

Must be treated by a medical practitioner or an authorised nurse practitioner experienced in the treatment of chronic hepatitis C infection; or in consultation with a gastroenterologist, hepatologist or infectious diseases physician experienced in the treatment of chronic hepatitis C infection.

The following information must be provided at the time of application:

- the patient's cirrhotic status (non-cirrhotic or cirrhotic)
- details of the previous treatment regimen (only for requests for sofosbuvir + velpatasvir + voxilaprevir (Vosevi®) or glecaprevir + pibrentasvir (Maviret®) for treatment in patients who have previously failed a treatment with a regimen containing an NS5A inhibitor).

The following information must be documented in the patient's medical records:

- evidence of chronic hepatitis C infection (repeatedly antibody to hepatitis C virus (anti-HCV) positive and hepatitis C virus ribonucleic acid (HCV RNA) positive); and
- where possible, evidence of the hepatitis C virus genotype

The following matrices identify the regimens which are available for PBS prescription for eligible patients, based on the hepatitis C virus genotype and treatment history.

Hepatitis C - Non-cirrhotic patients

	TREATMENT NAÏVE	TREATMENT EXPERIENCED
All genotypes (Pan-genotypic regimens)	SOFOBUVIR + VELPATASVIR [12 weeks] OR GLECAPREVIR + PIBRENTASVIR [8 weeks]	SOFOBUVIR + VELPATASVIR [12 weeks] OR SOFOBUVIR + VELPATASVIR + VOXILAPREVIR [12 weeks] ³ OR GLECAPREVIR + PIBRENTASVIR [8 or 12 or 16 weeks] ²
Genotype 1	LEDIPASVIR + SOFOBUVIR [8 or 12 weeks] ¹	LEDIPASVIR + SOFOBUVIR [12 weeks]
Genotype 2	Refer to treatment naïve pan-genotypic regimens above.	Refer to treatment experienced pan-genotypic regimens above.
Genotype 3	Refer to treatment naïve pan-genotypic regimens above.	Refer to treatment experienced pan-genotypic regimens above.
Genotype 4	Refer to treatment naïve pan-genotypic regimens above.	Refer to treatment experienced pan-genotypic regimens above.
Genotype 5 & 6	Refer to treatment naïve pan-genotypic regimens above.	Refer to treatment experienced pan-genotypic regimens above.

KEY

RBV - ribavirin

HEPATITIS C – CIRRHOTIC PATIENTS

	TREATMENT NAÏVE	TREATMENT EXPERIENCED
All genotypes (Pan-genotypic regimens)	SOFOSBUVIR + VELPATASVIR [12 weeks] ^{4, 6} OR GLECAPREVIR + PIBRENTASVIR [12 weeks]	SOFOSBUVIR + VELPATASVIR [12 weeks] ^{4, 6} OR SOFOSBUVIR + VELPATASVIR + VOXILAPREVIR [12 weeks] ³ OR GLECAPREVIR + PIBRENTASVIR [12 or 16 weeks] ⁵
Genotype 1	LEDIPASVIR + SOFOSBUVIR [12 weeks]	LEDIPASVIR + SOFOSBUVIR [24 weeks]
Genotype 2	Refer to treatment naïve pan-genotypic regimens above.	Refer to treatment experienced pan-genotypic regimens above.
Genotype 3	Refer to treatment naïve pan-genotypic regimens above.	Refer to treatment experienced pan-genotypic regimens above.
Genotype 4	Refer to treatment naïve pan-genotypic regimens above.	Refer to treatment experienced pan-genotypic regimens above.
Genotype 5 & 6	Refer to treatment naïve pan-genotypic regimens above.	Refer to treatment experienced pan-genotypic regimens above.

KEY

RBV – ribavirin

1. LEDIPASVIR + SOFOSBUVIR [8 or 12 weeks] for treatment-naïve, non-cirrhotic patients:

- consider treatment for 8 weeks where pre-treatment HCV RNA is less than 6 million IU/mL;
- otherwise treatment for 12 weeks where pre-treatment HCV RNA is 6 million IU/mL or greater.

2. GLECAPREVIR + PIBRENTASVIR [8 or 12 or 16 weeks] for non-cirrhotic patients:

- treatment for 8 weeks for treatment-experienced patients with genotypes 1, 2, 4, 5 or 6 who have failed regimens containing peginterferon, ribavirin, and/or sofosbuvir but no prior treatment experience with an HCV NS3/4A PI or NS5A inhibitor;
- treatment for 16 weeks for treatment-experienced patients with genotype 3 who have failed regimens containing peginterferon, ribavirin, and/or sofosbuvir but no prior treatment experience with an HCV NS3/4A PI or NS5A inhibitor;
- treatment for 12 weeks for treatment-experienced patients with genotype 1 who have failed regimens containing an NS3/4A PI;
- treatment for 16 weeks for treatment-experienced patients with genotype 1 who have failed regimens containing an NS5A inhibitor.

3. SOFOSBUVIR + VELPATASVIR + VOXILAPREVIR [12 weeks] only for patients who have failed an NS5A inhibitor.

4. SOFOSBUVIR + VELPATASVIR [12 weeks] for patients with decompensated cirrhosis. Use in combination with ribavirin.

5. GLECAPREVIR + PIBRENTASVIR [12 or 16 weeks] for cirrhotic patients:

- treatment for 12 weeks for treatment-experienced patients with genotypes 1, 2, 4, 5 or 6 who have failed regimens containing peginterferon, ribavirin, and/or sofosbuvir but no prior treatment experience with an HCV NS3/4A PI or NS5A inhibitor;
- treatment for 16 weeks for treatment-experienced patients with genotype 3 who have failed regimens containing peginterferon, ribavirin, and/or sofosbuvir but no prior treatment experience with an HCV NS3/4A PI or NS5A inhibitor;
- treatment for 12 weeks for treatment-experienced patients with genotype 1 who have failed regimens containing an NS3/4A PI;
- treatment for 16 weeks for treatment-experienced patients with genotype 1 who have failed regimens containing an NS5A inhibitor.

6. SOFOSBUVIR + VELPATASVIR [12 weeks] for patients with genotype 3 infection with compensated cirrhosis. Consider addition of ribavirin.

Pharmaceutical Benefits Schedules

Prescriber Bag

▪ ADRENALINE (EPINEPHRINE)

adrenaline (epinephrine) 1 in 1000 (1 mg/mL) injection, 5 x 1 mL ampoules

3451P	Max.Qty Packs	DPMQ \$	Brand Name and Manufacturer
NP	1	21.68	Link Medical Products Pty Ltd [LM]

▪ ATROPINE SULFATE

atropine sulfate monohydrate 600 microgram/mL injection, 10 x 1 mL ampoules

3453R	Max.Qty Packs	DPMQ \$	Brand Name and Manufacturer
NP	1	23.69	Pfizer Australia Pty Ltd [PF]

▪ BENZATHINE BENZYL PENICILLIN

benzathine benzylpenicillin tetrahydrate 1.2 million units (1016.6 mg)/2.3 mL injection, 10 x 2.3 mL syringes

11755Q	Max.Qty Packs	DPMQ \$	Brand Name and Manufacturer
NP	1	304.76	Bicillin L-A [PF]

▪ BENZATROPINE

benzatropine mesilate 2 mg/2 mL injection, 5 x 2 mL vials

11265X	Max.Qty Packs	DPMQ \$	Brand Name and Manufacturer
NP	1	117.17	Benzatropine Injection [FF]

▪ BENZYL PENICILLIN

benzylpenicillin 600 mg injection, 1 vial

3486L	Max.Qty Packs	DPMQ \$	Brand Name and Manufacturer
NP	5	*41.74	BenPen [CS]

OR

▪ PROCAINE BENZYL PENICILLIN (PROCAINE PENICILLIN)

procaine benzylpenicillin (procaine penicillin) 1.5 g/3.4 mL injection, 5 x 3.4 mL syringes

3485K	Max.Qty Packs	DPMQ \$	Brand Name and Manufacturer
NP	1	75.37	Cilicaine [AF]

▪ BENZYL PENICILLIN

benzylpenicillin 3 g injection, 1 vial

3487M	Max.Qty Packs	DPMQ \$	Brand Name and Manufacturer
NP	1	21.60	BenPen [CS]

▪ CHLORPROMAZINE

chlorpromazine hydrochloride 50 mg/2 mL injection, 10 x 2 mL ampoules

3455W	Max.Qty Packs	DPMQ \$	Brand Name and Manufacturer
NP	1	21.94	Largactil [SW]

OR

▪ HALOPERIDOL

haloperidol 5 mg/mL injection, 10 x 1 mL ampoules

3456X	Max.Qty Packs	DPMQ \$	Brand Name and Manufacturer
NP	1	23.22	Serenace [AS]

▪ CLONAZEPAM

clonazepam 2.5 mg/mL (0.1 mg/drop) oral liquid, 10 mL

3478C	Max.Qty Packs	DPMQ \$	Brand Name and Manufacturer
NP	±1	15.77	Rivotril [RO]

▪ DIPHTHERIA + TETANUS VACCINE

diphtheria 2 units + tetanus 20 units vaccine injection, 5 x 0.5 mL syringes

3463G	Max.Qty Packs	DPMQ \$	Brand Name and Manufacturer
NP	2	*129.74	ADT Booster [CS]

■ FUROSEMIDE (FRUSEMIDE)

furosemide (frusemide) 20 mg/2 mL injection, 5 x 2 mL ampoules

3466K	Max.Qty Packs	DPMQ \$	Brand Name and Manufacturer
NP	1	13.66	Lasix [SW]

■ FUROSEMIDE (FRUSEMIDE)

furosemide (frusemide) 20 mg tablet, 50

12222G	Max.Qty Packs	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	13.01	^a Frusemix-M [TY]	^a Urex-M [RW]
		14.13	^a Lasix-M [SW]	

■ GLUCAGON HYDROCHLORIDE

glucagon hydrochloride 1 mg injection [1 vial] (&) inert substance diluent [1 mL syringe], 1 pack

3467L	Max.Qty Packs	DPMQ \$	Brand Name and Manufacturer
NP	1	49.66	GlucaGen Hypokit [NO]

■ GLYCERYL TRINITRATE

glyceryl trinitrate 400 microgram/actuation spray, 200 actuations

3475X	Max.Qty Packs	DPMQ \$	Brand Name and Manufacturer
NP	±1	23.94	Nitrolingual Pumpspray [SW]

■ HYDROCORTISONE SODIUM SUCCINATE

hydrocortisone (as sodium succinate) 250 mg injection [1 chamber] (&) inert substance diluent [2 mL chamber], 1 dual chamber vial

3471Q	Max.Qty Packs	DPMQ \$	Brand Name and Manufacturer
NP	1	20.78	Solu-Cortef [PF]

OR

hydrocortisone (as sodium succinate) 100 mg injection [1 chamber] (&) inert substance diluent [2 mL chamber], 1 dual chamber vial

3470P	Max.Qty Packs	DPMQ \$	Brand Name and Manufacturer
NP	2	*21.82	Solu-Cortef [PF]

■ HYOSCINE BUTYLBROMIDE

hyoscine butylbromide 20 mg/mL injection, 5 x 1 mL ampoules

3473T	Max.Qty Packs	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	22.80	^a Buscopan [VZ]	^a HYOSCINE BUTYLBROMIDE MEDSURGE [DZ]
			^a HYOSCINE BUTYLBROMIDE SXP [XC]	

■ LIDOCAINE (LIGNOCAINE)

lidocaine (lignocaine) hydrochloride monohydrate 1% (50 mg/5 mL) injection, 5 x 5 mL ampoules

10209H	Max.Qty Packs	DPMQ \$	Brand Name and Manufacturer
NP	1	39.55	Pfizer Australia Pty Ltd [PF]

■ METHOXYFLURANE

methoxyflurane 99.9% (999 mg/g) inhalation solution, 3 mL bottle

3489P	Max.Qty Packs	DPMQ \$	Brand Name and Manufacturer
	1	40.98	Penthrox [DV]

▪ METOCLOPRAMIDE

metoclopramide hydrochloride 10 mg/2 mL injection, 10 x 2 mL ampoules

3476Y	Max.Qty Packs	DPMQ \$	Brand Name and Manufacturer
NP	1	17.73	Maxolon [IL]

OR

▪ PROCHLORPERAZINE

prochlorperazine mesilate 12.5 mg/mL injection, 10 x 1 mL ampoules

3477B	Max.Qty Packs	DPMQ \$	Brand Name and Manufacturer
NP	1	21.70	Stemetil [SW]

▪ MIDAZOLAM

midazolam 5 mg/mL injection, 10 x 1 mL ampoules

10178Q	Max.Qty Packs	DPMQ \$	Brand Name and Manufacturer
NP	1	39.79	Pfizer Australia Pty Ltd [PF]

▪ MORPHINE

morphine sulfate pentahydrate 15 mg/mL injection, 5 x 1 mL ampoules

3479D	Max.Qty Packs	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	24.54	^a DBL Morphine Sulfate Pentahydrate [PF]	^a MORPHINE SULFATE 15 mg/1 mL MEDSURGE [DZ]

OR

morphine hydrochloride trihydrate 10 mg/mL injection, 5 x 1 mL ampoules

10862Q	Max.Qty Packs	DPMQ \$	Brand Name and Manufacturer
NP	1	23.14	Morphine Juno [JU]

OR

morphine sulfate pentahydrate 30 mg/mL injection, 5 x 1 mL ampoules

3480E	Max.Qty Packs	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	26.64	^a DBL Morphine Sulfate Pentahydrate [PF]	^a MORPHINE SULFATE 30 mg/1 mL MEDSURGE [DZ]

OR

morphine hydrochloride trihydrate 20 mg/mL injection, 5 x 1 mL ampoules

10868B	Max.Qty Packs	DPMQ \$	Brand Name and Manufacturer
NP	1	26.50	Morphine Juno [JU]

▪ NALOXONE

naloxone hydrochloride 400 microgram/mL injection, 5 x 1 mL ampoules

10786Q	Max.Qty Packs	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	2	*66.36	^a Junalox [JO] ^a Naloxone Juno [JU]	^a Naloxone Hydrochloride (DBL) [PF] ^a NALOXONE SXP [XC]

OR

naloxone hydrochloride 400 microgram/mL injection, 10 x 1 mL ampoules

11233F	Max.Qty Packs	DPMQ \$	Brand Name and Manufacturer
NP	1	66.36	NARCAN [FF]

▪ PHYTOMENADIONE

phytomenadione 10 mg/mL injection, 5 x 1 mL ampoules

10213M	Max.Qty Packs	DPMQ \$	Brand Name and Manufacturer
	1	24.84	Konaktion MM [PB]

▪ PROMETHAZINE

promethazine hydrochloride 50 mg/2 mL injection, 5 x 2 mL ampoules

3488N	Max.Qty Packs	DPMQ \$	Brand Name and Manufacturer
NP	2	*35.60	DBL Promethazine Hydrochloride [PF]

SALBUTAMOL

salbutamol 5 mg/2.5 mL inhalation solution, 30 x 2.5 mL ampoules

3497C	Max.Qty Packs	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	±1	16.23	^a APO-Salbutamol [TX]	^a Salbutamol Actavis [EA]
			^a Salbutamol AN [ED]	^a Salbutamol Cipla [LR]
		16.48	^a Asmol 5 uni-dose [AF]	

OR

salbutamol 5 mg/2.5 mL inhalation solution, 20 x 2.5 mL ampoules

11088N	Max.Qty Packs	DPMQ \$	Brand Name and Manufacturer	
NP	±1	16.49	Ventolin Nebules [GK]	

SALBUTAMOL

salbutamol 2.5 mg/2.5 mL inhalation solution, 30 x 2.5 mL ampoules

3496B	Max.Qty Packs	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	±1	16.02	^a APO-Salbutamol [TX]	^a Salbutamol Actavis [EA]
			^a Salbutamol AN [ED]	^a Salbutamol Cipla [LR]
		16.27	^a Asmol 2.5 uni-dose [AF]	

OR

salbutamol 100 microgram/actuation inhalation, 200 actuations

12108G	Max.Qty Packs	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	±1	16.85	^a Zempreon CFC-Free with dose counter [GC]	
		19.35	^a Asmol CFC-Free with dose counter [AF]	^a Ventolin CFC-Free with dose counter [GK]

OR

salbutamol 100 microgram/actuation inhalation, 200 actuations

3495Y	Max.Qty Packs	DPMQ \$	Brand Name and Manufacturer	
NP	±1	16.21	^a Asmol CFC-free [AL]	
		18.04	^a Ventolin CFC-free [GK]	

OR

salbutamol 2.5 mg/2.5 mL inhalation solution, 20 x 2.5 mL ampoules

11125M	Max.Qty Packs	DPMQ \$	Brand Name and Manufacturer	
NP	±1	16.36	Ventolin Nebules [GK]	

TRAMADOL

tramadol hydrochloride 100 mg/2 mL injection, 5 x 2 mL ampoules

3484J	Max.Qty Packs	DPMQ \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	15.08	^a Tramadol ACT [JO]	^a Tramadol AN [JU]
			^a Tramadol Sandoz [SZ]	^a Tramal 100 [CS]

General Pharmaceutical Benefits

ALIMENTARY TRACT AND METABOLISM.....	29
STOMATOLOGICAL PREPARATIONS	29
STOMATOLOGICAL PREPARATIONS.....	29
DRUGS FOR ACID RELATED DISORDERS.....	29
DRUGS FOR PEPTIC ULCER AND GASTRO-OESOPHAGEAL REFLUX DISEASE (GORD)	29
DRUGS FOR FUNCTIONAL GASTROINTESTINAL DISORDERS	43
BELLADONNA AND DERIVATIVES, PLAIN	43
PROPULSIVES.....	44
ANTIEMETICS AND ANTINAUSEANTS.....	44
ANTIEMETICS AND ANTINAUSEANTS	44
BILE AND LIVER THERAPY	50
BILE THERAPY	50
DRUGS FOR CONSTIPATION	51
DRUGS FOR CONSTIPATION.....	51
ANTIDIARRHEALS, INTESTINAL ANTIINFLAMMATORY/ANTIINFECTIVE AGENTS	54
INTESTINAL ANTIINFECTIVES	54
ELECTROLYTES WITH CARBOHYDRATES	55
ANTIPROPULSIVES.....	55
INTESTINAL ANTIINFLAMMATORY AGENTS	56
DIGESTIVES, INCL. ENZYMES.....	59
DIGESTIVES, INCL. ENZYMES	59
DRUGS USED IN DIABETES	61
INSULINS AND ANALOGUES.....	61
BLOOD GLUCOSE LOWERING DRUGS, EXCL. INSULINS	63
VITAMINS	102
VITAMIN A AND D, INCL. COMBINATIONS OF THE TWO	102
VITAMIN B1, PLAIN AND IN COMBINATION WITH VITAMIN B6 AND B12.....	102
MINERAL SUPPLEMENTS	103
CALCIUM.....	103
POTASSIUM	103
OTHER MINERAL SUPPLEMENTS	103
OTHER ALIMENTARY TRACT AND METABOLISM PRODUCTS	103
OTHER ALIMENTARY TRACT AND METABOLISM PRODUCTS	103
BLOOD AND BLOOD FORMING ORGANS	106
ANTITHROMBOTIC AGENTS.....	106

ANTITHROMBOTIC AGENTS	106
ANTIHEMORRHAGICS	121
ANTIFIBRINOLYTICS	121
ANTIANEMIC PREPARATIONS	121
IRON PREPARATIONS	121
VITAMIN B12 AND FOLIC ACID	122
BLOOD SUBSTITUTES AND PERFUSION SOLUTIONS	123
BLOOD AND RELATED PRODUCTS	123
OTHER HEMATOLOGICAL AGENTS	123
OTHER HEMATOLOGICAL AGENTS	123
CARDIOVASCULAR SYSTEM	123
CARDIAC THERAPY	123
CARDIAC GLYCOSIDES	123
ANTIARRHYTHMICS, CLASS I AND III	124
CARDIAC STIMULANTS EXCL. CARDIAC GLYCOSIDES	125
VASODILATORS USED IN CARDIAC DISEASES	126
OTHER CARDIAC PREPARATIONS	127
ANTIHYPERTENSIVES	127
ANTIADRENERGIC AGENTS, CENTRALLY ACTING	127
ANTIADRENERGIC AGENTS, PERIPHERALLY ACTING	129
ARTERIOLAR SMOOTH MUSCLE, AGENTS ACTING ON.....	129
DIURETICS	130
LOW-CEILING DIURETICS, THIAZIDES	130
LOW-CEILING DIURETICS, EXCL. THIAZIDES	130
HIGH-CEILING DIURETICS	130
POTASSIUM-SPARING AGENTS	131
DIURETICS AND POTASSIUM-SPARING AGENTS IN COMBINATION	132
OTHER DIURETICS	132
PERIPHERAL VASODILATORS	133
PERIPHERAL VASODILATORS	133
BETA BLOCKING AGENTS	133
BETA BLOCKING AGENTS	133
CALCIUM CHANNEL BLOCKERS	136
SELECTIVE CALCIUM CHANNEL BLOCKERS WITH MAINLY VASCULAR EFFECTS	136
SELECTIVE CALCIUM CHANNEL BLOCKERS WITH DIRECT CARDIAC EFFECTS	137
AGENTS ACTING ON THE RENIN-ANGIOTENSIN SYSTEM	138
ACE INHIBITORS, PLAIN	138
ACE INHIBITORS, COMBINATIONS	142
ANGIOTENSIN II RECEPTOR BLOCKERS (ARBs), PLAIN	144
ANGIOTENSIN II RECEPTOR BLOCKERS (ARBs), COMBINATIONS	146
LIPID MODIFYING AGENTS	153
LIPID MODIFYING AGENTS, PLAIN	153
LIPID MODIFYING AGENTS, COMBINATIONS	167
DERMATOLOGICALS	173
ANTIFUNGALS FOR DERMATOLOGICAL USE	173
ANTIFUNGALS FOR TOPICAL USE	173
ANTIFUNGALS FOR SYSTEMIC USE	174

ANTIPSORIATICS.....	174
ANTIPSORIATICS FOR TOPICAL USE.....	174
ANTIPSORIATICS FOR SYSTEMIC USE.....	175
ANTIBIOTICS AND CHEMOTHERAPEUTICS FOR DERMATOLOGICAL USE.....	175
CHEMOTHERAPEUTICS FOR TOPICAL USE.....	175
CORTICOSTEROIDS, DERMATOLOGICAL PREPARATIONS.....	176
CORTICOSTEROIDS, PLAIN.....	176
ANTI-ACNE PREPARATIONS.....	185
ANTI-ACNE PREPARATIONS FOR TOPICAL USE.....	185
ANTI-ACNE PREPARATIONS FOR SYSTEMIC USE.....	185
OTHER DERMATOLOGICAL PREPARATIONS.....	186
OTHER DERMATOLOGICAL PREPARATIONS.....	186
GENITO URINARY SYSTEM AND SEX HORMONES.....	191
OTHER GYNECOLOGICALS.....	191
CONTRACEPTIVES FOR TOPICAL USE.....	191
OTHER GYNECOLOGICALS.....	191
SEX HORMONES AND MODULATORS OF THE GENITAL SYSTEM.....	192
HORMONAL CONTRACEPTIVES FOR SYSTEMIC USE.....	192
ANDROGENS.....	194
ESTROGENS.....	196
PROGESTOGENS.....	197
PROGESTOGENS AND ESTROGENS IN COMBINATION.....	198
GONADOTROPINS AND OTHER OVULATION STIMULANTS.....	199
ANTIANDROGENS.....	201
OTHER SEX HORMONES AND MODULATORS OF THE GENITAL SYSTEM.....	201
UROLOGICALS.....	202
UROLOGICALS.....	202
DRUGS USED IN BENIGN PROSTATIC HYPERTROPHY.....	203
SYSTEMIC HORMONAL PREPARATIONS, EXCL. SEX HORMONES AND INSULINS.....	203
PITUITARY AND HYPOTHALAMIC HORMONES AND ANALOGUES.....	203
ANTERIOR PITUITARY LOBE HORMONES AND ANALOGUES.....	203
POSTERIOR PITUITARY LOBE HORMONES.....	204
HYPOTHALAMIC HORMONES.....	205
CORTICOSTEROIDS FOR SYSTEMIC USE.....	206
CORTICOSTEROIDS FOR SYSTEMIC USE, PLAIN.....	206
THYROID THERAPY.....	209
THYROID PREPARATIONS.....	209
ANTITHYROID PREPARATIONS.....	210
PANCREATIC HORMONES.....	211
GLYCOGENOLYTIC HORMONES.....	211
CALCIUM HOMEOSTASIS.....	211
PARATHYROID HORMONES AND ANALOGUES.....	211
ANTI-PARATHYROID AGENTS.....	212
ANTIINFECTIVES FOR SYSTEMIC USE.....	212
ANTIBACTERIALS FOR SYSTEMIC USE.....	212
TETRACYCLINES.....	212
BETA-LACTAM ANTIBACTERIALS, PENICILLINS.....	215
OTHER BETA-LACTAM ANTIBACTERIALS.....	223

SULFONAMIDES AND TRIMETHOPRIM	229
MACROLIDES, LINCOSAMIDES AND STREPTOGRAMINS	230
AMINOGLYCOSIDE ANTIBACTERIALS	233
QUINOLONE ANTIBACTERIALS	234
OTHER ANTIBACTERIALS	236
ANTIMYCOTICS FOR SYSTEMIC USE	238
ANTIMYCOTICS FOR SYSTEMIC USE	238
ANTIMYCOBACTERIALS	244
DRUGS FOR TREATMENT OF TUBERCULOSIS	244
DRUGS FOR TREATMENT OF LEPROSY	245
ANTIVIRALS FOR SYSTEMIC USE	246
DIRECT ACTING ANTIVIRALS	246
VACCINES	253
BACTERIAL VACCINES	253
ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS	253
ANTINEOPLASTIC AGENTS	253
ALKYLATING AGENTS	253
ANTIMETABOLITES	255
PLANT ALKALOIDS AND OTHER NATURAL PRODUCTS	257
PROTEIN KINASE INHIBITORS	257
OTHER ANTINEOPLASTIC AGENTS	329
ENDOCRINE THERAPY	339
HORMONES AND RELATED AGENTS	339
HORMONE ANTAGONISTS AND RELATED AGENTS	342
IMMUNOSTIMULANTS	347
IMMUNOSTIMULANTS	347
IMMUNOSUPPRESSANTS	350
IMMUNOSUPPRESSANTS	350
MUSCULO-SKELETAL SYSTEM	760
ANTIINFLAMMATORY AND ANTIRHEUMATIC PRODUCTS	760
ANTIINFLAMMATORY AND ANTIRHEUMATIC PRODUCTS, NON- STEROIDS	760
SPECIFIC ANTIRHEUMATIC AGENTS	766
MUSCLE RELAXANTS	767
MUSCLE RELAXANTS, CENTRALLY ACTING AGENTS	767
MUSCLE RELAXANTS, DIRECTLY ACTING AGENTS	767
ANTIGOUT PREPARATIONS	768
ANTIGOUT PREPARATIONS	768
DRUGS FOR TREATMENT OF BONE DISEASES	768
DRUGS AFFECTING BONE STRUCTURE AND MINERALIZATION	768
NERVOUS SYSTEM	778
ANALGESICS	778
OPIOIDS	778
OTHER ANALGESICS AND ANTIPYRETICS	823
ANTIMIGRAINE PREPARATIONS	825
ANTIEPILEPTICS	827
ANTIEPILEPTICS	827
ANTI-PARKINSON DRUGS	842

ANTICHOLINERGIC AGENTS	842
DOPAMINERGIC AGENTS	842
PSYCHOLEPTICS.....	849
ANTIPSYCHOTICS.....	849
ANXIOLYTICS	865
HYPNOTICS AND SEDATIVES	867
PSYCHOANALEPTICS	868
ANTIDEPRESSANTS	868
PSYCHOSTIMULANTS, AGENTS USED FOR ADHD AND NOOTROPICS.....	878
ANTI-DEMENTIA DRUGS	884
OTHER NERVOUS SYSTEM DRUGS.....	894
PARASYMPATHOMIMETICS.....	894
DRUGS USED IN ADDICTIVE DISORDERS	894
OTHER NERVOUS SYSTEM DRUGS	898
ANTIPARASITIC PRODUCTS, INSECTICIDES AND REPELLENTS.....	898
ANTIPROTOZOALS	898
AGENTS AGAINST AMOEBIASIS AND OTHER PROTOZOAL DISEASES	898
ANTIMALARIALS.....	899
ANTHELMINTICS.....	900
ANTITREMATODALS.....	900
ANTINEMATODAL AGENTS.....	900
ECTOPARASITICIDES, INCL. SCABICIDES, INSECTICIDES AND REPELLENTS	901
ECTOPARASITICIDES, INCL. SCABICIDES.....	901
RESPIRATORY SYSTEM.....	901
NASAL PREPARATIONS.....	901
DECONGESTANTS AND OTHER NASAL PREPARATIONS FOR TOPICAL USE	901
DRUGS FOR OBSTRUCTIVE AIRWAY DISEASES	902
ADRENERGICS, INHALANTS.....	902
OTHER DRUGS FOR OBSTRUCTIVE AIRWAY DISEASES, INHALANTS	914
ADRENERGICS FOR SYSTEMIC USE	918
OTHER SYSTEMIC DRUGS FOR OBSTRUCTIVE AIRWAY DISEASES	919
ANTIHISTAMINES FOR SYSTEMIC USE	921
ANTIHISTAMINES FOR SYSTEMIC USE.....	921
SENSORY ORGANS	921
OPHTHALMOLOGICALS	921
ANTIINFECTIVES.....	921
ANTIINFLAMMATORY AGENTS.....	923
ANTIGLAUCOMA PREPARATIONS AND MIOTICS.....	927
MYDRIATICS AND CYCLOPLEGICS	933
OCULAR VASCULAR DISORDER AGENTS	933
OTHER OPHTHALMOLOGICALS.....	943
OTOLOGICALS	950
ANTIINFECTIVES.....	950
CORTICOSTEROIDS AND ANTIINFECTIVES IN COMBINATION	951

OPHTHALMOLOGICAL AND OTOLOGICAL PREPARATIONS	951
ANTIINFECTIVES	951
VARIOUS	951
ALLERGENS	951
ALLERGENS	951
ALL OTHER THERAPEUTIC PRODUCTS	952
ALL OTHER THERAPEUTIC PRODUCTS	952
DIAGNOSTIC AGENTS	955
URINE TESTS	955
GENERAL NUTRIENTS	955
OTHER NUTRIENTS	955

ALIMENTARY TRACT AND METABOLISM

STOMATOLOGICAL PREPARATIONS

STOMATOLOGICAL PREPARATIONS

Antifungives and antiseptics for local oral treatment

AMPHOTERICIN B

amphotericin B 10 mg lozenge, 20

2931G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	16.39	17.68	Fungilin [AS]

amphotericin B 10 mg lozenge, 20

3306B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	1	..	..	16.39	17.68	Fungilin [AS]

Other agents for local oral treatment

BENZYLAMINE

Restricted benefit

Mucositis

Clinical criteria:

- The condition must be radiation induced.

benzylamine hydrochloride 0.15% mouthwash, 500 mL

1121B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	1	..	23.21	24.50	Diffiam [IL]

BENZYLAMINE

Restricted benefit

Mucositis

Clinical criteria:

- The condition must be radiation induced.

benzylamine hydrochloride 0.15% mouthwash, 500 mL

5032W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	±1	..	..	23.21	24.50	Diffiam [IL]

DRUGS FOR ACID RELATED DISORDERS

DRUGS FOR PEPTIC ULCER AND GASTRO-OESOPHAGEAL REFLUX DISEASE (GORD)

H2-receptor antagonists

FAMOTIDINE

Note *Helicobacter pylori* eradication therapy should be considered prior to commencing initial treatment of peptic ulcer with this drug.

famotidine 20 mg tablet, 60

2487X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	16.01	17.30	^a Ausfam 20 [RW] ^a GenRx Famotidine [GX]	^a Famotidine AN [EA]

famotidine 40 mg tablet, 30

2488Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	16.01	17.30	^a Ausfam 40 [RW] ^a GenRx Famotidine [GX]	^a Famotidine AN [EA]

NIZATIDINE

Note *Helicobacter pylori* eradication therapy should be considered prior to commencing initial treatment of peptic ulcer with this drug.

nizatidine 150 mg capsule, 60

1505F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	.. ^B 6.61	22.46 29.07	23.75 23.75	^a Nizac [RF] ^a Tazac [RW]	^a Tacidine [AF]

nizatidine 300 mg capsule, 30

1504E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	.. ^B 6.61	22.46 29.07	23.75 23.75	^a Nizac [RF] ^a Tazac [RW]	^a Tacidine [AF]

■ RANITIDINE

Note *Helicobacter pylori* eradication therapy should be considered prior to commencing initial treatment of peptic ulcer with this drug.

ranitidine 150 mg/10 mL oral liquid, 300 mL

8162N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*29.26	30.55	Zantac Syrup [AS]

ranitidine 300 mg tablet, 30

1977C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	15.65	16.94	^a APO-Ranitidine [TX] ^a Rani 2 [AF]	^a Ausran [RW]
			^B 1.56	17.21	16.94	^a Zantac [AS]	

ranitidine 150 mg tablet, 60

1978D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP MW	1	5	..	15.65	16.94	^a APO-Ranitidine [TX] ^a Rani 2 [AF]	^a Ausran [RW] ^a Ranitidine AN [EA]
			^B 1.56	17.21	16.94	^a Zantac [AS]	

Proton pump inhibitors

■ ESOMEPRAZOLE

Note Pharmaceutical benefits that have the form esomeprazole tablet 40 mg and pharmaceutical benefits that have the form esomeprazole capsule 40 mg are equivalent for the purposes of substitution.

Note Check patient adherence to lower dose proton pump inhibitor before "stepping-up" therapy.

Note Standard dose proton pump inhibitors are appropriate step-down therapy from high dose proton pump inhibitors.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Gastro-oesophageal reflux disease

Clinical criteria:

- Patient must have symptoms which are inadequately controlled using a standard dose proton pump inhibitor.

esomeprazole 40 mg enteric capsule, 30

10330Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	1	..	22.74	24.03	^a Esomeprazole ACTAVIS [EA]	^a Noxicid Caps [AL]

esomeprazole 40 mg enteric tablet, 30

8601Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	1	..	22.74	24.03	^a APO-Esomeprazole [TY] ^a Esomeprazole GH [GQ] ^a Esomeprazole Mylan [AL] ^a Esomeprazole Sandoz [SZ] ^a Esopreze [BG] ^a Nexole [RF]	^a Esomeprazole Apotex [TX] ^a Esomeprazole GxP [AF] ^a Esomeprazole RBX [RA] ^a Esomeprazole SZ [HX] ^a Nexazole [RW] ^a NOUMED ESOMEPRAZOLE [VO]
			^B 3.58	26.32	24.03	^a Nexium [AP]	

■ ESOMEPRAZOLE

Note Pharmaceutical benefits that have the form esomeprazole tablet 40 mg and pharmaceutical benefits that have the form esomeprazole capsule 40 mg are equivalent for the purposes of substitution.

Note Check patient adherence to lower dose proton pump inhibitor before "stepping-up" therapy.

Note Standard dose proton pump inhibitors are appropriate step-down therapy from high dose proton pump inhibitors.

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required

Pathological hypersecretory conditions including Zollinger-Ellison syndrome and idiopathic hypersecretion

Clinical criteria:

- Patient must have symptoms which are inadequately controlled using a standard dose proton pump inhibitor.

Authority required

Scleroderma oesophagus

Clinical criteria:

- Patient must have symptoms which are inadequately controlled using a standard dose proton pump inhibitor.

esomeprazole 40 mg enteric capsule, 30

10331R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	22.74	24.03	^a Esomeprazole ACTAVIS [EA]	^a Noxicid Caps [AL]

esomeprazole 40 mg enteric tablet, 30

3401B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	22.74	24.03	^a APO-Esomeprazole [TY] ^a Esomeprazole GH [GQ] ^a Esomeprazole Mylan [AL] ^a Esomeprazole Sandoz [SZ] ^a Esopreze [BG] ^a Nexole [RF]	^a Esomeprazole Apotex [TX] ^a Esomeprazole GxP [AF] ^a Esomeprazole RBX [RA] ^a Esomeprazole SZ [HX] ^a Nexazole [RW] ^a NOUMED ESOMEPRAZOLE [VO]
			^B 3.58	26.32	24.03	^a Nexium [AP]	

■ ESOMEPRAZOLE

Note Pharmaceutical benefits that have the form esomeprazole tablet 20 mg and pharmaceutical benefits that have the form esomeprazole capsule 20 mg are equivalent for the purposes of substitution.

Note Check patient adherence to lower dose proton pump inhibitor before "stepping-up" therapy.

Note Low dose proton pump inhibitors are appropriate step-down therapy from standard dose proton pump inhibitors.

Note A low dose proton pump inhibitor includes: lansoprazole 15mg, omeprazole 10mg, pantoprazole 20mg and rabeprazole 10mg.

Note No increase in the maximum quantity or number of units may be authorised.

Authority required (STREAMLINED)**8780**

Scleroderma oesophagus

Authority required (STREAMLINED)**8827**

Pathological hypersecretory conditions including Zollinger-Ellison syndrome and idiopathic hypersecretion

esomeprazole 20 mg enteric tablet, 30

8600P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	18.33	19.62	^a APO-Esomeprazole [TY] ^a Esomeprazole GH [GQ] ^a Esomeprazole Mylan [AL] ^a Esomeprazole Sandoz [SZ] ^a Esopreze [BG] ^a Nexole [RF]	^a Esomeprazole Apotex [TX] ^a Esomeprazole GxP [AF] ^a Esomeprazole RBX [RA] ^a Esomeprazole SZ [HX] ^a Nexazole [RW] ^a NOUMED ESOMEPRAZOLE [VO]
			^B 3.58	21.91	19.62	^a Nexium [AP]	

esomeprazole 20 mg enteric capsule, 30

10343J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	18.33	19.62	^a Esomeprazole ACTAVIS [EA]	^a Noxicid Caps [AL]

■ ESOMEPRAZOLE

Note Pharmaceutical benefits that have the form esomeprazole tablet 20 mg and pharmaceutical benefits that have the form esomeprazole capsule 20 mg are equivalent for the purposes of substitution.

Note Check patient adherence to lower dose proton pump inhibitor before "stepping-up" therapy.

Note Low dose proton pump inhibitors are appropriate step-down therapy from standard dose proton pump inhibitors.

Note A low dose proton pump inhibitor includes: lansoprazole 15mg, omeprazole 10mg, pantoprazole 20mg and rabeprazole 10mg.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)**8776**

Gastro-oesophageal reflux disease

Clinical criteria:

- The treatment must be for long-term maintenance of gastro-oesophageal reflux disease in a patient with symptoms inadequately controlled using a low dose proton pump inhibitor.

esomeprazole 20 mg enteric tablet, 30

11692J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	18.33	19.62	^a APO-Esomeprazole [TY] ^a Esomeprazole GH [GQ] ^a Esomeprazole Mylan [AL] ^a Esomeprazole Sandoz [SZ] ^a Esopreze [BG] ^a Nexole [RF]	^a Esomeprazole Apotex [TX] ^a Esomeprazole GxP [AF] ^a Esomeprazole RBX [RA] ^a Esomeprazole SZ [HX] ^a Nexazole [RW] ^a NOUMED ESOMEPRAZOLE [VO]
			^B 3.58	21.91	19.62	^a Nexium [AP]	

esomeprazole 20 mg enteric capsule, 30

11687D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	18.33	19.62	^a Esomeprazole ACTAVIS [EA]	^a Noxicid Caps [AL]

■ ESOMEPRAZOLE

Note Pharmaceutical benefits that have the form esomeprazole tablet 20 mg and pharmaceutical benefits that have the form esomeprazole capsule 20 mg are equivalent for the purposes of substitution.

Note Check patient adherence to lower dose proton pump inhibitor before "stepping-up" therapy.

Note Low dose proton pump inhibitors are appropriate step-down therapy from standard dose proton pump inhibitors.

Note A low dose proton pump inhibitor includes: lansoprazole 15mg, omeprazole 10mg, pantoprazole 20mg and rabeprazole 10mg.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)**8775**

Peptic ulcer

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have tested negative for helicobacter pylori infection; OR
- Patient must have failed treatment with helicobacter pylori eradication therapy.

Authority required (STREAMLINED)**8774**

Gastro-oesophageal reflux disease

Clinical criteria:

- The treatment must be for initial treatment of symptomatic gastro-oesophageal reflux disease; OR
- The treatment must be for the short-term maintenance treatment of gastro-oesophageal reflux disease.

esomeprazole 20 mg enteric tablet, 30

8886Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	1	..	18.33	19.62	^a APO-Esomeprazole [TY] ^a Esomeprazole GH [GQ] ^a Esomeprazole Mylan [AL] ^a Esomeprazole Sandoz [SZ] ^a Esopreze [BG] ^a Nexole [RF]	^a Esomeprazole Apotex [TX] ^a Esomeprazole GxP [AF] ^a Esomeprazole RBX [RA] ^a Esomeprazole SZ [HX] ^a Nexazole [RW] ^a NOUMED ESOMEPRAZOLE [VO]
			^B 3.58	21.91	19.62	^a Nexium [AP]	

esomeprazole 20 mg enteric capsule, 30

10295W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	1	..	18.33	19.62	^a Esomeprazole ACTAVIS [EA]	^a Noxicid Caps [AL]

■ ESOMEPRAZOLE

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Note The treatment must not be prescribed under this restriction for any of the following PBS-indications: (i) gastro-oesophageal reflux disease (where the word 'complex' is absent), (ii) scleroderma oesophagus, (iii) pathological hypersecretory conditions including Zollinger-Ellison syndrome and idiopathic hypersecretion, (iv) peptic ulcer, (v) eradication of **Helicobacter pylori**.

Note A standard dose proton pump inhibitor is one of: esomeprazole 20 mg, lansoprazole 30 mg, omeprazole 20 mg, pantoprazole 40 mg, rabeprazole 20 mg.

Note A low dose proton pump inhibitor includes: lansoprazole 15mg, omeprazole 10mg, pantoprazole 20mg and rabeprazole 10mg.

Note Pharmaceutical benefits that have the form esomeprazole tablet 20 mg and pharmaceutical benefits that have the form esomeprazole capsule 20 mg are equivalent for the purposes of substitution.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Complex gastro-oesophageal reflux disease (GORD)

Treatment Phase: One of: (1) establishment of symptom control, (2) maintenance treatment, (3) re-establishment of symptom control

Treatment criteria:

- Must be treated by a gastroenterologist; OR
- Must be treated by a surgeon with expertise in the upper gastrointestinal tract; OR
- Must be treated by a medical practitioner who has consulted at least one of the above mentioned specialists in relation to this current PBS benefit being sought, with the specialist's name documented in the patient's medical records for auditing purposes; OR

- Must be treated by a medical practitioner who has not consulted a specialist, but only if treatment continues therapy initiated under this restriction with involvement by a specialist (i.e. continuing treatment initiated for non-complex GORD does not meet this criterion), with the specialist's name documented in the patient's medical records for auditing purposes.

Clinical criteria:

- The treatment must be: (i) the sole PBS-subsidised proton pump inhibitor (PPI) for this condition, (ii) the sole strength of this PPI, (iii) the sole form of PPI, **AND**
- Patient must have symptoms inadequately controlled with each of: (i) a standard dose proton pump inhibitor (PPI) administered once daily, (ii) a low dose PPI administered twice daily; treatment is for: (1) establishment of symptom control; OR
- Patient must be assessed for the risks/benefits of a step-down in dosing from standard dose PPI administered twice daily, with the determination being that the risks outweigh the benefits; treatment is for: (2) maintenance treatment; OR
- Patient must have trialled a step-down in dosing, yet symptoms have re-emerged/worsened; treatment is for: (3) re-establishment of symptom control; OR
- Patient must have trialled a step-down in dosing, with symptoms adequately managed with once daily dosing; treatment is for: (2) maintenance treatment, but with the quantity sought in this authority application being up to 1 pack per dispensing.

Check patient adherence to any preceding PPI treatment regimen. Exclude non-adherence as a cause of inadequate control before accessing treatment under this restriction.

esomeprazole 20 mg enteric tablet, 30

12287Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	5	..	*24.64	25.93	^a APO-Esomeprazole [TY] ^a Esomeprazole GH [GQ] ^a Esomeprazole Mylan [AL] ^a Esomeprazole Sandoz [SZ] ^a Esopreze [BG] ^a Nexole [RF]	^a Esomeprazole Apotex [TX] ^a Esomeprazole GxP [AF] ^a Esomeprazole RBX [RA] ^a Esomeprazole SZ [HX] ^a Nexazole [RW] ^a NOUMED ESOMEPRAZOLE [VO]
			^B 7.16	*31.80	25.93	^a Nexium [AP]	

esomeprazole 20 mg enteric capsule, 30

12275C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	5	..	*24.64	25.93	^a Esomeprazole ACTAVIS [EA]	^a Noxicid Caps [AL]

■ ESOMEPRAZOLE

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Note The treatment must not be prescribed under this restriction for any of the following PBS-indications: (i) gastro-oesophageal reflux disease (where the word 'complex' is absent), (ii) scleroderma oesophagus, (iii) pathological hypersecretory conditions including Zollinger-Ellison syndrome and idiopathic hypersecretion, (iv) peptic ulcer, (v) eradication of **Helicobacter pylori**.

Note A standard dose proton pump inhibitor is one of: esomeprazole 20 mg, lansoprazole 30 mg, omeprazole 20 mg, pantoprazole 40 mg, rabeprazole 20 mg.

Note A high dose proton pump inhibitor is: esomeprazole 40 mg.

Note Pharmaceutical benefits that have the form esomeprazole tablet 40 mg and pharmaceutical benefits that have the form esomeprazole capsule 40 mg are equivalent for the purposes of substitution.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Complex gastro-oesophageal reflux disease (GORD)

Treatment Phase: One of: (1) establishment of symptom control, (2) maintenance treatment, (3) re-establishment of symptom control

Treatment criteria:

- Must be treated by a gastroenterologist; OR
- Must be treated by a surgeon with expertise in the upper gastrointestinal tract.

Clinical criteria:

- The treatment must be: (i) the sole PBS-subsidised proton pump inhibitor (PPI) for this condition, (ii) the sole strength of this PPI, (iii) the sole form of PPI, **AND**
- Patient must have symptoms inadequately controlled with each of: (i) a high dose proton pump inhibitor (PPI) administered once daily, (ii) a standard dose PPI administered twice daily; treatment is for: (1) establishment of symptom control; OR
- Patient must be assessed for the risks/benefits of a step-down in dosing from a high dose PPI administered twice daily, with the determination being that the risks outweigh the benefits; treatment is for: (2) maintenance treatment; OR
- Patient must have trialled a step-down in dosing, yet symptoms have re-emerged/worsened; treatment is for: (3) re-establishment of symptom control; OR
- Patient must have trialled a step-down in dosing, with symptoms adequately managed with once daily dosing; treatment is for: (2) maintenance treatment, but with the quantity sought in this authority application being up to 1 pack per dispensing.

Check patient adherence to any preceding PPI treatment regimen. Exclude non-adherence as a cause of inadequate control before accessing treatment under this restriction.

esomeprazole 40 mg enteric capsule, 30

12290W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	5	..	*33.46	34.75	^a Esomeprazole ACTAVIS [EA]	^a Noxicid Caps [AL]

esomeprazole 40 mg enteric tablet, 30

12283L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	5	..	*33.46	34.75	^a APO-Esomeprazole [TY] ^a Esomeprazole GH [GQ] ^a Esomeprazole Mylan [AL] ^a Esomeprazole Sandoz [SZ] ^a Esopreze [BG] ^a Nexole [RF]	^a Esomeprazole Apotex [TX] ^a Esomeprazole GxP [AF] ^a Esomeprazole RBX [RA] ^a Esomeprazole SZ [HX] ^a Nexazole [RW] ^a NOUMED ESOMEPRAZOLE [VO]
			^B 7.16	*40.62	34.75	^a Nexium [AP]	

■ LANSOPRAZOLE

Note Check patient adherence to lower dose proton pump inhibitor before "stepping-up" therapy.

Restricted benefit

Gastro-oesophageal reflux disease

Restricted benefit

Scleroderma oesophagus

lansoprazole 15 mg enteric capsule, 30

8198L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	15.56	16.85	Zopral [AF]

lansoprazole 15 mg orally disintegrating tablet, 28

9331D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	15.35	16.64	^a APO-Lansoprazole ODT [TX] ^a Zopral ODT [AF]	^a Lansoprazole ODT GH [GQ]
			^B 4.62	19.97	16.64	^a Zoton FasTabs [PF]	

■ LANSOPRAZOLE

Note Pharmaceutical benefits that have the form lansoprazole capsule 30 mg and pharmaceutical benefits that have the form lansoprazole tablet 30 mg (orally disintegrating) are equivalent for the purposes of substitution.

Note Check patient adherence to lower dose proton pump inhibitor before "stepping-up" therapy.

Note Low dose proton pump inhibitors are appropriate step-down therapy from standard dose proton pump inhibitors.

Note A low dose proton pump inhibitor includes: lansoprazole 15mg, omeprazole 10mg, pantoprazole 20mg and rabeprazole 10mg.

Authority required (STREAMLINED)

8780

Scleroderma oesophagus

lansoprazole 30 mg enteric capsule, 28

2241Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	17.65	18.94	^a APO-Lansoprazole [TX] ^a NOUMED LANSOPRAZOLE [VO]	^a Lanzopran [RA] ^a Zopral [AF]

lansoprazole 30 mg orally disintegrating tablet, 28

9478W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	17.65	18.94	^a APO-Lansoprazole ODT [TX] ^a Zopral ODT [AF]	^a Lansoprazole ODT GH [GQ]
			^B 4.81	22.46	18.94	^a Zoton FasTabs [PF]	

■ LANSOPRAZOLE

Note Pharmaceutical benefits that have the form lansoprazole capsule 30 mg and pharmaceutical benefits that have the form lansoprazole tablet 30 mg (orally disintegrating) are equivalent for the purposes of substitution.

Note Check patient adherence to lower dose proton pump inhibitor before "stepping-up" therapy.

Note Low dose proton pump inhibitors are appropriate step-down therapy from standard dose proton pump inhibitors.

Note A low dose proton pump inhibitor includes: lansoprazole 15mg, omeprazole 10mg, pantoprazole 20mg and rabeprazole 10mg.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

8776

Gastro-oesophageal reflux disease

Clinical criteria:

- The treatment must be for long-term maintenance of gastro-oesophageal reflux disease in a patient with symptoms inadequately controlled using a low dose proton pump inhibitor.

lansoprazole 30 mg enteric capsule, 28

11669E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	17.65	18.94	^a APO-Lansoprazole [TX]	^a Lanzopran [RA]
						^a NOUMED LANSOPRAZOLE [VO]	^a Zopral [AF]

lansoprazole 30 mg orally disintegrating tablet, 28

11697P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	17.65	18.94	^a APO-Lansoprazole ODT [TX]	^a Lansoprazole ODT GH [GQ]
						^a Zopral ODT [AF]	
			^B 4.81	22.46	18.94	^a Zoton FasTabs [PF]	

■ LANSOPRAZOLE

Note Pharmaceutical benefits that have the form lansoprazole capsule 30 mg and pharmaceutical benefits that have the form lansoprazole tablet 30 mg (orally disintegrating) are equivalent for the purposes of substitution.

Note Check patient adherence to lower dose proton pump inhibitor before "stepping-up" therapy.

Note Low dose proton pump inhibitors are appropriate step-down therapy from standard dose proton pump inhibitors.

Note A low dose proton pump inhibitor includes: lansoprazole 15mg, omeprazole 10mg, pantoprazole 20mg and rabeprazole 10mg.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)**8775**

Peptic ulcer

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have tested negative for helicobacter pylori infection; OR
- Patient must have failed treatment with helicobacter pylori eradication therapy.

Authority required (STREAMLINED)**8774**

Gastro-oesophageal reflux disease

Clinical criteria:

- The treatment must be for initial treatment of symptomatic gastro-oesophageal reflux disease; OR
- The treatment must be for the short-term maintenance treatment of gastro-oesophageal reflux disease.

lansoprazole 30 mg enteric capsule, 28

2240X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	1	..	17.65	18.94	^a APO-Lansoprazole [TX]	^a Lanzopran [RA]
						^a NOUMED LANSOPRAZOLE [VO]	^a Zopral [AF]

lansoprazole 30 mg orally disintegrating tablet, 28

9477T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	1	..	17.65	18.94	^a APO-Lansoprazole ODT [TX]	^a Lansoprazole ODT GH [GQ]
						^a Zopral ODT [AF]	
			^B 4.81	22.46	18.94	^a Zoton FasTabs [PF]	

■ LANSOPRAZOLE

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Note The treatment must not be prescribed under this restriction for any of the following PBS-indications: (i) gastro-oesophageal reflux disease (where the word 'complex' is absent), (ii) scleroderma oesophagus, (iii) pathological hypersecretory conditions including Zollinger-Ellison syndrome and idiopathic hypersecretion, (iv) peptic ulcer, (v) eradication of **Helicobacter pylori**.

Note A standard dose proton pump inhibitor is one of: esomeprazole 20 mg, lansoprazole 30 mg, omeprazole 20 mg, pantoprazole 40 mg, rabeprazole 20 mg.

Note A low dose proton pump inhibitor includes: lansoprazole 15mg, omeprazole 10mg, pantoprazole 20mg and rabeprazole 10mg.

Note Pharmaceutical benefits that have the form lansoprazole capsule 30 mg and pharmaceutical benefits that have the form lansoprazole tablet 30 mg (orally disintegrating) are equivalent for the purposes of substitution.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Complex gastro-oesophageal reflux disease (GORD)

Treatment Phase: One of: (1) establishment of symptom control, (2) maintenance treatment, (3) re-establishment of symptom control

Treatment criteria:

- Must be treated by a gastroenterologist; OR

- Must be treated by a surgeon with expertise in the upper gastrointestinal tract; OR
- Must be treated by a medical practitioner who has consulted at least one of the above mentioned specialists in relation to this current PBS benefit being sought, with the specialist's name documented in the patient's medical records for auditing purposes; OR
- Must be treated by a medical practitioner who has not consulted a specialist, but only if treatment continues therapy initiated under this restriction with involvement by a specialist (i.e. continuing treatment initiated for non-complex GORD does not meet this criterion), with the specialist's name documented in the patient's medical records for auditing purposes.

Clinical criteria:

- The treatment must be: (i) the sole PBS-subsidised proton pump inhibitor (PPI) for this condition, (ii) the sole strength of this PPI, (iii) the sole form of PPI, **AND**
- Patient must have symptoms inadequately controlled with each of: (i) a standard dose proton pump inhibitor (PPI) administered once daily, (ii) a low dose PPI administered twice daily; treatment is for: (1) establishment of symptom control; OR
- Patient must be assessed for the risks/benefits of a step-down in dosing from standard dose PPI administered twice daily, with the determination being that the risks outweigh the benefits; treatment is for: (2) maintenance treatment; OR
- Patient must have trialled a step-down in dosing, yet symptoms have re-emerged/worsened; treatment is for: (3) re-establishment of symptom control; OR
- Patient must have trialled a step-down in dosing, with symptoms adequately managed with once daily dosing; treatment is for: (2) maintenance treatment, but with the quantity sought in this authority application being up to 1 pack per dispensing.

Check patient adherence to any preceding PPI treatment regimen. Exclude non-adherence as a cause of inadequate control before accessing treatment under this restriction.

lansoprazole 30 mg enteric capsule, 28

12284M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	5	..	*23.28	24.57	^a APO-Lansoprazole [TX] ^a NOUMED LANSOPRAZOLE [VO]	^a Lanzopran [RA] ^a Zopral [AF]

lansoprazole 30 mg orally disintegrating tablet, 28

12276D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	5	..	*23.28	24.57	^a APO-Lansoprazole ODT [TX] ^a Zopral ODT [AF]	^a Lansoprazole ODT GH [GQ]
			^B 9.62	*32.90	24.57	^a Zoton FasTabs [PF]	

■ OMEPRAZOLE

Note Check patient adherence to lower dose proton pump inhibitor before "stepping-up" therapy.

Restricted benefit

Gastro-oesophageal reflux disease

Restricted benefit

Scleroderma oesophagus

Restricted benefit

Zollinger-Ellison syndrome

omeprazole 10 mg enteric tablet, 30

8332M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	15.80	17.09	Losec Tablets [AP]

NP

■ OMEPRAZOLE

Note Pharmaceutical benefits that have the forms omeprazole tablet 20 mg, omeprazole capsule 20 mg and omeprazole tablet 20 mg (as magnesium) are equivalent for the purposes of substitution.

Note Check patient adherence to lower dose proton pump inhibitor before "stepping-up" therapy.

Note Low dose proton pump inhibitors are appropriate step-down therapy from standard dose proton pump inhibitors.

Note A low dose proton pump inhibitor includes: lansoprazole 15mg, omeprazole 10mg, pantoprazole 20mg and rabeprazole 10mg.

Authority required (STREAMLINED)

8780

Scleroderma oesophagus

Authority required (STREAMLINED)

8866

Zollinger-Ellison syndrome

omeprazole 20 mg enteric capsule, 30

1327W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	16.45	17.74	^a APO-Omeprazole [TX] ^a Omeprazole Sandoz [HX] ^a Pharmacor Omeprazole 20 [CR]	^a Maxor [AF] ^a Pemzo [RW] ^a Probitor [SZ]

NP

omeprazole 20 mg enteric tablet, 30

8333N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	16.45	17.84	^a APO-Omeprazole [TX] ^a Omeprazole AN [EA] ^a Ozmepr [ZP]	^a Maxor EC Tabs [AF] ^a Omeprazole generichealth [GQ]

omeprazole 20 mg enteric tablet, 30

9110L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	16.45	17.74	^a Acimax Tablets [AL] ^a Omeprazole Sandoz [SZ] ^B 7.36	^a Omepral [ZA]
				23.81	17.74	^a Losec Tablets [AP]	

■ OMEPRAZOLE

Note Pharmaceutical benefits that have the forms omeprazole tablet 20 mg, omeprazole capsule 20 mg and omeprazole tablet 20 mg (as magnesium) are equivalent for the purposes of substitution.

Note Check patient adherence to lower dose proton pump inhibitor before "stepping-up" therapy.

Note Low dose proton pump inhibitors are appropriate step-down therapy from standard dose proton pump inhibitors.

Note A low dose proton pump inhibitor includes: lansoprazole 15mg, omeprazole 10mg, pantoprazole 20mg and rabeprazole 10mg.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)**8776**

Gastro-oesophageal reflux disease

Clinical criteria:

- The treatment must be for long-term maintenance of gastro-oesophageal reflux disease in a patient with symptoms inadequately controlled using a low dose proton pump inhibitor.

omeprazole 20 mg enteric capsule, 30

11682W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	16.45	17.74	^a APO-Omeprazole [TX] ^a Omeprazole Sandoz [HX] ^a Pharmacor Omeprazole 20 [CR]	^a Maxor [AF] ^a Pemzo [RW] ^a Probitor [SZ]

omeprazole 20 mg enteric tablet, 30

11677N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	16.45	17.74	^a Acimax Tablets [AL] ^a Omeprazole Sandoz [SZ] ^B 7.36	^a Omepral [ZA]
				23.81	17.74	^a Losec Tablets [AP]	

omeprazole 20 mg enteric tablet, 30

11683X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	16.45	17.74	^a APO-Omeprazole [TX] ^a Omeprazole AN [EA] ^a Ozmepr [ZP]	^a Maxor EC Tabs [AF] ^a Omeprazole generichealth [GQ]

■ OMEPRAZOLE

Note Pharmaceutical benefits that have the forms omeprazole tablet 20 mg, omeprazole capsule 20 mg and omeprazole tablet 20 mg (as magnesium) are equivalent for the purposes of substitution.

Note Check patient adherence to lower dose proton pump inhibitor before "stepping-up" therapy.

Note Low dose proton pump inhibitors are appropriate step-down therapy from standard dose proton pump inhibitors.

Note A low dose proton pump inhibitor includes: lansoprazole 15mg, omeprazole 10mg, pantoprazole 20mg and rabeprazole 10mg.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)**8775**

Peptic ulcer

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have tested negative for helicobacter pylori infection; OR
- Patient must have failed treatment with helicobacter pylori eradication therapy.

Authority required (STREAMLINED)**8774**

Gastro-oesophageal reflux disease

Clinical criteria:

- The treatment must be for initial treatment of symptomatic gastro-oesophageal reflux disease; OR

- The treatment must be for the short-term maintenance treatment of gastro-oesophageal reflux disease.

omeprazole 20 mg enteric capsule, 30

1326T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	1	..	16.45	17.74	^a APO-Omeprazole [TX] ^a Omeprazole Sandoz [HX] ^a Pharmacor Omeprazole 20 [CR]	^a Maxor [AF] ^a Pemzo [RW] ^a Probitor [SZ]

omeprazole 20 mg enteric tablet, 30

8331L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	1	..	16.45	17.84	^a APO-Omeprazole [TX] ^a Omeprazole AN [EA] ^a Ozmepr [ZP]	^a Maxor EC Tabs [AF] ^a Omeprazole generichealth [GQ]

omeprazole 20 mg enteric tablet, 30

9109K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	1	..	16.45	17.74	^a Acimax Tablets [AL] ^a Omeprazole Sandoz [SZ] ^b 7.36	^a Omepral [ZA]
				23.81	17.74	^a Losec Tablets [AP]	

■ OMEPRAZOLE

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Note The treatment must not be prescribed under this restriction for any of the following PBS-indications: (i) gastro-oesophageal reflux disease (where the word 'complex' is absent), (ii) scleroderma oesophagus, (iii) pathological hypersecretory conditions including Zollinger-Ellison syndrome and idiopathic hypersecretion, (iv) peptic ulcer, (v) eradication of **Helicobacter pylori**.

Note A standard dose proton pump inhibitor is one of: esomeprazole 20 mg, lansoprazole 30 mg, omeprazole 20 mg, pantoprazole 40 mg, rabeprazole 20 mg.

Note A low dose proton pump inhibitor includes: lansoprazole 15mg, omeprazole 10mg, pantoprazole 20mg and rabeprazole 10mg.

Note Pharmaceutical benefits that have the forms omeprazole tablet 20 mg, omeprazole capsule 20 mg and omeprazole tablet 20 mg (as magnesium) are equivalent for the purposes of substitution.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Complex gastro-oesophageal reflux disease (GORD)

Treatment Phase: One of: (1) establishment of symptom control, (2) maintenance treatment, (3) re-establishment of symptom control

Treatment criteria:

- Must be treated by a gastroenterologist; OR
- Must be treated by a surgeon with expertise in the upper gastrointestinal tract; OR
- Must be treated by a medical practitioner who has consulted at least one of the above mentioned specialists in relation to this current PBS benefit being sought, with the specialist's name documented in the patient's medical records for auditing purposes; OR
- Must be treated by a medical practitioner who has not consulted a specialist, but only if treatment continues therapy initiated under this restriction with involvement by a specialist (i.e. continuing treatment initiated for non-complex GORD does not meet this criterion), with the specialist's name documented in the patient's medical records for auditing purposes.

Clinical criteria:

- The treatment must be: (i) the sole PBS-subsidised proton pump inhibitor (PPI) for this condition, (ii) the sole strength of this PPI, (iii) the sole form of PPI, **AND**
- Patient must have symptoms inadequately controlled with each of: (i) a standard dose proton pump inhibitor (PPI) administered once daily, (ii) a low dose PPI administered twice daily; treatment is for: (1) establishment of symptom control; OR
- Patient must be assessed for the risks/benefits of a step-down in dosing from standard dose PPI administered twice daily, with the determination being that the risks outweigh the benefits; treatment is for: (2) maintenance treatment; OR
- Patient must have trialled a step-down in dosing, yet symptoms have re-emerged/worsened; treatment is for: (3) re-establishment of symptom control; OR
- Patient must have trialled a step-down in dosing, with symptoms adequately managed with once daily dosing; treatment is for: (2) maintenance treatment, but with the quantity sought in this authority application being up to 1 pack per dispensing.

Check patient adherence to any preceding PPI treatment regimen. Exclude non-adherence as a cause of inadequate control before accessing treatment under this restriction.

omeprazole 20 mg enteric capsule, 30

12281J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	5	..	*20.88	22.17	^a APO-Omeprazole [TX] ^a Omeprazole Sandoz [HX] ^a Pharmacor Omeprazole 20 [CR]	^a Maxor [AF] ^a Pemzo [RW] ^a Probitor [SZ]

omeprazole 20 mg enteric tablet, 30

12270T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	5	..	*20.88	22.17	^a Acimax Tablets [AL] ^a Omeprazole Sandoz [SZ]	^a Omepral [ZA]
			^B 14.72	*35.60	22.17	^a Losec Tablets [AP]	

omeprazole 20 mg enteric tablet, 30

12272X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	5	..	*20.88	22.17	^a APO-Omeprazole [TX] ^a Omeprazole AN [EA] ^a Ozmepr [ZP]	^a Maxor EC Tabs [AF] ^a Omeprazole generichealth [GQ]

■ PANTOPRAZOLE

Note Check patient adherence to lower dose proton pump inhibitor before "stepping-up" therapy.

Restricted benefit

Gastro-oesophageal reflux disease

Restricted benefit

Scleroderma oesophagus

Restricted benefit

Zollinger-Ellison syndrome

pantoprazole 20 mg enteric tablet, 30

8399C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	13.59	14.88	^a APO-Pantoprazole [TX] ^a Ozpan [RA] ^a Pantoprazole AN [EA] ^a Pantoprazole generichealth [HQ] ^a Salpraz [AF] ^a Sozol [RW]	^a NOUMED PANTOPRAZOLE [VO] ^a Panthron [ER] ^a Pantoprazole APOTEX [TY] ^a Pantoprazole Sandoz [SZ] ^a Somac [NQ]

■ PANTOPRAZOLE

Note Check patient adherence to lower dose proton pump inhibitor before "stepping-up" therapy.

Note Low dose proton pump inhibitors are appropriate step-down therapy from standard dose proton pump inhibitors.

Note A low dose proton pump inhibitor includes: lansoprazole 15mg, omeprazole 10mg, pantoprazole 20mg and rabeprazole 10mg.

Authority required (STREAMLINED)

8780

Scleroderma oesophagus

Authority required (STREAMLINED)

8866

Zollinger-Ellison syndrome

pantoprazole 40 mg enteric coated granules, 30 sachets

9424B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	33.14	34.43	Somac [NQ]

pantoprazole 40 mg enteric tablet, 30

8008L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	14.60	15.89	^a APO-Pantoprazole [TX] ^a NOUMED PANTOPRAZOLE [VO] ^a Panthron [ER] ^a Pantoprazole AN [EA] ^a Pantoprazole generichealth [HQ] ^a Salpraz [AF] ^a Sozol [RW]	^a I-Pantoprazole [CR] ^a Ozpan [RA] ^a Pantoprazole Actavis [ED] ^a Pantoprazole APOTEX [TY] ^a Pantoprazole Sandoz [SZ] ^a Somac [NQ]

■ PANTOPRAZOLE

Note Check patient adherence to lower dose proton pump inhibitor before "stepping-up" therapy.

Note Low dose proton pump inhibitors are appropriate step-down therapy from standard dose proton pump inhibitors.

Note A low dose proton pump inhibitor includes: lansoprazole 15mg, omeprazole 10mg, pantoprazole 20mg and rabeprazole 10mg.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

8776

Gastro-oesophageal reflux disease

Clinical criteria:

- The treatment must be for long-term maintenance of gastro-oesophageal reflux disease in a patient with symptoms inadequately controlled using a low dose proton pump inhibitor.

pantoprazole 40 mg enteric coated granules, 30 sachets

11678P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	33.14	34.43	Somac [NQ]

pantoprazole 40 mg enteric tablet, 30

11681T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	14.60	15.89	^a APO-Pantoprazole [TX] ^a NOUMED PANTOPRAZOLE [VO] ^a Panthron [ER] ^a Pantoprazole AN [EA] ^a Pantoprazole generichealth [HQ] ^a Salpraz [AF] ^a Sozol [RW]	^a I-Pantoprazole [CR] ^a Ozpan [RA] ^a Pantoprazole Actavis [ED] ^a Pantoprazole APOTEX [TY] ^a Pantoprazole Sandoz [SZ] ^a Somac [NQ]

■ PANTOPRAZOLE

Note Check patient adherence to lower dose proton pump inhibitor before "stepping-up" therapy.

Note Low dose proton pump inhibitors are appropriate step-down therapy from standard dose proton pump inhibitors.

Note A low dose proton pump inhibitor includes: lansoprazole 15mg, omeprazole 10mg, pantoprazole 20mg and rabeprazole 10mg.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)**8775**

Peptic ulcer

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have tested negative for helicobacter pylori infection; OR
- Patient must have failed treatment with helicobacter pylori eradication therapy.

Authority required (STREAMLINED)**8774**

Gastro-oesophageal reflux disease

Clinical criteria:

- The treatment must be for initial treatment of symptomatic gastro-oesophageal reflux disease; OR
- The treatment must be for the short-term maintenance treatment of gastro-oesophageal reflux disease.

pantoprazole 40 mg enteric coated granules, 30 sachets

9423Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	33.14	34.43	Somac [NQ]

pantoprazole 40 mg enteric tablet, 30

8007K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	1	..	14.60	15.89	^a APO-Pantoprazole [TX] ^a NOUMED PANTOPRAZOLE [VO] ^a Panthron [ER] ^a Pantoprazole AN [EA] ^a Pantoprazole generichealth [HQ] ^a Salpraz [AF] ^a Sozol [RW]	^a I-Pantoprazole [CR] ^a Ozpan [RA] ^a Pantoprazole Actavis [ED] ^a Pantoprazole APOTEX [TY] ^a Pantoprazole Sandoz [SZ] ^a Somac [NQ]

■ PANTOPRAZOLE

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Note The treatment must not be prescribed under this restriction for any of the following PBS-indications: (i) gastro-oesophageal reflux disease (where the word 'complex' is absent), (ii) scleroderma oesophagus, (iii) pathological hypersecretory conditions including Zollinger-Ellison syndrome and idiopathic hypersecretion, (iv) peptic ulcer, (v) eradication of **Helicobacter pylori**.

Note A standard dose proton pump inhibitor is one of: esomeprazole 20 mg, lansoprazole 30 mg, omeprazole 20 mg, pantoprazole 40 mg, rabeprazole 20 mg.

Note A low dose proton pump inhibitor includes: lansoprazole 15mg, omeprazole 10mg, pantoprazole 20mg and rabeprazole 10mg.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Complex gastro-oesophageal reflux disease (GORD)

Treatment Phase: One of: (1) establishment of symptom control, (2) maintenance treatment, (3) re-establishment of symptom control

Treatment criteria:

- Must be treated by a gastroenterologist; OR
- Must be treated by a surgeon with expertise in the upper gastrointestinal tract; OR
- Must be treated by a medical practitioner who has consulted at least one of the above mentioned specialists in relation to this current PBS benefit being sought, with the specialist's name documented in the patient's medical records for auditing purposes; OR
- Must be treated by a medical practitioner who has not consulted a specialist, but only if treatment continues therapy initiated under this restriction with involvement by a specialist (i.e. continuing treatment initiated for non-complex GORD does not meet this criterion), with the specialist's name documented in the patient's medical records for auditing purposes.

Clinical criteria:

- The treatment must be: (i) the sole PBS-subsidised proton pump inhibitor (PPI) for this condition, (ii) the sole strength of this PPI, (iii) the sole form of PPI, **AND**
- Patient must have symptoms inadequately controlled with each of: (i) a standard dose proton pump inhibitor (PPI) administered once daily, (ii) a low dose PPI administered twice daily; treatment is for: (1) establishment of symptom control; OR
- Patient must be assessed for the risks/benefits of a step-down in dosing from standard dose PPI administered twice daily, with the determination being that the risks outweigh the benefits; treatment is for: (2) maintenance treatment; OR
- Patient must have trialled a step-down in dosing, yet symptoms have re-emerged/worsened; treatment is for: (3) re-establishment of symptom control; OR
- Patient must have trialled a step-down in dosing, with symptoms adequately managed with once daily dosing; treatment is for: (2) maintenance treatment, but with the quantity sought in this authority application being up to 1 pack per dispensing.

Check patient adherence to any preceding PPI treatment regimen. Exclude non-adherence as a cause of inadequate control before accessing treatment under this restriction.

pantoprazole 40 mg enteric coated granules, 30 sachets

12282K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*54.26	41.30	Somac [NQ]

pantoprazole 40 mg enteric tablet, 30

12277E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	5	..	*17.18	18.47	^a APO-Pantoprazole [TX] ^a NOUMED PANTOPRAZOLE [VO] ^a Panthron [ER] ^a Pantoprazole AN [EA] ^a Pantoprazole generichealth [HQ] ^a Salpraz [AF] ^a Sozol [RW]	^a I-Pantoprazole [CR] ^a Ozpan [RA] ^a Pantoprazole Actavis [ED] ^a Pantoprazole APOTEX [TY] ^a Pantoprazole Sandoz [SZ] ^a Somac [NQ]

■ RABEPRAZOLE**Note** Check patient adherence to lower dose proton pump inhibitor before "stepping-up" therapy.**Restricted benefit**

Gastro-oesophageal reflux disease

Restricted benefit

Scleroderma oesophagus

rabeprazole sodium 10 mg enteric tablet, 28

8507R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	15.26	16.55	^a APO-Rabeprazole [TX] ^a Rabeprazole AN [EA]	^a Parbezol [RW] ^a Rabeprazole Sandoz [SZ]
			^B 4.81	20.07	16.55	^a Pariet [JC]	

■ RABEPRAZOLE**Note** Check patient adherence to lower dose proton pump inhibitor before "stepping-up" therapy.**Note** Low dose proton pump inhibitors are appropriate step-down therapy from standard dose proton pump inhibitors.**Note** A low dose proton pump inhibitor includes: lansoprazole 15mg, omeprazole 10mg, pantoprazole 20mg and rabeprazole 10mg.**Authority required (STREAMLINED)****8780**

Scleroderma oesophagus

rabeprazole sodium 20 mg enteric tablet, 30

8508T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	15.06	16.35	^a APO-Rabeprazole [TX] ^a Rabeprazole AN [EA]	^a Parbezol [RW] ^a Rabeprazole Mylan [AF]

^a Rabeprazole Sandoz [SZ]^a Rabeprazole SUN [RN]^a Zabep [AL]^B5.01

20.07

16.35

^a Pariet [JC]**■ RABEPRAZOLE****Note** Check patient adherence to lower dose proton pump inhibitor before "stepping-up" therapy.**Note** Low dose proton pump inhibitors are appropriate step-down therapy from standard dose proton pump inhibitors.**Note** A low dose proton pump inhibitor includes: lansoprazole 15mg, omeprazole 10mg, pantoprazole 20mg and rabeprazole 10mg.**Note** No increase in the maximum quantity or number of units may be authorised.**Note** No increase in the maximum number of repeats may be authorised.**Authority required (STREAMLINED)****8776**

Gastro-oesophageal reflux disease

Clinical criteria:

- The treatment must be for long-term maintenance of gastro-oesophageal reflux disease in a patient with symptoms inadequately controlled using a low dose proton pump inhibitor.

rabeprazole sodium 20 mg enteric tablet, 30

11670F

NP

Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
1	5	..	15.06	16.35	^a APO-Rabeprazole [TX] ^a Rabeprazole AN [EA] ^a Rabeprazole Sandoz [SZ] ^a Zabep [AL]	^a Parbezol [RW] ^a Rabeprazole Mylan [AF] ^a Rabeprazole SUN [RN]
^B 5.01	20.07	16.35	^a Pariet [JC]			

■ RABEPRAZOLE**Note** Check patient adherence to lower dose proton pump inhibitor before "stepping-up" therapy.**Note** Low dose proton pump inhibitors are appropriate step-down therapy from standard dose proton pump inhibitors.**Note** A low dose proton pump inhibitor includes: lansoprazole 15mg, omeprazole 10mg, pantoprazole 20mg and rabeprazole 10mg.**Note** No increase in the maximum quantity or number of units may be authorised.**Note** No increase in the maximum number of repeats may be authorised.**Authority required (STREAMLINED)****8775**

Peptic ulcer

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have tested negative for helicobacter pylori infection; OR
- Patient must have failed treatment with helicobacter pylori eradication therapy.

Authority required (STREAMLINED)**8774**

Gastro-oesophageal reflux disease

Clinical criteria:

- The treatment must be for initial treatment of symptomatic gastro-oesophageal reflux disease; OR
- The treatment must be for the short-term maintenance treatment of gastro-oesophageal reflux disease.

rabeprazole sodium 20 mg enteric tablet, 30

8509W

NP

Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
1	1	..	15.06	16.35	^a APO-Rabeprazole [TX] ^a Rabeprazole AN [EA] ^a Rabeprazole Sandoz [SZ] ^a Zabep [AL]	^a Parbezol [RW] ^a Rabeprazole Mylan [AF] ^a Rabeprazole SUN [RN]
^B 5.01	20.07	16.35	^a Pariet [JC]			

■ RABEPRAZOLE**Note** Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.**Note** The treatment must not be prescribed under this restriction for any of the following PBS-indications: (i) gastro-oesophageal reflux disease (where the word 'complex' is absent), (ii) scleroderma oesophagus, (iii) pathological hypersecretory conditions including Zollinger-Ellison syndrome and idiopathic hypersecretion, (iv) peptic ulcer, (v) eradication of **Helicobacter pylori**.**Note** A standard dose proton pump inhibitor is one of: esomeprazole 20 mg, lansoprazole 30 mg, omeprazole 20 mg, pantoprazole 40 mg, rabeprazole 20 mg.**Note** A low dose proton pump inhibitor includes: lansoprazole 15mg, omeprazole 10mg, pantoprazole 20mg and rabeprazole 10mg.**Note** No increase in the maximum quantity or number of units may be authorised.**Note** No increase in the maximum number of repeats may be authorised.**Authority required**

Complex gastro-oesophageal reflux disease (GORD)

Treatment Phase: One of: (1) establishment of symptom control, (2) maintenance treatment, (3) re-establishment of symptom control

Treatment criteria:

- Must be treated by a gastroenterologist; OR
- Must be treated by a surgeon with expertise in the upper gastrointestinal tract; OR
- Must be treated by a medical practitioner who has consulted at least one of the above mentioned specialists in relation to this current PBS benefit being sought, with the specialist's name documented in the patient's medical records for auditing purposes; OR
- Must be treated by a medical practitioner who has not consulted a specialist, but only if treatment continues therapy initiated under this restriction with involvement by a specialist (i.e. continuing treatment initiated for non-complex GORD does not meet this criterion), with the specialist's name documented in the patient's medical records for auditing purposes.

Clinical criteria:

- The treatment must be: (i) the sole PBS-subsidised proton pump inhibitor (PPI) for this condition, (ii) the sole strength of this PPI, (iii) the sole form of PPI, **AND**
- Patient must have symptoms inadequately controlled with each of: (i) a standard dose proton pump inhibitor (PPI) administered once daily, (ii) a low dose PPI administered twice daily; treatment is for: (1) establishment of symptom control; OR
- Patient must be assessed for the risks/benefits of a step-down in dosing from standard dose PPI administered twice daily, with the determination being that the risks outweigh the benefits; treatment is for: (2) maintenance treatment; OR
- Patient must have trialled a step-down in dosing, yet symptoms have re-emerged/worsened; treatment is for: (3) re-establishment of symptom control; OR
- Patient must have trialled a step-down in dosing, with symptoms adequately managed with once daily dosing; treatment is for: (2) maintenance treatment, but with the quantity sought in this authority application being up to 1 pack per dispensing.

Check patient adherence to any preceding PPI treatment regimen. Exclude non-adherence as a cause of inadequate control before accessing treatment under this restriction.

rabeprazole sodium 20 mg enteric tablet, 30

12286P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	5	..	*18.10	19.39	^a APO-Rabeprazole [TX] ^a Rabeprazole AN [EA] ^a Rabeprazole Sandoz [SZ] ^a Zabep [AL]	^a Parbezol [RW] ^a Rabeprazole Mylan [AF] ^a Rabeprazole SUN [RN]
			^B 10.02	*28.12	19.39	^a Pariet [JC]	

Combinations for eradication of Helicobacter pylori

▪ **ESOMEPRAZOLE (&) CLARITHROMYCIN (&) AMOXICILLIN**

Note Pharmaceutical benefits that have the form pack containing 14 tablets (enteric coated) containing esomeprazole 20 mg (as magnesium trihydrate), 14 tablets clarithromycin 500 mg and 28 capsules amoxicillin 500 mg (as trihydrate) and pack containing 14 tablets (enteric coated) containing esomeprazole 20 mg (as magnesium), 14 tablets clarithromycin 500 mg and 28 capsules amoxicillin 500 mg (as trihydrate) are equivalent for the purposes of substitution.

Restricted benefit

Eradication of Helicobacter pylori

Clinical criteria:

- The condition must be associated with peptic ulcer disease.

esomeprazole 20 mg enteric tablet [14] (&) amoxicillin 500 mg capsule [28] (&) clarithromycin 500 mg tablet [14], 1 pack

10759G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	37.45	38.74	^a ESOMEPRAZOLE SANDOZ Hp7 [SZ]

esomeprazole 20 mg enteric tablet [14] (&) amoxicillin 500 mg capsule [28] (&) clarithromycin 500 mg tablet [14], 1 pack

8738X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	^B 3.58	41.03	38.74	^a Nexium Hp7 [AP]

▪ **DRUGS FOR FUNCTIONAL GASTROINTESTINAL DISORDERS**

BELLADONNA AND DERIVATIVES, PLAIN

Belladonna alkaloids, tertiary amines

▪ **ATROPINE SULFATE**

atropine sulfate monohydrate 600 microgram/mL injection, 10 x 1 mL ampoules

5022H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	23.69	24.98	Pfizer Australia Pty Ltd [PF]

▪ **ATROPINE SULFATE**

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical

practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

atropine sulfate monohydrate 600 microgram/mL injection, 10 x 1 mL ampoules

1089H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	23.69	24.98	Pfizer Australia Pty Ltd [PF]

PROPULSIVES*Propulsives***■ DOMPERIDONE****domperidone 10 mg tablet, 25**

1347X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	14.28	15.57	Motilium [JC]

■ METOCLOPRAMIDE**metoclopramide hydrochloride 10 mg tablet, 25**

1207M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP MW	1	..	..	13.48	14.77	^a APO-Metoclopramide [TX]	^a EMEXLON [RW]
			^B 1.78	15.26	14.77	^a Metoclopramide AN [EA]	^a Pramin [AF]
						^a Maxolon [IL]	

metoclopramide hydrochloride 10 mg tablet, 25

5151D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
DP	1	..	..	13.48	14.77	^a APO-Metoclopramide [TX]	^a EMEXLON [RW]
			^B 1.78	15.26	14.77	^a Metoclopramide AN [EA]	^a Pramin [AF]
						^a Maxolon [IL]	

metoclopramide hydrochloride 10 mg/2 mL injection, 10 x 2 mL ampoules

1206L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP MW	1	..	..	17.73	19.02	Maxolon [IL]

metoclopramide hydrochloride 10 mg/2 mL injection, 10 x 2 mL ampoules

5153F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	1	..	..	17.73	19.02	Maxolon [IL]

■ ANTIEMETICS AND ANTINAUSEANTS**ANTIEMETICS AND ANTINAUSEANTS***Serotonin (5HT₃) antagonists***■ GRANISETRON****Restricted benefit**

Nausea and vomiting

Clinical criteria:

- The condition must be associated with cytotoxic chemotherapy being used to treat malignancy which occurs within 48 hours of chemotherapy administration.

Increased maximum quantities will be limited to a maximum of 7 days per chemotherapy cycle.

granisetron 3 mg/3 mL injection, 3 mL ampoule

8729K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	14.35	15.64	^a Granisetron-AFT [AE]	^a Granisetron Kabi [PK]
						^a Kytril [IX]	

■ GRANISETRON**Authority required (STREAMLINED)**

4092

Nausea and vomiting

Clinical criteria:

- The condition must be associated with radiotherapy being used to treat malignancy.

granisetron 3 mg/3 mL injection, 3 mL ampoule

8730L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	14.35	15.64	^a Granisetron-AFT [AE]	^a Granisetron Kabi [PK]
						^a Kytril [IX]	

■ GRANISETRON**Authority required (STREAMLINED)**

10498

Nausea and vomiting

Clinical criteria:

- The condition must be associated with radiotherapy being used to treat malignancy; OR
- The condition must be associated with oral chemotherapy being used to treat malignancy.

granisetron 2 mg tablet, 5

8873B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	56.59	41.30	Kytril [IX]

▪ **GRANISETRON**

Restricted benefit

Nausea and vomiting

Clinical criteria:

- The condition must be associated with cytotoxic chemotherapy being used to treat malignancy which occurs within 48 hours of chemotherapy administration.

Increased maximum quantities will be limited to a maximum of 7 days per chemotherapy cycle.

granisetron 2 mg tablet, 1

8728J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	..	..	*29.84	31.13	Kytril [IX]

▪ **NETUPITANT + PALONOSETRON**

Note No increase in the maximum number of repeats may be authorised.

Note No increase in the maximum quantity or number of units may be authorised.

Note This medicine is not PBS-subsidised for nausea and vomiting associated with radiotherapy being used to treat malignancy.

Authority required (STREAMLINED)

5991

Nausea and vomiting

Clinical criteria:

- The condition must be associated with cytotoxic chemotherapy being used to treat malignancy, **AND**
- The treatment must be in combination with dexamethasone, **AND**
- Patient must be scheduled to be administered a chemotherapy regimen that includes any 1 of the following agents: altretamine; carmustine; cisplatin when a single dose constitutes a cycle of chemotherapy; cyclophosphamide at a dose of 1500 mg per square metre per day or greater; dacarbazine; procarbazine when a single dose constitutes a cycle of chemotherapy; streptozocin.

No more than 1 capsule of 300 mg netupitant/0.5 mg palonosetron fixed dose combination will be authorised per cycle of cytotoxic chemotherapy.

Authority required (STREAMLINED)

5994

Nausea and vomiting

Clinical criteria:

- The condition must be associated with cytotoxic chemotherapy being used to treat breast cancer, **AND**
- The treatment must be in combination with dexamethasone, **AND**
- Patient must be scheduled to be co-administered cyclophosphamide and an anthracycline.

No more than 1 capsule of 300 mg netupitant/0.5 mg palonosetron fixed dose combination will be authorised per cycle of cytotoxic chemotherapy.

Authority required (STREAMLINED)

6937

Nausea and vomiting

Clinical criteria:

- The condition must be associated with moderately emetogenic cytotoxic chemotherapy being used to treat malignancy, **AND**
- The treatment must be in combination with dexamethasone on day 1 of a chemotherapy cycle, **AND**
- Patient must have had a prior episode of chemotherapy induced nausea or vomiting, **AND**
- Patient must be scheduled to be administered a chemotherapy regimen that includes any 1 of the following intravenous chemotherapy agents: arsenic trioxide; azacitidine; cyclophosphamide at a dose of less than 1500 mg per square metre per day; cytarabine at a dose of greater than 1 g per square metre per day; dactinomycin; daunorubicin; doxorubicin; epirubicin; fotemustine; idarubicin; ifosfamide; irinotecan; melphalan; methotrexate at a dose of 250 mg to 1 g per square metre; raltitrexed.

No more than 1 capsule of 300 mg netupitant/0.5 mg palonosetron fixed dose combination will be authorised per cycle of cytotoxic chemotherapy.

Authority required (STREAMLINED)

6879

Nausea and vomiting

Clinical criteria:

- The condition must be associated with moderately emetogenic cytotoxic chemotherapy being used to treat malignancy, **AND**
- The treatment must be in combination with dexamethasone on day 1 of a chemotherapy cycle, **AND**

- Patient must be scheduled to be administered a chemotherapy regimen that includes either carboplatin or oxaliplatin. No more than 1 capsule of 300 mg netupitant/0.5 mg palonosetron fixed dose combination will be authorised per cycle of cytotoxic chemotherapy.

netupitant 300 mg + palonosetron 500 microgram capsule, 1

10731T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	116.71	41.30	Akynzeo [MF]

■ ONDANSETRON**Authority required (STREAMLINED)****10498**

Nausea and vomiting

Clinical criteria:

- The condition must be associated with radiotherapy being used to treat malignancy; OR
- The condition must be associated with oral chemotherapy being used to treat malignancy.

ondansetron 8 mg tablet, 10

1595Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	1	..	26.41	27.70	^a APO-Ondansetron [TX] ^a Ondansetron APOTEX [GX] ^a Ondansetron Mylan Tablets [AF] ^a Zofran [AS]	^a Ondansetron AN [EA] ^a Ondansetron-DRLA [RZ] ^a Ondansetron SZ [HX] ^a Zotren 8 [RF]

ondansetron 4 mg tablet, 10

1594X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	1	..	21.19	22.48	^a APO-Ondansetron [TX] ^a Ondansetron APOTEX [GX] ^a Ondansetron Mylan Tablets [AF] ^a Zofran [AS]	^a Ondansetron AN [EA] ^a Ondansetron-DRLA [RZ] ^a Ondansetron SZ [HX] ^a Zotren 4 [RF]

■ ONDANSETRON**Restricted benefit**

Nausea and vomiting

Clinical criteria:

- The condition must be associated with cytotoxic chemotherapy being used to treat malignancy which occurs within 48 hours of chemotherapy administration.
- Increased maximum quantities will be limited to a maximum of 7 days per chemotherapy cycle.

ondansetron 8 mg tablet, 4

8225X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	17.78	19.07	^a APO-Ondansetron [TX] ^a Ondansetron APOTEX [GX] ^a Ondansetron Mylan Tablets [AF] ^a Zofran [AS]	^a Ondansetron AN [EA] ^a Ondansetron-DRLA [RZ] ^a Ondansetron SZ [HX] ^a Zotren 8 [RF]

ondansetron 4 mg tablet, 4

8224W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	15.84	17.13	^a APO-Ondansetron [TX] ^a Ondansetron APOTEX [GX] ^a Ondansetron Mylan Tablets [AF] ^a Zofran [AS]	^a Ondansetron AN [EA] ^a Ondansetron-DRLA [RZ] ^a Ondansetron SZ [HX] ^a Zotren 4 [RF]

■ ONDANSETRON**Authority required (STREAMLINED)****4102**

Nausea and vomiting

Clinical criteria:

- The condition must be associated with radiotherapy being used to treat malignancy.

ondansetron 4 mg/5 mL oral liquid, 50 mL

8233H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	1	..	98.87	41.30	Zofran syrup 50 mL [AS]

■ ONDANSETRON**Restricted benefit**

Nausea and vomiting

Clinical criteria:

- The condition must be associated with cytotoxic chemotherapy being used to treat malignancy which occurs within 48 hours of chemotherapy administration.
- Increased maximum quantities will be limited to a maximum of 7 days per chemotherapy cycle.

ondansetron 4 mg/5 mL oral liquid, 50 mL

9441X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	..	..	98.87	41.30	Zofran syrup 50 mL [AS]

■ ONDANSETRON

Note Pharmaceutical benefits that have the form ondansetron tablet (orally disintegrating) 4 mg and pharmaceutical benefits that have the form ondansetron 4 mg wafer are equivalent for the purposes of substitution.

Authority required (STREAMLINED)**10498**

Nausea and vomiting

Clinical criteria:

- The condition must be associated with radiotherapy being used to treat malignancy; OR
- The condition must be associated with oral chemotherapy being used to treat malignancy.

ondansetron 4 mg wafer, 10

8412R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	^B 2.45	23.64	22.48	^a Zofran Zydys [AS]

ondansetron 4 mg orally disintegrating tablet, 10

5472B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	1	..	21.19	22.48	^a APO-Ondansetron ODT [TX] ^a Ondansetron Mylan ODT [AF] ^a Ondansetron ODT GH [GQ] ^a Zilfajim ODT 4 [DO]	^a Ondansetron AN ODT [EA] ^a Ondansetron ODT-DRLA [RZ] ^a Ondansetron SZ ODT [HX] ^a Zotren ODT [RF]

■ ONDANSETRON

Note Pharmaceutical benefits that have the form ondansetron tablet (orally disintegrating) 8 mg and pharmaceutical benefits that have the form ondansetron 8 mg wafer are equivalent for the purposes of substitution.

Authority required (STREAMLINED)**10498**

Nausea and vomiting

Clinical criteria:

- The condition must be associated with radiotherapy being used to treat malignancy; OR
- The condition must be associated with oral chemotherapy being used to treat malignancy.

ondansetron 8 mg orally disintegrating tablet, 10

5473C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	1	..	26.41	27.70	^a APO-Ondansetron ODT [TX] ^a Ondansetron Mylan ODT [AF] ^a Ondansetron ODT GH [GQ] ^a Zilfajim ODT 8 [DO]	^a Ondansetron AN ODT [EA] ^a Ondansetron ODT-DRLA [RZ] ^a Ondansetron SZ ODT [HX] ^a Zotren ODT [RF]

ondansetron 8 mg wafer, 10

8413T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	^B 2.45	28.86	27.70	^a Zofran Zydys [AS]

■ ONDANSETRON

Note Pharmaceutical benefits that have the form ondansetron tablet (orally disintegrating) 4 mg and pharmaceutical benefits that have the form ondansetron 4 mg wafer are equivalent for the purposes of substitution.

Note Pharmaceutical benefits that have the form ondansetron tablet (orally disintegrating) 8 mg and pharmaceutical benefits that have the form ondansetron 8 mg wafer are equivalent for the purposes of substitution.

Restricted benefit

Nausea and vomiting

Clinical criteria:

- The condition must be associated with cytotoxic chemotherapy being used to treat malignancy which occurs within 48 hours of chemotherapy administration.
- Increased maximum quantities will be limited to a maximum of 7 days per chemotherapy cycle.

ondansetron 4 mg orally disintegrating tablet, 4

5470X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	15.84	17.13	^a APO-Ondansetron ODT [TX] ^a Ondansetron Mylan ODT [AF] ^a Ondansetron ODT GH [GQ] ^a Zotren ODT [RF]	^a Ondansetron AN ODT [EA] ^a Ondansetron ODT-DRLA [RZ] ^a Ondansetron SZ ODT [HX]

ondansetron 8 mg wafer, 4

8411Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	^B 2.44	20.22	19.07	^a Zofran Zydis [AS]

ondansetron 8 mg orally disintegrating tablet, 4

5471Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	17.78	19.07	^a APO-Ondansetron ODT [TX] ^a Ondansetron Mylan ODT [AF] ^a Ondansetron ODT GH [GQ] ^a Zotren ODT [RF]	^a Ondansetron AN ODT [EA] ^a Ondansetron ODT-DRLA [RZ] ^a Ondansetron SZ ODT [HX]

ondansetron 4 mg wafer, 4

8410P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	^B 2.30	18.14	17.13	^a Zofran Zydis [AS]

■ PALONOSETRON

Note No increase in the maximum quantity or number of units may be authorised.

Note This drug is not PBS-subsidised for administration with oral 5-HT₃ antagonists.

Restricted benefit

Nausea and vomiting

Clinical criteria:

- The condition must be associated with cytotoxic chemotherapy being used to treat malignancy which occurs within 48 hours of chemotherapy administration.

palonosetron 250 microgram/5 mL injection, 5 mL vial

5295Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	43.61	41.30	Aloxi [MF]

■ TROPISETRON**Restricted benefit**

Nausea and vomiting

Clinical criteria:

- The condition must be associated with cytotoxic chemotherapy being used to treat malignancy which occurs within 48 hours of chemotherapy administration.

Increased maximum quantities will be limited to a maximum of 7 days per chemotherapy cycle.

tropisetron 5 mg/5 mL injection, 5 mL ampoule

2746M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	17.00	18.29	Tropisetron-AFT [AE]

Other antiemetics**■ APREPITANT**

Note Aprepitant is not PBS-subsidised for nausea and vomiting associated with radiotherapy being used to treat malignancy.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

4211

Nausea and vomiting

Clinical criteria:

- The condition must be associated with cytotoxic chemotherapy being used to treat malignancy, **AND**
- The treatment must be in combination with a 5-hydroxytryptamine receptor (5HT₃) antagonist and dexamethasone, **AND**
- Patient must be scheduled to be administered a chemotherapy regimen that includes any 1 of the following agents: altretamine; carmustine; cisplatin when a single dose constitutes a cycle of chemotherapy; cyclophosphamide at a dose of 1500 mg per square metre per day or greater; dacarbazine; procarbazine when a single dose constitutes a cycle of chemotherapy; streptozocin.

No more than 1 capsule of aprepitant 165 mg will be authorised per cycle of cytotoxic chemotherapy.

Authority required (STREAMLINED)

4215

Nausea and vomiting

Clinical criteria:

- The condition must be associated with cytotoxic chemotherapy being used to treat breast cancer, **AND**
- The treatment must be in combination with a 5-hydroxytryptamine receptor (5HT₃) antagonist and dexamethasone, **AND**
- Patient must be scheduled to be co-administered cyclophosphamide and an anthracycline.

No more than 1 capsule of aprepitant 165 mg will be authorised per cycle of cytotoxic chemotherapy.

Authority required (STREAMLINED)

6444

Nausea and vomiting

Clinical criteria:

- The condition must be associated with moderately emetogenic cytotoxic chemotherapy being used to treat malignancy, **AND**
- The treatment must be in combination with a 5-hydroxytryptamine receptor (5HT3) antagonist and dexamethasone on day 1 of a chemotherapy cycle, **AND**
- Patient must have had a prior episode of chemotherapy induced nausea or vomiting, **AND**
- Patient must be scheduled to be administered a chemotherapy regimen that includes any 1 of the following intravenous chemotherapy agents: arsenic trioxide; azacitidine; cyclophosphamide at a dose of less than 1500 mg per square metre per day; cytarabine at a dose of greater than 1 g per square metre per day; dactinomycin; daunorubicin; doxorubicin; epirubicin; fotemustine; idarubicin; ifosfamide; irinotecan; melphalan; methotrexate at a dose of 250 mg to 1 g per square metre; raltitrexed.

No more than 1 capsule of aprepitant 165 mg will be authorised per cycle of cytotoxic chemotherapy.

Concomitant use of a 5HT3 antagonist should not occur with aprepitant on days 2 and 3 of any chemotherapy cycle.

Authority required (STREAMLINED)**6370**

Nausea and vomiting

Clinical criteria:

- The condition must be associated with cytotoxic chemotherapy being used to treat malignancy, **AND**
- The treatment must be in combination with a 5-hydroxytryptamine receptor (5HT3) antagonist and dexamethasone on day 1 of a chemotherapy cycle, **AND**
- Patient must be scheduled to be administered a chemotherapy regimen that includes either carboplatin or oxaliplatin.

No more than 1 capsule of aprepitant 165 mg will be authorised per cycle of cytotoxic chemotherapy.

Concomitant use of a 5HT3 antagonist should not occur with aprepitant on days 2 and 3 of any chemotherapy cycle.

aprepitant 165 mg capsule, 1

2518M

NP

Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
1	5	..	79.00	41.30	Aprepitant APOTEX [TX]

■ FOSAPREPIANT

Note This medicine is not PBS-subsidised for nausea and vomiting associated with radiotherapy being used to treat malignancy.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)**6886**

Nausea and vomiting

Clinical criteria:

- The condition must be associated with cytotoxic chemotherapy being used to treat malignancy, **AND**
- The treatment must be in combination with a 5-hydroxytryptamine receptor (5HT3) antagonist and dexamethasone, **AND**
- Patient must be scheduled to be administered a chemotherapy regimen that includes any 1 of the following agents: altretamine; carmustine; cisplatin when a single dose constitutes a cycle of chemotherapy; cyclophosphamide at a dose of 1500 mg per square metre per day or greater; dacarbazine; procarbazine when a single dose constitutes a cycle of chemotherapy; streptozocin.

No more than 1 vial of fosaprepitant 150 mg injection will be authorised per cycle of cytotoxic chemotherapy.

Authority required (STREAMLINED)**6891**

Nausea and vomiting

Clinical criteria:

- The condition must be associated with cytotoxic chemotherapy being used to treat breast cancer, **AND**
- The treatment must be in combination with a 5-hydroxytryptamine receptor (5HT3) antagonist and dexamethasone, **AND**
- Patient must be scheduled to be co-administered cyclophosphamide and an anthracycline.

No more than 1 vial of fosaprepitant 150 mg injection will be authorised per cycle of cytotoxic chemotherapy.

Authority required (STREAMLINED)**6887**

Nausea and vomiting

Clinical criteria:

- The condition must be associated with moderately emetogenic cytotoxic chemotherapy being used to treat malignancy, **AND**
- The treatment must be in combination with a 5-hydroxytryptamine receptor (5HT3) antagonist and dexamethasone on day 1 of a chemotherapy cycle, **AND**
- Patient must have had a prior episode of chemotherapy induced nausea or vomiting, **AND**
- Patient must be scheduled to be administered a chemotherapy regimen that includes any 1 of the following intravenous chemotherapy agents: arsenic trioxide; azacitidine; cyclophosphamide at a dose of less than 1500 mg per square metre per day; cytarabine at a dose of greater than 1 g per square metre per day; dactinomycin; daunorubicin; doxorubicin; epirubicin; fotemustine; idarubicin; ifosfamide; irinotecan; melphalan; methotrexate at a dose of 250 mg to 1 g per square metre; raltitrexed.

No more than 1 vial of fosaprepitant 150 mg injection will be authorised per cycle of cytotoxic chemotherapy.

Concomitant use of a 5HT3 antagonist should not occur with fosaprepitant on days 2 and 3 of any chemotherapy cycle.

Authority required (STREAMLINED)**6852**

Nausea and vomiting

Clinical criteria:

- The condition must be associated with cytotoxic chemotherapy being used to treat malignancy, **AND**
- The treatment must be in combination with a 5-hydroxytryptamine receptor (5HT3) antagonist and dexamethasone on day 1 of a chemotherapy cycle, **AND**
- Patient must be scheduled to be administered a chemotherapy regimen that includes either carboplatin or oxaliplatin. No more than 1 vial of fosaprepitant 150 mg injection will be authorised per cycle of cytotoxic chemotherapy. Concomitant use of a 5HT3 antagonist should not occur with fosaprepitant on days 2 and 3 of any chemotherapy cycle.

fosaprepitant 150 mg injection, 1 vial

11107N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	116.71	41.30	Emend IV [MK]

PROCHLORPERAZINE

Caution Prochlorperazine may be associated with parkinsonism and tardive dyskinesia and should be used for short-term treatment only.

prochlorperazine mesilate 12.5 mg/mL injection, 10 x 1 mL ampoules

5206B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	1	..	..	21.70	22.99	Stemetil [SW]

prochlorperazine maleate 5 mg tablet, 25

5205Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
DP	1	..	..	13.33	14.62	^a APO-Prochlorperazine [TX] ^a Prochlorperazine AN [EA] ^a Stemetil [AV]	^a ProCalm [RW] ^a Prochlorperazine GH [GQ]
			^b 2.79	16.12	14.62	^a Stemetil [SW]	

PROCHLORPERAZINE

Caution Prochlorperazine may be associated with parkinsonism and tardive dyskinesia and should be used for short-term treatment only.

Note As prochlorperazine may be associated with parkinsonism and tardive dyskinesia it should be used for short-term treatment only. However, authorities for increased maximum quantities and/or repeats of prochlorperazine tablets will be granted for the treatment of emesis associated with malignant disease.

prochlorperazine mesilate 12.5 mg/mL injection, 10 x 1 mL ampoules

2369Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	21.70	22.99	Stemetil [SW]

prochlorperazine maleate 5 mg tablet, 25

2893G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	13.33	14.62	^a APO-Prochlorperazine [TX] ^a Prochlorperazine AN [EA] ^a Stemetil [AV]	^a ProCalm [RW] ^a Prochlorperazine GH [GQ]
			^b 2.79	16.12	14.62	^a Stemetil [SW]	

PROMETHAZINE**promethazine hydrochloride 50 mg/2 mL injection, 5 x 2 mL ampoules**

3374N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	2	..	..	*35.60	36.89	DBL Promethazine Hydrochloride [PF]

BILE AND LIVER THERAPY**BILE THERAPY***Bile acids and derivatives***URSODEOXYCHOLIC ACID**

Note Not for use in the treatment of sclerosing cholangitis or cholelithiasis.

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**9032**

Primary biliary cholangitis (previously known as Primary biliary cirrhosis)

ursodeoxycholic acid 250 mg capsule, 100

8448P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	2	2	..	*226.32	41.30	^a APO-Ursodeoxycholic acid [TX]	^a Ursodox GH [GQ]
						^a Ursofalk [FD]	^a Ursosan [BZ]

ursodeoxycholic acid 500 mg tablet, 100

11180K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	294.91	41.30	Ursofalk [FD]

DRUGS FOR CONSTIPATION

DRUGS FOR CONSTIPATION

Contact laxatives

■ BISACODYL

Restricted benefit

Constipation

Clinical criteria:

- Patient must be paraplegic or quadriplegic or have severe neurogenic impairment of bowel function.

Restricted benefit

Constipation

Clinical criteria:

- Patient must be receiving long-term nursing care on account of age, infirmity or other condition in a hospital, nursing home or residential facility.

Restricted benefit

Constipation

Clinical criteria:

- Patient must be receiving long-term nursing care and in respect of whom a Carer Allowance is payable as a disabled adult.

Restricted benefit

Constipation

Clinical criteria:

- Patient must be receiving palliative care.

Restricted benefit

Terminal malignant neoplasia

Restricted benefit

Anorectal congenital abnormalities

Restricted benefit

Megacolon

bisacodyl 5 mg enteric tablet, 200

1259G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	18.01	19.30	Lax-Tab [AE]

■ BISACODYL

Restricted benefit

Constipation

Clinical criteria:

- Patient must be paraplegic or quadriplegic or have severe neurogenic impairment of bowel function.

Restricted benefit

Constipation

Clinical criteria:

- Patient must be receiving palliative care.

Restricted benefit

Constipation

Clinical criteria:

- Patient must be receiving long-term nursing care on account of age, infirmity or other condition in a hospital, nursing home or residential facility.

Population criteria:

- Patient must identify as Aboriginal or Torres Strait Islander.

Restricted benefit

Constipation

Clinical criteria:

- Patient must be receiving long-term nursing care and in respect of whom a Carer Allowance is payable as a disabled adult.

Population criteria:

- Patient must identify as Aboriginal or Torres Strait Islander.

Restricted benefit

Terminal malignant neoplasia

Population criteria:

- Patient must identify as Aboriginal or Torres Strait Islander.

Restricted benefit

Anorectal congenital abnormalities

Population criteria:

- Patient must identify as Aboriginal or Torres Strait Islander.

Restricted benefit

Megacolon

Population criteria:

- Patient must identify as Aboriginal or Torres Strait Islander.

bisacodyl 10 mg suppository, 12

1258F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	3	4	..	*22.38	23.67	Petrus Bisacodyl Suppositories [PP]

bisacodyl 10 mg suppository, 10

1260H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	3	5	..	*24.66	25.95	^a Petrus Bisacodyl Suppositories [PP]
			^B 1.29	*25.95	25.95	^a Dulcolax [VZ]

Bulk-forming laxatives**■ RHAMNUS FRANGULA + STERCULIA****Restricted benefit**

Constipation

Clinical criteria:

- Patient must be paraplegic or quadriplegic or have severe neurogenic impairment of bowel function.

Restricted benefit

Constipation

Clinical criteria:

- Patient must be receiving long-term nursing care on account of age, infirmity or other condition in a hospital, nursing home or residential facility.

Restricted benefit

Constipation

Clinical criteria:

- Patient must be receiving long-term nursing care and in respect of whom a Carer Allowance is payable as a disabled adult.

Restricted benefit

Constipation

Clinical criteria:

- Patient must be receiving palliative care.

Restricted benefit

Terminal malignant neoplasia

Restricted benefit

Anorectal congenital abnormalities

Restricted benefit

Megacolon

rhhamnus frangula 80 mg/g + sterculia 620 mg/g granules, 500 g

1104D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	1	..	26.12	27.41	Normacol Plus [NE]

Osmotically acting laxatives**■ MACROGOL-3350****Restricted benefit**

Constipation

Clinical criteria:

- Patient must have malignant neoplasia.

Restricted benefit

Constipation

Clinical criteria:

- Patient must be paraplegic, quadriplegic or have severe neurogenic impairment of bowel function, **AND**
- The condition must be unresponsive to other oral therapies.

Restricted benefit

Constipation

Clinical criteria:

- Patient must be receiving palliative care.

Restricted benefit

Chronic constipation

Clinical criteria:

- The condition must be inadequately controlled with first line interventions such as bulk-forming agents.

Restricted benefit

Faecal impaction

Clinical criteria:

- The condition must be inadequately controlled with first line interventions such as bulk-forming agents.

macrogol-3350 1 g/g powder for oral liquid, 510 g

3416T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	19.59	20.88	OsmoLax [KY]

▪ MACROGOL-3350 + SODIUM CHLORIDE + BICARBONATE + POTASSIUM CHLORIDE**Restricted benefit**

Constipation

Clinical criteria:

- Patient must have malignant neoplasia.

Restricted benefit

Constipation

Clinical criteria:

- Patient must be paraplegic, quadriplegic or have severe neurogenic impairment of bowel function, **AND**
- The condition must be unresponsive to other oral therapies.

Restricted benefit

Constipation

Clinical criteria:

- Patient must be receiving palliative care.

Restricted benefit

Chronic constipation

Clinical criteria:

- The condition must be inadequately controlled with first line interventions such as bulk-forming agents.

Restricted benefit

Faecal impaction

Clinical criteria:

- The condition must be inadequately controlled with first line interventions such as bulk-forming agents.

macrogol-3350 13.125 g/25 mL + sodium chloride 350.7 mg/25 mL + sodium bicarbonate 178.5 mg/25 mL + potassium chloride 46.6 mg/25 mL oral liquid, 500 mL

10126Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*22.10	23.39	Movicol Liquid [NE]

macrogol-3350 13.125 g + sodium chloride 350.7 mg + sodium bicarbonate 178.5 mg + potassium chloride 46.6 mg powder for oral liquid, 30 sachets

8612G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	±1	5	..	19.59	20.88	^a APOHEALTH Macrogol with Electrolytes [GX] ^a Chemists' Own Macrogol with Electrolytes [RW] ^a lax-sachets [AE] ^a Molaxole [GO]	^a APO-MACROGOL plus ELECTROLYTES [TX] ^a LaxaCon [EA] ^a Macrovic [RF] ^a Movicol [NE]

Enemas**▪ BISACODYL****Restricted benefit**

Constipation

Clinical criteria:

- Patient must be paraplegic or quadriplegic or have severe neurogenic impairment of bowel function.

Restricted benefit

Constipation

Clinical criteria:

- Patient must be receiving long-term nursing care on account of age, infirmity or other condition in a hospital, nursing home or residential facility.

Restricted benefit

Constipation

Clinical criteria:

- Patient must be receiving long-term nursing care and in respect of whom a Carer Allowance is payable as a disabled adult.

Restricted benefit

Constipation

Clinical criteria:

- Patient must be receiving palliative care.

Restricted benefit

Terminal malignant neoplasia

Restricted benefit

Anorectal congenital abnormalities

Restricted benefit

Megacolon

bisacodyl 10 mg/5 mL enema, 25 x 5 mL

1263L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	39.43	40.72	Bisalax [AS]

■ CITRIC ACID + LAURYL SULFOACETATE SODIUM + SORBITOL**Restricted benefit**

Constipation

Clinical criteria:

- Patient must be paraplegic or quadriplegic or have severe neurogenic impairment of bowel function.

Restricted benefit

Constipation

Clinical criteria:

- Patient must be receiving long-term nursing care on account of age, infirmity or other condition in a hospital, nursing home or residential facility.

Restricted benefit

Constipation

Clinical criteria:

- Patient must be receiving long-term nursing care and in respect of whom a Carer Allowance is payable as a disabled adult.

Restricted benefit

Constipation

Clinical criteria:

- Patient must be receiving palliative care.

Restricted benefit

Terminal malignant neoplasia

Restricted benefit

Anorectal congenital abnormalities

Restricted benefit

Megacolon

sodium citrate dihydrate 450 mg/5 mL + lauryl sulfoacetate sodium 45 mg/5 mL + sorbitol 3.125 g/5 mL enema, 12 x 5 mL

2091C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	2	..	*30.28	31.57	Micolette [AE]

■ ANTIDIARRHEALS, INTESTINAL ANTIINFLAMMATORY/ANTIINFECTIVE AGENTS**INTESTINAL ANTIINFECTIVES***Antibiotics***■ NYSTATIN****nystatin 500 000 units capsule, 50**

1699K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	21.19	22.48	Nilstat [AS]

nystatin 500 000 units capsule, 50

3345C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	1	..	..	21.19	22.48	Nilstat [AS]

nystatin 500 000 units tablet, 50

1696G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	21.19	22.48	Nilstat [AS]

nystatin 500 000 units tablet, 50

3342X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	1	..	..	21.19	22.48	Nilstat [AS]

■ RIFAXIMIN

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Prevention of hepatic encephalopathy

Treatment criteria:

- Must be treated by a gastroenterologist or hepatologist or in consultation with a gastroenterologist or hepatologist.

Clinical criteria:

- The treatment must be in combination with lactulose, if lactulose is tolerated, **AND**
- Patient must have had prior episodes of hepatic encephalopathy.

rifaximin 550 mg tablet, 56

10001J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	475.40	41.30	Xifaxan [NE]

■ VANCOMYCIN

Note Metronidazole has similar efficacy to vancomycin but may have less selective pressure to vancomycin resistant enterococci and is therefore the preferred treatment.

Authority required

Antibiotic associated pseudomembranous colitis

Clinical criteria:

- The condition must be due to **Clostridium difficile**, **AND**
- The condition must be unresponsive to metronidazole.

Authority required

Antibiotic associated pseudomembranous colitis

Clinical criteria:

- The condition must be due to **Clostridium difficile**, **AND**
- Patient must have an intolerance to metronidazole.

vancomycin 125 mg capsule, 20

3113W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*225.26	41.30	Vancocin [AS]

vancomycin 250 mg capsule, 20

3114X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*443.42	41.30	Vancocin [AS]

ELECTROLYTES WITH CARBOHYDRATES*Oral rehydration salt formulations***■ SODIUM CHLORIDE + POTASSIUM CHLORIDE + GLUCOSE MONOHYDRATE + CITRIC ACID****Restricted benefit**

For treatment of a patient identifying as Aboriginal or Torres Strait Islander

sodium chloride 470 mg + potassium chloride 300 mg (potassium 4 mmol) + glucose monohydrate 3.56 g + sodium acid citrate 530 mg powder for oral liquid, 10 x 4.9 g sachets

3196F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	‡1	..	..	16.90	18.19	^a O.R.S. [AF]	^a restore O.R.S. [EA]

■ SODIUM CHLORIDE + POTASSIUM CHLORIDE + GLUCOSE MONOHYDRATE + CITRIC ACID**Authority required**

Rehydration in intestinal failure

sodium chloride 470 mg + potassium chloride 300 mg (potassium 4 mmol) + glucose monohydrate 3.56 g + sodium acid citrate 530 mg powder for oral liquid, 10 x 4.9 g sachets

11049M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	30	..	..	*158.64	41.30	restore O.R.S. [EA]

ANTIPROPULSIVES*Antipropulsives*

■ DIPHENOXYLATE + ATROPINE SULFATE

diphenoxylate hydrochloride 2.5 mg + atropine sulfate monohydrate 25 microgram tablet, 20

2501P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	14.09	15.38	^a Lofenoxal [IL]
			^B 1.40	15.49	15.38	^a Lomotil [IM]

■ LOPERAMIDE

Authority required (STREAMLINED)

6364

Diarrhoea

Population criteria:

- Patient must identify as Aboriginal or Torres Strait Islander.

loperamide hydrochloride 2 mg capsule, 12

1571Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	13.51	14.80	^a Gastrex [CR]	^a Gastro-Stop [AS]

■ LOPERAMIDE

Authority required

Diarrhoea

loperamide hydrochloride 2 mg capsule, 12

10889D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	5	..	..	*19.49	20.78	^a Gastrex [CR]	^a Gastro-Stop [AS]

INTESTINAL ANTIINFLAMMATORY AGENTS

Corticosteroids acting locally

■ BUDESONIDE

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

budesonide 2 mg/application enema, 2 x 14 applications

10034D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	3	..	166.42	41.30	Budenofalk [FD]

■ HYDROCORTISONE ACETATE

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Proctitis

Restricted benefit

Ulcerative colitis

hydrocortisone acetate 10% enema, 21.1 g

1502C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	3	..	*41.60	41.30	Colifoam [GO]

■ PREDNISOLONE SODIUM PHOSPHATE

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

prednisolone (as sodium phosphate) 20 mg/100 mL enema, 7 x 100 mL

1920C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	3	..	*203.30	41.30	Predsol [AS]

■ PREDNISOLONE SODIUM PHOSPHATE

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Proctitis

Restricted benefit

Ulcerative colitis

prednisolone (as sodium phosphate) 5 mg suppository, 10

2554K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	3	3	..	*42.81	41.30	Predsol [AS]

Aminosalicylic acid and similar agents**■ BALSALAZIDE****Note** Not for the treatment of Crohn disease**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**7621**

Ulcerative colitis

Clinical criteria:

- Patient must have had a documented hypersensitivity reaction to a sulphonamide; OR
- Patient must be intolerant to sulfasalazine.

balsalazide sodium 750 mg capsule, 280

11351K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	149.84	41.30	Colazide [PK]

■ MESALAZINE**Note** Not for the treatment of Crohn disease**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Ulcerative colitis

mesalazine 1 g modified release granules, 100 sachets

8599N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	133.25	41.30	Salofalk [FD]

mesalazine 1.2 g modified release tablet, 60

9353G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*195.22	41.30	Mezavant [TK]

mesalazine 3 g modified release granules, 30 sachets

10257W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	116.71	41.30	Salofalk [FD]

mesalazine 4 g modified release granules, 30 sachets

10254Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	150.66	41.30	Pentasa [FP]

mesalazine 500 mg modified release granules, 100 sachets

8598M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*140.86	41.30	Salofalk [FD]

mesalazine 1.5 g modified release granules, 60 sachets

9206M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	116.71	41.30	Salofalk [FD]

mesalazine 800 mg enteric tablet, 90

11210B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*179.40	41.30	Asacol [EU]

■ MESALAZINE**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Ulcerative colitis

Restricted benefit

Crohn disease

mesalazine 1 g modified release tablet, 60

3413P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*158.52	41.30	Pentasa [FP]

mesalazine 1 g modified release granules, 100 sachets

12203G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	133.25	41.30	Pentasa [FP]

mesalazine 500 mg modified release tablet, 100

2214M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*140.86	41.30	Pentasa [FP]

mesalazine 2 g modified release granules, 60 sachets

2287J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	150.66	41.30	Pentasa [FP]

mesalazine 250 mg enteric tablet, 100

1611T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	48.29	41.30	Mesasal [GO]

mesalazine 500 mg enteric tablet, 100

8731M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*140.86	41.30	Salofalk [FD]

mesalazine 1 g enteric tablet, 60

11554D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*150.66	41.30	Salofalk [FD]

■ MESALAZINE**Note** Not for the treatment of Crohn disease**Note** No increase in the maximum quantity or number of units may be authorised.**Note** No increase in the maximum number of repeats may be authorised.**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Acute episode of mild to moderate ulcerative proctitis

mesalazine 1 g suppository, 28

12198B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	101.59	41.30	Pentasa [FP]

mesalazine 1 g suppository, 30

5461K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	107.99	41.30	Salofalk [FD]

■ MESALAZINE**Note** Not for the treatment of Crohn disease**Note** No increase in the maximum quantity or number of units may be authorised.**Note** No increase in the maximum number of repeats may be authorised.**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**4888**

Acute episode of mild to moderate ulcerative colitis

mesalazine 1 g/100 mL enema, 7 x 100 mL

8753Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	1	..	*272.98	41.30	Pentasa [FP]

mesalazine 1 g/application enema, 14 applications

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8768L	4	1	..	*272.98	41.30	Salofalk [FD]

mesalazine 2 g/60 mL enema, 7 x 60 mL

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8616L	4	1	..	*272.98	41.30	Salofalk [FD]

mesalazine 4 g/60 mL enema, 7 x 60 mL

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8617M	4	1	..	*366.50	41.30	Salofalk [FD]

■ OLSALAZINE

Note Not for the treatment of Crohn disease

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**4824**

Ulcerative colitis

Clinical criteria:

- Patient must have had a documented hypersensitivity reaction to a sulphonamide; OR
- Patient must be intolerant to sulfasalazine.

olsalazine sodium 500 mg tablet, 100

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8086N	1	5	..	83.55	41.30	Dipentum [IX]

olsalazine sodium 250 mg capsule, 100

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
1728Y	1	5	..	52.63	41.30	Dipentum [IX]

■ SULFASALAZINE**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

sulfasalazine 500 mg enteric tablet, 100

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
2096H	2	5	..	*55.36	41.30	^a Pyralin EN [FZ]
			^B 4.00	*59.36	41.30	^a Salazopyrin-EN [PF]

sulfasalazine 500 mg tablet, 100

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
2093E	2	5	..	*51.38	41.30	Salazopyrin [PF]

■ SULFASALAZINE

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Restricted benefit

For use in patients who are receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.

sulfasalazine 500 mg enteric tablet, 100

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
9209Q	2	11	..	*55.36	41.30	^a Pyralin EN [FZ]
			^B 4.00	*59.36	41.30	^a Salazopyrin-EN [PF]

sulfasalazine 500 mg tablet, 100

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
9208P	2	11	..	*51.38	41.30	Salazopyrin [PF]

■ DIGESTIVES, INCL. ENZYMES**DIGESTIVES, INCL. ENZYMES***Enzyme preparations*

■ PANCREATIC EXTRACT

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

pancreatic extract 5000 units/100 mg enteric coated granules, 20 g

5453B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	3	10	..	*111.99	41.30	Creon Micro [GO]

pancreatic extract 25 000 units modified release capsule, 100

8021E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	10	..	*116.60	41.30	Creon 25,000 [GO]

pancreatic extract 40 000 units modified release capsule, 100

9412J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	10	..	*182.30	41.30	Creon 40,000 [GO]

pancreatic extract 10 000 units modified release capsule, 100

8020D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	5	10	..	*144.49	41.30	Creon 10,000 [GO]

■ PANCREATIC EXTRACT

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Restricted benefit

Cystic fibrosis

Clinical criteria:

- Patient must be receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.

pancreatic extract 5000 units/100 mg enteric coated granules, 20 g

5454C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	21	..	*111.99	41.30	Creon Micro [GO]

pancreatic extract 25 000 units modified release capsule, 100

9227P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	21	..	*116.60	41.30	Creon 25,000 [GO]

pancreatic extract 40 000 units modified release capsule, 100

9413K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	21	..	*182.30	41.30	Creon 40,000 [GO]

pancreatic extract 10 000 units modified release capsule, 100

9226N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	5	21	..	*144.49	41.30	Creon 10,000 [GO]

■ PANCRELIPASE

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

pancrelipase 25 000 units capsule, 100

8366H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	10	..	*109.08	41.30	Panzytrat 25000 [TM]

■ PANCRELIPASE

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Restricted benefit

Cystic fibrosis

Clinical criteria:

- Patient must be receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.

pancrelipase 25 000 units capsule, 100

9229R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	21	..	*109.08	41.30	Panzytrat 25000 [TM]

DRUGS USED IN DIABETES**INSULINS AND ANALOGUES***Insulins and analogues for injection, fast-acting***INSULIN ASPART****insulin aspart 100 units/mL injection, 1 x 10 mL vial**

8571D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	5	2	..	*125.59	41.30	NovoRapid [NO]

insulin aspart 100 units/mL fast acting injection, 5 x 3 mL pen devices

11706D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	5	1	..	*211.49	41.30	Fiasp FlexTouch [NO]

insulin aspart 100 units/mL fast acting injection, 1 x 10 mL vial

11705C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	5	2	..	*125.59	41.30	Fiasp [NO]

insulin aspart 100 units/mL injection, 5 x 3 mL cartridges

8435Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	5	1	..	*211.49	41.30	NovoRapid Penfill 3 mL [NO]

insulin aspart 100 units/mL injection, 5 x 3 mL pen devices

12254Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	5	1	..	*211.49	41.30	NovoRapid FlexPen [NF]

INSULIN GLULISINE**insulin glulisine 100 units/mL injection, 5 x 3 mL pen devices**

12268Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	5	1	..	*222.34	41.30	Apidra SoloStar [SW]

insulin glulisine 100 units/mL injection, 5 x 3 mL cartridges

1921D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	5	1	..	*222.34	41.30	Apidra [AV]

insulin glulisine 100 units/mL injection, 1 x 10 mL vial

9224L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	5	2	..	*131.74	41.30	Apidra [SW]

INSULIN LISPRO**insulin lispro 100 units/mL injection, 5 x 3 mL cartridges**

8212F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	5	1	..	*211.49	41.30	Humalog [LY]

insulin lispro 100 units/mL injection, 1 x 10 mL vial

8084L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	5	2	..	*125.59	41.30	Humalog [LY]

insulin lispro 100 units/mL injection, 5 x 3 mL pen devices

12237C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	5	1	..	*211.49	41.30	Humalog KwikPen [KP]

insulin lispro 200 units/mL injection, 5 x 3 mL pen devices

11645X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	5	1	..	*416.04	41.30	Humalog U200 Kwikpen [LY]

INSULIN NEUTRAL HUMAN**insulin neutral human 100 units/mL injection, 1 x 10 mL vial**

1531N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	5	2	..	*106.09	41.30	Actrapid [NO]	Humulin R [LY]

insulin neutral human 100 units/mL injection, 5 x 3 mL cartridges

1762R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	5	1	..	*177.49	41.30	Actrapid Penfill 3 mL [NO]	Humulin R [LY]

*Insulins and analogues for injection, intermediate-acting***■ INSULIN ISOPHANE HUMAN****insulin isophane human 100 units/mL injection, 1 x 10 mL vial**

1533Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	5	2	..	*106.09	41.30	Humulin NPH [LY]	Protaphane [NO]

insulin isophane human 100 units/mL injection, 5 x 3 mL pen devices

12262J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	5	1	..	*177.49	41.30	Protaphane InnoLet [NI]	

insulin isophane human 100 units/mL injection, 5 x 3 mL cartridges

1761Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	5	1	..	*177.49	41.30	Humulin NPH [LY]	Protaphane Penfill 3 mL [NO]

*Insulins and analogues for injection, intermediate- or long-acting combined with fast-acting***■ INSULIN ASPART + INSULIN ASPART PROTAMINE****insulin aspart 30 units/mL + insulin aspart protamine 70 units/mL injection, 5 x 3 mL pen devices**

12238D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	5	1	..	*211.49	41.30	NovoMix 30 FlexPen [NF]	

insulin aspart 30 units/mL + insulin aspart protamine 70 units/mL injection, 5 x 3 mL cartridges

8609D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	5	1	..	*211.49	41.30	NovoMix 30 Penfill 3 mL [NO]	

■ INSULIN DEGLUDEC + INSULIN ASPART**Note** Special Pricing Arrangements apply.**insulin degludec 70 units/mL + insulin aspart 30 units/mL injection, 5 x 3 mL pen devices**

11417X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	5	1	..	*379.54	41.30	Ryzodeg FlexTouch [NO]	

insulin degludec 70 units/mL + insulin aspart 30 units/mL injection, 5 x 3 mL cartridges

11426J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	5	1	..	*379.54	41.30	Ryzodeg Penfill [NO]	

■ INSULIN ISOPHANE HUMAN + INSULIN NEUTRAL HUMAN**insulin neutral human 30 units/mL + insulin isophane human 70 units/mL injection, 1 x 10 mL vial**

1426C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	5	2	..	*106.09	41.30	Humulin 30/70 [LY]	

insulin neutral human 50 units/mL + insulin isophane human 50 units/mL injection, 5 x 3 mL cartridges

2062M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	5	1	..	*185.84	41.30	Mixtard 50/50 Penfill 3 mL [NO]	

insulin isophane human 70 units/mL + insulin neutral human 30 units/mL injection, 5 x 3 mL pen devices

12255B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	5	1	..	*177.49	41.30	Mixtard 30/70 InnoLet [NI]	

insulin isophane human 70 units/mL + insulin neutral human 30 units/mL injection, 5 x 3 mL cartridges

1763T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	5	1	..	*177.49	41.30	Humulin 30/70 [LY]	Mixtard 30/70 Penfill 3 mL [NO]

■ INSULIN LISPRO + INSULIN LISPRO PROTAMINE**insulin lispro 50 units/mL + insulin lispro protamine 50 units/mL injection, 5 x 3 mL pen devices**

12261H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	5	1	..	*211.49	41.30	Humalog Mix50 KwikPen [KP]	

insulin lispro 50 units/mL + insulin lispro protamine 50 units/mL injection, 5 x 3 mL cartridges

8874C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	5	1	..	*211.49	41.30	Humalog Mix50 [LY]

insulin lispro 25 units/mL + insulin lispro protamine 75 units/mL injection, 5 x 3 mL cartridges

8390N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	5	1	..	*211.49	41.30	Humalog Mix25 [LY]

insulin lispro 25 units/mL + insulin lispro protamine 75 units/mL injection, 5 x 3 mL pen devices

12234X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	5	1	..	*211.49	41.30	Humalog Mix25 KwikPen [KP]

*Insulins and analogues for injection, long-acting***■ INSULIN DETEMIR****Note** Special Pricing Arrangements apply.**Restricted benefit**

Type 1 diabetes

insulin detemir 100 units/mL injection, 5 x 3 mL pen devices

12236B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	5	1	..	*354.79	41.30	Levemir FlexPen [NF]

insulin detemir 100 units/mL injection, 5 x 3 mL cartridges

9040T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	5	1	..	*354.79	41.30	Levemir Penfill [NO]

■ INSULIN GLARGINE**Note** Special Pricing Arrangements apply.**insulin glargine 300 units/mL injection, 5 x 1.5 mL pen devices**

11302W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	5	1	..	*419.39	41.30	Toujeo Solostar [SW]

insulin glargine 300 units/mL injection, 3 x 1.5 mL pen devices

11308E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	5	1	..	*254.49	41.30	Toujeo Solostar [SW]

■ INSULIN GLARGINE**Note Biosimilar prescribing policy** Prescribing of the biosimilar brand, Semglee, is encouraged for treatment naive patients.Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).**Note** Pharmaceutical benefits that have the brand Optisulin SoloStar 100 units/mL injection, 5 x 3 mL and pharmaceutical benefits that have the brand Semglee 100 units/mL injection, 5 x 3 mL are equivalent for the purposes of substitution.**insulin glargine 100 units/mL injection, 5 x 3 mL pen devices**

11815W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	5	1	..	*185.99	41.30	^a Optisulin SoloStar [WA]	^a Semglee [AF]

insulin glargine 100 units/mL injection, 5 x 3 mL cartridges

9039R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	5	1	..	*185.99	41.30	Optisulin [GZ]

BLOOD GLUCOSE LOWERING DRUGS, EXCL. INSULINS*Biguanides***■ METFORMIN****metformin hydrochloride 500 mg tablet, 100**

2430X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	14.93	16.22	^a APO-Metformin 500 [TX] ^a Diaformin [AF] ^a Glucobete 500 [DO] ^a Metformin GH [HQ]	^a APX-Metformin [TY] ^a FORMET 500 [RF] ^a Metformin AN [EA] ^a Metformin Sandoz [SZ]
			^B 4.73	19.66	16.22	^a Diabex [AL]	

metformin hydrochloride 850 mg tablet, 60

1801T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	14.93	16.22	^a APO-Metformin 850 [TX]	^a APX-Metformin [TY]

NP						^a Chem mart Metformin [CH]	^a Diaformin 850 [AF]
						^a FORMET 850 [RF]	^a Glucobete 850 [DO]
						^a Metformin AN [EA]	^a Metformin Sandoz [SZ]
						^a Terry White Chemists Metformin [TW]	
						^a Diabex 850 [AL]	
			^B 4.73	19.66	16.22		

metformin hydrochloride 1 g modified release tablet, 60

3439B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	16.44	17.73	^a APO-Metformin XR 1000 [TX]	^a Blooms the Chemist Metformin XR 1000 [IB]
						^a Diaformin XR 1000 [AF]	^a METEX XR [RF]
						^a METEX XR 1000 [RW]	^a Pharmacor Metformin XR [CR]
						^a Diabex XR 1000 [AL]	
			^B 4.84	21.28	17.73		

metformin hydrochloride 1 g tablet, 90

8607B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	16.79	18.08	^a APO-Metformin 1000 [TX]	^a APX-Metformin [TY]
						^a Diaformin 1000 [AF]	^a Formet 1000 [RW]
						^a Glucobete 1000 [DO]	^a Metformin AN [EA]
						^a Metformin GH [HQ]	^a Metformin Sandoz [SZ]
						^a Sandoz Metformin [HX]	
			^B 4.87	21.66	18.08	^a Diabex 1000 [AL]	

■ METFORMIN

Note Pharmaceutical benefits that have the form metformin hydrochloride 500 mg modified release tablet in a pack size of 120 can be substituted for a pack size of 56 in the case of a shortage.

metformin hydrochloride 500 mg modified release tablet, 56

12104C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2.1	5	..	*22.00	23.29	^a Metformin (Medsurge) [DZ]

metformin hydrochloride 500 mg modified release tablet, 120

9435N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	16.44	17.73	^a APO-Metformin XR 500 [TX]	^a Blooms the Chemist Metformin XR 500 [IB]
						^a Metex XR [RW]	^a Metformin XR 500 APOTEX [GX]
						^a Pharmacor Metformin XR [CR]	
						^a Diabex XR 500 [AL]	
			^B 4.84	21.28	17.73		

Sulfonylureas**■ GLIBENCLAMIDE**

Caution Sulfonylureas may cause hypoglycaemia, particularly in the elderly.

glibenclamide 5 mg tablet, 100

2939Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	16.45	17.74	Daonil [SW]

■ GLICLAZIDE

Caution Sulfonylureas may cause hypoglycaemia, particularly in the elderly.

gliclazide 60 mg modified release tablet, 60

9302N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	19.26	20.55	^a ARDIX GLICLAZIDE 60mg MR [RX]	^a Pharmacor Gliclazide MR [CR]
						^a Diamicon 60mg MR [SE]	
			^B 7.62	26.88	20.55		

gliclazide 80 mg tablet, 100

2449X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	17.88	19.17	^a APO-Gliclazide [TX]	^a Glyade [AF]
						^a Nidem [RW]	

gliclazide 30 mg modified release tablet, 100

8535F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	18.05	19.34	^a APO-Gliclazide MR [TX]	^a Glyade MR [AF]
						^a Pharmacor Gliclazide MR [CR]	

■ GLIMEPIRIDE

Caution Sulfonylureas may cause hypoglycaemia, particularly in the elderly.

glimepiride 2 mg tablet, 30

8451T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	14.15	15.44	^a APO-Glimepiride [TX] ^a Diapride 2 [RW] ^a Glimepiride AN [EA] ^a Glimepiride Sandoz [SZ]	^a Aylide 2 [AF] ^a Dimirel [AV] ^a Glimepiride APOTEX [GX]
			^b 2.03	16.18	15.44	^a Amaryl [SW]	

glimepiride 4 mg tablet, 30

8452W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	15.13	16.42	^a APO-Glimepiride [TX] ^a Diapride 4 [RW] ^a Glimepiride AN [EA] ^a Glimepiride Sandoz [SZ]	^a Aylide 4 [AF] ^a Dimirel [AV] ^a Glimepiride APOTEX [GX]
			^b 2.03	17.16	16.42	^a Amaryl [SW]	

glimepiride 3 mg tablet, 30

8533D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	14.62	15.91	^a APO-Glimepiride [TX] ^a Diapride 3 [RW] ^a Glimepiride AN [EA] ^a Glimepiride Sandoz [SZ]	^a Aylide 3 [AF] ^a Dimirel [AV] ^a Glimepiride APOTEX [GX]
			^b 2.03	16.65	15.91	^a Amaryl [SW]	

glimepiride 1 mg tablet, 30

8450R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	13.33	14.62	^a APO-Glimepiride [TX] ^a Diapride 1 [RW] ^a Glimepiride AN [EA] ^a Glimepiride Sandoz [SZ]	^a Aylide 1 [AF] ^a Dimirel [AV] ^a Glimepiride APOTEX [GX]
			^b 2.08	15.41	14.62	^a Amaryl [SW]	

GLIPIZIDE

Caution Sulfonylureas may cause hypoglycaemia, particularly in the elderly.

glipizide 5 mg tablet, 100

2440K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	23.81	25.10	^a Melizide [AF]	^a Minidiab [PF]

Combinations of oral blood glucose lowering drugs**ALOGLIPTIN + METFORMIN**

Note This fixed dose combination tablet is not PBS-subsidised for use in combination with a sulfonylurea (triple oral therapy), as initial therapy or in combination with a thiazolidinedione (glitazone), a glucagon-like peptide-1 or an SGLT2 inhibitor.

Authority required (STREAMLINED)**4423**

Diabetes mellitus type 2

Clinical criteria:

- Patient must have, or have had, a HbA1c measurement greater than 7% despite treatment with metformin; OR
- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period despite treatment with metformin.

The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

A patient whose diabetes was previously demonstrated unable to be controlled with metformin does not need to requalify on this criterion before being eligible for PBS-subsidised treatment with this fixed dose combination.

Authority required (STREAMLINED)**4427**

Diabetes mellitus type 2

Treatment Phase: Continuing

Clinical criteria:

- Patient must have previously received and been stabilised on a PBS-subsidised regimen of oral diabetic medicines which includes metformin and alogliptin.

alogliptin 12.5 mg + metformin hydrochloride 500 mg tablet, 56

10033C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	58.81	41.30	Nesina Met 12.5/500 [TK]

alogliptin 12.5 mg + metformin hydrochloride 1 g tablet, 56

10035E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	60.22	41.30	Nesina Met 12.5/1000 [TK]

alogliptin 12.5 mg + metformin hydrochloride 850 mg tablet, 56

10032B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	59.81	41.30	Nesina Met 12.5/850 [TK]

■ DAPAGLIFLOZIN + METFORMIN

Note This fixed dose combination is not PBS-subsidised for initiating dual oral combination treatment or in combination with a thiazolidinedione (glitazone), a glucagon-like peptide-1 analogue, or another SGLT2 inhibitor.

Note PBS-subsidised dual oral therapy does not include combination use of: a gliptin with an SGLT2 inhibitor; or

- a gliptin with a glitazone; or
- an SGLT2 inhibitor with a glitazone.

Authority required (STREAMLINED)**7498**

Diabetes mellitus type 2

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be in combination with a dipeptidyl peptidase 4 inhibitor (gliptin), **AND**
- Patient must have an HbA1c measurement greater than 7% despite treatment with a PBS-subsidised regimen of oral diabetic medicines which includes metformin and a gliptin for this condition; OR
- Patient must have, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation of triple oral therapy with a sodium-glucose co-transporter 2 (SGLT2) inhibitor, metformin and a gliptin.

The date and level of the qualifying HbA1c measurement must be documented in the patient's medical records at the time triple oral therapy with an SGLT2 inhibitor, metformin and a gliptin is initiated.

The HbA1c must be no more than 4 months old at the time triple oral therapy with an SGLT2 inhibitor, metformin and a gliptin is initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of triple oral therapy with an SGLT2 inhibitor, metformin and a gliptin, must be documented in the patient's medical records.

dapagliflozin 10 mg + metformin hydrochloride 1 g modified release tablet, 28

11313K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	58.35	41.30	Xigduo XR 10/1000 [AP]

dapagliflozin 5 mg + metformin hydrochloride 1 g modified release tablet, 56

11300R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	59.76	41.30	Xigduo XR 5/1000 [AP]

dapagliflozin 10 mg + metformin hydrochloride 500 mg modified release tablet, 28

11270E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	57.60	41.30	Xigduo XR 10/500 [AP]

■ DAPAGLIFLOZIN + METFORMIN**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**5631**

Diabetes mellitus type 2

Clinical criteria:

- Patient must have, or have had, a HbA1c measurement greater than 7% despite treatment with metformin; OR
- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period despite treatment with metformin.

The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

A patient whose diabetes was previously demonstrated unable to be controlled with metformin does not need to requalify on this criterion before being eligible for PBS-subsidised treatment with this fixed dose combination.

Note This fixed dose combination is not PBS-subsidised for use as initial therapy or in combination with a thiazolidinedione (glitazone) or a glucagon-like peptide-1.

Authority required (STREAMLINED)

5739

Diabetes mellitus type 2

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received and been stabilised on a PBS-subsidised regimen of oral diabetic medicines which includes metformin and dapagliflozin.

Note This fixed dose combination is not PBS-subsidised for use as initial therapy or in combination with a thiazolidinedione (glitazone) or a glucagon-like peptide-1.

Authority required (STREAMLINED)

5798

Diabetes mellitus type 2

Clinical criteria:

- The treatment must be in combination with a sulfonylurea, **AND**
- Patient must have, or have had, a HbA1c measurement greater than 7% prior to the initiation of a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor despite treatment with optimal doses of dual oral therapy; OR
- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 despite treatment with optimal doses of dual oral therapy.

The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

A patient whose diabetes was previously demonstrated unable to be controlled with metformin or a sulfonylurea does not need to requalify on this criterion before being eligible for PBS-subsidised treatment with this fixed dose combination.

Note This fixed dose combination is not PBS-subsidised for use as initial therapy or in combination with a thiazolidinedione (glitazone) or a glucagon-like peptide-1.

Note PBS subsidised dual oral therapy does not include concomitant use of a combination of: a gliptin, a glitazone or an SGLT2 inhibitor.

Authority required (STREAMLINED)

5657

Diabetes mellitus type 2

Clinical criteria:

- The treatment must be in combination with insulin, **AND**
- Patient must have, or have had, a HbA1c measurement greater than 7% prior to the initiation of a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor despite treatment with insulin and oral antidiabetic agents, or insulin alone where metformin is contraindicated; OR
- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor despite treatment with insulin and oral antidiabetic agents, or insulin alone where metformin is contraindicated.

The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

Note This fixed dose combination is not PBS-subsidised for use as initial therapy or in combination with a thiazolidinedione (glitazone) or a glucagon-like peptide-1.

Authority required (STREAMLINED)

7492

Diabetes mellitus type 2

Treatment Phase: Continuing treatment

Clinical criteria:

- The treatment must be in combination with a dipeptidyl peptidase 4 inhibitor (gliptin), **AND**
- Patient must have previously received a PBS-subsidised regimen of oral diabetic medicines which included a sodium-glucose co-transporter 2 (SGLT2) inhibitor, metformin and a gliptin for this condition.

Note This fixed dose combination is not PBS-subsidised for initiating dual oral combination treatment or in combination with a thiazolidinedione (glitazone), a glucagon-like peptide-1 analogue, or another SGLT2 inhibitor.

Note PBS-subsidised dual oral therapy does not include combination use of: a gliptin with an SGLT2 inhibitor; or

- a gliptin with a glitazone; or
- an SGLT2 inhibitor with a glitazone.

dapagliflozin 10 mg + metformin hydrochloride 1 g modified release tablet, 28

10515K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	58.35	41.30	Xigduo XR 10/1000 [AP]

dapagliflozin 5 mg + metformin hydrochloride 1 g modified release tablet, 56

10510E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	59.76	41.30	Xigduo XR 5/1000 [AP]

dapagliflozin 10 mg + metformin hydrochloride 500 mg modified release tablet, 28

10516L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	57.60	41.30	Xigduo XR 10/500 [AP]

▪ **EMPAGLIFLOZIN + LINAGLIPTIN**

Note This fixed dose combination is not PBS-subsidised for use as a sole therapy or in combination with a thiazolidinedione (glitazone), a glucagon-like peptide-1 analogue, an insulin, another dipeptidyl peptidase 4 inhibitor (gliptin), or another SGLT2 inhibitor.

Authority required (STREAMLINED)

7524

Diabetes mellitus type 2

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be in combination with metformin, **AND**
- Patient must have an HbA1c measurement greater than 7% despite treatment with dual oral combination therapy with metformin and a dipeptidyl peptidase 4 inhibitor (gliptin) or a sodium-glucose co-transporter 2 (SGLT2) inhibitor; OR
- Patient must have, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation of triple oral therapy with a sodium-glucose co-transporter 2 (SGLT2) inhibitor, metformin and a gliptin.

The date and level of the qualifying HbA1c measurement must be documented in the patient's medical records at the time triple oral therapy with an SGLT2 inhibitor, metformin and a gliptin is initiated.

The HbA1c must be no more than 4 months old at the time triple oral therapy with an SGLT2 inhibitor, metformin and a gliptin is initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of triple oral therapy with an SGLT2 inhibitor, metformin and a gliptin, must be documented in the patient's medical records.

empagliflozin 25 mg + linagliptin 5 mg tablet, 30

11303X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	83.37	41.30	Glyxambi [BY]

empagliflozin 10 mg + linagliptin 5 mg tablet, 30

11269D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	83.37	41.30	Glyxambi [BY]

■ EMPAGLIFLOZIN + LINAGLIPTIN

Note This fixed dose combination is not PBS-subsidised for use as a sole therapy or in combination with a thiazolidinedione (glitazone), a glucagon-like peptide-1 analogue, an insulin, another dipeptidyl peptidase 4 inhibitor (gliptin), or another SGLT2 inhibitor.

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)

7556

Diabetes mellitus type 2

Treatment Phase: Continuing treatment

Clinical criteria:

- The treatment must be in combination with metformin, **AND**
- Patient must have previously received a PBS-subsidised regimen of oral diabetic medicines which included a sodium-glucose co-transporter 2 (SGLT2) inhibitor, metformin and a gliptin for this condition.

empagliflozin 25 mg + linagliptin 5 mg tablet, 30

11298P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	83.37	41.30	Glyxambi [BY]

empagliflozin 10 mg + linagliptin 5 mg tablet, 30

11310G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	83.37	41.30	Glyxambi [BY]

■ EMPAGLIFLOZIN + METFORMIN

Authority required (STREAMLINED)

5953

Diabetes mellitus type 2

Clinical criteria:

- Patient must have, or have had, a HbA1c measurement greater than 7% despite treatment with metformin; OR
- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period despite treatment with metformin.

The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

Note A patient whose diabetes was previously demonstrated unable to be controlled with metformin does not need to requalify on this criterion before being eligible for PBS-subsidised treatment with this fixed dose combination.

Note This fixed dose combination is not PBS-subsidised for use as initial therapy or in combination with a thiazolidinedione (glitazone) or a glucagon-like peptide-1.

Authority required (STREAMLINED)

7498

Diabetes mellitus type 2

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be in combination with a dipeptidyl peptidase 4 inhibitor (gliptin), **AND**
- Patient must have an HbA1c measurement greater than 7% despite treatment with a PBS-subsidised regimen of oral diabetic medicines which includes metformin and a gliptin for this condition; OR
- Patient must have, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation of triple oral therapy with a sodium-glucose co-transporter 2 (SGLT2) inhibitor, metformin and a gliptin.

The date and level of the qualifying HbA1c measurement must be documented in the patient's medical records at the time triple oral therapy with an SGLT2 inhibitor, metformin and a gliptin is initiated.

The HbA1c must be no more than 4 months old at the time triple oral therapy with an SGLT2 inhibitor, metformin and a gliptin is initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of triple oral therapy with an SGLT2 inhibitor, metformin and a gliptin, must be documented in the patient's medical records.

Note This fixed dose combination is not PBS-subsidised for initiating dual oral combination treatment or in combination with a thiazolidinedione (glitazone), a glucagon-like peptide-1 analogue, or another SGLT2 inhibitor.

Note PBS-subsidised dual oral therapy does not include combination use of: a gliptin with an SGLT2 inhibitor; or

- a gliptin with a glitazone; or
- an SGLT2 inhibitor with a glitazone.

empagliflozin 5 mg + metformin hydrochloride 500 mg tablet, 60

10650M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	61.65	41.30	Jardiamet 5 mg/500 mg [BY]

empagliflozin 5 mg + metformin hydrochloride 1 g tablet, 60

10649L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	63.17	41.30	Jardiamet 5 mg/1000 mg [BY]

empagliflozin 12.5 mg + metformin hydrochloride 1 g tablet, 60

10640B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	63.17	41.30	Jardiamet 12.5 mg/1000 mg [BY]

empagliflozin 12.5 mg + metformin hydrochloride 500 mg tablet, 60

10639Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	61.65	41.30	Jardiamet 12.5 mg/500 mg [BY]

■ EMPAGLIFLOZIN + METFORMIN

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)

5966

Diabetes mellitus type 2

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received and been stabilised on a PBS-subsidised regimen of oral diabetic medicines which includes metformin and empagliflozin.

Note This fixed dose combination is not PBS-subsidised for use as initial therapy or in combination with a thiazolidinedione (glitazone) or a glucagon-like peptide-1.

Authority required (STREAMLINED)

5798

Diabetes mellitus type 2

Clinical criteria:

- The treatment must be in combination with a sulfonylurea, **AND**
- Patient must have, or have had, a HbA1c measurement greater than 7% prior to the initiation of a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor despite treatment with optimal doses of dual oral therapy; OR
- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 despite treatment with optimal doses of dual oral therapy.

The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

A patient whose diabetes was previously demonstrated unable to be controlled with metformin or a sulfonylurea does not need to requalify on this criterion before being eligible for PBS-subsidised treatment with this fixed dose combination.

Note This fixed dose combination is not PBS-subsidised for use as initial therapy or in combination with a thiazolidinedione (glitazone) or a glucagon-like peptide-1.

Note PBS subsidised dual oral therapy does not include concomitant use of a combination of: a gliptin, a glitazone or an SGLT2 inhibitor.

Authority required (STREAMLINED)

5657

Diabetes mellitus type 2

Clinical criteria:

- The treatment must be in combination with insulin, **AND**
- Patient must have, or have had, a HbA1c measurement greater than 7% prior to the initiation of a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor despite treatment with insulin and oral antidiabetic agents, or insulin alone where metformin is contraindicated; OR
- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor despite treatment with insulin and oral antidiabetic agents, or insulin alone where metformin is contraindicated.

The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

Note This fixed dose combination is not PBS-subsidised for use as initial therapy or in combination with a thiazolidinedione (glitazone) or a glucagon-like peptide-1.

Authority required (STREAMLINED)**7492**

Diabetes mellitus type 2

Treatment Phase: Continuing treatment

Clinical criteria:

- The treatment must be in combination with a dipeptidyl peptidase 4 inhibitor (gliptin), **AND**
- Patient must have previously received a PBS-subsidised regimen of oral diabetic medicines which included a sodium-glucose co-transporter 2 (SGLT2) inhibitor, metformin and a gliptin for this condition.

Note This fixed dose combination is not PBS-subsidised for initiating dual oral combination treatment or in combination with a thiazolidinedione (glitazone), a glucagon-like peptide-1 analogue, or another SGLT2 inhibitor.

Note PBS-subsidised dual oral therapy does not include combination use of: a gliptin with an SGLT2 inhibitor; or

- a gliptin with a glitazone; or
- an SGLT2 inhibitor with a glitazone.

empagliflozin 5 mg + metformin hydrochloride 500 mg tablet, 60

10626G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	61.65	41.30	Jardiamet 5 mg/500 mg [BY]

empagliflozin 5 mg + metformin hydrochloride 1 g tablet, 60

10627H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	63.17	41.30	Jardiamet 5 mg/1000 mg [BY]

empagliflozin 12.5 mg + metformin hydrochloride 1 g tablet, 60

10677Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	63.17	41.30	Jardiamet 12.5 mg/1000 mg [BY]

empagliflozin 12.5 mg + metformin hydrochloride 500 mg tablet, 60

10633P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	61.65	41.30	Jardiamet 12.5 mg/500 mg [BY]

■ ERTUGLIFLOZIN + METFORMIN

Note This fixed dose combination is not PBS-subsidised for initiating dual oral combination treatment or in combination with a thiazolidinedione (glitazone), insulin, a glucagon-like peptide-1 analogue, another dipeptidyl peptidase 4 inhibitor (gliptin), or another SGLT2 inhibitor.

Note PBS-subsidised dual oral therapy does not include combination use of: a gliptin with an SGLT2 inhibitor; or

- a gliptin with a glitazone; or
- an SGLT2 inhibitor with a glitazone.

Authority required (STREAMLINED)**7498**

Diabetes mellitus type 2

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be in combination with a dipeptidyl peptidase 4 inhibitor (gliptin), **AND**

- Patient must have an HbA1c measurement greater than 7% despite treatment with a PBS-subsidised regimen of oral diabetic medicines which includes metformin and a gliptin for this condition; OR
- Patient must have, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation of triple oral therapy with a sodium-glucose co-transporter 2 (SGLT2) inhibitor, metformin and a gliptin.

The date and level of the qualifying HbA1c measurement must be documented in the patient's medical records at the time triple oral therapy with an SGLT2 inhibitor, metformin and a gliptin is initiated.

The HbA1c must be no more than 4 months old at the time triple oral therapy with an SGLT2 inhibitor, metformin and a gliptin is initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of triple oral therapy with an SGLT2 inhibitor, metformin and a gliptin, must be documented in the patient's medical records.

ertugliflozin 2.5 mg + metformin hydrochloride 1 g tablet, 56

11564P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	62.11	41.30	Segluromet 2.5/1000 [MK]

ertugliflozin 2.5 mg + metformin hydrochloride 500 mg tablet, 56

11584Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	60.71	41.30	Segluromet 2.5/500 [MK]

ertugliflozin 7.5 mg + metformin hydrochloride 1 g tablet, 56

11569X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	62.11	41.30	Segluromet 7.5/1000 [MK]

ertugliflozin 7.5 mg + metformin hydrochloride 500 mg tablet, 56

11562M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	60.71	41.30	Segluromet 7.5/500 [MK]

■ ERTUGLIFLOZIN + METFORMIN

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)

5631

Diabetes mellitus type 2

Clinical criteria:

- Patient must have, or have had, a HbA1c measurement greater than 7% despite treatment with metformin; OR
- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period despite treatment with metformin.

The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

A patient whose diabetes was previously demonstrated unable to be controlled with metformin does not need to requalify on this criterion before being eligible for PBS-subsidised treatment with this fixed dose combination.

Note This fixed dose combination is not PBS-subsidised for use as initial therapy or in combination with a thiazolidinedione (glitazone), insulin or a glucagon-like peptide-1.

Authority required (STREAMLINED)

8249

Diabetes mellitus type 2

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received and been stabilised on a PBS-subsidised regimen of oral diabetic medicines which includes metformin and ertugliflozin.

Note This fixed dose combination is not PBS-subsidised for use as initial therapy or in combination with a thiazolidinedione (glitazone), insulin or a glucagon-like peptide-1.

Authority required (STREAMLINED)

7492

Diabetes mellitus type 2

Treatment Phase: Continuing treatment

Clinical criteria:

- The treatment must be in combination with a dipeptidyl peptidase 4 inhibitor (gliptin), **AND**
- Patient must have previously received a PBS-subsidised regimen of oral diabetic medicines which included a sodium-glucose co-transporter 2 (SGLT2) inhibitor, metformin and a gliptin for this condition.

Note This fixed dose combination is not PBS-subsidised for initiating dual oral combination treatment or in combination with a thiazolidinedione (glitazone), insulin, a glucagon-like peptide-1 analogue, another dipeptidyl peptidase 4 inhibitor (gliptin), or another SGLT2 inhibitor.

Note PBS-subsidised dual oral therapy does not include combination use of: a gliptin with an SGLT2 inhibitor; or

- a gliptin with a glitazone; or
- an SGLT2 inhibitor with a glitazone.

ertugliflozin 2.5 mg + metformin hydrochloride 1 g tablet, 56

11581M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	62.11	41.30	Segluromet 2.5/1000 [MK]

ertugliflozin 2.5 mg + metformin hydrochloride 500 mg tablet, 56

11575F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	60.71	41.30	Segluromet 2.5/500 [MK]

ertugliflozin 7.5 mg + metformin hydrochloride 1 g tablet, 56

11563N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	62.11	41.30	Segluromet 7.5/1000 [MK]

ertugliflozin 7.5 mg + metformin hydrochloride 500 mg tablet, 56

11568W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	60.71	41.30	Segluromet 7.5/500 [MK]

■ ERTUGLIFLOZIN + SITAGLIPTIN

Note This fixed dose combination is not PBS-subsidised for use as a sole therapy or in combination with a thiazolidinedione (glitazone), a glucagon-like peptide-1 analogue, an insulin, another dipeptidyl peptidase 4 inhibitor (gliptin), or another SGLT2 inhibitor.

Authority required (STREAMLINED)**7524**

Diabetes mellitus type 2

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be in combination with metformin, **AND**
- Patient must have an HbA1c measurement greater than 7% despite treatment with dual oral combination therapy with metformin and a dipeptidyl peptidase 4 inhibitor (gliptin) or a sodium-glucose co-transporter 2 (SGLT2) inhibitor; OR
- Patient must have, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation of triple oral therapy with a sodium-glucose co-transporter 2 (SGLT2) inhibitor, metformin and a gliptin.

The date and level of the qualifying HbA1c measurement must be documented in the patient's medical records at the time triple oral therapy with an SGLT2 inhibitor, metformin and a gliptin is initiated.

The HbA1c must be no more than 4 months old at the time triple oral therapy with an SGLT2 inhibitor, metformin and a gliptin is initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of triple oral therapy with an SGLT2 inhibitor, metformin and a gliptin, must be documented in the patient's medical records.

ertugliflozin 5 mg + sitagliptin 100 mg tablet, 28

11561L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	77.02	41.30	Steglujan 5/100 [MK]

ertugliflozin 15 mg + sitagliptin 100 mg tablet, 28

11583P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	77.02	41.30	Steglujan 15/100 [MK]

■ ERTUGLIFLOZIN + SITAGLIPTIN

Note This fixed dose combination is not PBS-subsidised for use as a sole therapy or in combination with a thiazolidinedione (glitazone), a glucagon-like peptide-1 analogue, an insulin, another dipeptidyl peptidase 4 inhibitor (gliptin), or another SGLT2 inhibitor.

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**7556**

Diabetes mellitus type 2

Treatment Phase: Continuing treatment

Clinical criteria:

- The treatment must be in combination with metformin, **AND**
- Patient must have previously received a PBS-subsidised regimen of oral diabetic medicines which included a sodium-glucose co-transporter 2 (SGLT2) inhibitor, metformin and a gliptin for this condition.

ertugliflozin 5 mg + sitagliptin 100 mg tablet, 28

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
11579K	1	5	..	77.02	41.30	Steglujan 5/100 [MK]

ertugliflozin 15 mg + sitagliptin 100 mg tablet, 28

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
11578J	1	5	..	77.02	41.30	Steglujan 15/100 [MK]

■ LINAGLIPTIN + METFORMIN

Note This fixed dose combination is not PBS-subsidised for initiating dual oral combination treatment or in combination with a thiazolidinedione (glitazone), a glucagon-like peptide-1 analogue, or another dipeptidyl peptidase 4 inhibitor (gliptin).

Note PBS-subsidised dual oral therapy does not include combination use of: a gliptin with an SGLT2 inhibitor; or

- a gliptin with a glitazone; or
- an SGLT2 inhibitor with a glitazone.

Authority required (STREAMLINED)**7507**

Diabetes mellitus type 2

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be in combination with a sodium-glucose co-transporter 2 (SGLT2) inhibitor, **AND**
- Patient must have an HbA1c measurement greater than 7% despite treatment with a PBS-subsidised regimen of oral diabetic medicines which includes metformin and an SGLT2 inhibitor for this condition; OR
- Patient must have, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation of triple oral therapy with a sodium-glucose co-transporter 2 (SGLT2) inhibitor, metformin and a gliptin.

The date and level of the qualifying HbA1c measurement must be documented in the patient's medical records at the time triple oral therapy with an SGLT2 inhibitor, metformin and a gliptin is initiated.

The HbA1c must be no more than 4 months old at the time triple oral therapy with an SGLT2 inhibitor, metformin and a gliptin is initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of triple oral therapy with an SGLT2 inhibitor, metformin and a gliptin, must be documented in the patient's medical records.

linagliptin 2.5 mg + metformin hydrochloride 500 mg tablet, 60

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
11274J	1	5	..	62.16	41.30	Trajentamet [BY]

linagliptin 2.5 mg + metformin hydrochloride 850 mg tablet, 60

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
11294K	1	5	..	63.23	41.30	Trajentamet [BY]

linagliptin 2.5 mg + metformin hydrochloride 1 g tablet, 60

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
11282T	1	5	..	63.67	41.30	Trajentamet [BY]

■ LINAGLIPTIN + METFORMIN**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**6333**

Diabetes mellitus type 2

Clinical criteria:

- Patient must have, or have had, a HbA1c measurement greater than 7% despite treatment with metformin; OR
- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period despite treatment with metformin.

The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

A patient whose diabetes was previously demonstrated unable to be controlled with metformin does not need to requalify on this criterion before being eligible for PBS-subsidised treatment with this fixed dose combination.

Note This fixed dose combination is not PBS-subsidised for use as initial therapy or in combination with a thiazolidinedione (glitazone) or a glucagon-like peptide-1.

Authority required (STREAMLINED)

6336

Diabetes mellitus type 2

Treatment Phase: Continuing

Clinical criteria:

- Patient must have previously received and been stabilised on a PBS-subsidised regimen of oral diabetic medicines which includes metformin and linagliptin.

Note This fixed dose combination is not PBS-subsidised for use as initial therapy or in combination with a thiazolidinedione (glitazone) or a glucagon-like peptide-1.

Authority required (STREAMLINED)

6344

Diabetes mellitus type 2

Clinical criteria:

- The treatment must be in combination with a sulfonylurea, **AND**
- Patient must have, or have had, a HbA1c measurement greater than 7% prior to the initiation of a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor despite treatment with optimal doses of dual oral therapy; OR
- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor despite treatment with optimal doses of dual oral therapy.

The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

A patient whose diabetes was previously demonstrated unable to be controlled with metformin or a sulfonylurea does not need to requalify on this criterion before being eligible for PBS-subsidised treatment with this fixed dose combination.

Note This fixed dose combination is not PBS-subsidised for use as initial therapy or in combination with a thiazolidinedione (glitazone) or a glucagon-like peptide-1.

Note PBS subsidised dual oral therapy does not include concomitant use of a combination of: a gliptin, a glitazone or an SGLT2 inhibitor.

Authority required (STREAMLINED)

6443

Diabetes mellitus type 2

Clinical criteria:

- The treatment must be in combination with insulin, **AND**
- Patient must have, or have had, a HbA1c measurement greater than 7% prior to the initiation of a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor despite treatment with insulin and oral antidiabetic agents, or insulin alone where metformin is contraindicated; OR
- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation with a gliptin, a glitazone, a glucagon-like

peptide-1 or an SGLT2 inhibitor despite treatment with insulin and oral antidiabetic agents, or insulin alone where metformin is contraindicated.

The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

Note This fixed dose combination is not PBS-subsidised for use as initial therapy or in combination with a thiazolidinedione (glitazone) or a glucagon-like peptide-1.

Authority required (STREAMLINED)

7530

Diabetes mellitus type 2

Treatment Phase: Continuing treatment

Clinical criteria:

- The treatment must be in combination with a sodium-glucose co-transporter 2 (SGLT2) inhibitor, **AND**
- Patient must have previously received a PBS-subsidised regimen of oral diabetic medicines which included a sodium-glucose co-transporter 2 (SGLT2) inhibitor, metformin and a gliptin for this condition.

Note This fixed dose combination is not PBS-subsidised for initiating dual oral combination treatment or in combination with a thiazolidinedione (glitazone), a glucagon-like peptide-1 analogue, or another dipeptidyl peptidase 4 inhibitor (gliptin).

Note PBS-subsidised dual oral therapy does not include combination use of: a gliptin with an SGLT2 inhibitor; or

- a gliptin with a glitazone; or
- an SGLT2 inhibitor with a glitazone.

linagliptin 2.5 mg + metformin hydrochloride 500 mg tablet, 60

10038H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	62.16	41.30	Trajentamet [BY]

linagliptin 2.5 mg + metformin hydrochloride 850 mg tablet, 60

10045Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	63.23	41.30	Trajentamet [BY]

linagliptin 2.5 mg + metformin hydrochloride 1 g tablet, 60

10044P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	63.67	41.30	Trajentamet [BY]

■ SAXAGLIPTIN + DAPAGLIFLOZIN

Note This fixed dose combination is not PBS-subsidised for use as a sole therapy or in combination with a thiazolidinedione (glitazone), a glucagon-like peptide-1 analogue, an insulin, another dipeptidyl peptidase 4 inhibitor (gliptin), or another SGLT2 inhibitor.

Authority required (STREAMLINED)

7524

Diabetes mellitus type 2

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be in combination with metformin, **AND**
- Patient must have an HbA1c measurement greater than 7% despite treatment with dual oral combination therapy with metformin and a dipeptidyl peptidase 4 inhibitor (gliptin) or a sodium-glucose co-transporter 2 (SGLT2) inhibitor; OR
- Patient must have, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation of triple oral therapy with a sodium-glucose co-transporter 2 (SGLT2) inhibitor, metformin and a gliptin.

The date and level of the qualifying HbA1c measurement must be documented in the patient's medical records at the time triple oral therapy with an SGLT2 inhibitor, metformin and a gliptin is initiated.

The HbA1c must be no more than 4 months old at the time triple oral therapy with an SGLT2 inhibitor, metformin and a gliptin is initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of triple oral therapy with an SGLT2 inhibitor, metformin and a gliptin, must be documented in the patient's medical records.

saxagliptin 5 mg + dapagliflozin 10 mg tablet, 28

11286B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	78.62	41.30	Qtern 5/10 [AP]

■ SAXAGLIPTIN + DAPAGLIFLOZIN

Note This fixed dose combination is not PBS-subsidised for use as a sole therapy or in combination with a thiazolidinedione (glitazone), a glucagon-like peptide-1 analogue, an insulin, another dipeptidyl peptidase 4 inhibitor (gliptin), or another SGLT2 inhibitor.

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**7556**

Diabetes mellitus type 2

Treatment Phase: Continuing treatment

Clinical criteria:

- The treatment must be in combination with metformin, **AND**
- Patient must have previously received a PBS-subsidised regimen of oral diabetic medicines which included a sodium-glucose co-transporter 2 (SGLT2) inhibitor, metformin and a gliptin for this condition.

saxagliptin 5 mg + dapagliflozin 10 mg tablet, 28

11305B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	78.62	41.30	Qtern 5/10 [AP]

NP

■ SAXAGLIPTIN + METFORMIN

Note This fixed dose combination is not PBS-subsidised for initiating dual oral combination treatment or in combination with a thiazolidinedione (glitazone), a glucagon-like peptide-1 analogue, or another dipeptidyl peptidase 4 inhibitor (gliptin).

Note PBS-subsidised dual oral therapy does not include combination use of: a gliptin with an SGLT2 inhibitor; or

- a gliptin with a glitazone; or
- an SGLT2 inhibitor with a glitazone.

Authority required (STREAMLINED)**7507**

Diabetes mellitus type 2

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be in combination with a sodium-glucose co-transporter 2 (SGLT2) inhibitor, **AND**
- Patient must have an HbA1c measurement greater than 7% despite treatment with a PBS-subsidised regimen of oral diabetic medicines which includes metformin and an SGLT2 inhibitor for this condition; OR
- Patient must have, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation of triple oral therapy with a sodium-glucose co-transporter 2 (SGLT2) inhibitor, metformin and a gliptin.

The date and level of the qualifying HbA1c measurement must be documented in the patient's medical records at the time triple oral therapy with an SGLT2 inhibitor, metformin and a gliptin is initiated.

The HbA1c must be no more than 4 months old at the time triple oral therapy with an SGLT2 inhibitor, metformin and a gliptin is initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of triple oral therapy with an SGLT2 inhibitor, metformin and a gliptin, must be documented in the patient's medical records.

saxagliptin 5 mg + metformin hydrochloride 1 g modified release tablet, 28

11299Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	58.81	41.30	Kombiglyze XR 5/1000 [AP]

saxagliptin 2.5 mg + metformin hydrochloride 1 g modified release tablet, 56

11285Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	60.22	41.30	Kombiglyze XR 2.5/1000 [AP]

saxagliptin 5 mg + metformin hydrochloride 500 mg modified release tablet, 28

11312J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	58.06	41.30	Kombiglyze XR 5/500 [AP]

■ SAXAGLIPTIN + METFORMIN**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a

patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)

6333

Diabetes mellitus type 2

Clinical criteria:

- Patient must have, or have had, a HbA1c measurement greater than 7% despite treatment with metformin; OR
- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period despite treatment with metformin.

The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

A patient whose diabetes was previously demonstrated unable to be controlled with metformin does not need to requalify on this criterion before being eligible for PBS-subsidised treatment with this fixed dose combination.

Note This fixed dose combination is not PBS-subsidised for use as initial therapy or in combination with a thiazolidinedione (glitazone) or a glucagon-like peptide-1.

Authority required (STREAMLINED)

6335

Diabetes mellitus type 2

Treatment Phase: Continuing

Clinical criteria:

- Patient must have previously received and been stabilised on a PBS-subsidised regimen of oral diabetic medicines which includes metformin and saxagliptin.

Note This fixed dose combination is not PBS-subsidised for use as initial therapy or in combination with a thiazolidinedione (glitazone) or a glucagon-like peptide-1.

Authority required (STREAMLINED)

6344

Diabetes mellitus type 2

Clinical criteria:

- The treatment must be in combination with a sulfonylurea, **AND**
- Patient must have, or have had, a HbA1c measurement greater than 7% prior to the initiation of a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor despite treatment with optimal doses of dual oral therapy; OR
- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor despite treatment with optimal doses of dual oral therapy.

The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

A patient whose diabetes was previously demonstrated unable to be controlled with metformin or a sulfonylurea does not need to requalify on this criterion before being eligible for PBS-subsidised treatment with this fixed dose combination.

Note This fixed dose combination is not PBS-subsidised for use as initial therapy or in combination with a thiazolidinedione (glitazone) or a glucagon-like peptide-1.

Note PBS subsidised dual oral therapy does not include concomitant use of a combination of: a gliptin, a glitazone or an SGLT2 inhibitor.

Authority required (STREAMLINED)

7530

Diabetes mellitus type 2

Treatment Phase: Continuing treatment

Clinical criteria:

- The treatment must be in combination with a sodium-glucose co-transporter 2 (SGLT2) inhibitor, **AND**

- Patient must have previously received a PBS-subsidised regimen of oral diabetic medicines which included a sodium-glucose co-transporter 2 (SGLT2) inhibitor, metformin and a gliptin for this condition.

Note This fixed dose combination is not PBS-subsidised for initiating dual oral combination treatment or in combination with a thiazolidinedione (glitazone), a glucagon-like peptide-1 analogue, or another dipeptidyl peptidase 4 inhibitor (gliptin).

Note PBS-subsidised dual oral therapy does not include combination use of: a gliptin with an SGLT2 inhibitor; or

- a gliptin with a glitazone; or
- an SGLT2 inhibitor with a glitazone.

saxagliptin 5 mg + metformin hydrochloride 1 g modified release tablet, 28

10051B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	58.81	41.30	Kombiglyze XR 5/1000 [AP]

saxagliptin 2.5 mg + metformin hydrochloride 1 g modified release tablet, 56

10048W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	60.22	41.30	Kombiglyze XR 2.5/1000 [AP]

saxagliptin 5 mg + metformin hydrochloride 500 mg modified release tablet, 28

10055F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	58.06	41.30	Kombiglyze XR 5/500 [AP]

■ SITAGLIPTIN + METFORMIN

Note This fixed dose combination is not PBS-subsidised for initiating dual oral combination treatment or in combination with a thiazolidinedione (glitazone), a glucagon-like peptide-1 analogue, or another dipeptidyl peptidase 4 inhibitor (gliptin).

Note PBS-subsidised dual oral therapy does not include combination use of: a gliptin with an SGLT2 inhibitor; or

- a gliptin with a glitazone; or
- an SGLT2 inhibitor with a glitazone.

Authority required (STREAMLINED)

7507

Diabetes mellitus type 2

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be in combination with a sodium-glucose co-transporter 2 (SGLT2) inhibitor, **AND**
- Patient must have an HbA1c measurement greater than 7% despite treatment with a PBS-subsidised regimen of oral diabetic medicines which includes metformin and an SGLT2 inhibitor for this condition; OR
- Patient must have, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation of triple oral therapy with a sodium-glucose co-transporter 2 (SGLT2) inhibitor, metformin and a gliptin.

The date and level of the qualifying HbA1c measurement must be documented in the patient's medical records at the time triple oral therapy with an SGLT2 inhibitor, metformin and a gliptin is initiated.

The HbA1c must be no more than 4 months old at the time triple oral therapy with an SGLT2 inhibitor, metformin and a gliptin is initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of triple oral therapy with an SGLT2 inhibitor, metformin and a gliptin, must be documented in the patient's medical records.

sitagliptin 50 mg + metformin hydrochloride 850 mg tablet, 56

11582N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	55.29	41.30	Janumet [MK]

sitagliptin 50 mg + metformin hydrochloride 500 mg tablet, 56

11586T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	54.29	41.30	Janumet [MK]

sitagliptin 50 mg + metformin hydrochloride 1 g modified release tablet, 56

11580L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	55.69	41.30	Janumet XR [MK]

sitagliptin 50 mg + metformin hydrochloride 1 g tablet, 56

11574E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	55.69	41.30	Janumet [MK]

sitagliptin 100 mg + metformin hydrochloride 1 g tablet: modified release, 28

11566R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	54.29	41.30	Janumet XR [MK]

■ SITAGLIPTIN + METFORMIN

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)

6333

Diabetes mellitus type 2

Clinical criteria:

- Patient must have, or have had, a HbA1c measurement greater than 7% despite treatment with metformin; OR
- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period despite treatment with metformin.

The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

A patient whose diabetes was previously demonstrated unable to be controlled with metformin does not need to requalify on this criterion before being eligible for PBS-subsidised treatment with this fixed dose combination.

Note This fixed dose combination is not PBS-subsidised for use as initial therapy or in combination with a thiazolidinedione (glitazone) or a glucagon-like peptide-1.

Authority required (STREAMLINED)

6334

Diabetes mellitus type 2

Treatment Phase: Continuing

Clinical criteria:

- Patient must have previously received and been stabilised on a PBS-subsidised regimen of oral diabetic medicines which includes metformin and sitagliptin.

Note This fixed dose combination is not PBS-subsidised for use as initial therapy or in combination with a thiazolidinedione (glitazone) or a glucagon-like peptide-1.

Authority required (STREAMLINED)

6344

Diabetes mellitus type 2

Clinical criteria:

- The treatment must be in combination with a sulfonylurea, **AND**
- Patient must have, or have had, a HbA1c measurement greater than 7% prior to the initiation of a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor despite treatment with optimal doses of dual oral therapy; OR
- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor despite treatment with optimal doses of dual oral therapy.

The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

A patient whose diabetes was previously demonstrated unable to be controlled with metformin or a sulfonylurea does not need to requalify on this criterion before being eligible for PBS-subsidised treatment with this fixed dose combination.

Note This fixed dose combination is not PBS-subsidised for use as initial therapy or in combination with a thiazolidinedione (glitazone) or a glucagon-like peptide-1.

Note PBS subsidised dual oral therapy does not include concomitant use of a combination of: a gliptin, a glitazone or an SGLT2 inhibitor.

Authority required (STREAMLINED)

6443

Diabetes mellitus type 2

Clinical criteria:

- The treatment must be in combination with insulin, **AND**
- Patient must have, or have had, a HbA1c measurement greater than 7% prior to the initiation of a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor despite treatment with insulin and oral antidiabetic agents, or insulin alone where metformin is contraindicated; OR
- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor despite treatment with insulin and oral antidiabetic agents, or insulin alone where metformin is contraindicated.

The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

Note This fixed dose combination is not PBS-subsidised for use as initial therapy or in combination with a thiazolidinedione (glitazone) or a glucagon-like peptide-1.

Authority required (STREAMLINED)

7530

Diabetes mellitus type 2

Treatment Phase: Continuing treatment

Clinical criteria:

- The treatment must be in combination with a sodium-glucose co-transporter 2 (SGLT2) inhibitor, **AND**
- Patient must have previously received a PBS-subsidised regimen of oral diabetic medicines which included a sodium-glucose co-transporter 2 (SGLT2) inhibitor, metformin and a gliptin for this condition.

Note This fixed dose combination is not PBS-subsidised for initiating dual oral combination treatment or in combination with a thiazolidinedione (glitazone), a glucagon-like peptide-1 analogue, or another dipeptidyl peptidase 4 inhibitor (gliptin).

Note PBS-subsidised dual oral therapy does not include combination use of: a gliptin with an SGLT2 inhibitor; or

- a gliptin with a glitazone; or
- an SGLT2 inhibitor with a glitazone.

sitagliptin 50 mg + metformin hydrochloride 850 mg tablet, 56

9450J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	55.29	41.30	Janumet [MK]

sitagliptin 50 mg + metformin hydrochloride 500 mg tablet, 56

9449H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	54.29	41.30	Janumet [MK]

sitagliptin 50 mg + metformin hydrochloride 1 g modified release tablet, 56

10090C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	55.69	41.30	Janumet XR [MK]

sitagliptin 50 mg + metformin hydrochloride 1 g tablet, 56

9451K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	55.69	41.30	Janumet [MK]

sitagliptin 100 mg + metformin hydrochloride 1 g tablet: modified release, 28

10089B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	54.29	41.30	Janumet XR [MK]

■ VILDAGLIPTIN + METFORMIN

Note This fixed dose combination tablet is not PBS-subsidised for use as initial therapy or in combination with a thiazolidinedione (glitazone), a glucagon-like peptide-1 or an SGLT2 inhibitor

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)

6333

Diabetes mellitus type 2

Clinical criteria:

- Patient must have, or have had, a HbA1c measurement greater than 7% despite treatment with metformin; OR

- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period despite treatment with metformin.

The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

A patient whose diabetes was previously demonstrated unable to be controlled with metformin does not need to requalify on this criterion before being eligible for PBS-subsidised treatment with this fixed dose combination.

Authority required (STREAMLINED)

6357

Diabetes mellitus type 2

Treatment Phase: Continuing

Clinical criteria:

- Patient must have previously received and been stabilised on a PBS-subsidised regimen of oral diabetic medicines which includes metformin and vildagliptin.

Authority required (STREAMLINED)

6344

Diabetes mellitus type 2

Clinical criteria:

- The treatment must be in combination with a sulfonylurea, **AND**
- Patient must have, or have had, a HbA1c measurement greater than 7% prior to the initiation of a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor despite treatment with optimal doses of dual oral therapy; OR
- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor despite treatment with optimal doses of dual oral therapy.

The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

A patient whose diabetes was previously demonstrated unable to be controlled with metformin or a sulfonylurea does not need to requalify on this criterion before being eligible for PBS-subsidised treatment with this fixed dose combination.

Note PBS subsidised dual oral therapy does not include concomitant use of a combination of: a gliptin, a glitazone or an SGLT2 inhibitor.

Authority required (STREAMLINED)

6443

Diabetes mellitus type 2

Clinical criteria:

- The treatment must be in combination with insulin, **AND**
- Patient must have, or have had, a HbA1c measurement greater than 7% prior to the initiation of a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor despite treatment with insulin and oral antidiabetic agents, or insulin alone where metformin is contraindicated; OR
- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor despite treatment with insulin and oral antidiabetic agents, or insulin alone where metformin is contraindicated.

The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

vildagliptin 50 mg + metformin hydrochloride 1 g tablet, 60

5476F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	56.63	41.30	Galvumet 50/1000 [NV]

vildagliptin 50 mg + metformin hydrochloride 850 mg tablet, 60

5475E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	56.19	41.30	Galvumet 50/850 [NV]

vildagliptin 50 mg + metformin hydrochloride 500 mg tablet, 60

5474D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	55.11	41.30	Galvumet 50/500 [NV]

Alpha glucosidase inhibitors

■ ACARBOSE

acarbose 100 mg tablet, 90

8189B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	35.63	36.92	^a Acarbose Mylan [AF]	^a GLYBOSAY [RW]
			^b 2.30	37.93	36.92	^a Glucobay 100 [BN]	

acarbose 50 mg tablet, 90

8188Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	28.70	29.99	^a Acarbose Mylan [AF]	^a GLYBOSAY [RW]

Thiazolidinediones

■ PIOGLITAZONE

Note This drug is not PBS-subsidised for use as monotherapy or in combination with a dipeptidyl peptidase 4 inhibitor (gliptin), a glucagon-like peptide-1 or an SGLT2 inhibitor.

Authority required (STREAMLINED)

4363

Diabetes mellitus type 2

Clinical criteria:

- The treatment must be in combination with metformin; OR
- The treatment must be in combination with a sulfonylurea, **AND**
- Patient must have a contraindication to a combination of metformin and a sulfonylurea; OR
- Patient must not have tolerated a combination of metformin and a sulfonylurea, **AND**
- Patient must have, or have had, a HbA1c measurement greater than 7% prior to the initiation of a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor despite treatment with either metformin or a sulfonylurea; OR
- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor despite treatment with either metformin or a sulfonylurea.

The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

Authority required (STREAMLINED)

4388

Diabetes mellitus type 2

Clinical criteria:

- The treatment must be in combination with insulin, **AND**
- Patient must have, or have had, a HbA1c measurement greater than 7% prior to the initiation of a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor despite treatment with insulin and oral antidiabetic agents, or insulin alone where metformin is contraindicated; OR
- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor despite treatment with insulin and oral antidiabetic agents, or insulin alone where metformin is contraindicated.

The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor is initiated. The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

Authority required (STREAMLINED)

4364

Diabetes mellitus type 2

Clinical criteria:

- The treatment must be in combination with metformin, **AND**
- The treatment must be in combination with a sulfonylurea, **AND**
- Patient must have, or have had, a HbA1c measurement greater than 7% prior to the initiation of a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor despite treatment with maximally tolerated doses of metformin and a sulfonylurea; OR
- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor despite treatment with maximally tolerated doses of metformin and a sulfonylurea.

The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

pioglitazone 15 mg tablet, 28

8694N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	19.78	21.07	^a Acpio 15 [RF] ^a Actos [TK] ^a Pioglitazone AN [EA]	^a Actaze [RW] ^a APOTEX-Pioglitazone [TX] ^a Vexazone [AF]

pioglitazone 45 mg tablet, 28

8696Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	27.55	28.84	^a Acpio 45 [RF] ^a Actos [TK] ^a NOUMED PIOGLITAZONE [VO] ^a Pioglitazone Sandoz [SZ]	^a Actaze [RW] ^a APOTEX-Pioglitazone [TX] ^a Pioglitazone AN [EA] ^a Vexazone [AF]

pioglitazone 30 mg tablet, 28

8695P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	23.97	25.26	^a Acpio 30 [RF] ^a Actos [TK] ^a NOUMED PIOGLITAZONE [VO] ^a Pioglitazone Sandoz [SZ]	^a Actaze [RW] ^a APOTEX-Pioglitazone [TX] ^a Pioglitazone AN [EA] ^a Vexazone [AF]

Dipeptidyl peptidase 4 (DPP-4) inhibitors

■ ALOGLIPTIN

Note Alogliptin is not PBS-subsidised for use in combination with metformin and a sulfonylurea (triple oral therapy), as monotherapy or in combination with a thiazolidinedione (glitazone), a glucagon-like peptide-1 or an SGLT2 inhibitor.

Authority required (STREAMLINED)

4349

Diabetes mellitus type 2

Clinical criteria:

- The treatment must be in combination with metformin; OR
- The treatment must be in combination with a sulfonylurea, **AND**
- Patient must have, or have had, a HbA1c measurement greater than 7% despite treatment with either metformin or a sulfonylurea; OR
- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period despite treatment with either metformin or a sulfonylurea.

The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

A patient whose diabetes was previously demonstrated unable to be controlled with metformin or a sulfonylurea does not need to requalify on this criterion before being eligible for PBS-subsidised treatment with alogliptin.

alogliptin 6.25 mg tablet, 28

2944Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	57.31	41.30	Nesina [TK]

alogliptin 12.5 mg tablet, 28

2933J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	57.31	41.30	Nesina [TK]

alogliptin 25 mg tablet, 28

2986E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	57.31	41.30	Nesina [TK]

■ LINAGLIPTIN

Note This drug is not PBS-subsidised for use as monotherapy or in combination with a thiazolidinedione (glitazone), or a glucagon-like peptide-1 analogue.

Note PBS-subsidised dual oral therapy does not include combination use of: a gliptin with an SGLT2 inhibitor; or

- a gliptin with a glitazone; or
- an SGLT2 inhibitor with a glitazone.

Authority required (STREAMLINED)

7541

Diabetes mellitus type 2

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be in combination with metformin, **AND**
- The treatment must be in combination with a sodium-glucose co-transporter 2 (SGLT2) inhibitor, **AND**
- Patient must have an HbA1c measurement greater than 7% despite treatment with dual oral combination therapy with metformin and an SGLT2 inhibitor; OR
- Patient must have, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation of triple oral therapy with a sodium-glucose co-transporter 2 (SGLT2) inhibitor, metformin and a gliptin.

The date and level of the qualifying HbA1c measurement must be documented in the patient's medical records at the time triple oral therapy with an SGLT2 inhibitor, metformin and a gliptin is initiated.

The HbA1c must be no more than 4 months old at the time triple oral therapy with an SGLT2 inhibitor, metformin and a gliptin is initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of triple oral therapy with an SGLT2 inhibitor, metformin and a gliptin, must be documented in the patient's medical records.

linagliptin 5 mg tablet, 30

11280Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	60.54	41.30	Trajenta [BY]

■ LINAGLIPTIN

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)

6346

Diabetes mellitus type 2

Clinical criteria:

- The treatment must be in combination with metformin; OR
- The treatment must be in combination with a sulfonylurea, **AND**
- Patient must have, or have had, a HbA1c measurement greater than 7% despite treatment with either metformin or a sulfonylurea; OR
- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period despite treatment with either metformin or a sulfonylurea. The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

A patient whose diabetes was previously demonstrated unable to be controlled with metformin or a sulfonylurea does not need to requalify on this criterion before being eligible for PBS-subsidised treatment with this drug.

Note This drug is not PBS-subsidised for use as monotherapy or in combination with a thiazolidinedione (glitazone), or a glucagon-like peptide-1.

Authority required (STREAMLINED)

6363

Diabetes mellitus type 2

Clinical criteria:

- The treatment must be in combination with metformin, **AND**
- The treatment must be in combination with a sulfonylurea, **AND**
- Patient must have, or have had, a HbA1c measurement greater than 7% prior to the initiation of a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor despite treatment with optimal doses of dual oral therapy; OR
- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor despite treatment with optimal doses of dual oral therapy.

The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

A patient whose diabetes was previously demonstrated unable to be controlled with metformin or a sulfonylurea does not need to requalify on this criterion before being eligible for PBS-subsidised treatment with this drug.

Note This drug is not PBS-subsidised for use as monotherapy or in combination with a thiazolidinedione (glitazone), or a glucagon-like peptide-1.

Note PBS subsidised dual oral therapy does not include concomitant use of a combination of: a gliptin, a glitazone or an SGLT2 inhibitor.

Authority required (STREAMLINED)

6376

Diabetes mellitus type 2

Clinical criteria:

- The treatment must be in combination with insulin, **AND**
- Patient must have, or have had, a HbA1c measurement greater than 7% prior to the initiation of a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor despite treatment with insulin and oral antidiabetic agents, or insulin alone where metformin is contraindicated; OR
- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor despite treatment with insulin and oral antidiabetic agents, or insulin alone where metformin is contraindicated.

The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or

(b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

Note This drug is not PBS-subsidised for use as monotherapy or in combination with a thiazolidinedione (glitazone), or a glucagon-like peptide-1.

Authority required (STREAMLINED)

7505

Diabetes mellitus type 2

Treatment Phase: Continuing treatment

Clinical criteria:

- The treatment must be in combination with metformin, **AND**
- The treatment must be in combination with a sodium-glucose co-transporter 2 (SGLT2) inhibitor, **AND**
- Patient must have previously received a PBS-subsidised regimen of oral diabetic medicines which included a sodium-glucose co-transporter 2 (SGLT2) inhibitor, metformin and a gliptin for this condition.

Note This drug is not PBS-subsidised for use as monotherapy or in combination with a thiazolidinedione (glitazone), or a glucagon-like peptide-1 analogue.

Note PBS-subsidised dual oral therapy does not include combination use of: a gliptin with an SGLT2 inhibitor; or

- a gliptin with a glitazone; or
- an SGLT2 inhibitor with a glitazone.

linagliptin 5 mg tablet, 30

3387G

NP

Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
1	5	..	60.54	41.30	Trajenta [BY]

▪ **SAXAGLIPTIN**

Note This drug is not PBS-subsidised for use as monotherapy or in combination with a thiazolidinedione (glitazone), or a glucagon-like peptide-1 analogue.

Note PBS-subsidised dual oral therapy does not include combination use of: a gliptin with an SGLT2 inhibitor; or

- a gliptin with a glitazone; or
- an SGLT2 inhibitor with a glitazone.

Authority required (STREAMLINED)

7541

Diabetes mellitus type 2

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be in combination with metformin, **AND**
- The treatment must be in combination with a sodium-glucose co-transporter 2 (SGLT2) inhibitor, **AND**
- Patient must have an HbA1c measurement greater than 7% despite treatment with dual oral combination therapy with metformin and an SGLT2 inhibitor; OR
- Patient must have, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation of triple oral therapy with a sodium-glucose co-transporter 2 (SGLT2) inhibitor, metformin and a gliptin.

The date and level of the qualifying HbA1c measurement must be documented in the patient's medical records at the time triple oral therapy with an SGLT2 inhibitor, metformin and a gliptin is initiated.

The HbA1c must be no more than 4 months old at the time triple oral therapy with an SGLT2 inhibitor, metformin and a gliptin is initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of triple oral therapy with an SGLT2 inhibitor, metformin and a gliptin, must be documented in the patient's medical records.

saxagliptin 5 mg tablet, 28

11311H

Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
1	5	..	57.31	41.30	Onglyza [AP]

saxagliptin 2.5 mg tablet, 28

11292H

Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
1	5	..	57.31	41.30	Onglyza [AP]

▪ **SAXAGLIPTIN**

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**6346**

Diabetes mellitus type 2

Clinical criteria:

- The treatment must be in combination with metformin; OR
- The treatment must be in combination with a sulfonylurea, **AND**
- Patient must have, or have had, a HbA1c measurement greater than 7% despite treatment with either metformin or a sulfonylurea; OR
- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period despite treatment with either metformin or a sulfonylurea. The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

A patient whose diabetes was previously demonstrated unable to be controlled with metformin or a sulfonylurea does not need to requalify on this criterion before being eligible for PBS-subsidised treatment with this drug.

Note This drug is not PBS-subsidised for use as monotherapy or in combination with a thiazolidinedione (glitazone), or a glucagon-like peptide-1.

Authority required (STREAMLINED)**6363**

Diabetes mellitus type 2

Clinical criteria:

- The treatment must be in combination with metformin, **AND**
- The treatment must be in combination with a sulfonylurea, **AND**
- Patient must have, or have had, a HbA1c measurement greater than 7% prior to the initiation of a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor despite treatment with optimal doses of dual oral therapy; OR
- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor despite treatment with optimal doses of dual oral therapy.

The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

A patient whose diabetes was previously demonstrated unable to be controlled with metformin or a sulfonylurea does not need to requalify on this criterion before being eligible for PBS-subsidised treatment with this drug.

Note This drug is not PBS-subsidised for use as monotherapy or in combination with a thiazolidinedione (glitazone), or a glucagon-like peptide-1.

Note PBS subsidised dual oral therapy does not include concomitant use of a combination of: a gliptin, a glitazone or an SGLT2 inhibitor.

Authority required (STREAMLINED)**7505**

Diabetes mellitus type 2

Treatment Phase: Continuing treatment

Clinical criteria:

- The treatment must be in combination with metformin, **AND**
- The treatment must be in combination with a sodium-glucose co-transporter 2 (SGLT2) inhibitor, **AND**
- Patient must have previously received a PBS-subsidised regimen of oral diabetic medicines which included a sodium-glucose co-transporter 2 (SGLT2) inhibitor, metformin and a gliptin for this condition.

Note This drug is not PBS-subsidised for use as monotherapy or in combination with a thiazolidinedione (glitazone), or a glucagon-like peptide-1 analogue.

Note PBS-subsidised dual oral therapy does not include combination use of: a gliptin with an SGLT2 inhibitor; or

- a gliptin with a glitazone; or
- an SGLT2 inhibitor with a glitazone.

saxagliptin 5 mg tablet, 28

8983T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	57.31	41.30	Onglyza [AP]

saxagliptin 2.5 mg tablet, 28

10128C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	57.31	41.30	Onglyza [AP]

■ SITAGLIPTIN

Note This drug is not PBS-subsidised for use as monotherapy or in combination with a thiazolidinedione (glitazone), or a glucagon-like peptide-1 analogue.

Note PBS-subsidised dual oral therapy does not include combination use of: a gliptin with an SGLT2 inhibitor; or

- a gliptin with a glitazone; or
- an SGLT2 inhibitor with a glitazone.

Authority required (STREAMLINED)**7541**

Diabetes mellitus type 2

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be in combination with metformin, **AND**
- The treatment must be in combination with a sodium-glucose co-transporter 2 (SGLT2) inhibitor, **AND**
- Patient must have an HbA1c measurement greater than 7% despite treatment with dual oral combination therapy with metformin and an SGLT2 inhibitor; OR
- Patient must have, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation of triple oral therapy with a sodium-glucose co-transporter 2 (SGLT2) inhibitor, metformin and a gliptin.

The date and level of the qualifying HbA1c measurement must be documented in the patient's medical records at the time triple oral therapy with an SGLT2 inhibitor, metformin and a gliptin is initiated.

The HbA1c must be no more than 4 months old at the time triple oral therapy with an SGLT2 inhibitor, metformin and a gliptin is initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of triple oral therapy with an SGLT2 inhibitor, metformin and a gliptin, must be documented in the patient's medical records.

sitagliptin 100 mg tablet, 28

11576G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	52.78	41.30	Januvia [MK]

sitagliptin 50 mg tablet, 28

11573D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	52.78	41.30	Januvia [MK]

sitagliptin 25 mg tablet, 28

11572C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	52.78	41.30	Januvia [MK]

■ SITAGLIPTIN**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**6346**

Diabetes mellitus type 2

Clinical criteria:

- The treatment must be in combination with metformin; OR
- The treatment must be in combination with a sulfonylurea, **AND**
- Patient must have, or have had, a HbA1c measurement greater than 7% despite treatment with either metformin or a sulfonylurea; OR
- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period despite treatment with either metformin or a sulfonylurea.

The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

A patient whose diabetes was previously demonstrated unable to be controlled with metformin or a sulfonylurea does not need to requalify on this criterion before being eligible for PBS-subsidised treatment with this drug.

Note This drug is not PBS-subsidised for use as monotherapy or in combination with a thiazolidinedione (glitazone), or a glucagon-like peptide-1.

Authority required (STREAMLINED)

6363

Diabetes mellitus type 2

Clinical criteria:

- The treatment must be in combination with metformin, **AND**
- The treatment must be in combination with a sulfonylurea, **AND**
- Patient must have, or have had, a HbA1c measurement greater than 7% prior to the initiation of a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor despite treatment with optimal doses of dual oral therapy; OR
- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor despite treatment with optimal doses of dual oral therapy.

The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

A patient whose diabetes was previously demonstrated unable to be controlled with metformin or a sulfonylurea does not need to requalify on this criterion before being eligible for PBS-subsidised treatment with this drug.

Note This drug is not PBS-subsidised for use as monotherapy or in combination with a thiazolidinedione (glitazone), or a glucagon-like peptide-1.

Note PBS subsidised dual oral therapy does not include concomitant use of a combination of: a gliptin, a glitazone or an SGLT2 inhibitor.

Authority required (STREAMLINED)

6376

Diabetes mellitus type 2

Clinical criteria:

- The treatment must be in combination with insulin, **AND**
- Patient must have, or have had, a HbA1c measurement greater than 7% prior to the initiation of a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor despite treatment with insulin and oral antidiabetic agents, or insulin alone where metformin is contraindicated; OR
- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor despite treatment with insulin and oral antidiabetic agents, or insulin alone where metformin is contraindicated.

The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

Note This drug is not PBS-subsidised for use as monotherapy or in combination with a thiazolidinedione (glitazone), or a glucagon-like peptide-1.

Authority required (STREAMLINED)**7505**

Diabetes mellitus type 2

Treatment Phase: Continuing treatment

Clinical criteria:

- The treatment must be in combination with metformin, **AND**
- The treatment must be in combination with a sodium-glucose co-transporter 2 (SGLT2) inhibitor, **AND**
- Patient must have previously received a PBS-subsidised regimen of oral diabetic medicines which included a sodium-glucose co-transporter 2 (SGLT2) inhibitor, metformin and a gliptin for this condition.

Note This drug is not PBS-subsidised for use as monotherapy or in combination with a thiazolidinedione (glitazone), or a glucagon-like peptide-1 analogue.

Note PBS-subsidised dual oral therapy does not include combination use of: a gliptin with an SGLT2 inhibitor; or

- a gliptin with a glitazone; or
- an SGLT2 inhibitor with a glitazone.

sitagliptin 100 mg tablet, 28

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
9182G	1	5	..	52.78	41.30	Januvia [MK]

sitagliptin 50 mg tablet, 28

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
9181F	1	5	..	52.78	41.30	Januvia [MK]

sitagliptin 25 mg tablet, 28

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
9180E	1	5	..	52.78	41.30	Januvia [MK]

■ VILDAGLIPTIN

Note This drug is not PBS-subsidised for use as monotherapy or in combination with a thiazolidinedione (glitazone), a glucagon-like peptide-1 or an SGLT2 inhibitor.

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**6346**

Diabetes mellitus type 2

Clinical criteria:

- The treatment must be in combination with metformin; OR
- The treatment must be in combination with a sulfonylurea, **AND**
- Patient must have, or have had, a HbA1c measurement greater than 7% despite treatment with either metformin or a sulfonylurea; OR
- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period despite treatment with either metformin or a sulfonylurea. The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

A patient whose diabetes was previously demonstrated unable to be controlled with metformin or a sulfonylurea does not need to requalify on this criterion before being eligible for PBS-subsidised treatment with this drug.

Authority required (STREAMLINED)**6363**

Diabetes mellitus type 2

Clinical criteria:

- The treatment must be in combination with metformin, **AND**
- The treatment must be in combination with a sulfonylurea, **AND**
- Patient must have, or have had, a HbA1c measurement greater than 7% prior to the initiation of a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor despite treatment with optimal doses of dual oral therapy; OR

- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor despite treatment with optimal doses of dual oral therapy.

The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

A patient whose diabetes was previously demonstrated unable to be controlled with metformin or a sulfonylurea does not need to requalify on this criterion before being eligible for PBS-subsidised treatment with this drug.

Note PBS subsidised dual oral therapy does not include concomitant use of a combination of: a gliptin, a glitazone or an SGLT2 inhibitor.

Authority required (STREAMLINED)

6376

Diabetes mellitus type 2

Clinical criteria:

- The treatment must be in combination with insulin, **AND**
- Patient must have, or have had, a HbA1c measurement greater than 7% prior to the initiation of a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor despite treatment with insulin and oral antidiabetic agents, or insulin alone where metformin is contraindicated; **OR**
- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor despite treatment with insulin and oral antidiabetic agents, or insulin alone where metformin is contraindicated.

The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

vildagliptin 50 mg tablet, 60

3415R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	55.69	41.30	Galvus [NV]

Glucagon-like peptide-1 (GLP-1) analogues

■ DULAGLUTIDE

Note This drug is not PBS-subsidised for use as monotherapy or in combination with a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone) or an SGLT2 inhibitor.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

7645

Diabetes mellitus type 2

Clinical criteria:

- The treatment must be in combination with metformin, **AND**
- Patient must have a contraindication to a combination of metformin and a sulfonylurea; **OR**
- Patient must not have tolerated a combination of metformin and a sulfonylurea, **AND**
- Patient must have, or have had, a HbA1c measurement greater than 7% prior to the initiation of a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor despite treatment with metformin; **OR**
- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor despite treatment with metformin.

The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or

(b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

Authority required (STREAMLINED)

5478

Diabetes mellitus type 2

Clinical criteria:

- The treatment must be in combination with metformin, **AND**
- The treatment must be in combination with a sulfonylurea, **AND**
- Patient must have, or have had, a HbA1c measurement greater than 7% prior to the initiation of a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor despite treatment with maximally tolerated doses of metformin and a sulfonylurea; OR
- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor despite treatment with maximally tolerated doses of metformin and a sulfonylurea.

The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

Authority required (STREAMLINED)

5469

Diabetes mellitus type 2

Clinical criteria:

- The treatment must be in combination with insulin, **AND**
- The treatment must be in combination with metformin unless contraindicated or not tolerated, **AND**
- Patient must have, or have had, a HbA1c measurement greater than 7% prior to the initiation of a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor despite treatment with insulin and oral antidiabetic agents, or insulin alone where metformin is contraindicated; OR
- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor despite treatment with insulin and oral antidiabetic agents, or insulin alone where metformin is contraindicated.

The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

dulaglutide 1.5 mg/0.5 mL injection, 4 x 0.5 mL pen devices

11364D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	132.83	41.30	Trulicity [LY]

▪ **EXENATIDE**

Note This drug is not PBS-subsidised for use as monotherapy or in combination with a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), an insulin or an SGLT2 inhibitor.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

6519

Diabetes mellitus type 2

Clinical criteria:

- The treatment must be in combination with metformin; OR
- The treatment must be in combination with a sulfonylurea, **AND**
- Patient must have a contraindication to a combination of metformin and a sulfonylurea; OR
- Patient must not have tolerated a combination of metformin and a sulfonylurea, **AND**

- Patient must have, or have had, a HbA1c measurement greater than 7% prior to the initiation of a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor despite treatment with either metformin or a sulfonylurea; OR
- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor despite treatment with either metformin or a sulfonylurea.

The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

Authority required (STREAMLINED)

6505

Diabetes mellitus type 2

Clinical criteria:

- The treatment must be in combination with metformin, **AND**
- The treatment must be in combination with a sulfonylurea, **AND**
- Patient must have, or have had, a HbA1c measurement greater than 7% prior to the initiation of a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor despite treatment with maximally tolerated doses of metformin and a sulfonylurea; OR
- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor despite treatment with maximally tolerated doses of metformin and a sulfonylurea.

The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

exenatide 2 mg modified release injection [1 chamber] (&) inert substance diluent [1 chamber], 4 dual chamber pen devices

10888C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	120.25	41.30	Bydureon [AP]

■ EXENATIDE

Note This drug is not PBS-subsidised for use as monotherapy or in combination with a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone) or an SGLT2 inhibitor.

Authority required (STREAMLINED)

5500

Diabetes mellitus type 2

Clinical criteria:

- The treatment must be in combination with metformin; OR
- The treatment must be in combination with a sulfonylurea, **AND**
- Patient must have a contraindication to a combination of metformin and a sulfonylurea; OR
- Patient must not have tolerated a combination of metformin and a sulfonylurea, **AND**
- Patient must have, or have had, a HbA1c measurement greater than 7% prior to the initiation of a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor despite treatment with either metformin or a sulfonylurea; OR
- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor despite treatment with either metformin or a sulfonylurea.

The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

Authority required (STREAMLINED)

5478

Diabetes mellitus type 2

Clinical criteria:

- The treatment must be in combination with metformin, **AND**
- The treatment must be in combination with a sulfonylurea, **AND**
- Patient must have, or have had, a HbA1c measurement greater than 7% prior to the initiation of a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor despite treatment with maximally tolerated doses of metformin and a sulfonylurea; OR
- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor despite treatment with maximally tolerated doses of metformin and a sulfonylurea.

The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

Authority required (STREAMLINED)

5469

Diabetes mellitus type 2

Clinical criteria:

- The treatment must be in combination with insulin, **AND**
- The treatment must be in combination with metformin unless contraindicated or not tolerated, **AND**
- Patient must have, or have had, a HbA1c measurement greater than 7% prior to the initiation of a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor despite treatment with insulin and oral antidiabetic agents, or insulin alone where metformin is contraindicated; OR
- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor despite treatment with insulin and oral antidiabetic agents, or insulin alone where metformin is contraindicated.

The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

exenatide 5 microgram/0.02 mL injection, 1.2 mL pen device

3423E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	61.44	41.30	Byetta 5 microgram [AP]

exenatide 10 microgram/0.04 mL injection, 2.4 mL pen device

3424F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	82.62	41.30	Byetta 10 microgram [AP]

■ SEMAGLUTIDE

Note This drug is not PBS-subsidised for use as monotherapy or in combination with a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), an insulin or an SGLT2 inhibitor.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

5500

Diabetes mellitus type 2

Clinical criteria:

- The treatment must be in combination with metformin; OR
- The treatment must be in combination with a sulfonylurea, **AND**

- Patient must have a contraindication to a combination of metformin and a sulfonylurea; OR
- Patient must not have tolerated a combination of metformin and a sulfonylurea, **AND**
- Patient must have, or have had, a HbA1c measurement greater than 7% prior to the initiation of a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor despite treatment with either metformin or a sulfonylurea; OR
- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor despite treatment with either metformin or a sulfonylurea.

The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

Authority required (STREAMLINED)

5478

Diabetes mellitus type 2

Clinical criteria:

- The treatment must be in combination with metformin, **AND**
- The treatment must be in combination with a sulfonylurea, **AND**
- Patient must have, or have had, a HbA1c measurement greater than 7% prior to the initiation of a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor despite treatment with maximally tolerated doses of metformin and a sulfonylurea; OR
- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor despite treatment with maximally tolerated doses of metformin and a sulfonylurea.

The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

semaglutide 1.34 mg/mL injection, 1 x 3 mL pen device

12075M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	132.83	41.30	Ozempic [NO]

semaglutide 1.34 mg/mL injection, 1 x 1.5 mL pen device

12080T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	132.83	41.30	Ozempic [NO]

Sodium-glucose co-transporter 2 (SGLT2) inhibitors

■ DAPAGLIFLOZIN

Note This drug is not PBS-subsidised for use as monotherapy or in combination with a thiazolidinedione (glitazone), or a glucagon-like peptide-1 analogue.

Note PBS-subsidised dual oral therapy does not include combination use of: a gliptin with an SGLT2 inhibitor; or

- a gliptin with a glitazone; or
- an SGLT2 inhibitor with a glitazone.

Authority required (STREAMLINED)

7528

Diabetes mellitus type 2

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be in combination with metformin, **AND**
- The treatment must be in combination with a dipeptidyl peptidase 4 inhibitor (gliptin), **AND**
- Patient must have an HbA1c measurement greater than 7% despite treatment with dual oral combination therapy with metformin and a gliptin; OR

- Patient must have, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation of triple oral therapy with a sodium-glucose co-transporter 2 (SGLT2) inhibitor, metformin and a gliptin.

The date and level of the qualifying HbA1c measurement must be documented in the patient's medical records at the time triple oral therapy with an SGLT2 inhibitor, metformin and a gliptin is initiated.

The HbA1c must be no more than 4 months old at the time triple oral therapy with an SGLT2 inhibitor, metformin and a gliptin is initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of triple oral therapy with an SGLT2 inhibitor, metformin and a gliptin, must be documented in the patient's medical records.

dapagliflozin 10 mg tablet, 28

11291G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	56.85	41.30	Forxiga [AP]

■ DAPAGLIFLOZIN

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)

7506

Diabetes mellitus type 2

Clinical criteria:

- The treatment must be in combination with metformin; OR
 - The treatment must be in combination with a sulfonylurea, **AND**
 - Patient must have, or have had, a HbA1c measurement greater than 7% despite treatment with either metformin or a sulfonylurea; OR
 - Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period despite treatment with either metformin or a sulfonylurea.
- The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of triple oral therapy with a gliptin and an SGLT2 inhibitor, must be documented in the patient's medical records.

A patient whose diabetes was previously demonstrated unable to be controlled with metformin or a sulfonylurea does not need to requalify on this criterion before being eligible for PBS-subsidised treatment with this drug.

Note This drug is not PBS-subsidised for use as monotherapy or in combination with a thiazolidinedione (glitazone), or a glucagon-like peptide-1.

Authority required (STREAMLINED)

4991

Diabetes mellitus type 2

Clinical criteria:

- The treatment must be in combination with insulin, **AND**
- Patient must have, or have had, a HbA1c measurement greater than 7% prior to the initiation of a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor despite treatment with insulin and oral antidiabetic agents, or insulin alone where metformin is contraindicated; OR
- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor despite treatment with insulin and oral antidiabetic agents, or insulin alone where metformin is contraindicated.

The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

Note This drug is not PBS-subsidised for use as monotherapy or in combination with a thiazolidinedione (glitazone), or a glucagon-like peptide-1.

Authority required (STREAMLINED)

5629

Diabetes mellitus type 2

Clinical criteria:

- The treatment must be in combination with metformin, **AND**
- The treatment must be in combination with a sulfonylurea, **AND**
- Patient must have, or have had, a HbA1c measurement greater than 7% prior to the initiation of a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor despite treatment with optimal doses of dual oral therapy; OR
- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 despite treatment with optimal doses of dual oral therapy.

The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

A patient whose diabetes was previously demonstrated unable to be controlled with metformin or a sulfonylurea does not need to requalify on this criterion before being eligible for PBS-subsidised treatment with this drug.

Note This drug is not PBS-subsidised for use as monotherapy or in combination with a thiazolidinedione (glitazone), or a glucagon-like peptide-1.

Note PBS subsidised dual oral therapy does not include concomitant use of a combination of: a gliptin, a glitazone or an SGLT2 inhibitor.

Authority required (STREAMLINED)

7495

Diabetes mellitus type 2

Treatment Phase: Continuing treatment

Clinical criteria:

- The treatment must be in combination with metformin, **AND**
- The treatment must be in combination with a dipeptidyl peptidase 4 inhibitor (gliptin), **AND**
- Patient must have previously received a PBS-subsidised regimen of oral diabetic medicines which included a sodium-glucose co-transporter 2 (SGLT2) inhibitor, metformin and a gliptin for this condition.

Note This drug is not PBS-subsidised for use as monotherapy or in combination with a thiazolidinedione (glitazone), or a glucagon-like peptide-1 analogue.

Note PBS-subsidised dual oral therapy does not include combination use of: a gliptin with an SGLT2 inhibitor; or

- a gliptin with a glitazone; or
- an SGLT2 inhibitor with a glitazone.

dapagliflozin 10 mg tablet, 28

10011X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	56.85	41.30	Forxiga [AP]

NP

▪ **EMPAGLIFLOZIN**

Note This drug is not PBS-subsidised for use as monotherapy or in combination with a thiazolidinedione (glitazone), or a glucagon-like peptide-1 analogue.

Note PBS-subsidised dual oral therapy does not include combination use of: a gliptin with an SGLT2 inhibitor; or

- a gliptin with a glitazone; or
- an SGLT2 inhibitor with a glitazone.

Authority required (STREAMLINED)

7528

Diabetes mellitus type 2

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be in combination with metformin, **AND**
- The treatment must be in combination with a dipeptidyl peptidase 4 inhibitor (gliptin), **AND**
- Patient must have an HbA1c measurement greater than 7% despite treatment with dual oral combination therapy with metformin and a gliptin; OR

- Patient must have, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation of triple oral therapy with a sodium-glucose co-transporter 2 (SGLT2) inhibitor, metformin and a gliptin.

The date and level of the qualifying HbA1c measurement must be documented in the patient's medical records at the time triple oral therapy with an SGLT2 inhibitor, metformin and a gliptin is initiated.

The HbA1c must be no more than 4 months old at the time triple oral therapy with an SGLT2 inhibitor, metformin and a gliptin is initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of triple oral therapy with an SGLT2 inhibitor, metformin and a gliptin, must be documented in the patient's medical records.

empagliflozin 10 mg tablet, 30

11314L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	60.04	41.30	Jardiance [BY]

empagliflozin 25 mg tablet, 30

11281R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	60.04	41.30	Jardiance [BY]

■ EMPAGLIFLOZIN

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)

7506

Diabetes mellitus type 2

Clinical criteria:

- The treatment must be in combination with metformin; OR
- The treatment must be in combination with a sulfonylurea, **AND**
- Patient must have, or have had, a HbA1c measurement greater than 7% despite treatment with either metformin or a sulfonylurea; OR
- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period despite treatment with either metformin or a sulfonylurea. The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor is initiated. The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of triple oral therapy with a gliptin and an SGLT2 inhibitor, must be documented in the patient's medical records.

A patient whose diabetes was previously demonstrated unable to be controlled with metformin or a sulfonylurea does not need to requalify on this criterion before being eligible for PBS-subsidised treatment with this drug.

Note This drug is not PBS-subsidised for use as monotherapy or in combination with a thiazolidinedione (glitazone), or a glucagon-like peptide-1.

Authority required (STREAMLINED)

4991

Diabetes mellitus type 2

Clinical criteria:

- The treatment must be in combination with insulin, **AND**
- Patient must have, or have had, a HbA1c measurement greater than 7% prior to the initiation of a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor despite treatment with insulin and oral antidiabetic agents, or insulin alone where metformin is contraindicated; OR
- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor despite treatment with insulin and oral antidiabetic agents, or insulin alone where metformin is contraindicated.

The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or

(b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

Note This drug is not PBS-subsidised for use as monotherapy or in combination with a thiazolidinedione (glitazone), or a glucagon-like peptide-1.

Authority required (STREAMLINED)

5629

Diabetes mellitus type 2

Clinical criteria:

- The treatment must be in combination with metformin, **AND**
- The treatment must be in combination with a sulfonylurea, **AND**
- Patient must have, or have had, a HbA1c measurement greater than 7% prior to the initiation of a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor despite treatment with optimal doses of dual oral therapy; OR
- Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 despite treatment with optimal doses of dual oral therapy.

The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor, must be documented in the patient's medical records.

A patient whose diabetes was previously demonstrated unable to be controlled with metformin or a sulfonylurea does not need to requalify on this criterion before being eligible for PBS-subsidised treatment with this drug.

Note This drug is not PBS-subsidised for use as monotherapy or in combination with a thiazolidinedione (glitazone), or a glucagon-like peptide-1.

Note PBS subsidised dual oral therapy does not include concomitant use of a combination of: a gliptin, a glitazone or an SGLT2 inhibitor.

Authority required (STREAMLINED)

7495

Diabetes mellitus type 2

Treatment Phase: Continuing treatment

Clinical criteria:

- The treatment must be in combination with metformin, **AND**
- The treatment must be in combination with a dipeptidyl peptidase 4 inhibitor (gliptin), **AND**
- Patient must have previously received a PBS-subsidised regimen of oral diabetic medicines which included a sodium-glucose co-transporter 2 (SGLT2) inhibitor, metformin and a gliptin for this condition.

Note This drug is not PBS-subsidised for use as monotherapy or in combination with a thiazolidinedione (glitazone), or a glucagon-like peptide-1 analogue.

Note PBS-subsidised dual oral therapy does not include combination use of: a gliptin with an SGLT2 inhibitor; or a gliptin with a glitazone; or an SGLT2 inhibitor with a glitazone.

empagliflozin 10 mg tablet, 30

10206E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	60.04	41.30	Jardiance [BY]

empagliflozin 25 mg tablet, 30

10202Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	60.04	41.30	Jardiance [BY]

ERTUGLIFLOZIN

Note This drug is not PBS-subsidised for use as monotherapy or in combination with a thiazolidinedione (glitazone), insulin or a glucagon-like peptide-1 analogue.

Note PBS-subsidised dual oral therapy does not include combination use of: a gliptin with an SGLT2 inhibitor; or a gliptin with a glitazone; or an SGLT2 inhibitor with a glitazone.

Authority required (STREAMLINED)

7528

Diabetes mellitus type 2

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be in combination with metformin, **AND**
- The treatment must be in combination with a dipeptidyl peptidase 4 inhibitor (gliptin), **AND**
- Patient must have an HbA1c measurement greater than 7% despite treatment with dual oral combination therapy with metformin and a gliptin; OR
- Patient must have, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period prior to initiation of triple oral therapy with a sodium-glucose co-transporter 2 (SGLT2) inhibitor, metformin and a gliptin.

The date and level of the qualifying HbA1c measurement must be documented in the patient's medical records at the time triple oral therapy with an SGLT2 inhibitor, metformin and a gliptin is initiated.

The HbA1c must be no more than 4 months old at the time triple oral therapy with an SGLT2 inhibitor, metformin and a gliptin is initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of triple oral therapy with an SGLT2 inhibitor, metformin and a gliptin, must be documented in the patient's medical records.

ertugliflozin 5 mg tablet, 28

11577H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	59.20	41.30	Steglatro 5 [MK]

ertugliflozin 15 mg tablet, 28

11570Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	59.20	41.30	Steglatro 15 [MK]

▪ **ERTUGLIFLOZIN**

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)

7506

Diabetes mellitus type 2

Clinical criteria:

- The treatment must be in combination with metformin; OR
 - The treatment must be in combination with a sulfonylurea, **AND**
 - Patient must have, or have had, a HbA1c measurement greater than 7% despite treatment with either metformin or a sulfonylurea; OR
 - Patient must have, or have had, where HbA1c measurement is clinically inappropriate, blood glucose levels greater than 10 mmol per L in more than 20% of tests over a 2 week period despite treatment with either metformin or a sulfonylurea.
- The date and level of the qualifying HbA1c measurement must be, or must have been, documented in the patient's medical records at the time treatment with a dipeptidyl peptidase 4 inhibitor (gliptin), a thiazolidinedione (glitazone), a glucagon-like peptide-1 or a sodium-glucose co-transporter 2 (SGLT2) inhibitor is initiated.

The HbA1c must be no more than 4 months old at the time treatment with a gliptin, a glitazone, a glucagon-like peptide-1 or an SGLT2 inhibitor was initiated.

Blood glucose monitoring may be used as an alternative assessment to HbA1c levels in the following circumstances:

- (a) A clinical condition with reduced red blood cell survival, including haemolytic anaemias and haemoglobinopathies; and/or
- (b) Had red cell transfusion within the previous 3 months.

The results of the blood glucose monitoring, which must be no more than 4 months old at the time of initiation of triple oral therapy with a gliptin and an SGLT2 inhibitor, must be documented in the patient's medical records.

A patient whose diabetes was previously demonstrated unable to be controlled with metformin or a sulfonylurea does not need to requalify on this criterion before being eligible for PBS-subsidised treatment with this drug.

Note This drug is not PBS-subsidised for use as monotherapy or in combination with a thiazolidinedione (glitazone), insulin or a glucagon-like peptide-1.

Authority required (STREAMLINED)

7495

Diabetes mellitus type 2

Treatment Phase: Continuing treatment

Clinical criteria:

- The treatment must be in combination with metformin, **AND**
- The treatment must be in combination with a dipeptidyl peptidase 4 inhibitor (gliptin), **AND**
- Patient must have previously received a PBS-subsidised regimen of oral diabetic medicines which included a sodium-glucose co-transporter 2 (SGLT2) inhibitor, metformin and a gliptin for this condition.

Note This drug is not PBS-subsidised for use as monotherapy or in combination with a thiazolidinedione (glitazone), insulin or a glucagon-like peptide-1 analogue.

Note PBS-subsidised dual oral therapy does not include combination use of: a gliptin with an SGLT2 inhibitor; or

- a gliptin with a glitazone; or
- an SGLT2 inhibitor with a glitazone.

ertugliflozin 5 mg tablet, 28

11585R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	59.20	41.30	Steglatro 5 [MK]

ertugliflozin 15 mg tablet, 28

11571B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	59.20	41.30	Steglatro 15 [MK]

VITAMINS

VITAMIN A AND D, INCL. COMBINATIONS OF THE TWO

Vitamin D and analogues

CALCITRIOL

Authority required (STREAMLINED)

5401

Hypocalcaemia

Clinical criteria:

- The condition must be due to renal disease.

Authority required (STREAMLINED)

5255

Hypoparathyroidism

Authority required (STREAMLINED)

5089

Hypophosphataemic rickets

Authority required (STREAMLINED)

5114

Vitamin D-resistant rickets

Authority required (STREAMLINED)

5402

Established osteoporosis

Clinical criteria:

- Patient must have fracture due to minimal trauma.

The fracture must have been demonstrated radiologically and the year of plain x-ray or computed tomography (CT) scan or magnetic resonance imaging (MRI) scan must be documented in the patient's medical records when treatment is initiated.

A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or, a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

calcitriol 0.25 microgram capsule, 100

2502Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	3	..	28.14	29.43	^a APO-Calcitriol [TX] ^a Calcitriol AN [EA] ^a Kosteo [RW]	^a Calciprox [ER] ^a CALITROL [ED] ^a Sical [AF]
			^b 2.29	30.43	29.43	^a Rocaltrol [IX]	

VITAMIN B1, PLAIN AND IN COMBINATION WITH VITAMIN B6 AND B12

Vitamin B1, plain

THIAMINE

Authority required (STREAMLINED)

5139

Thiamine deficiency

Clinical criteria:

- The treatment must be for prophylaxis.

Population criteria:

- Patient must be an Aboriginal or a Torres Strait Islander person.

thiamine hydrochloride 100 mg tablet, 100

1070H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	16.05	17.34	Betavit [PP]

MINERAL SUPPLEMENTS

CALCIUM

Calcium

CALCIUM

Authority required (STREAMLINED)

4586

Hyperphosphataemia

Clinical criteria:

- The condition must be associated with chronic renal failure.

calcium carbonate 1.5 g (calcium 600 mg) tablet, 240

3117C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	23.16	24.45	Calci-Tab 600 [AE]

calcium carbonate 1.25 g (calcium 500 mg) chewable tablet, 120

11726E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	1	..	*27.88	29.17	Cal-500 [PP]

POTASSIUM

Potassium

POTASSIUM CHLORIDE

potassium chloride 600 mg (potassium 8 mmol) modified release tablet, 200

1841X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	23.85	25.14	Span-K [AS]

POTASSIUM CHLORIDE + POTASSIUM BICARBONATE + POTASSIUM CARBONATE

potassium chloride 595 mg + potassium bicarbonate 384 mg + potassium carbonate 152 mg (total potassium 14 mmol) effervescent tablet, 60

3012M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	24.38	25.67	Chlorvescent [AS]

OTHER MINERAL SUPPLEMENTS

Magnesium

MAGNESIUM

Authority required (STREAMLINED)

5506

Hypomagnesaemia

Population criteria:

- Patient must be an Aboriginal or a Torres Strait Islander person.

Authority required (STREAMLINED)

5466

Chronic renal disease

Population criteria:

- Patient must be an Aboriginal or a Torres Strait Islander person.

magnesium 37.4 mg tablet, 50

5146W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	18.04	19.33	MagMin (PBS) [BB]	Mag-Sup [PP]

OTHER ALIMENTARY TRACT AND METABOLISM PRODUCTS

OTHER ALIMENTARY TRACT AND METABOLISM PRODUCTS

Amino acids and derivatives

BETAINE

Authority required

Homocystinuria

Clinical criteria:

- The treatment must be as adjunctive therapy to current standard care, **AND**
 - The condition must be treated by or in consultation with a metabolic physician.
- The name of the specialist must be included in the authority application.

betaine 1 g/g powder for oral liquid, 180 g

10119N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	5	..	547.77	41.30	Cystadane [RJ]

*Various alimentary tract and metabolism products***■ SAPROPTERIN**

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Authority required

Hyperphenylalaninaemia (HPA) due to tetrahydrobiopterin (BH4) deficiency

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a metabolic physician; OR
- Must be treated by a nurse practitioner experienced in the treatment of phenylketonuria in consultation with a metabolic physician.

Clinical criteria:

- Patient must have hyperphenylalaninaemia (HPA) due to tetrahydrobiopterin (BH4) deficiency, **AND**

- Patient must have previously received PBS-subsidised treatment with this drug for this condition.

Patient must have documented tetrahydrobiopterin (BH4) deficiency using tests for BH4 loading and/or urine pterin metabolites, blood spot dihydropteridine reductase (DHPR) and have cerebrospinal fluid neurotransmitter metabolites measured.

sapropterin dihydrochloride 500 mg powder for oral liquid, 30 sachets

11970B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	4246.16	41.30	Kuvan [IO]

NP

■ SAPROPTERIN

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Authority required

Hyperphenylalaninaemia

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a metabolic physician; OR
- Must be treated by a nurse practitioner experienced in the treatment of phenylketonuria in consultation with a metabolic physician.

Clinical criteria:

- Patient must have hyperphenylalaninaemia (HPA) due to tetrahydrobiopterin (BH4) deficiency, **AND**

- Patient must have previously received PBS-subsidised treatment with this drug for this condition.

Patient must have documented tetrahydrobiopterin (BH4) deficiency using tests for BH4 loading and/or urine pterin metabolites, blood spot dihydropteridine reductase (DHPR) and have cerebrospinal fluid neurotransmitter metabolites measured.

sapropterin dihydrochloride 100 mg soluble tablet, 30

10087X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	6	5	..	*5063.16	41.30	Kuvan [IO]

NP

■ SAPROPTERIN

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Note Patient will be eligible for a maximum of one PBS-subsidised prescription as initial therapy to enable their response to treatment with sapropterin to be assessed.

Authority required

Hyperphenylalaninaemia (HPA) due to tetrahydrobiopterin (BH4) deficiency

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a metabolic physician.

Clinical criteria:

- Patient must have hyperphenylalaninaemia (HPA) due to tetrahydrobiopterin (BH4) deficiency.

Patient must have documented tetrahydrobiopterin (BH4) deficiency using tests for BH4 loading and/or urine pterin metabolites, blood spot dihydropteridine reductase (DHPR) and have cerebrospinal fluid neurotransmitter metabolites measured.

sapropterin dihydrochloride 500 mg powder for oral liquid, 30 sachets

11973E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	4246.16	41.30	Kuvan [IO]

■ SAPROPTERIN

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Note Patient will be eligible for a maximum of one PBS-subsidised prescription as initial therapy to enable their response to treatment with sapropterin to be assessed.

Authority required

Hyperphenylalaninaemia

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a metabolic physician.

Clinical criteria:

- Patient must have hyperphenylalaninaemia (HPA) due to tetrahydrobiopterin (BH4) deficiency.
- Patient must have documented tetrahydrobiopterin (BH4) deficiency using tests for BH4 loading and/or urine pterin metabolites, blood spot dihydropteridine reductase (DHPR) and have cerebrospinal fluid neurotransmitter metabolites measured.

sapropterin dihydrochloride 100 mg soluble tablet, 30

10086W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	6	..	..	*5063.16	41.30	Kuvan [IO]

■ SAPROPTERIN

Note Special Pricing Arrangements apply.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Note Patient will be eligible for a maximum of one PBS-subsidised prescription as initial therapy to enable their response to treatment with sapropterin to be assessed.

Authority required

Hyperphenylalaninaemia (HPA) due to phenylketonuria (PKU)

Treatment Phase: Initial treatment - responsiveness testing

Treatment criteria:

- Must be treated by a metabolic physician.

Clinical criteria:

- Patient must not have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have a baseline blood phenylalanine level above 360 micromole per L and be less than one month of age; OR
- Patient must have a baseline blood phenylalanine level above 600 micromole per L and be more than one month of age, **AND**
- The treatment must be for the purpose of initial responsiveness testing for a period of 24 hours in a patient less than one month of age; OR
- The treatment must be for the purpose of initial responsiveness testing for a period of 7 days in a patient aged more than one month.

Population criteria:

- Patient must be under 18 years of age.

Dietary phenylalanine intake must be maintained at a constant level.

Patients or their parent/guardian should be assessed for their ability to comply with the sapropterin protocol and PKU diet prior to conducting initial responsiveness testing.

sapropterin dihydrochloride 100 mg soluble tablet, 30

11676M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	..	..	*2585.07	41.30	Kuvan [IO]

sapropterin dihydrochloride 500 mg powder for oral liquid, 30 sachets

11971C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	4246.16	41.30	Kuvan [IO]

■ SAPROPTERIN

Note Special Pricing Arrangements apply.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Authority required

Hyperphenylalaninaemia (HPA) due to phenylketonuria (PKU)

Treatment Phase: First continuing treatment

Treatment criteria:

- Must be treated by a metabolic physician; OR
- Must be treated by a nurse practitioner experienced in the treatment of phenylketonuria in consultation with a metabolic physician.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment under the Initial treatment - responsiveness testing restriction with this drug for this condition, **AND**
- Patient must have demonstrated a response to treatment with this drug of greater than or equal to a 30% reduction in phenylalanine levels from baseline during initial responsiveness testing.

Population criteria:

- Patient must have been under 18 years of age at the time treatment with this drug was initiated for this condition. Blood phenylalanine levels must be based on measurements taken during stable periods of the condition. Dietary phenylalanine intake must be maintained at a constant level.

Authority required

Hyperphenylalaninaemia (HPA) due to phenylketonuria (PKU)

Treatment Phase: Subsequent continuing

Treatment criteria:

- Must be treated by a metabolic physician; OR
- Must be treated by a nurse practitioner experienced in the treatment of phenylketonuria in consultation with a metabolic physician.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must be undergoing regular phenylalanine testing and assessment of adherence to dietary modifications.

Population criteria:

- Patient must have been under 18 years of age at the time treatment with this drug was initiated for this condition.

sapropterin dihydrochloride 100 mg soluble tablet, 30

11691H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	6	5	..	*5063.16	41.30	Kuvan [IO]

sapropterin dihydrochloride 500 mg powder for oral liquid, 30 sachets

11983Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	4246.16	41.30	Kuvan [IO]

■ SODIUM PHENYLBUTYRATE**Authority required (STREAMLINED)****9993**

Urea cycle disorders

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have elevated ammonia levels that are not controlled with diet alone and other adjunct care alone. An increase in the maximum quantity will be authorised to provide for up to one month's supply at a dose of up to 600 mg/kg/day in patients weighing less than 20 kg and up to 13 g/m²/day in patients weighing more than 20 kg.

Authority required (STREAMLINED)**9919**

Urea cycle disorders

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition. An increase in the maximum quantity will be authorised to provide for up to one month's supply at a dose of up to 600 mg/kg/day in patients weighing less than 20 kg and up to 13 g/m²/day in patients weighing more than 20 kg.

Authority required (STREAMLINED)**9888**

Urea cycle disorders

Treatment Phase: Grandfathered treatment

Clinical criteria:

- Patient must have previously received non-PBS subsidised treatment with this drug for this condition prior to 1 December 2019.

An increase in the maximum quantity will be authorised to provide for up to one month's supply at a dose of up to 600 mg/kg/day in patients weighing less than 20 kg and up to 13 g/m²/day in patients weighing more than 20 kg.

sodium phenylbutyrate 483 mg/g granules, 174 g

11865L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	3	5	..	*1638.87	41.30	Pheburane [OH]

■ BLOOD AND BLOOD FORMING ORGANS**■ ANTITHROMBOTIC AGENTS****ANTITHROMBOTIC AGENTS***Vitamin K antagonists*

■ WARFARIN

Caution The listed brands have NOT been shown to be bioequivalent and should not be interchanged.

warfarin sodium 2 mg tablet, 50

2209G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	16.96	18.25	Coumadin [GO]

warfarin sodium 3 mg tablet, 50

2844Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	17.02	18.31	Marevan [GT]

warfarin sodium 1 mg tablet, 50

2843P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	2	..	16.78	18.07	Coumadin [GO]	Marevan [GT]

warfarin sodium 5 mg tablet, 50

2211J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	2	..	17.43	18.72	Coumadin [GO]	Marevan [GT]

Heparin group

■ DALTEPARIN SODIUM

dalteparin sodium 5000 anti-Xa units/0.2 mL injection, 10 x 0.2 mL syringes

2816F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	..	..	*87.64	41.30	Fragmin [PF]

dalteparin sodium 2500 anti-Xa units/0.2 mL injection, 10 x 0.2 mL syringes

8603T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	..	..	*84.62	41.30	Fragmin [PF]

dalteparin sodium 7500 anti-Xa units/0.75 mL injection, 10 x 0.75 mL syringes

8271H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	57.77	41.30	Fragmin [PF]

dalteparin sodium 10 000 anti-Xa units/mL injection, 10 x 1 mL syringes

8269F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	74.47	41.30	Fragmin [PF]

dalteparin sodium 12 500 anti-Xa units/0.5 mL injection, 10 x 0.5 mL syringes

5445N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	100.10	41.30	Fragmin [PF]

■ DALTEPARIN SODIUM

Restricted benefit

Haemodialysis

dalteparin sodium 5000 anti-Xa units/0.2 mL injection, 10 x 0.2 mL syringes

8642W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	3	..	*87.64	41.30	Fragmin [PF]

dalteparin sodium 2500 anti-Xa units/0.2 mL injection, 10 x 0.2 mL syringes

8641T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	3	..	*84.62	41.30	Fragmin [PF]

dalteparin sodium 7500 anti-Xa units/0.75 mL injection, 10 x 0.75 mL syringes

8643X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	3	..	*103.52	41.30	Fragmin [PF]

dalteparin sodium 10 000 anti-Xa units/mL injection, 10 x 1 mL syringes

1229Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	3	..	*138.16	41.30	Fragmin [PF]

dalteparin sodium 12 500 anti-Xa units/0.5 mL injection, 10 x 0.5 mL syringes

1296F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	3	..	*191.98	41.30	Fragmin [PF]

■ DALTEPARIN SODIUM

Note No applications for increased maximum quantities will be authorised.

Restricted benefit

Symptomatic venous thromboembolism

Treatment Phase: Management

Clinical criteria:

- Patient must have a solid tumour(s).

dalteparin sodium 7500 anti-Xa units/0.75 mL injection, 10 x 0.75 mL syringes

8956J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	3	5	..	*151.14	41.30	Fragmin [PF]

dalteparin sodium 18 000 anti-Xa units/0.72 mL injection, 10 x 0.72 mL syringes

8960N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	3	5	..	*406.53	41.30	Fragmin [PF]

dalteparin sodium 10 000 anti-Xa units/mL injection, 10 x 1 mL syringes

8957K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	3	5	..	*203.73	41.30	Fragmin [PF]

dalteparin sodium 12 500 anti-Xa units/0.5 mL injection, 10 x 0.5 mL syringes

8958L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	3	5	..	*284.46	41.30	Fragmin [PF]

dalteparin sodium 15 000 anti-Xa units/0.6 mL injection, 10 x 0.6 mL syringes

8959M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	3	5	..	*339.90	41.30	Fragmin [PF]

▪ ENOXAPARIN SODIUM**enoxaparin sodium 60 mg/0.6 mL injection, 10 x 0.6 mL syringes**

8262W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	65.17	41.30	Clexane Safety-Lock [AV]

enoxaparin sodium 100 mg/mL injection, 10 x 1 mL syringes

8264Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	86.50	41.30	Clexane Safety-Lock [AV]

enoxaparin sodium 20 mg/0.2 mL injection, 10 x 0.2 mL syringes

8558K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	1	..	*83.32	41.30	Clexane Safety-Lock [AV]

enoxaparin sodium 80 mg/0.8 mL injection, 10 x 0.8 mL syringes

8263X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	73.17	41.30	Clexane Safety-Lock [AV]

enoxaparin sodium 40 mg/0.4 mL injection, 10 x 0.4 mL syringes

8510X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	1	..	*86.34	41.30	Clexane Safety-Lock [AV]

▪ ENOXAPARIN SODIUM**Restricted benefit**

Haemodialysis

enoxaparin sodium 60 mg/0.6 mL injection, 10 x 0.6 mL syringes

8640R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	3	..	*118.64	41.30	Clexane Safety-Lock [AV]

enoxaparin sodium 100 mg/mL injection, 10 x 1 mL syringes

5435C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	3	..	*163.42	41.30	Clexane Safety-Lock [AV]

enoxaparin sodium 20 mg/0.2 mL injection, 10 x 0.2 mL syringes

8716R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	3	..	*83.32	41.30	Clexane Safety-Lock [AV]

enoxaparin sodium 80 mg/0.8 mL injection, 10 x 0.8 mL syringes

5434B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	3	..	*135.44	41.30	Clexane Safety-Lock [AV]

enoxaparin sodium 40 mg/0.4 mL injection, 10 x 0.4 mL syringes

8639Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	3	..	*86.34	41.30	Clexane Safety-Lock [AV]

■ HEPARIN**heparin sodium 5000 units/0.2 mL injection, 5 x 0.2 mL ampoules**

1466E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	24.34	25.63	DBL Heparin Sodium [PF]

heparin sodium 5000 units/5 mL injection, 50 x 5 mL ampoules

1463B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	73.24	41.30	Pfizer Australia Pty Ltd [PF]

Platelet aggregation inhibitors excl. heparin**■ ASPIRIN****Restricted benefit**

For treatment of a patient identifying as Aboriginal or Torres Strait Islander

aspirin 100 mg tablet, 112

8202Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	14.14	15.43	Spren 100 [OW]

■ CLOPIDOGREL

Note Not for prophylaxis of deep vein thrombosis or peripheral arterial disease.

Note Pharmaceutical benefits that have the forms clopidogrel tablet 75 mg (as besilate) and clopidogrel tablet 75 mg (as hydrogen sulfate) are equivalent for the purposes of substitution.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**4166**

Acute coronary syndrome (myocardial infarction or unstable angina)

Clinical criteria:

- The treatment must be in combination with aspirin.

Authority required (STREAMLINED)**4165**

Cardiac stent insertion

Clinical criteria:

- The treatment must be in combination with aspirin, **AND**
- The treatment must follow insertion of a cardiac stent.

clopidogrel 75 mg tablet, 28

2275R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	16.23	17.52	^a BTC Clopidogrel [JB] ^a Clopidogrel-GA [EA] ^a Clovix 75 [RW]	^a Clopidogrel APOTEX [GX] ^a Clopidogrel GH [GQ] ^a Plidogrel [RF]

clopidogrel 75 mg tablet, 28

9317J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	16.23	17.52	^a APO-Clopidogrel [TX] ^a Clopidogrel AN [EA] ^a Clopidogrel Sandoz Pharma [HX] ^a Iscover [AV]	^a Blooms the Chemist Clopidogrel [IB] ^a Clopidogrel Sandoz [SZ] ^a Clopidogrel Winthrop [WA] ^a Piax [AF]

■ CLOPIDOGREL

Note Pharmaceutical benefits that have the forms clopidogrel tablet 75 mg (as besilate) and clopidogrel tablet 75 mg (as hydrogen sulfate) are equivalent for the purposes of substitution.

Note Not for prophylaxis of deep vein thrombosis or peripheral arterial disease.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**5517**

Prevention of recurrence of myocardial infarction or unstable angina

Clinical criteria:

- Patient must have a history of symptomatic cardiac ischaemic events while on therapy with low-dose aspirin.

Authority required (STREAMLINED)**5524**

Prevention of recurrence of myocardial infarction or unstable angina

Clinical criteria:

- Patient must be in one whom low-dose aspirin poses an unacceptable risk of gastrointestinal bleeding.

Authority required (STREAMLINED)**5525**

Prevention of recurrence of myocardial infarction or unstable angina

Clinical criteria:

- Patient must have a history of anaphylaxis, urticaria or asthma within 4 hours of ingestion of aspirin, other salicylates, or non-steroidal anti-inflammatory drugs (NSAIDs).

Authority required (STREAMLINED)**5459**

Prevention of recurrence of ischaemic stroke or transient cerebral ischaemic events

Clinical criteria:

- Patient must have a history of symptomatic cerebrovascular ischaemic episodes while on therapy with low-dose aspirin.

Authority required (STREAMLINED)**5436**

Prevention of recurrence of ischaemic stroke or transient cerebral ischaemic events

Clinical criteria:

- Patient must be in one whom low-dose aspirin poses an unacceptable risk of gastrointestinal bleeding.

Authority required (STREAMLINED)**5508**

Prevention of recurrence of ischaemic stroke or transient cerebral ischaemic events

Clinical criteria:

- Patient must have a history of anaphylaxis, urticaria or asthma within 4 hours of ingestion of aspirin, other salicylates, or non-steroidal anti-inflammatory drugs (NSAIDs).

clopidogrel 75 mg tablet, 28

8358X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	16.23	17.52	^a APO-Clopidogrel [TX]	^a Blooms the Chemist Clopidogrel [IB]
						^a Clopidogrel AN [EA]	^a Clopidogrel Sandoz [SZ]
						^a Clopidogrel Sandoz Pharma [HX]	^a Clopidogrel Winthrop [WA]
						^a Iscover [AV]	^a Piax [AF]
						^a Plavacor 75 [CR]	

clopidogrel 75 mg tablet, 28

9354H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	16.23	17.52	^a BTC Clopidogrel [JB]	^a Clopidogrel APOTEX [GX]
						^a Clopidogrel-GA [EA]	^a Clopidogrel GH [GQ]
						^a Clovis 75 [RW]	^a Plidogrel [RF]

■ CLOPIDOGREL + ASPIRIN**Note** Not for prophylaxis of deep vein thrombosis or peripheral arterial disease.**Note Shared Care Model:**

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**5488**

Acute coronary syndrome (myocardial infarction or unstable angina)

Authority required (STREAMLINED)**5443**

Cardiac stent insertion

Clinical criteria:

- The treatment must follow insertion of a cardiac stent.

Authority required (STREAMLINED)**5517**

Prevention of recurrence of myocardial infarction or unstable angina

Clinical criteria:

- Patient must have a history of symptomatic cardiac ischaemic events while on therapy with low-dose aspirin.

clopidogrel 75 mg + aspirin 100 mg tablet, 30

8296G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	16.50	17.79	^a APO-Clopidogrel/Aspirin 75/100 [TX]	^a APX-Clopidogrel/Aspirin 75/100 [TY]
						^a Clopidogrel/Aspirin Actavis 75/100 [EA]	^a CLOPIDOGREL/ASPIRIN AN 75/100 [ED]
						^a Clopidogrel Winthrop plus aspirin [WA]	^a CoPlavix [SW]
						^a DuoCover [AV]	^a DuoPlidogrel [GZ]
						^a Piax Plus Aspirin [AF]	

■ DIPYRIDAMOLE + ASPIRIN**Note Shared Care Model:**

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Prevention of recurrence of ischaemic stroke or transient cerebral ischaemic events

dipyridamole 200 mg + aspirin 25 mg modified release capsule, 60

8382E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	34.50	35.79	Diasp SR [RW]

■ EPTIFIBATIDE**Authority required (STREAMLINED)****6435**

Coronary artery disease

Treatment criteria:

- Patient must be undergoing non-urgent percutaneous intervention with intracoronary stenting.

eptifibatide 20 mg/10 mL injection, 10 mL vial

8683B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*210.10	41.30	Integrilin [MK]

eptifibatide 75 mg/100 mL injection, 100 mL vial

8684C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	..	..	*837.24	41.30	Integrilin [MK]

■ TICAGRELOR**Note Shared Care Model:**

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**5746**

Acute coronary syndrome (myocardial infarction or unstable angina)

Clinical criteria:

- The treatment must be in combination with aspirin.

ticagrelor 90 mg tablet, 56

1418P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	136.31	41.30	Brilinta [AP]

■ TIROFIBAN**Note Shared Care Model:**

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**5782**

High risk of unstable angina

Clinical criteria:

- Patient must have new transient or persistent ST-T ischaemic changes, **AND**
- Patient must have pain lasting longer than 20 minutes.

Authority required (STREAMLINED)**5809**

High risk of unstable angina

Clinical criteria:

- Patient must have new transient or persistent ST-T ischaemic changes, **AND**
- Patient must have repetitive episodes of angina at rest or during minimal exercise in the previous 12 hours.

Authority required (STREAMLINED)**5691**

Non-Q-wave myocardial infarction

tirofiban 12.5 mg/50 mL injection, 50 mL vial

8350L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	2	..	213.05	41.30	^a Aggrastat [AS] ^a Tirofiban Juno [JU]	^a Tirofiban AC [JO]

Enzymes**■ RETEPLASE****Note Shared Care Model:**

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Acute myocardial infarction

Clinical criteria:

- The treatment must be administered within 6 hours of the onset of attack.

reteplase 10 units (17.4 mg) injection [2 x 10 unit vials] (&) inert substance diluent [2 x 10 mL syringes], 1 pack

8253J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	..	..	1701.88	41.30	Rapilysin 10 U [TB]

■ TENECTEPLASE**Note Shared Care Model:**

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Acute myocardial infarction

Clinical criteria:

- The treatment must be administered within 12 hours of onset of attack.

tenecteplase 50 mg injection [1 vial] (&) inert substance diluent [10 mL syringe], 1 pack

8527T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	1969.80	41.30	Metalyse [BY]

tenecteplase 40 mg injection [1 vial] (&) inert substance diluent [8 mL syringe], 1 pack

8526R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	1873.73	41.30	Metalyse [BY]

Direct thrombin inhibitors**■ BIVALIRUDIN****Authority required (STREAMLINED)****4919**

Coronary artery disease

Treatment criteria:

- Patient must be undergoing percutaneous coronary intervention.

bivalirudin 250 mg injection, 1 vial

8844L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	474.28	41.30	^a Angiomax [XM]	^a Bivalirudin APOTEX [TX]

■ DABIGATRAN

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**4402**

Prevention of venous thromboembolism

Treatment criteria:

- Patient must be undergoing total hip replacement.

Clinical criteria:

- Patient must require up to 30 days supply to complete a course of treatment.

dabigatran etexilate 75 mg capsule, 60

9320M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	100.85	41.30	Pradaxa [BY]

dabigatran etexilate 110 mg capsule, 60

9321N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	81.56	41.30	Pradaxa [BY]

■ DABIGATRAN

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**4369**

Prevention of venous thromboembolism

Treatment criteria:

- Patient must be undergoing total hip replacement.

Clinical criteria:

- Patient must require up to 20 days supply to complete a course of treatment.

dabigatran etexilate 75 mg capsule, 10

9318K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	1	..	*41.64	41.30	Pradaxa [BY]

dabigatran etexilate 110 mg capsule, 10

9319L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	1	..	*35.20	36.49	Pradaxa [BY]

■ DABIGATRAN

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**4381**

Prevention of venous thromboembolism

Treatment criteria:

- Patient must be undergoing total knee replacement.

Clinical criteria:

- Patient must require up to 10 days of therapy.

dabigatran etexilate 75 mg capsule, 10

9322P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	..	..	*41.64	41.30	Pradaxa [BY]

dabigatran etexilate 110 mg capsule, 10

9323Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	..	..	*35.20	36.49	Pradaxa [BY]

■ DABIGATRAN

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)**4269**

Prevention of stroke or systemic embolism

Clinical criteria:

- Patient must have non-valvular atrial fibrillation, **AND**
- Patient must have one or more risk factors for developing stroke or systemic embolism.

Risk factors for developing stroke or systemic ischaemic embolism are:

- Prior stroke (ischaemic or unknown type), transient ischaemic attack or non-central nervous system (CNS) systemic embolism;
- age 75 years or older;
- hypertension;
- diabetes mellitus;
- heart failure and/or left ventricular ejection fraction 35% or less.

dabigatran etexilate 150 mg capsule, 60

2769R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	81.57	41.30	Pradaxa [BY]

dabigatran etexilate 110 mg capsule, 60

2753X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	81.56	41.30	Pradaxa [BY]

Direct factor Xa inhibitors

■ APIXABAN

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)

4402

Prevention of venous thromboembolism

Treatment criteria:

- Patient must be undergoing total hip replacement.

Clinical criteria:

- Patient must require up to 30 days supply to complete a course of treatment.

apixaban 2.5 mg tablet, 60

5061J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	93.85	41.30	Eliquis [BQ]

■ APIXABAN

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

4098

Deep vein thrombosis

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have confirmed acute symptomatic deep vein thrombosis, **AND**
- Patient must not have symptomatic pulmonary embolism.

Authority required (STREAMLINED)

5098

Pulmonary embolism

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have confirmed acute symptomatic pulmonary embolism.

apixaban 5 mg tablet, 28

10414D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	50.21	41.30	Eliquis [BQ]

■ APIXABAN

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical

practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)

4382

Prevention of venous thromboembolism

Treatment criteria:

- Patient must be undergoing total knee replacement.

Clinical criteria:

- Patient must require up to 15 days of therapy.

Authority required (STREAMLINED)

4409

Prevention of venous thromboembolism

Treatment criteria:

- Patient must be undergoing total hip replacement.

Clinical criteria:

- Patient must require up to 15 days supply to complete a course of treatment.

apixaban 2.5 mg tablet, 30

5054B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	52.94	41.30	Eliquis [BQ]

▪ **APIXABAN**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)

4381

Prevention of venous thromboembolism

Treatment criteria:

- Patient must be undergoing total knee replacement.

Clinical criteria:

- Patient must require up to 10 days of therapy.

Authority required (STREAMLINED)

4359

Prevention of venous thromboembolism

Treatment criteria:

- Patient must be undergoing total hip replacement.

Clinical criteria:

- Patient must require up to 10 days supply to complete a course of treatment.

apixaban 2.5 mg tablet, 20

5500L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	39.30	40.59	Eliquis [BQ]

▪ **APIXABAN**

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

4269

Prevention of stroke or systemic embolism

Clinical criteria:

- Patient must have non-valvular atrial fibrillation, **AND**
- Patient must have one or more risk factors for developing stroke or systemic embolism.

Risk factors for developing stroke or systemic ischaemic embolism are:

- Prior stroke (ischaemic or unknown type), transient ischaemic attack or non-central nervous system (CNS) systemic embolism;
- age 75 years or older;
- hypertension;
- diabetes mellitus;
- heart failure and/or left ventricular ejection fraction 35% or less.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

4132

Prevention of recurrent venous thromboembolism

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have a history of venous thromboembolism.

apixaban 2.5 mg tablet, 60

2744K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	93.85	41.30	Eliquis [BQ]

▪ **APIXABAN**

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

4269

Prevention of stroke or systemic embolism

Clinical criteria:

- Patient must have non-valvular atrial fibrillation, **AND**
- Patient must have one or more risk factors for developing stroke or systemic embolism.

Risk factors for developing stroke or systemic ischaemic embolism are:

- Prior stroke (ischaemic or unknown type), transient ischaemic attack or non-central nervous system (CNS) systemic embolism;
- age 75 years or older;
- hypertension;
- diabetes mellitus;
- heart failure and/or left ventricular ejection fraction 35% or less.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

4099

Deep vein thrombosis

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have confirmed acute symptomatic deep vein thrombosis, **AND**
- Patient must not have symptomatic pulmonary embolism.

Authority required (STREAMLINED)

5083

Pulmonary embolism

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have confirmed acute symptomatic pulmonary embolism.

apixaban 5 mg tablet, 60

2735Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	93.85	41.30	Eliquis [BQ]

▪ **RIVAROXABAN**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

11013

Chronic stable atherosclerotic disease

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be in combination with aspirin, but not with any other anti-platelet therapy, **AND**
- Patient must have a diagnosis of coronary artery disease in addition to at least one of the following risk factors: (i) diagnosed heart failure (left ventricular ejection fraction of at least 30% but less than 50%) (ii) diagnosed kidney disease classified by an eGFR in the range of 15-60 mL/min (iii) diabetes mellitus combined with at least one of the following: (a) age at least 60 years (b) concomitant microalbuminuria (c) Aboriginal/Torres Strait Islander descent; OR
- Patient must have a diagnosis of peripheral artery disease in addition to at least one of the following risk factors: (i) concomitant coronary artery disease (ii) diagnosed heart failure (left ventricular ejection fraction of at least 30% but less

than 50%) (iii) diagnosed kidney disease classified by an eGFR in the range of 15-60 mL/min (iv) diabetes mellitus combined with at least one of the following: (a) age at least 60 years (b) concomitant microalbuminuria (c) Aboriginal/Torres Strait Islander descent, **AND**

- Patient must have at least one of the following if coronary artery disease is present: (i) a previous multi-vessel coronary revascularisation procedure (ii) significant stenosis in at least 2 coronary arteries (iii) a previous single vessel coronary revascularisation procedure with significant stenosis in more than 1 coronary artery; OR
- Patient must have at least one of the following if peripheral arterial disease is present: (i) a previous peripheral/carotid artery revascularisation intervention (ii) intermittent claudication with an ankle-brachial index less than 0.9 (iii) asymptomatic carotid artery stenosis greater than 50%, **AND**
- The condition must be diagnosed by at least one of: (i) invasive (selective) angiography (ii) non-invasive imaging (i.e. CT scan, ultrasound) (iii) ankle-brachial index measurement in the case of peripheral arterial disease with intermittent claudication, **AND**
- Patient must have clinical findings/observations by the treating physician that exclude each of the following: (i) high risk of bleeding (ii) prior stroke within one month of treatment initiation (iii) prior haemorrhagic / lacunar stroke (iv) severe heart failure with a known ejection fraction less than 30% (v) New York Heart Association class III to IV heart failure symptoms (i.e. symptoms corresponding to moderate to severe limitation on physical activity, whereby any of fatigue/palpitations/dyspnoea occur upon zero to minimal activity) (vi) an estimated glomerular filtration rate less than 15 mL/minute (vii) a requirement for dual antiplatelet therapy (viii) a requirement for non-acetylsalicylic acid antiplatelet therapy (ix) a requirement for a higher dose of oral anticoagulant therapy.

Treatment criteria:

- Must be treated by a specialist physician; OR
- Must be treated by a physician who has consulted a specialist physician.

rivaroxaban 2.5 mg tablet, 60

12197Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	85.38	41.30	Xarelto [BN]

■ **RIVAROXABAN**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)

4369

Prevention of venous thromboembolism

Treatment criteria:

- Patient must be undergoing total hip replacement.

Clinical criteria:

- Patient must require up to 20 days supply to complete a course of treatment.

rivaroxaban 10 mg tablet, 10

9465E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	36.47	37.76	Xarelto [BN]

NP

■ **RIVAROXABAN**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)

4402

Prevention of venous thromboembolism

Treatment criteria:

- Patient must be undergoing total hip replacement.

Clinical criteria:

- Patient must require up to 30 days supply to complete a course of treatment.

rivaroxaban 10 mg tablet, 30

9467G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	85.37	41.30	Xarelto [BN]

NP

rivaroxaban 10 mg tablet, 15

9466F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	48.70	41.30	Xarelto [BN]

NP

■ RIVAROXABAN

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)

4381

Prevention of venous thromboembolism

Treatment criteria:

- Patient must be undergoing total knee replacement.

Clinical criteria:

- Patient must require up to 10 days of therapy.

rivaroxaban 10 mg tablet, 10

9468H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	36.47	37.76	Xarelto [BN]

■ RIVAROXABAN

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)

4382

Prevention of venous thromboembolism

Treatment criteria:

- Patient must be undergoing total knee replacement.

Clinical criteria:

- Patient must require up to 15 days of therapy.

rivaroxaban 10 mg tablet, 15

9469J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	48.70	41.30	Xarelto [BN]

■ RIVAROXABAN

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

4132

Prevention of recurrent venous thromboembolism

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have a history of venous thromboembolism.

rivaroxaban 10 mg tablet, 30

11633G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	85.37	41.30	Xarelto [BN]

■ RIVAROXABAN

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

4269

Prevention of stroke or systemic embolism

Clinical criteria:

- Patient must have non-valvular atrial fibrillation, **AND**
 - Patient must have one or more risk factors for developing stroke or systemic embolism.
- Risk factors for developing stroke or systemic ischaemic embolism are:
- (i) Prior stroke (ischaemic or unknown type), transient ischaemic attack or non-central nervous system (CNS) systemic embolism;
 - (ii) age 75 years or older;
 - (iii) hypertension;
 - (iv) diabetes mellitus;
 - (v) heart failure and/or left ventricular ejection fraction 35% or less.

rivaroxaban 15 mg tablet, 28

2691P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	80.49	41.30	Xarelto [BN]

■ RIVAROXABAN**Note Shared Care Model:**

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Treatment may be continued by a non-specialist prescriber without need for consultation with a specialist.

Authority required (STREAMLINED)**10992**

Chronic stable atherosclerotic disease

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be in combination with aspirin, but not with any other anti-platelet therapy.

rivaroxaban 2.5 mg tablet, 60

12192Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	85.38	41.30	Xarelto [BN]

■ RIVAROXABAN**Note Shared Care Model:**

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)**4098**

Deep vein thrombosis

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have confirmed acute symptomatic deep vein thrombosis, **AND**
- Patient must not have symptomatic pulmonary embolism.

Authority required (STREAMLINED)**4260**

Pulmonary embolism

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have confirmed acute symptomatic pulmonary embolism.

Note Special Pricing Arrangements apply.

rivaroxaban 15 mg tablet, 42

2160Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	114.86	41.30	Xarelto [BN]

■ RIVAROXABAN**Note Shared Care Model:**

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

4099

Deep vein thrombosis

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have confirmed acute symptomatic deep vein thrombosis, **AND**
- Patient must not have symptomatic pulmonary embolism.

Authority required (STREAMLINED)

4132

Prevention of recurrent venous thromboembolism

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have a history of venous thromboembolism.

Authority required (STREAMLINED)

4268

Pulmonary embolism

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have confirmed acute symptomatic pulmonary embolism.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

4269

Prevention of stroke or systemic embolism

Clinical criteria:

- Patient must have non-valvular atrial fibrillation, **AND**
- Patient must have one or more risk factors for developing stroke or systemic embolism.

Risk factors for developing stroke or systemic ischaemic embolism are:

- (i) Prior stroke (ischaemic or unknown type), transient ischaemic attack or non-central nervous system (CNS) systemic embolism;
- (ii) age 75 years or older;
- (iii) hypertension;
- (iv) diabetes mellitus;
- (v) heart failure and/or left ventricular ejection fraction 35% or less.

Note Special Pricing Arrangements apply.

rivaroxaban 20 mg tablet, 28

2268J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	80.49	41.30	Xarelto [BN]

Other antithrombotic agents

▪ **FONDAPARINUX**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)

5781

Prevention of venous thromboembolism

Treatment criteria:

- Patient must be undergoing major hip surgery.

Authority required (STREAMLINED)

5808

Prevention of venous thromboembolism

Treatment criteria:

- Patient must be undergoing total knee replacement.

fondaparinux sodium 2.5 mg/0.5 mL injection, 2 x 0.5 mL syringes

8775W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	3.5	..	..	*116.41	41.30	Arixtra [AS]

■ ANTIHEMORRHAGICS

ANTIFIBRINOLYTICS

Amino acids

■ TRANEXAMIC ACID

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

tranexamic acid 500 mg tablet, 100

2180R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	2	..	37.65	38.94	^a APO-Tranexamic Acid [TX]	^a Cyklokapron [PF]

■ ANTIANEMIC PREPARATIONS

IRON PREPARATIONS

Iron bivalent, oral preparations

■ FERROUS FUMARATE

Restricted benefit

For treatment of a patient identifying as Aboriginal or Torres Strait Islander

ferrous fumarate 200 mg (iron 65.7 mg) tablet, 60

8985X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	18.69	19.98	Ferro-tab [AE]

■ FERROUS SULFATE

ferrous sulfate heptahydrate 30 mg/mL (iron 6 mg/mL) oral liquid, 250 mL

8815Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	‡1	2	..	21.15	22.44	Ferro-Liquid [AE]

Iron, parenteral preparations

■ FERRIC CARBOXYMALTOSE

Note Special Pricing Arrangements apply.

iron (as ferric carboxymaltose) 500 mg/10 mL injection, 10 mL vial

10104T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	1	..	*298.76	41.30	Ferinject [VL]

iron (as ferric carboxymaltose) 1 g/20 mL injection, 20 mL vial

11702X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	298.74	41.30	Ferinject [VL]

■ FERRIC DERISOMALTOSE

Note Special Pricing Arrangements apply.

iron (as ferric derisomaltose) 1 g/10 mL injection, 10 mL vial

12049E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	298.74	41.30	Monofer [PF]

iron (as ferric derisomaltose) 500 mg/5 mL injection, 5 mL vial

11615H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	3	..	..	*444.63	41.30	Monofer [PF]

■ IRON POLYMALTOSE

iron (as polymaltose) 100 mg/2 mL injection, 5 x 2 mL ampoules

2593L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	31.26	32.55	Ferrosig [SI]

■ IRON POLYMALTOSE

Authority required (STREAMLINED)

4302

Iron deficiency anaemia

Treatment criteria:

- Patient must be undergoing chronic haemodialysis.

BLOOD AND BLOOD FORMING ORGANS

iron (as polymaltose) 100 mg/2 mL injection, 5 x 2 mL ampoules

2805P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	31.26	32.55	Ferrosig [SI]

■ IRON SUCROSE

iron (as sucrose) 100 mg/5 mL injection, 5 x 5 mL ampoules

10229J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	38.09	39.38	Venofer [VL]

■ IRON SUCROSE

Authority required (STREAMLINED)

4302

Iron deficiency anaemia

Treatment criteria:

- Patient must be undergoing chronic haemodialysis.

iron (as sucrose) 100 mg/5 mL injection, 5 x 5 mL ampoules

8807M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	38.09	39.38	Venofer [VL]

Iron in combination with folic acid

■ FERROUS FUMARATE + FOLIC ACID

Restricted benefit

For treatment of a patient identifying as Aboriginal or Torres Strait Islander

ferrous fumarate 310 mg (iron 100 mg) + folic acid 350 microgram tablet, 60

9011G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	19.71	21.00	Ferro-f-tab [AE]

VITAMIN B12 AND FOLIC ACID

Vitamin B12 (cyanocobalamin and analogues)

■ HYDROXOCOBALAMIN

Note One injection of hydroxocobalamin 1 mg every three months provides appropriate maintenance therapy in vitamin B₁₂ deficiencies.

Note Pharmaceutical benefits that have the form hydroxocobalamin injection 1 mg (as acetate) in 1 mL and pharmaceutical benefits that have the form hydroxocobalamin injection 1 mg (as chloride) in 1 mL are equivalent for the purposes of substitution.

Restricted benefit

Pernicious anaemia

Population criteria:

- Patient must identify as Aboriginal or Torres Strait Islander.

Restricted benefit

Proven vitamin B12 deficiencies other than pernicious anaemia

Population criteria:

- Patient must identify as Aboriginal or Torres Strait Islander.

Restricted benefit

Anaemias associated with vitamin B12 deficiency

Clinical criteria:

- Patient must have had a gastrectomy, **AND**
- The treatment must be for prophylaxis.

Population criteria:

- Patient must identify as Aboriginal or Torres Strait Islander.

hydroxocobalamin 1 mg/mL injection, 3 x 1 mL ampoules

2162T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	16.40	17.69	^a Cobal-B12 [JU]	^a Vita-B12 [GH]

hydroxocobalamin 1 mg/mL injection, 3 x 1 mL ampoules

9048F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	16.40	17.69	^a Hydroxo-B12 [AS]	^a Neo-B12 [PF]

Folic acid and derivatives

■ FOLIC ACID

Restricted benefit

For treatment of a patient identifying as Aboriginal or Torres Strait Islander

folic acid 500 microgram tablet, 100

2958Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	2	..	..	*16.22	17.51	^a Foltabs 500 [PP]	^a Megafol 0.5 [AF]

■ FOLIC ACID

Note The 5 mg strength tablet should be used in malabsorption states only.

Restricted benefit

For treatment of a patient identifying as Aboriginal or Torres Strait Islander

folic acid 5 mg tablet, 100

1437P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	1	..	*18.30	19.59	Megafol 5 [AF]

■ BLOOD SUBSTITUTES AND PERFUSION SOLUTIONS**BLOOD AND RELATED PRODUCTS**

Blood substitutes and plasma protein fractions

■ HYDROXYETHYL STARCH 130/0.4 + SODIUM CHLORIDE**HYDROXYETHYL STARCH 130/0.4 I.V. infusion 30 g per 500 mL, 500 mL, 1**

9487H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	3	..	..	*41.79	41.30	Voluven 6% [PK]

■ OTHER HEMATOLOGICAL AGENTS**OTHER HEMATOLOGICAL AGENTS**

Drugs used in hereditary angioedema

■ ICATIBANT

Note Icatibant should be provided in the framework of a comprehensive hereditary angioedema prophylaxis program and an emergency Action Plan including training in recognition of the symptoms of hereditary angioedema and the self-administration of icatibant. (For further information see the Australasian Society of Clinical Immunology and Allergy website at www.allergy.org.au)

Authority required

Anticipated emergency treatment of an acute attack of hereditary angioedema

Treatment Phase: Initial

Clinical criteria:

- Patient must have confirmed diagnosis of C1-esterase inhibitor deficiency, **AND**
- Patient must have been assessed to be at significant risk of an acute attack of hereditary angioedema, **AND**
- The condition must be assessed by a clinical immunologist; OR
- The condition must be assessed by a respiratory physician; OR
- The condition must be assessed by a specialist allergist; OR
- The condition must be assessed by a general physician experienced in the management of patients with hereditary angioedema.

The name of the specialist consulted must be provided at the time of application for initial supply.

The date of the pathology report and name of the Approved Pathology Authority must be provided at the time of application.

Increased maximum quantities will be limited to 12 injections per authority prescription.

Authority required

Anticipated emergency treatment of an acute attack of hereditary angioedema

Treatment Phase: Continuing

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition.
- Increased maximum quantities will be limited to 12 injections per authority prescription.

icatibant 30 mg/3 mL injection, 3 mL syringe

1976B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	850.66	41.30	Cipla Icatibant [LR]

■ CARDIOVASCULAR SYSTEM**■ CARDIAC THERAPY****CARDIAC GLYCOSIDES**

Digitalis glycosides

■ DIGOXIN**Note Shared Care Model:**

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical

practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

digoxin 250 microgram tablet, 100

1322N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	15.90	17.19	^a Sigmaxin [LN]
			^b 2.41	18.31	17.19	^a Lanoxin [AS]

digoxin 50 microgram/mL oral liquid, 60 mL

3164M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	3	..	*42.22	41.30	Lanoxin [AS]

digoxin 62.5 microgram tablet, 200

2605D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	15.67	16.96	^a Sigmaxin-PG [LN]
			^b 2.39	18.06	16.96	^a Lanoxin-PG [AS]

ANTIARRHYTHMICS, CLASS I AND III

Antiarrhythmics, class Ia

■ DISOPYRAMIDE

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

disopyramide 100 mg capsule, 100

2923W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	28.06	29.35	Rythmodan [SW]

Antiarrhythmics, class Ib

■ LIDOCAINE (LIGNOCAINE)

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

lidocaine (lignocaine) hydrochloride 10% (500 mg/5 mL) injection, 10 x 5 mL ampoules

2876J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	34.12	35.41	Xylocard 500 [AS]

Antiarrhythmics, class Ic

■ FLECAINIDE

Caution Flecainide acetate should be avoided in patients with poor cardiac function.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Serious supra-ventricular cardiac arrhythmias

Restricted benefit

Serious ventricular cardiac arrhythmias

Clinical criteria:

- The treatment must be initiated in a hospital.

flecainide acetate 50 mg tablet, 60

1088G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	34.90	36.19	^a APO-Flecainide [TX]	^a Flecainide Sandoz [SZ]
						^a Tambacor [IL]	

flecainide acetate 100 mg tablet, 60

1090J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	40.38	41.30	^a APO-Flecainide [TX]	^a Flecainide Sandoz [SZ]
						^a Flecatag [AF]	^a Tambacor [IL]

Antiarrhythmics, class III

■ AMIODARONE

Note This drug has been reported to cause frequent and potentially serious toxicity.

Note Regular monitoring of hepatic and thyroid function is recommended.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Severe cardiac arrhythmias

amiodarone hydrochloride 100 mg tablet, 30

2344J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	16.48	17.77	^a Aratac 100 [AF]	^a Cordarone X 100 [SW]

amiodarone hydrochloride 200 mg tablet, 30

2343H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	19.76	21.05	^a Amiodarone Sandoz [SZ] ^a Aratac 200 [AF] ^a Rithmik 200 [RW]	^a APO-Amiodarone [TX] ^a Cordarone X 200 [SW]

▪ **SOTALOL**

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Severe cardiac arrhythmias

sotalol hydrochloride 80 mg tablet, 60

8398B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	15.42	16.71	^a APO-Sotalol [TX] ^a Solavert [RF] ^a Sotacor [RW]	^a Cardol [AF] ^a Sotalol Sandoz [SZ]
			^B 4.14	19.56	16.71		

sotalol hydrochloride 160 mg tablet, 60

2043M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	18.93	20.22	^a APO-Sotalol [TX] ^a Solavert [RF] ^a Sotacor [RW]	^a Cardol [AF] ^a Sotalol Sandoz [SZ]
			^B 4.33	23.26	20.22		

CARDIAC STIMULANTS EXCL. CARDIAC GLYCOSIDES

Adrenergic and dopaminergic agents

▪ **ADRENALINE (EPINEPHRINE)**

adrenaline (epinephrine) 1 in 1000 (1 mg/mL) injection, 5 x 1 mL ampoules

1016L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	21.68	22.97	Link Medical Products Pty Ltd [LM]

adrenaline (epinephrine) 1 in 1000 (1 mg/mL) injection, 5 x 1 mL ampoules

5004J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	1	..	..	21.68	22.97	Link Medical Products Pty Ltd [LM]

▪ **ADRENALINE (EPINEPHRINE)**

Note The auto-injector should be provided in the framework of a comprehensive anaphylaxis prevention program and an emergency action plan including training in recognition of the symptoms of anaphylaxis and the use of the auto-injector device. (For further information see the Australasian Society of Clinical Immunology and Allergy website at www.allergy.org.au.)

Note Authority approvals will be limited to a maximum quantity of 2 auto-injectors at any one time.

Note No applications for repeats will be authorised.

Authority required

Acute allergic reaction with anaphylaxis

Treatment Phase: Initial sole PBS-subsidised supply for anticipated emergency treatment

Clinical criteria:

- Patient must have been assessed to be at significant risk of anaphylaxis by, or in consultation with a clinical immunologist; OR
 - Patient must have been assessed to be at significant risk of anaphylaxis by, or in consultation with an allergist; OR
 - Patient must have been assessed to be at significant risk of anaphylaxis by, or in consultation with a paediatrician; OR
 - Patient must have been assessed to be at significant risk of anaphylaxis by, or in consultation with a respiratory physician.
- The name of the specialist consulted must be provided at the time of application for initial supply.

Authority required

Acute allergic reaction with anaphylaxis

Treatment Phase: Initial sole PBS-subsidised supply for anticipated emergency treatment

Clinical criteria:

- Patient must have been discharged from hospital or an emergency department after treatment with adrenaline (epinephrine) for acute allergic reaction with anaphylaxis.

Authority required

Acute allergic reaction with anaphylaxis

Treatment Phase: Continuing sole PBS-subsidised supply for anticipated emergency treatment

Clinical criteria:

- Patient must have previously been issued with an authority prescription for this drug.

adrenaline (epinephrine) 150 microgram/0.3 mL injection, 0.3 mL pen device

8697R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	84.26	41.30	^a Adrenaline Jr Mylan [AF]	^a EpiPen Jr. [AL]

adrenaline (epinephrine) 300 microgram/0.3 mL injection, 0.3 mL pen device

8698T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	84.26	41.30	^a Adrenaline Mylan [AF]	^a EpiPen [AL]

VASODILATORS USED IN CARDIAC DISEASES

Organic nitrates

■ **GLYCERYL TRINITRATE**

glyceryl trinitrate 15 mg/24 hours patch, 30

8119H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	33.29	34.58	Minitran 15 [IL]

glyceryl trinitrate 5 mg/24 hours patch, 30

1515R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	28.24	29.53	Transiderm-Nitro 25 [SZ]

glyceryl trinitrate 5 mg/24 hours patch, 30

8027L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	28.24	29.53	Minitran 5 [IL]

glyceryl trinitrate 10 mg/24 hours patch, 30

1516T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	33.29	34.58	Transiderm-Nitro 50 [SZ]

glyceryl trinitrate 10 mg/24 hours patch, 30

8028M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	33.29	34.58	Minitran 10 [IL]

■ **GLYCERYL TRINITRATE**

Note The spray should not be inhaled.

glyceryl trinitrate 400 microgram/actuation spray, 200 actuations

8171C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	‡1	5	..	23.94	25.23	Nitrolingual Pumpspray [SW]

■ **ISOSORBIDE DINITRATE**

isosorbide dinitrate 5 mg sublingual tablet, 100

2588F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	2	..	*19.10	20.39	Isordil Sublingual [RW]

■ **ISOSORBIDE MONONITRATE**

isosorbide mononitrate 120 mg modified release tablet, 30

8273K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	19.34	20.63	^a Monodur 120 mg [IY]
			^B 3.37	22.71	20.63	^a Imdur 120 mg [IX]

isosorbide mononitrate 60 mg modified release tablet, 30

1558B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	15.85	17.14	^a APO-Isosorbide Mononitrate [TX]	^a Duride [AF]
						^a GenRx Isosorbide Mononitrate [GX]	^a ISOBIDE MR [RF]
						^a Isomonit [SZ]	^a Isosorbide AN [EA]

^B 2.33	18.18	17.14	^a Monodur 60 mg [IY]
^B 3.22	19.07	17.14	^a Imdur Durule [IX]

Other vasodilators used in cardiac diseases

■ **NICORANDIL**

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

nicorandil 10 mg tablet, 60

8228C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	±1	5	..	22.25	23.54	^a APO-Nicorandil [TX] ^a Ikotab [AF]	^a Ikorel [SW]

nicorandil 20 mg tablet, 60

8229D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	±1	5	..	26.27	27.56	^a APO-Nicorandil [TX] ^a Ikotab [AF]	^a Ikorel [SW]

■ **PERHEXILINE**

Note Regular monitoring of drug serum levels is recommended.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)

5592

Angina

Clinical criteria:

- The condition must not be responding to other therapy.

perhexiline maleate 100 mg tablet, 100

1822X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	53.51	41.30	Pexsig [AS]

OTHER CARDIAC PREPARATIONS

Other cardiac preparations

■ **IVABRADINE**

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)

4979

Chronic heart failure

Clinical criteria:

- Patient must be symptomatic with NYHA classes II or III, **AND**
- Patient must be in sinus rhythm, **AND**
- Patient must have a documented left ventricular ejection fraction (LVEF) of less than or equal to 35%, **AND**
- Patient must have a resting heart rate at or above 77 bpm at the time ivabradine treatment is initiated, **AND**
- Patient must receive concomitant optimal standard chronic heart failure treatment, which must include the maximum tolerated dose of a beta-blocker, unless contraindicated or not tolerated.

Resting heart rate should be measured by ECG or echocardiography, after 5 minutes rest.

The ECG or echocardiography, result must be documented in the patient's medical records when treatment is initiated.

ivabradine 7.5 mg tablet, 56

2960T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	45.38	41.30	^a APO-Ivabradine [TX]	^a Coralan [SE]

ivabradine 5 mg tablet, 56

10012Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	45.38	41.30	^a APO-Ivabradine [TX]	^a Coralan [SE]

■ **ANTIHYPERTENSIVES**

ANTIADRENERGIC AGENTS, CENTRALLY ACTING

Methyldopa

■ METHYLDOPA

methyldopa 250 mg tablet, 100

1629R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
(NP)	1	5	..	20.82	22.11	Aldomet [AS]

Imidazoline receptor agonists

■ CLONIDINE

clonidine hydrochloride 150 microgram tablet, 100

3141H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
(NP)	1	5	..	33.55	34.84	Catapres [IX]

clonidine hydrochloride 100 microgram tablet, 100

3145M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
(NP)	1	5	..	27.60	28.89	^a APO-Clonidine [TX]	^a Catapres 100 [IX]

■ GUANFACINE

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

9034

Attention deficit hyperactivity disorder

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a paediatrician or psychiatrist.

Clinical criteria:

- The condition must be or have been diagnosed according to the DSM-5 criteria, **AND**
- Patient must have a contraindication to dexamfetamine, methylphenidate or lisdexamfetamine as specified in TGA-approved product information; OR
- Patient must have a comorbid mood disorder that has developed or worsened as a result of dexamfetamine, methylphenidate or lisdexamfetamine treatment and is of a severity necessitating treatment withdrawal; OR
- Patient must be at an unacceptable medical risk of a severity necessitating permanent stimulant treatment withdrawal if given a stimulant treatment with another agent; OR
- Patient must have experienced adverse reactions of a severity necessitating permanent treatment withdrawal following treatment with dexamfetamine, methylphenidate and lisdexamfetamine (not simultaneously).

Population criteria:

- Patient must be or have been diagnosed between the ages of 6 and 17 years inclusive.

Authority required (STREAMLINED)

9031

Attention deficit hyperactivity disorder

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have a contraindication to dexamfetamine, methylphenidate or lisdexamfetamine as specified in TGA-approved product information; OR
- Patient must have a comorbid mood disorder that has developed or worsened as a result of dexamfetamine, methylphenidate or lisdexamfetamine treatment and is of a severity necessitating treatment withdrawal; OR
- Patient must be at an unacceptable medical risk of a severity necessitating permanent stimulant treatment withdrawal if given a stimulant treatment with another agent; OR
- Patient must have experienced adverse reactions of a severity necessitating permanent treatment withdrawal following treatment with dexamfetamine, methylphenidate and lisdexamfetamine (not simultaneously).

Authority required (STREAMLINED)

8544

Attention deficit hyperactivity disorder

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a paediatrician or psychiatrist.

Clinical criteria:

- The condition must be or have been diagnosed according to the DSM-5 criteria, **AND**
- Patient must be receiving a maximum tolerated dose (MTD) of stimulant (dexamfetamine, methylphenidate or lisdexamfetamine) which has been stable for at least four weeks, **AND**
- The treatment must be adjunctive to ongoing maximum tolerated dose (MTD) of stimulant (dexamfetamine, methylphenidate or lisdexamfetamine), **AND**
- Patient must be experiencing residual moderate to severe ADHD symptoms resulting in impaired functioning (social, academic or occupational), present in at least one setting (home, nursery/school/college/work, friends or family homes or other environment).

Population criteria:

- Patient must be or have been diagnosed between the ages of 6 and 17 years inclusive.

Authority required (STREAMLINED)**8585**

Attention deficit hyperactivity disorder

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be adjunctive to ongoing maximum tolerated dose (MTD) of stimulant (dexamfetamine, methylphenidate or lisdexamfetamine).

guanfacine 4 mg modified release tablet, 28

11441E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	125.28	41.30	Intuniv [TK]

guanfacine 3 mg modified release tablet, 28

11440D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	125.28	41.30	Intuniv [TK]

guanfacine 2 mg modified release tablet, 28

11451Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	125.28	41.30	Intuniv [TK]

guanfacine 1 mg modified release tablet, 28

11452R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	125.28	41.30	Intuniv [TK]

■ MOXONIDINE**Restricted benefit**

Hypertension

Clinical criteria:

- Patient must be receiving concurrent antihypertensive therapy.

moxonidine 400 microgram tablet, 30

9020R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	23.68	24.97	^a APO-Moxonidine [TX] ^a Moxonidine GX [SZ] ^a Moxotens [RF]	^a Moxonidine GH [GQ] ^a Moxonidine MYL [AF] ^a Physiotens [GO]

moxonidine 200 microgram tablet, 30

9019Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	18.86	20.15	^a APO-Moxonidine [TX] ^a Moxonidine GX [SZ] ^a Moxotens [RF]	^a Moxonidine GH [GQ] ^a Moxonidine MYL [AF] ^a Physiotens [GO]

ANTIADRENERGIC AGENTS, PERIPHERALLY ACTING*Alpha-adrenoreceptor antagonists***■ PRAZOSIN****prazosin 2 mg tablet, 100**

1480X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	17.90	19.19	^a APO-Prazosin [TX]	^a Minipress [PF]

prazosin 1 mg tablet, 100

1479W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	16.27	17.56	^a APO-Prazosin [TX]	^a Minipress [PF]

prazosin 5 mg tablet, 100

1478T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	22.58	23.87	^a APO-Prazosin [TX]	^a Minipress [PF]

ARTERIOLAR SMOOTH MUSCLE, AGENTS ACTING ON*Hydrazinophthalazine derivatives***■ HYDRALAZINE****hydralazine hydrochloride 50 mg tablet, 100**

1639G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	2	..	*20.94	22.23	Alphapress 50 [AF]

CARDIOVASCULAR SYSTEM

hydralazine hydrochloride 25 mg tablet, 100

1640H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	2	..	*19.70	20.99	Alphapress 25 [AF]

Pyrimidine derivatives

MINOXIDIL

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Severe refractory hypertension

Clinical criteria:

- The treatment must be initiated by a consultant physician.

minoxidil 10 mg tablet, 100

2313R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	51.78	41.30	Loniten [PF]

DIURETICS

LOW-CEILING DIURETICS, THIAZIDES

Thiazides, plain

HYDROCHLOROTHIAZIDE

hydrochlorothiazide 25 mg tablet, 100

1484D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	22.49	23.78	Dithiazide [FF]

LOW-CEILING DIURETICS, EXCL. THIAZIDES

Sulfonamides, plain

CHLORTALIDONE

chlortalidone 25 mg tablet, 50

1585K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	1	..	*19.90	21.19	Hygroton 25 [GH]

INDAPAMIDE

indapamide hemihydrate 1.5 mg modified release tablet, 90

8532C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	1	..	19.73	21.02	^a APO-Indapamide SR [TX]	^a INDAPAMIDE AN SR [EA]
						^a Odaplix SR [AF]	^a Tenaxil SR [RW]
			^B 7.15	26.88	21.02	^a Natrilix SR [SE]	

indapamide hemihydrate 2.5 mg tablet, 90

2436F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	1	..	17.34	18.63	^a Dapa-Tabs [AF]	^a GenRx Indapamide [GX]
						^a Indapamide AN [EA]	^a Insig [RW]
			^B 7.54	24.88	18.63	^a Natrilix [SE]	

HIGH-CEILING DIURETICS

Sulfonamides, plain

FUROSEMIDE (FRUSEMIDE)

furosemide (frusemide) 40 mg tablet, 100

2412Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	1	..	13.99	15.28	Urex [RW]	
						^a APO-Frusemide [TX]	^a Frusax [ER]
						^a Frusemix [TY]	^a FUROSEMIDE AN [EA]
						^a Uremide [AF]	
			^B 2.23	16.22	15.28	^a Lasix [SW]	

furosemide (frusemide) 500 mg tablet, 50

2415D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	3	..	21.83	23.12	Urex-Forte [RW]

furosemide (frusemide) 20 mg/2 mL injection, 5 x 2 mL ampoules

2413B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	13.66	14.95	Lasix [SW]

furosemide (frusemide) 10 mg/mL oral liquid, 30 mL

2411X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	3	..	28.04	29.33	Lasix [SW]

■ FUROSEMIDE (FRUSEMIDE)

Note For item codes 2414C and 1810G, pharmaceutical benefits that have the form tablet 20 mg are equivalent for the purposes of substitution.

furosemide (frusemide) 20 mg tablet, 100

2414C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	1	..	13.99	15.28	^a APO-Frusemide [TX] ^a FUROSEMIDE AN [EA]	^a Frusemix-M [TY]

furosemide (frusemide) 20 mg tablet, 50

1810G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	2	1	..	*14.00 ^B 2.24	15.29 15.29	^a Frusemix-M [TY] ^a Lasix-M [SW]	^a Urex-M [RW]

Aryloxyacetic acid derivatives**■ ETACRYNIC ACID****Restricted benefit**

Patients hypersensitive to other oral diuretics

etacrynic acid 25 mg tablet, 100

8748K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	1	..	*155.00	41.30	Edecrin [FK]

POTASSIUM-SPARING AGENTS***Aldosterone antagonists*****■ EPLERENONE**

Caution Serum electrolytes should be checked regularly

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**4937**

Heart failure with a left ventricular ejection fraction of 40% or less

Clinical criteria:

- The condition must occur within 3 to 14 days following an acute myocardial infarction, **AND**
- The treatment must be commenced within 14 days of an acute myocardial infarction.

The date of the acute myocardial infarction and the date of initiation of treatment with this drug must be documented in the patient's medical records when PBS-subsidised treatment is initiated

eplerenone 50 mg tablet, 30

8880J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	63.53	41.30	^a APO-Eplerenone [TX] ^a ESPLER [RW] ^a Inspra [UJ]	^a Eplerenone AN [EA] ^a Inpler [AF]

eplerenone 25 mg tablet, 30

8879H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	63.53	41.30	^a APO-Eplerenone [TX] ^a ESPLER [RW] ^a Inspra [UJ]	^a Eplerenone AN [EA] ^a Inpler [AF]

■ SPIRONOLACTONE

Caution Serum electrolytes should be checked regularly

Appropriate contraceptive measures should be taken by women of child-bearing age in whom spironolactone therapy has been initiated.

spironolactone 25 mg tablet, 100

2339D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	15.93 ^B 5.48	17.22 17.22	^a Spiractin 25 [AF] ^a Aldactone [PF]	^a Spironolactone Mylan 25 [AL]

spironolactone 100 mg tablet, 100

2340E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	26.83	28.12	^a Spiractin 100 [AF]	^a Spironolactone Mylan 100 [AL]
			^B 5.62	32.45	28.12	^a Aldactone [PF]	

DIURETICS AND POTASSIUM-SPARING AGENTS IN COMBINATION*Low-ceiling diuretics and potassium-sparing agents***■ AMILORIDE + HYDROCHLOROTHIAZIDE**

Caution Serum electrolytes should be checked regularly.

amiloride hydrochloride dihydrate 5 mg + hydrochlorothiazide 50 mg tablet, 50

1486F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	1	..	*18.18	19.47	Moduretic [AS]

OTHER DIURETICS*Vasopressin antagonists***■ TOLVAPTAN**

Caution Tolvaptan has been associated with idiosyncratic hepatic toxicity. Liver function monitoring is required.

Authority required

Autosomal dominant polycystic kidney disease (ADPKD)

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a nephrologist.

Clinical criteria:

- Patient must have an estimated glomerular filtration rate (eGFR) between 30 and 89 mL/min 1.73 m² at the initiation of treatment with this drug for this condition, **AND**
- Patient must have or have had rapidly progressing disease at the time of initiation of this drug for this condition.

Rapidly progressing disease is defined as either of the following:

A decline in eGFR of greater than or equal to 5 mL/min/1.73 m² within one year;

OR

An average decline in eGFR of greater than or equal to 2.5 mL/min/1.73 m² per year over a five year period.

tolvaptan 30 mg tablet [28] (&) tolvaptan 60 mg tablet [28], 56

11597J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1827.87	41.30	Jinarc [OS]

tolvaptan 30 mg tablet [28] (&) tolvaptan 90 mg tablet [28], 56

11588X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1827.87	41.30	Jinarc [OS]

tolvaptan 15 mg tablet [28] (&) tolvaptan 45 mg tablet [28], 56

11602P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1827.87	41.30	Jinarc [OS]

■ TOLVAPTAN

Caution Tolvaptan has been associated with idiosyncratic hepatic toxicity. Liver function monitoring is required.

Authority required (STREAMLINED)**8288**

Autosomal dominant polycystic kidney disease (ADPKD)

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a nephrologist or in consultation with a nephrologist.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have end-stage renal disease defined as an estimated glomerular filtration rate (eGFR) of less than 15 mL/min/1.73m², **AND**
- Patient must not have had a kidney transplant.

tolvaptan 30 mg tablet [28] (&) tolvaptan 60 mg tablet [28], 56

11593E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1827.87	41.30	Jinarc [OS]

tolvaptan 30 mg tablet [28] (&) tolvaptan 90 mg tablet [28], 56

11596H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1827.87	41.30	Jinarc [OS]

tolvaptan 15 mg tablet [28] (&) tolvaptan 45 mg tablet [28], 56

11600M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1827.87	41.30	Jinarc [OS]

PERIPHERAL VASODILATORS**PERIPHERAL VASODILATORS***Other peripheral vasodilators***PHENOXYBENZAMINE****Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Pheochromocytoma

Restricted benefit

Neurogenic urinary retention

phenoxybenzamine hydrochloride 10 mg capsule, 100

1862B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	953.89	41.30	Dibenyline [GH]

phenoxybenzamine hydrochloride 10 mg capsule, 30

1166J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	5	..	*864.90	41.30	Amdipharm Mercury (Australia) Pty Limited [GH]

BETA BLOCKING AGENTS**BETA BLOCKING AGENTS***Beta blocking agents, non-selective***OXPRENOLOL****oxprenolol hydrochloride 40 mg tablet, 100**

2961W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	42.33	41.30	Corbeton 40 [AF]

PINDOLOL

Note Pharmaceutical benefits that have the form pindolol tablet 5 mg are equivalent for the purposes of substitution.

pindolol 5 mg tablet, 100

12224J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	53.88	41.30	^a APO-PINDOL [DZ]

pindolol 5 mg tablet, 100

3062E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	32.90	34.19	^a Barbloc 5 [AF]

PROPRANOLOL**propranolol hydrochloride 40 mg tablet, 100**

2566C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	15.24	16.53	^a APO-Propranolol [TX]
			^B 2.79	18.03	16.53	^a Deralin 40 [AF]
			^B 8.28	23.52	16.53	^a Inderal [AP]

propranolol hydrochloride 160 mg tablet, 50

2899N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	18.74	20.03	Deralin 160 [AF]

propranolol hydrochloride 10 mg tablet, 100

2565B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	14.99	16.28	^a APO-Propranolol [TX]
			^B 2.78	17.77	16.28	^a Deralin 10 [AF]
			^B 8.27	23.26	16.28	^a Inderal [AP]

Beta blocking agents, selective

■ **ATENOLOL**

atenolol 50 mg tablet, 30

1081X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	13.49	14.78	^a APO-Atenolol [TX] ^a Atenolol-GA [ED] ^a Atenolol Sandoz [SZ] ^a Tenolten 50 [DO] ^a Tenormin [AP]	^a Atenolol Amneal [EF] ^a Atenolol GH [GQ] ^a Noten [AF] ^a Tensig [RW]
			^B 13.13	26.62	14.78		

■ **ATENOLOL**

Restricted benefit

For a patient who is unable to take a solid dose form of atenolol.

atenolol 50 mg/10 mL oral liquid, 300 mL

2243C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	30.30	31.59	Atenolol-AFT [AE]

■ **BISOPROLOL**

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Moderate to severe heart failure

Clinical criteria:

- Patient must be stabilised on conventional therapy, which must include an ACE inhibitor or Angiotensin II antagonist, if tolerated.

bisoprolol fumarate 5 mg tablet, 28

8605X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	20.45	21.74	^a APO-Bisoprolol [TX] ^a Bicard 5 [RW] ^a Bisoprolol generichealth [GQ] ^a Bispro 5 [AF] ^a Bicolor [AL]	^a Beprol 5 [DO] ^a Bisoprolol AN [EA] ^a Bisoprolol Sandoz [SZ]
			^B 6.70	27.15	21.74		

bisoprolol fumarate 2.5 mg tablet, 28

8604W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	18.77	20.06	^a APO-Bisoprolol [TX] ^a Bicard 2.5 [RW] ^a Bisoprolol generichealth [GQ] ^a Bispro 2.5 [AF] ^a Bicolor [AL]	^a Beprol 2.5 [DO] ^a Bisoprolol AN [EA] ^a Bisoprolol Sandoz [SZ]
			^B 6.70	25.47	20.06		

bisoprolol fumarate 10 mg tablet, 28

8606Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	22.56	23.85	^a APO-Bisoprolol [TX] ^a Bicard 10 [RW] ^a Bisoprolol generichealth [GQ] ^a Bispro 10 [AF] ^a Bicolor [AL]	^a Beprol 10 [DO] ^a Bisoprolol AN [EA] ^a Bisoprolol Sandoz [SZ]
			^B 6.70	29.26	23.85		

■ **METOPROLOL SUCCINATE**

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Moderate to severe heart failure

Clinical criteria:

- Patient must be stabilised on conventional therapy, which must include an ACE inhibitor or Angiotensin II antagonist, if tolerated.

METOPROLOL SUCCINATE Tablet 23.75 mg (controlled release), 15

8732N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	18.34	19.63	^a Metrol-XL 23.75 [RW]	^a Minax XL [AF]

METOPROLOL SUCCINATE Tablet 95 mg (controlled release), 30

8734Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	49.29	41.30	^a Metrol-XL 95 [RW]	^a Minax XL [AF]

^a Toprol-XL 95 [AP]

NP METOPROLOL SUCCINATE Tablet 190 mg (controlled release), 30

8735R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	58.61	41.30	^a Metrol-XL 190 [RW] ^a Toprol-XL 190 [AP]	^a Minax XL [AF]

METOPROLOL SUCCINATE Tablet 47.5 mg (controlled release), 30

8733P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	41.85	41.30	^a Metrol-XL 47.5 [RW] ^a Toprol-XL 47.5 [AP]	^a Minax XL [AF]

■ METOPROLOL TARTRATE

METOPROLOL TARTRATE Tablet 50 mg, 100

1324Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	14.20	15.49	^a Metoprolol AN [EA] ^b APO-Metoprolol [TX] ^b Metrol 50 [RW]	^a Mistrom [ER] ^b Metoprolol Sandoz [SZ] ^b Minax 50 [AF]
			^B 1.67	15.87	15.49	^a Lopresor 50 [NV]	
			^B 8.21	22.41	15.49	^b Betaloc [AP]	

METOPROLOL TARTRATE Tablet 100 mg, 60

1325R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	14.69	15.98	^a Metoprolol AN [EA] ^b APO-Metoprolol [TX] ^b Metrol 100 [RW]	^a Mistrom [ER] ^b Metoprolol Sandoz [SZ] ^b Minax 100 [AF]
			^B 1.67	16.36	15.98	^a Lopresor 100 [NV]	
			^B 8.25	22.94	15.98	^b Betaloc [AP]	

■ NEBIVOLOL

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Moderate to severe heart failure

Clinical criteria:

- Patient must be stabilised on conventional therapy, which must include an ACE inhibitor or Angiotensin II antagonist, if tolerated.

nebivolol 10 mg tablet, 28

9312D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	56.71	41.30	^a APO-Nebivolol [TX] ^a Nebivolol Sandoz [SZ]	^a Nebilet [FK] ^a Nepiten [AF]

nebivolol 1.25 mg tablet, 28

9316H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	2	5	..	*43.70	41.30	^a APO-Nebivolol [TX] ^a Nebivolol Sandoz [SZ]	^a Nebilet [FK] ^a Nepiten [AF]

nebivolol 5 mg tablet, 28

9311C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	51.57	41.30	^a APO-Nebivolol [TX] ^a Nebivolol Sandoz [SZ]	^a Nebilet [FK] ^a Nepiten [AF]

Alpha and beta blocking agents

■ CARVEDILOL

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Moderate to severe heart failure

Clinical criteria:

- Patient must be stabilised on conventional therapy, which must include an ACE inhibitor or Angiotensin II antagonist, if tolerated.

Restricted benefit

Patients receiving this drug as a pharmaceutical benefit prior to 1 August 2002

carvedilol 3.125 mg tablet, 30

8255L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	15.26	16.55	^a APO-Carvedilol [TX] ^a Carvidol [RF] ^a Volirop 3.125 [DO]	^a Carvedilol AN [EA] ^a Vedilol 3.125 [RW]

carvedilol 12.5 mg tablet, 60

8257N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	22.79	24.08	^a APO-Carvedilol [TX] ^a Carvedilol Sandoz [SZ] ^a Dicarz [AF] ^a Vedilol 12.5 [RW]	^a Carvedilol AN [EA] ^a Carvidol [RF] ^a Dilatrend 12.5 [PB] ^a Volirop 12.5 [DO]

carvedilol 6.25 mg tablet, 60

8256M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	20.63	21.92	^a APO-Carvedilol [TX] ^a Carvedilol Sandoz [SZ] ^a Dicarz [AF] ^a Vedilol 6.25 [RW]	^a Carvedilol AN [EA] ^a Carvidol [RF] ^a Dilatrend 6.25 [PB] ^a Volirop 6.25 [DO]

carvedilol 25 mg tablet, 60

8258P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	25.48	26.77	^a APO-Carvedilol [TX] ^a Carvedilol Sandoz [SZ] ^a Dicarz [AF] ^a Vedilol 25 [RW]	^a Carvedilol AN [EA] ^a Carvidol [RF] ^a Dilatrend 25 [PB] ^a Volirop 25 [DO]

■ **LABETALOL**

labetalol hydrochloride 200 mg tablet, 100

1567L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	25.27	26.56	Trandate [AS]

labetalol hydrochloride 100 mg tablet, 100

1566K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	25.27	26.56	^a Presolol 100 [AF]
			^B 3.50	28.77	26.56	^a Trandate [AS]

■ **CALCIUM CHANNEL BLOCKERS**

SELECTIVE CALCIUM CHANNEL BLOCKERS WITH MAINLY VASCULAR EFFECTS

Dihydropyridine derivatives

■ **AMLODIPINE**

amlodipine 5 mg tablet, 30

2751T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	13.26	14.55	^a Amlo 5 [RW] ^a Amlodipine AN [EA] ^a Amlodipine GH [GQ] ^a APO-Amlodipine [TX] ^a Blooms the Chemist Amlodipine [IB] ^a Nordip [AF] ^a Pharmacor Amlodipine [CR]	^a Amlodipine Amneal [EF] ^a Amlodipine APOTEX [GX] ^a Amlodipine Sandoz [SZ] ^a Auro-Amlodipine 5 [DO] ^a BTC Amlodipine [JB] ^a Norvapine [ED]
			^B 10.85	24.11	14.55	^a Norvasc [AS]	

amlodipine 10 mg tablet, 30

2752W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	13.90	15.19	^a Amlo 10 [RW] ^a Amlodipine AN [EA] ^a Amlodipine GH [GQ] ^a APO-Amlodipine [TX] ^a Blooms the Chemist Amlodipine [IB] ^a Nordip [AF] ^a Pharmacor Amlodipine [CR]	^a Amlodipine Amneal [EF] ^a Amlodipine APOTEX [GX] ^a Amlodipine Sandoz [SZ] ^a Auro-Amlodipine 10 [DO] ^a BTC Amlodipine [JB] ^a Norvapine [ED]
			^B 10.90	24.80	15.19	^a Norvasc [AS]	

■ FELODIPINE

felodipine 10 mg modified release tablet, 30

2367N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	19.32	20.61	^a Felodil XR 10 [RW]	^a Felodur ER 10 mg [TX]
						^a Fendex ER [AF]	
			^B 2.09	21.41	20.61	^a Plendil ER [GX]	

felodipine 2.5 mg modified release tablet, 30

2361G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	15.04	16.33	^a Felodur ER 2.5 mg [TX]	^a Fendex ER [AF]
			^B 1.94	16.98	16.33	^a Plendil ER [GX]	

felodipine 5 mg modified release tablet, 30

2366M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	16.18	17.47	^a Felodil XR 5 [RW]	^a Felodur ER 5 mg [TX]
						^a Fendex ER [AF]	
			^B 1.96	18.14	17.47	^a Plendil ER [GX]	

■ LERCANIDIPINE

lercanidipine hydrochloride 10 mg tablet, 28

8534E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	14.38	15.67	^a BTC Lercanidipine [JB]	^a Lercadip [EA]
						^a Lercan [RW]	^a Lercanidipine APOTEX [GX]
						^a Zircol 10 [AL]	
			^B 3.26	17.64	15.67	^a Zandip [GO]	

lercanidipine hydrochloride 20 mg tablet, 28

8679T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	15.68	16.97	^a BTC Lercanidipine [JB]	^a Lercadip [EA]
						^a Lercan [RW]	^a Lercanidipine APOTEX [GX]
						^a Zircol 20 [AL]	
			^B 3.33	19.01	16.97	^a Zandip [GO]	

■ NIFEDIPINE

nifedipine 30 mg modified release tablet, 30

1906H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	18.91	20.20	^a Addos XR 30 [RW]	^a Adefin XL 30 [AF]
						^a APO-Nifedipine XR [TX]	

nifedipine 60 mg modified release tablet, 30

1907J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	20.68	21.97	^a Addos XR 60 [RW]	^a Adefin XL 60 [AF]
						^a APO-Nifedipine XR [TX]	

nifedipine 20 mg tablet, 60

1695F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	18.16	19.45	Adefin 20 [AF]

nifedipine 10 mg tablet, 60

1694E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	17.07	18.36	Adefin 10 [AF]

SELECTIVE CALCIUM CHANNEL BLOCKERS WITH DIRECT CARDIAC EFFECTS

Phenylalkylamine derivatives

■ VERAPAMIL

Caution The myocardial depressant effects of this drug and of beta-blocking drugs are additive.

verapamil hydrochloride 80 mg tablet, 100

1250T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	19.49	20.78	^a Anpec 80 [AF]
			^B 3.50	22.99	20.78	^a Isoptin [GO]

verapamil hydrochloride 180 mg modified release tablet, 30

2208F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	18.05	19.34	^a Cordilox 180 SR [GT]
			^B 3.50	21.55	19.34	^a Isoptin 180 SR [GO]

verapamil hydrochloride 240 mg modified release tablet, 30

1241H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	20.06	21.35	^a Cordilox SR [GT]
			^B 3.50	23.56	21.35	^a Isoptin SR [GO]

Benzothiazepine derivatives

■ **DILTIAZEM**

Caution The myocardial depressant effects of this drug and of beta-blocking drugs are additive.

diltiazem hydrochloride 180 mg modified release capsule, 30

1312C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	18.74	20.03	^a Diltiazem Sandoz CD [SZ]	^a Vasocardol CD [AV]
			^B 1.90	20.64	20.03	^a Cardizem CD [SW]	

diltiazem hydrochloride 60 mg tablet, 90

1335G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	18.08	19.37	^a Diltiazem AN [EA]	^a Vasocardol [AV]
			^B 1.90	19.98	19.37	^a Cardizem [SW]	

diltiazem hydrochloride 360 mg modified release capsule, 30

8480H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	23.93	25.22	^a Diltiazem Sandoz CD [SZ]	^a Vasocardol CD [AV]
			^B 1.90	25.83	25.22	^a Cardizem CD [SW]	

diltiazem hydrochloride 240 mg modified release capsule, 30

1313D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	21.11	22.40	^a Diltiazem Sandoz CD [SZ]	^a Vasocardol CD [AV]
			^B 1.90	23.01	22.40	^a Cardizem CD [SW]	

■ **AGENTS ACTING ON THE RENIN-ANGIOTENSIN SYSTEM**

ACE INHIBITORS, PLAIN

ACE inhibitors, plain

■ **CAPTOPRIL**

Caution Use of ACE inhibitors during pregnancy is contraindicated since these drugs have been associated with foetal death in utero.

captopril 50 mg tablet, 90

1149L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	21.58	22.87	^a Captopril Sandoz [SZ]
			^B 3.90	25.48	22.87	^a Capoten [RW]

captopril 25 mg tablet, 90

1148K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	16.94	18.23	^a Captopril Sandoz [SZ]
			^B 4.60	21.54	18.23	^a Capoten [RW]

■ **CAPTOPRIL**

Caution Use of ACE inhibitors during pregnancy is contraindicated since these drugs have been associated with foetal death in utero.

Restricted benefit

Patients unable to take a solid dose form of an ACE inhibitor.

captopril 5 mg/mL oral liquid, 95 mL

8760C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	107.84	41.30	Capoten [RW]

■ **ENALAPRIL**

Caution Use of ACE inhibitors during pregnancy is contraindicated since these drugs have been associated with foetal death in utero.

enalapril maleate 5 mg tablet, 30

1370D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	14.17	15.46	^a Acetec [AL]	^a APO-Enalapril [TX]
						^a Enalapril Actavis [ED]	^a Enalapril generichealth [GQ]
						^a Enalapril Sandoz [SZ]	^a Malean [RW]

enalapril maleate 10 mg tablet, 30

1368B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	15.30	16.59	^a Acetec [AL]	^a APO-Enalapril [TX]
						^a Enalapril Actavis [ED]	^a Enalapril generichealth [GQ]
						^a Enalapril Sandoz [SZ]	^a Malean [RW]
			^B 6.21	21.51	16.59	^a Renitec [AF]	

enalapril maleate 20 mg tablet, 30

1369C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	16.08	17.37	^a Acetec [AL] ^a Enalapril Actavis [ED] ^a Enalapril Sandoz [SZ]	^a APO-Enalapril [TX] ^a Enalapril generichealth [GQ] ^a Malean [RW]
			^B 6.26	22.34	17.37	^a Renitec 20 [AF]	

▪ **FOSINOPRIL**

Caution Use of ACE inhibitors during pregnancy is contraindicated since these drugs have been associated with foetal death in utero.

fosinopril sodium 20 mg tablet, 30

1183G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	18.73	20.02	^a APO-Fosinopril [TX] ^a Monace 20 [AF]	^a Fosipril 20 [RW]

fosinopril sodium 10 mg tablet, 30

1182F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	16.61	17.90	^a APO-Fosinopril [TX] ^a Monace 10 [AF]	^a Fosipril 10 [RW]

▪ **LISINOPRIL**

Caution Use of ACE inhibitors during pregnancy is contraindicated since these drugs have been associated with foetal death in utero.

lisinopril 20 mg tablet, 30

2458J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	16.22	17.51	^a APO-Lisinopril [TX] ^a Fibsol 20 [RW] ^a Lisinopril generichealth [GQ] ^a Zinopril 20 [AL]	^a Auro-Lisinopril 20 [DO] ^a Lisinopril AN [EA] ^a Lisinopril Sandoz [SZ]
			^B 4.71	20.93	17.51	^a Zestril [AP]	

lisinopril 10 mg tablet, 30

2457H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	15.44	16.73	^a APO-Lisinopril [TX] ^a Fibsol 10 [RW] ^a Lisinopril generichealth [GQ] ^a Zinopril 10 [AL]	^a Auro-Lisinopril 10 [DO] ^a Lisinopril AN [EA] ^a Lisinopril Sandoz [SZ]
			^B 4.65	20.09	16.73	^a Zestril [AP]	

lisinopril 5 mg tablet, 30

2456G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	14.44	15.73	^a APO-Lisinopril [TX] ^a Fibsol 5 [RW] ^a Lisinopril generichealth [GQ] ^a Zinopril 5 [AL]	^a Auro-Lisinopril 5 [DO] ^a Lisinopril AN [EA] ^a Lisinopril Sandoz [SZ]
			^B 4.58	19.02	15.73	^a Zestril [AP]	

▪ **PERINDOPRIL**

Caution Use of ACE inhibitors during pregnancy is contraindicated since these drugs have been associated with foetal death in utero.

Note Pharmaceutical benefits that have the form perindopril erbumine 2 mg tablet and pharmaceutical benefits that have the form perindopril arginine 2.5 mg tablet are equivalent for the purposes of substitution.

perindopril erbumine 2 mg tablet, 30

3050M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	13.86	15.15	^a APO-Perindopril [TX] ^a BTC Perindopril [JB] ^a Indosyl Mono 2 [RW] ^a Perindo [AF] ^a Perindopril AN [EF]	^a Blooms the Chemist Perindopril [IB] ^a Idaprex 2 [SZ] ^a NOUMED PERINDOPRIL [VO] ^a Perindopril Actavis 2 [EA] ^a Perindopril APOTEX [TY]

perindopril arginine 2.5 mg tablet, 30

9006B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	13.86	15.15	^a APO-Perindopril Arginine [TX] ^a Coversyl 2.5mg [SE]	^a PREXUM 2.5 [RW]
			^B 9.46	23.32	15.15		

▪ **PERINDOPRIL**

Caution Use of ACE inhibitors during pregnancy is contraindicated since these drugs have been associated with foetal death in utero.

Note Pharmaceutical benefits that have the form perindopril erbumine 4 mg tablet and pharmaceutical benefits that have the form perindopril arginine 5 mg tablet are equivalent for the purposes of substitution.

perindopril erbumine 4 mg tablet, 30

3051N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	15.09	16.38	^a APO-Perindopril [TX]	^a Blooms the Chemist Perindopril [IB]
						^a BTC Perindopril [JB]	^a Idaprex 4 [SZ]
						^a Indosyl Mono 4 [RW]	^a NOUMED PERINDOPRIL [VO]
						^a Perindo [AF]	^a Perindopril Actavis 4 [ED]
						^a Perindopril AN [EF]	^a Perindopril APOTEX [TY]
						^a Perindopril generichealth [GQ]	

perindopril arginine 5 mg tablet, 30

9007C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	15.09	16.38	^a APO-Perindopril Arginine [TX]	^a PREXUM 5 [RW]
			^B 9.24	24.33	16.38	^a Coversyl 5mg [SE]	

■ **PERINDOPRIL**

Caution Use of ACE inhibitors during pregnancy is contraindicated since these drugs have been associated with foetal death in utero.

Note Pharmaceutical benefits that have the form perindopril erbumine 8 mg tablet and pharmaceutical benefits that have the form perindopril arginine 10 mg tablet are equivalent for the purposes of substitution.

perindopril arginine 10 mg tablet, 30

9008D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	16.50	17.79	^a APO-Perindopril Arginine [TX]	^a PREXUM 10 [RW]
			^B 9.82	26.32	17.79	^a Coversyl 10mg [SE]	

perindopril erbumine 8 mg tablet, 30

8704D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	16.50	17.79	^a APO-Perindopril [TX]	^a Blooms the Chemist Perindopril [IB]
						^a BTC Perindopril [JB]	^a Idaprex 8 [SZ]
						^a Indosyl Mono 8 [RW]	^a NOUMED PERINDOPRIL [VO]
						^a Perindo [AF]	^a Perindopril Actavis 8 [ED]
						^a Perindopril AN [EF]	^a Perindopril APOTEX [TY]
						^a Perindopril generichealth [GQ]	

■ **QUINAPRIL**

Caution Use of ACE inhibitors during pregnancy is contraindicated since these drugs have been associated with foetal death in utero.

quinapril 5 mg tablet, 30

1968N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	15.34	16.63	^a Accupril [PF]	^a ACQUIN [RF]
						^a Acquin Aspen 5 [RW]	^a Qpril 5 [AF]

quinapril 20 mg tablet, 30

1970Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	17.01	18.30	^a Accupril [PF]	^a ACQUIN [RF]
						^a Acquin Aspen 20 [RW]	^a APO-Quinapril [TX]
						^a Qpril 20 [AF]	^a Quinapril generichealth [GQ]

quinapril 10 mg tablet, 30

1969P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	16.14	17.43	^a ACQUIN [RF]	^a Acquin Aspen 10 [RW]
			^B 4.07	20.21	17.43	^a APO-Quinapril [TX]	^a Qpril 10 [AF]
						^a Accupril [PF]	

■ **RAMIPRIL**

Caution Use of ACE inhibitors during pregnancy is contraindicated since these drugs have been associated with foetal death in utero.

Note Pharmaceutical benefits that have the form ramipril 10 mg tablet and pharmaceutical benefits that have the form ramipril 10 mg capsule are equivalent for the purposes of substitution.

ramipril 10 mg tablet, 30

1316G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	15.50	16.79	^a APO-Ramipril [TX]	^a Ramipril AN [EA]
						^a Ramipril Sandoz [SZ]	^a Tritace [SW]
						^a Tryzan Tabs 10 [AF]	

ramipril 10 mg capsule, 30

8470T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	15.50	16.79	^a APO-Ramipril [TX]	^a Prilace [RF]
						^a Ramace 10 mg [AV]	^a Ramipril AN [EA]
						^a Ramipril Sandoz [SZ]	^a Ramipril Winthrop [WA]
						^a Tritace 10 mg [SW]	^a Tryzan Caps 10 [AF]

■ RAMIPRIL

Caution Use of ACE inhibitors during pregnancy is contraindicated since these drugs have been associated with foetal death in utero.

Note Pharmaceutical benefits that have the form ramipril 1.25 mg tablet and pharmaceutical benefits that have the form ramipril 1.25 mg capsule are equivalent for the purposes of substitution.

ramipril 1.25 mg tablet, 30

1944H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	13.38	14.67	^a Prilace [RF]	^a Ramace 1.25 mg [AV]
						^a Ramipril Sandoz [SZ]	^a Ramipril Winthrop [WA]
						^a Tritace 1.25 mg [SW]	^a Tryzan Tabs 1.25 [AF]

ramipril 1.25 mg capsule, 30

9120B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	13.38	14.67	^a APO-Ramipril [TX]	^a Tryzan Caps 1.25 [AF]

■ RAMIPRIL

Caution Use of ACE inhibitors during pregnancy is contraindicated since these drugs have been associated with foetal death in utero.

Note Pharmaceutical benefits that have the form ramipril 2.5 mg tablet and pharmaceutical benefits that have the form ramipril 2.5 mg capsule are equivalent for the purposes of substitution.

ramipril 2.5 mg capsule, 30

9121C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	13.82	15.11	^a APO-Ramipril [TX]	^a Tryzan Caps 2.5 [AF]

ramipril 2.5 mg tablet, 30

1945J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	13.82	15.11	^a APO-Ramipril [TX]	^a Prilace [RF]
						^a Ramace 2.5 mg [AV]	^a Ramipril AN [EA]
						^a Ramipril Sandoz [SZ]	^a Ramipril Winthrop [WA]
						^a Tritace 2.5 mg [SW]	^a Tryzan Tabs 2.5 [AF]

■ RAMIPRIL

Caution Use of ACE inhibitors during pregnancy is contraindicated since these drugs have been associated with foetal death in utero.

Note Pharmaceutical benefits that have the form ramipril 5 mg tablet and pharmaceutical benefits that have the form ramipril 5 mg capsule are equivalent for the purposes of substitution.

ramipril 5 mg tablet, 30

1946K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	14.17	15.46	^a APO-Ramipril [TX]	^a Prilace [RF]
						^a Ramace 5 mg [AV]	^a Ramipril AN [EA]
						^a Ramipril Sandoz [SZ]	^a Ramipril Winthrop [WA]
						^a Tritace 5 mg [SW]	^a Tryzan Tabs 5 [AF]

ramipril 5 mg capsule, 30

9122D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	14.17	15.46	^a APO-Ramipril [TX]	^a Tryzan Caps 5 [AF]

■ TRANDOLAPRIL

Caution Use of ACE inhibitors during pregnancy is contraindicated since these drugs have been associated with foetal death in utero.

trandolapril 500 microgram capsule, 28

2791X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	14.39	15.68	^a Dolapril 0.5 [RW]	^a Tranalpha [AF]
			^B 3.26	17.65	15.68	^a Gopten [GO]	

trandolapril 1 mg capsule, 28

2792Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	16.12	17.41	^a Dolapril 1 [RW]	^a Tranalpha [AF]
			^B 3.37	19.49	17.41	^a Gopten [GO]	

trandolapril 2 mg capsule, 28

2793B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	16.88	18.17	^a Dolapril 2 [RW]	^a Tranalpha [AF]
			^B 3.42	20.30	18.17	^a Gopten [GO]	

trandolapril 4 mg capsule, 28

8758Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	21.01	22.30	^a Dolapril 4 [RW]	^a Tranalpha [AF]
			^B 3.49	24.50	22.30	^a Gopten [GO]	

ACE INHIBITORS, COMBINATIONS

ACE inhibitors and diuretics

■ ENALAPRIL + HYDROCHLOROTHIAZIDE

Caution Use of ACE inhibitors during pregnancy is contraindicated since these drugs have been associated with foetal death in utero.

Restricted benefit

Hypertension

Clinical criteria:

- The treatment must not be for the initiation of anti-hypertensive therapy, **AND**
- The condition must be inadequately controlled with an ACE inhibitor; OR
- The condition must be inadequately controlled with a thiazide diuretic.

enalapril maleate 20 mg + hydrochlorothiazide 6 mg tablet, 30

8477E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	17.42	18.71	^a Enalapril/HCT Sandoz [SZ]
			^B 6.96	24.38	18.71	^a Renitec Plus 20/6 [AF]

■ FOSINOPRIL + HYDROCHLOROTHIAZIDE

Caution Use of ACE inhibitors during pregnancy is contraindicated since these drugs have been associated with foetal death in utero.

Restricted benefit

Hypertension

Clinical criteria:

- The treatment must not be for the initiation of anti-hypertensive therapy, **AND**
- The condition must be inadequately controlled with an ACE inhibitor; OR
- The condition must be inadequately controlled with a thiazide diuretic.

fosinopril sodium 20 mg + hydrochlorothiazide 12.5 mg tablet, 30

8401E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	20.42	21.71	^a APO-Fosinopril HCTZ 20/12.5 [TX]	^a Fosetic 20/12.5 [ZP]
						^a Fosinopril/HCT Actavis 20/12.5 [EA]	

■ PERINDOPRIL + INDAPAMIDE

Caution Use of ACE inhibitors during pregnancy is contraindicated since these drugs have been associated with foetal death in utero.

perindopril arginine 2.5 mg + indapamide hemihydrate 625 microgram tablet, 30

2190G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	14.27	15.56	^a PREXUM Combi LD 2.5/0.625 [RW]
			^B 7.61	21.88	15.56	^a Coversyl Plus LD 2.5mg/0.625mg [SE]

■ PERINDOPRIL + INDAPAMIDE

Caution Use of ACE inhibitors during pregnancy is contraindicated since these drugs have been associated with foetal death in utero.

Note Pharmaceutical benefits that have the form perindopril with indapamide hemihydrate tablet (containing 4 mg perindopril erbumine-1.25 mg indapamide hemihydrate) and pharmaceutical benefits that have the form perindopril with indapamide hemihydrate tablet (containing 5 mg perindopril arginine-1.25 mg indapamide hemihydrate) are equivalent for the purposes of substitution.

Restricted benefit

Hypertension

Clinical criteria:

- The treatment must not be for the initiation of anti-hypertensive therapy, **AND**
- The condition must be inadequately controlled with an ACE inhibitor; OR
- The condition must be inadequately controlled with a thiazide-like diuretic.

perindopril erbumine 4 mg + indapamide hemihydrate 1.25 mg tablet, 30

8449Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	15.91	17.20	^a GenRx Perindopril/ Indapamide 4/1.25 [GX]	^a Idaprex Combi 4/1.25 [SZ]
						^a Indosyl Combi 4/1.25 [RW]	^a Perindopril and Indapamide AN 4/1.25 [EF]
						^a Perindopril Combi Actavis 4/1.25 [ED]	^a Perindopril/ Indapamide GH 4/1.25 [GQ]
						^B 2.79	^a Perindo Combi 4/1.25 [AF]
				18.70	17.20		

perindopril arginine 5 mg + indapamide hemihydrate 1.25 mg tablet, 30

2845R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	15.91	17.20	^a Prexum Combi 5/1.25 [RW]
			^B 6.99	22.90	17.20	^a Coversyl Plus 5mg/1.25mg [SE]

■ QUINAPRIL + HYDROCHLOROTHIAZIDE

Caution Use of ACE inhibitors during pregnancy is contraindicated since these drugs have been associated with foetal death in utero.

Restricted benefit

Hypertension

Clinical criteria:

- The treatment must not be for the initiation of anti-hypertensive therapy, **AND**
- The condition must be inadequately controlled with an ACE inhibitor; OR
- The condition must be inadequately controlled with a thiazide diuretic.

quinapril 20 mg + hydrochlorothiazide 12.5 mg tablet, 30

8590D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	18.51	19.80	Accuretic 20/12.5mg [PF]

quinapril 10 mg + hydrochlorothiazide 12.5 mg tablet, 30

8589C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	17.60	18.89	Accuretic 10/12.5mg [PF]

ACE inhibitors and calcium channel blockers

■ LERCANIDIPINE + ENALAPRIL

Caution Use of ACE inhibitors during pregnancy is contraindicated since these drugs have been associated with foetal death in utero.

Restricted benefit

Hypertension

Clinical criteria:

- The treatment must not be for the initiation of anti-hypertensive therapy, **AND**
- The condition must be inadequately controlled with an ACE inhibitor; OR
- The condition must be inadequately controlled with a dihydropyridine calcium channel blocker.

lercanidipine hydrochloride 10 mg + enalapril maleate 10 mg tablet, 28

9144G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	17.06	18.35	Zan-Extra 10/10 [GO]

lercanidipine hydrochloride 10 mg + enalapril maleate 20 mg tablet, 28

9145H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	17.79	19.08	Zan-Extra 10/20 [GO]

■ PERINDOPRIL + AMLODIPINE

Caution Use of ACE inhibitors during pregnancy is contraindicated since these drugs have been associated with foetal death in utero.

Restricted benefit

Hypertension

Clinical criteria:

- The treatment must not be for the initiation of anti-hypertensive therapy, **AND**
- The condition must be inadequately controlled with an ACE inhibitor; OR
- The condition must be inadequately controlled with a dihydropyridine calcium channel blocker.

Restricted benefit

Stable coronary heart disease

Clinical criteria:

- The treatment must not be for the initiation of therapy for coronary heart disease, **AND**
- The condition must be stabilised by treatment with perindopril and amlodipine at the same doses.

perindopril arginine 5 mg + amlodipine 10 mg tablet, 30

9347Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	16.56	17.85	^a APO-Perindopril Arginine/Amlodipine 5/10 [TX]	^a Reaptan 5/10 [RW]
			^b 9.29	25.85	17.85	^a Coveram 5/10 [SE]	

perindopril arginine 5 mg + amlodipine 5 mg tablet, 30

9346X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	15.92	17.21	^a APO-Perindopril Arginine/Amlodipine 5/5 [TX]	^a Reaptan 5/5 [RW]
			^b 8.93	24.85	17.21	^a Coveram 5/5 [SE]	

perindopril arginine 10 mg + amlodipine 5 mg tablet, 30

9348B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	17.33	18.62	^a APO-Perindopril Arginine/Amlodipine 10/5 [TX]	^a Reaptan 10/5 [RW]
			^b 9.50	26.83	18.62	^a Coveram 10/5 [SE]	

perindopril arginine 10 mg + amlodipine 10 mg tablet, 30

9349C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	17.98	19.27	^a APO-Perindopril Arginine/Amlodipine 10/10 [TX]	^a Reaptan 10/10 [RW]
			^B 9.85	27.83	19.27	^a Coveram 10/10 [SE]	

■ RAMIPRIL + FELODIPINE

Caution Use of ACE inhibitors during pregnancy is contraindicated since these drugs have been associated with foetal death in utero.

Restricted benefit

Hypertension

Clinical criteria:

- The treatment must not be for the initiation of anti-hypertensive therapy, **AND**
- The condition must be inadequately controlled with an ACE inhibitor; OR
- The condition must be inadequately controlled with a dihydropyridine calcium channel blocker.

ramipril 5 mg + felodipine 5 mg modified release tablet, 30

2629J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	17.92	19.21	Triasyn 5.0/5.0 [SW]

ramipril 2.5 mg + felodipine 2.5 mg modified release tablet, 30

2626F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	16.43	17.72	Triasyn 2.5/2.5 [SW]

■ TRANDOLAPRIL + VERAPAMIL

Caution Use of ACE inhibitors during pregnancy is contraindicated since these drugs have been associated with foetal death in utero. The myocardial depressant effects of verapamil hydrochloride and of beta-blocking drugs are additive.

Restricted benefit

Hypertension

Clinical criteria:

- The treatment must not be for the initiation of anti-hypertensive therapy, **AND**
- The condition must be inadequately controlled with an ACE inhibitor; OR
- The condition must be inadequately controlled with verapamil.

trandolapril 4 mg + verapamil hydrochloride 240 mg modified release tablet, 28

2857J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	28.51	29.80	Tarka 4/240 [GO]

trandolapril 2 mg + verapamil hydrochloride 180 mg modified release tablet, 28

9387C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	22.44	23.73	Tarka 2/180 [GO]

ANGIOTENSIN II RECEPTOR BLOCKERS (ARBs), PLAIN*Angiotensin II receptor blockers (ARBs), plain***■ CANDESARTAN****candesartan cilexetil 8 mg tablet, 30**

8296P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	13.72	15.01	^a Adesan [AF]	^a APO-Candesartan [TX]
						^a Blooms the Chemist Candesartan [IB]	^a CANDESAN [RF]
						^a Candesartan AN [EA]	^a Candesartan Aspen 8 [RW]
						^a Candesartan Sandoz [SZ]	
			^B 13.16	26.88	15.01	^a Atacand [AP]	

candesartan cilexetil 32 mg tablet, 30

8889W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	17.07	18.36	^a Adesan [AF]	^a APO-Candesartan [TX]
						^a Blooms the Chemist Candesartan [IB]	^a CANDESAN [RF]
						^a Candesartan AN [EA]	^a Candesartan Aspen 32 [RW]
						^a Candesartan GH [GQ]	^a Candesartan Sandoz [SZ]
			^B 13.41	30.48	18.36	^a Atacand [AP]	

candesartan cilexetil 16 mg tablet, 30

8297Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	16.27	17.56	^a Adesan [AF]	^a APO-Candesartan [TX]
						^a Atacand [AP]	^a Blooms the Chemist Candesartan [IB]
						^a CANDESAN [RF]	^a Candesartan AN [EA]

^a Candesartan Aspen 16 [RW] ^a Candesartan GH [GQ]
^a Candesartan Sandoz [SZ]

candesartan cilexetil 4 mg tablet, 30

8295N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	12.86	14.15	^a Adesan [AF] ^a Blooms the Chemist Candesartan [IB] ^a Candesartan AN [EA] ^a Candesartan Sandoz [SZ]	^a APO-Candesartan [TX] ^a CANDESAN [RF] ^a Candesartan Aspen 4 [RW]
			^B 13.09	25.95	14.15	^a Atacand [AP]	

■ EPROSARTAN**eprosartan 400 mg tablet, 28**

8397Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	^T 7.00	*31.38	25.67	Teveten [GO]

eprosartan 600 mg tablet, 28

8447N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	^T 3.50	31.01	28.80	Teveten [GO]

■ EPROSARTAN**Authority required**

Adverse effects occurring with all of the base-priced drugs

Authority required

Drug interactions occurring with all of the base-priced drugs

Authority required

Drug interactions expected to occur with all of the base-priced drugs

Authority required

Transfer to a base-priced drug would cause patient confusion resulting in problems with compliance

eprosartan 400 mg tablet, 28

8951D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*31.38	32.67	Teveten [GO]

eprosartan 600 mg tablet, 28

5491B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	31.01	32.30	Teveten [GO]

■ IRBESARTAN**irbesartan 75 mg tablet, 30**

8246B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	13.63	14.92	^a Abisart 75 [AL] ^a AVSARTAN [RF] ^a Irbesartan Actavis 75 [ED] ^a Irbesartan AN [EA] ^a Irbesartan Sandoz [SZ]	^a APO-Irbesartan [TX] ^a Blooms the Chemist Irbesartan [IB] ^a Irbesartan AMNEAL [EF] ^a Irbesartan GH [GQ]
			^B 2.46	16.09	14.92	^a Avapro [AV]	^a Karvea [SW]

irbesartan 300 mg tablet, 30

8248D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	15.86	17.15	^a Abisart 300 [AL] ^a AVSARTAN [RF] ^a Irbesartan Actavis 300 [ED] ^a Irbesartan AN [EA] ^a Irbesartan Sandoz [SZ]	^a APO-Irbesartan [TX] ^a Blooms the Chemist Irbesartan [IB] ^a Irbesartan AMNEAL [EF] ^a Irbesartan GH [GQ]
			^B 2.50	18.36	17.15	^a Avapro [AV]	^a Karvea [SW]

irbesartan 150 mg tablet, 30

8247C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	14.14	15.43	^a Abisart 150 [AL] ^a AVSARTAN [RF] ^a Irbesartan Actavis 150 [ED] ^a Irbesartan AN [EA] ^a Irbesartan Sandoz [SZ]	^a APO-Irbesartan [TX] ^a Blooms the Chemist Irbesartan [IB] ^a Irbesartan AMNEAL [EF] ^a Irbesartan GH [GQ]
			^B 2.47	16.61	15.43	^a Avapro [AV]	^a Karvea [SW]

■ **LOSARTAN****losartan potassium 50 mg tablet, 30**

8203R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*27.66	28.95	Cozavan [AF]

losartan potassium 25 mg tablet, 30

5452Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	17.52	18.81	Cozavan [AF]

■ **OLMESARTAN****olmesartan medoxomil 40 mg tablet, 30**

2148C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	20.76	22.05	^a APO-Olmesartan [TX] ^a Olmesartan AN [EA] ^a Olmesartan Sandoz [SZ]	^a OLMERTAN [RW] ^a Olmesartan - MYL [AF] ^a Pharmacor Olmesartan 40 [CR]
			^B 3.51	24.27	22.05	^a Olmetec [AL]	

olmesartan medoxomil 20 mg tablet, 30

2147B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	16.26	17.55	^a APO-Olmesartan [TX] ^a Olmesartan AN [EA] ^a Olmesartan Sandoz [SZ]	^a OLMERTAN [RW] ^a Olmesartan - MYL [AF] ^a Pharmacor Olmesartan 20 [CR]
			^B 3.38	19.64	17.55	^a Olmetec [AL]	

■ **TELMISARTAN****telmisartan 80 mg tablet, 28**

8356T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	17.21	18.50	^a APO-Telmisartan [TX] ^a Pharmacor Telmisartan 80 [CR] ^a Telmisartan-DRLA [RZ] ^a Telmisartan Sandoz [SZ]	^a Mizart [AF] ^a Telmisartan AN [EA] ^a Telmisartan GH [GQ] ^a Teltartan [RW]
			^B 4.45	21.66	18.50	^a Micardis [BY]	

telmisartan 40 mg tablet, 28

8355R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	14.08	15.37	^a APO-Telmisartan [TX] ^a Pharmacor Telmisartan 40 [CR] ^a Telmisartan-DRLA [RZ] ^a Telmisartan Sandoz [SZ]	^a Mizart [AF] ^a Telmisartan AN [EA] ^a Telmisartan GH [GQ] ^a Teltartan [RW]
			^B 4.21	18.29	15.37	^a Micardis [BY]	

■ **VALSARTAN****valsartan 160 mg tablet, 28**

9370E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	21.20	22.49	^a Dilart [AF]	^a Diovan [NV]

valsartan 40 mg tablet, 28

9368C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	16.99	18.28	^a Dilart [AF]	^a Diovan [NV]

valsartan 80 mg tablet, 28

9369D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	19.17	20.46	^a Dilart [AF]	^a Diovan [NV]

■ **VALSARTAN**

Note No applications for increased maximum quantities and/or repeats will be authorised for the 320 mg tablet.

valsartan 320 mg tablet, 28

9371F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	23.79	25.08	^a Dilart [AF]	^a Diovan [NV]

ANGIOTENSIN II RECEPTOR BLOCKERS (ARBs), COMBINATIONS

Angiotensin II receptor blockers (ARBs) and diuretics

■ CANDESARTAN + HYDROCHLOROTHIAZIDE

Restricted benefit

Hypertension

Clinical criteria:

- The treatment must not be for the initiation of anti-hypertensive therapy, **AND**
- The condition must be inadequately controlled with an angiotensin II antagonist; OR
- The condition must be inadequately controlled with a thiazide diuretic.

candesartan cilexetil 32 mg + hydrochlorothiazide 25 mg tablet, 30

9315G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	18.03	19.32	^a Adesan HCT 32/25 [AF]	^a APO-Candesartan HCTZ 32/25 [TX]
						^a Asartan HCT 32/25 [DO]	^a Blooms the Chemist Candesartan HCTZ 32/25 [IB]
						^a CANDESAN COMBI 32/25 [RF]	^a Candesartan Combi Aspen 32/25 [RW]
						^a Candesartan HCT GH 32/25 [GQ]	^a Candesartan/HCT Sandoz [SZ]
						^a Candesartan HCTZ AN 32/25 [EA]	
			^B 13.47	31.50	19.32	^a Atacand Plus 32/25 [AP]	

candesartan cilexetil 32 mg + hydrochlorothiazide 12.5 mg tablet, 30

9314F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	17.25	18.54	^a Adesan HCT 32/12.5 [AF]	^a APO-Candesartan HCTZ 32/12.5 [TX]
						^a Asartan HCT 32/12.5 [DO]	^a Blooms the Chemist Candesartan HCTZ 32/12.5 [IB]
						^a CANDESAN COMBI 32/12.5 [RF]	^a Candesartan Combi Aspen 32/12.5 [RW]
						^a Candesartan HCT GH 32/12.5 [GQ]	^a Candesartan/HCT Sandoz [SZ]
						^a Candesartan HCTZ AN 32/12.5 [EA]	
			^B 13.42	30.67	18.54	^a Atacand Plus 32/12.5 [AP]	

candesartan cilexetil 16 mg + hydrochlorothiazide 12.5 mg tablet, 30

8504N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	16.55	17.84	^a Adesan HCT 16/12.5 [AF]	^a APO-Candesartan HCTZ 16/12.5 [TX]
						^a Asartan HCT 16/12.5 [DO]	^a Blooms the Chemist Candesartan HCTZ 16/12.5 [IB]
						^a CANDESAN COMBI 16/12.5 [RF]	^a Candesartan Combi Aspen 16/12.5 [RW]
						^a Candesartan HCT GH 16/12.5 [GQ]	^a Candesartan/HCT Sandoz [SZ]
						^a Candesartan HCTZ AN 16/12.5 [EA]	
			^B 13.37	29.92	17.84	^a Atacand Plus 16/12.5 [AP]	

■ EPROSARTAN + HYDROCHLOROTHIAZIDE

Restricted benefit

Hypertension

Clinical criteria:

- The treatment must not be for the initiation of anti-hypertensive therapy, **AND**
- The condition must be inadequately controlled with an angiotensin II antagonist; OR
- The condition must be inadequately controlled with a thiazide diuretic.

eprosartan 600 mg + hydrochlorothiazide 12.5 mg tablet, 28

8624X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	28.99	30.28	Teveten Plus 600/12.5 [GO]

■ IRBESARTAN + HYDROCHLOROTHIAZIDE

Restricted benefit

Hypertension

Clinical criteria:

- The treatment must not be for the initiation of anti-hypertensive therapy, **AND**
- The condition must be inadequately controlled with an angiotensin II antagonist; OR
- The condition must be inadequately controlled with a thiazide diuretic.

irbesartan 150 mg + hydrochlorothiazide 12.5 mg tablet, 30

8404H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	14.39	15.68	^a Abisart HCTZ 150/12.5 [AL] ^a AVSARTAN HCT 150/12.5 [RF] ^a Irbesartan HCT Actavis 150/12.5 [ED] ^a Irbesartan/HCT Sandoz [SZ] ^a KSART HCT 150/12.5 [RW]	^a APO-Irbesartan HCTZ [TX] ^a Blooms the Chemist Irbesartan HCTZ 150/12.5 [IB] ^a Irbesartan HCT GH 150/12.5 [GQ] ^a Irbesartan HCTZ AMNEAL [EF]
			^B 2.47	16.86	15.68	^a Avapro HCT 150/12.5 [AV]	^a Karvezide 150/12.5 [SW]

irbesartan 300 mg + hydrochlorothiazide 12.5 mg tablet, 30

8405J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	16.03	17.32	^a Abisart HCTZ 300/12.5 [AL] ^a AVSARTAN HCT 300/12.5 [RF] ^a Irbesartan HCT Actavis 300/12.5 [ED] ^a Irbesartan/HCT Sandoz [SZ] ^a KSART HCT 300/12.5 [RW]	^a APO-Irbesartan HCTZ [TX] ^a Blooms the Chemist Irbesartan HCTZ 300/12.5 [IB] ^a Irbesartan HCT GH 300/12.5 [GQ] ^a Irbesartan HCTZ AMNEAL [EF]
			^B 2.51	18.54	17.32	^a Avapro HCT 300/12.5 [AV]	^a Karvezide 300/12.5 [SW]

irbesartan 300 mg + hydrochlorothiazide 25 mg tablet, 30

2136K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	16.35	17.64	^a Abisart HCTZ 300/25 [AL] ^a AVSARTAN HCT 300/25 [RF] ^a Irbesartan HCT Actavis 300/25 [ED] ^a Irbesartan/HCT Sandoz [SZ] ^a KSART HCT 300/25 [RW]	^a APO-Irbesartan HCTZ [TX] ^a Blooms the Chemist Irbesartan HCTZ 300/25 [IB] ^a Irbesartan HCT GH 300/25 [GQ] ^a Irbesartan HCTZ AMNEAL [EF]
			^B 2.53	18.88	17.64	^a Avapro HCT 300/25 [AV]	^a Karvezide 300/25 [SW]

▪ **OLMESARTAN MEDOXOMIL + HYDROCHLOROTHIAZIDE**

Restricted benefit

Hypertension

Clinical criteria:

- The treatment must not be for the initiation of anti-hypertensive therapy, **AND**
- The condition must be inadequately controlled with an angiotensin II antagonist; OR
- The condition must be inadequately controlled with a thiazide diuretic.

olmesartan medoxomil 20 mg + hydrochlorothiazide 12.5 mg tablet, 30

2161R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	17.00	18.29	^a APO-Olmesartan/HCTZ 20/12.5 [TX] ^a Olmesartan HCT AN 20/12.5 [EA] ^a Olmesartan/HCT Sandoz [SZ] ^a Olmetec Plus [AL]	^a OLMERTAN COMBI 20/12.5 [RW] ^a Olmesartan HCT - MYL 20/12.5 [AF] ^a Pharmacor Olmesartan HCTZ 20/12.5 [CR]
			^B 1.51	18.51	18.29		

olmesartan medoxomil 40 mg + hydrochlorothiazide 25 mg tablet, 30

2170F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	22.31	23.60	^a APO-Olmesartan/HCTZ 40/25 [TX] ^a Olmesartan HCT AN 40/25 [EA] ^a Olmesartan/HCT Sandoz [SZ] ^a Olmetec Plus [AL]	^a OLMERTAN COMBI 40/25 [RW] ^a Olmesartan HCT - MYL 40/25 [AF] ^a Pharmacor Olmesartan HCTZ 40/25 [CR]
			^B 1.59	23.90	23.60		

olmesartan medoxomil 40 mg + hydrochlorothiazide 12.5 mg tablet, 30

2166B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	21.12	22.41	^a APO-Olmesartan/HCTZ 40/12.5 [TX] ^a Olmesartan HCT AN 40/12.5 [EA] ^a Olmesartan/HCT Sandoz [SZ] ^a Olmetec Plus [AL]	^a OLMERTAN COMBI 40/12.5 [RW] ^a Olmesartan HCT - MYL 40/12.5 [AF] ^a Pharmacor Olmesartan HCTZ 40/12.5 [CR]
			^B 5.00	26.12	22.41		

■ TELMISARTAN + HYDROCHLOROTHIAZIDE

Restricted benefit

Hypertension

Clinical criteria:

- The treatment must not be for the initiation of anti-hypertensive therapy, **AND**
- The condition must be inadequately controlled with an angiotensin II antagonist; OR
- The condition must be inadequately controlled with a thiazide diuretic.

telmisartan 80 mg + hydrochlorothiazide 12.5 mg tablet, 28

8623W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	17.72	19.01	^a APO-Telmisartan HCTZ 80/12.5 [TX]	^a Mizart HCT 80/12.5 mg [AF]
						^a Telmisartan HCT GH 80/12.5 [GQ]	^a Telmisartan/HCT Sandoz [SZ]
						^a Telmisartan HCTZ AN 80/12.5 [EA]	^a Teltartan HCT 80/12.5 [RW]
			^B 4.49	22.21	19.01	^a Micardis Plus 80/12.5 mg [BY]	

telmisartan 80 mg + hydrochlorothiazide 25 mg tablet, 28

9381R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	18.21	19.50	^a APO-Telmisartan HCTZ 80/25 [TX]	^a Mizart HCT 80/25 mg [AF]
						^a Telmisartan HCT GH 80/25 [GQ]	^a Telmisartan/HCT Sandoz [SZ]
						^a Telmisartan HCTZ AN 80/25 [EA]	^a Teltartan HCT 80/25 [RW]
			^B 4.50	22.71	19.50	^a Micardis Plus 80/25 mg [BY]	

telmisartan 40 mg + hydrochlorothiazide 12.5 mg tablet, 28

8622T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	14.55	15.84	^a APO-Telmisartan HCTZ 40/12.5 [TX]	^a Mizart HCT 40/12.5 mg [AF]
						^a Telmisartan HCT GH 40/12.5 [GQ]	^a Telmisartan/HCT Sandoz [SZ]
						^a Telmisartan HCTZ AN 40/12.5 [EA]	^a Teltartan HCT 40/12.5 [RW]
			^B 4.25	18.80	15.84	^a Micardis Plus 40/12.5 mg [BY]	

■ VALSARTAN + HYDROCHLOROTHIAZIDE

Restricted benefit

Hypertension

Clinical criteria:

- The treatment must not be for the initiation of anti-hypertensive therapy, **AND**
- The condition must be inadequately controlled with an angiotensin II antagonist; OR
- The condition must be inadequately controlled with a thiazide diuretic.

valsartan 160 mg + hydrochlorothiazide 12.5 mg tablet, 28

9373H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	21.27	22.56	^a Co-Diovan 160/12.5 [NV]	^a Dilart HCT 160/12.5 [AF]

valsartan 160 mg + hydrochlorothiazide 25 mg tablet, 28

9374J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	22.35	23.64	^a Co-Diovan 160/25 [NV]	^a Dilart HCT 160/25 [AF]

valsartan 80 mg + hydrochlorothiazide 12.5 mg tablet, 28

9372G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	19.46	20.75	^a Co-Diovan 80/12.5 [NV]	^a Dilart HCT 80/12.5 [AF]

■ VALSARTAN + HYDROCHLOROTHIAZIDE

Note No applications for increased maximum quantities and/or repeats will be authorised for the tablets containing 320 mg valsartan.

Restricted benefit

Hypertension

Clinical criteria:

- The treatment must not be for the initiation of anti-hypertensive therapy, **AND**
- The condition must be inadequately controlled with an angiotensin II antagonist; OR
- The condition must be inadequately controlled with a thiazide diuretic.

valsartan 320 mg + hydrochlorothiazide 12.5 mg tablet, 28

9481B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	23.58	24.87	^a Co-Diovan 320/12.5 [NV]	^a Dilart HCT 320/12.5 [AF]

valsartan 320 mg + hydrochlorothiazide 25 mg tablet, 28

9482C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	24.65	25.94	^a Co-Diovan 320/25 [NV]	^a Dilart HCT 320/25 [AF]

Angiotensin II receptor blockers (ARBs) and calcium channel blockers**■ AMLODIPINE + VALSARTAN****Restricted benefit**

Hypertension

Clinical criteria:

- The treatment must not be for the initiation of anti-hypertensive therapy, **AND**
- The condition must be inadequately controlled with an angiotensin II antagonist; OR
- The condition must be inadequately controlled with a dihydropyridine calcium channel blocker.

amlodipine 5 mg + valsartan 160 mg tablet, 28

9376L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	22.03	23.32	^a Valsartan/Amlodipine Novartis 160/5 [NM]
			^B 3.00	25.03	23.32	^a Exforge 5/160 [NV]

amlodipine 10 mg + valsartan 320 mg tablet, 28

5460J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	25.27	26.56	^a Valsartan/Amlodipine Novartis 320/10 [NM]
			^B 3.00	28.27	26.56	^a Exforge 10/320 [NV]

amlodipine 5 mg + valsartan 80 mg tablet, 28

9375K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	20.00	21.29	^a Valsartan/Amlodipine Novartis 80/5 [NM]
			^B 3.00	23.00	21.29	^a Exforge 5/80 [NV]

amlodipine 10 mg + valsartan 160 mg tablet, 28

9377M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	22.68	23.97	^a Valsartan/Amlodipine Novartis 160/10 [NM]
			^B 3.00	25.68	23.97	^a Exforge 10/160 [NV]

amlodipine 5 mg + valsartan 320 mg tablet, 28

5459H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	24.62	25.91	^a Valsartan/Amlodipine Novartis 320/5 [NM]
			^B 3.00	27.62	25.91	^a Exforge 5/320 [NV]

■ OLMESARTAN MEDOXOMIL + AMLODIPINE**Restricted benefit**

Hypertension

Clinical criteria:

- The treatment must not be for the initiation of anti-hypertensive therapy, **AND**
- The condition must be inadequately controlled with an angiotensin II antagonist; OR
- The condition must be inadequately controlled with a dihydropyridine calcium channel blocker.

olmesartan medoxomil 40 mg + amlodipine 5 mg tablet, 30

5293N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	20.52	21.81	^a OLMEKAR [RW]	^a Olmesartan/Amlodipine 40/5 APOTEX [TX]
						^a Olmesartan/Amlodipine - MYL 40/5 [AF]	^a Pharmacor Olmesartan Amlodipine 40/5 [CR]
			^B 2.65	23.17	21.81	^a Sevikar 40/5 [AL]	

olmesartan medoxomil 40 mg + amlodipine 10 mg tablet, 30

5294P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	21.14	22.43	^a OLMEKAR [RW]	^a Olmesartan/Amlodipine 40/10 APOTEX [TX]
						^a Olmesartan/Amlodipine - MYL 40/10 [AF]	^a Pharmacor Olmesartan Amlodipine 40/10 [CR]
			^B 2.64	23.78	22.43	^a Sevikar 40/10 [AL]	

olmesartan medoxomil 20 mg + amlodipine 5 mg tablet, 30

5292M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	16.54	17.83	^a OLMEKAR [RW]	^a Olmesartan/Amlodipine 20/5 APOTEX [TX]
						^a Olmesartan/Amlodipine - MYL 20/5 [AF]	^a Pharmacor Olmesartan Amlodipine 20/5 [CR]
						^a Sevikaar 20/5 [AL]	

▪ **TELMISARTAN + AMLODIPINE**

Restricted benefit

Hypertension

Clinical criteria:

- The treatment must not be for the initiation of anti-hypertensive therapy, **AND**
- The condition must be inadequately controlled with an angiotensin II antagonist; OR
- The condition must be inadequately controlled with a dihydropyridine calcium channel blocker.

telmisartan 80 mg + amlodipine 10 mg tablet, 28

8981Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	18.63	19.92	^a Pritor/Amlodipine [FI]
			^B 4.50	23.13	19.92	^a Twynsta [BY]

telmisartan 80 mg + amlodipine 5 mg tablet, 28

8980P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	17.99	19.28	^a Pritor/Amlodipine [FI]
			^B 4.50	22.49	19.28	^a Twynsta [BY]

telmisartan 40 mg + amlodipine 10 mg tablet, 28

8979N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	15.45	16.74	^a Pritor/Amlodipine [FI]
			^B 4.32	19.77	16.74	^a Twynsta [BY]

telmisartan 40 mg + amlodipine 5 mg tablet, 28

8978M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	14.85	16.14	^a Pritor/Amlodipine [FI]
			^B 4.27	19.12	16.14	^a Twynsta [BY]

Angiotensin II receptor blockers (ARBs), other combinations

▪ **AMLODIPINE + VALSARTAN + HYDROCHLOROTHIAZIDE**

Restricted benefit

Hypertension

Clinical criteria:

- The treatment must not be for the initiation of anti-hypertensive therapy, **AND**
- The condition must be inadequately controlled with concomitant treatment with two of the following: an angiotensin II antagonist, a dihydropyridine calcium channel blocker or a thiazide diuretic.

amlodipine 10 mg + valsartan 160 mg + hydrochlorothiazide 12.5 mg tablet, 28

5287G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	24.19	25.48	^a Valsartan/Amlodipine/HCT Novartis 160/10/12.5 [NM]
			^B 3.00	27.19	25.48	^a Exforge HCT 10/160/12.5 [NV]

amlodipine 10 mg + valsartan 160 mg + hydrochlorothiazide 25 mg tablet, 28

5288H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	25.71	27.00	^a Valsartan/Amlodipine/HCT Novartis 160/10/25 [NM]
			^B 3.00	28.71	27.00	^a Exforge HCT 10/160/25 [NV]

amlodipine 5 mg + valsartan 160 mg + hydrochlorothiazide 12.5 mg tablet, 28

5285E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	23.55	24.84	^a Valsartan/Amlodipine/HCT Novartis 160/5/12.5 [NM]
			^B 3.00	26.55	24.84	^a Exforge HCT 5/160/12.5 [NV]

amlodipine 5 mg + valsartan 160 mg + hydrochlorothiazide 25 mg tablet, 28

5286F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	25.06	26.35	^a Valsartan/Amlodipine/HCT Novartis 160/5/25 [NM]
			^B 3.00	28.06	26.35	^a Exforge HCT 5/160/25 [NV]

amlodipine 10 mg + valsartan 320 mg + hydrochlorothiazide 25 mg tablet, 28

5289J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	28.30	29.59	^a Valsartan/Amlodipine/HCT Novartis 320/10/25 [NM]

			^B 3.00	31.30	29.59	^a Exforge HCT 10/320/25 [NV]	
■ OLMESARTAN + AMLODIPINE + HYDROCHLOROTHIAZIDE							
Restricted benefit							
Hypertension							
Clinical criteria:							
- The treatment must not be for the initiation of anti-hypertensive therapy, AND - The condition must be inadequately controlled with concomitant treatment with two of the following: an angiotensin II antagonist, a dihydropyridine calcium channel blocker or a thiazide diuretic.							
olmesartan medoxomil 40 mg + amlodipine 10 mg + hydrochlorothiazide 12.5 mg tablet, 30							
2836G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	22.71	24.00	^a APO-Olmesartan/Amlodipine/HCTZ 40/10/12.5 [TX]	^a Olmekar HCT 40/10/12.5 [RF]
			^B 2.70	25.41	24.00	^a Sevikar HCT 40/10/12.5 [AF]	
olmesartan medoxomil 40 mg + amlodipine 5 mg + hydrochlorothiazide 12.5 mg tablet, 30							
2880N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	22.09	23.38	^a APO-Olmesartan/Amlodipine/HCTZ 40/5/12.5 tablet [TX]	^a Olmekar HCT 40/5/12.5 [RF]
			^B 2.70	24.79	23.38	^a Sevikar HCT 40/5/12.5 [AF]	
olmesartan medoxomil 40 mg + amlodipine 5 mg + hydrochlorothiazide 25 mg tablet, 30							
2864R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	23.66	24.95	^a APO-Olmesartan/Amlodipine/HCTZ 40/5/25 tablet [TX]	^a Olmekar HCT 40/5/25 [RF]
			^B 2.74	26.40	24.95	^a Sevikar HCT 40/5/25 [AF]	
olmesartan medoxomil 40 mg + amlodipine 10 mg + hydrochlorothiazide 25 mg tablet, 30							
2953K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	24.28	25.57	^a APO-Olmesartan/Amlodipine/HCTZ 40/10/25 [TX]	^a Olmekar HCT 40/10/25 [RF]
			^B 2.73	27.01	25.57	^a Sevikar HCT 40/10/25 [AF]	
olmesartan medoxomil 20 mg + amlodipine 5 mg + hydrochlorothiazide 12.5 mg tablet, 30							
10005N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	18.01	19.30	^a APO-Olmesartan/Amlodipine/HCTZ 20/5/12.5 [TX]	^a Olmekar HCT 20/5/12.5 [RF]
			^B 2.73	20.74	19.30	^a Sevikar HCT 20/5/12.5 [AF]	

■ SACUBITRIL + VALSARTAN

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

6915

Chronic heart failure

Clinical criteria:

- Patient must be symptomatic with NYHA classes II, III or IV, **AND**
- Patient must have a documented left ventricular ejection fraction (LVEF) of less than or equal to 40%, **AND**
- Patient must receive concomitant optimal standard chronic heart failure treatment, which must include the maximum tolerated dose of a beta-blocker, unless contraindicated or not tolerated, **AND**
- Patient must have been stabilised on an ACE inhibitor at the time of initiation with this drug, unless such treatment is contraindicated according to the TGA-approved Product Information or cannot be tolerated; OR
- Patient must have been stabilised on an angiotensin II antagonist at the time of initiation with this drug, unless such treatment is contraindicated according to the TGA-approved Product Information or cannot be tolerated, **AND**
- The treatment must not be co-administered with an ACE inhibitor or an angiotensin II antagonist.

sacubitril 48.6 mg + valsartan 51.4 mg tablet, 56

11131W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	202.89	41.30	Entresto [NV]

sacubitril 24.3 mg + valsartan 25.7 mg tablet, 56

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
11123K	1	5	..	202.89	41.30	Entresto [NV]

sacubitril 97.2 mg + valsartan 102.8 mg tablet, 56

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
11122J	1	5	..	202.89	41.30	Entresto [NV]

■ LIPID MODIFYING AGENTS**LIPID MODIFYING AGENTS, PLAIN***HMG CoA reductase inhibitors***■ ATORVASTATIN****atorvastatin 10 mg tablet, 30**

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
8213G	1	5	..	13.85	15.14	^a APO-Atorvastatin [TX] ^a Atorvastatin Amneal [EF] ^a Atorvastatin SZ [HX] ^a Lipitor [AS] ^a NOUMED ATORVASTATIN [VO] ^a Torvastat 10 [RW]	^a Atorvachol [RF] ^a Atorvastatin GH [GQ] ^a Blooms the Chemist Atorvastatin [IB] ^a Lorstat 10 [AF] ^a Pharmacor Atorvastatin [CR] ^a Trovas [RA]

atorvastatin 20 mg tablet, 30

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
8214H	1	5	..	14.53	15.82	^a APO-Atorvastatin [TX] ^a Atorvastatin Amneal [EF] ^a Atorvastatin SZ [HX] ^a Lipitor [AS] ^a NOUMED ATORVASTATIN [VO] ^a Torvastat 20 [RW]	^a Atorvachol [RF] ^a Atorvastatin GH [GQ] ^a Blooms the Chemist Atorvastatin [IB] ^a Lorstat 20 [AF] ^a Pharmacor Atorvastatin [CR] ^a Trovas [RA]

atorvastatin 80 mg tablet, 30

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
8521L	1	5	..	16.65	17.94	^a APO-Atorvastatin [TX] ^a Atorvastatin Amneal [EF] ^a Atorvastatin SZ [HX] ^a Lipitor [AS] ^a NOUMED ATORVASTATIN [VO] ^a Torvastat 80 [RW]	^a Atorvachol [RF] ^a Atorvastatin GH [GQ] ^a Blooms the Chemist Atorvastatin [IB] ^a Lorstat 80 [AF] ^a Pharmacor Atorvastatin [CR] ^a Trovas [RA]

atorvastatin 40 mg tablet, 30

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
8215J	1	5	..	15.38	16.67	^a APO-Atorvastatin [TX] ^a Atorvastatin Amneal [EF] ^a Atorvastatin SZ [HX] ^a Lipitor [AS] ^a NOUMED ATORVASTATIN [VO] ^a Torvastat 40 [RW]	^a Atorvachol [RF] ^a Atorvastatin GH [GQ] ^a Blooms the Chemist Atorvastatin [IB] ^a Lorstat 40 [AF] ^a Pharmacor Atorvastatin [CR] ^a Trovas [RA]

■ ATORVASTATIN**Note** No increase in the maximum quantity or number of units may be authorised.**Note** No increase in the maximum number of repeats may be authorised.**Restricted benefit**

For use in patients who are receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.

atorvastatin 10 mg tablet, 30

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
9230T	1	11	..	13.85	15.14	^a APO-Atorvastatin [TX] ^a Atorvastatin Amneal [EF] ^a Atorvastatin SZ [HX]	^a Atorvachol [RF] ^a Atorvastatin GH [GQ] ^a Blooms the Chemist Atorvastatin [IB]

^a Lipitor [AS]
^a NOUMED ATORVASTATIN [VO]
^a Torvastat 10 [RW]
^a Lorstat 10 [AF]
^a Pharmacor Atorvastatin [CR]
^a Trovas [RA]

atorvastatin 20 mg tablet, 30

9231W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	11	..	14.53	15.82	^a APO-Atorvastatin [TX] ^a Atorvastatin Amneal [EF] ^a Atorvastatin SZ [HX] ^a Lipitor [AS] ^a NOUMED ATORVASTATIN [VO] ^a Torvastat 20 [RW]	^a Atorvachol [RF] ^a Atorvastatin GH [GQ] ^a Blooms the Chemist Atorvastatin [IB] ^a Lorstat 20 [AF] ^a Pharmacor Atorvastatin [CR] ^a Trovas [RA]

atorvastatin 80 mg tablet, 30

9233Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	11	..	16.65	17.94	^a APO-Atorvastatin [TX] ^a Atorvastatin Amneal [EF] ^a Atorvastatin SZ [HX] ^a Lipitor [AS] ^a NOUMED ATORVASTATIN [VO] ^a Torvastat 80 [RW]	^a Atorvachol [RF] ^a Atorvastatin GH [GQ] ^a Blooms the Chemist Atorvastatin [IB] ^a Lorstat 80 [AF] ^a Pharmacor Atorvastatin [CR] ^a Trovas [RA]

atorvastatin 40 mg tablet, 30

9232X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	11	..	15.38	16.67	^a APO-Atorvastatin [TX] ^a Atorvastatin Amneal [EF] ^a Atorvastatin SZ [HX] ^a Lipitor [AS] ^a NOUMED ATORVASTATIN [VO] ^a Torvastat 40 [RW]	^a Atorvachol [RF] ^a Atorvastatin GH [GQ] ^a Blooms the Chemist Atorvastatin [IB] ^a Lorstat 40 [AF] ^a Pharmacor Atorvastatin [CR] ^a Trovas [RA]

■ **FLUVASTATIN**

fluvastatin 80 mg modified release tablet, 28

2863Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	46.52	41.30	Lescol XL [NV]

■ **FLUVASTATIN**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Restricted benefit

For use in patients who are receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.

fluvastatin 80 mg modified release tablet, 28

9236D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	11	..	46.52	41.30	Lescol XL [NV]

■ **PRAVASTATIN**

pravastatin sodium 40 mg tablet, 30

8197K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	15.96	17.25	^a APO-Pravastatin [TX] ^a Lipostat 40 [RF] ^a Pravastatin Sandoz [SZ]	^a Cholstat 40 [AF] ^a Pravastatin AN [EA]
			^b 2.87	18.83	17.25	^a Pravachol [RW]	

pravastatin sodium 80 mg tablet, 30

8829Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	17.86	19.15	^a APO-Pravastatin [TX] ^a Pravastatin AN [EA] ^a Pravastatin Sandoz [SZ]	^a Lipostat 80 [RF] ^a Pravastatin generichealth [GQ]
			^b 3.01	20.87	19.15	^a Pravachol [RW]	

pravastatin sodium 20 mg tablet, 30

2834E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	14.70	15.99	^a APO-Pravastatin [TX]	^a Cholstat 20 [AF]
						^a Lipostat 20 [RF]	^a Pravastatin AN [EA]
						^a Pravastatin generichealth [GQ]	^a Pravastatin Sandoz [SZ]
			^B 2.75	17.45	15.99	^a Pravachol [RW]	

pravastatin sodium 10 mg tablet, 30

2833D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	13.86	15.15	^a APO-Pravastatin [TX]	^a Lipostat 10 [RF]
						^a Pravastatin AN [EA]	^a Pravastatin Sandoz [SZ]
			^B 2.74	16.60	15.15	^a Pravachol [RW]	
			^B 2.75	16.61	15.15	^a Cholstat 10 [AF]	

■ **PRAVASTATIN**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Restricted benefit

For use in patients who are receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.

pravastatin sodium 40 mg tablet, 30

9239G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	11	..	15.96	17.25	^a APO-Pravastatin [TX]	^a Cholstat 40 [AF]
						^a Lipostat 40 [RF]	^a Pravastatin AN [EA]
						^a Pravastatin Sandoz [SZ]	
			^B 2.87	18.83	17.25	^a Pravachol [RW]	

pravastatin sodium 80 mg tablet, 30

9240H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	11	..	17.86	19.15	^a APO-Pravastatin [TX]	^a Lipostat 80 [RF]
						^a Pravastatin AN [EA]	^a Pravastatin generichealth [GQ]
						^a Pravastatin Sandoz [SZ]	
			^B 3.01	20.87	19.15	^a Pravachol [RW]	

pravastatin sodium 20 mg tablet, 30

9238F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	11	..	14.70	15.99	^a APO-Pravastatin [TX]	^a Cholstat 20 [AF]
						^a Lipostat 20 [RF]	^a Pravastatin AN [EA]
						^a Pravastatin generichealth [GQ]	^a Pravastatin Sandoz [SZ]
			^B 2.75	17.45	15.99	^a Pravachol [RW]	

pravastatin sodium 10 mg tablet, 30

9237E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	11	..	13.86	15.15	^a APO-Pravastatin [TX]	^a Lipostat 10 [RF]
						^a Pravastatin AN [EA]	^a Pravastatin Sandoz [SZ]
			^B 2.74	16.60	15.15	^a Pravachol [RW]	
			^B 2.75	16.61	15.15	^a Cholstat 10 [AF]	

■ **ROSUVASTATIN**

rosuvastatin 5 mg tablet, 30

2606E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	14.13	15.42	^a APO-Rosuvastatin [TX]	^a Blooms the Chemist Rosuvastatin [IB]
						^a BTC Rosuvastatin [JB]	^a Cavstat [AF]
						^a Crosuva 5 [RW]	^a Noumed Rosuvastatin [VO]
						^a Pharmacor Rosuvastatin 5 [CR]	^a Rostor 5 [DO]
						^a Rosuvastatin AMNEAL [EF]	^a Rosuvastatin APOTEX [GX]
						^a Rosuvastatin generichealth [HQ]	^a Rosuvastatin Lupin [GQ]
						^a Rosuvastatin RBX [RA]	^a Rosuvastatin Sandoz [SZ]
			^B 1.39	15.52	15.42	^a Crestor [AP]	

rosuvastatin 20 mg tablet, 30

2574L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	15.98	17.27	^a APO-Rosuvastatin [TX]	^a Blooms the Chemist Rosuvastatin [IB]
						^a BTC Rosuvastatin [JB]	^a Cavstat [AF]
						^a Crosuva 20 [RW]	^a Noumed Rosuvastatin [VO]

rosuvastatin 40 mg tablet, 30rosuvastatin 10 mg tablet, 30

▪ ROSUVASTATIN

Note No increase in the maximum number of repeats may be authorised.

For use in patients who are receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.

rosuvastatin 20 mg tablet, 30

Schedule of Pharmaceutical Benefits – April 2021

rosuvastatin 40 mg tablet, 30

2636R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	11	..	17.49	18.78	^a APO-Rosuvastatin [TX]	^a Blooms the Chemist Rosuvastatin [IB]
						^a BTC Rosuvastatin [JB]	^a Cavstat [AF]
						^a Crosuva 40 [RW]	^a Noumed Rosuvastatin [VO]
						^a Pharmacor Rosuvastatin 40 [CR]	^a Rostor 40 [DO]
						^a Rosuvastatin AMNEAL [EF]	^a Rosuvastatin APOTEX [GX]
						^a Rosuvastatin generichealth [HQ]	^a Rosuvastatin Lupin [GQ]
						^a Rosuvastatin RBX [RA]	^a Rosuvastatin Sandoz [SZ]
			^B 2.13	19.62	18.78	^a Crestor [AP]	

rosuvastatin 10 mg tablet, 30

2584B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	11	..	15.59	16.88	^a APO-Rosuvastatin [TX]	^a Blooms the Chemist Rosuvastatin [IB]
						^a BTC Rosuvastatin [JB]	^a Cavstat [AF]
						^a Crosuva 10 [RW]	^a Noumed Rosuvastatin [VO]
						^a Pharmacor Rosuvastatin 10 [CR]	^a Rostor 10 [DO]
						^a Rosuvastatin AMNEAL [EF]	^a Rosuvastatin APOTEX [GX]
						^a Rosuvastatin generichealth [HQ]	^a Rosuvastatin Lupin [GQ]
						^a Rosuvastatin RBX [RA]	^a Rosuvastatin Sandoz [SZ]
			^B 2.00	17.59	16.88	^a Crestor [AP]	

■ **SIMVASTATIN**

simvastatin 20 mg tablet, 30

2012X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	13.94	15.23	^a APO-Simvastatin [TX]	^a Simvar 20 [RW]
						^a Simvastatin AN [EA]	^a Simvastatin generichealth [GQ]
						^a Simvastatin Sandoz [SZ]	^a Zimstat [AF]
			^B 7.20	21.14	15.23	^a Lipex 20 [AL]	^a Zocor [MQ]

simvastatin 10 mg tablet, 30

2011W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	13.47	14.76	^a APO-Simvastatin [TX]	^a Ransim [RA]
						^a Simvar 10 [RW]	^a Simvastatin AN [EA]
						^a Simvastatin generichealth [GQ]	^a Simvastatin Sandoz [SZ]
						^a Zimstat [AF]	
			^B 5.00	18.47	14.76	^a Lipex 10 [AL]	^a Zocor [MQ]

simvastatin 40 mg tablet, 30

8173E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	14.60	15.89	^a APO-Simvastatin [TX]	^a Simvar 40 [RW]
						^a Simvastatin AN [EA]	^a Simvastatin generichealth [GQ]
						^a Simvastatin Sandoz [SZ]	^a Zimstat [AF]
			^B 7.25	21.85	15.89	^a Lipex 40 [AL]	^a Zocor [MQ]

simvastatin 5 mg tablet, 30

2013Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	13.19	14.48	^a Simvastatin Sandoz [SZ]	^a Zimstat [AF]

simvastatin 80 mg tablet, 30

8313M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	15.53	16.82	^a APO-Simvastatin [TX]	^a Ransim [RA]
						^a Simvar 80 [RW]	^a Simvastatin AN [EA]
						^a Simvastatin generichealth [GQ]	^a Simvastatin Sandoz [SZ]
						^a Zimstat [AF]	

■ **SIMVASTATIN**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Restricted benefit

CARDIOVASCULAR SYSTEM

For use in patients who are receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.

simvastatin 20 mg tablet, 30

9243L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	11	..	13.94	15.23	^a APO-Simvastatin [TX] ^a Simvastatin AN [EA] ^a Simvastatin Sandoz [SZ]	^a Simvar 20 [RW] ^a Simvastatin generichealth [GQ] ^a Zimstat [AF]
			^b 7.20	21.14	15.23	^a Lipex 20 [AL]	^a Zocor [MQ]

simvastatin 10 mg tablet, 30

9242K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	11	..	13.47	14.76	^a APO-Simvastatin [TX] ^a Simvar 10 [RW] ^a Simvastatin generichealth [GQ] ^a Zimstat [AF]	^a Ransim [RA] ^a Simvastatin AN [EA] ^a Simvastatin Sandoz [SZ]
			^b 5.00	18.47	14.76	^a Lipex 10 [AL]	^a Zocor [MQ]

simvastatin 40 mg tablet, 30

9244M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	11	..	14.60	15.89	^a APO-Simvastatin [TX] ^a Simvastatin AN [EA] ^a Simvastatin Sandoz [SZ]	^a Simvar 40 [RW] ^a Simvastatin generichealth [GQ] ^a Zimstat [AF]
			^b 7.25	21.85	15.89	^a Lipex 40 [AL]	^a Zocor [MQ]

simvastatin 5 mg tablet, 30

9241J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	11	..	13.19	14.48	^a Simvastatin Sandoz [SZ]	^a Zimstat [AF]

simvastatin 80 mg tablet, 30

9245N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	11	..	15.53	16.82	^a APO-Simvastatin [TX] ^a Simvar 80 [RW] ^a Simvastatin generichealth [GQ] ^a Zimstat [AF]	^a Ransim [RA] ^a Simvastatin AN [EA] ^a Simvastatin Sandoz [SZ]

Fibrates

■ FENOFIBRATE

Note The risk of serious muscle toxicity is increased if this drug is used concomitantly with HMG CoA reductase inhibitors or other fibrates. Such combination therapy should be used with caution in patients with severe combined dyslipidaemia and high cardiovascular risk without any history of muscular disease and patients monitored closely for chronic signs of muscle toxicity.

fenofibrate 145 mg tablet, 30

9203X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	28.44	29.73	^a APO-Fenofibrate [TX] ^a Fenocol [YC] ^a Fenofibrate Mylan [AF] ^a Fenofibrate Sandoz [SZ]	^a Blooms the Chemist Fenofibrate [IB] ^a Fenofibrate Cipla [LR] ^a FENOFIBRATE RBX [RA] ^a Lipidil [GO]

fenofibrate 48 mg tablet, 60

9202W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	22.97	24.26	^a APO-Fenofibrate [TX] ^a FENOFIBRATE RBX [RA]	^a Fenofibrate Mylan [AF] ^a Lipidil [GO]

■ FENOFIBRATE

Note The risk of serious muscle toxicity is increased if this drug is used concomitantly with HMG CoA reductase inhibitors or other fibrates. Such combination therapy should be used with caution in patients with severe combined dyslipidaemia and high cardiovascular risk without any history of muscular disease and patients monitored closely for chronic signs of muscle toxicity.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Restricted benefit

For use in patients who are receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.

fenofibrate 145 mg tablet, 30

9247Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	11	..	28.44	29.73	^a APO-Fenofibrate [TX]	^a Blooms the Chemist Fenofibrate [IB]
						^a Fenocol [YC]	^a Fenofibrate Cipla [LR]
						^a Fenofibrate Mylan [AF]	^a FENOFIBRATE RBX [RA]
						^a Fenofibrate Sandoz [SZ]	^a Lipidil [GO]

fenofibrate 48 mg tablet, 60

9246P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	11	..	22.97	24.26	^a APO-Fenofibrate [TX]	^a Fenofibrate Mylan [AF]
						^a FENOFIBRATE RBX [RA]	^a Lipidil [GO]

■ GEMFIBROZIL

Note The risk of serious muscle toxicity is increased if this drug is used concomitantly with HMG CoA reductase inhibitors or other fibrates. Such combination therapy should be used with caution in patients with severe combined dyslipidaemia and high cardiovascular risk without any history of muscular disease and patients monitored closely for chronic signs of muscle toxicity.

gemfibrozil 600 mg tablet, 60

1453L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	21.92	23.21	^a Ausgem [RW]	^a Lipigem [AF]

NP

■ GEMFIBROZIL

Note The risk of serious muscle toxicity is increased if this drug is used concomitantly with HMG CoA reductase inhibitors or other fibrates. Such combination therapy should be used with caution in patients with severe combined dyslipidaemia and high cardiovascular risk without any history of muscular disease and patients monitored closely for chronic signs of muscle toxicity.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Restricted benefit

For use in patients who are receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.

gemfibrozil 600 mg tablet, 60

9248R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	11	..	21.92	23.21	^a Ausgem [RW]	^a Lipigem [AF]

Bile acid sequestrants**■ COLESTYRAMINE****colestyramine 4 g powder for oral liquid, 50 sachets**

2967E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*81.44	41.30	Questran Lite [GO]

NP

■ COLESTYRAMINE

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Restricted benefit

Primary hypercholesterolaemia

Clinical criteria:

- Patient must be receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.

colestyramine 4 g powder for oral liquid, 50 sachets

9249T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	11	..	*81.44	41.30	Questran Lite [GO]

Other lipid modifying agents**■ EVOLOCUMAB**

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)**10388**

Familial homozygous hypercholesterolaemia

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be in conjunction with dietary therapy and exercise.

evolocumab 420 mg/3.5 mL injection, 3.5 mL cartridge

11972D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	539.12	41.30	Repatha [AN]

evolocumab 140 mg/mL injection, 1 mL pen device

11977J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	5	..	*743.22	41.30	Repatha [AN]

■ EVOLOCUMAB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)**10385**

Non-familial hypercholesterolaemia

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be in conjunction with dietary therapy and exercise.

Authority required (STREAMLINED)**10377**

Familial heterozygous hypercholesterolaemia

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be in conjunction with dietary therapy and exercise.

evolocumab 420 mg/3.5 mL injection, 3.5 mL cartridge

11986W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	539.12	41.30	Repatha [AN]

evolocumab 140 mg/mL injection, 1 mL pen device

11985T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*497.82	41.30	Repatha [AN]

■ EVOLOCUMAB

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Familial homozygous hypercholesterolaemia

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be in conjunction with dietary therapy and exercise, **AND**
- The condition must have been confirmed by genetic testing; OR
- The condition must have been confirmed by a Dutch Lipid Clinic Network Score of at least 7, **AND**
- Patient must have an LDL cholesterol level in excess of 2.6 millimoles per litre, **AND**
- Patient must have been treated with the maximum recommended dose of atorvastatin (80 mg daily) or rosuvastatin (40 mg daily) according to the TGA-approved Product Information or the maximum tolerated dose of atorvastatin or rosuvastatin for at least 12 consecutive weeks in conjunction with dietary therapy and exercise; OR
- Patient must have developed clinically important product-related adverse events necessitating withdrawal of statin treatment to trials of each of atorvastatin and rosuvastatin; OR
- Patient must be contraindicated to treatment with a HMG CoA reductase inhibitor (statin) as defined in the TGA-approved Product Information.

Treatment criteria:

- Must be treated by a specialist physician.

The qualifying LDL cholesterol level following at least 12 consecutive weeks of treatment with a statin (unless treatment with a statin is contraindicated or following completion of statin trials as described in these prescriber instructions in the event of clinically important adverse events) must be stated at the time of application, documented in the patient's medical records and must be no more than 8 weeks old.

A clinically important product-related adverse event is defined as follows:

- (i) Severe myalgia (muscle symptoms without creatine kinase elevation) which is proven to be temporally associated with statin treatment; or

(ii) Myositis (clinically important creatine kinase elevation, with or without muscle symptoms) demonstrated by results twice the upper limit of normal on a single reading or a rising pattern on consecutive measurements and which is unexplained by other causes; or

(iii) Unexplained, persistent elevations of serum transaminases (greater than 3 times the upper limit of normal) during treatment with a statin.

The following must be stated at the time of application and documented in the patient's medical records:

(i) the qualifying Dutch Lipid Clinic Network Score; or

(ii) the result of genetic testing confirming a diagnosis of familial homozygous hypercholesterolaemia

One of the following must be stated at the time of application and documented in the patient's medical records regarding prior statin treatment:

(i) the patient was treated with atorvastatin 80 mg or rosuvastatin 40 mg or the maximum tolerated dose of either for 12 consecutive weeks; or

(ii) the dose, duration of treatment and details of adverse events experienced with the trial of atorvastatin or rosuvastatin; or

(iii) the patient is contraindicated to treatment with a statin as defined in the TGA-approved Product Information

Authority required

Familial homozygous hypercholesterolaemia

Treatment Phase: Grandfather treatment

Clinical criteria:

- Patient must have previously received non-PBS-subsidised treatment with this drug for this condition prior to 1 May 2020, **AND**

- The treatment must be in conjunction with dietary therapy and exercise, **AND**

- The condition must have been confirmed by genetic testing prior to starting non-PBS-subsidised treatment with this drug for this condition; OR

- The condition must have been confirmed by a Dutch Lipid Clinic Network Score of at least 7 prior to starting non-PBS-subsidised treatment with this drug for this condition, **AND**

- Patient must have had an LDL cholesterol level in excess of 2.6 millimoles per litre prior to starting non-PBS-subsidised treatment with this drug for this condition, **AND**

- Patient must have been treated with the maximum recommended dose of atorvastatin (80 mg daily) or rosuvastatin (40 mg daily) according to the TGA-approved Product Information or the maximum tolerated dose of atorvastatin or rosuvastatin for at least 12 consecutive weeks in conjunction with dietary therapy and exercise prior to initiating non-PBS-subsidised treatment with this drug for this condition; OR

- Patient must have developed a clinically important product-related adverse event necessitating withdrawal of statin treatment to trials of each of atorvastatin and rosuvastatin prior to initiating non-PBS-subsidised treatment with this drug for this condition; OR

- Patient must be contraindicated to treatment with a HMG CoA reductase inhibitor (statin) as defined in the TGA-approved Product Information.

Treatment criteria:

- Must be treated by a specialist physician.

The qualifying LDL cholesterol level must have been measured following at least 12 consecutive weeks of treatment with a statin (unless treatment with a statin is contraindicated or following completion of statin trials as described in these prescriber instructions in the event of clinically important adverse events), must be stated at the time of application, documented in the patient's medical records and must have been no more than 8 weeks old at the time non-PBS-subsidised treatment with this drug for this condition was initiated.

A clinically important product-related adverse event is defined as follows:

(i) Severe myalgia (muscle symptoms without creatine kinase elevation) which is proven to be temporally associated with statin treatment; or

(ii) Myositis (clinically important creatine kinase elevation, with or without muscle symptoms) demonstrated by results twice the upper limit of normal on a single reading or a rising pattern on consecutive measurements and which is unexplained by other causes; or

(iii) Unexplained, persistent elevations of serum transaminases (greater than 3 times the upper limit of normal) during treatment with a statin.

The following must be stated at the time of application and documented in the patient's medical records:

(i) the qualifying Dutch Lipid Clinic Network Score; or

(ii) the result of genetic testing confirming a diagnosis of familial homozygous hypercholesterolaemia

One of the following must be stated at the time of application and documented in the patient's medical records regarding prior statin treatment:

(i) the patient was treated with atorvastatin 80 mg or rosuvastatin 40 mg or the maximum tolerated dose of either for 12 consecutive weeks; or

(ii) the dose, duration of treatment and details of adverse events experienced with the trial of atorvastatin or rosuvastatin; or

(iii) the patient is contraindicated to treatment with a statin as defined in the TGA-approved Product Information

A patient may qualify for PBS-subsidised treatment under this restriction once only.

For continuing PBS-subsidised treatment, a Grandfathered patient must qualify under the Continuing treatment criteria.

evolocumab 420 mg/3.5 mL injection, 3.5 mL cartridge

11193D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	539.12	41.30	Repatha [AN]

evolocumab 140 mg/mL injection, 1 mL pen device

10958R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	5	..	*743.22	41.30	Repatha [AN]

■ EVOLOCUMAB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Familial heterozygous hypercholesterolaemia

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be in conjunction with dietary therapy and exercise, **AND**
- The condition must have been confirmed by genetic testing; OR
- The condition must have been confirmed by a Dutch Lipid Clinic Network Score of at least 6, **AND**
- Patient must have an LDL cholesterol level in excess of 2.6 millimoles per litre in the presence of symptomatic atherosclerotic cardiovascular disease; OR
- Patient must have an LDL cholesterol level in excess of 5 millimoles per litre, **AND**
- Patient must have been treated with the maximum recommended dose of atorvastatin (80 mg daily) or rosuvastatin (40 mg daily) according to the TGA-approved Product Information or the maximum tolerated dose of atorvastatin or rosuvastatin for at least 12 consecutive weeks in conjunction with dietary therapy and exercise; OR
- Patient must have developed clinically important product-related adverse events necessitating withdrawal of statin treatment to trials of each of atorvastatin and rosuvastatin; OR
- Patient must be contraindicated to treatment with a HMG CoA reductase inhibitor (statin) as defined in the TGA-approved Product Information, **AND**
- Patient must have been treated with ezetimibe for at least 12 consecutive weeks in conjunction with a statin (if tolerated), dietary therapy and exercise.

Treatment criteria:

- Must be treated by a specialist physician.

Symptomatic atherosclerotic cardiovascular disease is defined as:

- (i) the presence of symptomatic coronary artery disease (prior myocardial infarction, prior revascularisation procedure, angina associated with demonstrated significant coronary artery disease (50% or greater stenosis in 1 or more coronary arteries on imaging), or positive functional testing (e.g. myocardial perfusion scanning or stress echocardiography); or
- (ii) the presence of symptomatic cerebrovascular disease (prior ischaemic stroke, prior revascularisation procedure, or transient ischaemic attack associated with 50% or greater stenosis in 1 or more cerebral arteries on imaging); or
- (iii) the presence of symptomatic peripheral arterial disease (prior acute ischaemic event due to atherosclerosis, prior revascularisation procedure, or symptoms of ischaemia with evidence of significant peripheral artery disease (50% or greater stenosis in 1 or more peripheral arteries on imaging)).

The qualifying LDL cholesterol level following at least 12 consecutive weeks of combined treatment with a statin, ezetimibe, dietary therapy and exercise (unless treatment with a statin is contraindicated, or following completion of statin trials as described in these prescriber instructions in the event of clinically important adverse events) must be stated at the time of application, documented in the patient's medical records and must be no more than 8 weeks old.

A clinically important product-related adverse event is defined as follows:

- (i) Severe myalgia (muscle symptoms without creatine kinase elevation) which is proven to be temporally associated with statin treatment; or
- (ii) Myositis (clinically important creatine kinase elevation, with or without muscle symptoms) demonstrated by results twice the upper limit of normal on a single reading or a rising pattern on consecutive measurements and which is unexplained by other causes; or
- (iii) Unexplained, persistent elevations of serum transaminases (greater than 3 times the upper limit of normal) during treatment with a statin.

If treatment with atorvastatin or rosuvastatin results in development of a clinically important product-related adverse event resulting in treatment withdrawal, the patient must be treated with the alternative statin (atorvastatin or rosuvastatin) unless there is a contraindication (e.g. prior rhabdomyolysis) to the alternative statin. This retreat should occur after a washout period of at least 4 weeks, or if the creatine kinase (CK) level is elevated, retreat should not occur until CK has returned to normal.

In the event of a trial of the alternative statin, it is recommended that the patient is started with the minimum dose of statin in conjunction with ezetimibe. The dose of the alternative statin should be increased not more often than every 4 weeks until the recommended or maximum tolerated dose has been reached or target LDL-c has been achieved.

The following must be stated at the time of application and documented in the patient's medical records:

- (i) the qualifying Dutch Lipid Clinic Network Score; or
- (ii) the result of genetic testing confirming a diagnosis of familial heterozygous hypercholesterolaemia

One of the following must be stated at the time of application and documented in the patient's medical records regarding prior statin treatment:

- (i) the patient was treated with atorvastatin 80 mg or rosuvastatin 40 mg or the maximum tolerated dose of either for 12 consecutive weeks; or

- (ii) the doses, duration of treatment and details of adverse events experienced with trials with each of atorvastatin and rosuvastatin; or
- (iii) the patient is contraindicated to treatment with a statin as defined in the TGA-approved Product Information.

Authority required

Non-familial hypercholesterolaemia

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be in conjunction with dietary therapy and exercise, **AND**
- Patient must have symptomatic atherosclerotic cardiovascular disease, **AND**
- Patient must have an LDL cholesterol level in excess of 2.6 millimoles per litre, **AND**
- Patient must have atherosclerotic disease in two or more vascular territories (coronary, cerebrovascular or peripheral vascular territories); OR
- Patient must have severe multi-vessel coronary heart disease defined as at least 50% stenosis in at least two large vessels; OR
- Patient must have had at least two major cardiovascular events (i.e. myocardial infarction, unstable angina, stroke or unplanned revascularisation) in the previous 5 years; OR
- Patient must have diabetes mellitus with microalbuminuria; OR
- Patient must have diabetes mellitus and be aged 60 years or more; OR
- Patient must be an Aboriginal or Torres Strait Islander with diabetes mellitus; OR
- Patient must have a Thrombolysis in Myocardial Infarction (TIMI) risk score for secondary prevention of 4 or higher, **AND**
- Patient must have been treated with the maximum recommended dose of atorvastatin (80 mg daily) or rosuvastatin (40 mg daily) according to the TGA-approved Product Information or the maximum tolerated dose of atorvastatin or rosuvastatin for at least 12 consecutive weeks in conjunction with dietary therapy and exercise; OR
- Patient must have developed clinically important product-related adverse events necessitating withdrawal of statin treatment to trials of each of atorvastatin and rosuvastatin; OR
- Patient must be contraindicated to treatment with a HMG CoA reductase inhibitor (statin) as defined in the TGA-approved Product Information, **AND**
- Patient must have been treated with ezetimibe for at least 12 consecutive weeks in conjunction with a statin (if tolerated), dietary therapy and exercise.

Treatment criteria:

- Must be treated by a specialist physician.

Symptomatic atherosclerotic cardiovascular disease is defined as:

- (i) the presence of symptomatic coronary artery disease (prior myocardial infarction, prior revascularisation procedure, angina associated with demonstrated significant coronary artery disease (50% or greater stenosis in 1 or more coronary arteries on imaging), or positive functional testing (e.g. myocardial perfusion scanning or stress echocardiography); or
- (ii) the presence of symptomatic cerebrovascular disease (prior ischaemic stroke, prior revascularisation procedure, or transient ischaemic attack associated with 50% or greater stenosis in 1 or more cerebral arteries on imaging); or
- (iii) the presence of symptomatic peripheral arterial disease (prior acute ischaemic event due to atherosclerosis, prior revascularisation procedure, or symptoms of ischaemia with evidence of significant peripheral artery disease (50% or greater stenosis in 1 or more peripheral arteries on imaging)).

The qualifying LDL cholesterol level following at least 12 consecutive weeks of combined treatment with a statin, ezetimibe, dietary therapy and exercise (unless treatment with a statin is contraindicated, or following completion of statin trials as described in these prescriber instructions in the event of clinically important adverse events) must be stated at the time of application, documented in the patient's medical records and must be no more than 8 weeks old.

A clinically important product-related adverse event is defined as follows:

- (i) Severe myalgia (muscle symptoms without creatine kinase elevation) which is proven to be temporally associated with statin treatment; or
- (ii) Myositis (clinically important creatine kinase elevation, with or without muscle symptoms) demonstrated by results twice the upper limit of normal on a single reading or a rising pattern on consecutive measurements and which is unexplained by other causes; or
- (iii) Unexplained, persistent elevations of serum transaminases (greater than 3 times the upper limit of normal) during treatment with a statin.

If treatment with atorvastatin or rosuvastatin results in development of a clinically important product-related adverse event resulting in treatment withdrawal, the patient must be treated with the alternative statin (atorvastatin or rosuvastatin) unless there is a contraindication (e.g. prior rhabdomyolysis) to the alternative statin. This retreat should occur after a washout period of at least 4 weeks, or if the creatine kinase (CK) level is elevated, retreat should not occur until CK has returned to normal.

In the event of a trial of the alternative statin, it is recommended that the patient is started with the minimum dose of statin in conjunction with ezetimibe. The dose of the alternative statin should be increased not more often than every 4 weeks until the recommended or maximum tolerated dose has been reached or target LDL-c has been achieved.

One of the following must be stated at the time of application and documented in the patient's medical records regarding prior statin treatment:

- (i) the patient was treated with atorvastatin 80 mg or rosuvastatin 40 mg or the maximum tolerated dose of either for 12 consecutive weeks; or
- (ii) the doses, duration of treatment and details of adverse events experienced with trials with each of atorvastatin and rosuvastatin; or
- (iii) the patient is contraindicated to treatment with a statin as defined in the TGA-approved Product Information.

One or more of the following must be stated at the time of application and documented in the patient's medical records regarding the presence of cardiovascular disease or high risk of experiencing a cardiovascular event:

- (i) atherosclerotic disease in two or more vascular territories (coronary, cerebrovascular or peripheral vascular territories); or
- (ii) severe multi-vessel coronary heart disease defined as at least 50% stenosis in at least two large vessels; or
- (iii) history of at least two major cardiovascular events (i.e. myocardial infarction, unstable angina, stroke or unplanned revascularisation) in the previous 5 years; or
- (iv) diabetes mellitus with microalbuminuria; or
- (v) diabetes mellitus and age 60 years or more; or
- (vi) Aboriginal or Torres Strait Islander with diabetes mellitus; or
- (vii) a Thrombolysis in Myocardial Infarction (TIMI) risk score for secondary prevention of 4 or higher

Authority required

Familial heterozygous hypercholesterolaemia

Treatment Phase: Grandfather treatment

Clinical criteria:

- Patient must have previously received non-PBS-subsidised treatment with this drug for this condition prior to 1 May 2020, **AND**
- The treatment must be in conjunction with dietary therapy and exercise, **AND**
- The condition must have been confirmed by genetic testing prior to starting non-PBS-subsidised treatment with this drug for this condition; OR
- The condition must have been confirmed by a Dutch Lipid Clinic Network Score of at least 6 prior to starting non-PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have had an LDL cholesterol level in excess of 2.6 millimoles per litre in the presence of symptomatic atherosclerotic cardiovascular disease at the time non-PBS-subsidised treatment with this drug for this condition was initiated; OR
- Patient must have had an LDL cholesterol level in excess of 5 millimoles per litre at the time non-PBS-subsidised treatment with this drug for this condition was initiated, **AND**
- Patient must have been treated with the maximum recommended dose of atorvastatin (80 mg daily) or rosuvastatin (40 mg daily) according to the TGA-approved Product Information or the maximum tolerated dose of atorvastatin or rosuvastatin for at least 12 consecutive weeks in conjunction with dietary therapy and exercise prior to initiating non-PBS-subsidised treatment with this drug for this condition; OR
- Patient must have developed a clinically important product-related adverse event necessitating withdrawal of statin treatment to trials of each of atorvastatin and rosuvastatin prior to initiating non-PBS-subsidised treatment with this drug for this condition; OR
- Patient must be contraindicated to treatment with a HMG CoA reductase inhibitor (statin) as defined in the TGA-approved Product Information, **AND**
- Patient must have been treated with ezetimibe for at least 12 consecutive weeks in conjunction with a statin (if tolerated), dietary therapy and exercise prior to initiating non-PBS-subsidised treatment with this drug for this condition.

Treatment criteria:

- Must be treated by a specialist physician.

Symptomatic atherosclerotic cardiovascular disease is defined as:

- (i) the presence of symptomatic coronary artery disease (prior myocardial infarction, prior revascularisation procedure, angina associated with demonstrated significant coronary artery disease (50% or greater stenosis in 1 or more coronary arteries on imaging), or positive functional testing (e.g. myocardial perfusion scanning or stress echocardiography); or
- (ii) the presence of symptomatic cerebrovascular disease (prior ischaemic stroke, prior revascularisation procedure, or transient ischaemic attack associated with 50% or greater stenosis in 1 or more cerebral arteries on imaging); or
- (iii) the presence of symptomatic peripheral arterial disease (prior acute ischaemic event due to atherosclerosis, prior revascularisation procedure, or symptoms of ischaemia with evidence of significant peripheral artery disease (50% or greater stenosis in 1 or more peripheral arteries on imaging)).

The qualifying LDL cholesterol level must have been measured following at least 12 consecutive weeks of combined treatment with a statin, ezetimibe, dietary therapy and exercise (unless treatment with a statin is contraindicated, or following completion of statin trials as described in these prescriber instructions in the event of clinically important adverse events), must be stated at the time of application, documented in the patient's medical records and must have been no more than 8 weeks old at the time non-PBS-subsidised treatment with this drug for this condition was initiated.

A clinically important product-related adverse event is defined as follows:

- (i) Severe myalgia (muscle symptoms without creatine kinase elevation) which is proven to be temporally associated with statin treatment; or
- (ii) Myositis (clinically important creatine kinase elevation, with or without muscle symptoms) demonstrated by results twice the upper limit of normal on a single reading or a rising pattern on consecutive measurements and which is unexplained by other causes; or
- (iii) Unexplained, persistent elevations of serum transaminases (greater than 3 times the upper limit of normal) during treatment with a statin.

If treatment with atorvastatin or rosuvastatin resulted in development of a clinically important product-related adverse event resulting in treatment withdrawal, the patient must have been treated with the alternative statin (atorvastatin or rosuvastatin) unless there was a contraindication (e.g. prior rhabdomyolysis) to the alternative statin. This retreatment should have occurred after a washout period of at least 4 weeks, or if the creatine kinase (CK) level was elevated, the retreatment should not have occurred until CK had returned to normal.

In the event of a trial of the alternative statin, the dose of the alternative statin should have been increased not more often than every 4 weeks until the maximum tolerated dose was reached or target LDL-c had been achieved.

The following must be stated at the time of application and documented in the patient's medical records:

- (i) the qualifying Dutch Lipid Clinic Network Score; or
- (ii) the result of genetic testing confirming a diagnosis of familial heterozygous hypercholesterolaemia

One of the following must be stated at the time of application and documented in the patient's medical records regarding prior statin treatment:

- (i) the patient was treated with atorvastatin 80 mg or rosuvastatin 40 mg or the maximum tolerated dose of either for 12 consecutive weeks; or
- (ii) the doses, duration of treatment and details of adverse events experienced with trials with each of atorvastatin and rosuvastatin; or
- (iii) the patient is contraindicated to treatment with a statin as defined in the TGA-approved Product Information.

A patient may qualify for PBS-subsidised treatment under this restriction once only.

For continuing PBS-subsidised treatment, a Grandfathered patient must qualify under the Continuing treatment criteria.

Authority required

Non-familial hypercholesterolaemia

Treatment Phase: Grandfather treatment

Clinical criteria:

- Patient must have previously received non-PBS-subsidised treatment with this drug for this condition prior to 1 May 2020, **AND**

- The treatment must be in conjunction with dietary therapy and exercise, **AND**
- Patient must have had symptomatic atherosclerotic cardiovascular disease prior to starting non-PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have had an LDL cholesterol level in excess of 2.6 millimoles per litre prior to starting non-PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have had atherosclerotic disease in two or more vascular territories (coronary, cerebrovascular or peripheral vascular territories) prior to starting non-PBS-subsidised treatment with this drug for this condition; OR
- Patient must have had severe multi-vessel coronary heart disease defined as at least 50% stenosis in at least two large vessels prior to starting non-PBS-subsidised treatment with this drug for this condition; OR
- Patient must have had at least two major cardiovascular events (i.e. myocardial infarction, unstable angina, stroke or unplanned revascularisation) in the previous 5 years prior to starting non-PBS-subsidised treatment with this drug for this condition; OR
- Patient must have had diabetes mellitus with microalbuminuria prior to starting non-PBS-subsidised treatment with this drug for this condition; OR
- Patient must have had diabetes mellitus and be aged 60 years or more prior to starting non-PBS-subsidised treatment with this drug for this condition; OR
- Patient must be an Aboriginal or Torres Strait Islander with diabetes mellitus that was present prior to starting non-PBS-subsidised treatment with this drug for this condition; OR
- Patient must have had a Thrombolysis in Myocardial Infarction (TIMI) Risk Score for Secondary Prevention of 4 or higher prior to starting non-PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have been treated with the maximum recommended dose of atorvastatin (80 mg daily) or rosuvastatin (40 mg daily) according to the TGA-approved Product Information or the maximum tolerated dose of atorvastatin or rosuvastatin for at least 12 consecutive weeks in conjunction with dietary therapy and exercise prior to initiating non-PBS-subsidised treatment with this drug for this condition; OR
- Patient must have developed a clinically important product-related adverse event necessitating withdrawal of statin treatment to trials of each of atorvastatin and rosuvastatin prior to initiating non-PBS-subsidised treatment with this drug for this condition; OR
- Patient must be contraindicated to treatment with a HMG CoA reductase inhibitor (statin) as defined in the TGA-approved Product Information, **AND**
- Patient must have been treated with ezetimibe for at least 12 consecutive weeks in conjunction with a statin (if tolerated), dietary therapy and exercise prior to initiating non-PBS-subsidised treatment with this drug for this condition.

Treatment criteria:

- Must be treated by a specialist physician.
- Symptomatic atherosclerotic cardiovascular disease is defined as:

- (i) the presence of symptomatic coronary artery disease (prior myocardial infarction, prior revascularisation procedure, angina associated with demonstrated significant coronary artery disease (50% or greater stenosis in 1 or more coronary arteries on imaging), or positive functional testing (e.g. myocardial perfusion scanning or stress echocardiography); or
- (ii) the presence of symptomatic cerebrovascular disease (prior ischaemic stroke, prior revascularisation procedure, or transient ischaemic attack associated with 50% or greater stenosis in 1 or more cerebral arteries on imaging); or
- (iii) the presence of symptomatic peripheral arterial disease (prior acute ischaemic event due to atherosclerosis, prior revascularisation procedure, or symptoms of ischaemia with evidence of significant peripheral artery disease (50% or greater stenosis in 1 or more peripheral arteries on imaging)).

The qualifying LDL cholesterol level must have been measured following at least 12 consecutive weeks of combined treatment with a statin, ezetimibe, dietary therapy and exercise (unless treatment with a statin is contraindicated, or following completion of statin trials as described in these prescriber instructions in the event of clinically important adverse events), must be stated at the time of application, documented in the patient's medical records and must have been no more than 8 weeks old at the time non-PBS-subsidised treatment with this drug for this condition was initiated.

A clinically important product-related adverse event is defined as follows:

- (i) Severe myalgia (muscle symptoms without creatine kinase elevation) which is proven to be temporally associated with statin treatment; or
- (ii) Myositis (clinically important creatine kinase elevation, with or without muscle symptoms) demonstrated by results twice the upper limit of normal on a single reading or a rising pattern on consecutive measurements and which is unexplained by other causes; or
- (iii) Unexplained, persistent elevations of serum transaminases (greater than 3 times the upper limit of normal) during treatment with a statin.

If treatment with atorvastatin or rosuvastatin resulted in development of a clinically important product-related adverse event resulting in treatment withdrawal, the patient must have been treated with the alternative statin (atorvastatin or rosuvastatin) unless there was a contraindication (e.g. prior rhabdomyolysis) to the alternative statin. This retreatment should have occurred after a washout period of at least 4 weeks, or if the creatine kinase (CK) level was elevated, the retreatment should not have occurred until CK had returned to normal.

In the event of a trial of the alternative statin, the dose of the alternative statin should have been increased not more often than every 4 weeks until the maximum tolerated dose was reached or target LDL-c had been achieved.

One of the following must be stated at the time of application and documented in the patient's medical records regarding prior statin treatment:

- (i) the patient was treated with atorvastatin 80 mg or rosuvastatin 40 mg or the maximum tolerated dose of either for 12 consecutive weeks; or
- (ii) the doses, duration of treatment and details of adverse events experienced with trials with each of atorvastatin and rosuvastatin; or
- (iii) the patient is contraindicated to treatment with a statin as defined in the TGA-approved Product Information.

One or more of the following must be stated at the time of application and documented in the patient's medical records regarding the presence of cardiovascular disease or high risk of experiencing a cardiovascular event:

- (i) atherosclerotic disease in two or more vascular territories (coronary, cerebrovascular or peripheral vascular territories); or
- (ii) severe multi-vessel coronary heart disease defined as at least 50% stenosis in at least two large vessels; or
- (iii) history of at least two major cardiovascular events (i.e. myocardial infarction, unstable angina, stroke or unplanned revascularisation) in the previous 5 years; or
- (iv) diabetes mellitus with microalbuminuria; or
- (v) diabetes mellitus and age 60 years or more; or
- (vi) Aboriginal or Torres Strait Islander with diabetes mellitus; or
- (vii) a Thrombolysis in Myocardial Infarction (TIMI) risk score for secondary prevention of 4 or higher

A patient may qualify for PBS-subsidised treatment under this restriction once only.

For continuing PBS-subsidised treatment, a Grandfathered patient must qualify under the Continuing treatment criteria.

evolocumab 420 mg/3.5 mL injection, 3.5 mL cartridge

11485L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	539.12	41.30	Repatha [AN]

evolocumab 140 mg/mL injection, 1 mL pen device

11484K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*497.82	41.30	Repatha [AN]

■ EZETIMIBE

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)

7990

Hypercholesterolaemia

Clinical criteria:

- Patient must have homozygous sitosterolaemia.

ezetimibe 10 mg tablet, 30

11408K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	23.66	24.95	^a EZEMICHOL [RW]	^a Pharmacor Ezetimibe 10 [CR]
						^a Zient 10mg [AF]	
				^b 2.56	26.22	^a Ezetrol [AL]	

■ EZETIMIBE

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)

7996

Hypercholesterolaemia

Clinical criteria:

- The treatment must be in conjunction with dietary therapy and exercise, **AND**
- The treatment must be co-administered with an HMG CoA reductase inhibitor (statin), **AND**
- Patient must have cholesterol concentrations that are inadequately controlled with an HMG CoA reductase inhibitor (statin), **AND**
- Patient must have coronary heart disease; OR
- Patient must have cerebrovascular disease; OR
- Patient must have peripheral vascular disease; OR
- Patient must have diabetes mellitus with microalbuminuria; OR

- Patient must be an Aboriginal or Torres Strait Islander with diabetes mellitus; OR
- Patient must have diabetes mellitus and be aged 60 years or more; OR
- Patient must have a family history of coronary heart disease in two or more first degree relatives before the age of 55 years; OR
- Patient must have a family history of coronary heart disease in one or more first degree relatives before the age of 45 years; OR
- Patient must have heterozygous familial hypercholesterolaemia; OR
- Patient must have homozygous familial hypercholesterolaemia; OR
- Patient must have a level of absolute risk of a cardiovascular event greater than 15% over 5 years as calculated using the Australian Absolute Cardiovascular Disease Risk Calculator (National Vascular Disease Prevention Alliance), as in force on 1 April 2018.

Inadequate control with a statin is defined as a LDL cholesterol concentration in excess of current target lipid levels for primary and secondary prevention after at least 3 months of treatment at a maximum tolerated dose of a statin.

The dose and duration of statin treatment and the cholesterol concentration which shows inadequate control must be documented in the patient's medical records when ezetimibe is initiated.

The cholesterol concentration which shows inadequate control must be no more than 2 months old when ezetimibe is initiated.

Microalbuminuria is defined as urinary albumin excretion rate of greater than 20mcg/min or urinary albumin to creatinine ratio of greater than 2.5 for males, or greater than 3.5 for females.

Note The Australian Absolute Cardiovascular Disease Risk Calculator is available at www.cvdcheck.org.au

Authority required (STREAMLINED)

7966

Hypercholesterolaemia

Clinical criteria:

- Patient must have developed a clinically important product-related adverse event during treatment with an HMG CoA reductase inhibitor (statin) necessitating a reduction in the statin dose; OR
- Patient must have developed a clinically important product-related adverse event during treatment with an HMG CoA reductase inhibitor (statin) necessitating a withdrawal of the statin treatment; OR
- Patient must be one in whom treatment with an HMG CoA reductase inhibitor (statin) is contraindicated, **AND**
- Patient must have coronary heart disease; OR
- Patient must have cerebrovascular disease; OR
- Patient must have peripheral vascular disease; OR
- Patient must have diabetes mellitus with microalbuminuria; OR
- Patient must be an Aboriginal or Torres Strait Islander with diabetes mellitus; OR
- Patient must have diabetes mellitus and be aged 60 years or more; OR
- Patient must have a family history of coronary heart disease in two or more first degree relatives before the age of 55 years; OR
- Patient must have a family history of coronary heart disease in one or more first degree relatives before the age of 45 years; OR
- Patient must have heterozygous familial hypercholesterolaemia; OR
- Patient must have homozygous familial hypercholesterolaemia; OR
- Patient must have a level of absolute risk of a cardiovascular event greater than 15% over 5 years as calculated using the Australian Absolute Cardiovascular Disease Risk Calculator (National Vascular Disease Prevention Alliance), as in force on 1 April 2018.

A clinically important product-related adverse event is defined as follows:

- Severe myalgia (muscle symptoms without creatine kinase elevation) which is proven to be temporally associated with statin treatment; or
- Myositis (clinically important creatine kinase elevation, with or without muscle symptoms) demonstrated by results twice the upper limit of normal on a single reading or a rising pattern on consecutive measurements and which is unexplained by other causes; or
- Unexplained, persistent elevations of serum transaminases (greater than 3 times the upper limit of normal) during treatment with a statin.

Microalbuminuria is defined as urinary albumin excretion rate of greater than 20mcg/min or urinary albumin to creatinine ratio of greater than 2.5 for males, or greater than 3.5 for females.

The type and severity of the adverse event or contraindication must be documented in the patient's medical records.

ezetimibe 10 mg tablet, 30

8757X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	23.66	24.95	^a APO-Ezetimibe [TX]	^a Blooms The Chemist Ezetimibe [IB]
						^a EZEMICHOL [RW]	^a Ezetimibe GH [GQ]
						^a Ezetimibe Sandoz [SZ]	^a Pharmacor Ezetimibe 10 [CR]
						^a Zient 10mg [AF]	
			^b 2.56	26.22	24.95	^a Ezetrol [AL]	

LIPID MODIFYING AGENTS, COMBINATIONS

Combinations of various lipid modifying agents

■ EZETIMIBE (&) ROSUVASTATIN

Note The Australian Absolute Cardiovascular Disease Risk Calculator is available at www.cvdcheck.org.au

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**7957**

Hypercholesterolaemia

Clinical criteria:

- The treatment must be in conjunction with dietary therapy and exercise, **AND**
- Patient must have cholesterol concentrations that are inadequately controlled with an HMG CoA reductase inhibitor (statin), **AND**
- Patient must have coronary heart disease; OR
- Patient must have cerebrovascular disease; OR
- Patient must have peripheral vascular disease; OR
- Patient must have diabetes mellitus with microalbuminuria; OR
- Patient must be an Aboriginal or Torres Strait Islander with diabetes mellitus; OR
- Patient must have diabetes mellitus and be aged 60 years or more; OR
- Patient must have a family history of coronary heart disease in two or more first degree relatives before the age of 55 years; OR
- Patient must have a family history of coronary heart disease in one or more first degree relatives before the age of 45 years; OR
- Patient must have heterozygous familial hypercholesterolaemia; OR
- Patient must have homozygous familial hypercholesterolaemia; OR
- Patient must have a level of absolute risk of a cardiovascular event greater than 15% over 5 years as calculated using the Australian Absolute Cardiovascular Disease Risk Calculator (National Vascular Disease Prevention Alliance), as in force on 1 April 2018.

Inadequate control with a statin is defined as a LDL cholesterol concentration in excess of current target lipid levels for primary and secondary prevention after at least 3 months of treatment at a maximum tolerated dose of a statin.

The dose and duration of statin treatment and the cholesterol concentration which shows inadequate control must be documented in the patient's medical records when ezetimibe is initiated.

The cholesterol concentration which shows inadequate control must be no more than 2 months old when ezetimibe is initiated.

Microalbuminuria is defined as urinary albumin excretion rate of greater than 20mcg/min or urinary albumin to creatinine ratio of greater than 2.5 for males, or greater than 3.5 for females.

ezetimibe 10 mg tablet [30] (&) rosuvastatin 20 mg tablet [30], 60

10201X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	25.87	27.16	^a Ezalo Composite Pack 10mg+20mg [AF]
			^B 2.42	28.29	27.16	^a Rosuzet Composite Pack [AL]

ezetimibe 10 mg tablet [30] (&) rosuvastatin 10 mg tablet [30], 60

10208G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	26.26	27.55	^a Ezalo Composite Pack 10mg+10mg [AF]
			^B 2.64	28.90	27.55	^a Rosuzet Composite Pack [AL]

ezetimibe 10 mg tablet [30] (&) rosuvastatin 40 mg tablet [30], 60

10207F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	26.84	28.13	^a Ezalo Composite Pack 10mg+40mg [AF]
			^B 2.38	29.22	28.13	^a Rosuzet Composite Pack [AL]

■ EZETIMIBE (&) ROSUVASTATIN**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note The Australian Absolute Cardiovascular Disease Risk Calculator is available at www.cvdcheck.org.au

Authority required (STREAMLINED)**7958**

Hypercholesterolaemia

Clinical criteria:

- The treatment must be in conjunction with dietary therapy and exercise, **AND**
- Patient must have cholesterol concentrations that are inadequately controlled with an HMG CoA reductase inhibitor (statin), **AND**
- Patient must have developed a clinically important product-related adverse event during treatment with an HMG CoA reductase inhibitor (statin) necessitating a reduction in the statin dose, **AND**
- Patient must have coronary heart disease; OR
- Patient must have cerebrovascular disease; OR

- Patient must have peripheral vascular disease; OR
- Patient must have diabetes mellitus with microalbuminuria; OR
- Patient must be an Aboriginal or Torres Strait Islander with diabetes mellitus; OR
- Patient must have diabetes mellitus and be aged 60 years or more; OR
- Patient must have a family history of coronary heart disease in two or more first degree relatives before the age of 55 years; OR
- Patient must have a family history of coronary heart disease in one or more first degree relatives before the age of 45 years; OR
- Patient must have heterozygous familial hypercholesterolaemia; OR
- Patient must have homozygous familial hypercholesterolaemia; OR
- Patient must have a level of absolute risk of a cardiovascular event greater than 15% over 5 years as calculated using the Australian Absolute Cardiovascular Disease Risk Calculator (National Vascular Disease Prevention Alliance), as in force on 1 April 2018.

A clinically important product-related adverse event is defined as follows:

- (i) Severe myalgia (muscle symptoms without creatine kinase elevation) which is proven to be temporally associated with statin treatment; or
- (ii) Myositis (clinically important creatine kinase elevation, with or without muscle symptoms) demonstrated by results twice the upper limit of normal on a single reading or a rising pattern on consecutive measurements and which is unexplained by other causes; or
- (iii) Unexplained, persistent elevations of serum transaminases (greater than 3 times the upper limit of normal) during treatment with a statin.

Microalbuminuria is defined as urinary albumin excretion rate of greater than 20mcg/min or urinary albumin to creatinine ratio of greater than 2.5 for males, or greater than 3.5 for females.

The type and severity of the adverse event or contraindication must be documented in the patient's medical records.

ezetimibe 10 mg tablet [30] (&) rosuvastatin 5 mg tablet [30], 60

10204C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	24.70	25.99	^a Ezalo Composite Pack 10mg+5mg [AF]
			^B 2.48	27.18	25.99	^a Rosuzet Composite Pack [AL]

■ **EZETIMIBE + ATORVASTATIN**

Note The Australian Absolute Cardiovascular Disease Risk Calculator is available at www.cvdcheck.org.au

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)

7957

Hypercholesterolaemia

Clinical criteria:

- The treatment must be in conjunction with dietary therapy and exercise, **AND**
- Patient must have cholesterol concentrations that are inadequately controlled with an HMG CoA reductase inhibitor (statin), **AND**
- Patient must have coronary heart disease; OR
- Patient must have cerebrovascular disease; OR
- Patient must have peripheral vascular disease; OR
- Patient must have diabetes mellitus with microalbuminuria; OR
- Patient must be an Aboriginal or Torres Strait Islander with diabetes mellitus; OR
- Patient must have diabetes mellitus and be aged 60 years or more; OR
- Patient must have a family history of coronary heart disease in two or more first degree relatives before the age of 55 years; OR
- Patient must have a family history of coronary heart disease in one or more first degree relatives before the age of 45 years; OR
- Patient must have heterozygous familial hypercholesterolaemia; OR
- Patient must have homozygous familial hypercholesterolaemia; OR
- Patient must have a level of absolute risk of a cardiovascular event greater than 15% over 5 years as calculated using the Australian Absolute Cardiovascular Disease Risk Calculator (National Vascular Disease Prevention Alliance), as in force on 1 April 2018.

Inadequate control with a statin is defined as a LDL cholesterol concentration in excess of current target lipid levels for primary and secondary prevention after at least 3 months of treatment at a maximum tolerated dose of a statin.

The dose and duration of statin treatment and the cholesterol concentration which shows inadequate control must be documented in the patient's medical records when ezetimibe is initiated.

The cholesterol concentration which shows inadequate control must be no more than 2 months old when ezetimibe is initiated.

Microalbuminuria is defined as urinary albumin excretion rate of greater than 20mcg/min or urinary albumin to creatinine ratio of greater than 2.5 for males, or greater than 3.5 for females.

ezetimibe 10 mg + atorvastatin 80 mg tablet, 30

10376D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	28.20	29.49	^a Atozet [AF]	^a Ezetimibe/Atorvastatin GH 10/80 [GQ]

ezetimibe 10 mg + atorvastatin 40 mg tablet, 30

10377E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	26.84	28.13	^a Atozet [AF]	^a Ezetimibe/Atorvastatin GH 10/40 [GQ]

ezetimibe 10 mg + atorvastatin 20 mg tablet, 30

10393B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	25.92	27.21	^a Atozet [AF]	^a Ezetimibe/Atorvastatin GH 10/20 [GQ]

■ EZETIMIBE + ATORVASTATIN**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note The Australian Absolute Cardiovascular Disease Risk Calculator is available at www.cvdcheck.org.au

Authority required (STREAMLINED)**7958**

Hypercholesterolaemia

Clinical criteria:

- The treatment must be in conjunction with dietary therapy and exercise, **AND**
- Patient must have cholesterol concentrations that are inadequately controlled with an HMG CoA reductase inhibitor (statin), **AND**
- Patient must have developed a clinically important product-related adverse event during treatment with an HMG CoA reductase inhibitor (statin) necessitating a reduction in the statin dose, **AND**
- Patient must have coronary heart disease; OR
- Patient must have cerebrovascular disease; OR
- Patient must have peripheral vascular disease; OR
- Patient must have diabetes mellitus with microalbuminuria; OR
- Patient must be an Aboriginal or Torres Strait Islander with diabetes mellitus; OR
- Patient must have diabetes mellitus and be aged 60 years or more; OR
- Patient must have a family history of coronary heart disease in two or more first degree relatives before the age of 55 years; OR
- Patient must have a family history of coronary heart disease in one or more first degree relatives before the age of 45 years; OR
- Patient must have heterozygous familial hypercholesterolaemia; OR
- Patient must have homozygous familial hypercholesterolaemia; OR
- Patient must have a level of absolute risk of a cardiovascular event greater than 15% over 5 years as calculated using the Australian Absolute Cardiovascular Disease Risk Calculator (National Vascular Disease Prevention Alliance), as in force on 1 April 2018.

A clinically important product-related adverse event is defined as follows:

- Severe myalgia (muscle symptoms without creatine kinase elevation) which is proven to be temporally associated with statin treatment; or
- Myositis (clinically important creatine kinase elevation, with or without muscle symptoms) demonstrated by results twice the upper limit of normal on a single reading or a rising pattern on consecutive measurements and which is unexplained by other causes; or
- Unexplained, persistent elevations of serum transaminases (greater than 3 times the upper limit of normal) during treatment with a statin.

Microalbuminuria is defined as urinary albumin excretion rate of greater than 20mcg/min or urinary albumin to creatinine ratio of greater than 2.5 for males, or greater than 3.5 for females.

The type and severity of the adverse event or contraindication must be documented in the patient's medical records.

ezetimibe 10 mg + atorvastatin 10 mg tablet, 30

10392Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	25.19	26.48	^a Atozet [AF]	^a Ezetimibe/Atorvastatin GH 10/10 [GQ]

■ EZETIMIBE + SIMVASTATIN

Note The Australian Absolute Cardiovascular Disease Risk Calculator is available at www.cvdcheck.org.au

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**7957**

Hypercholesterolaemia

Clinical criteria:

- The treatment must be in conjunction with dietary therapy and exercise, **AND**
- Patient must have cholesterol concentrations that are inadequately controlled with an HMG CoA reductase inhibitor (statin), **AND**
- Patient must have coronary heart disease; OR
- Patient must have cerebrovascular disease; OR
- Patient must have peripheral vascular disease; OR
- Patient must have diabetes mellitus with microalbuminuria; OR
- Patient must be an Aboriginal or Torres Strait Islander with diabetes mellitus; OR
- Patient must have diabetes mellitus and be aged 60 years or more; OR
- Patient must have a family history of coronary heart disease in two or more first degree relatives before the age of 55 years; OR
- Patient must have a family history of coronary heart disease in one or more first degree relatives before the age of 45 years; OR
- Patient must have heterozygous familial hypercholesterolaemia; OR
- Patient must have homozygous familial hypercholesterolaemia; OR
- Patient must have a level of absolute risk of a cardiovascular event greater than 15% over 5 years as calculated using the Australian Absolute Cardiovascular Disease Risk Calculator (National Vascular Disease Prevention Alliance), as in force on 1 April 2018.

Inadequate control with a statin is defined as a LDL cholesterol concentration in excess of current target lipid levels for primary and secondary prevention after at least 3 months of treatment at a maximum tolerated dose of a statin.

The dose and duration of statin treatment and the cholesterol concentration which shows inadequate control must be documented in the patient's medical records when ezetimibe is initiated.

The cholesterol concentration which shows inadequate control must be no more than 2 months old when ezetimibe is initiated.

Microalbuminuria is defined as urinary albumin excretion rate of greater than 20mcg/min or urinary albumin to creatinine ratio of greater than 2.5 for males, or greater than 3.5 for females.

ezetimibe 10 mg + simvastatin 80 mg tablet, 30

8882L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	27.00	28.29	^a APO-Ezetimibe/Simvastatin 10/80 [TX] ^a EZETORIN [RW] ^a Pharmacor Ezetimibe Simvastatin 10/80 [CR] ^a Zeklen 10/80 mg [AF]	^a EZETIMIBE/SIMVASTATIN SANDOZ [SZ] ^a EzSimva GH 10/80 [GQ] ^a Vytorin [AL]

ezetimibe 10 mg + simvastatin 40 mg tablet, 30

8881K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	26.00	27.29	^a APO-Ezetimibe/Simvastatin 10/40 [TX] ^a EZETORIN [RW] ^a Pharmacor Ezetimibe Simvastatin 10/40 [CR] ^a Zeklen 10/40 mg [AF]	^a EZETIMIBE/SIMVASTATIN SANDOZ [SZ] ^a EzSimva GH 10/40 [GQ] ^a Vytorin [AL]

▪ **EZETIMIBE + SIMVASTATIN**

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note The Australian Absolute Cardiovascular Disease Risk Calculator is available at www.cvdcheck.org.au

Authority required (STREAMLINED)

7958

Hypercholesterolaemia

Clinical criteria:

- The treatment must be in conjunction with dietary therapy and exercise, **AND**
- Patient must have cholesterol concentrations that are inadequately controlled with an HMG CoA reductase inhibitor (statin), **AND**
- Patient must have developed a clinically important product-related adverse event during treatment with an HMG CoA reductase inhibitor (statin) necessitating a reduction in the statin dose, **AND**
- Patient must have coronary heart disease; OR
- Patient must have cerebrovascular disease; OR
- Patient must have peripheral vascular disease; OR
- Patient must have diabetes mellitus with microalbuminuria; OR
- Patient must be an Aboriginal or Torres Strait Islander with diabetes mellitus; OR
- Patient must have diabetes mellitus and be aged 60 years or more; OR
- Patient must have a family history of coronary heart disease in two or more first degree relatives before the age of 55 years; OR
- Patient must have a family history of coronary heart disease in one or more first degree relatives before the age of 45 years; OR

- Patient must have heterozygous familial hypercholesterolaemia; OR
- Patient must have homozygous familial hypercholesterolaemia; OR
- Patient must have a level of absolute risk of a cardiovascular event greater than 15% over 5 years as calculated using the Australian Absolute Cardiovascular Disease Risk Calculator (National Vascular Disease Prevention Alliance), as in force on 1 April 2018.

A clinically important product-related adverse event is defined as follows:

- Severe myalgia (muscle symptoms without creatine kinase elevation) which is proven to be temporally associated with statin treatment; or
- Myositis (clinically important creatine kinase elevation, with or without muscle symptoms) demonstrated by results twice the upper limit of normal on a single reading or a rising pattern on consecutive measurements and which is unexplained by other causes; or
- Unexplained, persistent elevations of serum transaminases (greater than 3 times the upper limit of normal) during treatment with a statin.

Microalbuminuria is defined as urinary albumin excretion rate of greater than 20mcg/min or urinary albumin to creatinine ratio of greater than 2.5 for males, or greater than 3.5 for females.

The type and severity of the adverse event or contraindication must be documented in the patient's medical records.

ezetimibe 10 mg + simvastatin 10 mg tablet, 30

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
9483D	1	5	..	24.78	26.07	^a APO-Ezetimibe/Simvastatin 10/10 [TX]	^a EZETIMIBE/SIMVASTATIN SANDOZ [SZ]
NP						^a EZETORIN [RW]	^a EzSimva GH 10/10 [GQ]
						^a Pharmacor Ezetimibe Simvastatin 10/10 [CR]	^a Vytorin [AL]
						^a Zeklen 10/10 mg [AF]	

ezetimibe 10 mg + simvastatin 20 mg tablet, 30

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
9484E	1	5	..	25.29	26.58	^a APO-Ezetimibe/Simvastatin 10/20 [TX]	^a EZETIMIBE/SIMVASTATIN SANDOZ [SZ]
NP						^a EZETORIN [RW]	^a EzSimva GH 10/20 [GQ]
						^a Pharmacor Ezetimibe Simvastatin 10/20 [CR]	^a Vytorin [AL]
						^a Zeklen 10/20 mg [AF]	

Lipid modifying agents in combination with other drugs

■ AMLODIPINE + ATORVASTATIN

amlodipine 5 mg + atorvastatin 10 mg tablet, 30

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
9049G	1	5	..	14.68	15.97	Cadivast 5/10 [AF]
NP						

amlodipine 10 mg + atorvastatin 40 mg tablet, 30

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
9055N	1	5	..	16.85	18.14	^a Cadivast 10/40 [AF]
NP			^B 4.92	21.77	18.14	^a Caduet 10/40 [AS]

amlodipine 10 mg + atorvastatin 10 mg tablet, 30

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
9053L	1	5	..	15.32	16.61	^a Cadivast 10/10 [AF]
NP			^B 4.81	20.13	16.61	^a Caduet 10/10 [AS]

amlodipine 10 mg + atorvastatin 20 mg tablet, 30

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
9054M	1	5	..	16.00	17.29	^a Cadivast 10/20 [AF]
NP			^B 4.86	20.86	17.29	^a Caduet 10/20 [AS]

amlodipine 5 mg + atorvastatin 40 mg tablet, 30

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
9051J	1	5	..	16.21	17.50	^a Cadivast 5/40 [AF]
NP			^B 4.87	21.08	17.50	^a Caduet 5/40 [AS]

amlodipine 10 mg + atorvastatin 80 mg tablet, 30

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
9056P	1	5	..	18.14	19.43	^a Cadivast 10/80 [AF]
NP			^B 5.00	23.14	19.43	^a Caduet 10/80 [AS]

amlodipine 5 mg + atorvastatin 20 mg tablet, 30

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
9050H	1	5	..	15.36	16.65	Cadivast 5/20 [AF]
NP						

amlodipine 5 mg + atorvastatin 80 mg tablet, 30

9052K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	17.48	18.77	^a Cadivast 5/80 [AF]
			^B 4.97	22.45	18.77	^a Caduet 5/80 [AS]

■ **DERMATOLOGICALS**■ **ANTIFUNGALS FOR DERMATOLOGICAL USE****ANTIFUNGALS FOR TOPICAL USE***Imidazole and triazole derivatives*■ **KETOCONAZOLE**Authority required (STREAMLINED)**6434**

Fungal or yeast infection

Population criteria:

- Patient must be an Aboriginal or a Torres Strait Islander person.

ketoconazole 2% cream, 30 g

9024Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	2	..	26.56	27.85	Nizoral 2% Cream [JT]

ketoconazole 2% shampoo, 60 mL

1574W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	1	..	20.99	22.28	Nizoral 2% [JT]

■ **MICONAZOLE**Authority required (STREAMLINED)**6434**

Fungal or yeast infection

Population criteria:

- Patient must be an Aboriginal or a Torres Strait Islander person.

miconazole nitrate 2% cream, 70 g

9028E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	1	..	21.20	22.49	Daktarin [JT]

miconazole 2% solution, 30 mL

9031H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	2	..	23.49	24.78	Daktarin Tincture [JT]

miconazole nitrate 2% powder, 30 g

9029F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	2	..	20.25	21.54	Daktarin [JT]

miconazole nitrate 2% cream, 30 g

9027D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	2	..	19.52	20.81	Daktarin [JT]

Other antifungals for topical use■ **TERBINAFINE**Authority required (STREAMLINED)**6434**

Fungal or yeast infection

Population criteria:

- Patient must be an Aboriginal or a Torres Strait Islander person.

Authority required (STREAMLINED)**6412**

Fungal or yeast infection

Clinical criteria:

- The condition must be fungal; OR
- The condition must be due to yeast.

Population criteria:

- Patient must be 18 years of age or less.

DERMATOLOGICALS

terbinafine hydrochloride 1% cream, 15 g

9160D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	3	..	*38.92	40.21	Lamisil [GK]

ANTIFUNGALS FOR SYSTEMIC USE

Antifungals for systemic use

■ GRISEOFULVIN

griseofulvin 500 mg tablet, 28

2982Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	26.56	27.85	Grisovin 500 [AS]

griseofulvin 125 mg tablet, 100

1460W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	25.75	27.04	Grisovin [AS]

■ TERBINAFINE

Authority required

Dermatophyte infection

Clinical criteria:

- Patient must have failed to respond to topical treatment.

Population criteria:

- Patient must be an Aboriginal or a Torres Strait Islander person.

Authority required

Dermatophyte infection

Clinical criteria:

- Patient must have failed to respond to topical treatment, **AND**
- Patient must have failed to respond to griseofulvin.

Population criteria:

- Patient must be 18 years of age or less.

terbinafine 250 mg tablet, 42

2285G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	30.93	32.22	^a APO-Terbinafine [TX]	^a Lamisil (Novartis Pharmaceuticals Australia Pty Limited) [NV]
						^a NOUMED TERBINAFINE [VO]	^a Tamsil [RW]
						^a Terbinafine AN [EA]	^a Terbinafine-DRLA [RZ]
						^a Terbinafine Sandoz [SZ]	^a Tinasil [AF]

■ TERBINAFINE

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Onychomycosis

Clinical criteria:

- The condition must be proximal or extensive (greater than 80% nail involvement), **AND**
- Patient must have failed to respond to topical treatment, **AND**
- The condition must be due to dermatophyte infection proven by microscopy and confirmed by an Approved Pathology Provider; OR
- The condition must be due to dermatophyte infection proven by culture and confirmed by an Approved Pathology Provider.

The date of the pathology report must be provided at the time of application and must not be more than 12 months old

terbinafine 250 mg tablet, 42

2804N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	1	..	30.93	32.22	^a APO-Terbinafine [TX]	^a Lamisil (Novartis Pharmaceuticals Australia Pty Limited) [NV]
						^a NOUMED TERBINAFINE [VO]	^a Tamsil [RW]
						^a Terbinafine AN [EA]	^a Terbinafine-DRLA [RZ]
						^a Terbinafine Sandoz [SZ]	^a Tinasil [AF]

■ ANTIPSORIATICS

ANTIPSORIATICS FOR TOPICAL USE

Other antipsoriatics for topical use

■ CALCIPOTRIOL + BETAMETHASONE DIPROPIONATE

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)

7947

Chronic stable plaque type psoriasis vulgaris

Clinical criteria:

- The condition must be inadequately controlled by potent topical corticosteroid monotherapy.

calcipotriol 0.005% + betamethasone (as dipropionate) 0.05% gel, 60 g

10075G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	1	..	64.04	41.30	Daivobet 50/500 gel [LO]

■ CALCIPOTRIOL + BETAMETHASONE DIPROPIONATE

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Chronic stable plaque type psoriasis vulgaris

Clinical criteria:

- The condition must be inadequately controlled by potent topical corticosteroid monotherapy.

calcipotriol 0.005% + betamethasone (as dipropionate) 0.05% ointment, 30 g

9494Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	±1	1	..	38.03	39.32	^a Calcipotriol/Betamethasone Sandoz 50/500 [SZ]	^a Daivobet [LO]

calcipotriol 0.005% + betamethasone (as dipropionate) 0.05% foam, 60 g

11091R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	1	..	84.26	41.30	Enstilar [LO]

ANTIPSORIATICS FOR SYSTEMIC USE

Retinoids for treatment of psoriasis

■ ACITRETIN

Caution This drug is a potent teratogen - pregnancy should be avoided during therapy and for at least three years after cessation of therapy.

Note Care must be taken to comply with the provisions of State/Territory law when prescribing this drug.

Authority required (STREAMLINED)

5789

Severe intractable psoriasis

Authority required (STREAMLINED)

5727

Severe disorders of keratinisation

acitretin 25 mg capsule, 100

2020H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	2	..	212.55	41.30	^a Neotigason [TB] ^a ZETIN [RW]	^a Novatin [TX]

acitretin 10 mg capsule, 100

2019G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	2	..	108.27	41.30	^a Neotigason [TB] ^a ZETIN [RW]	^a Novatin [TX]

■ ANTIBIOTICS AND CHEMOTHERAPEUTICS FOR DERMATOLOGICAL USE

CHEMOTHERAPEUTICS FOR TOPICAL USE

Sulfonamides

■ SILVER SULFADIAZINE

Restricted benefit

Infection

Treatment Phase: Prevention and treatment

Clinical criteria:

- The condition must be in partial or full skin thickness loss due to burns; OR

- The condition must be in partial or full skin thickness loss due to epidermolysis bullosa.

Restricted benefit

Stasis ulcers

silver sulfadiazine 1% cream, 50 g

9479X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	‡1	..	..	21.49	22.78	Flamazine [SN]

CORTICOSTEROIDS, DERMATOLOGICAL PREPARATIONS**CORTICOSTEROIDS, PLAIN***Corticosteroids, weak (group I)***■ HYDROCORTISONE ACETATE****Restricted benefit**

Corticosteroid-responsive dermatoses

hydrocortisone acetate 1% ointment, 50 g

2882Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	‡1	1	..	14.16	15.45	^a Cortic-DS 1% [LN]
			^B 2.19	16.35	15.45	^a Sigmacort [AS]

hydrocortisone acetate 1% cream, 50 g

2881P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	‡1	1	..	14.16	15.45	^a Cortic-DS 1% [LN]
			^B 2.19	16.35	15.45	^a Sigmacort [AS]

■ HYDROCORTISONE ACETATE**Restricted benefit**

Corticosteroid-responsive dermatoses

hydrocortisone acetate 1% ointment, 50 g

5114E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	‡1	..	..	14.16	15.45	^a Cortic-DS 1% [LN]
			^B 2.19	16.35	15.45	^a Sigmacort [AS]

hydrocortisone acetate 1% cream, 50 g

5113D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	‡1	..	..	14.16	15.45	^a Cortic-DS 1% [LN]
			^B 2.19	16.35	15.45	^a Sigmacort [AS]

*Corticosteroids, moderately potent (group II)***■ TRIAMCINOLONE****Restricted benefit**

Corticosteroid-responsive dermatoses

triamcinolone acetonide 0.02% ointment, 100 g

2118L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	..	..	*18.96	20.25	^a Tricortone [LN]
			^B 3.28	*22.24	20.25	^a Aristocort 0.02% [AS]

triamcinolone acetonide 0.02% cream, 100 g

2117K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	..	..	*18.96	20.25	^a Tricortone [LN]
			^B 3.28	*22.24	20.25	^a Aristocort 0.02% [AS]

*Corticosteroids, potent (group III)***■ BETAMETHASONE DIPROPIONATE****Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Corticosteroid-responsive dermatoses

betamethasone (as dipropionate) 0.05% ointment, 15 g

1119X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	17.86	19.15	^a Elephrat [OV]
			^B 2.45	20.31	19.15	^a Diprosone [OQ]

betamethasone (as dipropionate) 0.05% cream, 15 g

1115Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	17.86	19.15	^a Eleuphrat [OV]
			^B 2.45	20.31	19.15	^a Diprosone [OQ]

■ BETAMETHASONE DIPROPIONATE**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**6232**

Corticosteroid-responsive dermatoses

Clinical criteria:

- The condition must cover 10-20% of the patient's body surface area.

betamethasone (as dipropionate) 0.05% ointment, 15 g

10795E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*23.70	24.99	^a Eleuphrat [OV]
			^B 4.90	*28.60	24.99	^a Diprosone [OQ]

betamethasone (as dipropionate) 0.05% cream, 15 g

10824Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*23.70	24.99	^a Eleuphrat [OV]
			^B 4.90	*28.60	24.99	^a Diprosone [OQ]

■ BETAMETHASONE DIPROPIONATE**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**6246**

Corticosteroid-responsive dermatoses

Clinical criteria:

- The condition must cover 20-40% of the patient's body surface area.

betamethasone (as dipropionate) 0.05% ointment, 15 g

10820L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	5	..	*35.38	36.67	^a Eleuphrat [OV]
			^B 9.80	*45.18	36.67	^a Diprosone [OQ]

betamethasone (as dipropionate) 0.05% cream, 15 g

10800K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	5	..	*35.38	36.67	^a Eleuphrat [OV]
			^B 9.80	*45.18	36.67	^a Diprosone [OQ]

■ BETAMETHASONE DIPROPIONATE**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**6218**

Corticosteroid-responsive dermatoses

Clinical criteria:

- The condition must cover 40-60% of the patient's body surface area.

betamethasone (as dipropionate) 0.05% ointment, 15 g

10821M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	6	5	..	*47.04	41.30	^a Eleuphrat [OV]
			^B 14.70	*61.74	41.30	^a Diprosone [OQ]

betamethasone (as dipropionate) 0.05% cream, 15 g

10813D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	6	5	..	*47.04	41.30	^a Eleuphrat [OV]
			^B 14.70	*61.74	41.30	^a Diprosone [OQ]

■ BETAMETHASONE DIPROPIONATE**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a

patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)

6263

Corticosteroid-responsive dermatoses

Clinical criteria:

- The condition must cover 60-80% of the patient's body surface area.

betamethasone (as dipropionate) 0.05% ointment, 15 g

10816G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	8	5	..	*58.78	41.30	^a Eleuphrat [OV]
			^B 19.60	*78.38	41.30	^a Diprosone [OQ]

betamethasone (as dipropionate) 0.05% cream, 15 g

10801L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	8	5	..	*58.78	41.30	^a Eleuphrat [OV]
			^B 19.60	*78.38	41.30	^a Diprosone [OQ]

▪ **BETAMETHASONE DIPROPIONATE**

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)

6231

Corticosteroid-responsive dermatoses

Clinical criteria:

- The condition must cover >80% of the patient's body surface area.

betamethasone (as dipropionate) 0.05% ointment, 15 g

10823P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	10	5	..	*70.44	41.30	^a Eleuphrat [OV]
			^B 24.50	*94.94	41.30	^a Diprosone [OQ]

betamethasone (as dipropionate) 0.05% cream, 15 g

10802M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	10	5	..	*70.44	41.30	^a Eleuphrat [OV]
			^B 24.50	*94.94	41.30	^a Diprosone [OQ]

▪ **BETAMETHASONE VALERATE**

Restricted benefit

Corticosteroid-responsive dermatoses

betamethasone (as valerate) 0.02% cream, 100 g

2812B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	..	..	*27.50	28.79	^a Antroquoril [OV]
						^b Cortival 1/5 [LN]
			^B 5.00	*32.50	28.79	^a Celestone-M [OQ]
			^B 5.98	*33.48	28.79	^b Betnovate 1/5 [AS]

▪ **BETAMETHASONE VALERATE**

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Corticosteroid-responsive dermatoses

betamethasone (as valerate) 0.05% cream, 15 g

2813C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	13.75	15.04	^a Cortival 1/2 [LN]
			^B 2.56	16.31	15.04	^a Betnovate 1/2 [AS]

▪ **BETAMETHASONE VALERATE**

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)

6232

Corticosteroid-responsive dermatoses

Clinical criteria:

- The condition must cover 10-20% of the patient's body surface area.

betamethasone (as valerate) 0.05% cream, 15 g

10799J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*15.48	16.77	^a Cortival 1/2 [LN]
			^B 5.12	*20.60	16.77	^a Betnovate 1/2 [AS]

■ BETAMETHASONE VALERATE**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**6246**

Corticosteroid-responsive dermatoses

Clinical criteria:

- The condition must cover 20-40% of the patient's body surface area.

betamethasone (as valerate) 0.05% cream, 15 g

10794D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	5	..	*18.94	20.23	^a Cortival 1/2 [LN]
			^B 10.24	*29.18	20.23	^a Betnovate 1/2 [AS]

■ BETAMETHASONE VALERATE**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**6218**

Corticosteroid-responsive dermatoses

Clinical criteria:

- The condition must cover 40-60% of the patient's body surface area.

betamethasone (as valerate) 0.05% cream, 15 g

10808W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	6	5	..	*22.38	23.67	^a Cortival 1/2 [LN]
			^B 15.36	*37.74	23.67	^a Betnovate 1/2 [AS]

■ BETAMETHASONE VALERATE**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**6263**

Corticosteroid-responsive dermatoses

Clinical criteria:

- The condition must cover 60-80% of the patient's body surface area.

betamethasone (as valerate) 0.05% cream, 15 g

10807T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	8	5	..	*25.90	27.19	^a Cortival 1/2 [LN]
			^B 20.48	*46.38	27.19	^a Betnovate 1/2 [AS]

■ BETAMETHASONE VALERATE**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**6231**

Corticosteroid-responsive dermatoses

Clinical criteria:

- The condition must cover >80% of the patient's body surface area.

betamethasone (as valerate) 0.05% cream, 15 g

10810Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	10	5	..	*29.34	30.63	^a Cortival 1/2 [LN]

^B25.60 *54.94 30.63 ^a Betnovate 1/2 [AS]

■ METHYLPREDNISOLONE

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Corticosteroid-responsive dermatoses

methylprednisolone aceponate 0.1% cream, 15 g

8054X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	18.59	19.88	Advantan [LO]

methylprednisolone aceponate 0.1% ointment, 15 g

8055Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	18.59	19.88	Advantan [LO]

methylprednisolone aceponate 0.1% ointment, 15 g

8128T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	18.59	19.88	Advantan (Fatty) [LO]

■ METHYLPREDNISOLONE

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Eczema

methylprednisolone aceponate 0.1% lotion, 20 g

8618N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	19.18	20.47	Advantan [LO]

■ METHYLPREDNISOLONE

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)

6232

Corticosteroid-responsive dermatoses

Clinical criteria:

- The condition must cover 10-20% of the patient's body surface area.

methylprednisolone aceponate 0.1% cream, 15 g

10842P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*25.16	26.45	Advantan [LO]

methylprednisolone aceponate 0.1% ointment, 15 g

10846W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*25.16	26.45	Advantan [LO]

methylprednisolone aceponate 0.1% ointment, 15 g

10848Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*25.16	26.45	Advantan (Fatty) [LO]

methylprednisolone aceponate 0.1% lotion, 20 g

10856J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*26.34	27.63	Advantan [LO]

■ METHYLPREDNISOLONE

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)

6246

Corticosteroid-responsive dermatoses

Clinical criteria:

- The condition must cover 20-40% of the patient's body surface area.

methyprednisolone aceponate 0.1% cream, 15 g

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
10855H	4	5	..	*38.30	39.59	Advantan [LO]

methyprednisolone aceponate 0.1% ointment, 15 g

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
10836H	4	5	..	*38.30	39.59	Advantan [LO]

methyprednisolone aceponate 0.1% ointment, 15 g

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
10840M	4	5	..	*38.30	39.59	Advantan (Fatty) [LO]

methyprednisolone aceponate 0.1% lotion, 20 g

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
10838K	3	5	..	*33.51	34.80	Advantan [LO]

■ METHYLPREDNISOLONE**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**6231**

Corticosteroid-responsive dermatoses

Clinical criteria:

- The condition must cover >80% of the patient's body surface area.

methyprednisolone aceponate 0.1% cream, 15 g

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
10833E	10	5	..	*77.74	41.30	Advantan [LO]

methyprednisolone aceponate 0.1% ointment, 15 g

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
10843Q	10	5	..	*77.74	41.30	Advantan (Fatty) [LO]

methyprednisolone aceponate 0.1% ointment, 15 g

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
10845T	10	5	..	*77.74	41.30	Advantan [LO]

methyprednisolone aceponate 0.1% lotion, 20 g

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
10830B	5	5	..	*47.84	41.30	Advantan [LO]

■ METHYLPREDNISOLONE**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**6218**

Corticosteroid-responsive dermatoses

Clinical criteria:

- The condition must cover 40-60% of the patient's body surface area.

methyprednisolone aceponate 0.1% cream, 15 g

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
10835G	6	5	..	*51.42	41.30	Advantan [LO]

methyprednisolone aceponate 0.1% ointment, 15 g

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
10844R	6	5	..	*51.42	41.30	Advantan (Fatty) [LO]

methylprednisolone aceponate 0.1% ointment, 15 g

10853F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	
NP	6	5	..	*51.42	41.30	Advantan [LO]	

■ METHYLPREDNISOLONE**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**6263**

Corticosteroid-responsive dermatoses

Clinical criteria:

- The condition must cover 60-80% of the patient's body surface area.

methylprednisolone aceponate 0.1% cream, 15 g

10851D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	
NP	8	5	..	*64.62	41.30	Advantan [LO]	

methylprednisolone aceponate 0.1% ointment, 15 g

10834F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	
NP	8	5	..	*64.62	41.30	Advantan [LO]	

methylprednisolone aceponate 0.1% ointment, 15 g

10839L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	
NP	8	5	..	*64.62	41.30	Advantan (Fatty) [LO]	

■ METHYLPREDNISOLONE**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**6263**

Corticosteroid-responsive dermatoses

Clinical criteria:

- The condition must cover 60-80% of the patient's body surface area.

Authority required (STREAMLINED)**6218**

Corticosteroid-responsive dermatoses

Clinical criteria:

- The condition must cover 40-60% of the patient's body surface area.

methylprednisolone aceponate 0.1% lotion, 20 g

10852E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	
NP	4	5	..	*40.66	41.30	Advantan [LO]	

■ MOMETASONE**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Corticosteroid-responsive dermatoses

mometasone furoate 0.1% ointment, 15 g

1915T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	
NP	1	..	..	15.33	16.62	^a Momasone [AS]	^a Novasone [AF]
			^b 3.53	18.86	16.62	^a Zatamil [EO]	
						^a Elocon [AL]	

mometasone furoate 0.1% cream, 15 g

1913Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	
NP	1	..	..	15.33	16.62	^a Momasone [AS]	^a Novasone [AF]
			^b 3.53	18.86	16.62	^a Elocon Alcohol Free [AL]	

mometasone furoate 0.1% lotion, 30 mL

8043H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	
	1	..	..	17.25	18.54	^a Momasone [AS]	^a Novasone [AF]

						^a Zatamil [EO]	
			^B 3.53	20.78	18.54	^a Elocon [AL]	
NP							
■ MOMETASONE							
Note Continuing Therapy Only:							
For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.							
Authority required (STREAMLINED)							
6232							
Corticosteroid-responsive dermatoses							
Clinical criteria:							
The condition must cover 10-20% of the patient's body surface area.							
mometasone furoate 0.1% ointment, 15 g							
10812C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	2	5	..	*18.64	19.93	^a Momasone [AS] ^a Zatamil [EO]	^a Novasone [AF]
			^B 7.06	*25.70	19.93	^a Elocon [AL]	
mometasone furoate 0.1% cream, 15 g							
10827W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	2	5	..	*18.64	19.93	^a Momasone [AS]	^a Novasone [AF]
			^B 7.06	*25.70	19.93	^a Elocon Alcohol Free [AL]	
mometasone furoate 0.1% lotion, 30 mL							
10819K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	2	5	..	*22.48	23.77	^a Momasone [AS] ^a Zatamil [EO]	^a Novasone [AF]
			^B 7.06	*29.54	23.77	^a Elocon [AL]	
■ MOMETASONE							
Note Continuing Therapy Only:							
For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.							
Authority required (STREAMLINED)							
6246							
Corticosteroid-responsive dermatoses							
Clinical criteria:							
The condition must cover 20-40% of the patient's body surface area.							
mometasone furoate 0.1% ointment, 15 g							
10814E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	4	5	..	*25.26	26.55	^a Momasone [AS] ^a Zatamil [EO]	^a Novasone [AF]
			^B 14.12	*39.38	26.55	^a Elocon [AL]	
mometasone furoate 0.1% cream, 15 g							
10809X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	4	5	..	*25.26	26.55	^a Momasone [AS]	^a Novasone [AF]
			^B 14.12	*39.38	26.55	^a Elocon Alcohol Free [AL]	
mometasone furoate 0.1% lotion, 30 mL							
10826T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	3	5	..	*27.72	29.01	^a Momasone [AS] ^a Zatamil [EO]	^a Novasone [AF]
			^B 10.59	*38.31	29.01	^a Elocon [AL]	
■ MOMETASONE							
Note Continuing Therapy Only:							
For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.							
Authority required (STREAMLINED)							
6218							
Corticosteroid-responsive dermatoses							
Clinical criteria:							
The condition must cover 40-60% of the patient's body surface area.							

mometasone furoate 0.1% ointment, 15 g

10828X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	6	5	..	*31.86	33.15	^a Momasone [AS] ^a Zatamil [EO]	^a Novasone [AF]
			^B 21.18	*53.04	33.15	^a Elocon [AL]	

mometasone furoate 0.1% cream, 15 g

10815F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	6	5	..	*31.86	33.15	^a Momasone [AS] ^a Elocon Alcohol Free [AL]	^a Novasone [AF]
			^B 21.18	*53.04	33.15		

■ MOMETASONE**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**6263**

Corticosteroid-responsive dermatoses

Clinical criteria:

- The condition must cover 60-80% of the patient's body surface area.

mometasone furoate 0.1% ointment, 15 g

10793C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	8	5	..	*38.54	39.83	^a Momasone [AS] ^a Zatamil [EO]	^a Novasone [AF]
			^B 28.24	*66.78	39.83	^a Elocon [AL]	

mometasone furoate 0.1% cream, 15 g

10818J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	8	5	..	*38.54	39.83	^a Momasone [AS] ^a Elocon Alcohol Free [AL]	^a Novasone [AF]
			^B 28.24	*66.78	39.83		

■ MOMETASONE**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**6231**

Corticosteroid-responsive dermatoses

Clinical criteria:

- The condition must cover >80% of the patient's body surface area.

mometasone furoate 0.1% ointment, 15 g

10791Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	10	5	..	*45.14	41.30	^a Momasone [AS] ^a Zatamil [EO]	^a Novasone [AF]
			^B 35.30	*80.44	41.30	^a Elocon [AL]	

mometasone furoate 0.1% cream, 15 g

10792B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	10	5	..	*45.14	41.30	^a Momasone [AS] ^a Elocon Alcohol Free [AL]	^a Novasone [AF]
			^B 35.30	*80.44	41.30		

mometasone furoate 0.1% lotion, 30 mL

10804P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	5	5	..	*38.19	39.48	^a Momasone [AS] ^a Zatamil [EO]	^a Novasone [AF]
			^B 17.65	*55.84	39.48	^a Elocon [AL]	

■ MOMETASONE**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**6263**

Corticosteroid-responsive dermatoses

Clinical criteria:

- The condition must cover 60-80% of the patient's body surface area.

Authority required (STREAMLINED)**6218**

Corticosteroid-responsive dermatoses

Clinical criteria:

- The condition must cover 40-60% of the patient's body surface area.

mometasone furoate 0.1% lotion, 30 mL

10805Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	4	5	..	*32.94	34.23	^a Momasone [AS]	^a Novasone [AF]
			^B 14.12	*47.06	34.23	^a Zatamil [EO]	
						^a Elocon [AL]	

*Corticosteroids, very potent (group IV)***■ CLOBETASOL****Authority required (STREAMLINED)****5461**

Moderate to severe scalp psoriasis

Clinical criteria:

- The condition must be inadequately controlled with either a vitamin D analogue or potent topical corticosteroid as monotherapy; OR
- The condition must be inadequately controlled with combination use of a vitamin D analogue and potent topical corticosteroid.

Population criteria:

- Patient must be aged 18 years or older.

clobetasol propionate 0.05% shampoo, 125 mL

10080M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	1	..	47.86	41.30	Clobex [GA]

■ ANTI-ACNE PREPARATIONS**ANTI-ACNE PREPARATIONS FOR TOPICAL USE***Retinoids for topical use in acne***■ ADAPALENE + BENZOYL PEROXIDE****Restricted benefit**

Severe acne vulgaris

Treatment Phase: Acute treatment

Clinical criteria:

- The treatment must be in combination with an oral antibiotic.

adapalene 0.1% + benzoyl peroxide 2.5% gel, 30 g

8954G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	1	..	34.71	36.00	Epiduo [GA]

■ ADAPALENE + BENZOYL PEROXIDE**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Severe acne vulgaris

Clinical criteria:

- The treatment must be maintenance therapy.

adapalene 0.1% + benzoyl peroxide 2.5% gel, 30 g

8955H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	3	..	34.71	36.00	Epiduo [GA]

ANTI-ACNE PREPARATIONS FOR SYSTEMIC USE*Retinoids for treatment of acne***■ ISOTRETINOIN****Caution** This drug causes birth defects.

This drug has been reported to cause other frequent and potentially serious toxicity.

Note Care must be taken to comply with the provisions of State/Territory law when prescribing this drug.**Authority required (STREAMLINED)****5224**

Severe cystic acne

Clinical criteria:

- The condition must be unresponsive to other therapy.

isotretinoin 30 mg capsule, 60

11921K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	61.98	41.30	Oratane [OU]

isotretinoin 5 mg capsule, 60

11716P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	23.86	25.15	Oratane [OU]

isotretinoin 40 mg capsule, 30

2549E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	3	..	45.33	41.30	^a Dermatane [ER]	^a Oratane [RF]

isotretinoin 10 mg capsule, 60

2591J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	3	..	35.70	36.99	^a APO-Isotretinoin [TX] ^a Isotretinoin AN [EA] ^a Isotretinoin Lupin [GQ] ^a Rocta 10 [RW]	^a Dermatane [ER] ^a Isotretinoin GX [SZ] ^a Oratane [RF]

ISOTRETINOIN

Caution This drug causes birth defects.

This drug has been reported to cause other frequent and potentially serious toxicity.

Note Pharmaceutical benefits that have form pack size isotretinoin 20 mg capsule, 60 and isotretinoin 20 mg capsule, 30 are equivalent for the purposes of substitution.

Note Care must be taken to comply with the provisions of State/Territory law when prescribing this drug.

Authority required (STREAMLINED)

5224

Severe cystic acne

Clinical criteria:

- The condition must be unresponsive to other therapy.

isotretinoin 20 mg capsule, 60

2592K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	3	..	48.92	41.30	^a APO-Isotretinoin [TX] ^a Isotretinoin AN [EA] ^a Isotretinoin Lupin [GQ] ^a Oratane [RF]	^a Dermatane [ER] ^a Isotretinoin GX [SZ] ^a Isotretinoin SCP 20 [CR] ^a Rocta 20 [RW]

isotretinoin 20 mg capsule, 30

11621P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	3	..	*48.92	41.30	^a Roaccutane [RO]

OTHER DERMATOLOGICAL PREPARATIONS**OTHER DERMATOLOGICAL PREPARATIONS**

Agents for dermatitis, excluding corticosteroids

DUPILUMAB

Note Instructions on the use of the Eczema Area and Severity Index and copyright details can be found here: <https://www.dupixent.co.uk/-/media/EMS/Conditions/Dermatology/Brands/Dupixent-UK/global/1051-EASI-Leaflet-v6-webready.pdf>

Note Instructions on the use of the Dermatology Life Quality Index and copyright details can be found here: <https://www.cardiff.ac.uk/medicine/resources/quality-of-life-questionnaires/dermatology-life-quality-index>

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Chronic severe atopic dermatitis

Treatment Phase: Initial treatment of the whole body

Clinical criteria:

- Patient must have a Physicians Global Assessment (PGA) (5-point scale) baseline score of at least 4 as evidence of severe disease despite treatment with daily topical therapy (corticosteroid of medium to high potency/calcineurin inhibitor), for at least 28 day, **AND**

- Patient must have an Eczema Area and Severity Index (EASI) baseline score of at least 20 despite treatment with daily topical therapy (corticosteroid of medium to high potency/calcineurin inhibitor), for at least 28 days, **AND**
- Patient must have an age appropriate Dermatology Life Quality Index (DLQI) baseline score (of any value) measured following treatment with daily topical therapy (corticosteroid of medium to high potency/calcineurin inhibitor), for at least 28 days, **AND**
- The condition must have had lesions for at least 6 months from the time of the initial diagnosis of chronic severe atopic dermatitis affecting either of: (i) the whole body, (ii) face/hands, **AND**
- The treatment must be the sole PBS-subsidised biological medicine for this PBS indication, **AND**
- Patient must not have experienced an inadequate response to this biological medicine in this PBS indication.

Treatment criteria:

- Must be treated by a dermatologist; OR
- Must be treated by a clinical immunologist.

Population criteria:

- Patient must be 12 years of age or older.

State each of the qualifying PGA, EASI and DLQI scores in the authority application. These baseline scores must have been measured within the past 4 weeks. The name/s of the medium to high potency topical corticosteroids trialled prior to commencing treatment with this biological medicine is/are to be documented in the patient's medical records.

The EASI and DLQI baseline measurements are to form the basis of determining if an adequate response to treatment has been achieved under the Continuing treatment restriction.

Note Instructions on the use of the Physician's Global Assessment (5-point scale) can be obtained from Sanofi Medical Information on 1800 818 806 or MedInfo.Australia@sanofi.com

Authority required

Chronic severe atopic dermatitis

Treatment Phase: Continuing or resuming treatment of the whole body

Clinical criteria:

- Patient must have received PBS-subsidised treatment with this biological medicine for the treatment of chronic severe atopic dermatitis affecting the whole body, **AND**
- Patient must have achieved an adequate response within the first 16 weeks of treatment; OR
- Patient must have maintained an adequate response to their most recent course of PBS-subsidised treatment with this biological medicine for this PBS indication if this is the second or subsequent Continuing treatment authority application; OR
- Patient must have temporarily ceased treatment for reasons other than lack of response (e.g. family planning, vaccination with live vaccines, adverse-effect investigation), thereby being unable to achieve/maintain an adequate response immediately prior to this authority application, **AND**
- The treatment must be the sole PBS-subsidised biological medicine for this PBS indication.

Treatment criteria:

- Must be treated by a dermatologist; OR
- Must be treated by a clinical immunologist.

For the purposes of this restriction, an adequate response to treatment is defined as:

(a) An improvement/maintenance in the Eczema Area and Severity Index (EASI) score of at least 50% compared to baseline; and

(b) An improvement/maintenance in Dermatology Life Quality Index (DLQI) score of at least 4 points compared to baseline

Where an initial baseline (post-topical corticosteroid, pre-biological medicine) DLQI score was not measured for a patient who had commenced treatment through a clinical trial, early access program or through private, non-PBS-subsidised supply, an absence of worsening in the current DLQI score compared to that measured at the time of the 'Grandfather listing' authority application will suffice as an adequate response for requirement (b) above.

State each of the current EASI and DLQI scores for this authority application.

Authority required

Chronic severe atopic dermatitis

Treatment Phase: Initial treatment of the face and/or hands

Clinical criteria:

- The condition must have at least 2 of the following Eczema Area and Severity Index (EASI) symptom sub-scores for erythema, oedema/papulation, excoriation, lichenification rated as severe despite treatment with daily topical therapy (corticosteroid of medium to high potency/calcineurin inhibitor), for at least 28 days; OR
- The condition must have affected at least 30% of the face/hands surface area despite treatment with daily topical therapy (corticosteroid of medium to high potency/calcineurin inhibitor), for at least 28 days, **AND**
- Patient must have an age appropriate Dermatology Life Quality Index (DLQI) baseline score (of any value) measured following treatment with daily topical therapy (corticosteroid of medium to high potency/calcineurin inhibitor), for at least 28 days, **AND**
- The condition must have had lesions for at least 6 months from the time of the initial diagnosis of chronic severe atopic dermatitis affecting either of: (i) the whole body, (ii) face/hands, **AND**
- The treatment must be the sole PBS-subsidised biological medicine for this PBS indication, **AND**
- Patient must not have experienced an inadequate response to this biological medicine in this PBS indication.

Treatment criteria:

- Must be treated by a dermatologist; OR
- Must be treated by a clinical immunologist.

Population criteria:

- Patient must be 12 years of age or older.

State each of the 4 Eczema Area and Severity Index (EASI) symptom sub-score ratings for erythema, oedema/papulation, excoriation, lichenification, in the authority application. These 4 symptom sub-score ratings must have been assessed within the past 4 weeks. The name/s of the medium to high potency topical corticosteroids trialled prior to commencing treatment with this biological medicine is/are to be documented in the patient's medical records.

Authority required

Chronic severe atopic dermatitis

Treatment Phase: Continuing or resuming treatment of the face and/or hands

Clinical criteria:

- Patient must have received PBS-subsidised treatment with this biological medicine for the treatment of chronic severe atopic dermatitis affecting the face/hands, **AND**
- Patient must have achieved an adequate response within the first 16 weeks of treatment; OR
- Patient must have maintained an adequate response to their most recent course of PBS-subsidised treatment with this biological medicine for this PBS indication if this is the second or subsequent Continuing treatment authority application; OR
- Patient must have temporarily ceased treatment for reasons other than lack of response (e.g. family planning, vaccination with live vaccines, adverse-effect investigation), thereby being unable to achieve/maintain an adequate response immediately prior to this authority application, **AND**
- The treatment must be the sole PBS-subsidised biological medicine for this PBS indication.

Treatment criteria:

- Must be treated by a dermatologist; OR
- Must be treated by a clinical immunologist.

For the purposes of this restriction, an adequate response to treatment of the face/hands is defined as:

(a) (i) A rating of either mild (1) to none (0) on at least 3 of the assessments of erythema, oedema/papulation, excoriation and lichenification mentioned in the Eczema Area and Severity Index (EASI); or

(ii) At least a 75% reduction in the skin area affected by this condition compared to baseline; and

(b) An improvement in Dermatology Life Quality Index (DLQI) score of at least 4 points compared to baseline

Where an initial baseline (post-topical corticosteroid, pre-biological medicine) DLQI score was not measured for a patient who had commenced treatment through a clinical trial, early access program or through private, non-PBS-subsidised supply, an absence of worsening in the current DLQI score compared to that measured at the time of the 'Grandfather listing' authority application will suffice as an adequate response for requirement (b) above.

Document each qualifying response measure in the patient's medical records for PBS compliance auditing purposes

Authority required

Chronic severe atopic dermatitis

Treatment Phase: Transitioning from non-PBS to PBS-subsidised supply - treatment of the whole body (Grandfather listing)

Clinical criteria:

- Patient must have been receiving treatment with this biological medicine for this PBS indication prior to 1 March 2021, **AND**
- Patient must have had a Physicians Global Assessment (PGA) baseline score of at least 4 as evidence of severe disease despite treatment with daily topical therapy (corticosteroid of medium to high potency/calcineurin inhibitor), for at least 28 days prior to commencing non-PBS-subsidised therapy with this biological medicine, **AND**
- Patient must have had an Eczema Area and Severity Index (EASI) baseline score of at least 20 despite treatment with daily topical therapy (corticosteroid of medium to high potency/calcineurin inhibitor), for at least 28 days prior to commencing non-PBS-subsidised therapy with this biological medicine, **AND**
- Patient must have an age appropriate Dermatology Life Quality Index (DLQI) baseline score (of any value) measured following treatment with daily topical therapy (corticosteroid of medium to high potency/calcineurin inhibitor), for at least 28 days, prior to having commenced non-PBS-subsidised therapy with this biological medicine; OR
- Patient must have, where the above baseline DLQI was not recorded in the patient's medical records, a current age-appropriate DLQI score (of any value) measured, **AND**
- The condition must have had lesions for at least 6 months from the time of the initial diagnosis of chronic severe atopic dermatitis affecting either of: (i) the whole body, (ii) face/hands, prior to commencing non-PBS-subsidised therapy with this biological medicine, **AND**
- Patient must not be experiencing an inadequate response to current non-PBS-subsidised therapy with this biological medicine, **AND**
- The treatment must be the sole PBS-subsidised biological medicine for this PBS indication, **AND**
- Patient must not have experienced an inadequate response to this biological medicine in this indication, prior to commencing non-PBS-subsidised therapy with this biological medicine.

Treatment criteria:

- Must be treated by a dermatologist; OR
- Must be treated by a clinical immunologist.

Population criteria:

- Patient must be 12 years of age or older.

State each of the qualifying PGA, EASI and DLQI scores in the authority application. The name/s of the medium to high potency topical corticosteroids trialled prior to commencing treatment with this biological medicine must be documented in the patient's medical records.

The EASI and DLQI baseline measurements are to form the basis of determining if an adequate response to treatment has been achieved under the Continuing treatment restriction.

A Grandfathered patient may qualify for PBS-subsidised treatment under this restriction once only.

For continuing PBS-subsidised treatment, a Grandfathered patient must qualify under the Continuing treatment criteria.

Note This grandfather restriction will cease to operate from 12 months after the date specified in the clinical criteria.

Note Instructions on the use of the Physician's Global Assessment (5-point scale) can be obtained from Sanofi Medical Information on 1800 818 806 or MedInfo.Australia@sanofi.com

Authority required

Chronic severe atopic dermatitis

Treatment Phase: Transitioning from non-PBS to PBS-subsidised supply - treatment of the face and/or hands (Grandfather listing)

Clinical criteria:

- Patient must have been receiving treatment with this biological medicine for this PBS indication prior to 1 March 2021, **AND**
- The condition must have had at least 2 of the following Eczema Area and Severity Index (EASI) symptom sub-scores for erythema, oedema/papulation, excoriation, lichenification rated as severe despite treatment with daily topical therapy (corticosteroid of medium to high potency/calcineurin inhibitor), for at least 28 days, prior to commencing non-PBS-subsidised therapy with this biological medicine; **OR**
- The condition must have affected at least 30% of the face/hands surface area despite treatment with daily topical therapy (corticosteroid of medium to high potency/calcineurin inhibitor), for at least 28 days, prior to commencing non-PBS-subsidised therapy with this biological medicine, **AND**
- Patient must have an age appropriate Dermatology Life Quality Index (DLQI) baseline score (of any value) measured following treatment with daily topical therapy (corticosteroid of medium to high potency/calcineurin inhibitor), for at least 28 days, prior to having commenced non-PBS-subsidised therapy with this biological medicine; **OR**
- Patient must have, where the above baseline DLQI was not recorded in the patient's medical records, a current age-appropriate DLQI score (of any value) measured, **AND**
- The condition must have had lesions for at least 6 months from the time of the initial diagnosis of chronic severe atopic dermatitis affecting either of: (i) the whole body, (ii) face/hands, prior to commencing non-PBS-subsidised therapy with this biological medicine, **AND**
- Patient must not be experiencing an inadequate response to current non-PBS-subsidised therapy with this biological medicine, **AND**
- The treatment must be the sole PBS-subsidised biological medicine for this condition, **AND**
- Patient must not have experienced an inadequate response to this biological medicine in this indication, prior to commencing non-PBS-subsidised therapy with this biological medicine.

Treatment criteria:

- Must be treated by a dermatologist; **OR**
- Must be treated by a clinical immunologist.

Population criteria:

- Patient must be 12 years of age or older.

State each of the 4 Eczema Area and Severity Index (EASI) symptom sub-score ratings for erythema, oedema/papulation, excoriation, lichenification that were present prior to having commenced non-PBS-subsidised therapy, in the authority application. The name/s of the medium to high potency topical corticosteroids trialed prior to commencing treatment with this biological medicine is/are to be documented in the patient's medical records.

A Grandfathered patient may qualify for PBS-subsidised treatment under this restriction once only.

For continuing PBS-subsidised treatment, a Grandfathered patient must qualify under the Continuing treatment criteria.

Note This grandfather restriction will cease to operate from 12 months after the date specified in the clinical criteria.

dupilumab 300 mg/2 mL injection, 2 x 2 mL syringes

12292Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1754.22	41.30	Dupixent [SW]

dupilumab 200 mg/1.14 mL injection, 2 x 1.14 mL syringes

12291X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1754.22	41.30	Dupixent [SW]

■ PIMECROLIMUS

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

5482

Atopic dermatitis

Population criteria:

- Patient must be at least 3 months of age.

Clinical criteria:

- The condition must be on the patient's face; **OR**
- The condition must be on the patient's eyelid, **AND**
- Patient must have 1 or more of the following contraindications to topical corticosteroids: (i) perioral dermatitis; (ii) periorbital dermatitis; (iii) rosacea; (iv) epidermal atrophy; (v) dermal atrophy; (vi) allergy to topical corticosteroids; (vii) cataracts; (viii) glaucoma; (ix) raised intraocular pressure, **AND**
- Patient must not receive more than two 15 g packs of PBS-subsidised pimecrolimus per 6-month period.

Authority required (STREAMLINED)

5472

Atopic dermatitis

Treatment Phase: Short-term (up to 3 weeks) intermittent treatment

Population criteria:

- Patient must be at least 3 months of age.

Clinical criteria:

- The condition must be on the patient's face; OR
 - The condition must be on the patient's eyelid, **AND**
 - Patient must have failed to achieve satisfactory disease control with intermittent topical corticosteroid therapy, **AND**
 - The condition must have been initially diagnosed more than three months prior to this treatment, **AND**
 - Patient must not receive more than two 15 g packs of PBS-subsidised pimecrolimus per 6-month period.
- Failure to achieve satisfactory disease control with intermittent topical corticosteroid therapy is manifest by:
- (i) failure of the facial skin to clear despite at least 2 weeks of topical hydrocortisone 1% applied every day; or
 - (ii) failure of the facial skin to clear despite at least 1 week of a moderate or potent topical corticosteroid applied every day; or
 - (iii) clearing of the facial skin with at least 2 weeks of topical hydrocortisone 1% applied every day, but almost immediate and significant flare in facial disease (within 48 hours) upon stopping topical corticosteroids, occurring on at least 2 consecutive occasions; or
 - (iv) clearing of the facial skin with at least 1 week of a moderate or potent topical corticosteroid applied every day, but almost immediate and significant flare in facial disease (within 48 hours) upon stopping topical corticosteroids, occurring on at least 2 consecutive occasions

pimecrolimus 1% cream, 15 g

8802G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	1	..	31.35	32.64	Elidel [GO]

Other dermatologicals**■ DAPSONE****Note Shared Care Model:**

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

dapsone 100 mg tablet, 100

1272Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	309.07	41.30	Link Medical Products Pty Ltd [LM]

dapsone 25 mg tablet, 100

8801F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	253.17	41.30	Link Medical Products Pty Ltd [LM]

■ IMIQUIMOD

Note The patient or carer must be able to understand and administer the imiquimod dosing regimen.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Treatment of recurrent (previously treated) lesions will not be authorised.

Note Pharmaceutical benefits that have the form imiquimod single use sachets and pharmaceutical benefits that have the form imiquimod multi-use pump are equivalent for the purposes of substitution.

Authority required

Superficial basal cell carcinoma

Clinical criteria:

- The condition must be previously untreated, **AND**
- The condition must be confirmed by biopsy, **AND**
- Patient must have normal immune function, **AND**
- The condition must not be suitable for treatment with surgical excision; OR
- The condition must not be suitable for treatment with cryotherapy; OR
- The condition must not be suitable for treatment with curettage with diathermy, **AND**
- Patient must require topical drug therapy.

The date of the pathology report and name of the Approved Pathology Authority must be provided at the time of application.

imiquimod 5% cream, 12 x 250 mg sachets

2546B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	1	..	87.55	41.30	^a Aldiq [AF]	^a APO-Imiquimod [TX]
			^B 2.28	89.83	41.30	^a Aldara [IL]	

imiquimod 5% cream, 2 x 2 g

2637T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	^B 4.55	92.10	41.30	^a Aldara Pump [IL]

GENITO URINARY SYSTEM AND SEX HORMONES

OTHER GYNECOLOGICALS

CONTRACEPTIVES FOR TOPICAL USE

Intrauterine contraceptives

LEVONORGESTREL

Restricted benefit

Contraception

levonorgestrel 19.5 mg intrauterine drug delivery system, 1 system

11909T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	170.35	41.30	Kyleena [SY]

LEVONORGESTREL

Restricted benefit

Contraception

Restricted benefit

Idiopathic menorrhagia

Clinical criteria:

- The treatment must be in a patient where oral treatments are ineffective.

Restricted benefit

Idiopathic menorrhagia

Clinical criteria:

- The treatment must be in a patient where oral treatments are contraindicated.

levonorgestrel 52 mg intrauterine drug delivery system, 1 system

8633J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	213.25	41.30	Mirena [BN]

OTHER GYNECOLOGICALS

Prolactine inhibitors

BROMOCRIPTINE

Restricted benefit

Prevention of the onset of lactation

Clinical criteria:

- The treatment must occur in the puerperium, **AND**
- The treatment must be for medical reasons.

bromocriptine 2.5 mg tablet, 30

1444B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	22.89	24.18	Parlodel [SZ]

BROMOCRIPTINE

Caution Care should be taken when treating patients with advanced age and significant cognitive impairment with dopamine agonists.

Restricted benefit

Acromegaly

Restricted benefit

Parkinson disease

Restricted benefit

Pathological hyperprolactinaemia

Clinical criteria:

- Patient must be one in whom surgery is not indicated.

Restricted benefit

Pathological hyperprolactinaemia

Clinical criteria:

- Patient must have had surgery for this condition with incomplete resolution.

Restricted benefit

Pathological hyperprolactinaemia

Clinical criteria:

- Patient must be one in whom radiotherapy is not indicated.

Restricted benefit

Pathological hyperprolactinaemia

Clinical criteria:

- Patient must have had radiotherapy for this condition with incomplete resolution.

bromocriptine 2.5 mg tablet, 30

1443Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*33.76	35.05	Parlodel [SZ]

■ CABERGOLINE
Restricted benefit

Prevention of the onset of lactation

Clinical criteria:

- The treatment must occur in the puerperium, **AND**
- The treatment must be for medical reasons.

cabergoline 500 microgram tablet, 2

8115D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	25.38	26.67	^a APO-Cabergoline [TX]	^a Dostinex [PF]

■ CABERGOLINE
Restricted benefit

Pathological hyperprolactinaemia

Clinical criteria:

- Patient must be one in whom surgery is not indicated.

Restricted benefit

Pathological hyperprolactinaemia

Clinical criteria:

- Patient must have had surgery for this condition with incomplete resolution.

Restricted benefit

Pathological hyperprolactinaemia

Clinical criteria:

- Patient must be one in whom radiotherapy is not indicated.

Restricted benefit

Pathological hyperprolactinaemia

Clinical criteria:

- Patient must have had radiotherapy for this condition with incomplete resolution.

cabergoline 500 microgram tablet, 8

8114C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	65.48	41.30	^a APO-Cabergoline [TX]	^a Dostinex [PF]

■ QUINAGOLIDE
Restricted benefit

Pathological hyperprolactinaemia

Clinical criteria:

- Patient must be one in whom surgery is not indicated.

Restricted benefit

Pathological hyperprolactinaemia

Clinical criteria:

- Patient must have had surgery for this condition with incomplete resolution.

Restricted benefit

Pathological hyperprolactinaemia

Clinical criteria:

- Patient must be one in whom radiotherapy is not indicated.

Restricted benefit

Pathological hyperprolactinaemia

Clinical criteria:

- Patient must have had radiotherapy for this condition with incomplete resolution.

quinagolide 75 microgram tablet, 30

8822H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	91.60	41.30	Norprolac [FP]

■ SEX HORMONES AND MODULATORS OF THE GENITAL SYSTEM
HORMONAL CONTRACEPTIVES FOR SYSTEMIC USE

Progestogens and estrogens, fixed combinations

■ **LEVONORGESTREL + ETHINYLESTRADIOL****levonorgestrel 150 microgram + ethinylestradiol 30 microgram tablet [21] (&) inert substance tablet [7], 4 x 28**

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
1394J	1	2	..	18.13	19.42	^a Monofeme 28 [FZ]	
						^b Eleanor 150/30 ED [EA]	^b Evelyn 150/30 ED [GQ]
						^b Femme-Tab ED 30/150 [AE]	^b Lenest 30 ED [AF]
						^b Micronelle 30 ED [TX]	
			^B 3.96	22.09	19.42	^b Levlen ED [SY]	
			^B 11.56	29.69	19.42	^a Nordette 28 [PF]	

levonorgestrel 125 microgram + ethinylestradiol 50 microgram tablet [21] (&) inert substance tablet [7], 4 x 28

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
1456P	1	2	..	21.89	23.18	Microgynon 50 ED [BN]

levonorgestrel 100 microgram + ethinylestradiol 20 microgram tablet [21] (&) inert substance tablet [7], 4 x 28

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
2416E	1	2	..	18.13	19.42	Femme-Tab ED 20/100 [AE]

■ **NORETHISTERONE + ETHINYLESTRADIOL****norethisterone 500 microgram + ethinylestradiol 35 microgram tablet [21] (&) inert substance tablet [7], 4 x 28**

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
2774B	1	2	..	20.75	22.04	^a Norimin 28 Day [FZ]
			^B 9.78	30.53	22.04	^a Brevinor [PF]

■ **NORETHISTERONE + ETHINYLESTRADIOL**

Note Pharmaceutical benefits that have the form norethisterone 1 mg + ethinylestradiol 35 microgram tablet in a 4 pack of 28 tablets can be substituted for a 3 pack of 28 tablets in the case of a shortage.

norethisterone 1 mg + ethinylestradiol 35 microgram tablet [21] (&) inert substance tablet [7], 3 x 28

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
12211Q	1.3	2	..	*63.84	41.30	^a Pirmella 1/35 [DZ]

norethisterone 1 mg + ethinylestradiol 35 microgram tablet [21] (&) inert substance tablet [7], 4 x 28

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
2775C	1	2	..	20.75	22.04	^a Norimin-1 28 Day [FZ]
			^B 9.78	30.53	22.04	^a Brevinor-1 [PF]

■ **NORETHISTERONE + MESTRANOL****norethisterone 1 mg + mestranol 50 microgram tablet [21] (&) inert substance tablet [7], 4 x 28**

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
3179H	1	2	..	21.65	22.94	Norinyl-1/28 [PF]

Progestogens and estrogens, sequential preparations■ **LEVONORGESTREL + ETHINYLESTRADIOL****levonorgestrel 50 microgram + ethinylestradiol 30 microgram tablet [6] (&) levonorgestrel 75 microgram + ethinylestradiol 40 microgram tablet [5] (&) levonorgestrel 125 microgram + ethinylestradiol 30 microgram tablet [10] (&) inert substance tablet [7], 4 x 28**

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
1392G	1	2	..	19.74	21.03	^a Trifeme 28 [FZ]
						^b Logynon ED [SY]
			^B 13.00	32.74	21.03	^a Triphasil 28 [PF]
			^B 13.56	33.30	21.03	^b Triquilar ED [BN]

Progestogens■ **ETONOGESTREL****etonogestrel 68 mg implant, 1**

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8487Q	1	..	..	170.35	41.30	Implanon NXT [OQ]

■ **LEVONORGESTREL****levonorgestrel 30 microgram tablet, 112 tablets [4 x 28]**

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
2913H	1	2	..	19.72	21.01	Microlut 28 [BN]

■ MEDROXYPROGESTERONE

medroxyprogesterone acetate 150 mg/mL injection, 1 mL vial

3118D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	26.15	27.44	^a Depo-Ralovera [FZ]
			^B 7.00	33.15	27.44	^a Depo-Provera [PF]

■ NORETHISTERONE

norethisterone 350 microgram tablet, 4 x 28

1967M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	20.75	22.04	Noriday 28 Day [PF]

ANDROGENS

3-oxoandrosten (4) derivatives

■ TESTOSTERONE

Authority required

Androgen deficiency

Clinical criteria:

- Patient must have an established pituitary or testicular disorder.

Treatment criteria:

- Must be treated by a specialist general paediatrician, specialist paediatric endocrinologist, specialist urologist, specialist endocrinologist or a Fellow of the Australasian Chapter of Sexual Health Medicine; or in consultation with one of these specialists; or have an appointment to be assessed by one of these specialists.

The name of the specialist must be included in the authority application.

Authority required

Androgen deficiency

Clinical criteria:

- Patient must not have an established pituitary or testicular disorder, **AND**
- The condition must not be due to age, obesity, cardiovascular diseases, infertility or drugs.

Population criteria:

- Patient must be aged 40 years or older.

Treatment criteria:

- Must be treated by a specialist urologist, specialist endocrinologist or a Fellow of the Australasian Chapter of Sexual Health Medicine; or in consultation with one of these specialists; or have an appointment to be assessed by one of these specialists.

Androgen deficiency is defined as:

(i) testosterone level of less than 6 nmol per litre; OR

(ii) testosterone level between 6 and 15 nmol per litre with high luteinising hormone (LH) (greater than 1.5 times the upper limit of the eugonadal reference range for young men, or greater than 14 IU per litre, whichever is higher).

Androgen deficiency must be confirmed by at least two morning blood samples taken on different mornings.

The dates and levels of the qualifying testosterone and LH measurements must be, or must have been provided in the authority application when treatment with this drug is or was initiated.

The name of the specialist must be included in the authority application.

Authority required

Micropenis

Population criteria:

- Patient must be under 18 years of age.

Treatment criteria:

- Must be treated by a specialist general paediatrician, specialist paediatric endocrinologist, specialist urologist, specialist endocrinologist or a Fellow of the Australasian Chapter of Sexual Health Medicine; or in consultation with one of these specialists; or have an appointment to be assessed by one of these specialists.

The name of the specialist must be included in the authority application.

Authority required

Pubertal induction

Population criteria:

- Patient must be under 18 years of age.

Treatment criteria:

- Must be treated by a specialist general paediatrician, specialist paediatric endocrinologist, specialist urologist, specialist endocrinologist or a Fellow of the Australasian Chapter of Sexual Health Medicine; or in consultation with one of these specialists; or have an appointment to be assessed by one of these specialists.

The name of the specialist must be included in the authority application.

Authority required

Constitutional delay of growth or puberty

Population criteria:

- Patient must be under 18 years of age.

Treatment criteria:

- Must be treated by a specialist general paediatrician, specialist paediatric endocrinologist, specialist urologist, specialist endocrinologist or a Fellow of the Australasian Chapter of Sexual Health Medicine; or in consultation with one of these specialists; or have an appointment to be assessed by one of these specialists.

The name of the specialist must be included in the authority application.

testosterone 2% (23 mg/actuation) gel, 56 actuations

11740X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	5	..	82.97	41.30	Testavan [FP]

testosterone 1% (12.5 mg/actuation) gel, 2 x 60 actuations

10380H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	4	..	77.52	41.30	Testogel [HB]

testosterone 5 mg/24 hours patch, 30

8619P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	5	..	78.05	41.30	Androderm [TB]

testosterone 1% (50 mg/5 g) gel, 30 x 5 g sachets

8830R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	5	..	77.52	41.30	Testogel [HB]

testosterone 5% (50 mg/mL) cream, 50 mL

10378F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	6	..	65.76	41.30	AndroForte 5 [LX]

testosterone 2.5 mg/24 hours patch, 60

8460G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	5	..	78.05	41.30	Androderm [TB]

■ TESTOSTERONE UNDECANOATE

Authority required

Androgen deficiency

Clinical criteria:

- Patient must have an established pituitary or testicular disorder.

Treatment criteria:

- Must be treated by a specialist general paediatrician, specialist paediatric endocrinologist, specialist urologist, specialist endocrinologist or a Fellow of the Australasian Chapter of Sexual Health Medicine; or in consultation with one of these specialists; or have an appointment to be assessed by one of these specialists.

The name of the specialist must be included in the authority application.

Authority required

Androgen deficiency

Clinical criteria:

- Patient must not have an established pituitary or testicular disorder, **AND**
- The condition must not be due to age, obesity, cardiovascular diseases, infertility or drugs.

Population criteria:

- Patient must be aged 40 years or older.

Treatment criteria:

- Must be treated by a specialist urologist, specialist endocrinologist or a Fellow of the Australasian Chapter of Sexual Health Medicine; or in consultation with one of these specialists; or have an appointment to be assessed by one of these specialists.

Androgen deficiency is defined as:

- (i) testosterone level of less than 6 nmol per litre; OR
- (ii) testosterone level between 6 and 15 nmol per litre with high luteinising hormone (LH) (greater than 1.5 times the upper limit of the eugonadal reference range for young men, or greater than 14 IU per litre, whichever is higher).

Androgen deficiency must be confirmed by at least two morning blood samples taken on different mornings.

The dates and levels of the qualifying testosterone and LH measurements must be, or must have been provided in the authority application when treatment with this drug is or was initiated.

The name of the specialist must be included in the authority application.

Authority required

Micropenis

Population criteria:

- Patient must be under 18 years of age.

Treatment criteria:

- Must be treated by a specialist general paediatrician, specialist paediatric endocrinologist, specialist urologist, specialist endocrinologist or a Fellow of the Australasian Chapter of Sexual Health Medicine; or in consultation with one of these specialists; or have an appointment to be assessed by one of these specialists.

The name of the specialist must be included in the authority application.

Authority required

Pubertal induction

Population criteria:

- Patient must be under 18 years of age.

Treatment criteria:

- Must be treated by a specialist general paediatrician, specialist paediatric endocrinologist, specialist urologist, specialist endocrinologist or a Fellow of the Australasian Chapter of Sexual Health Medicine; or in consultation with one of these specialists; or have an appointment to be assessed by one of these specialists.

The name of the specialist must be included in the authority application.

Authority required

Constitutional delay of growth or puberty

Population criteria:

- Patient must be under 18 years of age.

Treatment criteria:

- Must be treated by a specialist general paediatrician, specialist paediatric endocrinologist, specialist urologist, specialist endocrinologist or a Fellow of the Australasian Chapter of Sexual Health Medicine; or in consultation with one of these specialists; or have an appointment to be assessed by one of these specialists.

The name of the specialist must be included in the authority application.

testosterone undecanoate 1 g/4 mL modified release injection, 4 mL vial

10205D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	116.34	41.30	Reandron 1000 [BN]

testosterone undecanoate 40 mg capsule, 60

2115H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	33.99	35.28	Andriol Testocaps [MK]

ESTROGENS

Natural and semisynthetic estrogens, plain

■ ESTRADIOL

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

estradiol valerate 2 mg tablet, 56

1664N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	18.52	19.81	Prodynova [BN]

estradiol 2 mg tablet, 56

8274L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	18.22	19.51	Zumenon [GO]

estradiol valerate 1 mg tablet, 56

1663M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	16.68	17.97	Prodynova [BN]

estradiol 10 microgram modified release pessary, 18

10203B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	34.09	35.38	Vagifem Low [NO]

■ ESTRADIOL

Note Estradiol should be used in conjunction with an oral progestogen in women with an intact uterus.

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

estradiol 25 microgram/24 hours patch, 8

8311K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	21.30	22.59	Estraderm MX 25 [JU]

estradiol 25 microgram/24 hours patch, 8

8761D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	21.30	22.59	Estradot 25 [SZ]

estradiol 50 microgram/24 hours patch, 4

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8125P NP	±1	5	..	21.30	22.59	Climara 50 [BN]

estradiol 0.1% (1 mg/g) gel, 28 x 1 g sachets

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8286D NP	±1	5	..	21.30	22.59	Sandrena [AS]

estradiol 100 microgram/24 hours patch, 8

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8312L NP	±1	5	..	23.07	24.36	Estraderm MX 100 [JU]

estradiol 100 microgram/24 hours patch, 8

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8765H NP	±1	5	..	23.07	24.36	Estradot 100 [SZ]

estradiol 75 microgram/24 hours patch, 4

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8486P NP	±1	5	..	23.07	24.36	Climara 75 [BN]

estradiol 100 microgram/24 hours patch, 4

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8126Q NP	±1	5	..	23.07	24.36	Climara 100 [BN]

estradiol 75 microgram/24 hours patch, 8

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8764G NP	±1	5	..	23.07	24.36	Estradot 75 [SZ]

estradiol 25 microgram/24 hours patch, 4

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8485N NP	±1	5	..	21.30	22.59	Climara 25 [BN]

estradiol 50 microgram/24 hours patch, 8

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8140K NP	±1	5	..	21.30	22.59	Estraderm MX 50 [JU]

estradiol 50 microgram/24 hours patch, 8

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8763F NP	±1	5	..	21.30	22.59	Estradot 50 [SZ]

estradiol 37.5 microgram/24 hours patch, 8

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8762E NP	±1	5	..	21.30	22.59	Estradot 37.5 [SZ]

▪ ESTRIO**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

estriol 500 microgram pessary, 15

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
1771F NP	±1	2	..	24.83	26.12	Ovestin Ovula [AS]

estriol 0.1% (1 mg/g) cream, 15 g

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
1781R NP	±1	1	..	23.34	24.63	Ovestin [AS]

PROGESTOGENS*Pregnen (4) derivatives***▪ MEDROXYPROGESTERONE****Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

medroxyprogesterone acetate 5 mg tablet, 56

2323G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	20.02	21.31	^a Ralovera [FZ]
			^B 6.70	26.72	21.31	^a Provera [PF]

medroxyprogesterone acetate 10 mg tablet, 30

2321E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	18.89	20.18	^a Ralovera [FZ]
			^B 6.70	25.59	20.18	^a Provera [PF]

■ MEDROXYPROGESTERONE**Restricted benefit**

Endometriosis

medroxyprogesterone acetate 10 mg tablet, 100

2722G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	34.92	36.21	^a Ralovera [FZ]
			^B 6.70	41.62	36.21	^a Provera [PF]

*Estren derivatives***■ NORETHISTERONE****Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

norethisterone 5 mg tablet, 30

2993M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	34.23	35.52	Primolut N [BN]

PROGESTOGENS AND ESTROGENS IN COMBINATION*Progestogens and estrogens, fixed combinations***■ ESTRADIOL + DYDROGESTERONE****Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

estradiol 1 mg + dydrogesterone 5 mg tablet, 28

10142T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	22.75	24.04	Femoston-Conti [GO]

■ ESTRADIOL + NORETHISTERONE ACETATE**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

estradiol 50 microgram/24 hours + norethisterone acetate 250 microgram/24 hours patch, 8

8428N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	23.07	24.36	Estalis continuous 50/250 [SZ]

estradiol 50 microgram/24 hours + norethisterone acetate 140 microgram/24 hours patch, 8

8427M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	23.07	24.36	Estalis continuous 50/140 [SZ]

*Progestogens and estrogens, sequential preparations***■ ESTRADIOL (&) ESTRADIOL + DYDROGESTERONE****Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

estradiol 1 mg tablet [14] (&) estradiol 1 mg + dydrogesterone 10 mg tablet [14], 28

10146B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	22.75	24.04	Femoston 1/10 [GO]

estradiol 2 mg tablet [14] (&) estradiol 2 mg + dydrogesterone 10 mg tablet [14], 28

8244X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	22.75	24.04	Femoston 2/10 [GO]

▪ **NORETHISTERONE ACETATE + ESTRADIOL (&) ESTRADIOL**

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

estradiol 50 microgram/24 hours patch [4] (&) estradiol 50 microgram/24 hours + norethisterone acetate 140 microgram/24 hours patch [4], 8

8425K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	23.07	24.36	Estalis sequi 50/140 [SZ]

estradiol 50 microgram/24 hours patch [4] (&) estradiol 50 microgram/24 hours + norethisterone acetate 250 microgram/24 hours patch [4], 8

8426L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	23.07	24.36	Estalis sequi 50/250 [SZ]

GONADOTROPINS AND OTHER OVULATION STIMULANTS

Gonadotropins

▪ **FOLLITROPIN ALFA**

Note Biosimilar prescribing policy Prescribing of the biosimilar brand, Bemfola, is encouraged for treatment naive patients.

Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note Except in cases of hypopituitarism or primary amenorrhoea, the patient should have been adequately treated with clomifene citrate and/or gonadorelin and failed to have conceived.

Note Patients with hyperprolactinaemia should have had appropriate surgical or medical treatment prior to treatment.

Restricted benefit

Anovulatory infertility

Note Women who have had apparent ovulation induced by other agents and have failed to conceive should have laparoscopic evidence that there is no other impediment to conception.

Note Oligomenorrhoea should have been present for at least twelve months or amenorrhoea for at least six months prior to treatment.

Restricted benefit

Infertility

Clinical criteria:

- The condition must be due to hypogonadotrophic hypogonadism, **AND**
- The treatment must be following failure of 6 months' treatment with human chorionic gonadotrophin to achieve adequate spermatogenesis, **AND**
- The treatment must be administered with human chorionic gonadotrophin.

follitropin alfa 150 units (11 microgram)/0.25 mL injection, 5 x 0.25 mL pen devices

10877L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	1	..	*923.52	41.30	Bemfola [FX]

follitropin alfa 450 units (32.76 microgram)/0.75 mL injection, 0.75 mL pen device

8714P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	5	..	*561.57	41.30	Gonal-f Pen [SG]

follitropin alfa 300 units (21.84 microgram)/0.5 mL injection, 0.5 mL pen device

8713N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	5	..	*376.74	41.30	Gonal-f Pen [SG]

follitropin alfa 75 units (5.5 microgram)/0.125 mL injection, 5 x 0.125 mL pen devices

10865W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	1	..	*469.17	41.30	Bemfola [FX]

follitropin alfa 900 units (65.52 microgram)/1.5 mL injection, 1.5 mL pen device

8715Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*746.46	41.30	Gonal-f Pen [SG]

follitropin alfa 225 units (16.5 microgram)/0.375 mL injection, 5 x 0.375 mL pen devices

10876K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	1	..	*1353.33	41.30	Bemfola [FX]

▪ FOLLITROPIN BETA

Note Except in cases of hypopituitarism or primary amenorrhoea, the patient should have been adequately treated with clomifene citrate and/or gonadorelin and failed to have conceived.

Note Patients with hyperprolactinaemia should have had appropriate surgical or medical treatment prior to treatment.

Restricted benefit

Anovulatory infertility

Note Women who have had apparent ovulation induced by other agents and have failed to conceive should have laparoscopic evidence that there is no other impediment to conception.

Note Oligomenorrhoea should have been present for at least twelve months or amenorrhoea for at least six months prior to treatment.

Restricted benefit

Infertility

Clinical criteria:

- The condition must be due to hypogonadotrophic hypogonadism, **AND**
- The treatment must be following failure of 6 months' treatment with human chorionic gonadotrophin to achieve adequate spermatogenesis, **AND**
- The treatment must be administered with human chorionic gonadotrophin.

follitropin beta 600 units/0.72 mL injection, 0.72 mL cartridge

8566W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*543.86	41.30	Puregon 600 IU/0.72 mL [OQ]

follitropin beta 900 units/1.08 mL injection, 1.08 mL cartridge

8871X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*805.08	41.30	Puregon 900 IU/1.08 mL [OQ]

follitropin beta 300 units/0.36 mL injection, 0.36 mL cartridge

8565T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	5	..	*419.67	41.30	Puregon 300 IU/0.36 mL [OQ]

▪ HUMAN CHORIONIC GONADOTROPHIN

Note Patients with hyperprolactinaemia should have had appropriate surgical or medical treatment prior to treatment.

Restricted benefit

Anovulatory infertility

Note Except in cases of hypopituitarism or primary amenorrhoea, the patient should have been adequately treated with clomifene citrate and/or gonadorelin and failed to have conceived.

Note Women who have had apparent ovulation induced by other agents and have failed to conceive should have laparoscopic evidence that there is no other impediment to conception.

Note Oligomenorrhoea should have been present for at least twelve months or amenorrhoea for at least six months prior to treatment.

Restricted benefit

Infertility

Population criteria:

- Patient must be male.

Clinical criteria:

- The condition must be due to hypogonadotrophic hypogonadism.

Restricted benefit

Infertility

Population criteria:

- Patient must be male.

Clinical criteria:

- The condition must be associated with isolated luteinising hormone deficiency.

Restricted benefit

Combined deficiency of human growth hormone and gonadotrophins

Population criteria:

- Patient must be male.

Clinical criteria:

- Patient must be one in whom the absence of secondary sexual characteristics indicates a lag in maturation.

Restricted benefit

Hypogonadism or delayed puberty

Population criteria:

- Patient must be male, **AND**

- Patient must be aged 16 years or older.
- Clinical criteria:**
- Patient must show clinical evidence of the condition, **AND**
- The treatment must not extend beyond 6 months.

human chorionic gonadotrophin 1500 units injection [3 vials] (&) inert substance diluent [3 x 1 mL vials], 1 pack

11148R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	45.97	41.30	Pregnyl [OQ]

*Ovulation stimulants, synthetic***■ CLOMIFENE**

Note Care must be taken to comply with the provisions of State/Territory law when prescribing this drug.

Restricted benefit

Anovulatory infertility

Restricted benefit

Patients undergoing in-vitro fertilisation

clomifene citrate 50 mg tablet, 10

1211R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	36.45	37.74	Clomid [SW]

ANTIANDROGENS*Antiandrogens, plain***■ CYPROTERONE****cyproterone acetate 100 mg tablet, 50**

8019C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	67.80	41.30	^a ANTERONE 100 [RW] ^a Cyprone 100 [AF] ^a Cyproterone AN [EA] ^a GenRx Cyproterone Acetate [GX]	^a APO-Cyproterone [TX] ^a Cyprostat-100 [SY] ^a Cyproterone Sandoz [HX] ^a Pharmacor Cyproterone 100 [CR]
			^B 1.41	69.21	41.30	^a Androcur-100 [BN]	

cyproterone acetate 50 mg tablet, 50

1270W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	5	..	*83.62	41.30	^a ANTERONE 50 [RW] ^a Cyprone 50 [AL] ^a Cyproterone Sandoz [HX] ^a Pharmacor Cyproterone 50 [CR]	^a APO-Cyproterone [TX] ^a Cyproterone AN [EA] ^a GenRx Cyproterone Acetate [GX]
			^B 2.28	*85.90	41.30	^a Androcur [BN]	

■ CYPROTERONE

Caution This drug should not be used during pregnancy as it may result in feminisation of the male foetus.

Authority required (STREAMLINED)

5532

Moderate to severe androgenisation

Clinical criteria:

- The condition must not be indicated by acne alone, as this is not a sufficient indication of androgenisation.

Population criteria:

- Patient must be female.

Clinical criteria:

- Patient must not be pregnant.

cyproterone acetate 50 mg tablet, 20

1269T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	26.34	27.63	^a ANTERONE 50 [RW] ^a Cyprone 50 [AL] ^a Cyproterone Sandoz [HX] ^a Pharmacor Cyproterone 50 [CR]	^a APO-Cyproterone [TX] ^a Cyproterone AN [EA] ^a GenRx Cyproterone Acetate [GX]
			^B 2.40	28.74	27.63	^a Androcur [BN]	

OTHER SEX HORMONES AND MODULATORS OF THE GENITAL SYSTEM*Progesterone receptor modulators*

MIFEPRISTONE (&) MISOPROSTOL

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Termination of an intra-uterine pregnancy

Clinical criteria:

- The condition must be an intra-uterine pregnancy of up to 63 days of gestation.

Treatment criteria:

- Must be treated by a prescriber who is registered with the MS 2 Step Prescribing Program.

mifepristone 200 mg tablet [1] (&) misoprostol 200 microgram tablet [4], 1 pack

10211K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	318.47	41.30	MS-2 Step [XH]

UROLOGICALS

UROLOGICALS

Drugs for urinary frequency and incontinence

OXYBUTYNIN

Restricted benefit

Detrusor overactivity

oxybutynin hydrochloride 5 mg tablet, 100

8039D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	15.97	17.26	^a Ditropan [SW]	^a Oxybutynin Sandoz [SZ]

OXYBUTYNIN

Restricted benefit

Detrusor overactivity

Clinical criteria:

- Patient must be unable to tolerate oral oxybutynin; OR
- Patient must be unable to swallow oral oxybutynin.

oxybutynin 3.9 mg/24 hours patch, 8

9454N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	5	..	37.07	38.36	Oxytrol [TT]

PROPANTHELINE

Restricted benefit

Detrusor overactivity

propantheline bromide 15 mg tablet, 100

1953T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*26.16	27.45	Pro-Banthine [RW]

Other urologicals

BICARBONATE

sodium bicarbonate 840 mg capsule, 100

9470K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	20.21	21.50	Sodibic [AS]

PHENOXYBENZAMINE

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Phaeochromocytoma

Restricted benefit

Neurogenic urinary retention

phenoxybenzamine hydrochloride 10 mg capsule, 100

1862B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	953.89	41.30	Dibenylene [GH]

phenoxybenzamine hydrochloride 10 mg capsule, 30

1166J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	3	5	..	*864.90	41.30	Amdipharm Mercury (Australia) Pty Limited [GH]

DRUGS USED IN BENIGN PROSTATIC HYPERTROPHY*Alpha-adrenoreceptor antagonists***■ DUTASTERIDE + TAMSULOSIN****Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**6189**

Benign prostatic hyperplasia

Clinical criteria:

- Patient must have lower urinary tract symptoms, **AND**
- Patient must have moderate to severe benign prostatic hyperplasia.

dutasteride 500 microgram + tamsulosin hydrochloride 400 microgram modified release capsule, 30

5490Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	31.69	32.98	^a Doubluts [GC]
			^B 3.50	35.19	32.98	^a Duodart 500ug/400ug [GK]

*Testosterone-5-alpha reductase inhibitors***■ DUTASTERIDE****Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**6202**

Benign prostatic hyperplasia

Clinical criteria:

- Patient must have lower urinary tract symptoms, **AND**
- Patient must have moderate to severe benign prostatic hyperplasia, **AND**
- The treatment must be in combination with an alpha-antagonist.

dutasteride 500 microgram capsule, 30

5468T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	28.69	29.98	^a APO-Dutasteride [TX]
			^B 7.00	35.69	29.98	^a Avodart [GK]

■ SYSTEMIC HORMONAL PREPARATIONS, EXCL. SEX HORMONES AND INSULINS**■ PITUITARY AND HYPOTHALAMIC HORMONES AND ANALOGUES****ANTERIOR PITUITARY LOBE HORMONES AND ANALOGUES***ACTH***■ TETRACOSACTIDE (TETRACOSACTRIN)****Restricted benefit**

Hypsarrhythmia and/or infantile spasms

tetracosactide (tetracosactrin) 1 mg/mL modified release injection, 1 mL ampoule

2832C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	5	5	..	*393.09	41.30	Synacthen Depot 1 mg/1 mL [IX]

*Thyrotropin***■ THYROTROPIN ALFA****Restricted benefit**

Ablation of thyroid remnant tissue

Clinical criteria:

- Patient must have undergone a thyroidectomy, **AND**
- The treatment must be in combination with radioactive iodine, **AND**
- Patient must not have a known metastatic disease.

thyrotropin alfa 900 microgram injection, 2 vials

2700D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	1639.48	41.30	Thyrogen [GZ]

POSTERIOR PITUITARY LOBE HORMONES*Vasopressin and analogues***■ DESMOPRESSIN****Authority required (STREAMLINED)****5266**

Cranial diabetes insipidus

desmopressin acetate 100 microgram/mL nasal drops, 2.5 mL

2129C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	5	5	..	*127.04	41.30	Minirin [FP]

desmopressin acetate 200 microgram tablet, 30

8662X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	5	..	*141.54	41.30	Minirin [FP]

■ DESMOPRESSIN

Note Pharmaceutical benefits that have the form desmopressin acetate 10 microgram/actuation nasal spray, 50 actuations can be substituted for desmopressin acetate 10 microgram/actuation nasal spray, 60 actuations in the case of a shortage.

Authority required (STREAMLINED)**5266**

Cranial diabetes insipidus

desmopressin acetate 10 microgram/actuation nasal spray, 50 actuations

12288R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*193.90	41.30	^a Desmopressin Acetate (Medsurge) [DZ]

desmopressin acetate 10 microgram/actuation nasal spray, 60 actuations

8711L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*126.88	41.30	^a Minirin Nasal Spray [FP]

■ DESMOPRESSIN

Note Not to be used in preference to enuresis alarms.

Note Only one application per six months with no more than twice the maximum quantity will be authorised for the tablets.

Authority required (STREAMLINED)**5413**

Primary nocturnal enuresis

Population criteria:

- Patient must be 6 years of age or older.

Clinical criteria:

- Patient must be refractory to an enuresis alarm.

Authority required (STREAMLINED)**5295**

Primary nocturnal enuresis

Population criteria:

- Patient must be 6 years of age or older.

Clinical criteria:

- Patient must be one in whom an enuresis alarm is contraindicated.

The reason that an enuresis alarm is contraindicated must be documented in the patient's medical records when treatment is initiated

desmopressin acetate 200 microgram tablet, 30

8663Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	54.72	41.30	Minirin [FP]

NP

■ DESMOPRESSIN

Note Not to be used in preference to enuresis alarms.

Note Only one application per six months will be authorised for the wafers. No more than twice the maximum quantity for the 120 micrograms wafers and no applications for increased maximum quantities for the 240 micrograms wafers will be authorised.

Authority required (STREAMLINED)**5412**

Primary nocturnal enuresis

Population criteria:

- Patient must be 6 years of age or older.

Clinical criteria:

- Patient must be refractory to an enuresis alarm.

Authority required (STREAMLINED)**5226**

Primary nocturnal enuresis

Population criteria:

- Patient must be 6 years of age or older.

Clinical criteria:

- Patient must be one in whom an enuresis alarm is contraindicated.

The reason that an enuresis alarm is contraindicated must be documented in the patient's medical records when treatment is initiated

desmopressin 240 microgram sublingual wafer, 30

8975J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	92.89	41.30	Minirin Melt [FP]

desmopressin 120 microgram sublingual wafer, 30

9398P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	59.60	41.30	Minirin Melt [FP]

■ DESMOPRESSIN

Caution Desmopressin nasal spray may be associated with an increased risk of hyponatraemia compared to the oral formulations.

Note Pharmaceutical benefits that have the form desmopressin acetate 10 microgram/actuation nasal spray, 50 actuations can be substituted for desmopressin acetate 10 microgram/actuation nasal spray, 60 actuations in the case of a shortage.

Note Not to be used in preference to enuresis alarms.

Authority required (STREAMLINED)**5342**

Primary nocturnal enuresis

Population criteria:

- Patient must be 6 years of age or older.

Clinical criteria:

- Patient must be refractory to an enuresis alarm.

Authority required (STREAMLINED)**5267**

Primary nocturnal enuresis

Population criteria:

- Patient must be 6 years of age or older.

Clinical criteria:

- Patient must be one in whom an enuresis alarm is contraindicated.

The reason that an enuresis alarm is contraindicated must be documented in the patient's medical records when treatment is initiated

desmopressin acetate 10 microgram/actuation nasal spray, 50 actuations

12280H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	101.01	41.30	^a Desmopressin Acetate (Medsurge) [DZ]

desmopressin acetate 10 microgram/actuation nasal spray, 60 actuations

8712M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	69.10	41.30	^a Minirin Nasal Spray [FP]

HYPOTHALAMIC HORMONES*Gonadotropin-releasing hormones***■ NAFARELIN****Restricted benefit**

Endometriosis

Treatment Phase: Initial treatment, for up to 6 months

Clinical criteria:

- The condition must be visually proven.

Restricted benefit

Endometriosis

Treatment Phase: Subsequent treatment, for up to 6 months

Clinical criteria:

- The condition must be visually proven, **AND**
- The treatment must not be within 2 years of the end of the previous course of treatment with this drug, **AND**
- Patient must have had a recent bone density assessment.

SYSTEMIC HORMONAL PREPARATIONS, EXCL. SEX HORMONES AND INSULINS

The date of the bone density assessment must be recorded in the patient's medical records.

nafarelin 200 microgram/actuation nasal spray, 60 actuations

2962X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	‡1	5	..	122.86	41.30	Synarel [PF]

■ CORTICOSTEROIDS FOR SYSTEMIC USE

CORTICOSTEROIDS FOR SYSTEMIC USE, PLAIN

Mineralocorticoids

■ FLUDROCORTISONE ACETATE

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

fludrocortisone acetate 100 microgram tablet, 100

1433K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	1	..	*40.92	41.30	Florinef [AS]

NP

Glucocorticoids

■ BETAMETHASONE ACETATE + BETAMETHASONE SODIUM PHOSPHATE

Restricted benefit

Local intra-articular or peri-articular infiltration

Restricted benefit

Keloid

Restricted benefit

Lichen planus hypertrophic

betamethasone acetate 3 mg/mL + betamethasone sodium phosphate 3.9 mg/mL (total betamethasone 5.7 mg/mL) injection, 5 x 1 mL ampoules

5034Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	28.18	29.47	Celestone Chronodose [OQ]

DP

■ BETAMETHASONE ACETATE + BETAMETHASONE SODIUM PHOSPHATE

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Alopecia areata

Restricted benefit

Local intra-articular or peri-articular infiltration

Restricted benefit

Granulomata

Clinical criteria:

- The condition must be dermal.

Restricted benefit

Keloid

Restricted benefit

Lichen planus hypertrophic

Restricted benefit

Lichen simplex chronicus

Restricted benefit

Chronic discoid lupus erythematosus

Restricted benefit

Necrobiosis lipoidica

Restricted benefit

Uveitis

betamethasone acetate 3 mg/mL + betamethasone sodium phosphate 3.9 mg/mL (total betamethasone 5.7 mg/mL) injection, 5 x 1 mL ampoules

2694T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	28.18	29.47	Celestone Chronodose [OQ]

NP

■ CORTISONE

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

cortisone acetate 5 mg tablet, 50

1246N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	4	..	19.35	20.64	Cortate [AS]

cortisone acetate 25 mg tablet, 60

1247P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	4	..	24.48	25.77	Cortate [AS]

■ DEXAMETHASONE

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

dexamethasone 4 mg tablet, 30

2507Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	4	..	17.27	18.56	Dexmethsone [AS]

dexamethasone 500 microgram tablet, 30

1292B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	4	..	14.38	15.67	Dexmethsone [AS]

■ HYDROCORTISONE

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

hydrocortisone 4 mg tablet, 50

1499X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	4	..	26.48	27.77	^a Hydrocortisone Mylan 4 [AL]
			^B 3.00	29.48	27.77	^a Hysone 4 [AF]

hydrocortisone 20 mg tablet, 60

1500Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	4	..	32.66	33.95	^a Hydrocortisone Mylan 20 [AL]
			^B 3.00	35.66	33.95	^a Hysone 20 [AF]

■ HYDROCORTISONE SODIUM SUCCINATE

hydrocortisone (as sodium succinate) 250 mg injection [1 chamber] (&) inert substance diluent [2 mL chamber], 1 dual chamber vial

3096Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	20.78	22.07	Solu-Cortef [PF]

hydrocortisone (as sodium succinate) 100 mg injection [1 chamber] (&) inert substance diluent [2 mL chamber], 1 dual chamber vial

1501B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	..	..	*21.82	23.11	Solu-Cortef [PF]

■ HYDROCORTISONE SODIUM SUCCINATE

Restricted benefit

For use in a hospital

hydrocortisone (as sodium succinate) 250 mg injection [1 chamber] (&) inert substance diluent [2 mL chamber], 1 dual chamber vial

1511M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	6	..	..	*64.56	41.30	Solu-Cortef [PF]

hydrocortisone (as sodium succinate) 250 mg injection [1 chamber] (&) inert substance diluent [2 mL chamber], 1 dual chamber vial

5119K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	6	..	..	*64.56	41.30	Solu-Cortef [PF]

SYSTEMIC HORMONAL PREPARATIONS, EXCL. SEX HORMONES AND INSULINS

hydrocortisone (as sodium succinate) 100 mg injection [1 chamber] (&) inert substance diluent [2 mL chamber], 1 dual chamber vial

1510L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	6	..	..	*41.40	41.30	Solu-Cortef [PF]

hydrocortisone (as sodium succinate) 100 mg injection [1 chamber] (&) inert substance diluent [2 mL chamber], 1 dual chamber vial

5118J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	6	..	..	*41.40	41.30	Solu-Cortef [PF]

■ METHYLPREDNISOLONE

methylprednisolone 1 g injection, 1 vial

5264C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	44.52	41.30	^a Methylpred [AL]	^a Methylprednisolone Alphapharm [AF]
						^a Solu-Medrol [PF]	

■ METHYLPREDNISOLONE

Restricted benefit

Local intra-articular or peri-articular infiltration

methylprednisolone acetate 40 mg/mL injection, 5 x 1 mL vials

1928L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	23.40	24.69	^a Depo-Nisolone [FZ]
			^B 2.61	26.01	24.69	^a Depo-Medrol [PF]

■ METHYLPREDNISOLONE

Restricted benefit

Local intra-articular or peri-articular infiltration

methylprednisolone acetate 40 mg/mL injection, 5 x 1 mL vials

5148Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	1	..	..	23.40	24.69	^a Depo-Nisolone [FZ]
			^B 2.61	26.01	24.69	^a Depo-Medrol [PF]

■ METHYLPREDNISOLONE

Note Pharmaceutical benefits that have the form methylprednisolone powder for injection 40 mg (as sodium succinate) and pharmaceutical benefits that have the form methylprednisolone powder for injection 40 mg (as sodium succinate) with diluent are equivalent for the purposes of substitution.

methylprednisolone 40 mg injection [1 chamber] (&) inert substance diluent [1 mL chamber], 1 dual chamber vial

11739W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	5	..	..	*41.64	41.30	^a Solu-Medrol [PF]

methylprednisolone 40 mg injection, 5 vials

5263B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	24.16	25.45	^a Methylpred [AL]

■ PREDNISOLONE

prednisolone 25 mg tablet, 30

1916W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	4	..	16.41	17.70	Panafcortelone [AS]	Solone [IL]

prednisolone 5 mg tablet, 60

1917X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	4	..	15.22	16.51	Panafcortelone [AS]	Solone [IL]

prednisolone 1 mg tablet, 100

3152X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	4	..	14.81	16.10	^a Predsolone [LN]
			^B 0.93	15.74	16.10	^a Panafcortelone [AS]

■ PREDNISOLONE SODIUM PHOSPHATE

prednisolone (as sodium phosphate) 5 mg/mL oral liquid, 30 mL

8285C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	‡1	5	..	19.22	20.51	^a PredMix [LN]
			^B 2.35	21.57	20.51	^a Redipred [AS]

■ PREDNISONE

prednisone 5 mg tablet, 60

1935W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	4	..	15.14	16.43	Panafcort [AS]	Sone [IL]

prednisone 1 mg tablet, 100

1934T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	4	..	14.76	16.05	^a Predsone [LN]	
			^b 0.93	15.69	16.05	^a Panafcort [AS]	

prednisone 25 mg tablet, 30

1936X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	4	..	16.47	17.76	Panafcort [AS]	Sone [IL]

■ TRIAMCINOLONE

Restricted benefit

Local intra-articular or peri-articular infiltration

Restricted benefit

Keloid

Restricted benefit

Lichen planus hypertrophic

triamcinolone acetonide 10 mg/mL injection, 5 x 1 mL ampoules

5233K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
DP	1	..	..	28.18	29.47	Kenacort-A10 [AS]	

■ TRIAMCINOLONE

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Alopecia areata

Restricted benefit

Local intra-articular or peri-articular infiltration

Restricted benefit

Granulomata

Clinical criteria:

- The condition must be dermal.

Restricted benefit

Keloid

Restricted benefit

Lichen planus hypertrophic

Restricted benefit

Lichen simplex chronicus

Restricted benefit

Chronic discoid lupus erythematosus

Restricted benefit

Necrobiosis lipoidica

Restricted benefit

Psoriasis

triamcinolone acetonide 10 mg/mL injection, 5 x 1 mL ampoules

2990J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	28.18	29.47	Kenacort-A10 [AS]	

■ THYROID THERAPY

THYROID PREPARATIONS

Thyroid hormones

■ LEVOTHYROXINE

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

levothyroxine sodium 200 microgram tablet, 200

2173J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	29.92	31.21	^a Eutroxsig [LN]
			^B 1.93	31.85	31.21	^a Oroxine [AS]

levothyroxine sodium 100 microgram tablet, 200

2175L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	27.29	28.58	^a Eutroxsig [LN]
			^B 1.92	29.21	28.58	^a Oroxine [AS]

levothyroxine sodium 50 microgram tablet, 200

2174K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	26.76	28.05	^a Eutroxsig [LN]
			^B 1.91	28.67	28.05	^a Oroxine [AS]

levothyroxine sodium 75 microgram tablet, 200

9287T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	27.32	28.61	^a Eutroxsig [LN]
			^B 1.98	29.30	28.61	^a Oroxine [AS]

■ LIOTHYRONINE**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**6382**

Thyroid cancer

Authority required (STREAMLINED)**6410**

Hypothyroidism

Clinical criteria:

- The treatment must be for replacement therapy, **AND**
- Patient must have documented intolerance to levothyroxine sodium; OR
- Patient must have documented resistance to levothyroxine sodium.

Authority required (STREAMLINED)**6475**

Hypothyroidism

Clinical criteria:

- The condition must be severe hypothyroidism, **AND**
- The treatment must be for initiation of therapy only.

liothyronine sodium 20 microgram tablet, 100

2318B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	68.96	41.30	Tertroxin [AS]

ANTITHYROID PREPARATIONS*Thiouracils***■ PROPYLTHIOURACIL****Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

propylthiouracil 50 mg tablet, 100

1955X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	2	..	*43.48	41.30	PTU [FF]

*Sulfur-containing imidazole derivatives***■ CARBIMAZOLE****Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

carbimazole 5 mg tablet, 100

1153Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	2	..	*39.60	40.89	Neo-Mercazole [GH]

■ PANCREATIC HORMONES

GLYCOGENOLYTIC HORMONES

Glycogenolytic hormones

■ GLUCAGON HYDROCHLORIDE

glucagon hydrochloride 1 mg injection [1 vial] (&) inert substance diluent [1 mL syringe], 1 pack

1449G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	49.66	41.30	GlucaGen Hypokit [NO]

glucagon hydrochloride 1 mg injection [1 vial] (&) inert substance diluent [1 mL syringe], 1 pack

5105Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	1	..	..	49.66	41.30	GlucaGen Hypokit [NO]

■ CALCIUM HOMEOSTASIS

PARATHYROID HORMONES AND ANALOGUES

Parathyroid hormones and analogues

■ TERIPARATIDE

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Authority required

Severe established osteoporosis

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must be at very high risk of fracture, **AND**
- Patient must have a bone mineral density (BMD) T-score of -3.0 or less, **AND**
- Patient must have had 2 or more fractures due to minimal trauma, **AND**
- Patient must have experienced at least 1 symptomatic new fracture after at least 12 months continuous therapy with an anti-resorptive agent at adequate doses, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- The treatment must not exceed a lifetime maximum of 18 months therapy, **AND**
- Patient must not have received treatment with romosozumab; OR
- Patient must have developed intolerance to romosozumab of a severity necessitating permanent treatment withdrawal within the first 6 months of therapy.

Treatment criteria:

- Must be treated by a specialist; OR
- Must be treated by a consultant physician.

A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or, a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

If treatment with anti-resorptive therapy is contraindicated according to the relevant TGA-approved Product Information, details of the contraindication must be documented in the patient's medical record at the time treatment with this drug is initiated.

If an intolerance of a severity necessitating permanent treatment withdrawal develops during the relevant period of use of one anti-resorptive agent, alternate anti-resorptive agents must be trialled so that the patient achieves the minimum requirement of 12 months continuous therapy. Details must be documented in the patient's medical record at the time treatment with this drug is initiated.

Anti-resorptive therapies for osteoporosis and their adequate doses which will be accepted for the purposes of administering this restriction are alendronate sodium 10 mg per day or 70 mg once weekly, risedronate sodium 5 mg per day or 35 mg once weekly or 150 mg once monthly, raloxifene hydrochloride 60 mg per day (women only), denosumab 60 mg once every 6 months and zoledronic acid 5 mg per annum.

Details of prior anti-resorptive therapy, fracture history including the date(s), site(s), the symptoms associated with the fracture(s) which developed after at least 12 months continuous anti-resorptive therapy and the score of the qualifying BMD measurement must be provided at the time of application.

Authority required

Severe established osteoporosis

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must not exceed a lifetime maximum of 18 months therapy.

Treatment criteria:

- Must be treated by a specialist; OR

- Must be treated by a consultant physician.

Note Up to a maximum of 18 pens will be reimbursed through the PBS.

teriparatide 250 microgram/mL injection, 2.4 mL pen device

9411H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	378.65	41.30	Forteo [LY]

ANTI-PARATHYROID AGENTS

Calcitonin preparations

■ CALCITONIN SALMON (SALCATONIN)

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Symptomatic Paget disease of bone

Restricted benefit

Hypercalcaemia

Clinical criteria:

- The treatment must be initiated in a hospital.

calcitonin salmon (salcatonin) 100 units/mL injection, 5 x 1 mL ampoules

2997R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	5	..	*126.93	41.30	Miacalcic 100 [EU]

NP

Other anti-parathyroid agents

■ CINACALCET

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)

10068

Secondary hyperparathyroidism

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have chronic kidney disease, **AND**
- Patient must be on dialysis, **AND**
- Patient must have achieved a decrease of at least 30% in intact parathyroid hormone (iPTH) concentrations after 6 months treatment; OR
- Patient must have an intact parathyroid (iPTH) concentration greater than 15 pmol/L and an (adjusted) serum calcium concentration of less than 2.6 mmol/L after 6 months.

During the maintenance phase, iPTH should be monitored quarterly (measured at least 12 hours post dose) and dose adjusted as necessary to maintain an appropriate iPTH concentration.

During the maintenance phase, prescribers should request approval to allow sufficient supply for 4 weeks treatment up to a maximum of 6 months supply, with doses between 30 and 180 mg per day according to the patient's response and tolerability.

cinacalcet 30 mg tablet, 28

9157Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	46.77	41.30	^a Cinacalcet Mylan [AF]	^a Pharmacor Cinacalcet [CR]

NP

cinacalcet 90 mg tablet, 28

9159C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	116.48	41.30	^a Cinacalcet Mylan [AF]	^a Pharmacor Cinacalcet [CR]

NP

cinacalcet 60 mg tablet, 28

9158B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	81.52	41.30	^a Cinacalcet Mylan [AF]	^a Pharmacor Cinacalcet [CR]

NP

■ ANTIINFECTIVES FOR SYSTEMIC USE

■ ANTIBACTERIALS FOR SYSTEMIC USE

TETRACYCLINES

Tetracyclines

■ DOXYCYCLINE

Note Pharmaceutical benefits that have the forms doxycycline tablet 100 mg (as hyclate (hydrochloride)), doxycycline tablet 100 mg (as monohydrate) and doxycycline modified release capsule 100 mg (as hyclate (hydrochloride)) are equivalent for the purposes of substitution.

doxycycline 100 mg modified release capsule, 7

2708M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	^B 1.54	15.02	14.77	^a Mayne Pharma Doxycycline [YT]
			^B 2.96	16.44	14.77	^a Doryx [YN]

doxycycline 100 mg modified release capsule, 7

3322W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	1	..	^B 1.54	15.02	14.77	^a Mayne Pharma Doxycycline [YT]
			^B 2.96	16.44	14.77	^a Doryx [YN]

doxycycline 100 mg tablet, 7

2709N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	1	..	13.48	14.77	^a Doxsig [RW]	^a Doxycycline AN [EA]
						^a Doxylin 100 [AF]	

doxycycline 100 mg tablet, 7

3321T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
DP	1	..	..	13.48	14.77	^a Doxsig [RW]	^a Doxycycline AN [EA]
						^a Doxylin 100 [AF]	

doxycycline 100 mg tablet, 7

5082L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
DP	1	..	..	13.48	14.77	^a APO-Doxycycline [TX]	^a Doxycycline Sandoz [HX]

doxycycline 100 mg tablet, 7

9105F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	1	..	13.48	14.77	^a APO-Doxycycline [TX]	^a Doxycycline Sandoz [HX]

■ DOXYCYCLINE

Note Pharmaceutical benefits that have the forms doxycycline tablet 100 mg (as hyclate (hydrochloride)), doxycycline tablet 100 mg (as monohydrate) and doxycycline modified release capsule 100 mg (as hyclate (hydrochloride)) are equivalent for the purposes of substitution.

Restricted benefit

Urethritis

doxycycline 100 mg tablet, 21

10176N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	16.51	17.80	^a Doxsig [RW]	^a Doxycycline AN [EA]
						^a Doxylin 100 [AF]	

doxycycline 100 mg tablet, 21

1800R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	16.51	17.80	^a APO-Doxycycline [TX]

doxycycline 100 mg modified release capsule, 21

2715X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	^B 3.11	19.62	17.80	^a Mayne Pharma Doxycycline [YT]
			^B 8.90	25.41	17.80	^a Doryx [YN]

doxycycline 100 mg tablet, 7

2714W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	3	..	..	*16.41	17.70	^a Doxsig [RW]	^a Doxycycline AN [EA]
						^a Doxylin 100 [AF]	

doxycycline 100 mg tablet, 7

9108J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	3	..	..	*16.41	17.70	^a Doxycycline Sandoz [HX]

■ DOXYCYCLINE

Note Pharmaceutical benefits that have the forms doxycycline tablet 100 mg (as hyclate (hydrochloride)), doxycycline tablet 100 mg (as monohydrate) and doxycycline modified release capsule 100 mg (as hyclate (hydrochloride)) are equivalent for the purposes of substitution.

Restricted benefit

Severe acne

doxycycline 100 mg modified release capsule, 7

10777F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	4	5	^B 6.16	*24.02	19.15	^a Mayne Pharma Doxycycline [YT]
			^B 11.84	*29.70	19.15	^a Doryx [YN]

doxycycline 100 mg tablet, 7

10779H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	4	5	..	*17.86	19.15	^a Doxsig [RW]	^a Doxycycline AN [EA]
						^a Doxylin 100 [AF]	

doxycycline 100 mg tablet, 7

10781K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	4	5	..	*17.86	19.15	^a APO-Doxycycline [TX]	^a Doxycycline Sandoz [HX]

■ **DOXYCYCLINE**

Note Pharmaceutical benefits that have the forms doxycycline tablet 100 mg (as hyclate (hydrochloride)), doxycycline tablet 100 mg (as monohydrate) and doxycycline modified release capsule 100 mg (as hyclate (hydrochloride)) are equivalent for the purposes of substitution.

Restricted benefit

Pelvic inflammatory disease

doxycycline 100 mg modified release capsule, 7

2703G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	4	..	^B 6.16	*24.02	19.15	^a Mayne Pharma Doxycycline [YT]
			^B 11.84	*29.70	19.15	^a Doryx [YN]

doxycycline 100 mg tablet, 7

2702F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	4	..	..	*17.86	19.15	^a Doxsig [RW]	^a Doxycycline AN [EA]
						^a Doxylin 100 [AF]	

doxycycline 100 mg tablet, 7

9107H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	4	..	..	*17.86	19.15	^a APO-Doxycycline [TX]	^a Doxycycline Sandoz [HX]

■ **DOXYCYCLINE**

Note Pharmaceutical benefits that have the forms doxycycline tablet 50 mg (as hyclate (hydrochloride)), doxycycline tablet 50 mg (as monohydrate) and doxycycline modified release capsule 50 mg (as hyclate (hydrochloride)) are equivalent for the purposes of substitution.

Restricted benefit

Bronchiectasis

Population criteria:

- Patient must be aged 8 years or older.

Restricted benefit

Chronic bronchitis

Population criteria:

- Patient must be aged 8 years or older.

Restricted benefit

Severe acne

doxycycline 50 mg modified release capsule, 25

2707L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	^B 2.41	17.01	15.89	^a Mayne Pharma Doxycycline [YT]
			^B 4.76	19.36	15.89	^a Doryx [YN]

doxycycline 50 mg tablet, 25

2711Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	14.60	15.89	^a Doxsig [RW]	^a Doxycycline AN [EA]
						^a Doxylin 50 [AF]	

doxycycline 50 mg tablet, 25

9106G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	14.60	15.89	^a APO-Doxycycline [TX]	^a Doxycycline Sandoz [HX]
						^a Frakas [RW]	

■ **MINOCYCLINE**

Caution There are concerns about the incidence of benign intracranial hypertension associated with this drug.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Restricted benefit

Severe acne

Clinical criteria:

- The condition must not be responding to other tetracyclines.

minocycline 50 mg tablet, 60

1616C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	21.17	22.46	Minomycin-50 [AS]

BETA-LACTAM ANTIBACTERIALS, PENICILLINS

Penicillins with extended spectrum

AMOXICILLIN

amoxicillin 125 mg/5 mL powder for oral liquid, 100 mL

1886G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	‡1	1	..	#16.61	18.27	^a Alphamox 125 [AF] ^a NOUMED AMOXICILLIN [VO]	^a APO-Amoxycillin [TX]
			^B 3.22	#19.83	18.27	^a Amoxil [AS]	

amoxicillin 125 mg/5 mL powder for oral liquid, 100 mL

3302T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
DP	‡1	..	..	#16.61	18.27	^a Alphamox 125 [AF] ^a NOUMED AMOXICILLIN [VO]	^a APO-Amoxycillin [TX]
			^B 3.22	#19.83	18.27	^a Amoxil [AS]	

amoxicillin 250 mg capsule, 20

3301R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
DP	1	..	..	13.60	14.89	^a Alphamox 250 [AF] ^a Amoxycillin AN [EA] ^a APO-Amoxycillin [TX]	^a AMILOXYN [RF] ^a Amoxycillin Sandoz [SZ] ^a Cilamox [AL]
			^B 3.38	16.98	14.89	^a Amoxil [AS]	

amoxicillin 500 mg/5 mL powder for oral liquid, 100 mL

5225B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	‡1	..	..	#17.60	19.26	Maxamox [SZ]

amoxicillin 500 mg/5 mL powder for oral liquid, 100 mL

8705E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	‡1	1	..	#17.60	19.26	Maxamox [SZ]

amoxicillin 250 mg/5 mL powder for oral liquid, 100 mL

1887H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	‡1	1	..	#16.92	18.58	^a Alphamox 250 [AF] ^a APO-Amoxycillin [TX] ^a NOUMED AMOXICILLIN [VO]	^a Amoxycillin Sandoz [SZ] ^a Cilamox [AL]
			^B 3.31	#20.23	18.58	^a Amoxil Forte [AS]	

amoxicillin 250 mg/5 mL powder for oral liquid, 100 mL

3393N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
DP	‡1	..	..	#16.92	18.58	^a Alphamox 250 [AF] ^a APO-Amoxycillin [TX] ^a NOUMED AMOXICILLIN [VO]	^a Amoxycillin Sandoz [SZ] ^a Cilamox [AL]
			^B 3.31	#20.23	18.58	^a Amoxil Forte [AS]	

amoxicillin 100 mg/mL powder for oral liquid, 20 mL

1888J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	‡1	1	^S 0.53	#21.77	22.90	Amoxil [AS]

amoxicillin 100 mg/mL powder for oral liquid, 20 mL

3310F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	‡1	..	^S 0.53	#21.77	22.90	Amoxil [AS]

amoxicillin 500 mg capsule, 20

3300Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
DP	1	..	..	13.86	15.15	^a Alphamox 500 [AF] ^a Amoxycillin AN [EA] ^a Amoxycillin Sandoz [SZ]	^a AMILOXYN [RF] ^a Amoxycillin generichealth 500 [GQ] ^a APO-Amoxycillin [TX]

				^a Cilamox [AL]	^a NOUMED AMOXICILLIN [VO]
	^B 3.67	17.53	15.15	^a Amoxil [AS]	

■ AMOXICILLIN

Authority required (STREAMLINED)

10416

Community acquired pneumonia

Clinical criteria:

- Patient must have community acquired pneumonia.

amoxicillin 1 g tablet, 14

12002Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	13.90	15.19	^a Amoxycillin Sandoz [BG]
			^B 1.00	14.90	15.19	^a Maxamox [SZ]

■ AMOXICILLIN

Restricted benefit

Chronic bronchitis

Clinical criteria:

- Patient must have acute exacerbations of the condition.

amoxicillin 1 g tablet, 14

8581P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	13.90	15.19	^a Amoxycillin Sandoz [BG]
			^B 1.00	14.90	15.19	^a Maxamox [SZ]

■ AMOXICILLIN

Authority required

Infection suspected or proven to be due to a susceptible organism

Clinical criteria:

- The treatment must be for patients who require a liquid formulation and in whom the syrup formulations are unsuitable.

amoxicillin 100 mg/mL powder for oral liquid, 20 mL

9714G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	1	..	#21.77	23.43	Amoxil [AS]

■ AMOXICILLIN

Authority required (STREAMLINED)

10402

Infection

Clinical criteria:

- Patient must be a male with acute cystitis; OR
- Patient must have pyelonephritis; OR
- Patient must have a tooth avulsion; OR
- Patient must have salmonella enteritis; OR
- Patient must have community acquired pneumonia; OR
- Patient must have a condition requiring prolonged oral antibiotic therapy.

amoxicillin 500 mg capsule, 20

11947T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	2	..	..	*15.70	16.99	^a Alphamox 500 [AF]	^a AMILOXYN [RF]
						^a Amoxycillin AN [EA]	^a Amoxycillin generichealth 500 [GQ]
						^a Amoxycillin Sandoz [SZ]	^a APO-Amoxycillin [TX]
						^a Cilamox [AL]	^a NOUMED AMOXICILLIN [VO]
			^B 7.34	*23.04	16.99	^a Amoxil [AS]	

■ AMOXICILLIN

Authority required (STREAMLINED)

10404

Infection

Clinical criteria:

- Patient must have a condition requiring prolonged oral antibiotic therapy.

amoxicillin 250 mg capsule, 20

11998L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	2	..	..	*15.18	16.47	^a Alphamox 250 [AF]	^a AMILOXYN [RF]
						^a Amoxycillin AN [EA]	^a Amoxycillin Sandoz [SZ]
						^a APO-Amoxycillin [TX]	^a Cilamox [AL]
			^B 6.76	*21.94	16.47	^a Amoxil [AS]	

■ AMOXICILLIN

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

amoxicillin 250 mg capsule, 20

1884E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP MW	1	..	..	13.60	14.89	^a Alphamox 250 [AF] ^a Amoxycillin AN [EA] ^a APO-Amoxycillin [TX]	^a AMILOXYN [RF] ^a Amoxycillin Sandoz [SZ] ^a Cilamox [AL]
			^B 3.38	16.98	14.89	^a Amoxil [AS]	

amoxicillin 500 mg capsule, 20

1889K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP MW	1	..	..	13.86	15.15	^a Alphamox 500 [AF] ^a Amoxycillin AN [EA] ^a Amoxycillin Sandoz [SZ] ^a Cilamox [AL]	^a AMILOXYN [RF] ^a Amoxycillin generichealth 500 [GQ] ^a APO-Amoxycillin [TX] ^a NOUMED AMOXICILLIN [VO]
			^B 3.67	17.53	15.15	^a Amoxil [AS]	

■ AMPICILLIN

ampicillin 1 g injection, 5 vials

2977Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	18.16	19.45	Ampicyn [AF]

ampicillin 1 g injection, 5 vials

3314K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	1	..	..	18.16	19.45	Ampicyn [AF]

ampicillin 500 mg injection, 5 vials

2390T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	17.04	18.33	Austrapen [AL]

ampicillin 500 mg injection, 5 vials

3313J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	1	..	..	17.04	18.33	Austrapen [AL]

Beta-lactamase sensitive penicillins

■ BENZATHINE BENZYL PENICILLIN

benzathine benzylpenicillin tetrahydrate 600 000 units (517 mg)/1.17 mL injection, 10 x 1.17 mL syringes

11723B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	155.90	41.30	Bicillin L-A [PF]

benzathine benzylpenicillin tetrahydrate 600 000 units (517 mg)/1.17 mL injection, 10 x 1.17 mL syringes

11735P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	1	..	..	155.90	41.30	Bicillin L-A [PF]

benzathine benzylpenicillin tetrahydrate 1.2 million units (1016.6 mg)/2.3 mL injection, 10 x 2.3 mL syringes

2267H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	304.76	41.30	Bicillin L-A [PF]

benzathine benzylpenicillin tetrahydrate 1.2 million units (1016.6 mg)/2.3 mL injection, 10 x 2.3 mL syringes

5027N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	1	..	..	304.76	41.30	Bicillin L-A [PF]

■ BENZYL PENICILLIN

benzylpenicillin 600 mg injection, 1 vial

1775K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP MW	10	1	..	*71.44	41.30	BenPen [CS]

benzylpenicillin 600 mg injection, 1 vial

3398W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	10	..	..	*71.44	41.30	BenPen [CS]

ANTIINFECTIVES FOR SYSTEMIC USE

benzylpenicillin 3 g injection, 1 vial

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
2647H	10	..	..	*107.84	41.30	BenPen [CS]

benzylpenicillin 3 g injection, 1 vial

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
3399X	10	..	..	*107.84	41.30	BenPen [CS]

■ PHENOXYMETHYLPENICILLIN

phenoxymethylpenicillin 500 mg tablet, 25

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
3028J	2	..	..	*18.32	19.61	Aspecillin VK [AF]

phenoxymethylpenicillin 500 mg tablet, 25

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
3361X	2	..	..	*18.32	19.61	Aspecillin VK [AF]

phenoxymethylpenicillin 500 mg capsule, 50

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
2965C	1	..	..	18.15	19.44	Cilicaine VK [AF]	LPV [IL]

phenoxymethylpenicillin 500 mg capsule, 50

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
3364C	1	..	..	18.15	19.44	Cilicaine VK [AF]	LPV [IL]

phenoxymethylpenicillin 250 mg tablet, 25

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
1787C	2	..	..	*16.40	17.69	Aspecillin VK [AF]

phenoxymethylpenicillin 250 mg tablet, 25

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
3360W	2	..	..	*16.40	17.69	Aspecillin VK [AF]

phenoxymethylpenicillin 125 mg/5 mL powder for oral liquid, 100 mL

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
5024K	2	..	..	*#21.66	23.32	Phenoxymethylpenicillin-AFT [AE]

phenoxymethylpenicillin 125 mg/5 mL powder for oral liquid, 100 mL

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8976K	2	1	..	*#21.66	23.32	Phenoxymethylpenicillin-AFT [AE]

phenoxymethylpenicillin 150 mg/5 mL oral liquid, 100 mL

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
5012T	2	..	..	*25.22	26.51	Cilicaine V [AF]

phenoxymethylpenicillin 150 mg/5 mL oral liquid, 100 mL

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
9143F	2	1	..	*25.22	26.51	Cilicaine V [AF]

phenoxymethylpenicillin 250 mg capsule, 50

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
1789E	1	..	..	16.26	17.55	Cilicaine VK [AF]	LPV [IL]

phenoxymethylpenicillin 250 mg capsule, 50

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
3363B	1	..	..	16.26	17.55	Cilicaine VK [AF]	LPV [IL]

phenoxymethylpenicillin 250 mg/5 mL powder for oral liquid, 100 mL

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
5029Q	2	..	..	*#23.90	25.56	Phenoxymethylpenicillin-AFT [AE]

phenoxymethylpenicillin 250 mg/5 mL powder for oral liquid, 100 mL

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8977L	2	1	..	*#23.90	25.56	Phenoxymethylpenicillin-AFT [AE]

■ PHENOXYMETHYLPENICILLIN

Restricted benefit

Recurrent streptococcal infections (including rheumatic fever)

Clinical criteria:

- The treatment must be for prophylaxis.

phenoxymethylpenicillin 250 mg tablet, 25

1703P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*16.40	17.69	Aspecillin VK [AF]

phenoxymethylpenicillin 250 mg capsule, 50

1705R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	16.26	17.55	Cilicaine VK [AF]	LPV [IL]

■ PROCAINE BENZYL PENICILLIN (PROCAINE PENICILLIN)

procaine benzylpenicillin (procaine penicillin) 1.5 g/3.4 mL injection, 5 x 3.4 mL syringes

1794K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	75.37	41.30	Cilicaine [AF]

procaine benzylpenicillin (procaine penicillin) 1.5 g/3.4 mL injection, 5 x 3.4 mL syringes

3371K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	1	..	..	75.37	41.30	Cilicaine [AF]

Beta-lactamase resistant penicillins

■ DICLOXACILLIN

Restricted benefit

Serious staphylococcal infection

dicloxacillin 500 mg capsule, 24

5097G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	1	..	..	20.12	21.41	^a Dicloxacillin Mylan 500 [AL]
			^B 1.95	22.07	21.41	^a Distaph 500 [AF]

dicloxacillin 250 mg capsule, 24

5096F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	1	..	..	16.73	18.02	^a Dicloxacillin Mylan 250 [AL]
			^B 1.86	18.59	18.02	^a Distaph 250 [AF]

■ DICLOXACILLIN

Restricted benefit

Serious staphylococcal infection

dicloxacillin 500 mg capsule, 24

8122L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP MW	1	..	..	20.12	21.41	^a Dicloxacillin Mylan 500 [AL]
			^B 1.95	22.07	21.41	^a Distaph 500 [AF]

dicloxacillin 250 mg capsule, 24

8121K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP MW	1	..	..	16.73	18.02	^a Dicloxacillin Mylan 250 [AL]
			^B 1.86	18.59	18.02	^a Distaph 250 [AF]

■ DICLOXACILLIN

Authority required (STREAMLINED)

6188

Osteomyelitis

dicloxacillin 500 mg capsule, 24

10790X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	1	..	*28.22	29.51	^a Dicloxacillin Mylan 500 [AL]
			^B 3.90	*32.12	29.51	^a Distaph 500 [AF]

■ FLUCLOXACILLIN

Caution Severe cholestatic hepatitis has been reported with this drug. Significant risk factors are age, particularly greater than 55 years, and duration of treatment longer than 14 days.

Note Pharmaceutical benefits that have the form flucloxacillin 1 g injection in a pack size of 5 can be substituted for a pack size of 10 in the case of a shortage.

flucloxacillin 1 g injection, 10 vials

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
10605E NP	0.5	1	..	*25.24	26.53	^a Flubiclox [JU]

flucloxacillin 1 g injection, 10 vials

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
10609J DP	0.5	..	..	*25.24	26.53	^a Flubiclox [JU]

flucloxacillin 1 g injection, 5 vials

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
1525G NP	1	1	..	20.60	21.89	^a Flucil [AS]

flucloxacillin 1 g injection, 5 vials

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
5095E DP	1	..	..	20.60	21.89	^a Flucil [AS]

■ FLUCLOXACILLIN

Caution Severe cholestatic hepatitis has been reported with this drug. Significant risk factors are age, particularly greater than 55 years, and duration of treatment longer than 14 days.

Restricted benefit

Serious staphylococcal infection

flucloxacillin 250 mg/5 mL powder for oral liquid, 100 mL

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
5258R DP	±1	..	..	#24.12	25.78	Flucil [LN]

flucloxacillin 125 mg/5 mL powder for oral liquid, 100 mL

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
5257Q DP	±1	..	..	#21.05	22.71	Flucil [LN]

■ FLUCLOXACILLIN

Caution Severe cholestatic hepatitis has been reported with this drug. Significant risk factors are age, particularly greater than 55 years, and duration of treatment longer than 14 days.

Restricted benefit

Serious staphylococcal infection

flucloxacillin 250 mg/5 mL powder for oral liquid, 100 mL

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
9150N NP	±1	1	..	#24.12	25.78	Flucil [LN]

flucloxacillin 125 mg/5 mL powder for oral liquid, 100 mL

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
9149M NP	±1	1	..	#21.05	22.71	Flucil [LN]

■ FLUCLOXACILLIN

Caution Severe cholestatic hepatitis has been reported with this drug. Significant risk factors are age, particularly greater than 55 years, and duration of treatment longer than 14 days.

Note Pharmaceutical benefits that have the form flucloxacillin 250 mg capsule, 28 and pharmaceutical benefits that have the form flucloxacillin 250 mg capsule, 24 are equivalent for the purposes of substitution.

Restricted benefit

Serious staphylococcal infection

flucloxacillin 250 mg capsule, 24

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
1526H NP MW	1	..	..	16.29	17.58	^a APO-Flucloxacillin [TX] ^a Staphylex 250 [AF]	^a Flopen [AL]

■ FLUCLOXACILLIN

Caution Severe cholestatic hepatitis has been reported with this drug. Significant risk factors are age, particularly greater than 55 years, and duration of treatment longer than 14 days.

Note Pharmaceutical benefits that have the form flucloxacillin 500 mg capsule, 24 and pharmaceutical benefits that have the form flucloxacillin 500 mg capsule, 100 are equivalent for the purposes of substitution.

Restricted benefit

Serious staphylococcal infection

flucloxacillin 500 mg capsule, 24

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
1527J NP MW	1	..	..	20.71	22.00	^a APO-Flucloxacillin [TX] ^a Staphylex 500 [AF]	^a Flopen [AL]

■ FLUCLOXACILLIN

Caution Severe cholestatic hepatitis has been reported with this drug. Significant risk factors are age, particularly greater than 55 years, and duration of treatment longer than 14 days.

Note Pharmaceutical benefits that have the form flucloxacillin 250 mg capsule, 28 and pharmaceutical benefits that have the form flucloxacillin 250 mg capsule, 24 are equivalent for the purposes of substitution.

Restricted benefit

Serious staphylococcal infection

flucloxacillin 250 mg capsule, 24

5090X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
DP	1	..	..	16.29	17.58	^a APO-Flucloxacillin [TX] ^a Staphylex 250 [AF]	^a Flopen [AL]

■ FLUCLOXACILLIN

Caution Severe cholestatic hepatitis has been reported with this drug. Significant risk factors are age, particularly greater than 55 years, and duration of treatment longer than 14 days.

Note Pharmaceutical benefits that have the form flucloxacillin 500 mg capsule, 24 and pharmaceutical benefits that have the form flucloxacillin 500 mg capsule, 100 are equivalent for the purposes of substitution.

Restricted benefit

Serious staphylococcal infection

flucloxacillin 500 mg capsule, 24

5091Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
DP	1	..	..	20.71	22.00	^a APO-Flucloxacillin [TX] ^a Staphylex 500 [AF]	^a Flopen [AL]

■ FLUCLOXACILLIN

Caution Severe cholestatic jaundice has been reported with this drug. Significant risk factors are age, particularly greater than 55 years, and duration of treatment longer than 14 days.

Note Pharmaceutical benefits that have the form flucloxacillin 500 mg capsule, 24 and pharmaceutical benefits that have the form flucloxacillin 500 mg capsule, 100 are equivalent for the purposes of substitution.

Authority required (STREAMLINED)

6169

Osteomyelitis

flucloxacillin 500 mg capsule, 24

10788T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	1	..	*29.40	30.69	^a APO-Flucloxacillin [TX] ^a Staphylex 500 [AF]	^a Flopen [AL]

Combinations of penicillins, incl. beta-lactamase inhibitors

■ AMOXICILLIN + CLAVULANIC ACID

Authority required (STREAMLINED)

10413

Infection

Clinical criteria:

- Patient must have periorbital (preseptal) cellulitis; OR
- Patient must have postpartum endometritis; OR
- Patient must have an exacerbation of bronchiectasis; OR
- Patient must have pyelonephritis; OR
- Patient must have pneumonia acquired in hospital or aged care; OR
- Patient must have a diabetic foot infection; OR
- Patient must have a condition requiring prolonged oral antibiotic therapy.

amoxicillin 875 mg + clavulanic acid 125 mg tablet, 10

11933C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	2	..	..	*17.60	18.89	^a AlphaClav Duo Forte [AF] ^a AMOXICLAV AMNEAL 875/125 [ED] ^a AmoxyClav generichealth 875/125 [HQ] ^a Curam Duo Forte 875/125 [SZ] ^a Augmentin Duo forte [AS]	^a AMCLAVOX DUO FORTE 875/125 [RW] ^a Amoxyclav AN 875/125 [EA] ^a APO-Amoxycillin and Clavulanic Acid [TX]
			^B 12.44	*30.04	18.89		

■ AMOXICILLIN + CLAVULANIC ACID

Authority required (STREAMLINED)

10405

Infection

Clinical criteria:

- Patient must be a male with acute cystitis; OR
- Patient must have a condition requiring prolonged oral antibiotic therapy.

amoxicillin 500 mg + clavulanic acid 125 mg tablet, 10

11941L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	2	..	..	*17.04	18.33	^a AlphaClav Duo [AF]	^a AMCLAVOX DUO 500/125 [RW]
						^a AMOXICLAV AMNEAL 500/125 [ED]	^a Amoxycillin/Clavulanic Acid 500/125 APOTEX [TY]
						^a Amoxyclav AN 500/125 [EA]	^a APO-Amoxycillin/ Clavulanic Acid 500/125 [TX]
						^a Curam Duo 500/125 [SZ]	
			^B 9.66	*26.70	18.33	^a Augmentin Duo [AS]	

■ AMOXICILLIN + CLAVULANIC ACID

Caution Hepatotoxicity has been reported with this drug.

Restricted benefit

Infection where resistance to amoxicillin is suspected

Restricted benefit

Infections where resistance to amoxicillin is proven

amoxicillin 125 mg/5 mL + clavulanic acid 31.25 mg/5 mL powder for oral liquid, 75 mL

1892N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	1	..	#16.93	18.59	Curam [SZ]

amoxicillin 400 mg/5 mL + clavulanic acid 57 mg/5 mL powder for oral liquid, 60 mL

8319W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	1	..	#17.23	18.89	^a Curam Duo [SZ]
			^B 4.57	#21.80	18.89	^a Augmentin Duo 400 [AS]

■ AMOXICILLIN + CLAVULANIC ACID

Caution Hepatotoxicity has been reported with this drug.

Restricted benefit

Infection where resistance to amoxicillin is suspected

Restricted benefit

Infections where resistance to amoxicillin is proven

amoxicillin 875 mg + clavulanic acid 125 mg tablet, 10

5006L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
DP	1	..	..	14.81	16.10	^a AlphaClav Duo Forte [AF]	^a AMCLAVOX DUO FORTE 875/125 [RW]
						^a AMOXICLAV AMNEAL 875/125 [ED]	^a Amoxyclav AN 875/125 [EA]
						^a AmoxyClav generichealth 875/125 [HQ]	^a APO-Amoxycillin and Clavulanic Acid [TX]
						^a Curam Duo Forte 875/125 [SZ]	
			^B 6.22	21.03	16.10	^a Augmentin Duo forte [AS]	

amoxicillin 500 mg + clavulanic acid 125 mg tablet, 10

5008N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
DP	1	..	..	14.53	15.82	^a AlphaClav Duo [AF]	^a AMCLAVOX DUO 500/125 [RW]
						^a AMOXICLAV AMNEAL 500/125 [ED]	^a Amoxycillin/Clavulanic Acid 500/125 APOTEX [TY]
						^a Amoxyclav AN 500/125 [EA]	^a APO-Amoxycillin/ Clavulanic Acid 500/125 [TX]
						^a Curam Duo 500/125 [SZ]	
			^B 4.83	19.36	15.82	^a Augmentin Duo [AS]	

amoxicillin 125 mg/5 mL + clavulanic acid 31.25 mg/5 mL powder for oral liquid, 75 mL

5009P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	±1	..	..	#16.93	18.59	Curam [SZ]

amoxicillin 400 mg/5 mL + clavulanic acid 57 mg/5 mL powder for oral liquid, 60 mL

5011R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	±1	..	..	#17.23	18.89	^a Curam Duo [SZ]
			^B 4.57	#21.80	18.89	^a Augmentin Duo 400 [AS]

■ AMOXICILLIN + CLAVULANIC ACID

Caution Hepatotoxicity has been reported with this drug.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Restricted benefit

Infection where resistance to amoxicillin is suspected

Restricted benefit

Infections where resistance to amoxicillin is proven

amoxicillin 875 mg + clavulanic acid 125 mg tablet, 10

8254K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	14.81	16.10	^a AlphaClav Duo Forte [AF]	^a AMCLAVOX DUO FORTE 875/125 [RW]
						^a AMOXICLAV AMNEAL 875/125 [ED]	^a Amoxyclav AN 875/125 [EA]
						^a AmoxyClav generichealth 875/125 [HQ]	^a APO-Amoxycillin and Clavulanic Acid [TX]
						^a Curam Duo Forte 875/125 [SZ]	
			^B 6.22	21.03	16.10	^a Augmentin Duo forte [AS]	

amoxicillin 500 mg + clavulanic acid 125 mg tablet, 10

1891M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP MW	1	..	..	14.53	15.82	^a AlphaClav Duo [AF]	^a AMCLAVOX DUO 500/125 [RW]
						^a AMOXICLAV AMNEAL 500/125 [ED]	^a Amoxycillin/Clavulanic Acid 500/125 APOTEX [TY]
						^a Amoxyclav AN 500/125 [EA]	^a APO-Amoxycillin/ Clavulanic Acid 500/125 [TX]
						^a Curam Duo 500/125 [SZ]	
			^B 4.83	19.36	15.82	^a Augmentin Duo [AS]	

OTHER BETA-LACTAM ANTIBACTERIALS*First-generation cephalosporins***■ CEFALEXIN****cefalexin 250 mg/5 mL powder for oral liquid, 100 mL**

3095X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	‡1	1	..	#17.43	19.09	^a Cefalexin Sandoz [SZ]	^a Ibilex 250 [AF]
			^B 5.44	#22.87	19.09	^a Keflex [AS]	

cefalexin 250 mg/5 mL powder for oral liquid, 100 mL

3320R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
DP	‡1	..	..	#17.43	19.09	^a Cefalexin Sandoz [SZ]	^a Ibilex 250 [AF]
			^B 5.44	#22.87	19.09	^a Keflex [AS]	

cefalexin 125 mg/5 mL powder for oral liquid, 100 mL

3094W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	‡1	1	..	#17.15	18.81	^a Cefalexin Sandoz [SZ]	^a Ibilex 125 [AF]
			^B 3.88	#21.03	18.81	^a Keflex [AS]	

cefalexin 125 mg/5 mL powder for oral liquid, 100 mL

3319Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
DP	‡1	..	..	#17.15	18.81	^a Cefalexin Sandoz [SZ]	^a Ibilex 125 [AF]
			^B 3.88	#21.03	18.81	^a Keflex [AS]	

cefalexin 500 mg capsule, 20

3318P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
DP	1	..	..	13.96	15.25	^a APO-Cephalexin [TX]	^a Cefalexin Sandoz [SZ]
						^a Cephalax 500 [CR]	^a Cephalaxin AN [EA]
						^a Cephalaxin generichealth [GQ]	^a Ibilex 500 [AF]
						^a NOUMED CEFLEXIN [VO]	
			^B 5.39	19.35	15.25	^a Keflex [AS]	

cefalexin 250 mg capsule, 20

3317N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
DP	1	..	..	13.82	15.11	^a APO-Cephalexin [TX]	^a Cefalexin Sandoz [SZ]
						^a Cephalaxin AN [EA]	^a Ibilex 250 [AF]
			^B 3.67	17.49	15.11	^a Keflex [AS]	

■ CEFLEXIN**Authority required (STREAMLINED)****6188**

Osteomyelitis

cefaalexin 500 mg capsule, 20

10778G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	1	..	*15.90	17.19	^a APO-Cephalexin [TX] ^a Cephalex 500 [CR] ^a Cephalexin generichealth [GQ] ^a NOUMED CEFALEXIN [VO]	^a Cefalexin Sandoz [SZ] ^a Cephalexin AN [EA] ^a Ibilex 500 [AF]
			^B 10.78	*26.68	17.19	^a Keflex [AS]	

■ CEFALEXIN

Authority required (STREAMLINED)**10410**

Infection

Clinical criteria:

- Patient must have a pin-site infection; OR
- Patient must have an infection following cardiac device insertion; OR
- Patient must have acute otitis externa; OR
- Patient must have streptococcal pharyngitis or tonsillitis; OR
- Patient must have mastitis; OR
- Patient must have periorbital (preseptal) cellulitis; OR
- Patient must have acute rheumatic fever; OR
- Patient must have a diabetic foot infection; OR
- Patient must have a widespread infection of dermatitis; OR
- Patient must require treatment for prophylaxis for invasive group A streptococcal (iGAS) infection; OR
- Patient must have impetigo; OR
- Patient must have pyelonephritis; OR
- Patient must have a condition requiring prolonged oral antibiotic therapy.

Midwives may prescribe under this item for the treatment of mastitis only.

cefaalexin 500 mg capsule, 20

11934D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP MW	2	..	..	*15.90	17.19	^a APO-Cephalexin [TX] ^a Cephalex 500 [CR] ^a Cephalexin generichealth [GQ] ^a NOUMED CEFALEXIN [VO]	^a Cefalexin Sandoz [SZ] ^a Cephalexin AN [EA] ^a Ibilex 500 [AF]
			^B 10.78	*26.68	17.19	^a Keflex [AS]	

■ CEFALEXIN

Authority required (STREAMLINED)**10412**

Infection

Clinical criteria:

- Patient must have impaired renal function, **AND**
- Patient must have a pin-site infection; OR
- Patient must have an infection following cardiac device insertion; OR
- Patient must have acute otitis externa; OR
- Patient must have streptococcal pharyngitis or tonsillitis; OR
- Patient must have mastitis; OR
- Patient must have periorbital (preseptal) cellulitis; OR
- Patient must have acute rheumatic fever; OR
- Patient must have a diabetic foot infection; OR
- Patient must have a widespread infection of dermatitis; OR
- Patient must require treatment for prophylaxis for invasive group A streptococcal (iGAS) infection; OR
- Patient must have impetigo; OR
- Patient must have pyelonephritis; OR
- Patient must have a condition requiring prolonged oral antibiotic therapy.

Midwives may prescribe under this item for the treatment of mastitis only, where the patient has impaired renal function.

cefaalexin 250 mg capsule, 20

11963P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP MW	2	..	..	*15.62	16.91	^a APO-Cephalexin [TX] ^a Cephalexin AN [EA]	^a Cefalexin Sandoz [SZ] ^a Ibilex 250 [AF]
			^B 7.34	*22.96	16.91	^a Keflex [AS]	

■ CEFALEXIN

Authority required (STREAMLINED)**4243**

Prophylaxis of urinary tract infection

cefalexin 250 mg capsule, 20

2655R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	2	..	*15.62	16.91	^a APO-Cephalexin [TX]	^a Cefalexin Sandoz [SZ]
						^a Cephalexin AN [EA]	^a Ibilex 250 [AF]
			^B 7.34	*22.96	16.91	^a Keflex [AS]	

■ CEFALEXIN

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

cefalexin 500 mg capsule, 20

3119E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP MW	1	..	..	13.96	15.25	^a APO-Cephalexin [TX]	^a Cefalexin Sandoz [SZ]
						^a Cephalex 500 [CR]	^a Cephalexin AN [EA]
						^a Cephalexin generichealth [GQ]	^a Ibilex 500 [AF]
						^a NOUMED CEFALEXIN [VO]	
			^B 5.39	19.35	15.25	^a Keflex [AS]	

cefalexin 250 mg capsule, 20

3058Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP MW	1	..	..	13.82	15.11	^a APO-Cephalexin [TX]	^a Cefalexin Sandoz [SZ]
						^a Cephalexin AN [EA]	^a Ibilex 250 [AF]
			^B 3.67	17.49	15.11	^a Keflex [AS]	

■ CEFAZOLIN**Restricted benefit**

Cellulitis

cefazolin 1 g injection, 5 vials

1799Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	..	..	*21.80	23.09	Cefazolin-AFT [AE]

■ CEFAZOLIN**Restricted benefit**

Cellulitis

cefazolin 500 mg injection, 5 vials

5477G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	..	..	*18.32	19.61	Cefazolin-AFT [AE]

■ CEFAZOLIN

Note Pharmaceutical benefits that have the form cefazolin 2 g injection, 1 vial, and pharmaceutical benefits that have the form cefazolin 2 g injection, 10 vials are equivalent for the purposes of substitution.

Restricted benefit

Cellulitis

cefazolin 2 g injection, 10 vials

12115P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	39.87	41.16	^a Cephazolin Alphapharm [AF]

cefazolin 2 g injection, 1 vial

5479J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	10	..	..	*39.84	41.13	^a Cephazolin Alphapharm [AF]

■ CEFAZOLIN

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Infection where positive bacteriological evidence confirms that this antibiotic is an appropriate therapeutic agent

Restricted benefit

Septicaemia, suspected

Restricted benefit

Septicaemia, proven

cefazolin 500 mg injection, 5 vials

1256D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	..	..	*18.32	19.61	Cefazolin-AFT [AE]

■ CEFAZOLIN

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Infection where positive bacteriological evidence confirms that this antibiotic is an appropriate therapeutic agent

Restricted benefit

Septicaemia, suspected

Restricted benefit

Septicaemia, proven

cefazolin 1 g injection, 5 vials

1797N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	..	..	*21.80	23.09	Cefazolin-AFT [AE]

■ CEFAZOLIN

Note Pharmaceutical benefits that have the form cefazolin 2 g injection, 1 vial, and pharmaceutical benefits that have the form cefazolin 2 g injection, 10 vials are equivalent for the purposes of substitution.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Infection where positive bacteriological evidence confirms that this antibiotic is an appropriate therapeutic agent

Restricted benefit

Septicaemia, suspected

Restricted benefit

Septicaemia, proven

cefazolin 2 g injection, 10 vials

12118T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	39.87	41.16	^a Cephazolin Alphapharm [AF]

cefazolin 2 g injection, 1 vial

9326W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	10	..	..	*39.84	41.13	^a Cephazolin Alphapharm [AF]

Second-generation cephalosporins

■ CEFACLOR

Caution Serum sickness-like reactions have been reported with this drug, especially in children.

cefaclor 375 mg modified release tablet, 10

1169M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	1	..	15.93	17.22	^a APO-Cefaclor CD [TX]	^a Cefaclor GH [GQ]
						^a Karlor CD [MQ]	^a Keflor CD [AF]
			^B 7.85	23.78	17.22	^a Ceclor CD [AL]	

cefaclor 375 mg modified release tablet, 10

5045M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
DP	1	..	..	15.93	17.22	^a APO-Cefaclor CD [TX]	^a Cefaclor GH [GQ]
						^a Karlor CD [MQ]	^a Keflor CD [AF]
			^B 7.85	23.78	17.22	^a Ceclor CD [AL]	

cefaclor 125 mg/5 mL powder for oral liquid, 100 mL

2460L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	‡1	1	..	#18.44	20.10	^a Aclor 125 [MQ]	^a APO-Cefaclor [TX]
						^a Keflor [AF]	
			^B 7.82	#26.26	20.10	^a Ceclor [AL]	

cefaclor 125 mg/5 mL powder for oral liquid, 100 mL

5046N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
DP	‡1	..	..	#18.44	20.10	^a Aclor 125 [MQ]	^a APO-Cefaclor [TX]
						^a Keflor [AF]	
			^B 7.82	#26.26	20.10	^a Ceclor [AL]	

cefaclor 250 mg/5 mL powder for oral liquid, 75 mL

2461M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	‡1	1	..	#18.64	20.30	^a Aclor 250 [MQ]	^a APO-Cefaclor [TX]
						^a Keflor [AF]	

			^B 7.84	#26.48	20.30	^a Ceclor [AL]	
cefactor 250 mg/5 mL powder for oral liquid, 75 mL							
5047P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
DP	‡1	..	..	#18.64	20.30	^a Aclor 250 [MQ] ^a Keflor [AF]	^a APO-Cefactor [TX]
			^B 7.84	#26.48	20.30	^a Ceclor [AL]	

▪ CEFUROXIME

cefuroxime 250 mg tablet, 14

5052X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	
DP	1	..	..	20.58	21.87	Pharmacor Cefuroxime [CR]	

cefuroxime 250 mg tablet, 14

8292K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	
	1	1	..	20.58	21.87	Pharmacor Cefuroxime [CR]	

cefuroxime 125 mg/5 mL powder for oral liquid, 100 mL

11191B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	
DP	‡1	..	..	#27.21	28.87	Zinnat [AS]	

cefuroxime 125 mg/5 mL powder for oral liquid, 100 mL

11192C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	
	‡1	1	..	#27.21	28.87	Zinnat [AS]	

cefuroxime 250 mg tablet, 20

11227X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	1	..	24.25	25.54	^a Pharmacor Cefuroxime [CR]	^a Zinnat [AS]

cefuroxime 250 mg tablet, 20

11228Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
DP	1	..	..	24.25	25.54	^a Pharmacor Cefuroxime [CR]	^a Zinnat [AS]

Third-generation cephalosporins

▪ CEFOTAXIME

Restricted benefit

Infection where positive bacteriological evidence confirms that this antibiotic is an appropriate therapeutic agent

cefotaxime 1 g injection, 10 vials

1768C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	
DP	1	..	..	28.04	29.33	DBL Cefotaxime [PF]	

▪ CEFOTAXIME

Restricted benefit

Infection where positive bacteriological evidence confirms that this antibiotic is an appropriate therapeutic agent

cefotaxime 2 g injection, 10 vials

1769D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	
DP	1	..	..	35.67	36.96	DBL Cefotaxime [PF]	

▪ CEFOTAXIME

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Infection where positive bacteriological evidence confirms that this antibiotic is an appropriate therapeutic agent

Restricted benefit

Septicaemia, suspected

Restricted benefit

Septicaemia, proven

cefotaxime 2 g injection, 10 vials

1759N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	
NP	1	..	..	35.67	36.96	DBL Cefotaxime [PF]	

cefotaxime 1 g injection, 10 vials

1758M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	28.04	29.33	DBL Cefotaxime [PF]

■ CEFTRIAXONE**Restricted benefit**

Gonorrhoea

ceftriaxone 500 mg injection, 1 vial

9058R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	13.09	14.38	Ceftriaxone-AFT [AE]

■ CEFTRIAXONE**Note Shared Care Model:**

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Infection where positive bacteriological evidence confirms that this antibiotic is an appropriate therapeutic agent

Restricted benefit

Septicaemia, suspected

Restricted benefit

Septicaemia, proven

ceftriaxone 500 mg injection, 1 vial

1783W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	5	..	..	*17.39	18.68	Ceftriaxone-AFT [AE]

■ CEFTRIAXONE

Note Pharmaceutical benefits that have the form ceftriaxone 1 g injection, 1 vial, pharmaceutical benefits that have the form ceftriaxone 1 g injection, 5 vials, and pharmaceutical benefits that have the form ceftriaxone 1 g injection, 10 vials are equivalent for the purposes of substitution.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Infection where positive bacteriological evidence confirms that this antibiotic is an appropriate therapeutic agent

Restricted benefit

Septicaemia, suspected

Restricted benefit

Septicaemia, proven

ceftriaxone 1 g injection, 5 vials

1788D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	23.04	24.33	^a Ceftriaxone Alphapharm [AF]

ceftriaxone 1 g injection, 10 vials

12114N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	0.5	..	..	*28.26	29.55	^a Ceftriaxone Alphapharm [AF]

ceftriaxone 1 g injection, 1 vial

1784X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	5	..	..	*23.04	24.33	^a Ceftriaxone-AFT [AE]

■ CEFTRIAXONE

Note Pharmaceutical benefits that have the form ceftriaxone 2 g injection, 1 vial, pharmaceutical benefits that have the form ceftriaxone 2 g injection, 5 vials, and pharmaceutical benefits that have the form ceftriaxone 2 g injection, 10 vials are equivalent for the purposes of substitution.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Infection where positive bacteriological evidence confirms that this antibiotic is an appropriate therapeutic agent

Restricted benefit

Septicaemia, suspected

Restricted benefit

Septicaemia, proven

ceftriaxone 2 g injection, 10 vials

12112L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	0.5	..	..	*28.40	29.69	^a Ceftriaxone Alphapharm [AF]

ceftriaxone 2 g injection, 5 vials

11169W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	23.15	24.44	^a Ceftriaxone Alphapharm [AF]

ceftriaxone 2 g injection, 1 vial

1785Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	5	..	..	*23.19	24.48	^a Ceftriaxone-AFT [AE]

Fourth-generation cephalosporins**■ CEFEPIME****Note Shared Care Model:**

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required

Febrile neutropenia

cefepime 2 g injection, 1 vial

8316Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	10	..	..	*81.64	41.30	^a Cefepime-AFT [AE] ^a Cefepime Kabi [PK]	^a Cefepime Alphapharm [AF] ^a Omegapharm Pty Ltd [OE]

cefepime 1 g injection, 1 vial

8315P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	10	..	..	*52.14	41.30	^a Cefepime-AFT [AE] ^a Cefepime Kabi [PK]	^a Cefepime Alphapharm [AF] ^a Omegapharm Pty Ltd [OE]

SULFONAMIDES AND TRIMETHOPRIM**Trimethoprim and derivatives****■ TRIMETHOPRIM****trimethoprim 300 mg tablet, 7**

2922T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	1	..	13.72	15.01	^a Alprim [AF]	^a Trimethoprim Mylan [AL]
			^B 3.68	17.40	15.01	^a Triprim [RW]	

■ TRIMETHOPRIM**Authority required (STREAMLINED)****4243**

Prophylaxis of urinary tract infection

trimethoprim 300 mg tablet, 7

2666H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	2	..	*15.42	16.71	^a Alprim [AF]	^a Trimethoprim Mylan [AL]
			^B 7.36	*22.78	16.71	^a Triprim [RW]	

■ TRIMETHOPRIM**Restricted benefit**

Prostatitis

trimethoprim 300 mg tablet, 7

10785P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	4	..	..	*18.82	20.11	^a Alprim [AF]	^a Trimethoprim Mylan [AL]
			^B 14.72	*33.54	20.11	^a Triprim [RW]	

Combinations of sulfonamides and trimethoprim, incl. derivatives**■ TRIMETHOPRIM + SULFAMETHOXAZOLE**

Caution There is an increased risk of severe adverse reactions with this combination in the elderly.

trimethoprim 160 mg + sulfamethoxazole 800 mg tablet, 10

2951H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	13.92	15.21	^a Resprim Forte [AF]
			^B 4.16	18.08	15.21	^a Seprim Forte [RW]

ANTIINFECTIVES FOR SYSTEMIC USE

trimethoprim 160 mg + sulfamethoxazole 800 mg tablet, 10

3390K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	1	..	..	13.92	15.21	^a Resprim Forte [AF]
			^B 4.16	18.08	15.21	^a Septrin Forte [RW]

trimethoprim 40 mg/5 mL + sulfamethoxazole 200 mg/5 mL oral liquid, 100 mL

3103H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	1	..	20.32	21.61	Septrin [RW]

trimethoprim 40 mg/5 mL + sulfamethoxazole 200 mg/5 mL oral liquid, 100 mL

3391L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	±1	..	..	20.32	21.61	Septrin [RW]

■ TRIMETHOPRIM + SULFAMETHOXAZOLE

Caution There is an increased risk of severe adverse reactions with this combination in the elderly.

Authority required (STREAMLINED)

6201

Prophylaxis of Pneumocystis jiroveci pneumonia

trimethoprim 160 mg + sulfamethoxazole 800 mg tablet, 10

10784N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	2	..	*17.73	19.02	^a Resprim Forte [AF]
			^B 12.48	*30.21	19.02	^a Septrin Forte [RW]

MACROLIDES, LINCOSAMIDES AND STREPTOGRAMINS

Macrolides

■ AZITHROMYCIN

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Restricted benefit

Trachoma

azithromycin 500 mg tablet, 2

8336R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	2	..	16.41	17.70	^a APO-Azithromycin [TX] ^a Azithromycin Sandoz [SZ] ^a Zithromax [PF]	^a Azithromycin Mylan [AF] ^a ZITHRO [RW]

azithromycin 200 mg/5 mL powder for oral liquid, 15 mL

8201P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	..	..	#27.36	29.02	Zithromax [PF]

■ AZITHROMYCIN

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Restricted benefit

Urethritis

Clinical criteria:

- The condition must be uncomplicated and due to Chlamydia trachomatis.

Restricted benefit

Cervicitis

Clinical criteria:

- The condition must be uncomplicated and due to Chlamydia trachomatis.

azithromycin 500 mg tablet, 2

8200N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	16.41	17.70	^a APO-Azithromycin [TX] ^a Azithromycin Sandoz [SZ] ^a Zithromax [PF]	^a Azithromycin Mylan [AF] ^a ZITHRO [RW]

■ CLARITHROMYCIN

clarithromycin 250 mg tablet, 14

8318T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	1	..	15.18	16.47	^a Clarac [ED] ^a Clarithromycin AN [EA] ^a Kalixocin [AF]	^a Clarithro 250 [RW] ^a Clarithromycin Sandoz [SZ]
			^B 3.29	18.47	16.47	^a Klacid [GO]	

■ CLARITHROMYCIN

Restricted benefit

Bordetella pertussis

Restricted benefit

Atypical mycobacterial infections

clarithromycin 250 mg/5 mL powder for oral liquid, 50 mL

9192T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	..	..	#31.03	32.69	Klacid [GO]

■ ERYTHROMYCIN

erythromycin 250 mg enteric capsule, 25

1404X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	1	..	22.20	23.49	^a Eryc [YN]	^a Mayne Pharma Erythromycin [YT]

erythromycin 250 mg enteric capsule, 25

3325B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
DP	1	..	..	22.20	23.49	^a Eryc [YN]	^a Mayne Pharma Erythromycin [YT]

■ ERYTHROMYCIN

Authority required (STREAMLINED)**6160**

Severe acne

Clinical criteria:

- The condition must be one in which tetracycline therapy is inappropriate.

erythromycin 250 mg enteric capsule, 25

10780J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	5	..	*32.38	33.67	^a Eryc [YN]	^a Mayne Pharma Erythromycin [YT]

■ ERYTHROMYCIN ETHYLSUCCINATE

erythromycin (as ethylsuccinate) 400 mg/5 mL powder for oral liquid, 100 mL

2428T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	1	..	#21.03	22.69	E-Mycin 400 [AF]

erythromycin (as ethylsuccinate) 400 mg/5 mL powder for oral liquid, 100 mL

3337P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	±1	..	..	#21.03	22.69	E-Mycin 400 [AF]

erythromycin (as ethylsuccinate) 200 mg/5 mL powder for oral liquid, 100 mL

2424N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	1	..	#19.79	21.45	E-Mycin 200 [AF]

erythromycin (as ethylsuccinate) 200 mg/5 mL powder for oral liquid, 100 mL

3334L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	±1	..	..	#19.79	21.45	E-Mycin 200 [AF]

■ ROXITHROMYCIN

roxithromycin 150 mg tablet, 10

5260W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
DP	1	..	..	14.54	15.83	^a APO-Roxithromycin [TX] ^a Roxar 150 [RW] ^a Roxithromycin AN [EA] ^a Roxithromycin Sandoz [SZ]	^a Biaxsig [AV] ^a Roximycin [AF] ^a Roxithromycin-GA [ED]
			^b 2.26	16.80	15.83	^a Rulide [SW]	

roxithromycin 300 mg tablet, 5

5261X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
DP	1	..	..	14.54	15.83	^a APO-Roxithromycin [TX] ^a Roxar 300 [RW] ^a Roxithromycin AN [EA] ^a Roxithromycin Sandoz [SZ]	^a Biaxsig [AV] ^a Roximycin [AF] ^a Roxithromycin-GA [ED]
			^b 2.26	16.80	15.83	^a Rulide [SW]	

roxithromycin 50 mg dispersible tablet, 10

5259T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	1	..	..	17.67	18.96	Rulide D [SW]

■ ROXITHROMYCIN**Authority required (STREAMLINED)****10404**

Infection

Clinical criteria:

- Patient must have a condition requiring prolonged oral antibiotic therapy.

roxithromycin 150 mg tablet, 10

12001P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	2	..	..	*17.06	18.35	^a APO-Roxithromycin [TX] ^a Roxar 150 [RW] ^a Roxithromycin AN [EA] ^a Roxithromycin Sandoz [SZ]	^a Biaxsig [AV] ^a Roximycin [AF] ^a Roxithromycin-GA [ED]
			^b 4.52	*21.58	18.35	^a Rulide [SW]	

roxithromycin 300 mg tablet, 5

11993F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	2	..	..	*17.06	18.35	^a APO-Roxithromycin [TX] ^a Roxar 300 [RW] ^a Roxithromycin AN [EA] ^a Roxithromycin Sandoz [SZ]	^a Biaxsig [AV] ^a Roximycin [AF] ^a Roxithromycin-GA [ED]
			^b 4.52	*21.58	18.35	^a Rulide [SW]	

■ ROXITHROMYCIN**Note** No increase in the maximum quantity or number of units may be authorised.**Note** No increase in the maximum number of repeats may be authorised.**roxithromycin 150 mg tablet, 10**

1760P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	14.54	15.83	^a APO-Roxithromycin [TX] ^a Roxar 150 [RW] ^a Roxithromycin AN [EA] ^a Roxithromycin Sandoz [SZ]	^a Biaxsig [AV] ^a Roximycin [AF] ^a Roxithromycin-GA [ED]
			^b 2.26	16.80	15.83	^a Rulide [SW]	

roxithromycin 300 mg tablet, 5

8016X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	14.54	15.83	^a APO-Roxithromycin [TX] ^a Roxar 300 [RW] ^a Roxithromycin AN [EA] ^a Roxithromycin Sandoz [SZ]	^a Biaxsig [AV] ^a Roximycin [AF] ^a Roxithromycin-GA [ED]
			^b 2.26	16.80	15.83	^a Rulide [SW]	

roxithromycin 50 mg dispersible tablet, 10

8129W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	17.67	18.96	Rulide D [SW]

Lincosamides**■ CLINDAMYCIN****Restricted benefit**

Gram-positive coccal infections

Clinical criteria:

- The condition must not be able to be safely and effectively treated with a penicillin.

clindamycin 150 mg capsule, 24

5057E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
DP	1	..	..	18.89	20.18	^a APO-Clindamycin [TX] ^a Clindamycin BNM [BZ] ^a Clindamycin [AF]	^a Calindamin [RW] ^a Clindamycin LU [LV] ^a Dalacin C [PF]

■ CLINDAMYCIN**Restricted benefit**

Gram-positive coccal infections

Clinical criteria:

- The condition must not be able to be safely and effectively treated with a penicillin.

clindamycin 150 mg capsule, 24

3138E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP MW	2	1	..	*25.76	27.05	^a APO-Clindamycin [TX] ^a Clindamycin BNM [BZ] ^a Clindamylk [AF]	^a Calindamin [RW] ^a Clindamycin LU [LV] ^a Dalacin C [PF]

■ LINCOMYCIN

Note Pharmaceutical benefits that have the form lincomycin 600 mg/2 mL injection, 5 x 2 mL vials and pharmaceutical benefits that have the form lincomycin 600 mg/2 mL injection, 5 x 2 mL ampoules are equivalent for the purposes of substitution.

lincomycin 600 mg/2 mL injection, 5 x 2 mL vials

2530E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP MW	1	..	..	97.80	41.30	^a Lincocin [PF]

lincomycin 600 mg/2 mL injection, 5 x 2 mL vials

5144R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	1	..	..	97.80	41.30	^a Lincocin [PF]

lincomycin 600 mg/2 mL injection, 5 x 2 mL ampoules

11366F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	1	..	..	97.80	41.30	^a LINCOMYCIN SXP [XC]

lincomycin 600 mg/2 mL injection, 5 x 2 mL ampoules

11380Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP MW	1	..	..	97.80	41.30	^a LINCOMYCIN SXP [XC]

AMINOGLYCOSIDE ANTIBACTERIALS*Other aminoglycosides***■ GENTAMICIN****gentamicin 80 mg/2 mL injection, 10 x 2 mL ampoules**

2824P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	23.54	24.83	Pfizer Australia Pty Ltd [PF]

■ TOBRAMYCIN**Restricted benefit**

Pseudomonas aeruginosa infection

Clinical criteria:

- Patient must have cystic fibrosis, **AND**
- The treatment must be systemic.

tobramycin 500 mg/5 mL injection, 10 x 5 mL vials

9480Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	285.99	41.30	Tobra-Day [FF]

■ TOBRAMYCIN

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)**5520**

Proven Pseudomonas aeruginosa infection

Clinical criteria:

- Patient must have cystic fibrosis, **AND**
- The treatment must be for management.

tobramycin 300 mg/5 mL inhalation solution, 56 x 5 mL ampoules

5442K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	2	..	798.45	41.30	^a Tobr [GO] ^a Tobramycin WKT [LI]	^a TOBRAMYCIN SUN [RA] ^a TOBRAMYCIN WOCKHARDT [WC]

■ TOBRAMYCIN**Restricted benefit**

Infection where positive bacteriological evidence confirms that this antibiotic is an appropriate therapeutic agent

Restricted benefit

Septicaemia, suspected

Restricted benefit

Septicaemia, proven

tobramycin Injection 80 mg (base) in 2 mL (without preservative), 5

8872Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	1	..	*55.60	41.30	Pfizer Australia Pty Ltd [PF]

tobramycin 80 mg/2 mL injection, 5 x 2 mL vials

1356J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	2	1	..	*51.82	41.30	^a DBL Tobramycin [PF]	^a Tobramycin Mylan [AF]

■ TOBRAMYCIN**Note** No increase in the maximum quantity or number of units may be authorised.**Note** No increase in the maximum number of repeats may be authorised.**Note** Special Pricing Arrangements apply.**Authority required (STREAMLINED)****4456**Proven *Pseudomonas aeruginosa* infection

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have cystic fibrosis, **AND**
- Patient must have been assessed for bronchial hyperresponsiveness as per the TGA-approved Product Information, with a negative test result, **AND**
- Patient must be participating in a four week trial of tobramycin inhalation powder and will be assessed for ability to tolerate the dry powder formulation in order to qualify for continued PBS-subsidised therapy. The trial commencement date must be documented in the patient's medical records.

Population criteria:

- Patient must be 6 years of age or older.

tobramycin 28 mg powder for inhalation, 224 capsules

10066T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	2444.01	41.30	TOBI podhaler [GO]

■ TOBRAMYCIN**Note** No increase in the maximum quantity or number of units may be authorised.**Note** No increase in the maximum number of repeats may be authorised.**Note** Special Pricing Arrangements apply.**Authority required (STREAMLINED)****4513**Proven *Pseudomonas aeruginosa* infection

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have cystic fibrosis, **AND**
- Patient must have previously been issued with an authority prescription for tobramycin inhalation capsules, **AND**
- Patient must have demonstrated ability to tolerate the dry powder formulation following the initial 4-week treatment period, as agreed by the patient, the patient's family (in the case of paediatric patients) and the treating physician(s).

Population criteria:

- Patient must be 6 years of age or older.

tobramycin 28 mg powder for inhalation, 224 capsules

10074F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	2444.01	41.30	TOBI podhaler [GO]

QUINOLONE ANTIBACTERIALS***Fluoroquinolones*****■ CIPROFLOXACIN****Authority required**

Respiratory tract infection

Clinical criteria:

- The condition must be proven or suspected to be caused by *Pseudomonas aeruginosa*, **AND**
- Patient must be severely immunocompromised.

Authority required

Bacterial gastroenteritis

Clinical criteria:

- Patient must be severely immunocompromised.

Authority required

Infection

Clinical criteria:

- The condition must be proven to be due to *Pseudomonas aeruginosa* resistant to all other oral antimicrobials; OR
- The condition must be proven to be due to other gram-negative bacteria resistant to all other oral antimicrobials.

Authority required

Bone or joint infection

Clinical criteria:

- The condition must be suspected or proven to be caused by gram-negative bacteria resistant to all other appropriate antimicrobials; OR
- The condition must be suspected or proven to be caused by gram-positive bacteria resistant to all other appropriate antimicrobials.

Authority required

Epididymo-orchitis

Clinical criteria:

- The condition must be suspected or proven to be caused by gram-negative bacteria resistant to all other appropriate antimicrobials; OR
- The condition must be suspected or proven to be caused by gram-positive bacteria resistant to all other appropriate antimicrobials.

Authority required

Prostatitis

Clinical criteria:

- The condition must be suspected or proven to be caused by gram-negative bacteria resistant to all other appropriate antimicrobials; OR
- The condition must be suspected or proven to be caused by gram-positive bacteria resistant to all other appropriate antimicrobials.

Authority required

Perichondritis of the pinna

Clinical criteria:

- The condition must be suspected or proven to be caused by gram-negative bacteria resistant to all other appropriate antimicrobials; OR
- The condition must be suspected or proven to be caused by gram-positive bacteria resistant to all other appropriate antimicrobials.

ciprofloxacin 750 mg tablet, 14

1210Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	21.45	22.74	^a APO-Ciprofloxacin [TX] ^a Ciprofloxacin AN [EA] ^a Ciprol 750 [RW] ^a NOUMED CIPROFLOXACIN [VO]	^a C-Flox 750 [AL] ^a Ciprofloxacin Sandoz [SZ] ^a Loxip 750 [DO]

ciprofloxacin 500 mg tablet, 14

1209P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	18.28	19.57	^a APO-Ciprofloxacin [TX] ^a Cifran [RA] ^a Ciprofloxacin GH [HQ] ^a Ciprol 500 [RW] ^a NOUMED CIPROFLOXACIN [VO]	^a C-Flox 500 [AL] ^a Ciprofloxacin AN [EA] ^a Ciprofloxacin Sandoz [SZ] ^a Loxip 500 [DO]

▪ **CIPROFLOXACIN**

Authority required

Respiratory tract infection

Clinical criteria:

- The condition must be proven or suspected to be caused by *Pseudomonas aeruginosa*, **AND**
- Patient must be severely immunocompromised.

Authority required

Bacterial gastroenteritis

Clinical criteria:

- Patient must be severely immunocompromised.

Authority required

Infection

Clinical criteria:

- The condition must be proven to be due to *Pseudomonas aeruginosa* resistant to all other oral antimicrobials; OR
- The condition must be proven to be due to other gram-negative bacteria resistant to all other oral antimicrobials.

Authority required

Bone or joint infection

Clinical criteria:

- The condition must be suspected or proven to be caused by gram-negative bacteria resistant to all other appropriate antimicrobials; OR

- The condition must be suspected or proven to be caused by gram-positive bacteria resistant to all other appropriate antimicrobials.

Authority required

Epididymo-orchitis

Clinical criteria:

- The condition must be suspected or proven to be caused by gram-negative bacteria resistant to all other appropriate antimicrobials; OR
- The condition must be suspected or proven to be caused by gram-positive bacteria resistant to all other appropriate antimicrobials.

Authority required

Prostatitis

Clinical criteria:

- The condition must be suspected or proven to be caused by gram-negative bacteria resistant to all other appropriate antimicrobials; OR
- The condition must be suspected or proven to be caused by gram-positive bacteria resistant to all other appropriate antimicrobials.

Authority required

Perichondritis of the pinna

Clinical criteria:

- The condition must be suspected or proven to be caused by gram-negative bacteria resistant to all other appropriate antimicrobials; OR
- The condition must be suspected or proven to be caused by gram-positive bacteria resistant to all other appropriate antimicrobials.

Authority required

Gonorrhoea

ciprofloxacin 250 mg tablet, 14

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
1208N	1	..	..	15.41	16.70	^a APO-Ciprofloxacin [TX] ^a Ciprofloxacin Sandoz [SZ]	^a C-Flox 250 [AL] ^a Ciprol 250 [RW]

NP

■ NORFLOXACIN
Authority required

Acute bacterial enterocolitis

Authority required

Complicated urinary tract infection

norfloxacin 400 mg tablet, 14

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
3010K	1	1	..	15.76	17.05	^a APO-Norfloxacin [TX] ^a Nufloxib [AF]	^a GenRx Norfloxacin [GX] ^a Roxin [RW]

NP

OTHER ANTIBACTERIALS
Glycopeptide antibacterials
■ VANCOMYCIN
Restricted benefit

Endocarditis

Clinical criteria:

- The treatment must be for prophylaxis, **AND**
- Patient must be hypersensitive to penicillin.

vancomycin 1 g injection, 1 vial

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
2269K	1	..	..	20.08	21.37	Vancomycin Alphapharm [AF]

vancomycin 500 mg injection, 1 vial

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
3130R	2	..	..	*22.10	23.39	Vancomycin Alphapharm [AF]

■ VANCOMYCIN
Restricted benefit

Endocarditis

Clinical criteria:

- The treatment must be for prophylaxis, **AND**
- Patient must be hypersensitive to penicillin.

vancomycin 1 g injection, 1 vial

5083M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	1	..	..	20.08	21.37	Vancomycin Alphapharm [AF]

vancomycin 500 mg injection, 1 vial

3323X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	2	..	..	*22.10	23.39	Vancomycin Alphapharm [AF]

▪ **VANCOMYCIN**

Restricted benefit

Endophthalmitis

Restricted benefit

Infection

Clinical criteria:

- The treatment must be initiated in a hospital, **AND**
- The condition must be one in which vancomycin is an appropriate antibiotic.

vancomycin 1 g injection, 1 vial

2270L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	..	..	*36.21	37.50	Vancomycin Alphapharm [AF]

vancomycin 500 mg injection, 1 vial

3131T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	5	..	..	*37.24	38.53	Vancomycin Alphapharm [AF]

Steroid antibacterials

▪ **FUSIDATE**

Restricted benefit

Serious staphylococcal infections

Clinical criteria:

- The treatment must be used in combination with another antibiotic, **AND**
- The condition must be proven to be due to a staphylococcus.

sodium fusidate 250 mg tablet, 36

2312Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	84.97	41.30	Fucidin [LO]

▪ **FUSIDATE**

Authority required (STREAMLINED)

6133

Osteomyelitis

Clinical criteria:

- The condition must be methicillin-resistant staphylococcal aureus (MRSA), **AND**
- The treatment must be used in combination with other anti-staphylococcal antibiotics.

sodium fusidate 250 mg tablet, 36

10782L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	1	..	*160.22	41.30	Fucidin [LO]

Imidazole derivatives

▪ **METRONIDAZOLE**

metronidazole 200 mg tablet, 21

1636D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	1	..	13.61	14.90	^a Metrogyl 200 [AF]	^a Metronide 200 [AV]
			^B 1.86	15.47	14.90	^a Flagyl [SW]	

metronidazole 200 mg tablet, 21

3339R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
DP	1	..	..	13.61	14.90	^a Metrogyl 200 [AF]	^a Metronide 200 [AV]
			^B 1.86	15.47	14.90	^a Flagyl [SW]	

metronidazole 200 mg/5 mL oral liquid, 100 mL

1630T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	‡1	..	..	22.80	24.09	Flagyl S [SW]

metronidazole 200 mg/5 mL oral liquid, 100 mL

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
3341W	1	..	..	22.80	24.09	Flagyl S [SW]

metronidazole 500 mg suppository, 10

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
1642K	1	..	..	26.58	27.87	Flagyl [SW]

metronidazole 500 mg suppository, 10

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
5157K	1	..	..	26.58	27.87	Flagyl [SW]

■ METRONIDAZOLE**Restricted benefit**

Anaerobic infections

metronidazole 400 mg tablet, 21

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
1621H	1	1	..	15.20	16.49	^a Metrogyl 400 [AF]	^a Metronide 400 [AV]
			^B 1.86	17.06	16.49	^a Flagyl [SW]	

■ METRONIDAZOLE**Restricted benefit**

Anaerobic infections

metronidazole 400 mg tablet, 21

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
5155H	1	..	..	15.20	16.49	^a Metrogyl 400 [AF]	^a Metronide 400 [AV]
			^B 1.86	17.06	16.49	^a Flagyl [SW]	

*Nitrofurantoin derivatives***■ NITROFURANTOIN****Caution** Nitrofurantoin may cause peripheral neuritis and severe pulmonary reactions.**nitrofurantoin 50 mg capsule, 30**

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
1692C	1	1	..	23.27	24.56	^a APO-Nitrofurantoin [TX]	^a Macrochantin [PF]

nitrofurantoin 100 mg capsule, 30

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
1693D	1	1	..	27.79	29.08	^a APO-Nitrofurantoin [TX]	^a Macrochantin [PF]

*Other antibacterials***■ METHENAMINE HIPPURATE****methenamine hippurate 1 g tablet, 100**

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
3124K	1	5	..	33.20	34.49	^a Hiprex [IL]	^a Uramet [AS]

■ ANTIMYCOTICS FOR SYSTEMIC USE**ANTIMYCOTICS FOR SYSTEMIC USE***Triazole derivatives***■ FLUCONAZOLE****Note Shared Care Model:**

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**6002**

Cryptococcal meningitis

Authority required (STREAMLINED)**5978**

Cryptococcal meningitis

Clinical criteria:

- The treatment must be maintenance therapy, **AND**
- Patient must be immunosuppressed.

Authority required (STREAMLINED)

6023

Oropharyngeal candidiasis

Clinical criteria:

- Patient must be immunosuppressed.

Authority required (STREAMLINED)**5989**

Oesophageal candidiasis

Clinical criteria:

- Patient must be immunosuppressed.

Authority required (STREAMLINED)**6030**

Oropharyngeal candidiasis

Clinical criteria:

- The treatment must be for prophylaxis, **AND**
- Patient must be immunosuppressed.

Authority required (STREAMLINED)**7898**

Fungal infection

Clinical criteria:

- The condition must be serious or life-threatening.

fluconazole 400 mg/200 mL injection, 200 mL bag

1757L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	16.37	17.66	Fluconazole Alphapharm [AF]

■ FLUCONAZOLE**Note Shared Care Model:**

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**6956**

Cryptococcal meningitis

Authority required (STREAMLINED)**6978**

Cryptococcal meningitis

Clinical criteria:

- The treatment must be maintenance therapy, **AND**
- Patient must be immunosuppressed.

Authority required (STREAMLINED)**6974**

Oropharyngeal candidiasis

Clinical criteria:

- Patient must be immunosuppressed.

Authority required (STREAMLINED)**6969**

Oesophageal candidiasis

Clinical criteria:

- Patient must be immunosuppressed.

Authority required (STREAMLINED)**6965**

Oropharyngeal candidiasis

Clinical criteria:

- The treatment must be for prophylaxis, **AND**
- Patient must be immunosuppressed.

Authority required (STREAMLINED)**7897**

Fungal infection

Clinical criteria:

- The condition must be serious or life-threatening.

fluconazole 200 mg/100 mL injection, 100 mL bag

11139G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	7	..	..	*47.29	41.30	Fluconazole Alphapharm [AF]

■ FLUCONAZOLE

Note Not for use in vulvovaginal candida infections.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**6002**

Cryptococcal meningitis

Authority required (STREAMLINED)**5978**

Cryptococcal meningitis

Clinical criteria:

- The treatment must be maintenance therapy, **AND**
- Patient must be immunosuppressed.

Authority required (STREAMLINED)**6023**

Oropharyngeal candidiasis

Clinical criteria:

- Patient must be immunosuppressed.

Authority required (STREAMLINED)**5989**

Oesophageal candidiasis

Clinical criteria:

- Patient must be immunosuppressed.

Authority required (STREAMLINED)**6030**

Oropharyngeal candidiasis

Clinical criteria:

- The treatment must be for prophylaxis, **AND**
- Patient must be immunosuppressed.

Authority required (STREAMLINED)**7898**

Fungal infection

Clinical criteria:

- The condition must be serious or life-threatening.

fluconazole 50 mg capsule, 28

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
1471K	1	5	..	22.66	23.95	^a Dizole 50 [AF]	^a Fluconazole Sandoz [SZ]
						^a Ozole [RA]	
			^B 6.00	28.66	23.95	^a Diflucan [PF]	

fluconazole 200 mg capsule, 28

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
1475P	1	5	..	52.39	41.30	^a APO-Fluconazole [TX]	^a Diflucan [PF]
						^a Dizole 200 [AF]	^a Fluconazole APOTEX [GX]
						^a Fluconazole Sandoz [SZ]	^a Fluzole 200 [RW]
						^a Ozole [RA]	

fluconazole 100 mg capsule, 28

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
1472L	1	5	..	32.74	34.03	^a Diflucan [PF]	^a Dizole 100 [AF]
						^a Fluconazole Sandoz [SZ]	^a Ozole [RA]

■ FLUCONAZOLE

Note Not for use in vulvovaginal candida infections.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**6006**

Cryptococcal meningitis

Clinical criteria:

- Patient must be unable to take a solid dose form of fluconazole.

Authority required (STREAMLINED)**6045**

Cryptococcal meningitis

Clinical criteria:

- The treatment must be maintenance therapy, **AND**

- Patient must be immunosuppressed, **AND**
- Patient must be unable to take a solid dose form of fluconazole.

Authority required (STREAMLINED)

6031

Oropharyngeal candidiasis

Clinical criteria:

- Patient must be immunosuppressed, **AND**
- Patient must be unable to take a solid dose form of fluconazole.

Authority required (STREAMLINED)

6046

Oesophageal candidiasis

Clinical criteria:

- Patient must be immunosuppressed, **AND**
- Patient must be unable to take a solid dose form of fluconazole.

Authority required (STREAMLINED)

6032

Oropharyngeal candidiasis

Clinical criteria:

- The treatment must be for prophylaxis, **AND**
- Patient must be immunosuppressed, **AND**
- Patient must be unable to take a solid dose form of fluconazole.

Authority required (STREAMLINED)

7934

Fungal infection

Clinical criteria:

- The condition must be serious or life-threatening, **AND**
- Patient must be unable to take a solid dose form of fluconazole.

fluconazole 50 mg/5 mL powder for oral liquid, 35 mL

5446P

Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
1	..	..	#68.46	41.30	Diflucan [PF]

NP

■ **ITRACONAZOLE**

Note Not for use in vulvovaginal candida infections.

Note One capsule of itraconazole 50 mg (Lozanoc) is therapeutically equivalent to one 100 mg capsule of conventional itraconazole (Sporanox). The recommended dose of Lozanoc is therefore half the recommended dose for Sporanox. Lozanoc 50 mg capsules and Sporanox 100 mg capsules are not interchangeable.

Note Not for use in superficial mycoses

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)

6022

Systemic aspergillosis

Authority required (STREAMLINED)

6005

Systemic sporotrichosis

Authority required (STREAMLINED)

6057

Systemic histoplasmosis

Authority required (STREAMLINED)

5988

Disseminated pulmonary histoplasmosis infection

Treatment Phase: Treatment and maintenance therapy

Clinical criteria:

- Patient must be diagnosed with acquired immunodeficiency syndrome (AIDS).

Authority required (STREAMLINED)

6037

Chronic pulmonary histoplasmosis infection

Treatment Phase: Treatment and maintenance therapy

Clinical criteria:

- Patient must be diagnosed with acquired immunodeficiency syndrome (AIDS).

Authority required (STREAMLINED)

6016

Oropharyngeal candidiasis

Clinical criteria:

- Patient must be immunosuppressed.

Authority required (STREAMLINED)

6035

Oesophageal candidiasis

Clinical criteria:

- Patient must be immunosuppressed.

itraconazole 50 mg capsule, 60

10732W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	116.78	41.30	Lozanoc [YN]

itraconazole 100 mg capsule, 60

8196J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	116.78	41.30	^a APO-Itraconazole [TX] ^a ITRANOX [RW]	^a Itracap [AF] ^a Sporanox [JC]

▪ **POSACONAZOLE**

Note Application for an increased maximum quantity to allow for up to 1 month's treatment and repeats sufficient for up to 6 months' treatment may be authorised.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required

Invasive aspergillosis

Clinical criteria:

- Patient must be unable to tolerate alternative therapy; OR
- Patient must have disease refractory to alternative therapy.

Authority required

Prophylaxis of invasive fungal infections including both yeasts and moulds

Clinical criteria:

- Patient must be considered at high risk of developing an invasive fungal infection due to anticipated neutropenia (an absolute neutrophil count less than 500 cells per cubic millimetre), for at least 10 days whilst receiving chemotherapy for acute myeloid leukaemia or myelodysplastic syndrome; OR
- Patient must be considered at high risk of developing an invasive fungal infection due to having acute graft versus host disease (GVHD) grade II, III or IV, or extensive chronic GVHD, and receiving intensive immunosuppressive therapy after allogeneic haematopoietic stem cell transplant.

Treatment of neutropenia should continue until recovery of the neutrophil count to at least 500 cells per cubic millimetre.

Patients who have had a previous invasive fungal infection should have secondary prophylaxis during subsequent episodes of neutropenia.

No more than 6 months therapy per episode will be PBS-subsidised

Authority required

Fungal infection

Clinical criteria:

- The condition must be fusariosis; OR
- The condition must be zygomycosis; OR
- The condition must be coccidioidomycosis; OR
- The condition must be chromoblastomycosis; OR
- The condition must be mycetoma, **AND**
- Patient must be unable to tolerate alternative therapy; OR
- Patient must have disease refractory to alternative therapy.

posaconazole 100 mg modified release tablet, 24

10460M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	467.51	41.30	^a Pharmacor Posaconazole [CR] ^a POSACONAZOLE DR.REDDY'S [RZ] ^a Posaconazole Sandoz [SZ]	^a Posaconazole ARX [XT] ^a Posaconazole Juno [JU]

▪ **VORICONAZOLE**

Note For patients with graft versus host disease, acute myeloid leukaemia or myelodysplastic syndrome, applications for an increased maximum quantity to allow for up to 1 month's treatment and repeats sufficient for up to 6 months' treatment may be authorised.

Note For patients undergoing allogeneic haematopoietic stem cell transplant, applications for an increased maximum quantity to allow for up to 1 month's treatment and repeats sufficient for up to 2 months' treatment may be authorised.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required

Prophylaxis of invasive fungal infections including both yeasts and moulds

Clinical criteria:

- Patient must be considered at high risk of developing an invasive fungal infection due to anticipated neutropenia (an absolute neutrophil count less than 500 cells per cubic millimetre) for at least 10 days whilst receiving chemotherapy for acute myeloid leukaemia or myelodysplastic syndrome; OR
- Patient must be considered at high risk of developing an invasive fungal infection due to having acute graft versus host disease (GVHD) grade II, III or IV, or, extensive chronic GVHD, whilst receiving intensive immunosuppressive therapy after allogeneic haematopoietic stem cell transplant; OR
- Patient must be undergoing allogeneic haematopoietic stem cell transplant using either bone marrow from an unrelated donor or umbilical cord blood (related or unrelated), and, be considered to be at high risk of developing an invasive fungal infection during the neutropenic phase prior to engraftment.

voriconazole 40 mg/mL powder for oral liquid, 70 mL

10168E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	#505.69	41.30	Vfend [PF]

voriconazole 200 mg tablet, 56

10198R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	826.94	41.30	^a Vfend [PF] ^a Voriconazole APOTEX [TX] ^a Vttack [AF]	^a Voriconazole APO [GX] ^a Voriconazole Sandoz [SZ] ^a Vzole [RW]

voriconazole 50 mg tablet, 56

10173K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	212.14	41.30	^a Vfend [PF] ^a Voriconazole APOTEX [TX] ^a Vttack [AF]	^a Voriconazole APO [GX] ^a Voriconazole Sandoz [SZ] ^a Vzole [RW]

▪ **VORICONAZOLE**

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required

Definite or probable invasive aspergillosis

Treatment Phase: Treatment and maintenance therapy

Clinical criteria:

- Patient must be immunocompromised.

Authority required

Serious fungal infections

Treatment Phase: Treatment and maintenance therapy

Clinical criteria:

- The condition must be caused by *Scedosporium* species; OR
- The condition must be caused by *Fusarium* species.

Authority required

Serious Candida infections

Treatment Phase: Treatment and maintenance therapy

Clinical criteria:

- The condition must be caused by species not susceptible to fluconazole; OR
- The condition must be resistant to fluconazole; OR
- Patient must be unable to tolerate fluconazole.

Authority required

Serious invasive mycosis infections

Treatment Phase: Treatment and maintenance therapy

Clinical criteria:

- The treatment must be for invasive mycosis infections other than definite or probable invasive aspergillosis.

voriconazole 200 mg tablet, 56

9364W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	2	..	826.94	41.30	^a Vfend [PF] ^a Voriconazole APOTEX [TX] ^a Vttack [AF]	^a Voriconazole APO [GX] ^a Voriconazole Sandoz [SZ] ^a Vzole [RW]

voriconazole 50 mg tablet, 56

9363T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	2	..	212.14	41.30	^a Vfend [PF] ^a Voriconazole APOTEX [TX] ^a Vttack [AF]	^a Voriconazole APO [GX] ^a Voriconazole Sandoz [SZ] ^a Vzole [RW]

■ VORICONAZOLE

Note Application for an increased maximum quantity to allow for up to 1 month's treatment and repeats sufficient for up to 6 months' treatment may be authorised.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required

Definite or probable invasive aspergillosis

Treatment Phase: Treatment and maintenance therapy

Clinical criteria:

- Patient must be immunocompromised.

Authority required

Serious fungal infections

Treatment Phase: Treatment and maintenance therapy

Clinical criteria:

- The condition must be caused by *Scedosporium* species; OR
- The condition must be caused by *Fusarium* species.

Authority required

Serious *Candida* infections

Treatment Phase: Treatment and maintenance therapy

Clinical criteria:

- The condition must be caused by species not susceptible to fluconazole; OR
- The condition must be resistant to fluconazole; OR
- Patient must be unable to tolerate fluconazole.

Authority required

Serious invasive mycosis infections

Treatment Phase: Treatment and maintenance therapy

Clinical criteria:

- The treatment must be for invasive mycosis infections other than definite or probable invasive aspergillosis.

voriconazole 40 mg/mL powder for oral liquid, 70 mL

9452L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	#505.69	41.30	Vfend [PF]

■ ANTIMYCOBACTERIALS

DRUGS FOR TREATMENT OF TUBERCULOSIS

Antibiotics

■ RIFAMPICIN

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required

Mycobacterium ulcerans infection (Buruli ulcer)

Clinical criteria:

- The treatment must be used in combination with another antibiotic for the treatment of Buruli ulcer.

rifampicin 300 mg capsule, 10

12189M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	12	..	..	*138.42	41.30	Rimycin 300 [AF]

rifampicin 150 mg capsule, 100

12190N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	247.04	41.30	Rimycin 150 [AF]

rifampicin 300 mg capsule, 100

12215X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	116.52	41.30	Rimycin 300 [AF]

rifampicin 150 mg capsule, 10

12200D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	12	..	..	*295.02	41.30	Rimycin 150 [AF]

Hydrazides

■ ISONIAZID

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

isoniazid 100 mg tablet, 100

1554T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	22.65	23.94	Arrow Pharma Pty Ltd [RW]

DRUGS FOR TREATMENT OF LEPRA*Drugs for treatment of lepra*

■ DAPSONE

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

dapsone 100 mg tablet, 100

1272Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	309.07	41.30	Link Medical Products Pty Ltd [LM]

dapsone 25 mg tablet, 100

8801F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	253.17	41.30	Link Medical Products Pty Ltd [LM]

■ RIFAMPICIN

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required

Leprosy

Population criteria:

- Patient must be an adult.

rifampicin 150 mg capsule, 100

1982H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	247.04	41.30	Rimycin 150 [AF]

rifampicin 300 mg capsule, 100

1983J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	116.52	41.30	Rimycin 300 [AF]

■ RIFAMPICIN

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Meningococcal disease

Clinical criteria:

- The treatment must be for prophylaxis, **AND**
- Patient must be a carrier of the disease; OR
- Patient must be in close contact with people who have the disease.

Restricted benefit

Haemophilus influenzae type B

Clinical criteria:

- The treatment must be for prophylaxis, **AND**
- Patient must be in contact with people who have the disease.

rifampicin 100 mg/5 mL oral liquid, 60 mL

8025J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	..	..	28.91	30.20	Rifadin [SW]

rifampicin 300 mg capsule, 10

1984K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	22.45	23.74	Rimycin 300 [AF]

rifampicin 150 mg capsule, 10

1981G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	34.88	36.17	Rimycin 150 [AF]

■ **ANTIVIRALS FOR SYSTEMIC USE**

DIRECT ACTING ANTIVIRALS

Nucleosides and nucleotides excl. reverse transcriptase inhibitors

■ **ACICLOVIR**

Note Aciclovir 200 mg is not PBS-subsidised for chickenpox, herpes zoster or herpes simplex infections other than genital herpes.

Authority required (STREAMLINED)

5942

Recurrent moderate to severe genital herpes

Treatment Phase: Episodic treatment or suppressive therapy

Microbiological confirmation of diagnosis [viral culture, antigen detection or nucleic acid amplification by polymerase chain reaction (PCR)] is desirable but need not delay treatment.

aciclovir 200 mg tablet, 90

1007B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	36.56	37.85	^a Aciclovir AN [ED] ^a Aciclovir GH [GQ] ^a APO-Aciclovir [TX] ^a Lovir [EA]	^a Aciclovir APOTEX [TY] ^a Aciclovir Sandoz [HX] ^a GenRx Aciclovir [GX]

■ **ACICLOVIR**

Note This drug is only effective if commenced within 72 hours of onset of rash.

Note Aciclovir 800 mg is not PBS-subsidised for herpes simplex or chickenpox.

Note No applications for repeats will be authorised.

Authority required (STREAMLINED)

5967

Herpes zoster

Clinical criteria:

- The treatment must be administered within 72 hours of the onset of the rash.

Authority required (STREAMLINED)

5959

Herpes zoster ophthalmicus

aciclovir 800 mg tablet, 35

1052J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	41.72	41.30	^a Aciclovir APOTEX [TY] ^a APO-Aciclovir [TX]	^a Aciclovir Sandoz [HX] ^a GenRx Aciclovir [GX]

■ **ACICLOVIR**

Note Aciclovir 200 mg is not PBS-subsidised for chickenpox, herpes zoster or herpes simplex infections other than genital herpes.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note For item codes 1003T and 1555W, pharmaceutical benefits that have the form tablet 200 mg are equivalent for the purposes of substitution.

Authority required (STREAMLINED)

5936

Initial moderate to severe genital herpes

Microbiological confirmation of diagnosis [viral culture, antigen detection or nucleic acid amplification by polymerase chain reaction (PCR)] is desirable but need not delay treatment.

aciclovir 200 mg tablet, 50

1555W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	25.65	26.94	^a Aciclovir APOTEX [TY] ^a GenRx Aciclovir [GX]	^a APO-Aciclovir [TX]

aciclovir 200 mg tablet, 25

1003T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	..	..	*25.66	26.95	^a Aciclovir Sandoz [HX]

■ FAMCICLOVIR

Note Famciclovir 250 mg is not PBS-subsidised for chickenpox or herpes simplex infections other than genital herpes.

Authority required (STREAMLINED)

5937

Recurrent moderate to severe genital herpes

Treatment Phase: Episodic treatment

Microbiological confirmation of diagnosis [viral culture, antigen detection or nucleic acid amplification by polymerase chain reaction (PCR)] is desirable but need not delay treatment.

famciclovir 250 mg tablet, 20

2274Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	1	..	33.75	35.04	^a APO-Famciclovir [TX] ^a Famciclovir AN [EA] ^a Famvir [IX]	^a Ezovir [AF] ^a Famciclovir-GA [ED] ^a Favic 250 [RW]

■ FAMCICLOVIR

Note Famciclovir 125 mg is not PBS-subsidised for chickenpox, herpes zoster or herpes simplex infections other than genital herpes.

Authority required (STREAMLINED)

5937

Recurrent moderate to severe genital herpes

Treatment Phase: Episodic treatment

Microbiological confirmation of diagnosis [viral culture, antigen detection or nucleic acid amplification by polymerase chain reaction (PCR)] is desirable but need not delay treatment.

famciclovir 125 mg tablet, 40

8092X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	1	..	33.75	35.04	^a APO-Famciclovir [TX] ^a Ezovir [AF] ^a Famciclovir-GA [ED] ^a Favic 125 [RW]	^a Auro-Famciclovir 125 [DO] ^a Famciclovir AN [EA] ^a Famvir [IX]

■ FAMCICLOVIR

Note Famciclovir 250 mg is not PBS-subsidised for chickenpox or herpes simplex infections other than genital herpes.

Authority required (STREAMLINED)

5971

Recurrent moderate to severe genital herpes

Treatment Phase: Suppressive therapy

Microbiological confirmation of diagnosis [viral culture, antigen detection or nucleic acid amplification by polymerase chain reaction (PCR)] is desirable but need not delay treatment.

famciclovir 250 mg tablet, 56

8217L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	72.87	41.30	^a APO-Famciclovir [TX] ^a Ezovir [AF] ^a Famciclovir-GA [ED] ^a Famvir [IX]	^a Auro-Famciclovir 250 [DO] ^a Famciclovir AN [EA] ^a Famciclovir generichealth 250 [GQ] ^a Favic 250 [RW]

■ FAMCICLOVIR

Note This drug is only effective if commenced within 72 hours of onset of rash.

Note Famciclovir 250 mg is not PBS-subsidised for chickenpox or herpes simplex infections other than genital herpes.

Note No applications for repeats will be authorised.

Authority required (STREAMLINED)

5951

Herpes zoster

Clinical criteria:

- The treatment must be administered within 72 hours of the onset of the rash.

famciclovir 250 mg tablet, 21

8002E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	34.84	36.13	^a APO-Famciclovir [TX] ^a Ezovir [AF] ^a Famciclovir-GA [ED] ^a Famvir [IX]	^a Auro-Famciclovir 250 [DO] ^a Famciclovir AN [EA] ^a Famciclovir generichealth 250 [GQ] ^a Favic 250 [RW]

■ FAMCICLOVIR

Note This drug is only effective if commenced within 72 hours of onset of rash.

Note Famciclovir 500 mg is not PBS-subsidised for chickenpox.

Note Famciclovir 500 mg is not PBS-subsidised for herpes zoster, genital herpes or other herpes simplex infections in immunocompetent patients.

Note No applications for repeats will be authorised.

Authority required (STREAMLINED)

5943

Herpes zoster

Clinical criteria:

- Patient must be immunocompromised, **AND**
- The treatment must be administered within 72 hours of the onset of the rash.

famciclovir 500 mg tablet, 30

8897G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	44.62	41.30	^a APO-Famciclovir [TX] ^a Famciclovir AN [EA] ^a Favic 500 [RW]	^a Auro-Famciclovir 500 [DO] ^a Famvir [IX]

■ FAMCICLOVIR

Note Famciclovir 500 mg is not PBS-subsidised for chickenpox.

Note Famciclovir 500 mg is not PBS-subsidised for herpes zoster, genital herpes or other herpes simplex infections in immunocompetent patients.

Authority required (STREAMLINED)

5954

Recurrent moderate to severe genital herpes

Treatment Phase: Episodic treatment or suppressive therapy

Clinical criteria:

- Patient must be immunocompromised.
- Microbiological confirmation of diagnosis [viral culture, antigen detection or nucleic acid amplification by polymerase chain reaction (PCR)] is desirable but need not delay treatment.

Authority required (STREAMLINED)

5947

Recurrent moderate to severe oral or labial herpes

Treatment Phase: Episodic treatment

Clinical criteria:

- Patient must have HIV infection, **AND**
- Patient must have a CD4 cell count of less than 500 million per litre.

Microbiological confirmation of diagnosis [viral culture, antigen detection or nucleic acid amplification by polymerase chain reaction (PCR)] is desirable but need not delay treatment.

Authority required (STREAMLINED)

5948

Recurrent moderate to severe oral or labial herpes

Treatment Phase: Suppressive therapy

Clinical criteria:

- Patient must have HIV infection, **AND**
- Patient must have CD4 cell counts of less than 150 million per litre.

Microbiological confirmation of diagnosis [viral culture, antigen detection or nucleic acid amplification by polymerase chain reaction (PCR)] is desirable but need not delay treatment.

Authority required (STREAMLINED)

5949

Recurrent moderate to severe oral or labial herpes

Treatment Phase: Suppressive therapy

Clinical criteria:

- Patient must have HIV infection, **AND**
- Patient must present with other opportunistic infections or AIDS defining tumours.

Microbiological confirmation of diagnosis [viral culture, antigen detection or nucleic acid amplification by polymerase chain reaction (PCR)] is desirable but need not delay treatment.

famciclovir 500 mg tablet, 56

8896F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	72.88	41.30	^a APO-Famciclovir [TX] ^a Ezovir [AF] ^a Famciclovir-GA [ED] ^a Famvir [IX]	^a Auro-Famciclovir 500 [DO] ^a Famciclovir AN [EA] ^a Famciclovir generichealth 500 [GQ] ^a Favic 500 [RW]

■ VALACICLOVIR

Note Valaciclovir 500 mg is not PBS-subsidised for chickenpox or herpes simplex infections other than genital herpes.

Authority required (STREAMLINED)

5940

Recurrent moderate to severe genital herpes

Treatment Phase: Suppressive therapy

Microbiological confirmation of diagnosis [viral culture, antigen detection or nucleic acid amplification by polymerase chain reaction (PCR)] is desirable but need not delay treatment.

valaciclovir 500 mg tablet, 30

5480K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	26.28	27.57	^a APO-Valaciclovir [TX] ^a Valtrex [AF] ^a Valaciclovir APOTEX [GX] ^a Valaciclovir RBX [RA] ^a Valaciclovir SZ [HX] ^a Zelitrex [RF]	^a Shilova 500 [DO] ^a Valaciclovir AN [EA] ^a Valaciclovir generichealth [GQ] ^a Valaciclovir Sandoz [SZ] ^a Valacor 500 [CR]
			^b 2.26	28.54	27.57	^a Valtrex [RW]	

■ VALACICLOVIR

Note Valaciclovir 500 mg is not PBS-subsidised for chickenpox or herpes simplex infections other than genital herpes.

Authority required (STREAMLINED)

5961

Recurrent moderate to severe genital herpes

Treatment Phase: Episodic treatment

Microbiological confirmation of diagnosis [viral culture, antigen detection or nucleic acid amplification by polymerase chain reaction (PCR)] is desirable but need not delay treatment.

valaciclovir 500 mg tablet, 30

8134D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	26.28	27.57	^a APO-Valaciclovir [TX] ^a Valtrex [AF] ^a Valaciclovir APOTEX [GX] ^a Valaciclovir RBX [RA] ^a Valaciclovir SZ [HX] ^a Zelitrex [RF]	^a Shilova 500 [DO] ^a Valaciclovir AN [EA] ^a Valaciclovir generichealth [GQ] ^a Valaciclovir Sandoz [SZ] ^a Valacor 500 [CR]
			^b 2.26	28.54	27.57	^a Valtrex [RW]	

■ VALACICLOVIR

Note Valaciclovir 500 mg is not PBS-subsidised for chickenpox or herpes simplex infections other than genital herpes.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

5960

Initial moderate to severe genital herpes

Microbiological confirmation of diagnosis [viral culture, antigen detection or nucleic acid amplification by polymerase chain reaction (PCR)] is desirable but need not delay treatment.

valaciclovir 500 mg tablet, 10

8133C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	2	..	..	*21.52	22.81	^a APO-Valaciclovir [TX] ^a Valaciclovir AN [EA] ^a Valaciclovir Sandoz [SZ]	^a Valtrex [AF] ^a Valaciclovir APOTEX [GX] ^a Zelitrex [RF]
			^b 2.26	*23.78	22.81	^a Valtrex [RW]	

■ VALACICLOVIR

Note This drug is only effective if commenced within 72 hours of onset of rash.

Note Valaciclovir 500 mg is not PBS-subsidised for chickenpox or herpes simplex infections other than genital herpes.

Note No applications for repeats will be authorised.

Authority required (STREAMLINED)

5962

Herpes zoster

Clinical criteria:

- The treatment must be administered within 72 hours of the onset of the rash.

Authority required (STREAMLINED)

5968

Herpes zoster ophthalmicus

valaciclovir 500 mg tablet, 42

8064K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	31.98	33.27	^a APO-Valaciclovir [TX] ^a Valaciclovir AN [EA] ^a Valaciclovir generichealth [GQ] ^a Valaciclovir Sandoz [SZ]	^a Valtrex [AF] ^a Valaciclovir APOTEX [GX] ^a Valaciclovir RBX [RA] ^a Valacor 500 [CR]

^a Zelitrex [RF]
^b 2.26 34.24 33.27 ^a Valtrex [RW]

Antivirals for treatment of HCV infections**■ GLECAPREVIR + PIBRENTASVIR**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Chronic hepatitis C infection

Clinical criteria:

- Patient must meet the criteria set out in the General Statement for Drugs for the Treatment of Hepatitis C, **AND**
- Patient must be taking this drug as part of a regimen set out in the matrix in the General Statement for Drugs for the Treatment of Hepatitis C, based on the hepatitis C virus genotype, patient treatment history and cirrhotic status, **AND**
- The treatment must be limited to a maximum duration of 16 weeks.

The application must include details of the prior treatment regimen containing an NS5A inhibitor.

glecaprevir 100 mg + pibrentasvir 40 mg tablet, 84

11344C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	3	..	18827.83	41.30	Maviret [VE]

■ GLECAPREVIR + PIBRENTASVIR

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Chronic hepatitis C infection

Clinical criteria:

- Patient must meet the criteria set out in the General Statement for Drugs for the Treatment of Hepatitis C, **AND**
- Patient must be taking this drug as part of a regimen set out in the matrix in the General Statement for Drugs for the Treatment of Hepatitis C, based on the hepatitis C virus genotype, patient treatment history and cirrhotic status, **AND**
- The treatment must be limited to a maximum duration of 8 weeks.

glecaprevir 100 mg + pibrentasvir 40 mg tablet, 84

11353M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	18827.83	41.30	Maviret [VE]

■ GLECAPREVIR + PIBRENTASVIR

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Chronic hepatitis C infection

Clinical criteria:

- Patient must meet the criteria set out in the General Statement for Drugs for the Treatment of Hepatitis C, **AND**
- Patient must be taking this drug as part of a regimen set out in the matrix in the General Statement for Drugs for the Treatment of Hepatitis C, based on the hepatitis C virus genotype, patient treatment history and cirrhotic status, **AND**
- The treatment must be limited to a maximum duration of 12 weeks.

glecaprevir 100 mg + pibrentasvir 40 mg tablet, 84

11354N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	18827.83	41.30	Maviret [VE]

■ LEDIPASVIR + SOFOSBUVIR

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Chronic hepatitis C infection

Clinical criteria:

- Patient must meet the criteria set out in the General Statement for Drugs for the Treatment of Hepatitis C, **AND**
- Patient must be taking this drug as part of a regimen set out in the matrix in the General Statement for Drugs for the Treatment of Hepatitis C, based on the hepatitis C virus genotype, patient treatment history and cirrhotic status, **AND**
- The treatment must be limited to a maximum duration of 12 weeks.

ledipasvir 90 mg + sofosbuvir 400 mg tablet, 28

10628J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	12036.16	41.30	Harvoni [GI]

▪ LEDIPASVIR + SOFOSBUVIR

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Chronic hepatitis C infection

Clinical criteria:

- Patient must meet the criteria set out in the General Statement for Drugs for the Treatment of Hepatitis C, **AND**
- Patient must be taking this drug as part of a regimen set out in the matrix in the General Statement for Drugs for the Treatment of Hepatitis C, based on the hepatitis C virus genotype, patient treatment history and cirrhotic status, **AND**
- The treatment must be limited to a maximum duration of 8 weeks.

ledipasvir 90 mg + sofosbuvir 400 mg tablet, 28

10668L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	12036.16	41.30	Harvoni [GI]

▪ LEDIPASVIR + SOFOSBUVIR

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Chronic hepatitis C infection

Clinical criteria:

- Patient must meet the criteria set out in the General Statement for Drugs for the Treatment of Hepatitis C, **AND**
- Patient must be taking this drug as part of a regimen set out in the matrix in the General Statement for Drugs for the Treatment of Hepatitis C, based on the hepatitis C virus genotype, patient treatment history and cirrhotic status, **AND**
- The treatment must be limited to a maximum duration of 24 weeks.

ledipasvir 90 mg + sofosbuvir 400 mg tablet, 28

10670N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	12036.16	41.30	Harvoni [GI]

▪ RIBAVIRIN

Caution Ribavirin is a category X drug and must not be given to pregnant women. Pregnancy in female patients or in the partners of male patients must be avoided during treatment and during the 6 months period after cessation of treatment.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Chronic hepatitis C infection

Clinical criteria:

- Patient must meet the criteria set out in the General Statement for Drugs for the Treatment of Hepatitis C, **AND**
- Patient must be taking this drug as part of a regimen set out in the matrix in the General Statement for Drugs for the Treatment of Hepatitis C, based on the hepatitis C virus genotype, patient treatment history and cirrhotic status, **AND**
- The treatment must be limited to a maximum duration of 12 weeks.

Population criteria:

- Patient must not be pregnant or breastfeeding. Female partners of male patients must not be pregnant. Patients and their partners must each be using an effective form of contraception if of child-bearing age.

ribavirin 600 mg tablet, 28

10665H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	457.47	41.30	lbavyr [IX]

ribavirin 400 mg tablet, 28

10647J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	307.32	41.30	lbavyr [IX]

▪ RIBAVIRIN

Caution Ribavirin is a category X drug and must not be given to pregnant women. Pregnancy in female patients or in the partners of male patients must be avoided during treatment and during the 6 months period after cessation of treatment.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Chronic hepatitis C infection

Clinical criteria:

- Patient must meet the criteria set out in the General Statement for Drugs for the Treatment of Hepatitis C, **AND**
- Patient must be taking this drug as part of a regimen set out in the matrix in the General Statement for Drugs for the Treatment of Hepatitis C, based on the hepatitis C virus genotype, patient treatment history and cirrhotic status, **AND**
- The treatment must be limited to a maximum duration of 24 weeks.

Population criteria:

- Patient must not be pregnant or breastfeeding. Female partners of male patients must not be pregnant. Patients and their partners must each be using an effective form of contraception if of child-bearing age.

ribavirin 600 mg tablet, 28

10666J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	457.47	41.30	Ibavyr [IX]

ribavirin 400 mg tablet, 28

10673R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	307.32	41.30	Ibavyr [IX]

▪ **SOFOSBUVIR + VELPATASVIR**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Chronic hepatitis C infection

Clinical criteria:

- Patient must meet the criteria set out in the General Statement for Drugs for the Treatment of Hepatitis C, **AND**
- Patient must be taking this drug as part of a regimen set out in the matrix in the General Statement for Drugs for the Treatment of Hepatitis C, based on the hepatitis C virus genotype, patient treatment history and cirrhotic status, **AND**
- The treatment must be limited to a maximum duration of 12 weeks.

sofosbuvir 400 mg + velpatasvir 100 mg tablet, 28

11147Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	12661.16	41.30	Epclusa [GI]

▪ **SOFOSBUVIR + VELPATASVIR + VOXILAPREVIR**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Chronic hepatitis C infection

Clinical criteria:

- Patient must meet the criteria set out in the General Statement for Drugs for the Treatment of Hepatitis C, **AND**
- Patient must be taking this drug as part of a regimen set out in the matrix in the General Statement for Drugs for the Treatment of Hepatitis C, based on the hepatitis C virus genotype, patient treatment history and cirrhotic status, **AND**
- The treatment must be limited to a maximum duration of 12 weeks.

The application must include details of the prior treatment regimen containing an NS5A inhibitor.

sofosbuvir 400 mg + velpatasvir 100 mg + voxilaprevir 100 mg tablet, 28

11658N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	12661.16	41.30	Vosevi [GI]

Antivirals for treatment of HIV infections, combinations

▪ **TENOFOVIR DISOPROXIL + EMTRICITABINE**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Pharmaceutical benefits that have the forms tenofovir disoproxil phosphate 291 mg with emtricitabine 200 mg tablet, tenofovir disoproxil maleate 300 mg with emtricitabine 200 mg tablet, and tenofovir disoproxil fumarate 300 mg with emtricitabine 200 mg tablet are equivalent for the purposes of substitution.

Restricted benefit

Pre-exposure prophylaxis (PrEP) against human immunodeficiency virus (HIV) infection

Clinical criteria:

- Patient must have at least one of the following prior to having the latest PBS-subsidised prescription issued: (i) a negative HIV test result no older than 4 weeks, (ii) evidence that an HIV test has been conducted, but the result is still forthcoming.

tenofovir disoproxil maleate 300 mg + emtricitabine 200 mg tablet, 30

11296M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	69.88	41.30	^a Tenofovir Disoproxil Emtricitabine Mylan 300/200 [AF]

tenofovir disoproxil phosphate 291 mg + emtricitabine 200 mg tablet, 30

11306C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	69.88	41.30	^a Tenofovir EMT GH [GQ]

tenofovir disoproxil fumarate 300 mg + emtricitabine 200 mg tablet, 30

11276L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	69.88	41.30	^a Tenofovir/Emtricitabine 300/200 APOTEX [TX]

■ **VACCINES****BACTERIAL VACCINES***Tetanus vaccines*■ **DIPHTHERIA + TETANUS VACCINE**

Note For immunisation of adults and children aged greater than or equal to 8 years.

diphtheria 2 units + tetanus 20 units vaccine injection, 5 x 0.5 mL syringes

8783G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	70.46	41.30	ADT Booster [CS]

■ **ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS**■ **ANTINEOPLASTIC AGENTS****ALKYLATING AGENTS***Nitrogen mustard analogues*■ **CHLORAMBUCIL****chlorambucil 2 mg tablet, 25**

1163F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	4	2	..	*135.30	41.30	Leukeran [AS]

■ **CYCLOPHOSPHAMIDE****cyclophosphamide 50 mg tablet, 50**

1266P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	155.65	41.30	Cyclonex [ZX]

■ **MELPHALAN****melphalan 2 mg tablet, 25**

2547C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	63.97	41.30	Alkeran [AS]

Alkyl sulfonates■ **BUSULFAN****busulfan 2 mg tablet, 100**

1128J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	78.05	41.30	Myleran [AS]

Nitrosoureas■ **CARMUSTINE**

Note Carmustine is not PBS-subsidised for use in conjunction with PBS-subsidised temozolomide.

Restricted benefit

Glioblastoma multiforme

Clinical criteria:

- The condition must be suspected or confirmed at the time of initial surgery.

carmustine 7.7 mg implant, 8

8898H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	14288.59	41.30	Gliadel [EI]

Other alkylating agents

■ TEMOZOLOMIDE

temozolomide 180 mg capsule, 5

2438H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	208.10	41.30	^a APO-Temozolomide [TX] ^a Temolide [JU]	^a Temodal [MK] ^a Temozolomide Juno [JX]

temozolomide 100 mg capsule, 5

8380C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	127.42	41.30	^a APO-Temozolomide [TX] ^a Temodal [MK] ^a Temozolomide Amneal [JO]	^a Temizole 100 [AL] ^a Temozolomide Alphapharm [AF] ^a Temozolomide Juno [JX]

temozolomide 20 mg capsule, 5

8379B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	38.78	40.07	^a APO-Temozolomide [TX] ^a Temozolomide Alphapharm [AF]	^a Temizole 20 [AL] ^a Temozolomide Juno [JX]

temozolomide 250 mg capsule, 5

8381D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	293.10	41.30	^a APO-Temozolomide [TX] ^a Temodal [MK] ^a Temozolomide Juno [JX]	^a Temizole 250 [AL] ^a Temozolomide Alphapharm [AF]

temozolomide 140 mg capsule, 5

9362R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	173.07	41.30	^a APO-Temozolomide [TX] ^a Temodal [MK] ^a Temozolomide Juno [JX]	^a Temizole 140 [AL] ^a Temozolomide Alphapharm [AF]

temozolomide 5 mg capsule, 5

8378Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	21.13	22.42	^a APO-Temozolomide [TX] ^a Temozolomide Alphapharm [AF]	^a Temizole 5 [AL] ^a Temozolomide Juno [JX]

■ TEMOZOLOMIDE

Note Temozolomide is not PBS-subsidised for use in conjunction with PBS-subsidised carmustine.

Note No increase in the maximum number of repeats may be authorised.

Restricted benefit

Glioblastoma multiforme

Treatment criteria:

- Patient must be undergoing concomitant radiotherapy.

temozolomide 180 mg capsule, 5

10062N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	3	2	..	*610.26	41.30	^a APO-Temozolomide [TX] ^a Temolide [JU]	^a Temodal [MK] ^a Temozolomide Juno [JX]

temozolomide 100 mg capsule, 5

8821G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	3	2	..	*368.22	41.30	^a APO-Temozolomide [TX] ^a Temodal [MK] ^a Temozolomide Amneal [JO]	^a Temizole 100 [AL] ^a Temozolomide Alphapharm [AF] ^a Temozolomide Juno [JX]

temozolomide 20 mg capsule, 5

8820F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	3	2	..	*92.31	41.30	^a APO-Temozolomide [TX] ^a Temozolomide Alphapharm [AF]	^a Temizole 20 [AL] ^a Temozolomide Juno [JX]

temozolomide 140 mg capsule, 5

9361Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	3	2	..	*505.17	41.30	^a APO-Temozolomide [TX] ^a Temodal [MK] ^a Temozolomide Juno [JX]	^a Temizole 140 [AL] ^a Temozolomide Alphapharm [AF]

temozolomide 5 mg capsule, 5

8819E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	3	2	..	*39.36	40.65	^a APO-Temozolomide [TX] ^a Temozolomide Alphapharm [AF]	^a Temizole 5 [AL] ^a Temozolomide Juno [JX]

ANTIMETABOLITES
Folic acid analogues
METHOTREXATE
methotrexate 5 mg/2 mL injection, 5 x 2 mL vials

2396D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	40.87	41.30	DBL Methotrexate [PF]

methotrexate 2.5 mg tablet, 30

1622J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	17.85	19.14	Methoblastin [PF]

methotrexate 10 mg tablet, 15

2272N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	24.12	25.41	Methoblastin [PF]

METHOTREXATE
Restricted benefit

Patients requiring doses greater than 20 mg per week

methotrexate 10 mg tablet, 50

1623K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	52.34	41.30	Methoblastin [PF]

METHOTREXATE

Note For item codes 2395C and 1818Q, pharmaceutical benefits that have the form injection 50 mg in 2 mL are equivalent for the purposes of substitution.

methotrexate 50 mg/2 mL injection, 5 x 2 mL vials

2395C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	39.44	40.73	^a DBL Methotrexate [PF]

methotrexate 50 mg/2 mL injection, 2 mL vial

1818Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	5	5	..	*39.44	40.73	^a Methotrexate Accord [OD]

Purine analogues
FLUDARABINE
fludarabine phosphate 10 mg tablet, 20

9184J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	937.40	41.30	Fludara [GZ]

MERCAPTOPURINE
mercaptopurine monohydrate 50 mg tablet, 25

1598D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	4	2	..	*188.34	41.30	^a MERCAPTOPURINE-LINK [LM]	^a Purinethol [AS]

mercaptopurine monohydrate 20 mg/mL oral liquid, 100 mL

10214N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	2	..	380.24	41.30	Allmercap [LM]

TIOGUANINE
tioguanine 40 mg tablet, 25

1233X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	193.32	41.30	Lanvis [AS]

Pyrimidine analogues

■ CAPECITABINE

capecitabine 500 mg tablet, 120

8362D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	2	..	79.21	41.30	^a Capecitabine Alphapharm [AF]	^a Capecitabine-DRLA [RZ]
						^a Capecitabine Sandoz [SZ]	^a Xelabine [AL]

capecitabine 150 mg tablet, 60

8361C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	22.86	24.15	Capecitabine-DRLA [RZ]

■ TRIFLURIDINE + TIPIRACIL

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

10309

Metastatic colorectal cancer

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have a WHO performance status of 1 or less, **AND**
 - Patient must have previously received treatment with fluoropyrimidine, oxaliplatin, irinotecan-based chemotherapies, an anti-vascular endothelial growth factor (anti-VEGF) agent and an anti-epidermal growth factor receptor (anti-EGFR) agent for this condition; OR
 - Patient must not be a suitable candidate for treatment with fluoropyrimidine, oxaliplatin, irinotecan-based chemotherapies, an anti-VEGF agent and an anti-EGFR agent for this condition, **AND**
 - The treatment must be the sole PBS-subsidised therapy for this condition.
- The patient's WHO performance status and body weight must be documented in the patient's medical records at the time the treatment cycle is initiated.

Authority required (STREAMLINED)

8183

Metastatic colorectal cancer

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously been treated with PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not develop progressive disease whilst receiving PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

trifluridine 20 mg + tipiracil 8.19 mg tablet, 20

11524M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	4	2	..	*3932.62	41.30	Lonsurf 20/8.19 [SE]

trifluridine 15 mg + tipiracil 6.14 mg tablet, 20

11507P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	2	..	*2282.58	41.30	Lonsurf 15/6.14 [SE]

■ TRIFLURIDINE + TIPIRACIL

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

10252

Metastatic (Stage IV) adenocarcinoma of the stomach or gastro-oesophageal junction

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have a WHO performance status of 1 or less, **AND**
- Patient must have previously received at least two prior lines of chemotherapy that included a fluoropyrimidine, a platinum and either a taxane or irinotecan, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

The patient's WHO performance status and body weight must be documented in the patient's medical records at the time the treatment cycle is initiated.

Authority required (STREAMLINED)

10310

Metastatic (Stage IV) adenocarcinoma of the stomach or gastro-oesophageal junction

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not develop progressive disease whilst receiving PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

trifluridine 20 mg + tipiracil 8.19 mg tablet, 20

12033H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	4	2	..	*3932.62	41.30	Lonsurf 20/8.19 [SE]

trifluridine 15 mg + tipiracil 6.14 mg tablet, 20

12056M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	2	..	*2282.58	41.30	Lonsurf 15/6.14 [SE]

PLANT ALKALOIDS AND OTHER NATURAL PRODUCTS
Vinca alkaloids and analogues
■ VINORELBINE
Authority required

Advanced breast cancer

Clinical criteria:

- Patient must have failed standard prior therapy, which includes an anthracycline.

Authority required

Locally advanced or metastatic non-small cell lung cancer

vinorelbine 20 mg capsule, 1

9009E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	20	2	..	*1578.54	41.30	Navelbine [FB]

vinorelbine 30 mg capsule, 1

9010F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	16	2	..	*1891.58	41.30	Navelbine [FB]

Podophyllotoxin derivatives
■ ETOPOSIDE
etoposide 100 mg capsule, 10

1389D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	391.65	41.30	Vepesid [LM]

etoposide 50 mg capsule, 20

1396L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	448.57	41.30	Vepesid [LM]

PROTEIN KINASE INHIBITORS
BCR-ABL tyrosine kinase inhibitors
■ DASATINIB

Note The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of tyrosine kinase inhibitors (TKI) agents for all phases of chronic myeloid leukaemia. Where the term TKI agent appears in the following notes and restrictions it refers to dasatinib or nilotinib. Imatinib mesilate is not approved for use in second or third line treatment. Patients are eligible for PBS-subsidised treatment with only one of dasatinib or nilotinib at any one time and must not be receiving concomitant interferon alfa therapy. Eligible patients may only swap between these agents if they have not failed prior PBS-subsidised treatment with that agent. Nilotinib is not approved for patients in blast crisis. 1. Initial second line treatment From 1 April 2012, under the PBS, a patient will be able to be prescribed either dasatinib or nilotinib within the initial 18 month treatment period as second-line therapy, as long as only one agent is approved at a time and providing the patient did not fail that drug as first-line therapy. During the initial 18 month treatment period, switching between approved second-line agents may only occur for reasons of intolerance, not failure of response. 2. Initial third line treatment Third-line treatment with a TKI can only be approved when imatinib is used for first-line treatment. Patients will only be approved for PBS-subsidised treatment with one third-line agent. From 1 April 2012, under the PBS, a patient will be able to be prescribed either dasatinib or nilotinib providing the patient did not fail that drug as first or second line therapy and for nilotinib the patient is not in blast crisis. 3. Continuing treatment for second and third line treatment All continuing applications are to be written and must include a pathology report demonstrating the patient has responded to PBS-subsidised treatment as follows: (i) within 18 months of the commencement of treatment, at which time patients in whom a major cytogenetic response or peripheral blood BCR-ABL level of less than 1% has been demonstrated may receive authorisation for a further 12 months of treatment; and (ii) at no greater than 12 month intervals thereafter, to demonstrate that the major cytogenetic response or peripheral blood BCR-ABL level of less than 1% has been sustained. During second line continuing treatment beyond the initial 18 month treatment period, switching between approved second line TKI agents may only occur for reason of intolerance. Where there is failure of response, switching may only occur through application for prescription of a third line agent. 4. Authority approval requirements. Response criteria to initial treatment with dasatinib or nilotinib: For the purposes of assessing response to PBS-subsidised treatment with dasatinib or nilotinib, either cytogenetic analysis indicating the

number of Philadelphia positive [t (9;22)] cells in the bone marrow measured by standard karyotyping, or quantitative PCR indicating the relative level of BCR-ABL transcript in the peripheral blood using the international scale, must be submitted. For bone marrow analyses, where the standard karyotyping is not informative for technical reasons, a cytogenetic analysis performed on the bone marrow by the use of fluorescence in situ hybridisation (FISH) with BCR-ABL specific probe must be submitted. The cytogenetic or peripheral blood quantitative PCR analyses must be submitted within 18 months of the commencement of treatment with dasatinib or nilotinib (patients in whom a major cytogenetic response or peripheral blood BCR-ABL level of less than 1% is demonstrable by 18 months are eligible to receive continuing treatment with that agent).5. Definitions of response. A major cytogenetic response is defined as less than 35% Philadelphia positive bone marrow cells. A peripheral blood BCR-ABL level of less than 1% on the international scale (Blood 108: 28-37, 2006) also indicates a response, at least the biological equivalent of a major cytogenetic response.6. Definitions of loss of response. Loss of a previously documented major cytogenetic response (demonstrated by the presence of greater than 35% Ph positive cells on bone marrow biopsy), during ongoing tyrosine kinase inhibitor (TKI) therapy. Loss of a previously demonstrated molecular response (demonstrated by peripheral blood BCR-ABL levels increasing consecutively in value by at least 5 fold to a level of greater than 0.1% confirmed on a subsequent test), during ongoing tyrosine kinase inhibitor therapy.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Chronic Myeloid Leukaemia (CML)

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must not have failed PBS-subsidised first line treatment with this drug for this condition, **AND**
- Patient must have failed an adequate trial of PBS-subsidised first line treatment with imatinib for this condition; OR
- Patient must have failed an adequate trial of PBS-subsidised first line treatment with nilotinib for this condition; OR
- Patient must have experienced intolerance, not a failure of response, to PBS-subsidised second line treatment with nilotinib for this condition, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Failure of an adequate trial of imatinib or nilotinib is defined as: (i) Lack of response to initial imatinib or nilotinib therapy, defined as either: - failure to achieve a haematological response after a minimum of 3 months therapy with imatinib or nilotinib for patients initially treated in chronic phase; or - failure to achieve any cytogenetic response after a minimum of 6 months therapy with imatinib or nilotinib for patients initially treated in chronic phase as demonstrated on bone marrow biopsy by presence of greater than 95% Philadelphia chromosome positive cells; or - failure to achieve a major cytogenetic response or a peripheral blood BCR-ABL level of less than 1% after a minimum of 12 months therapy with imatinib or nilotinib; OR (ii) Loss of a previously documented major cytogenetic response (demonstrated by the presence of greater than 35% Ph positive cells on bone marrow biopsy), during ongoing imatinib or nilotinib therapy; OR (iii) Loss of a previously demonstrated molecular response (demonstrated by peripheral blood BCR-ABL levels increasing consecutively in value by at least 5 fold to a level of greater than 0.1% confirmed on a subsequent test), during ongoing imatinib or nilotinib therapy; OR (iv) Development of accelerated phase or blast crisis in a patient previously prescribed imatinib or nilotinib for any phase of chronic myeloid leukaemia.

Accelerated phase is defined by the presence of 1 or more of the following: (1) Percentage of blasts in the peripheral blood or bone marrow greater than or equal to 15% but less than 30%; or (2) Percentage of blasts plus promyelocytes in the peripheral blood or bone marrow greater than or equal to 30%, provided that blast count is less than 30%; or (3) Peripheral basophils greater than or equal to 20%; or (4) Progressive splenomegaly to a size greater than or equal to 10 cm below the left costal margin to be confirmed on 2 occasions at least 4 weeks apart, or a greater than or equal to 50% increase in size below the left costal margin over 4 weeks; or (5) Karyotypic evolution (chromosomal abnormalities in addition to a single Philadelphia chromosome); OR Blast crisis is defined as either: (1) Percentage of blasts in the peripheral blood or bone marrow greater than or equal to 30%; or (2) Extramedullary involvement other than spleen and liver; OR (v) Disease progression (defined as a greater than or equal to 50% increase in peripheral white blood cell count, blast count, basophils or platelets) during first-line imatinib or nilotinib therapy in patients with accelerated phase or blast crisis chronic myeloid leukaemia. Patients should be commenced on a dose of dasatinib of at least 100 mg (base) daily. Continuing therapy is dependent on patients demonstrating a major cytogenetic response to dasatinib therapy or a peripheral blood BCR-ABL level of less than 1% within 18 months and thereafter at 12 monthly intervals.

Applications for authorisation must be in writing and must include: (a) a completed authority prescription form; and (b) a completed Chronic Myeloid Leukaemia - Second and Third Line - Supporting Information Form; and (c) a signed patient acknowledgement; and (d) a bone marrow biopsy pathology report demonstrating the patient has active chronic myeloid leukaemia, either manifest as cytogenetic evidence of the Philadelphia chromosome, or RT-PCR level of BCR-ABL transcript greater than 0.1% on the international scale. (The date of the relevant pathology report needs to be provided); and (e) where there has been a loss of response to imatinib or nilotinib, a copy of the current confirming pathology report(s) from an Approved Pathology Authority or details of the dates of assessment in the case of progressive splenomegaly or extramedullary involvement

Authority required

Chronic Myeloid Leukaemia (CML)

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have received initial PBS- subsidised second line treatment with this drug for this condition; OR

- Patient must have experienced intolerance, not a failure to respond, to PBS-subsidised second line treatment with nilotinib for this condition, **AND**
- Patient must have demonstrated a major cytogenetic response in the preceding 18 months and thereafter at 12 monthly intervals; OR
- Patient must have achieved a peripheral blood level of BCR-ABL of less than 1% in the preceding 18 months and thereafter at 12 monthly intervals, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Applications for authorisation must be in writing and must include: (1) a completed authority prescription form; and (2) a completed Chronic Myeloid Leukaemia - Second and Third Line - Application Form for continuing treatment; and (3) demonstration of continued response to treatment as evidenced by either: (a) major cytogenetic response [see Note explaining definitions of response]. Where this has been supplied within the previous 12 months (or 18 months for the initial supply), only the date of the relevant pathology report need be provided; or (b) a peripheral blood level of BCR-ABL of less than 1% on the international scale on the international scale [see Note explaining definitions of response]. Where this has been supplied within the previous 12 months (or 18 months for the initial supply), only the date of the relevant pathology report need be provided.

dasatinib 70 mg tablet, 60

2485T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	5255.21	41.30	Sprycel [BQ]

dasatinib 50 mg tablet, 60

2482P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	4275.68	41.30	Sprycel [BQ]

dasatinib 100 mg tablet, 30

9342Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	4275.68	41.30	Sprycel [BQ]

dasatinib 20 mg tablet, 60

2478K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	2659.21	41.30	Sprycel [BQ]

■ DASATINIB

Note The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of tyrosine kinase inhibitors (TKI) agents for the chronic phase of chronic myeloid leukaemia. Where the term TKI agent appears in the following notes and restrictions it refers to imatinib mesilate, dasatinib or nilotinib.

Patients are eligible for PBS-subsidised treatment with only one TKI agent at any one time and must not be receiving concomitant interferon alfa therapy. Eligible patients may only swap between TKI agents if they have not failed prior PBS-subsidised treatment with that agent.

1. Initial First-line treatment From 1 April 2012, under the PBS, a patient will be able to be prescribed any of imatinib mesilate, dasatinib or nilotinib within the initial 18 month treatment period, as long as only one agent is used at a time and providing the patient has not failed to respond to any one of these TKIs. During the initial 18 month treatment period, switching between approved first-line agents may only occur for reasons of intolerance, not failure to respond

2. Continuing First-line treatment

Patients must maintain a major cytogenetic response or have a peripheral blood BCR-ABL of less than 1% to receive continuing therapy.

For imatinib mesilate:

First continuing applications may be made by telephoning the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

For dasatinib or nilotinib:

First continuing applications are to be written and must include a pathology report demonstrating the patient has responded to the initial course of treatment.

Second and subsequent authority applications for continuing therapy may be made by telephoning the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

During continuing therapy beyond the initial 18 month treatment period, switching between approved first-line agents may only occur for reason of intolerance. Where there is failure to respond, switching may only occur through application for prescription of second-line agents. Where a patient has previously received PBS-subsidised treatment with imatinib mesilate, dasatinib or nilotinib no approval will be granted for PBS-subsidised re-treatment in the chronic phase of chronic myeloid leukaemia, where that patient has at any time failed to meet the response criteria whilst on that TKI agent.

3. Authority approval requirements. Response criteria to initial first-line treatment with imatinib mesilate, dasatinib or nilotinib: For the purposes of assessing response to PBS-subsidised treatment with imatinib mesilate, dasatinib or nilotinib either cytogenetic analysis indicating the number of Philadelphia positive [t (9;22)] cells in the bone marrow measured by standard karyotyping, or quantitative PCR indicating the relative level of BCR-ABL transcript in the peripheral blood using the international scale, must be submitted. For bone marrow analyses, where the standard karyotyping is not informative for technical reasons, a cytogenetic analysis performed on the bone marrow by the use of fluorescence in situ hybridisation (FISH) with BCR-ABL specific probe must be submitted. The cytogenetic or peripheral blood quantitative PCR analyses must be submitted within 18 months of the commencement of treatment with dasatinib or nilotinib (patients in whom a major cytogenetic response or peripheral blood BCR-ABL level of less than 1% is demonstrable by 18 months are eligible to receive continuing treatment with that agent).

4. Definitions of response

A major cytogenetic response is defined as less than 35% Philadelphia positive bone marrow cells. A peripheral blood BCR-ABL level of less than 1% on the international scale (Blood 108: 28-37, 2006) also indicates a response, at least the

biological equivalent of a major cytogenetic response.

5. Definitions of loss of response

Loss of a previously documented major cytogenetic response (demonstrated by the presence of greater than 35% Ph positive cells on bone marrow biopsy), during ongoing tyrosine kinase inhibitor (TKI) therapy. Loss of a previously demonstrated molecular response (demonstrated by peripheral blood BCR-ABL levels increasing consecutively in value by at least 5 fold to a level of greater than 0.1% confirmed on a subsequent test), during ongoing tyrosine kinase inhibitor therapy.

Authority required

Chronic Myeloid Leukaemia (CML)

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be a primary diagnosis, **AND**
- The condition must be in the chronic phase, **AND**
- The condition must be expressing the Philadelphia chromosome; OR
- The condition must have the transcript BCR-ABL tyrosine kinase, **AND**
- The treatment must be for first line therapy for this condition, **AND**
- Patient must not have previously experienced a failure to respond to the PBS-subsidised first line treatment with this drug for this condition; OR
- Patient must have experienced intolerance, not a failure to respond, to initial PBS-subsidised treatment with imatinib as a first line therapy for this condition; OR
- Patient must have experienced intolerance, not a failure to respond, to initial PBS-subsidised treatment with nilotinib as a first line therapy for this condition, **AND**
- The treatment must not exceed a total maximum of 18 months of therapy with a PBS-subsidised treatment with a tyrosine kinase inhibitor for this condition, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Applications under this restriction will be limited to provide patients with a maximum of 18 months of therapy with dasatinib, imatinib or nilotinib from the date the first application for initial treatment was approved.

Patients should be commenced on a dose of dasatinib of 100 mg (base) daily. Continuing therapy is dependent on patients demonstrating a response to dasatinib therapy following the initial 18 months of treatment and at 12 monthly intervals thereafter.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Chronic Myeloid Leukaemia - Chronic Phase, First Line - Supporting Information form; and
- (3) a pathology cytogenetic report conducted on peripheral blood or bone marrow supporting the diagnosis of chronic myeloid leukaemia to confirm eligibility for treatment, or a qualitative PCR report documenting the presence of the BCR-ABL transcript in either peripheral blood or bone marrow.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Chronic Myeloid Leukaemia (CML)

Treatment Phase: First continuing treatment

Clinical criteria:

- The condition must be in the chronic phase, **AND**
- Patient must have received initial PBS-subsidised first line treatment with this drug for this condition; OR
- Patient must have experienced intolerance, not a failure to respond, to first continuing PBS-subsidised treatment with imatinib as a first line therapy for this condition; OR
- Patient must have experienced intolerance, not a failure to respond, to first continuing PBS-subsidised treatment with nilotinib as a first line therapy for this condition, **AND**
- Patient must have demonstrated a major cytogenetic response; OR
- Patient must have demonstrated a peripheral blood level of BCR-ABL of less than 1%, **AND**
- The treatment must not exceed a total maximum of 24 weeks of therapy with a PBS-subsidised treatment with a tyrosine kinase inhibitor for this condition under this restriction, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

First continuing applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) demonstration of continued response to treatment as evidenced by either:
 - (a) a major cytogenetic response [see Note explaining requirements]; or

(b) a peripheral blood level of BCR-ABL of less than 1% on the international scale [see Note explaining requirements]. Where this has been supplied within the previous 12 months, only the date of the relevant pathology report need be provided.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Chronic Myeloid Leukaemia (CML)

Treatment Phase: Subsequent continuing treatment

Clinical criteria:

- The condition must be in the chronic phase, **AND**
- Patient must have received the First continuing PBS-subsidised treatment with this drug as a first line therapy for this condition; OR
- Patient must have experienced intolerance, not a failure to respond, to subsequent continuing PBS-subsidised treatment with imatinib as a first line therapy for this condition; OR
- Patient must have experienced intolerance, not a failure to respond, to subsequent continuing PBS-subsidised treatment with nilotinib as a first line therapy for this condition, **AND**
- Patient must have maintained a major cytogenetic response; OR
- Patient must have maintained a peripheral blood level of BCR-ABL of less than 1%, **AND**
- The treatment must not exceed a total maximum of 24 weeks of therapy with a PBS-subsidised treatment with a tyrosine kinase inhibitor for this condition under this restriction, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Note Authority applications for continuing treatment may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

dasatinib 70 mg tablet, 60

1415L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	5255.21	41.30	Sprycel [BQ]

dasatinib 50 mg tablet, 60

1381Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	4275.68	41.30	Sprycel [BQ]

dasatinib 100 mg tablet, 30

1416M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	4275.68	41.30	Sprycel [BQ]

dasatinib 20 mg tablet, 60

1354G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	2659.21	41.30	Sprycel [BQ]

▪ **DASATINIB**

Note No increase in the maximum number of repeats may be authorised.

Authority required

Acute lymphoblastic leukaemia

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be expressing the Philadelphia chromosome; OR
- The condition must have the transcript BCR-ABL, **AND**
- Patient must have failed treatment with chemotherapy, **AND**
- Patient must have failed treatment with imatinib, **AND**
- Patient must have failed an allogeneic haemopoietic stem cell transplantation if applicable.

Failure of treatment is defined as either:

- Failure to achieve a complete morphological and cytogenetic remission after a minimum of 2 months treatment with intensive chemotherapy and imatinib;
 - Morphological or cytogenetic relapse of leukaemia after achieving a complete remission induced by chemotherapy and imatinib;
 - Morphological or cytogenetic relapse or persistence of leukaemia after allogeneic haemopoietic stem cell transplantation.
- Patients must have active leukaemia, as defined by presence on current pathology assessments of either morphological infiltration of the bone marrow (greater than 5% lymphoblasts) or cerebrospinal fluid or other sites; OR the presence of cells

expressing the Philadelphia chromosome on cytogenetic or FISH analysis in the bone marrow of patients in morphological remission.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Acute Lymphoblastic Leukaemia Dasatinib PBS Authority Application - Supporting Information Form; and
- (c) a pathology report demonstrating that the patient has active acute lymphoblastic leukaemia, either manifest as cytogenetic evidence of the Philadelphia chromosome, or morphological evidence of acute lymphoblastic leukaemia plus qualitative RT-PCR evidence of BCR-ABL transcript. The date of the relevant pathology report(s) need(s) to be provided.

Note Dasatinib will only be subsidised for patients with acute lymphoblastic leukaemia who are not receiving concomitant PBS-subsidised imatinib mesilate and who are not appropriate for an allogeneic haemopoietic stem cell transplant.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Acute lymphoblastic leukaemia
Treatment Phase: Continuing treatment

Clinical criteria:

- The condition must be expressing the Philadelphia chromosome; OR
- The condition must have the transcript BCR-ABL, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition as second-line therapy following treatment with imatinib, **AND**
- The condition must not have progressed.

Note Dasatinib will only be subsidised for patients with acute lymphoblastic leukaemia who are not receiving concomitant PBS-subsidised imatinib mesilate and who are not appropriate for an allogeneic haemopoietic stem cell transplant.

Note Authority applications for continuing treatment may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Acute lymphoblastic leukaemia
Treatment Phase: Initial treatment

Clinical criteria:

- Patient must be newly diagnosed, **AND**
- The condition must be expressing the Philadelphia chromosome; OR
- The condition must have the transcript BCR-ABL, **AND**
- The treatment must be for induction and consolidation therapy, **AND**
- The treatment must be in combination with chemotherapy or corticosteroids, **AND**
- Patient must not have previously experienced a failure to respond to the PBS-subsidised first line treatment with this drug for this condition; OR
- Patient must have experienced intolerance, not a failure to respond, to initial PBS-subsidised treatment with imatinib as a first-line therapy for this condition.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Acute Lymphoblastic Leukaemia Dasatinib PBS Authority Application - Supporting Information Form; and
- (c) a pathology cytogenetic report conducted on peripheral blood or bone marrow supporting the diagnosis of acute lymphoblastic leukaemia to confirm eligibility for treatment, with either cytogenetic evidence of the Philadelphia chromosome, or a qualitative PCR report documenting the presence of the BCR-ABL transcript in either peripheral blood or bone marrow. (The date of the relevant pathology report needs to be provided).

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Note Allogeneic stem cell transplantation is the preferred therapy for eligible patients achieving a complete remission of Philadelphia positive acute lymphoblastic leukaemia.

Authority required

Acute lymphoblastic leukaemia
Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition; OR
- Patient must have experienced intolerance, not a failure to respond, to continuing PBS-subsidised treatment with imatinib as a first-line therapy for this condition, **AND**
- The condition must be expressing the Philadelphia chromosome; OR
- The condition must have the transcript BCR-ABL, **AND**
- The treatment must be for maintenance of first complete remission, **AND**
- The treatment must be in combination with chemotherapy or corticosteroids.

Dasatinib and imatinib are available with a lifetime maximum of 24 months for continuing treatment for patients with acute lymphoblastic leukaemia reimbursed through the PBS in this treatment setting.

Note Any queries concerning the arrangements to prescribe this drug beyond 24 months may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note Allogeneic stem cell transplantation is the preferred therapy for eligible patients achieving a complete remission of Philadelphia positive acute lymphoblastic leukaemia.

dasatinib 70 mg tablet, 60

9127J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	5255.21	41.30	Sprycel [BQ]

dasatinib 50 mg tablet, 60

9126H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	4275.68	41.30	Sprycel [BQ]

dasatinib 100 mg tablet, 30

9343R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	4275.68	41.30	Sprycel [BQ]

dasatinib 20 mg tablet, 60

9125G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	2659.21	41.30	Sprycel [BQ]

■ IMATINIB

Note Authority applications for increased quantities/ repeats (where relevant) may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required (STREAMLINED)**9278**

Gastrointestinal stromal tumour
Treatment Phase: Continuing treatment

Clinical criteria:

- The treatment must be adjuvant to complete surgical resection of primary gastrointestinal stromal tumour (GIST), **AND**
- Patient must be at high risk of recurrence following complete surgical resection of primary GIST, **AND**
- The treatment must not exceed a dose of 400 mg per day for a period of 36 months in total (initial plus continuing therapy), **AND**
- Patient must have previously been issued with an authority prescription for imatinib for adjuvant treatment following complete resection of primary GIST.

imatinib 100 mg tablet, 60

11784F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	476.87	41.30	Glivec [AF]

imatinib 400 mg tablet, 30

11788K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	937.83	41.30	Glivec [AF]

■ IMATINIB

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Gastrointestinal stromal tumour
Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be adjuvant to complete surgical resection of primary gastrointestinal stromal tumour (GIST), **AND**
- Patient must be at high risk of recurrence following complete surgical resection of primary GIST, **AND**
- The condition must be histologically confirmed by the detection of CD117 on immunohistochemical staining, **AND**

- The treatment must not exceed a dose of 400 mg per day for a period of 36 months in total (initial plus continuing therapy).

High risk of recurrence is defined as:

Primary GIST greater than 5 cm with a mitotic count of greater than 5/50 high power fields (HPF); or

Primary GIST greater than 10 cm with any mitotic rate; or

Primary GIST with a mitotic count of greater than 10/50 HPF.

A pathology report from an Approved Pathology Authority supporting the diagnosis of a gastrointestinal stromal tumour and confirming the presence of CD117 on immunohistochemical staining must be documented in the patient's medical records.

The pathology report must include the size and mitotic rate of the tumour, and the date of tumour resection, which must not be more than 3 months prior to treatment initiation must be recorded in the patient's medical records.

imatinib 100 mg tablet, 60

5443L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	476.87	41.30	Glivec [AF]

imatinib 400 mg tablet, 30

5444M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	937.83	41.30	Glivec [AF]

■ IMATINIB

Note Pharmaceutical benefits that have the form imatinib tablet 100 mg and imatinib capsule 100 mg are equivalent for the purposes of substitution.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Myelodysplastic or myeloproliferative disorder

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have confirmed evidence of a platelet-derived growth factor receptor (PDGFR) gene re-arrangement by standard karyotyping; OR
- Patient must have confirmed evidence of a platelet-derived growth factor receptor (PDGFR) gene re-arrangement by fluorescence in situ hybridization (FISH); OR
- Patient must have confirmed evidence of a platelet-derived growth factor receptor (PDGFR) gene re-arrangement by PDGFRB fusion gene transcript, **AND**
- Patient must have previously failed an adequate trial of conventional therapy with cytarabine; OR
- Patient must have previously failed an adequate trial of conventional therapy with etoposide; OR
- Patient must have previously failed an adequate trial of conventional therapy with hydroxycarbamide (hydroxyurea), **AND**
- The treatment must not exceed a maximum dose of 400 mg per day.

A bone marrow biopsy report demonstrating the presence of a myelodysplastic or myeloproliferative disorder, a pathology report confirming the platelet-derived growth factor receptor (PDGFR) gene re-arrangement and details of the prior trialled therapy and the response must be documented in the patient's medical records.

imatinib 100 mg capsule, 60

10918P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	476.87	41.30	^a CIPLA IMATINIB ADULT [LR] ^a IMATINIB-DRLA [RZ]	^a Imatinib-APOTEX [TX] ^a Imatinib GH [GQ]

imatinib 100 mg tablet, 60

9176Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	476.87	41.30	^a Gilmat [CR] ^a IMATINIB RBX [RA]	^a Glivec [AF] ^a Imatinib-Teva [SZ]

■ IMATINIB

Note Pharmaceutical benefits that have the form imatinib tablet 400 mg and imatinib capsule 400 mg are equivalent for the purposes of substitution.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Aggressive systemic mastocytosis with eosinophilia

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have confirmed evidence of carrying the FIP1L1-PDGFR fusion gene, **AND**
- Patient must have previously failed an adequate trial of conventional therapy with corticosteroids; OR
- Patient must have previously failed an adequate trial of conventional therapy with hydroxycarbamide (hydroxyurea), **AND**
- The treatment must not exceed a maximum dose of 400 mg per day.

A pathology report confirming the presence of the FIP1L1-PDGFR fusion gene, a bone marrow biopsy report and/or other tissue biopsy report confirming the diagnosis of aggressive systemic mastocytosis and a full blood examination report demonstrating eosinophilia must be documented in the patient's medical records.

The details of symptomatic organ involvement requiring treatment, including radiology, nuclear medicine, respiratory function or anatomical pathology reports as appropriate must be documented in the patient's medical records.

imatinib 400 mg capsule, 30

10921T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	937.83	41.30	^a CIPLA IMATINIB ADULT [LR] ^a IMATINIB-DRLA [RZ]	^a Imatinib-APOTEX [TX] ^a Imatinib GH [GQ]

imatinib 400 mg tablet, 30

9179D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	937.83	41.30	^a Glivec [AF]	^a IMATINIB RBX [RA]

■ **IMATINIB**

Note Pharmaceutical benefits that have the form imatinib tablet 400 mg and imatinib capsule 400 mg are equivalent for the purposes of substitution.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Chronic eosinophilic leukaemia or Hypereosinophilic syndrome

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have confirmed evidence of carrying the FIP1L1-PDGFR fusion gene, **AND**
- The treatment must not exceed a maximum dose of 400 mg per day.

A pathology report confirming the presence of the FIP1L1-PDGFR fusion gene, a full blood examination report and details of organ involvement requiring treatment, including a copy of the radiology, nuclear medicine, respiratory function or anatomical pathology reports as appropriate must be documented in the patient's medical records.

imatinib 400 mg capsule, 30

10925B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	937.83	41.30	^a CIPLA IMATINIB ADULT [LR] ^a IMATINIB-DRLA [RZ]	^a Imatinib-APOTEX [TX] ^a Imatinib GH [GQ]

imatinib 400 mg tablet, 30

9175X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	937.83	41.30	^a Gilmat [CR] ^a IMATINIB RBX [RA]	^a Glivec [AF] ^a Imatinib-Teva [SZ]

■ **IMATINIB**

Note Pharmaceutical benefits that have the form imatinib tablet 400 mg and imatinib capsule 400 mg are equivalent for the purposes of substitution.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Dermatofibrosarcoma protuberans

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be unresectable; OR
- The condition must be locally recurrent; OR
- The condition must be metastatic, **AND**
- The treatment must not exceed a maximum dose of 800 mg per day.

Details of unresectable tumour or site of the local recurrence or site(s) of metastatic disease must be documented in the patient's medical records.

imatinib 400 mg capsule, 30

10933K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	937.83	41.30	^a CIPLA IMATINIB ADULT [LR] ^a IMATINIB-DRLA [RZ]	^a Imatinib-APOTEX [TX] ^a Imatinib GH [GQ]

imatinib 400 mg tablet, 30

9173T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	937.83	41.30	^a Gilmat [CR] ^a IMATINIB RBX [RA]	^a Glivec [AF] ^a Imatinib-Teva [SZ]

■ **IMATINIB**

Note Pharmaceutical benefits that have the form imatinib tablet 400 mg and imatinib capsule 400 mg are equivalent for the purposes of substitution.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Myelodysplastic or myeloproliferative disorder

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have confirmed evidence of a platelet-derived growth factor receptor (PDGFR) gene re-arrangement by standard karyotyping; OR
- Patient must have confirmed evidence of a platelet-derived growth factor receptor (PDGFR) gene re-arrangement by fluorescence in situ hybridization (FISH); OR

- Patient must have confirmed evidence of a platelet-derived growth factor receptor (PDGFR) gene re-arrangement by PDGFRB fusion gene transcript, **AND**
- Patient must have previously failed an adequate trial of conventional therapy with cytarabine; OR
- Patient must have previously failed an adequate trial of conventional therapy with etoposide; OR
- Patient must have previously failed an adequate trial of conventional therapy with hydroxycarbamide (hydroxyurea), **AND**
- The treatment must not exceed a maximum dose of 400 mg per day.

A bone marrow biopsy report demonstrating the presence of a myelodysplastic or myeloproliferative disorder, a pathology report confirming the platelet-derived growth factor receptor (PDGFR) gene re-arrangement and details of the prior trialled therapy and the response must be documented in the patient's medical records.

imatinib 400 mg capsule, 30

10939R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	937.83	41.30	^a CIPLA IMATINIB ADULT [LR] ^a IMATINIB-DRLA [RZ]	^a Imatinib-APOTEX [TX] ^a Imatinib GH [GQ]

imatinib 400 mg tablet, 30

9177B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	937.83	41.30	^a Gilmat [CR] ^a IMATINIB RBX [RA]	^a Glivec [AF] ^a Imatinib-Teva [SZ]

■ IMATINIB

Note Pharmaceutical benefits that have the form imatinib tablet 100 mg and imatinib capsule 100 mg are equivalent for the purposes of substitution.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Aggressive systemic mastocytosis with eosinophilia

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have confirmed evidence of carrying the FIP1L1-PDGFR fusion gene, **AND**
- Patient must have previously failed an adequate trial of conventional therapy with corticosteroids; OR
- Patient must have previously failed an adequate trial of conventional therapy with hydroxycarbamide (hydroxyurea), **AND**
- The treatment must not exceed a maximum dose of 400 mg per day.

A pathology report confirming the presence of the FIP1L1-PDGFR fusion gene, a bone marrow biopsy report and/or other tissue biopsy report confirming the diagnosis of aggressive systemic mastocytosis and a full blood examination report demonstrating eosinophilia must be documented in the patient's medical records.

The details of symptomatic organ involvement requiring treatment, including radiology, nuclear medicine, respiratory function or anatomical pathology reports as appropriate must be documented in the patient's medical records.

imatinib 100 mg capsule, 60

10940T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	476.87	41.30	^a CIPLA IMATINIB ADULT [LR] ^a IMATINIB-DRLA [RZ]	^a Imatinib-APOTEX [TX] ^a Imatinib GH [GQ]

imatinib 100 mg tablet, 60

9178C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	476.87	41.30	^a Glivec [AF]	^a IMATINIB RBX [RA]

■ IMATINIB

Note Pharmaceutical benefits that have the form imatinib tablet 100 mg and imatinib capsule 100 mg are equivalent for the purposes of substitution.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Chronic eosinophilic leukaemia or Hypereosinophilic syndrome

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have confirmed evidence of carrying the FIP1L1-PDGFR fusion gene, **AND**
- The treatment must not exceed a maximum dose of 400 mg per day.

A pathology report confirming the presence of the FIP1L1-PDGFR fusion gene, a full blood examination report and details of organ involvement requiring treatment, including a copy of the radiology, nuclear medicine, respiratory function or anatomical pathology reports as appropriate must be documented in the patient's medical records.

imatinib 100 mg capsule, 60

10941W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	476.87	41.30	^a CIPLA IMATINIB ADULT [LR] ^a IMATINIB-DRLA [RZ]	^a Imatinib-APOTEX [TX] ^a Imatinib GH [GQ]

imatinib 100 mg tablet, 60

9174W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	476.87	41.30	^a Gilmat [CR] ^a IMATINIB RBX [RA]	^a Glivec [AF] ^a Imatinib-Teva [SZ]

■ IMATINIB

Note Pharmaceutical benefits that have the form imatinib tablet 100 mg and imatinib capsule 100 mg are equivalent for the purposes of substitution.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Dermatofibrosarcoma protuberans

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be unresectable; OR
- The condition must be locally recurrent; OR
- The condition must be metastatic, **AND**
- The treatment must not exceed a maximum dose of 800 mg per day.

Details of unresectable tumour or site of the local recurrence or site(s) of metastatic disease must be documented in the patient's medical records.

imatinib 100 mg capsule, 60

10942X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	476.87	41.30	^a CIPLA IMATINIB ADULT [LR] ^a IMATINIB-DRLA [RZ]	^a Imatinib-APOTEX [TX] ^a Imatinib GH [GQ]

imatinib 100 mg tablet, 60

9172R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	476.87	41.30	^a Gilmat [CR] ^a IMATINIB RBX [RA]	^a Glivec [AF] ^a Imatinib-Teva [SZ]

■ IMATINIB

Note Pharmaceutical benefits that have the form imatinib tablet 100 mg and imatinib capsule 100 mg are equivalent for the purposes of substitution.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

9209

Dermatofibrosarcoma protuberans

Treatment Phase: Continuing treatment

Clinical criteria:

- The condition must be unresectable; OR
- The condition must be locally recurrent; OR
- The condition must be metastatic, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have demonstrated a response to the PBS-subsidised treatment, **AND**
- The condition must not have progressed while receiving PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must not exceed a maximum dose of 800 mg per day.

Evidence that the disease has not progressed on imatinib therapy must be documented in the patient's medical records.

imatinib 100 mg capsule, 60

11776T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	476.87	41.30	^a CIPLA IMATINIB ADULT [LR] ^a IMATINIB-DRLA [RZ]	^a Imatinib-APOTEX [TX] ^a Imatinib GH [GQ]

imatinib 100 mg tablet, 60

11753N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	476.87	41.30	^a Gilmat [CR] ^a IMATINIB RBX [RA]	^a Glivec [AF] ^a Imatinib-Teva [SZ]

■ IMATINIB

Note Pharmaceutical benefits that have the form imatinib tablet 400 mg and imatinib capsule 400 mg are equivalent for the purposes of substitution.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

9243

Myelodysplastic or myeloproliferative disorder

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The condition must be PDGFRB fusion gene-positive, **AND**
- Patient must have achieved and maintained a complete haematological response, **AND**
- The condition must not have progressed while receiving PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must not exceed a maximum dose of 400 mg per day.

A full blood examination report which demonstrates a complete haematological response and evidence that the disease has not progressed on imatinib therapy must be documented in the patient's medical records.

imatinib 400 mg capsule, 30

11756R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	937.83	41.30	^a CIPLA IMATINIB ADULT [LR] ^a IMATINIB-DRLA [RZ]	^a Imatinib-APOTEX [TX] ^a Imatinib GH [GQ]

imatinib 400 mg tablet, 30

11765F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	937.83	41.30	^a Gilmat [CR] ^a IMATINIB RBX [RA]	^a Glivec [AF] ^a Imatinib-Teva [SZ]

■ IMATINIB

Note Pharmaceutical benefits that have the form imatinib tablet 100 mg and imatinib capsule 100 mg are equivalent for the purposes of substitution.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)**9243**

Myelodysplastic or myeloproliferative disorder

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The condition must be PDGFRB fusion gene-positive, **AND**
- Patient must have achieved and maintained a complete haematological response, **AND**
- The condition must not have progressed while receiving PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must not exceed a maximum dose of 400 mg per day.

A full blood examination report which demonstrates a complete haematological response and evidence that the disease has not progressed on imatinib therapy must be documented in the patient's medical records.

imatinib 100 mg capsule, 60

11757T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	476.87	41.30	^a CIPLA IMATINIB ADULT [LR] ^a IMATINIB-DRLA [RZ]	^a Imatinib-APOTEX [TX] ^a Imatinib GH [GQ]

imatinib 100 mg tablet, 60

11769K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	476.87	41.30	^a Gilmat [CR] ^a IMATINIB RBX [RA]	^a Glivec [AF] ^a Imatinib-Teva [SZ]

■ IMATINIB

Note Pharmaceutical benefits that have the form imatinib tablet 400 mg and imatinib capsule 400 mg are equivalent for the purposes of substitution.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)**9296**

Chronic eosinophilic leukaemia or Hypereosinophilic syndrome

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have achieved and maintained a complete haematological response, **AND**
- The condition must not have progressed while receiving PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must not exceed a maximum dose of 400 mg per day.

A full blood examination report which demonstrates a complete haematological response, with a normal eosinophil count and a statement that the disease has not progressed on imatinib therapy must be documented in the patient's medical records.

imatinib 400 mg capsule, 30

11763D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	937.83	41.30	^a CIPLA IMATINIB ADULT [LR] ^a IMATINIB-DRLA [RZ]	^a Imatinib-APOTEX [TX] ^a Imatinib GH [GQ]

imatinib 400 mg tablet, 30

11758W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	937.83	41.30	^a Gilmat [CR] ^a IMATINIB RBX [RA]	^a Glivec [AF] ^a Imatinib-Teva [SZ]

■ IMATINIB

Note Pharmaceutical benefits that have the form imatinib tablet 100 mg and imatinib capsule 100 mg are equivalent for the purposes of substitution.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

9206

Aggressive systemic mastocytosis with eosinophilia

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have confirmed evidence of carrying the FIP1L1-PDGFR fusion gene, **AND**
- Patient must have achieved and maintained a complete haematological response, **AND**
- The condition must not have progressed while receiving PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must not exceed a maximum dose of 400 mg per day.

A full blood examination report which demonstrates a complete haematological response and evidence that the disease has not progressed on imatinib therapy must be documented in the patient's medical records.

imatinib 100 mg capsule, 60

11777W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	476.87	41.30	^a CIPLA IMATINIB ADULT [LR] ^a IMATINIB-DRLA [RZ]	^a Imatinib-APOTEX [TX] ^a Imatinib GH [GQ]

imatinib 100 mg tablet, 60

11762C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	476.87	41.30	^a Glivec [AF]	^a IMATINIB RBX [RA]

■ IMATINIB

Note Pharmaceutical benefits that have the form imatinib tablet 400 mg and imatinib capsule 400 mg are equivalent for the purposes of substitution.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)**9209**

Dermatofibrosarcoma protuberans

Treatment Phase: Continuing treatment

Clinical criteria:

- The condition must be unresectable; OR
- The condition must be locally recurrent; OR
- The condition must be metastatic, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have demonstrated a response to the PBS-subsidised treatment, **AND**
- The condition must not have progressed while receiving PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must not exceed a maximum dose of 800 mg per day.

Evidence that the disease has not progressed on imatinib therapy must be documented in the patient's medical records.

imatinib 400 mg capsule, 30

11764E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	937.83	41.30	^a CIPLA IMATINIB ADULT [LR] ^a IMATINIB-DRLA [RZ]	^a Imatinib-APOTEX [TX] ^a Imatinib GH [GQ]

imatinib 400 mg tablet, 30

11786H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	937.83	41.30	^a Gilmat [CR] ^a IMATINIB RBX [RA]	^a Glivec [AF] ^a Imatinib-Teva [SZ]

■ IMATINIB

Note Pharmaceutical benefits that have the form imatinib tablet 100 mg and imatinib capsule 100 mg are equivalent for the purposes of substitution.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)**9296**

Chronic eosinophilic leukaemia or Hypereosinophilic syndrome

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have achieved and maintained a complete haematological response, **AND**
- The condition must not have progressed while receiving PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must not exceed a maximum dose of 400 mg per day.

A full blood examination report which demonstrates a complete haematological response, with a normal eosinophil count and a statement that the disease has not progressed on imatinib therapy must be documented in the patient's medical records.

imatinib 100 mg capsule, 60

11770L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	476.87	41.30	^a CIPLA IMATINIB ADULT [LR]	^a Imatinib-APOTEX [TX]

^a IMATINIB-DRLA [RZ]^a Imatinib GH [GQ]**imatinib 100 mg tablet, 60**

11781C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	476.87	41.30	^a Gilmat [CR] ^a IMATINIB RBX [RA]	^a Glivec [AF] ^a Imatinib-Teva [SZ]

■ IMATINIB

Note Authority applications for increased quantities/ repeats (where relevant) may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)**9208**

Malignant gastrointestinal stromal tumour

Treatment Phase: Continuing treatment

Clinical criteria:

- The condition must be metastatic; OR
- The condition must be unresectable, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be given at a dose not exceeding 600 mg per day.

Patients who have failed to respond or are intolerant to imatinib are no longer eligible to receive PBS-subsidised imatinib. Patients with metastatic/unresectable disease who achieve a response to treatment at an imatinib dose of 400 mg per day should be continued at this dose and assessed for response at regular intervals. Patients who fail to achieve a response to 400 mg per day may have their dose increased to 600 mg per day. Authority applications for doses higher than 600 mg per day will not be approved.

A response to treatment is defined as a decrease from baseline in the sum of the products of the perpendicular diameters of all measurable lesions of 50% or greater. (Response definition based on the Southwest Oncology Group standard criteria, see Demetri et al. N Engl J Med 2002; 347: 472-80.)

imatinib 100 mg tablet, 60

11787J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	476.87	41.30	Glivec [AF]

imatinib 400 mg tablet, 30

11778X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	937.83	41.30	Glivec [AF]

■ IMATINIB

Note Pharmaceutical benefits that have the form imatinib tablet 400 mg and imatinib capsule 400 mg are equivalent for the purposes of substitution.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)**9206**

Aggressive systemic mastocytosis with eosinophilia

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have confirmed evidence of carrying the FIP1L1-PDGFR fusion gene, **AND**
- Patient must have achieved and maintained a complete haematological response, **AND**
- The condition must not have progressed while receiving PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must not exceed a maximum dose of 400 mg per day.

A full blood examination report which demonstrates a complete haematological response and evidence that the disease has not progressed on imatinib therapy must be documented in the patient's medical records.

imatinib 400 mg capsule, 30

11779Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	937.83	41.30	^a CIPLA IMATINIB ADULT [LR] ^a IMATINIB-DRLA [RZ]	^a Imatinib-APOTEX [TX] ^a Imatinib GH [GQ]

imatinib 400 mg tablet, 30

11785G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	937.83	41.30	^a Glivec [AF]	^a IMATINIB RBX [RA]

■ IMATINIB

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note No increase in the maximum number of repeats may be authorised.

Authority required

Malignant gastrointestinal stromal tumour

Treatment Phase: Initial Treatment

Clinical criteria:

- The condition must be metastatic; OR
- The condition must be unresectable, **AND**
- The condition must be histologically confirmed by the detection of CD117 on immunohistochemical staining, **AND**
- The treatment must be commenced at a dose not exceeding 400 mg per day, **AND**
- The treatment must not exceed 3 months under this restriction.

Authority prescriptions for a higher dose will not be approved during this initial 3 month treatment period.

Patients with metastatic/unresectable disease who achieve a response to treatment at an imatinib dose of 400 mg per day should be continued at this dose and assessed for response at regular intervals. Patients who fail to achieve a response to 400 mg per day may have their dose increased to 600 mg per day. Authority applications for doses higher than 600 mg per day will not be approved.

A response to treatment is defined as a decrease from baseline in the sum of the products of the perpendicular diameters of all measurable lesions of 50% or greater. (Response definition based on the Southwest Oncology Group standard criteria, see Demetri et al. N Engl J Med 2002; 347: 472-80.)

A pathology report from an Approved Pathology Authority supporting the diagnosis of a gastrointestinal stromal tumour and confirming the presence of CD117 on immunohistochemical staining must be documented in the patient's medical records. Details of the most recent (within 2 months of the application) computed tomography (CT) scan, magnetic resonance imaging (MRI) or ultrasound assessment of the tumour(s), including whether or not there is evidence of metastatic disease must be documented in the patient's medical records.

Where the application for authority to prescribe is being sought on the basis of an unresectable tumour, written evidence must be documented in the patient's medical records.

imatinib 100 mg tablet, 60

9111M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	476.87	41.30	Glivec [AF]

imatinib 400 mg tablet, 30

9112N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	937.83	41.30	Glivec [AF]

■ IMATINIB

Note Pharmaceutical benefits that have the form imatinib tablet 400 mg and imatinib capsule 400 mg are equivalent for the purposes of substitution.

Note Authority applications for increased quantities/ repeats (where relevant) may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of tyrosine kinase inhibitors (TKI) agents for the chronic phase of chronic myeloid leukaemia. Where the term TKI agent appears in the following notes and restrictions it refers to imatinib mesilate, dasatinib or nilotinib.

Patients are eligible for PBS-subsidised treatment with only one TKI agent at any one time and must not be receiving concomitant interferon alfa therapy. Eligible patients may only swap between TKI agents if they have not failed prior PBS-subsidised treatment with that agent.

1. Initial First-line treatment From 1 April 2012, under the PBS, a patient will be able to be prescribed any of imatinib mesilate, dasatinib or nilotinib within the initial 18 month treatment period, as long as only one agent is used at a time and providing the patient has not failed to respond to any one of these TKIs. During the initial 18 month treatment period, switching between approved first-line agents may only occur for reasons of intolerance, not failure to respond

2. Continuing First-line treatment

Patients must maintain a major cytogenetic response or have a peripheral blood BCR-ABL of less than 1% to receive continuing therapy.

For imatinib mesilate:

First continuing applications may be made by telephoning the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

For dasatinib or nilotinib:

First continuing applications are to be written and must include a pathology report demonstrating the patient has responded to the initial course of treatment.

Second and subsequent authority applications for continuing therapy may be made by telephoning the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

During continuing therapy beyond the initial 18 month treatment period, switching between approved first-line agents may only occur for reason of intolerance. Where there is failure to respond, switching may only occur through application for prescription of second-line agents. Where a patient has previously received PBS-subsidised treatment with imatinib mesilate, dasatinib or nilotinib no approval will be granted for PBS-subsidised re-treatment in the chronic phase of chronic myeloid leukaemia, where that patient has at any time failed to meet the response criteria whilst on that TKI agent.

3. Authority approval requirements. Response criteria to initial first-line treatment with imatinib mesilate, dasatinib or nilotinib: For the purposes of assessing response to PBS-subsidised treatment with imatinib mesilate, dasatinib or nilotinib either cytogenetic analysis indicating the number of Philadelphia positive [t (9;22)] cells in the bone marrow measured by standard karyotyping, or quantitative PCR indicating the relative level of BCR-ABL transcript in the peripheral blood using the international scale, must be submitted. For bone marrow analyses, where the standard karyotyping is not informative for technical reasons, a cytogenetic analysis performed on the bone marrow by the use of fluorescence in situ hybridisation (FISH) with BCR-ABL specific probe must be submitted. The cytogenetic or peripheral blood quantitative PCR analyses must be submitted within 18 months of the commencement of treatment with dasatinib or nilotinib (patients in whom a major

cytogenetic response or peripheral blood BCR-ABL level of less than 1% is demonstrable by 18 months are eligible to receive continuing treatment with that agent).

4. Definitions of response

A major cytogenetic response is defined as less than 35% Philadelphia positive bone marrow cells. A peripheral blood BCR-ABL level of less than 1% on the international scale (Blood 108: 28-37, 2006) also indicates a response, at least the biological equivalent of a major cytogenetic response.

5. Definitions of loss of response

Loss of a previously documented major cytogenetic response (demonstrated by the presence of greater than 35% Ph positive cells on bone marrow biopsy), during ongoing tyrosine kinase inhibitor (TKI) therapy. Loss of a previously demonstrated molecular response (demonstrated by peripheral blood BCR-ABL levels increasing consecutively in value by at least 5 fold to a level of greater than 0.1% confirmed on a subsequent test), during ongoing tyrosine kinase inhibitor therapy.

Authority required (STREAMLINED)

9295

Chronic Myeloid Leukaemia (CML)

Treatment Phase: Subsequent continuing

Clinical criteria:

- The condition must be in the chronic phase of chronic myeloid leukaemia, **AND**
- Patient must have received initial continuing PBS-subsidised treatment with this drug as a first line therapy for this condition; OR
- Patient must have experienced intolerance, not a failure to respond, to subsequent continuing PBS-subsidised treatment with dasatinib as a first line therapy for this condition; OR
- Patient must have experienced intolerance, not a failure to respond, to subsequent continuing PBS-subsidised treatment with nilotinib as a first line therapy for this condition, **AND**
- Patient must have maintained a major cytogenic response; OR
- Patient must have maintained a peripheral blood level of BCR-ABL of less than 1%, **AND**
- The treatment must not exceed a total maximum of 24 weeks of therapy with a PBS-subsidised treatment with a tyrosine kinase inhibitor for this condition under this restriction, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

imatinib 400 mg capsule, 30

11772N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	937.83	41.30	^a CIPLA IMATINIB ADULT [LR] ^a IMATINIB-DRLA [RZ]	^a Imatinib-APOTEX [TX] ^a Imatinib GH [GQ]

imatinib 400 mg tablet, 30

11752M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	937.83	41.30	^a Gilmat [CR] ^a IMATINIB RBX [RA]	^a Glivec [AF] ^a Imatinib-Teva [SZ]

■ IMATINIB

Note Pharmaceutical benefits that have the form imatinib tablet 100 mg and imatinib capsule 100 mg are equivalent for the purposes of substitution.

Note Authority applications for increased quantities/ repeats (where relevant) may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of tyrosine kinase inhibitors (TKI) agents for the chronic phase of chronic myeloid leukaemia. Where the term TKI agent appears in the following notes and restrictions it refers to imatinib mesilate, dasatinib or nilotinib.

Patients are eligible for PBS-subsidised treatment with only one TKI agent at any one time and must not be receiving concomitant interferon alfa therapy. Eligible patients may only swap between TKI agents if they have not failed prior PBS-subsidised treatment with that agent.

1. Initial First-line treatment From 1 April 2012, under the PBS, a patient will be able to be prescribed any of imatinib mesilate, dasatinib or nilotinib within the initial 18 month treatment period, as long as only one agent is used at a time and providing the patient has not failed to respond to any one of these TKIs. During the initial 18 month treatment period, switching between approved first-line agents may only occur for reasons of intolerance, not failure to respond

2. Continuing First-line treatment

Patients must maintain a major cytogenetic response or have a peripheral blood BCR-ABL of less than 1% to receive continuing therapy.

For imatinib mesilate:

First continuing applications may be made by telephoning the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

For dasatinib or nilotinib:

First continuing applications are to be written and must include a pathology report demonstrating the patient has responded to the initial course of treatment.

Second and subsequent authority applications for continuing therapy may be made by telephoning the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

During continuing therapy beyond the initial 18 month treatment period, switching between approved first-line agents may only occur for reason of intolerance. Where there is failure to respond, switching may only occur through application for prescription of second-line agents. Where a patient has previously received PBS-subsidised treatment with imatinib mesilate, dasatinib or nilotinib no approval will be granted for PBS-subsidised re-treatment in the chronic phase of chronic myeloid leukaemia, where that patient has at any time failed to meet the response criteria whilst on that TKI agent.

3. Authority approval requirements. Response criteria to initial first-line treatment with imatinib mesilate, dasatinib or

nilotinib: For the purposes of assessing response to PBS-subsidised treatment with imatinib mesilate, dasatinib or nilotinib either cytogenetic analysis indicating the number of Philadelphia positive [t (9;22)] cells in the bone marrow measured by standard karyotyping, or quantitative PCR indicating the relative level of BCR-ABL transcript in the peripheral blood using the international scale, must be submitted. For bone marrow analyses, where the standard karyotyping is not informative for technical reasons, a cytogenetic analysis performed on the bone marrow by the use of fluorescence in situ hybridisation (FISH) with BCR-ABL specific probe must be submitted. The cytogenetic or peripheral blood quantitative PCR analyses must be submitted within 18 months of the commencement of treatment with dasatinib or nilotinib (patients in whom a major cytogenetic response or peripheral blood BCR-ABL level of less than 1% is demonstrable by 18 months are eligible to receive continuing treatment with that agent).

4. Definitions of response

A major cytogenetic response is defined as less than 35% Philadelphia positive bone marrow cells. A peripheral blood BCR-ABL level of less than 1% on the international scale (Blood 108: 28-37, 2006) also indicates a response, at least the biological equivalent of a major cytogenetic response.

5. Definitions of loss of response

Loss of a previously documented major cytogenetic response (demonstrated by the presence of greater than 35% Ph positive cells on bone marrow biopsy), during ongoing tyrosine kinase inhibitor (TKI) therapy. Loss of a previously demonstrated molecular response (demonstrated by peripheral blood BCR-ABL levels increasing consecutively in value by at least 5 fold to a level of greater than 0.1% confirmed on a subsequent test), during ongoing tyrosine kinase inhibitor therapy.

Authority required (STREAMLINED)

9295

Chronic Myeloid Leukaemia (CML)

Treatment Phase: Subsequent continuing

Clinical criteria:

- The condition must be in the chronic phase of chronic myeloid leukaemia, **AND**
- Patient must have received initial continuing PBS-subsidised treatment with this drug as a first line therapy for this condition; OR
- Patient must have experienced intolerance, not a failure to respond, to subsequent continuing PBS-subsidised treatment with dasatinib as a first line therapy for this condition; OR
- Patient must have experienced intolerance, not a failure to respond, to subsequent continuing PBS-subsidised treatment with nilotinib as a first line therapy for this condition, **AND**
- Patient must have maintained a major cytogenetic response; OR
- Patient must have maintained a peripheral blood level of BCR-ABL of less than 1%, **AND**
- The treatment must not exceed a total maximum of 24 weeks of therapy with a PBS-subsidised treatment with a tyrosine kinase inhibitor for this condition under this restriction, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

imatinib 100 mg capsule, 60

11782D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	476.87	41.30	^a CIPLA IMATINIB ADULT [LR] ^a IMATINIB-DRLA [RZ]	^a Imatinib-APOTEX [TX] ^a Imatinib GH [GQ]

imatinib 100 mg tablet, 60

11775R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	476.87	41.30	^a Gilmat [CR] ^a IMATINIB RBX [RA]	^a Glivec [AF] ^a Imatinib-Teva [SZ]

■ IMATINIB

Note Pharmaceutical benefits that have the form imatinib tablet 100 mg and imatinib capsule 100 mg are equivalent for the purposes of substitution.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of tyrosine kinase inhibitors (TKI) agents for the chronic phase of chronic myeloid leukaemia. Where the term TKI agent appears in the following notes and restrictions it refers to imatinib mesilate, dasatinib or nilotinib.

Patients are eligible for PBS-subsidised treatment with only one TKI agent at any one time and must not be receiving concomitant interferon alfa therapy. Eligible patients may only swap between TKI agents if they have not failed prior PBS-subsidised treatment with that agent.

1. Initial First-line treatment From 1 April 2012, under the PBS, a patient will be able to be prescribed any of imatinib mesilate, dasatinib or nilotinib within the initial 18 month treatment period, as long as only one agent is used at a time and providing the patient has not failed to respond to any one of these TKIs. During the initial 18 month treatment period, switching between approved first-line agents may only occur for reasons of intolerance, not failure to respond

2. Continuing First-line treatment

Patients must maintain a major cytogenetic response or have a peripheral blood BCR-ABL of less than 1% to receive continuing therapy.

For imatinib mesilate:

First continuing applications may be made by telephoning the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

For dasatinib or nilotinib:

First continuing applications are to be written and must include a pathology report demonstrating the patient has responded to the initial course of treatment.

Second and subsequent authority applications for continuing therapy may be made by telephoning the Department of

Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

During continuing therapy beyond the initial 18 month treatment period, switching between approved first-line agents may only occur for reason of intolerance. Where there is failure to respond, switching may only occur through application for prescription of second-line agents. Where a patient has previously received PBS-subsidised treatment with imatinib mesilate, dasatinib or nilotinib no approval will be granted for PBS-subsidised re-treatment in the chronic phase of chronic myeloid leukaemia, where that patient has at any time failed to meet the response criteria whilst on that TKI agent.

3. Authority approval requirements. Response criteria to initial first-line treatment with imatinib mesilate, dasatinib or nilotinib: For the purposes of assessing response to PBS-subsidised treatment with imatinib mesilate, dasatinib or nilotinib either cytogenetic analysis indicating the number of Philadelphia positive [t (9;22)] cells in the bone marrow measured by standard karyotyping, or quantitative PCR indicating the relative level of BCR-ABL transcript in the peripheral blood using the international scale, must be submitted. For bone marrow analyses, where the standard karyotyping is not informative for technical reasons, a cytogenetic analysis performed on the bone marrow by the use of fluorescence in situ hybridisation (FISH) with BCR-ABL specific probe must be submitted. The cytogenetic or peripheral blood quantitative PCR analyses must be submitted within 18 months of the commencement of treatment with dasatinib or nilotinib (patients in whom a major cytogenetic response or peripheral blood BCR-ABL level of less than 1% is demonstrable by 18 months are eligible to receive continuing treatment with that agent).

4. Definitions of response

A major cytogenetic response is defined as less than 35% Philadelphia positive bone marrow cells. A peripheral blood BCR-ABL level of less than 1% on the international scale (Blood 108: 28-37, 2006) also indicates a response, at least the biological equivalent of a major cytogenetic response.

5. Definitions of loss of response

Loss of a previously documented major cytogenetic response (demonstrated by the presence of greater than 35% Ph positive cells on bone marrow biopsy), during ongoing tyrosine kinase inhibitor (TKI) therapy. Loss of a previously demonstrated molecular response (demonstrated by peripheral blood BCR-ABL levels increasing consecutively in value by at least 5 fold to a level of greater than 0.1% confirmed on a subsequent test), during ongoing tyrosine kinase inhibitor therapy.

Authority required

Chronic Myeloid Leukaemia (CML)

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have a primary diagnosis of chronic myeloid leukaemia, **AND**
- The condition must be in the chronic phase of chronic myeloid leukaemia, **AND**
- The condition must be expressing the Philadelphia chromosome; OR
- The condition must have the transcript BCR-ABL tyrosine kinase, **AND**
- The treatment must be for first line therapy for this condition, **AND**
- Patient must not have previously experienced a failure to respond to the PBS-subsidised treatment with this drug for this condition; OR
- Patient must have experienced intolerance, not a failure to respond, to initial PBS-subsidised treatment with dasatinib as a first line therapy for this condition; OR
- Patient must have experienced intolerance, not a failure to respond, to initial PBS-subsidised treatment with nilotinib as a first line therapy for this condition, **AND**
- The treatment must not exceed a total maximum of 18 months of therapy with a PBS-subsidised treatment with a tyrosine kinase inhibitor for this condition, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Applications under this restriction will be limited to provide patients with a maximum of 18 months of therapy with dasatinib, imatinib or nilotinib from the date the first application for initial treatment was approved.

Patients should be commenced on a dose of imatinib mesilate of 400 mg (base) daily. Continuing therapy is dependent on patients demonstrating a response to imatinib mesilate therapy following the initial 18 months of treatment and at 12 monthly intervals thereafter.

A pathology cytogenetic report conducted on peripheral blood or bone marrow supporting the diagnosis of chronic myeloid leukaemia to confirm eligibility for treatment, or a qualitative PCR report documenting the presence of the BCR-ABL transcript in either peripheral blood or bone marrow must be documented in the patient's medical records.

Authority required

Chronic Myeloid Leukaemia (CML)

Treatment Phase: First Continuing

Clinical criteria:

- The condition must be in the chronic phase of chronic myeloid leukaemia, **AND**
- Patient must have received initial PBS-subsidised treatment with this drug as a first line therapy for this condition; OR
- Patient must have experienced intolerance, not a failure to respond, to first continuing PBS-subsidised treatment with dasatinib as a first line therapy for this condition; OR
- Patient must have experienced intolerance, not a failure to respond, to first continuing PBS-subsidised treatment with nilotinib as a first line therapy for this condition, **AND**
- Patient must have demonstrated a major cytogenetic response; OR
- Patient must have demonstrated a peripheral blood level of BCR-ABL of less than 1%, **AND**
- The treatment must not exceed a total maximum of 24 weeks of therapy with a PBS-subsidised treatment with a tyrosine kinase inhibitor for this condition under this restriction, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

A major cytogenetic response [see Note explaining requirements] or a peripheral blood level of BCR-ABL of less than 1% on the international scale [see Note explaining requirements] must be documented in the patient's medical records.

imatinib 100 mg capsule, 60

10915L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	476.87	41.30	^a CIPLA IMATINIB ADULT [LR] ^a IMATINIB-DRLA [RZ]	^a Imatinib-APOTEX [TX] ^a Imatinib GH [GQ]

imatinib 100 mg tablet, 60

9113P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	476.87	41.30	^a Gilmat [CR] ^a IMATINIB RBX [RA]	^a Glivec [AF] ^a Imatinib-Teva [SZ]

■ **IMATINIB**

Note Pharmaceutical benefits that have the form imatinib tablet 400 mg and imatinib capsule 400 mg are equivalent for the purposes of substitution.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of tyrosine kinase inhibitors (TKI) agents for the chronic phase of chronic myeloid leukaemia. Where the term TKI agent appears in the following notes and restrictions it refers to imatinib mesilate, dasatinib or nilotinib.

Patients are eligible for PBS-subsidised treatment with only one TKI agent at any one time and must not be receiving concomitant interferon alfa therapy. Eligible patients may only swap between TKI agents if they have not failed prior PBS-subsidised treatment with that agent.

1. Initial First-line treatment From 1 April 2012, under the PBS, a patient will be able to be prescribed any of imatinib mesilate, dasatinib or nilotinib within the initial 18 month treatment period, as long as only one agent is used at a time and providing the patient has not failed to respond to any one of these TKIs. During the initial 18 month treatment period, switching between approved first-line agents may only occur for reasons of intolerance, not failure to respond

2. Continuing First-line treatment

Patients must maintain a major cytogenetic response or have a peripheral blood BCR-ABL of less than 1% to receive continuing therapy.

For imatinib mesilate:

First continuing applications may be made by telephoning the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

For dasatinib or nilotinib:

First continuing applications are to be written and must include a pathology report demonstrating the patient has responded to the initial course of treatment.

Second and subsequent authority applications for continuing therapy may be made by telephoning the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

During continuing therapy beyond the initial 18 month treatment period, switching between approved first-line agents may only occur for reason of intolerance. Where there is failure to respond, switching may only occur through application for prescription of second-line agents. Where a patient has previously received PBS-subsidised treatment with imatinib mesilate, dasatinib or nilotinib no approval will be granted for PBS-subsidised re-treatment in the chronic phase of chronic myeloid leukaemia, where that patient has at any time failed to meet the response criteria whilst on that TKI agent.

3. Authority approval requirements. Response criteria to initial first-line treatment with imatinib mesilate, dasatinib or nilotinib: For the purposes of assessing response to PBS-subsidised treatment with imatinib mesilate, dasatinib or nilotinib either cytogenetic analysis indicating the number of Philadelphia positive [t (9;22)] cells in the bone marrow measured by standard karyotyping, or quantitative PCR indicating the relative level of BCR-ABL transcript in the peripheral blood using the international scale, must be submitted. For bone marrow analyses, where the standard karyotyping is not informative for technical reasons, a cytogenetic analysis performed on the bone marrow by the use of fluorescence in situ hybridisation (FISH) with BCR-ABL specific probe must be submitted. The cytogenetic or peripheral blood quantitative PCR analyses must be submitted within 18 months of the commencement of treatment with dasatinib or nilotinib (patients in whom a major cytogenetic response or peripheral blood BCR-ABL level of less than 1% is demonstrable by 18 months are eligible to receive continuing treatment with that agent).

4. Definitions of response

A major cytogenetic response is defined as less than 35% Philadelphia positive bone marrow cells. A peripheral blood BCR-ABL level of less than 1% on the international scale (Blood 108: 28-37, 2006) also indicates a response, at least the biological equivalent of a major cytogenetic response.

5. Definitions of loss of response

Loss of a previously documented major cytogenetic response (demonstrated by the presence of greater than 35% Ph positive cells on bone marrow biopsy), during ongoing tyrosine kinase inhibitor (TKI) therapy. Loss of a previously demonstrated molecular response (demonstrated by peripheral blood BCR-ABL levels increasing consecutively in value by at least 5 fold to a level of greater than 0.1% confirmed on a subsequent test), during ongoing tyrosine kinase inhibitor therapy.

Authority required

Chronic Myeloid Leukaemia (CML)

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have a primary diagnosis of chronic myeloid leukaemia, **AND**
- The condition must be in the chronic phase of chronic myeloid leukaemia, **AND**
- The condition must be expressing the Philadelphia chromosome; OR
- The condition must have the transcript BCR-ABL tyrosine kinase, **AND**
- The treatment must be for first line therapy for this condition, **AND**

- Patient must not have previously experienced a failure to respond to the PBS-subsidised treatment with this drug for this condition; OR
- Patient must have experienced intolerance, not a failure to respond, to initial PBS-subsidised treatment with dasatinib as a first line therapy for this condition; OR
- Patient must have experienced intolerance, not a failure to respond, to initial PBS-subsidised treatment with nilotinib as a first line therapy for this condition, **AND**
- The treatment must not exceed a total maximum of 18 months of therapy with a PBS-subsidised treatment with a tyrosine kinase inhibitor for this condition, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Applications under this restriction will be limited to provide patients with a maximum of 18 months of therapy with dasatinib, imatinib or nilotinib from the date the first application for initial treatment was approved.

Patients should be commenced on a dose of imatinib mesilate of 400 mg (base) daily. Continuing therapy is dependent on patients demonstrating a response to imatinib mesilate therapy following the initial 18 months of treatment and at 12 monthly intervals thereafter.

A pathology cytogenetic report conducted on peripheral blood or bone marrow supporting the diagnosis of chronic myeloid leukaemia to confirm eligibility for treatment, or a qualitative PCR report documenting the presence of the BCR-ABL transcript in either peripheral blood or bone marrow must be documented in the patient's medical records.

Authority required

Chronic Myeloid Leukaemia (CML)

Treatment Phase: First Continuing

Clinical criteria:

- The condition must be in the chronic phase of chronic myeloid leukaemia, **AND**
- Patient must have received initial PBS-subsidised treatment with this drug as a first line therapy for this condition; OR
- Patient must have experienced intolerance, not a failure to respond, to first continuing PBS-subsidised treatment with dasatinib as a first line therapy for this condition; OR
- Patient must have experienced intolerance, not a failure to respond, to first continuing PBS-subsidised treatment with nilotinib as a first line therapy for this condition, **AND**
- Patient must have demonstrated a major cytogenetic response; OR
- Patient must have demonstrated a peripheral blood level of BCR-ABL of less than 1%, **AND**
- The treatment must not exceed a total maximum of 24 weeks of therapy with a PBS-subsidised treatment with a tyrosine kinase inhibitor for this condition under this restriction, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

A major cytogenetic response [see Note explaining requirements] or a peripheral blood level of BCR-ABL of less than 1% on the international scale [see Note explaining requirements] must be documented in the patient's medical records.

imatinib 400 mg capsule, 30

10916M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	937.83	41.30	^a CIPLA IMATINIB ADULT [LR] ^a IMATINIB-DRLA [RZ]	^a Imatinib-APOTEX [TX] ^a Imatinib GH [GQ]

imatinib 400 mg tablet, 30

9114Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	937.83	41.30	^a Gilmat [CR] ^a IMATINIB RBX [RA]	^a Glivec [AF] ^a Imatinib-Teva [SZ]

■ IMATINIB

Note Pharmaceutical benefits that have the form imatinib tablet 400 mg and imatinib capsule 400 mg are equivalent for the purposes of substitution.

Note No increase in the maximum number of repeats may be authorised.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note Allogeneic stem cell transplantation is the preferred therapy for eligible patients achieving a complete remission of Philadelphia positive acute lymphoblastic leukaemia.

Authority required

Acute Lymphoblastic leukaemia

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must be newly diagnosed, **AND**
- The condition must be expressing the Philadelphia chromosome; OR
- The condition must have the transcript BCR-ABL, **AND**
- The treatment must be for induction and consolidation therapy, **AND**
- The treatment must be in combination with chemotherapy or corticosteroids, **AND**
- Patient must not have previously experienced a failure to respond to the PBS-subsidised first line treatment with this drug for this condition; OR
- Patient must have experienced intolerance, not a failure to respond, to initial PBS-subsidised treatment with dasatinib as a first line therapy for this condition.

A pathology cytogenetic report conducted on peripheral blood or bone marrow supporting the diagnosis of acute lymphoblastic leukaemia with either cytogenetic evidence of the Philadelphia chromosome, or a qualitative PCR report

documenting the presence of the BCR-ABL transcript in either peripheral blood or bone marrow must be documented in the patient's medical records.

imatinib 400 mg capsule, 30

10917N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	2	..	937.83	41.30	^a CIPLA IMATINIB ADULT [LR] ^a IMATINIB-DRLA [RZ]	^a Imatinib-APOTEX [TX] ^a Imatinib GH [GQ]

imatinib 400 mg tablet, 30

9124F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	2	..	937.83	41.30	^a Gilmat [CR] ^a IMATINIB RBX [RA]	^a Glivec [AF] ^a Imatinib-Teva [SZ]

■ IMATINIB

Note Pharmaceutical benefits that have the form imatinib tablet 100 mg and imatinib capsule 100 mg are equivalent for the purposes of substitution.

Note No increase in the maximum number of repeats may be authorised.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Chronic Myeloid Leukaemia (CML)

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have a primary diagnosis of chronic myeloid leukaemia, **AND**
- The condition must be in the accelerated phase, **AND**
- The condition must be expressing the Philadelphia chromosome; OR
- The condition must have the transcript BCR-ABL tyrosine kinase.

Accelerated phase is defined by the presence of 1 or more of the following:

1. Percentage of blasts in the peripheral blood or bone marrow greater than or equal to 15% but less than 30%; or
2. Percentage of blasts plus promyelocytes in the peripheral blood or bone marrow greater than or equal to 30%, provided that blast count is less than 30%; or
3. Peripheral basophils greater than or equal to 20%; or
4. Progressive splenomegaly to a size greater than or equal to 10 cm below the left costal margin to be confirmed on 2 occasions at least 4 weeks apart, or a greater than or equal to 50% increase in size below the left costal margin over 4 weeks; or
5. Karyotypic evolution (chromosomal abnormalities in addition to a single Philadelphia chromosome).

Authority required

Chronic Myeloid Leukaemia (CML)

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must a primary diagnosis of chronic myeloid leukaemia, **AND**
- The condition must be in the blast phase, **AND**
- The condition must be expressing the Philadelphia chromosome; OR
- The condition must have the transcript BCR-ABL tyrosine kinase.

Blast crisis is defined as either:

1. Percentage of blasts in the peripheral blood or bone marrow greater than or equal to 30%; or
2. Extramedullary involvement other than spleen and liver.

imatinib 100 mg capsule, 60

10920R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	2	..	476.87	41.30	^a CIPLA IMATINIB ADULT [LR] ^a IMATINIB-DRLA [RZ]	^a Imatinib-APOTEX [TX] ^a Imatinib GH [GQ]

imatinib 100 mg tablet, 60

9115R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	2	..	476.87	41.30	^a Gilmat [CR] ^a IMATINIB RBX [RA]	^a Glivec [AF] ^a Imatinib-Teva [SZ]

■ IMATINIB

Note Pharmaceutical benefits that have the form imatinib tablet 100 mg and imatinib capsule 100 mg are equivalent for the purposes of substitution.

Note No increase in the maximum number of repeats may be authorised.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note Allogeneic stem cell transplantation is the preferred therapy for eligible patients achieving a complete remission of Philadelphia positive acute lymphoblastic leukaemia.

Authority required

Acute lymphoblastic leukaemia

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must be newly diagnosed, **AND**
- The condition must be expressing the Philadelphia chromosome; OR
- The condition must have the transcript BCR-ABL, **AND**
- The treatment must be for induction and consolidation therapy, **AND**
- The treatment must be in combination with chemotherapy or corticosteroids, **AND**
- Patient must not have previously experienced a failure to respond to the PBS-subsidised first line treatment with this drug for this condition; OR
- Patient must have experienced intolerance, not a failure to respond, to initial PBS-subsidised treatment with dasatinib as a first line therapy for this condition.

A pathology cytogenetic report conducted on peripheral blood or bone marrow supporting the diagnosis of acute lymphoblastic leukaemia with either cytogenetic evidence of the Philadelphia chromosome, or a qualitative PCR report documenting the presence of the BCR-ABL transcript in either peripheral blood or bone marrow must be documented in the patient's medical records.

imatinib 100 mg capsule, 60

10924Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	2	..	476.87	41.30	^a CIPLA IMATINIB ADULT [LR] ^a IMATINIB-DRLA [RZ]	^a Imatinib-APOTEX [TX] ^a Imatinib GH [GQ]

imatinib 100 mg tablet, 60

9123E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	2	..	476.87	41.30	^a Gilmat [CR] ^a IMATINIB RBX [RA]	^a Glivec [AF] ^a Imatinib-Teva [SZ]

■ IMATINIB

Note Pharmaceutical benefits that have the form imatinib tablet 400 mg and imatinib capsule 400 mg are equivalent for the purposes of substitution.

Note No increase in the maximum number of repeats may be authorised.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Chronic Myeloid Leukaemia (CML)

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have a primary diagnosis of chronic myeloid leukaemia, **AND**
- The condition must be in the accelerated phase, **AND**
- The condition must be expressing the Philadelphia chromosome; OR
- The condition must have the transcript BCR-ABL tyrosine kinase.

Accelerated phase is defined by the presence of 1 or more of the following:

1. Percentage of blasts in the peripheral blood or bone marrow greater than or equal to 15% but less than 30%; or
2. Percentage of blasts plus promyelocytes in the peripheral blood or bone marrow greater than or equal to 30%, provided that blast count is less than 30%; or
3. Peripheral basophils greater than or equal to 20%; or
4. Progressive splenomegaly to a size greater than or equal to 10 cm below the left costal margin to be confirmed on 2 occasions at least 4 weeks apart, or a greater than or equal to 50% increase in size below the left costal margin over 4 weeks; or
5. Karyotypic evolution (chromosomal abnormalities in addition to a single Philadelphia chromosome).

Authority required

Chronic Myeloid Leukaemia (CML)

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must a primary diagnosis of chronic myeloid leukaemia, **AND**
- The condition must be in the blast phase, **AND**
- The condition must be expressing the Philadelphia chromosome; OR
- The condition must have the transcript BCR-ABL tyrosine kinase.

Blast crisis is defined as either:

1. Percentage of blasts in the peripheral blood or bone marrow greater than or equal to 30%; or
2. Extramedullary involvement other than spleen and liver.

imatinib 400 mg capsule, 30

10935M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	2	..	937.83	41.30	^a CIPLA IMATINIB ADULT [LR] ^a IMATINIB-DRLA [RZ]	^a Imatinib-APOTEX [TX] ^a Imatinib GH [GQ]

imatinib 400 mg tablet, 30

9116T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	2	..	937.83	41.30	^a Gilmat [CR] ^a IMATINIB RBX [RA]	^a Glivec [AF] ^a Imatinib-Teva [SZ]

■ IMATINIB

Note Pharmaceutical benefits that have the form imatinib tablet 400 mg and imatinib capsule 400 mg are equivalent for the purposes of substitution.

Note No increase in the maximum number of repeats may be authorised.

Note Authority applications for increased quantities/ repeats (where relevant) may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note Allogeneic stem cell transplantation is the preferred therapy for eligible patients achieving a complete remission of Philadelphia positive acute lymphoblastic leukaemia.

Authority required (STREAMLINED)

9207

Acute lymphoblastic leukaemia

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition; OR
- Patient must have experienced intolerance, not a failure to respond, to continuing PBS-subsidised treatment with dasatinib as a first-line therapy for this condition, **AND**
- The condition must be expressing the Philadelphia chromosome; OR
- The condition must have the transcript BCR-ABL, **AND**
- The treatment must be for maintenance of first complete remission, **AND**
- The treatment must be in combination with chemotherapy or corticosteroids.

Dasatinib and imatinib are available with a lifetime maximum of 24 months for continuing treatment for patients with acute lymphoblastic leukaemia reimbursed through the PBS in this treatment setting.

imatinib 400 mg capsule, 30

11771M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	2	..	937.83	41.30	^a CIPLA IMATINIB ADULT [LR] ^a IMATINIB-DRLA [RZ]	^a Imatinib-APOTEX [TX] ^a Imatinib GH [GQ]

imatinib 400 mg tablet, 30

11789L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	2	..	937.83	41.30	^a Gilmat [CR] ^a IMATINIB RBX [RA]	^a Glivec [AF] ^a Imatinib-Teva [SZ]

■ IMATINIB

Note Pharmaceutical benefits that have the form imatinib tablet 100 mg and imatinib capsule 100 mg are equivalent for the purposes of substitution.

Note No increase in the maximum number of repeats may be authorised.

Note Authority applications for increased quantities/ repeats (where relevant) may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note Allogeneic stem cell transplantation is the preferred therapy for eligible patients achieving a complete remission of Philadelphia positive acute lymphoblastic leukaemia.

Authority required (STREAMLINED)

9207

Acute lymphoblastic leukaemia

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition; OR
- Patient must have experienced intolerance, not a failure to respond, to continuing PBS-subsidised treatment with dasatinib as a first-line therapy for this condition, **AND**
- The condition must be expressing the Philadelphia chromosome; OR
- The condition must have the transcript BCR-ABL, **AND**
- The treatment must be for maintenance of first complete remission, **AND**
- The treatment must be in combination with chemotherapy or corticosteroids.

Dasatinib and imatinib are available with a lifetime maximum of 24 months for continuing treatment for patients with acute lymphoblastic leukaemia reimbursed through the PBS in this treatment setting.

imatinib 100 mg capsule, 60

11783E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	2	..	476.87	41.30	^a CIPLA IMATINIB ADULT [LR] ^a IMATINIB-DRLA [RZ]	^a Imatinib-APOTEX [TX] ^a Imatinib GH [GQ]

imatinib 100 mg tablet, 60

11780B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	2	..	476.87	41.30	^a Gilmat [CR] ^a IMATINIB RBX [RA]	^a Glivec [AF] ^a Imatinib-Teva [SZ]

■ IMATINIB

Note Pharmaceutical benefits that have the form imatinib tablet 400 mg and imatinib capsule 400 mg are equivalent for the purposes of substitution.

Note Authority applications for increased quantities/ repeats (where relevant) may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

10026

Chronic Myeloid Leukaemia (CML)

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The condition must be in the accelerated phase, **AND**
- The condition must be expressing the Philadelphia chromosome; OR
- The condition must have the transcript BCR-ABL tyrosine kinase.

Authority required (STREAMLINED)

10048

Chronic Myeloid Leukaemia (CML)

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The condition must be in the blast phase, **AND**
- The condition must be expressing the Philadelphia chromosome; OR
- The condition must have the transcript BCR-ABL tyrosine kinase.

imatinib 400 mg capsule, 30

11870R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	2	..	937.83	41.30	^a CIPLA IMATINIB ADULT [LR] ^a IMATINIB-DRLA [RZ]	^a Imatinib-APOTEX [TX] ^a Imatinib GH [GQ]

imatinib 400 mg tablet, 30

11878E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	2	..	937.83	41.30	^a Gilmat [CR] ^a IMATINIB RBX [RA]	^a Glivec [AF] ^a Imatinib-Teva [SZ]

■ **IMATINIB**

Note Pharmaceutical benefits that have the form imatinib tablet 100 mg and imatinib capsule 100 mg are equivalent for the purposes of substitution.

Note Authority applications for increased quantities/ repeats (where relevant) may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

10026

Chronic Myeloid Leukaemia (CML)

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The condition must be in the accelerated phase, **AND**
- The condition must be expressing the Philadelphia chromosome; OR
- The condition must have the transcript BCR-ABL tyrosine kinase.

Authority required (STREAMLINED)

10048

Chronic Myeloid Leukaemia (CML)

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The condition must be in the blast phase, **AND**
- The condition must be expressing the Philadelphia chromosome; OR
- The condition must have the transcript BCR-ABL tyrosine kinase.

imatinib 100 mg capsule, 60

11875B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	2	..	476.87	41.30	^a CIPLA IMATINIB ADULT [LR] ^a IMATINIB-DRLA [RZ]	^a Imatinib-APOTEX [TX] ^a Imatinib GH [GQ]

imatinib 100 mg tablet, 60

11880G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	2	..	476.87	41.30	^a Gilmat [CR] ^a IMATINIB RBX [RA]	^a Glivec [AF] ^a Imatinib-Teva [SZ]

■ **NILOTINIB**

Authority required

Chronic Myeloid Leukaemia (CML)

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be a primary diagnosis, **AND**
- The condition must be in the chronic phase, **AND**
- The condition must be expressing the Philadelphia chromosome; OR
- The condition must have the transcript BCR-ABL tyrosine kinase, **AND**
- The treatment must be for first line therapy for this condition, **AND**
- Patient must not have previously experienced a failure to respond to the PBS-subsidised first line treatment with this drug for this condition; OR
- Patient must have experienced intolerance, not a failure to respond, to initial PBS-subsidised treatment with imatinib as a first line therapy for this condition; OR
- Patient must have experienced intolerance, not a failure to respond, to initial PBS-subsidised treatment with dasatinib as a first line therapy for this condition, **AND**
- The treatment must not exceed a total maximum of 18 months of therapy with a PBS-subsidised treatment with a tyrosine kinase inhibitor for this condition, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Applications under this restriction will be limited to provide patients with a maximum of 18 months of therapy with dasatinib, imatinib or nilotinib from the date the first application for initial treatment was approved. Patients should be commenced on a dose of nilotinib of 300 mg twice daily. Continuing therapy is dependent on patients demonstrating a response to nilotinib therapy following the initial 18 months of treatment and at 12 monthly intervals thereafter. Applications for authorisation must be in writing and must include: (1) a completed authority prescription form; and (2) a completed Chronic Myeloid Leukaemia - Chronic Phase, First Line - Supporting Information form; and (3) a pathology cytogenetic report conducted on peripheral blood or bone marrow supporting the diagnosis of chronic myeloid leukaemia to confirm eligibility for treatment, or a qualitative PCR report documenting the presence of the BCR-ABL transcript in either peripheral blood or bone marrow; and (4) a signed patient acknowledgement form. The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of tyrosine kinase inhibitors (TKI) agents for the chronic phase of chronic myeloid leukaemia. Where the term TKI agent appears in the following notes and restrictions it refers to imatinib mesilate, dasatinib or nilotinib.

Patients are eligible for PBS-subsidised treatment with only one TKI agent at any one time and must not be receiving concomitant interferon alfa therapy. Eligible patients may only swap between TKI agents if they have not failed prior PBS-subsidised treatment with that agent.

1. Initial First-line treatment From 1 April 2012, under the PBS, a patient will be able to be prescribed any of imatinib mesilate, dasatinib or nilotinib within the initial 18 month treatment period, as long as only one agent is used at a time and providing the patient has not failed to respond to any one of these TKIs. During the initial 18 month treatment period, switching between approved first-line agents may only occur for reasons of intolerance, not failure to respond. 2. Continuing First-line treatment -

Patients must maintain a major cytogenetic response or have a peripheral blood BCR-ABL of less than 1% to receive continuing therapy.

First continuing applications are to be written and must include a pathology report demonstrating the patient has responded to the initial course of treatment.

Second and subsequent authority applications for continuing therapy may be made by telephoning the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

During continuing therapy beyond the initial 18 month treatment period, switching between approved first-line agents may only occur for reason of intolerance. Where there is failure to respond, switching may only occur through application for prescription of second-line agents. Where a patient has previously received PBS-subsidised treatment with imatinib mesilate, dasatinib or nilotinib no approval will be granted for PBS-subsidised re-treatment in the chronic phase of chronic myeloid leukaemia, where that patient has at any time failed to meet the response criteria whilst on that TKI agent.

3. Authority approval requirements. Response criteria to initial first-line treatment with imatinib mesilate, dasatinib or nilotinib: For the purposes of assessing response to PBS-subsidised treatment with imatinib mesilate, dasatinib or nilotinib either cytogenetic analysis indicating the number of Philadelphia positive [t(9;22)] cells in the bone marrow measured by standard karyotyping, or quantitative PCR indicating the relative level of BCR-ABL transcript in the peripheral blood using the international scale, must be submitted. For bone marrow analyses, where the standard karyotyping is not informative for technical reasons, a cytogenetic analysis performed on the bone marrow by the use of fluorescence in situ hybridisation (FISH) with BCR-ABL specific probe must be submitted. The cytogenetic or peripheral blood quantitative PCR analyses must be submitted within 18 months of the commencement of treatment with imatinib mesilate, dasatinib or nilotinib (patients in whom a major cytogenetic response or peripheral blood BCR-ABL level of less than 1% is demonstrable by 18 months are eligible to receive continuing treatment with that agent).

4. Definitions of response. A major cytogenetic response is defined as less than 35% Philadelphia positive bone marrow cells. A peripheral blood BCR-ABL level of less than 1% on the international scale (Blood 108: 28-37, 2006) also indicates a response, at least the biological equivalent of a major cytogenetic response.

5. Definitions of loss of response. Loss of a previously documented major cytogenetic response (demonstrated by the presence of greater than 35% Ph positive cells on bone marrow biopsy), during ongoing tyrosine kinase inhibitor (TKI) therapy. Loss of a previously demonstrated molecular response (demonstrated by peripheral blood BCR-ABL levels increasing consecutively in value by at least 5 fold to a level of greater than 0.1% confirmed on a subsequent test), during ongoing tyrosine kinase inhibitor therapy.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Chronic Myeloid Leukaemia (CML)

Treatment Phase: First continuing treatment

Clinical criteria:

- The condition must be in the chronic phase, **AND**
- Patient must have received initial PBS-subsidised first line treatment with this drug for this condition; OR
- Patient must have experienced intolerance, not a failure to respond, to first continuing PBS-subsidised treatment with imatinib as a first line therapy for this condition; OR
- Patient must have experienced intolerance, not a failure to respond, to first continuing PBS-subsidised treatment with dasatinib as a first line therapy for this condition, **AND**
- Patient must have demonstrated a major cytogenetic response; OR
- Patient must have demonstrated a peripheral blood level of BCR-ABL of less than 1%, **AND**
- The treatment must not exceed a total maximum of 24 weeks of therapy with a PBS-subsidised treatment with a tyrosine kinase inhibitor for this condition under this restriction, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

First continuing applications for authorisation must be in writing and must include:(1) a completed authority prescription form; and(2) demonstration of continued response to treatment as evidenced by either:(a) a major cytogenetic response [see Note explaining requirements]; or(b) a peripheral blood level of BCR-ABL of less than 1% on the international scale [see Note explaining requirements].Where this has been supplied within the previous 12 months, only the date of the relevant pathology report need be provided.

Authority required

Chronic Myeloid Leukaemia (CML)

Treatment Phase: Subsequent continuing treatment

Clinical criteria:

- The condition must be in the chronic phase, **AND**
- Patient must have received the First continuing PBS-subsidised treatment with this drug as a first line therapy for this condition; OR
- Patient must have experienced intolerance, not a failure to respond, to subsequent continuing PBS-subsidised treatment with imatinib as a first line therapy for this condition; OR
- Patient must have experienced intolerance, not a failure to respond, to subsequent continuing PBS-subsidised treatment with dasatinib as a first line therapy for this condition, **AND**
- Patient must have maintained a major cytogenetic response; OR
- Patient must have maintained a peripheral blood level of BCR-ABL of less than 1%, **AND**
- The treatment must not exceed a total maximum of 24 weeks of therapy with a PBS-subsidised treatment with a tyrosine kinase inhibitor for this condition under this restriction, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Subsequent authority applications for continuing therapy with this drug may be made by telephoning the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

nilotinib 150 mg capsule, 120

1309X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	3856.05	41.30	Tasigna [NV]

■ NILOTINIB

Note Any queries concerning the arrangements to prescribe this drug may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Written applications for authority to prescribe this drug should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Chronic Myeloid Leukaemia (CML)

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be in the chronic phase; OR
- The condition must be in the accelerated phase, **AND**
- Patient must have failed an adequate trial of PBS-subsidised first line treatment with imatinib for this condition; OR
- Patient must have failed an adequate trial of PBS-subsidised first line treatment with dasatinib for this condition, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Failure of an adequate trial of imatinib or dasatinib is defined as:(i) Lack of response to initial imatinib or dasatinib therapy, defined as either:- failure to achieve a haematological response after a minimum of 3 months therapy with imatinib or

dasatinib for patients initially treated in chronic phase; or- failure to achieve any cytogenetic response after a minimum of 6 months therapy with imatinib or dasatinib for patients initially treated in chronic phase as demonstrated on bone marrow biopsy by presence of greater than 95% Philadelphia chromosome positive cells; or- failure to achieve a major cytogenetic response or a peripheral blood BCR-ABL level of less than 1% after a minimum of 12 months therapy with imatinib or dasatinib; OR(ii) Loss of a previously documented major cytogenetic response (demonstrated by the presence of greater than 35% Ph positive cells on bone marrow biopsy), during ongoing imatinib or dasatinib therapy; OR(iii) Loss of a previously demonstrated molecular response (demonstrated by peripheral blood BCR-ABL levels increasing consecutively in value by at least 5 fold to a level of greater than 0.1% confirmed on a subsequent test), during ongoing imatinib or dasatinib therapy; OR(iv) Development of accelerated phase in a patient previously prescribed imatinib or dasatinib for the chronic phase of chronic myeloid leukaemia.

Accelerated phase is defined by the presence of 1 or more of the following:(1) Percentage of blasts in the peripheral blood or bone marrow greater than or equal to 15% but less than 30%; or(2) Percentage of blasts plus promyelocytes in the peripheral blood or bone marrow greater than or equal to 30%, provided that blast count is less than 30%; or(3) Peripheral basophils greater than or equal to 20%; or(4) Progressive splenomegaly to a size greater than or equal to 10 cm below the left costal margin to be confirmed on 2 occasions at least 4 weeks apart, or a greater than or equal to 50% increase in size below the left costal margin over 4 weeks; or(5) Karyotypic evolution (chromosomal abnormalities in addition to a single Philadelphia chromosome); OR(v) Disease progression (defined as a greater than or equal to .50% increase in peripheral white blood cell count, blast count, basophils or platelets) during first-line imatinib or dasatinib therapy in patients with accelerated phase chronic myeloid leukaemia, provided that blast crisis has been excluded on bone marrow biopsy.

Patients should be commenced on a dose of nilotinib of 400 mg twice daily. Continuing therapy is dependent on patients demonstrating a major cytogenetic response to nilotinib therapy or a peripheral blood BCR-ABL level of less than 1% within 18 months and thereafter at 12 monthly intervals.

Applications for authorisation must be in writing and must include:(a) a completed authority prescription form; and(b) a completed Chronic Myeloid Leukaemia - Second and Third Line - Supporting Information Form; and(c) a signed patient acknowledgement; and(d) a bone marrow biopsy pathology report demonstrating the patient has active chronic myeloid leukaemia, either manifest as cytogenetic evidence of the Philadelphia chromosome, or RT-PCR level of BCR-ABL transcript greater than 0.1% on the international scale. (The date of the relevant pathology report needs to be provided); and(e) where there has been a loss of response to imatinib or dasatinib, a copy of the current confirming pathology report(s) from an Approved Pathology Authority or details of the dates of assessment in the case of progressive splenomegaly or extramedullary involvement.

Note The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of tyrosine kinase inhibitors (TKI) agents for all phases of chronic myeloid leukaemia. Where the term TKI agent appears in the following notes and restrictions it refers to dasatinib or nilotinib. Imatinib mesilate is not approved for use in second or third line treatment. Patients are eligible for PBS-subsidised treatment with only one of dasatinib or nilotinib at any one time and must not be receiving concomitant interferon alfa therapy. Eligible patients may only swap between these agents if they have not failed prior PBS-subsidised treatment with that agent. Nilotinib is not approved for patients in blast crisis. 1. Initial second line treatment From 1 April 2012, under the PBS, a patient will be able to be prescribed either dasatinib or nilotinib within the initial 18 month treatment period as second-line therapy, as long as only one agent is approved at a time and providing the patient did not fail that drug as first-line therapy. During the initial 18 month treatment period, switching between approved second-line agents may only occur for reasons of intolerance, not failure of response. 2. Initial third line treatment Third-line treatment with a TKI can only be approved when imatinib is used for first-line treatment. Patients will only be approved for PBS-subsidised treatment with one third-line agent. From 1 April 2012, under the PBS, a patient will be able to be prescribed either dasatinib or nilotinib providing the patient did not fail that drug as first or second line therapy and for nilotinib the patient is not in blast crisis. 3. Continuing treatment for second and third line treatment All continuing applications are to be written and must include a pathology report demonstrating the patient has responded to PBS-subsidised treatment as follows:(i) within 18 months of the commencement of treatment, at which time patients in whom a major cytogenetic response or peripheral blood BCR-ABL level of less than 1% has been demonstrated may receive authorisation for a further 12 months of treatment; and(ii) at no greater than 12 month intervals thereafter, to demonstrate that the major cytogenetic response or peripheral blood BCR-ABL level of less than 1% has been sustained. During second line continuing treatment beyond the initial 18 month treatment period, switching between approved second line TKI agents may only occur for reason of intolerance. Where there is failure of response, switching may only occur through application for prescription of a third line agent. 4. Authority approval requirements. Response criteria to initial treatment with dasatinib or nilotinib: For the purposes of assessing response to PBS-subsidised treatment with dasatinib or nilotinib, either cytogenetic analysis indicating the number of Philadelphia positive [t (9;22)] cells in the bone marrow measured by standard karyotyping, or quantitative PCR indicating the relative level of BCR-ABL transcript in the peripheral blood using the international scale, must be submitted. For bone marrow analyses, where the standard karyotyping is not informative for technical reasons, a cytogenetic analysis performed on the bone marrow by the use of fluorescence in situ hybridisation (FISH) with BCR-ABL specific probe must be submitted. The cytogenetic or peripheral blood quantitative PCR analyses must be submitted within 18 months of the commencement of treatment with dasatinib or nilotinib (patients in whom a major cytogenetic response or peripheral blood BCR-ABL level of less than 1% is demonstrable by 18 months are eligible to receive continuing treatment with that agent). 5. Definitions of response. A major cytogenetic response is defined as less than 35% Philadelphia positive bone marrow cells. A peripheral blood BCR-ABL level of less than 1% on the international scale (Blood 108: 28-37, 2006) also indicates a response, at least the biological equivalent of a major cytogenetic response. 6. Definitions of loss of response. Loss of a previously documented major cytogenetic response (demonstrated by the presence of greater than 35% Ph positive cells on bone marrow biopsy), during ongoing tyrosine kinase inhibitor (TKI) therapy. Loss of a previously demonstrated molecular response (demonstrated by peripheral blood BCR-ABL levels increasing consecutively in value by at least 5 fold to a level of greater than 0.1% confirmed on a subsequent test), during ongoing tyrosine kinase inhibitor therapy.

Authority required

Chronic Myeloid Leukaemia (CML)

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**

- Patient must have demonstrated a major cytogenetic response to nilotinib in the preceding 18 months and thereafter at 12 monthly intervals; OR
- Patient must have achieved a peripheral blood level of BCR-ABL of less than 1% to nilotinib in the preceding 18 months and thereafter at 12 monthly intervals, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Applications for authorisation must be in writing and must include: (1) a completed authority prescription form; and (2) a completed Chronic Myeloid Leukaemia - Second and Third Line - Application Form for continuing treatment; and (3) demonstration of continued response to treatment as evidenced by either: (a) major cytogenetic response [see Note explaining definitions of response]. Where this has been supplied within the previous 12 months (or 18 months for the initial supply), only the date of the relevant pathology report needs to be provided; or (b) a peripheral blood level of BCR-ABL of less than 1% on the international scale [see Note explaining definitions of response]. Where this has been supplied within the previous 12 months (or 18 months for the initial supply), only the date of the relevant pathology report needs to be provided

nilotinib 200 mg capsule, 120

9171Q	Max. Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	5056.70	41.30	Tasigna [NV]

■ PONATINIB**Authority required**

Acute lymphoblastic leukaemia

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be expressing the Philadelphia chromosome; OR
- The condition must have the transcript BCR-ABL, **AND**
- Patient must have failed prior treatment with PBS-subsidised dasatinib for this condition; OR
- Patient must have developed intolerance to PBS-subsidised dasatinib of a severity requiring treatment withdrawal.

Failure of treatment with dasatinib is defined as either:

1. Failure to achieve a complete morphological and cytogenetic remission after a minimum of 2 months treatment with PBS-subsidised dasatinib for this condition; or
2. Morphological or cytogenetic relapse of leukaemia after achieving a complete remission induced by PBS-subsidised dasatinib for this condition; or
3. Rising levels of BCR-ABL1 transcript on two consecutive occasions in a patient in complete remission while being treated with PBS-subsidised dasatinib for this condition.

Patients must have active leukaemia, as defined by presence on current pathology assessments of either morphological infiltration of the bone marrow (greater than 5% lymphoblasts) or cerebrospinal fluid or other sites; OR the presence of cells bearing the Philadelphia chromosome on cytogenetic or FISH analysis in the bone marrow of patients in morphological remission; OR rising levels of BCR-ABL1 transcript on two consecutive occasions in a patient in complete remission while being treated with PBS-subsidised dasatinib for this condition.

The authority application must be made in writing and must include:

1. a completed authority prescription form; and
2. a completed Acute Lymphoblastic Leukaemia ponatinib PBS Authority Application - Supporting Information Form; and
3. a pathology report demonstrating that the patient has active acute lymphoblastic leukaemia, manifest as cytogenetic evidence of the Philadelphia chromosome, or morphological evidence of acute lymphoblastic leukaemia plus qualitative RT-PCR evidence of BCR-ABL transcript. The date of the relevant pathology report(s) need(s) to be provided; or
4. pathology reports documenting rising levels of BCR-ABL1 transcript on two consecutive occasions in a patient in complete remission while being treated with PBS-subsidised dasatinib for this condition. The date of the relevant pathology report(s) need(s) to be provided

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Acute lymphoblastic leukaemia

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have progressive disease while receiving PBS-subsidised treatment with this drug for this condition.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

ponatinib 15 mg tablet, 60

11454W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	5489.34	41.30	Iclusig [TK]

ponatinib 45 mg tablet, 30

11453T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	6172.81	41.30	Iclusig [TK]

▪ **PONATINIB**

Authority required

Acute lymphoblastic leukaemia

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must be expressing the T315I mutation, **AND**
- Patient must have failed treatment with chemotherapy, with or without another tyrosine kinase inhibitor, **AND**
- Patient must have failed allogeneic haemopoietic stem cell transplantation (where appropriate).

Failure of treatment is defined as either:

1. Failure to achieve a complete morphological and cytogenetic remission after a minimum of 2 months treatment with intensive chemotherapy, with or without another tyrosine kinase inhibitor;
 2. Morphological or cytogenetic relapse of leukaemia after achieving a complete remission induced by chemotherapy, with or without another tyrosine kinase inhibitor;
 3. Morphological or cytogenetic relapse or persistence of leukaemia after allogeneic haemopoietic stem cell transplantation.
- Patients must have active leukaemia, as defined by presence on current pathology assessments of either morphological infiltration of the bone marrow (greater than 5% lymphoblasts) or cerebrospinal fluid or other sites; OR the presence of cells bearing the Philadelphia chromosome on cytogenetic or FISH analysis in the bone marrow of patients in morphological remission.

The authority application must be made in writing and must include:

1. a completed authority prescription form; and
2. a completed Acute Lymphoblastic Leukaemia - ponatinib Initial PBS authority application form; and
3. a signed patient acknowledgement; and
4. a pathology report demonstrating that the patient has active acute lymphoblastic leukaemia, either manifest as cytogenetic evidence of the Philadelphia chromosome, or morphological evidence of acute lymphoblastic leukaemia plus qualitative RT-PCR evidence of BCR-ABL transcript.; and evidence of the T315I mutation. The date of the relevant pathology report(s), which should be within the previous 6 months, need(s) to be provided

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Acute lymphoblastic leukaemia

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously been issued with an authority prescription for this drug for this condition, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must not have progressive disease.

Note Authority applications for continuing treatment may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au
Applications for authority to prescribe should be forwarded to:
Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

ponatinib 15 mg tablet, 60

10523W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	5489.34	41.30	Iclusig [TK]

ponatinib 45 mg tablet, 30

10524X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	6172.81	41.30	Iclusig [TK]

■ PONATINIB

Note Patients are eligible for PBS-subsidised treatment with only one of imatinib, dasatinib, nilotinib or ponatinib at any one time and must not be receiving concomitant interferon alfa therapy.

Note 1. Continuing treatment

First continuing applications are to be written, at which time patients in whom a major cytogenetic response or peripheral blood BCR-ABL level of less than 1% has been demonstrated may receive authorisation for a further 12 months of treatment. Subsequent authority applications for continuing therapy may be made by telephoning the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

2. Authority approval requirements.

Response criteria to treatment with ponatinib:

For the purposes of assessing response to PBS-subsidised treatment with ponatinib, either cytogenetic analysis indicating the number of Philadelphia positive [t (9;22)] cells in the bone marrow measured by standard karyotyping, or quantitative PCR indicating the relative level of BCR-ABL transcript in the peripheral blood using the international scale, must be submitted. For bone marrow analyses, where the standard karyotyping is not informative for technical reasons, a cytogenetic analysis performed on the bone marrow by the use of fluorescence in situ hybridisation (FISH) with BCR-ABL specific probe must be submitted. The cytogenetic or peripheral blood quantitative PCR analyses must be conducted within 18 months of the commencement of treatment with ponatinib (patients in whom a major cytogenetic response or peripheral blood BCR-ABL level of less than 1% is demonstrable by 18 months are eligible to receive first continuing treatment with this drug).

Thereafter, at no greater than 12 month intervals a major cytogenetic response or a peripheral blood BCR-ABL level of less than 1% must be sustained to receive subsequent continuing treatments with this drug.

3. Definitions of response.

A major cytogenetic response is defined as less than 35% Philadelphia positive bone marrow cells.

A peripheral blood BCR-ABL level of less than 1% on the international scale (Blood 108: 28-37, 2006).

4. Definitions of loss of response.

Loss of a previously documented major cytogenetic response (demonstrated by the presence of greater than 35% Ph positive cells on bone marrow biopsy), during ongoing tyrosine kinase inhibitor therapy.

Loss of a previously demonstrated molecular response (demonstrated by peripheral blood BCR-ABL levels increasing consecutively in value by at least 5 fold to a level of greater than 0.1% confirmed on a subsequent test), during ongoing tyrosine kinase inhibitor therapy.

Authority required

Chronic Myeloid Leukaemia (CML)

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have failed an adequate trial of dasatinib; OR
- Patient must have developed intolerance to dasatinib of a severity necessitating permanent treatment withdrawal, **AND**
- Patient must have failed an adequate trial of nilotinib; OR
- Patient must have developed intolerance to nilotinib of a severity necessitating permanent treatment withdrawal; OR
- Patient must not be eligible for PBS-subsidised treatment with nilotinib because the patient has a blast crisis.

Failure of an adequate trial of dasatinib or nilotinib is defined as:

1. Lack of response to dasatinib or nilotinib therapy, defined as either:

- (i) failure to achieve a haematological response after a minimum of 3 months therapy with dasatinib or nilotinib; or
- (ii) failure to achieve any cytogenetic response after a minimum of 6 months therapy with dasatinib or nilotinib as demonstrated on bone marrow biopsy by presence of greater than 95% Philadelphia chromosome positive cells; or
- (iii) failure to achieve a major cytogenetic response or a peripheral blood BCR-ABL level of less than 1% after a minimum of 12 months therapy with dasatinib or nilotinib; OR

2. Loss of a previously documented major cytogenetic response (demonstrated by the presence of greater than 35% Ph positive cells on bone marrow biopsy), during ongoing dasatinib or nilotinib therapy; OR

3. Loss of a previously demonstrated molecular response (demonstrated by peripheral blood BCR-ABL levels increasing consecutively in value by at least 5 fold to a level of greater than 0.1% confirmed on a subsequent test), during ongoing dasatinib or nilotinib therapy; OR

4. Development of accelerated phase or blast crisis in a patient previously prescribed dasatinib or nilotinib for any phase of chronic myeloid leukaemia; OR

5. Disease progression (defined as a greater than or equal to 50% increase in peripheral white blood cell count, blast count, basophils or platelets) during dasatinib or nilotinib therapy in patients with accelerated phase or blast crisis chronic myeloid leukaemia.

Accelerated phase is defined by the presence of 1 or more of the following:

- 1. Percentage of blasts in the peripheral blood or bone marrow greater than or equal to 15% but less than 30%; or
- 2. Percentage of blasts plus promyelocytes in the peripheral blood or bone marrow greater than or equal to 30%, provided that blast count is less than 30%; or
- 3. Peripheral basophils greater than or equal to 20%; or
- 4. Progressive splenomegaly to a size greater than or equal to 10 cm below the left costal margin to be confirmed on 2 occasions at least 4 weeks apart, or a greater than or equal to 50% increase in size below the left costal margin over 4 weeks; or

5. Karyotypic evolution (chromosomal abnormalities in addition to a single Philadelphia chromosome).

Blast crisis is defined as either:

1. Percentage of blasts in the peripheral blood or bone marrow greater than or equal to 30%; or
2. Extramedullary involvement other than spleen and liver.

The authority application must be made in writing and must include:

1. a completed authority prescription form; and
2. a completed Chronic Myeloid Leukaemia - ponatinib Initial PBS authority application form; and
3. a bone marrow biopsy pathology report demonstrating the patient has active chronic myeloid leukaemia, either manifest as cytogenetic evidence of the Philadelphia chromosome, or RT-PCR level of BCR-ABL transcript greater than 0.1% on the international scale. (The date of the relevant pathology report needs to be provided); and
4. where there has been a loss of response to dasatinib or nilotinib, a copy of the current confirming pathology report(s) from an Approved Pathology Authority or details of the dates of assessment in the case of progressive splenomegaly or extramedullary involvement.

Up to a maximum of 18 months of treatment may be authorised under this initial restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Chronic Myeloid Leukaemia (CML)

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must be expressing the T315I mutation, **AND**
- Patient must have failed an adequate trial of imatinib; OR
- Patient must have failed an adequate trial of dasatinib; OR
- Patient must have failed an adequate trial of nilotinib.

Failure of an adequate trial of imatinib or dasatinib or nilotinib is defined as:

1. Lack of response to imatinib or dasatinib or nilotinib therapy, defined as either:

(i) failure to achieve a haematological response after a minimum of 3 months therapy with imatinib or dasatinib or nilotinib; or

(ii) failure to achieve any cytogenetic response after a minimum of 6 months therapy with imatinib or dasatinib or nilotinib as demonstrated on bone marrow biopsy by presence of greater than 95% Philadelphia chromosome positive cells; or

(iii) failure to achieve a major cytogenetic response or a peripheral blood BCR-ABL level of less than 1% after a minimum of 12 months therapy with imatinib or dasatinib or nilotinib; OR

2. Loss of a previously documented major cytogenetic response (demonstrated by the presence of greater than 35% Ph positive cells on bone marrow biopsy), during ongoing imatinib or dasatinib or nilotinib therapy; OR

3. Loss of a previously demonstrated molecular response (demonstrated by peripheral blood BCR-ABL levels increasing consecutively in value by at least 5 fold to a level of greater than 0.1% confirmed on a subsequent test), during ongoing imatinib or dasatinib or nilotinib therapy; OR

4. Development of accelerated phase or blast crisis in a patient previously prescribed imatinib or dasatinib or nilotinib for any phase of chronic myeloid leukaemia; OR

5. Disease progression (defined as a greater than or equal to 50% increase in peripheral white blood cell count, blast count, basophils or platelets) during imatinib or dasatinib or nilotinib therapy in patients with accelerated phase or blast crisis chronic myeloid leukaemia.

Accelerated phase is defined by the presence of 1 or more of the following:

1. Percentage of blasts in the peripheral blood or bone marrow greater than or equal to 15% but less than 30%; or
2. Percentage of blasts plus promyelocytes in the peripheral blood or bone marrow greater than or equal to 30%, provided that blast count is less than 30%; or
3. Peripheral basophils greater than or equal to 20%; or
4. Progressive splenomegaly to a size greater than or equal to 10 cm below the left costal margin to be confirmed on 2 occasions at least 4 weeks apart, or a greater than or equal to 50% increase in size below the left costal margin over 4 weeks; or
5. Karyotypic evolution (chromosomal abnormalities in addition to a single Philadelphia chromosome).

Blast crisis is defined as either:

1. Percentage of blasts in the peripheral blood or bone marrow greater than or equal to 30%; or
2. Extramedullary involvement other than spleen and liver.

The authority application must be made in writing and must include:

1. a completed authority prescription form; and
2. a completed Chronic Myeloid Leukaemia - ponatinib Initial PBS authority application form; and

3. a bone marrow biopsy pathology report demonstrating the patient has active chronic myeloid leukaemia, either manifest as cytogenetic evidence of the Philadelphia chromosome, or RT-PCR level of BCR-ABL transcript greater than 0.1% on the international scale and evidence of the T315I mutation. (The date of the relevant pathology report(s) need to be provided); and

4. where there has been a loss of response to imatinib or dasatinib or nilotinib, a copy of the current confirming pathology report(s) from an Approved Pathology Authority or details of the dates of assessment in the case of progressive splenomegaly or extramedullary involvement.

Up to a maximum of 18 months of treatment may be authorised under this initial restriction.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs Programs

Reply Paid 9826

HOBART TAS 7001

Authority required

Chronic Myeloid Leukaemia (CML)

Treatment Phase: First continuing treatment

Clinical criteria:

- Patient must have received initial PBS subsidised treatment with this drug for this condition, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have demonstrated a major cytogenetic response; OR
- Patient must have demonstrated a peripheral blood level of BCR-ABL of less than 1%.

First continuing applications for authorisation must be in writing and must include:

1. a completed authority prescription form; and
2. a completed Chronic Myeloid Leukaemia Continuing PBS authority application Supporting information form; and
3. demonstration of continued response to treatment as evidenced by either:
 - (a) major cytogenetic response [see Note explaining definitions of response]; or
 - (b) a peripheral blood level of BCR-ABL of less than 1% on the international scale [see Note explaining definitions of response].

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Chronic Myeloid Leukaemia (CML)

Treatment Phase: Subsequent continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have maintained a major cytogenic response at 12 month intervals; OR
- Patient must have maintained a peripheral blood level of BCR-ABL of less than 1% at 12 month intervals.

ponatinib 15 mg tablet, 60

10520Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	5489.34	41.30	Iclusig [TK]

ponatinib 45 mg tablet, 30

10530F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	6172.81	41.30	Iclusig [TK]

Epidermal growth factor receptor (EGFR) tyrosine kinase inhibitors

■ AFATINIB

Note Special Pricing Arrangements apply.

Authority required

Stage IIIB (locally advanced) or Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be as monotherapy, **AND**
- The condition must be non-squamous type non-small cell lung cancer (NSCLC) or not otherwise specified type NSCLC, **AND**
- Patient must not have received previous PBS-subsidised treatment with another epidermal growth factor receptor (EGFR) tyrosine kinase inhibitor (TKI); OR
- Patient must have developed intolerance to another epidermal growth factor receptor (EGFR) tyrosine kinase inhibitor (TKI) of a severity necessitating permanent treatment withdrawal, **AND**
- Patient must have a WHO performance status of 2 or less.

Population criteria:

- Patient must have evidence of an activating epidermal growth factor receptor (EGFR) gene mutation known to confer sensitivity to treatment with EGFR tyrosine kinase inhibitors in tumour material.

afatinib 30 mg tablet, 28

11341X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	2885.16	41.30	Giotrif [BY]

afatinib 20 mg tablet, 28

11335N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	2885.16	41.30	Giotrif [BY]

afatinib 40 mg tablet, 28

11359W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	2885.16	41.30	Giotrif [BY]

afatinib 50 mg tablet, 28

11329G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	2885.16	41.30	Giotrif [BY]

■ AFATINIB

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)**7613**

Stage IIIB (locally advanced) or Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Continuing treatment

Clinical criteria:

- The treatment must be as monotherapy, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have progressive disease while receiving PBS-subsidised treatment with this drug for this condition.

Population criteria:

- Patient must have evidence of an activating epidermal growth factor receptor (EGFR) gene mutation known to confer sensitivity to treatment with EGFR tyrosine kinase inhibitors in tumour material.

afatinib 30 mg tablet, 28

11348G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	2885.16	41.30	Giotrif [BY]

afatinib 20 mg tablet, 28

11336P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	2885.16	41.30	Giotrif [BY]

afatinib 40 mg tablet, 28

11347F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	2885.16	41.30	Giotrif [BY]

afatinib 50 mg tablet, 28

11342Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	2885.16	41.30	Giotrif [BY]

■ ERLOTINIB**Authority required**

Stage IIIB (locally advanced) or Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be as monotherapy, **AND**
- The condition must be non-squamous type non-small cell lung cancer (NSCLC) or not otherwise specified type NSCLC, **AND**
- Patient must not have received previous PBS-subsidised treatment with another epidermal growth factor receptor (EGFR) tyrosine kinase inhibitor (TKI); OR

- Patient must have developed intolerance to another epidermal growth factor receptor (EGFR) tyrosine kinase inhibitor (TKI) of a severity necessitating permanent treatment withdrawal, **AND**
- Patient must have a WHO performance status of 2 or less.

Population criteria:

- Patient must have evidence of an activating epidermal growth factor receptor (EGFR) gene mutation known to confer sensitivity to treatment with EGFR tyrosine kinase inhibitors in tumour material.

erlotinib 25 mg tablet, 30

10022L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	215.12	41.30	Erlotinib Sandoz [SZ]

erlotinib 100 mg tablet, 30

10020J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	762.32	41.30	Erlotinib Sandoz [SZ]

erlotinib 150 mg tablet, 30

10014C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	932.91	41.30	Erlotinib Sandoz [SZ]

■ ERLOTINIB**Authority required (STREAMLINED)****4600**

Stage IIIB (locally advanced) or Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Continuing treatment

Clinical criteria:

- The treatment must be as monotherapy, **AND**
- Patient must have previously been issued with an authority prescription for this drug prior to 1 August 2014, **AND**
- Patient must not have progressive disease.

Population criteria:

- Patient must have a wild type epidermal growth factor receptor (EGFR) gene; OR
- Patient must have an epidermal growth factor receptor (EGFR) gene of unknown type.

erlotinib 25 mg tablet, 30

10028T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	215.12	41.30	Erlotinib Sandoz [SZ]

erlotinib 100 mg tablet, 30

10019H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	762.32	41.30	Erlotinib Sandoz [SZ]

erlotinib 150 mg tablet, 30

10025P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	932.91	41.30	Erlotinib Sandoz [SZ]

■ ERLOTINIB**Authority required (STREAMLINED)****7446**

Stage IIIB (locally advanced) or Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Continuing treatment

Clinical criteria:

- The treatment must be as monotherapy, **AND**
- Patient must have received an initial authority prescription for this drug for this condition, **AND**
- Patient must not have progressive disease.

Population criteria:

- Patient must have evidence of an activating epidermal growth factor receptor (EGFR) gene mutation known to confer sensitivity to treatment with EGFR tyrosine kinase inhibitors in tumour material.

erlotinib 25 mg tablet, 30

11263T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	215.12	41.30	Erlotinib Sandoz [SZ]

erlotinib 100 mg tablet, 30

11260P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	762.32	41.30	Erlotinib Sandoz [SZ]

erlotinib 150 mg tablet, 30

11259N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	932.91	41.30	Erlotinib Sandoz [SZ]

▪ **GEFITINIB****Authority required (STREAMLINED)****7447**

Stage IIIB (locally advanced) or Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Continuing treatment

Clinical criteria:

- The treatment must be as monotherapy, **AND**
- Patient must have received an initial authority prescription for this drug for this condition, **AND**
- Patient must not have progressive disease.

gefitinib 250 mg tablet, 30

11264W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	3	..	926.61	41.30	^a Cipla Gefitinib [LR]	^a Iressa [AP]

▪ **GEFITINIB****Authority required**

Stage IIIB (locally advanced) or Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be as monotherapy, **AND**
- The condition must be non-squamous type non-small cell lung cancer (NSCLC) or not otherwise specified type NSCLC, **AND**
- Patient must not have received previous PBS-subsidised treatment with another epidermal growth factor receptor (EGFR) tyrosine kinase inhibitor (TKI); OR
- Patient must have developed intolerance to another epidermal growth factor receptor (EGFR) tyrosine kinase inhibitor (TKI) of a severity necessitating permanent treatment withdrawal, **AND**
- Patient must have a WHO performance status of 2 or less.

Population criteria:

- Patient must have evidence of an activating epidermal growth factor receptor (EGFR) gene mutation known to confer sensitivity to treatment with EGFR tyrosine kinase inhibitors in tumour material.

gefitinib 250 mg tablet, 30

8769M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	3	..	926.61	41.30	^a Cipla Gefitinib [LR]	^a Iressa [AP]

▪ **OSIMERTINIB****Note** No increase in the maximum quantity or number of units may be authorised.**Note** No increase in the maximum number of repeats may be authorised.**Note** Special Pricing Arrangements apply.**Note** Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.**Authority required**

Stage IIIB (locally advanced) or Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Continuing treatment of second-line EGFR tyrosine kinase inhibitor therapy

Clinical criteria:

- The treatment must be as monotherapy, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while receiving treatment with this drug for this condition.

Treatment criteria:

- Patient must be undergoing continuing treatment with this drug as second-line therapy (i.e. there are 2 Continuing treatment listings for this drug - ensure the correct Continuing treatment restriction is being accessed).

osimertinib 40 mg tablet, 30

11620N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	7971.16	41.30	Tagrisso [AP]

▪ **OSIMERTINIB****Note** No increase in the maximum quantity or number of units may be authorised.**Note** No increase in the maximum number of repeats may be authorised.**Note** Special Pricing Arrangements apply.**Note** Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.**Authority required**

Stage IIIB (locally advanced) or Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Initial treatment as second-line EGFR tyrosine kinase inhibitor therapy

Clinical criteria:

- Patient must not have previously received this drug for this condition, **AND**

- The treatment must be as monotherapy, **AND**
- Patient must have a WHO performance status of 2 or less, **AND**
- The condition must have progressed on or after prior epidermal growth factor receptor (EGFR) tyrosine kinase inhibitor (TKI) therapy as first line treatment for this condition, **AND**
- Patient must have evidence of EGFR T790M mutation in tumour material at the point of progression on or after first line EGFR TKI treatment.

Authority required

Stage IIIB (locally advanced) or Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Continuing treatment of second-line EGFR tyrosine kinase inhibitor therapy

Clinical criteria:

- The treatment must be as monotherapy, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while receiving treatment with this drug for this condition.

Treatment criteria:

- Patient must be undergoing continuing treatment with this drug as second-line therapy (i.e. there are 2 Continuing treatment listings for this drug - ensure the correct Continuing treatment restriction is being accessed).

osimertinib 80 mg tablet, 30

11622Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	7971.16	41.30	Tagrisso [AP]

■ OSIMERTINIB

Note No increase in the maximum number of repeats may be authorised.

Note No increase in the maximum quantity or number of units may be authorised.

Note Special Pricing Arrangements apply.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Authority required

Stage IIIB (locally advanced) or Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Continuing treatment of first-line EGFR tyrosine kinase inhibitor therapy

Clinical criteria:

- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while receiving treatment with this drug for this condition.

Treatment criteria:

- Patient must be undergoing continuing treatment with this drug as first-line therapy (i.e. there are 2 Continuing treatment listings for this drug - ensure the correct Continuing treatment restriction is being accessed).

Authority required

Stage IIIB (locally advanced) or Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Transitioning from non-PBS to PBS-subsidised supply as first-line EGFR tyrosine kinase inhibitor therapy - 'Grandfather' treatment

Clinical criteria:

- Patient must have received non-PBS subsidised treatment with this drug as first-line EGFR tyrosine kinase inhibitor therapy for this PBS indication prior to 1 January 2021, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have had a WHO performance status of 2 or less prior to initiating non-PBS-subsidised treatment, **AND**
- Patient must not have developed disease progression while receiving non-PBS-subsidised treatment with this drug for this condition.

Population criteria:

- Patient must have evidence in tumour material of an activating epidermal growth factor receptor (EGFR) gene mutation known to confer sensitivity to treatment with EGFR tyrosine kinase inhibitors that was obtained prior to initiating non-PBS subsidised treatment.

A Grandfathered patient may qualify for PBS-subsidised treatment under this restriction once only. For continuing PBS-subsidised treatment, a Grandfathered patient must qualify under the continuing treatment criteria.

Note This grandfather restriction will cease to operate from 12 months after the date specified in the clinical criteria.

osimertinib 40 mg tablet, 30

12233W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	7971.16	41.30	Tagrisso [AP]

■ OSIMERTINIB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Authority required

Stage IIIB (locally advanced) or Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Initial treatment as first-line epidermal growth factor receptor tyrosine kinase inhibitor therapy

Clinical criteria:

- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have a WHO performance status of 2 or less, **AND**
- Patient must not have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have received previous PBS-subsidised treatment with another epidermal growth factor receptor (EGFR) tyrosine kinase inhibitor (TKI); OR
- Patient must have developed intolerance to another epidermal growth factor receptor (EGFR) tyrosine kinase inhibitor (TKI) of a severity necessitating permanent treatment withdrawal.

Population criteria:

- Patient must have evidence in tumour material of an activating epidermal growth factor receptor (EGFR) gene mutation known to confer sensitivity to treatment with EGFR tyrosine kinase inhibitors.

Authority required

Stage IIIB (locally advanced) or Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Continuing treatment of first-line EGFR tyrosine kinase inhibitor therapy

Clinical criteria:

- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while receiving treatment with this drug for this condition.

Treatment criteria:

- Patient must be undergoing continuing treatment with this drug as first-line therapy (i.e. there are 2 Continuing treatment listings for this drug - ensure the correct Continuing treatment restriction is being accessed).

Authority required

Stage IIIB (locally advanced) or Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Transitioning from non-PBS to PBS-subsidised supply as first-line EGFR tyrosine kinase inhibitor therapy - 'Grandfather' treatment

Clinical criteria:

- Patient must have received non-PBS subsidised treatment with this drug as first-line EGFR tyrosine kinase inhibitor therapy for this PBS indication prior to 1 January 2021, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have had a WHO performance status of 2 or less prior to initiating non-PBS-subsidised treatment, **AND**
- Patient must not have developed disease progression while receiving non-PBS-subsidised treatment with this drug for this condition.

Population criteria:

- Patient must have evidence in tumour material of an activating epidermal growth factor receptor (EGFR) gene mutation known to confer sensitivity to treatment with EGFR tyrosine kinase inhibitors that was obtained prior to initiating non-PBS subsidised treatment.

A Grandfathered patient may qualify for PBS-subsidised treatment under this restriction once only. For continuing PBS-subsidised treatment, a Grandfathered patient must qualify under the continuing treatment criteria.

Note This grandfather restriction will cease to operate from 12 months after the date specified in the clinical criteria.

osimertinib 80 mg tablet, 30

12232T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	7971.16	41.30	Tagrisso [AP]

B-Raf serine-threonine kinase (BRAF) inhibitors

▪ **DABRAFENIB**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note A patient who has had progressive disease when treated with another BRAF inhibitor is not eligible to receive PBS-subsidised treatment with this drug.

Authority required (STREAMLINED)

10157

Unresectable Stage III or Stage IV malignant melanoma

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be positive for a BRAF V600 mutation, **AND**
- The condition must not have been treated previously with PBS-subsidised BRAF inhibitor therapy for unresectable Stage III or Stage IV disease; OR
- Patient must have developed intolerance to other BRAF inhibitors of a severity necessitating permanent treatment withdrawal, **AND**
- Patient must not have experienced disease progression whilst on adjuvant BRAF inhibitor treatment or disease recurrence within 6 months of completion of adjuvant BRAF inhibitor with MEK inhibitor treatment if previously treated for resected Stage IIIB, IIIC or IIID melanoma, **AND**
- Patient must have a WHO performance status of 2 or less.

dabrafenib 75 mg capsule, 120

2846T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	7524.56	41.30	Tafinlar [NV]

dabrafenib 50 mg capsule, 120

2963Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	5070.10	41.30	Tafinlar [NV]

■ DABRAFENIB

Note A patient who has progressive disease when treated with this drug is no longer eligible for PBS-subsidised treatment with this drug.

Note A patient who has had progressive disease when treated with another BRAF inhibitor is not eligible to receive PBS-subsidised treatment with this drug.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)**6013**

Unresectable Stage III or Stage IV malignant melanoma

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously been issued with an authority prescription for this drug, **AND**
- Patient must have stable or responding disease.

dabrafenib 75 mg capsule, 120

10003L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	7524.56	41.30	Tafinlar [NV]

dabrafenib 50 mg capsule, 120

2954L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	5070.10	41.30	Tafinlar [NV]

■ DABRAFENIB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Resected Stage IIIB, Stage IIIC or Stage IIID malignant melanoma

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be adjuvant to complete surgical resection, **AND**
- The condition must be positive for a BRAF V600 mutation, **AND**
- Patient must have a WHO performance status of 1 or less, **AND**
- Patient must be receiving PBS-subsidised trametinib and dabrafenib concomitantly for this condition, **AND**
- Patient must not have received prior PBS-subsidised treatment for this condition, **AND**
- The treatment must commence within 12 weeks of complete resection, **AND**
- Patient must not receive more than 12 months of combined PBS-subsidised and non-PBS-subsidised adjuvant therapy.

Authority required

Resected Stage IIIB, Stage IIIC or Stage IIID malignant melanoma

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously been issued with an authority prescription for trametinib and dabrafenib concomitantly for adjuvant treatment following complete surgical resection, **AND**
- Patient must not have experienced disease recurrence, **AND**
- Patient must not receive more than 12 months of combined PBS-subsidised and non-PBS-subsidised adjuvant therapy.

Authority required

Resected Stage IIIB, Stage IIIC or Stage IIID malignant melanoma

Treatment Phase: Grandfather treatment

Clinical criteria:

- Patient must have previously received non-PBS subsidised drug for adjuvant treatment following complete surgical resection prior to 1 November 2019, **AND**
- The condition must be positive for a BRAF V600 mutation, **AND**
- Patient must have a WHO performance status of 1 or less prior to starting non-PBS treatment with this drug, **AND**
- Patient must not have evidence of recurrence, **AND**
- Patient must be receiving PBS-subsidised trametinib and dabrafenib concomitantly for this condition, **AND**

- Patient must not have received prior PBS-subsidised treatment for this condition, **AND**
- Patient must have commenced non-PBS-subsidised treatment within 12 weeks of complete surgical resection, **AND**
- Patient must not receive more than 12 months of combined PBS-subsidised and non-PBS-subsidised adjuvant therapy.

dabrafenib 75 mg capsule, 120

11823G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	7524.56	41.30	Tafinlar [NV]

dabrafenib 50 mg capsule, 120

11820D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	5070.10	41.30	Tafinlar [NV]

■ ENCORAFENIB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note A patient who has had progressive disease when treated with another BRAF inhibitor is not eligible to receive PBS-subsidised treatment with this drug.

Authority required (STREAMLINED)**10271**

Unresectable Stage III or Stage IV malignant melanoma

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be positive for a BRAF V600 mutation, **AND**
- The condition must not have been treated previously with PBS-subsidised BRAF inhibitor therapy for unresectable Stage III or Stage IV disease; OR
- Patient must have developed intolerance to other BRAF inhibitors of a severity necessitating permanent treatment withdrawal, **AND**
- Patient must not have experienced disease progression whilst on adjuvant BRAF inhibitor treatment or disease recurrence within 6 months of completion of adjuvant BRAF inhibitor with MEK inhibitor treatment if previously treated for resected Stage IIIB, IIIC or IIID melanoma, **AND**
- Patient must have a WHO performance status of 2 or less.

encorafenib 50 mg capsule, 28

11937G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	9	3	..	*7033.68	41.30	Braftovi [FB]

encorafenib 75 mg capsule, 42

11938H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	4	3	..	*7033.70	41.30	Braftovi [FB]

■ ENCORAFENIB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note A patient who has progressive disease when treated with this drug is no longer eligible for PBS-subsidised treatment with this drug.

Note A patient who has had progressive disease when treated with another BRAF inhibitor is not eligible to receive PBS-subsidised treatment with this drug.

Authority required (STREAMLINED)**6013**

Unresectable Stage III or Stage IV malignant melanoma

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously been issued with an authority prescription for this drug, **AND**
- Patient must have stable or responding disease.

encorafenib 50 mg capsule, 28

11954E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	9	5	..	*7033.68	41.30	Braftovi [FB]

encorafenib 75 mg capsule, 42

11949X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	4	5	..	*7033.70	41.30	Braftovi [FB]

■ VEMURAFENIB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note A patient who has had progressive disease when treated with another BRAF inhibitor is not eligible to receive PBS-subsidised treatment with this drug.

Authority required (STREAMLINED)

10157

Unresectable Stage III or Stage IV malignant melanoma

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be positive for a BRAF V600 mutation, **AND**
- The condition must not have been treated previously with PBS-subsidised BRAF inhibitor therapy for unresectable Stage III or Stage IV disease; OR
- Patient must have developed intolerance to other BRAF inhibitors of a severity necessitating permanent treatment withdrawal, **AND**
- Patient must not have experienced disease progression whilst on adjuvant BRAF inhibitor treatment or disease recurrence within 6 months of completion of adjuvant BRAF inhibitor with MEK inhibitor treatment if previously treated for resected Stage IIIB, IIIC or IIID melanoma, **AND**
- Patient must have a WHO performance status of 2 or less.

vemurafenib 240 mg tablet, 56

11076Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	4	3	..	*7033.70	41.30	Zelboraf [RO]

▪ **MEMURAFENIB**

Note A patient who has progressive disease when treated with this drug is no longer eligible for PBS-subsidised treatment with this drug.

Note A patient who has had progressive disease when treated with another BRAF inhibitor is not eligible to receive PBS-subsidised treatment with this drug.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

6013

Unresectable Stage III or Stage IV malignant melanoma

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously been issued with an authority prescription for this drug, **AND**
- Patient must have stable or responding disease.

vemurafenib 240 mg tablet, 56

11081F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	4	5	..	*7033.70	41.30	Zelboraf [RO]

Anaplastic lymphoma kinase (ALK) inhibitors

▪ **ALECTINIB**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Authority required

Stage IIIB (locally advanced) or Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be as monotherapy, **AND**
- The condition must be non-squamous type non-small cell lung cancer (NSCLC) or not otherwise specified type NSCLC, **AND**
- Patient must have a WHO performance status of 2 or less.

Population criteria:

- Patient must have evidence of an anaplastic lymphoma kinase (ALK) gene rearrangement in tumour material, defined as 15% (or greater) positive cells by fluorescence in situ hybridisation (FISH) testing.

Authority required

Stage IIIB (locally advanced) or Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Continuing treatment

Clinical criteria:

- The treatment must be as monotherapy, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not develop disease progression while receiving PBS-subsidised treatment with this drug for this condition.

alectinib 150 mg capsule, 4 x 56

11226W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	6814.24	41.30	Alecensa [RO]

■ BRIGATINIB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Stage IIIB (locally advanced) or Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Continuing treatment

Clinical criteria:

- The treatment must be as monotherapy, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not develop disease progression while receiving PBS-subsidised treatment with this drug for this condition.

brigatinib 180 mg tablet, 28

11984R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	6814.24	41.30	Alunbrig [TK]

brigatinib 30 mg tablet, 28

11980M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	4	3	..	*6814.26	41.30	Alunbrig [TK]

brigatinib 90 mg tablet, 28

11974F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	6814.24	41.30	Alunbrig [TK]

■ BRIGATINIB

Caution Careful monitoring of patients is required due to risk of developing pulmonary adverse events observed in patients within the first seven days of treatment with this drug. Patients must be instructed to report any new or worsening respiratory symptoms.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Stage IIIB (locally advanced) or Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be as monotherapy, **AND**
- The condition must be non-squamous type non-small cell lung cancer (NSCLC) or not otherwise specified type NSCLC, **AND**
- Patient must have a WHO performance status of 2 or less.

Population criteria:

- Patient must have evidence of an anaplastic lymphoma kinase (ALK) gene rearrangement in tumour material, defined as 15% (or greater) positive cells by fluorescence in situ hybridisation (FISH) testing.

brigatinib 90 mg tablet [7] (&) brigatinib 180 mg tablet [21], 1 pack

11976H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	6814.24	41.30	Alunbrig [TK]

■ CERITINIB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Stage IIIB (locally advanced) or Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be as monotherapy, **AND**
- The condition must be non-squamous type non-small cell lung cancer (NSCLC) or not otherwise specified type NSCLC, **AND**
- Patient must have a WHO performance status of 2 or less.

Population criteria:

- Patient must have evidence of an anaplastic lymphoma kinase (ALK) gene rearrangement in tumour material, defined as 15% (or greater) positive cells by fluorescence in situ hybridisation (FISH) testing.

Authority required

Stage IIIB (locally advanced) or Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Continuing treatment

Clinical criteria:

- The treatment must be as monotherapy, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not develop disease progression while receiving PBS-subsidised treatment with this drug for this condition.

ceritinib 150 mg capsule, 3 x 50

11056X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	7289.46	41.30	Zykadia [NV]

■ CRIZOTINIB**Note** Special Pricing Arrangements apply.**Authority required**

Stage IIIB (locally advanced) or Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be as monotherapy, **AND**
- The condition must be non-squamous type non-small cell lung cancer (NSCLC) or not otherwise specified type NSCLC, **AND**
- Patient must have a WHO performance status of 2 or less.

Population criteria:

- Patient must have evidence of an anaplastic lymphoma kinase (ALK) gene rearrangement in tumour material, defined as 15% (or greater) positive cells by fluorescence in situ hybridisation (FISH) testing.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Non-Small-Cell Lung Cancer Authority Application - Supporting Information Form.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos. Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Stage IIIB (locally advanced) or Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Continuing treatment

Clinical criteria:

- The treatment must be as monotherapy, **AND**
- Patient must have previously been issued with an authority prescription for this drug, **AND**
- Patient must not develop disease progression while receiving PBS-subsidised treatment with this drug for this condition.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

crizotinib 200 mg capsule, 60

10323H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	6933.05	41.30	Xalkori [PF]

crizotinib 250 mg capsule, 60

10322G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	6933.05	41.30	Xalkori [PF]

■ CRIZOTINIB**Note** Special Pricing Arrangements apply.**Authority required**

Stage IIIB (locally advanced) or Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be as monotherapy, **AND**
- The condition must be non-squamous type non-small cell lung cancer (NSCLC) or not otherwise specified type NSCLC, **AND**

- Patient must have a WHO performance status of 2 or less, **AND**
- Patient must have evidence of c-ROS proto-oncogene 1 (ROS1) gene rearrangement in tumour material, defined as 15% (or greater) positive cells by fluorescence in situ hybridisation (FISH) testing, **AND**
- Patient must not have received prior treatment with a c-ROS proto-oncogene 1 (ROS1) receptor tyrosine kinase inhibitor for this condition; OR
- Patient must have developed intolerance to a c-ROS proto-oncogene 1 (ROS1) receptor tyrosine kinase inhibitor necessitating permanent treatment withdrawal.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Non-Small-Cell Lung Cancer Authority Application - Supporting Information Form.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Stage IIIB (locally advanced) or Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Continuing treatment

Clinical criteria:

- The treatment must be as monotherapy, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while being treated with this drug for this condition.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

crizotinib 200 mg capsule, 60

11589Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	6933.05	41.30	Xalkori [PF]

crizotinib 250 mg capsule, 60

11594F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	6933.05	41.30	Xalkori [PF]

▪ LORLATINIB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Authority required

Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be as monotherapy, **AND**
- The condition must be non-squamous type non-small cell lung cancer (NSCLC) or not otherwise specified type NSCLC, **AND**
- Patient must have a WHO performance status of 2 or less, **AND**
- The condition must have progressed following treatment with an anaplastic lymphoma kinase (ALK) inhibitor other than crizotinib.

Population criteria:

- Patient must have evidence of an anaplastic lymphoma kinase (ALK) gene rearrangement.

Authority required

Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Continuing treatment

Clinical criteria:

- The treatment must be as monotherapy, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not develop disease progression while receiving PBS-subsidised treatment with this drug for this condition.

Authority required

Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Grandfather treatment

Clinical criteria:

- Patient must have received non-PBS-subsidised treatment with this drug for this PBS indication prior to 1 August 2020, **AND**
- The treatment must be as monotherapy, **AND**
- The condition must be non-squamous type non-small cell lung cancer (NSCLC) or not otherwise specified type NSCLC, **AND**
- Patient must have had a WHO performance status of 2 or less prior to initiating non-PBS-subsidised treatment, **AND**
- The condition must have progressed following treatment with an ALK inhibitor other than crizotinib prior to commencement of non-PBS-subsidised treatment with this drug for this PBS indication, **AND**
- Patient must not have progressive disease while receiving treatment with this drug for this condition.

Population criteria:

- Patient must have evidence of an anaplastic lymphoma kinase (ALK) gene rearrangement.
- A patient may qualify for PBS-subsidised treatment under this restriction once only.

For continuing PBS-subsidised treatment, a Grandfathered patient must qualify under the Continuing treatment criteria.

lorlatinib 25 mg tablet, 90

12096P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	7289.46	41.30	Lorviqua [PF]

lorlatinib 100 mg tablet, 30

12091J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	7289.46	41.30	Lorviqua [PF]

Mitogen-activated protein kinase (MEK) inhibitors

▪ **BINIMETINIB**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

10328

Unresectable Stage III or Stage IV malignant melanoma

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must be receiving PBS-subsidised encorafenib concomitantly for this condition.

binimetinib 15 mg tablet, 84

11948W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	3	..	*7395.38	41.30	Mektovi [FB]

▪ **BINIMETINIB**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note A patient who has progressive disease when treated with this drug is no longer eligible for PBS-subsidised treatment with this drug.

Authority required (STREAMLINED)

10306

Unresectable Stage III or Stage IV malignant melanoma

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously been issued with an authority prescription for this drug, **AND**
- Patient must be receiving PBS-subsidised encorafenib concomitantly for this condition, **AND**
- Patient must have stable or responding disease.

binimetinib 15 mg tablet, 84

11961M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*7395.38	41.30	Mektovi [FB]

▪ **COBIMETINIB**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

10033

Unresectable Stage III or Stage IV malignant melanoma

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must be receiving PBS subsidised vemurafenib concomitantly for this condition.

cobimetinib 20 mg tablet, 63

11074W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	7395.38	41.30	Cotellic [RO]

■ COBIMETINIB

Note A patient who has had progressive disease when treated with this drug is no longer eligible for PBS-subsidised treatment with this drug.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)**6803**

Unresectable Stage III or Stage IV malignant melanoma

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously been issued with an authority prescription for this drug, **AND**
- Patient must be receiving PBS-subsidised vemurafenib concomitantly for this condition, **AND**
- Patient must have stable or responding disease.

cobimetinib 20 mg tablet, 63

11075X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	7395.38	41.30	Cotellic [RO]

■ TRAMETINIB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)**10051**

Unresectable Stage III or Stage IV malignant melanoma

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must be receiving PBS-subsidised dabrafenib concomitantly for this condition.

trametinib 2 mg tablet, 30

10382K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	7524.56	41.30	Mekinist [NV]

trametinib 500 microgram tablet, 30

10403M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	3	..	*5683.71	41.30	Mekinist [NV]

■ TRAMETINIB

Note A patient who has progressive disease when treated with this drug is no longer eligible for PBS-subsidised treatment with this drug.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)**6752**

Unresectable Stage III or Stage IV malignant melanoma

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously been issued with an authority prescription for this drug, **AND**
- Patient must be receiving PBS-subsidised dabrafenib concomitantly for this condition, **AND**
- Patient must have stable or responding disease.

trametinib 2 mg tablet, 30

10405P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	7524.56	41.30	Mekinist [NV]

trametinib 500 microgram tablet, 30

10385N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	5	..	*5683.71	41.30	Mekinist [NV]

■ **TRAMETINIB**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Resected Stage IIIB, Stage IIIC or Stage IIID malignant melanoma

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be adjuvant to complete surgical resection, **AND**
- The condition must be positive for a BRAF V600 mutation, **AND**
- Patient must have a WHO performance status of 1 or less, **AND**
- Patient must be receiving PBS-subsidised trametinib and dabrafenib concomitantly for this condition, **AND**
- Patient must not have received prior PBS-subsidised treatment for this condition, **AND**
- The treatment must commence within 12 weeks of complete resection, **AND**
- Patient must not receive more than 12 months of combined PBS-subsidised and non-PBS-subsidised adjuvant therapy.

Authority required

Resected Stage IIIB, Stage IIIC or Stage IIID malignant melanoma

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously been issued with an authority prescription for trametinib and dabrafenib concomitantly for adjuvant treatment following complete surgical resection, **AND**
- Patient must not have experienced disease recurrence, **AND**
- Patient must not receive more than 12 months of combined PBS-subsidised and non-PBS-subsidised adjuvant therapy.

Authority required

Resected Stage IIIB, Stage IIIC or Stage IIID malignant melanoma

Treatment Phase: Grandfather treatment

Clinical criteria:

- Patient must have previously received non-PBS subsidised drug for adjuvant treatment following complete surgical resection prior to 1 November 2019, **AND**
- The condition must be positive for a BRAF V600 mutation, **AND**
- Patient must have a WHO performance status of 1 or less prior to starting non-PBS treatment with this drug, **AND**
- Patient must not have evidence of recurrence, **AND**
- Patient must be receiving PBS-subsidised trametinib and dabrafenib concomitantly for this condition, **AND**
- Patient must not have received prior PBS-subsidised treatment for this condition, **AND**
- Patient must have commenced non-PBS-subsidised treatment within 12 weeks of complete surgical resection, **AND**
- Patient must not receive more than 12 months of combined PBS-subsidised and non-PBS-subsidised adjuvant therapy.

trametinib 2 mg tablet, 30

11819C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	7524.56	41.30	Mekinist [NV]

trametinib 500 microgram tablet, 30

11821E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	3	..	*5683.71	41.30	Mekinist [NV]

Cyclin-dependent kinase (CDK) inhibitors■ **ABEMACICLIB**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Locally advanced or metastatic breast cancer

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must not have previously been treated with an aromatase inhibitor for advanced or metastatic breast cancer, **AND**
- Patient must not have previously been treated with palbociclib or ribociclib; OR
- Patient must have developed an intolerance to palbociclib or ribociclib of a severity necessitating permanent treatment withdrawal, **AND**
- The condition must be hormone receptor positive, **AND**
- The condition must be human epidermal growth factor receptor 2 (HER2) negative, **AND**
- The condition must be inoperable, **AND**
- Patient must have a World Health Organisation (WHO) Eastern Cooperative Oncology Group (ECOG) performance status score of 2 or less, **AND**
- The treatment must be in combination with anastrozole or letrozole, **AND**

- The treatment must not be in combination with palbociclib or ribociclib.

Population criteria:

- Patient must not be premenopausal.

Authority required

Locally advanced or metastatic breast cancer

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not develop disease progression while receiving treatment with this drug for this condition, **AND**
- Patient must have stable or responding disease, **AND**
- The treatment must be in combination with anastrozole or letrozole, **AND**
- The treatment must not be in combination with palbociclib or ribociclib.

Population criteria:

- Patient must not be premenopausal.

A patient who has progressive disease when treated with this drug is no longer eligible for PBS-subsidised treatment with this drug.

abemaciclib 150 mg tablet, 56

11868P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	4249.01	41.30	Verzenio [LY]

abemaciclib 100 mg tablet, 56

11871T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	4249.01	41.30	Verzenio [LY]

abemaciclib 50 mg tablet, 56

11876C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	4249.01	41.30	Verzenio [LY]

■ PALBOCICLIB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Authority required

Locally advanced or metastatic breast cancer

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must not have previously been treated with an aromatase inhibitor for advanced or metastatic breast cancer, **AND**
- Patient must not have previously been treated with abemaciclib or ribociclib; OR
- Patient must have developed an intolerance to abemaciclib or ribociclib of a severity necessitating permanent treatment withdrawal, **AND**
- The condition must be hormone receptor positive, **AND**
- The condition must be human epidermal growth factor receptor 2 (HER2) negative, **AND**
- The condition must be inoperable, **AND**
- Patient must have a World Health Organisation (WHO) Eastern Cooperative Oncology Group (ECOG) performance status score of 2 or less, **AND**
- The treatment must be in combination with anastrozole or letrozole, **AND**
- The treatment must not be in combination with abemaciclib or ribociclib.

Population criteria:

- Patient must not be premenopausal.

Authority required

Locally advanced or metastatic breast cancer

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not develop disease progression while receiving treatment with this drug for this condition, **AND**
- Patient must have stable or responding disease according to the Response Evaluation Criteria In Solid Tumours (RECIST), **AND**
- The treatment must be in combination with anastrozole or letrozole, **AND**
- The treatment must not be in combination with abemaciclib or ribociclib.

Population criteria:

- Patient must not be premenopausal.

A patient who has progressive disease when treated with this drug is no longer eligible for PBS-subsidised treatment with this drug.

palbociclib 125 mg capsule, 21

11698Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	4249.01	41.30	Ibrance [PF]

palbociclib 100 mg capsule, 21

11700T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	4249.01	41.30	Ibrance [PF]

palbociclib 75 mg capsule, 21

11699R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	4249.01	41.30	Ibrance [PF]

■ RIBOCICLIB

Caution QT interval monitoring is required for patients treated with this drug.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Note Special Pricing Arrangements apply.

Authority required

Locally advanced or metastatic breast cancer

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must be untreated with each of: (i) abemaciclib, (ii) palbociclib, (iii) ribociclib; OR
- Patient must have developed an intolerance to abemaciclib or palbociclib of a severity necessitating permanent treatment withdrawal, **AND**
- The condition must be hormone receptor positive, **AND**
- The condition must be human epidermal growth factor receptor 2 (HER2) negative, **AND**
- The condition must be inoperable, **AND**
- Patient must have a World Health Organisation (WHO) Eastern Cooperative Oncology Group (ECOG) performance status score of 2 or less, **AND**
- The treatment must be in combination, where the patient has never been treated with endocrine therapy for advanced/metastatic disease, with one of: (i) anastrozole, (ii) letrozole, (iii) fulvestrant; OR
- The treatment must be in combination, where the patient has recurrence/progressive disease despite being treated with endocrine therapy for advanced/metastatic disease, with fulvestrant only, **AND**
- The treatment must not be in combination with abemaciclib or palbociclib, **AND**
- Patient must require dosage reduction requiring a pack of 21 tablets.

Population criteria:

- Patient must not be premenopausal.

Authority required

Locally advanced or metastatic breast cancer

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while being treated with this drug for this condition, **AND**
- The treatment must be in combination with one of: (i) anastrozole, (ii) letrozole, (iii) fulvestrant, **AND**
- The treatment must not be in combination with abemaciclib or palbociclib, **AND**
- Patient must require dosage reduction requiring a pack of 21 tablets.

Population criteria:

- Patient must not be premenopausal.

Authority required

Locally advanced or metastatic breast cancer

Treatment Phase: Transitioning from non-PBS to PBS-subsidised supply - 'Grandfather' treatment

Clinical criteria:

- Patient must have received non-PBS-subsidised treatment with this drug for this PBS indication prior to 1 April 2021, **AND**
- Patient must not have developed disease progression while being treated with this drug for this condition, **AND**
- Patient must have been untreated with each of: (i) abemaciclib, (ii) palbociclib, (iii) ribociclib, at the time non-PBS supply was initiated; OR
- Patient must have developed an intolerance to abemaciclib or palbociclib of a severity necessitating permanent treatment withdrawal, **AND**
- The condition must be hormone receptor positive, **AND**
- The condition must be human epidermal growth factor receptor 2 (HER2) negative, **AND**

- The condition must be inoperable, **AND**
- Patient must have had a World Health Organisation (WHO) Eastern Cooperative Oncology Group (ECOG) performance status score no higher than 2 at the time of non-PBS supply was initiated, **AND**
- The treatment must be in combination with one of: (i) anastrozole, (ii) letrozole, (iii) fulvestrant, where the patient had never been treated with endocrine therapy for advanced/metastatic disease at the time non-PBS supply was initiated; OR
- The treatment must be in combination with fulvestrant only, where at the time non-PBS supply was initiated, the patient had recurrent/progressive disease despite being treated with endocrine therapy for advanced/metastatic disease, **AND**
- The treatment must not be in combination with abemaciclib or palbociclib, **AND**
- Patient must require dosage reduction requiring a pack of 21 tablets.

Population criteria:

- Patient must not be premenopausal.

Note This grandfather restriction will cease to operate from 12 months after the date specified in the clinical criteria.

Note Patients may qualify for PBS-subsidised treatment under this restriction once only. For continuing PBS-subsidised treatment, a 'Grandfathered' patient must qualify under the 'Continuing treatment' criteria.

ribociclib 200 mg tablet, 21

11385F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1939.87	41.30	Kisqali [NV]

▪ **RIBOCICLIB**

Caution QT interval monitoring is required for patients treated with this drug.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Note Special Pricing Arrangements apply.

Authority required

Locally advanced or metastatic breast cancer

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must be untreated with each of: (i) abemaciclib, (ii) palbociclib, (iii) ribociclib; OR
- Patient must have developed an intolerance to abemaciclib or palbociclib of a severity necessitating permanent treatment withdrawal, **AND**
- The condition must be hormone receptor positive, **AND**
- The condition must be human epidermal growth factor receptor 2 (HER2) negative, **AND**
- The condition must be inoperable, **AND**
- Patient must have a World Health Organisation (WHO) Eastern Cooperative Oncology Group (ECOG) performance status score of 2 or less, **AND**
- The treatment must be in combination, where the patient has never been treated with endocrine therapy for advanced/metastatic disease, with one of: (i) anastrozole, (ii) letrozole, (iii) fulvestrant; OR
- The treatment must be in combination, where the patient has recurrence/progressive disease despite being treated with endocrine therapy for advanced/metastatic disease, with fulvestrant only, **AND**
- The treatment must not be in combination with abemaciclib or palbociclib.

Population criteria:

- Patient must not be premenopausal.

Authority required

Locally advanced or metastatic breast cancer

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while being treated with this drug for this condition, **AND**
- The treatment must be in combination with one of: (i) anastrozole, (ii) letrozole, (iii) fulvestrant, **AND**
- The treatment must not be in combination with abemaciclib or palbociclib.

Population criteria:

- Patient must not be premenopausal.

Authority required

Locally advanced or metastatic breast cancer

Treatment Phase: Transitioning from non-PBS to PBS-subsidised supply - 'Grandfather' treatment

Clinical criteria:

- Patient must have received non-PBS-subsidised treatment with this drug for this PBS indication prior to 1 April 2021, **AND**
- Patient must not have developed disease progression while being treated with this drug for this condition, **AND**
- Patient must have been untreated with each of: (i) abemaciclib, (ii) palbociclib, (iii) ribociclib, at the time non-PBS supply was initiated; OR
- Patient must have developed an intolerance to abemaciclib or palbociclib of a severity necessitating permanent treatment withdrawal, **AND**

- The condition must be hormone receptor positive, **AND**
- The condition must be human epidermal growth factor receptor 2 (HER2) negative, **AND**
- The condition must be inoperable, **AND**
- Patient must have had a World Health Organisation (WHO) Eastern Cooperative Oncology Group (ECOG) performance status score no higher than 2 at the time of non-PBS supply was initiated, **AND**
- The treatment must be in combination with one of: (i) anastrozole, (ii) letrozole, (iii) fulvestrant, where the patient had never been treated with endocrine therapy for advanced/metastatic disease at the time non-PBS supply was initiated; OR
- The treatment must be in combination with fulvestrant only, where at the time non-PBS supply was initiated, the patient had recurrent/progressive disease despite being treated with endocrine therapy for advanced/metastatic disease, **AND**
- The treatment must not be in combination with abemaciclib or palbociclib.

Population criteria:

- Patient must not be premenopausal.

Note This grandfather restriction will cease to operate from 12 months after the date specified in the clinical criteria.

Note Patients may qualify for PBS-subsidised treatment under this restriction once only. For continuing PBS-subsidised treatment, a 'Grandfathered' patient must qualify under the 'Continuing treatment' criteria.

ribociclib 200 mg tablet, 63

11386G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	5521.17	41.30	Kisqali [NV]

■ RIBOCICLIB

Caution QT interval monitoring is required for patients treated with this drug.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Note Special Pricing Arrangements apply.

Authority required

Locally advanced or metastatic breast cancer

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must be untreated with each of: (i) abemaciclib, (ii) palbociclib, (iii) ribociclib; OR
- Patient must have developed an intolerance to abemaciclib or palbociclib of a severity necessitating permanent treatment withdrawal, **AND**
- The condition must be hormone receptor positive, **AND**
- The condition must be human epidermal growth factor receptor 2 (HER2) negative, **AND**
- The condition must be inoperable, **AND**
- Patient must have a World Health Organisation (WHO) Eastern Cooperative Oncology Group (ECOG) performance status score of 2 or less, **AND**
- The treatment must be in combination, where the patient has never been treated with endocrine therapy for advanced/metastatic disease, with one of: (i) anastrozole, (ii) letrozole, (iii) fulvestrant; OR
- The treatment must be in combination, where the patient has recurrence/progressive disease despite being treated with endocrine therapy for advanced/metastatic disease, with fulvestrant only, **AND**
- The treatment must not be in combination with abemaciclib or palbociclib, **AND**
- Patient must require dosage reduction requiring a pack of 42 tablets.

Population criteria:

- Patient must not be premenopausal.

Authority required

Locally advanced or metastatic breast cancer

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while being treated with this drug for this condition, **AND**
- The treatment must be in combination with one of: (i) anastrozole, (ii) letrozole, (iii) fulvestrant, **AND**
- The treatment must not be in combination with abemaciclib or palbociclib, **AND**
- Patient must require dosage reduction requiring a pack of 42 tablets.

Population criteria:

- Patient must not be premenopausal.

Authority required

Locally advanced or metastatic breast cancer

Treatment Phase: Transitioning from non-PBS to PBS-subsidised supply - 'Grandfather' treatment

Clinical criteria:

- Patient must have received non-PBS-subsidised treatment with this drug for this PBS indication prior to 1 April 2021, **AND**
- Patient must not have developed disease progression while being treated with this drug for this condition, **AND**

- Patient must have been untreated with each of: (i) abemaciclib, (ii) palbociclib, (iii) ribociclib, at the time non-PBS supply was initiated; OR
- Patient must have developed an intolerance to abemaciclib or palbociclib of a severity necessitating permanent treatment withdrawal, **AND**
- The condition must be hormone receptor positive, **AND**
- The condition must be human epidermal growth factor receptor 2 (HER2) negative, **AND**
- The condition must be inoperable, **AND**
- Patient must have had a World Health Organisation (WHO) Eastern Cooperative Oncology Group (ECOG) performance status score no higher than 2 at the time of non-PBS supply was initiated, **AND**
- The treatment must be in combination with one of: (i) anastrozole, (ii) letrozole, (iii) fulvestrant, where the patient had never been treated with endocrine therapy for advanced/metastatic disease at the time non-PBS supply was initiated; OR
- The treatment must be in combination with fulvestrant only, where at the time non-PBS supply was initiated, the patient had recurrent/progressive disease despite being treated with endocrine therapy for advanced/metastatic disease, **AND**
- The treatment must not be in combination with abemaciclib or palbociclib, **AND**
- Patient must require dosage reduction requiring a pack of 42 tablets.

Population criteria:

- Patient must not be premenopausal.

Note This grandfather restriction will cease to operate from 12 months after the date specified in the clinical criteria.

Note Patients may qualify for PBS-subsidised treatment under this restriction once only. For continuing PBS-subsidised treatment, a 'Grandfathered' patient must qualify under the 'Continuing treatment' criteria.

ribociclib 200 mg tablet, 42

11397W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	3734.50	41.30	Kisqali [NV]

Mammalian target of rapamycin (mTOR) kinase inhibitors**■ EVEROLIMUS****Authority required (STREAMLINED)****7431**

Tuberous sclerosis complex (TSC)

Treatment Phase: Continuing treatment

Clinical criteria:

- The condition must be subependymal giant cell astrocytomas (SEGAs) associated with TSC; OR
- The condition must be visceral tumours associated with TSC, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have received an initial authority prescription for this drug for this condition, **AND**
- Patient must have demonstrated a response to prior treatment.

everolimus 10 mg tablet, 30

11267B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1736.94	41.30	Afinitor [NV]

everolimus 5 mg tablet, 30

11254H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	900.40	41.30	Afinitor [NV]

everolimus 2.5 mg tablet, 30

11258M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	456.75	41.30	Afinitor [NV]

■ EVEROLIMUS**Authority required**

Tuberous sclerosis complex (TSC)

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be subependymal giant cell astrocytomas (SEGAs) associated with TSC; OR
- The condition must be visceral tumours associated with TSC, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must not be a candidate for curative surgical resection.

everolimus 2.5 mg tablet, 30

2818H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	456.75	41.30	Afinitor [NV]

■ EVEROLIMUS

Note No increase in the maximum number of repeats may be authorised.

Authority required

Refractory seizures associated with tuberous sclerosis complex

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have a confirmed diagnosis of tuberous sclerosis complex (TSC), **AND**
- Patient must be experiencing a minimum of two partial-onset seizures per week, **AND**
- The condition must have failed to be controlled satisfactorily at stable doses of at least two anti-epileptic drugs, **AND**
- The treatment must be in combination with at least one anti-epileptic drug, **AND**
- Patient must not be a candidate for curative surgery.

Population criteria:

- Patient must be at least 2 years of age.

everolimus 5 mg dispersible tablet, 30

11592D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	900.19	41.30	Afinitor [NV]

everolimus 3 mg dispersible tablet, 30

11599L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	546.55	41.30	Afinitor [NV]

everolimus 2 mg dispersible tablet, 30

11591C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	366.71	41.30	Afinitor [NV]

▪ **EVEROLIMUS**

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

8262

Refractory seizures associated with tuberous sclerosis complex

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have maintained a response to the PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be in combination with at least one anti-epileptic drug, **AND**
- Patient must not be a candidate for curative surgery.

everolimus 5 mg dispersible tablet, 30

11598K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	900.19	41.30	Afinitor [NV]

everolimus 3 mg dispersible tablet, 30

11608Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	546.55	41.30	Afinitor [NV]

everolimus 2 mg dispersible tablet, 30

11607X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	366.71	41.30	Afinitor [NV]

▪ **EVEROLIMUS**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Metastatic or unresectable, well-differentiated malignant pancreatic neuroendocrine tumour (pNET)

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously been issued with an authority prescription for this drug, **AND**
 - Patient must not have disease progression, **AND**
 - The treatment must be as monotherapy.
- Patients who have progressive disease with this drug are no longer eligible for PBS-subsidised treatment with this drug.

everolimus 10 mg tablet, 30

10135K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	1736.94	41.30	^a Afinitor [NV]	^a Everolimus Sandoz [SZ]

everolimus 5 mg tablet, 30

10131F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	900.40	41.30	^a Afinitor [NV]	^a Everolimus Sandoz [SZ]

■ EVEROLIMUS

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Metastatic or unresectable, well-differentiated malignant pancreatic neuroendocrine tumour (pNET)

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must be symptomatic (despite somatostatin analogues); OR
- Patient must have disease progression, **AND**

- The treatment must be as monotherapy.

Disease progression must be documented in the patient's medical records.

Patients who have developed progressive disease on sunitinib are not eligible to receive PBS-subsidised everolimus.

Patients who have developed intolerance to sunitinib of a severity necessitating permanent treatment withdrawal are eligible to receive PBS-subsidised everolimus.

everolimus 10 mg tablet, 30

11377T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	2	..	1736.94	41.30	^a Afinitor [NV]	^a Everolimus Sandoz [SZ]

everolimus 5 mg tablet, 30

11362B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	2	..	900.40	41.30	^a Afinitor [NV]	^a Everolimus Sandoz [SZ]

■ EVEROLIMUS

Authority required

Tuberous sclerosis complex (TSC)

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be subependymal giant cell astrocytomas (SEGAs) associated with TSC; OR
- The condition must be visceral tumours associated with TSC, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must not be a candidate for curative surgical resection.

Authority required

Metastatic (Stage IV) breast cancer

Clinical criteria:

- The condition must be hormone receptor positive, **AND**
- The condition must be human epidermal growth factor receptor 2 (HER2) negative, **AND**
- The condition must have acquired endocrine resistance as demonstrated by initial response and then recurrence or progression of disease after treatment with letrozole or anastrozole, **AND**
- The treatment must be in combination with exemestane.

Population criteria:

- Patient must not be pre-menopausal.

Note Patients who have progressive disease with everolimus are no longer eligible for PBS-subsidised everolimus.

everolimus 10 mg tablet, 30

2985D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1736.94	41.30	Afinitor [NV]

everolimus 5 mg tablet, 30

2819J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	900.40	41.30	Afinitor [NV]

■ EVEROLIMUS

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Response Evaluation Criteria In Solid Tumours (RECIST) is defined as follows:

Complete response (CR) is disappearance of all target lesions.

Partial response (PR) is a 30% decrease in the sum of the longest diameter of target lesions.

Progressive disease (PD) is a 20% increase in the sum of the longest diameter of target lesions.

Stable disease (SD) is small changes that do not meet above criteria.

Authority required

Stage IV clear cell variant renal cell carcinoma (RCC)

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have progressive disease according to the Response Evaluation Criteria in Solid Tumours (RECIST) following prior treatment with a tyrosine kinase inhibitor, **AND**

- Patient must have a WHO performance status of 2 or less, **AND**
 - The treatment must be the sole PBS-subsidised therapy for this condition.
- Patients who have developed intolerance to a tyrosine kinase inhibitor of a severity necessitating permanent treatment withdrawal are eligible to receive PBS-subsidised everolimus.

Patients who have progressive disease with everolimus are no longer eligible for PBS-subsidised everolimus.

everolimus 10 mg tablet, 30

10132G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	1736.94	41.30	Afinitor [NV]

everolimus 5 mg tablet, 30

10133H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	900.40	41.30	Afinitor [NV]

■ EVEROLIMUS

Note Response Evaluation Criteria In Solid Tumours (RECIST) is defined as follows:

Complete response (CR) is disappearance of all target lesions.

Partial response (PR) is a 30% decrease in the sum of the longest diameter of target lesions.

Progressive disease (PD) is a 20% increase in the sum of the longest diameter of target lesions.

Stable disease (SD) is small changes that do not meet above criteria.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)**7432**

Stage IV clear cell variant renal cell carcinoma (RCC)

Treatment Phase: Continuing treatment beyond 3 months

Clinical criteria:

- Patient must have received an initial authority prescription for this drug for this condition, **AND**
- Patient must have stable or responding disease according to the Response Evaluation Criteria In Solid Tumours (RECIST), **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

A patient who has progressive disease when treated with this drug is no longer eligible for PBS-subsidised treatment with this drug.

everolimus 10 mg tablet, 30

11262R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1736.94	41.30	Afinitor [NV]

everolimus 5 mg tablet, 30

11257L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	900.40	41.30	Afinitor [NV]

Human epidermal growth factor receptor 2 (HER2) tyrosine kinase inhibitors**■ LAPATINIB**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)**9360**

Metastatic (Stage IV) HER2 positive breast cancer

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have received an initial authority prescription for this drug for this condition, **AND**
- The treatment must be in combination with capecitabine, **AND**
- Patient must not develop disease progression while receiving PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be the sole PBS-subsidised anti-HER2 therapy for this condition, **AND**
- The treatment must not be used in a patient with a left ventricular ejection fraction (LVEF) of less than 45% and/or with symptomatic heart failure.

A patient who has progressive disease when treated with this drug is no longer eligible for PBS-subsidised treatment with this drug.

The treatment must not exceed a lifetime total of one continuous course.

lapatinib 250 mg tablet, 70

11251E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	2	..	*2932.30	41.30	Tykerb [NV]

■ LAPATINIB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Metastatic (Stage IV) HER2 positive breast cancer

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have evidence of human epidermal growth factor receptor 2 (HER2) gene amplification as demonstrated by in situ hybridisation (ISH) either in the primary tumour or a metastatic lesion, **AND**
- The treatment must be in combination with capecitabine, **AND**
- Patient must have received prior therapy with a taxane for at least 3 cycles; and experienced disease progression during or within 6 months of completing treatment with pertuzumab and trastuzumab in combination; OR
- Patient must have developed intolerance to treatment with a taxane of a severity necessitating permanent treatment withdrawal; and experienced disease progression during or within 6 months of completing treatment with pertuzumab and trastuzumab in combination; OR
- Patient must have experienced disease progression following treatment with trastuzumab emtansine in whom disease had relapsed during or within 6 months of completing prior adjuvant therapy with trastuzumab; OR
- Patient must have experienced disease relapsed during or within 6 months of completing prior adjuvant therapy with trastuzumab, **AND**
- The treatment must be the sole PBS-subsidised anti-HER2 therapy for this condition, **AND**
- The treatment must not be used in a patient with a left ventricular ejection fraction (LVEF) of less than 45% and/or with symptomatic heart failure.

Authority applications for initial treatment must be made in writing and must include:

- a completed authority prescription form; and
- a completed Late stage metastatic breast cancer Initial PBS authority application form which includes:
 - details of the pathology report from an Approved Pathology Authority confirming evidence of HER2 gene amplification in the primary tumour or a metastatic lesion by in situ hybridisation (ISH); and
 - date of last treatment with a taxane and total number of cycles; or
 - dates of treatment with trastuzumab and pertuzumab; or
 - date of demonstration of progression during or within 6 months of completing treatment with trastuzumab and pertuzumab; or
 - date of demonstration of progression during or within 6 months of completing treatment with trastuzumab.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, please provide details of the degree of this toxicity at the time of application.

Cardiac function must be tested by echocardiography (ECHO) or multigated acquisition (MUGA), prior to seeking the initial authority approval.

lapatinib 250 mg tablet, 70

9148L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	2	..	*2932.30	41.30	Tykerb [NV]

Janus-associated kinase (JAK) inhibitors

▪ **RUXOLITINIB**

Note Risk of myelofibrosis is defined in accordance with the Myelofibrosis International Prognostic Scoring System (IPSS) OR the Dynamic International Prognostic Scoring System (DIPSS) OR the Age-Adjusted DIPSS.

Note Authority applications for continuing treatment may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note No increase in the maximum quantity may be authorised for the 15 mg and 20 mg dose strengths.

Note Special Pricing Arrangements apply.

Authority required

High risk and intermediate-2 risk myelofibrosis

Treatment Phase: Continuing treatment

Clinical criteria:

- The condition must be primary myelofibrosis or post-polycythemia vera myelofibrosis or post-essential thrombocythemia myelofibrosis, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition.

Authority required

Intermediate-1 risk myelofibrosis

Treatment Phase: Continuing treatment

Clinical criteria:

- The condition must be primary myelofibrosis or post-polycythemia vera myelofibrosis or post-essential thrombocythemia myelofibrosis, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition.

ruxolitinib 10 mg tablet, 56

10927D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	4911.16	41.30	Jakavi [NV]

ruxolitinib 15 mg tablet, 56

10615Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	4911.16	41.30	Jakavi [NV]

ruxolitinib 5 mg tablet, 56

10616R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*4911.16	41.30	Jakavi [NV]

ruxolitinib 20 mg tablet, 56

10617T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	4911.16	41.30	Jakavi [NV]

■ RUXOLITINIB

Note Risk of myelofibrosis is defined in accordance with the Myelofibrosis International Prognostic Scoring System (IPSS) OR the Dynamic International Prognostic Scoring System (DIPSS) OR the Age-Adjusted DIPSS.

Note No increase in the maximum quantity may be authorised for the 15 mg and 20 mg dose strengths.

Note Special Pricing Arrangements apply.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

High risk and intermediate-2 risk myelofibrosis

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be primary myelofibrosis or post-polycythemia vera myelofibrosis or post-essential thrombocythemia myelofibrosis.

The authority application must be made in writing and must include:

- (1) A completed authority prescription form; and
- (2) A completed Myelofibrosis Authority Application Supporting Information Form, which includes all of the following:
 - (a) A copy of the bone marrow biopsy report confirming diagnosis of myelofibrosis; and
 - (b) A classification of risk of myelofibrosis according to either the IPSS, DIPSS, or the Age-Adjusted DIPSS.

Authority required

Intermediate-1 risk myelofibrosis

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be primary myelofibrosis or post-polycythemia vera myelofibrosis or post-essential thrombocythemia myelofibrosis, **AND**

- Patient must have severe disease-related symptoms that are resistant, refractory or intolerant to available therapy.

The authority application must be made in writing and must include:

- (1) A completed authority prescription form; and
- (2) A completed Myelofibrosis Authority Application Supporting Information Form, which includes all of the following:
 - a) A copy of the bone marrow biopsy report confirming diagnosis of myelofibrosis;
 - b) A classification of risk of myelofibrosis according to either the IPSS, DIPSS, or the Age-Adjusted DIPSS; and
 - c) A confirmation that the patient's disease related symptoms are resistant, refractory or intolerant to available therapy.

ruxolitinib 10 mg tablet, 56

10913J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	4911.16	41.30	Jakavi [NV]

ruxolitinib 15 mg tablet, 56

10619X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	4911.16	41.30	Jakavi [NV]

ruxolitinib 5 mg tablet, 56

10614P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*4911.16	41.30	Jakavi [NV]

ruxolitinib 20 mg tablet, 56

10618W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	4911.16	41.30	Jakavi [NV]

Vascular endothelial growth factor receptor (VEGFR) tyrosine kinase inhibitors

▪ **AXITINIB**

Note Response Evaluation Criteria In Solid Tumours (RECIST) is defined as follows:

Complete response (CR) is disappearance of all target lesions.

Partial response (PR) is a 30% decrease in the sum of the longest diameter of target lesions.

Progressive disease (PD) is a 20% increase in the sum of the longest diameter of target lesions.

Stable disease (SD) is small changes that do not meet above criteria.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

7433

Stage IV clear cell variant renal cell carcinoma (RCC)

Treatment Phase: Continuing treatment beyond 3 months

Clinical criteria:

- Patient must have received an initial authority prescription for this drug for this condition, **AND**
- Patient must have stable or responding disease according to the Response Evaluation Criteria In Solid Tumours (RECIST), **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Prescribers may request an increased maximum quantity sufficient to provide up to one month's supply for patients who require dose adjustment.

A patient who has progressive disease when treated with this drug is no longer eligible for PBS-subsidised treatment with this drug.

axitinib 1 mg tablet, 28

10539Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*1069.34	41.30	Inlyta [PF]

axitinib 5 mg tablet, 28

10556N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*4949.16	41.30	Inlyta [PF]

▪ **AXITINIB**

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Response Evaluation Criteria In Solid Tumours (RECIST) is defined as follows:

Complete response (CR) is disappearance of all target lesions.

Partial response (PR) is a 30% decrease in the sum of the longest diameter of target lesions.

Progressive disease (PD) is a 20% increase in the sum of the longest diameter of target lesions.

Stable disease (SD) is small changes that do not meet above criteria.

Authority required

Stage IV clear cell variant renal cell carcinoma (RCC)

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have progressive disease according to the Response Evaluation Criteria in Solid Tumours (RECIST) following prior treatment with a tyrosine kinase inhibitor, **AND**
- Patient must have a WHO performance status of 2 or less, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Patients who have developed intolerance to a tyrosine kinase inhibitor of a severity necessitating permanent treatment withdrawal are eligible to receive PBS-subsidised treatment with this drug.

A patient who has progressive disease when treated with this drug is no longer eligible for PBS-subsidised treatment with this drug.

Prescribers may request an increased maximum quantity sufficient to provide up to one month's supply for patients who require dose adjustment.

axitinib 1 mg tablet, 28

10572K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	2	..	*1069.34	41.30	Inlyta [PF]

axitinib 5 mg tablet, 28

10540R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	2	..	*4949.16	41.30	Inlyta [PF]

Bruton's tyrosine kinase (BTK) inhibitors**■ ACALABRUTINIB**

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Authority required

Relapsed or refractory chronic lymphocytic leukaemia (CLL) or small lymphocytic lymphoma (SLL)

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- The condition must have relapsed or be refractory to at least one prior therapy, **AND**
- Patient must have a WHO performance status of 1 or less, **AND**
- Patient must not have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must be considered unsuitable for treatment or retreatment with a purine analogue, **AND**
- Patient must not have received treatment with another Bruton's tyrosine kinase (BTK) inhibitor for any line of treatment of CLL/SLL (untreated or relapsed/refractory disease); OR
- Patient must have developed intolerance to another Bruton's tyrosine kinase (BTK) inhibitor of a severity necessitating permanent treatment withdrawal when being treated for relapsed or refractory CLL/SLL.

A patient is considered unsuitable for treatment or retreatment with a purine analogue as demonstrated by at least one of the following:

- Failure to respond (stable disease or disease progression on treatment), or a progression-free interval of less than 3 years from treatment with a purine analogue-based therapy and anti-CD20-containing chemoimmunotherapy regimen after at least two cycles;
- Age is 70 years or older;
- Age is 65 years or older and the presence of comorbidities (Cumulative Illness Rating Scale of 6 or greater, or creatinine clearance of less than 70 mL/min) that might place the patient at an unacceptable risk for treatment-related toxicity with purine analogue-based therapy, provided they have received one or more prior treatment including at least two cycles of an alkylating agent-based (or purine analogue-based) anti-CD20 antibody-containing chemoimmunotherapy regimen;
- History of purine analogue-associated autoimmune anaemia or autoimmune thrombocytopenia;
- Evidence of one or more 17p chromosomal deletions demonstrated by a Medical Benefits Schedule listed test.

Authority required

Relapsed or refractory chronic lymphocytic leukaemia (CLL) or small lymphocytic lymphoma (SLL)

Treatment Phase: Continuing treatment of relapsed or refractory CLL/SLL

Clinical criteria:

- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not develop disease progression while receiving PBS-subsidised treatment with this drug for this condition.

Authority required

Relapsed or refractory chronic lymphocytic leukaemia (CLL) or small lymphocytic lymphoma (SLL)

Treatment Phase: Grandfather treatment (initial treatment in a patient commenced on non-PBS-subsidised treatment)

Clinical criteria:

- Patient must have previously received non-PBS-subsidised treatment with this drug for relapsed or refractory CLL/SLL prior to 1 September 2020, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- The condition must have relapsed or be refractory to at least one prior therapy prior to initiating non-PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have a WHO performance status of 1 or less prior to starting non-PBS treatment with this drug, **AND**
- Patient must have been considered unsuitable for treatment or retreatment with a purine analogue prior to initiating non-PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must be considered unsuitable for treatment or retreatment with a purine analogue, **AND**
- Patient must not have received treatment with another Bruton's tyrosine kinase (BTK) inhibitor for any line of treatment of CLL/SLL (untreated or relapsed/refractory disease) prior to initiating non-PBS-subsidised treatment with this drug for this condition; OR
- Patient must have developed intolerance to another Bruton's tyrosine kinase (BTK) inhibitor of a severity necessitating permanent treatment withdrawal when being treated for relapsed or refractory CLL/SLL prior to initiating non-PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while receiving treatment with this drug for this condition.

A patient is considered unsuitable for treatment or retreatment with a purine analogue as demonstrated by at least one of the following being met prior to commencing non-PBS-subsidised treatment with this drug for this condition:

- Failure to respond (stable disease or disease progression on treatment), or a progression-free interval of less than 3 years from treatment with a purine analogue-based therapy and anti-CD20-containing chemoimmunotherapy regimen after at least two cycles;
- Age is 70 years or older;
- Age is 65 years or older and the presence of comorbidities (Cumulative Illness Rating Scale of 6 or greater, or creatinine clearance of less than 70 mL/min) that might place the patient at an unacceptable risk for treatment-related toxicity with purine analogue-based therapy, provided they have received one or more prior treatment including at least two cycles of an alkylating agent-based (or purine analogue-based) anti-CD20 antibody-containing chemoimmunotherapy regimen;
- History of purine analogue-associated autoimmune anaemia or autoimmune thrombocytopenia;
- Evidence of one or more 17p chromosomal deletions demonstrated by a Medical Benefits Schedule listed test.

Note Patients may qualify for PBS-subsidised treatment under this restriction once only. For continuing PBS-subsidised treatment, a 'Grandfathered' patient must qualify under the 'Continuing treatment' criteria.

Note This grandfather restriction will cease to operate from 12 months after the date specified in the clinical criteria.

acalabrutinib 100 mg capsule, 56

12117R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	8218.90	41.30	Calquence [AP]

■ IBRUTINIB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Chronic lymphocytic leukaemia (CLL) or small lymphocytic lymphoma (SLL)

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- The condition must have relapsed or be refractory to at least one prior therapy, **AND**
- Patient must have a WHO performance status of 0 or 1, **AND**
- Patient must not have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have received treatment with another Bruton's tyrosine kinase (BTK) inhibitor for any line of treatment of CLL/SLL (untreated or relapsed/refractory disease); OR
- Patient must have developed intolerance to another Bruton's tyrosine kinase (BTK) inhibitor of a severity necessitating permanent treatment withdrawal when being treated for relapsed or refractory CLL/SLL, **AND**
- Patient must be considered unsuitable for treatment or retreatment with a purine analogue.

A patient is considered unsuitable for treatment or retreatment with a purine analogue as demonstrated by at least one of the following:

- Failure to respond (stable disease or disease progression on treatment), or a progression-free interval of less than 3 years from treatment with a purine analogue-based therapy and anti-CD20-containing chemoimmunotherapy regimen after at least two cycles;
- Age is 70 years or older;
- Age is 65 years or older and the presence of comorbidities (Cumulative Illness Rating Scale of 6 or greater, or creatinine clearance of less than 70 mL/min) that might place the patient at an unacceptable risk for treatment-related toxicity with purine analogue-based therapy, provided they have received one or more prior treatment including at least two cycles of an alkylating agent-based (or purine analogue-based) anti-CD20 antibody-containing chemoimmunotherapy regimen;
- History of purine analogue-associated autoimmune anaemia or autoimmune thrombocytopenia;
- Evidence of one or more 17p chromosomal deletions demonstrated by fluorescence in situ hybridisation (FISH).

Authority required

Chronic lymphocytic leukaemia (CLL) or small lymphocytic lymphoma (SLL)

Treatment Phase: Continuing treatment

Clinical criteria:

- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not develop disease progression while receiving PBS-subsidised treatment with this drug for this condition.

ibrutinib 140 mg capsule, 90

11213E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	8794.45	41.30	Imbruvica [JC]

■ IBRUTINIB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Authority required

Mantle cell lymphoma

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must have relapsed or be refractory to at least one prior therapy, **AND**
- Patient must have a WHO performance status of 0 or 1, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must not have previously received PBS-subsidised treatment with this drug for this condition.

Authority required

Mantle cell lymphoma

Treatment Phase: Continuing treatment

Clinical criteria:

- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not develop disease progression while receiving PBS-subsidised treatment with this drug for this condition.

ibrutinib 140 mg capsule, 120

11419B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	11672.21	41.30	Imbruvica [JC]

Phosphatidylinositol-3-kinase (Pi3K) inhibitors
■ IDELALISIB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Chronic lymphocytic leukaemia (CLL) or small lymphocytic lymphoma (SLL)

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must not have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be in combination with rituximab for up to a maximum of 8 doses, followed by monotherapy, **AND**
- The condition must have relapsed or be refractory to at least one prior therapy, **AND**
- The condition must be CD20 positive, **AND**
- Patient must have a total cumulative illness rating scale (CIRS) score of greater than 6 (excluding CLL-induced illness or organ damage), **AND**
- Patient must be inappropriate for chemo-immunotherapy.

A patient can be considered inappropriate for chemo-immunotherapy when one or more of the following are experienced:

1. Severe neutropenia defined as absolute neutrophil count of less than or equal to $1.0 \times 10^9/L$; or
2. Severe thrombocytopenia defined as platelet count of less than or equal to $50 \times 10^9/L$; or
3. Evidence of one or more 17p chromosomal deletions demonstrated by fluorescence in situ hybridisation (FISH).

Full blood count results must be no more than 1 month old at the time of application.

The authority application must be made in writing and must include:

- a) A completed authority prescription form;
- b) A completed CLL/SLL PBS Authority Application - Supporting information form; and
- c) Pathology report indicating that the patient can be considered inappropriate for chemo-immunotherapy due to one or more of the following:
 - 1) Recent severe neutropenia; or
 - 2) Recent severe thrombocytopenia; or
 - 3) Presence of 17p chromosomal deletion using fluorescence in situ hybridisation (FISH).

A Grandfathered patient who has previously received non-PBS subsidised treatment with this drug for this condition prior to 1 September 2017 must have met all the initial restriction criteria prior to initiating non-PBS subsidised treatment. A Grandfathered patient may qualify for PBS-subsidised treatment under this restriction once only. For continuing PBS-subsidised treatment, a Grandfathered patient must qualify under the Continuing treatment criteria.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Chronic lymphocytic leukaemia (CLL) or small lymphocytic lymphoma (SLL)

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not develop disease progression while receiving PBS-subsidised treatment with this drug for this condition.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

idelalisib 150 mg tablet, 60

11162L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	5378.91	41.30	Zydelig [GI]

idelalisib 100 mg tablet, 60

11170X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	5378.91	41.30	Zydelig [GI]

■ IDEALISIB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Refractory follicular B-cell non-Hodgkin's Lymphoma

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be refractory to a prior therapy with rituximab, **AND**
 - The condition must be refractory to a prior therapy with an alkylating agent, **AND**
 - The treatment must be the sole PBS subsidised treatment for this condition.
- The condition is considered refractory to a prior therapy when the patient experiences less than a partial response or progression of disease within 6 months after completion of the prior therapy.
- The condition is considered refractory to both rituximab and an alkylating agent if the agents were administered together or in successive treatment regimens.

The authority application must be made in writing and must include:

- A completed authority prescription form; and
- A completed Refractory follicular B-cell non-Hodgkin's Lymphoma PBS Authority Application - Supporting information form which must include date of completion of prior therapies with rituximab and an alkylating agent.

Note Any queries concerning the arrangements to prescribe this drug may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Written applications for authority to prescribe this drug should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Refractory follicular B-cell non-Hodgkin's Lymphoma

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be the sole PBS subsidised treatment for this condition, **AND**
- Patient must not develop disease progression while receiving PBS-subsidised treatment with this drug for this condition.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

idelalisib 150 mg tablet, 60

11165P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	5378.91	41.30	Zydelig [GI]

idelalisib 100 mg tablet, 60

11171Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	5378.91	41.30	Zydelig [GI]

Other protein kinase inhibitors**■ CABOZANTINIB**

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Response Evaluation Criteria In Solid Tumours (RECIST) is defined as follows:

Complete response (CR) is disappearance of all target lesions.

Partial response (PR) is a 30% decrease in the sum of the longest diameter of target lesions.

Progressive disease (PD) is a 20% increase in the sum of the longest diameter of target lesions.
Stable disease (SD) is small changes that do not meet above criteria.

Authority required (STREAMLINED)**8572**

Stage IV clear cell variant renal cell carcinoma (RCC)

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have progressive disease according to the Response Evaluation Criteria in Solid Tumours (RECIST) following prior treatment with a tyrosine kinase inhibitor, **AND**
- Patient must have a WHO performance status of 2 or less, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

cabozantinib 20 mg tablet, 30

11371L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	9961.16	41.30	Cabometyx [IS]

cabozantinib 60 mg tablet, 30

11360X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	9961.16	41.30	Cabometyx [IS]

cabozantinib 40 mg tablet, 30

11369J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	9961.16	41.30	Cabometyx [IS]

■ CABOZANTINIB

Note No increase in the maximum number of repeats may be authorised.

Note Response Evaluation Criteria In Solid Tumours (RECIST) is defined as follows:

Complete response (CR) is disappearance of all target lesions.

Partial response (PR) is a 30% decrease in the sum of the longest diameter of target lesions.

Progressive disease (PD) is a 20% increase in the sum of the longest diameter of target lesions.

Stable disease (SD) is small changes that do not meet above criteria.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)**7631**

Stage IV clear cell variant renal cell carcinoma (RCC)

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have stable or responding disease according to the Response Evaluation Criteria In Solid Tumours (RECIST), **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must not receive PBS-subsidised treatment with this drug if progressive disease develops while on this drug.

cabozantinib 20 mg tablet, 30

11374P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	9961.16	41.30	Cabometyx [IS]

cabozantinib 60 mg tablet, 30

11367G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	9961.16	41.30	Cabometyx [IS]

cabozantinib 40 mg tablet, 30

11368H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	9961.16	41.30	Cabometyx [IS]

■ ENTRECTINIB

Note Special Pricing Arrangements apply.

Authority required

Stage IIIB (locally advanced) or Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be as monotherapy, **AND**
- The condition must be non-squamous type non-small cell lung cancer (NSCLC) or not otherwise specified type NSCLC, **AND**
- Patient must have a WHO performance status of 2 or less, **AND**
- Patient must not have received prior treatment with a c-ROS proto-oncogene 1 (ROS1) receptor tyrosine kinase inhibitor for this condition; OR

- Patient must have developed intolerance to a c-ROS proto-oncogene 1 (ROS1) receptor tyrosine kinase inhibitor necessitating permanent treatment withdrawal, **AND**
- Patient must have evidence of c-ROS proto-oncogene 1 (ROS1) gene rearrangement in tumour material, defined as 15% (or greater) positive cells by fluorescence in situ hybridisation (FISH) testing.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Non-Small-Cell Lung Cancer Authority Application - Supporting Information Form.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Stage IIIB (locally advanced) or Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Continuing treatment

Clinical criteria:

- The treatment must be as monotherapy, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while being treated with this drug for this condition.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Stage IIIB (locally advanced) or Stage IV (metastatic) non-small cell lung cancer (NSCLC)

Treatment Phase: Grandfather treatment

Clinical criteria:

- Patient must have previously received non-PBS-subsidised treatment with this drug for this condition prior to 1 August 2020, **AND**
- The condition must be non-squamous type non-small cell lung cancer (NSCLC) or not otherwise specified type NSCLC, **AND**
- The treatment must be as monotherapy, **AND**
- Patient must have had a WHO performance status of 2 or less prior to initiating non-PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have developed disease progression while being treated with this drug for this condition, **AND**
- Patient must not have received prior treatment with a c-ROS proto-oncogene 1 (ROS1) receptor tyrosine kinase inhibitor for this condition prior to initiating non-PBS-subsidised treatment with this drug for this condition; OR
- Patient must have developed intolerance to a c-ROS proto-oncogene 1 (ROS1) receptor tyrosine kinase inhibitor necessitating permanent treatment withdrawal prior to initiating non-PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have evidence of c-ROS proto-oncogene 1 (ROS1) gene rearrangement in tumour material, defined as 15% (or greater) positive cells by fluorescence in situ hybridisation (FISH) testing prior to initiating non-PBS subsidised treatment with this drug for this condition.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Non-Small-Cell Lung Cancer Authority Application - Supporting Information Form.

A patient may qualify for PBS-subsidised treatment under this restriction once only.

For continuing PBS-subsidised treatment, a Grandfathered patient must qualify under the Continuing treatment criteria.

Note This grandfather restriction will cease to operate from 12 months after the date specified in the clinical criteria.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

entrectinib 200 mg capsule, 90

12092K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	7289.46	41.30	Rozlytrek [RO]

■ LENVATINIB

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)**11168**

Advanced (unresectable) Barcelona Clinic Liver Cancer Stage B or Stage C hepatocellular carcinoma

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must not be suitable for transarterial chemoembolisation, **AND**
- Patient must have a WHO performance status of 2 or less, **AND**
- Patient must have Child Pugh class A, **AND**
- The condition must be untreated with systemic therapy; OR
- Patient must have developed intolerance of a severity necessitating permanent treatment withdrawal, in the absence of disease progression, to any of the following: (i) a vascular endothelial growth factor (VEGF) tyrosine kinase inhibitor (TKI), (ii) atezolizumab/bevacizumab combination therapy.

Authority required (STREAMLINED)**8584**

Advanced (unresectable) Barcelona Clinic Liver Cancer Stage B or Stage C hepatocellular carcinoma

Treatment Phase: Continuing treatment

Clinical criteria:

- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not develop disease progression while receiving treatment with this drug for this condition.

lenvatinib 4 mg capsule, 30

11638M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	2	..	*9626.16	41.30	Lenvima [EI]

■ LENVATINIB

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)**6604**

Locally advanced or metastatic differentiated thyroid cancer

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be refractory to radioactive iodine, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have symptomatic progressive disease prior to treatment; OR
- Patient must have progressive disease at critical sites with a high risk of morbidity or mortality where local control cannot be achieved by other measures, **AND**
- Patient must have thyroid stimulating hormone adequately repressed, **AND**
- Patient must be one in whom surgery is inappropriate, **AND**
- Patient must not be a candidate for radiotherapy with curative intent, **AND**
- Patient must have a WHO performance status of 2 or less.

Radioactive iodine refractory is defined as:

- a lesion without iodine uptake on a radioactive iodine (RAI) scan; or
- having received a cumulative RAI dose of greater than or equal to 600 mCi; or
- progression within 12 months of a single RAI treatment; or
- progression after two RAI treatments administered within 12 months of each other.

Authority required (STREAMLINED)**6578**

Locally advanced or metastatic differentiated thyroid cancer

Treatment Phase: Continuing treatment

Clinical criteria:

- The condition must be refractory to radioactive iodine, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have stable or responding disease according to the Response Evaluation Criteria In Solid Tumours (RECIST).

Note Response Evaluation Criteria In Solid Tumours (RECIST) is defined as follows:

Complete response (CR) is disappearance of all target lesions.

Partial response (PR) is a 30% decrease in the sum of the longest diameter of target lesions.

Progressive disease (PD) is a 20% increase in the sum of the longest diameter of target lesions.

Stable disease (SD) is small changes that do not meet above criteria.

lenvatinib 10 mg capsule, 30

10965D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	2	..	*6471.16	41.30	Lenvima [EI]

lenvatinib 4 mg capsule, 30

10952K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	3280.07	41.30	Lenvima [EI]

■ NINTEDANIB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Idiopathic pulmonary fibrosis

Treatment Phase: Initial treatment 1 - new patient

Clinical criteria:

- The condition must be diagnosed through a multidisciplinary team, **AND**
- Patient must have chest high resolution computed tomography (HRCT) consistent with diagnosis of idiopathic pulmonary fibrosis within the previous 12 months, **AND**
- Patient must have a forced vital capacity (FVC) greater than or equal to 50% predicted for age, gender and height, **AND**
- Patient must have a forced expiratory volume in 1 second to forced vital capacity ratio (FEV1/FVC) greater than 0.7, **AND**
- Patient must have diffusing capacity of the lungs for carbon monoxide (DLCO) corrected for haemoglobin equal to or greater than 30%, **AND**
- Patient must not have interstitial lung disease due to other known causes including domestic and occupational environmental exposures, connective tissue disease, or drug toxicity, **AND**
- The treatment must be the sole PBS subsidised treatment for this condition.

Treatment criteria:

- Must be treated by a respiratory physician or specialist physician, or in consultation with a respiratory physician or specialist physician.

A multidisciplinary team is defined as comprising of at least a specialist respiratory physician, a radiologist and where histological material is considered, a pathologist. If attendance is not possible because of geographical isolation, consultation with a multidisciplinary team is required for diagnosis.

Patient must not have an acute respiratory infection at the time of FVC testing.

Application for authorisation of initial treatment must be in writing and must include:

- a) a completed authority prescription form; and
- b) a completed IPF Authority Application Supporting Information Form; and
- c) a signed patient acknowledgement.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Idiopathic pulmonary fibrosis

Treatment Phase: Initial treatment 2 - change or re-commencement of treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with nintedanib or pirfenidone for this condition, **AND**
- The treatment must be the sole PBS subsidised treatment for this condition.

Treatment criteria:

- Must be treated by a respiratory physician or specialist physician, or in consultation with a respiratory physician or specialist physician.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Idiopathic pulmonary fibrosis

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be the sole PBS subsidised treatment for this condition.

Treatment criteria:

- Must be treated by a respiratory physician or specialist physician, or in consultation with a respiratory physician or specialist physician.

Note Authority applications for continuing treatment may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

nintedanib 100 mg capsule, 60

11100F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1763.61	41.30	Ofev [BY]

nintedanib 150 mg capsule, 60

11106M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	3398.76	41.30	Ofev [BY]

▪ **PAZOPANIB**

Note Response Evaluation Criteria In Solid Tumours (RECIST) is defined as follows:

Complete response (CR) is disappearance of all target lesions.

Partial response (PR) is a 30% decrease in the sum of the longest diameter of target lesions.

Progressive disease (PD) is a 20% increase in the sum of the longest diameter of target lesions.

Stable disease (SD) is small changes that do not meet above criteria.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

7422

Stage IV clear cell variant renal cell carcinoma (RCC)

Treatment Phase: Continuing treatment beyond 3 months

Clinical criteria:

- Patient must have received an initial authority prescription for this drug for this condition, **AND**
- Patient must have stable or responding disease according to the Response Evaluation Criteria In Solid Tumours (RECIST), **AND**
- Patient must require dose adjustment, **AND**
- The treatment must be the sole PBS-subsidised tyrosine kinase inhibitor therapy for this condition.

A patient who has progressive disease when treated with this drug is no longer eligible for PBS-subsidised treatment with this drug.

Patients who have progressive disease on sunitinib are not eligible to receive PBS-subsidised pazopanib.

pazopanib 400 mg tablet, 30

2201W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	2311.62	41.30	Votrient [NV]

pazopanib 200 mg tablet, 30

2232L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1192.86	41.30	Votrient [NV]

▪ **PAZOPANIB**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

9247

Advanced (unresectable and/or metastatic) soft tissue sarcoma

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have a WHO performance status of 2 or less, **AND**
- Patient must have received prior chemotherapy treatment including an anthracycline, **AND**
- Patient must not have received prior treatment with an angiogenesis inhibitor, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Patient must not have any of the following conditions:

adipocytic soft tissue sarcoma;

gastrointestinal stromal tumour (GIST);

rhabdomyosarcoma other than alveolar or pleomorphic;

chondrosarcoma;

osteosarcoma;

Ewings tumour/primitive neuroectodermal tumour;

dermofibromatosis sarcoma protuberans;
inflammatory myofibroblastic sarcoma;
malignant mesothelioma;
mixed mesodermal tumour of the uterus.

pazopanib 200 mg tablet, 90

10042M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	3386.85	41.30	Votrient [NV]

pazopanib 400 mg tablet, 60

10041L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	4462.08	41.30	Votrient [NV]

■ PAZOPANIB

Note Response Evaluation Criteria In Solid Tumours (RECIST) is defined as follows:

Complete response (CR) is disappearance of all target lesions.

Partial response (PR) is a 30% decrease in the sum of the longest diameter of target lesions.

Progressive disease (PD) is a 20% increase in the sum of the longest diameter of target lesions.

Stable disease (SD) is small changes that do not meet above criteria.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)**7458**

Advanced (unresectable and/or metastatic) soft tissue sarcoma

Treatment Phase: Continuing treatment beyond 3 months

Clinical criteria:

- Patient must have received an initial authority prescription for this drug for this condition, **AND**
- Patient must have stable or responding disease according to the Response Evaluation Criteria In Solid Tumours (RECIST), **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

pazopanib 200 mg tablet, 90

10047T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	3386.85	41.30	Votrient [NV]

pazopanib 400 mg tablet, 60

10043N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	4462.08	41.30	Votrient [NV]

■ PAZOPANIB

Note Response Evaluation Criteria In Solid Tumours (RECIST) is defined as follows:

Complete response (CR) is disappearance of all target lesions.

Partial response (PR) is a 30% decrease in the sum of the longest diameter of target lesions.

Progressive disease (PD) is a 20% increase in the sum of the longest diameter of target lesions.

Stable disease (SD) is small changes that do not meet above criteria.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)**7459**

Advanced (unresectable and/or metastatic) soft tissue sarcoma

Treatment Phase: Continuing treatment beyond 3 months

Clinical criteria:

- Patient must have received an initial authority prescription for this drug for this condition, **AND**
- Patient must have stable or responding disease according to the Response Evaluation Criteria In Solid Tumours (RECIST), **AND**
- Patient must require dose adjustment, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

pazopanib 400 mg tablet, 30

10052C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	2311.62	41.30	Votrient [NV]

pazopanib 200 mg tablet, 30

10054E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1192.86	41.30	Votrient [NV]

■ **PAZOPANIB**

Note Response Evaluation Criteria In Solid Tumours (RECIST) is defined as follows:

Complete response (CR) is disappearance of all target lesions.

Partial response (PR) is a 30% decrease in the sum of the longest diameter of target lesions.

Progressive disease (PD) is a 20% increase in the sum of the longest diameter of target lesions.

Stable disease (SD) is small changes that do not meet above criteria.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

7423

Stage IV clear cell variant renal cell carcinoma (RCC)

Treatment Phase: Continuing treatment beyond 3 months

Clinical criteria:

- Patient must have received an initial authority prescription for this drug for this condition, **AND**
- Patient must have stable or responding disease according to the Response Evaluation Criteria In Solid Tumours (RECIST), **AND**
- The treatment must be the sole PBS-subsidised tyrosine kinase inhibitor therapy for this condition.

A patient who has progressive disease when treated with this drug is no longer eligible for PBS-subsidised treatment with this drug.

Patients who have progressive disease on sunitinib are not eligible to receive PBS-subsidised pazopanib.

pazopanib 200 mg tablet, 90

11252F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	3386.85	41.30	Votrient [NV]

pazopanib 400 mg tablet, 60

11261Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	4462.08	41.30	Votrient [NV]

■ **PAZOPANIB**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Patients who have developed intolerance to sunitinib of a severity necessitating permanent treatment withdrawal are eligible to receive PBS-subsidised pazopanib.

Note Patients who have progressive disease with pazopanib are no longer eligible for PBS-subsidised pazopanib.

Authority required (STREAMLINED)

9281

Stage IV clear cell variant renal cell carcinoma (RCC)

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be classified as favourable to intermediate risk according to the International Metastatic Renal Cell Carcinoma Database Consortium (IMDC), **AND**
- Patient must have a WHO performance status of 2 or less, **AND**
- The treatment must be the sole PBS-subsidised tyrosine kinase inhibitor therapy for this condition.

Patients who have progressive disease on sunitinib are not eligible to receive PBS-subsidised pazopanib.

pazopanib 200 mg tablet, 90

2029T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	3386.85	41.30	Votrient [NV]

pazopanib 400 mg tablet, 60

2030W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	4462.08	41.30	Votrient [NV]

■ **SORAFENIB**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Response Evaluation Criteria In Solid Tumours (RECIST) is defined as follows:

Complete response (CR) is disappearance of all target lesions.

Partial response (PR) is a 30% decrease in the sum of the longest diameter of target lesions.

Progressive disease (PD) is a 20% increase in the sum of the longest diameter of target lesions.

Stable disease (SD) is small changes that do not meet above criteria.

Authority required

Stage IV clear cell variant renal cell carcinoma (RCC)

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have progressive disease according to the Response Evaluation Criteria in Solid Tumours (RECIST) following prior treatment with a tyrosine kinase inhibitor, **AND**
- Patient must have a WHO performance status of 2 or less, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Patients who have developed intolerance to a tyrosine kinase inhibitor of a severity necessitating permanent treatment withdrawal are eligible to receive PBS-subsidised treatment with this drug.

A patient who has progressive disease when treated with this drug is no longer eligible for PBS-subsidised treatment with this drug.

sorafenib 200 mg tablet, 60

10226F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	2	..	*5556.82	41.30	Nexavar [BN]

▪ **SORAFENIB**

Note Response Evaluation Criteria In Solid Tumours (RECIST) is defined as follows:

Complete response (CR) is disappearance of all target lesions.

Partial response (PR) is a 30% decrease in the sum of the longest diameter of target lesions.

Progressive disease (PD) is a 20% increase in the sum of the longest diameter of target lesions.

Stable disease (SD) is small changes that do not meet above criteria.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

7487

Stage IV clear cell variant renal cell carcinoma (RCC)

Treatment Phase: Continuing treatment beyond 3 months

Clinical criteria:

- Patient must have received an initial authority prescription for this drug for this condition, **AND**
- Patient must have stable or responding disease according to the Response Evaluation Criteria In Solid Tumours (RECIST), **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

A patient who has progressive disease when treated with this drug is no longer eligible for PBS-subsidised treatment with this drug.

sorafenib 200 mg tablet, 60

10242C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*5556.82	41.30	Nexavar [BN]

▪ **SORAFENIB**

Note Sorafenib is not PBS-subsidised for adjunctive treatment after resection, ablation or chemoembolization.

Sorafenib is not PBS-subsidised for maintenance therapy after disease progression.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

11160

Advanced Barcelona Clinic Liver Cancer Stage B or Stage C hepatocellular carcinoma

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have a WHO performance status of 2 or less, **AND**
- Patient must have Child Pugh class A, **AND**
- The condition must be untreated with systemic therapy; OR
- Patient must have developed intolerance of a severity necessitating permanent treatment withdrawal, in the absence of disease progression, to any of the following: (i) a vascular endothelial growth factor (VEGF) tyrosine kinase inhibitor (TKI), (ii) atezolizumab/bevacizumab combination therapy.

Authority required (STREAMLINED)

8617

Advanced Barcelona Clinic Liver Cancer Stage B or Stage C hepatocellular carcinoma

Treatment Phase: Continuing treatment

Clinical criteria:

- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not develop disease progression while receiving treatment with this drug for this condition.

sorafenib 200 mg tablet, 60

9380Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	2	..	*5556.82	41.30	Nexavar [BN]

■ SUNITINIB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Metastatic or unresectable, well-differentiated malignant pancreatic neuroendocrine tumour (pNET)

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must be symptomatic (despite somatostatin analogues); OR
- Patient must have disease progression, **AND**
- The treatment must be as monotherapy.

Disease progression must be documented in the patient's medical records.

Patients who have developed progressive disease on everolimus are not eligible to receive PBS-subsidised sunitinib for this condition.

Patients who have developed intolerance to everolimus of a severity necessitating permanent treatment withdrawal are eligible to receive PBS-subsidised sunitinib.

sunitinib 12.5 mg capsule, 28

10004M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	1579.13	41.30	Sutent [PF]

sunitinib 50 mg capsule, 28

2837H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	5933.34	41.30	Sutent [PF]

sunitinib 37.5 mg capsule, 28

10464R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	4490.23	41.30	Sutent [PF]

sunitinib 25 mg capsule, 28

2959R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	3047.13	41.30	Sutent [PF]

■ SUNITINIB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)**7471**

Metastatic or unresectable, well-differentiated malignant pancreatic neuroendocrine tumour (pNET)

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have received an initial authority prescription for this drug for this condition, **AND**
- Patient must not have disease progression, **AND**
- The treatment must be as monotherapy.

A patient who has progressive disease when treated with this drug is no longer eligible for PBS-subsidised treatment with this drug.

sunitinib 12.5 mg capsule, 28

10009T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1579.13	41.30	Sutent [PF]

sunitinib 50 mg capsule, 28

10010W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	5933.34	41.30	Sutent [PF]

sunitinib 37.5 mg capsule, 28

10473F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	4490.23	41.30	Sutent [PF]

sunitinib 25 mg capsule, 28

2842N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	3047.13	41.30	Sutent [PF]

■ SUNITINIB

Note Response Evaluation Criteria In Solid Tumours (RECIST) is defined as follows:

Complete response (CR) is disappearance of all target lesions.

Partial response (PR) is a 30% decrease in the sum of the longest diameter of target lesions.

Progressive disease (PD) is a 20% increase in the sum of the longest diameter of target lesions.

Stable disease (SD) is small changes that do not meet above criteria.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

7466

Stage IV clear cell variant renal cell carcinoma (RCC)

Treatment Phase: Continuing treatment beyond 3 months

Clinical criteria:

- Patient must have received an initial authority prescription for this drug for this condition, **AND**
- Patient must have stable or responding disease according to the Response Evaluation Criteria In Solid Tumours (RECIST), **AND**
- The treatment must be the sole PBS-subsidised tyrosine kinase inhibitor therapy for this condition.

A patient who has progressive disease when treated with this drug is no longer eligible for PBS-subsidised treatment with this drug.

Patients who have developed progressive disease on pazopanib are not eligible to receive PBS-subsidised sunitinib.

sunitinib 12.5 mg capsule, 28

9420T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	1579.13	41.30	Sutent [PF]

sunitinib 50 mg capsule, 28

9422X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	5933.34	41.30	Sutent [PF]

sunitinib 37.5 mg capsule, 28

10459L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	4490.23	41.30	Sutent [PF]

sunitinib 25 mg capsule, 28

9421W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	3047.13	41.30	Sutent [PF]

■ SUNITINIB

Note Sunitinib malate is not PBS-subsidised for the treatment of patients with resectable malignant gastrointestinal stromal tumours.

Note Written applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Metastatic or unresectable malignant gastrointestinal stromal tumour

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be as monotherapy, **AND**
- Patient must have a WHO performance status of 2 or less, **AND**
- Patient must have previously failed or be intolerant to imatinib mesilate.

Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Sunitinib Malate (Sutent) PBS Authority Application for Use in the Treatment of Gastrointestinal Stromal Tumour - Supporting Information Form; and
- (3) a signed patient acknowledgement.

Patients who have failed to respond or are intolerant to imatinib are no longer eligible to receive PBS-subsidised imatinib

sunitinib 12.5 mg capsule, 28

9488J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	1579.13	41.30	Sutent [PF]

sunitinib 50 mg capsule, 28

9490L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	5933.34	41.30	Sutent [PF]

sunitinib 37.5 mg capsule, 28

10503T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	4490.23	41.30	Sutent [PF]

sunitinib 25 mg capsule, 28

9489K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	3047.13	41.30	Sutent [PF]

■ SUNITINIB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Patients who have developed intolerance to pazopanib of a severity necessitating permanent treatment withdrawal are eligible to receive PBS-subsidised sunitinib.

Note Patients who have progressive disease with sunitinib are no longer eligible for PBS-subsidised sunitinib.

Authority required (STREAMLINED)**9210**

Stage IV clear cell variant renal cell carcinoma (RCC)

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be classified as favourable to intermediate risk according to the International Metastatic Renal Cell Carcinoma Database Consortium (IMDC), **AND**
 - Patient must have a WHO performance status of 2 or less, **AND**
 - The treatment must be the sole PBS-subsidised tyrosine kinase inhibitor therapy for this condition.
- Patients who have developed progressive disease on pazopanib are not eligible to receive PBS-subsidised sunitinib.

sunitinib 12.5 mg capsule, 28

9417P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	1579.13	41.30	Sutent [PF]

sunitinib 50 mg capsule, 28

9419R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	5933.34	41.30	Sutent [PF]

sunitinib 37.5 mg capsule, 28

10504W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	4490.23	41.30	Sutent [PF]

sunitinib 25 mg capsule, 28

9418Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	3047.13	41.30	Sutent [PF]

■ SUNITINIB

Note A patient who has progressive disease when treated with this drug is no longer eligible for PBS-subsidised treatment with this drug.

Note Patients who have failed to respond or are intolerant to imatinib are no longer eligible to receive PBS subsidised imatinib after progression on this drug

Note Sunitinib malate is not PBS-subsidised for the treatment of patients with resectable malignant gastrointestinal stromal tumours.

Note No increase in the maximum number of repeats may be authorised.

Note No increase in the maximum quantity or number of units may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)**7430**

Metastatic or unresectable malignant gastrointestinal stromal tumour

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have received an initial authority prescription for this drug for this condition, **AND**
- The treatment must be as monotherapy, **AND**
- Patient must have a WHO performance status of 2 or less, **AND**
- Patient must not have progressive disease.

sunitinib 12.5 mg capsule, 28

11266Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	1579.13	41.30	Sutent [PF]

sunitinib 50 mg capsule, 28

11250D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	5933.34	41.30	Sutent [PF]

sunitinib 37.5 mg capsule, 28

11256K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	4490.23	41.30	Sutent [PF]

sunitinib 25 mg capsule, 28

11253G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	3047.13	41.30	Sutent [PF]

OTHER ANTINEOPLASTIC AGENTS*Monoclonal antibodies***■ RITUXIMAB****Note** Special Pricing Arrangements apply.**Authority required (STREAMLINED)****7400**

Previously untreated or relapsed/refractory CD20 positive lymphoid cancer

Treatment Phase: Induction or re-induction therapy

Clinical criteria:

- The treatment must be for induction or re-induction for CD20 positive lymphoma; OR
- The treatment must be for induction or re-induction for CD20 positive chronic lymphocytic leukaemia; OR
- The treatment must be for induction or consolidation for CD20 positive acute lymphoblastic leukaemia, **AND**
- The treatment must be in combination with chemotherapy, **AND**
- Patient must not receive more than the number of cycles of treatment recommended by standard guidelines for the partner chemotherapy under this restriction.

An initial dose of rituximab must be administered with rituximab intravenous injection. Subsequent doses may be administered with either intravenous or subcutaneous rituximab.

No more than 8 doses in total as per course of treatment will be allowed for lymphoma or chronic lymphocytic leukaemia.

No more than 12 doses in total as per course of treatment will be allowed for acute lymphoblastic leukaemia for induction course (including consolidation course).

rituximab 1.4 g/11.7 mL injection, 11.7 mL vial

10703H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	6	..	1847.94	41.30	Mabthera SC [RO]

■ RITUXIMAB**Note** Special Pricing Arrangements apply.**Note** No increase in the maximum number of repeats may be authorised.**Authority required (STREAMLINED)****6011**

Relapsed or refractory Stage III or IV CD20 positive follicular B-cell non-Hodgkin's lymphoma

Treatment Phase: Maintenance therapy

Clinical criteria:

- The treatment must be maintenance therapy, **AND**
- Patient must have demonstrated a partial or complete response to re-induction treatment received immediately prior to this current Authority application, **AND**
- Patient must not receive more than 8 cycles or 2 years duration of treatment, whichever comes first, under this restriction.

rituximab 1.4 g/11.7 mL injection, 11.7 mL vial

10709P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	7	..	1847.94	41.30	Mabthera SC [RO]

■ RITUXIMAB**Note** Special Pricing Arrangements apply.**Note** No increase in the maximum number of repeats may be authorised.**Authority required (STREAMLINED)****7399**

Previously untreated or Relapsed/refractory CD20 positive acute lymphoblastic leukaemia

Treatment Phase: Maintenance therapy

Clinical criteria:

- The treatment must be maintenance therapy, **AND**
- The treatment must be in combination with chemotherapy, **AND**
- Patient must be in complete remission, **AND**
- Patient must not receive more than 6 doses in total under this restriction.

rituximab 1.4 g/11.7 mL injection, 11.7 mL vial

10719E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1847.94	41.30	Mabthera SC [RO]

■ RITUXIMAB

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)**10227**

Relapsed or refractory follicular B-cell non-Hodgkin's lymphoma

Treatment Phase: Re-induction therapy

Clinical criteria:

- The treatment must be for re-induction treatment purposes only, **AND**
- The condition must have relapsed or be refractory to treatment, **AND**
- Patient must not receive more than 4 doses of rituximab in total, including intravenous and subcutaneous injections, and no more than 3 doses of subcutaneous rituximab under this restriction.

An initial dose of rituximab must be administered with rituximab intravenous injection. Subsequent doses may be administered with either intravenous or subcutaneous rituximab with no more than 4 doses in total.

rituximab 1.4 g/11.7 mL injection, 11.7 mL vial

11988Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	1847.94	41.30	Mabthera SC [RO]

■ RITUXIMAB

Note Special Pricing Arrangements apply.

Note A patient may only qualify for PBS-subsidised treatment under this restriction once in a lifetime.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)**6161**

Stage III or IV CD20 positive follicular B-cell non-Hodgkin's lymphoma

Treatment Phase: Maintenance therapy

Clinical criteria:

- Patient must have demonstrated a partial or complete response to induction treatment with either R-CHOP or R-CVP regimens for previously untreated follicular B-cell Non-Hodgkin's lymphoma, received immediately prior to this current Authority application, **AND**
- Patient must not have received bendamustine induction therapy, **AND**
- The treatment must be maintenance therapy, **AND**
- Patient must not receive more than 12 doses or 2 years duration of treatment, whichever comes first, under this restriction.

rituximab 1.4 g/11.7 mL injection, 11.7 mL vial

10742J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	11	..	1847.94	41.30	Mabthera SC [RO]

■ TRASTUZUMAB

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)**10212**

Early HER2 positive breast cancer

Treatment Phase: 3 weekly treatment regimen

Clinical criteria:

- Patient must have undergone surgery (adjuvant) or be preparing for surgery (neoadjuvant), **AND**
- The treatment must not be used in a patient with a left ventricular ejection fraction (LVEF) of less than 45% and/or with symptomatic heart failure, **AND**
- Patient must not receive more than 52 weeks of combined PBS-subsidised and non-PBS-subsidised therapy; OR
- Patient must not receive more than 52 weeks of combined trastuzumab and trastuzumab emtansine therapy if adjuvant trastuzumab emtansine therapy has been discontinued due to intolerance.

Cardiac function must be tested by echocardiography (ECHO) or multigated acquisition (MUGA), prior to initiating treatment with this drug for this condition.

trastuzumab 600 mg/5 mL injection, 5 mL vial

10682F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	1607.60	41.30	Herceptin SC [RO]

▪ **TRASTUZUMAB**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required (STREAMLINED)

9353

Metastatic (Stage IV) HER2 positive breast cancer

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have evidence of human epidermal growth factor receptor 2 (HER2) gene amplification as demonstrated by in situ hybridisation (ISH) either in the primary tumour or a metastatic lesion, **AND**
- The treatment must not be in combination with nab-paclitaxel, **AND**
- The treatment must not be used in a patient with a left ventricular ejection fraction (LVEF) of less than 45% and/or with symptomatic heart failure.

Cardiac function must be tested by echocardiography (ECHO) or multigated acquisition (MUGA), prior to initiating treatment with this drug for this condition.

trastuzumab 600 mg/5 mL injection, 5 mL vial

10798H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	1607.60	41.30	Herceptin SC [RO]

▪ **TRASTUZUMAB**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required (STREAMLINED)

9462

Metastatic (Stage IV) HER2 positive breast cancer

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must not be used in a patient with a left ventricular ejection fraction (LVEF) of less than 45% and/or with symptomatic heart failure.

trastuzumab 600 mg/5 mL injection, 5 mL vial

10803N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	1607.60	41.30	Herceptin SC [RO]

Histone deacetylase (HDAC) inhibitors

▪ **VORINOSTAT**

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Cutaneous T-cell lymphoma

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have received systemic treatment with chemotherapy, **AND**
- Patient must demonstrate relapsed or chemotherapy-refractory disease, **AND**
- Patient must be ineligible for stem cell transplant, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Applications for authorisation of initial treatment must be in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed cutaneous T-cell lymphoma (CTCL) initial PBS Authority Application - Supporting Information Form.

vorinostat 100 mg capsule, 120

11138F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	4463.28	41.30	Zolinza [MK]

■ VORINOSTAT

Note Authority applications for continuing treatment may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Cutaneous T-cell lymphoma

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have progressive disease while receiving PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

vorinostat 100 mg capsule, 120

11141J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	4463.28	41.30	Zolinza [MK]

Hedgehog pathway inhibitors

■ SONIDEGIB

Caution Sonidegib is a category X drug and must not be given to pregnant women. Pregnancy in female patients or in the partners of male patients must be avoided during treatment and during the 20 months and 6 months period after cessation of treatment for female and male patients respectively, as according to the TGA approved Product Information.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Metastatic or locally advanced basal cell carcinoma

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be inappropriate for surgery, **AND**
- The condition must be inappropriate for curative radiotherapy, **AND**
- Patient must not have received previous PBS-subsidised treatment with another hedgehog (Hh) inhibitor for this condition; **OR**
- Patient must have developed intolerance to another hedgehog (Hh) inhibitor of a severity necessitating permanent treatment withdrawal, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

The authority application must be made in writing and must include:

- a) A completed authority prescription form; and
- b) A completed Basal Cell Carcinoma Initial PBS Authority Application Form - Supporting Information Form; and
- c) A histological confirmation of BCC and whether the condition is metastatic or locally advanced; and
- d) A letter from a surgically qualified clinician demonstrating inappropriateness for surgery for patients with locally advanced BCC; and
- e) A letter from a radiation oncologist demonstrating inappropriateness for curative radiotherapy for patients with locally advanced BCC; and
- f) A signed patient acknowledgement.

The assessment of the patient's response to this PBS-subsidised course of therapy must be made within the 4 weeks prior to completion of the course of treatment. It is recommended that an application is submitted to the Department of Human Services no less than 2 weeks prior to the date the next dose is due in order to ensure continuity of treatment for those patients who meet the continuation criteria.

Inappropriate for surgery is defined as:

- i/ Curative resection is unlikely, such as where BCC has recurred in the same location after two or more surgical procedures; **or**
- ii/ Anticipated substantial morbidity or deformity from surgery or requiring complicated reconstructive surgery (e.g. removal of all or part of a facial structure, such as nose, ear, eyelid, eye; or requirement for limb amputation or free tissue transfer); **or**
- iii/ Medical contraindication to surgery

Inappropriate for curative radiotherapy is defined as:

- i/Hypersensitivity to radiation due to genetic syndrome such as Gorlin Syndrome; **or**

- ii/ Limitations due to location of tumour; or
- iii/ Limitations due to cumulative prior radiotherapy dose; or
- iv/ Progressive disease despite prior irradiation of locally advanced BCC.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs Programs
Reply Paid 9826
HOBART TAS 7001

Authority required

Metastatic or locally advanced basal cell carcinoma

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not develop disease progression while receiving PBS-subsidised treatment with this drug for this condition, **AND**
- The condition must remain inappropriate for surgery, **AND**
- The condition must remain inappropriate for curative radiotherapy, **AND**
- Patient must not receive more than 16 weeks of treatment per continuing treatment under this restriction.

The authority application must be made in writing and must include:

- a) A completed authority prescription form; and
- b) A completed Basal Cell Carcinoma Continuing PBS Authority Application Form - Supporting Information Form; and
- c) A confirmation statement from the treating doctor that the disease has not progressed; and
- d) In patients with locally advanced BCC, a letter from a surgically qualified clinician demonstrating that the condition remains inappropriate for surgery; or a letter from a radiation oncologist demonstrating that the condition remains inappropriate for curative radiotherapy

The assessment of the patient's response to this PBS-subsidised course of therapy must be made within the 4 weeks prior to completion of the course of treatment. It is recommended that an application is submitted to the Department of Human Services no less than 2 weeks prior to the date the next dose is due in order to ensure continuity of treatment for those patients who meet the continuation criteria.

Inappropriate for surgery is defined as:

- i/ Curative resection is unlikely, such as where BCC has recurred in the same location after two or more surgical procedures; or
- ii/ Anticipated substantial morbidity or deformity from surgery or requiring complicated reconstructive surgery (e.g. removal of all or part of a facial structure, such as nose, ear, eyelid, eye; or requirement for limb amputation or free tissue transfer); or
- iii/ Medical contraindication to surgery

Inappropriate for curative radiotherapy is defined as:

- i/ Hypersensitivity to radiation due to genetic syndrome such as Gorlin Syndrome; or
- ii/ Limitations due to location of tumour; or
- iii/ Limitations due to cumulative prior radiotherapy dose; or
- iv/ Progressive disease despite prior irradiation of locally advanced BCC

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs Programs
Reply Paid 9826
HOBART TAS 7001

Authority required

Metastatic or locally advanced basal cell carcinoma

Treatment Phase: Initial treatment or Continuing treatment – balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the Initial treatment restriction to complete maximum of 16 weeks of treatment; OR
- Patient must have received insufficient therapy with this drug under the Continuing treatment restriction to complete maximum of 16 weeks of treatment, **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Note Authority approval for sufficient therapy to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

sonidegib 200 mg capsule, 30

11304Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	7982.59	41.30	Odomzo [RA]

■ VISMODEGIB

Caution Vismodegib is a category X drug and must not be given to pregnant women. Pregnancy in female patients or in the partners of male patients must be avoided during treatment and during the 24 months and 2 months period after cessation of treatment for female and male patients respectively, as according to the TGA approved Product Information.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Metastatic or locally advanced basal cell carcinoma

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be inappropriate for surgery, **AND**
- The condition must be inappropriate for curative radiotherapy, **AND**
- Patient must not have received previous PBS-subsidised treatment with another hedgehog (Hh) inhibitor for this condition; **OR**
- Patient must have developed intolerance to another hedgehog (Hh) inhibitor of a severity necessitating permanent treatment withdrawal, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

The authority application must be made in writing and must include:

- a) A completed authority prescription form; and
- b) A completed Basal Cell Carcinoma Initial PBS Authority Application Form - Supporting Information Form; and
- c) A histological confirmation of BCC and whether the condition is metastatic or locally advanced; and
- d) A letter from a surgically qualified clinician demonstrating inappropriateness for surgery for patients with locally advanced BCC; and
- e) A letter from a radiation oncologist demonstrating inappropriateness for curative radiotherapy for patients with locally advanced BCC; and
- f) A signed patient acknowledgement.

The assessment of the patient's response to this PBS-subsidised course of therapy must be made within the 4 weeks prior to completion of the course of treatment. It is recommended that an application is submitted to the Department of Human Services no less than 2 weeks prior to the date the next dose is due in order to ensure continuity of treatment for those patients who meet the continuation criteria.

Inappropriate for surgery is defined as:

- i/ Curative resection is unlikely, such as where BCC has recurred in the same location after two or more surgical procedures; **or**
- ii/ Anticipated substantial morbidity or deformity from surgery or requiring complicated reconstructive surgery (e.g. removal of all or part of a facial structure, such as nose, ear, eyelid, eye; or requirement for limb amputation or free tissue transfer); **or**
- iii/ Medical contraindication to surgery

Inappropriate for curative radiotherapy is defined as:

- i/ Hypersensitivity to radiation due to genetic syndrome such as Gorlin Syndrome; **or**
- ii/ Limitations due to location of tumour; **or**
- iii/ Limitations due to cumulative prior radiotherapy dose; **or**
- iv/ Progressive disease despite prior irradiation of locally advanced BCC.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs Programs

Reply Paid 9826

HOBART TAS 7001

Authority required

Metastatic or locally advanced basal cell carcinoma

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not develop disease progression while receiving PBS-subsidised treatment with this drug for this condition, **AND**
- The condition must remain inappropriate for surgery, **AND**
- The condition must remain inappropriate for curative radiotherapy, **AND**
- Patient must not receive more than 16 weeks of treatment per continuing treatment under this restriction.

The authority application must be made in writing and must include:

- a) A completed authority prescription form; and
- b) A completed Basal Cell Carcinoma Continuing PBS Authority Application Form - Supporting Information Form; and
- c) A confirmation statement from the treating doctor that the disease has not progressed; and
- d) In patients with locally advanced BCC, a letter from a surgically qualified clinician demonstrating that the condition remains inappropriate for surgery; or a letter from a radiation oncologist demonstrating that the condition remains inappropriate for curative radiotherapy

The assessment of the patient's response to this PBS-subsidised course of therapy must be made within the 4 weeks prior to completion of the course of treatment. It is recommended that an application is submitted to the Department of Human Services no less than 2 weeks prior to the date the next dose is due in order to ensure continuity of treatment for those patients who meet the continuation criteria.

Inappropriate for surgery is defined as:

- i/ Curative resection is unlikely, such as where BCC has recurred in the same location after two or more surgical procedures; or
- ii/ Anticipated substantial morbidity or deformity from surgery or requiring complicated reconstructive surgery (e.g. removal of all or part of a facial structure, such as nose, ear, eyelid, eye; or requirement for limb amputation or free tissue transfer); or
- iii/ Medical contraindication to surgery

Inappropriate for curative radiotherapy is defined as:

- i/ Hypersensitivity to radiation due to genetic syndrome such as Gorlin Syndrome; or
- ii/ Limitations due to location of tumour; or
- iii/ Limitations due to cumulative prior radiotherapy dose; or
- iv/ Progressive disease despite prior irradiation of locally advanced BCC

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs Programs

Reply Paid 9826

HOBART TAS 7001

Authority required

Metastatic or locally advanced basal cell carcinoma

Treatment Phase: Initial treatment or Continuing treatment – balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the Initial treatment restriction to complete maximum of 16 weeks of treatment; OR
- Patient must have received insufficient therapy with this drug under the Continuing treatment restriction to complete maximum of 16 weeks of treatment, **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Note Authority approval for sufficient therapy to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

vismodegib 150 mg capsule, 28

11070P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	7461.16	41.30	Erivedge [RO]

Poly (ADP-ribose) polymerase (PARP) inhibitors

▪ **OLAPARIB**

Caution Do not substitute olaparib 50 mg capsules with olaparib 100 mg or 150 mg tablets on a mg to mg basis due to difference in dosing and bioavailability of each formulation.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

10937

High grade epithelial ovarian, fallopian tube or primary peritoneal cancer

Treatment Phase: Continuing treatment - second line treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug as a second line therapy for this condition, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- The treatment must be maintenance therapy, **AND**
- Patient must not have developed disease progression while receiving treatment with this drug for this condition.

A response (complete or partial) to the platinum-based chemotherapy regimen is to be assessed using either Gynaecologic Cancer InterGroup (GCIG) or Response Evaluation Criteria in Solid Tumours (RECIST) guidelines.

olaparib 150 mg tablet, 56

11539H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*6971.16	41.30	Lynparza [AP]

olaparib 100 mg tablet, 56

11503K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*6971.16	41.30	Lynparza [AP]

olaparib 50 mg capsule, 4 x 112

11050N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	6971.16	41.30	Lynparza [AP]

■ OLAPARIB

Caution Do not substitute olaparib 50 mg capsules with olaparib 100 mg or 150 mg tablets on a mg to mg basis due to difference in dosing and bioavailability of each formulation.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Note Special Pricing Arrangements apply.

Authority required

High grade epithelial ovarian, fallopian tube or primary peritoneal cancer

Treatment Phase: Initial treatment - second line treatment

Clinical criteria:

- The condition must be associated with a class 4 or 5 BRCA1 or BRCA2 gene mutation, **AND**
- The condition must be platinum sensitive, **AND**
- Patient must have received at least two previous platinum-containing regimens, **AND**
- Patient must have relapsed following a previous platinum-containing regimen, **AND**
- Patient must be in partial or complete response to the immediately preceding platinum-based chemotherapy regimen, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- The treatment must be maintenance therapy, **AND**
- Patient must not have previously received PBS-subsidised treatment with this drug for this condition.

Platinum sensitivity is defined as disease progression greater than 6 months after completion of the penultimate platinum regimen.

A response (complete or partial) to the platinum-based chemotherapy regimen is to be assessed using either Gynaecologic Cancer InterGroup (GCIG) or Response Evaluation Criteria in Solid Tumours (RECIST) guidelines.

Evidence of a BRCA1 or BRCA2 gene mutation must be derived through germline or somatic mutation testing.

olaparib 150 mg tablet, 56

11528R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	2	..	*6971.16	41.30	Lynparza [AP]

olaparib 100 mg tablet, 56

11522K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	2	..	*6971.16	41.30	Lynparza [AP]

■ OLAPARIB

Caution Do not substitute olaparib 50 mg capsules with olaparib 100 mg or 150 mg tablets on a mg to mg basis due to difference in dosing and bioavailability of each formulation.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Note Special Pricing Arrangements apply.

Authority required

High grade stage III/IV epithelial ovarian, fallopian tube or primary peritoneal cancer

Treatment Phase: Initial treatment - first line treatment

Clinical criteria:

- The condition must be associated with a class 4 or 5 BRCA1 or BRCA2 gene mutation, **AND**
- Patient must be in partial or complete response to the immediately preceding platinum-based chemotherapy regimen prior to commencing treatment with this drug for this condition, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must not have previously received PBS-subsidised treatment with this drug for this condition.

A response (complete or partial) to the platinum-based chemotherapy regimen is to be assessed using either Gynaecologic Cancer InterGroup (GCIG) or Response Evaluation Criteria in Solid Tumours (RECIST) guidelines.

Evidence of a BRCA1 or BRCA2 gene mutation must be derived through germline or somatic mutation testing.

olaparib 150 mg tablet, 56

12157W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	2	..	*6971.16	41.30	Lynparza [AP]

olaparib 100 mg tablet, 56

12170M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	2	..	*6971.16	41.30	Lynparza [AP]

▪ **OLAPARIB**

Caution Do not substitute olaparib 50 mg capsules with olaparib 100 mg or 150 mg tablets on a mg to mg basis due to difference in dosing and bioavailability of each formulation.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Note Special Pricing Arrangements apply.

Authority required

High grade stage III/IV epithelial ovarian, fallopian tube or primary peritoneal cancer

Treatment Phase: Continuing treatment - first line treatment

Clinical criteria:

- Patient must have received previous PBS-subsidised treatment with this drug as first line maintenance therapy for this condition, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must not have developed disease progression while receiving treatment with this drug for this condition, **AND**
- The treatment must not exceed a total of 24 months of combined non-PBS-subsidised and PBS-subsidised treatment for patients who are in complete response.

olaparib 150 mg tablet, 56

12161C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*6971.16	41.30	Lynparza [AP]

olaparib 100 mg tablet, 56

12169L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*6971.16	41.30	Lynparza [AP]

Other antineoplastic agents

▪ **HYDROXYCARBAMIDE (HYDROXYUREA)**

hydroxycarbamide (hydroxyurea) 500 mg capsule, 100

3093T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	3	..	50.81	41.30	^a Hydrea [BQ]	^a HYDROXYCARBAMIDE MEDSURGE [DZ]

▪ **VENETOCLAX**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Chronic lymphocytic leukaemia (CLL)

Treatment Phase: Initial treatment - Dose titration

Clinical criteria:

- Patient must not have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must be considered unsuitable for treatment or retreatment with a purine analogue, **AND**
- The condition must have relapsed or be refractory to at least one prior therapy, **AND**
- Patient must have a WHO performance status of 0 or 1, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- The treatment must be used as monotherapy for this condition under this restriction.

A patient is considered unsuitable for treatment or retreatment with a purine analogue as demonstrated by at least one of the following:

- Failure to respond (stable disease or disease progression on treatment), or a progression-free interval of less than 3 years from treatment with a purine analogue-based therapy and anti-CD20-containing chemoimmunotherapy regimen after at least two cycles;
- Age is 70 years or older;
- Age is 65 years or older and the presence of comorbidities (Cumulative Illness Rating Scale of 6 or greater, or creatinine clearance of less than 70 mL/min) that might place the patient at an unacceptable risk for treatment-related toxicity with purine analogue-based therapy, provided they have received one or more prior treatment including at least two cycles of an alkylating agent-based (or purine analogue-based) anti-CD20 antibody-containing chemoimmunotherapy regimen;
- History of purine analogue-associated autoimmune anaemia or autoimmune thrombocytopenia;
- Evidence of one or more 17p chromosomal deletions demonstrated by fluorescence in situ hybridisation (FISH).

venetoclax 10 mg tablet [14] (&) venetoclax 50 mg tablet [7] (&) venetoclax 100 mg tablet [7] (&) venetoclax 100 mg tablet [14], 1 pack

11630D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	1791.43	41.30	Venclexta [VE]

▪ VENETOCLAX

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Authority required

Chronic lymphocytic leukaemia (CLL) or small lymphocytic lymphoma (SLL)

Treatment Phase: Dose modification

Clinical criteria:

- The treatment must be for dose titration purposes.

venetoclax 10 mg tablet, 14

11624T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	107.65	41.30	Venclexta [VE]

venetoclax 50 mg tablet, 7

11648C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	258.03	41.30	Venclexta [VE]

▪ VENETOCLAX

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Authority required

Chronic lymphocytic leukaemia (CLL)

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be in combination with rituximab for up to a maximum of 6 cycles, followed by monotherapy, **AND**
- The treatment must be ceased on disease progression or on completion of 24 months of PBS-subsidised treatment under this restriction with this drug for this condition, whichever comes first.

venetoclax 100 mg tablet, 120

11639N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	7784.18	41.30	Venclexta [VE]

▪ VENETOCLAX

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Authority required

Chronic lymphocytic leukaemia (CLL) or small lymphocytic lymphoma (SLL)

Treatment Phase: Initial treatment in first-line therapy - Dose titration (weeks 1 to 4 of a 5-week ramp-up schedule)

Clinical criteria:

- The condition must be untreated, **AND**
- Patient must be inappropriate for fludarabine based chemo-immunotherapy, **AND**
- The treatment must be in combination with obinutuzumab (refer to Product Information for timing of obinutuzumab and venetoclax doses), **AND**
- Patient must have a creatinine clearance 30 mL/min or greater, **AND**
- Patient must have a total cumulative illness rating scale (CIRS) score of greater than 6 (excluding CLL-induced illness or organ damage); OR
- Patient must have a creatinine clearance less than 70 mL/min.

venetoclax 10 mg tablet [14] (&) venetoclax 50 mg tablet [7] (&) venetoclax 100 mg tablet [7] (&) venetoclax 100 mg tablet [14], 1 pack

12188L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	1791.43	41.30	Venclexta [VE]

■ VENETOCLAX

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Authority required

Chronic lymphocytic leukaemia (CLL) or small lymphocytic lymphoma (SLL)

Treatment Phase: Second and final continuing treatment prescription (treatment cycles 7 to 12 inclusive) of first-line therapy

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must cease upon disease progression; OR
- The treatment must cease upon completion of 12 cycles of treatment with this drug for this condition, whichever comes first.

venetoclax 100 mg tablet, 120

12199C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	7784.18	41.30	Venclexta [VE]

■ VENETOCLAX

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Authority required

Chronic lymphocytic leukaemia (CLL) or small lymphocytic lymphoma (SLL)

Treatment Phase: First continuing treatment (treatment cycles 2 to 6 inclusive) of first-line therapy

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be in combination with obinutuzumab (refer to Product Information for timing of obinutuzumab and venetoclax doses), **AND**
- The treatment must cease upon disease progression.

venetoclax 100 mg tablet, 120

12205J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	4	..	7784.18	41.30	Venclexta [VE]

■ ENDOCRINE THERAPY
HORMONES AND RELATED AGENTS
Progestogens
■ MEDROXYPROGESTERONE
Restricted benefit

Advanced breast cancer

Clinical criteria:

- The condition must be hormone receptor positive.

medroxyprogesterone acetate 500 mg tablet, 30

2728N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	138.39	41.30	Provera [PF]

■ MEDROXYPROGESTERONE
Restricted benefit

Breast cancer

Clinical criteria:

- The condition must be hormone receptor positive.

Restricted benefit

Endometrial cancer

medroxyprogesterone acetate 250 mg tablet, 60

2727M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	138.39	41.30	Provera [PF]

medroxyprogesterone acetate 200 mg tablet, 60

2316X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	111.71	41.30	Provera [PF]

medroxyprogesterone acetate 100 mg tablet, 100

2725K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	99.65	41.30	Provera [PF]

Gonadotropin releasing hormone analogues**■ GOSERELIN****Restricted benefit**

Carcinoma of the prostate

Clinical criteria:

- The condition must be locally advanced (stage C); OR
- The condition must be metastatic (stage D).

goserelin 10.8 mg implant, 1

8093Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	908.20	41.30	Zoladex 10.8 Implant [AP]

■ GOSERELIN**Restricted benefit**

Carcinoma of the prostate

Clinical criteria:

- The condition must be locally advanced (stage C); OR
- The condition must be metastatic (stage D).

Restricted benefit

Endometriosis

Clinical criteria:

- The condition must be visually proven, **AND**
- The treatment must be for the short-term (up to 6 months).

Note Only 1 course of not more than 6 months' therapy will be authorised.**Restricted benefit**

Breast cancer

Clinical criteria:

- The condition must be hormone receptor positive.

Restricted benefit

Anticipated premature ovarian failure

Clinical criteria:

- Patient must be receiving treatment with an alkylating agent for a malignancy or an autoimmune disorder that has a high risk of causing premature ovarian failure, **AND**
- Patient must not receive more than 6 months' of treatment for this condition in a lifetime.

Population criteria:

- Patient must be pre-menopausal.

goserelin 3.6 mg implant, 1

1454M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	270.20	41.30	Zoladex Implant [AP]

■ GOSERELIN (&) BICALUTAMIDE**Note** No increase in the maximum quantity or number of units may be authorised.**Note** No increase in the maximum number of repeats may be authorised.**Restricted benefit**

Carcinoma of the prostate

Clinical criteria:

- The condition must be metastatic (stage D), **AND**
- Patient must require a combination of an antiandrogen and a GnRH (LH-RH) agonist.

goserelin 10.8 mg implant [1 implant] (&) bicalutamide 50 mg tablet [28], 1 pack

9065D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	1049.08	41.30	ZolaCos CP 10.8/50(28) [AP]

goserelin 3.6 mg implant [1] (&) bicalutamide 50 mg tablet [28], 1 pack

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
9064C	±1	5	..	421.68	41.30	ZolaCos CP 3.6/50 [AP]

goserelin 10.8 mg implant [1 implant] (&) bicalutamide 50 mg tablet [84], 1 pack

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
9066E	±1	1	..	1330.84	41.30	ZolaCos CP 10.8/50(84) [AP]

▪ **LEUPRORELIN**

Restricted benefit

Locally advanced (stage C) or metastatic (stage D) carcinoma of the prostate

leuprorelin acetate 7.5 mg modified release injection [1 chamber] (&) inert substance diluent [1 mL chamber], 1 dual chamber syringe

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8875D	1	5	..	344.56	41.30	Lucrin Depot 7.5mg PDS [VE]

leuprorelin acetate 45 mg modified release injection [1 syringe] (&) inert substance diluent [1 syringe], 1 pack

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8859G	1	..	..	1750.50	41.30	Eligard 6 month [MF]

leuprorelin acetate 30 mg modified release injection [1 syringe] (&) inert substance diluent [1 syringe], 1 pack

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8709J	1	1	..	1189.13	41.30	Eligard 4 month [MF]

leuprorelin acetate 22.5 mg modified release injection [1 syringe] (&) inert substance diluent [1 syringe], 1 pack

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8708H	1	1	..	908.20	41.30	Eligard 3 month [MF]

leuprorelin acetate 30 mg modified release injection [1 chamber] (&) inert substance diluent [1.5 mL chamber], 1 dual chamber syringe

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8877F	1	1	..	1189.13	41.30	Lucrin Depot 4 Month PDS [VE]

leuprorelin acetate 22.5 mg modified release injection [1 chamber] (&) inert substance diluent [1.5 mL chamber], 1 dual chamber syringe

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8876E	1	1	..	908.20	41.30	Lucrin Depot 3 Month PDS [VE]

leuprorelin acetate 7.5 mg modified release injection [1 syringe] (&) inert substance diluent [1 syringe], 1 pack

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8707G	1	5	..	344.56	41.30	Eligard 1 month [MF]

leuprorelin acetate 45 mg modified release injection [1 chamber] (&) inert substance diluent [1.5 mL chamber], 1 dual chamber syringe

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
11943N	1	..	..	1750.50	41.30	Lucrin Depot 6-Month [VE]

▪ **LEUPRORELIN**

Restricted benefit

Central precocious puberty

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a medical practitioner in consultation with a paediatric endocrinologist; OR
- Must be treated by a medical practitioner in consultation with an endocrinologist specialising in paediatrics.

Clinical criteria:

- Patient must have previously been issued with an authority prescription for this drug for this condition.

leuprorelin acetate 30 mg modified release injection [1 chamber] (&) inert substance diluent [1.5 mL chamber], 1 dual chamber syringe

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
11944P	1	1	..	1189.13	41.30	Lucrin Depot Paediatric 30 mg PDS [VE]

▪ **LEUPRORELIN**

Restricted benefit

Central precocious puberty

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a paediatric endocrinologist; OR
- Must be treated by an endocrinologist specialising in paediatrics.

Population criteria:

- Patient must be aged 10 years or younger (girls) or 11 years or younger (boys), **AND**
- Patient must have had onset of signs or symptoms of central precocious puberty prior to the age of 8 years (girls) or 9 years (boys).

Restricted benefit

Central precocious puberty

Treatment Phase: Initial - grandfather

Clinical criteria:

- Patient must have received treatment with a gonadotropin releasing hormone analogue (GnRHa) for this condition prior to 1 May 2015.

Treatment criteria:

- Must be treated by a paediatric endocrinologist; OR
- Must be treated by an endocrinologist specialising in paediatrics.

leuprorelin acetate 30 mg modified release injection [1 chamber] (&) inert substance diluent [1.5 mL chamber], 1 dual chamber syringe

11960L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	1189.13	41.30	Lucrin Depot Paediatric 30 mg PDS [VE]

▪ **LEUPRORELIN (&) BICALUTAMIDE**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Restricted benefit

Carcinoma of the prostate

Clinical criteria:

- The condition must be metastatic (stage D), **AND**
- Patient must require a combination of an antiandrogen and a GnRH (LH-RH) agonist.

leuprorelin acetate 7.5 mg modified release injection [1 syringe] (&) inert substance diluent [1 syringe] (&) bicalutamide 50 mg tablet [28], 1 pack

10962Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	5	..	423.95	41.30	Bi ELIGARD CP [MF]

leuprorelin acetate 22.5 mg modified release injection [1 syringe] (&) inert substance diluent [1 syringe] (&) bicalutamide 50 mg tablet [28], 1 pack

10963B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	982.04	41.30	Bi ELIGARD CP [MF]

leuprorelin acetate 22.5 mg modified release injection [1 syringe] (&) inert substance diluent [1 syringe] (&) bicalutamide 50 mg tablet [84], 1 pack

10969H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	1	..	1129.71	41.30	Bi ELIGARD CP [MF]

▪ **TRIPTORELIN**

Restricted benefit

Locally advanced (stage C) or metastatic (stage D) carcinoma of the prostate

triptorelin 11.25 mg injection [1 vial] (&) inert substance diluent [2 mL ampoule], 1 pack

9379P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	952.64	41.30	Diphereline [IS]

triptorelin 3.75 mg injection [1 vial] (&) inert substance diluent [2 mL ampoule], 1 pack

9378N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	362.34	41.30	Diphereline [IS]

triptorelin 22.5 mg injection [1 vial] (&) inert substance diluent [2 mL ampoule], 1 pack

5297T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	1839.28	41.30	Diphereline [IS]

HORMONE ANTAGONISTS AND RELATED AGENTS

Anti-estrogens

FULVESTRANT

Authority required (STREAMLINED)

11473

Locally advanced or metastatic breast cancer

Clinical criteria:

- The condition must be hormone receptor positive, **AND**
- The condition must be human epidermal growth factor receptor 2 (HER2) negative, **AND**
- The condition must be inoperable.

Population criteria:

- Patient must not be premenopausal.

A patient who has progressive disease when treated with this drug is no longer eligible for PBS-subsidised treatment with this drug.

fulvestrant 250 mg/5 mL injection, 2 x 5 mL syringes

12300J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	345.71	41.30	Fulvestrant Sandoz [SZ]

TAMOXIFEN

Note For item codes 2110C and 1880Y, pharmaceutical benefits that have the form tablet 20 mg (base) are equivalent for the purposes of substitution.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Breast cancer

Clinical criteria:

- The condition must be hormone receptor positive.

tamoxifen 20 mg tablet, 30

1880Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*27.54	28.83	^a Nolvadex-D [AP]

TAMOXIFEN

Note This pharmaceutical benefit is not PBS-subsidised for primary prevention of breast cancer.

Note For item codes 2110C and 1880Y, pharmaceutical benefits that have the form tablet 20 mg (base) are equivalent for the purposes of substitution.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Breast cancer

Clinical criteria:

- The condition must be hormone receptor positive.

tamoxifen 20 mg tablet, 60

2110C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	27.55	28.84	^a Genox 20 [AF] ^a Tamosin [AS]	^a GenRx Tamoxifen [GX] ^a Tamoxifen Sandoz [SZ]

TAMOXIFEN

Note A moderate risk of developing breast cancer is if the lifetime breast cancer risk is 1.5 to 3 times the population average. A high risk of developing breast cancer is if the lifetime breast cancer risk is more than 3 times the population average.

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Restricted benefit

Reduction of breast cancer risk

Clinical criteria:

- Patient must have a moderate or high risk of developing breast cancer, **AND**
- The treatment must not exceed a dose of 20 mg per day, **AND**
- The treatment must not exceed a lifetime maximum of 5 years for this condition.

tamoxifen 20 mg tablet, 30

10911G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	19.78	21.07	^a Genox 20 [AF]	^a Nolvadex-D [AP]

TOREMIFENE
toremifene 60 mg tablet, 30

8216K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	61.73	41.30	Fareston [AS]	

Anti-androgens
BICALUTAMIDE

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)

5729

Metastatic (stage D) carcinoma of the prostate

Clinical criteria:

- The treatment must be in combination with GnRH (LH-RH) analogue therapy.

bicalutamide 50 mg tablet, 28

8094B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	57.82	41.30	^a APO-Bicalutamide [TX] ^a Calutex [AS] ^a Cosudex [AP]	^a Bicalox [ER] ^a Cosamide 50 [AF]

CYPROTERONE
cyproterone acetate 100 mg tablet, 50

8019C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	67.80	41.30	^a ANTERONE 100 [RW] ^a Cyprone 100 [AF] ^a Cyproterone AN [EA] ^a GenRx Cyproterone Acetate [GX]	^a APO-Cyproterone [TX] ^a Cyprostat-100 [SY] ^a Cyproterone Sandoz [HX] ^a Pharmacor Cyproterone 100 [CR]
			^B 1.41	69.21	41.30	^a Androcur-100 [BN]	

cyproterone acetate 50 mg tablet, 50

1270W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	5	..	*83.62	41.30	^a ANTERONE 50 [RW] ^a Cyprone 50 [AL] ^a Cyproterone Sandoz [HX] ^a Pharmacor Cyproterone 50 [CR]	^a APO-Cyproterone [TX] ^a Cyproterone AN [EA] ^a GenRx Cyproterone Acetate [GX]
			^B 2.28	*85.90	41.30	^a Androcur [BN]	

ENZALUTAMIDE

Note Special Pricing Arrangements apply.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Castration resistant metastatic carcinoma of the prostate

Clinical criteria:

- The treatment must not be used in combination with chemotherapy, **AND**
- Patient must have failed treatment with docetaxel due to resistance or intolerance; OR
- Patient must be unsuitable for docetaxel treatment on the basis of predicted intolerance to docetaxel, **AND**
- Patient must have a WHO performance status of 2 or less, **AND**
- Patient must not receive PBS-subsidised treatment with this drug if progressive disease develops while on this drug, **AND**
- Patient must not have received prior treatment with abiraterone; OR
- Patient must have developed intolerance to abiraterone of a severity necessitating permanent treatment withdrawal.

enzalutamide 40 mg capsule, 112

10174L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	3536.80	41.30	Xtandi [LL]

FLUTAMIDE

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)

5816

Metastatic (stage D) carcinoma of the prostate

Clinical criteria:

- The treatment must be in combination with GnRH (LH-RH) analogue therapy.

flutamide 250 mg tablet, 100

1417N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	174.26	41.30	Flutamin [AF]

NILUTAMIDE

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)

5785

Locally advanced (stage C) or metastatic (stage D) carcinoma of the prostate

Clinical criteria:

- The treatment must be in combination with GnRH (LH-RH) analogue therapy.

Authority required (STREAMLINED)

5647

Locally advanced (stage C) or metastatic (stage D) carcinoma of the prostate

Clinical criteria:

- The treatment must be in conjunction with surgical orchidectomy.

nilutamide 150 mg tablet, 30

8131Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	187.95	41.30	Anandron [SW]

Aromatase inhibitors

ANASTROZOLE

Note This drug is not PBS-subsidised for primary prevention of breast cancer.

Note This drug is not PBS-subsidised for adjuvant hormonal treatment of early breast cancer where the total duration of this drug (or any other aromatase inhibitor) treatment extends beyond 5 years.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Breast cancer

Clinical criteria:

- The condition must be hormone receptor positive.

anastrozole 1 mg tablet, 30

8179L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	24.31	25.60	^a Anastro [AS] ^a Anastrozole GH [GQ] ^a APO-Anastrozole [TX] ^a Arimidex [AP]	^a Anastrozole FBM [FO] ^a Anastrozole Sandoz [SZ] ^a Arianna 1 [AF]

EXEMESTANE

Restricted benefit

Metastatic (Stage IV) breast cancer

Clinical criteria:

- The condition must be hormone receptor positive, **AND**
- The condition must be human epidermal growth factor receptor 2 (HER2) negative, **AND**

- Patient must be receiving PBS-subsidised everolimus concomitantly for this condition.

Population criteria:

- Patient must not be pre-menopausal.

exemestane 25 mg tablet, 30

10103R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	67.52	41.30	^a APO-Exemestane [TX]	^a Exemestane GH [GQ]
						^a Exemestane Sandoz [SZ]	
			^B 4.00	71.52	41.30	^a Aromasin [PF]	

■ **EXEMESTANE**

Note This drug is not PBS-subsidised for primary prevention of breast cancer.

Note This drug is not PBS-subsidised for adjuvant hormonal treatment of early breast cancer extended beyond 5 years, i.e. a patient who has received 2 years of tamoxifen therapy may only receive 3 years of PBS-subsidised treatment with exemestane.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Breast cancer

Clinical criteria:

- The condition must be hormone receptor positive.

exemestane 25 mg tablet, 30

8506Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	67.52	41.30	^a APO-Exemestane [TX]	^a Exemestane GH [GQ]
						^a Exemestane Sandoz [SZ]	
			^B 4.00	71.52	41.30	^a Aromasin [PF]	

■ **LETROZOLE**

Note This drug is not PBS-subsidised for primary prevention of breast cancer.

Note This drug is not PBS-subsidised for adjuvant hormonal treatment of early breast cancer where the total duration of this drug (or any other aromatase inhibitor) treatment extends beyond 5 years.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Breast cancer

Clinical criteria:

- The condition must be hormone receptor positive.

letrozole 2.5 mg tablet, 30

8245Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	31.81	33.10	^a APO-Letrozole [TX]	^a Femara 2.5 mg [NV]
						^a Femolet [AF]	^a Fera [AS]
						^a Gynotril [ER]	^a Letrozole APOTEX [GX]
						^a Letrozole FBM [FO]	^a Letrozole GH [HQ]
						^a Letrozole Sandoz [SZ]	^a Pharmacor Letrozole 2.5 [CR]

Other hormone antagonists and related agents■ **ABIRATERONE**

Note Special Pricing Arrangements apply.

Authority required

Castration resistant metastatic carcinoma of the prostate

Clinical criteria:

- The treatment must be used in combination with a corticosteroid, **AND**
- The treatment must not be used in combination with chemotherapy, **AND**
- Patient must have failed treatment with docetaxel due to resistance or intolerance; OR
- Patient must be unsuitable for docetaxel treatment on the basis of predicted intolerance to docetaxel, **AND**
- Patient must have a WHO performance status of 2 or less, **AND**
- Patient must not receive PBS-subsidised abiraterone if progressive disease develops while on abiraterone, **AND**
- Patient must not have received prior treatment with enzalutamide; OR
- Patient must have developed intolerance to enzalutamide of a severity necessitating permanent treatment withdrawal.

abiraterone acetate 250 mg tablet, 120

2698B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	3442.02	41.30	Zytiga [JC]

abiraterone acetate 500 mg tablet, 60

11206T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	3442.02	41.30	Zytiga [JC]

■ DEGARELIX**Restricted benefit**

Locally advanced (equivalent to stage C) or metastatic (equivalent to stage D) carcinoma of the prostate

degarelix 80 mg injection [1 vial] (&) inert substance diluent [1 syringe], 1 pack

2784M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	362.34	41.30	Firmagon 80mg [FP]

■ DEGARELIX

Note No applications for increased maximum quantities and/or repeats will be authorised for the 120 mg powder for injection.

Restricted benefit

Locally advanced (equivalent to stage C) or metastatic (equivalent to stage D) carcinoma of the prostate

degarelix 120 mg injection [2 vials] (&) inert substance diluent [2 syringes], 1 pack

2785N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	378.95	41.30	Firmagon 120mg [FP]

■ IMMUNOSTIMULANTS**IMMUNOSTIMULANTS***Interferons***■ INTERFERON ALFA-2A**

Caution Treatment with interferon alfa has been associated with depression and suicide in some patients. Patients with a history of suicidal ideation or depressive illness should be warned of the risks. Psychiatric status during therapy should be monitored.

Authority required

Myeloproliferative disease

Clinical criteria:

- Patient must have excessive thrombocytosis.

interferon alfa-2a 9 million units (33.333 microgram)/0.5 mL injection, 0.5 mL syringe

8553E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	5	4	..	*416.84	41.30	Roferon-A [RO]

■ INTERFERON ALFA-2A

Caution Treatment with interferon alfa has been associated with depression and suicide in some patients. Patients with a history of suicidal ideation or depressive illness should be warned of the risks. Psychiatric status during therapy should be monitored.

Authority required

Low grade non-Hodgkin's lymphoma

Clinical criteria:

- The condition must have clinical features suggestive of a poor prognosis, **AND**
- The treatment must be in combination with anthracycline-based chemotherapy.

interferon alfa-2a 9 million units (33.333 microgram)/0.5 mL injection, 0.5 mL syringe

8184R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	5	5	..	*416.84	41.30	Roferon-A [RO]

interferon alfa-2a 3 million units (11.111 microgram)/0.5 mL injection, 0.5 mL syringe

8181N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	15	5	..	*416.94	41.30	Roferon-A [RO]

■ INTERFERON ALFA-2A

Caution Treatment with interferon alfa has been associated with depression and suicide in some patients. Patients with a history of suicidal ideation or depressive illness should be warned of the risks. Psychiatric status during therapy should be monitored.

Authority required

Hairy cell leukaemia

Authority required

Myeloproliferative disease

Clinical criteria:

- Patient must have excessive thrombocytosis.

interferon alfa-2a 3 million units (11.111 microgram)/0.5 mL injection, 0.5 mL syringe

8180M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	15	4	..	*416.94	41.30	Roferon-A [RO]

INTERFERON BETA-1A

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

7695

Multiple sclerosis

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be diagnosed as clinically definite relapsing-remitting multiple sclerosis by magnetic resonance imaging of the brain and/or spinal cord; OR
- The condition must be diagnosed as clinically definite relapsing-remitting multiple sclerosis, with written certification provided by a radiologist that a magnetic resonance imaging scan is contraindicated because of the risk of physical (not psychological) injury to the patient, **AND**
- Patient must have experienced at least 2 documented attacks of neurological dysfunction, believed to be due to multiple sclerosis, in the preceding 2 years of commencing a PBS-subsidised disease modifying therapy for this condition, **AND**
- Patient must be ambulatory (without assistance or support).

Where applicable, the date of the magnetic resonance imaging scan must be recorded in the patient's medical records.

Authority required (STREAMLINED)

6860

Multiple sclerosis

Treatment Phase: Continuing treatment

Clinical criteria:

- The condition must be diagnosed as clinically definite relapsing-remitting multiple sclerosis, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not show continuing progression of disability while on treatment with this drug, **AND**
- Patient must have demonstrated compliance with, and an ability to tolerate this therapy.

interferon beta-1a 6 million units (30 microgram)/0.5 mL injection, 4 x 0.5 mL syringes

8805K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	865.56	41.30	Avonex [BD]

interferon beta-1a 12 million units (44 microgram)/0.5 mL injection, 12 x 0.5 mL pen devices

8968B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	865.56	41.30	Rebif 44 [SG]

interferon beta-1a 12 million units (44 microgram)/0.5 mL injection, 12 x 0.5 mL syringes

8403G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	865.56	41.30	Rebif 44 [SG]

interferon beta-1a 36 million units (132 microgram)/1.5 mL injection, 4 x 1.5 mL cartridges

9332E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	865.56	41.30	Rebif 44 [SG]

INTERFERON BETA-1B

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

7695

Multiple sclerosis

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be diagnosed as clinically definite relapsing-remitting multiple sclerosis by magnetic resonance imaging of the brain and/or spinal cord; OR
- The condition must be diagnosed as clinically definite relapsing-remitting multiple sclerosis, with written certification provided by a radiologist that a magnetic resonance imaging scan is contraindicated because of the risk of physical (not psychological) injury to the patient, **AND**
- Patient must have experienced at least 2 documented attacks of neurological dysfunction, believed to be due to multiple sclerosis, in the preceding 2 years of commencing a PBS-subsidised disease modifying therapy for this condition, **AND**
- Patient must be ambulatory (without assistance or support).

Where applicable, the date of the magnetic resonance imaging scan must be recorded in the patient's medical records.

Authority required (STREAMLINED)

6860

Multiple sclerosis

Treatment Phase: Continuing treatment

Clinical criteria:

- The condition must be diagnosed as clinically definite relapsing-remitting multiple sclerosis, **AND**

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not show continuing progression of disability while on treatment with this drug, **AND**
- Patient must have demonstrated compliance with, and an ability to tolerate this therapy.

interferon beta-1b 8 million units (250 microgram) injection [15 vials] (&) inert substance diluent [15 x 1.2 mL syringes], 1 pack

8101J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	997.60	41.30	Betaferon [BN]

▪ **PEGINTERFERON ALFA-2A**

peginterferon alfa-2a 135 microgram/0.5 mL injection, 4 x 0.5 mL syringes

11416W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	600.98	41.30	Pegasys [RO]

peginterferon alfa-2a 180 microgram/0.5 mL injection, 4 x 0.5 mL syringes

11037X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	694.89	41.30	Pegasys [RO]

▪ **PEGINTERFERON BETA-1A**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

7695

Multiple sclerosis

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be diagnosed as clinically definite relapsing-remitting multiple sclerosis by magnetic resonance imaging of the brain and/or spinal cord; OR
- The condition must be diagnosed as clinically definite relapsing-remitting multiple sclerosis, with written certification provided by a radiologist that a magnetic resonance imaging scan is contraindicated because of the risk of physical (not psychological) injury to the patient, **AND**
- Patient must have experienced at least 2 documented attacks of neurological dysfunction, believed to be due to multiple sclerosis, in the preceding 2 years of commencing a PBS-subsidised disease modifying therapy for this condition, **AND**
- Patient must be ambulatory (without assistance or support).

Where applicable, the date of the magnetic resonance imaging scan must be recorded in the patient's medical records.

peginterferon beta-1a 63 microgram/0.5 mL injection [0.5 mL pen device] (&) peginterferon beta-1a 94 microgram/0.5 mL injection [0.5 mL pen device], 1 pack

10218T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	1001.53	41.30	Plegridy [BD]

peginterferon beta-1a 125 microgram/0.5 mL injection, 2 x 0.5 mL pen devices

10212L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	4	..	1001.53	41.30	Plegridy [BD]

▪ **PEGINTERFERON BETA-1A**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

6860

Multiple sclerosis

Treatment Phase: Continuing treatment

Clinical criteria:

- The condition must be diagnosed as clinically definite relapsing-remitting multiple sclerosis, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not show continuing progression of disability while on treatment with this drug, **AND**
- Patient must have demonstrated compliance with, and an ability to tolerate this therapy.

peginterferon beta-1a 125 microgram/0.5 mL injection, 2 x 0.5 mL pen devices

10220X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1001.53	41.30	Plegridy [BD]

Other immunostimulants

▪ **GLATIRAMER ACETATE**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)
7695

Multiple sclerosis

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be diagnosed as clinically definite relapsing-remitting multiple sclerosis by magnetic resonance imaging of the brain and/or spinal cord; OR
- The condition must be diagnosed as clinically definite relapsing-remitting multiple sclerosis, with written certification provided by a radiologist that a magnetic resonance imaging scan is contraindicated because of the risk of physical (not psychological) injury to the patient, **AND**
- Patient must have experienced at least 2 documented attacks of neurological dysfunction, believed to be due to multiple sclerosis, in the preceding 2 years of commencing a PBS-subsidised disease modifying therapy for this condition, **AND**
- Patient must be ambulatory (without assistance or support).

Where applicable, the date of the magnetic resonance imaging scan must be recorded in the patient's medical records.

Authority required (STREAMLINED)
6860

Multiple sclerosis

Treatment Phase: Continuing treatment

Clinical criteria:

- The condition must be diagnosed as clinically definite relapsing-remitting multiple sclerosis, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not show continuing progression of disability while on treatment with this drug, **AND**
- Patient must have demonstrated compliance with, and an ability to tolerate this therapy.

glatiramer acetate 40 mg/mL injection, 12 x 1 mL syringes

10416F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	894.98	41.30	Copaxone [TB]

■ MYCOBACTERIUM BOVIS (BACILLUS CALMETTE AND GUERIN (BCG)) TICE STRAIN
Restricted benefit

Primary and relapsing superficial urothelial carcinoma of the bladder

Mycobacterium bovis (Bacillus Calmette and Guerin (BCG)) Tice strain 500 million CFU injection, 3 vials

1131M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	458.03	41.30	OncotICE [MK]

■ IMMUNOSUPPRESSANTS
IMMUNOSUPPRESSANTS
Selective immunosuppressants
■ ABATACEPT
Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for adults with severe active rheumatoid arthritis. Where the term biological medicine appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alfa antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab), the T-cell co-stimulation modulator (abatacept) and the Janus kinase (JAK) inhibitors (baricitinib, tofacitinib, upadacitinib).

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised biological medicine therapy may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy,

- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once, and

- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised biological medicines for the treatment of rheumatoid arthritis.

A serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Stevens Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered a treatment failure.

A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction.

A patient who has failed fewer than 5 biological medicines and who has a break in treatment of less than 24 months may commence a further course of treatment with a biological medicine under Initial 2 treatment restriction. A patient who has failed fewer than 5 biological medicines and who has had a break in therapy of longer than 24 months may commence a further course of treatment with a biological medicine under the Initial 3 treatment restriction. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine treatment is stopped to the date of the new application for treatment with a biological medicine.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 April 2019.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised biological medicine treatment and wishes to commence such therapy, excluding rituximab (Initial 1 - new patient); or

(ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to re-commence treatment with a specific biological medicine following a break of less than 24 months in PBS-subsidised therapy with that agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months).

(iv) a patient wishes to re-commence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 24 months (Initial 3 - re-commencement of treatment after a break in biological medicine of more than 24 months) Initial applications for a new patient (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6-month intensive DMARD trial, but prior to ceasing DMARD therapy.

Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, baricitinib, etanercept, golimumab, tocilizumab, tofacitinib and upadacitinib, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

Rituximab patients should be assessed following a minimum of 12 weeks after the first infusion, and the assessment should be submitted to Services Australia within 4 weeks to ensure continuity of treatment for those who meet the continuing restriction for PBS subsidised treatment with this drug for this condition.

Where a response assessment is not provided the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment. For second and subsequent courses of PBS-subsidised biological medicine (excluding rituximab) treatment, it is recommended that a patient be reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to Services Australia within sufficient time to allow processing.

Abatacept patients:

A patient is eligible to receive one I.V. loading dose when commencing treatment with the subcutaneous formulation. Two prescriptions are required, the first prescription for the I.V. loading dose for sufficient vials for one dose based on the patient's weight with no repeats. The second prescription for the subcutaneous formulation, with a maximum quantity of 4 and up to 3 repeats, must be submitted with the initial application.

Rituximab patients:

A further application may be submitted to Services Australia within sufficient time to allow processing. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

Assessment of the patient's response to treatment must be conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the cessation of the most recent course of biological medicine therapy.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction. Where a response assessment is not conducted within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

(2) Swapping therapy

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non- biological medicine therapy requirements except if the patient has had a break in therapy of more than 24 months who would need to requalify with respect to the indices of disease severity. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each biological medicine trialled.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent, unless they have experienced a serious adverse reaction of a severity necessitating permanent treatment withdrawal.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug.

Abatacept:

A patient swapping from I.V. abatacept to subcutaneous abatacept will not be eligible for an I.V. loading dose when commencing treatment with the subcutaneous formulation.

Rituximab:

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF-alfa antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they be assessed for response to every course of treatment, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological medicine during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological medicine therapy treatment-free period of at least 22 weeks, immediately following the second infusion,

before swapping to an alternate biological medicine. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate biological medicine may do so without having to have any treatment-free period.

(3) Baseline measurements to determine response.

Determination of whether a response to treatment has been demonstrated must be based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and the eligibility for continuing treatment must be assessed according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response must be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be determined on the total number of major joints.

Applications under the Initial 1 treatment restriction for a new patient must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. The results must be no more than 4 weeks old at the time of application.

Applications under the Initial 3 treatment restriction for re-commencement of treatment after a break in biological medicine of more than 24 months must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction, **AND**
- The treatment must be given concomitantly with methotrexate at a dose of at least 7.5 mg weekly.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient has either failed or ceased to respond to a PBS-subsidised biological medicine for this condition 5 times, they will not be eligible to receive further PBS-subsidised treatment with a biological medicine for this condition.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Continuing Treatment - balance of supply.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restriction.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

abatacept 125 mg/mL injection, 4 x 1 mL syringes

1221G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	958.76	41.30	Orencia [BQ]

abatacept 125 mg/mL injection, 4 x 1 mL pen devices

11684Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	958.76	41.30	Orencia ClickJect [BQ]

ABATACEPT

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for adults with severe active rheumatoid arthritis. Where the term biological medicine appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alfa antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab), the T-cell co-stimulation modulator (abatacept) and the Janus kinase (JAK) inhibitors (baricitinib, tofacitinib, upadacitinib).

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised biological medicine therapy may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once, and
- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised biological medicines for the treatment of rheumatoid arthritis.

A serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Stevens Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered a treatment failure.

A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction.

A patient who has failed fewer than 5 biological medicines and who has a break in treatment of less than 24 months may commence a further course of treatment with a biological medicine under Initial 2 treatment restriction. A patient who has failed fewer than 5 biological medicines and who has had a break in therapy of longer than 24 months may commence a further course of treatment with a biological medicine under the Initial 3 treatment restriction. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine treatment is stopped to the date of the new application for treatment with a biological medicine.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 April 2019.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised biological medicine treatment and wishes to commence such therapy, excluding rituximab (Initial 1 - new patient); or
 - (ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months) [further details are under 'Swapping therapy' below]; or
 - (iii) a patient wishes to re-commence treatment with a specific biological medicine following a break of less than 24 months in PBS-subsidised therapy with that agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months).
 - (iv) a patient wishes to re-commence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 24 months (Initial 3 - re-commencement of treatment after a break in biological medicine of more than 24 months) Initial applications for a new patient (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6-month intensive DMARD trial, but prior to ceasing DMARD therapy.
- Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, baricitinib, etanercept, golimumab, tocilizumab, tofacitinib and upadacitinib, 18 to 20 weeks of therapy with certolizumab

pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

Rituximab patients should be assessed following a minimum of 12 weeks after the first infusion, and the assessment should be submitted to Services Australia within 4 weeks to ensure continuity of treatment for those who meet the continuing restriction for PBS subsidised treatment with this drug for this condition.

Where a response assessment is not provided the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment. For second and subsequent courses of PBS-subsidised biological medicine (excluding rituximab) treatment, it is recommended that a patient be reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to Services Australia within sufficient time to allow processing.

Abatacept patients:

A patient is eligible to receive one I.V. loading dose when commencing treatment with the subcutaneous formulation. Two prescriptions are required, the first prescription for the I.V. loading dose for sufficient vials for one dose based on the patient's weight with no repeats. The second prescription for the subcutaneous formulation, with a maximum quantity of 4 and up to 3 repeats, must be submitted with the initial application.

Rituximab patients:

A further application may be submitted to Services Australia within sufficient time to allow processing. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

Assessment of the patient's response to treatment must be conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the cessation of the most recent course of biological medicine therapy.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction. Where a response assessment is not conducted within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

(2) Swapping therapy

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non- biological medicine therapy requirements except if the patient has had a break in therapy of more than 24 months who would need to requalify with respect to the indices of disease severity. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each biological medicine trialled.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent, unless they have experienced a serious adverse reaction of a severity necessitating permanent treatment withdrawal.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug.

Abatacept:

A patient swapping from I.V. abatacept to subcutaneous abatacept will not be eligible for an I.V. loading dose when commencing treatment with the subcutaneous formulation.

Rituximab:

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF-alfa antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they be assessed for response to every course of treatment, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological medicine during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological medicine therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate biological medicine may do so without having to have any treatment-free period.

(3) Baseline measurements to determine response.

Determination of whether a response to treatment has been demonstrated must be based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and the eligibility for continuing treatment must be assessed according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response must be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be determined on the total number of major joints.

Applications under the Initial 1 treatment restriction for a new patient must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. The results must be no more than 4 weeks old at the time of application.

Applications under the Initial 3 treatment restriction for re-commencement of treatment after a break in biological medicine of

more than 24 months must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with disease modifying anti-rheumatic drugs (DMARDs) which must include at least 3 months continuous treatment with each of at least 2 DMARDs, one of which must be methotrexate at a dose of at least 20 mg weekly and one of which must be: (i) hydroxychloroquine at a dose of at least 200 mg daily; or (ii) leflunomide at a dose of at least 10 mg daily; or (iii) sulfasalazine at a dose of at least 2 g daily; OR
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs which, if methotrexate is contraindicated according to the Therapeutic Goods Administration (TGA)-approved Product Information or cannot be tolerated at a 20 mg weekly dose, must include at least 3 months continuous treatment with each of at least 2 of the following DMARDs: (i) hydroxychloroquine at a dose of at least 200 mg daily; and/or (ii) leflunomide at a dose of at least 10 mg daily; and/or (iii) sulfasalazine at a dose of at least 2 g daily; OR
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs which, if 3 or more of methotrexate, hydroxychloroquine, leflunomide and sulfasalazine are contraindicated according to the relevant TGA-approved Product Information or cannot be tolerated at the doses specified above, must include at least 3 months continuous treatment with each of at least 2 DMARDs, with one or more of the following DMARDs being used in place of the DMARDs which are contraindicated or not tolerated: (i) azathioprine at a dose of at least 1 mg/kg per day; and/or (ii) cyclosporin at a dose of at least 2 mg/kg/day; and/or (iii) sodium aurothiomalate at a dose of 50 mg weekly, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction, **AND**
- The treatment must be given concomitantly with methotrexate at a dose of at least 7.5 mg weekly.

Population criteria:

- Patient must be aged 18 years or older.

If methotrexate is contraindicated according to the TGA-approved product information or cannot be tolerated at a 20 mg weekly dose, the application must include details of the contraindication or intolerance including severity to methotrexate. The maximum tolerated dose of methotrexate must be documented in the application, if applicable.

The application must include details of the DMARDs trialled, their doses and duration of treatment, and all relevant contraindications and/or intolerances including severity.

The requirement to trial at least 2 DMARDs for periods of at least 3 months each can be met using single agents sequentially or by using one or more combinations of DMARDs.

If the requirement to trial 6 months of intensive DMARD therapy with at least 2 DMARDs cannot be met because of contraindications and/or intolerances of a severity necessitating permanent treatment withdrawal to all of the DMARDs specified above, details of the contraindication or intolerance including severity and dose for each DMARD must be provided in the authority application.

The following criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; **AND** either

(a) a total active joint count of at least 20 active (swollen and tender) joints; or

(b) at least 4 active joints from the following list of major joints:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The joint count and ESR and/or CRP must be determined at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. All measures must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

At the time of authority application, medical practitioners should request the appropriate number of vials to provide sufficient drug, based on the weight of the patient, for a single infusion.

Up to a maximum of 4 repeats will be authorised.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

Initial treatment with an I.V. loading dose: Two completed authority prescriptions must be submitted with the initial application. One prescription must be for the I.V. loading dose for sufficient vials for one dose based on the patient's weight

with no repeats. The second prescription must be written for the subcutaneous formulation, with a maximum quantity of 4 and up to 3 repeats.

Initial treatment with no loading dose: One completed authority prescription must be submitted with the initial application. The prescription must be written with a maximum quantity of 4 and up to 3 repeats.

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note The Department of Human Services website (www.humanservices.gov.au) has details of the toxicities, including severity, which will be accepted for the following purposes:

- (a) exempting a patient from the requirement to undertake a minimum 3 month trial of methotrexate at a 20 mg weekly dose;
- (b) substituting azathioprine, cyclosporin or sodium aurothiomalate for another DMARD as part of the 6 month intensive DMARD trial;
- (c) exempting a patient from the requirement for a 6 month trial of intensive DMARD therapy.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 2 (change or re-commencement of treatment after a break in biological medicine of less than 24 months).

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must not have failed to respond to previous PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised biological medicine treatment for this condition 5 times, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction, **AND**
- The treatment must be given concomitantly with methotrexate at a dose of at least 7.5 mg weekly.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either of the Initial 1, Initial 2, Initial 3, or continuing treatment restrictions, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

Initial treatment with an I.V. loading dose: Two completed authority prescriptions must be submitted with the initial application. One prescription must be for the I.V. loading dose for sufficient vials for one dose based on the patient's weight with no repeats. The second prescription must be written for the subcutaneous formulation, with a maximum quantity of 4 and up to 3 repeats.

Initial treatment with no loading dose: One completed authority prescription must be submitted with the initial application. The prescription must be written with a maximum quantity of 4 and up to 3 repeats.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

A patient who has demonstrated a response to a course of rituximab must have a PBS-subsidised biological therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 3 (re-commencement of treatment after a break in biological medicine of more than 24 months)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 24 months or more from the most recent PBS-subsidised biological medicine for this condition, **AND**
- Patient must not have failed to respond to previous PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised biological medicine treatment for this condition 5 times, **AND**
- The condition must have an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; OR
- The condition must have a C-reactive protein (CRP) level greater than 15 mg per L, **AND**
- The condition must have either (a) a total active joint count of at least 20 active (swollen and tender) joints; or (b) at least 4 active major joints, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction, **AND**
- The treatment must be given concomitantly with methotrexate at a dose of at least 7.5 mg weekly.

Population criteria:

- Patient must be aged 18 years or older.

Major joints are defined as (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

All measures of joint count and ESR and/or CRP must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and

(2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

Initial treatment with an I.V. loading dose: Two completed authority prescriptions must be submitted with the initial application. One prescription must be for the I.V. loading dose for sufficient vials for one dose based on the patient's weight with no repeats. The second prescription must be written for the subcutaneous formulation, with a maximum quantity of 4 and up to 3 repeats.

Initial treatment with no loading dose: One completed authority prescription must be submitted with the initial application. The prescription must be written with a maximum quantity of 4 and up to 3 repeats.

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial 1 (new patient) or Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 24 months) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 24 months) - balance of supply

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 24 months) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 24 months) to complete 16 weeks of treatment, **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

abatacept 125 mg/mL injection, 4 x 1 mL syringes

1220F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	958.76	41.30	Orencia [BQ]

abatacept 125 mg/mL injection, 4 x 1 mL pen devices

11693K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	958.76	41.30	Orencia ClickJect [BQ]

■ APREMILAST

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

11115

Severe chronic plaque psoriasis

Clinical criteria:

- Patient must have failed to achieve an adequate response after at least 6 weeks of treatment with methotrexate prior to initiating treatment with this drug; OR

- Patient must have a contraindication to methotrexate according to the Therapeutic Goods Administration (TGA) approved Product Information; OR
- Patient must have demonstrated severe intolerance of, or toxicity due to, methotrexate, **AND**
- The condition must have caused significant interference with quality of life, **AND**
- Patient must not be undergoing concurrent PBS-subsidised treatment for psoriasis with each of: (i) a biological medicine, (ii) ciclosporin.

Treatment criteria:

- Must be treated by a dermatologist.

Population criteria:

- Patient must be aged 18 years or older.

apremilast 30 mg tablet, 56

12223H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	652.74	41.30	Otezla [AN]

apremilast 10 mg tablet [4] (&) apremilast 20 mg tablet [4] (&) apremilast 30 mg tablet [19], 27

12218C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	272.23	41.30	Otezla [AN]

■ BARICITINIB
Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for adults with severe active rheumatoid arthritis. Where the term biological medicine appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alfa antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab), the T-cell co-stimulation modulator (abatacept) and the Janus kinase (JAK) inhibitors (baricitinib, tofacitinib, upadacitinib).

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised biological medicine therapy may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once, and
- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised biological medicines for the treatment of rheumatoid arthritis.

A serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Stevens Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered a treatment failure.

A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction.

A patient who has failed fewer than 5 biological medicines and who has a break in treatment of less than 24 months may commence a further course of treatment with a biological medicine under Initial 2 treatment restriction. A patient who has failed fewer than 5 biological medicines and who has had a break in therapy of longer than 24 months may commence a further course of treatment with a biological medicine under the Initial 3 treatment restriction. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine treatment is stopped to the date of the new application for treatment with a biological medicine.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 April 2019.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised biological medicine treatment and wishes to commence such therapy, excluding rituximab (Initial 1 - new patient); or

(ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to re-commence treatment with a specific biological medicine following a break of less than 24 months in PBS-subsidised therapy with that agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months).

(iv) a patient wishes to re-commence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 24 months (Initial 3 - re-commencement of treatment after a break in biological medicine of more than 24 months) Initial applications for a new patient (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6-month intensive DMARD trial, but prior to ceasing DMARD therapy.

Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, baricitinib, etanercept, golimumab, tocilizumab, tofacitinib and upadacitinib, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

Rituximab patients should be assessed following a minimum of 12 weeks after the first infusion, and the assessment should be submitted to Services Australia within 4 weeks to ensure continuity of treatment for those who meet the continuing restriction for PBS subsidised treatment with this drug for this condition.

Where a response assessment is not provided the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for

permanent withdrawal of treatment. For second and subsequent courses of PBS-subsidised biological medicine (excluding rituximab) treatment, it is recommended that a patient be reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to Services Australia within sufficient time to allow processing. Abatacept patients:

A patient is eligible to receive one I.V. loading dose when commencing treatment with the subcutaneous formulation. Two prescriptions are required, the first prescription for the I.V. loading dose for sufficient vials for one dose based on the patient's weight with no repeats. The second prescription for the subcutaneous formulation, with a maximum quantity of 4 and up to 3 repeats, must be submitted with the initial application.

Rituximab patients:

A further application may be submitted to Services Australia within sufficient time to allow processing. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

Assessment of the patient's response to treatment must be conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the cessation of the most recent course of biological medicine therapy.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction. Where a response assessment is not conducted within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

(2) Swapping therapy

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non- biological medicine therapy requirements except if the patient has had a break in therapy of more than 24 months who would need to requalify with respect to the indices of disease severity. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each biological medicine trialled.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent, unless they have experienced a serious adverse reaction of a severity necessitating permanent treatment withdrawal.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug.

Abatacept:

A patient swapping from I.V. abatacept to subcutaneous abatacept will not be eligible for an I.V. loading dose when commencing treatment with the subcutaneous formulation.

Rituximab:

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF-alfa antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they be assessed for response to every course of treatment, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological medicine during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological medicine therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate biological medicine may do so without having to have any treatment-free period.

(3) Baseline measurements to determine response.

Determination of whether a response to treatment has been demonstrated must be based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and the eligibility for continuing treatment must be assessed according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response must be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be determined on the total number of major joints.

Applications under the Initial 1 treatment restriction for a new patient must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. The results must be no more than 4 weeks old at the time of application.

Applications under the Initial 3 treatment restriction for re-commencement of treatment after a break in biological medicine of more than 24 months must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment per continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient has either failed or ceased to respond to a PBS-subsidised biological medicine for this condition 5 times, they will not be eligible to receive further PBS-subsidised treatment with a biological medicine for this condition.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Continuing treatment - balance of supply

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the continuing treatment restriction to complete 24 weeks of treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restrictions.

Population criteria:

- Patient must be aged 18 years or older.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

baricitinib 4 mg tablet, 28

11443G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1271.34	41.30	Olumiant [LY]

baricitinib 2 mg tablet, 28

11442F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1271.34	41.30	Olumiant [LY]

■ BARICITINIB**Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS**

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for adults with severe active rheumatoid arthritis. Where the term biological medicine appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alfa antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab), the T-cell co-stimulation modulator (abatacept) and the Janus kinase (JAK) inhibitors (baricitinib, tofacitinib, upadacitinib).

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised biological medicine therapy may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once, and
- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised biological medicines for the treatment of rheumatoid arthritis.

A serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Stevens Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered a treatment failure.

A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction.

A patient who has failed fewer than 5 biological medicines and who has a break in treatment of less than 24 months may commence a further course of treatment with a biological medicine under Initial 2 treatment restriction. A patient who has failed fewer than 5 biological medicines and who has had a break in therapy of longer than 24 months may commence a further course of treatment with a biological medicine under the Initial 3 treatment restriction. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine treatment is stopped to the date of the new application for treatment with a biological medicine.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 April 2019.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised biological medicine treatment and wishes to commence such therapy, excluding rituximab (Initial 1 - new patient); or
 - (ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months) [further details are under 'Swapping therapy' below]; or
 - (iii) a patient wishes to re-commence treatment with a specific biological medicine following a break of less than 24 months in PBS-subsidised therapy with that agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months).
 - (iv) a patient wishes to re-commence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 24 months (Initial 3 - re-commencement of treatment after a break in biological medicine of more than 24 months)
- Initial applications for a new patient (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6-month intensive DMARD trial, but prior to ceasing DMARD therapy.

Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, baricitinib, etanercept, golimumab, tocilizumab, tofacitinib and upadacitinib, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

Rituximab patients should be assessed following a minimum of 12 weeks after the first infusion, and the assessment should be submitted to Services Australia within 4 weeks to ensure continuity of treatment for those who meet the continuing restriction for PBS subsidised treatment with this drug for this condition.

Where a response assessment is not provided the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment. For second and subsequent courses of PBS-subsidised biological medicine (excluding rituximab) treatment, it is recommended that a patient be reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to Services Australia within sufficient time to allow processing.

Abatacept patients:

A patient is eligible to receive one I.V. loading dose when commencing treatment with the subcutaneous formulation. Two prescriptions are required, the first prescription for the I.V. loading dose for sufficient vials for one dose based on the patient's weight with no repeats. The second prescription for the subcutaneous formulation, with a maximum quantity of 4 and up to 3 repeats, must be submitted with the initial application.

Rituximab patients:

A further application may be submitted to Services Australia within sufficient time to allow processing. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

Assessment of the patient's response to treatment must be conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the cessation of the most recent course of biological medicine therapy.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction. Where a response assessment is not conducted within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

(2) Swapping therapy

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non- biological medicine therapy requirements except if the patient has had a break in therapy of more than 24 months who would need to requalify with respect to the indices of disease severity. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each biological medicine trialled.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent, unless they have experienced a serious adverse reaction of a severity necessitating permanent treatment withdrawal.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug.

Abatacept:

A patient swapping from I.V. abatacept to subcutaneous abatacept will not be eligible for an I.V. loading dose when commencing treatment with the subcutaneous formulation.

Rituximab:

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF-alfa antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they be assessed for response to every course of treatment, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological medicine during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological medicine therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate biological medicine may do so without having to have any treatment-free period.

(3) Baseline measurements to determine response.

Determination of whether a response to treatment has been demonstrated must be based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and the eligibility for continuing treatment must be assessed according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response must be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be determined on the total number of major joints.

Applications under the Initial 1 treatment restriction for a new patient must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. The results must be no more than 4 weeks old at the time of application.

Applications under the Initial 3 treatment restriction for re-commencement of treatment after a break in biological medicine of more than 24 months must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with disease modifying anti-rheumatic drugs (DMARDs) which must include at least 3 months continuous treatment with each of at least 2 DMARDs, one of which must be

- methotrexate at a dose of at least 20 mg weekly and one of which must be: (i) hydroxychloroquine at a dose of at least 200 mg daily; or (ii) leflunomide at a dose of at least 10 mg daily; or (iii) sulfasalazine at a dose of at least 2 g daily; OR
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs which, if methotrexate is contraindicated according to the Therapeutic Goods Administration (TGA)-approved Product Information or cannot be tolerated at a 20 mg weekly dose, must include at least 3 months continuous treatment with each of at least 2 of the following DMARDs: (i) hydroxychloroquine at a dose of at least 200 mg daily; and/or (ii) leflunomide at a dose of at least 10 mg daily; and/or (iii) sulfasalazine at a dose of at least 2 g daily; OR
 - Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs which, if 3 or more of methotrexate, hydroxychloroquine, leflunomide and sulfasalazine are contraindicated according to the relevant TGA-approved Product Information or cannot be tolerated at the doses specified above, must include at least 3 months continuous treatment with each of at least 2 DMARDs, with one or more of the following DMARDs being used in place of the DMARDs which are contraindicated or not tolerated: (i) azathioprine at a dose of at least 1 mg/kg per day; and/or (ii) cyclosporin at a dose of at least 2 mg/kg/day; and/or (iii) sodium aurothiomalate at a dose of 50 mg weekly, **AND**
 - Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

If methotrexate is contraindicated according to the TGA-approved product information or cannot be tolerated at a 20 mg weekly dose, the application must include details of the contraindication or intolerance including severity to methotrexate. The maximum tolerated dose of methotrexate must be documented in the application, if applicable.

The application must include details of the DMARDs trialed, their doses and duration of treatment, and all relevant contraindications and/or intolerances including severity.

The requirement to trial at least 2 DMARDs for periods of at least 3 months each can be met using single agents sequentially or by using one or more combinations of DMARDs.

If the requirement to trial 6 months of intensive DMARD therapy with at least 2 DMARDs cannot be met because of contraindications and/or intolerances of a severity necessitating permanent treatment withdrawal to all of the DMARDs specified above, details of the contraindication or intolerance including severity and dose for each DMARD must be provided in the authority application.

The following criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; **AND** either

- (a) a total active joint count of at least 20 active (swollen and tender) joints; or
- (b) at least 4 active joints from the following list of major joints:
 - (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The joint count and ESR and/or CRP must be determined at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. All measures must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note The Department of Human Services website (www.humanservices.gov.au) has details of the toxicities, including severity, which will be accepted for the following purposes:

- (a) exempting a patient from the requirement to undertake a minimum 3 month trial of methotrexate at a 20 mg weekly dose;
- (b) substituting azathioprine, cyclosporin or sodium aurothiomalate for another DMARD as part of the 6 month intensive DMARD trial;
- (c) exempting a patient from the requirement for a 6 month trial of intensive DMARD therapy.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 2 (change or re-commencement of treatment after a break in biological medicine of less than 24 months)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must not have failed to respond to previous PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised biological medicine treatment for this condition 5 times, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either of the Initial 1, Initial 2, Initial 3, or continuing treatment restrictions, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

A patient who has demonstrated a response to a course of rituximab must have a PBS-subsidised biological therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 3 (re-commencement of treatment after a break in biological medicine of more than 24 months)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 24 months or more from the most recent PBS-subsidised biological medicine for this condition, **AND**
- Patient must not have failed to respond to previous PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised biological medicine treatment for this condition 5 times, **AND**
- The condition must have an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; OR
- The condition must have a C-reactive protein (CRP) level greater than 15 mg per L, **AND**
- The condition must have either (a) a total active joint count of at least 20 active (swollen and tender) joints; or (b) at least 4 active major joints, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Major joints are defined as (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

All measures of joint count and ESR and/or CRP must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial 1 (new patient) or Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 24 months) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 24 months) - balance of supply

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 24 months) restriction to complete 16 weeks treatment; OR

- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 24 months) to complete 16 weeks of treatment, **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

baricitinib 4 mg tablet, 28

11447L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	1271.34	41.30	Olumiant [LY]

baricitinib 2 mg tablet, 28

11437Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	1271.34	41.30	Olumiant [LY]

■ CLADRIBINE

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

10170

Relapsing remitting multiple sclerosis

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be diagnosed by a neurologist, **AND**
- The condition must be diagnosed as clinically definite relapsing-remitting multiple sclerosis by magnetic resonance imaging of the brain and/or spinal cord; OR
- The condition must be diagnosed as clinically definite relapsing-remitting multiple sclerosis, with written certification provided by a radiologist that a magnetic resonance imaging scan is contraindicated because of the risk of physical (not psychological) injury to the patient, **AND**
- The treatment must be the sole PBS-subsidised disease modifying therapy for this condition, **AND**
- Patient must have experienced at least 2 documented attacks of neurological dysfunction, believed to be due to multiple sclerosis, in the preceding 2 years of commencing a PBS-subsidised disease modifying therapy for this condition, **AND**
- Patient must be ambulatory (without assistance or support).

Where applicable, the date of the magnetic resonance imaging scan must be recorded in the patient's medical records.

The prescriber should write authority prescriptions for the appropriate combination of packs (1, 4 or 6 tablets) to provide sufficient drug for a treatment week based on the weight of the patient in accordance with the TGA approved Product Information. Separate authority prescriptions may be required where the dose for treatment week 5 is different to the dose for treatment week 1.

Authority required (STREAMLINED)

10171

Relapsing remitting multiple sclerosis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a neurologist.

Clinical criteria:

- The condition must be diagnosed as clinically definite relapsing-remitting multiple sclerosis by magnetic resonance imaging of the brain and/or spinal cord, **AND**
- The treatment must be the sole PBS-subsidised disease modifying therapy for this condition, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not show continuing progression of disability while on treatment with this drug, **AND**
- Patient must have demonstrated compliance with, and an ability to tolerate, this therapy.

The prescriber should request authority approval for the appropriate combination of packs (1, 4 or 6 tablets) to provide sufficient drug for a treatment week based on the weight of the patient in accordance with the TGA approved Product Information. Separate authority prescriptions may be required where the dose for treatment week 5 is different to the dose for treatment week 1.

cladribine 10 mg tablet, 1

11603Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	3994.20	41.30	Mavenclad [SG]

cladribine 10 mg tablet, 6

11611D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	23159.40	41.30	Mavenclad [SG]

cladribine 10 mg tablet, 4

11604R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	1	..	*30825.48	41.30	Mavenclad [SG]

■ EVEROLIMUS

Caution Careful monitoring of patients is mandatory.

everolimus 250 microgram tablet, 60

8840G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	223.51	41.30	Certican [NV]

everolimus 1 mg tablet, 60

9352F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	3	..	*1674.66	41.30	Certican [NV]

everolimus 500 microgram tablet, 60

8841H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	440.00	41.30	Certican [NV]

everolimus 750 microgram tablet, 60

8842J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	3	..	*1271.96	41.30	Certican [NV]

■ FINGOLIMOD

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

10198

Multiple sclerosis

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be diagnosed as clinically definite relapsing-remitting multiple sclerosis by magnetic resonance imaging of the brain and/or spinal cord; OR
- The condition must be diagnosed as clinically definite relapsing-remitting multiple sclerosis by accompanying written certification provided by a radiologist that a magnetic resonance imaging scan is contraindicated because of the risk of physical (not psychological) injury to the patient, **AND**
- The treatment must be the sole PBS-subsidised disease modifying therapy for this condition, **AND**
- Patient must have experienced at least 2 documented attacks of neurological dysfunction, believed to be due to multiple sclerosis, in the preceding 2 years of commencing a PBS-subsidised disease modifying therapy for this condition, **AND**
- Patient must be ambulatory (without assistance or support).

Population criteria:

- Patient must weigh 40 kg or less.

Where applicable, the date of the magnetic resonance imaging scan must be recorded in the patient's medical records.

Authority required (STREAMLINED)

10093

Multiple sclerosis

Treatment Phase: Continuing treatment

Clinical criteria:

- The condition must be diagnosed as clinically definite relapsing-remitting multiple sclerosis, **AND**
- The treatment must be the sole PBS-subsidised disease modifying therapy for this condition, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not show continuing progression of disability while on treatment with this drug, **AND**
- Patient must have demonstrated compliance with, and an ability to tolerate this therapy.

Population criteria:

- Patient must weigh 40 kg or less.

fingolimod 250 microgram capsule, 28

11818B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	2219.45	41.30	Gilenya [NV]

■ FINGOLIMOD

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

10162

Multiple sclerosis

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be diagnosed as clinically definite relapsing-remitting multiple sclerosis by magnetic resonance imaging of the brain and/or spinal cord; OR
- The condition must be diagnosed as clinically definite relapsing-remitting multiple sclerosis by accompanying written certification provided by a radiologist that a magnetic resonance imaging scan is contraindicated because of the risk of physical (not psychological) injury to the patient, **AND**
- The treatment must be the sole PBS-subsidised disease modifying therapy for this condition, **AND**
- Patient must have experienced at least 2 documented attacks of neurological dysfunction, believed to be due to multiple sclerosis, in the preceding 2 years of commencing a PBS-subsidised disease modifying therapy for this condition, **AND**
- Patient must be ambulatory (without assistance or support).

Where applicable, the date of the magnetic resonance imaging scan must be recorded in the patient's medical records.

Authority required (STREAMLINED)

10172

Multiple sclerosis

Treatment Phase: Continuing treatment

Clinical criteria:

- The condition must be diagnosed as clinically definite relapsing-remitting multiple sclerosis, **AND**
- The treatment must be the sole PBS-subsidised disease modifying therapy for this condition, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not show continuing progression of disability while on treatment with this drug, **AND**
- Patient must have demonstrated compliance with, and an ability to tolerate this therapy.

fingolimod 500 microgram capsule, 28

5262Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	2219.45	41.30	Gilenya [NV]

▪ **LEFLUNOMIDE**

Caution Leflunomide is a category X drug and must not be given to pregnant women. Pregnancy should be avoided for two years after cessation of therapy, unless special wash-out procedures are carried out.

Authority required (STREAMLINED)

5766

Severe active psoriatic arthritis

Clinical criteria:

- Patient must have previously received, and failed to achieve an adequate response to, one or more disease modifying anti-rheumatic drugs including methotrexate; OR
- Patient must be clinically inappropriate for treatment with one or more disease modifying anti-rheumatic drugs including methotrexate, **AND**
- The treatment must be initiated by a physician.

leflunomide 10 mg tablet, 30

5449T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	33.58	34.87	^a Arabloc [AV] ^a Ataris 10 [AF] ^a Leflunomide generichealth [HQ]	^a Arava [SW] ^a Leflunomide APOTEX [GX] ^a Leflunomide Sandoz [SZ]

leflunomide 20 mg tablet, 30

5450W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	44.86	41.30	^a Arabloc [AV] ^a Ataris 20 [AF] ^a Leflunomide generichealth [HQ]	^a Arava [SW] ^a Leflunomide APOTEX [GX] ^a Leflunomide Sandoz [SZ]

▪ **LEFLUNOMIDE**

Caution Leflunomide is a category X drug and must not be given to pregnant women. Pregnancy should be avoided for two years after cessation of therapy, unless special wash-out procedures are carried out.

Authority required (STREAMLINED)

5681

Severe active rheumatoid arthritis

Clinical criteria:

- Patient must have previously received, and failed to achieve an adequate response to, one or more disease modifying anti-rheumatic drugs including methotrexate; OR
- Patient must be clinically inappropriate for treatment with one or more disease modifying anti-rheumatic drugs including methotrexate, **AND**
- The treatment must be initiated by a physician.

leflunomide 10 mg tablet, 30

8374R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	33.58	34.87	^a Arabloc [AV] ^a Ataris 10 [AF]	^a Arava [SW] ^a Leflunomide AN [EA]

^a Leflunomide APOTEX [GX]^a Leflunomide generichealth [HQ]^a Leflunomide Sandoz [SZ]^a Lunava 10 [ZP]**leflunomide 20 mg tablet, 30**

8375T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	44.86	41.30	^a Arabloc [AV] ^a Ataris 20 [AF] ^a Leflunomide APOTEX [GX] ^a Leflunomide Sandoz [SZ]	^a Arava [SW] ^a Leflunomide AN [EA] ^a Leflunomide generichealth [HQ] ^a Lunava 20 [ZP]

■ MYCOPHENOLATE**Caution** Careful monitoring of patients is mandatory.**mycophenolate mofetil 1 g/5 mL powder for oral liquid, 165 mL**

8651H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	5	..	#285.97	41.30	CellCept [RO]

mycophenolate 180 mg enteric tablet, 120

2150E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	104.26	41.30	Myfortic [NV]

mycophenolate mofetil 500 mg tablet, 50

8650G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	3	5	..	*155.91	41.30	^a APO-Mycophenolate [TX] ^a Ceptolate [AF] ^a Mycophenolate AN [EA] ^a Mycophenolate GH [GQ] ^a Pharmacor Mycophenolate 500 [CR]	^a CellCept [RO] ^a MycoCept [RF] ^a Mycophenolate APOTEX [GX] ^a Mycophenolate Sandoz [SZ]

mycophenolate 360 mg enteric tablet, 120

2193K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	200.70	41.30	Myfortic [NV]

■ MYCOPHENOLATE**Caution** Careful monitoring of patients is mandatory.**Note** For item codes 8649F and 1836P, pharmaceutical benefits that have the form capsule 250 mg are equivalent for the purposes of substitution.**mycophenolate mofetil 250 mg capsule, 50**

1836P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	6	5	..	*155.94	41.30	^a Ceptolate [AF]

mycophenolate mofetil 250 mg capsule, 100

8649F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	3	5	..	*155.97	41.30	^a APO-Mycophenolate [TX] ^a Mycophenolate Sandoz [SZ]	^a CellCept [RO] ^a Pharmacor Mycophenolate 250 [CR]

■ OZANIMOD**Note** The initiation pack is intended for use at the commencement of treatment and for re-initiation of therapy following treatment interruption. The first prescription for the 920 microgram capsules should occur under the Initial treatment restriction. Subsequent prescriptions should then occur under the Continuing treatment restriction. If treatment interruption of more than 14 consecutive days occurs for reasons other than lack of efficacy, apply under the Continuing treatment restriction for each of the Initiation pack and 920 microgram capsules. If treatment interruption occurs within the first 14 days, or, for more than 7 consecutive doses between treatment Days 15 to 28, re-apply for an initiation pack under the 'Initial treatment' treatment phase.**Note** No increase in the maximum quantity or number of units may be authorised.**Note** No increase in the maximum number of repeats may be authorised.**Note** Special Pricing Arrangements apply.**Authority required (STREAMLINED)****10162**

Multiple sclerosis

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be diagnosed as clinically definite relapsing-remitting multiple sclerosis by magnetic resonance imaging of the brain and/or spinal cord; OR

- The condition must be diagnosed as clinically definite relapsing-remitting multiple sclerosis by accompanying written certification provided by a radiologist that a magnetic resonance imaging scan is contraindicated because of the risk of physical (not psychological) injury to the patient, **AND**
- The treatment must be the sole PBS-subsidised disease modifying therapy for this condition, **AND**
- Patient must have experienced at least 2 documented attacks of neurological dysfunction, believed to be due to multiple sclerosis, in the preceding 2 years of commencing a PBS-subsidised disease modifying therapy for this condition, **AND**
- Patient must be ambulatory (without assistance or support).

Where applicable, the date of the magnetic resonance imaging scan must be recorded in the patient's medical records.

Authority required (STREAMLINED)

10172

Multiple sclerosis

Treatment Phase: Continuing treatment

Clinical criteria:

- The condition must be diagnosed as clinically definite relapsing-remitting multiple sclerosis, **AND**
- The treatment must be the sole PBS-subsidised disease modifying therapy for this condition, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not show continuing progression of disability while on treatment with this drug, **AND**
- Patient must have demonstrated compliance with, and an ability to tolerate this therapy.

ozanimod 230 microgram capsule [4] (&) ozanimod 460 microgram capsule [3], 7

12278F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	587.95	41.30	Zeposia [CJ]

ozanimod 920 microgram capsule, 28

12271W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	2219.45	41.30	Zeposia [CJ]

▪ **SIPONIMOD**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

10955

Multiple sclerosis

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be/have previously been diagnosed as clinically definite relapsing-remitting multiple sclerosis by magnetic resonance imaging of at least one of the brain/spinal cord; OR
- The condition must be/have previously been diagnosed as clinically definite relapsing-remitting multiple sclerosis supported by written certification, which is documented in the patient's medical records, from a radiologist that a magnetic resonance imaging scan is contraindicated because of the risk of physical (not psychological) injury to the patient, **AND**
- The treatment must be the sole PBS-subsidised disease modifying therapy for this condition, **AND**
- Patient must be ambulatory, with/without assistance/support, **AND**
- Patient must have mild disability in at least 3 functional systems; OR
- Patient must have moderate disability in at least 1 functional system.

Functional systems referred to in this restriction are the: visual, brain stem, pyramidal, cerebellar, sensory, bowel/bladder and cerebral/cognitive systems.

Select a dose and pack size appropriate for the patient's CYP2C9 metabolising enzyme status.

Note There is no specific Medical Benefits Schedule item for CYP2C9 metabolising enzyme status testing.

Authority required (STREAMLINED)

10953

Multiple sclerosis

Treatment Phase: Continuing treatment (including recommencement of treatment)

Clinical criteria:

- The treatment must be the sole PBS-subsidised disease modifying therapy for this condition, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not show continuing progression of disability while on treatment with this drug, **AND**
- Patient must be ambulatory, with/without assistance/support, **AND**
- Patient must have demonstrated compliance with, and an ability to tolerate this therapy.

Authority required (STREAMLINED)

10956

Multiple sclerosis

Treatment Phase: Grandfather treatment (transition from non-PBS-subsidised to PBS-subsidised treatment)

Clinical criteria:

- Patient must have commenced non-PBS-subsidised treatment with this drug for this PBS indication prior to 1 November 2020, **AND**

- The condition must have previously been diagnosed as clinically definite relapsing-remitting multiple sclerosis by magnetic resonance imaging of at least one of the brain/spinal cord prior to initiating non-PBS-subsidised treatment with this drug for this condition; OR
 - The condition must have previously been diagnosed as clinically definite relapsing-remitting multiple sclerosis supported by written certification, which is documented in the patient's medical records, from a radiologist that a magnetic resonance imaging scan is contraindicated because of the risk of physical (not psychological) injury to the patient prior to initiating non-PBS-subsidised treatment with this drug for this condition, **AND**
 - The treatment must be the sole PBS-subsidised disease modifying therapy for this condition, **AND**
 - Patient must have mild disability in at least 3 functional systems prior to initiating non-PBS-subsidised treatment with this drug for this condition; OR
 - Patient must have moderate disability in at least 1 functional system prior to initiating non-PBS-subsidised treatment with this drug for this condition, **AND**
 - Patient must not show continuing progression of disability while on treatment with this drug, **AND**
 - Patient must be ambulatory, with/without assistance/support, **AND**
 - Patient must have demonstrated compliance with, and an ability to tolerate this therapy.
- Functional systems referred to in this restriction are the: visual, brain stem, pyramidal, cerebellar, sensory, bowel/bladder and cerebral/cognitive systems.

A Grandfathered patient may qualify for PBS-subsidised treatment under this restriction once only.

Note This grandfather restriction will cease to operate from 12 months after the date specified in the clinical criteria.

siponimod 250 microgram tablet, 120

12160B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	2219.46	41.30	Mayzent [NV]

siponimod 2 mg tablet, 28

12158X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	2219.45	41.30	Mayzent [NV]

siponimod 250 microgram tablet, 12

12172P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	239.40	41.30	Mayzent [NV]

■ SIROLIMUS

Caution Careful monitoring of patients is mandatory.

sirolimus 500 microgram tablet, 100

8984W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	338.67	41.30	Rapamune [PF]

sirolimus 2 mg tablet, 100

8833X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	1297.68	41.30	Rapamune [PF]

sirolimus 1 mg/mL oral liquid, 60 mL

8725F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	3	..	528.61	41.30	Rapamune [PF]

sirolimus 1 mg tablet, 100

8724E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	670.31	41.30	Rapamune [PF]

■ TERIFLUNOMIDE

Caution Teriflunomide is a category X drug and must not be given to pregnant women or women of childbearing potential who are not currently using reliable contraception.

Pregnancy should be avoided for two years after cessation of therapy, unless special wash-out procedures are carried out.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

10150

Multiple sclerosis

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be diagnosed as clinically definite relapsing-remitting multiple sclerosis by magnetic resonance imaging of the brain and/or spinal cord; OR
- The condition must be diagnosed as clinically definite relapsing-remitting multiple sclerosis by accompanying written certification provided by a radiologist that a magnetic resonance imaging scan is contraindicated because of the risk of physical (not psychological) injury to the patient, **AND**
- The treatment must be the sole PBS-subsidised disease modifying therapy for this condition, **AND**

- Patient must have experienced at least 2 documented attacks of neurological dysfunction, believed to be due to multiple sclerosis, in the preceding 2 years of commencing a PBS-subsidised disease modifying therapy for this condition, **AND**
- Patient must be ambulatory (without assistance or support).

Where applicable, the date of the magnetic resonance imaging scan must be recorded in the patient's medical records.

Authority required (STREAMLINED)

10199

Multiple sclerosis

Treatment Phase: Continuing treatment

Clinical criteria:

- The condition must be diagnosed as clinically definite relapsing-remitting multiple sclerosis by magnetic resonance imaging of the brain and/or spinal cord; OR
- The condition must be diagnosed as clinically definite relapsing-remitting multiple sclerosis by accompanying written certification provided by a radiologist that a magnetic resonance imaging scan is contraindicated because of the risk of physical (not psychological) injury to the patient, **AND**
- The treatment must be the sole PBS-subsidised disease modifying therapy for this condition, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not show continuing progression of disability while on treatment with this drug.

Where applicable, the date of the magnetic resonance imaging scan must be recorded in the patient's medical records.

teriflunomide 14 mg tablet, 28

2898M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	584.13	41.30	^a APO-TERIFLUNOMIDE [TX]	^a Aubagio [GZ]
						^a Pharmacor Teriflunomide [CR]	^a TERIFLAGIO [RW]
						^a Teriflunomide Dr.Reddy's [RZ]	^a Teriflunomide GH [GQ]
						^a Teriflunomide Sandoz [SZ]	^a Terimide [AF]

▪ **TOFACITINIB**

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for adults with severe active rheumatoid arthritis. Where the term biological medicine appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alfa antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab), the T-cell co-stimulation modulator (abatacept) and the Janus kinase (JAK) inhibitors (baricitinib, tofacitinib, upadacitinib).

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised biological medicine therapy may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once, and
- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised biological medicines for the treatment of rheumatoid arthritis.

A serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Stevens Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered a treatment failure.

A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction.

A patient who has failed fewer than 5 biological medicines and who has a break in treatment of less than 24 months may commence a further course of treatment with a biological medicine under Initial 2 treatment restriction. A patient who has failed fewer than 5 biological medicines and who has had a break in therapy of longer than 24 months may commence a further course of treatment with a biological medicine under the Initial 3 treatment restriction. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine treatment is stopped to the date of the new application for treatment with a biological medicine.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 April 2019.

(a) Initial treatment.

Applications for initial treatment should be made where:

- a patient has received no prior PBS-subsidised biological medicine treatment and wishes to commence such therapy, excluding rituximab (Initial 1 - new patient); or
 - a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months) [further details are under 'Swapping therapy' below]; or
 - a patient wishes to re-commence treatment with a specific biological medicine following a break of less than 24 months in PBS-subsidised therapy with that agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months).
 - a patient wishes to re-commence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 24 months (Initial 3 - re-commencement of treatment after a break in biological medicine of more than 24 months)
- Initial applications for a new patient (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6-month intensive DMARD trial, but prior to ceasing DMARD therapy.
- Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab,

baricitinib, etanercept, golimumab, tocilizumab, tofacitinib and upadacitinib, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab. Rituximab patients should be assessed following a minimum of 12 weeks after the first infusion, and the assessment should be submitted to Services Australia within 4 weeks to ensure continuity of treatment for those who meet the continuing restriction for PBS subsidised treatment with this drug for this condition.

Where a response assessment is not provided the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment. For second and subsequent courses of PBS-subsidised biological medicine (excluding rituximab) treatment, it is recommended that a patient be reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to Services Australia within sufficient time to allow processing. Abatacept patients:

A patient is eligible to receive one I.V. loading dose when commencing treatment with the subcutaneous formulation. Two prescriptions are required, the first prescription for the I.V. loading dose for sufficient vials for one dose based on the patient's weight with no repeats. The second prescription for the subcutaneous formulation, with a maximum quantity of 4 and up to 3 repeats, must be submitted with the initial application.

Rituximab patients:

A further application may be submitted to Services Australia within sufficient time to allow processing. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

Assessment of the patient's response to treatment must be conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the cessation of the most recent course of biological medicine therapy.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction. Where a response assessment is not conducted within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

(2) Swapping therapy

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non- biological medicine therapy requirements except if the patient has had a break in therapy of more than 24 months who would need to requalify with respect to the indices of disease severity. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each biological medicine trialled.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent, unless they have experienced a serious adverse reaction of a severity necessitating permanent treatment withdrawal.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug.

Abatacept:

A patient swapping from I.V. abatacept to subcutaneous abatacept will not be eligible for an I.V. loading dose when commencing treatment with the subcutaneous formulation.

Rituximab:

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF- α antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they be assessed for response to every course of treatment, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological medicine during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological medicine therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate biological medicine may do so without having to have any treatment-free period.

(3) Baseline measurements to determine response.

Determination of whether a response to treatment has been demonstrated must be based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and the eligibility for continuing treatment must be assessed according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response must be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be determined on the total number of major joints.

Applications under the Initial 1 treatment restriction for a new patient must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. The results must be no more than 4 weeks old at the time of application.

Applications under the Initial 3 treatment restriction for re-commencement of treatment after a break in biological medicine of more than 24 months must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment per continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient has either failed or ceased to respond to a PBS-subsidised biological medicine for this condition 5 times, they will not be eligible to receive further PBS-subsidised treatment with a biological medicine for this condition.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Continuing Treatment - balance of supply.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restriction.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

tofacitinib 5 mg tablet, 56

10511F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1210.96	41.30	Xeljanz [PF]

■ TOFACITINIB
Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE PSORIATIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab for adult patients with severe active psoriatic arthritis. Therefore, where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient receiving PBS-subsidised treatment for psoriatic arthritis is able to commence a treatment cycle where they may trial biological medicines without having to experience a disease flare when swapping to the alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

A patient who received PBS-subsidised adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab or ustekinumab treatment prior to 1 May 2019 is considered to start their first cycle as of 1 May 2019.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence another cycle [further details are under '(5) Recommencement of treatment after a 5-year break in PBS-subsidised therapy' below].

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe active psoriatic arthritis.

(1) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an alternate medicine (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years) or An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, golimumab secukinumab and tofacitinib, 18 to 20 weeks of therapy for certolizumab pegol (depending upon the dosing regimen), 20 weeks of therapy for ixekizumab, 22 weeks of therapy for infliximab, and 28 weeks of therapy for ustekinumab. It is recommended that a patient be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of PBS-subsidised initial treatment following a minimum of 12 weeks of therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

Grandfather patients (ixekizumab only).

A patient who commenced treatment with ixekizumab for severe psoriatic arthritis prior to 1 March 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

Grandfather patients (tofacitinib only).

A patient who commenced treatment with Tofacitinib for severe psoriatic arthritis prior to 1 May 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment under the Initial 1 or Initial 2 treatment restrictions.

For the second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to re-qualify with respect to either the indices of disease severity (i.e. erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP) level, and active joint count) or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify under the Initial 3 treatment restriction with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application.

However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

Within a treatment cycle a patient may alternate between therapy with any biological medicine of their choice (1 at a time) providing:

- (i) they have not received PBS-subsidised treatment with that particular biological medicine previously; or
- (ii) they have demonstrated an adequate response to that particular biological medicine if they have previously trialled it on the PBS; and
- (iii) they have not previously failed to respond to treatment 3 times in this treatment cycle with that biological medicine.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(4) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the indices of disease severity submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial or change or commencement treatment application is submitted within a treatment cycle and these revised baseline measurements will be used to assess response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to determine response for all subsequent continuing treatments. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. 20 or more active joints), response will be determined according to a reduction in the total number of active joints.

(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment restriction according to the criteria of the relevant restriction and index of disease severity. The application must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial treatment - Grandfather treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must have received non-PBS subsidised treatment with this drug for this condition prior to 1 May 2019, **AND**
- Patient must be receiving treatment with this drug for this condition at the time of application, **AND**

- Patient must have failed to achieve an adequate response to methotrexate at a dose of at least 20 mg weekly for a minimum period of 3 months prior to initiating non-PBS subsidised treatment with this drug for this condition, **AND**
- Patient must have failed to achieve an adequate response to sulfasalazine at a dose of at least 2 g per day for a minimum period of 3 months prior to initiating non-PBS subsidised treatment with this drug for this condition; OR
- Patient must have failed to achieve an adequate response to leflunomide at a dose of up to 20 mg daily for a minimum period of 3 months prior to initiating non-PBS subsidised treatment with this drug for this condition, **AND**
- Patient must have demonstrated an adequate response following at least 12 weeks of non-PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

The following initiation criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; and

either

(a) an active joint count of at least 20 active (swollen and tender) joints; or

(b) at least 4 active joints from the following list of major joints:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

An adequate response to treatment is defined as:

an erythrocyte sedimentation rate (ESR) no greater than 25 mm per hour or a C-reactive protein (CRP) level no greater than 15 mg per L or either marker reduced by at least 20% from baseline; and

either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following major active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to determine response for all subsequent continuing treatments.

The assessment of the patient's response to this PBS-subsidised course of therapy must be conducted no later than 4 weeks from the cessation of the treatment course.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

A patient may qualify for PBS-subsidised treatment under this restriction once only.

For continuing PBS-subsidised treatment, a Grandfathered patient must qualify under the Continuing treatment criteria.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Severe Psoriatic Arthritis PBS Authority Application - Supporting Information Form; and
- (3) the date of commencement of this drug; and
- (4) results of the baseline patient assessment prior to initiation of non-PBS subsidised therapy with this drug.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an erythrocyte sedimentation rate (ESR) no greater than 25 mm per hour or a C-reactive protein (CRP) level no greater than 15 mg per L or either marker reduced by at least 20% from baseline; and

either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following major active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to determine response for all subsequent continuing treatments.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Severe Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either Initial 1, Initial 2, or Initial 3 treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Continuing treatment or Grandfathered patients - balance of supply

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Continuing treatment restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Grandfathered treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restriction.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

tofacitinib 5 mg tablet, 56

11675L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1210.96	41.30	Xeljanz [PF]

▪ TOFACITINIB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for adults with severe active rheumatoid arthritis. Where the term biological medicine appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alpha antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab), the T-cell co-stimulation modulator (abatacept) and the Janus kinase (JAK) inhibitors (baricitinib, tofacitinib, upadacitinib).

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF- α antagonist.

A patient receiving PBS-subsidised biological medicine therapy may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once, and
- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised biological medicines for the treatment of rheumatoid arthritis.

A serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Stevens Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered a treatment failure.

A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction.

A patient who has failed fewer than 5 biological medicines and who has a break in treatment of less than 24 months may commence a further course of treatment with a biological medicine under Initial 2 treatment restriction. A patient who has failed fewer than 5 biological medicines and who has had a break in therapy of longer than 24 months may commence a further course of treatment with a biological medicine under the Initial 3 treatment restriction. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine treatment is stopped to the date of the new application for treatment with a biological medicine.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 April 2019.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised biological medicine treatment and wishes to commence such therapy, excluding rituximab (Initial 1 - new patient); or
 - (ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months) [further details are under 'Swapping therapy' below]; or
 - (iii) a patient wishes to re-commence treatment with a specific biological medicine following a break of less than 24 months in PBS-subsidised therapy with that agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months).
 - (iv) a patient wishes to re-commence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 24 months (Initial 3 - re-commencement of treatment after a break in biological medicine of more than 24 months)
- Initial applications for a new patient (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6-month intensive DMARD trial, but prior to ceasing DMARD therapy.

Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, baricitinib, etanercept, golimumab, tocilizumab, tofacitinib and upadacitinib, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

Rituximab patients should be assessed following a minimum of 12 weeks after the first infusion, and the assessment should be submitted to Services Australia within 4 weeks to ensure continuity of treatment for those who meet the continuing restriction for PBS subsidised treatment with this drug for this condition.

Where a response assessment is not provided the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment. For second and subsequent courses of PBS-subsidised biological medicine (excluding rituximab) treatment, it is recommended that a patient be reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to Services Australia within sufficient time to allow processing.

Abatacept patients:

A patient is eligible to receive one I.V. loading dose when commencing treatment with the subcutaneous formulation. Two prescriptions are required, the first prescription for the I.V. loading dose for sufficient vials for one dose based on the patient's weight with no repeats. The second prescription for the subcutaneous formulation, with a maximum quantity of 4 and up to 3 repeats, must be submitted with the initial application.

Rituximab patients:

A further application may be submitted to Services Australia within sufficient time to allow processing. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

Assessment of the patient's response to treatment must be conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the cessation of the most recent course of biological medicine therapy.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction. Where a response assessment is not conducted within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

(2) Swapping therapy

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non- biological medicine

therapy requirements except if the patient has had a break in therapy of more than 24 months who would need to requalify with respect to the indices of disease severity. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each biological medicine trialled.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent, unless they have experienced a serious adverse reaction of a severity necessitating permanent treatment withdrawal.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug.

Abatacept:

A patient swapping from I.V. abatacept to subcutaneous abatacept will not be eligible for an I.V. loading dose when commencing treatment with the subcutaneous formulation.

Rituximab:

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF-alfa antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they be assessed for response to every course of treatment, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological medicine during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological medicine therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate biological medicine may do so without having to have any treatment-free period.

(3) Baseline measurements to determine response.

Determination of whether a response to treatment has been demonstrated must be based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and the eligibility for continuing treatment must be assessed according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response must be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be determined on the total number of major joints.

Applications under the Initial 1 treatment restriction for a new patient must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. The results must be no more than 4 weeks old at the time of application.

Applications under the Initial 3 treatment restriction for re-commencement of treatment after a break in biological medicine of more than 24 months must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with disease modifying anti-rheumatic drugs (DMARDs) which must include at least 3 months continuous treatment with each of at least 2 DMARDs, one of which must be methotrexate at a dose of at least 20 mg weekly and one of which must be: (i) hydroxychloroquine at a dose of at least 200 mg daily; or (ii) leflunomide at a dose of at least 10 mg daily; or (iii) sulfasalazine at a dose of at least 2 g daily; OR
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs which, if methotrexate is contraindicated according to the Therapeutic Goods Administration (TGA)-approved Product Information or cannot be tolerated at a 20 mg weekly dose, must include at least 3 months continuous treatment with each of at least 2 of the following DMARDs: (i) hydroxychloroquine at a dose of at least 200 mg daily; and/or (ii) leflunomide at a dose of at least 10 mg daily; and/or (iii) sulfasalazine at a dose of at least 2 g daily; OR
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs which, if 3 or more of methotrexate, hydroxychloroquine, leflunomide and sulfasalazine are contraindicated according to the relevant TGA-approved Product Information or cannot be tolerated at the doses specified above, must include at least 3 months continuous treatment with each of at least 2 DMARDs, with one or more of the following DMARDs being used in place of the DMARDs which are contraindicated or not tolerated: (i) azathioprine at a dose of at least 1 mg/kg per day; and/or (ii) cyclosporin at a dose of at least 2 mg/kg/day; and/or (iii) sodium aurothiomalate at a dose of 50 mg weekly, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

If methotrexate is contraindicated according to the TGA-approved product information or cannot be tolerated at a 20 mg weekly dose, the application must include details of the contraindication or intolerance including severity to methotrexate. The maximum tolerated dose of methotrexate must be documented in the application, if applicable.

The application must include details of the DMARDs trialled, their doses and duration of treatment, and all relevant contraindications and/or intolerances including severity.

The requirement to trial at least 2 DMARDs for periods of at least 3 months each can be met using single agents sequentially or by using one or more combinations of DMARDs.

If the requirement to trial 6 months of intensive DMARD therapy with at least 2 DMARDs cannot be met because of contraindications and/or intolerances of a severity necessitating permanent treatment withdrawal to all of the DMARDs specified above, details of the contraindication or intolerance including severity and dose for each DMARD must be provided in the authority application.

The following criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; AND either

(a) a total active joint count of at least 20 active (swollen and tender) joints; or

(b) at least 4 active joints from the following list of major joints:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The joint count and ESR and/or CRP must be determined at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. All measures must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note The Department of Human Services website (www.humanservices.gov.au) has details of the toxicities, including severity, which will be accepted for the following purposes:

- (a) exempting a patient from the requirement to undertake a minimum 3 month trial of methotrexate at a 20 mg weekly dose;
- (b) substituting azathioprine, cyclosporin or sodium aurothiomalate for another DMARD as part of the 6 month intensive DMARD trial;
- (c) exempting a patient from the requirement for a 6 month trial of intensive DMARD therapy.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 2 (change or re-commencement of treatment after a break in biological medicine of less than 24 months)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition, **AND**

- Patient must not have failed to respond to previous PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have already failed , or ceased to respond to, PBS-subsidised biological medicine treatment for this condition 5 times, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either of the Initial 1, Initial 2, Initial 3, or continuing treatment restrictions, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

A patient who has demonstrated a response to a course of rituximab must have a PBS-subsidised biological therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 3 (re-commencement of treatment after a break in biological medicine of more than 24 months)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 24 months or more from the most recent PBS-subsidised biological medicine for this condition, **AND**
- Patient must not have failed to respond to previous PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have already failed , or ceased to respond to, PBS-subsidised biological medicine treatment for this condition 5 times, **AND**
- The condition must have an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; OR
- The condition must have a C-reactive protein (CRP) level greater than 15 mg per L, **AND**
- The condition must have either (a) a total active joint count of at least 20 active (swollen and tender) joints; or (b) at least 4 active major joints, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Major joints are defined as (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

All measures of joint count and ESR and/or CRP must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial 1 (new patient) or Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 24 months) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 24 months) - balance of supply

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 24 months) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 24 months) to complete 16 weeks of treatment, **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

tofacitinib 5 mg tablet, 56

10517M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	1210.96	41.30	Xeljanz [PF]

■ TOFACITINIB**Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE PSORIATIC ARTHRITIS**

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab for adult patients with severe active psoriatic arthritis. Therefore, where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient receiving PBS-subsidised treatment for psoriatic arthritis is able to commence a treatment cycle where they may trial biological medicines without having to experience a disease flare when swapping to the alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

A patient who received PBS-subsidised adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab or ustekinumab treatment prior to 1 May 2019 is considered to start their first cycle as of 1 May 2019.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence another cycle [further details are under '(5) Recommencement of treatment after a 5-year break in PBS-subsidised therapy' below].

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe active psoriatic arthritis.

(1) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an alternate medicine (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years) or An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, golimumab secukinumab and tofacitinib, 18 to 20 weeks of therapy for certolizumab pegol (depending upon the dosing regimen), 20 weeks of therapy for ixekizumab, 22 weeks of therapy for infliximab, and 28 weeks of therapy for ustekinumab. It is recommended that a patient be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of PBS-subsidised initial treatment following a minimum of 12 weeks of therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

Grandfather patients (ixekizumab only).

A patient who commenced treatment with ixekizumab for severe psoriatic arthritis prior to 1 March 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

Grandfather patients (tofacitinib only).

A patient who commenced treatment with Tofacitinib for severe psoriatic arthritis prior to 1 May 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment under the Initial 1 or Initial 2 treatment restrictions.

For the second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to re-qualify with respect to either the indices of disease severity (i.e. erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP) level, and active joint count) or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify under the Initial 3 treatment restriction with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application.

However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

Within a treatment cycle a patient may alternate between therapy with any biological medicine of their choice (1 at a time) providing:

- (i) they have not received PBS-subsidised treatment with that particular biological medicine previously; or
- (ii) they have demonstrated an adequate response to that particular biological medicine if they have previously trialled it on the PBS; and
- (iii) they have not previously failed to respond to treatment 3 times in this treatment cycle with that biological medicine.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(4) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the indices of disease severity submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and these revised baseline measurements will be used to assess response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to determine response for all subsequent continuing treatments. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. 20 or more active joints), response will be determined according to a reduction in the total number of active joints.

(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment restriction according to the criteria of the relevant restriction and index of disease severity. The application must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response to methotrexate at a dose of at least 20 mg weekly for a minimum period of 3 months, **AND**
- Patient must have failed to achieve an adequate response to sulfasalazine at a dose of at least 2 g per day for a minimum period of 3 months; OR
- Patient must have failed to achieve an adequate response to leflunomide at a dose of up to 20 mg daily for a minimum period of 3 months, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Where treatment with methotrexate, sulfasalazine or leflunomide is contraindicated according to the relevant TGA-approved Product Information, details must be provided at the time of application.

Where intolerance to treatment with methotrexate, sulfasalazine or leflunomide developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

The following initiation criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; and
either

- (a) an active joint count of at least 20 active (swollen and tender) joints; or
 - (b) at least 4 active joints from the following list of major joints:
 - (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).
- If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Severe Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

An assessment of a patient's response to an initial course of treatment must be conducted following a minimum of 12 weeks of therapy. An application for the continuing treatment must be accompanied with the assessment of response and submitted to the Department of Human Services no later than 4 weeks from the date of completion of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 3 months treatment with methotrexate and 3 months treatment with sulfasalazine or leflunomide can be found on the Department of Human Services website (www.humanservices.gov.au)

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial treatment - Initial 2 (change or recommencement of treatment after a break in in biological medicine of less than 5 years)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an erythrocyte sedimentation rate (ESR) no greater than 25 mm per hour or a C-reactive protein (CRP) level no greater than 15 mg per L or either marker reduced by at least 20% from baseline; and
either of the following:

- (a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or
- (b) a reduction in the number of the following major active joints, from at least 4, by at least 50%:
 - (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Severe Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to change or recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial treatment - Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must have an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; OR
- The condition must have a C-reactive protein (CRP) level greater than 15 mg per L, **AND**
- The condition must have either (a) a total active joint count of at least 20 active (swollen and tender) joints; or (b) at least 4 active major joints, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Major joints are defined as (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

All measures of joint count and ESR and/or CRP must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Severe Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date

of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial 1 (new patient) or Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years) - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 16 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

tofacitinib 5 mg tablet, 56

11690G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	1210.96	41.30	Xeljanz [PF]

■ UPADACITINIB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for adults with severe active rheumatoid arthritis. Where the term biological medicine appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alfa antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab), the T-cell co-stimulation modulator (abatacept) and the Janus kinase (JAK) inhibitors (baricitinib, tofacitinib, upadacitinib).

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised biological medicine therapy may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once, and
- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised biological medicines for the treatment of rheumatoid arthritis.

A serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Stevens Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered a treatment failure.

A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction.

A patient who has failed fewer than 5 biological medicines and who has a break in treatment of less than 24 months may commence a further course of treatment with a biological medicine under Initial 2 treatment restriction. A patient who has failed fewer than 5 biological medicines and who has had a break in therapy of longer than 24 months may commence a further course of treatment with a biological medicine under the Initial 3 treatment restriction. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine treatment is stopped to the

date of the new application for treatment with a biological medicine.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 April 2019.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised biological medicine treatment and wishes to commence such therapy, excluding rituximab (Initial 1 - new patient); or

(ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to re-commence treatment with a specific biological medicine following a break of less than 24 months in PBS-subsidised therapy with that agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months).

(iv) a patient wishes to re-commence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 24 months (Initial 3 - re-commencement of treatment after a break in biological medicine of more than 24 months). Initial applications for a new patient (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6-month intensive DMARD trial, but prior to ceasing DMARD therapy.

Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, baricitinib, etanercept, golimumab, tocilizumab, tofacitinib and upadacitinib, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

Rituximab patients should be assessed following a minimum of 12 weeks after the first infusion, and the assessment should be submitted to Services Australia within 4 weeks to ensure continuity of treatment for those who meet the continuing restriction for PBS subsidised treatment with this drug for this condition.

Where a response assessment is not provided the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment. For second and subsequent courses of PBS-subsidised biological medicine (excluding rituximab) treatment, it is recommended that a patient be reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to Services Australia within sufficient time to allow processing.

Abatacept patients:

A patient is eligible to receive one I.V. loading dose when commencing treatment with the subcutaneous formulation. Two prescriptions are required, the first prescription for the I.V. loading dose for sufficient vials for one dose based on the patient's weight with no repeats. The second prescription for the subcutaneous formulation, with a maximum quantity of 4 and up to 3 repeats, must be submitted with the initial application.

Rituximab patients:

A further application may be submitted to Services Australia within sufficient time to allow processing. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

Assessment of the patient's response to treatment must be conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the cessation of the most recent course of biological medicine therapy.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction. Where a response assessment is not conducted within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

(2) Swapping therapy

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non- biological medicine therapy requirements except if the patient has had a break in therapy of more than 24 months who would need to requalify with respect to the indices of disease severity. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each biological medicine trialled.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent, unless they have experienced a serious adverse reaction of a severity necessitating permanent treatment withdrawal.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug.

Abatacept:

A patient swapping from I.V. abatacept to subcutaneous abatacept will not be eligible for an I.V. loading dose when commencing treatment with the subcutaneous formulation.

Rituximab:

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF-alfa antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they be assessed for response to every course of treatment, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological medicine during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-

subsidised biological medicine therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate biological medicine may do so without having to have any treatment-free period.

(3) Baseline measurements to determine response.

Determination of whether a response to treatment has been demonstrated must be based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and the eligibility for continuing treatment must be assessed according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response must be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be determined on the total number of major joints.

Applications under the Initial 1 treatment restriction for a new patient must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. The results must be no more than 4 weeks old at the time of application.

Applications under the Initial 3 treatment restriction for re-commencement of treatment after a break in biological medicine of more than 24 months must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment per continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response must be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be determined on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker must be used to determine response.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient has either failed or ceased to respond to a PBS-subsidised biological medicine for this condition 5 times, they will not be eligible to receive further PBS-subsidised treatment with a biological medicine for this condition.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Grandfathered patients

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have previously received non-PBS-subsidised therapy with this drug for this condition prior to 1 May 2020, **AND**
- Patient must be receiving treatment with this drug for this condition at the time of application, **AND**
- Patient must have failed to achieve an adequate response to a trial of at least 6 months of intensive treatment with disease modifying anti-rheumatic drugs (DMARDs) prior to initiating non-PBS-subsidised treatment with this drug for this condition. This must have included at least 3 months continuous treatment with each of at least 2 DMARDs, one of which must have been methotrexate at a dose of at least 20 mg weekly and one of which must have been: (i) hydroxychloroquine at a dose of at least 200 mg daily; or (ii) leflunomide at a dose of at least 10 mg daily; or (iii) sulfasalazine at a dose of at least 2 g daily; OR
- Patient must have failed to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs prior to initiating non-PBS-subsidised treatment with this drug for this condition. If methotrexate was contraindicated according to the Therapeutic Goods Administration (TGA)-approved Product Information or could not be tolerated at a 20 mg weekly dose, this intensive treatment must have included at least 3 months continuous treatment with each of at least 2 of the following DMARDs: (i) hydroxychloroquine at a dose of at least 200 mg daily; and/or (ii) leflunomide at a dose of at least 10 mg daily; and/or (iii) sulfasalazine at a dose of at least 2 g daily; OR
- Patient must have failed to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs prior to initiating non-PBS-subsidised treatment with this drug for this condition. If 3 or more of methotrexate, hydroxychloroquine, leflunomide and sulfasalazine were contraindicated according to the relevant TGA-approved Product Information or could not be tolerated at the doses specified above, the intensive treatment must have included at least 3 months continuous treatment with each of at least 2 DMARDs, with one or more of the following DMARDs used in place of the DMARDs which were contraindicated or not tolerated: (i) azathioprine at a dose of at least 1 mg/kg per day; and/or (ii) cyclosporin at a dose of at least 2 mg/kg/day; and/or (iii) sodium aurothiomalate at a dose of 50 mg weekly, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised biological medicine treatment for this condition 5 times, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

The application must include details of the DMARDs trialed, their doses and duration of treatment, and all relevant contraindications and/or intolerances including severity.

The requirement to trial at least 2 DMARDs for periods of at least 3 months each can be met using single agents sequentially or by using one or more combinations of DMARDs.

If the requirement to trial 6 months of intensive DMARD therapy with at least 2 DMARDs cannot be met because of contraindications and/or intolerances of a severity necessitating permanent treatment withdrawal to all of the DMARDs specified above, details of the contraindication or intolerance including severity and dose for each DMARD must be provided in the authority application.

The following criteria indicate failure to have achieved an adequate response to DMARD treatment prior to initiating non-PBS-subsidised treatment with this drug for this condition:

an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour and/or a C-reactive protein (CRP) level greater than 15 mg per L; **AND** either

(a) a total active joint count of at least 20 active (swollen and tender) joints; or

(b) at least 4 active joints from the following list of major joints:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response must be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be determined on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker must be used to determine response.

All applications for treatment with this drug for this condition under this restriction must include baseline joint count and ESR and/or CRP as determined at the completion of a 6 month intensive DMARD trial but prior to ceasing DMARD therapy.

If the requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied. Treatment with prednisolone dosed at 7.5 mg or higher daily (or equivalent) or a parenteral steroid within the past month (intramuscular or intravenous methylprednisolone or equivalent) is an acceptable reason.

A Grandfathered patient may qualify for PBS-subsidised treatment under this restriction once only. For continuing PBS-subsidised treatment, a Grandfathered patient must qualify under the continuing treatment criteria.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Continuing and Initial Grandfathered patients treatment - balance of supply

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the continuing treatment restriction to complete 24 weeks of treatment; OR
- Patient must have received insufficient treatment with this drug to complete 24 weeks of treatment under the Initial treatment - Grandfathered patients, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restrictions.

Population criteria:

- Patient must be aged 18 years or older.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

upadacitinib 15 mg modified release tablet, 28

11979L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1271.34	41.30	Rinvoq [VE]

■ UPADACITINIB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for adults with severe active rheumatoid arthritis. Where the term biological medicine appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alfa antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab), the T-cell co-stimulation modulator (abatacept) and the Janus kinase (JAK) inhibitors (baricitinib, tofacitinib, upadacitinib).

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised biological medicine therapy may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once, and
- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised biological medicines for the treatment of rheumatoid arthritis.

A serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Stevens Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered a treatment failure.

A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction.

A patient who has failed fewer than 5 biological medicines and who has a break in treatment of less than 24 months may commence a further course of treatment with a biological medicine under Initial 2 treatment restriction. A patient who has failed fewer than 5 biological medicines and who has had a break in therapy of longer than 24 months may commence a further course of treatment with a biological medicine under the Initial 3 treatment restriction. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine treatment is stopped to the date of the new application for treatment with a biological medicine.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 April 2019.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised biological medicine treatment and wishes to commence such therapy, excluding rituximab (Initial 1 - new patient); or
- (ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an

alternate agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to re-commence treatment with a specific biological medicine following a break of less than 24 months in PBS-subsidised therapy with that agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months).

(iv) a patient wishes to re-commence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 24 months (Initial 3 - re-commencement of treatment after a break in biological medicine of more than 24 months) Initial applications for a new patient (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6-month intensive DMARD trial, but prior to ceasing DMARD therapy.

Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, baricitinib, etanercept, golimumab, tocilizumab, tofacitinib and upadacitinib, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

Rituximab patients should be assessed following a minimum of 12 weeks after the first infusion, and the assessment should be submitted to Services Australia within 4 weeks to ensure continuity of treatment for those who meet the continuing restriction for PBS subsidised treatment with this drug for this condition.

Where a response assessment is not provided the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment. For second and subsequent courses of PBS-subsidised biological medicine (excluding rituximab) treatment, it is recommended that a patient be reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to Services Australia within sufficient time to allow processing.

Abatacept patients:

A patient is eligible to receive one I.V. loading dose when commencing treatment with the subcutaneous formulation. Two prescriptions are required, the first prescription for the I.V. loading dose for sufficient vials for one dose based on the patient's weight with no repeats. The second prescription for the subcutaneous formulation, with a maximum quantity of 4 and up to 3 repeats, must be submitted with the initial application.

Rituximab patients:

A further application may be submitted to Services Australia within sufficient time to allow processing. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

Assessment of the patient's response to treatment must be conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the cessation of the most recent course of biological medicine therapy.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction. Where a response assessment is not conducted within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

(2) Swapping therapy

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non- biological medicine therapy requirements except if the patient has had a break in therapy of more than 24 months who would need to requalify with respect to the indices of disease severity. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each biological medicine trialled.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent, unless they have experienced a serious adverse reaction of a severity necessitating permanent treatment withdrawal.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug.

Abatacept:

A patient swapping from I.V. abatacept to subcutaneous abatacept will not be eligible for an I.V. loading dose when commencing treatment with the subcutaneous formulation.

Rituximab:

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF-alfa antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they be assessed for response to every course of treatment, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological medicine during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological medicine therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate biological medicine may do so without having to have any treatment-free period.

(3) Baseline measurements to determine response.

Determination of whether a response to treatment has been demonstrated must be based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and the

eligibility for continuing treatment must be assessed according to these revised baseline measurements. To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response must be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be determined on the total number of major joints.

Applications under the Initial 1 treatment restriction for a new patient must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. The results must be no more than 4 weeks old at the time of application.

Applications under the Initial 3 treatment restriction for re-commencement of treatment after a break in biological medicine of more than 24 months must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with disease modifying anti-rheumatic drugs (DMARDs) which must include at least 3 months continuous treatment with each of at least 2 DMARDs, one of which must be methotrexate at a dose of at least 20 mg weekly and one of which must be: (i) hydroxychloroquine at a dose of at least 200 mg daily; or (ii) leflunomide at a dose of at least 10 mg daily; or (iii) sulfasalazine at a dose of at least 2 g daily; OR
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs which, if methotrexate is contraindicated according to the Therapeutic Goods Administration (TGA)-approved Product Information or cannot be tolerated at a 20 mg weekly dose, must include at least 3 months continuous treatment with each of at least 2 of the following DMARDs: (i) hydroxychloroquine at a dose of at least 200 mg daily; and/or (ii) leflunomide at a dose of at least 10 mg daily; and/or (iii) sulfasalazine at a dose of at least 2 g daily; OR
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs which, if 3 or more of methotrexate, hydroxychloroquine, leflunomide and sulfasalazine are contraindicated according to the relevant TGA-approved Product Information or cannot be tolerated at the doses specified above, must include at least 3 months continuous treatment with each of at least 2 DMARDs, with one or more of the following DMARDs being used in place of the DMARDs which are contraindicated or not tolerated: (i) azathioprine at a dose of at least 1 mg/kg per day; and/or (ii) cyclosporin at a dose of at least 2 mg/kg/day; and/or (iii) sodium aurothiomalate at a dose of 50 mg weekly, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

The application must include details of the DMARDs trialled, their doses and duration of treatment, and all relevant contraindications and/or intolerances including severity.

The requirement to trial at least 2 DMARDs for periods of at least 3 months each can be met using single agents sequentially or by using one or more combinations of DMARDs.

If the requirement to trial 6 months of intensive DMARD therapy with at least 2 DMARDs cannot be met because of contraindications and/or intolerances of a severity necessitating permanent treatment withdrawal to all of the DMARDs specified above, details of the contraindication or intolerance including severity and dose for each DMARD must be provided in the authority application.

The following criteria indicate failure to achieve an adequate response to DMARD treatment and must be demonstrated in all patients at the time of the initial application:

an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour and/or a C-reactive protein (CRP) level greater than 15 mg per L; **AND** either

(a) a total active joint count of at least 20 active (swollen and tender) joints; or

(b) at least 4 active joints from the following list of major joints:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The joint count and ESR and/or CRP must be determined at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. All measures must be no more than one month old at the time of initial application.

If the requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied. Treatment with prednisolone dosed at 7.5 mg or higher daily (or equivalent) or a parenteral steroid within the past month (intramuscular or intravenous methylprednisolone or equivalent) is an acceptable reason.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response must be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number

of major joints, the response must be determined on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker must be used to determine response.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note The Services Australia website (www.servicesaustralia.gov.au) has details of the toxicities, including severity, which will be accepted for the following purposes:

- (a) exempting a patient from the requirement to undertake a minimum 3 month trial of methotrexate at a 20 mg weekly dose;
- (b) substituting azathioprine, cyclosporin or sodium aurothiomalate for another DMARD as part of the 6 month intensive DMARD trial;
- (c) exempting a patient from the requirement for a 6 month trial of intensive DMARD therapy.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 2 (change or re-commencement of treatment after a break in biological medicine of less than 24 months)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must not have failed to respond to previous PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised biological medicine treatment for this condition 5 times, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

- (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
- (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, conducted within the timeframes specified below.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response must be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be determined on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker must be used to determine response.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

A patient who has demonstrated a response to a course of rituximab must have a PBS-subsidised biological therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 3 (re-commencement of treatment after a break in biological medicine of more than 24 months)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 24 months or more from the most recent PBS-subsidised biological medicine for this condition, **AND**
- Patient must not have failed to respond to previous PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised biological medicine treatment for this condition 5 times, **AND**
- The condition must have an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; OR
- The condition must have a C-reactive protein (CRP) level greater than 15 mg per L, **AND**
- The condition must have either (a) a total active joint count of at least 20 active (swollen and tender) joints; or (b) at least 4 active major joints, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Major joints are defined as (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

All measures of joint count, ESR and/or CRP must be no more than 4 weeks old at the time of application.

If the requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied. Treatment with prednisolone dosed at 7.5 mg or higher daily (or equivalent) or a parenteral steroid within the past month (intramuscular or intravenous methylprednisolone or equivalent) is an acceptable reason.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response must be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be determined on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker must be used to determine response.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial 1 (new patient) or Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 24 months) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 24 months) - balance of supply

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 24 months) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 24 months) to complete 16 weeks of treatment, **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

upadacitinib 15 mg modified release tablet, 28

11989B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	1271.34	41.30	Rinvoq [VE]

Tumor necrosis factor alpha (TNF-alpha) inhibitors

■ ADALIMUMAB

Note TREATMENT OF PAEDIATRIC PATIENTS WITH REFRACTORY CROHN DISEASE

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) for paediatric patients with adalimumab for severe refractory Crohn disease and infliximab for moderate to severe refractory Crohn disease. Where the term "biological medicines" appears in the following NOTES and restrictions, it refers to adalimumab and infliximab only. A patient is eligible for PBS-subsidised treatment with only one PBS-subsidised biological medicine at any one time. For paediatric patients with Crohn disease, infliximab is PBS-subsidised for moderate to severe disease while adalimumab is PBS-subsidised for severe disease.

From 1 August 2015, under the PBS, patients commencing on adalimumab will be able to commence a treatment cycle where they may trial each PBS-subsidised biological medicine without having to experience a disease flare when swapping to infliximab. Patients on infliximab will be able to commence a treatment cycle where they may trial each PBS-subsidised biological medicine but will need to meet a PCDAI score of greater than or equal to 40 when swapping to adalimumab.

Under these arrangements, within a single treatment cycle and depending on the disease severity, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy.

A patient who received PBS-subsidised biological medicine treatment prior to 1 August 2015 is considered to have started their treatment cycle as of 1 August 2015.

Within the same treatment cycle, a paediatric patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than twice.

Once a patient has either failed, or ceased to respond to treatment for this condition 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy for this condition before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle.

A patient who has failed 3 trials or fewer of biological medicine therapy in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 August 2015.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised biological medicine therapy in this treatment cycle and wishes to commence such therapy - Initial 1 (new patient); or

(ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an

alternate agent - Initial 2 (Change or Re-commencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or
(iii) a patient wishes to re-commence treatment with a specific biological medicine following a break in PBS-subsidised therapy with that agent - Initial 2 (Change or Re-commencement of treatment after a break in therapy of less than 5 years); or

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years - Initial 3 (Recommencement of treatment after a break in biological medicine of more than 5 years). Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy for adalimumab and 14 weeks of therapy for infliximab.

From 1 August 2015, a patient must be assessed for response to any course of initial PBS-subsidised biological therapy following a minimum of 12 weeks of therapy for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab, and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

For first and subsequent continuing courses of PBS-subsidised biological medicine therapy, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

Adalimumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats and the second prescription should be written for 2 doses of 40 mg and 2 repeats for patients weighing 40 kg or greater. For patients weighing less than 40 kg, one prescription should be written for 2 doses of 40 mg with no repeats and the second prescription should be written for 2 doses of 20 mg with 3 repeats.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient remains eligible to receive up to 24 weeks per course of continuing treatment under the First continuing treatment and Subsequent Continuing treatment restrictions with that drug providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted supply of treatment.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be conducted within 1 month of the last dose. Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine therapy is approved, a patient with severe disease may swap if eligible to the alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Paediatric Crohn Disease Activity Index (PCDAI) Score, confirmation of Crohn disease), or the prior conventional therapies of corticosteroid therapy, immunosuppressive therapy or enteral nutrition. Patients on infliximab may swap to adalimumab within the same treatment cycle provided that their disease severity has progressed to severe disease (i.e. they have a current PCDAI score of 40 or more).

A patient cannot swap to a biological medicine if they have failed to respond to prior treatment with that drug twice within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment within the timeframes specified in the relevant restriction.

A patient who is not able to complete an initial treatment course for a biological medicine will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the PCDAI submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used to assess response to all subsequent treatments.

(4) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to recommence treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity under the Initial 3 restriction. A retrieval of conventional therapies is not required.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe Crohn disease

Treatment Phase: Continuing treatment of Crohn disease in a paediatric patient assessed by PCDAI

Clinical criteria:

- Patient must have a documented history of severe Crohn disease, **AND**
- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have a reduction in PCDAI Score by at least 15 points from baseline value, **AND**
- Patient must have a total PCDAI score of 40 points or less, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 6 to 17 years inclusive.

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician; OR
- Must be treated by a specialist paediatric gastroenterologist.

Applications for authorisation must be made in writing and must include:

- a completed authority prescription form; and
- a completed Paediatric Crohn Disease PBS Authority Application - Supporting Information Form, which includes the completed Paediatric Crohn Disease Activity Index (PCDAI) calculation sheet along with the date of the assessment of the patient's condition.

The PCDAI assessment must be no more than 1 month old at the time of application.

An application for continuing treatment with this drug must include a measurement of response to the most recent course of PBS-subsidised therapy. This assessment must be conducted no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment with this drug, it must be accompanied by an assessment of response to a minimum of 12 weeks of treatment with the initial treatment course.

The assessment of the patient's response to a continuing course of therapy must be made within the 4 weeks prior to completion of that course and posted to the Department of Human Services no less than 2 weeks prior to the date the next dose is scheduled, in order to ensure continuity of treatment for those patients who meet the continuation criterion.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Patients are only eligible to receive subsequent continuing PBS-subsidised treatment with this drug in courses of up to 24 weeks providing they continue to sustain the response.

Where fewer than 5 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment with this drug may be requested through the balance of supply restriction.

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL syringes

10412B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1092.41	41.30	Humira [VE]

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL pen devices

10420K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1092.41	41.30	Humira [VE]

adalimumab 20 mg/0.4 mL injection, 2 x 0.4 mL syringes

10396E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1092.41	41.30	Humira [VE]

▪ **ADALIMUMAB**

Note TREATMENT OF PAEDIATRIC PATIENTS WITH REFRACTORY CROHN DISEASE

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) for paediatric patients with adalimumab for severe refractory Crohn disease and infliximab for moderate to severe refractory Crohn disease. Where the term "biological medicines" appears in the following NOTES and restrictions, it refers to adalimumab and infliximab only. A patient is eligible for PBS-subsidised treatment with only one PBS-subsidised biological medicine at any one time. For paediatric patients with Crohn disease, infliximab is PBS-subsidised for moderate to severe disease while adalimumab is PBS-subsidised for severe disease.

From 1 August 2015, under the PBS, patients commencing on adalimumab will be able to commence a treatment cycle where they may trial each PBS-subsidised biological medicine without having to experience a disease flare when swapping to infliximab. Patients on infliximab will be able to commence a treatment cycle where they may trial each PBS-subsidised biological medicine but will need to meet a PCDAI score of greater than or equal to 40 when swapping to adalimumab.

Under these arrangements, within a single treatment cycle and depending on the disease severity, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy.

A patient who received PBS-subsidised biological medicine treatment prior to 1 August 2015 is considered to have started their treatment cycle as of 1 August 2015.

Within the same treatment cycle, a paediatric patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than twice.

Once a patient has either failed, or ceased to respond to treatment for this condition 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine

therapy for this condition before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle.

A patient who has failed 3 trials or fewer of biological medicine therapy in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 August 2015.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised biological medicine therapy in this treatment cycle and wishes to commence such therapy - Initial 1 (new patient); or

(ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent - Initial 2 (Change or Re-commencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to re-commence treatment with a specific biological medicine following a break in PBS-subsidised therapy with that agent - Initial 2 (Change or Re-commencement of treatment after a break in therapy of less than 5 years); or

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years - Initial 3 (Recommencement of treatment after a break in biological medicine of more than 5 years). Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy for adalimumab and 14 weeks of therapy for infliximab.

From 1 August 2015, a patient must be assessed for response to any course of initial PBS-subsidised biological therapy following a minimum of 12 weeks of therapy for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab, and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

For first and subsequent continuing courses of PBS-subsidised biological medicine therapy, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

Adalimumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats and the second prescription should be written for 2 doses of 40 mg and 2 repeats for patients weighing 40 kg or greater. For patients weighing less than 40 kg, one prescription should be written for 2 doses of 40 mg with no repeats and the second prescription should be written for 2 doses of 20 mg with 3 repeats.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient remains eligible to receive up to 24 weeks per course of continuing treatment under the First continuing treatment and Subsequent Continuing treatment restrictions with that drug providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted supply of treatment.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be conducted within 1 month of the last dose. Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine therapy is approved, a patient with severe disease may swap if eligible to the alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Paediatric Crohn Disease Activity Index (PCDAI) Score, confirmation of Crohn disease), or the prior conventional therapies of corticosteroid therapy, immunosuppressive therapy or enteral nutrition. Patients on infliximab may swap to adalimumab within the same treatment cycle provided that their disease severity has progressed to severe disease (i.e. they have a current PCDAI score of 40 or more).

A patient cannot swap to a biological medicine if they have failed to respond to prior treatment with that drug twice within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment within the timeframes specified in the relevant restriction.

A patient who is not able to complete an initial treatment course for a biological medicine will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the PCDAI submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used to assess response to all subsequent treatments.

(4) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to recommence treatment following a break in PBS-subsidised biological medicine therapy of at least

5 years, must requalify for initial treatment with respect to the indices of disease severity under the Initial 3 restriction. A retreatal of conventional therapies is not required.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Severe Crohn disease

Treatment Phase: Balance of supply for paediatric patient

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the continuing treatment restriction to complete 24 weeks of treatment, **AND**
- The treatment must provide no more than the balance of up to 16 weeks therapy available under Initial 1, 2 or 3 treatment; OR
- The treatment must provide no more than the balance of up to 24 weeks therapy available under Continuing treatment.

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician; OR
- Must be treated by a specialist paediatric gastroenterologist.

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL syringes

10399H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	1092.41	41.30	Humira [VE]

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL pen devices

10400J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	1092.41	41.30	Humira [VE]

adalimumab 20 mg/0.4 mL injection, 2 x 0.4 mL syringes

10422M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	1092.41	41.30	Humira [VE]

■ ADALIMUMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CROHN DISEASE

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for adult patients with severe Crohn disease. Where the term 'biological medicine' appears in the following NOTES and restrictions, it refers to the tumour necrosis factor (TNF) alfa-antagonists (adalimumab and infliximab), the alpha-4 beta-7 integrin inhibitor (vedolizumab) and the human IgG1kappa monoclonal antibody (ustekinumab). Patients are eligible for PBS-subsidised treatment with only 1 of the above PBS-subsidised biological medicines at any one time.

From 1 September 2017, under the PBS, all patients will be able to commence a treatment cycle where they may trial PBS-subsidised a biological medicine without having to experience a disease flare when swapping to the alternate agent. Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy.

A patient who received PBS-subsidised treatment with a biological medicine prior to 1 September 2017 is considered to have started their treatment cycle as of 1 September 2017.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once.

Once a patient has either failed or ceased to respond to treatment for this condition 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy for this condition before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of less than 5 years, may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab therapy after 1 September 2017.

(a) Initial treatment.

Applications for initial treatment should be made where:

- a patient has received no prior PBS-subsidised biological medicine treatment in this treatment cycle and wishes to commence such therapy Initial 1 (new patient); or
- a patient has received prior PBS-subsidised (initial or continuing) biological medicine and wishes to trial an alternate

agent - Initial 2 (change or recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy with that agent - Initial 2 (change or recommencement of treatment after a break in therapy of less than 5 years); or
(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years - Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years).

From 1 September 2017, a patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy for adalimumab or ustekinumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab or vedolizumab, and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

For subsequent courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

Adalimumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats and the second prescription should be written for 2 doses of 40 mg and 2 repeats for patients weighing 40 kg or greater. For patients weighing less than 40 kg, one prescription should be written for 2 doses of 40 mg with no repeats and the second prescription should be written for 2 doses of 20 mg with 3 repeats.

Ustekinumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be written under S100 (Highly Specialised Drugs) for a weight-based loading dose, containing a quantity of up to 4 vials of 130 mg and no repeats. The second prescription should be written under S85 (General) for the subsequent first dose, containing a quantity of 2 vials of 45 mg and no repeats.

(b) Continuing treatment.

Following the completion of an initial treatment course with a biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted supply of treatment.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be conducted within 4 weeks of the last dose. Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine therapy is approved, a patient may swap if eligible to the alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Crohn Disease Activity Index (CDAI) Score, confirmation of Crohn disease), or the prior conventional therapies of corticosteroid therapy and immunosuppressive therapy.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug once within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment, within the timeframes specified in the relevant restriction.

A patient who is not able to complete an initial treatment course for a biological medicine will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the CDAI or evidence of intestinal inflammation submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used to assess response to all subsequent treatments.

(4) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to recommence treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity under the Initial 3 restriction. A re-trial of systemic therapy is not required.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe Crohn disease

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have an adequate response to this drug defined as a reduction in Crohn Disease Activity Index (CDAI) Score to a level no greater than 150 if assessed by CDAI or if affected by extensive small intestine disease; OR
- Patient must have an adequate response to this drug defined as (a) an improvement of intestinal inflammation as demonstrated by: (i) blood: normalisation of the platelet count, or an erythrocyte sedimentation rate (ESR) level no greater than 25 mm per hour, or a C-reactive protein (CRP) level no greater than 15 mg per L; or (ii) faeces: normalisation of lactoferrin or calprotectin level; or (iii) evidence of mucosal healing, as demonstrated by diagnostic imaging findings, compared to the baseline assessment; or (b) reversal of high faecal output state; or (c) avoidance of the need for surgery or total parenteral nutrition (TPN), if affected by short gut syndrome, extensive small intestine or is an ostomy patient, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Applications for authorisation must be made in writing and must include:

(a) a completed authority prescription form; and

(b) a completed Crohn Disease PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed Crohn Disease Activity Index (CDAI) Score calculation sheet including the date of the assessment of the patient's condition, if relevant; or

(ii) the reports and dates of the pathology test or diagnostic imaging test(s) used to assess response to therapy for patients with short gut syndrome, extensive small intestine disease or an ostomy, if relevant; and

(iii) the date of clinical assessment.

All assessments, pathology tests, and diagnostic imaging studies must be made within 1 month of the date of application.

An application for continuing treatment with this drug must include a measurement of response to the most recent course of PBS-subsidised therapy. This assessment must be conducted no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment with this drug, it must be accompanied by an assessment of response to a minimum of 12 weeks of treatment with the initial treatment course.

The assessment of the patient's response to a continuing course of therapy must be made within the 4 weeks prior to completion of that course and posted to the Department of Human Services no less than 2 weeks prior to the date the next dose is scheduled, in order to ensure continuity of treatment for those patients who meet the continuation criterion.

Where an assessment is not submitted to the Department of Human Services within these timeframes, patients will be deemed to have failed to respond, or to have failed to sustain a response, to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Patients are eligible to receive continuing treatment with this drug in courses of up to 24 weeks providing they continue to sustain a response.

At the time of the authority application, medical practitioners should request the appropriate quantity and number of repeats to provide sufficient dose. Up to a maximum of 5 repeats will be authorised.

Where fewer than 5 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 24 weeks of treatment with this drug may be requested through the balance of supply restriction.

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL syringes

9189P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1092.41	41.30	Humira [VE]

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL pen devices

9191R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1092.41	41.30	Humira [VE]

■ ADALIMUMAB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Moderate to severe hidradenitis suppurativa

Treatment Phase: Initial treatment 1 - New patient or Initial treatment 2 - Recommencement of treatment – balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial treatment 1 - New patient restriction to complete a maximum of 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial treatment 2 - Recommencement of treatment restriction to complete a maximum of 16 weeks treatment.

Treatment criteria:

- Must be treated by a dermatologist.
- A maximum of 12 weeks of treatment will be authorised under this restriction.

adalimumab 40 mg/0.8 mL injection, 4 x 0.8 mL pen devices

11133Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	2120.28	41.30	Humira [VE]

■ ADALIMUMAB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Moderate to severe hidradenitis suppurativa

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have demonstrated a response to treatment with this drug for this condition.

Treatment criteria:

- Must be treated by a dermatologist.

A response to treatment is defined as:

Achieving Hidradenitis Suppurativa Clinical Response (HiSCR) of a 50% reduction in AN count compared to baseline with no increase in abscesses or draining fistulae.

For the first application for continuing treatment a Hidradenitis Suppurativa Clinical Response (HiSCR) assessment must be made following a minimum of 12 weeks of treatment. For subsequent continuing treatment a HiSCR assessment must be made every 24 weeks.

The assessment of the patient's response to a continuing course of therapy must be made within the 4 weeks prior to completion of that course and must be provided no less than 2 weeks prior to the date the next dose is scheduled, in order to ensure continuity of treatment for those patients who meet the continuation criterion.

Where an assessment is not submitted within these timeframes, patients will be deemed to have failed to respond, or to have failed to sustain a response, to treatment with this drug.

A maximum of 24 weeks treatment will be authorised under this restriction per continuing treatment.

The authority application must be made in writing and must include:

- a completed authority prescription form; and
- a completed hidradenitis suppurativa PBS authority application supporting Information form which must include the Hidradenitis Suppurativa Clinical Response (HiSCR) result.

adalimumab 40 mg/0.8 mL injection, 4 x 0.8 mL pen devices

11137E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	2120.28	41.30	Humira [VE]

■ ADALIMUMAB**Note TREATMENT OF PAEDIATRIC PATIENTS WITH REFRACTORY CROHN DISEASE**

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) for paediatric patients with adalimumab for severe refractory Crohn disease and infliximab for moderate to severe refractory Crohn disease. Where the term "biological medicines" appears in the following NOTES and restrictions, it refers to adalimumab and infliximab only. A patient is eligible for PBS-subsidised treatment with only one PBS-subsidised biological medicine at any one time. For paediatric patients with Crohn disease, infliximab is PBS-subsidised for moderate to severe disease while adalimumab is PBS-subsidised for severe disease.

From 1 August 2015, under the PBS, patients commencing on adalimumab will be able to commence a treatment cycle where they may trial each PBS-subsidised biological medicine without having to experience a disease flare when swapping to infliximab. Patients on infliximab will be able to commence a treatment cycle where they may trial each PBS-subsidised biological medicine but will need to meet a PCDAI score of greater than or equal to 40 when swapping to adalimumab. Under these arrangements, within a single treatment cycle and depending on the disease severity, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy.

A patient who received PBS-subsidised biological medicine treatment prior to 1 August 2015 is considered to have started their treatment cycle as of 1 August 2015.

Within the same treatment cycle, a paediatric patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than twice.

Once a patient has either failed, or ceased to respond to treatment for this condition 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy for this condition before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle.

A patient who has failed 3 trials or fewer of biological medicine therapy in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 August 2015.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised biological medicine therapy in this treatment cycle and wishes to commence such therapy - Initial 1 (new patient); or

(ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent - Initial 2 (Change or Re-commencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to re-commence treatment with a specific biological medicine following a break in PBS-subsidised therapy with that agent - Initial 2 (Change or Re-commencement of treatment after a break in therapy of less than 5 years); or

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years - Initial 3 (Recommencement of treatment after a break in biological medicine of more than 5 years).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy for adalimumab and 14 weeks of therapy for infliximab.

From 1 August 2015, a patient must be assessed for response to any course of initial PBS-subsidised biological therapy following a minimum of 12 weeks of therapy for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab, and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

For first and subsequent continuing courses of PBS-subsidised biological medicine therapy, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

Adalimumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats and the second prescription should be written for 2 doses of 40 mg and 2 repeats for patients weighing 40 kg or greater. For patients weighing less than 40 kg, one prescription should be written for 2 doses of 40 mg with no repeats and the second prescription should be written for 2 doses of 20 mg with 3 repeats.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient remains eligible to receive up to 24 weeks per course of continuing treatment under the First continuing treatment and Subsequent Continuing treatment restrictions with that drug providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted supply of treatment.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be conducted within 1 month of the last dose. Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine therapy is approved, a patient with severe disease may swap if eligible to the alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Paediatric Crohn Disease Activity Index (PCDAI) Score, confirmation of Crohn disease), or the prior conventional therapies of corticosteroid therapy, immunosuppressive therapy or enteral nutrition.

Patients on infliximab may swap to adalimumab within the same treatment cycle provided that their disease severity has progressed to severe disease (i.e. they have a current PCDAI score of 40 or more).

A patient cannot swap to a biological medicine if they have failed to respond to prior treatment with that drug twice within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment within the timeframes specified in the relevant restriction.

A patient who is not able to complete an initial treatment course for a biological medicine will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the PCDAI submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to these revised baseline

measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used to assess response to all subsequent treatments.

(4) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to recommence treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity under the Initial 3 restriction. A retreatal of conventional therapies is not required.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe Crohn disease

Treatment Phase: Initial treatment - Initial 1 (new patient)

Clinical criteria:

- Patient must have confirmed diagnosis of Crohn disease, defined by standard clinical, endoscopic and/or imaging features including histological evidence, **AND**
- Patient must have failed to achieve an adequate response to 2 of the following 3 conventional prior therapies including: (i) a tapered course of steroids, starting at a dose of at least 1 mg per kg or 40 mg (whichever is the lesser) prednisolone (or equivalent), over a 6 week period; (ii) an 8 week course of enteral nutrition; or (iii) immunosuppressive therapy including azathioprine at a dose of at least 2 mg per kg daily for 3 or more months, or, 6-mercaptopurine at a dose of at least 1 mg per kg daily for 3 or more months, or, methotrexate at a dose of at least 10 mg per square metre weekly for 3 or more months; **OR**
- Patient must have a documented intolerance of a severity necessitating permanent treatment withdrawal or a contra-indication to each of prednisolone (or equivalent), azathioprine, 6-mercaptopurine and methotrexate, **AND**
- Patient must have, at the time of application, disease severity considered to be severe as demonstrated by a Paediatric Crohn Disease Activity Index (PCDAI) Score greater than or equal to 40 preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior conventional treatment and which is no more than 1 month old at the time of application, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 6 to 17 years inclusive.

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); **OR**
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; **OR**
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)]; **OR**
- Must be treated by a paediatrician; **OR**
- Must be treated by a specialist paediatric gastroenterologist.

Applications for authorisation of initial treatment must be in writing and must include:

- two completed authority prescription forms; and
- a completed paediatric Crohn Disease PBS Authority Application -Supporting Information Form [may be downloaded from the Department of Human Services website (www.humanservices.gov.au)] which includes the following:
 - the completed current Paediatric Crohn Disease Activity Index (PCDAI) calculation sheet including the date of assessment of the patient's condition; and
 - details of previous systemic drug therapy [dosage, date of commencement and duration of therapy] or dates of enteral nutrition.

If treatment with any of the specified prior conventional drugs is contraindicated according to the relevant TGA-approved Product Information, please provide details at the time of application. If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, please provide details of the degree of this toxicity at the time of application. Details of the accepted toxicities including severity can be found on the Human Services website (www.humanservices.gov.au).

A maximum quantity and number of repeats to provide for an initial 16 week course of this drug consisting of a 160 mg dose at week 0, 80 mg dose at week 2 and 40 mg dose at weeks 4, 6, 8, 10, 12 and 14 for patients 40 kg or greater (for patients 40 kg or less, the course is a 80 mg dose at week 0, 40 mg dose at week 2 and a 20 mg dose at weeks 4, 6, 8, 10, 12 and 14) will be authorised.

Two completed authority prescriptions should be submitted with every initial application for this drug. For patients weighing 40 kg or greater: one prescription should be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats and the second prescription should be written for 2 doses of 40 mg and 2 repeats. For patients weighing less than 40 kg: one prescription should be written for 2 doses of 40 mg with no repeats and the second prescription should be written for 2 doses of 20 mg with 3 repeats.

If fewer than 2 repeats (for patients 40 kg or greater) or 3 repeats (for patients less than 40 kg) are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment with this drug may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8

a.m. to 5 p.m. EST Monday to Friday) and authorised through the Balance of Supply treatment phase PBS restriction. Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period.

A PCDAI assessment of the patient's response to this initial course of treatment must be made following a minimum of 12 weeks therapy so that there is adequate time for a response to be demonstrated.

This assessment, which will be used to determine eligibility for the first continuing treatment, must be submitted to the Department of Human Services no later than 1 month from the date of completion of this initial course of treatment.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Authority required

Severe Crohn disease

Treatment Phase: Initial treatment - Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have a documented history of severe Crohn disease, **AND**
- Patient must in this treatment cycle, have received prior PBS-subsidised treatment with this drug for this condition; OR
- Patient must in this treatment cycle, have received prior PBS-subsidised treatment with infliximab for this condition and have a current PCDAI score of 40 or greater, **AND**
- Patient must not have failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition more than once in the current treatment cycle, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 6 to 17 years inclusive.

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician; OR
- Must be treated by a specialist paediatric gastroenterologist.

Applications for authorisation of initial treatment must be in writing and must include:

(a) two completed authority prescription forms; and

(b) a completed paediatric Crohn Disease PBS Authority Application -Supporting Information Form [may be downloaded from the Department of Human Services website (www.humanservices.gov.au)] which includes the following:

- (i) the completed current Paediatric Crohn Disease Activity Index (PCDAI) Score calculation sheet; and
- (ii) details of prior TNF-alfa antagonist treatment including details of date and duration of treatment.

The PCDAI assessment must be no more than 1 month old at the time of application.

Two completed authority prescriptions should be submitted with every initial application for this drug. For patients weighing 40 kg or greater: one prescription should be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats and the second prescription should be written for 2 doses of 40 mg and 2 repeats. For patients weighing less than 40 kg: one prescription should be written for 2 doses of 40 mg with no repeats and the second prescription should be written for 2 doses of 20 mg with 3 repeats.

If fewer than 2 repeats (for patients 40 kg or greater) or 3 repeats (for patients less than 40 kg) are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment with this drug may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) and authorised through the Balance of Supply treatment phase PBS restriction. Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to change or recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under an initial treatment restriction, the patient must have been assessed for response to that course following a minimum of 12 weeks therapy for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

If the response assessment to the previous course of biological medicine treatment is not submitted as detailed above, the patient will be deemed to have failed therapy with that particular course of biological medicine.

A PCDAI assessment of the patient's response to this initial course of treatment must be made following a minimum of 12 weeks therapy so that there is adequate time for a response to be demonstrated.

This assessment, which will be used to determine eligibility for the first continuing treatment, must be submitted to the Department of Human Services no later than 1 month from the date of completion of this initial course of treatment.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Authority required

Severe Crohn disease

Treatment Phase: Initial treatment - Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- Patient must have confirmed Crohn disease, defined by standard clinical, endoscopic and/or imaging features, including histological evidence, with the diagnosis confirmed by a gastroenterologist, consultant physician, paediatrician or specialist paediatric gastroenterologist, **AND**
- Patient must have, at the time of application, disease severity considered to be severe as demonstrated by a Paediatric Crohn Disease Activity Index (PCDAI) Score greater than or equal to 40, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 6 to 17 years inclusive.

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician; OR
- Must be treated by a specialist paediatric gastroenterologist.

Applications for authorisation must be made in writing and must include:

(a) two completed authority prescription forms; and

(b) a completed paediatric Crohn Disease PBS Authority Application - Supporting Information Form [may be downloaded from the Department of Human Services website (www.humanservices.gov.au)] which includes the following:

(i) the completed current Paediatric Crohn Disease Activity Index (PCDAI) calculation sheet including the date of assessment of the patient's condition.

The PCDAI assessment must be no more than 1 month old at the time of application.

A maximum quantity and number of repeats to provide for an initial 16 week course of this drug consisting of a 160 mg dose at week 0, 80 mg dose at week 2 and 40 mg dose at weeks 4, 6, 8, 10, 12 and 14 for patients 40 kg or greater (for patients 40 kg or less, the course is a 80 mg dose at week 0, 40 mg dose at week 2 and a 20 mg dose at weeks 4, 6, 8, 10, 12 and 14) will be authorised.

Two completed authority prescriptions should be submitted with every initial application for this drug. For patients weighing 40 kg or greater: one prescription should be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats and the second prescription should be written for 2 doses of 40 mg and 2 repeats. For patients weighing less than 40 kg: one prescription should be written for 2 doses of 40 mg with no repeats and the second prescription should be written for 2 doses of 20 mg with 3 repeats.

If fewer than 2 repeats (for patients 40 kg or greater) or 3 repeats (for patients less than 40 kg) are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment with this drug may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday) and authorised through the Balance of Supply treatment phase PBS restriction. Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period.

A PCDAI assessment of the patient's response to this initial course of treatment must be made following a minimum of 12 weeks therapy so that there is adequate time for a response to be demonstrated.

This assessment, which will be used to determine eligibility for the first continuing treatment, must be submitted to the Department of Human Services no later than 1 month from the date of completion of this initial course of treatment.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

adalimumab 40 mg/0.8 mL injection, 6 x 0.8 mL pen devices

10397F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	3099.84	41.30	Humira [VE]

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL syringes

10419J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	1092.41	41.30	Humira [VE]

adalimumab 40 mg/0.8 mL injection, 6 x 0.8 mL syringes

10404N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	3099.84	41.30	Humira [VE]

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL pen devices

10413C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	1092.41	41.30	Humira [VE]

adalimumab 20 mg/0.4 mL injection, 2 x 0.4 mL syringes

10389T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	1092.41	41.30	Humira [VE]

■ **ADALIMUMAB**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Moderate to severe hidradenitis suppurativa

Treatment Phase: Initial treatment 1 - New patient

Clinical criteria:

- Patient must have, at the time of application, a Hurley stage II or III grading with an abscess and inflammatory nodule (AN) count greater than or equal to 3, **AND**
- Patient must have failed to achieve an adequate response to 2 courses of different antibiotics each for 3 months prior to initiation of PBS subsidised treatment with this drug for this condition; OR
- Patient must have had an adverse reaction to an antibiotic of a severity necessitating permanent treatment withdrawal resulting in the patient being unable to complete treatment with 2 different courses of antibiotics each for 3 months prior to initiation of PBS-subsidised treatment with this drug for this condition; OR
- Patient must be contraindicated to treatment with an antibiotic due to an allergic reaction of a severity necessitating permanent treatment withdrawal resulting in the patient being unable to complete treatment with 2 different courses of antibiotics each for 3 months prior to initiation of PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be limited to a maximum duration of 16 weeks.

Treatment criteria:

- Must be treated by a dermatologist.

Assessment of disease severity must be no more than 1 month old at the time of application.

An assessment of the patient's response to this recommencement course of treatment must be made following a minimum of 12 weeks of treatment.

At the time of authority application the prescriber must request the first 4 weeks of treatment under this restriction; and weeks 5 to 16 of treatment under Initial treatment 1 - New patient or Initial treatment 2 - Recommencement of treatment - balance of supply

The authority application must be made in writing and must include:

- a completed authority prescription form; and
- a completed hidradenitis suppurativa PBS authority application supporting Information form which must include:
 - the Hurley stage grading; and
 - the AN count; and
 - the name of the antibiotic/s received for two separate courses each of three months; or
 - confirmation that the adverse reaction or allergy to an antibiotic necessitated permanent treatment withdrawal resulting in the patient being unable to complete a three month course of antibiotics. The name of the one course of antibiotics of three months duration must be provided. Where the patient is unable to be treated with any courses of antibiotics the prescriber must confirm that the patient has a history of adverse reaction or allergy necessitating permanent treatment withdrawal to two different antibiotics.

Authority required

Moderate to severe hidradenitis suppurativa

Treatment Phase: Initial treatment 2 - Recommencement of treatment

Clinical criteria:

- Patient must have, at the time of application, a Hurley stage II or III grading with an abscess and inflammatory nodule (AN) count greater than or equal to 3, **AND**
- Patient must have demonstrated a response to the most recent PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be limited to a maximum duration of 16 weeks.

Treatment criteria:

- Must be treated by a dermatologist.

Assessment of disease severity must be no more than 1 month old at the time of application.

A response to treatment is defined as:

Achieving Hidradenitis Suppurativa Clinical Response (HiSCR) of a 50% reduction in AN count compared to baseline with no increase in abscesses or draining fistulae.

An assessment of the patient's response to this recommencement course of treatment must be made following a minimum of 12 weeks of treatment.

At the time of authority application the prescriber must request the first 4 weeks of treatment under this restriction; and weeks 5 to 16 of treatment under Initial treatment 1 - New patient or Initial treatment 2 - Recommencement of treatment - balance of supply

The authority application must be made in writing and must include:

- a completed authority prescription form; and
- a completed hidradenitis suppurativa PBS authority application supporting Information form which must include:

- (i) the Hurley stage grading; and
- (ii) the AN count.

adalimumab 40 mg/0.8 mL injection, 6 x 0.8 mL pen devices

11132X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	3099.84	41.30	Humira [VE]

ADALIMUMAB

Note TREATMENT OF ADULT PATIENTS WITH A HISTORY OF JUVENILE IDIOPATHIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, etanercept and tocilizumab for a patient over 18 years who has a history of juvenile idiopathic arthritis with onset prior to the age of 18 years. Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, etanercept and tocilizumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 3 biological medicines at any one time.

From 1 April 2014, a patient receiving PBS-subsidised biological medicine therapy is considered to be in a treatment cycle where they may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements, within a single treatment cycle, a patient may:

(i) continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy; and

(ii) fail to respond or to sustain a response to each PBS-subsidised biological medicine once only. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment.

Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single treatment cycle and they must have, at a minimum, a 5 year break in PBS-subsidised biological medicine therapy before they are eligible to receive further PBS-subsidised biological medicine therapy.

The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine was approved to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who received PBS-subsidised biological medicine treatment immediately prior to 1 April 2014 is considered to be in their first cycle as of 1 April 2014. A patient who has had a break in biological medicine treatment of at least 24 months immediately prior to making a new application, on or after 1 April 2014, will commence a new treatment cycle under the Initial 3 treatment restriction.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of less than 24 months may commence a further course of treatment within the same treatment cycle under the Initial 2 treatment restriction.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of more than 24 months must commence a new treatment cycle under the Initial 3 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine therapy after 1 April 2014.

(1) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an alternate medicine (Initial 2 - Change or recommencement of treatment after a break in biological medicine of less than 24 months) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 24 months with the same medicine (Initial 2 - Change or recommencement of treatment after a break in biological medicine therapy of less than 24 months); or

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 24 months (Initial 3 - recommencement of treatment after a break in biological medicine of more than 24 months). Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy.

A patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not conducted within these timeframes, the patient will be deemed to have failed this course of treatment.

For second and subsequent cycles of PBS-subsidised biological medicine, it is recommended that a patient is reviewed in the 4 weeks prior to completing their current course of treatment and that an application is posted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services where applicable. Where a response assessment is not submitted, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (joint count and ESR/CRP) or the prior non- biological medicine therapy requirements, except if the patient has had a break in therapy of more than 24 months who would then need to requalify under the Initial 3 restrictions with respect to the indices of disease severity. A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle. To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, an application for a patient who wishes to swap to an alternate biological medicine should be accompanied by the approved authority prescription or remaining repeats for the biological medicine the patient is ceasing. (3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count submitted with the first authority application for a biological medicine. However, prescribers may provide a new baseline measurement any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to the revised baseline measurement.

(4) Recommencement of treatment after a 24 months break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological medicine therapy of at least 24 months, must qualify under the Initial 3 restriction and meet the relevant criteria and index of disease severity.

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment per continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

- (a) an active joint count of fewer than 10 active (swollen and tender) joints; or
- (b) a reduction in the active (swollen and tender) joint count by at least 50% from baseline; or
- (c) a reduction in the number of the following active joints, from at least 4, by at least 50%:
 - (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - (ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

- (1) completed authority prescription form(s); and
- (2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either Initial 1, Initial 2, or Initial 3 treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

If a patient fails to respond to PBS-subsidised biological medicine treatment 3 times (once with each agent) they will not be eligible to receive further PBS-subsidised biological medicine therapy in this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis
Treatment Phase: Continuing Treatment - balance of supply

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restriction.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL syringes

5283C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1092.41	41.30	Humira [VE]

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL pen devices

5284D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1092.41	41.30	Humira [VE]

■ ADALIMUMAB

Note TREATMENT OF ADULT PATIENTS WITH ANKYLOSING SPONDYLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab for adult patients with ankylosing spondylitis.

Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 7 biological medicines at any 1 time.

Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy.

A patient who has been receiving PBS-subsidised adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab treatment prior to 1 December 2020 is considered to start their first cycle as of 1 December 2020.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last prescription for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised biological medicine treatment with adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised biological medicine treatment in this treatment cycle and wishes to commence such therapy (Initial 1 - New patient)
- (ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or
- (iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years); or

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years). A patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy.

(b) Continuing treatment.

For the first continuing treatment course of PBS-subsidised biological medicine, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1, Initial 2 or Initial 3 treatment restrictions. For second and subsequent continuing courses of PBS-subsidised biological medicine it is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply. Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be conducted no later than 4 weeks from the cessation of that treatment course. Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI), or the prior NSAID therapy and exercise program requirements.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(3) Baseline measurements to determine response.

Services Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the BASDAI, ESR and/or CRP submitted with the first authority application for a biological medicine. For a new patient, the BASDAI used to determine the baseline must be measured while the patient is receiving NSAID therapy and completing their exercise program.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response.

Prescribers may provide new baseline measurements any time an 'Initial treatment' authority application is submitted within a treatment cycle and the eligibility for continuing treatment must be assessed according to these revised baseline measurements.

(4) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must qualify under the Initial 3 treatment restriction. The same clinical criteria and indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI) as for the Initial 1 (New patient) restriction will need to be met, but a re-trial of NSAID therapy and exercise therapy is not required.

Authority required

Ankylosing spondylitis

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Ankylosing Spondylitis PBS Authority Application - Supporting Information Form.

An adequate response is defined as an improvement from baseline of at least 2 of the BASDAI and 1 of the following:

- (a) an ESR measurement no greater than 25 mm per hour; or
- (b) a CRP measurement no greater than 10 mg per L; or
- (c) an ESR or CRP measurement reduced by at least 20% from baseline.

Where only 1 acute phase reactant measurement is supplied in the first application for PBS-subsidised treatment, that same marker must be measured and supplied in all subsequent continuing treatment applications.

All measurements provided must be no more than 1 month old at the time of application.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Ankylosing spondylitis

Treatment Phase: Continuing treatment - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the Continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restriction.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL syringes

9078T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1092.41	41.30	Humira [VE]

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL pen devices

9104E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1092.41	41.30	Humira [VE]

■ ADALIMUMAB

Note TREATMENT OF ADULT PATIENTS WITH MODERATE TO SEVERE ULCERATIVE COLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, golimumab, infliximab and vedolizumab for adult patients with ulcerative colitis. Patients are eligible for PBS-subsidised treatment with either adalimumab, golimumab, infliximab or vedolizumab at any one time.

Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, golimumab, infliximab and vedolizumab only.

From 1 June 2018, under the PBS, all adult patients will be able to commence a treatment cycle where they may trial PBS-subsidised biological medicine without having to experience a disease flare when swapping to one of the alternate agents. Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a PBS subsidised biological medicine while they continue to show a response to therapy.

A patient who received PBS-subsidised biological medicine treatment prior to 1 June 2018 is considered to start their first cycle as of 1 June 2018. Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a PBS-subsidised biological medicine under the new treatment cycle.

Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under the Initial 2 treatment restriction.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle under the Initial 3 treatment restrictions.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment after 1 June 2018.

(1) Initial treatment.

Applications for initial treatment should be made where:

- (i) an adult patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy (Initial 1 - New patient); or
 - (ii) an adult patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an alternate medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or
 - (iii) an adult patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same agent (Initial 2 - Change or Recommencement of treatment after a break in biological medicine therapy of less than 5 years); or
 - (iv) an adult patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - recommencement of treatment after a break in biological medicine of more than 5 years).
- Treatment authorisations under Initial 1, Initial 2 and Initial 3 will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, 14 weeks of therapy for golimumab, infliximab and vedolizumab.

A patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of treatment for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for golimumab, infliximab and vedolizumab. For the second and subsequent courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine treatment is approved, a patient may swap to an alternate biological medicine treatment within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Mayo clinic score or partial Mayo clinic score), or the prior corticosteroid therapy and immunosuppressive therapy.

A patient may trial an alternate biological medicine treatment at any time, regardless of whether they are receiving therapy (initial or continuing) at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug once within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(4) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological medicine therapy of at least 5 years, must requalify under Initial 3 treatment restriction and meet the relevant criteria with respect to the scores of disease severity.

Note TREATMENT OF PAEDIATRIC PATIENTS WITH MODERATE TO SEVERE ULCERATIVE COLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) for paediatric patients with infliximab or adalimumab for moderate to severe ulcerative colitis; and infliximab for acute severe ulcerative colitis.

Where the term 'biological medicine' appears in the following NOTES and restrictions, it refers to infliximab and adalimumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 2 biological medicines at any one time.

From 1 June 2017, under the PBS, all paediatric patients will be able to commence a treatment cycle where they may trial each PBS-subsidised biological medicine without having to experience a disease flare when swapping to the alternate agent.

Under these arrangements, within a single treatment cycle and depending on the disease severity, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy. A patient who received PBS-subsidised biological medicine treatment prior to 1 June 2017 is considered to have started their treatment cycle as of 1 June 2017. Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than twice. Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under the Initial 2 treatment restriction.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under the Initial 3 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine therapy after 1 June 2017.

(1) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy - Initial 1 treatment (new patient); or,
- (ii) a patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an

alternate agent - Initial 2 treatment (Change or Recommencement of treatment after a break in biological medicine therapy of less than 5 years) [further details are under 'Swapping treatment' below]; or
 (iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy with the same agent - Initial 2 treatment (Change or Recommencement of treatment after a break in therapy of less than 5 years); or
 (iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - recommencement of treatment after a break in biological medicine of more than 5 years). Treatment authorisations under Initial 1, Initial 2 and Initial 3 will be limited to provide for a maximum of 16 weeks of treatment for adalimumab and 14 weeks of treatment for infliximab. From 1 June 2017, a patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of treatment for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab. For the second and subsequent courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment.

Adalimumab only:

Two completed authority prescriptions should be submitted with every initial application for this drug. For patients weighing 40 kg or greater, one prescription should be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats and the second prescription should be written for 2 doses of 40 mg and 2 repeats. For patients weighing less than 40 kg, one prescription should be written for 2 doses of 40 mg with no repeats and the second prescription should be written for 2 doses of 20 mg with 3 repeats.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient remains eligible to receive up to 24 weeks per course of continuing treatment with that drug under the continuing treatment restriction providing they continue to sustain the response. It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure PBS subsidy criteria are met.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Paediatric Ulcerative Colitis Activity Index (PUCAI) Score, confirmation of ulcerative colitis disease), or the prior conventional therapies of corticosteroids or immunosuppressives.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving treatment (initial or continuing) at the time of the application. However, a patient cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug twice within the same treatment cycle. To ensure a patient receives the maximum treatment opportunities allowed under these swapping arrangements, it is important that they are assessed for response to every course of treatment.

(4) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to recommence treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must requalify under Initial 3 treatment restriction and meet the relevant criteria with respect to the indices of disease severity.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician; OR
- Must be treated by a specialist paediatric gastroenterologist.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have demonstrated or sustained an adequate response to treatment by having a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1 while receiving treatment with this drug; OR
- Patient must have demonstrated or sustained an adequate response to treatment by having a Paediatric Ulcerative Colitis Activity Index (PUCAI) score less than 10 while receiving treatment with this drug if aged 6 to 17 years.

Population criteria:

- Patient must be 6 years of age or older.

Patients who have failed to maintain a partial Mayo clinic score of less than or equal to 2, with no subscore greater than 1, or, patients who have failed to maintain a Paediatric Ulcerative Colitis Activity Index (PUCAI) score of less than 10 (if aged 6 to 17 years) with continuing treatment with this drug, will not be eligible to receive further PBS-subsidised treatment with this drug.

Patients are eligible to receive continuing treatment with this drug in courses of up to 24 weeks providing they continue to sustain a response.

At the time of the authority application, medical practitioners should request sufficient quantity for up to 24 weeks of treatment under this restriction.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

If patients aged 6 to 17 years fail to respond to PBS-subsidised biological medicine treatment 3 times (twice with one agent) they will not be eligible to receive further PBS-subsidised biological medicine therapy in this treatment cycle.

Note Authority applications for continuing treatment may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Continuing treatment - balance of supply

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician; OR
- Must be treated by a specialist paediatric gastroenterologist.

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the Continuing treatment restriction to complete 24 weeks of treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under this restriction.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL syringes

10960W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1092.41	41.30	Humira [VE]

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL pen devices

10961X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1092.41	41.30	Humira [VE]

adalimumab 20 mg/0.4 mL injection, 2 x 0.4 mL syringes

11121H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1092.41	41.30	Humira [VE]

ADALIMUMAB

Note TREATMENT OF COMPLEX REFRACTORY FISTULISING CROHN DISEASE

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for patients with complex refractory fistulising Crohn disease. Where the term 'biological medicine' appears in the following NOTES and restrictions, it refers to adalimumab and infliximab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the PBS-subsidised biological medicines for this condition at any one time.

From 1 April 2011, under the PBS, all patients will be able to commence a treatment cycle where they may trial PBS-subsidised adalimumab or infliximab without having to experience a disease flare when swapping to the alternate agent. Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with adalimumab or infliximab while they continue to show a response to therapy.

A patient who received PBS-subsidised adalimumab or infliximab treatment prior to 1 April 2011 is considered to have started their treatment cycle as of 1 April 2011.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised adalimumab or infliximab more than twice.

Once a patient has either failed or ceased to respond to treatment for this condition 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy for this condition before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised adalimumab or infliximab treatment in the most recent cycle to the date of the first application for initial treatment with adalimumab or infliximab under the new treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of less than 5 years, may commence a further course of treatment within the same treatment cycle.

A patient who has failed 3 trials or fewer of biological medicine therapy in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised adalimumab or infliximab therapy after 1 April 2011.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised adalimumab or infliximab therapy in this treatment cycle and wishes to commence such therapy (Initial treatment (new patient or Recommencement of treatment after more than 5 years break in therapy - Initial 1); or

(ii) a patient has received prior PBS-subsidised (initial or continuing) adalimumab or infliximab therapy and wishes to trial an

alternate agent (Initial 2 - Change or recommencement) [further details are under 'Swapping therapy' below]; or
(iii) a patient wishes to recommence treatment with adalimumab or infliximab following a break in PBS-subsidised therapy with that agent (Change or Recommencement of treatment after a break in therapy of less than 5 years (Initial 2)
Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy for adalimumab and 14 weeks of therapy for infliximab.

From 1 April 2011, a patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab, and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

For subsequent courses of PBS-subsidised adalimumab or infliximab treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

Adalimumab only: Two completed authority prescriptions must be submitted with every initial application for adalimumab. One prescription must be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats. The second prescription must be written for 2 doses of 40 mg and 2 repeats.

(b) Continuing treatment.

Adalimumab patients:

Following the completion of an initial treatment course with adalimumab, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

Infliximab patients:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment under the Initial 1 or Initial 2 treatment restrictions.

For the second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted supply of treatment.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be conducted within 4 weeks of the last dose. Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap if eligible to the alternate biological medicine within the same treatment cycle.

A patient may trial the alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with adalimumab or infliximab at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug two times within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment within the timeframes specified in the relevant restriction.

A patient who is not able to complete an initial treatment course for a biological medicine will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements submitted with the first authority application for adalimumab or infliximab. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used to assess response to all subsequent treatments.

(4) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to recommence treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Complex refractory Fistulising Crohn disease

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug for this condition.

An adequate response is defined as:

(a) a decrease from baseline in the number of open draining fistulae of greater than or equal to 50%; and/or
(b) a marked reduction in drainage of all fistula(e) from baseline, together with less pain and induration as reported by the patient.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Fistulising Crohn Disease PBS Authority Application - Supporting Information Form which includes a completed Fistula Assessment form including the date of the assessment of the patient's condition.

The most recent fistula assessment must be no more than 1 month old at the time of application.

To demonstrate a response to treatment the application must be accompanied by the results of the most recent course of biological medicine therapy following a minimum of 12 weeks of therapy.

It is recommended that an application for continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of this initial course of treatment to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Patients are eligible to receive continuing treatment with this drug in courses of up to 24 weeks providing they continue to sustain a response.

At the time of the authority application, medical practitioners should request the appropriate quantity and number of repeats to provide sufficient dose. Up to a maximum of 5 repeats will be authorised.

A maximum of 24 weeks treatment will be authorised under this restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Complex refractory Fistulising Crohn disease

Treatment Phase: Continuing treatment - balance of supply

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the Continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restriction.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL syringes

8964T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1092.41	41.30	Humira [VE]

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL pen devices

8966X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1092.41	41.30	Humira [VE]

■ ADALIMUMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE PSORIATIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab for adult patients with severe active psoriatic arthritis. Therefore, where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient receiving PBS-subsidised treatment for psoriatic arthritis is able to commence a treatment cycle where they may trial biological medicines without having to experience a disease flare when swapping to the alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

A patient who received PBS-subsidised adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab or ustekinumab treatment prior to 1 May 2019 is considered to start their first cycle as of 1 May 2019.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological

medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence another cycle [further details are under '(5) Recommencement of treatment after a 5-year break in PBS-subsidised therapy' below].

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe active psoriatic arthritis.

(1) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an alternate medicine (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years) or An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, golimumab secukinumab and tofacitinib, 18 to 20 weeks of therapy for certolizumab pegol (depending upon the dosing regimen), 20 weeks of therapy for ixekizumab, 22 weeks of therapy for infliximab, and 28 weeks of therapy for ustekinumab. It is recommended that a patient be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of PBS-subsidised initial treatment following a minimum of 12 weeks of therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

Grandfather patients (ixekizumab only).

A patient who commenced treatment with ixekizumab for severe psoriatic arthritis prior to 1 March 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

Grandfather patients (tofacitinib only).

A patient who commenced treatment with Tofacitinib for severe psoriatic arthritis prior to 1 May 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment under the Initial 1 or Initial 2 treatment restrictions.

For the second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to re-qualify with respect to either the indices of disease severity (i.e. erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP) level, and active joint count) or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify under the Initial 3 treatment restriction with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application.

However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

Within a treatment cycle a patient may alternate between therapy with any biological medicine of their choice (1 at a time) providing:

- (i) they have not received PBS-subsidised treatment with that particular biological medicine previously; or
- (ii) they have demonstrated an adequate response to that particular biological medicine if they have previously trialled it on the PBS; and
- (iii) they have not previously failed to respond to treatment 3 times in this treatment cycle with that biological medicine.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(4) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the indices of disease severity submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and these revised baseline measurements will be used to assess response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to determine response for all subsequent continuing treatments. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. 20 or more active joints), response will be determined according to a reduction in the total number of active joints.

(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment restriction according to the criteria of the relevant restriction and index of disease severity. The application must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Severe psoriatic arthritis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an erythrocyte sedimentation rate (ESR) no greater than 25 mm per hour or a C-reactive protein (CRP) level no greater than 15 mg per L or either marker reduced by at least 20% from baseline; and

either of the following:

- (a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or
- (b) a reduction in the number of the following major active joints, from at least 4, by at least 50%:
 - (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to determine response for all subsequent continuing treatments.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Severe Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either Initial 1, Initial 2, or Initial 3 treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Continuing treatment - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restriction.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL syringes

9034L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1092.41	41.30	Humira [VE]

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL pen devices

9102C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1092.41	41.30	Humira [VE]

■ ADALIMUMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CROHN DISEASE

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for adult patients with severe Crohn disease. Where the term 'biological medicine' appears in the following NOTES and restrictions, it refers to the tumour necrosis factor (TNF) alfa-antagonists (adalimumab and infliximab), the alpha-4 beta-7 integrin inhibitor (vedolizumab) and the human IgG1kappa monoclonal antibody (ustekinumab).

Patients are eligible for PBS-subsidised treatment with only 1 of the above PBS-subsidised biological medicines at any one time.

From 1 September 2017, under the PBS, all patients will be able to commence a treatment cycle where they may trial PBS-subsidised a biological medicine without having to experience a disease flare when swapping to the alternate agent. Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy.

A patient who received PBS-subsidised treatment with a biological medicine prior to 1 September 2017 is considered to have started their treatment cycle as of 1 September 2017.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once.

Once a patient has either failed or ceased to respond to treatment for this condition 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy for this condition before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of less than 5 years, may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab therapy after 1 September 2017.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised biological medicine treatment in this treatment cycle and wishes to commence such therapy Initial 1 (new patient); or
- (ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine and wishes to trial an alternate agent - Initial 2 (change or recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or
- (iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy with that agent - Initial 2 (change or recommencement of treatment after a break in therapy of less than 5 years); or
- (iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years - Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years).

From 1 September 2017, a patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy for adalimumab or ustekinumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab or vedolizumab, and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

For subsequent courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

Adalimumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats and the second prescription should be written for 2 doses of 40 mg and 2 repeats for patients weighing 40 kg or greater. For patients weighing less than 40 kg, one prescription should be written for 2 doses of 40 mg with no repeats and the second prescription should be written for 2 doses of 20 mg with 3 repeats.

Ustekinumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be written under S100 (Highly Specialised Drugs) for a weight-based loading dose, containing a quantity of up to 4 vials of 130 mg and no repeats. The second prescription should be written under S85 (General) for the subsequent first dose, containing a quantity of 2 vials of 45 mg and no repeats.

(b) Continuing treatment.

Following the completion of an initial treatment course with a biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted supply of treatment.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be conducted within 4 weeks of the last dose. Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine therapy is approved, a patient may swap if eligible to the alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Crohn Disease Activity Index (CDAI) Score, confirmation of Crohn disease), or the prior conventional therapies of corticosteroid therapy and immunosuppressive therapy.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug once within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment, within the timeframes specified in the relevant restriction.

A patient who is not able to complete an initial treatment course for a biological medicine will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the CDAI or evidence of intestinal inflammation submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used to assess response to all subsequent treatments.

(4) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to recommence treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity under the Initial 3 restriction. A re-trial of systemic therapy is not required.

Authority required

Severe Crohn disease

Treatment Phase: Initial treatment - Initial 1 (new patient)

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Population criteria:

- Patient must be aged 18 years or older.

Clinical criteria:

- Patient must have confirmed severe Crohn disease, defined by standard clinical, endoscopic and/or imaging features, including histological evidence, with the diagnosis confirmed by a gastroenterologist or a consultant physician, **AND**
- Patient must have failed to achieve an adequate response to prior systemic therapy with a tapered course of steroids, starting at a dose of at least 40 mg prednisolone (or equivalent), over a 6 week period, **AND**
- Patient must have failed to achieve adequate response to prior systemic immunosuppressive therapy with azathioprine at a dose of at least 2 mg per kg daily for 3 or more consecutive months; OR
- Patient must have failed to achieve adequate response to prior systemic immunosuppressive therapy with 6-mercaptopurine at a dose of at least 1 mg per kg daily for 3 or more consecutive months; OR
- Patient must have failed to achieve adequate response to prior systemic immunosuppressive therapy with methotrexate at a dose of at least 15 mg weekly for 3 or more consecutive months, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction, **AND**
- Patient must have a Crohn Disease Activity Index (CDAI) Score greater than or equal to 300 as evidence of failure to achieve an adequate response to prior systemic therapy; OR
- Patient must have short gut syndrome with diagnostic imaging or surgical evidence, or have had an ileostomy or colostomy; and must have evidence of intestinal inflammation; and must have evidence of failure to achieve an adequate response to prior systemic therapy as specified below; OR
- Patient must have extensive intestinal inflammation affecting more than 50 cm of the small intestine as evidenced by radiological imaging; and must have a Crohn Disease Activity Index (CDAI) Score greater than or equal to 220; and must have evidence of failure to achieve an adequate response to prior systemic therapy as specified below.

Applications for authorisation must be made in writing and must include:

- two completed authority prescription forms; and
- a completed Crohn Disease PBS Authority Application - Supporting Information Form which includes the following:
 - the completed current Crohn Disease Activity Index (CDAI) calculation sheet including the date of assessment of the patient's condition if relevant; and
 - details of prior systemic drug therapy [dosage, date of commencement and duration of therapy]; and
 - the reports and dates of the pathology or diagnostic imaging test(s) nominated as the response criterion, if relevant; and
 - the date of the most recent clinical assessment.

Evidence of failure to achieve an adequate response to prior therapy must include at least one of the following:

- patient must have evidence of intestinal inflammation;
- patient must be assessed clinically as being in a high faecal output state;
- patient must be assessed clinically as requiring surgery or total parenteral nutrition (TPN) as the next therapeutic option, in the absence of this drug, if affected by short gut syndrome, extensive small intestine disease or is an ostomy patient.

Evidence of intestinal inflammation includes:

- blood: higher than normal platelet count, or, an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour, or, a C-reactive protein (CRP) level greater than 15 mg per L; or
- faeces: higher than normal lactoferrin or calprotectin level; or
- diagnostic imaging: demonstration of increased uptake of intravenous contrast with thickening of the bowel wall or mesenteric lymphadenopathy or fat streaking in the mesentery.

Two completed authority prescriptions must be submitted with every initial application for adalimumab. One prescription must be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats. The second prescription must be written for 2 doses of 40 mg and 2 repeats.

Where fewer than 2 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment with adalimumab may be requested by telephone by contacting the Department of Human Services.

Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period.

All assessments, pathology tests and diagnostic imaging studies must be made within 1 month of the date of application and should be performed preferably whilst still on conventional treatment, but no longer than 1 month following cessation of the most recent prior treatment

If treatment with any of the specified prior conventional drugs is contraindicated according to the relevant TGA-approved Product Information, please provide details at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, details of this toxicity must be provided at the time of application.

Details of the accepted toxicities including severity can be found on the Department of Human Services website.

Any one of the baseline criteria may be used to determine response to an initial course of treatment and eligibility for continued therapy, according to the criteria included in the continuing treatment restriction. However, the same criterion must be used for any subsequent determination of response to treatment, for the purpose of eligibility for continuing PBS-subsidised therapy.

An assessment of a patient's response to an initial course of treatment must be conducted following a minimum of 12 weeks of therapy. An application for the continuing treatment must be accompanied with the assessment of response and submitted to the Department of Human Services no later than 4 weeks from the date of completion of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos. Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe Crohn disease

Treatment Phase: Initial treatment - Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years)

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Applications for authorisation must be made in writing and must include:

(a) two completed authority prescription forms; and

(b) a completed Crohn Disease PBS Authority Application - Supporting Information Form, which includes the following:

- (i) the completed Crohn Disease Activity Index (CDAI) Score calculation sheet including the date of the assessment of the patient's condition, if relevant; or
- (ii) the reports and dates of the pathology or diagnostic imaging test(s) used to assess response to therapy for patients with short gut syndrome, extensive small intestine disease or an ostomy, if relevant; and
- (iii) the date of clinical assessment; and
- (iv) the details of prior biological medicine treatment including the details of date and duration of treatment.

Two completed authority prescriptions must be submitted with every initial application for adalimumab. One prescription must be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats. The second prescription must be written for 2 doses of 40 mg and 2 repeats.

Where fewer than 2 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment with adalimumab may be requested by telephone by contacting the Department of Human Services.

Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period.

To demonstrate a response to treatment the application must be accompanied by the results of the most recent course of biological medicine therapy within the timeframes specified in the relevant restriction.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy for adalimumab or ustekinumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab and vedolizumab and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An assessment of a patient's response to an initial course of treatment must be conducted following a minimum of 12 weeks of therapy. An application for the continuing treatment must be accompanied with the assessment of response and submitted to the Department of Human Services no later than 4 weeks from the date of completion of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos.

Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe Crohn disease

Treatment Phase: Initial treatment - Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years)

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- Patient must have confirmed severe Crohn disease, defined by standard clinical, endoscopic and/or imaging features, including histological evidence, with the diagnosis confirmed by a gastroenterologist or a consultant physician, **AND**
- Patient must have a Crohn Disease Activity Index (CDAI) Score of greater than or equal to 300 that is no more than 4 weeks old at the time of application; OR
- Patient must have a documented history of intestinal inflammation and have diagnostic imaging or surgical evidence of short gut syndrome if affected by the syndrome or has an ileostomy or colostomy; OR
- Patient must have a documented history and radiological evidence of intestinal inflammation if the patient has extensive small intestinal disease affecting more than 50 cm of the small intestine, together with a Crohn Disease Activity Index (CDAI) Score greater than or equal to 220 and that is no more than 4 weeks old at the time of application, **AND**
- Patient must have evidence of intestinal inflammation; OR
- Patient must be assessed clinically as being in a high faecal output state; OR
- Patient must be assessed clinically as requiring surgery or total parenteral nutrition (TPN) as the next therapeutic option, in the absence of this drug, if affected by short gut syndrome, extensive small intestine disease or is an ostomy patient,

AND

- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Applications for authorisation must be made in writing and must include:

- (a) two completed authority prescription forms; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current Crohn Disease Activity Index (CDAI) calculation sheet including the date of assessment of the patient's condition if relevant; and
 - (ii) the reports and dates of the pathology or diagnostic imaging test(s) nominated as the response criterion, if relevant; and
 - (iii) the date of the most recent clinical assessment.

Evidence of intestinal inflammation includes:

- (i) blood: higher than normal platelet count, or, an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour, or, a C-reactive protein (CRP) level greater than 15 mg per L; or
- (ii) faeces: higher than normal lactoferrin or calprotectin level; or
- (iii) diagnostic imaging: demonstration of increased uptake of intravenous contrast with thickening of the bowel wall or mesenteric lymphadenopathy or fat streaking in the mesentery.

Two completed authority prescriptions must be submitted with every initial application for adalimumab. One prescription must be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats. The second prescription must be written for 2 doses of 40 mg and 2 repeats.

Where fewer than 2 repeats are requested at the time of the application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment with adalimumab may be requested by telephone by contacting the Department of Human Services.

Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period.

Any one of the baseline criteria may be used to determine response to an initial course of treatment and eligibility for continued therapy, according to the criteria included in the continuing treatment restriction. However, the same criterion must be used for any subsequent determination of response to treatment, for the purpose of eligibility for continuing PBS-subsidised therapy.

An assessment of a patient's response to an initial course of treatment must be conducted following a minimum of 12 weeks of therapy. An application for the continuing treatment must be accompanied with the assessment of response and submitted to the Department of Human Services no later than 4 weeks from the date of completion of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe Crohn disease

Treatment Phase: Balance of supply

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the Continuing treatment restriction to complete 24 weeks of treatment, **AND**
- The treatment must provide no more than the balance of up to 16 weeks therapy available under Initial 1, 2 or 3 treatment; OR
- The treatment must provide no more than the balance of up to 24 weeks therapy available under Continuing treatment.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

adalimumab 40 mg/0.8 mL injection, 6 x 0.8 mL pen devices

9187M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	3099.84	41.30	Humira [VE]

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL syringes

9188N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	1092.41	41.30	Humira [VE]

adalimumab 40 mg/0.8 mL injection, 6 x 0.8 mL syringes

9186L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	3099.84	41.30	Humira [VE]

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL pen devices

9190Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	1092.41	41.30	Humira [VE]

ADALIMUMAB

Note TREATMENT OF ADULT PATIENTS WITH A HISTORY OF JUVENILE IDIOPATHIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, etanercept and tocilizumab for a patient over 18 years who has a history of juvenile idiopathic arthritis with onset prior to the age of 18 years. Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, etanercept and tocilizumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 3 biological medicines at any one time.

From 1 April 2014, a patient receiving PBS-subsidised biological medicine therapy is considered to be in a treatment cycle where they may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements, within a single treatment cycle, a patient may:

- (i) continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy; and
- (ii) fail to respond or to sustain a response to each PBS-subsidised biological medicine once only. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment.

Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a

single treatment cycle and they must have, at a minimum, a 5 year break in PBS-subsidised biological medicine therapy before they are eligible to receive further PBS-subsidised biological medicine therapy.

The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine was approved to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who received PBS-subsidised biological medicine treatment immediately prior to 1 April 2014 is considered to be in their first cycle as of 1 April 2014. A patient who has had a break in biological medicine treatment of at least 24 months immediately prior to making a new application, on or after 1 April 2014, will commence a new treatment cycle under the Initial 3 treatment restriction.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of less than 24 months may commence a further course of treatment within the same treatment cycle under the Initial 2 treatment restriction.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of more than 24 months must commence a new treatment cycle under the Initial 3 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine therapy after 1 April 2014.

(1) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy (Initial 1 - New patient); or
- (ii) a patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an alternate medicine (Initial 2 - Change or recommencement of treatment after a break in biological medicine of less than 24 months) [further details are under 'Swapping therapy' below]; or
- (iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 24 months with the same medicine (Initial 2 - Change or recommencement of treatment after a break in biological medicine therapy of less than 24 months); or
- (iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 24 months (Initial 3 - recommencement of treatment after a break in biological medicine of more than 24 months).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy.

A patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not conducted within these timeframes, the patient will be deemed to have failed this course of treatment.

For second and subsequent cycles of PBS-subsidised biological medicine, it is recommended that a patient is reviewed in the 4 weeks prior to completing their current course of treatment and that an application is posted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services where applicable. Where a response assessment is not submitted, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (joint count and ESR/CRP) or the prior non- biological medicine therapy requirements, except if the patient has had a break in therapy of more than 24 months who would then need to requalify under the Initial 3 restrictions with respect to the indices of disease severity.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, an application for a patient who wishes to swap to an alternate biological medicine should be accompanied by the approved authority prescription or remaining repeats for the biological medicine the patient is ceasing.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count submitted with the first authority application for a biological medicine. However, prescribers may provide a new baseline measurement any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to the revised baseline measurement.

(4) Recommencement of treatment after a 24 months break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological medicine therapy of at least 24 months, must qualify under the Initial 3 restriction and meet the relevant criteria and index of disease severity.

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have a documented history of severe active juvenile idiopathic arthritis with onset prior to the age of 18 years, **AND**
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with disease modifying anti-rheumatic drugs (DMARDs) which must include at least 3 months continuous treatment with each of at least 2 DMARDs, one of which must be methotrexate at a dose of at least 20 mg weekly and one of which must be: (i) hydroxychloroquine at a dose of at least 200 mg daily; or (ii) leflunomide at a dose of at least 10 mg daily; or (iii) sulfasalazine at a dose of at least 2 g daily; OR
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs which, if methotrexate is contraindicated according to the Therapeutic Goods Administration (TGA)-approved Product Information or cannot be tolerated at a 20 mg weekly dose, must include at least 3 months continuous treatment with each of at least 2 of the following DMARDs: (i) hydroxychloroquine at a dose of at least 200 mg daily; and/or (ii) leflunomide at a dose of at least 10 mg daily; and/or (iii) sulfasalazine at a dose of at least 2 g daily; OR
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs which, if 3 or more of methotrexate, hydroxychloroquine, leflunomide and sulfasalazine are contraindicated according to the relevant TGA-approved Product Information or cannot be tolerated at the doses specified above, must include at least 3 months continuous treatment with each of at least 2 DMARDs, with one or more of the following DMARDs being used in place of the DMARDs which are contraindicated or not tolerated: (i) azathioprine at a dose of at least 1 mg/kg per day; and/or (ii) cyclosporin at a dose of at least 2 mg/kg/day; and/or (iii) sodium aurothiomalate at a dose of 50 mg weekly, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

If methotrexate is contraindicated according to the TGA-approved Product Information or cannot be tolerated at a 20 mg weekly dose, the application must include details of the contraindication or intolerance to methotrexate. The maximum tolerated dose of methotrexate must be documented in the application, if applicable.

The application must include details of the DMARDs trialed, their doses and duration of treatment, and all relevant contraindications and/or intolerances.

The requirement to trial at least 2 DMARDs for periods of at least 3 months each can be met using single agents sequentially or by using one or more combinations of DMARDs.

If the requirement to trial 6 months of intensive DMARD therapy with at least 2 DMARDs cannot be met because of contraindications and/or intolerances of a severity necessitating permanent treatment withdrawal to all of the DMARDs specified above, details of the contraindication or intolerance and dose for each DMARD must be provided in the authority application.

The following criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; **AND** either

(a) an active joint count of at least 20 active (swollen and tender) joints; or

(b) at least 4 active joints from the following list:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The joint count and ESR and/or CRP must be determined at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. All measures must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

The authority application must be made in writing and must include:

(1) completed authority prescription form(s); and

(2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

An assessment of a patient's response to an initial course of treatment must be conducted following a minimum of 12 weeks of therapy. An application for the continuing treatment must be accompanied with the assessment of response and submitted to the Department of Human Services no later than 4 weeks from the date of completion of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note The Department of Human Services website (www.humanservices.gov.au) has details of the toxicities, including severity, which will be accepted for the following purposes:

- (a) exempting a patient from the requirement to undertake a minimum 3 month trial of methotrexate at a 20 mg weekly dose;
- (b) substituting azathioprine, cyclosporin or sodium aurothiomalate for another DMARD as part of the 6 month intensive DMARD trial;
- (c) exempting a patient from the requirement for a 6 month trial of intensive DMARD therapy.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 2 (change or recommencement of treatment after break in biological medicine of less than 24 months)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have a documented history of severe active juvenile idiopathic arthritis with onset prior to the age of 18 years, **AND**
- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

- (a) an active joint count of fewer than 10 active (swollen and tender) joints; or
- (b) a reduction in the active (swollen and tender) joint count by at least 50% from baseline; or
- (c) a reduction in the number of the following active joints, from at least 4, by at least 50%:
 - (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - (ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The authority application must be made in writing and must include:

- (1) completed authority prescription form(s); and
- (2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to change or recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient who fails to demonstrate a response to treatment with this drug under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug in this treatment cycle. A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the initial 3 treatment restriction.

If a patient fails to respond to PBS-subsidised biological medicine treatment 3 times (once with each agent) they will not be eligible to receive further PBS-subsidised biological medicine therapy in this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online

Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 3 (recommencement of treatment after a break in biological medicine of more than 24 months)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 24 months or more from the most recently approved PBS-subsidised biological medicine for this condition; OR
- Patient must not have received PBS-subsidised biological medicine for at least 5 years if they failed or ceased to respond to PBS-subsidised biological medicine treatment 3 times in their last treatment cycle, **AND**
- The condition must have an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; OR
- The condition must have a C-reactive protein (CRP) level greater than 15 mg per L, **AND**
- The condition must have either (a) a total active joint count of at least 20 active (swollen and tender) joints; or (b) at least 4 active major joints, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Active joints are defined as:

- (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
- (ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

All measures of joint count must be no more than 4 weeks old at the time of this application.

The authority application must be made in writing and must include:

- (1) completed authority prescription form(s); and
- (2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos. Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient) or Initial 2 (change or recommencement of treatment after break of less than 24 months) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 24 months) - balance of supply

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 24 months) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 24 months) to complete 16 weeks of treatment, **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL syringes

5281Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	1092.41	41.30	Humira [VE]

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL pen devices

5282B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	1092.41	41.30	Humira [VE]

■ ADALIMUMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for adults with severe active rheumatoid arthritis. Where the term biological medicine appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alfa antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab), the T-cell co-stimulation modulator (abatacept) and the Janus kinase (JAK) inhibitors (baricitinib, tofacitinib, upadacitinib).

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised biological medicine therapy may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once, and
- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised biological medicines for the treatment of rheumatoid arthritis.

A serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Stevens Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered a treatment failure.

A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction.

A patient who has failed fewer than 5 biological medicines and who has a break in treatment of less than 24 months may commence a further course of treatment with a biological medicine under Initial 2 treatment restriction. A patient who has failed fewer than 5 biological medicines and who has had a break in therapy of longer than 24 months may commence a further course of treatment with a biological medicine under the Initial 3 treatment restriction. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine treatment is stopped to the date of the new application for treatment with a biological medicine.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 April 2019.

(a) Initial treatment.

Applications for initial treatment should be made where:

- a patient has received no prior PBS-subsidised biological medicine treatment and wishes to commence such therapy, excluding rituximab (Initial 1 - new patient); or
- a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months) [further details are under 'Swapping therapy' below]; or
- a patient wishes to re-commence treatment with a specific biological medicine following a break of less than 24 months in PBS-subsidised therapy with that agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months).
- a patient wishes to re-commence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 24 months (Initial 3 - re-commencement of treatment after a break in biological medicine of more than 24 months)

Initial applications for a new patient (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6-month intensive DMARD trial, but prior to ceasing DMARD therapy.

Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, baricitinib, etanercept, golimumab, tocilizumab, tofacitinib and upadacitinib, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

Rituximab patients should be assessed following a minimum of 12 weeks after the first infusion, and the assessment should be submitted to Services Australia within 4 weeks to ensure continuity of treatment for those who meet the continuing restriction for PBS subsidised treatment with this drug for this condition.

Where a response assessment is not provided the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for

permanent withdrawal of treatment. For second and subsequent courses of PBS-subsidised biological medicine (excluding rituximab) treatment, it is recommended that a patient be reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to Services Australia within sufficient time to allow processing. Abatacept patients:

A patient is eligible to receive one I.V. loading dose when commencing treatment with the subcutaneous formulation. Two prescriptions are required, the first prescription for the I.V. loading dose for sufficient vials for one dose based on the patient's weight with no repeats. The second prescription for the subcutaneous formulation, with a maximum quantity of 4 and up to 3 repeats, must be submitted with the initial application.

Rituximab patients:

A further application may be submitted to Services Australia within sufficient time to allow processing. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

Assessment of the patient's response to treatment must be conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the cessation of the most recent course of biological medicine therapy.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction. Where a response assessment is not conducted within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

(2) Swapping therapy

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non- biological medicine therapy requirements except if the patient has had a break in therapy of more than 24 months who would need to requalify with respect to the indices of disease severity. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each biological medicine trialled.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent, unless they have experienced a serious adverse reaction of a severity necessitating permanent treatment withdrawal.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug.

Abatacept:

A patient swapping from I.V. abatacept to subcutaneous abatacept will not be eligible for an I.V. loading dose when commencing treatment with the subcutaneous formulation.

Rituximab:

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF-alfa antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they be assessed for response to every course of treatment, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological medicine during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological medicine therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate biological medicine may do so without having to have any treatment-free period.

(3) Baseline measurements to determine response.

Determination of whether a response to treatment has been demonstrated must be based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and the eligibility for continuing treatment must be assessed according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response must be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be determined on the total number of major joints.

Applications under the Initial 1 treatment restriction for a new patient must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. The results must be no more than 4 weeks old at the time of application.

Applications under the Initial 3 treatment restriction for re-commencement of treatment after a break in biological medicine of more than 24 months must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Applications for treatment with this drug where the dosing frequency exceeds 40 mg per fortnight will not be approved.

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment per continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient has either failed or ceased to respond to a PBS-subsidised biological medicine for this condition 5 times, they will not be eligible to receive further PBS-subsidised treatment with a biological medicine for this condition.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Continuing Treatment - balance of supply.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restriction.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL syringes

8741C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1092.41	41.30	Humira [VE]

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL pen devices

9100Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1092.41	41.30	Humira [VE]

ADALIMUMAB
Note TREATMENT OF COMPLEX REFRACTORY FISTULISING CROHN DISEASE

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for patients with complex refractory fistulising Crohn disease. Where the term 'biological medicine' appears in the following NOTES and restrictions, it refers to adalimumab and infliximab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the PBS-subsidised biological medicines for this condition at any one time.

From 1 April 2011, under the PBS, all patients will be able to commence a treatment cycle where they may trial PBS-subsidised adalimumab or infliximab without having to experience a disease flare when swapping to the alternate agent. Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with adalimumab or infliximab while they continue to show a response to therapy.

A patient who received PBS-subsidised adalimumab or infliximab treatment prior to 1 April 2011 is considered to have started their treatment cycle as of 1 April 2011.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised adalimumab or infliximab more than twice.

Once a patient has either failed or ceased to respond to treatment for this condition 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy for this condition before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised adalimumab or infliximab treatment in the most recent cycle to the date of the first application for initial treatment with adalimumab or infliximab under the new treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of less than 5 years, may commence a further course of treatment within the same treatment cycle.

A patient who has failed 3 trials or fewer of biological medicine therapy in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised adalimumab or infliximab therapy after 1 April 2011.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised adalimumab or infliximab therapy in this treatment cycle and wishes to commence such therapy (Initial treatment (new patient or Recommencement of treatment after more than 5 years break in therapy - Initial 1); or

(ii) a patient has received prior PBS-subsidised (initial or continuing) adalimumab or infliximab therapy and wishes to trial an alternate agent (Initial 2 - Change or recommencement) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with adalimumab or infliximab following a break in PBS-subsidised therapy with that agent (Change or Recommencement of treatment after a break in therapy of less than 5 years (Initial 2)

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy for adalimumab and 14 weeks of therapy for infliximab.

From 1 April 2011, a patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab, and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

For subsequent courses of PBS-subsidised adalimumab or infliximab treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

Adalimumab only: Two completed authority prescriptions must be submitted with every initial application for adalimumab.

One prescription must be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats. The second prescription must be written for 2 doses of 40 mg and 2 repeats.

(b) Continuing treatment.

Adalimumab patients:

Following the completion of an initial treatment course with adalimumab, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

Infliximab patients:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment under the Initial 1 or Initial 2 treatment restrictions.

For the second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted supply of treatment.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be conducted within 4 weeks of the last dose. Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious

adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap if eligible to the alternate biological medicine within the same treatment cycle.

A patient may trial the alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with adalimumab or infliximab at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug two times within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment within the timeframes specified in the relevant restriction.

A patient who is not able to complete an initial treatment course for a biological medicine will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements submitted with the first authority application for adalimumab or infliximab. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used to assess response to all subsequent treatments.

(4) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to recommence treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Complex refractory Fistulising Crohn disease

Treatment Phase: Initial treatment (new patient or Recommencement of treatment after more than 5 years break in therapy - Initial 1)

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have confirmed Crohn disease, defined by standard clinical, endoscopic and/or imaging features, including histological evidence, with the diagnosis confirmed by a gastroenterologist or a consultant physician, **AND**
- Patient must have an externally draining enterocutaneous or rectovaginal fistula.

Applications for authorisation must be made in writing and must include:

(a) two completed authority prescription forms; and

(b) a completed Fistulising Crohn Disease PBS Authority Application - Supporting Information Form which includes a completed current Fistula Assessment Form including the date of assessment of the patient's condition of no more than 1 month old at the time of application.

A maximum of 16 weeks of treatment with this drug will be approved under this criterion.

Two completed authority prescriptions must be submitted with every initial application for adalimumab. One prescription must be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats. The second prescription must be written for 2 doses of 40 mg and 2 repeats.

The assessment of the patient's response to this initial course of treatment must be made following a minimum of 12 weeks of therapy so that there is adequate time for a response to be demonstrated.

It is recommended that an application for continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of this initial course of treatment to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Complex refractory Fistulising Crohn disease

Treatment Phase: Change or Recommencement of treatment after a break in therapy of less than 5 years (Initial 2)

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR

- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have failed PBS-subsidised therapy with this drug for this condition more than once in the current treatment cycle.

To demonstrate a response to treatment the application must be accompanied by the results of the most recent course of biological medicine therapy following a minimum of 12 weeks of therapy.

It is recommended that an application for continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of this initial course of treatment to ensure continuity of treatment for those patients who meet the continuation criterion for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Applications for authorisation must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Fistulising Crohn Disease PBS Authority Application - Supporting Information Form which includes the following:
 - (i) a completed current Fistula Assessment Form including the date of assessment of the patient's condition; and
 - (ii) details of prior biological medicine treatment including details of date and duration of treatment.

The most recent fistula assessment must be no more than 1 month old at the time of application.

A maximum of 16 weeks of treatment with this drug will be approved under this criterion.

Two completed authority prescriptions must be submitted with every initial application for adalimumab. One prescription must be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats. The second prescription must be written for 2 doses of 40 mg and 2 repeats.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Complex refractory Fistulising Crohn disease

Treatment Phase: Initial 1 (new patient or Recommencement of treatment after more than 5 years break in therapy), Initial 2 (Change or Recommencement of treatment after a break in therapy of less than 5 years) - Balance of supply

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient or patient recommencing treatment after a break of 5 years or more) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break of less than 5 years) restriction to complete 16 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

adalimumab 40 mg/0.8 mL injection, 6 x 0.8 mL pen devices

8962Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	3099.84	41.30	Humira [VE]

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL syringes

8963R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	1092.41	41.30	Humira [VE]

adalimumab 40 mg/0.8 mL injection, 6 x 0.8 mL syringes

8961P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	3099.84	41.30	Humira [VE]

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL pen devices

8965W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	1092.41	41.30	Humira [VE]

■ ADALIMUMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years); or

(iii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or

(iv) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient be reviewed in the 4 weeks prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

Grandfather patients (risankizumab only).

A patient who commenced treatment with risankizumab for chronic plaque psoriasis prior to 1 December 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the Initial treatment Grandfather treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment. This assessment must be conducted within 4 weeks of the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24

weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the 4 weeks prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions. For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the 4 weeks prior to completing their current course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure patients receive the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

Services Australia will determine whether a response to treatment has been demonstrated based on the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or commencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 the PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Whole body

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

The authority application must be made in writing and must include:

- a completed authority prescription form(s); and
- a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Face, hand, foot

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or
- (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet and face, hand, foot area diagrams including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

The PASI assessment for continuing treatment must be performed on the same affected area assessed at baseline.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Whole body or Continuing treatment, Face, hand, foot - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the continuing treatment, Whole body restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the continuing treatment, Face, hand, foot restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restrictions, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate).

Treatment criteria:

- Must be treated by a dermatologist.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL syringes

9427E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1092.41	41.30	Humira [VE]

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL pen devices

9428F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1092.41	41.30	Humira [VE]

■ **ADALIMUMAB**

Note TREATMENT OF ADULT PATIENTS WITH ANKYLOSING SPONDYLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab for adult patients with ankylosing spondylitis.

Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 7 biological medicines at any 1 time.

Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy.

A patient who has been receiving PBS-subsidised adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab treatment prior to 1 December 2020 is considered to start their first cycle as of 1 December 2020.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last prescription for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised biological medicine treatment with adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised biological medicine treatment in this treatment cycle and wishes to commence such therapy (Initial 1 - New patient)

(ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years); or

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

A patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy.

(b) Continuing treatment.

For the first continuing treatment course of PBS-subsidised biological medicine, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1, Initial 2 or Initial 3 treatment restrictions. For second and subsequent continuing courses of PBS-subsidised biological medicine it is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply. Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24

weeks providing they continue to sustain the response.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be conducted no later than 4 weeks from the cessation of that treatment course. Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI), or the prior NSAID therapy and exercise program requirements.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(3) Baseline measurements to determine response.

Services Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the BASDAI, ESR and/or CRP submitted with the first authority application for a biological medicine. For a new patient, the BASDAI used to determine the baseline must be measured while the patient is receiving NSAID therapy and completing their exercise program.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response.

Prescribers may provide new baseline measurements any time an 'Initial treatment' authority application is submitted within a treatment cycle and the eligibility for continuing treatment must be assessed according to these revised baseline measurements.

(4) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must qualify under the Initial 3 treatment restriction. The same clinical criteria and indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI) as for the Initial 1 (New patient) restriction will need to be met, but a re-trial of NSAID therapy and exercise therapy is not required.

Authority required

Ankylosing spondylitis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Clinical criteria:

- The condition must be radiographically (plain X-ray) confirmed Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have at least 2 of the following: (i) low back pain and stiffness for 3 or more months that is relieved by exercise but not by rest; or (ii) limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by a score of at least 1 on each of the lumbar flexion and lumbar side flexion measurements of the Bath Ankylosing Spondylitis Metrology Index (BASMI); or (iii) limitation of chest expansion relative to normal values for age and gender,

AND

- Patient must have failed to achieve an adequate response following treatment with at least 2 non-steroidal anti-inflammatory drugs (NSAIDs), whilst completing an appropriate exercise program, for a total period of 3 months, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

The application must include details of the NSAIDs trialled, their doses and duration of treatment.

If the NSAID dose is less than the maximum recommended dose in the relevant TGA-approved Product Information, the application must include the reason a higher dose cannot be used.

If treatment with NSAIDs is contraindicated according to the relevant TGA-approved Product Information, the application must provide details of the contraindication.

If intolerance to NSAID treatment develops during the relevant period of use which is of a severity to necessitate permanent treatment withdrawal, the application must provide details of the nature and severity of this intolerance.

The following criteria indicate failure to achieve an adequate response and must be demonstrated at the time of the initial application:

- (a) a Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 4 on a 0-10 scale; AND
- (b) an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 10 mg per L.

The BASDAI must be determined at the completion of the 3 month NSAID and exercise trial, but prior to ceasing NSAID treatment. The BASDAI must be no more than 1 month old at the time of initial application.

Both ESR and CRP measures should be provided with the initial treatment application and both must be no more than 1 month old. If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reason this criterion cannot be satisfied.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Ankylosing Spondylitis PBS Authority Application - Supporting Information Form which includes the following:

- (i) a copy of the radiological report confirming Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis; and
- (ii) a completed BASDAI Assessment Form; and
- (iii) a completed Exercise Program Self Certification Form included in the supporting information form.

An assessment of a patient's response to an initial course of treatment must be conducted following a minimum of 12 weeks of therapy. An application for the continuing treatment must be accompanied with the assessment of response and submitted to the Department of Human Services no later than 4 weeks from the date of completion of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Details of the toxicities, including severity, which will be accepted for the purposes of administering this restriction can be found on the Department of Human Services website at www.humanservices.gov.au

Note For details on the appropriate minimum exercise program that will be accepted for the purposes of administering this restriction, please refer to the Department of Human Services website at www.humanservices.gov.au

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Ankylosing spondylitis

Treatment Phase: Initial treatment - Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Ankylosing Spondylitis PBS Authority Application - Supporting Information Form.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to change or recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

An adequate response is defined as an improvement from baseline of at least 2 of the BASDAI and 1 of the following:

- (a) an ESR measurement no greater than 25 mm per hour; or
- (b) a CRP measurement no greater than 10 mg per L; or
- (c) an ESR or CRP measurement reduced by at least 20% from baseline.

Where only 1 acute phase reactant measurement is supplied in the first application for PBS-subsidised treatment, that same marker must be measured and supplied in all subsequent continuing treatment applications.

All measurements provided must be no more than 1 month old at the time of application.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Ankylosing spondylitis

Treatment Phase: Initial treatment - Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must be radiographically (plain X-ray) confirmed Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis, **AND**
- Patient must have at least 2 of the following: (i) low back pain and stiffness for 3 or more months that is relieved by exercise but not by rest; or (ii) limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by a score of at least 1 on each of the lumbar flexion and lumbar side flexion measurements of the Bath Ankylosing Spondylitis Metrology Index (BASMI); or (iii) limitation of chest expansion relative to normal values for age and gender,

AND

- Patient must have a Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 4 on a 0-10 scale that is no more than 4 weeks old at the time of application, **AND**
- Patient must have an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour that is no more than 4 weeks old at the time of application; OR
- Patient must have a C-reactive protein (CRP) level greater than 10 mg per L that is no more than 4 weeks old at the time of application; OR
- Patient must have a clinical reason as to why demonstration of an elevated ESR or CRP cannot be met and the application must state the reason, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Ankylosing Spondylitis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) a copy of the radiological report confirming Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis; and
 - (ii) a completed BASDAI Assessment Form.

An assessment of a patient's response to an initial course of treatment must be conducted following a minimum of 12 weeks of therapy. An application for the continuing treatment must be accompanied with the assessment of response and submitted to the Department of Human Services no later than 4 weeks from the date of completion of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online

Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Ankylosing spondylitis

Treatment Phase: Initial treatment - Initial 1 (new patient), Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years) - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 16 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL syringes

9077R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	1092.41	41.30	Humira [VE]

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL pen devices

9103D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	1092.41	41.30	Humira [VE]

ADALIMUMAB

Note TREATMENT OF ADULT PATIENTS WITH MODERATE TO SEVERE ULCERATIVE COLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, golimumab, infliximab and vedolizumab for adult patients with ulcerative colitis. Patients are eligible for PBS-subsidised treatment with either adalimumab, golimumab, infliximab or vedolizumab at any one time.

Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, golimumab, infliximab and vedolizumab only.

From 1 June 2018, under the PBS, all adult patients will be able to commence a treatment cycle where they may trial PBS-subsidised biological medicine without having to experience a disease flare when swapping to one of the alternate agents. Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a PBS subsidised biological medicine while they continue to show a response to therapy.

A patient who received PBS-subsidised biological medicine treatment prior to 1 June 2018 is considered to start their first cycle as of 1 June 2018. Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a PBS-subsidised biological medicine under the new treatment cycle.

Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under the Initial 2 treatment restriction.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle under the Initial 3 treatment restrictions.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment after 1 June 2018.

(1) Initial treatment.

Applications for initial treatment should be made where:

- an adult patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy (Initial 1 - New patient); or
- an adult patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an alternate medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or
- an adult patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised

therapy of less than 5 years with the same agent (Initial 2 - Change or Recommencement of treatment after a break in biological medicine therapy of less than 5 years); or
 (iv) an adult patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - recommencement of treatment after a break in biological medicine of more than 5 years). Treatment authorisations under Initial 1, Initial 2 and Initial 3 will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, 14 weeks of therapy for golimumab, infliximab and vedolizumab.

A patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of treatment for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for golimumab, infliximab and vedolizumab. For the second and subsequent courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine treatment is approved, a patient may swap to an alternate biological medicine treatment within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Mayo clinic score or partial Mayo clinic score), or the prior corticosteroid therapy and immunosuppressive therapy.

A patient may trial an alternate biological medicine treatment at any time, regardless of whether they are receiving therapy (initial or continuing) at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug once within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(4) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological medicine therapy of at least 5 years, must requalify under Initial 3 treatment restriction and meet the relevant criteria with respect to the scores of disease severity.

Note TREATMENT OF PAEDIATRIC PATIENTS WITH MODERATE TO SEVERE ULCERATIVE COLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) for paediatric patients with infliximab or adalimumab for moderate to severe ulcerative colitis; and infliximab for acute severe ulcerative colitis.

Where the term 'biological medicine' appears in the following NOTES and restrictions, it refers to infliximab and adalimumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 2 biological medicines at any one time.

From 1 June 2017, under the PBS, all paediatric patients will be able to commence a treatment cycle where they may trial each PBS-subsidised biological medicine without having to experience a disease flare when swapping to the alternate agent. Under these arrangements, within a single treatment cycle and depending on the disease severity, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy. A patient who received PBS-subsidised biological medicine treatment prior to 1 June 2017 is considered to have started their treatment cycle as of 1 June 2017. Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than twice. Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under the Initial 2 treatment restriction.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under the Initial 3 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine therapy after 1 June 2017.

(1) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy - Initial 1 treatment (new patient); or,

(ii) a patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an alternate agent - Initial 2 treatment (Change or Recommencement of treatment after a break in biological medicine therapy of less than 5 years) [further details are under 'Swapping treatment' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy with the same agent - Initial 2 treatment (Change or Recommencement of treatment after a break in therapy of less than 5 years); or

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - recommencement of treatment after a break in biological medicine of more than 5 years).

Treatment authorisations under Initial 1, Initial 2 and Initial 3 will be limited to provide for a maximum of 16 weeks of treatment for adalimumab and 14 weeks of treatment for infliximab. From 1 June 2017, a patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of treatment for adalimumab and

up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab. For the second and subsequent courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment.

Adalimumab only:

Two completed authority prescriptions should be submitted with every initial application for this drug. For patients weighing 40 kg or greater, one prescription should be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats and the second prescription should be written for 2 doses of 40 mg and 2 repeats. For patients weighing less than 40 kg, one prescription should be written for 2 doses of 40 mg with no repeats and the second prescription should be written for 2 doses of 20 mg with 3 repeats.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient remains eligible to receive up to 24 weeks per course of continuing treatment with that drug under the continuing treatment restriction providing they continue to sustain the response. It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure PBS subsidy criteria are met.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Paediatric Ulcerative Colitis Activity Index (PUCAI) Score, confirmation of ulcerative colitis disease), or the prior conventional therapies of corticosteroids or immunosuppressives.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving treatment (initial or continuing) at the time of the application. However, a patient cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug twice within the same treatment cycle. To ensure a patient receives the maximum treatment opportunities allowed under these swapping arrangements, it is important that they are assessed for response to every course of treatment.

(4) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to recommence treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must requalify under Initial 3 treatment restriction and meet the relevant criteria with respect to the indices of disease severity.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician; OR
- Must be treated by a specialist paediatric gastroenterologist.

Clinical criteria:

- Patient must have failed to achieve an adequate response to a 5-aminosalicylate oral preparation in a standard dose for induction of remission for 3 or more consecutive months or have intolerance necessitating permanent treatment withdrawal, **AND**
- Patient must have failed to achieve an adequate response to azathioprine at a dose of at least 2 mg per kg daily for 3 or more consecutive months or have intolerance necessitating permanent treatment withdrawal; OR
- Patient must have failed to achieve an adequate response to 6-mercaptopurine at a dose of at least 1 mg per kg daily for 3 or more consecutive months or have intolerance necessitating permanent treatment withdrawal; OR
- Patient must have failed to achieve an adequate response to a tapered course of oral steroids, starting at a dose of at least 40 mg (for a child, 1 to 2 mg/kg up to 40 mg) prednisolone (or equivalent), over a 6 week period or have intolerance necessitating permanent treatment withdrawal, and followed by a failure to achieve an adequate response to 3 or more consecutive months of treatment of an appropriately dosed thiopurine agent, **AND**
- Patient must have a Mayo clinic score greater than or equal to 6 if an adult patient; OR
- Patient must have a partial Mayo clinic score greater than or equal to 6, provided the rectal bleeding and stool frequency subscores are both greater than or equal to 2 (endoscopy subscore is not required for a partial Mayo clinic score); OR
- Patient must have a Paediatric Ulcerative Colitis Activity Index (PUCAI) Score greater than or equal to 30 if aged 6 to 17 years.

Population criteria:

- Patient must be 6 years of age or older.

Applications for authorisation of change or recommencement treatment must be in writing and must include:

(a) two completed authority prescription forms; and

(b) a completed Ulcerative Colitis PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed current Mayo clinic or partial Mayo clinic or Paediatric Ulcerative Colitis Activity Index (PUCAI) calculation sheet including the date of assessment of the patient's condition; and

(ii) details of prior systemic drug therapy [dosage, date of commencement and duration of therapy].

For patients weighing 40 kg or greater, a maximum quantity and number of repeats to provide for an initial 16 weeks course of this drug consisting of a 160 mg dose at week 0, 80 mg dose at week 2 and 40 mg dose at weeks 4, 6, 8, 10, 12 and 14 will be authorised.

For patients weighing less than 40 kg, a maximum quantity and number of repeats to provide for an initial 16 weeks of this drug consisting of a 80 mg dose at week 0, 40 mg dose at week 2 and a 20 mg dose at weeks 4, 6, 8, 10, 12 and 14 will be authorised.

Two completed authority prescriptions must be submitted with every initial application for this drug. For patients weighing 40 kg or greater, one prescription should be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats and the second prescription must be written for 2 doses of 40 mg and 2 repeats.

For patients weighing less than 40 kg, one prescription should be written for 2 doses of 40 mg with no repeats and the second prescription should be written for 2 doses of 20 mg with 3 repeats.

All tests and assessments should be performed preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior conventional treatment.

The most recent Mayo clinic, partial Mayo clinic or Paediatric Ulcerative Colitis Activity Index (PUCAI) score must be no more than 4 weeks old at the time of application.

A partial Mayo clinic or Paediatric Ulcerative Colitis Activity Index (PUCAI) assessment of the patient's response to this initial course of treatment must be made following a minimum of 12 weeks of treatment for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for golimumab, infliximab and vedolizumab so that there is adequate time for a response to be demonstrated.

The measurement of response to the prior course of therapy must be documented in the patient's medical notes.

If treatment with any of the above-mentioned drugs is contraindicated according to the relevant TGA-approved Product Information, details must be provided at the time of application.

An application for the continuing treatment must be accompanied with the assessment of response and submitted to the Department of Human Services no later than 4 weeks from the date of completion of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Details of the accepted toxicities including severity can be found on the Department of Human Services website.

Note At the time of the authority application, medical practitioners should request sufficient quantity for up to 16 weeks of treatment under this restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Initial treatment - Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years)

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician; OR
- Must be treated by a specialist paediatric gastroenterologist.

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle; OR
- Patient must have previously received PBS-subsidised treatment with a biological medicine (adalimumab or infliximab) for this condition in this treatment cycle if aged 6 to 17 years, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle; OR
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle more than once if aged 6 to 17 years.

Population criteria:

- Patient must be 6 years of age or older.

Applications for authorisation must be in writing and must include:

(a) two completed authority prescription forms; and

(b) a completed Ulcerative Colitis PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed current Mayo clinic or partial Mayo clinic or Paediatric Ulcerative Colitis Activity Index (PUCAI) calculation sheet including the date of assessment of the patient's condition if relevant; and

(ii) the details of prior biological medicine treatment including the details of date and duration of treatment.

Two completed authority prescriptions must be submitted with every initial application for this drug. For patients weighing 40 kg or greater, one prescription should be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats and the second prescription must be written for 2 doses of 40 mg and 2 repeats.

For patients weighing less than 40 kg, one prescription should be written for 2 doses of 40 mg with no repeats and the second prescription should be written for 2 doses of 20 mg with 3 repeats.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to change or recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3, or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for golimumab, infliximab and vedolizumab and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient who fails to demonstrate a response to treatment with this drug under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug in this treatment cycle. A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the initial 3 treatment restriction.

If patients aged 6 to 17 years fail to respond to PBS-subsidised biological medicine treatment 3 times (twice with one agent) they will not be eligible to receive further PBS-subsidised biological medicine therapy in this treatment cycle.

Note At the time of the authority application, medical practitioners should request sufficient quantity for up to 16 weeks of treatment under this restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Initial treatment - Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years)

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician; OR
- Must be treated by a specialist paediatric gastroenterologist.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have had a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- Patient must have a Mayo clinic score greater than or equal to 6 if an adult patient; OR
- Patient must have a partial Mayo clinic score greater than or equal to 6, provided the rectal bleeding and stool frequency subscores are both greater than or equal to 2 (endoscopy subscore is not required for a partial Mayo clinic score); OR
- Patient must have a Paediatric Ulcerative Colitis Activity Index (PUCAI) Score greater than or equal to 30 if aged 6 to 17 years.

Population criteria:

- Patient must be 6 years of age or older.

Applications for authorisation must be in writing and must include:

(a) two completed authority prescription forms; and

(b) a completed Ulcerative Colitis PBS Authority Application - Supporting Information Form which includes the following:

- (i) the completed current Mayo clinic or partial Mayo clinic or Paediatric Ulcerative Colitis Activity Index (PUCAI) calculation sheet including the date of assessment of the patient's condition; and
- (ii) the details of prior biological medicine treatment including the details of date and duration of treatment.

For patients weighing 40 kg or greater, a maximum quantity and number of repeats to provide for an initial 16 weeks course of this drug consisting of a 160 mg dose at week 0, 80 mg dose at week 2 and 40 mg dose at weeks 4, 6, 8, 10, 12 and 14 will be authorised.

For patients weighing less than 40 kg, a maximum quantity and number of repeats to provide for an initial 16 weeks of this drug consisting of a 80 mg dose at week 0, 40 mg dose at week 2 and a 20 mg dose at weeks 4, 6, 8, 10, 12 and 14 will be authorised.

Two completed authority prescriptions must be submitted with every initial application for this drug. For patients weighing 40 kg or greater, one prescription should be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats and the second prescription must be written for 2 doses of 40 mg and 2 repeats.

For patients weighing less than 40 kg, one prescription should be written for 2 doses of 40 mg with no repeats and the second prescription should be written for 2 doses of 20 mg with 3 repeats.

All tests and assessments should be performed preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior conventional treatment.

The most recent Mayo clinic, partial Mayo clinic or Paediatric Ulcerative Colitis Activity Index (PUCAI) score must be no more than 4 weeks old at the time of application.

A partial Mayo clinic or Paediatric Ulcerative Colitis Activity Index (PUCAI) assessment of the patient's response to this initial course of treatment must be made following a minimum of 12 weeks of treatment for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for golimumab, infliximab and vedolizumab so that there is adequate time for a response to be demonstrated.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Details of the accepted toxicities including severity can be found on the Department of Human Services website.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Initial 1 (new patient) or Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years) - balance of supply

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)]; OR
- Must be treated by a paediatrician; OR
- Must be treated by a specialist paediatric gastroenterologist.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 16 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

adalimumab 40 mg/0.8 mL injection, 6 x 0.8 mL pen devices

10945C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	3099.84	41.30	Humira [VE]

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL syringes

10944B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	1092.41	41.30	Humira [VE]

adalimumab 40 mg/0.8 mL injection, 6 x 0.8 mL syringes

10972L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	3099.84	41.30	Humira [VE]

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL pen devices

10955N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	1092.41	41.30	Humira [VE]

adalimumab 20 mg/0.4 mL injection, 2 x 0.4 mL syringes

11127P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	1092.41	41.30	Humira [VE]

■ ADALIMUMAB**Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE PSORIATIC ARTHRITIS**

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab for adult patients with severe active psoriatic arthritis. Therefore, where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient receiving PBS-subsidised treatment for psoriatic arthritis is able to commence a treatment cycle where they may trial biological medicines without having to experience a disease flare when swapping to the alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

A patient who received PBS-subsidised adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab or ustekinumab treatment prior to 1 May 2019 is considered to start their first cycle as of 1 May 2019.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence another cycle [further details are under '(5) Recommencement of treatment after a 5-year break in PBS-subsidised therapy' below].

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe active psoriatic arthritis.

(1) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an alternate medicine (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years) or An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, golimumab secukinumab and tofacitinib, 18 to 20 weeks of therapy for certolizumab pegol (depending upon the dosing regimen), 20 weeks of therapy for ixekizumab, 22 weeks of therapy for infliximab, and 28 weeks of therapy for ustekinumab. It is recommended that a patient be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of PBS-subsidised initial treatment following a minimum of 12 weeks of therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

Grandfather patients (ixekizumab only).

A patient who commenced treatment with ixekizumab for severe psoriatic arthritis prior to 1 March 2019 and who continues

to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction. A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

Grandfather patients (tofacitinib only).

A patient who commenced treatment with Tofacitinib for severe psoriatic arthritis prior to 1 May 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment under the Initial 1 or Initial 2 treatment restrictions.

For the second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to re-qualify with respect to either the indices of disease severity (i.e. erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP) level, and active joint count) or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify under the Initial 3 treatment restriction with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application.

However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

Within a treatment cycle a patient may alternate between therapy with any biological medicine of their choice (1 at a time) providing:

- (i) they have not received PBS-subsidised treatment with that particular biological medicine previously; or
- (ii) they have demonstrated an adequate response to that particular biological medicine if they have previously trialled it on the PBS; and
- (iii) they have not previously failed to respond to treatment 3 times in this treatment cycle with that biological medicine.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(4) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the indices of disease severity submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial or change or commencement treatment application is submitted within a treatment cycle and these revised baseline measurements will be used to assess response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to determine response for all subsequent continuing treatments. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. 20 or more active joints), response will be determined according to a reduction in the total number of active joints.

(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment restriction according to the criteria of the relevant restriction and index of disease severity. The application must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response to methotrexate at a dose of at least 20 mg weekly for a minimum period of 3 months, **AND**
- Patient must have failed to achieve an adequate response to sulfasalazine at a dose of at least 2 g per day for a minimum period of 3 months; OR
- Patient must have failed to achieve an adequate response to leflunomide at a dose of up to 20 mg daily for a minimum period of 3 months, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Where treatment with methotrexate, sulfasalazine or leflunomide is contraindicated according to the relevant TGA-approved Product Information, details must be provided at the time of application.

Where intolerance to treatment with methotrexate, sulfasalazine or leflunomide developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

The following initiation criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; and

either

- (a) an active joint count of at least 20 active (swollen and tender) joints; or
- (b) at least 4 active joints from the following list of major joints:
 - (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Severe Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

An assessment of a patient's response to an initial course of treatment must be conducted following a minimum of 12 weeks of therapy. An application for the continuing treatment must be accompanied with the assessment of response and submitted to the Department of Human Services no later than 4 weeks from the date of completion of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 3 months treatment with methotrexate and 3 months treatment with sulfasalazine or leflunomide can be found on the Department of Human Services website (www.humanservices.gov.au)

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial treatment - Initial 2 (change or recommencement of treatment after a break in in biological medicine of less than 5 years)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an erythrocyte sedimentation rate (ESR) no greater than 25 mm per hour or a C-reactive protein (CRP) level no greater than 15 mg per L or either marker reduced by at least 20% from baseline; and

either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following major active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Severe Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to change or recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial treatment - Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must have an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; OR
- The condition must have a C-reactive protein (CRP) level greater than 15 mg per L, **AND**
- The condition must have either (a) a total active joint count of at least 20 active (swollen and tender) joints; or (b) at least 4 active major joints, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Major joints are defined as (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

All measures of joint count and ESR and/or CRP must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Severe Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial 1 (new patient) or Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years) - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 16 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL syringes

9033K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	1092.41	41.30	Humira [VE]

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL pen devices

9101B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	1092.41	41.30	Humira [VE]

■ ADALIMUMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for adults with severe active rheumatoid arthritis. Where the term biological medicine appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alpha antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab), the T-cell co-stimulation modulator (abatacept) and the Janus kinase (JAK) inhibitors (baricitinib, tofacitinib, upadacitinib).

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised biological medicine therapy may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once, and
- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised biological medicines for the treatment of rheumatoid arthritis.

A serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Stevens Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered a treatment failure.

A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction.

A patient who has failed fewer than 5 biological medicines and who has a break in treatment of less than 24 months may commence a further course of treatment with a biological medicine under Initial 2 treatment restriction. A patient who has failed fewer than 5 biological medicines and who has had a break in therapy of longer than 24 months may commence a further course of treatment with a biological medicine under the Initial 3 treatment restriction. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine treatment is stopped to the date of the new application for treatment with a biological medicine.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 April 2019.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised biological medicine treatment and wishes to commence such therapy, excluding rituximab (Initial 1 - new patient); or
 - (ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months) [further details are under 'Swapping therapy' below]; or
 - (iii) a patient wishes to re-commence treatment with a specific biological medicine following a break of less than 24 months in PBS-subsidised therapy with that agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months).
 - (iv) a patient wishes to re-commence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 24 months (Initial 3 - re-commencement of treatment after a break in biological medicine of more than 24 months)
- Initial applications for a new patient (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6-month intensive DMARD trial, but prior to ceasing DMARD therapy.

Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, baricitinib, etanercept, golimumab, tocilizumab, tofacitinib and upadacitinib, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

Rituximab patients should be assessed following a minimum of 12 weeks after the first infusion, and the assessment should be submitted to Services Australia within 4 weeks to ensure continuity of treatment for those who meet the continuing restriction for PBS subsidised treatment with this drug for this condition.

Where a response assessment is not provided the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment. For second and subsequent courses of PBS-subsidised biological medicine (excluding rituximab) treatment, it is recommended that a patient be reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to Services Australia within sufficient time to allow processing.

Abatacept patients:
A patient is eligible to receive one I.V. loading dose when commencing treatment with the subcutaneous formulation. Two prescriptions are required, the first prescription for the I.V. loading dose for sufficient vials for one dose based on the patient's weight with no repeats. The second prescription for the subcutaneous formulation, with a maximum quantity of 4 and up to 3 repeats, must be submitted with the initial application.

Rituximab patients:

A further application may be submitted to Services Australia within sufficient time to allow processing. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

Assessment of the patient's response to treatment must be conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the cessation of the most recent course of biological medicine therapy.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have

demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction. Where a response assessment is not conducted within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

(2) Swapping therapy

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non- biological medicine therapy requirements except if the patient has had a break in therapy of more than 24 months who would need to requalify with respect to the indices of disease severity. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each biological medicine trialled.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent, unless they have experienced a serious adverse reaction of a severity necessitating permanent treatment withdrawal.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug.

Abatacept:

A patient swapping from I.V. abatacept to subcutaneous abatacept will not be eligible for an I.V. loading dose when commencing treatment with the subcutaneous formulation.

Rituximab:

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF-alfa antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they be assessed for response to every course of treatment, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological medicine during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological medicine therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate biological medicine may do so without having to have any treatment-free period.

(3) Baseline measurements to determine response.

Determination of whether a response to treatment has been demonstrated must be based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and the eligibility for continuing treatment must be assessed according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response must be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be determined on the total number of major joints.

Applications under the Initial 1 treatment restriction for a new patient must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. The results must be no more than 4 weeks old at the time of application.

Applications under the Initial 3 treatment restriction for re-commencement of treatment after a break in biological medicine of more than 24 months must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Applications for treatment with this drug where the dosing frequency exceeds 40 mg per fortnight will not be approved.

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with disease modifying anti-rheumatic drugs (DMARDs) which must include at least 3 months continuous treatment with each of at least 2 DMARDs, one of which must be methotrexate at a dose of at least 20 mg weekly and one of which must be: (i) hydroxychloroquine at a dose of at least 200 mg daily; or (ii) leflunomide at a dose of at least 10 mg daily; or (iii) sulfasalazine at a dose of at least 2 g daily; OR
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs which, if methotrexate is contraindicated according to the Therapeutic Goods Administration (TGA)-approved Product Information or cannot be tolerated at a 20 mg weekly dose, must include at least 3 months continuous treatment with each of at least 2 of the following DMARDs: (i) hydroxychloroquine at a dose of at least 200 mg daily; and/or (ii) leflunomide at a dose of at least 10 mg daily; and/or (iii) sulfasalazine at a dose of at least 2 g daily; OR

- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs which, if 3 or more of methotrexate, hydroxychloroquine, leflunomide and sulfasalazine are contraindicated according to the relevant TGA-approved Product Information or cannot be tolerated at the doses specified above, must include at least 3 months continuous treatment with each of at least 2 DMARDs, with one or more of the following DMARDs being used in place of the DMARDs which are contraindicated or not tolerated: (i) azathioprine at a dose of at least 1 mg/kg per day; and/or (ii) cyclosporin at a dose of at least 2 mg/kg/day; and/or (iii) sodium aurothiomalate at a dose of 50 mg weekly, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

If methotrexate is contraindicated according to the TGA-approved product information or cannot be tolerated at a 20 mg weekly dose, the application must include details of the contraindication or intolerance including severity to methotrexate. The maximum tolerated dose of methotrexate must be documented in the application, if applicable.

The application must include details of the DMARDs trialled, their doses and duration of treatment, and all relevant contraindications and/or intolerances including severity.

The requirement to trial at least 2 DMARDs for periods of at least 3 months each can be met using single agents sequentially or by using one or more combinations of DMARDs.

If the requirement to trial 6 months of intensive DMARD therapy with at least 2 DMARDs cannot be met because of contraindications and/or intolerances of a severity necessitating permanent treatment withdrawal to all of the DMARDs specified above, details of the contraindication or intolerance including severity and dose for each DMARD must be provided in the authority application.

The following criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; **AND** either

(a) a total active joint count of at least 20 active (swollen and tender) joints; or

(b) at least 4 active joints from the following list of major joints:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The joint count and ESR and/or CRP must be determined at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. All measures must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note The Department of Human Services website (www.humanservices.gov.au) has details of the toxicities, including severity, which will be accepted for the following purposes:

- (a) exempting a patient from the requirement to undertake a minimum 3 month trial of methotrexate at a 20 mg weekly dose;
- (b) substituting azathioprine, cyclosporin or sodium aurothiomalate for another DMARD as part of the 6 month intensive DMARD trial;

(c) exempting a patient from the requirement for a 6 month trial of intensive DMARD therapy.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 2 (change or re-commencement of treatment after a break in biological medicine of less than 24 months)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must not have failed to respond to previous PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised biological medicine treatment for this condition 5 times, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

- (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
- (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either of the Initial 1, Initial 2, Initial 3, or continuing treatment restrictions, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

A patient who has demonstrated a response to a course of rituximab must have a PBS-subsidised biological therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos. Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 3 (re-commencement of treatment after a break in biological medicine of more than 24 months)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**

- Patient must have a break in treatment of 24 months or more from the most recent PBS-subsidised biological medicine for this condition, **AND**
- Patient must not have failed to respond to previous PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised biological medicine treatment for this condition 5 times, **AND**
- The condition must have an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; OR
- The condition must have a C-reactive protein (CRP) level greater than 15 mg per L, **AND**
- The condition must have either (a) a total active joint count of at least 20 active (swollen and tender) joints; or (b) at least 4 active major joints, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Major joints are defined as (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

All measures of joint count and ESR and/or CRP must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial 1 (new patient) or Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 24 months) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 24 months) - balance of supply

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 24 months) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 24 months) to complete 16 weeks of treatment, **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL syringes

8737W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	1092.41	41.30	Humira [VE]

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL pen devices

9099X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	1092.41	41.30	Humira [VE]

■ ADALIMUMAB**Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS**

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years); or

(iii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or

(iv) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient be reviewed in the 4 weeks prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

Grandfather patients (risankizumab only).

A patient who commenced treatment with risankizumab for chronic plaque psoriasis prior to 1 December 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the Initial treatment Grandfather treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment. This assessment must be conducted within 4 weeks of the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the 4 weeks prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the 4 weeks prior to completing their current course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure patients receive the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

Services Australia will determine whether a response to treatment has been demonstrated based on the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 the PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Whole body (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis where lesions have been present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 5 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; (iii) ciclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks; (v) apremilast at a dose of 30 mg twice a day for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, ciclosporin, apremilast or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application.

Where intolerance to treatment with phototherapy, methotrexate, ciclosporin, apremilast or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, ciclosporin, apremilast, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

- (a) A current Psoriasis Area and Severity Index (PASI) score of greater than 15, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment.
- (b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 4 weeks following cessation of each course of treatment.
- (c) The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and
 - (ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, ciclosporin, apremilast or acitretin can be found on the Services Australia website (www.servicesaustralia.gov.au).

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos. Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Whole body (change or re-commencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and
 - (ii) details of prior biological treatment, including dosage, date and duration of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos. Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 3, Whole body (re-commencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must have a current Psoriasis Area and Severity Index (PASI) score of greater than 15, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos. Or mailed to:
Services Australia

Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Face, hand, foot (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 5 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; (iii) ciclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks; (v) apremilast at a dose of 30 mg twice a day for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, ciclosporin, apremilast or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application. Where intolerance to treatment with phototherapy, methotrexate, ciclosporin, apremilast or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, ciclosporin, apremilast, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

(a) Chronic plaque psoriasis classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:

(i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment; or

(ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment;

(b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 4 weeks following cessation of each course of treatment.

(c) The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition; and

(ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, ciclosporin, apremilast or acitretin can be found on the Services Australia website (www.servicesaustralia.gov.au).

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Face, hand, foot (change or re-commencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or
- (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition; and
 - (ii) details of prior biological treatment, including dosage, date and duration of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 3, Face, hand, foot (re-commencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**

- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must be classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where: (i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or (ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Whole body or Face, hand, foot (new patient) or Initial 2, Whole body or Face, hand, foot (change or re-commencement of treatment after a break in biological medicine of less than 5 years) or Initial 3, Whole body or Face, hand, foot (re-commencement of treatment after a break in biological medicine of more than 5 years) - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1, Whole body (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2, Whole body (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3, Whole body (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 1, Face, hand, foot (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2, Face, hand, foot (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3, Face, hand, foot (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 16 weeks treatment, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a dermatologist.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL syringes

9425C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	4	..	1092.41	41.30	Humira [VE]

adalimumab 40 mg/0.8 mL injection, 2 x 0.8 mL pen devices

9426D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	4	..	1092.41	41.30	Humira [VE]

■ **CERTOLIZUMAB PEGOL**

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial 1 (new patient) or Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 24 months) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 24 months - balance of supply).

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 18 to 20 weeks treatment, depending on the dosage regimen; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 24 months) restriction to complete 18 to 20 weeks treatment, depending on the dosage regimen; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 24 months) restriction to complete 18 to 20 weeks treatment, depending on the dosage regimen, **AND**
- The treatment must provide no more than the balance of up to 18 to 20 weeks treatment available under the above restrictions.

certolizumab pegol 200 mg/mL injection, 2 x 1 mL syringes

10892G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	1025.71	41.30	Cimzia [UC]

certolizumab pegol 200 mg/mL injection, 2 x 1 mL pen devices

11321W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	1025.71	41.30	Cimzia [UC]

■ **CERTOLIZUMAB PEGOL**

Note TREATMENT OF ADULT PATIENTS WITH ANKYLOSING SPONDYLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab for adult patients with ankylosing spondylitis.

Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 7 biological medicines at any 1 time.

Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy.

A patient who has been receiving PBS-subsidised adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab treatment prior to 1 December 2020 is considered to start their first cycle as of 1 December 2020.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last prescription for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised biological medicine treatment with adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised biological medicine treatment in this treatment cycle and wishes to commence such therapy (Initial 1 - New patient)

(ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years); or

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years). A patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy.

(b) Continuing treatment.

For the first continuing treatment course of PBS-subsidised biological medicine, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1, Initial 2 or Initial 3 treatment restrictions. For second and subsequent continuing courses of PBS-subsidised biological medicine it is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply. Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be conducted no later than 4 weeks from the cessation of that treatment course. Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI), or the prior NSAID therapy and exercise program requirements.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(3) Baseline measurements to determine response.

Services Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the BASDAI, ESR and/or CRP submitted with the first authority application for a biological medicine. For a new patient, the BASDAI used to determine the baseline must be measured while the patient is receiving NSAID therapy and completing their exercise program.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response.

Prescribers may provide new baseline measurements any time an 'Initial treatment' authority application is submitted within a treatment cycle and the eligibility for continuing treatment must be assessed according to these revised baseline measurements.

(4) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must qualify under the Initial 3 treatment restriction. The same clinical criteria and indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI) as for the Initial 1 (New patient) restriction will need to be met, but a re-trial of NSAID therapy and exercise therapy is not required.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Ankylosing spondylitis

Treatment Phase: Initial treatment - Initial 1 (new patient), Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years) - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 18 to 20 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 18 to 20 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 18 to 20 weeks treatment,

AND

- The treatment must provide no more than the balance of up to 18 to 20 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

certolizumab pegol 200 mg/mL injection, 2 x 1 mL syringes

10897M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	1025.71	41.30	Cimzia [UC]

certolizumab pegol 200 mg/mL injection, 2 x 1 mL pen devices

11318Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	1025.71	41.30	Cimzia [UC]

■ CERTOLIZUMAB PEGOL**Note TREATMENT OF ADULT PATIENTS WITH ANKYLOSING SPONDYLITIS**

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab for adult patients with ankylosing spondylitis.

Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 7 biological medicines at any 1 time.

Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy.

A patient who has been receiving PBS-subsidised adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab treatment prior to 1 December 2020 is considered to start their first cycle as of 1 December 2020.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last prescription for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised biological medicine treatment with adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised biological medicine treatment in this treatment cycle and wishes to commence such therapy (Initial 1 - New patient)

(ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years); or

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

A patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy.

(b) Continuing treatment.

For the first continuing treatment course of PBS-subsidised biological medicine, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1, Initial 2 or Initial 3 treatment restrictions. For second and subsequent continuing courses of PBS-subsidised biological medicine it is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply. Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be conducted no later than 4 weeks from the cessation of that treatment course. Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI), or the prior NSAID therapy and exercise program requirements.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological

medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(3) Baseline measurements to determine response.

Services Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the BASDAI, ESR and/or CRP submitted with the first authority application for a biological medicine. For a new patient, the BASDAI used to determine the baseline must be measured while the patient is receiving NSAID therapy and completing their exercise program.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response.

Prescribers may provide new baseline measurements any time an 'Initial treatment' authority application is submitted within a treatment cycle and the eligibility for continuing treatment must be assessed according to these revised baseline measurements.

(4) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must qualify under the Initial 3 treatment restriction. The same clinical criteria and indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI) as for the Initial 1 (New patient) restriction will need to be met, but a re-trial of NSAID therapy and exercise therapy is not required.

Authority required

Ankylosing spondylitis

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Ankylosing Spondylitis PBS Authority Application - Supporting Information Form.

An adequate response is defined as an improvement from baseline of at least 2 of the BASDAI and 1 of the following:

- (a) an ESR measurement no greater than 25 mm per hour; or
- (b) a CRP measurement no greater than 10 mg per L; or
- (c) an ESR or CRP measurement reduced by at least 20% from baseline.

Where only 1 acute phase reactant measurement is supplied in the first application for PBS-subsidised treatment, that same marker must be measured and supplied in all subsequent continuing treatment applications.

All measurements provided must be no more than 1 month old at the time of application.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Ankylosing spondylitis

Treatment Phase: Continuing treatment - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the Continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restriction.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

certolizumab pegol 200 mg/mL injection, 2 x 1 mL syringes

10137M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1025.71	41.30	Cimzia [UC]

certolizumab pegol 200 mg/mL injection, 2 x 1 mL pen devices

11320T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1025.71	41.30	Cimzia [UC]

▪ **CERTOLIZUMAB PEGOL**

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE PSORIATIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab for adult patients with severe active psoriatic arthritis. Therefore, where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient receiving PBS-subsidised treatment for psoriatic arthritis is able to commence a treatment cycle where they may trial biological medicines without having to experience a disease flare when swapping to the alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

A patient who received PBS-subsidised adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab or ustekinumab treatment prior to 1 May 2019 is considered to start their first cycle as of 1 May 2019.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence another cycle [further details are under '(5) Recommencement of treatment after a 5-year break in PBS-subsidised therapy' below].

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe active psoriatic arthritis.

(1) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an alternate medicine (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years) or An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, golimumab secukinumab and tofacitinib, 18 to 20 weeks of therapy for certolizumab pegol (depending upon the dosing regimen), 20 weeks of therapy for ixekizumab, 22 weeks of therapy for infliximab, and 28 weeks of therapy for ustekinumab. It is recommended that a patient be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of PBS-subsidised initial treatment following a minimum of 12 weeks of

therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

Grandfather patients (ixekizumab only).

A patient who commenced treatment with ixekizumab for severe psoriatic arthritis prior to 1 March 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

Grandfather patients (tofacitinib only).

A patient who commenced treatment with Tofacitinib for severe psoriatic arthritis prior to 1 May 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment under the Initial 1 or Initial 2 treatment restrictions.

For the second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to re-qualify with respect to either the indices of disease severity (i.e. erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP) level, and active joint count) or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify under the Initial 3 treatment restriction with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application.

However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

Within a treatment cycle a patient may alternate between therapy with any biological medicine of their choice (1 at a time) providing:

- (i) they have not received PBS-subsidised treatment with that particular biological medicine previously; or
- (ii) they have demonstrated an adequate response to that particular biological medicine if they have previously trialled it on the PBS; and
- (iii) they have not previously failed to respond to treatment 3 times in this treatment cycle with that biological medicine.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(4) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the indices of disease severity submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and these revised baseline measurements will be used to assess response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to determine response for all subsequent continuing treatments. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. 20 or more active joints), response will be determined according to a reduction in the total number of active joints.

(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment restriction according to the criteria of the relevant restriction and index of disease severity. The application must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial 1 (new patient) or Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years) - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 18 to 20 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 18 to 20 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 18 to 20 weeks treatment,

AND

- The treatment must provide no more than the balance of up to 18 to 20 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

certolizumab pegol 200 mg/mL injection, 2 x 1 mL syringes

10896L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	1025.71	41.30	Cimzia [UC]

certolizumab pegol 200 mg/mL injection, 2 x 1 mL pen devices

11326D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	1025.71	41.30	Cimzia [UC]

■ CERTOLIZUMAB PEGOL

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for adults with severe active rheumatoid arthritis. Where the term biological medicine appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alfa antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab), the T-cell co-stimulation modulator (abatacept) and the Janus kinase (JAK) inhibitors (baricitinib, tofacitinib, upadacitinib).

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised biological medicine therapy may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once, and
- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised biological medicines for the treatment of rheumatoid arthritis.

A serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Stevens Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered a treatment failure.

A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction.

A patient who has failed fewer than 5 biological medicines and who has a break in treatment of less than 24 months may commence a further course of treatment with a biological medicine under Initial 2 treatment restriction. A patient who has failed fewer than 5 biological medicines and who has had a break in therapy of longer than 24 months may commence a further course of treatment with a biological medicine under the Initial 3 treatment restriction. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine treatment is stopped to the date of the new application for treatment with a biological medicine.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 April 2019.

(a) Initial treatment.

Applications for initial treatment should be made where:

- a patient has received no prior PBS-subsidised biological medicine treatment and wishes to commence such therapy, excluding rituximab (Initial 1 - new patient); or
- a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months) [further details are under 'Swapping therapy' below]; or
- a patient wishes to re-commence treatment with a specific biological medicine following a break of less than 24 months in PBS-subsidised therapy with that agent (Initial 2 - change or re-commencement of treatment after a break in biological

medicine of less than 24 months).

(iv) a patient wishes to re-commence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 24 months (Initial 3 - re-commencement of treatment after a break in biological medicine of more than 24 months) Initial applications for a new patient (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6-month intensive DMARD trial, but prior to ceasing DMARD therapy.

Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, baricitinib, etanercept, golimumab, tocilizumab, tofacitinib and upadacitinib, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

Rituximab patients should be assessed following a minimum of 12 weeks after the first infusion, and the assessment should be submitted to Services Australia within 4 weeks to ensure continuity of treatment for those who meet the continuing restriction for PBS subsidised treatment with this drug for this condition.

Where a response assessment is not provided the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment. For second and subsequent courses of PBS-subsidised biological medicine (excluding rituximab) treatment, it is recommended that a patient be reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to Services Australia within sufficient time to allow processing.

Abatacept patients:

A patient is eligible to receive one I.V. loading dose when commencing treatment with the subcutaneous formulation. Two prescriptions are required, the first prescription for the I.V. loading dose for sufficient vials for one dose based on the patient's weight with no repeats. The second prescription for the subcutaneous formulation, with a maximum quantity of 4 and up to 3 repeats, must be submitted with the initial application.

Rituximab patients:

A further application may be submitted to Services Australia within sufficient time to allow processing. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

Assessment of the patient's response to treatment must be conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the cessation of the most recent course of biological medicine therapy.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction. Where a response assessment is not conducted within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

(2) Swapping therapy

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non- biological medicine therapy requirements except if the patient has had a break in therapy of more than 24 months who would need to requalify with respect to the indices of disease severity. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each biological medicine trialled.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent, unless they have experienced a serious adverse reaction of a severity necessitating permanent treatment withdrawal.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug.

Abatacept:

A patient swapping from I.V. abatacept to subcutaneous abatacept will not be eligible for an I.V. loading dose when commencing treatment with the subcutaneous formulation.

Rituximab:

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF-alfa antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they be assessed for response to every course of treatment, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological medicine during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological medicine therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate biological medicine may do so without having to have any treatment-free period.

(3) Baseline measurements to determine response.

Determination of whether a response to treatment has been demonstrated must be based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and the eligibility for continuing treatment must be assessed according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint

count is based on total active joints (i.e. more than 20 active joints), response must be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be determined on the total number of major joints.

Applications under the Initial 1 treatment restriction for a new patient must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. The results must be no more than 4 weeks old at the time of application.

Applications under the Initial 3 treatment restriction for re-commencement of treatment after a break in biological medicine of more than 24 months must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment per continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient has either failed or ceased to respond to a PBS-subsidised biological medicine for this condition 5 times, they will not be eligible to receive further PBS-subsidised treatment with a biological medicine for this condition.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Continuing Treatment - balance of supply.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restriction.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

certolizumab pegol 200 mg/mL injection, 2 x 1 mL syringes

3425G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1025.71	41.30	Cimzia [UC]

certolizumab pegol 200 mg/mL injection, 2 x 1 mL pen devices

11325C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1025.71	41.30	Cimzia [UC]

■ CERTOLIZUMAB PEGOL

Note TREATMENT OF ADULT PATIENTS WITH NON-RADIOGRAPHIC AXIAL SPONDYLOARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of secukinumab, certolizumab pegol and golimumab for adult patients with non-radiographic axial spondyloarthritis.

Where the term 'biological medicine' appears in notes and restrictions, it refers to secukinumab, certolizumab pegol and golimumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 3 biological medicines at any one time.

Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy.

A serious adverse reaction requiring permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once.

Once a patient has either failed or ceased to respond to treatment 3 times (once with any biological medicine) within the same treatment cycle, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last prescription for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle.

A patient who has failed 3 trials or fewer of biological medicine therapy in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment with secukinumab, certolizumab pegol and golimumab

(1) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised biological medicine treatment in this treatment cycle and wishes to commence such therapy (Initial 1 - New patient)

(ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years); or

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

With the exception of grandfathered patients, a patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy.

(2) Continuing treatment.

For continuing courses of PBS-subsidised biological medicine it is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that biological medicine provided they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment in courses of up to 24 weeks provided they continue to sustain an adequate response.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be conducted no later than 4 weeks from the cessation of that treatment course. Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction requiring permanent treatment withdrawal.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. the C-reactive protein (CRP) level and the BASDAI score), or the prior NSAID therapy and exercise program requirements.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction requiring permanent

treatment withdrawal.

(4) Baseline measurements to determine response.

A response to treatment is based on the baseline BASDAI score and CRP level documented in the patient's medical records. For a new patient, the BASDAI score used to determine baseline disease severity must be measured while the patient is receiving NSAID therapy and completing their exercise program.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications.

Prescribers may provide new baseline measurements any time that an 'Initial treatment' authority application is submitted and the eligibility for continuing treatment must be assessed according to these revised baseline measurements.

(5) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must qualify under the Initial 3 treatment restriction. The same clinical criteria and indices of disease severity (i.e. the C-reactive protein (CRP) levels and the BASDAI) as for the Initial 1 (New patient) restriction will need to be met, but a re-trial of NSAID therapy and exercise therapy is not required.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Non-radiographic axial spondyloarthritis

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug for this condition, **AND**
- The treatment must not exceed a maximum of 24 weeks with this drug per authorised course under this restriction.

Treatment criteria:

- Must be treated by a rheumatologist; OR
 - Must be treated by a clinical immunologist with expertise in the management of non-radiographic axial spondyloarthritis.
- An adequate response to therapy with this biological medicine is defined as a reduction from baseline in the Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) score by 2 or more units (on a scale of 0-10) and 1 of the following:
- (a) a CRP measurement no greater than 10 mg per L; or
 - (b) a CRP measurement reduced by at least 20% from baseline.

If the requirement to demonstrate an elevated CRP level could not be met under an initial treatment restriction, a reduction in the BASDAI score from baseline will suffice for the purposes of administering this continuing treatment restriction.

The patient remains eligible to receive continuing treatment with the same biological medicine in courses of up to 24 weeks providing they continue to sustain an adequate response. It is recommended that a patient be reviewed in the month prior to completing their current course of treatment.

certolizumab pegol 200 mg/mL injection, 2 x 1 mL syringes

12005W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1025.71	41.30	Cimzia [UC]

certolizumab pegol 200 mg/mL injection, 2 x 1 mL pen devices

12028C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1025.71	41.30	Cimzia [UC]

■ CERTOLIZUMAB PEGOL

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for adults with severe active rheumatoid arthritis. Where the term biological medicine appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alfa antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab), the T-cell co-stimulation modulator (abatacept) and the Janus kinase (JAK) inhibitors (baricitinib, tofacitinib, upadacitinib).

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised biological medicine therapy may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once, and
- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised biological medicines for the treatment of rheumatoid arthritis.

A serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Stevens Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered a treatment failure.

A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is

sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction. A patient who has failed fewer than 5 biological medicines and who has a break in treatment of less than 24 months may commence a further course of treatment with a biological medicine under Initial 2 treatment restriction. A patient who has failed fewer than 5 biological medicines and who has had a break in therapy of longer than 24 months may commence a further course of treatment with a biological medicine under the Initial 3 treatment restriction. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine treatment is stopped to the date of the new application for treatment with a biological medicine.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 April 2019.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised biological medicine treatment and wishes to commence such therapy, excluding rituximab (Initial 1 - new patient); or

(ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to re-commence treatment with a specific biological medicine following a break of less than 24 months in PBS-subsidised therapy with that agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months).

(iv) a patient wishes to re-commence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 24 months (Initial 3 - re-commencement of treatment after a break in biological medicine of more than 24 months). Initial applications for a new patient (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6-month intensive DMARD trial, but prior to ceasing DMARD therapy.

Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, baricitinib, etanercept, golimumab, tocilizumab, tofacitinib and upadacitinib, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

Rituximab patients should be assessed following a minimum of 12 weeks after the first infusion, and the assessment should be submitted to Services Australia within 4 weeks to ensure continuity of treatment for those who meet the continuing restriction for PBS subsidised treatment with this drug for this condition.

Where a response assessment is not provided the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment. For second and subsequent courses of PBS-subsidised biological medicine (excluding rituximab) treatment, it is recommended that a patient be reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to Services Australia within sufficient time to allow processing.

Abatacept patients:

A patient is eligible to receive one I.V. loading dose when commencing treatment with the subcutaneous formulation. Two prescriptions are required, the first prescription for the I.V. loading dose for sufficient vials for one dose based on the patient's weight with no repeats. The second prescription for the subcutaneous formulation, with a maximum quantity of 4 and up to 3 repeats, must be submitted with the initial application.

Rituximab patients:

A further application may be submitted to Services Australia within sufficient time to allow processing. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

Assessment of the patient's response to treatment must be conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the cessation of the most recent course of biological medicine therapy.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction. Where a response assessment is not conducted within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

(2) Swapping therapy

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non-biological medicine therapy requirements except if the patient has had a break in therapy of more than 24 months who would need to requalify with respect to the indices of disease severity. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each biological medicine trialled.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent, unless they have experienced a serious adverse reaction of a severity necessitating permanent treatment withdrawal.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug.

Abatacept:

A patient swapping from I.V. abatacept to subcutaneous abatacept will not be eligible for an I.V. loading dose when commencing treatment with the subcutaneous formulation.

Rituximab:

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF-alfa antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they be assessed for response to every course of treatment, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological medicine during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological medicine therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate biological medicine may do so without having to have any treatment-free period.

(3) Baseline measurements to determine response.

Determination of whether a response to treatment has been demonstrated must be based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and the eligibility for continuing treatment must be assessed according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response must be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be determined on the total number of major joints.

Applications under the Initial 1 treatment restriction for a new patient must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. The results must be no more than 4 weeks old at the time of application.

Applications under the Initial 3 treatment restriction for re-commencement of treatment after a break in biological medicine of more than 24 months must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos. Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with disease modifying anti-rheumatic drugs (DMARDs) which must include at least 3 months continuous treatment with each of at least 2 DMARDs, one of which must be methotrexate at a dose of at least 20 mg weekly and one of which must be: (i) hydroxychloroquine at a dose of at least 200 mg daily; or (ii) leflunomide at a dose of at least 10 mg daily; or (iii) sulfasalazine at a dose of at least 2 g daily; OR
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs which, if methotrexate is contraindicated according to the Therapeutic Goods Administration (TGA)-approved Product Information or cannot be tolerated at a 20 mg weekly dose, must include at least 3 months continuous treatment with each of at least 2 of the following DMARDs: (i) hydroxychloroquine at a dose of at least 200 mg daily; and/or (ii) leflunomide at a dose of at least 10 mg daily; and/or (iii) sulfasalazine at a dose of at least 2 g daily; OR
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs which, if 3 or more of methotrexate, hydroxychloroquine, leflunomide and sulfasalazine are contraindicated according to the relevant TGA-approved Product Information or cannot be tolerated at the doses specified above, must include at least 3 months continuous treatment with each of at least 2 DMARDs, with one or more of the following DMARDs being used in place of the DMARDs which are contraindicated or not tolerated: (i) azathioprine at a dose of at least 1 mg/kg per day; and/or (ii) cyclosporin at a dose of at least 2 mg/kg/day; and/or (iii) sodium aurothiomalate at a dose of 50 mg weekly, **AND**
- Patient must not receive more than 18 to 20 weeks of treatment, depending on the dosage regimen, under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

If methotrexate is contraindicated according to the TGA-approved product information or cannot be tolerated at a 20 mg weekly dose, the application must include details of the contraindication or intolerance including severity to methotrexate. The maximum tolerated dose of methotrexate must be documented in the application, if applicable.

The application must include details of the DMARDs trialled, their doses and duration of treatment, and all relevant contraindications and/or intolerances including severity.

The requirement to trial at least 2 DMARDs for periods of at least 3 months each can be met using single agents sequentially or by using one or more combinations of DMARDs.

If the requirement to trial 6 months of intensive DMARD therapy with at least 2 DMARDs cannot be met because of contraindications and/or intolerances of a severity necessitating permanent treatment withdrawal to all of the DMARDs specified above, details of the contraindication or intolerance including severity and dose for each DMARD must be provided in the authority application.

The following criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; AND either

- (a) a total active joint count of at least 20 active (swollen and tender) joints; or
- (b) at least 4 active joints from the following list of major joints:
 - (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The joint count and ESR and/or CRP must be determined at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. All measures must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note The Department of Human Services website (www.humanservices.gov.au) has details of the toxicities, including severity, which will be accepted for the following purposes:

- (a) exempting a patient from the requirement to undertake a minimum 3 month trial of methotrexate at a 20 mg weekly dose;
- (b) substituting azathioprine, cyclosporin or sodium aurothiomalate for another DMARD as part of the 6 month intensive DMARD trial;
- (c) exempting a patient from the requirement for a 6 month trial of intensive DMARD therapy.

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 2 (change or re-commencement of treatment after a break in biological medicine of less than 24 months).

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must not have failed to respond to previous PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised biological medicine treatment for this condition 5 times, **AND**
- Patient must not receive more than 18 to 20 weeks of treatment, depending on the dosage regimen, under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

- (a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or
- (b) a reduction in the number of the following active joints, from at least 4, by at least 50%:
 - (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either of the Initial 1, Initial 2, Initial 3, or continuing treatment restrictions, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

A patient who has demonstrated a response to a course of rituximab must have a PBS-subsidised biological therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine.

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 3 (re-commencement of treatment after a break in biological medicine of more than 24 months)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 24 months or more from the most recent PBS-subsidised biological medicine for this condition, **AND**
- Patient must not have failed to respond to previous PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised biological medicine treatment for this condition 5 times, **AND**
- The condition must have an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; OR
- The condition must have a C-reactive protein (CRP) level greater than 15 mg per L, **AND**
- The condition must have either (a) a total active joint count of at least 20 active (swollen and tender) joints; or (b) at least 4 active major joints, **AND**
- Patient must not receive more than 18 to 20 weeks of treatment, depending on the dosage regimen, under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Major joints are defined as (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

All measures of joint count and ESR and/or CRP must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

certolizumab pegol 200 mg/mL injection, 2 x 1 mL syringes

10905Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	..	..	*3017.55	41.30	Cimzia [UC]

certolizumab pegol 200 mg/mL injection, 2 x 1 mL pen devices

11322X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	..	..	*3017.55	41.30	Cimzia [UC]

■ CERTOLIZUMAB PEGOL**Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE PSORIATIC ARTHRITIS**

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab for adult patients with severe active psoriatic arthritis. Therefore, where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient receiving PBS-subsidised treatment for psoriatic arthritis is able to commence a treatment cycle where they may trial biological medicines without having to experience a disease flare when swapping to the alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

A patient who received PBS-subsidised adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab or ustekinumab treatment prior to 1 May 2019 is considered to start their first cycle as of 1 May 2019.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence another cycle [further details are under '(5) Recommencement of treatment after a 5-year break in PBS-subsidised therapy' below].

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe active psoriatic arthritis.

(1) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an alternate medicine (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years) or An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, golimumab secukinumab and tofacitinib, 18 to 20 weeks of therapy for certolizumab pegol (depending upon the dosing regimen), 20 weeks of therapy for ixekizumab, 22 weeks of therapy for infliximab, and 28 weeks of therapy for ustekinumab. It is recommended that a patient be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of PBS-subsidised initial treatment following a minimum of 12 weeks of therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

Grandfather patients (ixekizumab only).

A patient who commenced treatment with ixekizumab for severe psoriatic arthritis prior to 1 March 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

Grandfather patients (tofacitinib only).

A patient who commenced treatment with Tofacitinib for severe psoriatic arthritis prior to 1 May 2019 and who continues to

receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction. A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment under the Initial 1 or Initial 2 treatment restrictions.

For the second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to re-qualify with respect to either the indices of disease severity (i.e. erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP) level, and active joint count) or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify under the Initial 3 treatment restriction with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application.

However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

Within a treatment cycle a patient may alternate between therapy with any biological medicine of their choice (1 at a time) providing:

- (i) they have not received PBS-subsidised treatment with that particular biological medicine previously; or
- (ii) they have demonstrated an adequate response to that particular biological medicine if they have previously trialled it on the PBS; and
- (iii) they have not previously failed to respond to treatment 3 times in this treatment cycle with that biological medicine.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(4) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the indices of disease severity submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial or change or commencement treatment application is submitted within a treatment cycle and these revised baseline measurements will be used to assess response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to determine response for all subsequent continuing treatments. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. 20 or more active joints), response will be determined according to a reduction in the total number of active joints.

(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment restriction according to the criteria of the relevant restriction and index of disease severity. The application must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Severe psoriatic arthritis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**

- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an erythrocyte sedimentation rate (ESR) no greater than 25 mm per hour or a C-reactive protein (CRP) level no greater than 15 mg per L or either marker reduced by at least 20% from baseline; and

either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following major active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to determine response for all subsequent continuing treatments.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Severe Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either Initial 1, Initial 2, or Initial 3 treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Continuing treatment - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restriction.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

certolizumab pegol 200 mg/mL injection, 2 x 1 mL syringes

10238W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1025.71	41.30	Cimzia [UC]

certolizumab pegol 200 mg/mL injection, 2 x 1 mL pen devices

11324B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1025.71	41.30	Cimzia [UC]

■ CERTOLIZUMAB PEGOL

Note TREATMENT OF ADULT PATIENTS WITH NON-RADIOGRAPHIC AXIAL SPONDYLOARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of secukinumab, certolizumab pegol and golimumab for adult patients with non-radiographic axial spondyloarthritis.

Where the term 'biological medicine' appears in notes and restrictions, it refers to secukinumab, certolizumab pegol and golimumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 3 biological medicines at any one time.

Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy.

A serious adverse reaction requiring permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once.

Once a patient has either failed or ceased to respond to treatment 3 times (once with any biological medicine) within the same treatment cycle, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last prescription for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle.

A patient who has failed 3 trials or fewer of biological medicine therapy in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment with secukinumab, certolizumab pegol and golimumab

(1) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised biological medicine treatment in this treatment cycle and wishes to commence such therapy (Initial 1 - New patient)

(ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years); or

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

With the exception of grandfathered patients, a patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy.

(2) Continuing treatment.

For continuing courses of PBS-subsidised biological medicine it is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that biological medicine provided they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment in courses of up to 24 weeks provided they continue to sustain an adequate response.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be conducted no later than 4 weeks from the cessation of that treatment course. Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction requiring permanent treatment withdrawal.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. the C-reactive protein (CRP) level and the BASDAI score), or the prior NSAID therapy and exercise program requirements.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction requiring permanent treatment withdrawal.

(4) Baseline measurements to determine response.

A response to treatment is based on the baseline BASDAI score and CRP level documented in the patient's medical records. For a new patient, the BASDAI score used to determine baseline disease severity must be measured while the patient is receiving NSAID therapy and completing their exercise program.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications.

Prescribers may provide new baseline measurements any time that an 'Initial treatment' authority application is submitted and the eligibility for continuing treatment must be assessed according to these revised baseline measurements.

(5) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must qualify under the Initial 3 treatment restriction. The same clinical criteria and indices of disease severity (i.e. the C-reactive protein (CRP) levels and the BASDAI) as for the Initial 1 (New patient) restriction will need to be met, but a re-trial of NSAID therapy and exercise therapy is not required.

Note No increase in the maximum quantity or number of units may be authorised.

Authority required

Non-radiographic axial spondyloarthritis

Treatment Phase: Initial 1 (New patient), Initial 2 (Change or re-commencement of treatment after a break in biological medicine of less than 5 years) or Initial 3 (Recommencement of treatment after a break in biological medicine of more than 5 years) - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 18 to 20 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 18 to 20 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 18 to 20 weeks treatment,

AND

- The treatment must provide no more than the balance of up to 20 weeks treatment.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of non-radiographic axial spondyloarthritis.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Non-radiographic axial spondyloarthritis

Treatment Phase: Continuing treatment or Grandfather patient - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Continuing treatment restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Grandfathered treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the continuing treatment restriction or the grandfather restriction.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of non-radiographic axial spondyloarthritis.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Non-radiographic axial spondyloarthritis

Treatment Phase: Grandfather treatment

Clinical criteria:

- Patient must have received non-PBS-subsidised treatment with this drug for this PBS indication prior to 1 June 2020, **AND**
- Patient must have had chronic lower back pain and stiffness for 3 or more months that was relieved by exercise but not rest, prior to initiating non-PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have had failed to achieve an adequate response following treatment with at least 2 non-steroidal anti-inflammatory drugs (NSAIDs), whilst completing an appropriate exercise program, for a total period of 3 months, prior to initiating non-PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have had one or more of the following: (a) enthesitis (heel); (b) uveitis; (c) dactylitis; (d) psoriasis; (e) inflammatory bowel disease; or (f) positive for Human Leukocyte Antigen B27 (HLA-B27); prior to initiating non-PBS-subsidised treatment with this drug for this condition, **AND**
- The condition must not be radiographically evidenced on plain x-ray of Grade II bilateral sacroiliitis or Grade III or IV unilateral sacroiliitis prior to commencing non-PBS-subsidised treatment with this biological medicine, **AND**
- The condition must have been diagnosed as non-radiographic axial spondyloarthritis, as defined by Assessment of Spondyloarthritis International Society (ASAS) criteria, prior to having commenced non-PBS-subsidised treatment with this biological medicine, **AND**
- The condition must have been sacroiliitis with active inflammation and/or oedema on non-contrast Magnetic Resonance Imaging (MRI) prior to commencing non-PBS-subsidised treatment with this biological medicine, **AND**
- The condition must have had presence of Bone Marrow Oedema (BMO) depicted as a hyperintense signal on a Short Tau Inversion Recovery (STIR) image (or equivalent) prior to commencing non-PBS-subsidised treatment with this biological medicine, **AND**
- The condition must have had BMO depicted as a hypointense signal on a T1 weighted image (without gadolinium) prior to commencing non-PBS-subsidised treatment with this biological medicine, **AND**
- The treatment must not exceed a maximum of 24 weeks with this drug under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR

- Must be treated by a clinical immunologist with expertise in the management of non-radiographic axial spondyloarthritis. The application must include details of the NSAIDs trialled, their doses and duration of treatment.

If the NSAID dose is less than the maximum recommended dose in the relevant TGA-approved Product Information, the application must include the reason a higher dose cannot be used.

If treatment with NSAIDs is contraindicated according to the relevant TGA-approved Product Information, the application must provide details of the contraindication.

If intolerance to NSAID treatment develops during the relevant period of use which is of a severity to necessitate permanent treatment withdrawal, the application must provide details of the nature and severity of this intolerance.

The following criteria indicate failure to achieve an adequate response to NSAIDs and must have been demonstrated prior to initiation of non-PBS subsidised treatment with this biological medicine for this condition:

- (a) a Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) score of at least 4 on a 0-10 scale; and
- (b) C-reactive protein (CRP) level greater than 10 mg per L.

The BASDAI score and CRP level must have been determined at the completion of the 3-month NSAID and exercise trial, but prior to ceasing NSAID treatment. All measures must have been no more than 1 month old at the time of initiating non-PBS subsidised treatment with this biological medicine for this condition.

If the requirement to demonstrate an elevated CRP level could not be met, the reason must be stated in the application. Treatment with prednisolone dosed at 7.5 mg or higher daily (or equivalent) or a parenteral steroid within the past month (intramuscular or intravenous methylprednisolone or equivalent) is an acceptable reason.

The assessment of the patient's response to the initial course of treatment must be conducted following a minimum of 12 weeks of treatment and no later than 4 weeks from the cessation of that treatment course. If the response assessment is not conducted within these timeframes, the patient will be deemed to have failed this course of treatment in this treatment cycle.

A Grandfathered patient may qualify for PBS-subsidised treatment under this restriction once only. For continuing PBS-subsidised treatment, a Grandfathered patient must qualify under the continuing treatment criteria.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Non-radiographic axial spondyloarthritis Grandfathered PBS Authority Application - Supporting Information Form which seeks details of:
 - (i) a copy of the radiological report confirming the absence of Grade II bilateral sacroiliitis or Grade III or IV unilateral sacroiliitis; and
 - (ii) a BASDAI score and CRP level that substantiates failure to achieve an adequate response to NSAIDs prior to initiating non-PBS subsidised treatment with this biological medicine for this condition; and
 - (iii) the MRI report; and
 - (iv) the NSAIDs trialled, their doses and duration of treatment. If applicable, the reason a higher dose cannot be used where the NSAID dose is less than the maximum recommended dose in the relevant TGA-approved Product Information or details of the contraindication or intolerance according to the relevant TGA-approved Product Information must be included.

The baseline BASDAI score and CRP level must also be documented in the patient's medical records.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

certolizumab pegol 200 mg/mL injection, 2 x 1 mL syringes

12040Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	1025.71	41.30	Cimzia [UC]

certolizumab pegol 200 mg/mL injection, 2 x 1 mL pen devices

12013G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	1025.71	41.30	Cimzia [UC]

■ CERTOLIZUMAB PEGOL

Note TREATMENT OF ADULT PATIENTS WITH ANKYLOSING SPONDYLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab for adult patients with ankylosing spondylitis.

Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 7 biological medicines at any 1 time.

Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy.

A patient who has been receiving PBS-subsidised adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab treatment prior to 1 December 2020 is considered to start their first cycle as of 1 December 2020.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological

medicine more than once.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last prescription for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised biological medicine treatment with adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised biological medicine treatment in this treatment cycle and wishes to commence such therapy (Initial 1 - New patient)

(ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years); or

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

A patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy.

(b) Continuing treatment.

For the first continuing treatment course of PBS-subsidised biological medicine, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1, Initial 2 or Initial 3 treatment restrictions. For second and subsequent continuing courses of PBS-subsidised biological medicine it is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply. Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be conducted no later than 4 weeks from the cessation of that treatment course. Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI), or the prior NSAID therapy and exercise program requirements.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(3) Baseline measurements to determine response.

Services Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the BASDAI, ESR and/or CRP submitted with the first authority application for a biological medicine. For a new patient, the BASDAI used to determine the baseline must be measured while the patient is receiving NSAID therapy and completing their exercise program.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response.

Prescribers may provide new baseline measurements any time an 'Initial treatment' authority application is submitted within a treatment cycle and the eligibility for continuing treatment must be assessed according to these revised baseline measurements.

(4) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must qualify under the Initial 3 treatment restriction. The same clinical criteria and indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI) as for the Initial 1 (New patient) restriction will need to be met, but a re-trial of NSAID therapy and exercise therapy is not required.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available

on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Ankylosing spondylitis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Clinical criteria:

- The condition must be radiographically (plain X-ray) confirmed Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have at least 2 of the following: (i) low back pain and stiffness for 3 or more months that is relieved by exercise but not by rest; or (ii) limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by a score of at least 1 on each of the lumbar flexion and lumbar side flexion measurements of the Bath Ankylosing Spondylitis Metrology Index (BASMI); or (iii) limitation of chest expansion relative to normal values for age and gender, **AND**
- Patient must have failed to achieve an adequate response following treatment with at least 2 non-steroidal anti-inflammatory drugs (NSAIDs), whilst completing an appropriate exercise program, for a total period of 3 months, **AND**
- Patient must not receive more than 18 to 20 weeks of treatment, depending on the dosage regimen, under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

The application must include details of the NSAIDs trialed, their doses and duration of treatment.

If the NSAID dose is less than the maximum recommended dose in the relevant TGA-approved Product Information, the application must include the reason a higher dose cannot be used.

If treatment with NSAIDs is contraindicated according to the relevant TGA-approved Product Information, the application must provide details of the contraindication.

If intolerance to NSAID treatment develops during the relevant period of use which is of a severity to necessitate permanent treatment withdrawal, the application must provide details of the nature and severity of this intolerance.

The following criteria indicate failure to achieve an adequate response and must be demonstrated at the time of the initial application:

- (a) a Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 4 on a 0-10 scale; AND
- (b) an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 10 mg per L.

The BASDAI must be determined at the completion of the 3 month NSAID and exercise trial, but prior to ceasing NSAID treatment. The BASDAI must be no more than 1 month old at the time of initial application.

Both ESR and CRP measures should be provided with the initial treatment application and both must be no more than 1 month old. If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reason this criterion cannot be satisfied.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Ankylosing Spondylitis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) a copy of the radiological report confirming Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis; and
 - (ii) a completed BASDAI Assessment Form; and
 - (iii) a completed Exercise Program Self Certification Form included in the supporting information form.

An assessment of a patient's response to an initial course of treatment must be conducted following a minimum of 12 weeks of therapy. An application for the continuing treatment must be accompanied with the assessment of response and submitted to the Department of Human Services no later than 4 weeks from the date of completion of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Details of the toxicities, including severity, which will be accepted for the purposes of administering this restriction can be found on the Department of Human Services website at www.humanservices.gov.au

Note For details on the appropriate minimum exercise program that will be accepted for the purposes of administering this restriction, please refer to the Department of Human Services website at www.humanservices.gov.au

Authority required

Ankylosing spondylitis

Treatment Phase: Initial treatment - Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- Patient must not receive more than 18 to 20 weeks of treatment, depending on the dosage regimen, under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Ankylosing Spondylitis PBS Authority Application - Supporting Information Form.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to change or recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

An adequate response is defined as an improvement from baseline of at least 2 of the BASDAI and 1 of the following:

- (a) an ESR measurement no greater than 25 mm per hour; or
- (b) a CRP measurement no greater than 10 mg per L; or
- (c) an ESR or CRP measurement reduced by at least 20% from baseline.

Where only 1 acute phase reactant measurement is supplied in the first application for PBS-subsidised treatment, that same marker must be measured and supplied in all subsequent continuing treatment applications.

All measurements provided must be no more than 1 month old at the time of application.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Authority required

Ankylosing spondylitis

Treatment Phase: Initial treatment - Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must be radiographically (plain X-ray) confirmed Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis, **AND**
- Patient must have at least 2 of the following: (i) low back pain and stiffness for 3 or more months that is relieved by exercise but not by rest; or (ii) limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by a score of at least 1 on each of the lumbar flexion and lumbar side flexion measurements of the Bath Ankylosing Spondylitis Metrology Index (BASMI); or (iii) limitation of chest expansion relative to normal values for age and gender,

AND

- Patient must have a Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 4 on a 0-10 scale that is no more than 4 weeks old at the time of application, **AND**
- Patient must have an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour that is no more than 4 weeks old at the time of application; OR
- Patient must have a C-reactive protein (CRP) level greater than 10 mg per L that is no more than 4 weeks old at the time of application; OR
- Patient must have a clinical reason as to why demonstration of an elevated ESR or CRP cannot be met and the application must state the reason, **AND**
- Patient must not receive more than 18 to 20 weeks of treatment, depending on the dosage regimen, under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR

- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis. The authority application must be made in writing and must include:
 - (a) a completed authority prescription form; and
 - (b) a completed Ankylosing Spondylitis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) a copy of the radiological report confirming Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis; and
 - (ii) a completed BASDAI Assessment Form.

An assessment of a patient's response to an initial course of treatment must be conducted following a minimum of 12 weeks of therapy. An application for the continuing treatment must be accompanied with the assessment of response and submitted to the Department of Human Services no later than 4 weeks from the date of completion of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

certolizumab pegol 200 mg/mL injection, 2 x 1 mL syringes

10904X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	..	..	*3017.55	41.30	Cimzia [UC]

certolizumab pegol 200 mg/mL injection, 2 x 1 mL pen devices

11319R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	..	..	*3017.55	41.30	Cimzia [UC]

■ CERTOLIZUMAB PEGOL

Note TREATMENT OF ADULT PATIENTS WITH NON-RADIOGRAPHIC AXIAL SPONDYLOARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of secukinumab, certolizumab pegol and golimumab for adult patients with non-radiographic axial spondyloarthritis.

Where the term 'biological medicine' appears in notes and restrictions, it refers to secukinumab, certolizumab pegol and golimumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 3 biological medicines at any one time.

Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy.

A serious adverse reaction requiring permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once.

Once a patient has either failed or ceased to respond to treatment 3 times (once with any biological medicine) within the same treatment cycle, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last prescription for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle.

A patient who has failed 3 trials or fewer of biological medicine therapy in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment with secukinumab, certolizumab pegol and golimumab
(1) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised biological medicine treatment in this treatment cycle and wishes to commence such therapy (Initial 1 - New patient)
 - (ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or
 - (iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years); or
 - (iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).
- With the exception of grandfathered patients, a patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy.

(2) Continuing treatment.

For continuing courses of PBS-subsidised biological medicine it is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that biological medicine provided they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment in courses of up to 24 weeks provided they continue to sustain an adequate response.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be conducted no later than 4 weeks from the cessation of that treatment course. Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction requiring permanent treatment withdrawal.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. the C-reactive protein (CRP) level and the BASDAI score), or the prior NSAID therapy and exercise program requirements.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction requiring permanent treatment withdrawal.

(4) Baseline measurements to determine response.

A response to treatment is based on the baseline BASDAI score and CRP level documented in the patient's medical records. For a new patient, the BASDAI score used to determine baseline disease severity must be measured while the patient is receiving NSAID therapy and completing their exercise program.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications.

Prescribers may provide new baseline measurements any time that an 'Initial treatment' authority application is submitted and the eligibility for continuing treatment must be assessed according to these revised baseline measurements.

(5) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must qualify under the Initial 3 treatment restriction. The same clinical criteria and indices of disease severity (i.e. the C-reactive protein (CRP) levels and the BASDAI) as for the Initial 1 (New patient) restriction will need to be met, but a re-trial of NSAID therapy and exercise therapy is not required.

Note No increase in the maximum quantity or number of units may be authorised.

Authority required

Non-radiographic axial spondyloarthritis

Treatment Phase: Initial treatment - Initial 1 (New patient)

Clinical criteria:

- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have had chronic lower back pain and stiffness for 3 or more months that is relieved by exercise but not rest, **AND**
- Patient must have failed to achieve an adequate response following treatment with at least 2 non-steroidal anti-inflammatory drugs (NSAIDs), whilst completing an appropriate exercise program, for a total period of 3 months, **AND**
- Patient must have one or more of the following: (a) enthesitis (heel); (b) uveitis; (c) dactylitis; (d) psoriasis; (e) inflammatory bowel disease; or (f) positive for Human Leukocyte Antigen B27 (HLA-B27), **AND**
- The condition must not be radiographically evidenced on plain x-ray of Grade II bilateral sacroiliitis or Grade III or IV unilateral sacroiliitis, **AND**
- The condition must be non-radiographic axial spondyloarthritis, as defined by Assessment of Spondyloarthritis International Society (ASAS) criteria, **AND**
- The condition must be sacroiliitis with active inflammation and/or oedema on non-contrast Magnetic Resonance Imaging (MRI), **AND**
- The condition must have presence of Bone Marrow Oedema (BMO) depicted as a hyperintense signal on a Short Tau Inversion Recovery (STIR) image (or equivalent), **AND**
- The condition must have BMO depicted as a hypointense signal on a T1 weighted image (without gadolinium), **AND**
- Patient must not receive more than 18 to 20 weeks of treatment, depending on the dosage regimen, under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of non-radiographic axial spondyloarthritis.

The application must include details of the NSAIDs trialled, their doses and duration of treatment.

If the NSAID dose is less than the maximum recommended dose in the relevant TGA-approved Product Information, the application must include the reason a higher dose cannot be used.

If treatment with NSAIDs is contraindicated according to the relevant TGA-approved Product Information, the application must provide details of the contraindication.

If intolerance to NSAID treatment develops during the relevant period of use which is of a severity to necessitate permanent treatment withdrawal, the application must provide details of the nature and severity of this intolerance.

The following criteria indicate failure to achieve an adequate response to NSAIDs and must be demonstrated at the time of the initial application:

- (a) a Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) score of at least 4 on a 0-10 scale; and
- (b) C-reactive protein (CRP) level greater than 10 mg per L.

The baseline BASDAI score and CRP level must be determined at the completion of the 3-month NSAID and exercise trial, but prior to ceasing NSAID treatment. All measures must be no more than 4 weeks old at the time of initial application.

If the requirement to demonstrate an elevated CRP level could not be met, the reason must be stated in the application. Treatment with prednisolone dosed at 7.5 mg or higher daily (or equivalent) or a parenteral steroid within the past month (intramuscular or intravenous methylprednisolone or equivalent) is an acceptable reason.

The assessment of the patient's response to the initial course of treatment must be conducted following a minimum of 12 weeks of treatment and no later than 4 weeks from the cessation of that treatment course. If the response assessment is not conducted within these timeframes, the patient will be deemed to have failed this course of treatment in this treatment cycle.

The authority application must include:

- (a) a completed authority prescription; and
- (b) a completed PBS authority application form relevant to the indication and treatment phase (the latest version located at the website mentioned in the administrative note).

The baseline BASDAI score and CRP level must also be documented in the patient's medical records.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Non-radiographic axial spondyloarthritis

Treatment Phase: Initial treatment - Initial 2 (Change or re-commencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have failed, or ceased to respond to, PBS-subsidised treatment with biological medicines more than three times for this PBS-indication during the current treatment cycle, **AND**
- Patient must not have failed PBS-subsidised therapy with this biological medicine for this PBS indication more than once in the current treatment cycle, **AND**
- Patient must not receive more than 18 to 20 weeks of treatment, depending on the dosage regimen, under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
 - Must be treated by a clinical immunologist with expertise in the management of non-radiographic axial spondyloarthritis.
- An application for Initial 2 treatment must indicate whether the patient has demonstrated an adequate response (an absence of treatment failure), failed or experienced an intolerance to the most recent supply of biological medicine treatment.

A new baseline Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) score and C-reactive protein (CRP) level may be provided at the time of this application.

An adequate response to therapy with this biological medicine is defined as a reduction from baseline in the Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) score by 2 or more units (on a scale of 0-10) and 1 of the following:

- (a) a CRP measurement no greater than 10 mg per L; or
- (b) a CRP measurement reduced by at least 20% from baseline.

The assessment of the patient's response to the most recent supply of biological medicine must be conducted following a minimum of 12 weeks of treatment.

BASDAI scores and CRP levels must be documented in the patient's medical records.

The assessment of the patient's response to the initial course of treatment must be conducted following a minimum of 12 weeks of treatment and no later than 4 weeks from the cessation of that treatment course. If the response assessment is not conducted within these timeframes, the patient will be deemed to have failed this course of treatment in this treatment cycle.

The following must be provided at the time of application and documented in the patient's medical records:

- (a) the BASDAI score; and
- (b) the C-reactive protein (CRP) level.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Non-radiographic axial spondyloarthritis

Treatment Phase: Initial treatment - Initial 3 (Recommencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have had chronic lower back pain and stiffness for 3 or more months that is relieved by exercise but not rest, **AND**
- Patient must have had a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- Patient must have one or more of the following: (a) enthesitis (heel); (b) uveitis; (c) dactylitis; (d) psoriasis; (e) inflammatory bowel disease; or (f) positive for Human Leukocyte Antigen B27 (HLA-B27), **AND**

- The condition must not be radiographically evidenced on plain x-ray of Grade II bilateral sacroiliitis or Grade III or IV unilateral sacroiliitis, **AND**
- The condition must be non-radiographic axial spondyloarthritis, as defined by Assessment of Spondyloarthritis International Society (ASAS) criteria, **AND**
- The condition must be sacroiliitis with active inflammation and/or oedema on non-contrast Magnetic Resonance Imaging (MRI), **AND**
- The condition must have presence of Bone Marrow Oedema (BMO) depicted as a hyperintense signal on a Short Tau Inversion Recovery (STIR) image (or equivalent), **AND**
- The condition must have BMO depicted as a hypointense signal on a T1 weighted image (without gadolinium), **AND**
- Patient must not receive more than 18 to 20 weeks of treatment, depending on the dosage regimen, under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of non-radiographic axial spondyloarthritis.

The following must be provided at the time of application and documented in the patient's medical records:

- (a) a Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) score of at least 4 on a 0-10 scale; and
- (b) C-reactive protein (CRP) level greater than 10 mg per L.

The BASDAI score and CRP level must be no more than 4 weeks old at the time of this application.

If the requirement to demonstrate an elevated CRP level could not be met, the reason must be stated in the application.

Treatment with prednisolone dosed at 7.5 mg or higher daily (or equivalent) or a parenteral steroid within the past month (intramuscular or intravenous methylprednisolone or equivalent) is an acceptable reason.

The assessment of the patient's response to the initial course of treatment must be conducted following a minimum of 12 weeks of treatment and no later than 4 weeks from the cessation of that treatment course. If the response assessment is not conducted within these timeframes, the patient will be deemed to have failed this course of treatment in this treatment cycle.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

certolizumab pegol 200 mg/mL injection, 2 x 1 mL syringes

12063X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	..	..	*3017.55	41.30	Cimzia [UC]

certolizumab pegol 200 mg/mL injection, 2 x 1 mL pen devices

12027B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	..	..	*3017.55	41.30	Cimzia [UC]

■ CERTOLIZUMAB PEGOL**Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE PSORIATIC ARTHRITIS**

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab for adult patients with severe active psoriatic arthritis. Therefore, where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient receiving PBS-subsidised treatment for psoriatic arthritis is able to commence a treatment cycle where they may trial biological medicines without having to experience a disease flare when swapping to the alternate biological medicine.

Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

A patient who received PBS-subsidised adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab or ustekinumab treatment prior to 1 May 2019 is considered to start their first cycle as of 1 May 2019.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence another cycle [further details are under '(5) Recommencement of treatment after a 5-year break in PBS-subsidised therapy' below].

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe active psoriatic arthritis.

(1) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy (Initial 1 - New patient); or
- (ii) a patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an alternate medicine (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or
- (iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).
- (iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years) or An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, golimumab secukinumab and tofacitinib, 18 to 20 weeks of therapy for certolizumab pegol (depending upon the dosing regimen), 20 weeks of therapy for ixekizumab, 22 weeks of therapy for infliximab, and 28 weeks of therapy for ustekinumab. It is recommended that a patient be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of PBS-subsidised initial treatment following a minimum of 12 weeks of therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

Grandfather patients (ixekizumab only).

A patient who commenced treatment with ixekizumab for severe psoriatic arthritis prior to 1 March 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

Grandfather patients (tofacitinib only).

A patient who commenced treatment with Tofacitinib for severe psoriatic arthritis prior to 1 May 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment under the Initial 1 or Initial 2 treatment restrictions.

For the second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to re-qualify with respect to either the indices of disease severity (i.e. erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP) level, and active joint count) or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify under the Initial 3 treatment restriction with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application.

However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

Within a treatment cycle a patient may alternate between therapy with any biological medicine of their choice (1 at a time) providing:

- (i) they have not received PBS-subsidised treatment with that particular biological medicine previously; or
- (ii) they have demonstrated an adequate response to that particular biological medicine if they have previously trialled it on the PBS; and
- (iii) they have not previously failed to respond to treatment 3 times in this treatment cycle with that biological medicine.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(4) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the indices of disease severity submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and these revised baseline measurements will be used to assess response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to determine response for all subsequent continuing treatments. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. 20 or more active joints), response will be determined according to a reduction in the total number of active joints.

(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment restriction according to the criteria of the relevant restriction and index of disease severity. The application must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Clinical criteria:

- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response to methotrexate at a dose of at least 20 mg weekly for a minimum period of 3 months, **AND**
- Patient must have failed to achieve an adequate response to sulfasalazine at a dose of at least 2 g per day for a minimum period of 3 months; OR
- Patient must have failed to achieve an adequate response to leflunomide at a dose of up to 20 mg daily for a minimum period of 3 months, **AND**
- Patient must not receive more than 18 to 20 weeks of treatment, depending on the dosage regimen, under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Where treatment with methotrexate, sulfasalazine or leflunomide is contraindicated according to the relevant TGA-approved Product Information, details must be provided at the time of application.

Where intolerance to treatment with methotrexate, sulfasalazine or leflunomide developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

The following initiation criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; and

either

(a) an active joint count of at least 20 active (swollen and tender) joints; or

(b) at least 4 active joints from the following list of major joints:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Severe Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

An assessment of a patient's response to an initial course of treatment must be conducted following a minimum of 12 weeks of therapy. An application for the continuing treatment must be accompanied with the assessment of response and

submitted to the Department of Human Services no later than 4 weeks from the date of completion of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 3 months treatment with methotrexate and 3 months treatment with sulfasalazine or leflunomide can be found on the Department of Human Services website (www.humanservices.gov.au)

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial treatment - Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- Patient must not receive more than 18 to 20 weeks of treatment, depending on the dosage regimen, under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

An adequate response to treatment is defined as:

an erythrocyte sedimentation rate (ESR) no greater than 25 mm per hour or a C-reactive protein (CRP) level no greater than 15 mg per L or either marker reduced by at least 20% from baseline; and

either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following major active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Severe Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to change or recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial treatment - Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must have an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; OR
- The condition must have a C-reactive protein (CRP) level greater than 15 mg per L, **AND**
- The condition must have either (a) a total active joint count of at least 20 active (swollen and tender) joints; or (b) at least 4 active major joints, **AND**
- Patient must not receive more than 18 to 20 weeks of treatment, depending on the dosage regimen, under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Major joints are defined as (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

All measures of joint count and ESR and/or CRP must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Severe Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

certolizumab pegol 200 mg/mL injection, 2 x 1 mL syringes

10909E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	..	..	*3017.55	41.30	Cimzia [UC]

certolizumab pegol 200 mg/mL injection, 2 x 1 mL pen devices

11323Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	..	..	*3017.55	41.30	Cimzia [UC]

■ ETANERCEPT**Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE PSORIATIC ARTHRITIS**

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab for adult patients with severe active psoriatic arthritis. Therefore, where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient receiving PBS-subsidised treatment for psoriatic arthritis is able to commence a treatment cycle where they may trial biological medicines without having to experience a disease flare when swapping to the alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

A patient who received PBS-subsidised adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab or ustekinumab treatment prior to 1 May 2019 is considered to start their first cycle as of 1 May 2019.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they

are eligible to commence another cycle [further details are under '(5) Recommencement of treatment after a 5-year break in PBS-subsidised therapy' below].

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe active psoriatic arthritis.

(1) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an alternate medicine (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years) or An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, golimumab secukinumab and tofacitinib, 18 to 20 weeks of therapy for certolizumab pegol (depending upon the dosing regimen), 20 weeks of therapy for ixekizumab, 22 weeks of therapy for infliximab, and 28 weeks of therapy for ustekinumab. It is recommended that a patient be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of PBS-subsidised initial treatment following a minimum of 12 weeks of therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

Grandfather patients (ixekizumab only).

A patient who commenced treatment with ixekizumab for severe psoriatic arthritis prior to 1 March 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

Grandfather patients (tofacitinib only).

A patient who commenced treatment with Tofacitinib for severe psoriatic arthritis prior to 1 May 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment under the Initial 1 or Initial 2 treatment restrictions.

For the second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to re-qualify with respect to either the indices of disease severity (i.e. erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP) level, and active joint count) or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify under the Initial 3 treatment restriction with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting

in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application.

However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

Within a treatment cycle a patient may alternate between therapy with any biological medicine of their choice (1 at a time) providing:

- (i) they have not received PBS-subsidised treatment with that particular biological medicine previously; or
- (ii) they have demonstrated an adequate response to that particular biological medicine if they have previously trialled it on the PBS; and
- (iii) they have not previously failed to respond to treatment 3 times in this treatment cycle with that biological medicine.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(4) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the indices of disease severity submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and these revised baseline measurements will be used to assess response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to determine response for all subsequent continuing treatments. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. 20 or more active joints), response will be determined according to a reduction in the total number of active joints.

(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment restriction according to the criteria of the relevant restriction and index of disease severity. The application must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

9156

Severe psoriatic arthritis

Treatment Phase: Subsequent continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment per subsequent continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

An adequate response to treatment is defined as:

an erythrocyte sedimentation rate (ESR) no greater than 25 mm per hour or a C-reactive protein (CRP) level no greater than 15 mg per L or either marker reduced by at least 20% from baseline; and

either of the following:

- (a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or
- (b) a reduction in the number of the following major active joints, from at least 4, by at least 50%:
 - (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to determine response for all subsequent continuing treatments.

The measurement of response to the prior course of therapy must have been conducted following a minimum of 12 weeks of therapy with this drug and must be documented in the patient's medical records.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

etanercept 50 mg/mL injection, 4 x 1 mL pen devices

11202N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1050.08	41.30	Brenzys [RF]

etanercept 50 mg/mL injection, 4 x 1 mL syringes

11216H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1050.08	41.30	Brenzys [RF]

■ ETANERCEPT
Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for adults with severe active rheumatoid arthritis. Where the term biological medicine appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alfa antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab), the T-cell co-stimulation modulator (abatacept) and the Janus kinase (JAK) inhibitors (baricitinib, tofacitinib, upadacitinib).

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised biological medicine therapy may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once, and
- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised biological medicines for the treatment of rheumatoid arthritis.

A serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Stevens Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered a treatment failure.

A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction.

A patient who has failed fewer than 5 biological medicines and who has a break in treatment of less than 24 months may commence a further course of treatment with a biological medicine under Initial 2 treatment restriction. A patient who has failed fewer than 5 biological medicines and who has had a break in therapy of longer than 24 months may commence a further course of treatment with a biological medicine under the Initial 3 treatment restriction. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine treatment is stopped to the date of the new application for treatment with a biological medicine.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 April 2019.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised biological medicine treatment and wishes to commence such therapy, excluding rituximab (Initial 1 - new patient); or
 - (ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months) [further details are under 'Swapping therapy' below]; or
 - (iii) a patient wishes to re-commence treatment with a specific biological medicine following a break of less than 24 months in PBS-subsidised therapy with that agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months).
 - (iv) a patient wishes to re-commence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 24 months (Initial 3 - re-commencement of treatment after a break in biological medicine of more than 24 months)
- Initial applications for a new patient (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6-month intensive DMARD trial, but prior to ceasing DMARD therapy.

Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, baricitinib, etanercept, golimumab, tocilizumab, tofacitinib and upadacitinib, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

Rituximab patients should be assessed following a minimum of 12 weeks after the first infusion, and the assessment should be submitted to Services Australia within 4 weeks to ensure continuity of treatment for those who meet the continuing restriction for PBS subsidised treatment with this drug for this condition.

Where a response assessment is not provided the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment. For second and subsequent courses of PBS-subsidised biological medicine (excluding rituximab) treatment, it is recommended that a patient be reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to Services Australia within sufficient time to allow processing. Abatacept patients:

A patient is eligible to receive one I.V. loading dose when commencing treatment with the subcutaneous formulation. Two prescriptions are required, the first prescription for the I.V. loading dose for sufficient vials for one dose based on the patient's weight with no repeats. The second prescription for the subcutaneous formulation, with a maximum quantity of 4 and up to 3 repeats, must be submitted with the initial application.

Rituximab patients:

A further application may be submitted to Services Australia within sufficient time to allow processing. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the

month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply. Assessment of the patient's response to treatment must be conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the cessation of the most recent course of biological medicine therapy.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction. Where a response assessment is not conducted within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

(2) Swapping therapy

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non- biological medicine therapy requirements except if the patient has had a break in therapy of more than 24 months who would need to requalify with respect to the indices of disease severity. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each biological medicine trialled.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent, unless they have experienced a serious adverse reaction of a severity necessitating permanent treatment withdrawal.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug.

Abatacept:

A patient swapping from I.V. abatacept to subcutaneous abatacept will not be eligible for an I.V. loading dose when commencing treatment with the subcutaneous formulation.

Rituximab:

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF-alfa antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they be assessed for response to every course of treatment, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological medicine during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological medicine therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate biological medicine may do so without having to have any treatment-free period.

(3) Baseline measurements to determine response.

Determination of whether a response to treatment has been demonstrated must be based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and the eligibility for continuing treatment must be assessed according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response must be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be determined on the total number of major joints.

Applications under the Initial 1 treatment restriction for a new patient must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. The results must be no more than 4 weeks old at the time of application.

Applications under the Initial 3 treatment restriction for re-commencement of treatment after a break in biological medicine of more than 24 months must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

7276

Severe active rheumatoid arthritis

Treatment Phase: Subsequent continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Population criteria:

- Patient must be aged 18 years or older.

Clinical criteria:

- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must have received this drug as their most recent course of PBS-subsidised biological disease modifying anti-rheumatic drug (bDMARD) treatment for this condition, **AND**
- Patient must not receive more than 24 weeks of treatment per subsequent continuing treatment course authorised under this restriction.

For the purposes of this restriction bDMARD means abatacept, adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, rituximab, tocilizumab or tofacitinib.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The measurement of response to the prior course of therapy must be documented in the patient's medical notes.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

etanercept 50 mg/mL injection, 4 x 1 mL pen devices

11218K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1050.08	41.30	Brenzys [RF]

etanercept 50 mg/mL injection, 4 x 1 mL syringes

11211C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1050.08	41.30	Brenzys [RF]

■ ETANERCEPT

Note TREATMENT OF ADULT PATIENTS WITH ANKYLOSING SPONDYLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab for adult patients with ankylosing spondylitis.

Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 7 biological medicines at any 1 time.

Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy.

A patient who has been receiving PBS-subsidised adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab treatment prior to 1 December 2020 is considered to start their first cycle as of 1 December 2020.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last prescription for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised biological medicine treatment with adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised biological medicine treatment in this treatment cycle and wishes to commence such therapy (Initial 1 - New patient)

(ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years); or

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

A patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy.

(b) Continuing treatment.

For the first continuing treatment course of PBS-subsidised biological medicine, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1, Initial 2 or Initial 3 treatment restrictions. For

second and subsequent continuing courses of PBS-subsidised biological medicine it is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply. Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be conducted no later than 4 weeks from the cessation of that treatment course. Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI), or the prior NSAID therapy and exercise program requirements.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(3) Baseline measurements to determine response.

Services Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the BASDAI, ESR and/or CRP submitted with the first authority application for a biological medicine. For a new patient, the BASDAI used to determine the baseline must be measured while the patient is receiving NSAID therapy and completing their exercise program.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response.

Prescribers may provide new baseline measurements any time an 'Initial treatment' authority application is submitted within a treatment cycle and the eligibility for continuing treatment must be assessed according to these revised baseline measurements.

(4) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must qualify under the Initial 3 treatment restriction. The same clinical criteria and indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI) as for the Initial 1 (New patient) restriction will need to be met, but a re-trial of NSAID therapy and exercise therapy is not required.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

9481

Ankylosing spondylitis

Treatment Phase: Subsequent continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response is defined as an improvement from baseline of at least 2 of the BASDAI and 1 of the following:

- (a) an ESR measurement no greater than 25 mm per hour; or
- (b) a CRP measurement no greater than 10 mg per L; or
- (c) an ESR or CRP measurement reduced by at least 20% from baseline.

Where only 1 acute phase reactant measurement is supplied in the first application for PBS-subsidised treatment, that same marker must be used to determine response for all subsequent continuing treatments.

The measurement of response to the prior course of therapy must be documented in the patient's medical notes.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

etanercept 50 mg/mL injection, 4 x 1 mL pen devices

11215G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1050.08	41.30	Brenzys [RF]

etanercept 50 mg/mL injection, 4 x 1 mL syringes

11217J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1050.08	41.30	Brenzys [RF]

■ ETANERCEPT
Note TREATMENT OF ADULT PATIENTS WITH A HISTORY OF JUVENILE IDIOPATHIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, etanercept and tocilizumab for a patient over 18 years who has a history of juvenile idiopathic arthritis with onset prior to the age of 18 years. Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, etanercept and tocilizumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 3 biological medicines at any one time.

From 1 April 2014, a patient receiving PBS-subsidised biological medicine therapy is considered to be in a treatment cycle where they may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements, within a single treatment cycle, a patient may:

- (i) continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy; and
- (ii) fail to respond or to sustain a response to each PBS-subsidised biological medicine once only. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment.

Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single treatment cycle and they must have, at a minimum, a 5 year break in PBS-subsidised biological medicine therapy before they are eligible to receive further PBS-subsidised biological medicine therapy.

The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine was approved to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who received PBS-subsidised biological medicine treatment immediately prior to 1 April 2014 is considered to be in their first cycle as of 1 April 2014. A patient who has had a break in biological medicine treatment of at least 24 months immediately prior to making a new application, on or after 1 April 2014, will commence a new treatment cycle under the Initial 3 treatment restriction.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of less than 24 months may commence a further course of treatment within the same treatment cycle under the Initial 2 treatment restriction.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of more than 24 months must commence a new treatment cycle under the Initial 3 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine therapy after 1 April 2014.

(1) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy (Initial 1 - New patient); or
- (ii) a patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an alternate medicine (Initial 2 - Change or recommencement of treatment after a break in biological medicine of less than 24 months) [further details are under 'Swapping therapy' below]; or
- (iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 24 months with the same medicine (Initial 2 - Change or recommencement of treatment after a break in biological medicine therapy of less than 24 months); or
- (iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 24 months (Initial 3 - recommencement of treatment after a break in biological medicine of more than 24 months). Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy.

A patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not conducted within these timeframes, the patient will be deemed to have failed this course of treatment.

For second and subsequent cycles of PBS-subsidised biological medicine, it is recommended that a patient is reviewed in the 4 weeks prior to completing their current course of treatment and that an application is posted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services where applicable. Where a response assessment is not submitted, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious

adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (joint count and ESR/CRP) or the prior non- biological medicine therapy requirements, except if the patient has had a break in therapy of more than 24 months who would then need to requalify under the Initial 3 restrictions with respect to the indices of disease severity.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, an application for a patient who wishes to swap to an alternate biological medicine should be accompanied by the approved authority prescription or remaining repeats for the biological medicine the patient is ceasing.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count submitted with the first authority application for a biological medicine. However, prescribers may provide a new baseline measurement any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to the revised baseline measurement.

(4) Recommencement of treatment after a 24 months break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological medicine therapy of at least 24 months, must qualify under the Initial 3 restriction and meet the relevant criteria and index of disease severity.

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment per continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

- (a) an active joint count of fewer than 10 active (swollen and tender) joints; or
- (b) a reduction in the active (swollen and tender) joint count by at least 50% from baseline; or
- (c) a reduction in the number of the following active joints, from at least 4, by at least 50%:
 - (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - (ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

- (1) completed authority prescription form(s); and
- (2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either Initial 1, Initial 2, or Initial 3 treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

If a patient fails to respond to PBS-subsidised biological medicine treatment 3 times (once with each agent) they will not be eligible to receive further PBS-subsidised biological medicine therapy in this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Continuing Treatment - balance of supply

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restriction.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

etanercept 50 mg/mL injection, 4 x 1 mL pen devices

3450N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1050.08	41.30	Enbrel [PF]

etanercept 25 mg injection [4 vials] (&) inert substance diluent [4 x 1 mL syringes], 1 pack

3448L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*1050.10	41.30	Enbrel [PF]

etanercept 50 mg/mL injection, 4 x 1 mL syringes

3449M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1050.08	41.30	Enbrel [PF]

■ ETANERCEPT

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction. There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years); or

(iii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or

(iv) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient be reviewed in the 4 weeks prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

Grandfather patients (risankizumab only).

A patient who commenced treatment with risankizumab for chronic plaque psoriasis prior to 1 December 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the Initial treatment Grandfather treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment. This assessment must be conducted within 4 weeks of the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the 4 weeks prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the 4 weeks prior to completing their current course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure patients receive the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

Services Australia will determine whether a response to treatment has been demonstrated based on the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised

treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 the PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Authority applications for increased quantities/ repeats (where relevant) may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required (STREAMLINED)

8887

Severe chronic plaque psoriasis

Treatment Phase: Subsequent continuing treatment, whole body

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per subsequent continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

The measurement of response to the prior course of therapy must be documented in the patient's medical notes.

Determination of response must be based on the PASI assessment of response to the most recent course of treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Authority required (STREAMLINED)

8955

Severe chronic plaque psoriasis

Treatment Phase: Subsequent continuing treatment, face, hand, foot

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to their most recent course of treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per subsequent continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or
- (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

The measurement of response to the prior course of therapy must be documented in the patient's medical notes.

Determination of response must be based on the PASI assessment of response to the most recent course of treatment with this drug.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

etanercept 50 mg/mL injection, 4 x 1 mL pen devices

11221N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1050.08	41.30	Brenzys [RF]

etanercept 50 mg/mL injection, 4 x 1 mL syringes

11225T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1050.08	41.30	Brenzys [RF]

■ ETANERCEPT

Note TREATMENT OF ADULT PATIENTS WITH ANKYLOSING SPONDYLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab for adult patients with ankylosing spondylitis.

Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 7 biological medicines at any 1 time.

Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy.

A patient who has been receiving PBS-subsidised adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab treatment prior to 1 December 2020 is considered to start their first cycle as of 1 December 2020.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last prescription for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised biological medicine treatment with adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised biological medicine treatment in this treatment cycle and wishes to commence such therapy (Initial 1 - New patient)

(ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years); or

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

A patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy.

(b) Continuing treatment.

For the first continuing treatment course of PBS-subsidised biological medicine, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1, Initial 2 or Initial 3 treatment restrictions. For second and subsequent continuing courses of PBS-subsidised biological medicine it is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply. Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be conducted no later than 4 weeks from the cessation of that treatment course. Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI), or the prior NSAID therapy and exercise program requirements.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or

continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(3) Baseline measurements to determine response.

Services Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the BASDAI, ESR and/or CRP submitted with the first authority application for a biological medicine. For a new patient, the BASDAI used to determine the baseline must be measured while the patient is receiving NSAID therapy and completing their exercise program.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response.

Prescribers may provide new baseline measurements any time an 'Initial treatment' authority application is submitted within a treatment cycle and the eligibility for continuing treatment must be assessed according to these revised baseline measurements.

(4) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must qualify under the Initial 3 treatment restriction. The same clinical criteria and indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI) as for the Initial 1 (New patient) restriction will need to be met, but a re-trial of NSAID therapy and exercise therapy is not required.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Ankylosing spondylitis

Treatment Phase: Subsequent continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Ankylosing Spondylitis PBS Authority Application - Supporting Information Form.

An adequate response is defined as an improvement from baseline of at least 2 of the BASDAI and 1 of the following:

- (a) an ESR measurement no greater than 25 mm per hour; or
- (b) a CRP measurement no greater than 10 mg per L; or
- (c) an ESR or CRP measurement reduced by at least 20% from baseline.

Where only 1 acute phase reactant measurement is supplied in the first application for PBS-subsidised treatment, that same marker must be measured and supplied in all subsequent continuing treatment applications.

All measurements provided must be no more than 1 month old at the time of application.

Each application for subsequent continuing treatment with this drug must include an assessment of the patient's response to the prior course of therapy. If the response assessment is not provided at the time of application the patient will be deemed to have failed this course of treatment, unless the patient has experienced serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Ankylosing spondylitis

Treatment Phase: Continuing treatment - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the first continuing treatment restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the subsequent continuing Authority Required (in writing) treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

etanercept 50 mg/mL injection, 4 x 1 mL pen devices

11201M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	1050.08	41.30	^a Brenzys [RF]	^a Enbrel [PF]

etanercept 25 mg injection [4 vials] (&) inert substance diluent [4 x 1 mL syringes], 1 pack

11204Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	5	..	*1050.10	41.30	Enbrel [PF]	

etanercept 50 mg/mL injection, 4 x 1 mL syringes

11196G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	1050.08	41.30	^a Brenzys [RF]	^a Enbrel [PF]

■ ETANERCEPT
Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for adults with severe active rheumatoid arthritis. Where the term biological medicine appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alpha antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab), the T-cell co-stimulation modulator (abatacept) and the Janus kinase (JAK) inhibitors (baricitinib, tofacitinib, upadacitinib).

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alpha antagonist.

A patient receiving PBS-subsidised biological medicine therapy may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once, and
- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised biological medicines for the treatment of rheumatoid arthritis.

A serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Stevens Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered a treatment failure.

A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction.

A patient who has failed fewer than 5 biological medicines and who has a break in treatment of less than 24 months may commence a further course of treatment with a biological medicine under Initial 2 treatment restriction. A patient who has failed fewer than 5 biological medicines and who has had a break in therapy of longer than 24 months may commence a further course of treatment with a biological medicine under the Initial 3 treatment restriction. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine treatment is stopped to the date of the new application for treatment with a biological medicine.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 April 2019.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised biological medicine treatment and wishes to commence such therapy, excluding rituximab (Initial 1 - new patient); or
 - (ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months) [further details are under 'Swapping therapy' below]; or
 - (iii) a patient wishes to re-commence treatment with a specific biological medicine following a break of less than 24 months in PBS-subsidised therapy with that agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months).
 - (iv) a patient wishes to re-commence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 24 months (Initial 3 - re-commencement of treatment after a break in biological medicine of more than 24 months)
- Initial applications for a new patient (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6-month intensive DMARD trial, but prior to ceasing DMARD therapy.
- Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, baricitinib, etanercept, golimumab, tocilizumab, tofacitinib and upadacitinib, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

Rituximab patients should be assessed following a minimum of 12 weeks after the first infusion, and the assessment should be submitted to Services Australia within 4 weeks to ensure continuity of treatment for those who meet the continuing restriction for PBS subsidised treatment with this drug for this condition.

Where a response assessment is not provided the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment. For second and subsequent courses of PBS-subsidised biological medicine (excluding rituximab) treatment, it is recommended that a patient be reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to Services Australia within sufficient time to allow processing.

Abatacept patients:

A patient is eligible to receive one I.V. loading dose when commencing treatment with the subcutaneous formulation. Two prescriptions are required, the first prescription for the I.V. loading dose for sufficient vials for one dose based on the patient's weight with no repeats. The second prescription for the subcutaneous formulation, with a maximum quantity of 4 and up to 3 repeats, must be submitted with the initial application.

Rituximab patients:

A further application may be submitted to Services Australia within sufficient time to allow processing. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

Assessment of the patient's response to treatment must be conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the cessation of the most recent course of biological medicine therapy.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction. Where a response assessment is not conducted within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

(2) Swapping therapy

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non- biological medicine therapy requirements except if the patient has had a break in therapy of more than 24 months who would need to requalify with respect to the indices of disease severity. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each biological medicine trialled.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent, unless they have experienced a serious adverse reaction of a severity necessitating permanent treatment withdrawal.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug.

Abatacept:

A patient swapping from I.V. abatacept to subcutaneous abatacept will not be eligible for an I.V. loading dose when commencing treatment with the subcutaneous formulation.

Rituximab:

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF-alfa antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they be assessed for response to every course of treatment, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological medicine during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological medicine therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate biological medicine may do so without having to have any treatment-free period.

(3) Baseline measurements to determine response.

Determination of whether a response to treatment has been demonstrated must be based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and the eligibility for continuing treatment must be assessed according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response must be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be determined on the total number of major joints.

Applications under the Initial 1 treatment restriction for a new patient must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. The results must be no more than 4 weeks old at the time of application.

Applications under the Initial 3 treatment restriction for re-commencement of treatment after a break in biological medicine of

more than 24 months must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Subsequent continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment per subsequent continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient has either failed or ceased to respond to a PBS-subsidised biological medicine for this condition 5 times, they will not be eligible to receive further PBS-subsidised treatment with a biological medicine for this condition.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Continuing treatment - balance of supply

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the first continuing treatment restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the subsequent continuing Authority Required (in writing) treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restrictions.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

etanercept 50 mg/mL injection, 4 x 1 mL pen devices

11220M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	1050.08	41.30	^a Brenzys [RF]	^a Enbrel [PF]

etanercept 25 mg injection [4 vials] (&) inert substance diluent [4 x 1 mL syringes], 1 pack

11197H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	5	..	*1050.10	41.30	Enbrel [PF]	

etanercept 50 mg/mL injection, 4 x 1 mL syringes

11219L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	1050.08	41.30	^a Brenzys [RF]	^a Enbrel [PF]

■ **ETANERCEPT**

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for adults with severe active rheumatoid arthritis. Where the term biological medicine appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alfa antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab), the T-cell co-stimulation modulator (abatacept) and the Janus kinase (JAK) inhibitors (baricitinib, tofacitinib, upadacitinib).

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised biological medicine therapy may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy,

- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once, and

- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised biological medicines for the treatment of rheumatoid arthritis.

A serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Stevens Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered a treatment failure.

A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction.

A patient who has failed fewer than 5 biological medicines and who has a break in treatment of less than 24 months may commence a further course of treatment with a biological medicine under Initial 2 treatment restriction. A patient who has failed fewer than 5 biological medicines and who has had a break in therapy of longer than 24 months may commence a further course of treatment with a biological medicine under the Initial 3 treatment restriction. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine treatment is stopped to the date of the new application for treatment with a biological medicine.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 April 2019.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised biological medicine treatment and wishes to commence such therapy, excluding rituximab (Initial 1 - new patient); or

(ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to re-commence treatment with a specific biological medicine following a break of less than 24 months in PBS-subsidised therapy with that agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months).

(iv) a patient wishes to re-commence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 24 months (Initial 3 - re-commencement of treatment after a break in biological medicine of more than 24 months) Initial applications for a new patient (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6-month intensive DMARD trial, but prior to ceasing DMARD therapy.

Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, baricitinib, etanercept, golimumab, tocilizumab, tofacitinib and upadacitinib, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

Rituximab patients should be assessed following a minimum of 12 weeks after the first infusion, and the assessment should be submitted to Services Australia within 4 weeks to ensure continuity of treatment for those who meet the continuing restriction for PBS subsidised treatment with this drug for this condition.

Where a response assessment is not provided the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment. For second and subsequent courses of PBS-subsidised biological medicine (excluding rituximab) treatment, it is recommended that a patient be reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to Services Australia within sufficient time to allow processing. Abatacept patients:

A patient is eligible to receive one I.V. loading dose when commencing treatment with the subcutaneous formulation. Two prescriptions are required, the first prescription for the I.V. loading dose for sufficient vials for one dose based on the patient's weight with no repeats. The second prescription for the subcutaneous formulation, with a maximum quantity of 4

and up to 3 repeats, must be submitted with the initial application.

Rituximab patients:

A further application may be submitted to Services Australia within sufficient time to allow processing. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

Assessment of the patient's response to treatment must be conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the cessation of the most recent course of biological medicine therapy.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction. Where a response assessment is not conducted within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

(2) Swapping therapy

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non- biological medicine therapy requirements except if the patient has had a break in therapy of more than 24 months who would need to requalify with respect to the indices of disease severity. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each biological medicine trialled.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent, unless they have experienced a serious adverse reaction of a severity necessitating permanent treatment withdrawal.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug.

Abatacept:

A patient swapping from I.V. abatacept to subcutaneous abatacept will not be eligible for an I.V. loading dose when commencing treatment with the subcutaneous formulation.

Rituximab:

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF-alfa antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they be assessed for response to every course of treatment, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological medicine during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological medicine therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate biological medicine may do so without having to have any treatment-free period.

(3) Baseline measurements to determine response.

Determination of whether a response to treatment has been demonstrated must be based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and the eligibility for continuing treatment must be assessed according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response must be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be determined on the total number of major joints.

Applications under the Initial 1 treatment restriction for a new patient must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. The results must be no more than 4 weeks old at the time of application.

Applications under the Initial 3 treatment restriction for re-commencement of treatment after a break in biological medicine of more than 24 months must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Severe active rheumatoid arthritis

Treatment Phase: First Continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient has either failed or ceased to respond to a PBS-subsidised biological medicine for this condition 5 times, they will not be eligible to receive further PBS-subsidised treatment with a biological medicine for this condition.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Continuing treatment - balance of supply

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the first continuing treatment restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the subsequent continuing Authority Required (in writing) treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restrictions.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

etanercept 50 mg/mL injection, 4 x 1 mL pen devices

9460X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	1050.08	41.30	^a Brenzys [RF]	^a Enbrel [PF]

etanercept 25 mg injection [4 vials] (&) inert substance diluent [4 x 1 mL syringes], 1 pack

8638P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*1050.10	41.30	Enbrel [PF]

etanercept 50 mg/mL injection, 4 x 1 mL syringes

9090K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	1050.08	41.30	^a Brenzys [RF]	^a Enbrel [PF]

■ ETANERCEPT**Note TREATMENT OF ADULT PATIENTS WITH ANKYLOSING SPONDYLITIS**

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab for adult patients with ankylosing spondylitis.

Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 7 biological medicines at any 1 time.

Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy.

A patient who has been receiving PBS-subsidised adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab treatment prior to 1 December 2020 is considered to start their first cycle as of 1 December 2020.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last prescription for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised biological medicine treatment with adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised biological medicine treatment in this treatment cycle and wishes to commence such therapy (Initial 1 - New patient)

(ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years); or

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

A patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy.

(b) Continuing treatment.

For the first continuing treatment course of PBS-subsidised biological medicine, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1, Initial 2 or Initial 3 treatment restrictions. For second and subsequent continuing courses of PBS-subsidised biological medicine it is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be conducted no later than 4 weeks from the cessation of that treatment course. Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI), or the prior NSAID therapy and exercise program requirements.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(3) Baseline measurements to determine response.

Services Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the BASDAI, ESR and/or CRP submitted with the first authority application for a biological medicine. For a new patient, the BASDAI used to determine the baseline must be measured while the patient is receiving NSAID therapy and completing their exercise program.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response.

Prescribers may provide new baseline measurements any time an 'Initial treatment' authority application is submitted within a treatment cycle and the eligibility for continuing treatment must be assessed according to these revised baseline measurements.

(4) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must qualify under the Initial 3 treatment restriction. The same clinical criteria and indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI) as for the Initial 1 (New patient) restriction will need to be met, but a re-trial of NSAID therapy and exercise therapy is not required.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Ankylosing spondylitis

Treatment Phase: First continuing treatment

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Ankylosing Spondylitis PBS Authority Application - Supporting Information Form.

An adequate response is defined as an improvement from baseline of at least 2 of the BASDAI and 1 of the following:

- (a) an ESR measurement no greater than 25 mm per hour; or
- (b) a CRP measurement no greater than 10 mg per L; or
- (c) an ESR or CRP measurement reduced by at least 20% from baseline.

Where only 1 acute phase reactant measurement is supplied in the first application for PBS-subsidised treatment, that same marker must be measured and supplied in all subsequent continuing treatment applications.

All measurements provided must be no more than 1 month old at the time of application.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Ankylosing spondylitis

Treatment Phase: Continuing treatment - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the first continuing treatment restriction to complete 24 weeks treatment; OR
 - Patient must have received insufficient therapy with this drug for this condition under the subsequent continuing Authority Required (in writing) treatment restriction to complete 24 weeks treatment, **AND**
 - The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restrictions.
- Treatment criteria:**
- Must be treated by a rheumatologist; OR
 - Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

etanercept 50 mg/mL injection, 4 x 1 mL pen devices

9456Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	1050.08	41.30	^a Brenzys [RF]	^a Enbrel [PF]

etanercept 25 mg injection [4 vials] (&) inert substance diluent [4 x 1 mL syringes], 1 pack

8779C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	5	..	*1050.10	41.30	Enbrel [PF]	

etanercept 50 mg/mL injection, 4 x 1 mL syringes

9086F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	1050.08	41.30	^a Brenzys [RF]	^a Enbrel [PF]

■ ETANERCEPT

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE PSORIATIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab for adult patients with severe active psoriatic arthritis. Therefore, where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient receiving PBS-subsidised treatment for psoriatic arthritis is able to commence a treatment cycle where they may trial biological medicines without having to experience a disease flare when swapping to the alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

A patient who received PBS-subsidised adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab or ustekinumab treatment prior to 1 May 2019 is considered to start their first cycle as of 1 May 2019.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence another cycle [further details are under '(5) Recommencement of treatment after a 5-year break in PBS-subsidised therapy' below].

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe active psoriatic arthritis.

(1) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an alternate medicine (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years) or

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, golimumab secukinumab and tofacitinib, 18 to 20 weeks of therapy for certolizumab pegol (depending upon the dosing regimen), 20 weeks of therapy for ixekizumab, 22 weeks of therapy for infliximab, and 28 weeks of therapy for

ustekinumab. It is recommended that a patient be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of PBS-subsidised initial treatment following a minimum of 12 weeks of therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

Grandfather patients (ixekizumab only).

A patient who commenced treatment with ixekizumab for severe psoriatic arthritis prior to 1 March 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

Grandfather patients (tofacitinib only).

A patient who commenced treatment with Tofacitinib for severe psoriatic arthritis prior to 1 May 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment under the Initial 1 or Initial 2 treatment restrictions.

For the second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to re-qualify with respect to either the indices of disease severity (i.e. erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP) level, and active joint count) or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify under the Initial 3 treatment restriction with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application.

However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

Within a treatment cycle a patient may alternate between therapy with any biological medicine of their choice (1 at a time) providing:

- (i) they have not received PBS-subsidised treatment with that particular biological medicine previously; or
- (ii) they have demonstrated an adequate response to that particular biological medicine if they have previously trialled it on the PBS; and
- (iii) they have not previously failed to respond to treatment 3 times in this treatment cycle with that biological medicine.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(4) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the indices of disease severity submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and these revised baseline measurements will be used to assess response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to determine response for all subsequent continuing treatments. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. 20 or more active joints), response will be determined according to a reduction in the total number of active joints.

(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological

therapy of at least 5 years, must qualify under Initial 3 treatment restriction according to the criteria of the relevant restriction and index of disease severity. The application must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Severe psoriatic arthritis

Treatment Phase: Subsequent continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment per subsequent continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

An adequate response to treatment is defined as:

an erythrocyte sedimentation rate (ESR) no greater than 25 mm per hour or a C-reactive protein (CRP) level no greater than 15 mg per L or either marker reduced by at least 20% from baseline; and

either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following major active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to determine response for all subsequent continuing treatments.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Severe Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

Where the most recent course of PBS-subsidised treatment with this drug was approved under the first continuing treatment restriction, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Authority required

Severe psoriatic arthritis

Treatment Phase: Continuing treatment - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the first continuing treatment restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the subsequent continuing Authority Required (in writing) treatment restriction to complete 24 weeks treatment, **AND**

- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

etanercept 50 mg/mL injection, 4 x 1 mL pen devices

11198J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	1050.08	41.30	^a Brenzys [RF]	^a Enbrel [PF]

etanercept 25 mg injection [4 vials] (&) inert substance diluent [4 x 1 mL syringes], 1 pack

11207W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*1050.10	41.30	Enbrel [PF]

etanercept 50 mg/mL injection, 4 x 1 mL syringes

11208X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	1050.08	41.30	^a Brenzys [RF]	^a Enbrel [PF]

▪ **ETANERCEPT**

Note TREATMENT OF ADULT PATIENTS WITH A HISTORY OF JUVENILE IDIOPATHIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, etanercept and tocilizumab for a patient over 18 years who has a history of juvenile idiopathic arthritis with onset prior to the age of 18 years. Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, etanercept and tocilizumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 3 biological medicines at any one time.

From 1 April 2014, a patient receiving PBS-subsidised biological medicine therapy is considered to be in a treatment cycle where they may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements, within a single treatment cycle, a patient may:

- continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy; and
- fail to respond or to sustain a response to each PBS-subsidised biological medicine once only. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment.

Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single treatment cycle and they must have, at a minimum, a 5 year break in PBS-subsidised biological medicine therapy before they are eligible to receive further PBS-subsidised biological medicine therapy.

The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine was approved to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who received PBS-subsidised biological medicine treatment immediately prior to 1 April 2014 is considered to be in their first cycle as of 1 April 2014. A patient who has had a break in biological medicine treatment of at least 24 months immediately prior to making a new application, on or after 1 April 2014, will commence a new treatment cycle under the Initial 3 treatment restriction.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of less than 24 months may commence a further course of treatment within the same treatment cycle under the Initial 2 treatment restriction.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of more than 24 months must commence a new treatment cycle under the Initial 3 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine therapy after 1 April 2014.

(1) Initial treatment.

Applications for initial treatment should be made where:

- a patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy (Initial 1 - New patient); or
- a patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an alternate medicine (Initial 2 - Change or recommencement of treatment after a break in biological medicine of less than 24 months) [further details are under 'Swapping therapy' below]; or
- a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 24 months with the same medicine (Initial 2 - Change or recommencement of treatment after a break in biological medicine therapy of less than 24 months); or
- a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 24 months (Initial 3 - recommencement of treatment after a break in biological medicine of more than 24 months).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy.

A patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not conducted within these timeframes, the patient will be deemed to have failed this course of treatment.

For second and subsequent cycles of PBS-subsidised biological medicine, it is recommended that a patient is reviewed in

the 4 weeks prior to completing their current course of treatment and that an application is posted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services where applicable. Where a response assessment is not submitted, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (joint count and ESR/CRP) or the prior non- biological medicine therapy requirements, except if the patient has had a break in therapy of more than 24 months who would then need to requalify under the Initial 3 restrictions with respect to the indices of disease severity.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, an application for a patient who wishes to swap to an alternate biological medicine should be accompanied by the approved authority prescription or remaining repeats for the biological medicine the patient is ceasing.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count submitted with the first authority application for a biological medicine. However, prescribers may provide a new baseline measurement any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to the revised baseline measurement.

(4) Recommencement of treatment after a 24 months break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological medicine therapy of at least 24 months, must qualify under the Initial 3 restriction and meet the relevant criteria and index of disease severity.

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have a documented history of severe active juvenile idiopathic arthritis with onset prior to the age of 18 years, **AND**
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with disease modifying anti-rheumatic drugs (DMARDs) which must include at least 3 months continuous treatment with each of at least 2 DMARDs, one of which must be methotrexate at a dose of at least 20 mg weekly and one of which must be: (i) hydroxychloroquine at a dose of at least 200 mg daily; or (ii) leflunomide at a dose of at least 10 mg daily; or (iii) sulfasalazine at a dose of at least 2 g daily; OR
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs which, if methotrexate is contraindicated according to the Therapeutic Goods Administration (TGA)-approved Product Information or cannot be tolerated at a 20 mg weekly dose, must include at least 3 months continuous treatment with each of at least 2 of the following DMARDs: (i) hydroxychloroquine at a dose of at least 200 mg daily; and/or (ii) leflunomide at a dose of at least 10 mg daily; and/or (iii) sulfasalazine at a dose of at least 2 g daily; OR
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs which, if 3 or more of methotrexate, hydroxychloroquine, leflunomide and sulfasalazine are contraindicated according to the relevant TGA-approved Product Information or cannot be tolerated at the doses specified above, must include at least 3 months continuous treatment with each of at least 2 DMARDs, with one or more of the following DMARDs being used in place of the DMARDs which are contraindicated or not tolerated: (i) azathioprine at a dose of at least 1 mg/kg per day; and/or (ii) cyclosporin at a dose of at least 2 mg/kg/day; and/or (iii) sodium aurothiomalate at a dose of 50 mg weekly, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

If methotrexate is contraindicated according to the TGA-approved Product Information or cannot be tolerated at a 20 mg weekly dose, the application must include details of the contraindication or intolerance to methotrexate. The maximum tolerated dose of methotrexate must be documented in the application, if applicable.

The application must include details of the DMARDs trialed, their doses and duration of treatment, and all relevant contraindications and/or intolerances.

The requirement to trial at least 2 DMARDs for periods of at least 3 months each can be met using single agents sequentially or by using one or more combinations of DMARDs.

If the requirement to trial 6 months of intensive DMARD therapy with at least 2 DMARDs cannot be met because of contraindications and/or intolerances of a severity necessitating permanent treatment withdrawal to all of the DMARDs specified above, details of the contraindication or intolerance and dose for each DMARD must be provided in the authority application.

The following criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; AND either

(a) an active joint count of at least 20 active (swollen and tender) joints; or

(b) at least 4 active joints from the following list:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The joint count and ESR and/or CRP must be determined at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. All measures must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

The authority application must be made in writing and must include:

(1) completed authority prescription form(s); and

(2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

An assessment of a patient's response to an initial course of treatment must be conducted following a minimum of 12 weeks of therapy. An application for the continuing treatment must be accompanied with the assessment of response and submitted to the Department of Human Services no later than 4 weeks from the date of completion of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note The Department of Human Services website (www.humanservices.gov.au) has details of the toxicities, including severity, which will be accepted for the following purposes:

(a) exempting a patient from the requirement to undertake a minimum 3 month trial of methotrexate at a 20 mg weekly dose;

(b) substituting azathioprine, cyclosporin or sodium aurothiomalate for another DMARD as part of the 6 month intensive DMARD trial;

(c) exempting a patient from the requirement for a 6 month trial of intensive DMARD therapy.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 24 months)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have a documented history of severe active juvenile idiopathic arthritis with onset prior to the age of 18 years, **AND**
- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

- (a) an active joint count of fewer than 10 active (swollen and tender) joints; or
- (b) a reduction in the active (swollen and tender) joint count by at least 50% from baseline; or
- (c) a reduction in the number of the following active joints, from at least 4, by at least 50%:
 - (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - (ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The authority application must be made in writing and must include:

- (1) completed authority prescription form(s); and
- (2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to change or recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient who fails to demonstrate a response to treatment with this drug under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug in this treatment cycle. A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the initial 3 treatment restriction.

If a patient fails to respond to PBS-subsidised biological medicine treatment 3 times (once with each agent) they will not be eligible to receive further PBS-subsidised biological medicine therapy in this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 3 (recommencement of treatment after a break in biological medicine of more than 24 months)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 24 months or more from the most recently approved PBS-subsidised biological medicine for this condition; OR
- Patient must not have received PBS-subsidised biological medicine for at least 5 years if they failed or ceased to respond to PBS-subsidised biological medicine treatment 3 times in their last treatment cycle, **AND**
- The condition must have an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; OR
- The condition must have a C-reactive protein (CRP) level greater than 15 mg per L, **AND**
- The condition must have either (a) a total active joint count of at least 20 active (swollen and tender) joints; or (b) at least 4 active major joints, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Active joints are defined as:

- (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

All measures of joint count must be no more than 4 weeks old at the time of this application.

The authority application must be made in writing and must include:

(1) completed authority prescription form(s); and

(2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient) or Initial 2 (change or recommencement of treatment after break of less than 24 months) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 24 months) - balance of supply

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 24 months) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 24 months) to complete 16 weeks of treatment, **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

etanercept 50 mg/mL injection, 4 x 1 mL pen devices

3447K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	1050.08	41.30	Enbrel [PF]

etanercept 25 mg injection [4 vials] (&) inert substance diluent [4 x 1 mL syringes], 1 pack

3445H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	3	..	*1050.10	41.30	Enbrel [PF]

etanercept 50 mg/mL injection, 4 x 1 mL syringes

3446J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	1050.08	41.30	Enbrel [PF]

■ ETANERCEPT

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE PSORIATIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab for adult patients with severe active psoriatic arthritis. Therefore, where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient receiving PBS-subsidised treatment for psoriatic arthritis is able to commence a treatment cycle where they may trial biological medicines without having to experience a disease flare when swapping to the alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

A patient who received PBS-subsidised adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab or ustekinumab treatment prior to 1 May 2019 is considered to start their first cycle as of 1 May 2019.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence another cycle [further details are under '(5) Recommencement of treatment after a 5-year break in PBS-subsidised therapy' below].

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe active psoriatic arthritis.

(1) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an alternate medicine (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years) or An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, golimumab secukinumab and tofacitinib, 18 to 20 weeks of therapy for certolizumab pegol (depending upon the dosing regimen), 20 weeks of therapy for ixekizumab, 22 weeks of therapy for infliximab, and 28 weeks of therapy for ustekinumab. It is recommended that a patient be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of PBS-subsidised initial treatment following a minimum of 12 weeks of therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

Grandfather patients (ixekizumab only).

A patient who commenced treatment with ixekizumab for severe psoriatic arthritis prior to 1 March 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

Grandfather patients (tofacitinib only).

A patient who commenced treatment with Tofacitinib for severe psoriatic arthritis prior to 1 May 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has

experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment under the Initial 1 or Initial 2 treatment restrictions.

For the second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to re-qualify with respect to either the indices of disease severity (i.e. erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP) level, and active joint count) or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify under the Initial 3 treatment restriction with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application.

However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

Within a treatment cycle a patient may alternate between therapy with any biological medicine of their choice (1 at a time) providing:

(i) they have not received PBS-subsidised treatment with that particular biological medicine previously; or
(ii) they have demonstrated an adequate response to that particular biological medicine if they have previously trialled it on the PBS; and

(iii) they have not previously failed to respond to treatment 3 times in this treatment cycle with that biological medicine.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(4) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the indices of disease severity submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and these revised baseline measurements will be used to assess response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to determine response for all subsequent continuing treatments. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. 20 or more active joints), response will be determined according to a reduction in the total number of active joints.

(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment restriction according to the criteria of the relevant restriction and index of disease severity. The application must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Severe psoriatic arthritis

Treatment Phase: First continuing treatment

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

An adequate response to treatment is defined as:

an erythrocyte sedimentation rate (ESR) no greater than 25 mm per hour or a C-reactive protein (CRP) level no greater than 15 mg per L or either marker reduced by at least 20% from baseline; and

either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following major active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to determine response for all subsequent continuing treatments.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Severe Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either Initial 1, Initial 2, or Initial 3 treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Continuing treatment - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the first continuing treatment restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the subsequent continuing Authority Required (in writing) treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

etanercept 50 mg/mL injection, 4 x 1 mL pen devices

9458T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	1050.08	41.30	^a Brenzys [RF]	^a Enbrel [PF]

etanercept 25 mg injection [4 vials] (&) inert substance diluent [4 x 1 mL syringes], 1 pack

9036N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*1050.10	41.30	Enbrel [PF]

etanercept 50 mg/mL injection, 4 x 1 mL syringes

9088H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	1050.08	41.30	^a Brenzys [RF]	^a Enbrel [PF]

■ ETANERCEPT

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of

1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years); or

(iii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or

(iv) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient be reviewed in the 4 weeks prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

Grandfather patients (risankizumab only).

A patient who commenced treatment with risankizumab for chronic plaque psoriasis prior to 1 December 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the Initial treatment Grandfather treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment. This assessment must be conducted within 4 weeks of the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the 4 weeks prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a

patient is reviewed in the 4 weeks prior to completing their current course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure patients receive the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

Services Australia will determine whether a response to treatment has been demonstrated based on the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 the PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Subsequent continuing treatment, whole body

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per subsequent continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

The authority application must be made in writing and must include:

- a completed authority prescription form(s); and
- a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet including the date of the assessment of the patient's condition.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available

on the Department of Human Services website at www.humanservices.gov.au
Applications for authority to prescribe should be forwarded to:
Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Subsequent continuing treatment, Face, hand, foot

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition under the First continuing treatment restriction, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per subsequent continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or
- (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet including the date of the assessment of the patient's condition.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Whole body or Continuing treatment, Face, hand, foot - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the first continuing treatment, Whole body restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the first continuing treatment, Face, hand, foot restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the subsequent continuing treatment Authority Required (in writing), Whole body restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the subsequent continuing treatment Authority Required (in writing), Face, hand, foot restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restrictions, **AND**

- The treatment must be as systemic monotherapy (other than methotrexate).

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

etanercept 50 mg/mL injection, 4 x 1 mL pen devices

11222P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	1050.08	41.30	^a Brenzys [RF]	^a Enbrel [PF]

etanercept 25 mg injection [4 vials] (&) inert substance diluent [4 x 1 mL syringes], 1 pack

11223Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*1050.10	41.30	Enbrel [PF]

etanercept 50 mg/mL injection, 4 x 1 mL syringes

11224R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	1050.08	41.30	^a Brenzys [RF]	^a Enbrel [PF]

■ ETANERCEPT
Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

- a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or
- a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years); or
- a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or
- a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab and 28

weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient be reviewed in the 4 weeks prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

Grandfather patients (risankizumab only).

A patient who commenced treatment with risankizumab for chronic plaque psoriasis prior to 1 December 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the Initial treatment Grandfather treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment. This assessment must be conducted within 4 weeks of the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the 4 weeks prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the 4 weeks prior to completing their current course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal.

To ensure patients receive the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

Services Australia will determine whether a response to treatment has been demonstrated based on the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 the PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: First continuing treatment, Whole body

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**

- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: First continuing treatment, Face, hand, foot

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or
- (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet and face, hand, foot area diagrams including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

The PASI assessment for first continuing or subsequent continuing treatment must be performed on the same affected area assessed at baseline.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Whole body or Continuing treatment, Face, hand, foot - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the first continuing treatment, Whole body restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the first continuing treatment, Face, hand, foot restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restrictions, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate).

Treatment criteria:

- Must be treated by a dermatologist.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the

Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

etanercept 50 mg/mL injection, 4 x 1 mL pen devices

9462B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	1050.08	41.30	^a Brenzys [RF]	^a Enbrel [PF]

etanercept 25 mg injection [4 vials] (&) inert substance diluent [4 x 1 mL syringes], 1 pack

9429G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*1050.10	41.30	Enbrel [PF]

etanercept 50 mg/mL injection, 4 x 1 mL syringes

9431J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	1050.08	41.30	^a Brenzys [RF]	^a Enbrel [PF]

■ ETANERCEPT

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for adults with severe active rheumatoid arthritis. Where the term biological medicine appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alfa antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab), the T-cell co-stimulation modulator (abatacept) and the Janus kinase (JAK) inhibitors (baricitinib, tofacitinib, upadacitinib).

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised biological medicine therapy may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once, and
- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised biological medicines for the treatment of rheumatoid arthritis.

A serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Stevens Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered a treatment failure.

A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction. A patient who has failed fewer than 5 biological medicines and who has a break in treatment of less than 24 months may commence a further course of treatment with a biological medicine under Initial 2 treatment restriction. A patient who has failed fewer than 5 biological medicines and who has had a break in therapy of longer than 24 months may commence a further course of treatment with a biological medicine under the Initial 3 treatment restriction. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine treatment is stopped to the date of the new application for treatment with a biological medicine.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 April 2019.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised biological medicine treatment and wishes to commence such therapy, excluding rituximab (Initial 1 - new patient); or

(ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to re-commence treatment with a specific biological medicine following a break of less than 24 months in PBS-subsidised therapy with that agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months).

(iv) a patient wishes to re-commence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 24 months (Initial 3 - re-commencement of treatment after a break in biological medicine of more than 24 months). Initial applications for a new patient (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6-month intensive DMARD trial, but prior to ceasing DMARD therapy.

Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, baricitinib, etanercept, golimumab, tocilizumab, tofacitinib and upadacitinib, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

Rituximab patients should be assessed following a minimum of 12 weeks after the first infusion, and the assessment should be submitted to Services Australia within 4 weeks to ensure continuity of treatment for those who meet the continuing restriction for PBS subsidised treatment with this drug for this condition.

Where a response assessment is not provided the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment. For second and subsequent courses of PBS-subsidised biological medicine (excluding rituximab) treatment, it is recommended that a patient be reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to Services Australia within sufficient time to allow processing. Abatacept patients:

A patient is eligible to receive one I.V. loading dose when commencing treatment with the subcutaneous formulation. Two prescriptions are required, the first prescription for the I.V. loading dose for sufficient vials for one dose based on the patient's weight with no repeats. The second prescription for the subcutaneous formulation, with a maximum quantity of 4 and up to 3 repeats, must be submitted with the initial application.

Rituximab patients:

A further application may be submitted to Services Australia within sufficient time to allow processing. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

Assessment of the patient's response to treatment must be conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the cessation of the most recent course of biological medicine therapy.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction. Where a response assessment is not conducted within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

(2) Swapping therapy

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non- biological medicine therapy requirements except if the patient has had a break in therapy of more than 24 months who would need to requalify with respect to the indices of disease severity. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each biological medicine trialled.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent, unless they have experienced a serious adverse reaction of a severity necessitating permanent treatment withdrawal.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug.

Abatacept:

A patient swapping from I.V. abatacept to subcutaneous abatacept will not be eligible for an I.V. loading dose when commencing treatment with the subcutaneous formulation.

Rituximab:

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised

TNF- α antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they be assessed for response to every course of treatment, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological medicine during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological medicine therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate biological medicine may do so without having to have any treatment-free period.

(3) Baseline measurements to determine response.

Determination of whether a response to treatment has been demonstrated must be based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and the eligibility for continuing treatment must be assessed according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response must be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be determined on the total number of major joints.

Applications under the Initial 1 treatment restriction for a new patient must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. The results must be no more than 4 weeks old at the time of application.

Applications under the Initial 3 treatment restriction for re-commencement of treatment after a break in biological medicine of more than 24 months must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with disease modifying anti-rheumatic drugs (DMARDs) which must include at least 3 months continuous treatment with each of at least 2 DMARDs, one of which must be methotrexate at a dose of at least 20 mg weekly and one of which must be: (i) hydroxychloroquine at a dose of at least 200 mg daily; or (ii) leflunomide at a dose of at least 10 mg daily; or (iii) sulfasalazine at a dose of at least 2 g daily; OR
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs which, if methotrexate is contraindicated according to the Therapeutic Goods Administration (TGA)-approved Product Information or cannot be tolerated at a 20 mg weekly dose, must include at least 3 months continuous treatment with each of at least 2 of the following DMARDs: (i) hydroxychloroquine at a dose of at least 200 mg daily; and/or (ii) leflunomide at a dose of at least 10 mg daily; and/or (iii) sulfasalazine at a dose of at least 2 g daily; OR
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs which, if 3 or more of methotrexate, hydroxychloroquine, leflunomide and sulfasalazine are contraindicated according to the relevant TGA-approved Product Information or cannot be tolerated at the doses specified above, must include at least 3 months continuous treatment with each of at least 2 DMARDs, with one or more of the following DMARDs being used in place of the DMARDs which are contraindicated or not tolerated: (i) azathioprine at a dose of at least 1 mg/kg per day; and/or (ii) cyclosporin at a dose of at least 2 mg/kg/day; and/or (iii) sodium aurothiomalate at a dose of 50 mg weekly, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

If methotrexate is contraindicated according to the TGA-approved product information or cannot be tolerated at a 20 mg weekly dose, the application must include details of the contraindication or intolerance including severity to methotrexate. The maximum tolerated dose of methotrexate must be documented in the application, if applicable.

The application must include details of the DMARDs trialled, their doses and duration of treatment, and all relevant contraindications and/or intolerances including severity.

The requirement to trial at least 2 DMARDs for periods of at least 3 months each can be met using single agents sequentially or by using one or more combinations of DMARDs.

If the requirement to trial 6 months of intensive DMARD therapy with at least 2 DMARDs cannot be met because of contraindications and/or intolerances of a severity necessitating permanent treatment withdrawal to all of the DMARDs specified above, details of the contraindication or intolerance including severity and dose for each DMARD must be provided in the authority application.

The following criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; AND either

(a) a total active joint count of at least 20 active (swollen and tender) joints; or

(b) at least 4 active joints from the following list of major joints:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The joint count and ESR and/or CRP must be determined at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. All measures must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note Biosimilar prescribing policy Prescribing of the biosimilar brand Brenzys is encouraged for treatment naive patients.

Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note The Department of Human Services website (www.humanservices.gov.au) has details of the toxicities, including severity, which will be accepted for the following purposes:

(a) exempting a patient from the requirement to undertake a minimum 3 month trial of methotrexate at a 20 mg weekly dose;

(b) substituting azathioprine, cyclosporin or sodium aurothiomalate for another DMARD as part of the 6 month intensive DMARD trial;

(c) exempting a patient from the requirement for a 6 month trial of intensive DMARD therapy.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 2 (change or re-commencement of treatment after a break in biological medicine of less than 24 months)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must not have failed to respond to previous PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised biological medicine treatment for this condition 5 times, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either of the Initial 1, Initial 2, Initial 3, first or subsequent continuing treatment restrictions, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

A patient who has demonstrated a response to a course of rituximab must have a PBS-subsidised biological therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 3 (re-commencement of treatment after a break in biological medicine of more than 24 months)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 24 months or more from the most recent PBS-subsidised biological medicine for this condition, **AND**
- Patient must not have failed to respond to previous PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised biological medicine treatment for this condition 5 times, **AND**
- The condition must have an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; OR
- The condition must have a C-reactive protein (CRP) level greater than 15 mg per L, **AND**
- The condition must have either (a) a total active joint count of at least 20 active (swollen and tender) joints; or (b) at least 4 active major joints, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Major joints are defined as (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

All measures of joint count and ESR and/or CRP must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note Biosimilar prescribing policy Prescribing of the biosimilar brand Brenzys is encouraged for treatment naive patients. Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos. Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial 1 (new patient) or Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 24 months) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 24 months) - balance of supply

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 24 months) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 24 months) to complete 16 weeks of treatment, **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

etanercept 50 mg/mL injection, 4 x 1 mL pen devices

9459W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	3	..	1050.08	41.30	^a Brenzys [RF]	^a Enbrel [PF]

etanercept 25 mg injection [4 vials] (&) inert substance diluent [4 x 1 mL syringes], 1 pack

8637N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	3	..	*1050.10	41.30	Enbrel [PF]

etanercept 50 mg/mL injection, 4 x 1 mL syringes

9089J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	3	..	1050.08	41.30	^a Brenzys [RF]	^a Enbrel [PF]

■ ETANERCEPT

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE PSORIATIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and

ustekinumab for adult patients with severe active psoriatic arthritis. Therefore, where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient receiving PBS-subsidised treatment for psoriatic arthritis is able to commence a treatment cycle where they may trial biological medicines without having to experience a disease flare when swapping to the alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

A patient who received PBS-subsidised adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab or ustekinumab treatment prior to 1 May 2019 is considered to start their first cycle as of 1 May 2019.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence another cycle [further details are under '(5) Recommencement of treatment after a 5-year break in PBS-subsidised therapy' below].

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe active psoriatic arthritis.

(1) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an alternate medicine (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years) or An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, golimumab secukinumab and tofacitinib, 18 to 20 weeks of therapy for certolizumab pegol (depending upon the dosing regimen), 20 weeks of therapy for ixekizumab, 22 weeks of therapy for infliximab, and 28 weeks of therapy for ustekinumab. It is recommended that a patient be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of PBS-subsidised initial treatment following a minimum of 12 weeks of therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

Grandfather patients (ixekizumab only).

A patient who commenced treatment with ixekizumab for severe psoriatic arthritis prior to 1 March 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

Grandfather patients (tofacitinib only).

A patient who commenced treatment with Tofacitinib for severe psoriatic arthritis prior to 1 May 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment. Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment under the Initial 1 or Initial 2 treatment restrictions.

For the second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to re-qualify with respect to either the indices of disease severity (i.e. erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP) level, and active joint count) or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify under the Initial 3 treatment restriction with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application.

However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

Within a treatment cycle a patient may alternate between therapy with any biological medicine of their choice (1 at a time) providing:

- (i) they have not received PBS-subsidised treatment with that particular biological medicine previously; or
- (ii) they have demonstrated an adequate response to that particular biological medicine if they have previously trialled it on the PBS; and
- (iii) they have not previously failed to respond to treatment 3 times in this treatment cycle with that biological medicine.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(4) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the indices of disease severity submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and these revised baseline measurements will be used to assess response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to determine response for all subsequent continuing treatments. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. 20 or more active joints), response will be determined according to a reduction in the total number of active joints.

(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment restriction according to the criteria of the relevant restriction and index of disease severity. The application must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Clinical criteria:

- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response to methotrexate at a dose of at least 20 mg weekly for a minimum period of 3 months, **AND**
- Patient must have failed to achieve an adequate response to sulfasalazine at a dose of at least 2 g per day for a minimum period of 3 months; **OR**
- Patient must have failed to achieve an adequate response to leflunomide at a dose of up to 20 mg daily for a minimum period of 3 months, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; **OR**
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Where treatment with methotrexate, sulfasalazine or leflunomide is contraindicated according to the relevant TGA-approved Product Information, details must be provided at the time of application.

Where intolerance to treatment with methotrexate, sulfasalazine or leflunomide developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

The following initiation criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; and

either

(a) an active joint count of at least 20 active (swollen and tender) joints; or

(b) at least 4 active joints from the following list of major joints:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Severe Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

An assessment of a patient's response to an initial course of treatment must be conducted following a minimum of 12 weeks of therapy. An application for the continuing treatment must be accompanied with the assessment of response and submitted to the Department of Human Services no later than 4 weeks from the date of completion of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 3 months treatment with methotrexate and 3 months treatment with sulfasalazine or leflunomide can be found on the Department of Human Services website (www.humanservices.gov.au)

Note Biosimilar prescribing policy Prescribing of the biosimilar brand Brenzys is encouraged for treatment naive patients. Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial treatment - Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

An adequate response to treatment is defined as:

an erythrocyte sedimentation rate (ESR) no greater than 25 mm per hour or a C-reactive protein (CRP) level no greater than 15 mg per L or either marker reduced by at least 20% from baseline; and

either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following major active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth). The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Severe Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to change or recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3, first or subsequent continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Biosimilar prescribing policy Prescribing of the biosimilar brand Brenzys is encouraged for treatment naive patients.

Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial treatment - Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must have an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; OR
- The condition must have a C-reactive protein (CRP) level greater than 15 mg per L, **AND**
- The condition must have either (a) a total active joint count of at least 20 active (swollen and tender) joints; or (b) at least 4 active major joints, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Major joints are defined as (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

All measures of joint count and ESR and/or CRP must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Severe Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3, first or subsequent continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Biosimilar prescribing policy Prescribing of the biosimilar brand Brenzys is encouraged for treatment naive patients. Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial 1 (new patient) or Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years) - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 16 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

etanercept 50 mg/mL injection, 4 x 1 mL pen devices

9457R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	3	..	1050.08	41.30	^a Brenzys [RF]	^a Enbrel [PF]

etanercept 25 mg injection [4 vials] (&) inert substance diluent [4 x 1 mL syringes], 1 pack

9035M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	3	..	*1050.10	41.30	Enbrel [PF]

etanercept 50 mg/mL injection, 4 x 1 mL syringes

9087G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	3	..	1050.08	41.30	^a Brenzys [RF]	^a Enbrel [PF]

■ ETANERCEPT

Note TREATMENT OF ADULT PATIENTS WITH ANKYLOSING SPONDYLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab for adult patients with ankylosing

spondylitis.

Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 7 biological medicines at any 1 time.

Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy.

A patient who has been receiving PBS-subsidised adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab treatment prior to 1 December 2020 is considered to start their first cycle as of 1 December 2020.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last prescription for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised biological medicine treatment with adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised biological medicine treatment in this treatment cycle and wishes to commence such therapy (Initial 1 - New patient)

(ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years); or

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

A patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy.

(b) Continuing treatment.

For the first continuing treatment course of PBS-subsidised biological medicine, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1, Initial 2 or Initial 3 treatment restrictions. For second and subsequent continuing courses of PBS-subsidised biological medicine it is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply. Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be conducted no later than 4 weeks from the cessation of that treatment course. Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI), or the prior NSAID therapy and exercise program requirements.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(3) Baseline measurements to determine response.

Services Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the BASDAI, ESR and/or CRP submitted with the first authority application for a biological medicine. For a new patient, the BASDAI used to determine the baseline must be measured while the patient is receiving NSAID therapy and completing their exercise program.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response.

Prescribers may provide new baseline measurements any time an 'Initial treatment' authority application is submitted within a treatment cycle and the eligibility for continuing treatment must be assessed according to these revised baseline

measurements.

(4) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must qualify under the Initial 3 treatment restriction. The same clinical criteria and indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI) as for the Initial 1 (New patient) restriction will need to be met, but a re-trial of NSAID therapy and exercise therapy is not required.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Ankylosing spondylitis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Clinical criteria:

- The condition must be radiographically (plain X-ray) confirmed Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have at least 2 of the following: (i) low back pain and stiffness for 3 or more months that is relieved by exercise but not by rest; or (ii) limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by a score of at least 1 on each of the lumbar flexion and lumbar side flexion measurements of the Bath Ankylosing Spondylitis Metrology Index (BASMI); or (iii) limitation of chest expansion relative to normal values for age and gender,

AND

- Patient must have failed to achieve an adequate response following treatment with at least 2 non-steroidal anti-inflammatory drugs (NSAIDs), whilst completing an appropriate exercise program, for a total period of 3 months, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

The application must include details of the NSAIDs trialled, their doses and duration of treatment.

If the NSAID dose is less than the maximum recommended dose in the relevant TGA-approved Product Information, the application must include the reason a higher dose cannot be used.

If treatment with NSAIDs is contraindicated according to the relevant TGA-approved Product Information, the application must provide details of the contraindication.

If intolerance to NSAID treatment develops during the relevant period of use which is of a severity to necessitate permanent treatment withdrawal, the application must provide details of the nature and severity of this intolerance.

The following criteria indicate failure to achieve an adequate response and must be demonstrated at the time of the initial application:

- (a) a Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 4 on a 0-10 scale; **AND**
- (b) an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 10 mg per L.

The BASDAI must be determined at the completion of the 3 month NSAID and exercise trial, but prior to ceasing NSAID treatment. The BASDAI must be no more than 1 month old at the time of initial application.

Both ESR and CRP measures should be provided with the initial treatment application and both must be no more than 1 month old. If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reason this criterion cannot be satisfied.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Ankylosing Spondylitis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) a copy of the radiological report confirming Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis; and
 - (ii) a completed BASDAI Assessment Form; and
 - (iii) a completed Exercise Program Self Certification Form included in the supporting information form.

An assessment of a patient's response to an initial course of treatment must be conducted following a minimum of 12 weeks of therapy. An application for the continuing treatment must be accompanied with the assessment of response and submitted to the Department of Human Services no later than 4 weeks from the date of completion of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Biosimilar prescribing policy Prescribing of the biosimilar brand Brenzys is encouraged for treatment naive patients. Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Details of the toxicities, including severity, which will be accepted for the purposes of administering this restriction can be found on the Department of Human Services website at www.humanservices.gov.au

Note For details on the appropriate minimum exercise program that will be accepted for the purposes of administering this restriction, please refer to the Department of Human Services website at www.humanservices.gov.au

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Ankylosing spondylitis

Treatment Phase: Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Ankylosing Spondylitis PBS Authority Application - Supporting Information Form.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to change or recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

An adequate response is defined as an improvement from baseline of at least 2 of the BASDAI and 1 of the following:

- (a) an ESR measurement no greater than 25 mm per hour; or
- (b) a CRP measurement no greater than 10 mg per L; or
- (c) an ESR or CRP measurement reduced by at least 20% from baseline.

Where only 1 acute phase reactant measurement is supplied in the first application for PBS-subsidised treatment, that same marker must be measured and supplied in all subsequent continuing treatment applications.

All measurements provided must be no more than 1 month old at the time of application.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Ankylosing spondylitis

Treatment Phase: Initial treatment - Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must be radiographically (plain X-ray) confirmed Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis, **AND**
- Patient must have at least 2 of the following: (i) low back pain and stiffness for 3 or more months that is relieved by exercise but not by rest; or (ii) limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by a score of at least 1 on each of the lumbar flexion and lumbar side flexion measurements of the Bath Ankylosing Spondylitis Metrology Index (BASMI); or (iii) limitation of chest expansion relative to normal values for age and gender,

AND

- Patient must have a Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 4 on a 0-10 scale that is no more than 4 weeks old at the time of application, **AND**
- Patient must have an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour that is no more than 4 weeks old at the time of application; OR
- Patient must have a C-reactive protein (CRP) level greater than 10 mg per L that is no more than 4 weeks old at the time of application; OR
- Patient must have a clinical reason as to why demonstration of an elevated ESR or CRP cannot be met and the application must state the reason, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Ankylosing Spondylitis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) a copy of the radiological report confirming Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis; and
 - (ii) a completed BASDAI Assessment Form.

An assessment of a patient's response to an initial course of treatment must be conducted following a minimum of 12 weeks of therapy. An application for the continuing treatment must be accompanied with the assessment of response and submitted to the Department of Human Services no later than 4 weeks from the date of completion of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Biosimilar prescribing policy Prescribing of the biosimilar brand Brenzys is encouraged for treatment naive patients.

Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Ankylosing spondylitis

Treatment Phase: Initial treatment - Initial 1 (new patient), Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years) - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 16 weeks treatment; OR

- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 16 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

etanercept 50 mg/mL injection, 4 x 1 mL pen devices

9455P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	3	..	1050.08	41.30	^a Brenzys [RF]	^a Enbrel [PF]

etanercept 25 mg injection [4 vials] (&) inert substance diluent [4 x 1 mL syringes], 1 pack

8778B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	3	..	*1050.10	41.30	Enbrel [PF]

etanercept 50 mg/mL injection, 4 x 1 mL syringes

9085E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	3	..	1050.08	41.30	^a Brenzys [RF]	^a Enbrel [PF]

■ ETANERCEPT
Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

- a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or
- a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years); or
- a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or
- a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised

therapy of less than 5 years with the same medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient be reviewed in the 4 weeks prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

Grandfather patients (risankizumab only).

A patient who commenced treatment with risankizumab for chronic plaque psoriasis prior to 1 December 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the Initial treatment Grandfather treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment. This assessment must be conducted within 4 weeks of the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the 4 weeks prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the 4 weeks prior to completing their current course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure patients receive the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

Services Australia will determine whether a response to treatment has been demonstrated based on the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 the PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Whole body (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis where lesions have been present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 5 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; (iii) ciclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks; (v) apremilast at a dose of 30 mg twice a day for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, ciclosporin, apremilast or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application.

Where intolerance to treatment with phototherapy, methotrexate, ciclosporin, apremilast or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, ciclosporin, apremilast, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

- (a) A current Psoriasis Area and Severity Index (PASI) score of greater than 15, as assessed, preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior treatment.
- (b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 1 month following cessation of each course of treatment.
- (c) The most recent PASI assessment must be no more than 1 month old at the time of application.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and
 - (ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to Services Australia no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition. Demonstration of response should be provided within this timeframe.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Biosimilar prescribing policy Prescribing of the biosimilar brand Brenzys is encouraged for treatment naive patients.

Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, ciclosporin, apremilast or acitretin can be found on the Services Australia website (www.servicesaustralia.gov.au).

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Whole body (change or recommencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either of the Initial 1, Initial 2, Initial 3, first or subsequent continuing treatment restrictions, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition. Demonstration of response should be provided within this timeframe.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and
 - (ii) details of prior biological treatment, including dosage, date and duration of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Biosimilar prescribing policy Prescribing of the biosimilar brand Brenzys is encouraged for treatment naive patients.

Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 3, Whole body (re-commencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**

- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must have a current Psoriasis Area and Severity Index (PASI) score of greater than 15, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

The most recent PASI assessment must be no more than 1 month old at the time of application.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition.

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition. Demonstration of response should be provided within this timeframe.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Biosimilar prescribing policy Prescribing of the biosimilar brand Brenzys is encouraged for treatment naive patients.

Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Face, hand, foot (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 5 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; (iii) ciclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks; (v) apremilast at a dose of 30 mg twice a day for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, ciclosporin, apremilast or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application.

Where intolerance to treatment with phototherapy, methotrexate, ciclosporin, apremilast or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, ciclosporin, apremilast, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

(a) Chronic plaque psoriasis classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:

(i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe, as assessed, preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior treatment; or

(ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed, preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior treatment;

(b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 1 month following cessation of each course of treatment.

(c) The most recent PASI assessment must be no more than 1 month old at the time of application.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition; and

(ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to Services Australia no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition. Demonstration of response should be provided within this timeframe.

The PASI assessment for first continuing or subsequent continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, ciclosporin, apremilast or acitretin can be found on the Services Australia website (www.servicesaustralia.gov.au).

Note Biosimilar prescribing policy Prescribing of the biosimilar brand Brenzys is encouraged for treatment naive patients. Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos. Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Face, hand, foot (change or recommencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**

- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or
- (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either of the Initial 1, Initial 2, Initial 3, first or subsequent continuing treatment restrictions, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition. Demonstration of response should be provided within this timeframe. The PASI assessment for first continuing or subsequent continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition; and
 - (ii) details of prior biological treatment, including dosage, date and duration of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Biosimilar prescribing policy Prescribing of the biosimilar brand Brenzys is encouraged for treatment naive patients. Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos. Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 3, Face, hand, foot (re-commencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must be classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:
 - (i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or (ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

The most recent PASI assessment must be no more than 1 month old at the time of application.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition.

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition. Demonstration of response should be provided within this timeframe.

The PASI assessment for first continuing or subsequent continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Biosimilar prescribing policy Prescribing of the biosimilar brand Brenzys is encouraged for treatment naive patients.

Encouraging biosimilar prescribing for treatment naive patients is Government policy. A viable biosimilar market is expected to result in reduced costs for biological medicines, allowing the Government to reinvest in new treatments. Further information can be found on the Biosimilar Awareness Initiative webpage (www.health.gov.au/biosimilars).

Note Prescribers must include the proprietary name (brand) on the prescription to ensure the appropriate item is approved.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Whole body or Face, hand, foot (new patient) or Initial 2, Whole body or Face, hand, foot (change or re-commencement of treatment after a break in biological medicine of less than 5 years) or Initial 3, Whole body or Face, hand, foot (re-commencement of treatment after a break in biological medicine of more than 5 years) - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1, Whole body (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2, Whole body (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3, Whole body (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 1, Face, hand, foot (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2, Face, hand, foot (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3, Face, hand, foot (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 16 weeks treatment, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a dermatologist.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

etanercept 50 mg/mL injection, 4 x 1 mL pen devices

9461Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	3	..	1050.08	41.30	^a Brenzys [RF]	^a Enbrel [PF]

etanercept 25 mg injection [4 vials] (&) inert substance diluent [4 x 1 mL syringes], 1 pack

9037P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	3	..	*1050.10	41.30	Enbrel [PF]	

etanercept 50 mg/mL injection, 4 x 1 mL syringes

9091L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	3	..	1050.08	41.30	^a Brenzys [RF]	^a Enbrel [PF]

■ ETANERCEPT

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment (whole body)

TREATMENT OF PATIENTS UNDER 18 YEARS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing of etanercept under the Pharmaceutical Benefits Scheme (PBS) for patients under 18 years with severe chronic plaque psoriasis.

Applications for treatment of this condition will be limited to provide patients with a maximum of 24 weeks of therapy per course of treatment. A maximum of 16 weeks treatment with etanercept will be authorised for the primary application. The balance, a further 8 weeks treatment, will be authorised if the submitted Psoriasis Area and Severity Index (PASI) assessment demonstrates an adequate response to treatment. Where fewer than 3 repeats are requested at the time of the authority application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone.

Once a patient has failed to respond to treatment 2 times, they must have, at a minimum, a 12 month break. The length of a treatment break is measured from the date the most recent treatment was stopped to the date of the first application for initial treatment.

There are separate restrictions for treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Application for approval for initial treatment.

Applications for a course of initial treatment should be made for patients who have received no prior PBS-subsidised biological treatment and wish to commence such therapy.

(2) Applications for approval for re-treatment.

Applications for re-treatment with etanercept should be made in the following situations:

(i) a patient who has received prior PBS-subsidised etanercept and experiences a disease flare, and wishes to start a second or subsequent treatment course with etanercept following a break of less than 12 months in PBS-subsidised therapy; or

(ii) a patient who has received and failed to respond to prior PBS-subsidised etanercept and wishes to start a second or subsequent treatment course following a break of less than 12 months in PBS-subsidised therapy.

For psoriasis affecting the whole body:

Patients are eligible for re-treatment due to disease flare if there is a 50% or greater change in the patients PASI score or the patient has a current PASI score of greater than 15, compared to the most recent response assessment following cessation of the most recent 24 weeks of PBS-subsidised etanercept.

For psoriasis affecting the face, hand or foot:

Patients are eligible for re-treatment due to disease flare if:

(i) all subscores are rated moderate to severe or 2 of the three subscores are rated severe to very severe; OR

(ii) the skin area affected is a 50% or greater change or the area affected is 30% or more of the face, palm of a hand or sole of a foot, compared to the most recent response assessment following cessation of the most recent 24 weeks of PBS-subsidised etanercept.

(3) Applications for approval for completion of a course

Applications for a further 8 weeks of treatment to allow for completion of 24 weeks of therapy should be submitted with a PASI assessment.

The PASI assessment must be conducted after at least 12 weeks of treatment.

This assessment must be submitted to Department of Human Services (the Department) within 1 month of the completion of 12 weeks of treatment. Where a response assessment is not undertaken and submitted to the Department within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological agent. In circumstances where it is not possible to submit a response assessment within these timeframes, please call the Department on 1800 700 270 to discuss.

(4) Baseline measurements to determine response.

The Department will determine whether a response to treatment has been demonstrated, based on the baseline PASI assessment submitted with the first authority application for etanercept. However, prescribers may provide new baseline

measurements any time that an initial or re-treatment authority is submitted and subsequent response will be assessed according to this revised PASI score.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of gaining approval for the remainder of 24 weeks treatment.

(5) Re-commencement of treatment after a 12 month break in PBS-subsidised therapy.

A patient who wishes to start a second or subsequent treatment course with etanercept following a break in PBS-subsidised etanercept therapy of at least 12 months, must requalify for treatment under the initial treatment restriction. The most recent PASI assessment must be no more than 1 month old at the time of application.

Treatment criteria:

- Must be treated by a dermatologist.

Population criteria:

- Patient must be under 18 years of age and a parent or authorised guardian must have signed a patient acknowledgement.

Clinical criteria:

- The treatment must be as systemic monotherapy; OR
- The treatment must be in combination with methotrexate, **AND**
- Patient must have lesions present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received any prior PBS-subsidised treatment with etanercept for this condition; OR
- Patient must not have received any PBS-subsidised treatment with etanercept for this condition for at least 12 months, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 3 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; and/or (ii) methotrexate at a dose of at least 10 mg or 10 mg per square metre weekly (whichever is lowest) for at least 6 weeks; and/or (iii) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks, **AND**
- Patient must not receive more than 16 weeks of treatment with etanercept under this restriction.

Where treatment with any of the above-mentioned drugs was contraindicated according to the relevant TGA-approved Product Information, or where phototherapy was contraindicated, details must be provided at the time of application.

Where intolerance to phototherapy, methotrexate and/or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

- (a) A current Psoriasis Area and Severity Index (PASI) score of greater than 15, as assessed, preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior treatment.
- (b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 1 month following cessation of each course of treatment.
- (c) The most recent PASI assessment must be no more than 1 month old at the time of application.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Severe Chronic Plaque Psoriasis in Patients Less Than 18 Years PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition and
 - (ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy]; and
 - (iii) the parent or authorised guardian signed patient and prescriber acknowledgements.

Where a patient has had a 12 month treatment break, the length of the break is measured from the date the most recent treatment was stopped to the date of the application to re-commence treatment.

Note Details of acceptable toxicities including severity, associated with phototherapy, methotrexate and acitretin, can be found on the Department of Human Services website at www.humanservices.gov.au

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment or Re-treatment (Whole body) - balance of first supply

TREATMENT OF PATIENTS UNDER 18 YEARS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing of etanercept under the Pharmaceutical Benefits Scheme (PBS) for patients under 18 years with severe chronic plaque psoriasis.

Applications for treatment of this condition will be limited to provide patients with a maximum of 24 weeks of therapy per course of treatment. A maximum of 16 weeks treatment with etanercept will be authorised for the primary application. The balance, a further 8 weeks treatment, will be authorised if the submitted Psoriasis Area and Severity Index (PASI) assessment demonstrates an adequate response to treatment. Where fewer than 3 repeats are requested at the time of the authority application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone.

Once a patient has failed to respond to treatment 2 times, they must have, at a minimum, a 12 month break. The length of a treatment break is measured from the date the most recent treatment was stopped to the date of the first application for initial treatment.

There are separate restrictions for treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Application for approval for initial treatment.

Applications for a course of initial treatment should be made for patients who have received no prior PBS-subsidised biological treatment and wish to commence such therapy.

(2) Applications for approval for re-treatment.

Applications for re-treatment with etanercept should be made in the following situations:

- (i) a patient who has received prior PBS-subsidised etanercept and experiences a disease flare, and wishes to start a second or subsequent treatment course with etanercept following a break of less than 12 months in PBS-subsidised therapy; or
- (ii) a patient who has received and failed to respond to prior PBS-subsidised etanercept and wishes to start a second or subsequent treatment course following a break of less than 12 months in PBS-subsidised therapy.

For psoriasis affecting the whole body:

Patients are eligible for re-treatment due to disease flare if there is a 50% or greater change in the patients PASI score or the patient has a current PASI score of greater than 15, compared to the most recent response assessment following cessation of the most recent 24 weeks of PBS-subsidised etanercept.

For psoriasis affecting the face, hand or foot:

Patients are eligible for re-treatment due to disease flare if:

- (i) all subscores are rated moderate to severe or 2 of the three subscores are rated severe to very severe; OR
- (ii) the skin area affected is a 50% or greater change or the area affected is 30% or more of the face, palm of a hand or sole of a foot, compared to the most recent response assessment following cessation of the most recent 24 weeks of PBS-subsidised etanercept.

(3) Applications for approval for completion of a course

Applications for a further 8 weeks of treatment to allow for completion of 24 weeks of therapy should be submitted with a PASI assessment.

The PASI assessment must be conducted after at least 12 weeks of treatment.

This assessment must be submitted to Department of Human Services (the Department) within 1 month of the completion of 12 weeks of treatment. Where a response assessment is not undertaken and submitted to the Department within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological agent. In circumstances where it is not possible to submit a response assessment within these timeframes, please call the Department on 1800 700 270 to discuss.

(4) Baseline measurements to determine response.

The Department will determine whether a response to treatment has been demonstrated, based on the baseline PASI assessment submitted with the first authority application for etanercept. However, prescribers may provide new baseline measurements any time that an initial or re-treatment authority is submitted and subsequent response will be assessed according to this revised PASI score.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of gaining approval for the remainder of 24 weeks treatment.

(5) Re-commencement of treatment after a 12 month break in PBS-subsidised therapy.

A patient who wishes to start a second or subsequent treatment course with etanercept following a break in PBS-subsidised etanercept therapy of at least 12 months, must requalify for treatment under the initial treatment restriction. The most recent PASI assessment must be no more than 1 month old at the time of application.

Treatment criteria:

- Must be treated by a dermatologist.

Clinical criteria:

- The treatment must be as systemic monotherapy; OR
- The treatment must be in combination with methotrexate, **AND**
- Patient must have received insufficient therapy under the Initial treatment (whole body) restriction for severe chronic plaque psoriasis to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy under the Re-treatment (whole body) restriction for severe chronic plaque psoriasis to complete 16 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Note Authority approval for sufficient therapy to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written application for authority approval for sufficient therapy to complete a maximum of 16 weeks of treatment should be forwarded to:

Department of Human Services
Complex Drugs

Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment or Re-treatment (Whole body) - completion of course

TREATMENT OF PATIENTS UNDER 18 YEARS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing of etanercept under the Pharmaceutical Benefits Scheme (PBS) for patients under 18 years with severe chronic plaque psoriasis.

Applications for treatment of this condition will be limited to provide patients with a maximum of 24 weeks of therapy per course of treatment. A maximum of 16 weeks treatment with etanercept will be authorised for the primary application. The balance, a further 8 weeks treatment, will be authorised if the submitted Psoriasis Area and Severity Index (PASI) assessment demonstrates an adequate response to treatment. Where fewer than 3 repeats are requested at the time of the authority application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone.

Once a patient has failed to respond to treatment 2 times, they must have, at a minimum, a 12 month break. The length of a treatment break is measured from the date the most recent treatment was stopped to the date of the first application for initial treatment.

There are separate restrictions for treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Application for approval for initial treatment.

Applications for a course of initial treatment should be made for patients who have received no prior PBS-subsidised biological treatment and wish to commence such therapy.

(2) Applications for approval for re-treatment.

Applications for re-treatment with etanercept should be made in the following situations:

(i) a patient who has received prior PBS-subsidised etanercept and experiences a disease flare, and wishes to start a second or subsequent treatment course with etanercept following a break of less than 12 months in PBS-subsidised therapy; or

(ii) a patient who has received and failed to respond to prior PBS-subsidised etanercept and wishes to start a second or subsequent treatment course following a break of less than 12 months in PBS-subsidised therapy.

For psoriasis affecting the whole body:

Patients are eligible for re-treatment due to disease flare if there is a 50% or greater change in the patients PASI score or the patient has a current PASI score of greater than 15, compared to the most recent response assessment following cessation of the most recent 24 weeks of PBS-subsidised etanercept.

For psoriasis affecting the face, hand or foot:

Patients are eligible for re-treatment due to disease flare if:

- (i) all subscores are rated moderate to severe or 2 of the three subscores are rated severe to very severe; OR
- (ii) the skin area affected is a 50% or greater change or the area affected is 30% or more of the face, palm of a hand or sole of a foot, compared to the most recent response assessment following cessation of the most recent 24 weeks of PBS-subsidised etanercept.

(3) Applications for approval for completion of a course

Applications for a further 8 weeks of treatment to allow for completion of 24 weeks of therapy should be submitted with a PASI assessment.

The PASI assessment must be conducted after at least 12 weeks of treatment.

This assessment must be submitted to Department of Human Services (the Department) within 1 month of the completion of 12 weeks of treatment. Where a response assessment is not undertaken and submitted to the Department within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological agent. In circumstances where it is not possible to submit a response assessment within these timeframes, please call the Department on 1800 700 270 to discuss.

(4) Baseline measurements to determine response.

The Department will determine whether a response to treatment has been demonstrated, based on the baseline PASI assessment submitted with the first authority application for etanercept. However, prescribers may provide new baseline measurements any time that an initial or re-treatment authority is submitted and subsequent response will be assessed according to this revised PASI score.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of gaining approval for the remainder of 24 weeks treatment.

(5) Re-commencement of treatment after a 12 month break in PBS-subsidised therapy.

A patient who wishes to start a second or subsequent treatment course with etanercept following a break in PBS-subsidised etanercept therapy of at least 12 months, must requalify for treatment under the initial treatment restriction. The most recent PASI assessment must be no more than 1 month old at the time of application.

Treatment criteria:

- Must be treated by a dermatologist.

Clinical criteria:

- The treatment must be as systemic monotherapy; OR
- The treatment must be in combination with methotrexate, **AND**
- Patient must have received 16 weeks treatment under the Initial treatment (whole body) restriction for severe chronic plaque psoriasis; OR
- Patient must have received 16 weeks treatment under the Re-treatment (whole body) restriction for severe chronic plaque psoriasis, **AND**

- Patient must have demonstrated an adequate response to treatment, **AND**
- Patient must not receive more than 8 weeks of treatment with etanercept under this restriction.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, when compared with the pre-etanercept treatment baseline value.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) the completed current Psoriasis Area and Severity Index (PASI) calculation sheet including the date of assessment of the patient's condition.

The same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of gaining approval for the remainder of 24 weeks treatment.

A PASI assessment of the patient's response to the initial course of treatment must be made after at least 12 weeks of treatment so that there is adequate time for a response to be demonstrated. This assessment, which will be used to determine eligibility for a further 8 weeks of treatment, must be submitted no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted within these timeframes, the patient will be deemed to have failed to respond to treatment with etanercept.

Note It is recommended that an application is posted to the Department of Human Services no later than 2 weeks prior to the patient completing their initial 16 week treatment course to ensure continuity of treatment for those patients who meet the eligibility criterion for a further 8 weeks of PBS-subsidised etanercept treatment.

Note In circumstances where it is not possible to submit a response assessment after 12 weeks of treatment, please call the Department of Human Services on 1800 700 270 to discuss.

Note The Department of Human Services will determine whether a response to treatment has been demonstrated, based on the baseline PASI assessment submitted with the first authority application for etanercept.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Re-treatment (Whole body)

TREATMENT OF PATIENTS UNDER 18 YEARS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing of etanercept under the Pharmaceutical Benefits Scheme (PBS) for patients under 18 years with severe chronic plaque psoriasis.

Applications for treatment of this condition will be limited to provide patients with a maximum of 24 weeks of therapy per course of treatment. A maximum of 16 weeks treatment with etanercept will be authorised for the primary application. The balance, a further 8 weeks treatment, will be authorised if the submitted Psoriasis Area and Severity Index (PASI) assessment demonstrates an adequate response to treatment. Where fewer than 3 repeats are requested at the time of the authority application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone.

Once a patient has failed to respond to treatment 2 times, they must have, at a minimum, a 12 month break. The length of a treatment break is measured from the date the most recent treatment was stopped to the date of the first application for initial treatment.

There are separate restrictions for treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Application for approval for initial treatment.

Applications for a course of initial treatment should be made for patients who have received no prior PBS-subsidised biological treatment and wish to commence such therapy.

(2) Applications for approval for re-treatment.

Applications for re-treatment with etanercept should be made in the following situations:

- (i) a patient who has received prior PBS-subsidised etanercept and experiences a disease flare, and wishes to start a second or subsequent treatment course with etanercept following a break of less than 12 months in PBS-subsidised therapy; or
- (ii) a patient who has received and failed to respond to prior PBS-subsidised etanercept and wishes to start a second or subsequent treatment course following a break of less than 12 months in PBS-subsidised therapy.

For psoriasis affecting the whole body:

Patients are eligible for re-treatment due to disease flare if there is a 50% or greater change in the patients PASI score or the patient has a current PASI score of greater than 15, compared to the most recent response assessment following cessation of the most recent 24 weeks of PBS-subsidised etanercept.

For psoriasis affecting the face, hand or foot:

Patients are eligible for re-treatment due to disease flare if:

- (i) all subscores are rated moderate to severe or 2 of the three subscores are rated severe to very severe; OR

(ii) the skin area affected is a 50% or greater change or the area affected is 30% or more of the face, palm of a hand or sole of a foot, compared to the most recent response assessment following cessation of the most recent 24 weeks of PBS-subsidised etanercept.

(3) Applications for approval for completion of a course

Applications for a further 8 weeks of treatment to allow for completion of 24 weeks of therapy should be submitted with a PASI assessment.

The PASI assessment must be conducted after at least 12 weeks of treatment.

This assessment must be submitted to Department of Human Services (the Department) within 1 month of the completion of 12 weeks of treatment. Where a response assessment is not undertaken and submitted to the Department within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological agent. In circumstances where it is not possible to submit a response assessment within these timeframes, please call the Department on 1800 700 270 to discuss.

(4) Baseline measurements to determine response.

The Department will determine whether a response to treatment has been demonstrated, based on the baseline PASI assessment submitted with the first authority application for etanercept. However, prescribers may provide new baseline measurements any time that an initial or re-treatment authority is submitted and subsequent response will be assessed according to this revised PASI score.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of gaining approval for the remainder of 24 weeks treatment.

(5) Re-commencement of treatment after a 12 month break in PBS-subsidised therapy.

A patient who wishes to start a second or subsequent treatment course with etanercept following a break in PBS-subsidised etanercept therapy of at least 12 months, must requalify for treatment under the initial treatment restriction. The most recent PASI assessment must be no more than 1 month old at the time of application.

Treatment criteria:

- Must be treated by a dermatologist.

Population criteria:

- Patient must be under 18 years of age.

Clinical criteria:

- The treatment must be as systemic monotherapy; OR
- The treatment must be in combination with methotrexate, **AND**
- Patient must have a documented history of severe chronic plaque psoriasis of the whole body, **AND**
- Patient must have received prior PBS-subsidised treatment with etanercept for this condition in the past 12 months, **AND**
- Patient must have demonstrated a response to etanercept and experienced a disease flare; OR
- Patient must not have failed more than once to achieve an adequate response with etanercept, **AND**
- Patient must not receive more than 16 weeks of treatment with etanercept under this restriction.

A patient is eligible for re-treatment due to disease flare if there is a 50% or greater change in the patient's PASI score or the patient has a current PASI score of greater than 15, compared to the most recent response assessment following cessation of the most recent 24 weeks of PBS-subsidised etanercept.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Severe Chronic Plaque Psoriasis in Patients Less Than 18 Years PBS Authority Application - Supporting Information which includes the following:
 - (i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and
 - (ii) details of prior etanercept treatment, including date ceased.

Where a patient has had a treatment break the length of the break is measured from the date the most recent treatment was stopped to the date of the application for further treatment.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment (Face, hand, foot)

TREATMENT OF PATIENTS UNDER 18 YEARS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing of etanercept under the Pharmaceutical Benefits Scheme (PBS) for patients under 18 years with severe chronic plaque psoriasis.

Applications for treatment of this condition will be limited to provide patients with a maximum of 24 weeks of therapy per course of treatment. A maximum of 16 weeks treatment with etanercept will be authorised for the primary application. The balance, a further 8 weeks treatment, will be authorised if the submitted Psoriasis Area and Severity Index (PASI) assessment demonstrates an adequate response to treatment. Where fewer than 3 repeats are requested at the time of the

authority application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone.

Once a patient has failed to respond to treatment 2 times, they must have, at a minimum, a 12 month break. The length of a treatment break is measured from the date the most recent treatment was stopped to the date of the first application for initial treatment.

There are separate restrictions for treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Application for approval for initial treatment.

Applications for a course of initial treatment should be made for patients who have received no prior PBS-subsidised biological treatment and wish to commence such therapy.

(2) Applications for approval for re-treatment.

Applications for re-treatment with etanercept should be made in the following situations:

(i) a patient who has received prior PBS-subsidised etanercept and experiences a disease flare, and wishes to start a second or subsequent treatment course with etanercept following a break of less than 12 months in PBS-subsidised therapy; or

(ii) a patient who has received and failed to respond to prior PBS-subsidised etanercept and wishes to start a second or subsequent treatment course following a break of less than 12 months in PBS-subsidised therapy.

For psoriasis affecting the whole body:

Patients are eligible for re-treatment due to disease flare if there is a 50% or greater change in the patients PASI score or the patient has a current PASI score of greater than 15, compared to the most recent response assessment following cessation of the most recent 24 weeks of PBS-subsidised etanercept.

For psoriasis affecting the face, hand or foot:

Patients are eligible for re-treatment due to disease flare if:

(i) all subscores are rated moderate to severe or 2 of the three subscores are rated severe to very severe; OR

(ii) the skin area affected is a 50% or greater change or the area affected is 30% or more of the face, palm of a hand or sole of a foot, compared to the most recent response assessment following cessation of the most recent 24 weeks of PBS-subsidised etanercept.

(3) Applications for approval for completion of a course

Applications for a further 8 weeks of treatment to allow for completion of 24 weeks of therapy should be submitted with a PASI assessment.

The PASI assessment must be conducted after at least 12 weeks of treatment.

This assessment must be submitted to Department of Human Services (the Department) within 1 month of the completion of 12 weeks of treatment. Where a response assessment is not undertaken and submitted to the Department within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological agent. In circumstances where it is not possible to submit a response assessment within these timeframes, please call the Department on 1800 700 270 to discuss.

(4) Baseline measurements to determine response.

The Department will determine whether a response to treatment has been demonstrated, based on the baseline PASI assessment submitted with the first authority application for etanercept. However, prescribers may provide new baseline measurements any time that an initial or re-treatment authority is submitted and subsequent response will be assessed according to this revised PASI score.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of gaining approval for the remainder of 24 weeks treatment.

(5) Re-commencement of treatment after a 12 month break in PBS-subsidised therapy.

A patient who wishes to start a second or subsequent treatment course with etanercept following a break in PBS-subsidised etanercept therapy of at least 12 months, must requalify for treatment under the initial treatment restriction. The most recent PASI assessment must be no more than 1 month old at the time of application.

Treatment criteria:

- Must be treated by a dermatologist.

Population criteria:

- Patient must be under 18 years of age and a parent or authorised guardian must have signed a patient acknowledgement.

Clinical criteria:

- The treatment must be as systemic monotherapy; OR
- The treatment must be in combination with methotrexate, **AND**
- Patient must have the plaque or plaques of the face, or palm of hand or sole of foot present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received any prior PBS-subsidised treatment with etanercept for this condition; OR
- Patient must not have received any PBS-subsidised treatment with etanercept for this condition for at least 12 months, **AND**

- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 3 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; and/or (ii) methotrexate at a dose of at least 10 mg or 10 mg per square metre weekly (whichever is lowest) for at least 6 weeks; and/or (iii) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks, **AND**
- Patient must not receive more than 16 weeks of treatment with etanercept under this restriction.

Where treatment with any of the above-mentioned drugs was contraindicated according to the relevant TGA-approved Product Information, or where phototherapy was contraindicated, details must be provided at the time of application.

Where intolerance to phototherapy, methotrexate and/or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

(a) Chronic plaque psoriasis classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:

(i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe, as assessed, preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior treatment; or

(ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed, preferably whilst still on treatment, but no longer than 1 month following cessation of the most recent prior treatment;

(b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 1 month following cessation of each course of treatment.

(c) The most recent PASI assessment must be no more than 1 month old at the time of application.

The authority application must be made in writing and must include:

(a) a completed authority prescription form; and

(b) a completed Severe Chronic Plaque Psoriasis in Patients Less Than 18 Years PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets, and face, hand, foot area diagrams including the dates of assessment of the patient's condition

(ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy]; and

(iii) the parent or authorised guardian signed patient and prescriber acknowledgements.

Where a patient has had a 12 month treatment break, the length of the break is measured from the date the most recent treatment was stopped to the date of the application to re-commence treatment.

Note Details of acceptable toxicities including severity, associated with phototherapy, methotrexate and acitretin, can be found on the Department of Human Services website at www.humanservices.gov.au

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment of Re-treatment (Face, hand, foot) - balance of first supply

TREATMENT OF PATIENTS UNDER 18 YEARS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing of etanercept under the Pharmaceutical Benefits Scheme (PBS) for patients under 18 years with severe chronic plaque psoriasis.

Applications for treatment of this condition will be limited to provide patients with a maximum of 24 weeks of therapy per course of treatment. A maximum of 16 weeks treatment with etanercept will be authorised for the primary application. The balance, a further 8 weeks treatment, will be authorised if the submitted Psoriasis Area and Severity Index (PASI) assessment demonstrates an adequate response to treatment. Where fewer than 3 repeats are requested at the time of the authority application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone.

Once a patient has failed to respond to treatment 2 times, they must have, at a minimum, a 12 month break. The length of a treatment break is measured from the date the most recent treatment was stopped to the date of the first application for initial treatment.

There are separate restrictions for treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Application for approval for initial treatment.

Applications for a course of initial treatment should be made for patients who have received no prior PBS-subsidised biological treatment and wish to commence such therapy.

(2) Applications for approval for re-treatment.

Applications for re-treatment with etanercept should be made in the following situations:

(i) a patient who has received prior PBS-subsidised etanercept and experiences a disease flare, and wishes to start a second or subsequent treatment course with etanercept following a break of less than 12 months in PBS-subsidised therapy; or

(ii) a patient who has received and failed to respond to prior PBS-subsidised etanercept and wishes to start a second or subsequent treatment course following a break of less than 12 months in PBS-subsidised therapy.

For psoriasis affecting the whole body:

Patients are eligible for re-treatment due to disease flare if there is a 50% or greater change in the patients PASI score or the patient has a current PASI score of greater than 15, compared to the most recent response assessment following cessation of the most recent 24 weeks of PBS-subsidised etanercept.

For psoriasis affecting the face, hand or foot:

Patients are eligible for re-treatment due to disease flare if:

- (i) all subscores are rated moderate to severe or 2 of the three subscores are rated severe to very severe; OR
- (ii) the skin area affected is a 50% or greater change or the area affected is 30% or more of the face, palm of a hand or sole of a foot, compared to the most recent response assessment following cessation of the most recent 24 weeks of PBS-subsidised etanercept.

(3) Applications for approval for completion of a course

Applications for a further 8 weeks of treatment to allow for completion of 24 weeks of therapy should be submitted with a PASI assessment.

The PASI assessment must be conducted after at least 12 weeks of treatment.

This assessment must be submitted to Department of Human Services (the Department) within 1 month of the completion of 12 weeks of treatment. Where a response assessment is not undertaken and submitted to the Department within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological agent. In circumstances where it is not possible to submit a response assessment within these timeframes, please call the Department on 1800 700 270 to discuss.

(4) Baseline measurements to determine response.

The Department will determine whether a response to treatment has been demonstrated, based on the baseline PASI assessment submitted with the first authority application for etanercept. However, prescribers may provide new baseline measurements any time that an initial or re-treatment authority is submitted and subsequent response will be assessed according to this revised PASI score.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of gaining approval for the remainder of 24 weeks treatment.

(5) Re-commencement of treatment after a 12 month break in PBS-subsidised therapy.

A patient who wishes to start a second or subsequent treatment course with etanercept following a break in PBS-subsidised etanercept therapy of at least 12 months, must requalify for treatment under the initial treatment restriction. The most recent PASI assessment must be no more than 1 month old at the time of application.

Treatment criteria:

- Must be treated by a dermatologist.

Clinical criteria:

- The treatment must be as systemic monotherapy; OR
- The treatment must be in combination with methotrexate, **AND**
- Patient must have received insufficient therapy under the Initial treatment (Face, hand, foot) restriction for severe chronic plaque psoriasis to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy under the Re-treatment (Face, hand, foot) restriction for severe chronic plaque psoriasis to complete 16 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Note Authority approval for sufficient therapy to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Written application for authority approval for sufficient therapy to complete a maximum of 16 weeks of treatment should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment or Re-treatment (Face, hand, foot) - completion of course

TREATMENT OF PATIENTS UNDER 18 YEARS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing of etanercept under the Pharmaceutical Benefits Scheme (PBS) for patients under 18 years with severe chronic plaque psoriasis.

Applications for treatment of this condition will be limited to provide patients with a maximum of 24 weeks of therapy per course of treatment. A maximum of 16 weeks treatment with etanercept will be authorised for the primary application. The balance, a further 8 weeks treatment, will be authorised if the submitted Psoriasis Area and Severity Index (PASI) assessment demonstrates an adequate response to treatment. Where fewer than 3 repeats are requested at the time of the authority application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone.

Once a patient has failed to respond to treatment 2 times, they must have, at a minimum, a 12 month break. The length of a treatment break is measured from the date the most recent treatment was stopped to the date of the first application for initial treatment.

There are separate restrictions for treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Application for approval for initial treatment.

Applications for a course of initial treatment should be made for patients who have received no prior PBS-subsidised biological treatment and wish to commence such therapy.

(2) Applications for approval for re-treatment.

Applications for re-treatment with etanercept should be made in the following situations:

(i) a patient who has received prior PBS-subsidised etanercept and experiences a disease flare, and wishes to start a second or subsequent treatment course with etanercept following a break of less than 12 months in PBS-subsidised therapy; or

(ii) a patient who has received and failed to respond to prior PBS-subsidised etanercept and wishes to start a second or subsequent treatment course following a break of less than 12 months in PBS-subsidised therapy.

For psoriasis affecting the whole body:

Patients are eligible for re-treatment due to disease flare if there is a 50% or greater change in the patients PASI score or the patient has a current PASI score of greater than 15, compared to the most recent response assessment following cessation of the most recent 24 weeks of PBS-subsidised etanercept.

For psoriasis affecting the face, hand or foot:

Patients are eligible for re-treatment due to disease flare if:

(i) all subscores are rated moderate to severe or 2 of the three subscores are rated severe to very severe; OR

(ii) the skin area affected is a 50% or greater change or the area affected is 30% or more of the face, palm of a hand or sole of a foot, compared to the most recent response assessment following cessation of the most recent 24 weeks of PBS-subsidised etanercept.

(3) Applications for approval for completion of a course

Applications for a further 8 weeks of treatment to allow for completion of 24 weeks of therapy should be submitted with a PASI assessment.

The PASI assessment must be conducted after at least 12 weeks of treatment.

This assessment must be submitted to Department of Human Services (the Department) within 1 month of the completion of 12 weeks of treatment. Where a response assessment is not undertaken and submitted to the Department within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological agent. In circumstances where it is not possible to submit a response assessment within these timeframes, please call the Department on 1800 700 270 to discuss.

(4) Baseline measurements to determine response.

The Department will determine whether a response to treatment has been demonstrated, based on the baseline PASI assessment submitted with the first authority application for etanercept. However, prescribers may provide new baseline measurements any time that an initial or re-treatment authority is submitted and subsequent response will be assessed according to this revised PASI score.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of gaining approval for the remainder of 24 weeks treatment.

(5) Re-commencement of treatment after a 12 month break in PBS-subsidised therapy.

A patient who wishes to start a second or subsequent treatment course with etanercept following a break in PBS-subsidised etanercept therapy of at least 12 months, must requalify for treatment under the initial treatment restriction. The most recent PASI assessment must be no more than 1 month old at the time of application.

Treatment criteria:

- Must be treated by a dermatologist.

Clinical criteria:

- The treatment must be as systemic monotherapy; OR
- The treatment must be in combination with methotrexate, **AND**
- Patient must have received 16 weeks treatment under the Initial treatment (Face, hand, foot) restriction for severe chronic plaque psoriasis; OR
- Patient must have received 16 weeks treatment under the Re-treatment (Face, hand, foot) restriction for severe chronic plaque psoriasis, **AND**
- Patient must have demonstrated an adequate response to treatment, **AND**
- Patient must not receive more than 8 weeks of treatment with etanercept under this restriction.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the pre-biological treatment baseline values; or
- (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the pre-biological treatment baseline value.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) the completed current Psoriasis Area and Severity Index (PASI) calculation sheet including the date of assessment of the patient's condition.

The same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of gaining approval for the remainder of 24 weeks treatment.

A PASI assessment of the patient's response to the initial course of treatment must be made after at least 12 weeks of treatment so that there is adequate time for a response to be demonstrated. This assessment, which will be used to determine eligibility for a further 8 weeks of treatment, must be submitted no later than 1 month from the date of completion of this initial course of treatment. Where a response assessment is not undertaken and submitted within these timeframes, the patient will be deemed to have failed to respond to treatment with etanercept.

Note It is recommended that an application is posted to the Department of Human Services no later than 2 weeks prior to the patient completing their initial 16 week treatment course to ensure continuity of treatment for those patients who meet the eligibility criterion for a further 8 weeks of PBS-subsidised etanercept treatment.

- Note** In circumstances where it is not possible to submit a response assessment after 12 weeks of treatment, please call the Department of Human Services on 1800 700 270 to discuss.
- Note** The Department of Human Services will determine whether a response to treatment has been demonstrated, based on the baseline PASI assessment submitted with the first authority application for etanercept.
- Note** Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Re-treatment (Face, hand, foot)

TREATMENT OF PATIENTS UNDER 18 YEARS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing of etanercept under the Pharmaceutical Benefits Scheme (PBS) for patients under 18 years with severe chronic plaque psoriasis.

Applications for treatment of this condition will be limited to provide patients with a maximum of 24 weeks of therapy per course of treatment. A maximum of 16 weeks treatment with etanercept will be authorised for the primary application. The balance, a further 8 weeks treatment, will be authorised if the submitted Psoriasis Area and Severity Index (PASI) assessment demonstrates an adequate response to treatment. Where fewer than 3 repeats are requested at the time of the authority application, authority approvals for sufficient repeats to complete a maximum of 16 weeks of treatment may be requested by telephone.

Once a patient has failed to respond to treatment 2 times, they must have, at a minimum, a 12 month break. The length of a treatment break is measured from the date the most recent treatment was stopped to the date of the first application for initial treatment.

There are separate restrictions for treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Application for approval for initial treatment.

Applications for a course of initial treatment should be made for patients who have received no prior PBS-subsidised biological treatment and wish to commence such therapy.

(2) Applications for approval for re-treatment.

Applications for re-treatment with etanercept should be made in the following situations:

(i) a patient who has received prior PBS-subsidised etanercept and experiences a disease flare, and wishes to start a second or subsequent treatment course with etanercept following a break of less than 12 months in PBS-subsidised therapy; or

(ii) a patient who has received and failed to respond to prior PBS-subsidised etanercept and wishes to start a second or subsequent treatment course following a break of less than 12 months in PBS-subsidised therapy.

For psoriasis affecting the whole body:

Patients are eligible for re-treatment due to disease flare if there is a 50% or greater change in the patients PASI score or the patient has a current PASI score of greater than 15, compared to the most recent response assessment following cessation of the most recent 24 weeks of PBS-subsidised etanercept.

For psoriasis affecting the face, hand or foot:

Patients are eligible for re-treatment due to disease flare if:

(i) all subscores are rated moderate to severe or 2 of the three subscores are rated severe to very severe; OR

(ii) the skin area affected is a 50% or greater change or the area affected is 30% or more of the face, palm of a hand or sole of a foot, compared to the most recent response assessment following cessation of the most recent 24 weeks of PBS-subsidised etanercept.

(3) Applications for approval for completion of a course

Applications for a further 8 weeks of treatment to allow for completion of 24 weeks of therapy should be submitted with a PASI assessment.

The PASI assessment must be conducted after at least 12 weeks of treatment.

This assessment must be submitted to Department of Human Services (the Department) within 1 month of the completion of 12 weeks of treatment. Where a response assessment is not undertaken and submitted to the Department within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological agent. In circumstances where it is not possible to submit a response assessment within these timeframes, please call the Department on 1800 700 270 to discuss.

(4) Baseline measurements to determine response.

The Department will determine whether a response to treatment has been demonstrated, based on the baseline PASI assessment submitted with the first authority application for etanercept. However, prescribers may provide new baseline measurements any time that an initial or re-treatment authority is submitted and subsequent response will be assessed according to this revised PASI score.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of gaining approval for the remainder of 24 weeks treatment.

(5) Re-commencement of treatment after a 12 month break in PBS-subsidised therapy.

A patient who wishes to start a second or subsequent treatment course with etanercept following a break in PBS-subsidised etanercept therapy of at least 12 months, must requalify for treatment under the initial treatment restriction. The most recent PASI assessment must be no more than 1 month old at the time of application.

Treatment criteria:

- Must be treated by a dermatologist.

Population criteria:

- Patient must be under 18 years of age.

Clinical criteria:

- The treatment must be as systemic monotherapy; OR
- The treatment must be in combination with methotrexate, **AND**
- Patient must have a documented history of severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, **AND**
- Patient must have received prior PBS-subsidised treatment with etanercept for this condition in the past 12 months, **AND**
- Patient must have demonstrated a response to etanercept and experienced a disease flare; OR
- Patient must not have failed more than once to achieve an adequate response with etanercept, **AND**
- Patient must not receive more than 16 weeks of treatment with etanercept under this restriction.

A patient is eligible for re-treatment due to disease flare if:

- (i) all subscores are rated moderate to severe or 2 of the 3 subscores are rated severe to very severe; or
- (ii) the skin area affected is a 50% or greater change or the area affected is 30% or more of the face, palm of a hand or sole of a foot, compared to the most recent response assessment following cessation of the most recent 24 weeks of PBS-subsidised etanercept.

The authority application must be made in writing and must include :

- (a) a completed authority prescription form; and
- (b) a completed Severe Chronic Plaque Psoriasis in Patients Less Than 18 Years PBS Authority Application - Supporting Information which includes the following:
 - (i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition; and
 - (ii) details of prior etanercept treatment, including date ceased.

Where a patient has had a treatment break the length of the break is measured from the date the most recent treatment was stopped to the date of the application for further treatment.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

etanercept 50 mg/mL injection, 4 x 1 mL pen devices

1964J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	1050.08	41.30	Enbrel [PF]

etanercept 25 mg injection [4 vials] (&) inert substance diluent [4 x 1 mL syringes], 1 pack

1954W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	3	..	*1050.10	41.30	Enbrel [PF]

etanercept 50 mg/mL injection, 4 x 1 mL syringes

1963H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	1050.08	41.30	Enbrel [PF]

▪ **GOLIMUMAB**

Note TREATMENT OF ADULT PATIENTS WITH MODERATE TO SEVERE ULCERATIVE COLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, golimumab, infliximab and vedolizumab for adult patients with ulcerative colitis. Patients are eligible for PBS-subsidised treatment with either adalimumab, golimumab, infliximab or vedolizumab at any one time.

Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, golimumab, infliximab and vedolizumab only.

From 1 June 2018, under the PBS, all adult patients will be able to commence a treatment cycle where they may trial PBS-subsidised biological medicine without having to experience a disease flare when swapping to one of the alternate agents. Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a PBS subsidised biological medicine while they continue to show a response to therapy.

A patient who received PBS-subsidised biological medicine treatment prior to 1 June 2018 is considered to start their first cycle as of 1 June 2018. Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the

last approval for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a PBS-subsidised biological medicine under the new treatment cycle.

Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under the Initial 2 treatment restriction.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle under the Initial 3 treatment restrictions.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment after 1 June 2018.

(1) Initial treatment.

Applications for initial treatment should be made where:

- (i) an adult patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy (Initial 1 - New patient); or
- (ii) an adult patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an alternate medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or
- (iii) an adult patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same agent (Initial 2 - Change or Recommencement of treatment after a break in biological medicine therapy of less than 5 years); or
- (iv) an adult patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - recommencement of treatment after a break in biological medicine of more than 5 years).

Treatment authorisations under Initial 1, Initial 2 and Initial 3 will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, 14 weeks of therapy for golimumab, infliximab and vedolizumab.

A patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of treatment for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for golimumab, infliximab and vedolizumab. For the second and subsequent courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine treatment is approved, a patient may swap to an alternate biological medicine treatment within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Mayo clinic score or partial Mayo clinic score), or the prior corticosteroid therapy and immunosuppressive therapy.

A patient may trial an alternate biological medicine treatment at any time, regardless of whether they are receiving therapy (initial or continuing) at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug once within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(4) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological medicine therapy of at least 5 years, must requalify under Initial 3 treatment restriction and meet the relevant criteria with respect to the scores of disease severity.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Initial 1 (new patient) or Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years) - balance of supply

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 14 weeks of treatment (weeks 0, 2, 6 and 10); OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 14 weeks of treatment (weeks 0, 2, 6 and 10); OR

- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 14 weeks of treatment (weeks 0, 2, 6 and 10), **AND**
- The treatment must provide no more than the balance of up to 14 weeks therapy available under Initial 1, 2 or 3 treatment.

golimumab 100 mg/mL injection, 1 mL pen device

11502J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	1122.62	41.30	Simponi [JC]

■ GOLIMUMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for adults with severe active rheumatoid arthritis. Where the term biological medicine appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alfa antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab), the T-cell co-stimulation modulator (abatacept) and the Janus kinase (JAK) inhibitors (baricitinib, tofacitinib, upadacitinib).

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised biological medicine therapy may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy,

- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once, and - once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised biological medicines for the treatment of rheumatoid arthritis.

A serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Stevens Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered a treatment failure.

A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction.

A patient who has failed fewer than 5 biological medicines and who has a break in treatment of less than 24 months may commence a further course of treatment with a biological medicine under Initial 2 treatment restriction. A patient who has failed fewer than 5 biological medicines and who has had a break in therapy of longer than 24 months may commence a further course of treatment with a biological medicine under the Initial 3 treatment restriction. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine treatment is stopped to the date of the new application for treatment with a biological medicine.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 April 2019.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised biological medicine treatment and wishes to commence such therapy, excluding rituximab (Initial 1 - new patient); or

(ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to re-commence treatment with a specific biological medicine following a break of less than 24 months in PBS-subsidised therapy with that agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months).

(iv) a patient wishes to re-commence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 24 months (Initial 3 - re-commencement of treatment after a break in biological medicine of more than 24 months) Initial applications for a new patient (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6-month intensive DMARD trial, but prior to ceasing DMARD therapy.

Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, baricitinib, etanercept, golimumab, tocilizumab, tofacitinib and upadacitinib, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

Rituximab patients should be assessed following a minimum of 12 weeks after the first infusion, and the assessment should be submitted to Services Australia within 4 weeks to ensure continuity of treatment for those who meet the continuing restriction for PBS subsidised treatment with this drug for this condition.

Where a response assessment is not provided the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment. For second and subsequent courses of PBS-subsidised biological medicine (excluding rituximab) treatment, it is recommended that a patient be reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to Services Australia within sufficient time to allow processing. Abatacept patients:

A patient is eligible to receive one I.V. loading dose when commencing treatment with the subcutaneous formulation. Two prescriptions are required, the first prescription for the I.V. loading dose for sufficient vials for one dose based on the patient's weight with no repeats. The second prescription for the subcutaneous formulation, with a maximum quantity of 4 and up to 3 repeats, must be submitted with the initial application.

Rituximab patients:

A further application may be submitted to Services Australia within sufficient time to allow processing. New baselines may

be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

Assessment of the patient's response to treatment must be conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the cessation of the most recent course of biological medicine therapy.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction. Where a response assessment is not conducted within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

(2) Swapping therapy

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non- biological medicine therapy requirements except if the patient has had a break in therapy of more than 24 months who would need to requalify with respect to the indices of disease severity. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each biological medicine trialled.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent, unless they have experienced a serious adverse reaction of a severity necessitating permanent treatment withdrawal.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug.

Abatacept:

A patient swapping from I.V. abatacept to subcutaneous abatacept will not be eligible for an I.V. loading dose when commencing treatment with the subcutaneous formulation.

Rituximab:

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF-alfa antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they be assessed for response to every course of treatment, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological medicine during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological medicine therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate biological medicine may do so without having to have any treatment-free period.

(3) Baseline measurements to determine response.

Determination of whether a response to treatment has been demonstrated must be based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and the eligibility for continuing treatment must be assessed according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response must be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be determined on the total number of major joints.

Applications under the Initial 1 treatment restriction for a new patient must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. The results must be no more than 4 weeks old at the time of application.

Applications under the Initial 3 treatment restriction for re-commencement of treatment after a break in biological medicine of more than 24 months must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note Special Pricing Arrangements apply.

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction, **AND**

- The treatment must be given concomitantly with methotrexate at a dose of at least 7.5 mg weekly.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient has either failed or ceased to respond to a PBS-subsidised biological medicine for this condition 5 times, they will not be eligible to receive further PBS-subsidised treatment with a biological medicine for this condition.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Continuing Treatment - balance of supply

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restriction.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

golimumab 50 mg/0.5 mL injection, 0.5 mL pen device

11375Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1160.52	41.30	Simponi [JC]

golimumab 50 mg/0.5 mL injection, 0.5 mL syringe

3428K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1160.52	41.30	Simponi [JC]

■ GOLIMUMAB**Note TREATMENT OF ADULT PATIENTS WITH ANKYLOSING SPONDYLITIS**

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab for adult patients with ankylosing spondylitis.

Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 7 biological medicines at any 1 time.

Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy.

A patient who has been receiving PBS-subsidised adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab treatment prior to 1 December 2020 is considered to start their first cycle as of 1 December 2020.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last prescription for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised biological medicine treatment with adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised biological medicine treatment in this treatment cycle and wishes to commence such therapy (Initial 1 - New patient)

(ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years); or

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

A patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy.

(b) Continuing treatment.

For the first continuing treatment course of PBS-subsidised biological medicine, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1, Initial 2 or Initial 3 treatment restrictions. For second and subsequent continuing courses of PBS-subsidised biological medicine it is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply. Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be conducted no later than 4 weeks from the cessation of that treatment course. Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI), or the prior NSAID therapy and exercise program requirements.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(3) Baseline measurements to determine response.

Services Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the BASDAI, ESR and/or CRP submitted with the first authority application for a biological medicine. For a new patient, the BASDAI used to determine the baseline must be measured while the patient is receiving NSAID therapy and completing their exercise program.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response.

Prescribers may provide new baseline measurements any time an 'Initial treatment' authority application is submitted within a treatment cycle and the eligibility for continuing treatment must be assessed according to these revised baseline measurements.

(4) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological

medicine therapy of at least 5 years, must qualify under the Initial 3 treatment restriction. The same clinical criteria and indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI) as for the Initial 1 (New patient) restriction will need to be met, but a re-trial of NSAID therapy and exercise therapy is not required.

Note Special Pricing Arrangements apply.

Authority required

Ankylosing spondylitis

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

The authority application must be made in writing and must include:

- a completed authority prescription form; and
- a completed Ankylosing Spondylitis PBS Authority Application - Supporting Information Form.

An adequate response is defined as an improvement from baseline of at least 2 of the BASDAI and 1 of the following:

- an ESR measurement no greater than 25 mm per hour; or
- a CRP measurement no greater than 10 mg per L; or
- an ESR or CRP measurement reduced by at least 20% from baseline.

Where only 1 acute phase reactant measurement is supplied in the first application for PBS-subsidised treatment, that same marker must be measured and supplied in all subsequent continuing treatment applications.

All measurements provided must be no more than 1 month old at the time of application.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Ankylosing spondylitis

Treatment Phase: Continuing treatment - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the Continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restriction.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

golimumab 50 mg/0.5 mL injection, 0.5 mL pen device

11376R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1160.52	41.30	Simponi [JC]

golimumab 50 mg/0.5 mL injection, 0.5 mL syringe

3436W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1160.52	41.30	Simponi [JC]

■ GOLIMUMAB**Note TREATMENT OF ADULT PATIENTS WITH NON-RADIOGRAPHIC AXIAL SPONDYLOARTHRITIS**

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of secukinumab, certolizumab pegol and golimumab for adult patients with non-radiographic axial spondyloarthritis.

Where the term 'biological medicine' appears in notes and restrictions, it refers to secukinumab, certolizumab pegol and golimumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 3 biological medicines at any one time.

Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy.

A serious adverse reaction requiring permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once.

Once a patient has either failed or ceased to respond to treatment 3 times (once with any biological medicine) within the same treatment cycle, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last prescription for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle.

A patient who has failed 3 trials or fewer of biological medicine therapy in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment with secukinumab, certolizumab pegol and golimumab

(1) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised biological medicine treatment in this treatment cycle and wishes to commence such therapy (Initial 1 - New patient)

(ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years); or

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

With the exception of grandfathered patients, a patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy.

(2) Continuing treatment.

For continuing courses of PBS-subsidised biological medicine it is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that biological medicine provided they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment in courses of up to 24 weeks provided they continue to sustain an adequate response.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be conducted no later than 4 weeks from the cessation of that treatment course. Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction requiring permanent treatment withdrawal.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. the C-reactive protein (CRP) level and the BASDAI score), or the prior NSAID therapy and exercise program requirements.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction requiring permanent treatment withdrawal.

(4) Baseline measurements to determine response.

A response to treatment is based on the baseline BASDAI score and CRP level documented in the patient's medical records. For a new patient, the BASDAI score used to determine baseline disease severity must be measured while the patient is receiving NSAID therapy and completing their exercise program.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications.

Prescribers may provide new baseline measurements any time that an 'Initial treatment' authority application is submitted and the eligibility for continuing treatment must be assessed according to these revised baseline measurements.

(5) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must qualify under the Initial 3 treatment restriction. The same clinical criteria and indices of disease severity (i.e. the C-reactive protein (CRP) levels and the BASDAI) as for the Initial 1 (New patient) restriction will need to be met, but a re-trial of NSAID therapy and exercise therapy is not required.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Non-radiographic axial spondyloarthritis

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug for this condition, **AND**
- The treatment must not exceed a maximum of 24 weeks with this drug per authorised course under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
 - Must be treated by a clinical immunologist with expertise in the management of non-radiographic axial spondyloarthritis.
- An adequate response to therapy with this biological medicine is defined as a reduction from baseline in the Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) score by 2 or more units (on a scale of 0-10) and 1 of the following:
- (a) a CRP measurement no greater than 10 mg per L; or
 - (b) a CRP measurement reduced by at least 20% from baseline.

If the requirement to demonstrate an elevated CRP level could not be met under an initial treatment restriction, a reduction in the BASDAI score from baseline will suffice for the purposes of administering this continuing treatment restriction.

The patient remains eligible to receive continuing treatment with the same biological medicine in courses of up to 24 weeks providing they continue to sustain an adequate response. It is recommended that a patient be reviewed in the month prior to completing their current course of treatment.

Authority required

Non-radiographic axial spondyloarthritis

Treatment Phase: Continuing treatment - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the Continuing treatment restriction to complete 24 weeks of treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of non-radiographic axial spondyloarthritis.

golimumab 50 mg/0.5 mL injection, 0.5 mL pen device

11521J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1160.52	41.30	Simponi [JC]

golimumab 50 mg/0.5 mL injection, 0.5 mL syringe

11516D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1160.52	41.30	Simponi [JC]

■ GOLIMUMAB

Note TREATMENT OF ADULT PATIENTS WITH MODERATE TO SEVERE ULCERATIVE COLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, golimumab, infliximab and vedolizumab for adult patients with ulcerative colitis. Patients are eligible for PBS-subsidised treatment with either adalimumab, golimumab, infliximab or vedolizumab at any one time.

Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, golimumab, infliximab and vedolizumab only.

From 1 June 2018, under the PBS, all adult patients will be able to commence a treatment cycle where they may trial PBS-subsidised biological medicine without having to experience a disease flare when swapping to one of the alternate agents. Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a PBS subsidised biological medicine while they continue to show a response to therapy.

A patient who received PBS-subsidised biological medicine treatment prior to 1 June 2018 is considered to start their first cycle as of 1 June 2018. Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a PBS-subsidised biological medicine under the new treatment cycle.

Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under the Initial 2 treatment restriction.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle under the Initial 3 treatment restrictions.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment after 1 June 2018.

(1) Initial treatment.

Applications for initial treatment should be made where:

- (i) an adult patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy (Initial 1 - New patient); or
- (ii) an adult patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an alternate medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or
- (iii) an adult patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same agent (Initial 2 - Change or Recommencement of treatment after a break in biological medicine therapy of less than 5 years); or
- (iv) an adult patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - recommencement of treatment after a break in biological medicine of more than 5 years).

Treatment authorisations under Initial 1, Initial 2 and Initial 3 will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, 14 weeks of therapy for golimumab, infliximab and vedolizumab.

A patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of treatment for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for golimumab, infliximab and vedolizumab. For the second and subsequent courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine treatment is approved, a patient may swap to an alternate biological medicine treatment within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Mayo clinic score or partial Mayo clinic score), or the prior corticosteroid therapy and immunosuppressive therapy.

A patient may trial an alternate biological medicine treatment at any time, regardless of whether they are receiving therapy (initial or continuing) at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug once within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(4) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological medicine therapy of at least 5 years, must requalify under Initial 3 treatment restriction and meet the relevant criteria with respect to the scores of disease severity.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have failed to achieve an adequate response to a 5-aminosalicylate oral preparation in a standard dose for induction of remission for 3 or more consecutive months or have intolerance necessitating permanent treatment withdrawal, **AND**
- Patient must have failed to achieve an adequate response to azathioprine at a dose of at least 2 mg per kg daily for 3 or more consecutive months or have intolerance necessitating permanent treatment withdrawal; OR
- Patient must have failed to achieve an adequate response to 6-mercaptopurine at a dose of at least 1 mg per kg daily for 3 or more consecutive months or have intolerance necessitating permanent treatment withdrawal; OR
- Patient must have failed to achieve an adequate response to a tapered course of oral steroids, starting at a dose of at least 40 mg prednisolone (or equivalent), over a 6 week period or have intolerance necessitating permanent treatment withdrawal, and followed by a failure to achieve an adequate response to 3 or more consecutive months of treatment of an appropriately dosed thiopurine agent, **AND**
- Patient must have a Mayo clinic score greater than or equal to 6; OR
- Patient must have a partial Mayo clinic score greater than or equal to 6, provided the rectal bleeding and stool frequency subscores are both greater than or equal to 2 (endoscopy subscore is not required for a partial Mayo clinic score).

Population criteria:

- Patient must be aged 18 years or older.

Application for authorisation of initial treatment must be in writing and must include:

(a) a completed authority prescription form; and

(b) a completed Ulcerative Colitis PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed current Mayo clinic or partial Mayo clinic calculation sheet including the date of assessment of the patient's condition; and

(ii) details of prior systemic drug therapy [dosage, date of commencement and duration of therapy].

All tests and assessments should be performed preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior conventional treatment.

The most recent Mayo clinic or partial Mayo clinic score must be no more than 4 weeks old at the time of application.

A partial Mayo clinic assessment of the patient's response to this initial course of treatment must be following a minimum of 12 weeks of treatment for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for golimumab, infliximab and vedolizumab so that there is adequate time for a response to be demonstrated.

A maximum of 14 weeks of treatment with this drug will be approved under this criterion. A loading dose of 200 mg at week 0 and a dose of 100 mg at weeks 2, 6 and 10.

If treatment with any of the above-mentioned drugs is contraindicated according to the relevant TGA-approved Product Information, details must be provided at the time of application.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Details of the accepted toxicities including severity can be found on the Department of Human Services website.

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Initial treatment - Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years)

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle.

Population criteria:

- Patient must be aged 18 years or older.

Application for authorisation must be made in writing and must include:

(a) a completed authority prescription form; and

(b) a completed Ulcerative Colitis PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed current Mayo clinic or partial Mayo clinic calculation sheet including the date of assessment of the patient's condition if relevant; and

(ii) the details of prior biological medicine treatment including the details of date and duration of treatment.

A maximum of 14 weeks of treatment with this drug will be approved under this criterion. A loading dose of 200 mg at week 0 and a dose of 100 mg at weeks 2, 6 and 10.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to change or recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3, or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for golimumab, infliximab and vedolizumab and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient who fails to demonstrate a response to treatment with this drug under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug in this treatment cycle. A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the initial 3 treatment restriction.

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Initial treatment - Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years)

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have had a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- Patient must have a Mayo clinic score greater than or equal to 6; OR
- Patient must have a partial Mayo clinic score greater than or equal to 6, provided the rectal bleeding and stool frequency subscores are both greater than or equal to 2 (endoscopy subscore is not required for a partial Mayo clinic score).

Population criteria:

- Patient must be aged 18 years or older.

Application for authorisation must be made in writing and must include:

(a) a completed authority prescription form; and

(b) a completed Ulcerative Colitis PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed current Mayo clinic or partial Mayo clinic calculation sheet including the date of assessment of the patient's condition; and

(ii) the details of prior biological medicine treatment including the details of date and duration of treatment.

A maximum of 14 weeks of treatment with this drug will be approved under this criterion. A loading dose of 200 mg at week 0 and a dose of 100 mg at weeks 2, 6 and 10.

All tests and assessments should be performed preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior conventional treatment.

The most recent Mayo clinic or partial Mayo clinic score must be no more than 4 weeks old at the time of application.

A partial Mayo clinic assessment of the patient's response to this initial course of treatment must be following a minimum of 12 weeks of treatment for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for golimumab, infliximab and vedolizumab so that there is adequate time for a response to be demonstrated.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Details of the accepted toxicities including severity can be found on the Department of Human Services website.

golimumab 100 mg/mL injection, 1 mL pen device

11382C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	..	..	*3294.45	41.30	Simponi [JC]

■ GOLIMUMAB
Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE PSORIATIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab for adult patients with severe active psoriatic arthritis. Therefore, where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient receiving PBS-subsidised treatment for psoriatic arthritis is able to commence a treatment cycle where they may trial biological medicines without having to experience a disease flare when swapping to the alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

A patient who received PBS-subsidised adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab or ustekinumab treatment prior to 1 May 2019 is considered to start their first cycle as of 1 May 2019.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence another cycle [further details are under '(5) Recommencement of treatment after a 5-year break in PBS-subsidised therapy' below].

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction. There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe active psoriatic arthritis.

(1) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an alternate medicine (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years) or An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, golimumab secukinumab and tofacitinib, 18 to 20 weeks of therapy for certolizumab pegol (depending upon the dosing regimen), 20 weeks of therapy for ixekizumab, 22 weeks of therapy for infliximab, and 28 weeks of therapy for ustekinumab. It is recommended that a patient be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of PBS-subsidised initial treatment following a minimum of 12 weeks of therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

Grandfather patients (ixekizumab only).

A patient who commenced treatment with ixekizumab for severe psoriatic arthritis prior to 1 March 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

Grandfather patients (tofacitinib only).

A patient who commenced treatment with Tofacitinib for severe psoriatic arthritis prior to 1 May 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will

only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment under the Initial 1 or Initial 2 treatment restrictions.

For the second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to re-qualify with respect to either the indices of disease severity (i.e. erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP) level, and active joint count) or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify under the Initial 3 treatment restriction with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application.

However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

Within a treatment cycle a patient may alternate between therapy with any biological medicine of their choice (1 at a time) providing:

- (i) they have not received PBS-subsidised treatment with that particular biological medicine previously; or
- (ii) they have demonstrated an adequate response to that particular biological medicine if they have previously trialled it on the PBS; and
- (iii) they have not previously failed to respond to treatment 3 times in this treatment cycle with that biological medicine.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(4) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the indices of disease severity submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial or change or commencement treatment application is submitted within a treatment cycle and these revised baseline measurements will be used to assess response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to determine response for all subsequent continuing treatments. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. 20 or more active joints), response will be determined according to a reduction in the total number of active joints.

(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment restriction according to the criteria of the relevant restriction and index of disease severity. The application must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Severe psoriatic arthritis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an erythrocyte sedimentation rate (ESR) no greater than 25 mm per hour or a C-reactive protein (CRP) level no greater than 15 mg per L or either marker reduced by at least 20% from baseline; and

either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following major active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to determine response for all subsequent continuing treatments.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Severe Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either Initial 1, Initial 2, or Initial 3 treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Continuing treatment - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restriction.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

golimumab 50 mg/0.5 mL injection, 0.5 mL pen device

11373N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1160.52	41.30	Simponi [JC]

golimumab 50 mg/0.5 mL injection, 0.5 mL syringe

3432P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1160.52	41.30	Simponi [JC]

■ GOLIMUMAB

Note TREATMENT OF ADULT PATIENTS WITH MODERATE TO SEVERE ULCERATIVE COLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, golimumab, infliximab and vedolizumab for adult patients with ulcerative colitis. Patients are eligible for PBS-subsidised treatment with either adalimumab, golimumab, infliximab or vedolizumab at any one time.

Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, golimumab, infliximab and vedolizumab only.

From 1 June 2018, under the PBS, all adult patients will be able to commence a treatment cycle where they may trial PBS-subsidised biological medicine without having to experience a disease flare when swapping to one of the alternate agents. Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a PBS subsidised biological medicine while they continue to show a response to therapy.

A patient who received PBS-subsidised biological medicine treatment prior to 1 June 2018 is considered to start their first cycle as of 1 June 2018. Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a PBS-subsidised biological medicine under the new treatment cycle.

Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under the Initial 2 treatment restriction.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle under the Initial 3 treatment restrictions.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment after 1 June 2018.

(1) Initial treatment.

Applications for initial treatment should be made where:

- (i) an adult patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy (Initial 1 - New patient); or
- (ii) an adult patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an alternate medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or
- (iii) an adult patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same agent (Initial 2 - Change or Recommencement of treatment after a break in biological medicine therapy of less than 5 years); or
- (iv) an adult patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - recommencement of treatment after a break in biological medicine of more than 5 years).

Treatment authorisations under Initial 1, Initial 2 and Initial 3 will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, 14 weeks of therapy for golimumab, infliximab and vedolizumab.

A patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of treatment for adalimumab and up to 12 weeks after the first dose (6 weeks following the third dose) for golimumab, infliximab and vedolizumab. For the second and subsequent courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine treatment is approved, a patient may swap to an alternate biological medicine treatment within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Mayo clinic score or partial Mayo clinic score), or the prior corticosteroid therapy and immunosuppressive therapy.

A patient may trial an alternate biological medicine treatment at any time, regardless of whether they are receiving therapy (initial or continuing) at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug once within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(4) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological medicine therapy of at least 5 years, must requalify under Initial 3 treatment restriction and meet the relevant criteria with respect to the scores of disease severity.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have demonstrated or sustained an adequate response to treatment by having a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1 while receiving treatment with this drug.

Population criteria:

- Patient must be aged 18 years or older.

Patients who have failed to maintain a partial Mayo clinic score less than or equal to 2, with no subscore greater than 1 with continuing treatment with this drug, will not be eligible to receive further PBS-subsidised treatment with this drug.

Patients are eligible to receive continuing treatment with this drug in courses of up to 24 weeks providing they continue to sustain a response.

At the time of the authority application, medical practitioners should request sufficient quantity for up to 24 weeks of treatment under this restriction.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Authority applications for continuing treatment may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Moderate to severe ulcerative colitis

Treatment Phase: Continuing treatment - balance of supply

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under this restriction.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

golimumab 100 mg/mL injection, 1 mL pen device

11381B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1122.62	41.30	Simponi [JC]

▪ **GOLIMUMAB**

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for adults with severe active rheumatoid arthritis. Where the term biological medicine appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alfa antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab), the T-cell co-stimulation modulator (abatacept) and the Janus kinase (JAK) inhibitors (baricitinib, tofacitinib, upadacitinib).

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised biological medicine therapy may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once, and
- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised biological medicines for the treatment of rheumatoid arthritis.

A serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Stevens Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered a treatment failure.

A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction.

A patient who has failed fewer than 5 biological medicines and who has a break in treatment of less than 24 months may commence a further course of treatment with a biological medicine under Initial 2 treatment restriction. A patient who has

failed fewer than 5 biological medicines and who has had a break in therapy of longer than 24 months may commence a further course of treatment with a biological medicine under the Initial 3 treatment restriction. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine treatment is stopped to the date of the new application for treatment with a biological medicine.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 April 2019.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised biological medicine treatment and wishes to commence such therapy, excluding rituximab (Initial 1 - new patient); or

(ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to re-commence treatment with a specific biological medicine following a break of less than 24 months in PBS-subsidised therapy with that agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months).

(iv) a patient wishes to re-commence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 24 months (Initial 3 - re-commencement of treatment after a break in biological medicine of more than 24 months) Initial applications for a new patient (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6-month intensive DMARD trial, but prior to ceasing DMARD therapy.

Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, baricitinib, etanercept, golimumab, tocilizumab, tofacitinib and upadacitinib, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

Rituximab patients should be assessed following a minimum of 12 weeks after the first infusion, and the assessment should be submitted to Services Australia within 4 weeks to ensure continuity of treatment for those who meet the continuing restriction for PBS subsidised treatment with this drug for this condition.

Where a response assessment is not provided the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment. For second and subsequent courses of PBS-subsidised biological medicine (excluding rituximab) treatment, it is recommended that a patient be reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to Services Australia within sufficient time to allow processing.

Abatacept patients:

A patient is eligible to receive one I.V. loading dose when commencing treatment with the subcutaneous formulation. Two prescriptions are required, the first prescription for the I.V. loading dose for sufficient vials for one dose based on the patient's weight with no repeats. The second prescription for the subcutaneous formulation, with a maximum quantity of 4 and up to 3 repeats, must be submitted with the initial application.

Rituximab patients:

A further application may be submitted to Services Australia within sufficient time to allow processing. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

Assessment of the patient's response to treatment must be conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the cessation of the most recent course of biological medicine therapy.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction. Where a response assessment is not conducted within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

(2) Swapping therapy

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non- biological medicine therapy requirements except if the patient has had a break in therapy of more than 24 months who would need to requalify with respect to the indices of disease severity. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each biological medicine trialled.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent, unless they have experienced a serious adverse reaction of a severity necessitating permanent treatment withdrawal.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug.

Abatacept:

A patient swapping from I.V. abatacept to subcutaneous abatacept will not be eligible for an I.V. loading dose when commencing treatment with the subcutaneous formulation.

Rituximab:

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF-alfa antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they be assessed for response to every course of treatment, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological medicine during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological medicine therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate biological medicine may do so without having to have any treatment-free period.

(3) Baseline measurements to determine response.

Determination of whether a response to treatment has been demonstrated must be based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and the eligibility for continuing treatment must be assessed according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response must be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be determined on the total number of major joints.

Applications under the Initial 1 treatment restriction for a new patient must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. The results must be no more than 4 weeks old at the time of application.

Applications under the Initial 3 treatment restriction for re-commencement of treatment after a break in biological medicine of more than 24 months must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note Special Pricing Arrangements apply.

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with disease modifying anti-rheumatic drugs (DMARDs) which must include at least 3 months continuous treatment with each of at least 2 DMARDs, one of which must be methotrexate at a dose of at least 20 mg weekly and one of which must be: (i) hydroxychloroquine at a dose of at least 200 mg daily; or (ii) leflunomide at a dose of at least 10 mg daily; or (iii) sulfasalazine at a dose of at least 2 g daily; OR
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs which, if methotrexate is contraindicated according to the Therapeutic Goods Administration (TGA)-approved Product Information or cannot be tolerated at a 20 mg weekly dose, must include at least 3 months continuous treatment with each of at least 2 of the following DMARDs: (i) hydroxychloroquine at a dose of at least 200 mg daily; and/or (ii) leflunomide at a dose of at least 10 mg daily; and/or (iii) sulfasalazine at a dose of at least 2 g daily; OR
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs which, if 3 or more of methotrexate, hydroxychloroquine, leflunomide and sulfasalazine are contraindicated according to the relevant TGA-approved Product Information or cannot be tolerated at the doses specified above, must include at least 3 months continuous treatment with each of at least 2 DMARDs, with one or more of the following DMARDs being used in place of the DMARDs which are contraindicated or not tolerated: (i) azathioprine at a dose of at least 1 mg/kg per day; and/or (ii) cyclosporin at a dose of at least 2 mg/kg/day; and/or (iii) sodium aurothiomalate at a dose of 50 mg weekly, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction, **AND**
- The treatment must be given concomitantly with methotrexate at a dose of at least 7.5 mg weekly.

Population criteria:

- Patient must be aged 18 years or older.

If methotrexate is contraindicated according to the TGA-approved product information or cannot be tolerated at a 20 mg weekly dose, the application must include details of the contraindication or intolerance including severity to methotrexate. The maximum tolerated dose of methotrexate must be documented in the application, if applicable.

The application must include details of the DMARDs trialled, their doses and duration of treatment, and all relevant contraindications and/or intolerances including severity.

The requirement to trial at least 2 DMARDs for periods of at least 3 months each can be met using single agents sequentially or by using one or more combinations of DMARDs.

If the requirement to trial 6 months of intensive DMARD therapy with at least 2 DMARDs cannot be met because of contraindications and/or intolerances of a severity necessitating permanent treatment withdrawal to all of the DMARDs specified above, details of the contraindication or intolerance including severity and dose for each DMARD must be provided in the authority application.

The following criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; **AND** either

- (a) a total active joint count of at least 20 active (swollen and tender) joints; or
- (b) at least 4 active joints from the following list of major joints:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).
 The joint count and ESR and/or CRP must be determined at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. All measures must be no more than one month old at the time of initial application.
 If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note The Department of Human Services website (www.humanservices.gov.au) has details of the toxicities, including severity, which will be accepted for the following purposes:

- (a) exempting a patient from the requirement to undertake a minimum 3 month trial of methotrexate at a 20 mg weekly dose;
- (b) substituting azathioprine, cyclosporin or sodium aurothiomalate for another DMARD as part of the 6 month intensive DMARD trial;
- (c) exempting a patient from the requirement for a 6 month trial of intensive DMARD therapy.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
 Complex Drugs
 Reply Paid 9826
 HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 2 (change or re-commencement of treatment after a break in biological medicine of less than 24 months)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must not have failed to respond to previous PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised biological medicine treatment for this condition 5 times, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction, **AND**
- The treatment must be given concomitantly with methotrexate at a dose of at least 7.5 mg weekly.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either of the Initial 1, Initial 2, Initial 3, or continuing treatment restrictions, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

A patient who has demonstrated a response to a course of rituximab must have a PBS-subsidised biological therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 3 (re-commencement of treatment after a break in biological medicine of more than 24 months)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 24 months or more from the most recent PBS-subsidised biological medicine for this condition, **AND**
- Patient must not have failed to respond to previous PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised biological medicine treatment for this condition 5 times, **AND**
- The condition must have an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; OR
- The condition must have a C-reactive protein (CRP) level greater than 15 mg per L, **AND**
- The condition must have either (a) a total active joint count of at least 20 active (swollen and tender) joints; or (b) at least 4 active major joints, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction, **AND**
- The treatment must be given concomitantly with methotrexate at a dose of at least 7.5 mg weekly.

Population criteria:

- Patient must be aged 18 years or older.

Major joints are defined as (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

All measures of joint count and ESR and/or CRP must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial 1 (new patient) or Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 24 months) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 24 months) - balance of supply

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 24 months) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 24 months) to complete 16 weeks of treatment, **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

golimumab 50 mg/0.5 mL injection, 0.5 mL pen device

11372M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	1160.52	41.30	Simponi [JC]

golimumab 50 mg/0.5 mL injection, 0.5 mL syringe

3426H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	1160.52	41.30	Simponi [JC]

■ GOLIMUMAB

Note TREATMENT OF ADULT PATIENTS WITH ANKYLOSING SPONDYLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab for adult patients with ankylosing spondylitis.

Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 7 biological medicines at any 1 time.

Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy.

A patient who has been receiving PBS-subsidised adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab treatment prior to 1 December 2020 is considered to start their first cycle as of 1 December 2020.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last prescription for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle.

A patient who has failed more than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised biological medicine treatment with adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised biological medicine treatment in this treatment cycle and wishes to commence such therapy (Initial 1 - New patient)

(ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years); or

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

A patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy.

(b) Continuing treatment.

For the first continuing treatment course of PBS-subsidised biological medicine, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1, Initial 2 or Initial 3 treatment restrictions. For second and subsequent continuing courses of PBS-subsidised biological medicine it is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply. Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be conducted no later than 4 weeks from the cessation of that treatment course. Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI), or the prior NSAID therapy and exercise program requirements.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(3) Baseline measurements to determine response.

Services Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the BASDAI, ESR and/or CRP submitted with the first authority application for a biological medicine. For a new patient, the BASDAI used to determine the baseline must be measured while the patient is receiving NSAID therapy and completing their exercise program.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response.

Prescribers may provide new baseline measurements any time an 'Initial treatment' authority application is submitted within a treatment cycle and the eligibility for continuing treatment must be assessed according to these revised baseline measurements.

(4) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must qualify under the Initial 3 treatment restriction. The same clinical criteria and indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI) as for the Initial 1 (New patient) restriction will need to be met, but a re-trial of NSAID therapy and exercise therapy is not required.

Note Special Pricing Arrangements apply.

Authority required

Ankylosing spondylitis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Clinical criteria:

- The condition must be radiographically (plain X-ray) confirmed Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**

- Patient must have at least 2 of the following: (i) low back pain and stiffness for 3 or more months that is relieved by exercise but not by rest; or (ii) limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by a score of at least 1 on each of the lumbar flexion and lumbar side flexion measurements of the Bath Ankylosing Spondylitis Metrology Index (BASMI); or (iii) limitation of chest expansion relative to normal values for age and gender,

AND

- Patient must have failed to achieve an adequate response following treatment with at least 2 non-steroidal anti-inflammatory drugs (NSAIDs), whilst completing an appropriate exercise program, for a total period of 3 months, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

The application must include details of the NSAIDs trialled, their doses and duration of treatment.

If the NSAID dose is less than the maximum recommended dose in the relevant TGA-approved Product Information, the application must include the reason a higher dose cannot be used.

If treatment with NSAIDs is contraindicated according to the relevant TGA-approved Product Information, the application must provide details of the contraindication.

If intolerance to NSAID treatment develops during the relevant period of use which is of a severity to necessitate permanent treatment withdrawal, the application must provide details of the nature and severity of this intolerance.

The following criteria indicate failure to achieve an adequate response and must be demonstrated at the time of the initial application:

- (a) a Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 4 on a 0-10 scale; AND
- (b) an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 10 mg per L.

The BASDAI must be determined at the completion of the 3 month NSAID and exercise trial, but prior to ceasing NSAID treatment. The BASDAI must be no more than 1 month old at the time of initial application.

Both ESR and CRP measures should be provided with the initial treatment application and both must be no more than 1 month old. If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reason this criterion cannot be satisfied.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Ankylosing Spondylitis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) a copy of the radiological report confirming Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis; and
 - (ii) a completed BASDAI Assessment Form; and
 - (iii) a completed Exercise Program Self Certification Form included in the supporting information form.

An assessment of a patient's response to an initial course of treatment must be conducted following a minimum of 12 weeks of therapy. An application for the continuing treatment must be accompanied with the assessment of response and submitted to the Department of Human Services no later than 4 weeks from the date of completion of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Details of the toxicities, including severity, which will be accepted for the purposes of administering this restriction can be found on the Department of Human Services website at www.humanservices.gov.au

Note For details on the appropriate minimum exercise program that will be accepted for the purposes of administering this restriction, please refer to the Department of Human Services website at www.humanservices.gov.au

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Ankylosing spondylitis

Treatment Phase: Initial treatment - Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**

- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Ankylosing Spondylitis PBS Authority Application - Supporting Information Form.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to change or recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

An adequate response is defined as an improvement from baseline of at least 2 of the BASDAI and 1 of the following:

- (a) an ESR measurement no greater than 25 mm per hour; or
- (b) a CRP measurement no greater than 10 mg per L; or
- (c) an ESR or CRP measurement reduced by at least 20% from baseline.

Where only 1 acute phase reactant measurement is supplied in the first application for PBS-subsidised treatment, that same marker must be measured and supplied in all subsequent continuing treatment applications.

All measurements provided must be no more than 1 month old at the time of application.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos. Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Ankylosing spondylitis

Treatment Phase: Initial treatment - Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must be radiographically (plain X-ray) confirmed Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis, **AND**
- Patient must have at least 2 of the following: (i) low back pain and stiffness for 3 or more months that is relieved by exercise but not by rest; or (ii) limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by a score of at least 1 on each of the lumbar flexion and lumbar side flexion measurements of the Bath Ankylosing Spondylitis Metrology Index (BASMI); or (iii) limitation of chest expansion relative to normal values for age and gender,

AND

- Patient must have a Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 4 on a 0-10 scale that is no more than 4 weeks old at the time of application, **AND**
- Patient must have an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour that is no more than 4 weeks old at the time of application; OR
- Patient must have a C-reactive protein (CRP) level greater than 10 mg per L that is no more than 4 weeks old at the time of application; OR
- Patient must have a clinical reason as to why demonstration of an elevated ESR or CRP cannot be met and the application must state the reason, **AND**

- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

The authority application must be made in writing and must include:

(a) a completed authority prescription form; and

(b) a completed Ankylosing Spondylitis PBS Authority Application - Supporting Information Form which includes the following:

(i) a copy of the radiological report confirming Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis; and

(ii) a completed BASDAI Assessment Form.

An assessment of a patient's response to an initial course of treatment must be conducted following a minimum of 12 weeks of therapy. An application for the continuing treatment must be accompanied with the assessment of response and submitted to the Department of Human Services no later than 4 weeks from the date of completion of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos. Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Ankylosing spondylitis

Treatment Phase: Initial treatment - Initial 1 (new patient), Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years) - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 16 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

golimumab 50 mg/0.5 mL injection, 0.5 mL pen device

11361Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	1160.52	41.30	Simponi [JC]

golimumab 50 mg/0.5 mL injection, 0.5 mL syringe

3434R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	1160.52	41.30	Simponi [JC]

■ GOLIMUMAB**Note TREATMENT OF ADULT PATIENTS WITH NON-RADIOGRAPHIC AXIAL SPONDYLOARTHRITIS**

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of secukinumab, certolizumab pegol and golimumab for adult patients with non-radiographic axial spondyloarthritis. Where the term 'biological medicine' appears in notes and restrictions, it refers to secukinumab, certolizumab pegol and golimumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 3 biological medicines at any one time.

Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy.

A serious adverse reaction requiring permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once.

Once a patient has either failed or ceased to respond to treatment 3 times (once with any biological medicine) within the same treatment cycle, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last prescription for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle.

A patient who has failed 3 trials or fewer of biological medicine therapy in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment with secukinumab, certolizumab pegol and golimumab

(1) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised biological medicine treatment in this treatment cycle and wishes to commence such therapy (Initial 1 - New patient)
 - (ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or
 - (iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years); or
 - (iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).
- With the exception of grandfathered patients, a patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy.

(2) Continuing treatment.

For continuing courses of PBS-subsidised biological medicine it is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that biological medicine provided they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment in courses of up to 24 weeks provided they continue to sustain an adequate response.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be conducted no later than 4 weeks from the cessation of that treatment course. Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction requiring permanent treatment withdrawal.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. the C-reactive protein (CRP) level and the BASDAI score), or the prior NSAID therapy and exercise program requirements.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction requiring permanent treatment withdrawal.

(4) Baseline measurements to determine response.

A response to treatment is based on the baseline BASDAI score and CRP level documented in the patient's medical records.

For a new patient, the BASDAI score used to determine baseline disease severity must be measured while the patient is receiving NSAID therapy and completing their exercise program.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications.

Prescribers may provide new baseline measurements any time that an 'Initial treatment' authority application is submitted and the eligibility for continuing treatment must be assessed according to these revised baseline measurements.

(5) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must qualify under the Initial 3 treatment restriction. The same clinical criteria and indices of disease severity (i.e. the C-reactive protein (CRP) levels and the BASDAI) as for the Initial 1 (New patient) restriction will need to be met, but a re-trial of NSAID therapy and exercise therapy is not required.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Non-radiographic axial spondyloarthritis

Treatment Phase: Initial treatment - Initial 1 (New patient)

Clinical criteria:

- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**

- Patient must have had chronic lower back pain and stiffness for 3 or more months that is relieved by exercise but not rest, **AND**
- Patient must have failed to achieve an adequate response following treatment with at least 2 non-steroidal anti-inflammatory drugs (NSAIDs), whilst completing an appropriate exercise program, for a total period of 3 months, **AND**
- Patient must have one or more of the following: (a) enthesitis (heel); (b) uveitis; (c) dactylitis; (d) psoriasis; (e) inflammatory bowel disease; or (f) positive for Human Leukocyte Antigen B27 (HLA-B27), **AND**
- The condition must not be radiographically evidenced on plain x-ray of Grade II bilateral sacroiliitis or Grade III or IV unilateral sacroiliitis, **AND**
- The condition must be non-radiographic axial spondyloarthritis, as defined by Assessment of Spondyloarthritis International Society (ASAS) criteria, **AND**
- The condition must be sacroiliitis with active inflammation and/or oedema on non-contrast Magnetic Resonance Imaging (MRI), **AND**
- The condition must have presence of Bone Marrow Oedema (BMO) depicted as a hyperintense signal on a Short Tau Inversion Recovery (STIR) image (or equivalent), **AND**
- The condition must have BMO depicted as a hypointense signal on a T1 weighted image (without gadolinium), **AND**
- The treatment must not exceed a maximum of 16 weeks with this drug under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of non-radiographic axial spondyloarthritis. The application must include details of the NSAIDs trialled, their doses and duration of treatment.

If the NSAID dose is less than the maximum recommended dose in the relevant TGA-approved Product Information, the application must include the reason a higher dose cannot be used.

If treatment with NSAIDs is contraindicated according to the relevant TGA-approved Product Information, the application must provide details of the contraindication.

If intolerance to NSAID treatment develops during the relevant period of use which is of a severity to necessitate permanent treatment withdrawal, the application must provide details of the nature and severity of this intolerance.

The following criteria indicate failure to achieve an adequate response to NSAIDs and must be demonstrated at the time of the initial application:

- (a) a Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) score of at least 4 on a 0-10 scale; and
- (b) C-reactive protein (CRP) level greater than 10 mg per L.

The baseline BASDAI score and CRP level must be determined at the completion of the 3-month NSAID and exercise trial, but prior to ceasing NSAID treatment. All measures must be no more than 4 weeks old at the time of initial application.

If the requirement to demonstrate an elevated CRP level could not be met, the reason must be stated in the application.

Treatment with prednisolone dosed at 7.5 mg or higher daily (or equivalent) or a parenteral steroid within the past month (intramuscular or intravenous methylprednisolone or equivalent) is an acceptable reason.

The assessment of the patient's response to the initial course of treatment must be conducted following a minimum of 12 weeks of treatment and no later than 4 weeks from the cessation of that treatment course. If the response assessment is not conducted within these timeframes, the patient will be deemed to have failed this course of treatment in this treatment cycle.

The authority application must include:

- (a) a completed authority prescription; and
- (b) a completed PBS authority application form relevant to the indication and treatment phase (the latest version located at the website mentioned in the administrative note).

The baseline BASDAI score and CRP level must also be documented in the patient's medical records.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Non-radiographic axial spondyloarthritis

Treatment Phase: Initial treatment - Initial 2 (Change or re-commencement of treatment after a break of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have failed, or ceased to respond to, PBS-subsidised treatment with biological medicines more than three times for this PBS-indication during the current treatment cycle, **AND**
- The treatment must not exceed a maximum of 16 weeks with this drug under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of non-radiographic axial spondyloarthritis.

Clinical criteria:

- Patient must not have failed PBS-subsidised therapy with this biological medicine for this PBS indication more than once in the current treatment cycle.

An application for Initial 2 treatment must indicate whether the patient has demonstrated an adequate response (an absence of treatment failure), failed or experienced an intolerance to the most recent supply of biological medicine treatment.

A new baseline Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) score and C-reactive protein (CRP) level may be provided at the time of this application.

An adequate response to therapy with this biological medicine is defined as a reduction from baseline in the Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) score by 2 or more units (on a scale of 0-10) and 1 of the following:

- (a) a CRP measurement no greater than 10 mg per L; or
- (b) a CRP measurement reduced by at least 20% from baseline.

The assessment of the patient's response to the most recent supply of biological medicine must be conducted following a minimum of 12 weeks of treatment.

BASDAI scores and CRP levels must be documented in the patient's medical records.

The assessment of the patient's response to the initial course of treatment must be conducted following a minimum of 12 weeks of treatment and no later than 4 weeks from the cessation of that treatment course. If the response assessment is not conducted within these timeframes, the patient will be deemed to have failed this course of treatment in this treatment cycle.

The following must be provided at the time of application and documented in the patient's medical records:

- (a) the BASDAI score; and
- (b) the C-reactive protein (CRP) level.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Non-radiographic axial spondyloarthritis

Treatment Phase: Initial treatment - Initial 3 (Recommendation of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have had chronic lower back pain and stiffness for 3 or more months that is relieved by exercise but not rest, **AND**
- Patient must have had a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- Patient must have one or more of the following: (a) enthesitis (heel); (b) uveitis; (c) dactylitis; (d) psoriasis; (e) inflammatory bowel disease; or (f) positive for Human Leukocyte Antigen B27 (HLA-B27), **AND**
- The condition must not be radiographically evidenced on plain x-ray of Grade II bilateral sacroiliitis or Grade III or IV unilateral sacroiliitis, **AND**
- The condition must be non-radiographic axial spondyloarthritis, as defined by Assessment of Spondyloarthritis International Society (ASAS) criteria, **AND**
- The condition must be sacroiliitis with active inflammation and/or oedema on non-contrast Magnetic Resonance Imaging (MRI), **AND**
- The condition must have presence of Bone Marrow Oedema (BMO) depicted as a hyperintense signal on a Short Tau Inversion Recovery (STIR) image (or equivalent), **AND**
- The condition must have BMO depicted as a hypointense signal on a T1 weighted image (without gadolinium), **AND**
- The treatment must not exceed a maximum of 16 weeks duration under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of non-radiographic axial spondyloarthritis.

The following must be provided at the time of application and documented in the patient's medical records:

- (a) a Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) score of at least 4 on a 0-10 scale; and
- (b) C-reactive protein (CRP) level greater than 10 mg per L.

The BASDAI score and CRP level must be no more than 4 weeks old at the time of this application.

If the requirement to demonstrate an elevated CRP level could not be met, the reason must be stated in the application.

Treatment with prednisolone dosed at 7.5 mg or higher daily (or equivalent) or a parenteral steroid within the past month (intramuscular or intravenous methylprednisolone or equivalent) is an acceptable reason.

The assessment of the patient's response to the initial course of treatment must be conducted following a minimum of 12 weeks of treatment and no later than 4 weeks from the cessation of that treatment course. If the response assessment is not conducted within these timeframes, the patient will be deemed to have failed this course of treatment in this treatment cycle.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Non-radiographic axial spondyloarthritis

Treatment Phase: Initial 1 (New patient), Initial 2 (Change or re-commencement of treatment after a break in biological medicine of less than 5 years) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years) - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 16 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of non-radiographic axial spondyloarthritis.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

golimumab 50 mg/0.5 mL injection, 0.5 mL pen device

11538G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	1160.52	41.30	Simponi [JC]

golimumab 50 mg/0.5 mL injection, 0.5 mL syringe

11560K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	1160.52	41.30	Simponi [JC]

▪ **GOLIMUMAB**

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE PSORIATIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab for adult patients with severe active psoriatic arthritis. Therefore, where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient receiving PBS-subsidised treatment for psoriatic arthritis is able to commence a treatment cycle where they may trial biological medicines without having to experience a disease flare when swapping to the alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

A patient who received PBS-subsidised adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab or ustekinumab treatment prior to 1 May 2019 is considered to start their first cycle as of 1 May 2019.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence another cycle [further details are under '(5) Recommencement of treatment after a 5-year break in PBS-subsidised therapy' below].

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction. There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe active psoriatic arthritis.

(1) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an alternate medicine (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of

more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years) or An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, golimumab secukinumab and tofacitinib, 18 to 20 weeks of therapy for certolizumab pegol (depending upon the dosing regimen), 20 weeks of therapy for ixekizumab, 22 weeks of therapy for infliximab, and 28 weeks of therapy for ustekinumab. It is recommended that a patient be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of PBS-subsidised initial treatment following a minimum of 12 weeks of therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

Grandfather patients (ixekizumab only).

A patient who commenced treatment with ixekizumab for severe psoriatic arthritis prior to 1 March 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

Grandfather patients (tofacitinib only).

A patient who commenced treatment with Tofacitinib for severe psoriatic arthritis prior to 1 May 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment under the Initial 1 or Initial 2 treatment restrictions.

For the second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to re-qualify with respect to either the indices of disease severity (i.e. erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP) level, and active joint count) or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify under the Initial 3 treatment restriction with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application.

However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

Within a treatment cycle a patient may alternate between therapy with any biological medicine of their choice (1 at a time) providing:

- (i) they have not received PBS-subsidised treatment with that particular biological medicine previously; or
- (ii) they have demonstrated an adequate response to that particular biological medicine if they have previously trialled it on the PBS; and
- (iii) they have not previously failed to respond to treatment 3 times in this treatment cycle with that biological medicine.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(4) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the indices of disease severity submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and these revised baseline measurements will be used to assess response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to determine response for all subsequent continuing treatments. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level

respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. 20 or more active joints), response will be determined according to a reduction in the total number of active joints. (5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment restriction according to the criteria of the relevant restriction and index of disease severity. The application must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response to methotrexate at a dose of at least 20 mg weekly for a minimum period of 3 months, **AND**
- Patient must have failed to achieve an adequate response to sulfasalazine at a dose of at least 2 g per day for a minimum period of 3 months; OR
- Patient must have failed to achieve an adequate response to leflunomide at a dose of up to 20 mg daily for a minimum period of 3 months, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Where treatment with methotrexate, sulfasalazine or leflunomide is contraindicated according to the relevant TGA-approved Product Information, details must be provided at the time of application.

Where intolerance to treatment with methotrexate, sulfasalazine or leflunomide developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

The following initiation criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; and

either

- (a) an active joint count of at least 20 active (swollen and tender) joints; or
- (b) at least 4 active joints from the following list of major joints:
 - (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Severe Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

An assessment of a patient's response to an initial course of treatment must be conducted following a minimum of 12 weeks of therapy. An application for the continuing treatment must be accompanied with the assessment of response and submitted to the Department of Human Services no later than 4 weeks from the date of completion of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 3 months treatment with methotrexate and 3 months treatment with sulfasalazine or leflunomide can be found on the Department of Human Services website (www.humanservices.gov.au)

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos Or mailed to:
Services Australia
Complex Drugs

Reply Paid 9826
HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial treatment - Initial 2 (change or recommencement of treatment after a break in in biological medicine of less than 5 years)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an erythrocyte sedimentation rate (ESR) no greater than 25 mm per hour or a C-reactive protein (CRP) level no greater than 15 mg per L or either marker reduced by at least 20% from baseline; and

either of the following:

- (a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or
- (b) a reduction in the number of the following major active joints, from at least 4, by at least 50%:
 - (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Severe Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to change or recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial treatment - Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must have an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; OR
- The condition must have a C-reactive protein (CRP) level greater than 15 mg per L, **AND**
- The condition must have either (a) a total active joint count of at least 20 active (swollen and tender) joints; or (b) at least 4 active major joints, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Major joints are defined as (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

All measures of joint count and ESR and/or CRP must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Severe Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial 1 (new patient) or Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years) - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 16 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

golimumab 50 mg/0.5 mL injection, 0.5 mL pen device

11365E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	1160.52	41.30	Simponi [JC]

golimumab 50 mg/0.5 mL injection, 0.5 mL syringe

3430M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	1160.52	41.30	Simponi [JC]

Interleukin inhibitors

■ **GUSELKUMAB**

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years); or

(iii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or

(iv) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient be reviewed in the 4 weeks prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

Grandfather patients (risankizumab only).

A patient who commenced treatment with risankizumab for chronic plaque psoriasis prior to 1 December 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the Initial treatment Grandfather treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised

treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment. This assessment must be conducted within 4 weeks of the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the 4 weeks prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the 4 weeks prior to completing their current course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure patients receive the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

Services Australia will determine whether a response to treatment has been demonstrated based on the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 the PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Whole body (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis where lesions have been present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 5 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; (iii) ciclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks; (v) apremilast at a dose of 30 mg twice a day for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 20 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, ciclosporin, apremilast or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application. Where intolerance to treatment with phototherapy, methotrexate, ciclosporin, apremilast or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, ciclosporin, apremilast, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

- (a) A current Psoriasis Area and Severity Index (PASI) score of greater than 15, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment.
- (b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 4 weeks following cessation of each course of treatment.
- (c) The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and
 - (ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, ciclosporin, apremilast or acitretin can be found on the Services Australia website (www.servicesaustralia.gov.au).

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos. Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Whole body (change or re-commencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 20 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and
 - (ii) details of prior biological treatment, including dosage, date and duration of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 3, Whole body (re-commencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must have a current Psoriasis Area and Severity Index (PASI) score of greater than 15, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 20 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Face, hand, foot (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 5 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; (iii) ciclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks; (v) apremilast at a dose of 30 mg twice a day for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 20 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, ciclosporin, apremilast or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application. Where intolerance to treatment with phototherapy, methotrexate, ciclosporin, apremilast or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, ciclosporin, apremilast, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

(a) Chronic plaque psoriasis classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:

(i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment; or

(ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment;

(b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 4 weeks following cessation of each course of treatment.

(c) The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition; and

(ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, ciclosporin, apremilast or acitretin can be found on the Services Australia website (www.servicesaustralia.gov.au).

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Face, hand, foot (change or re-commencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 20 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or
- (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:

- (i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition; and
- (ii) details of prior biological treatment, including dosage, date and duration of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 3, Face, hand, foot (re-commencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must be classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where: (i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or (ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 20 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Whole body or Face, hand, foot (new patient) or Initial 2, Whole body or Face, hand, foot (change or re-commencement of treatment after a break in biological medicine of less than 5 years) or Initial 3, Whole body or Face, hand, foot (re-commencement of treatment after a break in biological medicine of more than 5 years) - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1, Whole body (new patient) restriction to complete 20 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2, Whole body (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 20 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3, Whole body (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 20 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 1, Face, hand, foot (new patient) restriction to complete 20 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2, Face, hand, foot (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 20 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3, Face, hand, foot (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 20 weeks treatment, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- The treatment must provide no more than the balance of up to 20 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a dermatologist.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Whole body

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

The authority application must be made in writing and must include:

- a completed authority prescription form(s); and
- a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

An application for the continuing treatment must be accompanied with the assessment of response conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos. Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Face, hand, foot

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or
- a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet and face, hand, foot area diagrams including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

An application for the continuing treatment must be accompanied with the assessment of response conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Whole body or Continuing treatment, Face, hand, foot - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the continuing treatment, Whole body restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the continuing treatment, Face, hand, foot restriction to complete 24 weeks treatment, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a dermatologist.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

guselkumab 100 mg/mL injection, 1 x 1 mL syringe

11614G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	3795.78	41.30	Tremfya [JC]

■ IXEKIZUMAB

Note TREATMENT OF ADULT PATIENTS WITH ANKYLOSING SPONDYLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab for adult patients with ankylosing spondylitis.

Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 7 biological medicines at any 1 time.

Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy.

A patient who has been receiving PBS-subsidised adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab treatment prior to 1 December 2020 is considered to start their first cycle as of 1 December 2020.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last prescription for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than

5 years may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised biological medicine treatment with adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised biological medicine treatment in this treatment cycle and wishes to commence such therapy (Initial 1 - New patient)

(ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years); or

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

A patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy.

(b) Continuing treatment.

For the first continuing treatment course of PBS-subsidised biological medicine, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1, Initial 2 or Initial 3 treatment restrictions. For second and subsequent continuing courses of PBS-subsidised biological medicine it is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply. Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be conducted no later than 4 weeks from the cessation of that treatment course. Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI), or the prior NSAID therapy and exercise program requirements.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(3) Baseline measurements to determine response.

Services Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the BASDAI, ESR and/or CRP submitted with the first authority application for a biological medicine. For a new patient, the BASDAI used to determine the baseline must be measured while the patient is receiving NSAID therapy and completing their exercise program.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response.

Prescribers may provide new baseline measurements any time an 'Initial treatment' authority application is submitted within a treatment cycle and the eligibility for continuing treatment must be assessed according to these revised baseline measurements.

(4) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must qualify under the Initial 3 treatment restriction. The same clinical criteria and indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI) as for the Initial 1 (New patient) restriction will need to be met, but a re-trial of NSAID therapy and exercise therapy is not required.

Note Special Pricing Arrangements apply.

Authority required

Ankylosing spondylitis

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Ankylosing Spondylitis PBS Authority Application Form.

An adequate response is defined as an improvement from baseline of at least 2 units (on a scale of 0-10) in the BASDAI score combined with at least 1 of the following:

- (a) an ESR measurement no greater than 25 mm per hour; or
- (b) a CRP measurement no greater than 10 mg per L; or
- (c) an ESR or CRP measurement reduced by at least 20% from baseline.

Where only 1 acute phase reactant measurement is supplied in the first application for PBS-subsidised treatment, that same marker must be measured and supplied in all subsequent continuing treatment applications.

All measurements provided must be no more than 4 weeks old at the time of application.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where a response assessment is not conducted within these timeframes, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Ankylosing spondylitis

Treatment Phase: Continuing treatment - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the Continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restriction.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

ixekizumab 80 mg/mL injection, 2 x 1 mL pen devices

12209N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	3421.16	41.30	Taltz [LY]

■ IXEKIZUMAB**Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS**

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological

medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years); or

(iii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or

(iv) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient be reviewed in the 4 weeks prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

Grandfather patients (risankizumab only).

A patient who commenced treatment with risankizumab for chronic plaque psoriasis prior to 1 December 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the Initial treatment Grandfather treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment. This assessment must be conducted within 4 weeks of the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the 4 weeks prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the 4 weeks prior to completing their current course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate

biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure patients receive the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

Services Australia will determine whether a response to treatment has been demonstrated based on the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 the PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Whole body

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

The authority application must be made in writing and must include:

- a completed authority prescription form(s); and
- a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online

Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Face, hand, foot

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or
- (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet and face, hand, foot area diagrams including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

The PASI assessment for continuing treatment must be performed on the same affected area assessed at baseline.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Whole body or Continuing treatment, Face, hand, foot - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the continuing treatment, Whole body restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the continuing treatment, Face, hand, foot restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restrictions, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate).

Treatment criteria:

- Must be treated by a dermatologist.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

ixekizumab 80 mg/mL injection, 2 x 1 mL pen devices

11033Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	3421.16	41.30	Taltz [LY]

■ IXEKIZUMAB

Note TREATMENT OF ADULT PATIENTS WITH ANKYLOSING SPONDYLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab for adult patients with ankylosing spondylitis.

Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 7 biological medicines at any 1 time.

Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy.

A patient who has been receiving PBS-subsidised adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab treatment prior to 1 December 2020 is considered to start their first cycle as of 1 December 2020.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last prescription for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised biological medicine treatment with adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised biological medicine treatment in this treatment cycle and wishes to commence such therapy (Initial 1 - New patient)

(ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years); or

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

A patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy.

(b) Continuing treatment.

For the first continuing treatment course of PBS-subsidised biological medicine, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1, Initial 2 or Initial 3 treatment restrictions. For second and subsequent continuing courses of PBS-subsidised biological medicine it is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply. Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be conducted no later than 4 weeks from the cessation of that treatment course. Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI), or the prior NSAID therapy and exercise program requirements.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed

treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(3) Baseline measurements to determine response.

Services Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the BASDAI, ESR and/or CRP submitted with the first authority application for a biological medicine. For a new patient, the BASDAI used to determine the baseline must be measured while the patient is receiving NSAID therapy and completing their exercise program.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response.

Prescribers may provide new baseline measurements any time an 'Initial treatment' authority application is submitted within a treatment cycle and the eligibility for continuing treatment must be assessed according to these revised baseline measurements.

(4) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must qualify under the Initial 3 treatment restriction. The same clinical criteria and indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI) as for the Initial 1 (New patient) restriction will need to be met, but a re-trial of NSAID therapy and exercise therapy is not required.

Note Special Pricing Arrangements apply.

Authority required

Ankylosing spondylitis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Clinical criteria:

- The condition must be radiographically (plain X-ray) confirmed Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have at least 2 of the following: (i) low back pain and stiffness for 3 or more months that is relieved by exercise but not by rest; or (ii) limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by a score of at least 1 on each of the lumbar flexion and lumbar side flexion measurements of the Bath Ankylosing Spondylitis Metrology Index (BASMI); or (iii) limitation of chest expansion relative to normal values for age and gender, **AND**

AND

- Patient must have failed to achieve an adequate response following treatment with at least 2 non-steroidal anti-inflammatory drugs (NSAIDs), whilst completing an appropriate exercise program, for a total period of 3 months, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

The application must include details of the NSAIDs trialled, their doses and duration of treatment.

If the NSAID dose is less than the maximum recommended dose in the relevant TGA-approved Product Information, the application must include the reason a higher dose cannot be used.

If treatment with NSAIDs is contraindicated according to the relevant TGA-approved Product Information, the application must provide details of the contraindication.

If intolerance to NSAID treatment develops during the relevant period of use which is of a severity to necessitate permanent treatment withdrawal, the application must provide details of the nature and severity of this intolerance.

The following criteria indicate failure to achieve an adequate response and must be demonstrated at the time of the initial application:

- (a) a Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 4 on a 0-10 scale; and
- (b) an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 10 mg per L.

The baseline BASDAI score and ESR or CRP level must be determined at the completion of the 3 month NSAID and exercise trial, but prior to ceasing NSAID treatment. All measurements must be no more than 4 weeks old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reason this criterion cannot be satisfied.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Ankylosing Spondylitis PBS Authority Application Form which includes the following:
 - (i) details of the radiological report confirming Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis; and
 - (ii) a baseline BASDAI score; and
 - (iii) a completed Exercise Program Self Certification Form included in the supporting information form; and
 - (iv) baseline ESR and/or CRP level

An assessment of a patient's response to an initial course of treatment must be conducted following a minimum of 12 weeks of therapy. An application for the continuing treatment must be accompanied with the assessment of response and submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where a response assessment is not conducted within these timeframes, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Details of the toxicities, including severity, which will be accepted for the purposes of administering this restriction can be found on the Services Australia website at www.servicesaustralia.gov.au

Note For details on the appropriate minimum exercise program that will be accepted for the purposes of administering this restriction, please refer to the Services Australia website at www.servicesaustralia.gov.au

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Ankylosing spondylitis

Treatment Phase: Initial treatment - Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Ankylosing Spondylitis PBS Authority Application Form.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to change or recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

An adequate response is defined as an improvement from baseline of at least 2 units (on a scale of 0-10) in the BASDAI score combined with at least 1 of the following:

- (a) an ESR measurement no greater than 25 mm per hour; or
- (b) a CRP measurement no greater than 10 mg per L; or
- (c) an ESR or CRP measurement reduced by at least 20% from baseline.

Where only 1 acute phase reactant measurement is supplied in the first application for PBS-subsidised treatment, that same marker must be measured and supplied in all subsequent continuing treatment applications.

All measurements provided must be no more than 4 weeks old at the time of application.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs

Reply Paid 9826
HOBART TAS 7001

Authority required

Ankylosing spondylitis

Treatment Phase: Initial treatment - Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must be radiographically (plain X-ray) confirmed Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis, **AND**
- Patient must have at least 2 of the following: (i) low back pain and stiffness for 3 or more months that is relieved by exercise but not by rest; or (ii) limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by a score of at least 1 on each of the lumbar flexion and lumbar side flexion measurements of the Bath Ankylosing Spondylitis Metrology Index (BASMI); or (iii) limitation of chest expansion relative to normal values for age and gender,

AND

- Patient must have a Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 4 on a 0-10 scale that is no more than 4 weeks old at the time of application, **AND**
- Patient must have an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour that is no more than 4 weeks old at the time of application; OR
- Patient must have a C-reactive protein (CRP) level greater than 10 mg per L that is no more than 4 weeks old at the time of application; OR
- Patient must have a clinical reason as to why demonstration of an elevated ESR or CRP cannot be met and the application must state the reason, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Ankylosing Spondylitis PBS Authority Application Form which includes the following:
 - (i) details of the radiological report confirming Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis; and
 - (ii) a BASDAI score.

An assessment of a patient's response to an initial course of treatment must be conducted following a minimum of 12 weeks of therapy. An application for the continuing treatment must be accompanied with the assessment of response and submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where a response assessment is not conducted within these timeframes, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Ankylosing spondylitis

Treatment Phase: Initial treatment - Initial 1 (new patient), Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years) - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 16 weeks treatment, **AND**

- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

ixekizumab 80 mg/mL injection, 2 x 1 mL pen devices

12217B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	3421.16	41.30	Taltz [LY]

IXEKIZUMAB
Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

- a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or
- a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years); or
- a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or
- a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient be reviewed in the 4 weeks prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

Grandfather patients (risankizumab only).

A patient who commenced treatment with risankizumab for chronic plaque psoriasis prior to 1 December 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the Initial treatment Grandfather treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised

treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment. This assessment must be conducted within 4 weeks of the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the 4 weeks prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the 4 weeks prior to completing their current course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure patients receive the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

Services Australia will determine whether a response to treatment has been demonstrated based on the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 the PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Whole body (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis where lesions have been present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 5 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; (iii) ciclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks; (v) apremilast at a dose of 30 mg twice a day for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, ciclosporin, apremilast or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application. Where intolerance to treatment with phototherapy, methotrexate, ciclosporin, apremilast or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, ciclosporin, apremilast, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

- (a) A current Psoriasis Area and Severity Index (PASI) score of greater than 15, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment.
- (b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 4 weeks following cessation of each course of treatment.
- (c) The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and
 - (ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, ciclosporin, apremilast or acitretin can be found on the Services Australia website (www.servicesaustralia.gov.au).

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos. Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Whole body (change or re-commencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and
 - (ii) details of prior biological treatment, including dosage, date and duration of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 3, Whole body (re-commencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must have a current Psoriasis Area and Severity Index (PASI) score of greater than 15, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Face, hand, foot (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 5 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; (iii) ciclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks; (v) apremilast at a dose of 30 mg twice a day for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, ciclosporin, apremilast or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application. Where intolerance to treatment with phototherapy, methotrexate, ciclosporin, apremilast or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, ciclosporin, apremilast, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

(a) Chronic plaque psoriasis classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:

(i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment; or

(ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment;

(b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 4 weeks following cessation of each course of treatment.

(c) The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition; and

(ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, ciclosporin, apremilast or acitretin can be found on the Services Australia website (www.servicesaustralia.gov.au).

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Face, hand, foot (change or re-commencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or
- (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:

- (i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition; and
- (ii) details of prior biological treatment, including dosage, date and duration of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 3, Face, hand, foot (re-commencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must be classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where: (i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or (ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Whole body or Face, hand, foot (new patient) or Initial 2, Whole body or Face, hand, foot (change or re-commencement of treatment after a break in biological medicine of less than 5 years) or Initial 3, Whole body or Face, hand, foot (re-commencement of treatment after a break in biological medicine of more than 5 years) - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1, Whole body (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2, Whole body (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3, Whole body (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 1, Face, hand, foot (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2, Face, hand, foot (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3, Face, hand, foot (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 16 weeks treatment, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a dermatologist.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

ixekizumab 80 mg/mL injection, 2 x 1 mL pen devices

11032P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	3421.16	41.30	Taltz [LY]

■ IXEKIZUMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE PSORIATIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab for adult patients with severe active psoriatic arthritis. Therefore, where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient receiving PBS-subsidised treatment for psoriatic arthritis is able to commence a treatment cycle where they may trial biological medicines without having to experience a disease flare when swapping to the alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

A patient who received PBS-subsidised adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab or ustekinumab treatment prior to 1 May 2019 is considered to start their first cycle as of 1 May 2019.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence another cycle [further details are under '(5) Recommencement of treatment after a 5-year break in PBS-subsidised therapy' below].

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe active psoriatic arthritis.

(1) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an alternate medicine (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years) or An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, golimumab secukinumab and tofacitinib, 18 to 20 weeks of therapy for certolizumab pegol (depending upon the dosing regimen), 20 weeks of therapy for ixekizumab, 22 weeks of therapy for infliximab, and 28 weeks of therapy for ustekinumab. It is recommended that a patient be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of PBS-subsidised initial treatment following a minimum of 12 weeks of therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

Grandfather patients (ixekizumab only).

A patient who commenced treatment with ixekizumab for severe psoriatic arthritis prior to 1 March 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

Grandfather patients (tofacitinib only).

A patient who commenced treatment with Tofacitinib for severe psoriatic arthritis prior to 1 May 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction. A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment under the Initial 1 or Initial 2 treatment restrictions.

For the second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to re-qualify with respect to either the indices of disease severity (i.e. erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP) level, and active joint count) or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify under the Initial 3 treatment restriction with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application.

However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

Within a treatment cycle a patient may alternate between therapy with any biological medicine of their choice (1 at a time) providing:

- (i) they have not received PBS-subsidised treatment with that particular biological medicine previously; or
- (ii) they have demonstrated an adequate response to that particular biological medicine if they have previously trialled it on the PBS; and
- (iii) they have not previously failed to respond to treatment 3 times in this treatment cycle with that biological medicine.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(4) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the indices of disease severity submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and these revised baseline measurements will be used to assess response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to determine response for all subsequent continuing treatments. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. 20 or more active joints), response will be determined according to a reduction in the total number of active joints.

(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment restriction according to the criteria of the relevant restriction and index of disease severity. The application must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response to methotrexate at a dose of at least 20 mg weekly for a minimum period of 3 months, **AND**
- Patient must have failed to achieve an adequate response to sulfasalazine at a dose of at least 2 g per day for a minimum period of 3 months; OR
- Patient must have failed to achieve an adequate response to leflunomide at a dose of up to 20 mg daily for a minimum period of 3 months, **AND**
- Patient must not receive more than 20 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Where treatment with methotrexate, sulfasalazine or leflunomide is contraindicated according to the relevant TGA-approved Product Information, details must be provided at the time of application.

Where intolerance to treatment with methotrexate, sulfasalazine or leflunomide developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

The following initiation criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; and

either

(a) an active joint count of at least 20 active (swollen and tender) joints; or

(b) at least 4 active joints from the following list of major joints:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Severe Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

An assessment of a patient's response to an initial course of treatment must be conducted following a minimum of 12 weeks of therapy. An application for the continuing treatment must be accompanied with the assessment of response and submitted to the Department of Human Services no later than 4 weeks from the date of completion of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 3 months treatment with methotrexate and 3 months treatment with sulfasalazine or leflunomide can be found on the Department of Human Services website (www.humanservices.gov.au)

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial treatment - Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- Patient must not receive more than 20 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an erythrocyte sedimentation rate (ESR) no greater than 25 mm per hour or a C-reactive protein (CRP) level no greater than 15 mg per L or either marker reduced by at least 20% from baseline; and

either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following major active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Severe Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to change or recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial treatment - Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must have an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; OR
- The condition must have a C-reactive protein (CRP) level greater than 15 mg per L, **AND**
- The condition must have either (a) a total active joint count of at least 20 active (swollen and tender) joints; or (b) at least 4 active major joints, **AND**
- Patient must not receive more than 20 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Major joints are defined as (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

All measures of joint count and ESR and/or CRP must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Severe Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos. Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial 1 (new patient) or Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years) - balance of supply

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 20 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 20 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 20 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 20 weeks treatment available under the above restrictions.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial treatment - Grandfather treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must have received non-PBS subsidised treatment with this drug for this condition prior to 1 March 2019, **AND**
- Patient must be receiving treatment with this drug for this condition at the time of application, **AND**
- Patient must have demonstrated an adequate response following at least 12 weeks of non-PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have failed to achieve an adequate response to methotrexate at a dose of at least 20 mg weekly for a minimum period of 3 months prior to initiating non-PBS subsidised treatment with this drug for this condition, **AND**
- Patient must have failed to achieve an adequate response to sulfasalazine at a dose of at least 2 g per day for a minimum period of 3 months prior to initiating non-PBS subsidised treatment with this drug for this condition; OR
- Patient must have failed to achieve an adequate response to leflunomide at a dose of up to 20 mg daily for a minimum period of 3 months prior to initiating non-PBS subsidised treatment with this drug for this condition, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

The following initiation criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; and

either

(a) an active joint count of at least 20 active (swollen and tender) joints; or

(b) at least 4 active joints from the following list of major joints:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

An adequate response to treatment is defined as:

an erythrocyte sedimentation rate (ESR) no greater than 25 mm per hour or a C-reactive protein (CRP) level no greater than 15 mg per L or either marker reduced by at least 20% from baseline; and

either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following major active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to determine response for all subsequent continuing treatments.

The assessment of the patient's response to this PBS-subsidised course of therapy must be conducted no later than 4 weeks from the cessation of the treatment course.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

A patient may qualify for PBS-subsidised treatment under this restriction once only.

For continuing PBS-subsidised treatment, a Grandfathered patient must qualify under the Continuing treatment criteria.

The authority application must be made in writing and must include:

(1) a completed authority prescription form; and

(2) a completed Severe Psoriatic Arthritis PBS Authority Application - Supporting Information Form; and

(3) the date of commencement of this drug; and

(4) results of the baseline patient assessment prior to initiation of non-PBS subsidised therapy with this drug.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an erythrocyte sedimentation rate (ESR) no greater than 25 mm per hour or a C-reactive protein (CRP) level no greater than 15 mg per L or either marker reduced by at least 20% from baseline; and

either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following major active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to determine response for all subsequent continuing treatments.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Severe Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either Initial 1, Initial 2, or Initial 3 treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Continuing treatment or Grandfathered patients - balance of supply

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Continuing treatment restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Grandfathered treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restriction.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

ixekizumab 80 mg/mL injection, 2 x 1 mL pen devices

11623R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	3421.16	41.30	Taltz [LY]

■ RISANKIZUMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious

infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years); or

(iii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or

(iv) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient be reviewed in the 4 weeks prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

Grandfather patients (risankizumab only).

A patient who commenced treatment with risankizumab for chronic plaque psoriasis prior to 1 December 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the Initial treatment Grandfather treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment. This assessment must be conducted within 4 weeks of the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the 4 weeks prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the 4 weeks prior to completing their current course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure patients receive the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

Services Australia will determine whether a response to treatment has been demonstrated based on the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 the PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Whole body

Treatment criteria:

- Must be treated by a dermatologist.

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

The authority application must be made in writing and must include:

- a completed authority prescription form(s); and
- a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Face, hand, foot

Treatment criteria:

- Must be treated by a dermatologist.

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or
- (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet and face, hand, foot area diagrams including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos. Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Whole body or Continuing treatment, Face, hand, foot or Grandfathered patients - balance of supply

Treatment criteria:

- Must be treated by a dermatologist.

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the continuing treatment, Whole body restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the continuing treatment, Face, hand, foot restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Grandfathered treatment, Whole body restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Grandfathered treatment, Face, hand, foot restriction to complete 24 weeks treatment, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restrictions.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Written applications for authority approval for sufficient therapy to complete the balance of supply should be forwarded to:
Department of Human Services
Complex Drugs

Reply Paid 9826
HOBART TAS 7001

risankizumab 75 mg/0.83 mL injection, 2 x 0.83 mL syringes

11858D	Max. Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	5400.45	41.30	Skyrizi [VE]

■ RISANKIZUMAB**Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS**

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years); or

(iii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or

(iv) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient be reviewed in the 4 weeks prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

Grandfather patients (risankizumab only).

A patient who commenced treatment with risankizumab for chronic plaque psoriasis prior to 1 December 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the Initial treatment Grandfather treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment. This assessment must be conducted within 4 weeks of the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the 4 weeks prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the 4 weeks prior to completing their current course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure patients receive the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

Services Australia will determine whether a response to treatment has been demonstrated based on the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 the PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Whole body (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis where lesions have been present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 5 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; (iii) ciclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks; (v) apremilast at a dose of 30 mg twice a day for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 28 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, ciclosporin, apremilast or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application.

Where intolerance to treatment with phototherapy, methotrexate, ciclosporin, apremilast or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, ciclosporin, apremilast, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

- A current Psoriasis Area and Severity Index (PASI) score of greater than 15, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment.
- A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 4 weeks following cessation of each course of treatment.
- The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

- a completed authority prescription form(s); and
- a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
 - the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and
 - details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

At the time of the authority application, medical practitioners should request to provide for an initial course of this drug for this condition sufficient for up to 28 weeks of therapy, at a dose of 150 mg for weeks 0 and 4, then 150 mg every 12 weeks thereafter.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, ciclosporin, apremilast or acitretin can be found on the Services Australia website (www.servicesaustralia.gov.au).

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Whole body (change or re-commencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 28 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and
 - (ii) details of prior biological treatment, including dosage, date and duration of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

At the time of the authority application, medical practitioners should request to provide for an initial course of this drug for this condition sufficient for up to 28 weeks of therapy, at a dose of 150 mg for weeks 0 and 4, then 150 mg every 12 weeks thereafter.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 3, Whole body (re-commencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must have a current Psoriasis Area and Severity Index (PASI) score of greater than 15, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 28 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Face, hand, foot (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 5 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; (iii) ciclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks; (v) apremilast at a dose of 30 mg twice a day for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 28 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, ciclosporin, apremilast or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application.

Where intolerance to treatment with phototherapy, methotrexate, ciclosporin, apremilast or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, ciclosporin, apremilast, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

(a) Chronic plaque psoriasis classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:

(i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment; or

(ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment;

(b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 4 weeks following cessation of each course of treatment.

(c) The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition; and

(ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

At the time of the authority application, medical practitioners should request to provide for an initial course of this drug for this condition sufficient for up to 28 weeks of therapy, at a dose of 150 mg for weeks 0 and 4, then 150 mg every 12 weeks thereafter.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, ciclosporin, apremilast or acitretin can be found on the Services Australia website (www.servicesaustralia.gov.au).

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Face, hand, foot (change or re-commencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 28 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or
- (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition; and
 - (ii) details of prior biological treatment, including dosage, date and duration of treatment.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

At the time of the authority application, medical practitioners should request to provide for an initial course of this drug for this condition sufficient for up to 28 weeks of therapy, at a dose of 150 mg for weeks 0 and 4, then 150 mg every 12 weeks thereafter.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available

on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 3, Face, hand, foot (re-commencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must be classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:
 - (i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or
 - (ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 28 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Whole body or Face, hand, foot (new patient) or Initial 2, Whole body or Face, hand, foot (change or re-commencement of treatment after a break in biological medicine of less than 5 years) or Initial 3, Whole body or Face, hand, foot (re-commencement of treatment after a break in biological medicine of more than 5 years) - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1, Whole body (new patient) restriction to complete 28 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2, Whole body (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 28 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3, Whole body (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 28 weeks treatment; OR

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1, Face, hand, foot (new patient) restriction to complete 28 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2, Face, hand, foot (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 28 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3, Face, hand, foot (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 28 weeks treatment, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- The treatment must provide no more than the balance of up to 28 weeks treatment available under the above restriction.

Treatment criteria:

- Must be treated by a dermatologist.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

risankizumab 75 mg/0.83 mL injection, 2 x 0.83 mL syringes

11827L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	5400.45	41.30	Skyrizi [VE]

■ SECUKINUMAB
Note TREATMENT OF ADULT PATIENTS WITH ANKYLOSING SPONDYLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab for adult patients with ankylosing spondylitis.

Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 7 biological medicines at any 1 time.

Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy.

A patient who has been receiving PBS-subsidised adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab treatment prior to 1 December 2020 is considered to start their first cycle as of 1 December 2020.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last prescription for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised biological medicine treatment with adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised biological medicine treatment in this treatment cycle and wishes to commence such therapy (Initial 1 - New patient)

(ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years); or

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

A patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy.

(b) Continuing treatment.

For the first continuing treatment course of PBS-subsidised biological medicine, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1, Initial 2 or Initial 3 treatment restrictions. For second and subsequent continuing courses of PBS-subsidised biological medicine it is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply. Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be conducted no later than 4 weeks from the cessation of that treatment course. Where a response assessment is not conducted within the

required timeframe, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI), or the prior NSAID therapy and exercise program requirements.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(3) Baseline measurements to determine response.

Services Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the BASDAI, ESR and/or CRP submitted with the first authority application for a biological medicine. For a new patient, the BASDAI used to determine the baseline must be measured while the patient is receiving NSAID therapy and completing their exercise program.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response.

Prescribers may provide new baseline measurements any time an 'Initial treatment' authority application is submitted within a treatment cycle and the eligibility for continuing treatment must be assessed according to these revised baseline measurements.

(4) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must qualify under the Initial 3 treatment restriction. The same clinical criteria and indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI) as for the Initial 1 (New patient) restriction will need to be met, but a re-trial of NSAID therapy and exercise therapy is not required.

Note Special Pricing Arrangements apply.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Ankylosing spondylitis

Treatment Phase: Initial treatment - Initial 1 (new patient), Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years) - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 16 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

secukinumab 150 mg/mL injection, 1 mL pen device

10893H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	709.59	41.30	Cosentyx [NV]

▪ **SECUKINUMAB**

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological

medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

- (i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or
- (ii) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years); or
- (iii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or
- (iv) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient be reviewed in the 4 weeks prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

Grandfather patients (risankizumab only).

A patient who commenced treatment with risankizumab for chronic plaque psoriasis prior to 1 December 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the Initial treatment Grandfather treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment. This assessment must be conducted within 4 weeks of the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the 4 weeks prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the 4 weeks prior to completing their current course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater

than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure patients receive the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

Services Australia will determine whether a response to treatment has been demonstrated based on the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 the PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Authority approval for sufficient therapy to complete a maximum of 16 weeks of treatment may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial 1, Whole body or Face, hand, foot (new patient) or Initial 2, Whole body or Face, hand, foot (change or recommencement of treatment after a break in biological medicine of less than 5 years) or Initial 3, Whole body or Face, hand, foot (re-commencement of treatment after a break in biological medicine of more than 5 years) - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1, Whole body (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2, Whole body (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3, Whole body (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 1, Face, hand, foot (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2, Face, hand, foot (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3, Face, hand, foot (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 16 weeks treatment, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a dermatologist.

secukinumab 150 mg/mL injection, 2 x 1 mL pen devices

10494H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	1403.63	41.30	Cosentyx [NV]

▪ SECUKINUMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE PSORIATIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab for adult patients with severe active psoriatic arthritis. Therefore, where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient receiving PBS-subsidised treatment for psoriatic arthritis is able to commence a treatment cycle where they may trial biological medicines without having to experience a disease flare when swapping to the alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

A patient who received PBS-subsidised adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab or ustekinumab treatment prior to 1 May 2019 is considered to start their first cycle as of 1 May 2019.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence another cycle [further details are under '(5) Recommencement of treatment after a 5-year break in PBS-subsidised therapy' below].

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe active psoriatic arthritis.

(1) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an alternate medicine (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years) or An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, golimumab secukinumab and tofacitinib, 18 to 20 weeks of therapy for certolizumab pegol (depending upon the dosing regimen), 20 weeks of therapy for ixekizumab, 22 weeks of therapy for infliximab, and 28 weeks of therapy for ustekinumab. It is recommended that a patient be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of PBS-subsidised initial treatment following a minimum of 12 weeks of therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

Grandfather patients (ixekizumab only).

A patient who commenced treatment with ixekizumab for severe psoriatic arthritis prior to 1 March 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

Grandfather patients (tofacitinib only).

A patient who commenced treatment with Tofacitinib for severe psoriatic arthritis prior to 1 May 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment under the Initial 1 or Initial 2 treatment restrictions.

For the second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to re-qualify with respect to either the indices of disease severity (i.e. erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP) level, and active joint count) or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify under the Initial 3 treatment restriction with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application.

However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

Within a treatment cycle a patient may alternate between therapy with any biological medicine of their choice (1 at a time) providing:

- (i) they have not received PBS-subsidised treatment with that particular biological medicine previously; or
- (ii) they have demonstrated an adequate response to that particular biological medicine if they have previously trialled it on the PBS; and
- (iii) they have not previously failed to respond to treatment 3 times in this treatment cycle with that biological medicine.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(4) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the indices of disease severity submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and these revised baseline measurements will be used to assess response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to determine response for all subsequent continuing treatments. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. 20 or more active joints), response will be determined according to a reduction in the total number of active joints.

(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment restriction according to the criteria of the relevant restriction and index of disease severity. The application must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial 1 (new patient) or Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years) - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 16 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

secukinumab 150 mg/mL injection, 2 x 1 mL pen devices

10901R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	1403.63	41.30	Cosentyx [NV]

secukinumab 150 mg/mL injection, 1 mL pen device

10898N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	709.59	41.30	Cosentyx [NV]

■ SECUKINUMAB
Note TREATMENT OF ADULT PATIENTS WITH ANKYLOSING SPONDYLITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab for adult patients with ankylosing spondylitis.

Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 7 biological medicines at any 1 time.

Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy.

A patient who has been receiving PBS-subsidised adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab treatment prior to 1 December 2020 is considered to start their first cycle as of 1 December 2020.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last prescription for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised biological medicine treatment with adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised biological medicine treatment in this treatment cycle and wishes to commence such therapy (Initial 1 - New patient)

(ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years); or

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

A patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy.

(b) Continuing treatment.

For the first continuing treatment course of PBS-subsidised biological medicine, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1, Initial 2 or Initial 3 treatment restrictions. For second and subsequent continuing courses of PBS-subsidised biological medicine it is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be conducted no later than 4 weeks from the cessation of that treatment course. Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI), or the prior NSAID therapy and exercise program requirements.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(3) Baseline measurements to determine response.

Services Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the BASDAI, ESR and/or CRP submitted with the first authority application for a biological medicine. For a new patient, the BASDAI used to determine the baseline must be measured while the patient is receiving NSAID therapy and completing their exercise program.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response.

Prescribers may provide new baseline measurements any time an 'Initial treatment' authority application is submitted within a treatment cycle and the eligibility for continuing treatment must be assessed according to these revised baseline measurements.

(4) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must qualify under the Initial 3 treatment restriction. The same clinical criteria and indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI) as for the Initial 1 (New patient) restriction will need to be met, but a re-trial of NSAID therapy and exercise therapy is not required.

Note Special Pricing Arrangements apply.

Authority required

Ankylosing spondylitis

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Ankylosing Spondylitis PBS Authority Application - Supporting Information Form.

An adequate response is defined as an improvement from baseline of at least 2 of the BASDAI and 1 of the following:

- (a) an ESR measurement no greater than 25 mm per hour; or
- (b) a CRP measurement no greater than 10 mg per L; or
- (c) an ESR or CRP measurement reduced by at least 20% from baseline.

Where only 1 acute phase reactant measurement is supplied in the first application for PBS-subsidised treatment, that same marker must be measured and supplied in all subsequent continuing treatment applications.

All measurements provided must be no more than 1 month old at the time of application.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Ankylosing spondylitis

Treatment Phase: Continuing treatment - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the Continuing treatment restriction to complete 24 weeks treatment, **AND**

- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restriction.
- Treatment criteria:**
- Must be treated by a rheumatologist; OR
 - Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

secukinumab 150 mg/mL injection, 1 mL pen device

10906B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	709.59	41.30	Cosentyx [NV]

■ SECUKINUMAB

Note TREATMENT OF ADULT PATIENTS WITH NON-RADIOGRAPHIC AXIAL SPONDYLOARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of secukinumab, certolizumab pegol and golimumab for adult patients with non-radiographic axial spondyloarthritis.

Where the term 'biological medicine' appears in notes and restrictions, it refers to secukinumab, certolizumab pegol and golimumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 3 biological medicines at any one time.

Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy.

A serious adverse reaction requiring permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once.

Once a patient has either failed or ceased to respond to treatment 3 times (once with any biological medicine) within the same treatment cycle, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last prescription for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle.

A patient who has failed 3 trials or fewer of biological medicine therapy in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment with secukinumab, certolizumab pegol and golimumab

(1) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised biological medicine treatment in this treatment cycle and wishes to commence such therapy (Initial 1 - New patient)

(ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years); or

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

With the exception of grandfathered patients, a patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy.

(2) Continuing treatment.

For continuing courses of PBS-subsidised biological medicine it is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that biological medicine provided they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment in courses of up to 24 weeks provided they continue to sustain an adequate response.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be conducted no later than 4 weeks from the cessation of that treatment course. Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction requiring permanent treatment withdrawal.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. the C-reactive protein (CRP) level and the BASDAI score), or the prior NSAID therapy and exercise program requirements.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction requiring permanent treatment withdrawal.

(4) Baseline measurements to determine response.

A response to treatment is based on the baseline BASDAI score and CRP level documented in the patient's medical records.

For a new patient, the BASDAI score used to determine baseline disease severity must be measured while the patient is receiving NSAID therapy and completing their exercise program.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications.

Prescribers may provide new baseline measurements any time that an 'Initial treatment' authority application is submitted and the eligibility for continuing treatment must be assessed according to these revised baseline measurements.

(5) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must qualify under the Initial 3 treatment restriction. The same clinical criteria and indices of disease severity (i.e. the C-reactive protein (CRP) levels and the BASDAI) as for the Initial 1 (New patient) restriction will need to be met, but a re-trial of NSAID therapy and exercise therapy is not required.

Note No increase in the maximum quantity or number of units may be authorised.

Note Special Pricing Arrangements apply.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Non-radiographic axial spondyloarthritis

Treatment Phase: Initial 1 (New patient), Initial 2 (Change or recommencement of treatment after a break in biological medicine of less than 5 years) or Initial 3 (Recommencement of treatment after a break in biological medicine of more than 5 years) - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patients) restriction to complete 20 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 20 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 20 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 20 weeks treatment.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of non-radiographic axial spondyloarthritis.

Authority required

Non-radiographic axial spondyloarthritis

Treatment Phase: Continuing treatment or Grandfather patient - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Continuing treatment restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Grandfathered treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the continuing treatment restriction or the grandfather restriction.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of non-radiographic axial spondyloarthritis.

secukinumab 150 mg/mL injection, 1 mL pen device

12297F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	709.59	41.30	Cosentyx [NV]

■ SECUKINUMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE PSORIATIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab for adult patients with severe active psoriatic arthritis. Therefore, where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient receiving PBS-subsidised treatment for psoriatic arthritis is able to commence a treatment cycle where they may trial biological medicines without having to experience a disease flare when swapping to the alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

A patient who received PBS-subsidised adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab or ustekinumab treatment prior to 1 May 2019 is considered to start their first cycle as of 1 May 2019.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a

treatment failure.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence another cycle [further details are under '(5) Recommencement of treatment after a 5-year break in PBS-subsidised therapy' below].

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe active psoriatic arthritis.

(1) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an alternate medicine (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years) or An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, golimumab secukinumab and tofacitinib, 18 to 20 weeks of therapy for certolizumab pegol (depending upon the dosing regimen), 20 weeks of therapy for ixekizumab, 22 weeks of therapy for infliximab, and 28 weeks of therapy for ustekinumab. It is recommended that a patient be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of PBS-subsidised initial treatment following a minimum of 12 weeks of therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

Grandfather patients (ixekizumab only).

A patient who commenced treatment with ixekizumab for severe psoriatic arthritis prior to 1 March 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

Grandfather patients (tofacitinib only).

A patient who commenced treatment with Tofacitinib for severe psoriatic arthritis prior to 1 May 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment under the Initial 1 or Initial 2 treatment restrictions.

For the second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to re-qualify with respect to either the indices of disease severity (i.e. erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP) level, and active joint count) or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify under the

Initial 3 treatment restriction with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application.

However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

Within a treatment cycle a patient may alternate between therapy with any biological medicine of their choice (1 at a time) providing:

(i) they have not received PBS-subsidised treatment with that particular biological medicine previously; or
(ii) they have demonstrated an adequate response to that particular biological medicine if they have previously trialled it on the PBS; and

(iii) they have not previously failed to respond to treatment 3 times in this treatment cycle with that biological medicine.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(4) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the indices of disease severity submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and these revised baseline measurements will be used to assess response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to determine response for all subsequent continuing treatments. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. 20 or more active joints), response will be determined according to a reduction in the total number of active joints.

(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment restriction according to the criteria of the relevant restriction and index of disease severity. The application must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Severe psoriatic arthritis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an erythrocyte sedimentation rate (ESR) no greater than 25 mm per hour or a C-reactive protein (CRP) level no greater than 15 mg per L or either marker reduced by at least 20% from baseline; and

either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following major active joints, from at least 4, by at least 50%:

- (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
- (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to determine response for all subsequent continuing treatments.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Severe Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either Initial 1, Initial 2, or Initial 3 treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date

of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Continuing treatment - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restriction.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

secukinumab 150 mg/mL injection, 2 x 1 mL pen devices

10899P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1403.63	41.30	Cosentyx [NV]

secukinumab 150 mg/mL injection, 1 mL pen device

10895K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	709.59	41.30	Cosentyx [NV]

SECUKINUMAB

Note TREATMENT OF ADULT PATIENTS WITH NON-RADIOGRAPHIC AXIAL SPONDYLOARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of secukinumab, certolizumab pegol and golimumab for adult patients with non-radiographic axial spondyloarthritis.

Where the term 'biological medicine' appears in notes and restrictions, it refers to secukinumab, certolizumab pegol and golimumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 3 biological medicines at any one time.

Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy.

A serious adverse reaction requiring permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once.

Once a patient has either failed or ceased to respond to treatment 3 times (once with any biological medicine) within the same treatment cycle, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last prescription for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle.

A patient who has failed 3 trials or fewer of biological medicine therapy in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment with secukinumab, certolizumab pegol and golimumab

(1) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised biological medicine treatment in this treatment cycle and wishes to commence such therapy (Initial 1 - New patient)

(ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years); or

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years). With the exception of grandfathered patients, a patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy.

(2) Continuing treatment.

For continuing courses of PBS-subsidised biological medicine it is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that biological medicine provided they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment in courses of up to 24 weeks provided they continue to sustain an adequate response.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be conducted no later than 4 weeks from the cessation of that treatment course. Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction requiring permanent treatment withdrawal.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. the C-reactive protein (CRP) level and the BASDAI score), or the prior NSAID therapy and exercise program requirements.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction requiring permanent treatment withdrawal.

(4) Baseline measurements to determine response.

A response to treatment is based on the baseline BASDAI score and CRP level documented in the patient's medical records. For a new patient, the BASDAI score used to determine baseline disease severity must be measured while the patient is receiving NSAID therapy and completing their exercise program.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications.

Prescribers may provide new baseline measurements any time that an 'Initial treatment' authority application is submitted and the eligibility for continuing treatment must be assessed according to these revised baseline measurements.

(5) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must qualify under the Initial 3 treatment restriction. The same clinical criteria and indices of disease severity (i.e. the C-reactive protein (CRP) levels and the BASDAI) as for the Initial 1 (New patient) restriction will need to be met, but a re-trial of NSAID therapy and exercise therapy is not required.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Non-radiographic axial spondyloarthritis

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug for this condition, **AND**
- The treatment must not exceed a maximum of 24 weeks with this drug per authorised course under this restriction.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of non-radiographic axial spondyloarthritis. An adequate response to therapy with this biological medicine is defined as a reduction from baseline in the Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) score by 2 or more units (on a scale of 0-10) and 1 of the following:
 - (a) a CRP measurement no greater than 10 mg per L; or
 - (b) a CRP measurement reduced by at least 20% from baseline.

If the requirement to demonstrate an elevated CRP level could not be met under an initial treatment restriction, a reduction in the BASDAI score from baseline will suffice for the purposes of administering this continuing treatment restriction.

The patient remains eligible to receive continuing treatment with the same biological medicine in courses of up to 24 weeks providing they continue to sustain an adequate response. It is recommended that a patient be reviewed in the month prior to completing their current course of treatment.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Non-radiographic axial spondyloarthritis

Treatment Phase: Grandfather treatment

Clinical criteria:

- Patient must have previously received non-PBS-subsidised treatment with this drug for this indication prior to 1 April 2021, **AND**
- Patient must have had chronic lower back pain and stiffness for 3 or more months that was relieved by exercise but not rest, prior to initiating non-PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have had failed to achieve an adequate response following treatment with at least 2 non-steroidal anti-inflammatory drugs (NSAIDs), whilst completing an appropriate exercise program, for a total period of 3 months, prior to initiating non-PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have had one or more of the following: (a) enthesitis (heel); (b) uveitis; (c) dactylitis; (d) psoriasis; (e) inflammatory bowel disease; or (f) positive for Human Leukocyte Antigen B27 (HLA-B27); prior to initiating non-PBS-subsidised treatment with this drug for this condition, **AND**
- The condition must not be radiographically evidenced on plain x-ray of Grade II bilateral sacroiliitis or Grade III or IV unilateral sacroiliitis prior to commencing non-PBS-subsidised treatment with this biological medicine, **AND**
- The condition must have been diagnosed as non-radiographic axial spondyloarthritis, as defined by Assessment of Spondyloarthritis International Society (ASAS) criteria, prior to having commenced non-PBS-subsidised treatment with this biological medicine, **AND**
- The condition must have been sacroiliitis with active inflammation and/or oedema on non-contrast Magnetic Resonance Imaging (MRI) prior to commencing non-PBS-subsidised treatment with this biological medicine, **AND**
- The condition must have had presence of Bone Marrow Oedema (BMO) depicted as a hyperintense signal on a Short Tau Inversion Recovery (STIR) image (or equivalent) prior to commencing non-PBS-subsidised treatment with this biological medicine, **AND**
- The condition must have had BMO depicted as a hypointense signal on a T1 weighted image (without gadolinium) prior to commencing non-PBS-subsidised treatment with this biological medicine, **AND**
- The treatment must not exceed a maximum of 24 weeks with this drug under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of non-radiographic axial spondyloarthritis.

The application must include details of the NSAIDs trialled, their doses and duration of treatment.

If the NSAID dose is less than the maximum recommended dose in the relevant TGA-approved Product Information, the application must include the reason a higher dose cannot be used.

If treatment with NSAIDs is contraindicated according to the relevant TGA-approved Product Information, the application must provide details of the contraindication.

If intolerance to NSAID treatment develops during the relevant period of use which is of a severity to necessitate permanent treatment withdrawal, the application must provide details of the nature and severity of this intolerance.

The following criteria indicate failure to achieve an adequate response to NSAIDs and must have been demonstrated prior to initiation of non-PBS subsidised treatment with this biological medicine for this condition:

- (a) a Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) score of at least 4 on a 0-10 scale; and
- (b) C-reactive protein (CRP) level greater than 10 mg per L.

The BASDAI score and CRP level must have been determined at the completion of the 3-month NSAID and exercise trial, but prior to ceasing NSAID treatment. All measures must have been no more than 1 month old at the time of initiating non-PBS subsidised treatment with this biological medicine for this condition.

If the requirement to demonstrate an elevated CRP level could not be met, the reason must be stated in the application. Treatment with prednisolone dosed at 7.5 mg or higher daily (or equivalent) or a parenteral steroid within the past month (intramuscular or intravenous methylprednisolone or equivalent) is an acceptable reason.

The assessment of the patient's response to the initial course of treatment must be conducted following a minimum of 12 weeks of treatment and no later than 4 weeks from the cessation of that treatment course. If the response assessment is not conducted within these timeframes, the patient will be deemed to have failed this course of treatment in this treatment cycle.

A Grandfathered patient may qualify for PBS-subsidised treatment under this restriction once only. For continuing PBS-subsidised treatment, a Grandfathered patient must qualify under the continuing treatment criteria.

The authority application must include:

- (a) a completed authority prescription; and
- (b) a completed PBS authority application form relevant to the indication and treatment phase (the latest version located at the website mentioned in the administrative note).

The baseline BASDAI score and CRP level must also be documented in the patient's medical records.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

secukinumab 150 mg/mL injection, 1 mL pen device

12307R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	709.59	41.30	Cosentyx [NV]

■ SECUKINUMAB**Note TREATMENT OF ADULT PATIENTS WITH ANKYLOSING SPONDYLITIS**

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab for adult patients with ankylosing spondylitis.

Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 7 biological medicines at any 1 time.

Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy.

A patient who has been receiving PBS-subsidised adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab treatment prior to 1 December 2020 is considered to start their first cycle as of 1 December 2020.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once.

Once a patient has either failed or ceased to respond to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last prescription for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised biological medicine treatment with adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab and secukinumab.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised biological medicine treatment in this treatment cycle and wishes to commence such therapy (Initial 1 - New patient)

(ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years); or

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years).

A patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy.

(b) Continuing treatment.

For the first continuing treatment course of PBS-subsidised biological medicine, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1, Initial 2 or Initial 3 treatment restrictions. For second and subsequent continuing courses of PBS-subsidised biological medicine it is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be conducted no later than 4 weeks from the cessation of that treatment course. Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI), or the prior NSAID therapy and exercise program requirements.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(3) Baseline measurements to determine response.

Services Australia will determine whether a response to treatment has been demonstrated based on the baseline measurements of the BASDAI, ESR and/or CRP submitted with the first authority application for a biological medicine. For a new patient, the BASDAI used to determine the baseline must be measured while the patient is receiving NSAID therapy and completing their exercise program.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response.

Prescribers may provide new baseline measurements any time an 'Initial treatment' authority application is submitted within a treatment cycle and the eligibility for continuing treatment must be assessed according to these revised baseline measurements.

(4) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must qualify under the Initial 3 treatment restriction. The same clinical criteria and indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the BASDAI) as for the Initial 1 (New patient) restriction will need to be met, but a re-trial of NSAID therapy and exercise therapy is not required.

Note Special Pricing Arrangements apply.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Ankylosing spondylitis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Clinical criteria:

- The condition must be radiographically (plain X-ray) confirmed Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have at least 2 of the following: (i) low back pain and stiffness for 3 or more months that is relieved by exercise but not by rest; or (ii) limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by a score of at least 1 on each of the lumbar flexion and lumbar side flexion measurements of the Bath Ankylosing Spondylitis Metrology Index (BASMI); or (iii) limitation of chest expansion relative to normal values for age and gender,

AND

- Patient must have failed to achieve an adequate response following treatment with at least 2 non-steroidal anti-inflammatory drugs (NSAIDs), whilst completing an appropriate exercise program, for a total period of 3 months, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

The application must include details of the NSAIDs trialled, their doses and duration of treatment.

If the NSAID dose is less than the maximum recommended dose in the relevant TGA-approved Product Information, the application must include the reason a higher dose cannot be used.

If treatment with NSAIDs is contraindicated according to the relevant TGA-approved Product Information, the application must provide details of the contraindication.

If intolerance to NSAID treatment develops during the relevant period of use which is of a severity to necessitate permanent treatment withdrawal, the application must provide details of the nature and severity of this intolerance.

The following criteria indicate failure to achieve an adequate response and must be demonstrated at the time of the initial application:

- (a) a Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 4 on a 0-10 scale; AND
- (b) an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 10 mg per L.

The BASDAI must be determined at the completion of the 3 month NSAID and exercise trial, but prior to ceasing NSAID treatment. The BASDAI must be no more than 1 month old at the time of initial application.

Both ESR and CRP measures should be provided with the initial treatment application and both must be no more than 1 month old. If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reason this criterion cannot be satisfied.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form; and
- (b) a completed Ankylosing Spondylitis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) a copy of the radiological report confirming Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis; and

(ii) a completed BASDAI Assessment Form; and

(iii) a completed Exercise Program Self Certification Form included in the supporting information form.

An assessment of a patient's response to an initial course of treatment must be conducted following a minimum of 12 weeks of therapy. An application for the continuing treatment must be accompanied with the assessment of response and submitted to the Department of Human Services no later than 4 weeks from the date of completion of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Details of the toxicities, including severity, which will be accepted for the purposes of administering this restriction can be found on the Department of Human Services website at www.humanservices.gov.au

Note For details on the appropriate minimum exercise program that will be accepted for the purposes of administering this restriction, please refer to the Department of Human Services website at www.humanservices.gov.au

Authority required

Ankylosing spondylitis

Treatment Phase: Initial treatment - Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

The authority application must be made in writing and must include:

(a) a completed authority prescription form; and

(b) a completed Ankylosing Spondylitis PBS Authority Application - Supporting Information Form.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to change or recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

An adequate response is defined as an improvement from baseline of at least 2 of the BASDAI and 1 of the following:

- (a) an ESR measurement no greater than 25 mm per hour; or
- (b) a CRP measurement no greater than 10 mg per L; or
- (c) an ESR or CRP measurement reduced by at least 20% from baseline.

Where only 1 acute phase reactant measurement is supplied in the first application for PBS-subsidised treatment, that same marker must be measured and supplied in all subsequent continuing treatment applications.

All measurements provided must be no more than 1 month old at the time of application.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Authority required

Ankylosing spondylitis

Treatment Phase: Initial treatment - Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must be radiographically (plain X-ray) confirmed Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis, **AND**

- Patient must have at least 2 of the following: (i) low back pain and stiffness for 3 or more months that is relieved by exercise but not by rest; or (ii) limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by a score of at least 1 on each of the lumbar flexion and lumbar side flexion measurements of the Bath Ankylosing Spondylitis Metrology Index (BASMI); or (iii) limitation of chest expansion relative to normal values for age and gender,

AND

- Patient must have a Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 4 on a 0-10 scale that is no more than 4 weeks old at the time of application, **AND**
- Patient must have an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour that is no more than 4 weeks old at the time of application; OR
- Patient must have a C-reactive protein (CRP) level greater than 10 mg per L that is no more than 4 weeks old at the time of application; OR
- Patient must have a clinical reason as to why demonstration of an elevated ESR or CRP cannot be met and the application must state the reason, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of ankylosing spondylitis.

The authority application must be made in writing and must include:

(a) a completed authority prescription form; and

(b) a completed Ankylosing Spondylitis PBS Authority Application - Supporting Information Form which includes the following:

(i) a copy of the radiological report confirming Grade II bilateral sacroiliitis or Grade III unilateral sacroiliitis; and

(ii) a completed BASDAI Assessment Form.

An assessment of a patient's response to an initial course of treatment must be conducted following a minimum of 12 weeks of therapy. An application for the continuing treatment must be accompanied with the assessment of response and submitted to the Department of Human Services no later than 4 weeks from the date of completion of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

secukinumab 150 mg/mL injection, 1 mL pen device

10890E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	4	..	..	*2783.46	41.30	Cosentyx [NV]

■ SECUKINUMAB**Note TREATMENT OF ADULT PATIENTS WITH NON-RADIOGRAPHIC AXIAL SPONDYLOARTHRITIS**

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of secukinumab, certolizumab pegol and golimumab for adult patients with non-radiographic axial spondyloarthritis.

Where the term 'biological medicine' appears in notes and restrictions, it refers to secukinumab, certolizumab pegol and golimumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 3 biological medicines at any one time.

Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy.

A serious adverse reaction requiring permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once.

Once a patient has either failed or ceased to respond to treatment 3 times (once with any biological medicine) within the same treatment cycle, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last prescription for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle.

A patient who has failed 3 trials or fewer of biological medicine therapy in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment with secukinumab, certolizumab pegol and golimumab

(1) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised biological medicine treatment in this treatment cycle and wishes to commence such therapy (Initial 1 - New patient)

(ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further

details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same agent (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years); or

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years). With the exception of grandfathered patients, a patient must be assessed for response to a course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy.

(2) Continuing treatment.

For continuing courses of PBS-subsidised biological medicine it is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that biological medicine provided they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment in courses of up to 24 weeks provided they continue to sustain an adequate response.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be conducted no later than 4 weeks from the cessation of that treatment course. Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction requiring permanent treatment withdrawal.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. the C-reactive protein (CRP) level and the BASDAI score), or the prior NSAID therapy and exercise program requirements.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction requiring permanent treatment withdrawal.

(4) Baseline measurements to determine response.

A response to treatment is based on the baseline BASDAI score and CRP level documented in the patient's medical records. For a new patient, the BASDAI score used to determine baseline disease severity must be measured while the patient is receiving NSAID therapy and completing their exercise program.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications.

Prescribers may provide new baseline measurements any time that an 'Initial treatment' authority application is submitted and the eligibility for continuing treatment must be assessed according to these revised baseline measurements.

(5) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must qualify under the Initial 3 treatment restriction. The same clinical criteria and indices of disease severity (i.e. the C-reactive protein (CRP) levels and the BASDAI) as for the Initial 1 (New patient) restriction will need to be met, but a re-trial of NSAID therapy and exercise therapy is not required.

Note No increase in the maximum quantity or number of units may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Non-radiographic axial spondyloarthritis

Treatment Phase: Initial treatment - Initial 1 (New patient)

Clinical criteria:

- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have had chronic lower back pain and stiffness for 3 or more months that is relieved by exercise but not rest, **AND**
- Patient must have failed to achieve an adequate response following treatment with at least 2 non-steroidal anti-inflammatory drugs (NSAIDs), whilst completing an appropriate exercise program, for a total period of 3 months, **AND**
- Patient must have one or more of the following: (a) enthesitis (heel); (b) uveitis; (c) dactylitis; (d) psoriasis; (e) inflammatory bowel disease; or (f) positive for Human Leukocyte Antigen B27 (HLA-B27), **AND**
- The condition must not be radiographically evidenced on plain x-ray of Grade II bilateral sacroiliitis or Grade III or IV unilateral sacroiliitis, **AND**
- The condition must be non-radiographic axial spondyloarthritis, as defined by Assessment of Spondyloarthritis International Society (ASAS) criteria, **AND**
- The condition must be sacroiliitis with active inflammation and/or oedema on non-contrast Magnetic Resonance Imaging (MRI), **AND**
- The condition must have presence of Bone Marrow Oedema (BMO) depicted as a hyperintense signal on a Short Tau Inversion Recovery (STIR) image (or equivalent), **AND**
- The condition must have BMO depicted as a hypointense signal on a T1 weighted image (without gadolinium), **AND**
- Patient must not receive more than 20 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of non-radiographic axial spondyloarthritis.

The stated maximum quantity of 5 with zero repeats is intended for a patient undergoing the loading dose regimen of 150 mg administered at weeks 0, 1, 2, 3, and 4 (a total of 5 doses) followed by monthly administration thereafter.

State in the application whether a loading dose regimen is intended or not.

Where a loading dose regimen is intended, request a maximum quantity of 5 and zero repeats to cover doses at weeks 0, 1, 2, 3 and 4. Doses at week 8, 12, and 16 can be sought under the relevant 'Balance of supply' listing.

Where no loading dose regimen is intended, request a maximum quantity of 1 and seek an increase in the number of repeats from zero to 4 repeats to cover dosing at weeks 4, 8, 12 and 16. Where increased repeats are sought, the maximum quantity sought must not be greater than 1.

The application must include details of the NSAIDs trialled, their doses and duration of treatment.

If the NSAID dose is less than the maximum recommended dose in the relevant TGA-approved Product Information, the application must include the reason a higher dose cannot be used.

If treatment with NSAIDs is contraindicated according to the relevant TGA-approved Product Information, the application must provide details of the contraindication.

If intolerance to NSAID treatment develops during the relevant period of use which is of a severity to necessitate permanent treatment withdrawal, the application must provide details of the nature and severity of this intolerance.

The following criteria indicate failure to achieve an adequate response to NSAIDs and must be demonstrated at the time of the initial application:

- (a) a Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) score of at least 4 on a 0-10 scale; and
- (b) C-reactive protein (CRP) level greater than 10 mg per L.

The baseline BASDAI score and CRP level must be determined at the completion of the 3-month NSAID and exercise trial, but prior to ceasing NSAID treatment. All measures must be no more than 4 weeks old at the time of initial application.

If the requirement to demonstrate an elevated CRP level could not be met, the reason must be stated in the application.

Treatment with prednisolone dosed at 7.5 mg or higher daily (or equivalent) or a parenteral steroid within the past month (intramuscular or intravenous methylprednisolone or equivalent) is an acceptable reason.

The assessment of the patient's response to the initial course of treatment must be conducted following a minimum of 12 weeks of treatment and no later than 4 weeks from the cessation of that treatment course. If the response assessment is not conducted within these timeframes, the patient will be deemed to have failed this course of treatment in this treatment cycle.

The authority application must include:

- (a) a completed authority prescription; and
- (b) a completed PBS authority application form relevant to the indication and treatment phase (the latest version located at the website mentioned in the administrative note).

The baseline BASDAI score and CRP level must also be documented in the patient's medical records.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Non-radiographic axial spondyloarthritis

Treatment Phase: Initial treatment - Initial 2 (Change or recommencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have failed, or ceased to respond to, PBS-subsidised treatment with biological medicines more than three times for this PBS-indication during the current treatment cycle, **AND**
- Patient must not have failed PBS-subsidised therapy with this biological medicine for this PBS indication more than once in the current treatment cycle, **AND**
- Patient must not receive more than 20 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of non-radiographic axial spondyloarthritis.

An application for Initial 2 treatment must indicate whether the patient has demonstrated an adequate response (an absence of treatment failure), failed or experienced an intolerance to the most recent supply of biological medicine treatment.

A new baseline Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) score and C-reactive protein (CRP) level may be provided at the time of this application.

An adequate response to therapy with this biological medicine is defined as a reduction from baseline in the Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) score by 2 or more units (on a scale of 0-10) and 1 of the following:

- (a) a CRP measurement no greater than 10 mg per L; or
- (b) a CRP measurement reduced by at least 20% from baseline.

The assessment of the patient's response to the most recent supply of biological medicine must be conducted following a minimum of 12 weeks of treatment.

BASDAI scores and CRP levels must be documented in the patient's medical records.

The assessment of the patient's response to the initial course of treatment must be conducted following a minimum of 12 weeks of treatment and no later than 4 weeks from the cessation of that treatment course. If the response assessment is not conducted within these timeframes, the patient will be deemed to have failed this course of treatment in this treatment cycle. The following must be provided at the time of application and documented in the patient's medical records:

(a) the BASDAI score; and

(b) the C-reactive protein (CRP) level.

The stated maximum quantity of 5 with zero repeats is intended for a patient undergoing the loading dose regimen of 150 mg administered at weeks 0, 1, 2, 3, and 4 (a total of 5 doses) followed by monthly administration thereafter.

State in the application whether a loading dose regimen is intended or not.

Where a loading dose regimen is intended, request a maximum quantity of 5 and zero repeats to cover doses at weeks 0, 1, 2, 3 and 4. Doses at week 8, 12, and 16 can be sought under the relevant 'Balance of supply' listing.

Where no loading dose regimen is intended, request a maximum quantity of 1 and seek an increase in the number of repeats from zero to 4 repeats to cover dosing at weeks 4, 8, 12 and 16. Where increased repeats are sought, the maximum quantity sought must not be greater than 1.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Non-radiographic axial spondyloarthritis

Treatment Phase: Initial treatment - Initial 3 (Recommencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have had chronic lower back pain and stiffness for 3 or more months that is relieved by exercise but not rest, **AND**
- Patient must have had a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- Patient must have one or more of the following: (a) enthesitis (heel); (b) uveitis; (c) dactylitis; (d) psoriasis; (e) inflammatory bowel disease; or (f) positive for Human Leukocyte Antigen B27 (HLA-B27), **AND**
- The condition must not be radiographically evidenced on plain x-ray of Grade II bilateral sacroiliitis or Grade III or IV unilateral sacroiliitis, **AND**
- The condition must be non-radiographic axial spondyloarthritis, as defined by Assessment of Spondyloarthritis International Society (ASAS) criteria, **AND**
- The condition must be sacroiliitis with active inflammation and/or oedema on non-contrast Magnetic Resonance Imaging (MRI), **AND**
- The condition must have presence of Bone Marrow Oedema (BMO) depicted as a hyperintense signal on a Short Tau Inversion Recovery (STIR) image (or equivalent), **AND**
- The condition must have BMO depicted as a hypointense signal on a T1 weighted image (without gadolinium), **AND**
- Patient must not receive more than 20 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of non-radiographic axial spondyloarthritis.

The following must be provided at the time of application and documented in the patient's medical records:

(a) a Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) score of at least 4 on a 0-10 scale; and

(b) C-reactive protein (CRP) level greater than 10 mg per L.

The BASDAI score and CRP level must be no more than 4 weeks old at the time of this application.

If the requirement to demonstrate an elevated CRP level could not be met, the reason must be stated in the application.

Treatment with prednisolone dosed at 7.5 mg or higher daily (or equivalent) or a parenteral steroid within the past month (intramuscular or intravenous methylprednisolone or equivalent) is an acceptable reason.

The assessment of the patient's response to the initial course of treatment must be conducted following a minimum of 12 weeks of treatment and no later than 4 weeks from the cessation of that treatment course. If the response assessment is not conducted within these timeframes, the patient will be deemed to have failed this course of treatment in this treatment cycle.

The stated maximum quantity of 5 with zero repeats is intended for a patient undergoing the loading dose regimen of 150 mg administered at weeks 0, 1, 2, 3, and 4 (a total of 5 doses) followed by monthly administration thereafter.

State in the application whether a loading dose regimen is intended or not.

Where a loading dose regimen is intended, request a maximum quantity of 5 and zero repeats to cover doses at weeks 0, 1, 2, 3 and 4. Doses at week 8, 12, and 16 can be sought under the relevant 'Balance of supply' listing.

Where no loading dose regimen is intended, request a maximum quantity of 1 and seek an increase in the number of repeats from zero to 4 repeats to cover dosing at weeks 4, 8, 12 and 16. Where increased repeats are sought, the maximum quantity sought must not be greater than 1.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

secukinumab 150 mg/mL injection, 1 mL pen device

12321L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	5	..	..	*3452.59	41.30	Cosentyx [NV]

■ SECUKINUMAB
Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years); or

(iii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or

(iv) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient be reviewed in the 4 weeks prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

Grandfather patients (risankizumab only).

A patient who commenced treatment with risankizumab for chronic plaque psoriasis prior to 1 December 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the Initial treatment Grandfather treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment. This assessment must be conducted within 4 weeks of the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the 4 weeks prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions. For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the 4 weeks prior to completing their current course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure patients receive the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

Services Australia will determine whether a response to treatment has been demonstrated based on the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 the PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Whole body

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

The authority application must be made in writing and must include:

- a completed authority prescription form(s); and
- a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Face, hand, foot

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or
- (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet and face, hand, foot area diagrams including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

The PASI assessment for continuing treatment must be performed on the same affected area assessed at baseline.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia

Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Whole body or Continuing treatment, Face, hand, foot - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the continuing treatment, Whole body restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the continuing treatment, Face, hand, foot restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restrictions, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate).

Treatment criteria:

- Must be treated by a dermatologist.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

secukinumab 150 mg/mL injection, 2 x 1 mL pen devices

10425Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1403.63	41.30	Cosentyx [NV]

■ SECUKINUMAB**Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE PSORIATIC ARTHRITIS**

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab for adult patients with severe active psoriatic arthritis. Therefore, where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient receiving PBS-subsidised treatment for psoriatic arthritis is able to commence a treatment cycle where they may trial biological medicines without having to experience a disease flare when swapping to the alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

A patient who received PBS-subsidised adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab or ustekinumab treatment prior to 1 May 2019 is considered to start their first cycle as of 1 May 2019.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence another cycle [further details are under '(5) Recommencement of treatment after a 5-year break in PBS-subsidised therapy' below].

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe active psoriatic arthritis.

(1) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an alternate medicine (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years) or An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, golimumab secukinumab and tofacitinib, 18 to 20 weeks of therapy for certolizumab pegol (depending upon the dosing regimen), 20 weeks of therapy for ixekizumab, 22 weeks of therapy for infliximab, and 28 weeks of therapy for

ustekinumab. It is recommended that a patient be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of PBS-subsidised initial treatment following a minimum of 12 weeks of therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

Grandfather patients (ixekizumab only).

A patient who commenced treatment with ixekizumab for severe psoriatic arthritis prior to 1 March 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

Grandfather patients (tofacitinib only).

A patient who commenced treatment with Tofacitinib for severe psoriatic arthritis prior to 1 May 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment under the Initial 1 or Initial 2 treatment restrictions.

For the second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to re-qualify with respect to either the indices of disease severity (i.e. erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP) level, and active joint count) or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify under the Initial 3 treatment restriction with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application.

However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

Within a treatment cycle a patient may alternate between therapy with any biological medicine of their choice (1 at a time) providing:

- (i) they have not received PBS-subsidised treatment with that particular biological medicine previously; or
- (ii) they have demonstrated an adequate response to that particular biological medicine if they have previously trialled it on the PBS; and
- (iii) they have not previously failed to respond to treatment 3 times in this treatment cycle with that biological medicine.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(4) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the indices of disease severity submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and these revised baseline measurements will be used to assess response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to determine response for all subsequent continuing treatments. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. 20 or more active joints), response will be determined according to a reduction in the total number of active joints.

(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological

therapy of at least 5 years, must qualify under Initial 3 treatment restriction according to the criteria of the relevant restriction and index of disease severity. The application must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response to methotrexate at a dose of at least 20 mg weekly for a minimum period of 3 months, **AND**
- Patient must have failed to achieve an adequate response to sulfasalazine at a dose of at least 2 g per day for a minimum period of 3 months; OR
- Patient must have failed to achieve an adequate response to leflunomide at a dose of up to 20 mg daily for a minimum period of 3 months, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Where treatment with methotrexate, sulfasalazine or leflunomide is contraindicated according to the relevant TGA-approved Product Information, details must be provided at the time of application.

Where intolerance to treatment with methotrexate, sulfasalazine or leflunomide developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

The following initiation criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; and
either

- (a) an active joint count of at least 20 active (swollen and tender) joints; or
- (b) at least 4 active joints from the following list of major joints:
 - (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

The authority application must be in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Severe Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

An assessment of a patient's response to an initial course of treatment must be conducted following a minimum of 12 weeks of therapy. An application for the continuing treatment must be accompanied with the assessment of response and submitted to the Department of Human Services no later than 4 weeks from the date of completion of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 3 months treatment with methotrexate and 3 months treatment with sulfasalazine or leflunomide can be found on the Department of Human Services website (www.humanservices.gov.au)

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial treatment - Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an erythrocyte sedimentation rate (ESR) no greater than 25 mm per hour or a C-reactive protein (CRP) level no greater than 15 mg per L or either marker reduced by at least 20% from baseline; and

either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following major active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Severe Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to change or recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note The assessment of the patient's response to this initial course of treatment must be made following a minimum of 12 weeks of treatment and submitted to the Department of Human Services no later than 4 weeks from the cessation of the treatment course. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial treatment - Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must have an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; OR
- The condition must have a C-reactive protein (CRP) level greater than 15 mg per L, **AND**
- The condition must have either (a) a total active joint count of at least 20 active (swollen and tender) joints; or (b) at least 4 active major joints, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Major joints are defined as (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

All measures of joint count and ESR and/or CRP must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Severe Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

secukinumab 150 mg/mL injection, 2 x 1 mL pen devices

10894J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	4	..	..	*5427.42	41.30	Cosentyx [NV]

secukinumab 150 mg/mL injection, 1 mL pen device

10900Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	4	..	..	*2783.46	41.30	Cosentyx [NV]

■ SECUKINUMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than

5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction. There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

- (i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or
- (ii) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years); or
- (iii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or
- (iv) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient be reviewed in the 4 weeks prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

Grandfather patients (risankizumab only).

A patient who commenced treatment with risankizumab for chronic plaque psoriasis prior to 1 December 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the Initial treatment Grandfather treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment. This assessment must be conducted within 4 weeks of the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the 4 weeks prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the 4 weeks prior to completing their current course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure patients receive the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

Services Australia will determine whether a response to treatment has been demonstrated based on the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 the PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Whole body (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis where lesions have been present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 5 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; (iii) ciclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks; (v) apremilast at a dose of 30 mg twice a day for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, ciclosporin, apremilast or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application.

Where intolerance to treatment with phototherapy, methotrexate, ciclosporin, apremilast or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, ciclosporin, apremilast, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

- (a) A current Psoriasis Area and Severity Index (PASI) score of greater than 15, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment.
- (b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 4 weeks following cessation of each course of treatment.
- (c) The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and
 - (ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, ciclosporin, apremilast or acitretin can be found on the Services Australia website (www.servicesaustralia.gov.au).

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Whole body (change or re-commencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and
 - (ii) details of prior biological treatment, including dosage, date and duration of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 3, Whole body (re-commencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must have a current Psoriasis Area and Severity Index (PASI) score of greater than 15, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Face, hand, foot (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 5 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; (iii) ciclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks; (v) apremilast at a dose of 30 mg twice a day for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, ciclosporin, apremilast or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application. Where intolerance to treatment with phototherapy, methotrexate, ciclosporin, apremilast or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, ciclosporin, apremilast, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

(a) Chronic plaque psoriasis classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:

(i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment; or

(ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment;

(b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 4 weeks following cessation of each course of treatment.

(c) The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition; and

(ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, ciclosporin, apremilast or acitretin can be found on the Services Australia website (www.servicesaustralia.gov.au).

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Face, hand, foot (change or re-commencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or
- (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition; and
 - (ii) details of prior biological treatment, including dosage, date and duration of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 3, Face, hand, foot (re-commencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must be classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:
 - (i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or (ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

secukinumab 150 mg/mL injection, 2 x 1 mL pen devices

10910F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	4	..	..	*5427.42	41.30	Cosentyx [NV]

■ TILDRAKIZUMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

- a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or
- a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years); or
- a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or
- a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient be reviewed in the 4 weeks prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

Grandfather patients (risankizumab only).

A patient who commenced treatment with risankizumab for chronic plaque psoriasis prior to 1 December 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the Initial treatment Grandfather treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment. This assessment must be conducted within 4 weeks of the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the 4 weeks prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the 4 weeks prior to completing their current course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure patients receive the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

Services Australia will determine whether a response to treatment has been demonstrated based on the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 the PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Whole body

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**

- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

An application for the continuing treatment must be accompanied with the assessment of response conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Face, hand, foot

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or
- (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet and face, hand, foot area diagrams including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

An application for the continuing treatment must be accompanied with the assessment of response conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Whole body or Continuing treatment, Face, hand, foot - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the continuing treatment, Whole body restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the continuing treatment, Face, hand, foot restriction to complete 24 weeks treatment, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restrictions.

Treatment criteria:

- Must be treated by a dermatologist.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

tildrakizumab 100 mg/mL injection, 1 mL syringe

11613F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	3271.21	41.30	Ilumya [RA]

■ TILDRAKIZUMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction. There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years); or

(iii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or

(iv) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient be reviewed in the 4 weeks prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

Grandfather patients (risankizumab only).

A patient who commenced treatment with risankizumab for chronic plaque psoriasis prior to 1 December 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the Initial treatment Grandfather treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment. This assessment must be conducted within 4 weeks of the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the 4 weeks prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the 4 weeks prior to completing their current course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure patients receive the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

Services Australia will determine whether a response to treatment has been demonstrated based on the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be

assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 the PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Whole body (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis where lesions have been present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 5 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; (iii) ciclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks; (v) apremilast at a dose of 30 mg twice a day for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 28 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, ciclosporin, apremilast or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application.

Where intolerance to treatment with phototherapy, methotrexate, ciclosporin, apremilast or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, ciclosporin, apremilast, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

- (a) A current Psoriasis Area and Severity Index (PASI) score of greater than 15, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment.
- (b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 4 weeks following cessation of each course of treatment.
- (c) The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and
 - (ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

At the time of the authority application, medical practitioners should request to provide for an initial course of this drug for this condition sufficient for up to 28 weeks of therapy, at a dose of 100 mg for weeks 0 and 4, then 100 mg every 12 weeks thereafter.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, ciclosporin, apremilast or acitretin can be found on the Services Australia website (www.servicesaustralia.gov.au).

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Whole body (change or re-commencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 28 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:

- (i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and
- (ii) details of prior biological treatment, including dosage, date and duration of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

At the time of the authority application, medical practitioners should request to provide for an initial course of this drug for this condition sufficient for up to 28 weeks of therapy, at a dose of 100 mg for weeks 0 and 4, then 100 mg every 12 weeks thereafter.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 3, Whole body (re-commencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must have a current Psoriasis Area and Severity Index (PASI) score of greater than 15, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 28 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Face, hand, foot (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 5 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; (iii) ciclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks; (v) apremilast at a dose of 30 mg twice a day for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 28 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, ciclosporin, apremilast or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application.

Where intolerance to treatment with phototherapy, methotrexate, ciclosporin, apremilast or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, ciclosporin, apremilast, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

(a) Chronic plaque psoriasis classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:

(i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment; or

(ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment;

(b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 4 weeks following cessation of each course of treatment.

(c) The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition; and

(ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

At the time of the authority application, medical practitioners should request to provide for an initial course of this drug for this condition sufficient for up to 28 weeks of therapy, at a dose of 100 mg for weeks 0 and 4, then 100 mg every 12 weeks thereafter.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, ciclosporin, apremilast or acitretin can be found on the Services Australia website (www.servicesaustralia.gov.au).

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Face, hand, foot (change or re-commencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 28 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

(i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or

(ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition; and
 - (ii) details of prior biological treatment, including dosage, date and duration of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

At the time of the authority application, medical practitioners should request to provide for an initial course of this drug for this condition sufficient for up to 28 weeks of therapy, at a dose of 100 mg for weeks 0 and 4, then 100 mg every 12 weeks thereafter.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 3, Face, hand, foot (re-commencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must be classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:
 - (i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or (ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 28 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.
- The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Whole body or Face, hand, foot (new patient) or Initial 2, Whole body or Face, hand, foot (change or re-commencement of treatment after a break in biological medicine of less than 5 years) or Initial 3, Whole body or Face, hand, foot (re-commencement of treatment after a break in biological medicine of more than 5 years) - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1, Whole body (new patient) restriction to complete 28 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2, Whole body (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 28 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3, Whole body (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 28 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 1, Face, hand, foot (new patient) restriction to complete 28 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2, Face, hand, foot (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 28 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3, Face, hand, foot (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 28 weeks treatment, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- The treatment must provide no more than the balance of up to 28 weeks treatment available under the above restriction.

Treatment criteria:

- Must be treated by a dermatologist.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

tildrakizumab 100 mg/mL injection, 1 mL syringe

11616J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	3271.21	41.30	Ilumya [RA]

■ TOCILIZUMAB

Note TREATMENT OF PATIENTS WITH SEVERE ACTIVE JUVENILE IDIOPATHIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, etanercept and tocilizumab for a patient who has severe active juvenile idiopathic arthritis. Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, etanercept and tocilizumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 3 biological medicines at any one time.

From 1 April 2014, a patient receiving PBS-subsidised biological medicine is considered to be in a treatment cycle where they may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements, within a single treatment cycle, a patient may:

- continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy; and
 - fail to respond or to sustain a response to each PBS-subsidised biological medicine once only. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.
- Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single treatment cycle and they must have, at a minimum, a 12 month break in PBS-subsidised biological medicine therapy before they are eligible to commence another cycle.

The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine was approved to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who received PBS-subsidised biological medicine immediately prior to 1 April 2014 is considered to be in their first cycle as of 1 April 2014. A patient who has had a break in biological medicine treatment of at least 12 months immediately prior to making a new application, on or after 1 April 2014, will commence a new treatment cycle under the Initial 3 treatment restriction.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of less than 12 months may commence a further course of treatment within the same treatment cycle under the Initial 2 treatment restriction.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of more than 12 months must commence a new treatment cycle under the Initial 3 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment therapy after 1 April 2014.

(1) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an alternate medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine therapy of less than 12 months) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 12 months with the same medicine (initial 2 - Change or Recommencement of treatment after a break in biological medicine therapy of less than 12 months).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 12 months (initial 3 - Recommencement of treatment after a break in biological medicine of more than 12 months). Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy.

A patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not conducted within these timeframes, the patient will be deemed to have failed this course of treatment.

For the second and subsequent cycles of PBS-subsidised biological medicine, it is recommended that a patient is reviewed in the 4 weeks prior to completing their current course of treatment and that an application is posted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services where applicable. Where a response assessment is not submitted, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (joint count) or the prior non-biological medicine therapy requirements, except if the patient has had a break in therapy of more than 12 months who would then need to requalify under the initial 3 restrictions with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application.

However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, an application for a patient who wishes to swap to an alternate biological medicine should be accompanied by the approved authority prescription or remaining repeats for the biological medicine the patient is ceasing.

(4) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count submitted with the first authority application for a biological medicine. However, prescribers may provide a new baseline measurement any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to the revised baseline measurement.

(5) Recommencement of treatment after a 12 months break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological medicine therapy of at least 12 months, must qualify under the initial 3 restriction and meet the relevant criteria and index of disease severity.

(6) Withdrawal of treatment after sustained remission.

Withdrawal of treatment with biological medicine should be considered in a patient who has achieved and sustained

complete remission of disease for 12 months. A demonstration of response to the current treatment should be submitted to the Department of Human Services at the time treatment is ceased.

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must be 30kg or over, **AND**
- Patient must not receive more than 24 weeks of treatment per continuing treatment course authorised under this restriction.

An adequate response to treatment is defined as:

- (a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or
- (b) a reduction in the number of the following active joints, from at least 4, by at least 50%:
 - (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - (ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

Determination of whether a response has been demonstrated to initial and subsequent courses of treatment will be based on the baseline measurement of joint count submitted with the initial treatment application.

The authority application must be made in writing and must include:

- (1) completed authority prescription form(s); and
- (2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either Initial 1, Initial 2, or Initial 3 treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 12 months have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

If a patient fails to respond to PBS-subsidised biological medicine treatment 3 times (once with each agent) they will not be eligible to receive further PBS-subsidised biological medicine therapy in this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Continuing treatment - balance of supply

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restriction.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

tocilizumab 162 mg/0.9 mL injection, 4 x 0.9 mL syringes

11720W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	811.83	41.30	Actemra Subcutaneous Injection [RO]

tocilizumab 162 mg/0.9 mL injection, 4 x 0.9 mL pen devices

11742B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	811.83	41.30	Actemra ACTPen [RO]

■ TOCILIZUMAB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note A patient may only qualify for PBS-subsidised treatment under this restriction once in a lifetime.

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Active giant cell arteritis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist, clinical immunologist or neurologist experienced in the management of giant cell arteritis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must not exceed 52 weeks in total including initial and continuing applications.

tocilizumab 162 mg/0.9 mL injection, 4 x 0.9 mL syringes

11721X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	6	..	811.83	41.30	Actemra Subcutaneous Injection [RO]

tocilizumab 162 mg/0.9 mL injection, 4 x 0.9 mL pen devices

11722Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	6	..	811.83	41.30	Actemra ACTPen [RO]

■ TOCILIZUMAB
Note TREATMENT OF ADULT PATIENTS WITH A HISTORY OF JUVENILE IDIOPATHIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, etanercept and tocilizumab for a patient over 18 years who has a history of juvenile idiopathic arthritis with onset prior to the age of 18 years. Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, etanercept and tocilizumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 3 biological medicines at any one time.

From 1 April 2014, a patient receiving PBS-subsidised biological medicine therapy is considered to be in a treatment cycle where they may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements, within a single treatment cycle, a patient may:

(i) continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy; and

(ii) fail to respond or to sustain a response to each PBS-subsidised biological medicine once only. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment.

Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single treatment cycle and they must have, at a minimum, a 5 year break in PBS-subsidised biological medicine therapy before they are eligible to receive further PBS-subsidised biological medicine therapy.

The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine was approved to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who received PBS-subsidised biological medicine treatment immediately prior to 1 April 2014 is considered to be in their first cycle as of 1 April 2014. A patient who has had a break in biological medicine treatment of at least 24 months immediately prior to making a new application, on or after 1 April 2014, will commence a new treatment cycle under the Initial 3 treatment restriction.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of less than 24 months may commence a further course of treatment within the same treatment cycle under the Initial 2 treatment restriction.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of more than 24 months must commence a new treatment cycle under the Initial 3 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine therapy after 1 April 2014.

(1) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy (Initial 1 - New patient); or
 - (ii) a patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an alternate medicine (Initial 2 - Change or recommencement of treatment after a break in biological medicine of less than 24 months) [further details are under 'Swapping therapy' below]; or
 - (iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 24 months with the same medicine (Initial 2 - Change or recommencement of treatment after a break in biological medicine therapy of less than 24 months); or
 - (iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 24 months (Initial 3 - recommencement of treatment after a break in biological medicine of more than 24 months).
- Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy.

A patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not conducted within these timeframes, the patient will be deemed to have failed this course of treatment.

For second and subsequent cycles of PBS-subsidised biological medicine, it is recommended that a patient is reviewed in the 4 weeks prior to completing their current course of treatment and that an application is posted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services where applicable. Where a response assessment is not submitted, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (joint count and ESR/CRP) or the prior non- biological medicine therapy requirements, except if the patient has had a break in therapy of more than 24 months who would then need to requalify under the Initial 3 restrictions with respect to the indices of disease severity.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, an application for a patient who wishes to swap to an alternate biological medicine should be accompanied by the approved authority prescription or remaining repeats for the biological medicine the patient is ceasing.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count submitted with the first authority application for a biological medicine. However, prescribers may provide a new baseline measurement any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to the revised baseline measurement.

(4) Recommencement of treatment after a 24 months break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological medicine therapy of at least 24 months, must qualify under the Initial 3 restriction and meet the relevant criteria and index of disease severity.

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment per continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

- (a) an active joint count of fewer than 10 active (swollen and tender) joints; or
- (b) a reduction in the active (swollen and tender) joint count by at least 50% from baseline; or
- (c) a reduction in the number of the following active joints, from at least 4, by at least 50%:
 - (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - (ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

- (1) completed authority prescription form(s); and
- (2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either Initial 1, Initial 2, or Initial 3 treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

If a patient fails to respond to PBS-subsidised biological medicine treatment 3 times (once with each agent) they will not be eligible to receive further PBS-subsidised biological medicine therapy in this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Continuing Treatment - balance of supply

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restriction.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

tocilizumab 162 mg/0.9 mL injection, 4 x 0.9 mL syringes

11730J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	811.83	41.30	Actemra Subcutaneous Injection [RO]

tocilizumab 162 mg/0.9 mL injection, 4 x 0.9 mL pen devices

11750K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	811.83	41.30	Actemra ACTPen [RO]

■ TOCILIZUMAB

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note A patient may only qualify for PBS-subsidised treatment under this restriction once in a lifetime.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Active giant cell arteritis

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a rheumatologist, clinical immunologist or neurologist experienced in the management of giant cell arteritis.

Clinical criteria:

- Patient must have clinical symptoms of active giant cell arteritis in the absence of any other identifiable cause, **AND**
- Patient must have an ESR equal to or greater than 30 mm/hour within the past 6 weeks; OR
- Patient must have a CRP equal to or greater than 10 mg/L within the past 6 weeks; OR
- Patient must have active giant cell arteritis confirmed by positive temporal artery biopsy or imaging, **AND**
- Patient must have had a history of an ESR equal to or greater than 50 mm/hour or a CRP equal to or greater than 24.5 mg/L at diagnosis, **AND**
- Patient must have had temporal artery biopsy revealing features of giant cell arteritis at diagnosis; OR
- Patient must have had evidence of large-vessel vasculitis by magnetic resonance (MR) or computed tomography (CT) angiography or PET/CT at diagnosis, **AND**
- The treatment must be in combination with a tapering course of corticosteroids, **AND**
- The treatment must not exceed 52 weeks in total including initial and continuing applications.

Population criteria:

- Patient must be aged 50 years or older.

Clinical symptoms of giant cell arteritis at diagnosis include unequivocal cranial symptoms of giant cell arteritis (new onset localized headache, scalp tenderness, temporal artery tenderness or decreased pulsation, ischemia related vision loss, or otherwise unexplained mouth or jaw pain upon mastication); or symptoms of polymyalgia rheumatica, defined as shoulder and/or hip girdle pain associated with inflammatory morning stiffness.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form;
- (2) a completed Giant Cell Arteritis - Supporting Information Form; and
- (3) documentation that the patient has active giant cell arteritis including pathology reports outlining the patient's ESR or CRP levels within the last 6 weeks, or positive temporal artery biopsy or imaging;
- (4) documentation that the patient has been diagnosed with giant cell arteritis with a history of an ESR equal to or greater than 50 mm/hour or a CRP equal to or greater than 24.5 mg/L at diagnosis.

tocilizumab 162 mg/0.9 mL injection, 4 x 0.9 mL syringes

11743C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	811.83	41.30	Actemra Subcutaneous Injection [RO]

tocilizumab 162 mg/0.9 mL injection, 4 x 0.9 mL pen devices

11744D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	811.83	41.30	Actemra ACTPen [RO]

■ TOCILIZUMAB

Caution Inadvertent muscular injection in patients aged less than 12 years may occur with the pen device.

Note TREATMENT OF PATIENTS WITH SEVERE ACTIVE SYSTEMIC JUVENILE IDIOPATHIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of tocilizumab for a patient who has severe active systemic juvenile idiopathic arthritis (sJIA).

From 1 May 2012, a patient receiving PBS-subsidised tocilizumab therapy is considered to be in a treatment cycle. Under these arrangements, within a single treatment cycle, a patient may:

- (i) continue to receive long-term treatment with PBS-subsidised tocilizumab while they continue to show an adequate response to therapy, and
- (ii) fail to respond, or to sustain a response, to PBS-subsidised tocilizumab twice.

Once a patient has either failed or ceased to sustain an adequate response to 2 courses of treatment, they are deemed to have completed a single treatment cycle and they must have, at a minimum, a 12 month break in PBS-subsidised tocilizumab therapy before they are eligible to commence another cycle. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised tocilizumab treatment was approved to the date of the first application for initial treatment with tocilizumab under the new treatment cycle.

A serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, will not be considered as a

treatment failure.

How to prescribe PBS-subsidised tocilizumab treatment therapy after 1 May 2012.

(1) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has not received prior PBS-subsidised tocilizumab treatment and wishes to commence such therapy (Initial 1 - New patient); or
- (ii) a patient has received the first course of PBS-subsidised (initial or continuing) tocilizumab therapy in a treatment cycle and is deemed to have failed to respond or sustain an adequate response and the treating physician wishes to trial a second course, provided any break in therapy is less than 12 months (Initial 2 - retreat or recommencement of treatment after a break of less than 12 months); or
- (iii) a patient wishes to recommence treatment with tocilizumab following a break in PBS-subsidised therapy of less than 12 months (Initial 2 - retreat or recommencement of treatment after a break of less than 12 months).
- (iv) a patient wishes to recommence treatment following a break in PBS-subsidised tocilizumab therapy of more than 12 months (Initial 3 - recommencement of a new treatment cycle after a break of more than 12 months).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy.

A patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not conducted within these timeframes, the patient will be deemed to have failed this course of treatment.

(2) Continuing treatment.

Following the completion of an initial treatment course with tocilizumab, a patient may qualify to receive up to 24 weeks of continuing treatment with this drug provided they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing tocilizumab treatment in courses of up to 24 weeks provided they continue to sustain the response. It is recommended that a patient be reviewed in the 4 weeks prior to completing their current course of treatment to ensure uninterrupted tocilizumab supply.

Assessments of response to a course of PBS-subsidised therapy must be conducted after a minimum of 12 weeks of treatment and no later than 4 weeks from the completion of treatment. Where a response assessment is not conducted within these timeframes, the patient will be deemed to have failed to respond to treatment with tocilizumab.

For the second and subsequent courses of PBS-subsidised tocilizumab, it is recommended that a patient be assessed for response in the 4 weeks prior to completing their current course of treatment and that an application is posted to Services Australia no later than 2 weeks prior to the patient completing their current treatment course.

(3) Treatment cycle.

Once initial treatment with PBS-subsidised tocilizumab is approved, a patient deemed to have failed to respond to the first course of treatment may have a second course without having to requalify with respect to the indices of disease severity (joint count, fever and/or CRP level and platelet count), except if the patient has had a break in therapy of more than 12 months.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

(4) Baseline measurements to determine response.

Whether an adequate response to treatment has been demonstrated or not will be based on the relative change from relevant baseline measurements of the joint count, fever and/or CRP level and platelet count submitted with the first authority application for tocilizumab.

Where a patient is deemed to have failed to respond or to sustain an adequate response to the first course of therapy in a treatment cycle, prescribers may provide new baseline measurements for the second course of treatment within that cycle. Assessment of response according to these revised baseline measurements may then occur. If new baseline measurements are not submitted with the initial application for the second course of treatment, then those submitted with the first course will be used to assess response to the second course.

(5) Recommencement of treatment after a 12 month break in PBS-subsidised therapy.

A patient who wishes to start a second or subsequent treatment cycle following a break in PBS-subsidised tocilizumab therapy of at least 12 months, must requalify for treatment under the Initial 3 treatment restriction.

(6) Withdrawal of treatment after sustained remission.

Withdrawal of treatment with tocilizumab should be considered in a patient who has achieved and sustained complete remission of disease for 12 months. A demonstration of response to the current treatment should be submitted to Services Australia at the time treatment is ceased.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Systemic juvenile idiopathic arthritis

Treatment Phase: Balance of supply for Initial treatment - Initial 1 (new patient) or Initial 2 (retreat or recommencement of treatment after a break of less than 12 months) or Initial 3 (recommencement of treatment after a break of more than 12 months) - in a patient of any weight being administered a subcutaneous form of this biological medicine

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (retreat or recommencement of treatment after a break of less than 12 months) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under Initial 3 (recommencement of treatment after a break of more than 12 months) restriction to complete 16 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 16 weeks therapy available under Initial 1, 2 or 3 treatment.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Authority required

Systemic juvenile idiopathic arthritis

Treatment Phase: Balance of supply - Continuing treatment in a patient of any weight being administered a subcutaneous form of this biological medicine

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks therapy available under Continuing treatment.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

tocilizumab 162 mg/0.9 mL injection, 4 x 0.9 mL syringes

12102Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	811.83	41.30	Actemra Subcutaneous Injection [RO]

tocilizumab 162 mg/0.9 mL injection, 4 x 0.9 mL pen devices

12094M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	811.83	41.30	Actemra ACTPen [RO]

■ TOCILIZUMAB**Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS**

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for adults with severe active rheumatoid arthritis. Where the term biological medicine appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alfa antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab), the T-cell co-stimulation modulator (abatacept) and the Janus kinase (JAK) inhibitors (baricitinib, tofacitinib, upadacitinib).

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised biological medicine therapy may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once, and
- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised biological medicines for the treatment of rheumatoid arthritis.

A serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Stevens Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered a treatment failure.

A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction.

A patient who has failed fewer than 5 biological medicines and who has a break in treatment of less than 24 months may commence a further course of treatment with a biological medicine under Initial 2 treatment restriction. A patient who has failed fewer than 5 biological medicines and who has had a break in therapy of longer than 24 months may commence a further course of treatment with a biological medicine under the Initial 3 treatment restriction. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine treatment is stopped to the date of the new application for treatment with a biological medicine.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 April 2019.

(a) Initial treatment.

Applications for initial treatment should be made where:

- a patient has received no prior PBS-subsidised biological medicine treatment and wishes to commence such therapy, excluding rituximab (Initial 1 - new patient); or
- a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months) [further details are under 'Swapping therapy' below]; or
- a patient wishes to re-commence treatment with a specific biological medicine following a break of less than 24 months in PBS-subsidised therapy with that agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months).
- a patient wishes to re-commence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 24 months (Initial 3 - re-commencement of treatment after a break in biological medicine of more than 24 months) Initial applications for a new patient (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6-month intensive DMARD trial, but prior to ceasing DMARD therapy.

Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, baricitinib, etanercept, golimumab, tocilizumab, tofacitinib and upadacitinib, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

Rituximab patients should be assessed following a minimum of 12 weeks after the first infusion, and the assessment should be submitted to Services Australia within 4 weeks to ensure continuity of treatment for those who meet the continuing

restriction for PBS subsidised treatment with this drug for this condition.

Where a response assessment is not provided the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment. For second and subsequent courses of PBS-subsidised biological medicine (excluding rituximab) treatment, it is recommended that a patient be reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to Services Australia within sufficient time to allow processing.

Abatacept patients:

A patient is eligible to receive one I.V. loading dose when commencing treatment with the subcutaneous formulation. Two prescriptions are required, the first prescription for the I.V. loading dose for sufficient vials for one dose based on the patient's weight with no repeats. The second prescription for the subcutaneous formulation, with a maximum quantity of 4 and up to 3 repeats, must be submitted with the initial application.

Rituximab patients:

A further application may be submitted to Services Australia within sufficient time to allow processing. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

Assessment of the patient's response to treatment must be conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the cessation of the most recent course of biological medicine therapy.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction. Where a response assessment is not conducted within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

(2) Swapping therapy

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non- biological medicine therapy requirements except if the patient has had a break in therapy of more than 24 months who would need to requalify with respect to the indices of disease severity. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each biological medicine trialled.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent, unless they have experienced a serious adverse reaction of a severity necessitating permanent treatment withdrawal.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug.

Abatacept:

A patient swapping from I.V. abatacept to subcutaneous abatacept will not be eligible for an I.V. loading dose when commencing treatment with the subcutaneous formulation.

Rituximab:

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF-alfa antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they be assessed for response to every course of treatment, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological medicine during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological medicine therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate biological medicine may do so without having to have any treatment-free period.

(3) Baseline measurements to determine response.

Determination of whether a response to treatment has been demonstrated must be based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and the eligibility for continuing treatment must be assessed according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response must be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be determined on the total number of major joints.

Applications under the Initial 1 treatment restriction for a new patient must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. The results must be no more than 4 weeks old at the time of application.

Applications under the Initial 3 treatment restriction for re-commencement of treatment after a break in biological medicine of more than 24 months must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment per continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient has either failed or ceased to respond to a PBS-subsidised biological medicine for this condition 5 times, they will not be eligible to receive further PBS-subsidised treatment with a biological medicine for this condition.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Continuing Treatment - balance of supply.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restriction.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

tocilizumab 162 mg/0.9 mL injection, 4 x 0.9 mL syringes

10954M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	811.83	41.30	Actemra Subcutaneous Injection [RO]

tocilizumab 162 mg/0.9 mL injection, 4 x 0.9 mL pen devices

11567T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	811.83	41.30	Actemra ACTPen [RO]

■ TOCILIZUMAB

Caution Inadvertent muscular injection in patients aged less than 12 years may occur with the pen device.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note TREATMENT OF PATIENTS WITH SEVERE ACTIVE SYSTEMIC JUVENILE IDIOPATHIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of tocilizumab for a patient who has severe active systemic juvenile idiopathic arthritis (sJIA).

From 1 May 2012, a patient receiving PBS-subsidised tocilizumab therapy is considered to be in a treatment cycle. Under these arrangements, within a single treatment cycle, a patient may:

(i) continue to receive long-term treatment with PBS-subsidised tocilizumab while they continue to show an adequate response to therapy, and

(ii) fail to respond, or to sustain a response, to PBS-subsidised tocilizumab twice.

Once a patient has either failed or ceased to sustain an adequate response to 2 courses of treatment, they are deemed to have completed a single treatment cycle and they must have, at a minimum, a 12 month break in PBS-subsidised tocilizumab therapy before they are eligible to commence another cycle. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised tocilizumab treatment was approved to the date of the first application for initial treatment with tocilizumab under the new treatment cycle.

A serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, will not be considered as a treatment failure.

How to prescribe PBS-subsidised tocilizumab treatment therapy after 1 May 2012.

(1) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised tocilizumab treatment and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient has received the first course of PBS-subsidised (initial or continuing) tocilizumab therapy in a treatment cycle and is deemed to have failed to respond or sustain an adequate response and the treating physician wishes to trial a second course, provided any break in therapy is less than 12 months (Initial 2 - retreat or recommencement of treatment after a break of less than 12 months); or

(iii) a patient wishes to recommence treatment with tocilizumab following a break in PBS-subsidised therapy of less than 12 months (Initial 2 - retreat or recommencement of treatment after a break of less than 12 months).

(iv) a patient wishes to recommence treatment following a break in PBS-subsidised tocilizumab therapy of more than 12 months (Initial 3 - recommencement of a new treatment cycle after a break of more than 12 months).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy.

A patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not conducted within these timeframes, the patient will be deemed to have failed this course of treatment.

(2) Continuing treatment.

Following the completion of an initial treatment course with tocilizumab, a patient may qualify to receive up to 24 weeks of continuing treatment with this drug provided they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing tocilizumab treatment in courses of up to 24 weeks provided they continue to sustain the response. It is recommended that a patient be reviewed in the 4 weeks prior to completing their current course of treatment to ensure uninterrupted tocilizumab supply.

Assessments of response to a course of PBS-subsidised therapy must be conducted after a minimum of 12 weeks of treatment and no later than 4 weeks from the completion of treatment. Where a response assessment is not conducted within these timeframes, the patient will be deemed to have failed to respond to treatment with tocilizumab.

For the second and subsequent courses of PBS-subsidised tocilizumab, it is recommended that a patient be assessed for response in the 4 weeks prior to completing their current course of treatment and that an application is posted to Services Australia no later than 2 weeks prior to the patient completing their current treatment course.

(3) Treatment cycle.

Once initial treatment with PBS-subsidised tocilizumab is approved, a patient deemed to have failed to respond to the first course of treatment may have a second course without having to requalify with respect to the indices of disease severity (joint count, fever and/or CRP level and platelet count), except if the patient has had a break in therapy of more than 12 months.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

(4) Baseline measurements to determine response.

Whether an adequate response to treatment has been demonstrated or not will be based on the relative change from relevant baseline measurements of the joint count, fever and/or CRP level and platelet count submitted with the first authority application for tocilizumab.

Where a patient is deemed to have failed to respond or to sustain an adequate response to the first course of therapy in a treatment cycle, prescribers may provide new baseline measurements for the second course of treatment within that cycle.

Assessment of response according to these revised baseline measurements may then occur. If new baseline measurements are not submitted with the initial application for the second course of treatment, then those submitted with the first course will be used to assess response to the second course.

(5) Recommencement of treatment after a 12 month break in PBS-subsidised therapy.

A patient who wishes to start a second or subsequent treatment cycle following a break in PBS-subsidised tocilizumab

therapy of at least 12 months, must requalify for treatment under the Initial 3 treatment restriction.

(6) Withdrawal of treatment after sustained remission.

Withdrawal of treatment with tocilizumab should be considered in a patient who has achieved and sustained complete remission of disease for 12 months. A demonstration of response to the current treatment should be submitted to Services Australia at the time treatment is ceased.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Systemic juvenile idiopathic arthritis

Treatment Phase: Continuing treatment in a patient weighing at least 30 kg

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

An adequate response to treatment is defined as:

(a) in a patient with polyarticular course disease:

(i) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(ii) a reduction in the number of the following major active joints, from at least 4, by at least 50%:

- elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

- shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

(b) in a patient with refractory systemic symptoms:

(i) absence of fever greater than 38 degrees Celsius in the preceding seven days; and/or

(ii) a reduction in the C-reactive protein (CRP) level and platelet count by at least 30% from baseline; and/or

(iii) a reduction in the dose of corticosteroid by at least 30% from baseline.

Determination of whether a response has been demonstrated to initial and subsequent courses of treatment will be based on the baseline measurements of disease severity submitted with the initial treatment application.

The authority application must be made in writing and must include:

(1) completed authority prescription form(s); and

(2) a completed Systemic Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form which includes baseline and current pathology reports detailing CRP and platelet count where appropriate.

The most recent systemic juvenile idiopathic arthritis assessment must be no more than 4 weeks old at the time of application.

The assessment of the patient's response to the most recent course of biological medicine must be conducted following a minimum of 12 weeks of treatment and no later than 4 weeks from the cessation of that treatment course. If the response assessment is not conducted within these timeframes, the patient will be deemed to have failed that most recent course of treatment in this treatment cycle.

The patient remains eligible to receive continuing treatment with the same biological medicine in courses of up to 24 weeks providing they continue to sustain an adequate response. It is recommended that a patient be reviewed in the month prior to completing their current course of treatment.

If a patient fails to demonstrate a response to 2 courses of treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition in the current treatment cycle. A serious adverse reaction of a severity requiring permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 12 months have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

tocilizumab 162 mg/0.9 mL injection, 4 x 0.9 mL syringes

12099T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	811.83	41.30	Actemra Subcutaneous Injection [RO]

tocilizumab 162 mg/0.9 mL injection, 4 x 0.9 mL pen devices

12084B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	811.83	41.30	Actemra ACTPen [RO]

■ TOCILIZUMAB

Caution Inadvertent muscular injection in patients aged less than 12 years may occur with the pen device.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note TREATMENT OF PATIENTS WITH SEVERE ACTIVE SYSTEMIC JUVENILE IDIOPATHIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of tocilizumab for a patient who has severe active systemic juvenile idiopathic arthritis (sJIA).

From 1 May 2012, a patient receiving PBS-subsidised tocilizumab therapy is considered to be in a treatment cycle. Under these arrangements, within a single treatment cycle, a patient may:

- (i) continue to receive long-term treatment with PBS-subsidised tocilizumab while they continue to show an adequate response to therapy, and
- (ii) fail to respond, or to sustain a response, to PBS-subsidised tocilizumab twice.

Once a patient has either failed or ceased to sustain an adequate response to 2 courses of treatment, they are deemed to have completed a single treatment cycle and they must have, at a minimum, a 12 month break in PBS-subsidised tocilizumab therapy before they are eligible to commence another cycle. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised tocilizumab treatment was approved to the date of the first application for initial treatment with tocilizumab under the new treatment cycle.

A serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, will not be considered as a treatment failure.

How to prescribe PBS-subsidised tocilizumab treatment therapy after 1 May 2012.

(1) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has not received prior PBS-subsidised tocilizumab treatment and wishes to commence such therapy (Initial 1 - New patient); or
- (ii) a patient has received the first course of PBS-subsidised (initial or continuing) tocilizumab therapy in a treatment cycle and is deemed to have failed to respond or sustain an adequate response and the treating physician wishes to trial a second course, provided any break in therapy is less than 12 months (Initial 2 - retreat or recommencement of treatment after a break of less than 12 months); or
- (iii) a patient wishes to recommence treatment with tocilizumab following a break in PBS-subsidised therapy of less than 12 months (Initial 2 - retreat or recommencement of treatment after a break of less than 12 months).
- (iv) a patient wishes to recommence treatment following a break in PBS-subsidised tocilizumab therapy of more than 12 months (Initial 3 - recommencement of a new treatment cycle after a break of more than 12 months).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy.

A patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not conducted within these timeframes, the patient will be deemed to have failed this course of treatment.

(2) Continuing treatment.

Following the completion of an initial treatment course with tocilizumab, a patient may qualify to receive up to 24 weeks of continuing treatment with this drug provided they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing tocilizumab treatment in courses of up to 24 weeks provided they continue to sustain the response. It is recommended that a patient be reviewed in the 4 weeks prior to completing their current course of treatment to ensure uninterrupted tocilizumab supply.

Assessments of response to a course of PBS-subsidised therapy must be conducted after a minimum of 12 weeks of treatment and no later than 4 weeks from the completion of treatment. Where a response assessment is not conducted within these timeframes, the patient will be deemed to have failed to respond to treatment with tocilizumab.

For the second and subsequent courses of PBS-subsidised tocilizumab, it is recommended that a patient be assessed for response in the 4 weeks prior to completing their current course of treatment and that an application is posted to Services Australia no later than 2 weeks prior to the patient completing their current treatment course.

(3) Treatment cycle.

Once initial treatment with PBS-subsidised tocilizumab is approved, a patient deemed to have failed to respond to the first course of treatment may have a second course without having to requalify with respect to the indices of disease severity (joint count, fever and/or CRP level and platelet count), except if the patient has had a break in therapy of more than 12 months.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

(4) Baseline measurements to determine response.

Whether an adequate response to treatment has been demonstrated or not will be based on the relative change from relevant baseline measurements of the joint count, fever and/or CRP level and platelet count submitted with the first authority application for tocilizumab.

Where a patient is deemed to have failed to respond or to sustain an adequate response to the first course of therapy in a treatment cycle, prescribers may provide new baseline measurements for the second course of treatment within that cycle. Assessment of response according to these revised baseline measurements may then occur. If new baseline measurements are not submitted with the initial application for the second course of treatment, then those submitted with the first course will be used to assess response to the second course.

(5) Recommencement of treatment after a 12 month break in PBS-subsidised therapy.

A patient who wishes to start a second or subsequent treatment cycle following a break in PBS-subsidised tocilizumab therapy of at least 12 months, must requalify for treatment under the Initial 3 treatment restriction.

(6) Withdrawal of treatment after sustained remission.

Withdrawal of treatment with tocilizumab should be considered in a patient who has achieved and sustained complete

remission of disease for 12 months. A demonstration of response to the current treatment should be submitted to Services Australia at the time treatment is ceased.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos. Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Systemic juvenile idiopathic arthritis

Treatment Phase: Continuing treatment in a patient weighing less than 30 kg

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

An adequate response to treatment is defined as:

(a) in a patient with polyarticular course disease:

(i) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(ii) a reduction in the number of the following major active joints, from at least 4, by at least 50%:

- elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

- shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

(b) in a patient with refractory systemic symptoms:

(i) absence of fever greater than 38 degrees Celsius in the preceding seven days; and/or

(ii) a reduction in the C-reactive protein (CRP) level and platelet count by at least 30% from baseline; and/or

(iii) a reduction in the dose of corticosteroid by at least 30% from baseline.

Determination of whether a response has been demonstrated to initial and subsequent courses of treatment will be based on the baseline measurements of disease severity submitted with the initial treatment application.

The authority application must be made in writing and must include:

(1) completed authority prescription form(s); and

(2) a completed Systemic Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form which includes baseline and current pathology reports detailing CRP and platelet count where appropriate.

The most recent systemic juvenile idiopathic arthritis assessment must be no more than 4 weeks old at the time of application.

The assessment of the patient's response to the most recent course of biological medicine must be conducted following a minimum of 12 weeks of treatment and no later than 4 weeks from the cessation of that treatment course. If the response assessment is not conducted within these timeframes, the patient will be deemed to have failed that most recent course of treatment in this treatment cycle.

The patient remains eligible to receive continuing treatment with the same biological medicine in courses of up to 24 weeks providing they continue to sustain an adequate response. It is recommended that a patient be reviewed in the month prior to completing their current course of treatment.

If a patient fails to demonstrate a response to 2 courses of treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition in the current treatment cycle. A serious adverse reaction of a severity requiring permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 12 months have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

tocilizumab 162 mg/0.9 mL injection, 4 x 0.9 mL syringes

12086D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	811.83	41.30	Actemra Subcutaneous Injection [RO]

tocilizumab 162 mg/0.9 mL injection, 4 x 0.9 mL pen devices

12090H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	811.83	41.30	Actemra ACTPen [RO]

■ TOCILIZUMAB

Note TREATMENT OF ADULT PATIENTS WITH A HISTORY OF JUVENILE IDIOPATHIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab,

etanercept and tocilizumab for a patient over 18 years who has a history of juvenile idiopathic arthritis with onset prior to the age of 18 years. Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, etanercept and tocilizumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 3 biological medicines at any one time.

From 1 April 2014, a patient receiving PBS-subsidised biological medicine therapy is considered to be in a treatment cycle where they may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements, within a single treatment cycle, a patient may:

- (i) continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy; and
- (ii) fail to respond or to sustain a response to each PBS-subsidised biological medicine once only. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment.

Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single treatment cycle and they must have, at a minimum, a 5 year break in PBS-subsidised biological medicine therapy before they are eligible to receive further PBS-subsidised biological medicine therapy.

The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine was approved to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who received PBS-subsidised biological medicine treatment immediately prior to 1 April 2014 is considered to be in their first cycle as of 1 April 2014. A patient who has had a break in biological medicine treatment of at least 24 months immediately prior to making a new application, on or after 1 April 2014, will commence a new treatment cycle under the Initial 3 treatment restriction.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of less than 24 months may commence a further course of treatment within the same treatment cycle under the Initial 2 treatment restriction.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of more than 24 months must commence a new treatment cycle under the Initial 3 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine therapy after 1 April 2014.

(1) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy (Initial 1 - New patient); or
- (ii) a patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an alternate medicine (Initial 2 - Change or recommencement of treatment after a break in biological medicine of less than 24 months) [further details are under 'Swapping therapy' below]; or
- (iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 24 months with the same medicine (Initial 2 - Change or recommencement of treatment after a break in biological medicine therapy of less than 24 months); or
- (iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 24 months (Initial 3 - recommencement of treatment after a break in biological medicine of more than 24 months).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy.

A patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not conducted within these timeframes, the patient will be deemed to have failed this course of treatment.

For second and subsequent cycles of PBS-subsidised biological medicine, it is recommended that a patient is reviewed in the 4 weeks prior to completing their current course of treatment and that an application is posted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services where applicable. Where a response assessment is not submitted, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (joint count and ESR/CRP) or the prior non- biological medicine therapy requirements, except if the patient has had a break in therapy of more than 24 months who would then need to requalify under the Initial 3 restrictions with respect to the indices of disease severity.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, an application for a patient who wishes to swap to an alternate biological medicine should be accompanied by the approved authority prescription or remaining repeats for the biological medicine the patient is ceasing.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count submitted with the first authority application for a biological medicine. However, prescribers may provide a new baseline measurement any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to the revised baseline measurement.

(4) Recommencement of treatment after a 24 months break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological medicine therapy of at least 24 months, must qualify under the Initial 3 restriction and meet the relevant criteria and index of disease severity.

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have a documented history of severe active juvenile idiopathic arthritis with onset prior to the age of 18 years, **AND**
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with disease modifying anti-rheumatic drugs (DMARDs) which must include at least 3 months continuous treatment with each of at least 2 DMARDs, one of which must be methotrexate at a dose of at least 20 mg weekly and one of which must be: (i) hydroxychloroquine at a dose of at least 200 mg daily; or (ii) leflunomide at a dose of at least 10 mg daily; or (iii) sulfasalazine at a dose of at least 2 g daily; OR
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs which, if methotrexate is contraindicated according to the Therapeutic Goods Administration (TGA)-approved Product Information or cannot be tolerated at a 20 mg weekly dose, must include at least 3 months continuous treatment with each of at least 2 of the following DMARDs: (i) hydroxychloroquine at a dose of at least 200 mg daily; and/or (ii) leflunomide at a dose of at least 10 mg daily; and/or (iii) sulfasalazine at a dose of at least 2 g daily; OR
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs which, if 3 or more of methotrexate, hydroxychloroquine, leflunomide and sulfasalazine are contraindicated according to the relevant TGA-approved Product Information or cannot be tolerated at the doses specified above, must include at least 3 months continuous treatment with each of at least 2 DMARDs, with one or more of the following DMARDs being used in place of the DMARDs which are contraindicated or not tolerated: (i) azathioprine at a dose of at least 1 mg/kg per day; and/or (ii) cyclosporin at a dose of at least 2 mg/kg/day; and/or (iii) sodium aurothiomalate at a dose of 50 mg weekly, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

If methotrexate is contraindicated according to the TGA-approved Product Information or cannot be tolerated at a 20 mg weekly dose, the application must include details of the contraindication or intolerance to methotrexate. The maximum tolerated dose of methotrexate must be documented in the application, if applicable.

The application must include details of the DMARDs trialed, their doses and duration of treatment, and all relevant contraindications and/or intolerances.

The requirement to trial at least 2 DMARDs for periods of at least 3 months each can be met using single agents sequentially or by using one or more combinations of DMARDs.

If the requirement to trial 6 months of intensive DMARD therapy with at least 2 DMARDs cannot be met because of contraindications and/or intolerances of a severity necessitating permanent treatment withdrawal to all of the DMARDs specified above, details of the contraindication or intolerance and dose for each DMARD must be provided in the authority application.

The following criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; **AND** either

(a) an active joint count of at least 20 active (swollen and tender) joints; or

(b) at least 4 active joints from the following list:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The joint count and ESR and/or CRP must be determined at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. All measures must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

The authority application must be made in writing and must include:

- (1) completed authority prescription form(s); and
- (2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

An assessment of a patient's response to an initial course of treatment must be conducted following a minimum of 12 weeks of therapy. An application for the continuing treatment must be accompanied with the assessment of response and submitted to the Department of Human Services no later than 4 weeks from the date of completion of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note The Department of Human Services website (www.humanservices.gov.au) has details of the toxicities, including severity, which will be accepted for the following purposes:

(a) exempting a patient from the requirement to undertake a minimum 3 month trial of methotrexate at a 20 mg weekly dose; (b) substituting azathioprine, cyclosporin or sodium aurothiomalate for another DMARD as part of the 6 month intensive DMARD trial;

(c) exempting a patient from the requirement for a 6 month trial of intensive DMARD therapy.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 24 months)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have a documented history of severe active juvenile idiopathic arthritis with onset prior to the age of 18 years, **AND**
- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

- (a) an active joint count of fewer than 10 active (swollen and tender) joints; or
- (b) a reduction in the active (swollen and tender) joint count by at least 50% from baseline; or
- (c) a reduction in the number of the following active joints, from at least 4, by at least 50%:
 - (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - (ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The authority application must be made in writing and must include:

- (1) completed authority prescription form(s); and
- (2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to change or recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient who fails to demonstrate a response to treatment with this drug under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug in this treatment cycle. A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the initial 3 treatment restriction.

If a patient fails to respond to PBS-subsidised biological medicine treatment 3 times (once with each agent) they will not be eligible to receive further PBS-subsidised biological medicine therapy in this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos. Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 3 (recommencement of treatment after a break in biological medicine of more than 24 months)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 24 months or more from the most recently approved PBS-subsidised biological medicine for this condition; OR
- Patient must not have received PBS-subsidised biological medicine for at least 5 years if they failed or ceased to respond to PBS-subsidised biological medicine treatment 3 times in their last treatment cycle, **AND**
- The condition must have an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; OR
- The condition must have a C-reactive protein (CRP) level greater than 15 mg per L, **AND**
- The condition must have either (a) a total active joint count of at least 20 active (swollen and tender) joints; or (b) at least 4 active major joints, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Active joints are defined as:

- (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
- (ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

All measures of joint count must be no more than 4 weeks old at the time of this application.

The authority application must be made in writing and must include:

- (1) completed authority prescription form(s); and
- (2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online

Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient) or Initial 2 (change or recommencement of treatment after break of less than 24 months) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 24 months) - balance of supply

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 24 months) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 24 months) to complete 16 weeks of treatment, **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

tocilizumab 162 mg/0.9 mL injection, 4 x 0.9 mL syringes

11741Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	811.83	41.30	Actemra Subcutaneous Injection [RO]

tocilizumab 162 mg/0.9 mL injection, 4 x 0.9 mL pen devices

11725D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	811.83	41.30	Actemra ACTPen [RO]

■ TOCILIZUMAB

Caution Inadvertent muscular injection in patients aged less than 12 years may occur with the pen device.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note TREATMENT OF PATIENTS WITH SEVERE ACTIVE SYSTEMIC JUVENILE IDIOPATHIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of tocilizumab for a patient who has severe active systemic juvenile idiopathic arthritis (sJIA).

From 1 May 2012, a patient receiving PBS-subsidised tocilizumab therapy is considered to be in a treatment cycle. Under these arrangements, within a single treatment cycle, a patient may:

- continue to receive long-term treatment with PBS-subsidised tocilizumab while they continue to show an adequate response to therapy, and
- fail to respond, or to sustain a response, to PBS-subsidised tocilizumab twice.

Once a patient has either failed or ceased to sustain an adequate response to 2 courses of treatment, they are deemed to have completed a single treatment cycle and they must have, at a minimum, a 12 month break in PBS-subsidised tocilizumab therapy before they are eligible to commence another cycle. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised tocilizumab treatment was approved to the date of the first application for initial treatment with tocilizumab under the new treatment cycle.

A serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, will not be considered as a treatment failure.

How to prescribe PBS-subsidised tocilizumab treatment therapy after 1 May 2012.

(1) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised tocilizumab treatment and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient has received the first course of PBS-subsidised (initial or continuing) tocilizumab therapy in a treatment cycle and is deemed to have failed to respond or sustain an adequate response and the treating physician wishes to trial a second course, provided any break in therapy is less than 12 months (Initial 2 - retreat or recommencement of treatment after a break of less than 12 months); or

(iii) a patient wishes to recommence treatment with tocilizumab following a break in PBS-subsidised therapy of less than 12 months (Initial 2 - retreat or recommencement of treatment after a break of less than 12 months).

(iv) a patient wishes to recommence treatment following a break in PBS-subsidised tocilizumab therapy of more than 12 months (Initial 3 - recommencement of a new treatment cycle after a break of more than 12 months).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy.

A patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks

of therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not conducted within these timeframes, the patient will be deemed to have failed this course of treatment.

(2) Continuing treatment.

Following the completion of an initial treatment course with tocilizumab, a patient may qualify to receive up to 24 weeks of continuing treatment with this drug provided they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing tocilizumab treatment in courses of up to 24 weeks provided they continue to sustain the response. It is recommended that a patient be reviewed in the 4 weeks prior to completing their current course of treatment to ensure uninterrupted tocilizumab supply.

Assessments of response to a course of PBS-subsidised therapy must be conducted after a minimum of 12 weeks of treatment and no later than 4 weeks from the completion of treatment. Where a response assessment is not conducted within these timeframes, the patient will be deemed to have failed to respond to treatment with tocilizumab.

For the second and subsequent courses of PBS-subsidised tocilizumab, it is recommended that a patient be assessed for response in the 4 weeks prior to completing their current course of treatment and that an application is posted to Services Australia no later than 2 weeks prior to the patient completing their current treatment course.

(3) Treatment cycle.

Once initial treatment with PBS-subsidised tocilizumab is approved, a patient deemed to have failed to respond to the first course of treatment may have a second course without having to requalify with respect to the indices of disease severity (joint count, fever and/or CRP level and platelet count), except if the patient has had a break in therapy of more than 12 months.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

(4) Baseline measurements to determine response.

Whether an adequate response to treatment has been demonstrated or not will be based on the relative change from relevant baseline measurements of the joint count, fever and/or CRP level and platelet count submitted with the first authority application for tocilizumab.

Where a patient is deemed to have failed to respond or to sustain an adequate response to the first course of therapy in a treatment cycle, prescribers may provide new baseline measurements for the second course of treatment within that cycle. Assessment of response according to these revised baseline measurements may then occur. If new baseline measurements are not submitted with the initial application for the second course of treatment, then those submitted with the first course will be used to assess response to the second course.

(5) Recommencement of treatment after a 12 month break in PBS-subsidised therapy.

A patient who wishes to start a second or subsequent treatment cycle following a break in PBS-subsidised tocilizumab therapy of at least 12 months, must requalify for treatment under the Initial 3 treatment restriction.

(6) Withdrawal of treatment after sustained remission.

Withdrawal of treatment with tocilizumab should be considered in a patient who has achieved and sustained complete remission of disease for 12 months. A demonstration of response to the current treatment should be submitted to Services Australia at the time treatment is ceased.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos. Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Systemic juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient weighing at least 30 kg)

Clinical criteria:

- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have polyarticular course disease which has failed to respond adequately to oral or parenteral methotrexate at a dose of at least 15 mg per square metre weekly, alone or in combination with oral or intra-articular corticosteroids, for a minimum of 3 months; OR
- Patient must have polyarticular course disease and have demonstrated severe intolerance of, or toxicity due to, methotrexate; OR
- Patient must have refractory systemic symptoms, demonstrated by an inability to decrease and maintain the dose of prednisolone (or equivalent) below 0.5 mg per kg per day following a minimum of 2 months of therapy, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be under 18 years of age.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

The following criteria indicate failure to achieve an adequate response to prior methotrexate therapy in a patient with polyarticular course disease and must be demonstrated in the patient at the time of the initial application:

- (a) an active joint count of at least 20 active (swollen and tender) joints; or
- (b) at least 4 active joints from the following list of major joints:
 - (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The following criteria indicate failure to achieve an adequate response to prior therapy in a patient with refractory systemic symptoms and must be demonstrated in the patient at the time of the initial application:

- (a) an active joint count of at least 2 active joints; and
- (b) persistent fever greater than 38 degrees Celsius for at least 5 out of 14 consecutive days; and/or
- (c) a C-reactive protein (CRP) level and platelet count above the upper limits of normal (ULN).

The baseline measurements of joint count, fever and/or CRP level and platelet count must be performed preferably whilst on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment.

The same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to determine response for all subsequent continuing treatments.

Severe intolerance to methotrexate is defined as intractable nausea and vomiting and general malaise unresponsive to manoeuvres, including reducing or omitting concomitant non-steroidal anti-inflammatory drugs (NSAIDs) on the day of methotrexate administration, use of folic acid supplementation, or administering the dose of methotrexate in 2 divided doses over 24 hours.

Toxicity due to methotrexate is defined as evidence of hepatotoxicity with repeated elevations of transaminases, bone marrow suppression temporally related to methotrexate use, pneumonitis, or serious sepsis.

If treatment with methotrexate alone or in combination with other treatments is contraindicated according to the relevant TGA-approved Product Information, details must be provided at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, details of this toxicity must be provided at the time of application.

The authority application must be made in writing and must include:

- (1) completed authority prescription form(s); and
- (2) a completed Systemic Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the date of assessment of severe active systemic juvenile idiopathic arthritis;
 - (ii) details of prior treatment including dose and duration of treatment;
 - (iii) pathology reports detailing CRP and platelet count where appropriate.

The assessment of the patient's response to the initial course of treatment must be conducted following a minimum of 12 weeks of treatment and no later than 4 weeks from the cessation of that treatment course. If the response assessment is not conducted within these timeframes, the patient will be deemed to have failed this course of treatment in this treatment cycle.

Authority required

Systemic juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 2 (retial or recommencement of treatment after a break of less than 12 months in a patient weighing at least 30 kg)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with this drug for this condition in the previous 12 months, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug more than once during the current treatment cycle, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be under 18 years of age.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

An adequate response to treatment is defined as:

(a) in a patient with polyarticular course disease:

(i) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(ii) a reduction in the number of the following major active joints, from at least 4, by at least 50%:

- elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

- shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

(b) in a patient with refractory systemic symptoms:

- (i) absence of fever greater than 38 degrees Celsius in the preceding seven days; and/or
- (ii) a reduction in the C-reactive protein (CRP) level and platelet count by at least 30% from baseline; and/or
- (iii) a reduction in the dose of corticosteroid by at least 30% from baseline.

The authority application must be made in writing and must include:

- (1) completed authority prescription form(s); and
- (2) a completed Systemic Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form which includes pathology reports detailing C-reactive protein (CRP) level and platelet count where appropriate.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to retial or recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

The assessment of the patient's response to the most recent course of biological medicine must be conducted following a minimum of 12 weeks of treatment and no later than 4 weeks from the cessation of that treatment course. If the response assessment is not conducted within these timeframes, the patient will be deemed to have failed that most recent course of treatment in this treatment cycle.

If a patient fails to demonstrate a response to 2 courses of treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition in the current treatment cycle. A serious adverse reaction of a severity requiring permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 12 months have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Authority required

Systemic juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 3 (recommencement of treatment after a break of more than 12 months in a patient weighing at least 30 kg)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have had a break in treatment of 12 months or more from this drug for this condition, **AND**
- Patient must have polyarticular course disease and the condition must have (a) an active joint count of at least 20 active (swollen and tender) joints; or (b) at least 4 active joints from the following list of major joints: i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or (ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth); OR
- Patient must have refractory systemic symptoms and the condition must have (a) an active joint count of at least 2 active joints; and (b) persistent fever greater than 38 degrees Celsius for at least 5 out of 14 consecutive days; and/or (c) a C-reactive protein (CRP) level and platelet count above the upper limits of normal (ULN), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Population criteria:

- Patient must be under 18 years of age.

The authority application must be made in writing and must include:

(1) completed authority prescription form(s); and

(2) a completed Systemic Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form which includes the following:

(i) the date of assessment of severe active systemic juvenile idiopathic arthritis;

(ii) pathology reports detailing C-reactive protein (CRP) level and platelet count where appropriate.

The most recent systemic juvenile idiopathic arthritis assessment must be no more than 4 weeks old at the time of application.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

The assessment of the patient's response to the most recent course of biological medicine must be conducted following a minimum of 12 weeks of treatment and no later than 4 weeks from the cessation of that treatment course. If the response assessment is not conducted within these timeframes, the patient will be deemed to have failed that most recent course of treatment in this treatment cycle.

If a patient fails to demonstrate a response to 2 courses of treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition in the current treatment cycle. A serious adverse reaction of a severity requiring permanent withdrawal of treatment is not considered as a treatment failure.

tocilizumab 162 mg/0.9 mL injection, 4 x 0.9 mL syringes

12095N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	811.83	41.30	Actemra Subcutaneous Injection [RO]

tocilizumab 162 mg/0.9 mL injection, 4 x 0.9 mL pen devices

12083Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	811.83	41.30	Actemra ACTPen [RO]

■ TOCILIZUMAB

Caution Inadvertent muscular injection in patients aged less than 12 years may occur with the pen device.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note TREATMENT OF PATIENTS WITH SEVERE ACTIVE SYSTEMIC JUVENILE IDIOPATHIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of tocilizumab for a patient who has severe active systemic juvenile idiopathic arthritis (sJIA).

From 1 May 2012, a patient receiving PBS-subsidised tocilizumab therapy is considered to be in a treatment cycle. Under these arrangements, within a single treatment cycle, a patient may:

- (i) continue to receive long-term treatment with PBS-subsidised tocilizumab while they continue to show an adequate response to therapy, and

(ii) fail to respond, or to sustain a response, to PBS-subsidised tocilizumab twice.

Once a patient has either failed or ceased to sustain an adequate response to 2 courses of treatment, they are deemed to have completed a single treatment cycle and they must have, at a minimum, a 12 month break in PBS-subsidised tocilizumab therapy before they are eligible to commence another cycle. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised tocilizumab treatment was approved to the date of the first application for initial treatment with tocilizumab under the new treatment cycle.

A serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, will not be considered as a treatment failure.

How to prescribe PBS-subsidised tocilizumab treatment therapy after 1 May 2012.

(1) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised tocilizumab treatment and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient has received the first course of PBS-subsidised (initial or continuing) tocilizumab therapy in a treatment cycle and is deemed to have failed to respond or sustain an adequate response and the treating physician wishes to trial a second course, provided any break in therapy is less than 12 months (Initial 2 - retreat or recommencement of treatment after a break of less than 12 months); or

(iii) a patient wishes to recommence treatment with tocilizumab following a break in PBS-subsidised therapy of less than 12 months (Initial 2 - retreat or recommencement of treatment after a break of less than 12 months).

(iv) a patient wishes to recommence treatment following a break in PBS-subsidised tocilizumab therapy of more than 12 months (Initial 3 - recommencement of a new treatment cycle after a break of more than 12 months).

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy.

A patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not conducted within these timeframes, the patient will be deemed to have failed this course of treatment.

(2) Continuing treatment.

Following the completion of an initial treatment course with tocilizumab, a patient may qualify to receive up to 24 weeks of continuing treatment with this drug provided they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing tocilizumab treatment in courses of up to 24 weeks provided they continue to sustain the response. It is recommended that a patient be reviewed in the 4 weeks prior to completing their current course of treatment to ensure uninterrupted tocilizumab supply.

Assessments of response to a course of PBS-subsidised therapy must be conducted after a minimum of 12 weeks of treatment and no later than 4 weeks from the completion of treatment. Where a response assessment is not conducted within these timeframes, the patient will be deemed to have failed to respond to treatment with tocilizumab.

For the second and subsequent courses of PBS-subsidised tocilizumab, it is recommended that a patient be assessed for response in the 4 weeks prior to completing their current course of treatment and that an application is posted to Services Australia no later than 2 weeks prior to the patient completing their current treatment course.

(3) Treatment cycle.

Once initial treatment with PBS-subsidised tocilizumab is approved, a patient deemed to have failed to respond to the first course of treatment may have a second course without having to requalify with respect to the indices of disease severity (joint count, fever and/or CRP level and platelet count), except if the patient has had a break in therapy of more than 12 months.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

(4) Baseline measurements to determine response.

Whether an adequate response to treatment has been demonstrated or not will be based on the relative change from relevant baseline measurements of the joint count, fever and/or CRP level and platelet count submitted with the first authority application for tocilizumab.

Where a patient is deemed to have failed to respond or to sustain an adequate response to the first course of therapy in a treatment cycle, prescribers may provide new baseline measurements for the second course of treatment within that cycle.

Assessment of response according to these revised baseline measurements may then occur. If new baseline measurements are not submitted with the initial application for the second course of treatment, then those submitted with the first course will be used to assess response to the second course.

(5) Recommencement of treatment after a 12 month break in PBS-subsidised therapy.

A patient who wishes to start a second or subsequent treatment cycle following a break in PBS-subsidised tocilizumab therapy of at least 12 months, must requalify for treatment under the Initial 3 treatment restriction.

(6) Withdrawal of treatment after sustained remission.

Withdrawal of treatment with tocilizumab should be considered in a patient who has achieved and sustained complete remission of disease for 12 months. A demonstration of response to the current treatment should be submitted to Services Australia at the time treatment is ceased.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Systemic juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient weighing less than 30 kg)

Clinical criteria:

- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have polyarticular course disease which has failed to respond adequately to oral or parenteral methotrexate at a dose of at least 15 mg per square metre weekly, alone or in combination with oral or intra-articular corticosteroids, for a minimum of 3 months; OR
- Patient must have polyarticular course disease and have demonstrated severe intolerance of, or toxicity due to, methotrexate; OR
- Patient must have refractory systemic symptoms, demonstrated by an inability to decrease and maintain the dose of prednisolone (or equivalent) below 0.5 mg per kg per day following a minimum of 2 months of therapy, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be under 18 years of age.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

The following criteria indicate failure to achieve an adequate response to prior methotrexate therapy in a patient with polyarticular course disease and must be demonstrated in the patient at the time of the initial application:

(a) an active joint count of at least 20 active (swollen and tender) joints; or

(b) at least 4 active joints from the following list of major joints:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The following criteria indicate failure to achieve an adequate response to prior therapy in a patient with refractory systemic symptoms and must be demonstrated in the patient at the time of the initial application:

(a) an active joint count of at least 2 active joints; and

(b) persistent fever greater than 38 degrees Celsius for at least 5 out of 14 consecutive days; and/or

(c) a C-reactive protein (CRP) level and platelet count above the upper limits of normal (ULN).

The baseline measurements of joint count, fever and/or CRP level and platelet count must be performed preferably whilst on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment.

The same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to determine response for all subsequent continuing treatments.

Severe intolerance to methotrexate is defined as intractable nausea and vomiting and general malaise unresponsive to manoeuvres, including reducing or omitting concomitant non-steroidal anti-inflammatory drugs (NSAIDs) on the day of methotrexate administration, use of folic acid supplementation, or administering the dose of methotrexate in 2 divided doses over 24 hours.

Toxicity due to methotrexate is defined as evidence of hepatotoxicity with repeated elevations of transaminases, bone marrow suppression temporally related to methotrexate use, pneumonitis, or serious sepsis.

If treatment with methotrexate alone or in combination with other treatments is contraindicated according to the relevant TGA-approved Product Information, details must be provided at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, details of this toxicity must be provided at the time of application.

The authority application must be made in writing and must include:

(1) completed authority prescription form(s); and

(2) a completed Systemic Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form which includes the following:

(i) the date of assessment of severe active systemic juvenile idiopathic arthritis;

(ii) details of prior treatment including dose and duration of treatment;

(iii) pathology reports detailing CRP and platelet count where appropriate.

The assessment of the patient's response to the initial course of treatment must be conducted following a minimum of 12 weeks of treatment and no later than 4 weeks from the cessation of that treatment course. If the response assessment is not conducted within these timeframes, the patient will be deemed to have failed this course of treatment in this treatment cycle.

Authority required

Systemic juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 2 (retial or recommencement of treatment after a break of less than 12 months in a patient weighing less than 30 kg)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with this drug for this condition in the previous 12 months, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug more than once during the current treatment cycle, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be under 18 years of age.

Treatment criteria:

- Must be treated by a rheumatologist; OR
 - Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.
- An adequate response to treatment is defined as:
- (a) in a patient with polyarticular course disease:
- (i) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or
- (ii) a reduction in the number of the following major active joints, from at least 4, by at least 50%:
- elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
 - shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).
- (b) in a patient with refractory systemic symptoms:
- (i) absence of fever greater than 38 degrees Celsius in the preceding seven days; and/or
- (ii) a reduction in the C-reactive protein (CRP) level and platelet count by at least 30% from baseline; and/or
- (iii) a reduction in the dose of corticosteroid by at least 30% from baseline.

The authority application must be made in writing and must include:

- (1) completed authority prescription form(s); and
- (2) a completed Systemic Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form which includes pathology reports detailing C-reactive protein (CRP) level and platelet count where appropriate.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to retreat or recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

The assessment of the patient's response to the most recent course of biological medicine must be conducted following a minimum of 12 weeks of treatment and no later than 4 weeks from the cessation of that treatment course. If the response assessment is not conducted within these timeframes, the patient will be deemed to have failed that most recent course of treatment in this treatment cycle.

If a patient fails to demonstrate a response to 2 courses of treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition in the current treatment cycle. A serious adverse reaction of a severity requiring permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 12 months have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Authority required

Systemic juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 3 (recommencement of a new treatment cycle after a break of more than 12 months in a patient weighing less than 30 kg)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must have had a break in treatment of 12 months or more from this drug for this condition, **AND**
- Patient must have polyarticular course disease and the condition must have (a) an active joint count of at least 20 active (swollen and tender) joints; or (b) at least 4 active joints from the following list of major joints: i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or (ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth); OR
- Patient must have refractory systemic symptoms and the condition must have (a) an active joint count of at least 2 active joints; and (b) persistent fever greater than 38 degrees Celsius for at least 5 out of 14 consecutive days; and/or (c) a C-reactive protein (CRP) level and platelet count above the upper limits of normal (ULN), **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Population criteria:

- Patient must be under 18 years of age.

The authority application must be made in writing and must include:

- (1) completed authority prescription form(s); and
- (2) a completed Systemic Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the date of assessment of severe active systemic juvenile idiopathic arthritis;
 - (ii) pathology reports detailing C-reactive protein (CRP) level and platelet count where appropriate.

The most recent systemic juvenile idiopathic arthritis assessment must be no more than 4 weeks old at the time of application.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

The assessment of the patient's response to the most recent course of biological medicine must be conducted following a minimum of 12 weeks of treatment and no later than 4 weeks from the cessation of that treatment course. If the response assessment is not conducted within these timeframes, the patient will be deemed to have failed that most recent course of treatment in this treatment cycle.

If a patient fails to demonstrate a response to 2 courses of treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition in the current treatment cycle. A serious adverse reaction of a severity requiring permanent withdrawal of treatment is not considered as a treatment failure.

tocilizumab 162 mg/0.9 mL injection, 4 x 0.9 mL syringes

12105D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	811.83	41.30	Actemra Subcutaneous Injection [RO]

tocilizumab 162 mg/0.9 mL injection, 4 x 0.9 mL pen devices

12085C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	811.83	41.30	Actemra ACTPen [RO]

■ TOCILIZUMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE RHEUMATOID ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for adults with severe active rheumatoid arthritis. Where the term biological medicine appears in the following notes and restrictions it refers to the tumour necrosis factor (TNF) alfa antagonists (adalimumab, certolizumab pegol, etanercept, golimumab, infliximab), the chimeric anti-CD20 monoclonal antibody (rituximab), the interleukin-6 inhibitor (tocilizumab), the T-cell co-stimulation modulator (abatacept) and the Janus kinase (JAK) inhibitors (baricitinib, tofacitinib, upadacitinib).

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

In order to be eligible to receive PBS-subsidised treatment with rituximab, a patient must have already failed to demonstrate a response to at least 1 course of treatment with a PBS-subsidised TNF-alfa antagonist.

A patient receiving PBS-subsidised biological medicine therapy may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements:

- a patient may continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy,
- a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once, and
- once a patient has either failed or ceased to respond to treatment 5 times, they will not be eligible to receive further PBS-subsidised biological medicines for the treatment of rheumatoid arthritis.

A serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Stevens Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered a treatment failure.

A patient whose most recent course of PBS-subsidised therapy was with rituximab and whose response to this treatment is sustained for more than 12 months, may apply for a further course of rituximab under the Continuing treatment restriction.

A patient who has failed fewer than 5 biological medicines and who has a break in treatment of less than 24 months may commence a further course of treatment with a biological medicine under Initial 2 treatment restriction. A patient who has failed fewer than 5 biological medicines and who has had a break in therapy of longer than 24 months may commence a further course of treatment with a biological medicine under the Initial 3 treatment restriction. The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine treatment is stopped to the date of the new application for treatment with a biological medicine.

(1) How to prescribe PBS-subsidised biological medicine therapy after 1 April 2019.

(a) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has received no prior PBS-subsidised biological medicine treatment and wishes to commence such therapy, excluding rituximab (Initial 1 - new patient); or
 - (ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine therapy and wishes to trial an alternate agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months) [further details are under 'Swapping therapy' below]; or
 - (iii) a patient wishes to re-commence treatment with a specific biological medicine following a break of less than 24 months in PBS-subsidised therapy with that agent (Initial 2 - change or re-commencement of treatment after a break in biological medicine of less than 24 months).
 - (iv) a patient wishes to re-commence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 24 months (Initial 3 - re-commencement of treatment after a break in biological medicine of more than 24 months)
- Initial applications for a new patient (Initial 1) must include a joint count and ESR and/or CRP measured at the completion of the 6-month intensive DMARD trial, but prior to ceasing DMARD therapy.

Initial treatment authorisations will be limited to provide a maximum of 16 weeks of therapy for abatacept, adalimumab, baricitinib, etanercept, golimumab, tocilizumab, tofacitinib and upadacitinib, 18 to 20 weeks of therapy with certolizumab pegol (depending upon the dosing regimen), 22 weeks of therapy for infliximab and 2 infusions of rituximab.

Rituximab patients should be assessed following a minimum of 12 weeks after the first infusion, and the assessment should be submitted to Services Australia within 4 weeks to ensure continuity of treatment for those who meet the continuing restriction for PBS subsidised treatment with this drug for this condition.

Where a response assessment is not provided the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment. For second and subsequent courses of PBS-subsidised biological medicine (excluding rituximab) treatment, it is recommended that a patient be reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to Services Australia within sufficient time to allow processing. Abatacept patients:

A patient is eligible to receive one I.V. loading dose when commencing treatment with the subcutaneous formulation. Two prescriptions are required, the first prescription for the I.V. loading dose for sufficient vials for one dose based on the patient's weight with no repeats. The second prescription for the subcutaneous formulation, with a maximum quantity of 4 and up to 3 repeats, must be submitted with the initial application.

Rituximab patients:

A further application may be submitted to Services Australia within sufficient time to allow processing. New baselines may be submitted with this application if appropriate.

(b) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine (excluding rituximab), a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

Assessment of the patient's response to treatment must be conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the cessation of the most recent course of biological medicine therapy.

Rituximab patients:

A patient may qualify to receive a further course of treatment (every 24 weeks) with this agent providing they have demonstrated an adequate response to treatment following a minimum of 12 weeks after the first infusion of their most recent treatment with rituximab. The patient remains eligible to receive a course of rituximab every 24 weeks providing they continue to demonstrate a response as specified in the restriction. Where a response assessment is not conducted within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine.

(2) Swapping therapy

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. the erythrocyte sedimentation rate (ESR), the C-reactive protein (CRP) levels and the joint count) or the prior non- biological medicine therapy requirements except if the patient has had a break in therapy of more than 24 months who would need to requalify with respect to the indices of disease severity. However the requirement for concomitant treatment with methotrexate, where it applies, must be met for each biological medicine trialled.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that agent, unless they have experienced a serious adverse reaction of a severity necessitating permanent treatment withdrawal.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug.

Abatacept:

A patient swapping from I.V. abatacept to subcutaneous abatacept will not be eligible for an I.V. loading dose when commencing treatment with the subcutaneous formulation.

Rituximab:

In order to trial rituximab, a patient must have trialled and failed to demonstrate a response to at least 1 PBS-subsidised TNF-alfa antagonist treatment.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they be assessed for response to every course of treatment, within the timeframes specified in the relevant restriction.

PBS subsidy does not allow for patients to receive treatment with another PBS-subsidised biological medicine during the required treatment-free period applying to patients who have demonstrated a response to their most recent course of rituximab. This means that patients who have demonstrated a response to a course of rituximab must have a PBS-subsidised biological medicine therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine. Patients who fail to respond to rituximab and who qualify and wish to trial a course of an alternate biological medicine may do so without having to have any treatment-free period.

(3) Baseline measurements to determine response.

Determination of whether a response to treatment has been demonstrated must be based on the baseline measurements of the joint count, ESR and/or CRP submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted and the eligibility for continuing treatment must be assessed according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used for all subsequent continuing treatment applications. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response must be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be determined on the total number of major joints.

Applications under the Initial 1 treatment restriction for a new patient must include a joint count and ESR and/or CRP measured at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. The results must be no more than 4 weeks old at the time of application.

Applications under the Initial 3 treatment restriction for re-commencement of treatment after a break in biological medicine of more than 24 months must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**

- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with disease modifying anti-rheumatic drugs (DMARDs) which must include at least 3 months continuous treatment with each of at least 2 DMARDs, one of which must be methotrexate at a dose of at least 20 mg weekly and one of which must be: (i) hydroxychloroquine at a dose of at least 200 mg daily; or (ii) leflunomide at a dose of at least 10 mg daily; or (iii) sulfasalazine at a dose of at least 2 g daily; OR
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs which, if methotrexate is contraindicated according to the Therapeutic Goods Administration (TGA)-approved Product Information or cannot be tolerated at a 20 mg weekly dose, must include at least 3 months continuous treatment with each of at least 2 of the following DMARDs: (i) hydroxychloroquine at a dose of at least 200 mg daily; and/or (ii) leflunomide at a dose of at least 10 mg daily; and/or (iii) sulfasalazine at a dose of at least 2 g daily; OR
- Patient must have failed, in the 24 months immediately prior to the date of the application, to achieve an adequate response to a trial of at least 6 months of intensive treatment with DMARDs which, if 3 or more of methotrexate, hydroxychloroquine, leflunomide and sulfasalazine are contraindicated according to the relevant TGA-approved Product Information or cannot be tolerated at the doses specified above, must include at least 3 months continuous treatment with each of at least 2 DMARDs, with one or more of the following DMARDs being used in place of the DMARDs which are contraindicated or not tolerated: (i) azathioprine at a dose of at least 1 mg/kg per day; and/or (ii) cyclosporin at a dose of at least 2 mg/kg/day; and/or (iii) sodium aurothiomalate at a dose of 50 mg weekly, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

If methotrexate is contraindicated according to the TGA-approved product information or cannot be tolerated at a 20 mg weekly dose, the application must include details of the contraindication or intolerance including severity to methotrexate. The maximum tolerated dose of methotrexate must be documented in the application, if applicable.

The application must include details of the DMARDs trialed, their doses and duration of treatment, and all relevant contraindications and/or intolerances including severity.

The requirement to trial at least 2 DMARDs for periods of at least 3 months each can be met using single agents sequentially or by using one or more combinations of DMARDs.

If the requirement to trial 6 months of intensive DMARD therapy with at least 2 DMARDs cannot be met because of contraindications and/or intolerances of a severity necessitating permanent treatment withdrawal to all of the DMARDs specified above, details of the contraindication or intolerance including severity and dose for each DMARD must be provided in the authority application.

The following criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; AND either

(a) a total active joint count of at least 20 active (swollen and tender) joints; or

(b) at least 4 active joints from the following list of major joints:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The joint count and ESR and/or CRP must be determined at the completion of the 6 month intensive DMARD trial, but prior to ceasing DMARD therapy. All measures must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note The Department of Human Services website (www.humanservices.gov.au) has details of the toxicities, including severity, which will be accepted for the following purposes:

- (a) exempting a patient from the requirement to undertake a minimum 3 month trial of methotrexate at a 20 mg weekly dose;
- (b) substituting azathioprine, cyclosporin or sodium aurothiomalate for another DMARD as part of the 6 month intensive DMARD trial;
- (c) exempting a patient from the requirement for a 6 month trial of intensive DMARD therapy.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 2 (change or re-commencement of treatment after a break in biological medicine of less than 24 months)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must not have failed to respond to previous PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised biological medicine treatment for this condition 5 times, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an ESR no greater than 25 mm per hour or a CRP level no greater than 15 mg per L or either marker reduced by at least 20% from baseline;

AND either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either of the Initial 1, Initial 2, Initial 3, or continuing treatment restrictions, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

A patient who has demonstrated a response to a course of rituximab must have a PBS-subsidised biological therapy treatment-free period of at least 22 weeks, immediately following the second infusion, before swapping to an alternate biological medicine.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial treatment - Initial 3 (re-commencement of treatment after a break in biological medicine of more than 24 months)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 24 months or more from the most recent PBS-subsidised biological medicine for this condition, **AND**
- Patient must not have failed to respond to previous PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised biological medicine treatment for this condition 5 times, **AND**
- The condition must have an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; OR
- The condition must have a C-reactive protein (CRP) level greater than 15 mg per L, **AND**
- The condition must have either (a) a total active joint count of at least 20 active (swollen and tender) joints; or (b) at least 4 active major joints, **AND**
- Patient must not receive more than 16 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Major joints are defined as (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

All measures of joint count and ESR and/or CRP must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Rheumatoid Arthritis PBS Authority Application - Supporting Information Form.

It is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

To demonstrate a response to treatment the application must be accompanied with the assessment of response from the most recent course of biological medicine therapy following a minimum of 12 weeks in therapy. It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not provided within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos. Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active rheumatoid arthritis

Treatment Phase: Initial 1 (new patient) or Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 24 months) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 24 months) - balance of supply

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of rheumatoid arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 16 weeks treatment; OR

- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 24 months) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 24 months) to complete 16 weeks of treatment, **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

tocilizumab 162 mg/0.9 mL injection, 4 x 0.9 mL syringes

10951J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	811.83	41.30	Actemra Subcutaneous Injection [RO]

tocilizumab 162 mg/0.9 mL injection, 4 x 0.9 mL pen devices

11565Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	811.83	41.30	Actemra ACTPen [RO]

■ TOCILIZUMAB

Note TREATMENT OF PATIENTS WITH SEVERE ACTIVE JUVENILE IDIOPATHIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of adalimumab, etanercept and tocilizumab for a patient who has severe active juvenile idiopathic arthritis. Where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, etanercept and tocilizumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the 3 biological medicines at any one time.

From 1 April 2014, a patient receiving PBS-subsidised biological medicine is considered to be in a treatment cycle where they may swap to an alternate biological medicine without having to experience a disease flare. Under these interchangeability arrangements, within a single treatment cycle, a patient may:

- continue to receive long-term treatment with a PBS-subsidised biological medicine while they continue to show a response to therapy; and
- fail to respond or to sustain a response to each PBS-subsidised biological medicine once only. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single treatment cycle and they must have, at a minimum, a 12 month break in PBS-subsidised biological medicine therapy before they are eligible to commence another cycle.

The length of a treatment break is measured from the date the most recent treatment with PBS-subsidised biological medicine was approved to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who received PBS-subsidised biological medicine immediately prior to 1 April 2014 is considered to be in their first cycle as of 1 April 2014. A patient who has had a break in biological medicine treatment of at least 12 months immediately prior to making a new application, on or after 1 April 2014, will commence a new treatment cycle under the Initial 3 treatment restriction.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of less than 12 months may commence a further course of treatment within the same treatment cycle under the Initial 2 treatment restriction.

A patient who has failed fewer than 3 trials of a biological medicine in a treatment cycle and who has a break in therapy of more than 12 months must commence a new treatment cycle under the Initial 3 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment therapy after 1 April 2014.

(1) Initial treatment.

Applications for initial treatment should be made where:

- a patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy (Initial 1 - New patient); or
- a patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an alternate medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine therapy of less than 12 months) [further details are under 'Swapping therapy' below]; or
- a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 12 months with the same medicine (initial 2 - Change or Recommencement of treatment after a break in biological medicine therapy of less than 12 months).
- a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 12 months (initial 3 - Recommencement of treatment after a break in biological medicine of more than 12 months)

Initial treatment authorisations will be limited to provide for a maximum of 16 weeks of therapy.

A patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not conducted within these timeframes, the patient will be deemed to have failed this course of treatment.

For the second and subsequent cycles of PBS-subsidised biological medicine, it is recommended that a patient is reviewed in the 4 weeks prior to completing their current course of treatment and that an application is posted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up

to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services where applicable. Where a response assessment is not submitted, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (joint count) or the prior non-biological medicine therapy requirements, except if the patient has had a break in therapy of more than 12 months who would then need to requalify under the initial 3 restrictions with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application.

However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under the interchangeability arrangements, it is important that they are assessed for response to every course of treatment approved, within the timeframes specified in the relevant restriction.

To avoid confusion, an application for a patient who wishes to swap to an alternate biological medicine should be accompanied by the approved authority prescription or remaining repeats for the biological medicine the patient is ceasing.

(4) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the joint count submitted with the first authority application for a biological medicine. However, prescribers may provide a new baseline measurement any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to the revised baseline measurement.

(5) Recommencement of treatment after a 12 months break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological medicine therapy of at least 12 months, must qualify under the initial 3 restriction and meet the relevant criteria and index of disease severity.

(6) Withdrawal of treatment after sustained remission.

Withdrawal of treatment with biological medicine should be considered in a patient who has achieved and sustained complete remission of disease for 12 months. A demonstration of response to the current treatment should be submitted to the Department of Human Services at the time treatment is ceased.

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Treatment criteria:

- Must be treated by a paediatric rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Clinical criteria:

- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have demonstrated severe intolerance of, or toxicity due to, methotrexate; OR
- Patient must have demonstrated failure to achieve an adequate response to 1 or more of the following treatment regimens: (i) oral or parenteral methotrexate at a dose of at least 20 mg per square metre weekly, alone or in combination with oral or intra-articular corticosteroids, for a minimum of 3 months; or (ii) oral methotrexate at a dose of at least 10 mg per square metre weekly together with at least 1 other disease modifying anti-rheumatic drug (DMARD), alone or in combination with corticosteroids, for a minimum of 3 months.

Population criteria:

- Patient must be under 18 years of age.

Severe intolerance to methotrexate is defined as intractable nausea and vomiting and general malaise unresponsive to manoeuvres, including reducing or omitting concomitant non-steroidal anti-inflammatory drugs (NSAIDs) on the day of methotrexate administration, use of folic acid supplementation, or administering the dose of methotrexate in 2 divided doses over 24 hours.

Toxicity due to methotrexate is defined as evidence of hepatotoxicity with repeated elevations of transaminases, bone marrow suppression temporally related to methotrexate use, pneumonitis, or serious sepsis.

If treatment with methotrexate alone or in combination with another DMARD is contraindicated according to the relevant TGA-approved Product Information, details must be provided at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, details of this toxicity must be provided at the time of application.

The following criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

- (a) an active joint count of at least 20 active (swollen and tender) joints; OR
- (b) at least 4 active joints from the following list:
 - (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The joint count assessment must be performed preferably whilst still on DMARD treatment, but no longer than 4 weeks following cessation of the most recent prior treatment.

The authority application must be made in writing and must include:

- (1) completed authority prescription form(s); and
- (2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

Patients under 30 kg may receive up to 24 weeks of treatment under this restriction. Patients 30 kg and over may receive up to 16 weeks of treatment under this restriction.

An assessment of a patient's response to an initial course of treatment must be conducted following a minimum of 12 weeks of therapy. An application for the continuing treatment must be accompanied with the assessment of response and submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Use of alternative DMARDs in children is dependent on approval by the Therapeutic Goods Administration as age restrictions may apply.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos. Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 12 months)

Treatment criteria:

- Must be treated by a paediatric rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle.

An adequate response to treatment is defined as:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

- (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
- (ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

Patients under 30 kg may receive up to 24 weeks of treatment under this restriction. Patients 30 kg and over may receive up to 16 weeks of treatment under this restriction.

The authority application must be made in writing and must include:

- (1) completed authority prescription form(s); and
- (2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to change or recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient who fails to demonstrate a response to treatment with this drug under this restriction will not be eligible to receive further PBS-subsidised treatment with this drug in this treatment cycle. A patient may re-trial this drug after a minimum of 12 months have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the initial 3 treatment restriction.

If a patient fails to respond to PBS-subsidised biological medicine treatment 3 times (once with each agent) they will not be eligible to receive further PBS-subsidised biological medicine therapy in this treatment cycle.

Note Use of alternative DMARDs in children is dependent on approval by the Therapeutic Goods Administration as age restrictions may apply.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 3 (recommencement of treatment after a break in biological medicine of more than 12 months)

Treatment criteria:

- Must be treated by a paediatric rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have had a break in treatment of 12 months or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must have either (a) a total active joint count of at least 20 active (swollen and tender) joints; or (b) at least 4 active major joints.

Active joints are defined as:

- (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or
- (ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

All measures of joint count must be no more than 4 weeks old at the time of this application.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of active joints, the response must be demonstrated on the total number of active joints.

Patients under 30 kg may receive up to 24 weeks of treatment under this restriction. Patients 30 kg and over may receive up to 16 weeks of treatment under this restriction.

The authority application must be made in writing and must include:

- (1) completed authority prescription form(s); and
- (2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient) or Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 12 months) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 12 months) - balance of supply

Treatment criteria:

- Must be treated by a paediatric rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the Initial 1 (new patient) restriction to complete 16 or 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 12 months) restriction to complete 16 or 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 12 months) restriction to complete 16 or 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 16 weeks treatment available under the above restrictions for patients 30 kg or over; OR
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restrictions for patients under 30 kg.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Severe active juvenile idiopathic arthritis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must be under 30kg, **AND**
- Patient must not receive more than 24 weeks of treatment per continuing treatment course authorised under this restriction.

An adequate response to treatment is defined as:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder, cervical spine and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

Determination of whether a response has been demonstrated to initial and subsequent courses of treatment will be based on the baseline measurement of joint count submitted with the initial treatment application.

The authority application must be made in writing and must include:

- (1) completed authority prescription form(s); and
- (2) a completed Juvenile Idiopathic Arthritis PBS Authority Application - Supporting Information Form.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either Initial 1, Initial 2, or Initial 3 treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 12 months have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

If a patient fails to respond to PBS-subsidised biological medicine treatment 3 times (once with each agent) they will not be eligible to receive further PBS-subsidised biological medicine therapy in this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe active juvenile idiopathic arthritis
Treatment Phase: Continuing treatment - balance of supply

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Patient must be undergoing treatment under the supervision of a paediatric rheumatology treatment centre.

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restriction.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

tocilizumab 162 mg/0.9 mL injection, 4 x 0.9 mL syringes

11748H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	811.83	41.30	Actemra Subcutaneous Injection [RO]

tocilizumab 162 mg/0.9 mL injection, 4 x 0.9 mL pen devices

11734N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	811.83	41.30	Actemra ACTPen [RO]

■ USTEKINUMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE PSORIATIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab for adult patients with severe active psoriatic arthritis. Therefore, where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient receiving PBS-subsidised treatment for psoriatic arthritis is able to commence a treatment cycle where they may trial biological medicines without having to experience a disease flare when swapping to the alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

A patient who received PBS-subsidised adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab or ustekinumab treatment prior to 1 May 2019 is considered to start their first cycle as of 1 May 2019.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment.

Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence another cycle [further details are under '(5) Recommencement of treatment after a 5-year break in PBS-subsidised therapy' below].

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe active psoriatic arthritis.

(1) Initial treatment.

Applications for initial treatment should be made where:

- (i) a patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy (Initial 1 - New patient); or
- (ii) a patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an

alternate medicine (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years) or An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, golimumab secukinumab and tofacitinib, 18 to 20 weeks of therapy for certolizumab pegol (depending upon the dosing regimen), 20 weeks of therapy for ixekizumab, 22 weeks of therapy for infliximab, and 28 weeks of therapy for ustekinumab. It is recommended that a patient be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of PBS-subsidised initial treatment following a minimum of 12 weeks of therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

Grandfather patients (ixekizumab only).

A patient who commenced treatment with ixekizumab for severe psoriatic arthritis prior to 1 March 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

Grandfather patients (tofacitinib only).

A patient who commenced treatment with Tofacitinib for severe psoriatic arthritis prior to 1 May 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment under the Initial 1 or Initial 2 treatment restrictions.

For the second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to re-qualify with respect to either the indices of disease severity (i.e. erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP) level, and active joint count) or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify under the Initial 3 treatment restriction with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application.

However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

Within a treatment cycle a patient may alternate between therapy with any biological medicine of their choice (1 at a time) providing:

(i) they have not received PBS-subsidised treatment with that particular biological medicine previously; or

(ii) they have demonstrated an adequate response to that particular biological medicine if they have previously trialled it on the PBS; and

(iii) they have not previously failed to respond to treatment 3 times in this treatment cycle with that biological medicine.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(4) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the indices of disease severity submitted with the first authority application for a biological

medicine. However, prescribers may provide new baseline measurements any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and these revised baseline measurements will be used to assess response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to determine response for all subsequent continuing treatments. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. 20 or more active joints), response will be determined according to a reduction in the total number of active joints. (5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment restriction according to the criteria of the relevant restriction and index of disease severity. The application must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Severe psoriatic arthritis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an erythrocyte sedimentation rate (ESR) no greater than 25 mm per hour or a C-reactive protein (CRP) level no greater than 15 mg per L or either marker reduced by at least 20% from baseline; and

either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following major active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to determine response for all subsequent continuing treatments.

The authority application must be made in writing and must include:

(1) a completed authority prescription form(s); and

(2) a completed Severe Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

Where the most recent course of PBS-subsidised treatment with this drug was approved under either Initial 1, Initial 2, or Initial 3 treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826
HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Continuing treatment - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the continuing treatment restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restriction.

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

ustekinumab 45 mg/0.5 mL injection, 0.5 mL vial

10767Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	3943.17	41.30	Stelara [JC]

▪ **USTEKINUMAB**

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

- a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or
- a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years); or
- a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or
- a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient be reviewed in the 4 weeks prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

Grandfather patients (risankizumab only).

A patient who commenced treatment with risankizumab for chronic plaque psoriasis prior to 1 December 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the Initial treatment Grandfather treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment. This assessment must be conducted within 4 weeks of the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the 4 weeks prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the 4 weeks prior to completing their current course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure patients receive the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

Services Australia will determine whether a response to treatment has been demonstrated based on the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 the PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Whole body

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**

- Patient must not receive more than 24 weeks of treatment per continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single injection. Up to a maximum of 1 repeat will be authorised.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos. Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Face, hand, foot

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have demonstrated an adequate response to treatment with this drug, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 24 weeks of treatment per continuing treatment course authorised under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

(i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or

(ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single injection. Up to a maximum of 1 repeat will be authorised.

The authority application must be made in writing and must include:

(a) a completed authority prescription form(s); and

(b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed Psoriasis Area and Severity Index (PASI) calculation sheet and face, hand, foot area diagrams including the date of the assessment of the patient's condition.

The most recent PASI assessment must be no more than 1 month old at the time of application.

Approval will be based on the PASI assessment of response to the most recent course of treatment with this drug.

The PASI assessment for continuing treatment must be performed on the same affected area assessed at baseline.

It is recommended that an application for the continuing treatment is submitted to the Department of Human Services no later than 1 month from the date of completion of the most recent course of treatment. This is to ensure continuity of treatment for those who meet the continuing restriction for PBS-subsidised treatment with this drug for this condition.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Continuing treatment, Whole body or Continuing treatment, Face, hand, foot - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the continuing treatment, Whole body restriction to complete 24 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the continuing treatment, Face, hand, foot restriction to complete 24 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 24 weeks treatment available under the above restrictions, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate).

Treatment criteria:

- Must be treated by a dermatologist.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

ustekinumab 45 mg/0.5 mL injection, 0.5 mL vial

9305R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	3943.17	41.30	Stelara [JC]

■ USTEKINUMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CROHN DISEASE

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines for adult patients with severe Crohn disease. Where the term 'biological medicine' appears in the following NOTES and restrictions, it refers to the tumour necrosis factor (TNF) alfa-antagonists (adalimumab and infliximab), the alpha-4 beta-7 integrin inhibitor (vedolizumab) and the human IgG1kappa monoclonal antibody (ustekinumab). Patients are eligible for PBS-subsidised treatment with only 1 of the above PBS-subsidised biological medicines at any one time.

From 1 September 2017, under the PBS, all patients will be able to commence a treatment cycle where they may trial PBS-subsidised a biological medicine without having to experience a disease flare when swapping to the alternate agent. Under these arrangements, within a single treatment cycle, a patient may continue to receive long-term treatment with a biological medicine while they continue to show a response to therapy.

A patient who received PBS-subsidised treatment with a biological medicine prior to 1 September 2017 is considered to have started their treatment cycle as of 1 September 2017.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once.

Once a patient has either failed or ceased to respond to treatment for this condition 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy for this condition before they are eligible to commence the next cycle. The 5-year break is measured from the date of the last approval for PBS-subsidised biological medicine treatment in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of less than 5 years, may commence a further course of treatment within the same treatment cycle.

A patient who has failed fewer than 3 trials of biological medicine therapy in a treatment cycle and who has a break in therapy of more than 5 years, may commence a new treatment cycle.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

(1) How to prescribe PBS-subsidised adalimumab, infliximab, vedolizumab or ustekinumab therapy after 1 September 2017.

(a) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has received no prior PBS-subsidised biological medicine treatment in this treatment cycle and wishes to commence such therapy Initial 1 (new patient); or
 (ii) a patient has received prior PBS-subsidised (initial or continuing) biological medicine and wishes to trial an alternate agent - Initial 2 (change or recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or
 (iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy with that agent - Initial 2 (change or recommencement of treatment after a break in therapy of less than 5 years); or
 (iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years - Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years).
 From 1 September 2017, a patient must be assessed for response to any course of initial PBS-subsidised treatment following a minimum of 12 weeks of therapy for adalimumab or ustekinumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab or vedolizumab, and this assessment must be submitted to the Department of Human Services no later than 4 weeks from the date that course was ceased.

Where a response assessment is not submitted to the Department of Human Services within these timeframes, the patient will be deemed to have failed to respond to treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

For subsequent courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the month prior to completing their current course of treatment and that where required an application is submitted to the Department of Human Services no later than 2 weeks prior to the patient completing their current treatment course.

Adalimumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be for the induction pack containing a quantity of 6 doses of 40 mg and no repeats and the second prescription should be written for 2 doses of 40 mg and 2 repeats for patients weighing 40 kg or greater. For patients weighing less than 40 kg, one prescription should be written for 2 doses of 40 mg with no repeats and the second prescription should be written for 2 doses of 20 mg with 3 repeats.

Ustekinumab only: Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be written under S100 (Highly Specialised Drugs) for a weight-based loading dose, containing a quantity of up to 4 vials of 130 mg and no repeats. The second prescription should be written under S85 (General) for the subsequent first dose, containing a quantity of 2 vials of 45 mg and no repeats.

(b) Continuing treatment.

Following the completion of an initial treatment course with a biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response.

It is recommended that a patient be reviewed in the month prior to completing their current course of treatment to ensure uninterrupted supply of treatment.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be conducted within 4 weeks of the last dose. Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(2) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine therapy is approved, a patient may swap if eligible to the alternate biological medicine within the same treatment cycle without having to requalify with respect to the indices of disease severity (i.e. Crohn Disease Activity Index (CDAI) Score, confirmation of Crohn disease), or the prior conventional therapies of corticosteroid therapy and immunosuppressive therapy.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug once within the same treatment cycle.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment, within the timeframes specified in the relevant restriction.

A patient who is not able to complete an initial treatment course for a biological medicine will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

(3) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the CDAI or evidence of intestinal inflammation submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial treatment authority application is submitted within a treatment cycle and the Department of Human Services will assess response according to these revised baseline measurements.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline must be used to assess response to all subsequent treatments.

(4) Recommencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to recommence treatment following a break in PBS-subsidised biological medicine therapy of at least 5 years, must requalify for initial treatment with respect to the indices of disease severity under the Initial 3 restriction. A re-trial of systemic therapy is not required.

Authority required

Severe Crohn disease

Treatment Phase: Initial treatment - Initial 1 (new patient)

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Population criteria:

- Patient must be aged 18 years or older.

Clinical criteria:

- Patient must have confirmed severe Crohn disease, defined by standard clinical, endoscopic and/or imaging features, including histological evidence, with the diagnosis confirmed by a gastroenterologist or a consultant physician, **AND**
- Patient must have failed to achieve an adequate response to prior systemic therapy with a tapered course of steroids, starting at a dose of at least 40 mg prednisolone (or equivalent), over a 6 week period, **AND**
- Patient must have failed to achieve adequate response to prior systemic immunosuppressive therapy with azathioprine at a dose of at least 2 mg per kg daily for 3 or more consecutive months; OR
- Patient must have failed to achieve adequate response to prior systemic immunosuppressive therapy with 6-mercaptopurine at a dose of at least 1 mg per kg daily for 3 or more consecutive months; OR
- Patient must have failed to achieve adequate response to prior systemic immunosuppressive therapy with methotrexate at a dose of at least 15 mg weekly for 3 or more consecutive months, **AND**
- The treatment must not exceed a total of 2 doses to be administered at weeks 0 and 8 under this restriction, **AND**
- Patient must have a Crohn Disease Activity Index (CDAI) Score greater than or equal to 300 as evidence of failure to achieve an adequate response to prior systemic therapy; OR
- Patient must have short gut syndrome with diagnostic imaging or surgical evidence, or have had an ileostomy or colostomy; and must have evidence of intestinal inflammation; and must have evidence of failure to achieve an adequate response to prior systemic therapy as specified below; OR
- Patient must have extensive intestinal inflammation affecting more than 50 cm of the small intestine as evidenced by radiological imaging; and must have a Crohn Disease Activity Index (CDAI) Score greater than or equal to 220; and must have evidence of failure to achieve an adequate response to prior systemic therapy as specified below.

Applications for authorisation must be made in writing and must include:

- (a) two completed authority prescription forms; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current Crohn Disease Activity Index (CDAI) calculation sheet including the date of assessment of the patient's condition if relevant; and
 - (ii) details of prior systemic drug therapy [dosage, date of commencement and duration of therapy]; and
 - (iii) the reports and dates of the pathology or diagnostic imaging test(s) nominated as the response criterion, if relevant; and
 - (iv) the date of the most recent clinical assessment.

Evidence of failure to achieve an adequate response to prior therapy must include at least one of the following:

- (a) patient must have evidence of intestinal inflammation;
- (b) patient must be assessed clinically as being in a high faecal output state;
- (c) patient must be assessed clinically as requiring surgery or total parenteral nutrition (TPN) as the next therapeutic option, in the absence of this drug, if affected by short gut syndrome, extensive small intestine disease or is an ostomy patient.

Evidence of intestinal inflammation includes:

- (i) blood: higher than normal platelet count, or, an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour, or, a C-reactive protein (CRP) level greater than 15 mg per L; or
- (ii) faeces: higher than normal lactoferrin or calprotectin level; or
- (iii) diagnostic imaging: demonstration of increased uptake of intravenous contrast with thickening of the bowel wall or mesenteric lymphadenopathy or fat streaking in the mesentery.

Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be written under S100 (Highly Specialised Drugs) for a weight-based loading dose, containing a quantity of up to 4 vials of 130 mg and no repeats. The second prescription should be written under S85 (General) for 2 vials of 45 mg and no repeats.

A maximum quantity of a weight based loading dose is up to 4 vials with no repeats and the subsequent first dose of 90 mg (2 vials of 45 mg) with no repeats provide for an initial 16 week course of this drug will be authorised.

Where fewer than 6 vials in total are requested at the time of the application, authority approvals for a sufficient number of vials based on the patient's weight to complete dosing at weeks 0 and 8 may be requested by telephone through the balance of supply restriction.

Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period.

All assessments, pathology tests and diagnostic imaging studies must be made within 1 month of the date of application and should be performed preferably whilst still on conventional treatment, but no longer than 1 month following cessation of the most recent prior treatment

If treatment with any of the specified prior conventional drugs is contraindicated according to the relevant TGA-approved Product Information, please provide details at the time of application.

If intolerance to treatment develops during the relevant period of use, which is of a severity necessitating permanent treatment withdrawal, details of this toxicity must be provided at the time of application.

Details of the accepted toxicities including severity can be found on the Department of Human Services website.

Any one of the baseline criteria may be used to determine response to an initial course of treatment and eligibility for continued therapy, according to the criteria included in the continuing treatment restriction. However, the same criterion must be used for any subsequent determination of response to treatment, for the purpose of eligibility for continuing PBS-subsidised therapy.

An assessment of a patient's response to an initial course of treatment must be conducted following a minimum of 12 weeks of therapy. An application for the continuing treatment must be accompanied with the assessment of response and submitted to the Department of Human Services no later than 4 weeks from the date of completion of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Increase in the maximum quantity or number of units up to 4 may be authorised for the purpose of weight-based loading dose.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos. Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe Crohn disease

Treatment Phase: Initial treatment - Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years)

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- The treatment must not exceed a total of 2 doses to be administered at weeks 0 and 8 under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Applications for authorisation must be made in writing and must include:

- (a) two completed authority prescription forms; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form, which includes the following:
 - (i) the completed Crohn Disease Activity Index (CDAI) Score calculation sheet including the date of the assessment of the patient's condition, if relevant; or
 - (ii) the reports and dates of the pathology or diagnostic imaging test(s) used to assess response to therapy for patients with short gut syndrome, extensive small intestine disease or an ostomy, if relevant; and
 - (iii) the date of clinical assessment; and
 - (iv) the details of prior biological medicine treatment including the details of date and duration of treatment.

Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be written under S100 (Highly Specialised Drugs) for a weight-based loading dose, containing a quantity of up to 4 vials of 130 mg and no repeats. The second prescription should be written under S85 (General) for 2 vials of 45 mg and no repeats.

A maximum quantity of a weight based loading dose is up to 4 vials with no repeats and the subsequent first dose of 90 mg (2 vials of 45 mg) with no repeats provide for an initial 16 week course of this drug will be authorised.

Where fewer than 6 vials in total are requested at the time of the application, authority approvals for a sufficient number of vials based on the patient's weight to complete dosing at weeks 0 and 8 may be requested by telephone through the balance of supply restriction.

Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period.

To demonstrate a response to treatment the application must be accompanied by the results of the most recent course of biological medicine therapy within the timeframes specified in the relevant restriction.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy for adalimumab or ustekinumab and up to 12 weeks after the first dose (6 weeks following the third dose) for infliximab and vedolizumab and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An assessment of a patient's response to an initial course of treatment must be conducted following a minimum of 12 weeks of therapy. An application for the continuing treatment must be accompanied with the assessment of response and submitted to the Department of Human Services no later than 4 weeks from the date of completion of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Increase in the maximum quantity or number of units up to 4 may be authorised for the purpose of weight-based loading dose.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe Crohn disease

Treatment Phase: Initial treatment - Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years)

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- Patient must have confirmed severe Crohn disease, defined by standard clinical, endoscopic and/or imaging features, including histological evidence, with the diagnosis confirmed by a gastroenterologist or a consultant physician, **AND**
- Patient must have a Crohn Disease Activity Index (CDAI) Score of greater than or equal to 300 that is no more than 4 weeks old at the time of application; OR
- Patient must have a documented history of intestinal inflammation and have diagnostic imaging or surgical evidence of short gut syndrome if affected by the syndrome or has an ileostomy or colostomy; OR
- Patient must have a documented history and radiological evidence of intestinal inflammation if the patient has extensive small intestinal disease affecting more than 50 cm of the small intestine, together with a Crohn Disease Activity Index (CDAI) Score greater than or equal to 220 and that is no more than 4 weeks old at the time of application, **AND**
- Patient must have evidence of intestinal inflammation; OR
- Patient must be assessed clinically as being in a high faecal output state; OR
- Patient must be assessed clinically as requiring surgery or total parenteral nutrition (TPN) as the next therapeutic option, in the absence of this drug, if affected by short gut syndrome, extensive small intestine disease or is an ostomy patient,

AND

- The treatment must not exceed a total of 2 doses to be administered at weeks 0 and 8 under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Applications for authorisation must be made in writing and must include:

- (a) two completed authority prescription forms; and
- (b) a completed Crohn Disease PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current Crohn Disease Activity Index (CDAI) calculation sheet including the date of assessment of the patient's condition if relevant; and
 - (ii) the reports and dates of the pathology or diagnostic imaging test(s) nominated as the response criterion, if relevant; and
 - (iii) the date of the most recent clinical assessment.

Evidence of intestinal inflammation includes:

- (i) blood: higher than normal platelet count, or, an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour, or, a C-reactive protein (CRP) level greater than 15 mg per L; or
- (ii) faeces: higher than normal lactoferrin or calprotectin level; or
- (iii) diagnostic imaging: demonstration of increased uptake of intravenous contrast with thickening of the bowel wall or mesenteric lymphadenopathy or fat streaking in the mesentery.

Two completed authority prescriptions should be submitted with every initial application for this drug. One prescription should be written under S100 (Highly Specialised Drugs) for a weight-based loading dose, containing a quantity of up to 4 vials of 130 mg and no repeats. The second prescription should be written under S85 (General) for 2 vials of 45 mg and no repeats.

A maximum quantity of a weight based loading dose is up to 4 vials with no repeats and the subsequent first dose of 90 mg (2 vials of 45 mg) with no repeats provide for an initial 16 week course of this drug will be authorised.

Where fewer than 6 vials in total are requested at the time of the application, authority approvals for a sufficient number of vials based on the patient's weight to complete dosing at weeks 0 and 8 may be requested by telephone through the balance of supply restriction.

Under no circumstances will telephone approvals be granted for initial authority applications, or for treatment that would otherwise extend the initial treatment period.

Any one of the baseline criteria may be used to determine response to an initial course of treatment and eligibility for continued therapy, according to the criteria included in the continuing treatment restriction. However, the same criterion

must be used for any subsequent determination of response to treatment, for the purpose of eligibility for continuing PBS-subsidised therapy.

An assessment of a patient's response to an initial course of treatment must be conducted following a minimum of 12 weeks of therapy. An application for the continuing treatment must be accompanied with the assessment of response and submitted to the Department of Human Services no later than 4 weeks from the date of completion of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Increase in the maximum quantity or number of units up to 4 may be authorised for the purpose of weight-based loading dose.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos. Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe Crohn disease

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have received this drug as their most recent course of PBS-subsidised biological medicine treatment for this condition, **AND**
- Patient must have an adequate response to this drug defined as a reduction in Crohn Disease Activity Index (CDAI) Score to a level no greater than 150 if assessed by CDAI or if affected by extensive small intestine disease; OR
- Patient must have an adequate response to this drug defined as (a) an improvement of intestinal inflammation as demonstrated by: (i) blood: normalisation of the platelet count, or an erythrocyte sedimentation rate (ESR) level no greater than 25 mm per hour, or a C-reactive protein (CRP) level no greater than 15 mg per L; or (ii) faeces: normalisation of lactoferrin or calprotectin level; or (iii) evidence of mucosal healing, as demonstrated by diagnostic imaging findings, compared to the baseline assessment; or (b) reversal of high faecal output state; or (c) avoidance of the need for surgery or total parenteral nutrition (TPN), if affected by short gut syndrome, extensive small intestine or is an ostomy patient, **AND**
- Patient must not receive more than 24 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Applications for authorisation must be made in writing and must include:

(a) a completed authority prescription form; and

(b) a completed Crohn Disease PBS Authority Application - Supporting Information Form which includes the following:

(i) the completed Crohn Disease Activity Index (CDAI) Score calculation sheet including the date of the assessment of the patient's condition, if relevant; or

(ii) the reports and dates of the pathology test or diagnostic imaging test(s) used to assess response to therapy for patients with short gut syndrome, extensive small intestine disease or an ostomy, if relevant; and

(iii) the date of clinical assessment.

All assessments, pathology tests, and diagnostic imaging studies must be made within 1 month of the date of application.

An application for continuing treatment with this drug must include a measurement of response to the most recent course of PBS-subsidised therapy. This assessment must be conducted no later than 4 weeks from the cessation of that treatment course. If the application is the first application for continuing treatment with this drug, it must be accompanied by an assessment of response to a minimum of 12 weeks of treatment with the initial treatment course.

The assessment of the patient's response to a continuing course of therapy must be made within the 4 weeks prior to completion of that course and posted to the Department of Human Services no less than 2 weeks prior to the date the next dose is scheduled, in order to ensure continuity of treatment for those patients who meet the continuation criterion.

Where an assessment is not submitted to the Department of Human Services within these timeframes, patients will be deemed to have failed to respond, or to have failed to sustain a response, to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Patients are eligible to receive continuing treatment with this drug in courses of up to 24 weeks providing they continue to sustain a response.

At the time of the authority application, medical practitioners should request the appropriate quantity and number of repeats; up to 1 repeat will be authorised for patients whose dosing frequency is every 12 weeks. Up to a maximum of 2 repeats will be authorised for patients whose dosing frequency is every 8 weeks.

Where an inadequate number of repeats are requested at the time of the application to complete a course of 24 weeks treatment, authority approvals for sufficient repeats to complete 24 weeks of treatment may be requested by telephone by contacting the Department of Human Services and applying through the Balance of Supply restriction. Under no circumstances will telephone approvals be granted for treatment that would otherwise extend continuing treatment beyond 24 months.

Note Increase in the maximum number of repeats of up to 2 may be authorised in patients whose dosing frequency is every 8 weeks.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe Crohn disease

Treatment Phase: Balance of supply

Treatment criteria:

- Must be treated by a gastroenterologist (code 87); OR
- Must be treated by a consultant physician [internal medicine specialising in gastroenterology (code 81)]; OR
- Must be treated by a consultant physician [general medicine specialising in gastroenterology (code 82)].

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1 (new patient) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 16 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the Continuing treatment restriction to complete 24 weeks of treatment, **AND**
- The treatment must provide no more than the balance of up to 16 weeks therapy available under Initial 1, 2 or 3 treatment; OR
- The treatment must provide no more than the balance of up to 24 weeks therapy available under Continuing treatment.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

ustekinumab 45 mg/0.5 mL injection, 0.5 mL vial

11178H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*7779.32	41.30	Stelara [JC]

■ USTEKINUMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE ACTIVE PSORIATIC ARTHRITIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab for adult patients with severe active psoriatic arthritis. Therefore, where the term 'biological medicine' appears in notes and restrictions, it refers to adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab, tofacitinib and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient receiving PBS-subsidised treatment for psoriatic arthritis is able to commence a treatment cycle where they may trial biological medicines without having to experience a disease flare when swapping to the alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

A patient who received PBS-subsidised adalimumab, certolizumab pegol, etanercept, golimumab, infliximab, ixekizumab, secukinumab or ustekinumab treatment prior to 1 May 2019 is considered to start their first cycle as of 1 May 2019.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological

medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a single cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence another cycle [further details are under '(5) Recommencement of treatment after a 5-year break in PBS-subsidised therapy' below].

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe active psoriatic arthritis.

(1) Initial treatment.

Applications for initial treatment should be made where:

(i) a patient has not received prior PBS-subsidised biological medicine treatment and wishes to commence such therapy (Initial 1 - New patient); or

(ii) a patient has received prior PBS-subsidised biological medicine therapy (initial or continuing) and wishes to trial an alternate medicine (Initial 2 - Change or Recommencement of treatment after a break in therapy of less than 5 years) [further details are under 'Swapping therapy' below]; or

(iii) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Recommencement of treatment after a break in biological medicine of less than 5 years).

(iv) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Recommencement of treatment after a break in biological medicine of more than 5 years) or An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, golimumab secukinumab and tofacitinib, 18 to 20 weeks of therapy for certolizumab pegol (depending upon the dosing regimen), 20 weeks of therapy for ixekizumab, 22 weeks of therapy for infliximab, and 28 weeks of therapy for ustekinumab. It is recommended that a patient be reviewed in the month prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of PBS-subsidised initial treatment following a minimum of 12 weeks of therapy and conducted no later than 4 weeks from the cessation of the treatment course. If the response assessment is not submitted within these timeframes, the patient will be deemed to have failed this course of treatment.

Grandfather patients (ixekizumab only).

A patient who commenced treatment with ixekizumab for severe psoriatic arthritis prior to 1 March 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

Grandfather patients (tofacitinib only).

A patient who commenced treatment with Tofacitinib for severe psoriatic arthritis prior to 1 May 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the 'Grandfather' treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment.

The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the month prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to the Department of Human Services where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment under the Initial 1 or Initial 2 treatment restrictions.

For the second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that an assessment of a patient's response is conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from the completion of the most recent course of treatment.

(3) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to re-qualify with respect to either the indices of disease severity (i.e. erythrocyte sedimentation rate (ESR) or C-reactive protein (CRP) level, and active joint count) or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify under the Initial 3 treatment restriction with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application.

However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that drug within the same treatment cycle.

Within a treatment cycle a patient may alternate between therapy with any biological medicine of their choice (1 at a time) providing:

- (i) they have not received PBS-subsidised treatment with that particular biological medicine previously; or
- (ii) they have demonstrated an adequate response to that particular biological medicine if they have previously trialled it on the PBS; and

- (iii) they have not previously failed to respond to treatment 3 times in this treatment cycle with that biological medicine.

To ensure a patient receives the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(4) Baseline measurements to determine response.

The Department of Human Services will determine whether a response to treatment has been demonstrated based on the baseline measurements of the indices of disease severity submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline measurements any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and these revised baseline measurements will be used to assess response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same indices of disease severity used to establish baseline at the commencement of treatment with each initial treatment application must be used to determine response for all subsequent continuing treatments. Therefore, where only an ESR or CRP level is provided at baseline, an ESR or CRP level respectively must be used to determine response. Similarly, where the baseline active joint count is based on total active joints (i.e. 20 or more active joints), response will be determined according to a reduction in the total number of active joints.

(5) Re-commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent course of treatment following a break in PBS-subsidised biological therapy of at least 5 years, must qualify under Initial 3 treatment restriction according to the criteria of the relevant restriction and index of disease severity. The application must include a joint count and ESR and/or CRP measurement that is no more than one month old at the time of application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response to methotrexate at a dose of at least 20 mg weekly for a minimum period of 3 months, **AND**
- Patient must have failed to achieve an adequate response to sulfasalazine at a dose of at least 2 g per day for a minimum period of 3 months; OR
- Patient must have failed to achieve an adequate response to leflunomide at a dose of up to 20 mg daily for a minimum period of 3 months, **AND**
- Patient must not receive more than 28 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Where treatment with methotrexate, sulfasalazine or leflunomide is contraindicated according to the relevant TGA-approved Product Information, details must be provided at the time of application.

Where intolerance to treatment with methotrexate, sulfasalazine or leflunomide developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

The following initiation criteria indicate failure to achieve an adequate response and must be demonstrated in all patients at the time of the initial application:

an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour or a C-reactive protein (CRP) level greater than 15 mg per L; and

either

- (a) an active joint count of at least 20 active (swollen and tender) joints; or
- (b) at least 4 active joints from the following list of major joints:
 - (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth). If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Severe Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

An assessment of a patient's response to an initial course of treatment must be conducted following a minimum of 12 weeks of therapy. An application for the continuing treatment must be accompanied with the assessment of response and submitted to the Department of Human Services no later than 4 weeks from the date of completion of the most recent course of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 3 months treatment with methotrexate and 3 months treatment with sulfasalazine or leflunomide can be found on the Department of Human Services website (www.humanservices.gov.au)

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial treatment - Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- Patient must not receive more than 28 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

An adequate response to treatment is defined as:

an erythrocyte sedimentation rate (ESR) no greater than 25 mm per hour or a C-reactive protein (CRP) level no greater than 15 mg per L or either marker reduced by at least 20% from baseline; and

either of the following:

(a) a reduction in the total active (swollen and tender) joint count by at least 50% from baseline, where baseline is at least 20 active joints; or

(b) a reduction in the number of the following major active joints, from at least 4, by at least 50%:

(i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or

(ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Severe Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to change or recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 3 months treatment with methotrexate and 3 months treatment with sulfasalazine or leflunomide can be found on the Department of Human Services website (www.humanservices.gov.au)

Note Any queries concerning the arrangements to prescribe may be directed to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Department of Human Services website at www.humanservices.gov.au

Applications for authority to prescribe should be forwarded to:

Department of Human Services

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial treatment - Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years)

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must have an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; OR
- The condition must have a C-reactive protein (CRP) level greater than 15 mg per L, **AND**
- The condition must have either (a) a total active joint count of at least 20 active (swollen and tender) joints; or (b) at least 4 active major joints, **AND**
- Patient must not receive more than 28 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Major joints are defined as (i) elbow, wrist, knee and/or ankle (assessed as swollen and tender); and/or (ii) shoulder and/or hip (assessed as pain in passive movement and restriction of passive movement, where pain and limitation of movement are due to active disease and not irreversible damage such as joint destruction or bony overgrowth).

All measures of joint count and ESR and/or CRP must be no more than one month old at the time of initial application.

If the above requirement to demonstrate an elevated ESR or CRP cannot be met, the application must state the reasons why this criterion cannot be satisfied.

Where the baseline active joint count is based on total active joints (i.e. more than 20 active joints), response will be determined according to the reduction in the total number of active joints. Where the baseline is determined on total number of major joints, the response must be demonstrated on the total number of major joints. If only an ESR or CRP level is provided with the initial application, the same marker will be used to determine response.

The authority application must be made in writing and must include:

- (1) a completed authority prescription form(s); and
- (2) a completed Severe Psoriatic Arthritis PBS Authority Application - Supporting Information Form.

An application for a patient who has received PBS-subsidised biological medicine treatment for this condition who wishes to recommence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised biological medicine treatment, within the timeframes specified below.

Where the most recent course of PBS-subsidised biological medicine treatment was approved under either Initial 1, Initial 2, Initial 3 or continuing treatment restrictions, an assessment of a patient's response must have been conducted following a minimum of 12 weeks of therapy and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment.

An application for the continuing treatment must be accompanied with the assessment of response following a minimum of 12 weeks of therapy with this drug and submitted to the Department of Human Services no later than 4 weeks from the date of completion of treatment. This will enable ongoing treatment for those who meet the continuing restriction for PBS-subsidised treatment.

Where the response assessment is not submitted within this timeframe, the patient will be deemed to have failed to respond to treatment with this drug.

If a patient fails to demonstrate a response to treatment with this drug they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment is not considered as a treatment failure.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe psoriatic arthritis

Treatment Phase: Initial treatment - Initial 1 (new patient), Initial 2 (change or recommencement of treatment after a break in medicine of less than 5 years) or Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years) - balance of supply

Treatment criteria:

- Must be treated by a rheumatologist; OR
- Must be treated by a clinical immunologist with expertise in the management of psoriatic arthritis.

Clinical criteria:

- Patient must have received insufficient therapy with this drug under the Initial 1 (new patient) restriction to complete 28 weeks treatment; OR
- Patient must have received insufficient therapy with this drug under the Initial 2 (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 28 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3 (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 28 weeks treatment, **AND**
- The treatment must provide no more than the balance of up to 28 weeks treatment available under the above restrictions.

Note Authority approval for sufficient therapy to complete the balance of supply may be requested by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

ustekinumab 45 mg/0.5 mL injection, 0.5 mL vial

10774C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	3943.17	41.30	Stelara [JC]

■ USTEKINUMAB

Note TREATMENT OF ADULT PATIENTS WITH SEVERE CHRONIC PLAQUE PSORIASIS

The following information applies to the prescribing under the Pharmaceutical Benefits Scheme (PBS) of the biological medicines adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab for adult patients with severe chronic plaque psoriasis. Therefore, where the term 'biological medicines' appears in notes and restrictions, it refers to adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab only.

A patient is eligible for PBS-subsidised treatment with only 1 of the above biological medicines at any 1 time.

A patient who received PBS-subsidised adalimumab, etanercept, guselkumab, infliximab, ixekizumab, risankizumab, secukinumab, tildrakizumab and ustekinumab treatment prior to 1 February 2019 is considered to start their first cycle as of 1 February 2019.

A patient receiving PBS-subsidised treatment for chronic plaque psoriasis is able to commence a 'treatment cycle', where they may trial biological medicines without having to experience a disease flare, when swapping to an alternate biological medicine. Under these arrangements, within a single cycle, a patient may receive long-term treatment with a biological medicine as long as they sustain a response to therapy.

Within the same treatment cycle, a patient cannot trial and fail, or cease to respond to, the same PBS-subsidised biological medicine more than once. Therefore, once a patient fails to meet the response criteria for a PBS-subsidised biological medicine, they must change to an alternate biological medicine if they wish to continue PBS-subsidised biological treatment. Serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment, including serious infusion or injection related reactions, Steven's Johnson Syndrome, development of a demyelinating lesion, progressive multifocal leukoencephalopathy and malignancy related to treatment with the biological medicine, is not considered as a treatment failure.

A patient must be assessed for response to each course of treatment according to the criteria included in the relevant continuing treatment restriction.

Once a patient has either failed or ceased to sustain a response to treatment 3 times, they are deemed to have completed a treatment cycle and they must have, at a minimum, a 5-year break in PBS-subsidised biological medicine therapy before they are eligible to commence the next cycle.

The duration of the break in therapy will be measured from the date the last prescription for PBS-subsidised treatment was approved in the most recent cycle to the date of the first application for initial treatment with a biological medicine under the new cycle.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of more than 5 years may commence a new treatment cycle under Initial 3 treatment restriction.

A patient who has failed fewer than 3 biological medicines in a treatment cycle and who has a break in therapy of less than 5 years may commence a further course of treatment within the same treatment cycle under Initial 2 treatment restriction.

There is no limit to the number of treatment cycles a patient may undertake in their lifetime.

How to prescribe PBS-subsidised biological medicine treatment for severe chronic plaque psoriasis.

There are separate restrictions for both the initial and continuing treatment for psoriasis affecting the whole body, versus psoriasis affecting the face, hands and feet.

(1) Initial treatment.

An application for initial treatment should be made where:

- (i) a patient has not received prior PBS-subsidised biological medicine treatment for this condition and wishes to commence such therapy (Initial 1 - New patient); or
- (ii) a patient wishes to recommence treatment with a biological medicine following a break in PBS-subsidised therapy of more than 5 years (Initial 3 - Re commencement of treatment after a break in biological medicine of more than 5 years); or
- (iii) a patient who has received prior PBS-subsidised biological medicine therapy for this condition (initial or continuing) and wishes to trial an alternate biological medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years) [further details are under (4) 'Swapping therapy' below]; or
- (iv) a patient wishes to recommence treatment with a specific biological medicine following a break in PBS-subsidised therapy of less than 5 years with the same medicine (Initial 2 - Change or Re-commencement of treatment after a break in biological medicine of less than 5 years).

An application for initial treatment will be limited to provide for a maximum of 16 weeks of therapy for adalimumab, etanercept, ixekizumab and secukinumab, 20 weeks of therapy for guselkumab, 22 weeks of therapy for infliximab and 28 weeks of therapy for risankizumab, tildrakizumab and ustekinumab.

It is recommended that a patient be reviewed in the 4 weeks prior to completing their course of initial treatment to ensure uninterrupted biological medicine supply.

Grandfather patients (risankizumab only).

A patient who commenced treatment with risankizumab for chronic plaque psoriasis prior to 1 December 2019 and who continues to receive treatment at the time of application, may qualify for treatment under the Initial treatment Grandfather treatment restriction.

A patient may only qualify for PBS-subsidised treatment under this restriction once. A maximum of 24 weeks of treatment will be authorised under this criterion. Following completion of the initial PBS-subsidised course, further subsidised treatment must be prescribed under the continuing treatment restriction of the relevant drug. 'Grandfather' arrangements will only apply for the first treatment cycle.

For the second and subsequent cycles, a 'grandfather' patient must qualify for continuing treatment under the criteria that apply to a continuing patient.

(2) Assessment of response to initial treatment.

When prescribing initial treatment with a biological medicine, it is recommended that a PASI assessment is conducted after at least 12 weeks of treatment. This assessment must be conducted within 4 weeks of the completion of this initial treatment course.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

(3) Continuing treatment.

Following the completion of an initial treatment course with a specific biological medicine, a patient may qualify to receive up to 24 weeks of continuing treatment with that drug providing they have demonstrated an adequate response to treatment. The patient remains eligible to receive continuing biological medicine treatment with the same drug in courses of up to 24 weeks providing they continue to sustain the response. It is recommended that a patient be reviewed the 4 weeks prior to completing their current course of treatment to ensure uninterrupted biological medicine supply.

A patient must be assessed for response to a course of continuing therapy, and the assessment must be submitted to Services Australia where applicable. Where a response assessment is not submitted where applicable, the patient will be deemed to have failed to respond to treatment with that biological medicine, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

Infliximab and etanercept only:

For the first continuing treatment course of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed for response following a minimum of 12 weeks of therapy under the Initial 1 or Initial 2 treatment restrictions.

For second and subsequent continuing courses of PBS-subsidised biological medicine treatment, it is recommended that a patient is reviewed in the 4 weeks prior to completing their current course of treatment.

(4) Swapping therapy.

Once initial treatment with the first PBS-subsidised biological medicine is approved, a patient may swap to an alternate biological medicine without having to requalify with respect to the indices of disease severity (i.e. a PASI score of greater than 15), or the prior non-biological therapy requirements, except if the patient has had a break in therapy of more than 5 years who would need to requalify with respect to the indices of disease severity.

A patient who is not able to complete a minimum of 12 weeks of an initial treatment course will be deemed to have failed treatment with that biological medicine unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

A patient may trial an alternate biological medicine at any time, regardless of whether they are receiving therapy (initial or continuing) with a biological medicine at the time of the application. However, they cannot swap to a particular biological medicine if they have failed to respond to prior treatment with that particular biological medicine within the same cycle or have experienced treatment failure with that particular biological medicine that required permanent treatment withdrawal. To ensure patients receive the maximum treatment opportunities allowed under these arrangements, it is important that they are assessed for response to every course of treatment.

(5) Baseline measurements to determine response.

Services Australia will determine whether a response to treatment has been demonstrated based on the baseline PASI assessment submitted with the first authority application for a biological medicine. However, prescribers may provide new baseline PASI assessments any time that an initial or change or recommencement treatment application is submitted within a treatment cycle and this revised baseline PASI score will be used to assess the patient's response to the PBS-subsidised treatment.

To ensure consistency in determining response, the same body area assessed at the baseline PASI assessment must be assessed for demonstration of response to treatment for the purposes of all continuing treatments.

(6) Re commencement of treatment after a 5-year break in PBS-subsidised therapy.

A patient who wishes to trial a second or subsequent treatment cycle following a break in PBS-subsidised biological therapy

of at least 5 years, must qualify under Initial 3 treatment according to the criteria of the relevant restriction and index of disease severity. A PASI assessment must be conducted prior to application and patient must have a PASI score of greater than 15 for whole body. For the face, hand, foot at least 2 of the 3 the PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. The PASI assessment must be no older than 4 weeks at the time of application.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Whole body (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis where lesions have been present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 5 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; (iii) ciclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks; (v) apremilast at a dose of 30 mg twice a day for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 28 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, ciclosporin, apremilast or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application.

Where intolerance to treatment with phototherapy, methotrexate, ciclosporin, apremilast or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, ciclosporin, apremilast, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

- (a) A current Psoriasis Area and Severity Index (PASI) score of greater than 15, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment.
- (b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 4 weeks following cessation of each course of treatment.
- (c) The most recent PASI assessment must be no more than 4 weeks old at the time of application.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and
 - (ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single injection. Up to a maximum of 2 repeats will be authorised.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, ciclosporin, apremilast or acitretin can be found on the Services Australia website (www.servicesaustralia.gov.au).

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Whole body (change or re-commencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 28 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as:

A Psoriasis Area and Severity Index (PASI) score which is reduced by 75% or more, or is sustained at this level, when compared with the baseline value for this treatment cycle.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single injection. Up to a maximum of 2 repeats will be authorised.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition; and
 - (ii) details of prior biological treatment, including dosage, date and duration of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 3, Whole body (re-commencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**

- The condition must have a current Psoriasis Area and Severity Index (PASI) score of greater than 15, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 28 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single injection. Up to a maximum of 2 repeats will be authorised.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets including the dates of assessment of the patient's condition.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos. Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Face, hand, foot (new patient)

Clinical criteria:

- Patient must have severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis, **AND**
- Patient must not have received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have failed to achieve an adequate response, as demonstrated by a Psoriasis Area and Severity Index (PASI) assessment, to at least 2 of the following 5 treatments: (i) phototherapy (UVB or PUVA) for 3 treatments per week for at least 6 weeks; (ii) methotrexate at a dose of at least 10 mg weekly for at least 6 weeks; (iii) ciclosporin at a dose of at least 2 mg per kg per day for at least 6 weeks; (iv) acitretin at a dose of at least 0.4 mg per kg per day for at least 6 weeks; (v) apremilast at a dose of 30 mg twice a day for at least 6 weeks, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 28 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

Where treatment with methotrexate, ciclosporin, apremilast or acitretin is contraindicated according to the relevant TGA-approved Product Information, or where phototherapy is contraindicated, details must be provided at the time of application.

Where intolerance to treatment with phototherapy, methotrexate, ciclosporin, apremilast or acitretin developed during the relevant period of use, which was of a severity to necessitate permanent treatment withdrawal, details of the degree of this toxicity must be provided at the time of application.

Regardless of if a patient has a contraindication to treatment with either methotrexate, ciclosporin, apremilast, acitretin or phototherapy, the patient is still required to trial 2 of these prior therapies until a failure to achieve an adequate response is met.

The following criterion indicates failure to achieve an adequate response to prior treatment and must be demonstrated in the patient at the time of the application:

(a) Chronic plaque psoriasis classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:

(i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment; or

- (ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed, preferably whilst still on treatment, but no longer than 4 weeks following cessation of the most recent prior treatment;
- (b) A PASI assessment must be completed for each prior treatment course, preferably whilst still on treatment, but no longer than 4 weeks following cessation of each course of treatment.

(c) The most recent PASI assessment must be no more than 4 weeks old at the time of application.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single injection. Up to a maximum of 2 repeats will be authorised.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current and previous Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition; and
 - (ii) details of previous phototherapy and systemic drug therapy [dosage (where applicable), date of commencement and duration of therapy].

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Details of the toxicities, including severity, which will be accepted as a reason for exempting a patient from the requirement for 6 weeks treatment with phototherapy, methotrexate, ciclosporin, apremilast or acitretin can be found on the Services Australia website (www.servicesaustralia.gov.au).

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 2, Face, hand, foot (change or re-commencement of treatment after a break in biological medicine of less than 5 years)

Clinical criteria:

- Patient must have received prior PBS-subsidised treatment with a biological medicine for this condition in this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with 3 biological medicines for this condition within this treatment cycle, **AND**
- Patient must not have already failed, or ceased to respond to, PBS-subsidised treatment with this drug for this condition during the current treatment cycle, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 28 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

An adequate response to treatment is defined as the plaque or plaques assessed prior to biological treatment showing:

- (i) a reduction in the Psoriasis Area and Severity Index (PASI) symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the baseline values; or
- (ii) a reduction by 75% or more in the skin area affected, or sustained at this level, as compared to the baseline value for this treatment cycle.

An application for a patient who has received PBS-subsidised treatment with this drug and who wishes to re-commence therapy with this drug, must be accompanied by evidence of a response to the patient's most recent course of PBS-subsidised treatment with this drug, within the timeframes specified below.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks

from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single injection. Up to a maximum of 2 repeats will be authorised.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the following:
 - (i) the completed current Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition; and
 - (ii) details of prior biological treatment, including dosage, date and duration of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

A patient may re-trial this drug after a minimum of 5 years have elapsed between the date the last prescription for a PBS-subsidised biological medicine was approved in this cycle and the date of the first application under a new cycle under the Initial 3 treatment restriction.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial 3, Face, hand, foot (re-commencement of treatment after a break in biological medicine of more than 5 years)

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with a biological medicine for this condition, **AND**
- Patient must have a break in treatment of 5 years or more from the most recently approved PBS-subsidised biological medicine for this condition, **AND**
- The condition must be classified as severe due to a plaque or plaques on the face, palm of a hand or sole of a foot where:
 - (i) at least 2 of the 3 Psoriasis Area and Severity Index (PASI) symptom subscores for erythema, thickness and scaling are rated as severe or very severe; or
 - (ii) the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- Patient must not receive more than 28 weeks of treatment under this restriction.

Population criteria:

- Patient must be aged 18 years or older.

Treatment criteria:

- Must be treated by a dermatologist.

The most recent PASI assessment must be no more than 4 weeks old at the time of application.

At the time of the authority application, medical practitioners should request the appropriate number of vials, based on the weight of the patient, to provide sufficient for a single injection. Up to a maximum of 2 repeats will be authorised.

The authority application must be made in writing and must include:

- (a) a completed authority prescription form(s); and
- (b) a completed Severe Chronic Plaque Psoriasis PBS Authority Application - Supporting Information Form which includes the completed current Psoriasis Area and Severity Index (PASI) calculation sheets and face, hand, foot area diagrams including the dates of assessment of the patient's condition.

To demonstrate a response to treatment the application must be accompanied with the assessment of response, conducted following a minimum of 12 weeks of therapy and no later than 4 weeks from cessation of the most recent course of biological medicine. It is recommended that an application for the continuing treatment be submitted no later than 4 weeks from the date of completion of the most recent course of treatment. This is to ensure treatment continuity for those who meet the continuing restriction.

The PASI assessment for continuing treatment must be performed on the same affected area as assessed at baseline.

Where a response assessment is not conducted within the required timeframe, the patient will be deemed to have failed to respond to treatment with this drug, unless the patient has experienced a serious adverse reaction of a severity resulting in the necessity for permanent withdrawal of treatment.

If a patient fails to demonstrate a response to treatment with this drug under this restriction they will not be eligible to receive further PBS-subsidised treatment with this drug for this condition within this treatment cycle.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
 Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
 Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
 Or mailed to:
 Services Australia
 Complex Drugs
 Reply Paid 9826
 HOBART TAS 7001

Authority required

Severe chronic plaque psoriasis

Treatment Phase: Initial treatment - Initial 1, Whole body or Face, hand, foot (new patient) or Initial 2, Whole body or Face, hand, foot (change or re-commencement of treatment after a break in biological medicine of less than 5 years) or Initial 3, Whole body or Face, hand, foot (re-commencement of treatment after a break in biological medicine of more than 5 years) - balance of supply

Clinical criteria:

- Patient must have received insufficient therapy with this drug for this condition under the Initial 1, Whole body (new patient) restriction to complete 28 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2, Whole body (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 28 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3, Whole body (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 28 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 1, Face, hand, foot (new patient) restriction to complete 28 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 2, Face, hand, foot (change or recommencement of treatment after a break in biological medicine of less than 5 years) restriction to complete 28 weeks treatment; OR
- Patient must have received insufficient therapy with this drug for this condition under the Initial 3, Face, hand, foot (recommencement of treatment after a break in biological medicine of more than 5 years) restriction to complete 28 weeks treatment, **AND**
- The treatment must be as systemic monotherapy (other than methotrexate), **AND**
- The treatment must provide no more than the balance of up to 28 weeks treatment available under the above restriction.

Treatment criteria:

- Must be treated by a dermatologist.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

ustekinumab 45 mg/0.5 mL injection, 0.5 mL vial

9304Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	3943.17	41.30	Stelara [JC]

Calcineurin inhibitors**■ CICLOSPORIN**

Caution Careful monitoring of patients is mandatory.

ciclosporin 10 mg capsule, 60

8657P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	3	..	*92.02	41.30	Neoral 10 [NV]

ciclosporin 25 mg capsule, 30

8658Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	3	..	*81.30	41.30	^a Cyclosporin Sandoz [SZ]	^a Neoral 25 [NV]

ciclosporin 100 mg/mL oral liquid, 50 mL

8661W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	3	..	*720.06	41.30	Neoral [NV]

ciclosporin 50 mg capsule, 30

8659R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	3	..	*158.38	41.30	^a Cyclosporin Sandoz [SZ]	^a Neoral 50 [NV]

ciclosporin 100 mg capsule, 30

8660T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	3	..	*315.44	41.30	^a Cyclosporin Sandoz [SZ]	^a Neoral 100 [NV]

■ TACROLIMUS

Caution Careful monitoring of patients is mandatory.

tacrolimus 5 mg modified release capsule, 30

5451X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	430.63	41.30	ADVAGRAF XL [LQ]

tacrolimus 500 microgram modified release capsule, 30

5299X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	52.36	41.30	ADVAGRAF XL [LQ]

tacrolimus 500 microgram capsule, 100

8646C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	3	..	130.87	41.30	^a Pacrolim [AF]	^a Pharmacor Tacrolimus 0.5 [CR]
						^a Prograf [LL]	^a Tacrograf [RW]
						^a TACROLIMUS APOTEX [TX]	^a Tacrolimus Sandoz [SZ]

tacrolimus 750 microgram capsule, 100

10870D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	218.70	41.30	Tacrolimus Sandoz [SZ]

tacrolimus 2 mg capsule, 100

10871E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	571.50	41.30	Tacrolimus Sandoz [SZ]

tacrolimus 1 mg modified release capsule, 60

5300Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	155.64	41.30	ADVAGRAF XL [LQ]

tacrolimus 3 mg modified release capsule, 50

11914C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	378.56	41.30	ADVAGRAF XL [LQ]

tacrolimus 1 mg capsule, 100

8647D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	3	..	254.72	41.30	^a Pacrolim [AF]	^a Pharmacor Tacrolimus 1 [CR]
						^a Prograf [LL]	^a Tacrograf [RW]
						^a TACROLIMUS APOTEX [TX]	^a Tacrolimus Sandoz [SZ]

tacrolimus 5 mg capsule, 50

8648E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	3	..	625.94	41.30	^a Pacrolim [AF]	^a Pharmacor Tacrolimus 5 [CR]
						^a Prograf [LL]	^a Tacrograf [RW]
						^a TACROLIMUS APOTEX [TX]	^a Tacrolimus Sandoz [SZ]

Other immunosuppressants

■ AZATHIOPRINE

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

azathioprine 25 mg tablet, 100

2688L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	24.18	25.47	^a APO-Azathioprine [TX]	^a Azathioprine GH [GQ]
						^a Azathioprine Sandoz [SZ]	^a Imuran [AS]

azathioprine 50 mg tablet, 100

2687K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	32.73	34.02	^a APO-Azathioprine [TX]	^a Azapin [RW]
						^a Azathioprine AN [EA]	^a Azathioprine GH [GQ]
						^a Azathioprine Sandoz [SZ]	^a Imazan [ER]
						^a Imuran [AS]	^a Thioprine 50 [AF]

■ DIMETHYL FUMARATE

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

10139

Multiple sclerosis

Treatment Phase: Continuing treatment

Clinical criteria:

- The condition must be diagnosed as clinically definite relapsing-remitting multiple sclerosis by magnetic resonance imaging of the brain and/or spinal cord; OR
- The condition must be diagnosed as clinically definite relapsing-remitting multiple sclerosis by accompanying written certification provided by a radiologist that a magnetic resonance imaging scan is contraindicated because of the risk of physical (not psychological) injury to the patient, **AND**
- The treatment must be the sole PBS-subsidised disease modifying therapy for this condition, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must not show continuing progression of disability while on treatment with this drug.

Where applicable, the date of the magnetic resonance imaging scan must be recorded in the patient's medical records.

dimethyl fumarate 240 mg enteric capsule, 56

2966D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1291.74	41.30	Tecfidera [BD]

dimethyl fumarate 120 mg enteric capsule, 14

2943X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*649.90	41.30	Tecfidera [BD]

▪ **DIMETHYL FUMARATE**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required (STREAMLINED)

10140

Multiple sclerosis

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be diagnosed as clinically definite relapsing-remitting multiple sclerosis by magnetic resonance imaging of the brain and/or spinal cord; OR
- The condition must be diagnosed as clinically definite relapsing-remitting multiple sclerosis by accompanying written certification provided by a radiologist that a magnetic resonance imaging scan is contraindicated because of the risk of physical (not psychological) injury to the patient, **AND**
- The treatment must be the sole PBS-subsidised disease modifying therapy for this condition, **AND**
- Patient must have experienced at least 2 documented attacks of neurological dysfunction, believed to be due to multiple sclerosis, in the preceding 2 years of commencing a PBS-subsidised disease modifying therapy for this condition, **AND**
- Patient must be ambulatory (without assistance or support).

Where applicable, the date of the magnetic resonance imaging scan must be recorded in the patient's medical records.

dimethyl fumarate 120 mg enteric capsule, 14

2896K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*649.90	41.30	Tecfidera [BD]

▪ **METHOTREXATE**

methotrexate 2.5 mg tablet, 30

1622J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	17.85	19.14	Methoblastin [PF]

methotrexate 10 mg tablet, 15

2272N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	3	..	24.12	25.41	Methoblastin [PF]

▪ **METHOTREXATE**

Restricted benefit

Patients requiring doses greater than 20 mg per week

methotrexate 10 mg tablet, 50

1623K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	52.34	41.30	Methoblastin [PF]

▪ **METHOTREXATE**

Note Pharmaceutical benefits that have the form methotrexate Injection 15 mg/0.3 mL pre-filled syringe and pharmaceutical benefits that have the form methotrexate Injection 15 mg/0.6 mL pre-filled syringe are equivalent for the purposes of substitution.

Authority required (STREAMLINED)

7488

Severe active rheumatoid arthritis

Clinical criteria:

- Patient must be unsuitable for administration of an oral form of methotrexate for this condition.

Authority required (STREAMLINED)

7518

Severe psoriasis

Clinical criteria:

- The condition must not have adequately responded to topical treatment, **AND**
- Patient must be unsuitable for administration of an oral form of methotrexate for this condition.

methotrexate 15 mg/0.3 mL injection, 0.3 mL syringe

11268C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	4	5	..	*89.42	41.30	^a Trexject [LM]

methotrexate 15 mg/0.6 mL injection, 4 x 0.6 mL syringes

11508Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	89.43	41.30	^a Methoblastin PFS [PF]

▪ **METHOTREXATE**

Note Pharmaceutical benefits that have the form methotrexate Injection 7.5 mg/0.15 mL pre-filled syringe and pharmaceutical benefits that have the form methotrexate Injection 7.5 mg/0.3 mL pre-filled syringe are equivalent for the purposes of substitution.

Authority required (STREAMLINED)

7488

Severe active rheumatoid arthritis

Clinical criteria:

- Patient must be unsuitable for administration of an oral form of methotrexate for this condition.

Authority required (STREAMLINED)

7518

Severe psoriasis

Clinical criteria:

- The condition must not have adequately responded to topical treatment, **AND**
- Patient must be unsuitable for administration of an oral form of methotrexate for this condition.

methotrexate 7.5 mg/0.15 mL injection, 0.15 mL syringe

11275K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	4	5	..	*89.42	41.30	^a Trexject [LM]

methotrexate 7.5 mg/0.3 mL injection, 4 x 0.3 mL syringes

11525N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	89.43	41.30	^a Methoblastin PFS [PF]

▪ **METHOTREXATE**

Note Pharmaceutical benefits that have the form methotrexate Injection 10 mg/0.2 mL pre-filled syringe and pharmaceutical benefits that have the form methotrexate Injection 10 mg/0.4 mL pre-filled syringe are equivalent for the purposes of substitution.

Authority required (STREAMLINED)

7488

Severe active rheumatoid arthritis

Clinical criteria:

- Patient must be unsuitable for administration of an oral form of methotrexate for this condition.

Authority required (STREAMLINED)

7518

Severe psoriasis

Clinical criteria:

- The condition must not have adequately responded to topical treatment, **AND**
- Patient must be unsuitable for administration of an oral form of methotrexate for this condition.

methotrexate 10 mg/0.2 mL injection, 0.2 mL syringe

11283W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	4	5	..	*89.42	41.30	^a Trexject [LM]

methotrexate 10 mg/0.4 mL injection, 4 x 0.4 mL syringes

11526P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	89.43	41.30	^a Methoblastin PFS [PF]

■ METHOTREXATE

Note Pharmaceutical benefits that have the form methotrexate Injection 20 mg/0.4 mL pre-filled syringe and pharmaceutical benefits that have the form methotrexate Injection 20 mg/0.8 mL pre-filled syringe are equivalent for the purposes of substitution.

Authority required (STREAMLINED)**7488**

Severe active rheumatoid arthritis

Clinical criteria:

- Patient must be unsuitable for administration of an oral form of methotrexate for this condition.

Authority required (STREAMLINED)**7518**

Severe psoriasis

Clinical criteria:

- The condition must not have adequately responded to topical treatment, **AND**
- Patient must be unsuitable for administration of an oral form of methotrexate for this condition.

methotrexate 20 mg/0.8 mL injection, 4 x 0.8 mL syringes

11509R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	89.43	41.30	^a Methoblastin PFS [PF]

methotrexate 20 mg/0.4 mL injection, 0.4 mL syringe

11288D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	4	5	..	*89.42	41.30	^a Trexject [LM]

■ METHOTREXATE

Note Pharmaceutical benefits that have the form methotrexate Injection 25 mg/0.5 mL pre-filled syringe and pharmaceutical benefits that have the form methotrexate Injection 25 mg/mL pre-filled syringe are equivalent for the purposes of substitution.

Authority required (STREAMLINED)**7488**

Severe active rheumatoid arthritis

Clinical criteria:

- Patient must be unsuitable for administration of an oral form of methotrexate for this condition.

Authority required (STREAMLINED)**7518**

Severe psoriasis

Clinical criteria:

- The condition must not have adequately responded to topical treatment, **AND**
- Patient must be unsuitable for administration of an oral form of methotrexate for this condition.

methotrexate 25 mg/0.5 mL injection, 0.5 mL syringe

11295L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	4	5	..	*89.42	41.30	^a Trexject [LM]

methotrexate 25 mg/mL injection, 4 x 1 mL syringes

11544N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	89.43	41.30	^a Methoblastin PFS [PF]

■ PIRFENIDONE

Note Authority applications for continuing treatment may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Idiopathic pulmonary fibrosis

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be the sole PBS subsidised treatment for this condition.

Treatment criteria:

- Must be treated by a respiratory physician or specialist physician, or in consultation with a respiratory physician or specialist physician.

pirfenidone 801 mg tablet, 90

11410M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	3076.16	41.30	Esbriet [RO]

■ PIRFENIDONE

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Idiopathic pulmonary fibrosis

Treatment Phase: Initial treatment 1 - new patient

Clinical criteria:

- The condition must be diagnosed through a multidisciplinary team, **AND**
- Patient must have chest high resolution computed tomography (HRCT) consistent with diagnosis of idiopathic pulmonary fibrosis within the previous 12 months, **AND**
- Patient must have a forced vital capacity (FVC) greater than or equal to 50% predicted for age, gender and height, **AND**
- Patient must have a forced expiratory volume in 1 second to forced vital capacity ratio (FEV1/FVC) greater than 0.7, **AND**
- Patient must have diffusing capacity of the lungs for carbon monoxide (DLCO) corrected for haemoglobin equal to or greater than 30%, **AND**
- Patient must not have interstitial lung disease due to other known causes including domestic and occupational environmental exposures, connective tissue disease, or drug toxicity, **AND**
- The treatment must be the sole PBS subsidised treatment for this condition.

Treatment criteria:

- Must be treated by a respiratory physician or specialist physician, or in consultation with a respiratory physician or specialist physician.

A multidisciplinary team is defined as comprising of at least a specialist respiratory physician, a radiologist and where histological material is considered, a pathologist. If attendance is not possible because of geographical isolation, consultation with a multidisciplinary team is required for diagnosis.

Patient must not have an acute respiratory infection at the time of FVC testing.

Application for authorisation of initial treatment must be in writing and must include:

- a completed authority prescription form; and
- a completed IPF Authority Application Supporting Information Form; and
- a signed patient acknowledgement.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Idiopathic pulmonary fibrosis

Treatment Phase: Initial treatment 2 - change or re-commencement of treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with nintedanib or pirfenidone for this condition, **AND**
- The treatment must be the sole PBS subsidised treatment for this condition.

Treatment criteria:

- Must be treated by a respiratory physician or specialist physician, or in consultation with a respiratory physician or specialist physician.

Note Applications for authorisation under this criterion may be made by telephone by contacting the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Idiopathic pulmonary fibrosis

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be the sole PBS subsidised treatment for this condition.

Treatment criteria:

- Must be treated by a respiratory physician or specialist physician, or in consultation with a respiratory physician or specialist physician.

Note Authority applications for continuing treatment may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

pirfenidone 267 mg tablet, 90

11406H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	5	..	*3076.17	41.30	Esbriet [RO]

MUSCULO-SKELETAL SYSTEM

pirfenidone 267 mg capsule, 270

11136D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	3076.16	41.30	Esbriet [RO]

■ MUSCULO-SKELETAL SYSTEM

■ ANTIINFLAMMATORY AND ANTIRHEUMATIC PRODUCTS

ANTIINFLAMMATORY AND ANTIRHEUMATIC PRODUCTS, NON-STERIODS

Acetic acid derivatives and related substances

■ DICLOFENAC

diclofenac sodium 100 mg suppository, 20

1302M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP MW	2	3	..	*28.44	29.73	Voltaren 100 [NV]

diclofenac sodium 100 mg suppository, 20

5079H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	2	..	..	*28.44	29.73	Voltaren 100 [NV]

■ DICLOFENAC

Restricted benefit

Chronic arthropathies (including osteoarthritis)

Clinical criteria:

- The condition must have an inflammatory component.

Restricted benefit

Bone pain

Clinical criteria:

- The condition must be due to malignant disease.

diclofenac sodium 50 mg enteric tablet, 50

1300K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	3	..	14.34	15.63	^a APO-Diclofenac [TX] ^a Diclofenac Amneal [ED] ^a Diclofenac Sandoz [SZ]	^a Clonac 50 [RW] ^a Diclofenac AN [EA] ^a Fenac [AF]
			^B 3.21	17.55	15.63	^a Voltaren 50 [NV]	

diclofenac sodium 25 mg enteric tablet, 50

1299J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	2	3	..	*15.18	16.47	^a APO-Diclofenac [TX] ^a Diclofenac Amneal [ED] ^a Diclofenac Sandoz [SZ]	^a Clonac 25 [RW] ^a Diclofenac AN [EA] ^a Fenac 25 [AF]
			^B 3.22	*18.40	16.47	^a Voltaren 25 [NV]	

■ DICLOFENAC

Restricted benefit

Chronic arthropathies (including osteoarthritis)

Clinical criteria:

- The condition must have an inflammatory component.

Restricted benefit

Bone pain

Clinical criteria:

- The condition must be due to malignant disease.

diclofenac sodium 50 mg enteric tablet, 50

5077F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
DP	1	..	..	14.34	15.63	^a APO-Diclofenac [TX] ^a Diclofenac Amneal [ED] ^a Diclofenac Sandoz [SZ]	^a Clonac 50 [RW] ^a Diclofenac AN [EA] ^a Fenac [AF]
			^B 3.21	17.55	15.63	^a Voltaren 50 [NV]	

diclofenac sodium 25 mg enteric tablet, 50

5076E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
DP	2	..	..	*15.18	16.47	^a APO-Diclofenac [TX] ^a Diclofenac Amneal [ED] ^a Diclofenac Sandoz [SZ]	^a Clonac 25 [RW] ^a Diclofenac AN [EA] ^a Fenac 25 [AF]
			^B 3.22	*18.40	16.47	^a Voltaren 25 [NV]	

■ **INDOMETACIN****indometacin 100 mg suppository, 20**

2757D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	3	..	*26.00	27.29	Indocid [AS]

indometacin 100 mg suppository, 20

5128X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	2	..	..	*26.00	27.29	Indocid [AS]

■ **INDOMETACIN****Restricted benefit**

Chronic arthropathies (including osteoarthritis)

Clinical criteria:

- The condition must have an inflammatory component.

Restricted benefit

Bone pain

Clinical criteria:

- The condition must be due to malignant disease.

indometacin 25 mg capsule, 50

2454E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	3	..	*17.64	18.93	^a Arthrexin [AF]
			^B 4.02	*21.66	18.93	^a Indocid [AS]

■ **INDOMETACIN****Restricted benefit**

Chronic arthropathies (including osteoarthritis)

Clinical criteria:

- The condition must have an inflammatory component.

Restricted benefit

Bone pain

Clinical criteria:

- The condition must be due to malignant disease.

indometacin 25 mg capsule, 50

5126T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	2	..	..	*17.64	18.93	^a Arthrexin [AF]
			^B 4.02	*21.66	18.93	^a Indocid [AS]

Oxicams■ **MELOXICAM**

Note Pharmaceutical benefits that have the form meloxicam tablet 7.5 mg and pharmaceutical benefits that have the form meloxicam capsule 7.5 mg are equivalent for the purposes of substitution.

Note The use of this drug for the treatment of the following conditions is not subsidised through the PBS:

- (a) acute pain;
- (b) soft tissue injury;
- (c) arthrosis without an inflammatory component.

Restricted benefit

Osteoarthritis

Clinical criteria:

- The treatment must be for symptomatic treatment.

Restricted benefit

Rheumatoid arthritis

Clinical criteria:

- The treatment must be for symptomatic treatment.

meloxicam 7.5 mg tablet, 30

8561N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	3	..	15.03	16.32	^a APO-Meloxicam [TX] ^a CIPLA MELOXICAM 7.5 [LR] ^a Meloxiauro 7.5 [DO] ^a Meloxicam AN [EA] ^a Meloxicam Sandoz [SZ] ^a Moxicam 7.5 [AF]	^a APX-Meloxicam [TY] ^a MELOBIC [RF] ^a Meloxibell [GQ] ^a Meloxicam-GA [ED] ^a Movalis 7.5 [RW] ^a Pharmacor Meloxicam 7.5 [CR]
			^B 2.79	17.82	16.32	^a Mobic [BY]	

meloxicam 7.5 mg capsule, 30

8887R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	3	..	15.03	16.32	^a APO-Meloxicam [TX] ^a MELOBIC [RF] ^a Movalis 7.5 [RW] ^a Terry White Chemists Meloxicam [TW]	^a Chem mart Meloxicam [CH] ^a Meloxicam Sandoz [SZ] ^a Moxicam [AF]
			^b 2.79	17.82	16.32	^a Mobic [BY]	

■ MELOXICAM

Note Pharmaceutical benefits that have the form meloxicam tablet 15 mg and pharmaceutical benefits that have the form meloxicam capsule 15 mg are equivalent for the purposes of substitution.

Note The use of this drug for the treatment of the following conditions is not subsidised through the PBS:

- (a) acute pain;
- (b) soft tissue injury;
- (c) arthrosis without an inflammatory component.

Restricted benefit

Osteoarthritis

Clinical criteria:

- The treatment must be for symptomatic treatment.

Restricted benefit

Rheumatoid arthritis

Clinical criteria:

- The treatment must be for symptomatic treatment.

meloxicam 15 mg capsule, 30

8888T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	3	..	16.23	17.52	^a APO-Meloxicam [TX] ^a MELOBIC [RF] ^a Movalis 15 [RW] ^a Terry White Chemists Meloxicam [TW]	^a Chem mart Meloxicam [CH] ^a Meloxicam Sandoz [SZ] ^a Moxicam [AF]
			^b 2.88	19.11	17.52	^a Mobic [BY]	

meloxicam 15 mg tablet, 30

8562P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	3	..	16.23	17.52	^a APO-Meloxicam [TX] ^a CIPLA MELOXICAM 15 [LR] ^a Meloxiauro 15 [DO] ^a Meloxicam AN [EA] ^a Meloxicam Sandoz [SZ] ^a Moxicam 15 [AF]	^a APX-Meloxicam [TY] ^a MELOBIC [RF] ^a Meloxibell [GQ] ^a Meloxicam-GA [ED] ^a Movalis 15 [RW] ^a Pharmacor Meloxicam 15 [CR]
			^b 2.88	19.11	17.52	^a Mobic [BY]	

■ PIROXICAM**Restricted benefit**

Chronic arthropathies (including osteoarthritis)

Clinical criteria:

- The condition must have an inflammatory component.

piroxicam 10 mg capsule, 50

1897W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	3	..	17.11	18.40	^a APO-Piroxicam [TX]	^a Mobilis 10 [AF]
			^b 7.94	25.05	18.40	^a Feldene [PF]	

piroxicam 10 mg capsule, 50

5203W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
DP	1	..	..	17.11	18.40	^a APO-Piroxicam [TX]	^a Mobilis 10 [AF]
			^b 7.94	25.05	18.40	^a Feldene [PF]	

piroxicam 20 mg dispersible tablet, 25

1896T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	3	..	16.88	18.17	^a Mobilis D-20 [AF]	
			^b 7.92	24.80	18.17	^a Feldene-D [PF]	

piroxicam 20 mg dispersible tablet, 25

5202T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
DP	1	..	..	16.88	18.17	^a Mobilis D-20 [AF]	
			^b 7.92	24.80	18.17	^a Feldene-D [PF]	

piroxicam 10 mg dispersible tablet, 50

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
1895R	1	3	..	17.11	18.40	Mobilis D-10 [AF]

piroxicam 10 mg dispersible tablet, 50

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
5201R	1	..	..	17.11	18.40	Mobilis D-10 [AF]

piroxicam 20 mg capsule, 25

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
1898X	1	3	..	16.88	18.17	^a APO-Piroxicam [TX]	^a Mobilis 20 [AF]
			^b 7.92	24.80	18.17	^a Feldene [PF]	

piroxicam 20 mg capsule, 25

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
5204X	1	..	..	16.88	18.17	^a APO-Piroxicam [TX]	^a Mobilis 20 [AF]
			^b 7.92	24.80	18.17	^a Feldene [PF]	

Propionic acid derivatives

■ **IBUPROFEN**

ibuprofen 400 mg tablet, 30

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
3192B	1	..	..	13.66	14.95	^a APO-Ibuprofen 400 [TX]
			^b 2.49	16.15	14.95	^a Brufen [GO]

ibuprofen 400 mg tablet, 30

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
5124Q	1	..	..	13.66	14.95	^a APO-Ibuprofen 400 [TX]
			^b 2.49	16.15	14.95	^a Brufen [GO]

■ **IBUPROFEN**

Restricted benefit

Chronic arthropathies (including osteoarthritis)

Clinical criteria:

- The condition must have an inflammatory component.

Restricted benefit

Bone pain

Clinical criteria:

- The condition must be due to malignant disease.

ibuprofen 400 mg tablet, 30

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
3190X	3	3	..	*16.95	18.24	^a APO-Ibuprofen 400 [TX]
			^b 7.47	*24.42	18.24	^a Brufen [GO]

■ **IBUPROFEN**

Restricted benefit

Chronic arthropathies (including osteoarthritis)

Clinical criteria:

- The condition must have an inflammatory component.

Restricted benefit

Bone pain

Clinical criteria:

- The condition must be due to malignant disease.

ibuprofen 400 mg tablet, 30

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
5123P	3	..	..	*16.95	18.24	^a APO-Ibuprofen 400 [TX]
			^b 7.47	*24.42	18.24	^a Brufen [GO]

■ **KETOPROFEN**

Restricted benefit

Chronic arthropathies (including osteoarthritis)

Clinical criteria:

- The condition must have an inflammatory component.

ketoprofen 200 mg modified release capsule, 28

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
1590Q	1	3	..	23.05	24.34	^a Oruvail SR [AV]

			^B 1.92	24.97	24.34	^a Orudis SR 200 [SW]
NP	etoprofen 200 mg modified release capsule, 28					
5136H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	1	..	..	23.05	24.34	^a Oruvail SR [AV]
			^B 1.92	24.97	24.34	^a Orudis SR 200 [SW]

■ **NAPROXEN****Restricted benefit**

Chronic arthropathies (including osteoarthritis)

Clinical criteria:

- The condition must have an inflammatory component.

Restricted benefit

Bone pain

Clinical criteria:

- The condition must be due to malignant disease.

naproxen 500 mg tablet, 50

1659H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	3	..	17.42	18.71	Naprosyn [IX]

naproxen 750 mg modified release tablet, 28

1614Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	3	..	17.01	18.30	^a Proxen SR 750 [IY]
			^B 0.99	18.00	18.30	^a Naprosyn SR750 [IX]

naproxen 1 g modified release tablet, 28

1615B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	3	..	18.58	19.87	^a Proxen SR 1000 [IY]
			^B 1.12	19.70	19.87	^a Naprosyn SR1000 [IX]

naproxen 250 mg tablet, 50

1674D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	3	..	*21.52	22.81	Naprosyn [IX]

■ **NAPROXEN****Authority required (STREAMLINED)****4159**

Chronic arthropathies (including osteoarthritis)

Clinical criteria:

- The condition must have an inflammatory component, **AND**
- Patient must be unable to take a solid dose form of a non-steroidal anti-inflammatory agent.

Authority required (STREAMLINED)**4124**

Bone pain

Clinical criteria:

- The condition must be due to malignant disease, **AND**
- Patient must be unable to take a solid dose form of a non-steroidal anti-inflammatory agent.

naproxen 125 mg/5 mL oral liquid, 474 mL

1658G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	3	..	122.71	41.30	Phebra Naproxen Suspension [FF]

■ **NAPROXEN****Restricted benefit**

Chronic arthropathies (including osteoarthritis)

Clinical criteria:

- The condition must have an inflammatory component.

Restricted benefit

Bone pain

Clinical criteria:

- The condition must be due to malignant disease.

naproxen 500 mg tablet, 50

5177L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	1	..	..	17.42	18.71	Naprosyn [IX]

naproxen 750 mg modified release tablet, 28

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
5178M	1	..	..	17.01	18.30	^a Proxen SR 750 [IY]
DP			^B 0.99	18.00	18.30	^a Naprosyn SR750 [IX]

naproxen 1 g modified release tablet, 28

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
5179N	1	..	..	18.58	19.87	^a Proxen SR 1000 [IY]
DP			^B 1.12	19.70	19.87	^a Naprosyn SR1000 [IX]

naproxen 250 mg tablet, 50

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
5176K	2	..	..	*21.52	22.81	Naprosyn [IX]
DP						

■ NAPROXEN

Note Naproxen sodium 550 mg is approximately equivalent to 500 mg of naproxen acid.

Restricted benefit

Chronic arthropathies (including osteoarthritis)

Clinical criteria:

- The condition must have an inflammatory component.

Restricted benefit

Bone pain

Clinical criteria:

- The condition must be due to malignant disease.

naproxen sodium 550 mg tablet, 50

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
1795L	1	3	..	17.56	18.85	^a Crysanal [IY]
NP			^B 1.87	19.43	18.85	^a Anaprox 550 [IX]

■ NAPROXEN

Note Naproxen sodium 550 mg is approximately equivalent to 500 mg of naproxen acid.

Restricted benefit

Chronic arthropathies (including osteoarthritis)

Clinical criteria:

- The condition must have an inflammatory component.

Restricted benefit

Bone pain

Clinical criteria:

- The condition must be due to malignant disease.

naproxen sodium 550 mg tablet, 50

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
5186Y	1	..	..	17.56	18.85	^a Crysanal [IY]
DP			^B 1.87	19.43	18.85	^a Anaprox 550 [IX]

Fenamates**■ MEFENAMIC ACID****Restricted benefit**

Dysmenorrhoea

Restricted benefit

Menorrhagia

mefenamic acid 250 mg capsule, 50

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
1824B	1	2	..	20.32	21.61	Ponstan [PF]
NP						

Coxibs**■ CELECOXIB**

Note The use of this drug for the treatment of the following conditions is not subsidised through the PBS:

- acute pain;
- soft tissue injury;
- arthrosis without an inflammatory component.

Restricted benefit

Osteoarthritis

Clinical criteria:

- The treatment must be for symptomatic treatment.

Restricted benefit

Rheumatoid arthritis

Clinical criteria:

- The treatment must be for symptomatic treatment.

celecoxib 200 mg capsule, 30

8440F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	3	..	17.03	18.32	^a APO-Celecoxib [TX]	^a Blooms the Chemist Celecoxib [IB]
						^a Celaxib [AF]	^a Celebrex [UJ]
						^a Celecoxib AN [EA]	^a Celecoxib APOTEX [TY]
						^a Celecoxib BTC [JB]	^a Celecoxib GH [GQ]
						^a Celecoxib Sandoz [SZ]	^a Celexi [RW]
						^a GenRx Celecoxib [GX]	

celecoxib 100 mg capsule, 60

8439E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	3	..	17.03	18.32	^a APO-Celecoxib [TX]	^a Blooms the Chemist Celecoxib [IB]
						^a Celaxib [AF]	^a Celebrex [UJ]
						^a Celecoxib AN [EA]	^a Celecoxib APOTEX [TY]
						^a Celecoxib BTC [JB]	^a Celecoxib GH [GQ]
						^a Celecoxib Sandoz [SZ]	^a Celexi [RW]
						^a GenRx Celecoxib [GX]	

SPECIFIC ANTIRHEUMATIC AGENTS**Quinolines****■ HYDROXYCHLOROQUINE****Authority required (STREAMLINED)****10417**

Autoimmune disorder

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be authorised by a medical practitioner who is recognised under State or Territory legislation that forms part of the Health Practitioner Regulation National Law, as a specialist in dermatology, intensive care medicine, paediatrics and child health, physician, or emergency medicine, **AND**
- Patient must have an autoimmune disorder.

Authority required (STREAMLINED)**10418**

Malaria

Treatment Phase: Initial treatment

Clinical criteria:

- The treatment must be authorised by a medical practitioner who is recognised under State or Territory legislation that forms part of the Health Practitioner Regulation National Law, as a specialist in dermatology, intensive care medicine, paediatrics and child health, physician, or emergency medicine, **AND**
- Patient must require treatment of acute attack; OR
- Patient must require suppressive therapy.

hydroxychloroquine sulfate 200 mg tablet, 100

12003R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	1	..	26.08	27.37	^a APO- Hydroxychloroquine [TX]	^a Hequinel [RW]
						^a Hydroxychloroquine AN [EA]	^a Hydroxychloroquine GH [GQ]
						^a Plaquenil [SW]	

■ HYDROXYCHLOROQUINE**Note Shared Care Model:**

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**10419**

Autoimmune disorder

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously been treated with PBS-subsidised therapy with this drug for this condition.

Authority required (STREAMLINED)**10420**

Malaria

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously been treated with PBS-subsidised therapy with this drug for this condition.

hydroxychloroquine sulfate 200 mg tablet, 100

1512N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	1	..	26.08	27.37	^a APO- Hydroxychloroquine [TX] ^a Hydroxychloroquine AN [EA] ^a Plaquenil [SW]	^a Hequinel [RW] ^a Hydroxychloroquine GH [GQ]

*Gold preparations***■ AURANOFIN**

Caution Regular blood and urine checks are essential.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

auranofin 3 mg capsule, 60

2022K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	903.87	41.30	Ridaura [BZ]

auranofin 3 mg tablet, 60

1095P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	209.30	41.30	Ridaura [GH]

*Penicillamine and similar agents***■ PENICILLAMINE**

Caution Regular blood and urine checks are essential.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

penicillamine 250 mg tablet, 100

2838J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	55.03	41.30	D-Penamine [AL]

penicillamine 125 mg tablet, 100

2721F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	49.65	41.30	D-Penamine [AL]

■ MUSCLE RELAXANTS**MUSCLE RELAXANTS, CENTRALLY ACTING AGENTS***Other centrally acting agents***■ BACLOFEN****baclofen 25 mg tablet, 100**

2730Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	33.83	35.12	^a APO-Baclofen [TX] ^a Lioresal 25 [NV]	^a Clofen 25 [AF] ^a Stelax 25 [RW]

baclofen 10 mg tablet, 100

2729P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	22.00	23.29	^a APO-Baclofen [TX] ^a Lioresal 10 [NV]	^a Clofen 10 [AF] ^a Stelax 10 [RW]

MUSCLE RELAXANTS, DIRECTLY ACTING AGENTS*Dantrolene and derivatives***■ DANTROLENE****Restricted benefit**

Chronic spasticity

dantrolene sodium hemiheptahydrate 50 mg capsule, 100

1780Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	77.13	41.30	Dantrium [PF]

dantrolene sodium hemiheptahydrate 25 mg capsule, 100

1779P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	122.73	41.30	Dantrium [PF]

■ ANTIGOUT PREPARATIONS**ANTIGOUT PREPARATIONS***Preparations inhibiting uric acid production***■ ALLOPURINOL****Note** The dose should be adjusted in accordance with renal function.**allopurinol 100 mg tablet, 200**

2600W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	2	..	16.95	18.24	^a Allopurinol APOTEX [GX] ^a Allosig [RF] ^a Progout 100 [AF]	^a Allopurinol Sandoz [SZ] ^a NOUMED ALLOPURINOL [VO]
			^B 3.40	20.35	18.24	^a Zyloprim [RW]	

allopurinol 300 mg tablet, 60

2604C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	2	..	15.70	16.99	^a Allopurinol APOTEX [GX] ^a Allosig [RF] ^a Progout 300 [AF]	^a Allopurinol Sandoz [SZ] ^a NOUMED ALLOPURINOL [VO]
			^B 3.32	19.02	16.99	^a Zyloprim [RW]	

■ FEBUXOSTAT**Note Shared Care Model:**

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**8921**

Chronic gout

Clinical criteria:

- The condition must be either chronic gouty arthritis or chronic tophaceous gout, **AND**
- Patient must have a medical contraindication to allopurinol; OR
- Patient must have a documented history of allopurinol hypersensitivity syndrome; OR
- Patient must have an intolerance to allopurinol necessitating permanent treatment discontinuation.

febuxostat 80 mg tablet, 28

10445R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	49.88	41.30	Adenuric [FK]

*Preparations increasing uric acid excretion***■ PROBENECID****probenecid 500 mg tablet, 100**

1940D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	63.17	41.30	Pro-Cid [FF]

*Preparations with no effect on uric acid metabolism***■ COLCHICINE****colchicine 500 microgram tablet, 30**

3410L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	16.13	17.42	^a Lengout [LN] ^B 2.77 18.90 17.42 ^a Colgout [AS]

■ DRUGS FOR TREATMENT OF BONE DISEASES**DRUGS AFFECTING BONE STRUCTURE AND MINERALIZATION***Bisphosphonates***■ ALENDRONATE****Restricted benefit**

Corticosteroid-induced osteoporosis

Clinical criteria:

- Patient must currently be on long-term (at least 3 months), high-dose (at least 7.5 mg per day prednisolone or equivalent) corticosteroid therapy, **AND**
- Patient must have a Bone Mineral Density (BMD) T-score of -1.5 or less, **AND**
- Patient must not receive concomitant treatment with any other PBS-subsidised anti-resorptive agent for this condition. The duration and dose of corticosteroid therapy together with the date, site (femoral neck or lumbar spine) and score of the qualifying BMD measurement must be documented in the patient's medical records when treatment is initiated.

Note Anti-resorptive agents in osteoporosis include alendronate sodium, risedronate sodium, denosumab, raloxifene hydrochloride and zoledronic acid.

Restricted benefit

Osteoporosis

Population criteria:

- Patient must be aged 70 years or older.

Clinical criteria:

- Patient must have a Bone Mineral Density (BMD) T-score of -2.5 or less, **AND**
- Patient must not receive concomitant treatment with any other PBS-subsidised anti-resorptive agent for this condition. The date, site (femoral neck or lumbar spine) and score of the qualifying BMD measurement must be documented in the patient's medical records when treatment is initiated.

Note Anti-resorptive agents in osteoporosis include alendronate sodium, risedronate sodium, denosumab, raloxifene hydrochloride and zoledronic acid.

Restricted benefit

Established osteoporosis

Clinical criteria:

- Patient must have fracture due to minimal trauma, **AND**
- Patient must not receive concomitant treatment with any other PBS-subsidised anti-resorptive agent for this condition. The fracture must have been demonstrated radiologically and the year of plain x-ray or computed tomography (CT) scan or magnetic resonance imaging (MRI) scan must be documented in the patient's medical records when treatment is initiated. A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or, a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

Note Anti-resorptive agents in established osteoporosis include alendronate sodium, risedronate sodium, denosumab, raloxifene hydrochloride and zoledronic acid.

alendronate 70 mg tablet, 4

8511Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	16.07	17.36	^a Alendrobell 70mg [GQ] ^a Alendro Once Weekly [RW] ^a Densate 70 [DO]	^a Alendronate Sandoz [SZ] ^a APO-Alendronate [TX] ^a Fonat [AL]

■ IBANDRONATE

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Bone metastases

Clinical criteria:

- The condition must be due to breast cancer.

ibandronate 50 mg tablet, 28

9357L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	292.43	41.30	Bondronat [IX]

■ PAMIDRONATE DISODIUM

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Symptomatic Paget disease of bone

pamidronate disodium 30 mg/10 mL injection, 10 mL vial

8462J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	..	..	*71.44	41.30	Pamisol [PF]

pamidronate disodium 60 mg/10 mL injection, 10 mL vial

8463K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	71.44	41.30	Pamisol [PF]

pamidronate disodium 15 mg/5 mL injection, 5 mL vial

8461H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	..	..	*71.42	41.30	Pamisol [PF]

■ RISEDRONATE**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Symptomatic Paget disease of bone

risedronate sodium 30 mg tablet, 28

8482K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	179.22	41.30	Actonel [TT]

■ RISEDRONATE**Restricted benefit**

Corticosteroid-induced osteoporosis

Clinical criteria:

- Patient must currently be on long-term (at least 3 months), high-dose (at least 7.5 mg per day prednisolone or equivalent) corticosteroid therapy, **AND**
- Patient must have a Bone Mineral Density (BMD) T-score of -1.5 or less, **AND**
- Patient must not receive concomitant treatment with any other PBS-subsidised anti-resorptive agent for this condition. The duration and dose of corticosteroid therapy together with the date, site (femoral neck or lumbar spine) and score of the qualifying BMD measurement must be documented in the patient's medical records when treatment is initiated.

Note Anti-resorptive agents in osteoporosis include alendronate sodium, risedronate sodium, denosumab, raloxifene hydrochloride and zoledronic acid.

Restricted benefit

Osteoporosis

Population criteria:

- Patient must be aged 70 years or older.

Clinical criteria:

- Patient must have a Bone Mineral Density (BMD) T-score of -2.5 or less, **AND**
- Patient must not receive concomitant treatment with any other PBS-subsidised anti-resorptive agent for this condition. The date, site (femoral neck or lumbar spine) and score of the qualifying BMD measurement must be documented in the patient's medical records when treatment is initiated.

Note Anti-resorptive agents in osteoporosis include alendronate sodium, risedronate sodium, denosumab, raloxifene hydrochloride and zoledronic acid.

Restricted benefit

Established osteoporosis

Clinical criteria:

- Patient must have fracture due to minimal trauma, **AND**
- Patient must not receive concomitant treatment with any other PBS-subsidised anti-resorptive agent for this condition. The fracture must have been demonstrated radiologically and the year of plain x-ray or computed tomography (CT) scan or magnetic resonance imaging (MRI) scan must be documented in the patient's medical records when treatment is initiated. A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or, a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

Note Anti-resorptive agents in established osteoporosis include alendronate sodium, risedronate sodium, denosumab, raloxifene hydrochloride and zoledronic acid.

risedronate sodium 35 mg enteric tablet, 4

8972F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	36.03	37.32	Actonel EC [TT]

risedronate sodium 150 mg tablet, 1

9391G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	38.05	39.34	^a Acris Once-a-Month [AF] ^a APO-Risedronate [TX]	^a Actonel Once-a-Month [TT]

risedronate sodium 35 mg tablet, 4

8621R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	36.03	37.32	^a Acris Once-a-Week [AF] ^a Risedronate AN [EA] ^a Risedro once a week [RW]	^a APO-Risedronate [TX] ^a Risedronate Sandoz [SZ]

risedronate sodium 5 mg tablet, 28

8481J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	36.03	37.32	Actonel [TT]

■ ZOLEDRONIC ACID

Note Pharmaceutical benefits that have the form zoledronic acid injection 5 mg/100 mL vial and pharmaceutical benefits that have the form zoledronic acid injection 5 mg/100 mL bag are equivalent for the purposes of substitution.

Authority required (STREAMLINED)**5710**

Symptomatic Paget disease of bone

Only 1 treatment each year per patient will be PBS-subsidised

zoledronic acid 5 mg/100 mL injection, 100 mL vial

9350D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	117.06	41.30	^a Aclasta [HX] ^a Zoledasta [TX]	^a Osteovan [SZ] ^a Zoledronic Acid SUN [RA]

■ ZOLEDRONIC ACID

Note Anti-resorptive agents in established osteoporosis include alendronate sodium, risedronate sodium, denosumab, raloxifene hydrochloride and zoledronic acid.

Note Pharmaceutical benefits that have the form zoledronic acid injection 5 mg/100 mL vial and pharmaceutical benefits that have the form zoledronic acid injection 5 mg/100 mL bag are equivalent for the purposes of substitution.

Authority required (STREAMLINED)**6308**

Corticosteroid-induced osteoporosis

Clinical criteria:

- Patient must currently be on long-term (at least 3 months), high-dose (at least 7.5 mg per day prednisolone or equivalent) corticosteroid therapy, **AND**
 - Patient must have a Bone Mineral Density (BMD) T-score of -1.5 or less, **AND**
 - Patient must not receive concomitant treatment with any other PBS-subsidised anti-resorptive agent for this condition, **AND**
 - Patient must not receive more than one PBS-subsidised treatment per year.
- The duration and dose of corticosteroid therapy together with the date, site (femoral neck or lumbar spine) and score of the qualifying BMD measurement must be documented in the patient's medical records when treatment is initiated.

Authority required (STREAMLINED)**6313**

Osteoporosis

Population criteria:

- Patient must be aged 70 years or older.

Clinical criteria:

- Patient must have a Bone Mineral Density (BMD) T-score of -3.0 or less, **AND**
 - Patient must not receive concomitant treatment with any other PBS-subsidised anti-resorptive agent for this condition, **AND**
 - Patient must not receive more than one PBS-subsidised treatment per year.
- The date, site (femoral neck or lumbar spine) and score of the qualifying BMD measurement must be documented in the patient's medical records when treatment is initiated.

Authority required (STREAMLINED)**6318**

Established osteoporosis

Clinical criteria:

- Patient must have fracture due to minimal trauma, **AND**
- Patient must not receive concomitant treatment with any other PBS-subsidised anti-resorptive agent for this condition, **AND**
- Patient must not receive more than one PBS-subsidised treatment per year.

The fracture must have been demonstrated radiologically and the year of plain x-ray or computed tomography (CT) scan or magnetic resonance imaging (MRI) scan must be documented in the patient's medical records when treatment is initiated.

A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or, a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

zoledronic acid 5 mg/100 mL injection, 100 mL vial

9288W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	117.06	41.30	^a Aclasta [HX] ^a Zoledasta [TX]	^a Osteovan [SZ] ^a Zoledronic Acid SUN [RA]

Bisphosphonates, combinations

■ ALENDRONATE + COLECALCIFEROL

Note Anti-resorptive agents in established osteoporosis include alendronate sodium, risedronate sodium, denosumab, raloxifene hydrochloride and zoledronic acid.

Authority required (STREAMLINED)

6306

Corticosteroid-induced osteoporosis

Clinical criteria:

- Patient must currently be on long-term (at least 3 months), high-dose (at least 7.5 mg per day prednisolone or equivalent) corticosteroid therapy, **AND**
- Patient must have a Bone Mineral Density (BMD) T-score of -1.5 or less, **AND**
- Patient must not receive concomitant treatment with any other PBS-subsidised anti-resorptive agent for this condition. The duration and dose of corticosteroid therapy together with the date, site (femoral neck or lumbar spine) and score of the qualifying BMD measurement must be documented in the patient's medical records when treatment is initiated.

Authority required (STREAMLINED)

6325

Osteoporosis

Population criteria:

- Patient must be aged 70 years or older.

Clinical criteria:

- Patient must have a Bone Mineral Density (BMD) T-score of -2.5 or less, **AND**
- Patient must not receive concomitant treatment with any other PBS-subsidised anti-resorptive agent for this condition. The date, site (femoral neck or lumbar spine) and score of the qualifying BMD measurement must be documented in the patient's medical records when treatment is initiated.

Authority required (STREAMLINED)

6319

Established osteoporosis

Clinical criteria:

- Patient must have fracture due to minimal trauma, **AND**
- Patient must not receive concomitant treatment with any other PBS-subsidised anti-resorptive agent for this condition. The fracture must have been demonstrated radiologically and the year of plain x-ray or computed tomography (CT) scan or magnetic resonance imaging (MRI) scan must be documented in the patient's medical records when treatment is initiated. A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or, a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

alendronate 70 mg + colecalciferol 140 microgram (5600 units) tablet, 4

9183H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	18.87	20.16	^a ALENDRONATE PLUS D3 70mg/140ug APOTEX [GX]	^a Alendronate plus D3-DRLA [RZ]
						^a Alendronate Plus D3 Sandoz [SZ]	^a FonatPlus [AF]
			^B 4.00	22.87	20.16	^a Fosamax Plus 70 mg/140 mcg [MQ]	

■ ALENDRONATE + COLECALCIFEROL

Note Anti-resorptive agents in established osteoporosis include alendronate sodium, risedronate sodium, denosumab, raloxifene hydrochloride and zoledronic acid.

Note Fosamax Plus provides a supplemental intake of vitamin D. The amount of colecalciferol present in Fosamax Plus is not sufficient to use as the sole treatment for correction of vitamin D deficiency.

Authority required (STREAMLINED)

6307

Corticosteroid-induced osteoporosis

Clinical criteria:

- Patient must currently be on long-term (at least 3 months), high-dose (at least 7.5 mg per day prednisolone or equivalent) corticosteroid therapy, **AND**
- Patient must have a Bone Mineral Density (BMD) T-score of -1.5 or less, **AND**
- Patient must not receive concomitant treatment with any other PBS-subsidised anti-resorptive agent for this condition. The duration and dose of corticosteroid therapy together with the date, site (femoral neck or lumbar spine) and score of the qualifying BMD measurement must be documented in the patient's medical records when treatment is initiated.

Authority required (STREAMLINED)

6320

Osteoporosis

Population criteria:

- Patient must be aged 70 years or older.

Clinical criteria:

- Patient must have a Bone Mineral Density (BMD) T-score of -2.5 or less, **AND**
- Patient must not receive concomitant treatment with any other PBS-subsidised anti-resorptive agent for this condition.

The date, site (femoral neck or lumbar spine) and score of the qualifying BMD measurement must be documented in the patient's medical records when treatment is initiated.

Authority required (STREAMLINED)

6315

Established osteoporosis

Clinical criteria:

- Patient must have fracture due to minimal trauma, **AND**
- Patient must not receive concomitant treatment with any other PBS-subsidised anti-resorptive agent for this condition. The fracture must have been demonstrated radiologically and the year of plain x-ray or computed tomography (CT) scan or magnetic resonance imaging (MRI) scan must be documented in the patient's medical records when treatment is initiated. A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or, a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

alendronate 70 mg + colecalciferol 70 microgram (2800 units) tablet, 4

9012H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	18.87	20.16	^a ALENDRONATE PLUS D3 70mg/70ug APOTEX [GX]	^a Alendronate plus D3-DRLA [RZ]
						^a Alendronate Plus D3 Sandoz [SZ]	^a FonatPlus [AF]
			^B 4.00	22.87	20.16	^a Fosamax Plus [MQ]	

■ ALENDRONATE + COLECALCIFEROL (&) CALCIUM CARBONATE

Note Anti-resorptive agents in established osteoporosis include alendronate sodium, risedronate sodium, denosumab, raloxifene hydrochloride and zoledronic acid.

Authority required (STREAMLINED)

6306

Corticosteroid-induced osteoporosis

Clinical criteria:

- Patient must currently be on long-term (at least 3 months), high-dose (at least 7.5 mg per day prednisolone or equivalent) corticosteroid therapy, **AND**
- Patient must have a Bone Mineral Density (BMD) T-score of -1.5 or less, **AND**
- Patient must not receive concomitant treatment with any other PBS-subsidised anti-resorptive agent for this condition. The duration and dose of corticosteroid therapy together with the date, site (femoral neck or lumbar spine) and score of the qualifying BMD measurement must be documented in the patient's medical records when treatment is initiated.

Authority required (STREAMLINED)

6325

Osteoporosis

Population criteria:

- Patient must be aged 70 years or older.

Clinical criteria:

- Patient must have a Bone Mineral Density (BMD) T-score of -2.5 or less, **AND**
- Patient must not receive concomitant treatment with any other PBS-subsidised anti-resorptive agent for this condition. The date, site (femoral neck or lumbar spine) and score of the qualifying BMD measurement must be documented in the patient's medical records when treatment is initiated.

Authority required (STREAMLINED)

6319

Established osteoporosis

Clinical criteria:

- Patient must have fracture due to minimal trauma, **AND**
- Patient must not receive concomitant treatment with any other PBS-subsidised anti-resorptive agent for this condition. The fracture must have been demonstrated radiologically and the year of plain x-ray or computed tomography (CT) scan or magnetic resonance imaging (MRI) scan must be documented in the patient's medical records when treatment is initiated. A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or, a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

alendronate 70 mg + colecalciferol 140 microgram tablet [4] (&) calcium (as carbonate) 500 mg tablet [48], 1 pack

9351E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	23.76	25.05	ReddyMax Plus D-Cal [RZ]

■ RISEDRONATE (&) CALCIUM CARBONATE

Note Anti-resorptive agents in established osteoporosis include alendronate sodium, risedronate sodium, denosumab, raloxifene hydrochloride and zoledronic acid.

Authority required (STREAMLINED)

6306

Corticosteroid-induced osteoporosis

Clinical criteria:

- Patient must currently be on long-term (at least 3 months), high-dose (at least 7.5 mg per day prednisolone or equivalent) corticosteroid therapy, **AND**
- Patient must have a Bone Mineral Density (BMD) T-score of -1.5 or less, **AND**
- Patient must not receive concomitant treatment with any other PBS-subsidised anti-resorptive agent for this condition. The duration and dose of corticosteroid therapy together with the date, site (femoral neck or lumbar spine) and score of the qualifying BMD measurement must be documented in the patient's medical records when treatment is initiated.

Authority required (STREAMLINED)**6325**

Osteoporosis

Population criteria:

- Patient must be aged 70 years or older.

Clinical criteria:

- Patient must have a Bone Mineral Density (BMD) T-score of -2.5 or less, **AND**
- Patient must not receive concomitant treatment with any other PBS-subsidised anti-resorptive agent for this condition. The date, site (femoral neck or lumbar spine) and score of the qualifying BMD measurement must be documented in the patient's medical records when treatment is initiated.

Authority required (STREAMLINED)**6319**

Established osteoporosis

Clinical criteria:

- Patient must have fracture due to minimal trauma, **AND**
- Patient must not receive concomitant treatment with any other PBS-subsidised anti-resorptive agent for this condition. The fracture must have been demonstrated radiologically and the year of plain x-ray or computed tomography (CT) scan or magnetic resonance imaging (MRI) scan must be documented in the patient's medical records when treatment is initiated. A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or, a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

risedronate sodium 35 mg tablet [4] (&) calcium (as carbonate) 500 mg tablet [24], 28

8899J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	36.03	37.32	Acris Combi [AF]

*Other drugs affecting bone structure and mineralization***■ CALCITRIOL****Authority required (STREAMLINED)****5401**

Hypocalcaemia

Clinical criteria:

- The condition must be due to renal disease.

Authority required (STREAMLINED)**5255**

Hypoparathyroidism

Authority required (STREAMLINED)**5089**

Hypophosphataemic rickets

Authority required (STREAMLINED)**5114**

Vitamin D-resistant rickets

Authority required (STREAMLINED)**5402**

Established osteoporosis

Clinical criteria:

- Patient must have fracture due to minimal trauma. The fracture must have been demonstrated radiologically and the year of plain x-ray or computed tomography (CT) scan or magnetic resonance imaging (MRI) scan must be documented in the patient's medical records when treatment is initiated. A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or, a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

calcitriol 0.25 microgram capsule, 100

2502Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	3	..	28.14	29.43	^a APO-Calcitriol [TX]	^a Calciprox [ER]
						^a Calcitriol AN [EA]	^a CALITROL [ED]
						^a Kosteo [RW]	^a Sical [AF]
			^B 2.29	30.43	29.43	^a Rocaltrol [IX]	

■ DENOSUMAB

Note Denosumab is not PBS-subsidised for use in patients who have undergone curative surgical resection.

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**4504**

Giant cell tumour of bone

Clinical criteria:

- Patient must be one in whom surgical resection is not feasible; OR
- Patient must be one in whom surgical resection is possible but surgery would result in significant morbidity.

Population criteria:

- Patient must be an adult; OR
- Patient must be a skeletally mature adolescent.

denosumab 120 mg/1.7 mL injection, 1.7 mL vial

10061M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	460.69	41.30	Xgeva [AN]

■ DENOSUMAB**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**4158**

Bone metastases

Clinical criteria:

- The condition must be due to breast cancer.

Authority required (STREAMLINED)**4150**

Bone metastases

Clinical criteria:

- The condition must be due to castration-resistant prostate cancer.

denosumab 120 mg/1.7 mL injection, 1.7 mL vial

5110Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	460.69	41.30	Xgeva [AN]

■ DENOSUMAB

Note Anti-resorptive agents in established osteoporosis include alendronate sodium, risedronate sodium, denosumab, raloxifene hydrochloride and zoledronic acid.

Authority required (STREAMLINED)**6548**

Osteoporosis

Population criteria:

- Patient must be aged 70 years or older.

Clinical criteria:

- Patient must have a Bone Mineral Density (BMD) T-score of -2.5 or less, **AND**
- Patient must not receive concomitant treatment with any other PBS-subsidised anti-resorptive agent for this condition. The date, site (femoral neck or lumbar spine) and score of the qualifying BMD measurement must be documented in the patient's medical records when treatment is initiated.

Authority required (STREAMLINED)**6524**

Established osteoporosis

Clinical criteria:

- Patient must have fracture due to minimal trauma, **AND**
- Patient must not receive concomitant treatment with any other PBS-subsidised anti-resorptive agent for this condition. The fracture must have been demonstrated radiologically and the year of plain x-ray or computed tomography (CT) scan or magnetic resonance imaging (MRI) scan must be documented in the patient's medical records when treatment is initiated. A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or, a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

denosumab 60 mg/mL injection, 1 mL syringe

5457F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	250.53	41.30	Prolia [AN]

■ RALOXIFENE

Note Anti-resorptive agents in established osteoporosis include alendronate sodium, risedronate sodium, denosumab, raloxifene hydrochloride and zoledronic acid.

Authority required (STREAMLINED)

6314

Established post-menopausal osteoporosis

Clinical criteria:

- Patient must have fracture due to minimal trauma, **AND**
- Patient must not receive concomitant treatment with any other PBS-subsidised anti-resorptive agent for this condition. The fracture must have been demonstrated radiologically and the year of plain x-ray or computed tomography (CT) scan or magnetic resonance imaging (MRI) scan must be documented in the patient's medical records when treatment is initiated. A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or, a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

raloxifene hydrochloride 60 mg tablet, 28

8363E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	32.58	33.87	^a APO-Raloxifene [TX] ^a Evista [LY] ^a RALOVISTA [RF] ^a Raloxifene AN [EA]	^a Evifyne [EL] ^a Fixta 60 [DO] ^a Raloxifene AMNEAL [ED] ^a Raloxifene GH [GQ]

■ ROMOSOZUMAB

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Authority required

Severe established osteoporosis

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must be at very high risk of fracture, **AND**
- Patient must have a bone mineral density (BMD) T-score of -3.0 or less, **AND**
- Patient must have had 2 or more fractures due to minimal trauma, **AND**
- Patient must have experienced at least 1 symptomatic new fracture after at least 12 months continuous therapy with an anti-resorptive agent at adequate doses, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- The treatment must not exceed a lifetime maximum of 12 months therapy, **AND**
- Patient must not have received treatment with teriparatide; OR
- Patient must have developed intolerance to teriparatide of a severity necessitating permanent treatment withdrawal within the first 6 months of therapy.

Treatment criteria:

- Must be treated by a specialist; OR
- Must be treated by a consultant physician.

A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or, a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

If treatment with anti-resorptive therapy is contraindicated according to the relevant TGA-approved Product Information, details of the contraindication must be documented in the patient's medical record at the time treatment with this drug is initiated.

If an intolerance of a severity necessitating permanent treatment withdrawal develops during the relevant period of use of one anti-resorptive agent, alternate anti-resorptive agents must be trialled so that the patient achieves the minimum requirement of 12 months continuous therapy. Details must be documented in the patient's medical record at the time treatment with this drug is initiated.

Anti-resorptive therapies for osteoporosis and their adequate doses which will be accepted for the purposes of administering this restriction are alendronate sodium 10 mg per day or 70 mg once weekly, risedronate sodium 5 mg per day or 35 mg once weekly or 150 mg once monthly, raloxifene hydrochloride 60 mg per day (women only), denosumab 60 mg once every 6 months and zoledronic acid 5 mg per annum.

Details of prior anti-resorptive therapy, fracture history including the date(s), site(s), the symptoms associated with the fracture(s) which developed after at least 12 months continuous anti-resorptive therapy and the score of the qualifying BMD measurement must be provided at the time of application.

Authority required

Severe established osteoporosis

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must not exceed a lifetime maximum of 12 months therapy.

Treatment criteria:

- Must be treated by a specialist; OR
- Must be treated by a consultant physician.

romosozumab 105 mg/1.17 mL injection, 2 x 1.17 mL syringes

12301K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	628.81	41.30	Evenity [AN]

■ TERIPARATIDE

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Authority required

Severe established osteoporosis

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must be at very high risk of fracture, **AND**
- Patient must have a bone mineral density (BMD) T-score of -3.0 or less, **AND**
- Patient must have had 2 or more fractures due to minimal trauma, **AND**
- Patient must have experienced at least 1 symptomatic new fracture after at least 12 months continuous therapy with an anti-resorptive agent at adequate doses, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- The treatment must not exceed a lifetime maximum of 18 months therapy, **AND**
- Patient must not have received treatment with romosozumab; OR
- Patient must have developed intolerance to romosozumab of a severity necessitating permanent treatment withdrawal within the first 6 months of therapy.

Treatment criteria:

- Must be treated by a specialist; OR
- Must be treated by a consultant physician.

A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or, a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

If treatment with anti-resorptive therapy is contraindicated according to the relevant TGA-approved Product Information, details of the contraindication must be documented in the patient's medical record at the time treatment with this drug is initiated.

If an intolerance of a severity necessitating permanent treatment withdrawal develops during the relevant period of use of one anti-resorptive agent, alternate anti-resorptive agents must be trialled so that the patient achieves the minimum requirement of 12 months continuous therapy. Details must be documented in the patient's medical record at the time treatment with this drug is initiated.

Anti-resorptive therapies for osteoporosis and their adequate doses which will be accepted for the purposes of administering this restriction are alendronate sodium 10 mg per day or 70 mg once weekly, risedronate sodium 5 mg per day or 35 mg once weekly or 150 mg once monthly, raloxifene hydrochloride 60 mg per day (women only), denosumab 60 mg once every 6 months and zoledronic acid 5 mg per annum.

Details of prior anti-resorptive therapy, fracture history including the date(s), site(s), the symptoms associated with the fracture(s) which developed after at least 12 months continuous anti-resorptive therapy and the score of the qualifying BMD measurement must be provided at the time of application.

Authority required

Severe established osteoporosis

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must not exceed a lifetime maximum of 18 months therapy.

Treatment criteria:

- Must be treated by a specialist; OR
- Must be treated by a consultant physician.

Note Up to a maximum of 18 pens will be reimbursed through the PBS.

teriparatide 250 microgram/mL injection, 2.4 mL pen device

9411H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	378.65	41.30	Forteo [LY]

NERVOUS SYSTEM

ANALGESICS

OPIOIDS

Natural opium alkaloids

CODEINE

Caution The risk of drug dependence is high.

Note Prescribing of drugs of addiction by dentists is not permitted in some States/Territories.

Restricted benefit

Severe pain

Clinical criteria:

- The treatment must be for short term therapy of acute severe pain, **AND**
- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid analgesics; OR
- Patient must be unable to use non-opioid analgesics due to contraindications or intolerance.

codeine phosphate hemihydrate 30 mg tablet, 20

12054K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	0.5	..	..	*22.13	23.42	Aspen Pharma Pty Ltd [AS]

CODEINE

Caution The risk of drug dependence is high.

Note Prescribing of drugs of addiction by dentists is not permitted in some States/Territories.

Restricted benefit

Severe pain

Clinical criteria:

- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid analgesics; OR
- Patient must be unable to use non-opioid analgesics due to contraindications or intolerance.

codeine phosphate hemihydrate 30 mg tablet, 20

5063L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	1	..	..	22.32	23.61	Aspen Pharma Pty Ltd [AS]

CODEINE

Caution The risk of drug dependence is high.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Restricted benefit

Severe pain

Clinical criteria:

- The treatment must be for short term therapy of acute severe pain, **AND**
- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid analgesics; OR
- Patient must be unable to use non-opioid analgesics due to contraindications or intolerance.

codeine phosphate hemihydrate 30 mg tablet, 20

12065B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	0.5	..	..	*22.13	23.42	Aspen Pharma Pty Ltd [AS]

CODEINE

Caution The risk of drug dependence is high.

Note Real time online applications for increased maximum quantities/repeats may be made using the Online PBS Authorities system (see www.servicesaustralia.gov.au/organisations/health-professionals/services/medicare/hpos/services/request-authority-using-online-pbs-authorities-hpos).

Phone applications for increased maximum quantities/repeats may be made by calling 1800 888 333.

Written authority applications for increased maximum quantities/repeats can be uploaded online through HPOS form upload or mailed to:

Pharmaceutical Benefits Scheme

Reply Paid 9857

[Your capital city]

Restricted benefit

Severe pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for less than 12 months

Clinical criteria:

- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid analgesics; OR
- Patient must be unable to use non-opioid analgesics due to contraindications or intolerance.

Authorities for increased maximum quantities and/or repeats under this restriction must only be considered for severe disabling pain associated with malignant neoplasia or chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment is less than 12 months.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Restricted benefit

Severe pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for more than 12 months

Clinical criteria:

- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid analgesics; OR
- Patient must be unable to use non-opioid analgesics due to contraindications or intolerance.

Authorities for increased maximum quantities and/or repeats must only be considered for:

- severe disabling pain associated with proven malignant neoplasia; or
- palliative care patients with chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment exceeds 12 months and the patient is unable to have annual pain management review due to their clinical condition; or
- chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or
- chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment has exceeded 12 months prior to 1 June 2020 and the patient's clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Restricted benefit

Severe pain

Treatment Phase: Continuing PBS treatment after 1 June 2020

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this form of this drug for this condition after 1 June 2020.

Authorities for increased maximum quantities and/or repeats must only be considered where the patient has received initial authority approval for:

- severe disabling pain associated with malignant neoplasia; or
- chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment is less than 12 months; or
- palliative care patients with chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment exceeds 12 months and the patient is unable to have annual pain management review due to their clinical condition; or
- chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or
- chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment has exceeded 12 months prior to 1 June 2020 and the patient's clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

codeine phosphate hemihydrate 30 mg tablet, 20

1214X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	22.32	23.61	Aspen Pharma Pty Ltd [AS]

■ HYDROMORPHONE

Caution The risk of drug dependence is high.

Note Prescribing of drugs of addiction by dentists is not permitted in some States/Territories.

Restricted benefit

Severe pain

Clinical criteria:

- The treatment must be for short term therapy of acute severe pain, **AND**
- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid and other opioid analgesics; OR
- Patient must be unable to use non-opioid and other opioid analgesics due to contraindications or intolerance.

hydromorphone hydrochloride 2 mg tablet, 20

12045Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	0.5	..	..	*22.80	24.09	Dilaudid [MF]

hydromorphone hydrochloride 4 mg tablet, 20

12032G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	0.5	..	..	*24.21	25.50	Dilaudid [MF]

hydromorphone hydrochloride 8 mg tablet, 20

12010D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	0.5	..	..	*29.42	30.71	Dilaudid [MF]

■ HYDROMORPHONE

Caution The risk of drug dependence is high.

Note Prescribing of drugs of addiction by dentists is not permitted in some States/Territories.

Restricted benefit

Severe pain

Clinical criteria:

- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid and other opioid analgesics; OR
- Patient must be unable to use non-opioid and other opioid analgesics due to contraindications or intolerance.

hydromorphone hydrochloride 2 mg tablet, 20

5115F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	1	..	..	23.41	24.70	Dilaudid [MF]

hydromorphone hydrochloride 4 mg tablet, 20

5116G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	1	..	..	25.68	26.97	Dilaudid [MF]

hydromorphone hydrochloride 8 mg tablet, 20

5117H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	1	..	..	34.09	35.38	Dilaudid [MF]

hydromorphone hydrochloride 1 mg/mL oral liquid, 200 mL

11479E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	1	..	..	48.11	41.30	Dilaudid [MF]

■ HYDROMORPHONE

Caution The risk of drug dependence is high.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Restricted benefit

Severe pain

Clinical criteria:

- The treatment must be for short term therapy of acute severe pain, **AND**
- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid and other opioid analgesics; OR
- Patient must be unable to use non-opioid and other opioid analgesics due to contraindications or intolerance.

hydromorphone hydrochloride 2 mg tablet, 20

12047C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	0.5	..	..	*22.80	24.09	Dilaudid [MF]

hydromorphone hydrochloride 4 mg tablet, 20

12046B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	0.5	..	..	*24.21	25.50	Dilaudid [MF]

hydromorphone hydrochloride 8 mg tablet, 20

12016K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	0.5	..	..	*29.42	30.71	Dilaudid [MF]

■ HYDROMORPHONE

Caution The risk of drug dependence is high.

Note Consider consultation with a multidisciplinary pain service prior to, or after commencement of this medication.

Note Real time online applications for increased maximum quantities/repeats may be made using the Online PBS Authorities system (see www.servicesaustralia.gov.au/organisations/health-professionals/services/medicare/hpos/services/request-authority-using-online-pbs-authorities-hpos).

Phone applications for increased maximum quantities/repeats may be made by calling 1800 888 333.

Written authority applications for increased maximum quantities/repeats can be uploaded online through HPOS form upload or mailed to:

Pharmaceutical Benefits Scheme

Reply Paid 9857

[Your capital city]

Restricted benefit

Severe pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for less than 12 months

Clinical criteria:

- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid and other opioid analgesics; OR
- Patient must be unable to use non-opioid and other opioid analgesics due to contraindications or intolerance.

Authorities for increased maximum quantities and/or repeats under this restriction must only be considered for severe disabling pain associated with malignant neoplasia or chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment is less than 12 months.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Restricted benefit

Severe pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for more than 12 months

Clinical criteria:

- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid and other opioid analgesics; OR
- Patient must be unable to use non-opioid and other opioid analgesics due to contraindications or intolerance.

Authorities for increased maximum quantities and/or repeats must only be considered for:

- severe disabling pain associated with proven malignant neoplasia; or
- palliative care patients with chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment exceeds 12 months and the patient is unable to have annual pain management review due to their clinical condition; or
- chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or
- chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment has exceeded 12 months prior to 1 June 2020 and the patient's clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Restricted benefit

Severe pain

Treatment Phase: Continuing PBS treatment after 1 June 2020

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this form of this drug for this condition after 1 June 2020.

Authorities for increased maximum quantities and/or repeats must only be considered where the patient has received initial authority approval for:

- (i) severe disabling pain associated with malignant neoplasia; or
- (ii) chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment is less than 12 months; or
- (iii) palliative care patients with chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment exceeds 12 months and the patient is unable to have annual pain management review due to their clinical condition; or
- (iv) chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or
- (v) chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment has exceeded 12 months prior to 1 June 2020 and the patient's clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

hydromorphone hydrochloride 2 mg tablet, 20

8541M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	23.41	24.70	Dilaudid [MF]

hydromorphone hydrochloride 4 mg tablet, 20

8542N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	25.68	26.97	Dilaudid [MF]

hydromorphone hydrochloride 8 mg tablet, 20

8543P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	34.09	35.38	Dilaudid [MF]

hydromorphone hydrochloride 1 mg/mL oral liquid, 200 mL

11467M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	48.11	41.30	Dilaudid [MF]

hydromorphone hydrochloride 2 mg/mL injection, 5 x 1 mL ampoules

8420E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	27.65	28.94	^a Dilaudid [MF]	^a HYDROMORPHONE JUNO [JU]
						^a MEDSURGE HYDROMORPHONE 2 mg/1 mL [DZ]	

hydromorphone hydrochloride 10 mg/mL injection, 5 x 1 mL ampoules

8421F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	31.26	32.55	^a Dilaudid-HP [MF]	^a HYDROMORPHONE JUNO-HP [JU]
						^a MEDSURGE HYDROMORPHONE HP 10 mg/1 mL [DZ]	

■ HYDROMORPHONE

Caution The risk of drug dependence is high.

Note This treatment is not suitable for 'as-required' pain relief.

Note Consider consultation with a multidisciplinary pain service prior to, or after commencement of this medication.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note Real time online applications for increased maximum quantities/repeats may be made using the Online PBS Authorities system (see www.servicesaustralia.gov.au/organisations/health-professionals/services/medicare/hpos/services/request-authority-using-online-pbs-authorities-hpos).

Phone applications for increased maximum quantities/repeats may be made by calling 1800 888 333.

Written authority applications for increased maximum quantities/repeats can be uploaded online through HPOS form upload or mailed to:

Pharmaceutical Benefits Scheme

Reply Paid 9857

[Your capital city]

Authority required (STREAMLINED)**10754**

Chronic severe pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for less than 12 months

Clinical criteria:

- The condition must require daily, continuous, long term opioid treatment, **AND**
- Patient must not be opioid naive, **AND**
- Patient must have cancer pain; OR
- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid and other opioid analgesics; OR
- Patient must be unable to use non-opioid and other opioid analgesics due to contraindications or intolerance.

Authorities for increased maximum quantities and/or repeats under this restriction must only be considered for chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment is less than 12 months.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Authority required (STREAMLINED)**10753**

Chronic severe pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for more than 12 months

Clinical criteria:

- The condition must require daily, continuous, long term opioid treatment, **AND**
- Patient must not be opioid naive, **AND**
- Patient must have cancer pain; OR
- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid and other opioid analgesics; OR
- Patient must be unable to use non-opioid and other opioid analgesics due to contraindications or intolerance.

Authorities for increased maximum quantities and/or repeats must only be considered for chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment:

(i) exceeds 12 months and the palliative care patient is unable to have annual pain management review due to their clinical condition; or

(ii) exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through

consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or

(iii) has exceeded 12 months prior to 1 June 2020 and the patient's clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Authority required (STREAMLINED)**10752**

Chronic severe pain

Treatment Phase: Continuing PBS treatment after 1 June 2020

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this form of this drug for this condition after 1 June 2020.

Authorities for increased maximum quantities and/or repeats must only be considered for chronic severe disabling pain where the patient has received initial authority approval and the total duration of non-PBS and PBS opioid analgesic treatment:

(i) is less than 12 months; or

(ii) exceeds 12 months and the palliative care patient is unable to have annual pain management review due to their clinical condition; or

(iii) exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through

consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or

(iv) has exceeded 12 months prior to 1 June 2020 and the patient's pain management and clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

hydromorphone hydrochloride 16 mg modified release tablet, 14

9407D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	48.66	41.30	Jurnista [JC]

hydromorphone hydrochloride 32 mg modified release tablet, 14

9408E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	75.58	41.30	Jurnista [JC]

hydromorphone hydrochloride 8 mg modified release tablet, 14

9406C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	36.09	37.38	Jurnista [JC]

hydromorphone hydrochloride 4 mg modified release tablet, 14

9299K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	32.23	33.52	Jurnista [JC]

hydromorphone hydrochloride 64 mg modified release tablet, 14

9409F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	120.56	41.30	Jurnista [JC]

■ MORPHINE

Caution The risk of drug dependence is high.

Note Prescribing of drugs of addiction by dentists is not permitted in some States/Territories.

Restricted benefit

Severe pain

Clinical criteria:

- The treatment must be for short term therapy of acute severe pain, **AND**
- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid and other opioid analgesics; OR
- Patient must be unable to use non-opioid and other opioid analgesics due to contraindications or intolerance.

morphine sulfate pentahydrate 30 mg tablet, 20

12067D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	0.5	..	..	*21.43	22.72	Anamorph [RW]

■ MORPHINE

Caution The risk of drug dependence is high.

Note Prescribing of drugs of addiction by dentists is not permitted in some States/Territories.

Restricted benefit

Severe pain

Clinical criteria:

- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid and other opioid analgesics; OR
- Patient must be unable to use non-opioid and other opioid analgesics due to contraindications or intolerance; OR
- The treatment must be part of pre-operative care; OR
- The treatment must be used as an analgesic adjunct in general anaesthesia.

morphine sulfate pentahydrate 15 mg/mL injection, 5 x 1 mL ampoules

5169C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
DP	1	..	..	24.54	25.83	^a DBL Morphine Sulfate Pentahydrate [PF]	^a MORPHINE SULFATE 15 mg/1 mL MEDSURGE [DZ]

morphine sulfate pentahydrate 30 mg/mL injection, 5 x 1 mL ampoules

5170D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
DP	1	..	..	26.64	27.93	^a DBL Morphine Sulfate Pentahydrate [PF]	^a MORPHINE SULFATE 30 mg/1 mL MEDSURGE [DZ]

morphine hydrochloride trihydrate 20 mg/mL injection, 5 x 1 mL ampoules

10858L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	1	..	..	26.50	27.79	Morphine Juno [JU]

■ MORPHINE

Caution The risk of drug dependence is high.

Note Prescribing of drugs of addiction by dentists is not permitted in some States/Territories.

Restricted benefit

Severe pain

Clinical criteria:

- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid and other opioid analgesics; OR
- Patient must be unable to use non-opioid and other opioid analgesics due to contraindications or intolerance.

morphine hydrochloride trihydrate 2 mg/mL oral liquid, 200 mL

5237P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	1	..	..	26.56	27.85	Ordine 2 [MF]

morphine hydrochloride trihydrate 10 mg/mL oral liquid, 200 mL

5239R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	1	..	..	32.24	33.53	Ordine 10 [MF]

morphine sulfate pentahydrate 30 mg tablet, 20

5163R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	1	..	..	21.19	22.48	Anamorph [RW]

morphine hydrochloride trihydrate 5 mg/mL oral liquid, 200 mL

5238Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	1	..	..	28.65	29.94	Ordine 5 [MF]

■ MORPHINE**Caution** The risk of drug dependence is high.**Note** No increase in the maximum quantity or number of units may be authorised.**Note** No increase in the maximum number of repeats may be authorised.**Restricted benefit**

Severe pain

Clinical criteria:

- The treatment must be for short term therapy of acute severe pain, **AND**
- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid and other opioid analgesics; OR
- Patient must be unable to use non-opioid and other opioid analgesics due to contraindications or intolerance.

morphine sulfate pentahydrate 30 mg tablet, 20

12009C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	0.5	..	..	*21.43	22.72	Anamorph [RW]

■ MORPHINE**Caution** The risk of drug dependence is high.**Note** Prescribing of drugs of addiction by dentists is not permitted in some States/Territories.**Note** Pharmaceutical benefits that have the forms morphine sulfate 10 mg/mL injection and morphine hydrochloride 10 mg/mL injection are equivalent for the purposes of substitution.**Restricted benefit**

Severe pain

Clinical criteria:

- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid and other opioid analgesics; OR
- Patient must be unable to use non-opioid and other opioid analgesics due to contraindications or intolerance; OR
- The treatment must be part of pre-operative care; OR
- The treatment must be used as an analgesic adjunct in general anaesthesia.

morphine sulfate pentahydrate 10 mg/mL injection, 5 x 1 mL ampoules

5168B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
DP	1	..	..	23.14	24.43	^a DBL Morphine Sulfate Pentahydrate [PF]	^a MORPHINE SULFATE 10 mg/1 mL MEDSURGE [DZ]

morphine hydrochloride trihydrate 10 mg/mL injection, 5 x 1 mL ampoules

10863R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	1	..	..	23.14	24.43	^a Morphine Juno [JU]

■ MORPHINE**Caution** The risk of drug dependence is high.**Note** Consider consultation with a multidisciplinary pain service prior to, or after commencement of this medication.**Note** Real time online applications for increased maximum quantities/repeats may be made using the Online PBS Authorities system (see www.servicesaustralia.gov.au/organisations/health-professionals/services/medicare/hpos/services/request-authority-using-online-pbs-authorities-hpos).

Phone applications for increased maximum quantities/repeats may be made by calling 1800 888 333.

Written authority applications for increased maximum quantities/repeats can be uploaded online through HPOS form upload

or mailed to:
Pharmaceutical Benefits Scheme
Reply Paid 9857
[Your capital city]

Restricted benefit

Severe pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for less than 12 months

Clinical criteria:

- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid and other opioid analgesics; OR
- Patient must be unable to use non-opioid and other opioid analgesics due to contraindications or intolerance; OR
- The treatment must be part of pre-operative care; OR
- The treatment must be used as an analgesic adjunct in general anaesthesia.

Authorities for increased maximum quantities and/or repeats under this restriction must only be considered for severe disabling pain associated with malignant neoplasia or chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment is less than 12 months.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Restricted benefit

Severe pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for more than 12 months

Clinical criteria:

- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid and other opioid analgesics; OR
- Patient must be unable to use non-opioid and other opioid analgesics due to contraindications or intolerance; OR
- The treatment must be part of pre-operative care; OR
- The treatment must be used as an analgesic adjunct in general anaesthesia.

Authorities for increased maximum quantities and/or repeats must only be considered for:

- severe disabling pain associated with proven malignant neoplasia; or
- palliative care patients with chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment exceeds 12 months and the patient is unable to have annual pain management review due to their clinical condition; or
- chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or
- chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment has exceeded 12 months prior to 1 June 2020 and the patient's clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Restricted benefit

Severe pain

Treatment Phase: Continuing PBS treatment after 1 June 2020

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this form of this drug for this condition after 1 June 2020.

Authorities for increased maximum quantities and/or repeats must only be considered where the patient has received initial authority approval for:

- severe disabling pain associated with malignant neoplasia; or
- chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment is less than 12 months; or
- palliative care patients with chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment exceeds 12 months and the patient is unable to have annual pain management review due to their clinical condition; or
- chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or
- chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment has exceeded 12 months prior to 1 June 2020 and the patient's clinical need for continuing opioid treatment has not been confirmed through

consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

morphine sulfate pentahydrate 15 mg/mL injection, 5 x 1 mL ampoules

1645N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP MW	1	..	..	24.54	25.83	^a DBL Morphine Sulfate Pentahydrate [PF]	^a MORPHINE SULFATE 15 mg/1 mL MEDSURGE [DZ]

morphine sulfate pentahydrate 30 mg/mL injection, 5 x 1 mL ampoules

1647Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	26.64	27.93	^a DBL Morphine Sulfate Pentahydrate [PF]	^a MORPHINE SULFATE 30 mg/1 mL MEDSURGE [DZ]

morphine hydrochloride trihydrate 50 mg/5 mL injection, 5 x 5 mL ampoules

10869C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	32.95	34.24	Morphine Juno [JU]

morphine hydrochloride trihydrate 20 mg/mL injection, 5 x 1 mL ampoules

10874H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	26.50	27.79	Morphine Juno [JU]

morphine hydrochloride trihydrate 100 mg/5 mL injection, 5 x 5 mL ampoules

10878M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	43.70	41.30	Morphine Juno [JU]

■ MORPHINE

Caution The risk of drug dependence is high.

Note Consider consultation with a multidisciplinary pain service prior to, or after commencement of this medication.

Note Real time online applications for increased maximum quantities/repeats may be made using the Online PBS Authorities system (see www.servicesaustralia.gov.au/organisations/health-professionals/services/medicare/hpos/services/request-authority-using-online-pbs-authorities-hpos).

Phone applications for increased maximum quantities/repeats may be made by calling 1800 888 333.

Written authority applications for increased maximum quantities/repeats can be uploaded online through HPOS form upload or mailed to:

Pharmaceutical Benefits Scheme

Reply Paid 9857

[Your capital city]

Restricted benefit

Severe pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for less than 12 months

Clinical criteria:

- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid and other opioid analgesics; OR

- Patient must be unable to use non-opioid and other opioid analgesics due to contraindications or intolerance.

Authorities for increased maximum quantities and/or repeats under this restriction must only be considered for severe disabling pain associated with malignant neoplasia or chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment is less than 12 months.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Restricted benefit

Severe pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for more than 12 months

Clinical criteria:

- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid and other opioid analgesics; OR

- Patient must be unable to use non-opioid and other opioid analgesics due to contraindications or intolerance.

Authorities for increased maximum quantities and/or repeats must only be considered for:

- (i) severe disabling pain associated with proven malignant neoplasia; or

(ii) palliative care patients with chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment exceeds 12 months and the patient is unable to have annual pain management review due to their clinical condition; or

(iii) chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or

(iv) chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment has exceeded 12 months prior to 1 June 2020 and the patient's clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Restricted benefit

Severe pain

Treatment Phase: Continuing PBS treatment after 1 June 2020

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this form of this drug for this condition after 1 June 2020.

Authorities for increased maximum quantities and/or repeats must only be considered where the patient has received initial authority approval for:

(i) severe disabling pain associated with malignant neoplasia; or

(ii) chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment is less than 12 months; or

(iii) palliative care patients with chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment exceeds 12 months and the patient is unable to have annual pain management review due to their clinical condition; or

(iv) chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or

(v) chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment has exceeded 12 months prior to 1 June 2020 and the patient's clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

morphine hydrochloride trihydrate 2 mg/mL oral liquid, 200 mL

2122Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	26.56	27.85	Ordine 2 [MF]

morphine hydrochloride trihydrate 10 mg/mL oral liquid, 200 mL

2124T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	32.24	33.53	Ordine 10 [MF]

morphine sulfate pentahydrate 30 mg tablet, 20

1646P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	21.19	22.48	Anamorph [RW]

morphine hydrochloride trihydrate 5 mg/mL oral liquid, 200 mL

2123R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	28.65	29.94	Ordine 5 [MF]

■ MORPHINE

Caution The risk of drug dependence is high.

Note Consider consultation with a multidisciplinary pain service prior to, or after commencement of this medication.

Note Real time online applications for increased maximum quantities/repeats may be made using the Online PBS Authorities system (see www.servicesaustralia.gov.au/organisations/health-professionals/services/medicare/hpos/services/request-authority-using-online-pbs-authorities-hpos).

Phone applications for increased maximum quantities/repeats may be made by calling 1800 888 333.

Written authority applications for increased maximum quantities/repeats can be uploaded online through HPOS form upload or mailed to:

Pharmaceutical Benefits Scheme

Reply Paid 9857
[Your capital city]

Restricted benefit

Cancer pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for less than 12 months

Clinical criteria:

- Patient must have cancer pain, **AND**
- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid and other opioid analgesics; OR
- Patient must be unable to use non-opioid and other opioid analgesics due to contraindications or intolerance.

Authorities for increased maximum quantities and/or repeats under this restriction must only be considered for chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment is less than 12 months.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Restricted benefit

Cancer pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for more than 12 months

Clinical criteria:

- Patient must have cancer pain, **AND**
- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid and other opioid analgesics; OR
- Patient must be unable to use non-opioid and other opioid analgesics due to contraindications or intolerance.

Authorities for increased maximum quantities and/or repeats must only be considered for:

(i) palliative care patients with chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment exceeds 12 months and the patient is unable to have annual pain management review due to their clinical condition; or

(ii) chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or

(iii) chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment has exceeded 12 months prior to 1 June 2020 and the patient's clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Restricted benefit

Cancer pain

Treatment Phase: Continuing PBS treatment after 1 June 2020

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this form of this drug for this condition after 1 June 2020.

Authorities for increased maximum quantities and/or repeats must only be considered for chronic severe disabling pain where the patient has received initial authority approval and the total duration of non-PBS and PBS opioid analgesic treatment:

(i) is less than 12 months; or

(ii) exceeds 12 months and the palliative care patient is unable to have annual pain management review due to their clinical condition; or

(iii) exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or

(iv) has exceeded 12 months prior to 1 June 2020 and the patient's pain management and clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

morphine sulfate pentahydrate 10 mg tablet, 20

8669G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	21.42	22.71	Sevredol [MF]

morphine sulfate pentahydrate 20 mg tablet, 20

8670H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	22.19	23.48	Sevredol [MF]

■ MORPHINE

Caution The risk of drug dependence is high.

Note Consider consultation with a multidisciplinary pain service prior to, or after commencement of this medication.

Note Pharmaceutical benefits that have the forms morphine sulfate 10 mg/mL injection and morphine hydrochloride 10 mg/mL injection are equivalent for the purposes of substitution.

Note Real time online applications for increased maximum quantities/repeats may be made using the Online PBS Authorities system (see www.servicesaustralia.gov.au/organisations/health-professionals/services/medicare/hpos/services/request-authority-using-online-pbs-authorities-hpos).

Phone applications for increased maximum quantities/repeats may be made by calling 1800 888 333.

Written authority applications for increased maximum quantities/repeats can be uploaded online through HPOS form upload or mailed to:

Pharmaceutical Benefits Scheme

Reply Paid 9857

[Your capital city]

Restricted benefit

Severe pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for less than 12 months

Clinical criteria:

- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid and other opioid analgesics; OR
- Patient must be unable to use non-opioid and other opioid analgesics due to contraindications or intolerance; OR
- The treatment must be part of pre-operative care; OR
- The treatment must be used as an analgesic adjunct in general anaesthesia.

Authorities for increased maximum quantities and/or repeats under this restriction must only be considered for severe disabling pain associated with malignant neoplasia or chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment is less than 12 months.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Restricted benefit

Severe pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for more than 12 months

Clinical criteria:

- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid and other opioid analgesics; OR
- Patient must be unable to use non-opioid and other opioid analgesics due to contraindications or intolerance; OR
- The treatment must be part of pre-operative care; OR
- The treatment must be used as an analgesic adjunct in general anaesthesia.

Authorities for increased maximum quantities and/or repeats must only be considered for:

- severe disabling pain associated with proven malignant neoplasia; or
- palliative care patients with chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment exceeds 12 months and the patient is unable to have annual pain management review due to their clinical condition; or
- chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or
- chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment has exceeded 12 months prior to 1 June 2020 and the patient's clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Restricted benefit

Severe pain

Treatment Phase: Continuing PBS treatment after 1 June 2020

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this form of this drug for this condition after 1 June 2020.

Authorities for increased maximum quantities and/or repeats must only be considered where the patient has received initial authority approval for:

- (i) severe disabling pain associated with malignant neoplasia; or
- (ii) chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment is less than 12 months; or
- (iii) palliative care patients with chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment exceeds 12 months and the patient is unable to have annual pain management review due to their clinical condition; or
- (iv) chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or
- (v) chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment has exceeded 12 months prior to 1 June 2020 and the patient's clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

morphine sulfate pentahydrate 10 mg/mL injection, 5 x 1 mL ampoules

1644M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
 	1	..	..	23.14	24.43	^a DBL Morphine Sulfate Pentahydrate [PF]	^a MORPHINE SULFATE 10 mg/1 mL MEDSURGE [DZ]

morphine hydrochloride trihydrate 10 mg/mL injection, 5 x 1 mL ampoules

10864T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
 	1	..	..	23.14	24.43	^a Morphine Juno [JU]

■ **MORPHINE**

Caution The risk of drug dependence is high.

Note This treatment is not suitable for 'as-required' pain relief.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note Real time online applications for increased maximum quantities/repeats may be made using the Online PBS Authorities system (see www.servicesaustralia.gov.au/organisations/health-professionals/services/medicare/hpos/services/request-authority-using-online-pbs-authorities-hpos).

Phone applications for increased maximum quantities/repeats may be made by calling 1800 888 333.

Written authority applications for increased maximum quantities/repeats can be uploaded online through HPOS form upload or mailed to:

Pharmaceutical Benefits Scheme

Reply Paid 9857

[Your capital city]

Authority required

Chronic severe disabling pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for less than 12 months

Clinical criteria:

- The condition must require daily, continuous, long term opioid treatment, **AND**
- Patient must have cancer pain; OR
- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid or other opioid analgesics; OR
- Patient must be unable to use non-opioid or other opioid analgesics due to contraindications or intolerance.

Authorities for increased maximum quantities and/or repeats under this restriction must only be considered for chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment is less than 12 months.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Authority required

Chronic severe disabling pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for more than 12 months

Clinical criteria:

- The condition must require daily, continuous, long term opioid treatment, **AND**
- Patient must have cancer pain; OR
- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid or other opioid analgesics; OR
- Patient must be unable to use non-opioid or other opioid analgesics due to contraindications or intolerance.

Authorities for increased maximum quantities and/or repeats must only be considered for chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment:

(i) exceeds 12 months and the palliative care patient is unable to have annual pain management review due to their clinical condition; or

(ii) exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or

(iii) has exceeded 12 months prior to 1 June 2020 and the patient's clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Authority required

Chronic severe disabling pain

Treatment Phase: Continuing PBS treatment after 1 June 2020

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this form of this drug for this condition after 1 June 2020.

Authorities for increased maximum quantities and/or repeats must only be considered for chronic severe disabling pain where the patient has received initial authority approval and the total duration of non-PBS and PBS opioid analgesic treatment:

(i) is less than 12 months; or

(ii) exceeds 12 months and the palliative care patient is unable to have annual pain management review due to their clinical condition; or

(iii) exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or

(iv) has exceeded 12 months prior to 1 June 2020 and the patient's pain management and clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

morphine sulfate pentahydrate 200 mg modified release granules, 28 sachets

12053J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	159.41	41.30	MS Contin Suspension 200 mg [MF]

morphine sulfate pentahydrate 200 mg modified release tablet, 28

12055L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	119.42	41.30	MS Contin [MF]

■ **MORPHINE**

Caution The risk of drug dependence is high.

Note This treatment is not suitable for 'as-required' pain relief.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note Real time online applications for increased maximum quantities/repeats may be made using the Online PBS Authorities system (see www.servicesaustralia.gov.au/organisations/health-professionals/services/medicare/hpos/services/request-authority-using-online-pbs-authorities-hpos).

Phone applications for increased maximum quantities/repeats may be made by calling 1800 888 333.

Written authority applications for increased maximum quantities/repeats can be uploaded online through HPOS form upload or mailed to:

Pharmaceutical Benefits Scheme
Reply Paid 9857
[Your capital city]

Authority required (STREAMLINED)

10755

Chronic severe pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for less than 12 months

Clinical criteria:

- The condition must require daily, continuous, long term opioid treatment, **AND**
- Patient must have cancer pain; OR
- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid or other opioid analgesics; OR
- Patient must be unable to use non-opioid or other opioid analgesics due to contraindications or intolerance.

Authorities for increased maximum quantities and/or repeats under this restriction must only be considered for chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment is less than 12 months.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Authority required (STREAMLINED)

10748

Chronic severe pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for more than 12 months

Clinical criteria:

- The condition must require daily, continuous, long term opioid treatment, **AND**
- Patient must have cancer pain; OR
- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid or other opioid analgesics; OR
- Patient must be unable to use non-opioid or other opioid analgesics due to contraindications or intolerance.

Authorities for increased maximum quantities and/or repeats must only be considered for chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment:

(i) exceeds 12 months and the palliative care patient is unable to have annual pain management review due to their clinical condition; or

(ii) exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or

(iii) has exceeded 12 months prior to 1 June 2020 and the patient's clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Authority required (STREAMLINED)

10752

Chronic severe pain

Treatment Phase: Continuing PBS treatment after 1 June 2020

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this form of this drug for this condition after 1 June 2020.

Authorities for increased maximum quantities and/or repeats must only be considered for chronic severe disabling pain where the patient has received initial authority approval and the total duration of non-PBS and PBS opioid analgesic treatment:

(i) is less than 12 months; or

(ii) exceeds 12 months and the palliative care patient is unable to have annual pain management review due to their clinical condition; or

(iii) exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or

(iv) has exceeded 12 months prior to 1 June 2020 and the patient's pain management and clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

morphine sulfate pentahydrate 10 mg modified release capsule, 28

8349K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	26.30	27.59	Kapanol [YN]

morphine sulfate pentahydrate 60 mg modified release tablet, 28

1655D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	57.67	41.30	^a Momex SR 60 [RW] ^a MORPHINE MR APOTEX [TX] ^a MS Contin [MF]	^a Morphine MR AN [EA] ^a Morphine MR Mylan [AF]

morphine sulfate pentahydrate 20 mg modified release granules, 28 sachets

8490W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	63.33	41.30	MS Contin Suspension 20 mg [MF]

morphine sulfate pentahydrate 100 mg modified release granules, 28 sachets

8306E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	87.04	41.30	MS Contin Suspension 100 mg [MF]

morphine sulfate pentahydrate 100 mg modified release tablet, 28

1656E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	74.44	41.30	^a Momex SR 100 [RW] ^a MORPHINE MR APOTEX [TX] ^a MS Contin [MF]	^a Morphine MR AN [EA] ^a Morphine MR Mylan [AF]

morphine sulfate pentahydrate 20 mg modified release capsule, 28

2839K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	30.63	31.92	Kapanol [YN]

morphine sulfate pentahydrate 30 mg modified release tablet, 28

1654C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	40.09	41.30	^a Momex SR 30 [RW] ^a MORPHINE MR APOTEX [TX] ^a MS Contin [MF]	^a Morphine MR AN [EA] ^a Morphine MR Mylan [AF]

morphine sulfate pentahydrate 30 mg modified release granules, 28 sachets

8146R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	64.95	41.30	MS Contin Suspension 30 mg [MF]

morphine sulfate pentahydrate 120 mg modified release capsule, 14

8494C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	57.66	41.30	MS Mono [MF]

morphine sulfate pentahydrate 30 mg modified release capsule, 14

8491X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	29.94	31.23	MS Mono [MF]

morphine sulfate pentahydrate 90 mg modified release capsule, 14

8493B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	44.90	41.30	MS Mono [MF]

morphine sulfate pentahydrate 10 mg modified release tablet, 28

1653B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	26.31	27.60	^a Momex SR 10 [RW] ^a MORPHINE MR APOTEX [TX] ^a MS Contin [MF]	^a Morphine MR AN [EA] ^a Morphine MR Mylan [AF]

morphine sulfate pentahydrate 60 mg modified release capsule, 14

8492Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	40.07	41.30	MS Mono [MF]

morphine sulfate pentahydrate 50 mg modified release capsule, 28

2840L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	46.54	41.30	Kapanol [YN]

morphine sulfate pentahydrate 100 mg modified release capsule, 28

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
2841M	1	..	..	72.58	41.30	Kapanol [YN]

morphine sulfate pentahydrate 5 mg modified release tablet, 28

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8035X	1	..	..	24.19	25.48	MS Contin [MF]

morphine sulfate pentahydrate 15 mg modified release tablet, 28

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8489T	1	..	..	29.95	31.24	MS Contin [MF]

morphine sulfate pentahydrate 60 mg modified release granules, 28 sachets

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8305D	1	..	..	72.04	41.30	MS Contin Suspension 60 mg [MF]

■ MORPHINE

Caution The risk of drug dependence is high.

Note This treatment is not suitable for 'as-required' pain relief.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Authority required

Chronic severe disabling pain

Clinical criteria:

- The condition must require daily, continuous, long term opioid treatment, **AND**
- Patient must have cancer pain; OR
- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid or other opioid analgesics; OR
- Patient must be unable to use non-opioid or other opioid analgesics due to contraindications or intolerance.

morphine sulfate pentahydrate 200 mg modified release granules, 28 sachets

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8454Y	1	..	..	159.41	41.30	MS Contin Suspension 200 mg [MF]

morphine sulfate pentahydrate 200 mg modified release tablet, 28

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8453X	1	..	..	119.42	41.30	MS Contin [MF]

■ OXYCODONE

Caution The risk of drug dependence is high.

Note Prescribing of drugs of addiction by dentists is not permitted in some States/Territories.

Restricted benefit

Severe pain

Clinical criteria:

- The treatment must be for short term therapy of acute severe pain, **AND**
- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid analgesics; OR
- Patient must be unable to use non-opioid analgesics due to contraindications or intolerance.

oxycodone hydrochloride 5 mg capsule, 10

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
12311Y	1	..	..	19.17	20.46	OxyNorm [MF]

oxycodone hydrochloride 5 mg tablet, 20

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
12048D	0.5	..	..	*20.09	21.38	^a Mayne Pharma Oxycodone IR [YN]	^a Oxycodone Mylan [AL]
			^B 1.16	*21.25	21.38	^a Endone [AF]	

oxycodone hydrochloride 10 mg capsule, 20

12074L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
DP	0.5	..	..	*21.63	22.92	^a Oxycodone BNM [BZ]	^a OxyNorm [MF]

■ OXYCODONE

Caution The risk of drug dependence is high.

Note Prescribing of drugs of addiction by dentists is not permitted in some States/Territories.

Restricted benefit

Severe pain

Clinical criteria:

- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid analgesics; OR
- Patient must be unable to use non-opioid analgesics due to contraindications or intolerance.

oxycodone hydrochloride 5 mg tablet, 20

5195K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
DP	1	..	..	19.04	20.33	^a Mayne Pharma Oxycodone IR [YN]	^a Oxycodone Mylan [AL]
			^b 1.86	20.90	20.33	^a Endone [AF]	

oxycodone hydrochloride 5 mg capsule, 20

5191F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
DP	1	..	..	21.11	22.40	^a Oxycodone BNM [BZ]	^a OxyNorm [MF]

oxycodone hydrochloride 10 mg capsule, 20

5197M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
DP	1	..	..	21.51	22.80	^a Oxycodone BNM [BZ]	^a OxyNorm [MF]

oxycodone hydrochloride 1 mg/mL oral liquid, 250 mL

5190E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	1	..	..	30.06	31.35	OxyNorm Liquid 1mg/mL [MF]

■ OXYCODONE

Caution The risk of drug dependence is high.

Note Prescribing of drugs of addiction by dentists is not permitted in some States/Territories.

Restricted benefit

Severe pain

Clinical criteria:

- The treatment must be for post-operative pain following a major operative procedure, **AND**
- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid analgesics; OR
- Patient must be unable to use non-opioid analgesics due to contraindications or intolerance.

oxycodone 30 mg suppository, 12

5194J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	1	..	..	49.17	41.30	Proladone [FF]

■ OXYCODONE

Caution The risk of drug dependence is high.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Restricted benefit

Severe pain

Clinical criteria:

- The treatment must be for short term therapy of acute severe pain, **AND**
- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid analgesics; OR
- Patient must be unable to use non-opioid analgesics due to contraindications or intolerance.

oxycodone hydrochloride 5 mg capsule, 10

12314D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	19.17	20.46	OxyNorm [MF]

oxycodone hydrochloride 5 mg tablet, 20

12023T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	0.5	..	..	*20.09	21.38	^a Mayne Pharma Oxycodone IR [YN]	^a Oxycodone Mylan [AL]
			^b 1.16	*21.25	21.38	^a Endone [AF]	

oxycodone hydrochloride 10 mg capsule, 20

12031F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	0.5	..	..	*21.63	22.92	^a Oxycodone BNM [BZ]	^a OxyNorm [MF]

■ OXYCODONE

Caution The risk of drug dependence is high.

Note Real time online applications for increased maximum quantities/repeats may be made using the Online PBS Authorities system (see www.servicesaustralia.gov.au/organisations/health-professionals/services/medicare/hpos/services/request-authority-using-online-pbs-authorities-hpos).

Phone applications for increased maximum quantities/repeats may be made by calling 1800 888 333.

Written authority applications for increased maximum quantities/repeats can be uploaded online through HPOS form upload or mailed to:

Pharmaceutical Benefits Scheme

Reply Paid 9857

[Your capital city]

Restricted benefit

Severe pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for less than 12 months

Clinical criteria:

- Patient must have cancer pain; OR
- The treatment must be for post-operative pain following a major operative procedure, **AND**
- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid analgesics; OR
- Patient must be unable to use non-opioid analgesics due to contraindications or intolerance.

Authorities for increased maximum quantities and/or repeats under this restriction must only be considered for severe disabling pain associated with malignant neoplasia or chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment is less than 12 months.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Restricted benefit

Severe pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for more than 12 months

Clinical criteria:

- Patient must have cancer pain; OR
- The treatment must be for post-operative pain following a major operative procedure, **AND**
- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid analgesics; OR
- Patient must be unable to use non-opioid analgesics due to contraindications or intolerance.

Authorities for increased maximum quantities and/or repeats must only be considered for:

- severe disabling pain associated with proven malignant neoplasia; or
- palliative care patients with chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment exceeds 12 months and the patient is unable to have annual pain management review due to their clinical condition; or
- chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or
- chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment has exceeded 12 months prior to 1 June 2020 and the patient's clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Restricted benefit

Severe pain

Treatment Phase: Continuing PBS treatment after 1 June 2020

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this form of this drug for this condition after 1 June 2020.

Authorities for increased maximum quantities and/or repeats must only be considered where the patient has received initial authority approval for:

- (i) severe disabling pain associated with malignant neoplasia; or
- (ii) chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment is less than 12 months; or
- (iii) palliative care patients with chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment exceeds 12 months and the patient is unable to have annual pain management review due to their clinical condition; or
- (iv) chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or
- (v) chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment has exceeded 12 months prior to 1 June 2020 and the patient's clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

oxycodone 30 mg suppository, 12

2481N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	49.17	41.30	Proladone [FF]

■ OXYCODONE

Caution The risk of drug dependence is high.

Note Real time online applications for increased maximum quantities/repeats may be made using the Online PBS Authorities system (see www.servicesaustralia.gov.au/organisations/health-professionals/services/medicare/hpos/services/request-authority-using-online-pbs-authorities-hpos).

Phone applications for increased maximum quantities/repeats may be made by calling 1800 888 333.

Written authority applications for increased maximum quantities/repeats can be uploaded online through HPOS form upload or mailed to:

Pharmaceutical Benefits Scheme

Reply Paid 9857

[Your capital city]

Restricted benefit

Severe pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for less than 12 months

Clinical criteria:

- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid analgesics; OR
 - Patient must be unable to use non-opioid analgesics due to contraindications or intolerance.
- Authorities for increased maximum quantities and/or repeats under this restriction must only be considered for severe disabling pain associated with malignant neoplasia or chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment is less than 12 months.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Restricted benefit

Severe pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for more than 12 months

Clinical criteria:

- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid analgesics; OR
- Patient must be unable to use non-opioid analgesics due to contraindications or intolerance.

Authorities for increased maximum quantities and/or repeats must only be considered for:

- (i) severe disabling pain associated with proven malignant neoplasia; or
- (ii) palliative care patients with chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment exceeds 12 months and the patient is unable to have annual pain management review due to their clinical condition; or
- (iii) chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or

(iv) chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment has exceeded 12 months prior to 1 June 2020 and the patient's clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Restricted benefit

Severe pain

Treatment Phase: Continuing PBS treatment after 1 June 2020

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this form of this drug for this condition after 1 June 2020.

Authorities for increased maximum quantities and/or repeats must only be considered where the patient has received initial authority approval for:

(i) severe disabling pain associated with malignant neoplasia; or

(ii) chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment is less than 12 months; or

(iii) palliative care patients with chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment exceeds 12 months and the patient is unable to have annual pain management review due to their clinical condition; or

(iv) chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or

(v) chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment has exceeded 12 months prior to 1 June 2020 and the patient's clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

oxycodone hydrochloride 20 mg capsule, 20

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
8502L	1	..	..	24.86	26.15	^a Oxycodone BNM [BZ]	^a OxyNorm [MF]

oxycodone hydrochloride 5 mg tablet, 20

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
2622B	1	..	..	19.04	20.33	^a Mayne Pharma Oxycodone IR [YN]	^a Oxycodone Mylan [AL]
			^b 1.86	20.90	20.33	^a Endone [AF]	

oxycodone hydrochloride 5 mg capsule, 20

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
8464L	1	..	..	21.11	22.40	^a Oxycodone BNM [BZ]	^a OxyNorm [MF]

oxycodone hydrochloride 10 mg capsule, 20

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
8501K	1	..	..	21.51	22.80	^a Oxycodone BNM [BZ]	^a OxyNorm [MF]

oxycodone hydrochloride 1 mg/mL oral liquid, 250 mL

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8644Y	1	..	..	30.06	31.35	OxyNorm Liquid 1mg/mL [MF]

■ OXYCODONE

Caution The risk of drug dependence is high.

Note This treatment is not suitable for 'as-required' pain relief.

Note OxyContin and Novacodone modified release tablets are intended to be crush-deterrent and to reduce the rapid release of oxycodone upon accidental or intentional misuse.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note Real time online applications for increased maximum quantities/repeats may be made using the Online PBS Authorities system (see www.servicesaustralia.gov.au/organisations/health-professionals/services/medicare/hpos/services/request-authority-using-online-pbs-authorities-hpos).
Phone applications for increased maximum quantities/repeats may be made by calling 1800 888 333.
Written authority applications for increased maximum quantities/repeats can be uploaded online through HPOS form upload or mailed to:
Pharmaceutical Benefits Scheme
Reply Paid 9857
[Your capital city]

Authority required (STREAMLINED)

10755

Chronic severe pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for less than 12 months

Clinical criteria:

- The condition must require daily, continuous, long term opioid treatment, **AND**
- Patient must have cancer pain; OR
- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid or other opioid analgesics; OR
- Patient must be unable to use non-opioid or other opioid analgesics due to contraindications or intolerance.

Authorities for increased maximum quantities and/or repeats under this restriction must only be considered for chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment is less than 12 months.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Authority required (STREAMLINED)

10748

Chronic severe pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for more than 12 months

Clinical criteria:

- The condition must require daily, continuous, long term opioid treatment, **AND**
- Patient must have cancer pain; OR
- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid or other opioid analgesics; OR
- Patient must be unable to use non-opioid or other opioid analgesics due to contraindications or intolerance.

Authorities for increased maximum quantities and/or repeats must only be considered for chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment:

(i) exceeds 12 months and the palliative care patient is unable to have annual pain management review due to their clinical condition; or

(ii) exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or

(iii) has exceeded 12 months prior to 1 June 2020 and the patient's clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Authority required (STREAMLINED)

10752

Chronic severe pain

Treatment Phase: Continuing PBS treatment after 1 June 2020

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this form of this drug for this condition after 1 June 2020.

Authorities for increased maximum quantities and/or repeats must only be considered for chronic severe disabling pain where the patient has received initial authority approval and the total duration of non-PBS and PBS opioid analgesic treatment:

(i) is less than 12 months; or

(ii) exceeds 12 months and the palliative care patient is unable to have annual pain management review due to their clinical condition; or

(iii) exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or

(iv) has exceeded 12 months prior to 1 June 2020 and the patient's pain management and clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

oxycodone hydrochloride 80 mg modified release tablet, 28

8388L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	72.87	41.30	^a Novacodone [HX] ^a OxyContin [MF]	^a Oxycodone Sandoz [SZ]

oxycodone hydrochloride 40 mg modified release tablet, 28

8387K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	51.03	41.30	^a Novacodone [HX] ^a OxyContin [MF]	^a Oxycodone Sandoz [SZ]

oxycodone hydrochloride 20 mg modified release tablet, 28

8386J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	36.30	37.59	^a Novacodone [HX] ^a OxyContin [MF]	^a Oxycodone Sandoz [SZ]

oxycodone hydrochloride 10 mg modified release tablet, 28

8385H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	27.82	29.11	^a Novacodone [HX] ^a OxyContin [MF]	^a Oxycodone Sandoz [SZ]

■ OXYCODONE

Caution The risk of drug dependence is high.

Note This treatment is not suitable for 'as-required' pain relief.

Note OxyContin modified release tablets are intended to be crush-deterrent and to reduce the rapid release of oxycodone upon accidental or intentional misuse.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note Real time online applications for increased maximum quantities/repeats may be made using the Online PBS Authorities system (see www.servicesaustralia.gov.au/organisations/health-professionals/services/medicare/hpos/services/request-authority-using-online-pbs-authorities-hpos).

Phone applications for increased maximum quantities/repeats may be made by calling 1800 888 333.

Written authority applications for increased maximum quantities/repeats can be uploaded online through HPOS form upload or mailed to:

Pharmaceutical Benefits Scheme

Reply Paid 9857

[Your capital city]

Authority required (STREAMLINED)

10755

Chronic severe pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for less than 12 months

Clinical criteria:

- The condition must require daily, continuous, long term opioid treatment, **AND**
- Patient must have cancer pain; OR
- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid or other opioid analgesics; OR
- Patient must be unable to use non-opioid or other opioid analgesics due to contraindications or intolerance.

Authorities for increased maximum quantities and/or repeats under this restriction must only be considered for chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment is less than 12 months.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Authority required (STREAMLINED)

10748

Chronic severe pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for more than 12 months

Clinical criteria:

- The condition must require daily, continuous, long term opioid treatment, **AND**
- Patient must have cancer pain; OR
- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid or other opioid analgesics; OR
- Patient must be unable to use non-opioid or other opioid analgesics due to contraindications or intolerance.

Authorities for increased maximum quantities and/or repeats must only be considered for chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment:

(i) exceeds 12 months and the palliative care patient is unable to have annual pain management review due to their clinical condition; or

(ii) exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or

(iii) has exceeded 12 months prior to 1 June 2020 and the patient's clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Authority required (STREAMLINED)**10752**

Chronic severe pain

Treatment Phase: Continuing PBS treatment after 1 June 2020

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this form of this drug for this condition after 1 June 2020.

Authorities for increased maximum quantities and/or repeats must only be considered for chronic severe disabling pain where the patient has received initial authority approval and the total duration of non-PBS and PBS opioid analgesic treatment:

(i) is less than 12 months; or

(ii) exceeds 12 months and the palliative care patient is unable to have annual pain management review due to their clinical condition; or

(iii) exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or

(iv) has exceeded 12 months prior to 1 June 2020 and the patient's pain management and clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

oxycodone hydrochloride 15 mg modified release tablet, 28

9399Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	36.06	37.35	OxyContin [MF]

oxycodone hydrochloride 30 mg modified release tablet, 28

9400R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	49.64	41.30	OxyContin [MF]

■ OXYCODONE + NALOXONE

Caution The risk of drug dependence is high.

Note This treatment is not suitable for 'as-required' pain relief.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note Real time online applications for increased maximum quantities/repeats may be made using the Online PBS Authorities system (see www.servicesaustralia.gov.au/organisations/health-professionals/services/medicare/hpos/services/request-authority-using-online-pbs-authorities-hpos).

Phone applications for increased maximum quantities/repeats may be made by calling 1800 888 333.

Written authority applications for increased maximum quantities/repeats can be uploaded online through HPOS form upload or mailed to:

Pharmaceutical Benefits Scheme

Reply Paid 9857

[Your capital city]

Authority required (STREAMLINED)**10755**

Chronic severe pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for less than 12 months

Clinical criteria:

- The condition must require daily, continuous, long term opioid treatment, **AND**
- Patient must have cancer pain; OR
- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid or other opioid analgesics; OR
- Patient must be unable to use non-opioid or other opioid analgesics due to contraindications or intolerance.

Authorities for increased maximum quantities and/or repeats under this restriction must only be considered for chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment is less than 12 months.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Authority required (STREAMLINED)**10748**

Chronic severe pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for more than 12 months

Clinical criteria:

- The condition must require daily, continuous, long term opioid treatment, **AND**
- Patient must have cancer pain; OR
- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid or other opioid analgesics; OR
- Patient must be unable to use non-opioid or other opioid analgesics due to contraindications or intolerance.

Authorities for increased maximum quantities and/or repeats must only be considered for chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment:

(i) exceeds 12 months and the palliative care patient is unable to have annual pain management review due to their clinical condition; or

(ii) exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or

(iii) has exceeded 12 months prior to 1 June 2020 and the patient's clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Authority required (STREAMLINED)**10752**

Chronic severe pain

Treatment Phase: Continuing PBS treatment after 1 June 2020

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this form of this drug for this condition after 1 June 2020.

Authorities for increased maximum quantities and/or repeats must only be considered for chronic severe disabling pain where the patient has received initial authority approval and the total duration of non-PBS and PBS opioid analgesic treatment:

(i) is less than 12 months; or

(ii) exceeds 12 months and the palliative care patient is unable to have annual pain management review due to their clinical condition; or

(iii) exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or

(iv) has exceeded 12 months prior to 1 June 2020 and the patient's pain management and clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

oxycodone hydrochloride 2.5 mg + naloxone hydrochloride 1.25 mg modified release tablet, 28

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
10776E	1	..	..	26.68	27.97	Targin 2.5/1.25 mg [MF]

oxycodone hydrochloride 20 mg + naloxone hydrochloride 10 mg modified release tablet, 28

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8935G	1	..	..	50.04	41.30	Targin 20/10mg [MF]

oxycodone hydrochloride 10 mg + naloxone hydrochloride 5 mg modified release tablet, 28

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8934F	1	..	..	35.55	36.84	Targin 10/5mg [MF]

oxycodone hydrochloride 80 mg + naloxone hydrochloride 40 mg modified release tablet, 28

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
11111T	1	..	..	97.08	41.30	Targin 80/40 [MF]

oxycodone hydrochloride 60 mg + naloxone hydrochloride 30 mg modified release tablet, 28

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
11102H	1	..	..	82.44	41.30	Targin 60/30 [MF]

oxycodone hydrochloride 40 mg + naloxone hydrochloride 20 mg modified release tablet, 28

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8936H	1	..	..	75.14	41.30	Targin 40/20mg [MF]

oxycodone hydrochloride 30 mg + naloxone hydrochloride 15 mg modified release tablet, 28

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
10758F	1	..	..	56.19	41.30	Targin 30/15 mg [MF]

oxycodone hydrochloride 15 mg + naloxone hydrochloride 7.5 mg modified release tablet, 28

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
10757E	1	..	..	39.90	41.19	Targin 15/7.5mg [MF]

oxycodone hydrochloride 5 mg + naloxone hydrochloride 2.5 mg modified release tablet, 28

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8000C	1	..	..	34.42	35.71	Targin 5/2.5mg [MF]

Phenylpiperidine derivatives**■ FENTANYL**

Caution The risk of drug dependence is high.

Note This treatment is not suitable for 'as-required' pain relief.

Note Consider consultation with a multidisciplinary pain service prior to, or after commencement of this medication.

Note Fentanyl transdermal patches are not recommended in opioid naive patients with non-cancer pain because of a high incidence of adverse events in these patients. Patients with cancer pain may be initiated on the lowest strength patch (12 micrograms per hour).

Note Pharmaceutical benefits that have the form fentanyl 12 microgram/hour patch are equivalent for the purposes of substitution.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note Real time online applications for increased maximum quantities/repeats may be made using the Online PBS Authorities system (see www.servicesaustralia.gov.au/organisations/health-professionals/services/medicare/hpos/services/request-authority-using-online-pbs-authorities-hpos).

Phone applications for increased maximum quantities/repeats may be made by calling 1800 888 333.

Written authority applications for increased maximum quantities/repeats can be uploaded online through HPOS form upload or mailed to:

Pharmaceutical Benefits Scheme

Reply Paid 9857

[Your capital city]

Authority required (STREAMLINED)**10745**

Chronic severe disabling pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for less than 12 months

Clinical criteria:

- The condition must require daily, continuous, long term opioid treatment, **AND**
- Patient must not be opioid naive, **AND**
- Patient must have cancer pain; OR

- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid and other opioid analgesics; OR
 - Patient must be unable to use non-opioid and other opioid analgesics due to contraindications or intolerance. Authorities for increased maximum quantities and/or repeats under this restriction must only be considered for chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment is less than 12 months. Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.
- Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Authority required (STREAMLINED)**10747**

Chronic severe disabling pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for more than 12 months

Clinical criteria:

- The condition must require daily, continuous, long term opioid treatment, **AND**
- Patient must not be opioid naive, **AND**
- Patient must have cancer pain; OR
- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid and other opioid analgesics; OR
- Patient must be unable to use non-opioid and other opioid analgesics due to contraindications or intolerance.

Authorities for increased maximum quantities and/or repeats must only be considered for chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment:

- (i) exceeds 12 months and the palliative care patient is unable to have annual pain management review due to their clinical condition; or
- (ii) exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or
- (iii) has exceeded 12 months prior to 1 June 2020 and the patient's clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Authority required (STREAMLINED)**10751**

Chronic severe disabling pain

Treatment Phase: Continuing PBS treatment after 1 June 2020

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this form of this drug for this condition after 1 June 2020.

Authorities for increased maximum quantities and/or repeats must only be considered for chronic severe disabling pain where the patient has received initial authority approval and the total duration of non-PBS and PBS opioid analgesic treatment:

- (i) is less than 12 months; or
- (ii) exceeds 12 months and the palliative care patient is unable to have annual pain management review due to their clinical condition; or
- (iii) exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or
- (iv) has exceeded 12 months prior to 1 June 2020 and the patient's pain management and clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

fentanyl 12 microgram/hour patch, 5

5265D

NP

Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
1	..	..	25.28	26.57	^a Denpax [AF]

fentanyl 12 microgram/hour patch, 5

5437E

NP

Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
1	..	..	25.28	26.57	^a Dutran 12 [EA]	^a Fenpatch 12 [ZP]

fantanyl 12 microgram/hour patch, 5

8878G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	25.28	26.57	^a APO-Fentanyl [TX] ^a Fentanyl Sandoz [SZ]	^a Durogesic 12 [JC]

■ FENTANYL

Caution The risk of drug dependence is high.

Note This treatment is not suitable for 'as-required' pain relief.

Note Consider consultation with a multidisciplinary pain service prior to, or after commencement of this medication.

Note Fentanyl transdermal patches are not recommended in opioid naive patients with non-cancer pain because of a high incidence of adverse events in these patients. Patients with cancer pain may be initiated on the lowest strength patch (12 micrograms per hour).

Note Pharmaceutical benefits that have the form fentanyl 25 microgram/hour patch are equivalent for the purposes of substitution.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note Real time online applications for increased maximum quantities/repeats may be made using the Online PBS Authorities system (see www.servicesaustralia.gov.au/organisations/health-professionals/services/medicare/hpos/services/request-authority-using-online-pbs-authorities-hpos).

Phone applications for increased maximum quantities/repeats may be made by calling 1800 888 333.

Written authority applications for increased maximum quantities/repeats can be uploaded online through HPOS form upload or mailed to:

Pharmaceutical Benefits Scheme

Reply Paid 9857

[Your capital city]

Authority required (STREAMLINED)**10745**

Chronic severe disabling pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for less than 12 months

Clinical criteria:

- The condition must require daily, continuous, long term opioid treatment, **AND**
- Patient must not be opioid naive, **AND**
- Patient must have cancer pain; OR
- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid and other opioid analgesics; OR
- Patient must be unable to use non-opioid and other opioid analgesics due to contraindications or intolerance.

Authorities for increased maximum quantities and/or repeats under this restriction must only be considered for chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment is less than 12 months.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Authority required (STREAMLINED)**10747**

Chronic severe disabling pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for more than 12 months

Clinical criteria:

- The condition must require daily, continuous, long term opioid treatment, **AND**
- Patient must not be opioid naive, **AND**
- Patient must have cancer pain; OR
- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid and other opioid analgesics; OR
- Patient must be unable to use non-opioid and other opioid analgesics due to contraindications or intolerance.

Authorities for increased maximum quantities and/or repeats must only be considered for chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment:

(i) exceeds 12 months and the palliative care patient is unable to have annual pain management review due to their clinical condition; or

(ii) exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or

(iii) has exceeded 12 months prior to 1 June 2020 and the patient's clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Authority required (STREAMLINED)

10751

Chronic severe disabling pain

Treatment Phase: Continuing PBS treatment after 1 June 2020

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this form of this drug for this condition after 1 June 2020.

Authorities for increased maximum quantities and/or repeats must only be considered for chronic severe disabling pain where the patient has received initial authority approval and the total duration of non-PBS and PBS opioid analgesic treatment:

- (i) is less than 12 months; or
- (ii) exceeds 12 months and the palliative care patient is unable to have annual pain management review due to their clinical condition; or
- (iii) exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or
- (iv) has exceeded 12 months prior to 1 June 2020 and the patient's pain management and clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

fentanyl 25 microgram/hour patch, 5

5277R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	27.58	28.87	^a Denpax [AF]

fentanyl 25 microgram/hour patch, 5

5438F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	27.58	28.87	^a Dutran 25 [EA]	^a Fenpatch 25 [ZP]

fentanyl 25 microgram/hour patch, 5

8891Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	27.58	28.87	^a APO-Fentanyl [TX] ^a Fentanyl Sandoz [SZ]	^a Durogesic 25 [JC]

▪ **FENTANYL**

Caution The risk of drug dependence is high.

Note This treatment is not suitable for 'as-required' pain relief.

Note Consider consultation with a multidisciplinary pain service prior to, or after commencement of this medication.

Note Fentanyl transdermal patches are not recommended in opioid naive patients with non-cancer pain because of a high incidence of adverse events in these patients. Patients with cancer pain may be initiated on the lowest strength patch (12 micrograms per hour).

Note Pharmaceutical benefits that have the form fentanyl 50 microgram/hour patch are equivalent for the purposes of substitution.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note Real time online applications for increased maximum quantities/repeats may be made using the Online PBS Authorities system (see www.servicesaustralia.gov.au/organisations/health-professionals/services/medicare/hpos/services/request-authority-using-online-pbs-authorities-hpos).

Phone applications for increased maximum quantities/repeats may be made by calling 1800 888 333.

Written authority applications for increased maximum quantities/repeats can be uploaded online through HPOS form upload or mailed to:

Pharmaceutical Benefits Scheme
Reply Paid 9857
[Your capital city]

Authority required (STREAMLINED)

10745

Chronic severe disabling pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for less than 12 months

Clinical criteria:

- The condition must require daily, continuous, long term opioid treatment, **AND**
- Patient must not be opioid naive, **AND**

- Patient must have cancer pain; OR
- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid and other opioid analgesics; OR
- Patient must be unable to use non-opioid and other opioid analgesics due to contraindications or intolerance.

Authorities for increased maximum quantities and/or repeats under this restriction must only be considered for chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment is less than 12 months.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Authority required (STREAMLINED)

10747

Chronic severe disabling pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for more than 12 months

Clinical criteria:

- The condition must require daily, continuous, long term opioid treatment, **AND**
- Patient must not be opioid naive, **AND**
- Patient must have cancer pain; OR
- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid and other opioid analgesics; OR
- Patient must be unable to use non-opioid and other opioid analgesics due to contraindications or intolerance.

Authorities for increased maximum quantities and/or repeats must only be considered for chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment:

(i) exceeds 12 months and the palliative care patient is unable to have annual pain management review due to their clinical condition; or

(ii) exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or

(iii) has exceeded 12 months prior to 1 June 2020 and the patient's clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Authority required (STREAMLINED)

10751

Chronic severe disabling pain

Treatment Phase: Continuing PBS treatment after 1 June 2020

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this form of this drug for this condition after 1 June 2020.

Authorities for increased maximum quantities and/or repeats must only be considered for chronic severe disabling pain where the patient has received initial authority approval and the total duration of non-PBS and PBS opioid analgesic treatment:

(i) is less than 12 months; or

(ii) exceeds 12 months and the palliative care patient is unable to have annual pain management review due to their clinical condition; or

(iii) exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or

(iv) has exceeded 12 months prior to 1 June 2020 and the patient's pain management and clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

fentanyl 50 microgram/hour patch, 5

5278T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	36.60	37.89	^a Denpax [AF]

fentanyl 50 microgram/hour patch, 5

5439G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	36.60	37.89	^a Dutran 50 [EA]	^a Fenpatch 50 [ZP]

fentanyl 50 microgram/hour patch, 5

8892B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	36.60	37.89	^a APO-Fentanyl [TX] ^a Fentanyl Sandoz [SZ]	^a Durogesic 50 [JC]

■ FENTANYL

Caution The risk of drug dependence is high.

Note This treatment is not suitable for 'as-required' pain relief.

Note Consider consultation with a multidisciplinary pain service prior to, or after commencement of this medication.

Note Fentanyl transdermal patches are not recommended in opioid naive patients with non-cancer pain because of a high incidence of adverse events in these patients. Patients with cancer pain may be initiated on the lowest strength patch (12 micrograms per hour).

Note Pharmaceutical benefits that have the form fentanyl 75 microgram/hour patch are equivalent for the purposes of substitution.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note Real time online applications for increased maximum quantities/repeats may be made using the Online PBS Authorities system (see www.servicesaustralia.gov.au/organisations/health-professionals/services/medicare/hpos/services/request-authority-using-online-pbs-authorities-hpos).

Phone applications for increased maximum quantities/repeats may be made by calling 1800 888 333.

Written authority applications for increased maximum quantities/repeats can be uploaded online through HPOS form upload or mailed to:

Pharmaceutical Benefits Scheme

Reply Paid 9857

[Your capital city]

Authority required (STREAMLINED)**10745**

Chronic severe disabling pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for less than 12 months

Clinical criteria:

- The condition must require daily, continuous, long term opioid treatment, **AND**
- Patient must not be opioid naive, **AND**
- Patient must have cancer pain; OR
- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid and other opioid analgesics; OR
- Patient must be unable to use non-opioid and other opioid analgesics due to contraindications or intolerance.

Authorities for increased maximum quantities and/or repeats under this restriction must only be considered for chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment is less than 12 months.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Authority required (STREAMLINED)**10747**

Chronic severe disabling pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for more than 12 months

Clinical criteria:

- The condition must require daily, continuous, long term opioid treatment, **AND**
- Patient must not be opioid naive, **AND**
- Patient must have cancer pain; OR
- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid and other opioid analgesics; OR
- Patient must be unable to use non-opioid and other opioid analgesics due to contraindications or intolerance.

Authorities for increased maximum quantities and/or repeats must only be considered for chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment:

(i) exceeds 12 months and the palliative care patient is unable to have annual pain management review due to their clinical condition; or

(ii) exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or

(iii) has exceeded 12 months prior to 1 June 2020 and the patient's clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Authority required (STREAMLINED)

10751

Chronic severe disabling pain

Treatment Phase: Continuing PBS treatment after 1 June 2020

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this form of this drug for this condition after 1 June 2020.

Authorities for increased maximum quantities and/or repeats must only be considered for chronic severe disabling pain where the patient has received initial authority approval and the total duration of non-PBS and PBS opioid analgesic treatment:

(i) is less than 12 months; or

(ii) exceeds 12 months and the palliative care patient is unable to have annual pain management review due to their clinical condition; or

(iii) exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or

(iv) has exceeded 12 months prior to 1 June 2020 and the patient's pain management and clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

fentanyl 75 microgram/hour patch, 5

5279W

Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
1	..	..	43.93	41.30	^a Denpax [AF]

NP

fentanyl 75 microgram/hour patch, 5

5440H

Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
1	..	..	43.93	41.30	^a Dutran 75 [EA]	^a Fenpatch 75 [ZP]

NP

fentanyl 75 microgram/hour patch, 5

8893C

Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
1	..	..	43.93	41.30	^a APO-Fentanyl [TX]	^a Durogesic 75 [JC]

NP

^a Fentanyl Sandoz [SZ]

■ FENTANYL

Caution The risk of drug dependence is high.

Note This treatment is not suitable for 'as-required' pain relief.

Note Consider consultation with a multidisciplinary pain service prior to, or after commencement of this medication.

Note Fentanyl transdermal patches are not recommended in opioid naive patients with non-cancer pain because of a high incidence of adverse events in these patients. Patients with cancer pain may be initiated on the lowest strength patch (12 micrograms per hour).

Note Pharmaceutical benefits that have the form fentanyl 100 microgram/hour patch are equivalent for the purposes of substitution.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note Real time online applications for increased maximum quantities/repeats may be made using the Online PBS Authorities system (see www.servicesaustralia.gov.au/organisations/health-professionals/services/medicare/hpos/services/request-authority-using-online-pbs-authorities-hpos).

Phone applications for increased maximum quantities/repeats may be made by calling 1800 888 333.

Written authority applications for increased maximum quantities/repeats can be uploaded online through HPOS form upload or mailed to:

Pharmaceutical Benefits Scheme

Reply Paid 9857

[Your capital city]

Authority required (STREAMLINED)

10745

Chronic severe disabling pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for less than 12 months

Clinical criteria:

- The condition must require daily, continuous, long term opioid treatment, **AND**
- Patient must not be opioid naive, **AND**
- Patient must have cancer pain; OR
- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid and other opioid analgesics; OR
- Patient must be unable to use non-opioid and other opioid analgesics due to contraindications or intolerance.

Authorities for increased maximum quantities and/or repeats under this restriction must only be considered for chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment is less than 12 months.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Authority required (STREAMLINED)**10747**

Chronic severe disabling pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for more than 12 months

Clinical criteria:

- The condition must require daily, continuous, long term opioid treatment, **AND**
- Patient must not be opioid naive, **AND**
- Patient must have cancer pain; OR
- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid and other opioid analgesics; OR
- Patient must be unable to use non-opioid and other opioid analgesics due to contraindications or intolerance.

Authorities for increased maximum quantities and/or repeats must only be considered for chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment:

(i) exceeds 12 months and the palliative care patient is unable to have annual pain management review due to their clinical condition; or

(ii) exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or

(iii) has exceeded 12 months prior to 1 June 2020 and the patient's clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Authority required (STREAMLINED)**10751**

Chronic severe disabling pain

Treatment Phase: Continuing PBS treatment after 1 June 2020

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this form of this drug for this condition after 1 June 2020.

Authorities for increased maximum quantities and/or repeats must only be considered for chronic severe disabling pain where the patient has received initial authority approval and the total duration of non-PBS and PBS opioid analgesic treatment:

(i) is less than 12 months; or

(ii) exceeds 12 months and the palliative care patient is unable to have annual pain management review due to their clinical condition; or

(iii) exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or

(iv) has exceeded 12 months prior to 1 June 2020 and the patient's pain management and clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

fentanyl 100 microgram/hour patch, 5

5280X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	50.45	41.30	^a Denpax [AF]

fentanyl 100 microgram/hour patch, 5

5441J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	50.45	41.30	^a Dutran 100 [EA]	^a Fenpatch 100 [ZP]

fentanyl 100 microgram/hour patch, 5

8894D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	50.45	41.30	^a APO-Fentanyl [TX] ^a Fentanyl Sandoz [SZ]	^a Durogesic 100 [JC]

*Diphenylpropylamine derivatives***■ METHADONE**

Caution The risk of drug dependence is high.

Note This treatment is not suitable for 'as-required' pain relief.

Note This treatment is not recommended for use in ambulant patients.

Note Consider consultation with a multidisciplinary pain service prior to, or after commencement of this medication.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note Real time online applications for increased maximum quantities/repeats may be made using the Online PBS Authorities system (see www.servicesaustralia.gov.au/organisations/health-professionals/services/medicare/hpos/services/request-authority-using-online-pbs-authorities-hpos).

Phone applications for increased maximum quantities/repeats may be made by calling 1800 888 333.

Written authority applications for increased maximum quantities/repeats can be uploaded online through HPOS form upload or mailed to:

Pharmaceutical Benefits Scheme

Reply Paid 9857

[Your capital city]

Authority required (STREAMLINED)**10745**

Chronic severe disabling pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for less than 12 months

Clinical criteria:

- The condition must require daily, continuous, long term opioid treatment, **AND**
 - Patient must not be opioid naive, **AND**
 - Patient must have cancer pain; OR
 - Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid and other opioid analgesics; OR
 - Patient must be unable to use non-opioid and other opioid analgesics due to contraindications or intolerance.
- Authorities for increased maximum quantities and/or repeats under this restriction must only be considered for chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment is less than 12 months.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Authority required (STREAMLINED)**10747**

Chronic severe disabling pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for more than 12 months

Clinical criteria:

- The condition must require daily, continuous, long term opioid treatment, **AND**
- Patient must not be opioid naive, **AND**
- Patient must have cancer pain; OR
- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid and other opioid analgesics; OR
- Patient must be unable to use non-opioid and other opioid analgesics due to contraindications or intolerance.

Authorities for increased maximum quantities and/or repeats must only be considered for chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment:

(i) exceeds 12 months and the palliative care patient is unable to have annual pain management review due to their clinical condition; or

- (ii) exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or
- (iii) has exceeded 12 months prior to 1 June 2020 and the patient's clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Authority required (STREAMLINED)

10751

Chronic severe disabling pain

Treatment Phase: Continuing PBS treatment after 1 June 2020

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this form of this drug for this condition after 1 June 2020.

Authorities for increased maximum quantities and/or repeats must only be considered for chronic severe disabling pain where the patient has received initial authority approval and the total duration of non-PBS and PBS opioid analgesic treatment:

- (i) is less than 12 months; or
- (ii) exceeds 12 months and the palliative care patient is unable to have annual pain management review due to their clinical condition; or
- (iii) exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or
- (iv) has exceeded 12 months prior to 1 June 2020 and the patient's pain management and clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

methadone hydrochloride 10 mg tablet, 20

1609Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	22.16	23.45	Physeptone [AS]

methadone hydrochloride 10 mg/mL injection, 5 x 1 mL ampoules

1606M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	52.50	41.30	Physeptone [AS]

Oripavine derivatives

■ BUPRENORPHINE

Caution The risk of drug dependence is high.

Note This treatment is not suitable for 'as-required' pain relief.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note Real time online applications for increased maximum quantities/repeats may be made using the Online PBS Authorities system (see www.servicesaustralia.gov.au/organisations/health-professionals/services/medicare/hpos/services/request-authority-using-online-pbs-authorities-hpos).

Phone applications for increased maximum quantities/repeats may be made by calling 1800 888 333.

Written authority applications for increased maximum quantities/repeats can be uploaded online through HPOS form upload or mailed to:

Pharmaceutical Benefits Scheme

Reply Paid 9857

[Your capital city]

Authority required (STREAMLINED)

10755

Chronic severe pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for less than 12 months

Clinical criteria:

- The condition must require daily, continuous, long term opioid treatment, **AND**
- Patient must have cancer pain; OR

- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid or other opioid analgesics; OR
- Patient must be unable to use non-opioid or other opioid analgesics due to contraindications or intolerance.

Authorities for increased maximum quantities and/or repeats under this restriction must only be considered for chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment is less than 12 months.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Authority required (STREAMLINED)

10748

Chronic severe pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for more than 12 months

Clinical criteria:

- The condition must require daily, continuous, long term opioid treatment, **AND**
- Patient must have cancer pain; OR
- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid or other opioid analgesics; OR
- Patient must be unable to use non-opioid or other opioid analgesics due to contraindications or intolerance.

Authorities for increased maximum quantities and/or repeats must only be considered for chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment:

(i) exceeds 12 months and the palliative care patient is unable to have annual pain management review due to their clinical condition; or

(ii) exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or

(iii) has exceeded 12 months prior to 1 June 2020 and the patient's clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Authority required (STREAMLINED)

10752

Chronic severe pain

Treatment Phase: Continuing PBS treatment after 1 June 2020

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this form of this drug for this condition after 1 June 2020.

Authorities for increased maximum quantities and/or repeats must only be considered for chronic severe disabling pain where the patient has received initial authority approval and the total duration of non-PBS and PBS opioid analgesic treatment:

(i) is less than 12 months; or

(ii) exceeds 12 months and the palliative care patient is unable to have annual pain management review due to their clinical condition; or

(iii) exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or

(iv) has exceeded 12 months prior to 1 June 2020 and the patient's pain management and clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

buprenorphine 25 microgram/hour patch, 2

10756D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	56.34	41.30	^a Bupredermal [TX] ^a Norspan [MF]	^a Burprenorphine Sandoz [SZ]

buprenorphine 30 microgram/hour patch, 2

10755C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	61.99	41.30	^a Bupredermal [TX] ^a Norspan [MF]	^a Buprenorphine Sandoz [SZ]

buprenorphine 5 microgram/hour patch, 2

8865N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	29.01	30.30	^a Bupredermal [TX] ^a Norspan [MF]	^a Buprenorphine Sandoz [SZ]

buprenorphine 10 microgram/hour patch, 2

8866P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	38.78	40.07	^a Bupredermal [TX] ^a Norspan [MF]	^a Buprenorphine Sandoz [SZ]

buprenorphine 20 microgram/hour patch, 2

8867Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	50.70	41.30	^a Bupredermal [TX] ^a Norspan [MF]	^a Buprenorphine Sandoz [SZ]

buprenorphine 40 microgram/hour patch, 2

10746N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	73.28	41.30	^a Bupredermal [TX] ^a Norspan [MF]	^a Buprenorphine Sandoz [SZ]

buprenorphine 15 microgram/hour patch, 2

10770W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	44.73	41.30	^a Bupredermal [TX] ^a Norspan [MF]	^a Buprenorphine Sandoz [SZ]

Opioids in combination with non-opioid analgesics**■ PARACETAMOL + CODEINE****Caution** The risk of drug dependence is high.**Note** Prescribing of drugs of addiction by dentists is not permitted in some States/Territories.**Restricted benefit**

Severe pain

Clinical criteria:

- The treatment must be for short term therapy of acute severe pain, **AND**
- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid analgesics; OR
- Patient must be unable to use non-opioid analgesics due to contraindications or intolerance.

paracetamol 500 mg + codeine phosphate hemihydrate 30 mg tablet, 20

12066C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
DP	0.5	..	..	*14.95	16.24	^a APO- Paracetamol/Codeine 500/30 [TX] ^a Codalgin Forte [AF] ^a Comfarol Forte [SZ] ^a Prodeine Forte [AV]	^a APX-Paracetamol/Codeine [TY] ^a Codapane Forte 500/30 [AL] ^a Paracetamol/Codeine GH 500/30 [GQ]
			^B 1.21	*16.16	16.24	^a Panadeine Forte [SW]	

■ PARACETAMOL + CODEINE**Caution** The risk of drug dependence is high.**Note** Prescribing of drugs of addiction by dentists is not permitted in some States/Territories.**Restricted benefit**

Severe pain

Clinical criteria:

- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid analgesics; OR
- Patient must be unable to use non-opioid analgesics due to contraindications or intolerance.

paracetamol 500 mg + codeine phosphate hemihydrate 30 mg tablet, 20

3316M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
DP	1	..	..	13.69	14.98	^a APO- Paracetamol/Codeine 500/30 [TX] ^a Codalgin Forte [AF] ^a Comfarol Forte [SZ] ^a Prodeine Forte [AV]	^a APX-Paracetamol/Codeine [TY] ^a Codapane Forte 500/30 [AL] ^a Paracetamol/Codeine GH 500/30 [GQ]
			^B 1.95	15.64	14.98	^a Panadeine Forte [SW]	

■ PARACETAMOL + CODEINE**Caution** The risk of drug dependence is high.**Note** No increase in the maximum quantity or number of units may be authorised.**Note** No increase in the maximum number of repeats may be authorised.

Restricted benefit

Severe pain

Clinical criteria:

- The treatment must be for short term therapy of acute severe pain, **AND**
- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid analgesics; OR
- Patient must be unable to use non-opioid analgesics due to contraindications or intolerance.

paracetamol 500 mg + codeine phosphate hemihydrate 30 mg tablet, 20

12022R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	0.5	..	..	*14.95	16.24	^a APO- Paracetamol/Codeine 500/30 [TX]	^a APX-Paracetamol/Codeine [TY]
						^a Codalgin Forte [AF]	^a Codapane Forte 500/30 [AL]
						^a Comfarol Forte [SZ]	^a Paracetamol/Codeine GH 500/30 [GQ]
						^a Prodeine Forte [AV]	
						^a Panadeine Forte [SW]	
			^B 1.21	*16.16	16.24		

■ PARACETAMOL + CODEINE**Caution** The risk of drug dependence is high.

Note Real time online applications for increased maximum quantities/repeats may be made using the Online PBS Authorities system (see www.servicesaustralia.gov.au/organisations/health-professionals/services/medicare/hpos/services/request-authority-using-online-pbs-authorities-hpos).

Phone applications for increased maximum quantities/repeats may be made by calling 1800 888 333.

Written authority applications for increased maximum quantities/repeats can be uploaded online through HPOS form upload or mailed to:

Pharmaceutical Benefits Scheme

Reply Paid 9857

[Your capital city]

Restricted benefit

Severe pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for less than 12 months

Clinical criteria:

- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid analgesics; OR
- Patient must be unable to use non-opioid analgesics due to contraindications or intolerance.

Authorities for increased maximum quantities and/or repeats under this restriction must only be considered for severe disabling pain associated with malignant neoplasia or chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment is less than 12 months.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Restricted benefit

Severe pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for more than 12 months

Clinical criteria:

- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid analgesics; OR
- Patient must be unable to use non-opioid analgesics due to contraindications or intolerance.

Authorities for increased maximum quantities and/or repeats must only be considered for:

- severe disabling pain associated with proven malignant neoplasia; or
- palliative care patients with chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment exceeds 12 months and the patient is unable to have annual pain management review due to their clinical condition; or
- chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or
- chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment has exceeded 12 months prior to 1 June 2020 and the patient's clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Restricted benefit

Severe pain

Treatment Phase: Continuing PBS treatment after 1 June 2020

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this form of this drug for this condition after 1 June 2020.

Authorities for increased maximum quantities and/or repeats must only be considered where the patient has received initial authority approval for:

- (i) severe disabling pain associated with malignant neoplasia; or
- (ii) chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment is less than 12 months; or
- (iii) palliative care patients with chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment exceeds 12 months and the patient is unable to have annual pain management review due to their clinical condition; or
- (iv) chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or
- (v) chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment has exceeded 12 months prior to 1 June 2020 and the patient's clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

paracetamol 500 mg + codeine phosphate hemihydrate 30 mg tablet, 20

1215Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	13.69	14.98	^a APO- Paracetamol/Codeine 500/30 [TX]	^a APX-Paracetamol/Codeine [TY]
						^a Codalgin Forte [AF]	^a Codapane Forte 500/30 [AL]
						^a Comfarol Forte [SZ]	^a Paracetamol/Codeine GH 500/30 [GQ]
						^a Prodeine Forte [AV]	
			^B 1.95	15.64	14.98	^a Panadeine Forte [SW]	

Other opioids

■ TAPENTADOL

Caution The risk of drug dependence is high.

Note This treatment is not suitable for 'as-required' pain relief.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note Real time online applications for increased maximum quantities/repeats may be made using the Online PBS Authorities system (see www.servicesaustralia.gov.au/organisations/health-professionals/services/medicare/hpos/services/request-authority-using-online-pbs-authorities-hpos).

Phone applications for increased maximum quantities/repeats may be made by calling 1800 888 333.

Written authority applications for increased maximum quantities/repeats can be uploaded online through HPOS form upload or mailed to:

Pharmaceutical Benefits Scheme

Reply Paid 9857

[Your capital city]

Authority required (STREAMLINED)

10755

Chronic severe pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for less than 12 months

Clinical criteria:

- The condition must require daily, continuous, long term opioid treatment, **AND**
- Patient must have cancer pain; OR
- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid or other opioid analgesics; OR
- Patient must be unable to use non-opioid or other opioid analgesics due to contraindications or intolerance.

Authorities for increased maximum quantities and/or repeats under this restriction must only be considered for chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment is less than 12 months. Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Authority required (STREAMLINED)

10748

Chronic severe pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for more than 12 months

Clinical criteria:

- The condition must require daily, continuous, long term opioid treatment, **AND**
- Patient must have cancer pain; OR
- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid or other opioid analgesics; OR
- Patient must be unable to use non-opioid or other opioid analgesics due to contraindications or intolerance.

Authorities for increased maximum quantities and/or repeats must only be considered for chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment:

(i) exceeds 12 months and the palliative care patient is unable to have annual pain management review due to their clinical condition; or

(ii) exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or

(iii) has exceeded 12 months prior to 1 June 2020 and the patient's clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Authority required (STREAMLINED)

10752

Chronic severe pain

Treatment Phase: Continuing PBS treatment after 1 June 2020

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this form of this drug for this condition after 1 June 2020.

Authorities for increased maximum quantities and/or repeats must only be considered for chronic severe disabling pain where the patient has received initial authority approval and the total duration of non-PBS and PBS opioid analgesic treatment:

(i) is less than 12 months; or

(ii) exceeds 12 months and the palliative care patient is unable to have annual pain management review due to their clinical condition; or

(iii) exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or

(iv) has exceeded 12 months prior to 1 June 2020 and the patient's pain management and clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

tapentadol 100 mg modified release tablet, 28

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
10094G	1	..	..	35.14	36.43	Palexia SR [CS]

NP

tapentadol 200 mg modified release tablet, 28

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
10091D	1	..	..	48.67	41.30	Palexia SR [CS]

NP

tapentadol 150 mg modified release tablet, 28

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
10100N	1	..	..	42.32	41.30	Palexia SR [CS]

NP

tapentadol 250 mg modified release tablet, 28

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
10092E NP	1	..	..	54.04	41.30	Palexia SR [CS]

tapentadol 50 mg modified release tablet, 28

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
10096J NP	1	..	..	27.67	28.96	Palexia SR [CS]

■ TRAMADOL**Caution** The risk of drug dependence is high.**Note** Prescribing of drugs of addiction by dentists is not permitted in some States/Territories.**Restricted benefit**

Severe pain

Clinical criteria:

- The treatment must be for short term therapy of acute severe pain, **AND**
- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid analgesics; OR
- Patient must be unable to use non-opioid analgesics due to contraindications or intolerance.

tramadol hydrochloride 50 mg capsule, 20

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
12024W DP	0.5	..	..	*15.32	16.61	^a APO-Tramadol [TX] ^a Tramadol AN [EA] ^a Tramedo [AF]	^a Tramadol AMNEAL [EF] ^a Tramadol Sandoz [SZ] ^a Zydol [RW]
			^B 1.40	*16.72	16.61	^a Tramal [CS]	

■ TRAMADOL**Caution** The risk of drug dependence is high.**Note** Prescribing of drugs of addiction by dentists is not permitted in some States/Territories.**Restricted benefit**

Severe pain

Clinical criteria:

- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid analgesics; OR
- Patient must be unable to use non-opioid analgesics due to contraindications or intolerance.

tramadol hydrochloride 50 mg capsule, 20

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
5232J DP	1	..	..	14.29	15.58	^a APO-Tramadol [TX] ^a Tramadol AN [EA] ^a Tramedo [AF]	^a Tramadol AMNEAL [EF] ^a Tramadol Sandoz [SZ] ^a Zydol [RW]
			^B 2.25	16.54	15.58	^a Tramal [CS]	

tramadol hydrochloride 100 mg/mL oral liquid, 10 mL

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
5150C DP	‡1	..	..	18.36	19.65	Tramal [CS]

tramadol hydrochloride 100 mg/2 mL injection, 5 x 2 mL ampoules

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
5231H DP	1	..	..	15.08	16.37	^a Tramadol ACT [JO] ^a Tramadol Sandoz [SZ]	^a Tramadol AN [JU] ^a Tramal 100 [CS]

■ TRAMADOL**Caution** The risk of drug dependence is high.**Note** No increase in the maximum quantity or number of units may be authorised.**Note** No increase in the maximum number of repeats may be authorised.**Restricted benefit**

Severe pain

Clinical criteria:

- The treatment must be for short term therapy of acute severe pain, **AND**
- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid analgesics; OR
- Patient must be unable to use non-opioid analgesics due to contraindications or intolerance.

tramadol hydrochloride 50 mg capsule, 20

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
12008B NP	0.5	..	..	*15.32	16.61	^a APO-Tramadol [TX] ^a Tramadol AN [EA] ^a Tramedo [AF]	^a Tramadol AMNEAL [EF] ^a Tramadol Sandoz [SZ] ^a Zydol [RW]

^B1.40 *16.72 16.61 ^a Tramal [CS]

■ TRAMADOL

Caution The risk of drug dependence is high.

Note Real time online applications for increased maximum quantities/repeats may be made using the Online PBS Authorities system (see www.servicesaustralia.gov.au/organisations/health-professionals/services/medicare/hpos/services/request-authority-using-online-pbs-authorities-hpos).
Phone applications for increased maximum quantities/repeats may be made by calling 1800 888 333.
Written authority applications for increased maximum quantities/repeats can be uploaded online through HPOS form upload or mailed to:
Pharmaceutical Benefits Scheme
Reply Paid 9857
[Your capital city]

Restricted benefit

Severe pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for less than 12 months

Clinical criteria:

- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid analgesics; OR
- Patient must be unable to use non-opioid analgesics due to contraindications or intolerance.

Authorities for increased maximum quantities and/or repeats under this restriction must only be considered for severe disabling pain associated with malignant neoplasia or chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment is less than 12 months.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Restricted benefit

Severe pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for more than 12 months

Clinical criteria:

- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid analgesics; OR
- Patient must be unable to use non-opioid analgesics due to contraindications or intolerance.

Authorities for increased maximum quantities and/or repeats must only be considered for:

- severe disabling pain associated with proven malignant neoplasia; or
- palliative care patients with chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment exceeds 12 months and the patient is unable to have annual pain management review due to their clinical condition; or
- chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or
- chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment has exceeded 12 months prior to 1 June 2020 and the patient's clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Restricted benefit

Severe pain

Treatment Phase: Continuing PBS treatment after 1 June 2020

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this form of this drug for this condition after 1 June 2020.

Authorities for increased maximum quantities and/or repeats must only be considered where the patient has received initial authority approval for:

- severe disabling pain associated with malignant neoplasia; or
- chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment is less than 12 months; or
- palliative care patients with chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment exceeds 12 months and the patient is unable to have annual pain management review due to their clinical condition; or

(iv) chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or

(v) chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment has exceeded 12 months prior to 1 June 2020 and the patient's clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

tramadol hydrochloride 50 mg capsule, 20

8455B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	14.29	15.58	^a APO-Tramadol [TX] ^a Tramadol AN [EA] ^a Tramedo [AF]	^a Tramadol AMNEAL [EF] ^a Tramadol Sandoz [SZ] ^a Zydol [RW]
			^B 2.25	16.54	15.58	^a Tramal [CS]	

tramadol hydrochloride 100 mg/mL oral liquid, 10 mL

8843K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	‡1	..	..	18.36	19.65	Tramal [CS]

tramadol hydrochloride 100 mg/2 mL injection, 5 x 2 mL ampoules

8582Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	15.08	16.37	^a Tramadol ACT [JO] ^a Tramadol Sandoz [SZ]	^a Tramadol AN [JU] ^a Tramal 100 [CS]

■ TRAMADOL

Caution The risk of drug dependence is high.

Note This treatment is not suitable for 'as-required' pain relief.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note Real time online applications for increased maximum quantities/repeats may be made using the Online PBS Authorities system (see www.servicesaustralia.gov.au/organisations/health-professionals/services/medicare/hpos/services/request-authority-using-online-pbs-authorities-hpos).

Phone applications for increased maximum quantities/repeats may be made by calling 1800 888 333.

Written authority applications for increased maximum quantities/repeats can be uploaded online through HPOS form upload or mailed to:

Pharmaceutical Benefits Scheme

Reply Paid 9857

[Your capital city]

Authority required (STREAMLINED)

10755

Chronic severe pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for less than 12 months

Clinical criteria:

- The condition must require daily, continuous, long term opioid treatment, **AND**
- Patient must have cancer pain; OR
- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid or other opioid analgesics; OR
- Patient must be unable to use non-opioid or other opioid analgesics due to contraindications or intolerance.

Authorities for increased maximum quantities and/or repeats under this restriction must only be considered for chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment is less than 12 months.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Authority required (STREAMLINED)

10748

Chronic severe pain

Treatment Phase: Initial PBS treatment after 1 June 2020 where patient has been treated with opioids for more than 12 months

Clinical criteria:

- The condition must require daily, continuous, long term opioid treatment, **AND**
- Patient must have cancer pain; OR
- Patient must have had or would have inadequate pain management with maximum tolerated doses of non-opioid or other opioid analgesics; OR
- Patient must be unable to use non-opioid or other opioid analgesics due to contraindications or intolerance.

Authorities for increased maximum quantities and/or repeats must only be considered for chronic severe disabling pain where the total duration of non-PBS and PBS opioid analgesic treatment:

(i) exceeds 12 months and the palliative care patient is unable to have annual pain management review due to their clinical condition; or

(ii) exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or

(iii) has exceeded 12 months prior to 1 June 2020 and the patient's clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

Authority required (STREAMLINED)

10752

Chronic severe pain

Treatment Phase: Continuing PBS treatment after 1 June 2020

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this form of this drug for this condition after 1 June 2020.

Authorities for increased maximum quantities and/or repeats must only be considered for chronic severe disabling pain where the patient has received initial authority approval and the total duration of non-PBS and PBS opioid analgesic treatment:

(i) is less than 12 months; or

(ii) exceeds 12 months and the palliative care patient is unable to have annual pain management review due to their clinical condition; or

(iii) exceeds 12 months and the patient's clinical need for continuing opioid treatment has been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months; or

(iv) has exceeded 12 months prior to 1 June 2020 and the patient's pain management and clinical need for continuing opioid treatment has not been confirmed through consultation with the patient by another medical practitioner or a palliative care nurse practitioner in the past 12 months, but is planned in the next 3 months.

Palliative care nurses may conduct annual review under this item for the treatment of palliative care patients only.

Authority requests extending treatment duration up to 1 month may be requested through the Online PBS Authorities system or by calling Services Australia.

Authority requests extending treatment duration beyond 1 month may be requested through the Online PBS Authorities system or in writing and must not provide a treatment duration exceeding 3 months (quantity sufficient for up to 1 month treatment and sufficient repeats).

tramadol hydrochloride 100 mg modified release tablet, 20

8523N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	14.67	15.96	^a APO-Tramadol SR [TX] ^a Tramadol Sandoz SR [SZ] ^a Tramedo SR [AL] ^a Tramal SR 100 [CS]	^a Tramadol AN SR [EA] ^a Tramadol SR generichealth [GQ] ^a Zydol SR 100 [RW]
			^b 4.25	18.92	15.96		

tramadol hydrochloride 50 mg modified release tablet, 20

2527B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	15.36	16.65	Tramal SR 50 [CS]

tramadol hydrochloride 200 mg modified release tablet, 20

8525Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	16.09	17.38	^a APO-Tramadol SR [TX] ^a Tramadol Sandoz SR [SZ] ^a Tramedo SR [AL] ^a Tramal SR 200 [CS]	^a Tramadol AN SR [EA] ^a Tramadol SR generichealth [GQ] ^a Zydol SR 200 [RW]
			^b 5.95	22.04	17.38		

tramadol hydrochloride 150 mg modified release tablet, 20

8524P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	15.43	16.72	^a APO-Tramadol SR [TX] ^a Tramadol Sandoz SR [SZ] ^a Tramedo SR [AL]	^a Tramadol AN SR [EA] ^a Tramadol SR generichealth [GQ] ^a Zydol SR 150 [RW]

^B5.19 20.62 16.72 ^a Tramal SR 150 [CS]

OTHER ANALGESICS AND ANTIPYRETICS

Anilides

■ PARACETAMOL

Restricted benefit

For treatment of a patient identifying as Aboriginal or Torres Strait Islander

paracetamol 500 mg tablet, 100

1746X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	1	..	13.69	14.98	^a APO-Paracetamol [TX] ^a Mendelev Paracetamol [HX] ^a Paracetamol (Sandoz) [SZ] ^a Parapane [AF]	^a Febridol [EA] ^a Panamax [SW] ^a Paralgin [OW] ^a PHARMACY CARE PARACETAMOL [SI]

paracetamol 240 mg/5 mL oral liquid, 200 mL

1770E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	2	..	15.87	17.16	Panamax 240 Elixir [SW]

paracetamol 120 mg/5 mL oral liquid, 100 mL

1747Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	2	..	14.82	16.11	Panamax [SW]

■ PARACETAMOL

Restricted benefit

For treatment of a patient identifying as Aboriginal or Torres Strait Islander

paracetamol 500 mg tablet, 100

5196L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
DP	1	..	..	13.69	14.98	^a APO-Paracetamol [TX] ^a Mendelev Paracetamol [HX] ^a Paracetamol (Sandoz) [SZ] ^a Parapane [AF]	^a Febridol [EA] ^a Panamax [SW] ^a Paralgin [OW] ^a PHARMACY CARE PARACETAMOL [SI]

paracetamol 240 mg/5 mL oral liquid, 200 mL

3349G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	±1	..	..	15.87	17.16	Panamax 240 Elixir [SW]

paracetamol 120 mg/5 mL oral liquid, 100 mL

3348F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	±1	..	..	14.82	16.11	Panamax [SW]

■ PARACETAMOL

Restricted benefit

Chronic arthropathies

Population criteria:

- Patient must identify as Aboriginal or Torres Strait Islander.

paracetamol 500 mg tablet, 100

5224Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
DP	3	..	..	*17.04	18.33	^a APO-Paracetamol [TX] ^a Mendelev Paracetamol [HX] ^a Paracetamol (Sandoz) [SZ] ^a Parapane [AF]	^a Febridol [EA] ^a Panamax [SW] ^a Paralgin [OW] ^a PHARMACY CARE PARACETAMOL [SI]

■ PARACETAMOL

Restricted benefit

Chronic arthropathies

Population criteria:

- Patient must identify as Aboriginal or Torres Strait Islander.

paracetamol 500 mg tablet, 100

8784H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	3	4	..	*17.04	18.33	^a APO-Paracetamol [TX] ^a Mendelev Paracetamol [HX] ^a Paracetamol (Sandoz) [SZ]	^a Febridol [EA] ^a Panamax [SW] ^a Paralgin [OW]

^a Parapane [AF]^a PHARMACY CARE
PARACETAMOL [SI]

■ PARACETAMOL

Note Pharmaceutical benefits that have the form paracetamol 665 mg tablet: modified release, 96 and pharmaceutical benefits that have the form paracetamol 665 mg tablet: modified release, 192 are equivalent for the purposes of substitution.

Restricted benefit

Persistent pain

Clinical criteria:

- The condition must be associated with osteoarthritis.

Population criteria:

- Patient must identify as Aboriginal or Torres Strait Islander.

paracetamol 665 mg modified release tablet, 192

10797G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	19.48	20.77	^a Osteomol 665 Paracetamol [CR]

paracetamol 665 mg modified release tablet, 96

8814X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	2	5	..	*19.48	20.77	^a APOHEALTH Osteo Relief Paracetamol 665 mg [TX]	^a Osteomol 665 Paracetamol [CR]

Other analgesics and antipyretics

■ PREGABALIN

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**4172**

Neuropathic pain

Clinical criteria:

- The condition must be refractory to treatment with other drugs.

pregabalin 25 mg capsule, 56

2348N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	16.69	17.98	^a BTC PREGABALIN [JB] ^a LYPRALIN [RW] ^a Lyzalon [AF] ^a Pregabalin AMNEAL [EA] ^a PREGABALIN-DRLA [RZ] ^a Pregabalin Sandoz [SZ]	^a Cipla Pregabalin [LR] ^a Lyrica [UJ] ^a Neuroccord [CR] ^a Pregabalin APOTEX [GX] ^a Pregabalin GH [GQ] ^a Pregabalin-Teva [TB]

pregabalin 150 mg capsule, 56

2355Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	27.58	28.87	^a BTC PREGABALIN [JB] ^a LYPRALIN [RW] ^a Lyzalon [AF] ^a Pregabalin AMNEAL [EA] ^a PREGABALIN-DRLA [RZ] ^a Pregabalin Sandoz [SZ]	^a Cipla Pregabalin [LR] ^a Lyrica [UJ] ^a Neuroccord [CR] ^a Pregabalin APOTEX [GX] ^a Pregabalin GH [GQ] ^a Pregabalin-Teva [TB]

pregabalin 75 mg capsule, 56

2335X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	22.16	23.45	^a BTC PREGABALIN [JB] ^a LYPRALIN [RW] ^a Lyzalon [AF] ^a Pregabalin AMNEAL [EA] ^a PREGABALIN-DRLA [RZ] ^a Pregabalin Sandoz [SZ]	^a Cipla Pregabalin [LR] ^a Lyrica [UJ] ^a Neuroccord [CR] ^a Pregabalin APOTEX [GX] ^a Pregabalin GH [GQ] ^a Pregabalin-Teva [TB]

pregabalin 300 mg capsule, 56

2363J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	35.23	36.52	^a BTC PREGABALIN [JB] ^a LYPRALIN [RW] ^a Lyzalon [AF] ^a Pregabalin AMNEAL [EA] ^a PREGABALIN-DRLA [RZ] ^a Pregabalin Sandoz [SZ]	^a Cipla Pregabalin [LR] ^a Lyrica [UJ] ^a Neuroccord [CR] ^a Pregabalin APOTEX [GX] ^a Pregabalin GH [GQ] ^a Pregabalin-Teva [TB]

ANTIMIGRAINE PREPARATIONS

Selective serotonin (5HT₁) agonists

■ ELETRIPTAN

Caution Selective serotonin (5HT₁) agonists are contraindicated in patients with known or suspected coronary artery disease. The drug should not be used within 24 hours of ergotamine or dihydroergotamine use.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Migraine attack

Clinical criteria:

- The condition must have usually failed to respond to analgesics in the past.

eletriptan 40 mg tablet, 4

5290K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	25.65	26.94	Relpax [UJ]

eletriptan 80 mg tablet, 4

5291L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	25.65	26.94	Relpax [UJ]

■ NARATRIPTAN

Caution Selective serotonin (5HT₁) agonists are contraindicated in patients with known or suspected coronary artery disease. The drug should not be used within 24 hours of ergotamine or dihydroergotamine use.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required

Migraine attack

Clinical criteria:

- The condition must have usually failed to respond to analgesics in the past.

naratriptan 2.5 mg tablet, 2

8298R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	^s 1.96	*27.74	27.07	Naramig [AS]

■ NARATRIPTAN

Caution Selective serotonin (5HT₁) agonists are contraindicated in patients with known or suspected coronary artery disease. The drug should not be used within 24 hours of ergotamine or dihydroergotamine use.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required

Migraine attack

Clinical criteria:

- The condition must have usually failed to respond to analgesics in the past, **AND**
- Patient must be one in whom adverse events have occurred with other suitable PBS-listed drugs.

Authority required

Migraine attack

Clinical criteria:

- The condition must have usually failed to respond to analgesics in the past, **AND**
- Patient must be one in whom drug interactions have occurred with other suitable PBS-listed drugs.

Authority required

Migraine attack

Clinical criteria:

- The condition must have usually failed to respond to analgesics in the past, **AND**
- Patient must be one in whom drug interactions are expected to occur with other suitable PBS-listed drugs.

Authority required

Migraine attack

Clinical criteria:

- The condition must have usually failed to respond to analgesics in the past, **AND**
- Patient must be one in whom transfer to another suitable PBS-listed drug would cause patient confusion resulting in problems with compliance.

Authority required

Migraine attack

Clinical criteria:

- The condition must have usually failed to respond to analgesics in the past, **AND**
- Patient must be one in whom transfer to another suitable PBS-listed drug is likely to result in adverse clinical consequences.

naratriptan 2.5 mg tablet, 2

9734H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*27.74	29.03	Naramig [AS]

■ RIZATRIPTAN

Caution Selective serotonin (5HT₁) agonists are contraindicated in patients with known or suspected coronary artery disease. The drug should not be used within 24 hours of ergotamine or dihydroergotamine use.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note Pharmaceutical benefits that have the form rizatriptan wafer 10 mg (as benzoate) and pharmaceutical benefits that have the form rizatriptan tablet (orally disintegrating) 10 mg (as benzoate) are equivalent for the purposes of substitution.

Restricted benefit

Migraine attack

Clinical criteria:

- The condition must have usually failed to respond to analgesics in the past.

rizatriptan 10 mg wafer, 2

9313E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	2	5	..	*20.82	22.11	^a Maxalt [AL]	^a Rizatriptan Wafers-10mg [AF]

rizatriptan 10 mg orally disintegrating tablet, 2

10551H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	2	5	..	*20.82	22.11	^a APO-Rizatriptan [TX] ^a Rizatriptan AN ODT [EA] ^a Rizatriptan ODT APOTEX [GX]	^a RIXALT [RF] ^a Rizatriptan-AU [DZ] ^a Rizatriptan ODT GH [GQ]

■ SUMATRIPTAN

Caution Selective serotonin (5HT₁) agonists are contraindicated in patients with known or suspected coronary artery disease. The drug should not be used within 24 hours of ergotamine or dihydroergotamine use.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Pharmaceutical benefits that have the form sumatriptan tablet 50 mg (as succinate) and pharmaceutical benefits that have the form sumatriptan tablet (fast disintegrating) 50 mg (as succinate) are equivalent for the purposes of substitution.

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Migraine attack

Clinical criteria:

- The condition must have usually failed to respond to analgesics in the past.

SUMATRIPTAN Tablet 50 mg (base) (fast disintegrating), 4

10694W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	^B 4.14	21.37	18.52	^a Imigran FDT [AS]

sumatriptan 50 mg tablet, 4

1849H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	17.23	18.52	^a APO-Sumatriptan [TX] ^a Pharmacor Sumatriptan 50 [CR]	^a Iptam [AL] ^a Sumatran [OW]

^a Sumatriptan AN [EA]^a Sumatriptan generichealth [GQ]^a Sumatriptan Sandoz [SZ]^B4.14 21.37 18.52 ^a Imigran [LN]**SUMATRIPTAN Tablet (fast disintegrating) 50 mg (as succinate), 2**

8885P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	^B 4.14	*21.38	18.53	^a Imigran FDT [AS]

sumatriptan 50 mg tablet, 2

8144P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	2	5	..	*17.24	18.53	^a APO-Sumatriptan [TX]	^a Iptam [AL]
						^a Sumatran [OW]	^a Sumatriptan Sandoz [SZ]
			^B 4.14	*21.38	18.53	^a Imigran [LN]	

sumatriptan 20 mg/actuation nasal spray, 2 x 1 actuation

8341B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	23.86	25.15	Imigran [AS]

■ ZOLMITRIPTAN

Caution Selective serotonin (5HT₁) agonists are contraindicated in patients with known or suspected coronary artery disease. The drug should not be used within 24 hours of ergotamine or dihydroergotamine use.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Migraine attack

Clinical criteria:

- The condition must have usually failed to respond to analgesics in the past.

zolmitriptan 2.5 mg tablet, 2

8266C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	2	5	..	*23.90	25.19	^a APO-Zolmitriptan [TX]	^a Zoltrip [RW]
			^B 2.76	*26.66	25.19	^a Zomig [AP]	

Other antimigraine preparations**■ PIZOTIFEN****Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

pizotifen 500 microgram tablet, 100

3074T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	23.63	24.92	Sandomigran 0.5 [AE]

■ ANTIEPILEPTICS**ANTIEPILEPTICS****Barbiturates and derivatives****■ PHENOBARBITAL (PHENOBARBITONE)****Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Epilepsy

phenobarbital (phenobarbitone) 30 mg tablet, 200

1850J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	4	..	19.21	20.50	Phenobarb [RW]

phenobarbital (phenobarbitone) sodium 219 mg/mL injection, 5 x 1 mL ampoules

2138M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	35.05	36.34	Phenobarbitone Injection (Aspen Pharmacare Australia Pty Ltd) [AS]

■ PRIMIDONE**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

primidone 250 mg tablet, 100

11883K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	2	..	*87.40	41.30	APO-Primidone [LM]

primidone 250 mg tablet, 200

1939C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	68.92	41.30	Mysoline [LM]

*Hydantoin derivatives***■ PHENYTOIN****Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

phenytoin 50 mg chewable tablet, 200

1249R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	79.11	41.30	Dilantin Infatabs [UJ]

phenytoin 30 mg/5 mL oral liquid, 500 mL

2692Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	3	..	31.74	33.03	Dilantin [UJ]

phenytoin sodium 30 mg capsule, 200

1873N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	51.59	41.30	Dilantin Sodium [UJ]

phenytoin sodium 100 mg capsule, 200

1874P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	32.34	33.63	Dilantin Sodium [UJ]

*Succinimide derivatives***■ ETHOSUXIMIDE****Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

ethosuximide 250 mg/5 mL oral liquid, 200 mL

1414K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	72.61	41.30	Zarontin [IX]

■ ETHOSUXIMIDE**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note Pharmaceutical benefits that have the form ethosuximide 250 mg capsule, 200 and pharmaceutical benefits that have the form ethosuximide 250 mg capsule, 100 are equivalent for the purposes of substitution.

ethosuximide 250 mg capsule, 100

11703Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	2	..	*294.88	41.30	^a Zarontin [IX]

ethosuximide 250 mg capsule, 200

1413J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	294.88	41.30	^a Zarontin [IX]

Benzodiazepine derivatives**CLONAZEPAM****Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Epilepsy

clonazepam 1 mg/mL injection [5 x 1 mL ampoules] (& inert substance diluent [5 x 1 mL ampoules], 1 pack

1807D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	22.59	23.88	Rivotril [RO]

CLONAZEPAM

Caution Abuse of clonazepam has been reported. Refer to the current product information.

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required

Epilepsy

Clinical criteria:

- The condition must be neurologically proven.

clonazepam 2 mg tablet, 100

1806C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	2	..	*33.44	34.73	Paxam 2 [AF]

clonazepam 2.5 mg/mL (0.1 mg/drop) oral liquid, 10 mL

1808E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	..	..	*19.52	20.81	Rivotril [RO]

CLONAZEPAM

Caution Abuse of clonazepam has been reported. Refer to the current product information.

Note Pharmaceutical benefits that have form pack size clonazepam 500 microgram tablet, 100 and clonazepam 500 microgram tablet, 50 are equivalent for the purposes of substitution.

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required

Epilepsy

Clinical criteria:

- The condition must be neurologically proven.

clonazepam 500 microgram tablet, 100

1805B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	2	..	*23.42	24.71	^a Paxam 0.5 [AF]

clonazepam 500 microgram tablet, 50

11559J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	2	^B 3.68	*27.10	24.71	^a Rivotril [RO]

NITRAZEPAM**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required

Myoclonic epilepsy

Authority required

Malignant neoplasia (late stage)

Authority required

Insomnia

Clinical criteria:

- Patient must be receiving this drug for the management of insomnia, **AND**
- Patient must be receiving long-term nursing care on account of age, infirmity or other condition in a hospital, nursing home or residential facility, **AND**
- Patient must have demonstrated, within the past 6 months, benzodiazepine dependence by an unsuccessful attempt at gradual withdrawal.

Authority required

Insomnia

Clinical criteria:

- Patient must be receiving this drug for the management of insomnia, **AND**
- Patient must be receiving long-term nursing care, **AND**
- Patient must be one in respect of whom a Carer Allowance is payable as a disabled adult, **AND**
- Patient must have demonstrated, within the past 6 months, benzodiazepine dependence by an unsuccessful attempt at gradual withdrawal.

nitrazepam 5 mg tablet, 25

2732T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*15.44	16.73	Mogadon [IL]

Carboxamide derivatives**■ CARBAMAZEPINE****carbamazepine 100 mg/5 mL oral liquid, 300 mL**

5041H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	‡1	..	..	25.00	26.29	Tegretol Liquid [NV]

carbamazepine 200 mg modified release tablet, 200

5038E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	1	..	..	32.07	33.36	Tegretol CR 200 [NV]

carbamazepine 200 mg tablet, 100

1724R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	2	..	..	*31.66	32.95	^a Carbamazepine Sandoz [SZ]
			^B 2.96	*34.62	32.95	^a Tegretol 200 [NV]

carbamazepine 100 mg tablet, 100

5039F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	2	..	..	*23.78	25.07	^a Carbamazepine Sandoz [SZ]
			^B 3.00	*26.78	25.07	^a Tegretol 100 [NV]

carbamazepine 400 mg modified release tablet, 200

5037D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	1	..	..	50.13	41.30	Tegretol CR 400 [NV]

■ CARBAMAZEPINE**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

carbamazepine 100 mg/5 mL oral liquid, 300 mL

2427R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	‡1	5	..	25.00	26.29	Tegretol Liquid [NV]

carbamazepine 200 mg modified release tablet, 200

2426Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	32.07	33.36	Tegretol CR 200 [NV]

carbamazepine 200 mg tablet, 100

1706T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	2	..	*31.66	32.95	^a Carbamazepine Sandoz [SZ]
			^B 2.96	*34.62	32.95	^a Tegretol 200 [NV]

carbamazepine 100 mg tablet, 100

2422L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	2	..	*23.78	25.07	^a Carbamazepine Sandoz [SZ]
			^B 3.00	*26.78	25.07	^a Tegretol 100 [NV]

carbamazepine 400 mg modified release tablet, 200

2431Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	50.13	41.30	Tegretol CR 400 [NV]

■ OXCARBAZEPINE**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**5183**

Seizures

Clinical criteria:

- Patient must have partial epileptic seizures; OR
- Patient must have primary generalised tonic-clonic seizures, **AND**
- The condition must have failed to be controlled satisfactorily by other anti-epileptic drugs.

oxcarbazepine 600 mg tablet, 100

8586X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	147.78	41.30	Trileptal [NV]

oxcarbazepine 300 mg tablet, 100

8585W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	92.25	41.30	Trileptal [NV]

oxcarbazepine 150 mg tablet, 100

8584T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	60.64	41.30	Trileptal [NV]

oxcarbazepine 60 mg/mL oral liquid, 250 mL

8588B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*109.26	41.30	Trileptal [NV]

Fatty acid derivatives**■ TIAGABINE****Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**4928**

Partial epileptic seizures

Clinical criteria:

- The condition must have failed to be controlled satisfactorily by other anti-epileptic drugs.

tiagabine 10 mg tablet, 50

8222R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*109.80	41.30	Gabitril [TB]

tiagabine 5 mg tablet, 50

8221Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*60.90	41.30	Gabitril [TB]

tiagabine 15 mg tablet, 50

8223T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*154.70	41.30	Gabitril [TB]

■ VALPROATE

Caution There are reports of fatal hepatotoxicity, particularly in children.

There is increasing evidence of dose-related teratogenesis from this drug.

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

valproate sodium 500 mg enteric tablet, 100

2290M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	2	..	*35.26	36.55	^a Sodium Valproate Sandoz [SZ]	^a Valprease 500 [RW]

						^a Valproate Winthrop EC 500 [WA]	^a Valpro EC 500 [AF]
NP			^B 2.00	*37.26	36.55	^a Epilim EC [SW]	
valproate sodium 200 mg/5 mL oral liquid, 300 mL							
2293Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	
NP	2	2	..	*39.98	41.27	Epilim Liquid [SW]	
valproate sodium 200 mg/5 mL oral liquid, 300 mL							
2295T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	
NP	2	2	..	*39.98	41.27	Epilim Syrup [SW]	
valproate sodium 200 mg enteric tablet, 100							
2289L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	2	2	..	*23.76	25.05	^a Sodium Valproate Sandoz [SZ]	^a Valprease 200 [RW]
						^a Valproate Winthrop EC 200 [WA]	^a Valpro EC 200 [AF]
			^B 2.00	*25.76	25.05	^a Epilim EC [SW]	
valproate sodium 100 mg tablet, 100							
2294R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	
NP	2	2	..	*34.26	35.55	Epilim [SW]	

■ VIGABATRIN

Caution Visual field defects have been reported with this drug.

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)

4929

Epileptic seizures

Clinical criteria:

- The condition must have failed to be controlled satisfactorily by other anti-epileptic drugs.

vigabatrin 500 mg powder for oral liquid, 60 sachets

2668K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	
NP	1	5	..	65.73	41.30	Sabril [SW]	

vigabatrin 500 mg tablet, 100

2667J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	
NP	1	5	..	94.98	41.30	Sabril [SW]	

Other antiepileptics

■ BRIVARACETAM

Authority required (STREAMLINED)

10210

Intractable partial epileptic seizures

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a neurologist.

Clinical criteria:

- The treatment must be in combination with two or more anti-epileptic drugs which includes one second-line adjunctive agent, **AND**
- The condition must have failed to be controlled satisfactorily by other anti-epileptic drugs, which includes at least one first-line anti-epileptic agent and at least two second-line adjunctive anti-epileptic agents, **AND**
- The treatment must not be given concomitantly with levetiracetam, except for cross titration.

brivaracetam 100 mg tablet, 56

11339T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	
	1	5	..	163.69	41.30	Briviact [UC]	

brivaracetam 25 mg tablet, 56

11328F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	
	1	5	..	163.69	41.30	Briviact [UC]	

brivaracetam 50 mg tablet, 56

11334M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	
	1	5	..	163.69	41.30	Briviact [UC]	

brivaracetam 75 mg tablet, 56

11356Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	163.69	41.30	Briviact [UC]

■ BRIVARACETAM**Authority required (STREAMLINED)****10251**

Intractable partial epileptic seizures

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a neurologist.

Clinical criteria:

- The treatment must be in combination with two or more anti-epileptic drugs which includes one second-line adjunctive agent, **AND**
- The condition must have failed to be controlled satisfactorily by other anti-epileptic drugs, which includes at least one first-line anti-epileptic agent and at least two second-line adjunctive anti-epileptic agents, **AND**
- Patient must be unable to take a solid dose form of this drug, **AND**
- The treatment must not be given concomitantly with levetiracetam, except for cross titration.

brivaracetam 10 mg/mL oral liquid, 300 mL

11349H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	202.85	41.30	Briviact [UC]

■ BRIVARACETAM**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**10208**

Intractable partial epileptic seizures

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously been treated with PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must not be given concomitantly with levetiracetam.

brivaracetam 100 mg tablet, 56

11357R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	163.69	41.30	Briviact [UC]

brivaracetam 25 mg tablet, 56

11327E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	163.69	41.30	Briviact [UC]

brivaracetam 50 mg tablet, 56

11338R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	163.69	41.30	Briviact [UC]

brivaracetam 75 mg tablet, 56

11350J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	163.69	41.30	Briviact [UC]

■ BRIVARACETAM**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**10330**

Intractable partial epileptic seizures

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously been treated with PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must be unable to take a solid dose form of this drug, **AND**
- The treatment must not be given concomitantly with levetiracetam.

brivaracetam 10 mg/mL oral liquid, 300 mL

11358T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	202.85	41.30	Briviact [UC]

■ GABAPENTIN**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**4928**

Partial epileptic seizures

Clinical criteria:

- The condition must have failed to be controlled satisfactorily by other anti-epileptic drugs.

gabapentin 800 mg tablet, 100

8389M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	50.99	41.30	^a APO-Gabapentin [TX] ^a Gabapentin APOTEX [TY] ^a GAPENTIN [RF] ^a Pharmacor Gabapentin 800 [CR]	^a Gabapentin AN [EA] ^a Gabapentin Aspen 800 [RW] ^a Neurontin [UJ]

gabapentin 600 mg tablet, 100

8559L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	41.24	41.30	^a APO-Gabapentin [TX] ^a Gabapentin APOTEX [TY] ^a GAPENTIN [RF] ^a Pharmacor Gabapentin 600 [CR]	^a Gabapentin AN [EA] ^a Gabapentin Aspen 600 [RW] ^a Neurontin [UJ]

gabapentin 400 mg capsule, 100

1835N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	30.43	31.72	^a APO-Gabapentin [TX] ^a Gabacor [CR] ^a Gabapentin APOTEX [TY] ^a Gabapentin generichealth [HQ] ^a GAPENTIN [RF] ^a Nupentin 400 [AF]	^a APX-Gabapentin [GX] ^a Gabapentin AN [EA] ^a Gabapentin Aspen 400 [RW] ^a Gabapentin Sandoz [SZ] ^a Neurontin [UJ]

gabapentin 300 mg capsule, 100

1834M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	25.52	26.81	^a APO-Gabapentin [TX] ^a Gabacor [CR] ^a Gabapentin APOTEX [TY] ^a Gabapentin generichealth [HQ] ^a GAPENTIN [RF] ^a Nupentin 300 [AF]	^a APX-Gabapentin [GX] ^a Gabapentin AN [EA] ^a Gabapentin Aspen 300 [RW] ^a Gabapentin Sandoz [SZ] ^a Neurontin [UJ]

gabapentin 100 mg capsule, 100

8505P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	16.19	17.48	^a APO-Gabapentin [TX] ^a Gabacor [CR] ^a Gabapentin Aspen 100 [RW] ^a Neurontin [UJ]	^a APX-Gabapentin [GX] ^a Gabapentin APOTEX [TY] ^a GAPENTIN [RF] ^a Nupentin 100 [AF]

■ LACOSAMIDE**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**8813**

Intractable partial epileptic seizures

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a neurologist.

Clinical criteria:

- The treatment must be in combination with two or more anti-epileptic drugs which includes one second-line adjunctive agent, **AND**

- The condition must have failed to be controlled satisfactorily by other anti-epileptic drugs, which includes at least one first-line anti-epileptic agent and at least two second-line adjunctive anti-epileptic agents, **AND**
- The treatment must be for dose titration purposes.

lacosamide 150 mg tablet, 14

9336J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	65.08	41.30	Vimpat [UC]

lacosamide 50 mg tablet, 14

9333F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	29.71	31.00	Vimpat [UC]

lacosamide 100 mg tablet, 14

9334G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	47.39	41.30	Vimpat [UC]

■ LACOSAMIDE**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**8815**

Intractable partial epileptic seizures

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition.

lacosamide 50 mg tablet, 14

10293R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	5	..	*82.78	41.30	Vimpat [UC]

■ LACOSAMIDE**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**8770**

Intractable partial epileptic seizures

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a neurologist.

Clinical criteria:

- The treatment must be in combination with two or more anti-epileptic drugs which includes one second-line adjunctive agent, **AND**
- The condition must have failed to be controlled satisfactorily by other anti-epileptic drugs, which includes at least one first-line anti-epileptic agent and at least two second-line adjunctive anti-epileptic agents.

Authority required (STREAMLINED)**8815**

Intractable partial epileptic seizures

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition.

lacosamide 100 mg tablet, 56

9335H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	155.60	41.30	Vimpat [UC]

■ LACOSAMIDE**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note No increase in the maximum quantity or number of units may be authorised.

Authority required (STREAMLINED)**8770**

Intractable partial epileptic seizures

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a neurologist.

Clinical criteria:

- The treatment must be in combination with two or more anti-epileptic drugs which includes one second-line adjunctive agent, **AND**
- The condition must have failed to be controlled satisfactorily by other anti-epileptic drugs, which includes at least one first-line anti-epileptic agent and at least two second-line adjunctive anti-epileptic agents.

Authority required (STREAMLINED)**8815**

Intractable partial epileptic seizures

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition.

lacosamide 150 mg tablet, 56

9337K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	229.87	41.30	Vimpat [UC]

lacosamide 200 mg tablet, 56

9338L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	304.14	41.30	Vimpat [UC]

lacosamide 10 mg/mL oral liquid, 200 mL

11694L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±6	5	..	*357.30	41.30	Vimpat [UC]

■ LAMOTRIGINE**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**11081**

Epileptic seizures

Clinical criteria:

- The condition must have failed to be controlled satisfactorily by other anti-epileptic drugs; OR
- Patient must be a woman of childbearing potential.

lamotrigine 5 mg tablet, 56

8063J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	18.34	19.63	^a Lamictal [AS]	^a Lamotrigine Aspen 5 [RW]

lamotrigine 25 mg tablet, 56

2848X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	17.61	18.90	^a APO-Lamotrigine [TX] ^a Lamotrigine AN [EA] ^a Lamotrigine GH [GQ] ^a Logem [AL] ^a Reedos 25 [DO] ^a Lamictal [AS]	^a LAMITAN [RF] ^a Lamotrigine Aspen 25 [RW] ^a Lamotrigine Sandoz [SZ] ^a NOUMED LAMOTRIGINE [VO] ^a Sandoz Lamotrigine [HX]
			^B 1.59	19.20	18.90		

lamotrigine 200 mg tablet, 56

2851C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	37.99	39.28	^a APO-Lamotrigine [TX] ^a Lamotrigine AN [EA] ^a Lamotrigine GH [GQ] ^a Logem [AL] ^a Reedos 200 [DO] ^a Lamictal [AS]	^a LAMITAN [RF] ^a Lamotrigine Aspen 200 [RW] ^a Lamotrigine Sandoz [SZ] ^a NOUMED LAMOTRIGINE [VO] ^a Sandoz Lamotrigine [HX]
			^B 1.49	39.48	39.28		

lamotrigine 100 mg tablet, 56

2850B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	27.47	28.76	^a APO-Lamotrigine [TX] ^a Lamotrigine AN [EA] ^a Lamotrigine GH [GQ] ^a Logem [AL]	^a LAMITAN [RF] ^a Lamotrigine Aspen 100 [RW] ^a Lamotrigine Sandoz [SZ] ^a NOUMED LAMOTRIGINE [VO]

						^a Reedos 100 [DO]	^a Sandoz Lamotrigine [HX]
			^B 1.51	28.98	28.76	^a Lamictal [AS]	
lamotrigine 50 mg tablet, 56							
2849Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	21.30	22.59	^a APO-Lamotrigine [TX] ^a Lamotrigine AN [EA] ^a Lamotrigine GH [GQ] ^a Logem [AL]	^a LAMITAN [RF] ^a Lamotrigine Aspen 50 [RW] ^a Lamotrigine Sandoz [SZ] ^a NOUMED LAMOTRIGINE [VO]
						^a Reedos 50 [DO]	^a Sandoz Lamotrigine [HX]
			^B 1.44	22.74	22.59	^a Lamictal [AS]	

■ LEVETIRACETAM

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)

11116

Partial epileptic seizures

Clinical criteria:

- The condition must have failed to be controlled satisfactorily by other anti-epileptic drugs; OR
- Patient must be a woman of childbearing potential, **AND**
- The treatment must not be given concomitantly with brivaracetam, except for cross titration.

levetiracetam 500 mg tablet, 60

8655M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	31.22	32.51	^a APO-Levetiracetam [TX] ^a Kerron 500 [DO] ^a Levactam [ER] ^a Levetiracetam AN [EA] ^a Levetiracetam Mylan [AL] ^a Levi 500 [RW]	^a Keppra [UC] ^a Kevtam 500 [AF] ^a Levecetam 500 [RZ] ^a Levetiracetam GH [GQ] ^a Levetiracetam SZ [SZ] ^a NOUMED LEVETIRACETAM [VO]

levetiracetam 250 mg tablet, 60

8654L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	23.55	24.84	^a APO-Levetiracetam [TX] ^a Kerron 250 [DO] ^a Levactam [ER] ^a Levetiracetam AN [EA] ^a Levetiracetam Mylan [AL] ^a Levi 250 [RW]	^a Keppra [UC] ^a Kevtam 250 [AF] ^a Levecetam 250 [RZ] ^a Levetiracetam GH [GQ] ^a Levetiracetam SZ [SZ] ^a NOUMED LEVETIRACETAM [VO]

levetiracetam 1 g tablet, 60

8656N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	44.02	41.30	^a APO-Levetiracetam [TX] ^a Kerron 1000 [DO] ^a Levactam [ER] ^a Levetiracetam AN [EA] ^a Levetiracetam Mylan [AL] ^a Levi 1000 [RW]	^a Keppra [UC] ^a Kevtam 1000 [AF] ^a Levecetam 1000 [RZ] ^a Levetiracetam GH [GQ] ^a Levetiracetam SZ [SZ] ^a NOUMED LEVETIRACETAM [VO]

■ LEVETIRACETAM

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)

11077

Partial epileptic seizures

Clinical criteria:

- The condition must have failed to be controlled satisfactorily by other anti-epileptic drugs; OR
- Patient must be a woman of childbearing potential, **AND**
- Patient must be unable to take a solid dose form of levetiracetam, **AND**
- The treatment must not be given concomitantly with brivaracetam, except for cross titration.

levetiracetam 100 mg/mL oral liquid, 300 mL

9169N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	‡1	5	..	74.38	41.30	^a APO-Levetiracetam [TX] ^a Kerron [DO]	^a Keppra [UC] ^a Levetiracetam-AFT [AE]

■ PERAMPANEL**Authority required (STREAMLINED)****4656**

Intractable partial epileptic seizures

Treatment Phase: Initial

Clinical criteria:

- The treatment must be in combination with two or more anti-epileptic drugs which includes one second-line adjunctive agent, **AND**
- The condition must have failed to be controlled satisfactorily by other anti-epileptic drugs, which includes at least one first-line anti-epileptic agent and at least two second-line adjunctive anti-epileptic agents.

Treatment criteria:

- Must be treated by a neurologist.

perampanel 2 mg tablet, 7

10157N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	1	..	*47.16	41.30	Fycompa [EI]

■ PERAMPANEL**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**4658**

Intractable partial epileptic seizures

Treatment Phase: Continuing

Clinical criteria:

- Patient must have previously been issued with an authority prescription for this drug.

perampanel 4 mg tablet, 28

10162W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	154.62	41.30	Fycompa [EI]

perampanel 6 mg tablet, 28

10163X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	228.42	41.30	Fycompa [EI]

perampanel 12 mg tablet, 28

10159Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	302.19	41.30	Fycompa [EI]

perampanel 10 mg tablet, 28

10151G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	302.19	41.30	Fycompa [EI]

perampanel 8 mg tablet, 28

10160R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	302.19	41.30	Fycompa [EI]

■ PERAMPANEL

Note No applications for increased maximum quantities will be authorised.

Authority required (STREAMLINED)**7815**

Idiopathic generalised epilepsy with primary generalised tonic-clonic seizures

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a neurologist.

Clinical criteria:

- The condition must have failed to be controlled satisfactorily by at least two anti-epileptic drugs, **AND**
- The treatment must be in combination with at least one PBS-subsidised anti-epileptic drug, **AND**
- The treatment must be for dose titration purposes.

Population criteria:

- Patient must be aged 12 years or older.

perampanel 2 mg tablet, 7

11436X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	1	..	*47.16	41.30	Fycompa [EI]

■ PERAMPANEL

Note No applications for increased maximum quantities will be authorised.

Note Special Pricing Arrangements apply.

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**7789**

Idiopathic generalised epilepsy with primary generalised tonic-clonic seizures

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition.

Population criteria:

- Patient must be aged 12 years or older.

perampanel 4 mg tablet, 28

11418Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	154.62	41.30	Fycompa [EI]

perampanel 6 mg tablet, 28

11407J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	228.42	41.30	Fycompa [EI]

perampanel 12 mg tablet, 28

11409L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	302.19	41.30	Fycompa [EI]

perampanel 10 mg tablet, 28

11428L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	302.19	41.30	Fycompa [EI]

perampanel 8 mg tablet, 28

11429M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	302.19	41.30	Fycompa [EI]

■ STIRIPENTOL**Authority required (STREAMLINED)****10632**

Severe myoclonic epilepsy in infancy (Dravet syndrome)

Clinical criteria:

- Patient must have, or have had, generalised tonic-clonic seizures or generalised clonic seizures that are not adequately controlled with a benzodiazepine and valproate, **AND**
- The treatment must be as adjunctive therapy to a benzodiazepine and valproate.

Treatment criteria:

- Must be treated by a neurologist if treatment is being initiated; OR
- Must be treated by a neurologist if treatment is being continued or re-initiated; OR
- Must be treated by a paediatrician in consultation with a neurologist if treatment is being continued; OR
- Must be treated by a general practitioner in consultation with a neurologist if treatment is being continued.

stiripentol 250 mg capsule, 60

12103B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	3	..	*668.58	41.30	Diacomit [EU]

stiripentol 500 mg powder for oral liquid, 60 sachets

12088F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	3	..	*1294.46	41.30	Diacomit [EU]

stiripentol 500 mg capsule, 60

12107F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	3	..	*1294.46	41.30	Diacomit [EU]

stiripentol 250 mg powder for oral liquid, 60 sachets

12106E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	3	..	*668.58	41.30	Diacomit [EU]

■ SULTHIAME**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

sulthiame 50 mg tablet, 200

2099L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	68.18	41.30	Ospolot [FF]

sulthiame 200 mg tablet, 200

2100M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	161.88	41.30	Ospolot [FF]

■ TOPIRAMATE**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**5516**

Seizures

Clinical criteria:

- Patient must have partial epileptic seizures; OR
- Patient must have primary generalised tonic-clonic seizures; OR
- Patient must have seizures of the Lennox-Gastaut syndrome, **AND**
- The condition must have failed to be controlled satisfactorily by other anti-epileptic drugs.

topiramate 200 mg tablet, 60

8166T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	41.41	41.30	^a APO-Topiramate [TX] ^a RBX Topiramate [RA] ^a Topamax [JC] ^a Topiramate Sandoz [SZ]	^a Epiramax 200 [RW] ^a Tamate [AF] ^a Topiramate AN [EA]

topiramate 100 mg tablet, 60

8165R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	29.51	30.80	^a APO-Topiramate [TX] ^a RBX Topiramate [RA] ^a Topamax [JC] ^a Topiramate Sandoz [SZ]	^a Epiramax 100 [RW] ^a Tamate [AF] ^a Topiramate AN [EA]

■ TOPIRAMATE**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**5173**

Seizures

Clinical criteria:

- Patient must have partial epileptic seizures; OR
- Patient must have primary generalised tonic-clonic seizures; OR
- Patient must have seizures of the Lennox-Gastaut syndrome, **AND**
- The condition must have failed to be controlled satisfactorily by other anti-epileptic drugs, **AND**
- Patient must be unable to take a solid dose form of topiramate.

topiramate 50 mg capsule, 60

8520K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	27.55	28.84	Topamax Sprinkle [JC]

topiramate 25 mg capsule, 60

8372P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	21.33	22.62	Topamax Sprinkle [JC]

topiramate 15 mg capsule, 60

8371N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	23.33	24.62	Topamax Sprinkle [JC]

■ TOPIRAMATE**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**5516**

Seizures

Clinical criteria:

- Patient must have partial epileptic seizures; OR
- Patient must have primary generalised tonic-clonic seizures; OR
- Patient must have seizures of the Lennox-Gastaut syndrome, **AND**
- The condition must have failed to be controlled satisfactorily by other anti-epileptic drugs.

Authority required (STREAMLINED)**5325**

Migraine

Clinical criteria:

- The treatment must be for prophylaxis, **AND**
- Patient must have experienced an average of 3 or more migraines per month over a period of at least 6 months, **AND**
- Patient must have a contraindication to beta-blockers, as described in the relevant TGA-approved Product Information; OR
- Patient must have experienced intolerance of a severity necessitating permanent withdrawal during treatment with a beta-blocker, **AND**
- Patient must have a contraindication to pizotifen because the weight gain associated with this drug poses an unacceptable risk; OR
- Patient must have experienced intolerance of a severity necessitating permanent withdrawal during treatment with pizotifen.

Details of the contraindication and/or intolerance(s) must be documented in the patient's medical records when treatment is initiated.

topiramate 50 mg tablet, 60

8164Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	23.14	24.43	^a APO-Topiramate [TX] ^a RBX Topiramate [RA] ^a Topamax [JC] ^a Topiramate Sandoz [SZ]	^a Epiramax 50 [RW] ^a Tamate [AF] ^a Topiramate AN [EA]

topiramate 25 mg tablet, 60

8163P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	18.79	20.08	^a APO-Topiramate [TX] ^a RBX Topiramate [RA] ^a Topamax [JC] ^a Topiramate Sandoz [SZ]	^a Epiramax 25 [RW] ^a Tamate [AF] ^a Topiramate AN [EA]

■ ZONISAMIDE**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**4928**

Partial epileptic seizures

Clinical criteria:

- The condition must have failed to be controlled satisfactorily by other anti-epileptic drugs.

zonisamide 50 mg capsule, 56

9389E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	30.97	32.26	^a APO-Zonisamide [TX]	^a Zonegran [EI]

zonisamide 100 mg capsule, 56

9390F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	2	5	..	*75.16	41.30	^a APO-Zonisamide [TX]	^a Zonegran [EI]

zonisamide 25 mg capsule, 56

9388D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	23.38	24.67	^a APO-Zonisamide [TX]	^a Zonegran [EI]

ANTI-PARKINSON DRUGS**ANTICHOLINERGIC AGENTS***Tertiary amines***TRIHEXYPHENIDYL (BENZHEXOL)****trihexyphenidyl (benzhexol) hydrochloride 5 mg tablet, 200**

1110K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	23.48	24.77	Artane [RW]

TRIHEXYPHENIDYL (BENZHEXOL)

Note Pharmaceutical benefits that have the form trihexyphenidyl hydrochloride 2 mg tablet, 200 and pharmaceutical benefits that have the form trihexyphenidyl hydrochloride 2 mg tablet, 100 are equivalent for the purposes of substitution.

trihexyphenidyl (benzhexol) hydrochloride 2 mg tablet, 100

12250R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	2	..	*69.68	41.30	^a Trihexyphenidyl hydrochloride USP (Medsurge) [DZ]

trihexyphenidyl (benzhexol) hydrochloride 2 mg tablet, 200

1109J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	18.65	19.94	^a Artane [RW]

*Ethers of tropine or tropine derivatives***BENZATROPINE****benzatropine mesilate 2 mg tablet, 60**

2362H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	18.17	19.46	Benzotrop [FF]

benzatropine mesilate 2 mg/2 mL injection, 5 x 2 mL vials

11249C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	1	..	..	117.17	41.30	Benzatropine Injection [FF]

benzatropine mesilate 2 mg/2 mL injection, 5 x 2 mL vials

11255J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	117.17	41.30	Benzatropine Injection [FF]

DOPAMINERGIC AGENTS*Dopa and dopa derivatives***LEVODOPA + BENSERAZIDE****Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

levodopa 200 mg + benserazide 50 mg capsule, 100

2226E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	43.77	41.30	Madopar [RO]

levodopa 50 mg + benserazide 12.5 mg dispersible tablet, 100

8218M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	23.73	25.02	Madopar Rapid 62.5 [RO]

levodopa 100 mg + benserazide 25 mg capsule, 100

2225D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	34.98	36.27	Madopar 125 [RO]

levodopa 100 mg + benserazide 25 mg modified release capsule, 100

2231K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	37.27	38.56	Madopar HBS [RO]

levodopa 100 mg + benserazide 25 mg dispersible tablet, 100

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8219N	1	5	..	34.98	36.27	Madopar Rapid 125 [RO]

levodopa 100 mg + benserazide 25 mg tablet, 100

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
2229H	1	5	..	34.98	36.27	Madopar 125 [RO]

levodopa 200 mg + benserazide 50 mg tablet, 100

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
2228G	1	5	..	43.77	41.30	Madopar [RO]

levodopa 50 mg + benserazide 12.5 mg capsule, 100

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
2227F	1	5	..	23.73	25.02	Madopar 62.5 [RO]

■ LEVODOPA + CARBIDOPA**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

levodopa 250 mg + carbidopa 25 mg tablet, 100

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
1245M	1	5	..	37.65	38.94	^a APO-Levodopa/Carbidopa [TX]	^a SINADOPA 250/25 [RW]
						^a Sinemet [AL]	

levodopa 100 mg + carbidopa 25 mg tablet, 100

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
1242J	1	5	..	32.80	34.09	^a APO-Levodopa/Carbidopa [TX]	^a Kinson [AF]
						^a SINADOPA 100/25 [RW]	^a Sinemet 100/25 [AL]

■ LEVODOPA + CARBIDOPA

Note Pharmaceutical benefits that have the form levodopa 200 mg + carbidopa 50 mg modified release tablet, 100 and pharmaceutical benefits that have the form levodopa 200 mg + carbidopa 50 mg modified release tablet, 60 are equivalent for the purposes of substitution.

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Parkinson disease

Clinical criteria:

- The condition must be one in which fluctuations in motor function are not adequately controlled by frequent dosing with conventional formulations of levodopa with decarboxylase inhibitor.

levodopa 200 mg + carbidopa 50 mg modified release tablet, 60

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
11890T	1.7	5	..	*72.92	41.30	^a Sinemet CR Prolonged-Release Tablets [OQ]

levodopa 200 mg + carbidopa 50 mg modified release tablet, 100

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
1255C	1	5	..	67.88	41.30	^a Sinemet CR [AL]

■ LEVODOPA + CARBIDOPA

Note Special Pricing Arrangements apply.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note Patients should have adequate cognitive function to manage administration with a portable continuous infusion pump.

Authority required (STREAMLINED)

10197

Advanced Parkinson disease

Treatment Phase: Maintenance therapy

Clinical criteria:

- Patient must have severe disabling motor fluctuations not adequately controlled by oral therapy, **AND**
- Patient must have been commenced on treatment in a hospital-based movement disorder clinic.

levodopa 20 mg/mL + carbidopa monohydrate 5 mg/mL intestinal gel, 7 x 100 mL

11919H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	5	..	*5902.10	41.30	Duodopa [VE]

■ LEVODOPA + CARBIDOPA

Note Special Pricing Arrangements apply.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note Patients should have adequate cognitive function to manage administration with a portable continuous infusion pump.

Authority required (STREAMLINED)**10386**

Advanced Parkinson disease

Treatment Phase: Maintenance therapy

Clinical criteria:

- Patient must have severe disabling motor fluctuations not adequately controlled by oral therapy, **AND**
- Patient must have been commenced on treatment in a hospital-based movement disorder clinic, **AND**
- Patient must require continuous administration of levodopa without an overnight break; OR
- Patient must require a total daily dose of more than 2000 mg of levodopa.

levodopa 20 mg/mL + carbidopa monohydrate 5 mg/mL intestinal gel, 7 x 100 mL

8970D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	8	5	..	*11697.18	41.30	Duodopa [VE]

■ LEVODOPA + CARBIDOPA + ENTACAPONE**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Parkinson disease

Clinical criteria:

- Patient must be being treated with levodopa decarboxylase inhibitor combinations, **AND**
- Patient must be experiencing fluctuations in motor function due to end-of-dose effect.

Restricted benefit

Parkinson disease

Clinical criteria:

- Patient must be stabilised on concomitant treatment with levodopa decarboxylase inhibitor combinations and entacapone.

levodopa 75 mg + carbidopa 18.75 mg + entacapone 200 mg tablet, 100

9344T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	2	4	..	*243.82	41.30	^a Carlevent [TX] ^a Lecteva [TB]	^a L.C.E. Sandoz [HX] ^a Stalevo 75/18.75/200mg [SZ]

levodopa 100 mg + carbidopa 25 mg + entacapone 200 mg tablet, 100

8798C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	2	4	..	*257.06	41.30	^a Carlevent [TX] ^a Lecteva [TB]	^a L.C.E. Sandoz [HX] ^a Stalevo 100/25/200mg [SZ]

levodopa 150 mg + carbidopa 37.5 mg + entacapone 200 mg tablet, 100

8799D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	2	4	..	*280.74	41.30	^a Carlevent [TX] ^a Lecteva [TB]	^a L.C.E. Sandoz [HX] ^a Stalevo 150/37.5/200mg [SZ]

levodopa 50 mg + carbidopa 12.5 mg + entacapone 200 mg tablet, 100

8797B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	2	4	..	*233.40	41.30	^a Carlevent [TX] ^a Lecteva [TB]	^a L.C.E. Sandoz [HX] ^a Stalevo 50/12.5/200mg [SZ]

levodopa 200 mg + carbidopa 50 mg + entacapone 200 mg tablet, 100

9292C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	2	4	..	*302.50	41.30	^a Carlevent [TX] ^a Lecteva [TB]	^a L.C.E. Sandoz [HX] ^a Stalevo 200/50/200mg [SZ]

levodopa 125 mg + carbidopa 31.25 mg + entacapone 200 mg tablet, 100

9345W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	2	4	..	*266.54	41.30	^a Carlevent [TX] ^a Lecteva [TB]	^a L.C.E. Sandoz [HX] ^a Stalevo 125/31.25/200mg [SZ]

Adamantane derivatives

■ AMANTADINE

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Parkinson disease

Clinical criteria:

- The condition must not be drug induced.

amantadine hydrochloride 100 mg capsule, 100

3016R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	39.14	40.43	Symmetrel 100 [NV]

Dopamine agonists

■ APOMORPHINE

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)

10844

Parkinson disease

Treatment Phase: Maintenance therapy

Clinical criteria:

- Patient must have experienced severely disabling motor fluctuations which have not responded to other therapy, **AND**
- Patient must have been commenced on treatment in a specialist unit in a hospital setting.

apomorphine hydrochloride hemihydrate 50 mg/10 mL injection, 5 x 10 mL syringes

12319J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	36	5	..	*8178.30	41.30	Movapo PFS [TD]

apomorphine hydrochloride hemihydrate 20 mg/2 mL injection, 5 x 2 mL ampoules

12320K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	72	5	..	*6530.22	41.30	Movapo [TD]

apomorphine hydrochloride hemihydrate 100 mg/20 mL injection, 5 x 20 mL vials

12142C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	18	5	..	*7588.62	41.30	Apomine Solution for Infusion [PF]

apomorphine hydrochloride hemihydrate 50 mg/5 mL injection, 5 x 5 mL ampoules

12306Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	36	5	..	*8178.30	41.30	Movapo [TD]

■ APOMORPHINE

Note No increase in the maximum quantity or number of units may be authorised.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note Pharmaceutical benefits that have the form apomorphine injection 30 mg/3 mL pen device and pharmaceutical benefits that have the form apomorphine injection 30 mg/3 mL cartridge are equivalent for the purposes of substitution.

Authority required (STREAMLINED)

10844

Parkinson disease

Treatment Phase: Maintenance therapy

Clinical criteria:

- Patient must have experienced severely disabling motor fluctuations which have not responded to other therapy, **AND**
- Patient must have been commenced on treatment in a specialist unit in a hospital setting.

apomorphine hydrochloride hemihydrate 30 mg/3 mL injection, 5 x 3 mL cartridges

12133N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	20	5	..	*2749.14	41.30	^a Apomine Intermittent [PF]

apomorphine hydrochloride hemihydrate 30 mg/3 mL injection, 5 x 3 mL pen devices

12137T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	20	5	..	*2749.14	41.30	^a Movapo Pen [TD]

■ BROMOCRIPTINE

Caution Care should be taken when treating patients with advanced age and significant cognitive impairment with dopamine agonists.

Restricted benefit

Acromegaly

Restricted benefit

Parkinson disease

Restricted benefit

Pathological hyperprolactinaemia

Clinical criteria:

- Patient must be one in whom surgery is not indicated.

Restricted benefit

Pathological hyperprolactinaemia

Clinical criteria:

- Patient must have had surgery for this condition with incomplete resolution.

Restricted benefit

Pathological hyperprolactinaemia

Clinical criteria:

- Patient must be one in whom radiotherapy is not indicated.

Restricted benefit

Pathological hyperprolactinaemia

Clinical criteria:

- Patient must have had radiotherapy for this condition with incomplete resolution.

bromocriptine 2.5 mg tablet, 30

1443Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*33.76	35.05	Parlodel [SZ]

■ CABERGOLINE

Caution Care should be taken when treating patients with advanced age and significant cognitive impairment with dopamine agonists.

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Parkinson disease

cabergoline 1 mg tablet, 30

8393R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	60.53	41.30	Cabaser [PF]

cabergoline 2 mg tablet, 30

8394T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	77.04	41.30	Cabaser [PF]

■ PRAMIPEXOLE

Caution Episodes of sudden onset of sleep without warning, during activity, have been reported with this drug.

Care should be taken when treating patients with advanced age and significant cognitive impairment with dopamine agonists.

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Parkinson disease

pramipexole dihydrochloride monohydrate 125 microgram tablet, 30

9151P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	14.97	16.26	^a APO-Pramipexole [TX] ^a Sifrol [BY] ^a Simpral [AF]	^a Pramipexole AN [EA] ^a Simipex 0.125 [RW]

pramipexole dihydrochloride monohydrate 1 mg tablet, 100

9153R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	90.07	41.30	^a APO-Pramipexole [TX] ^a Sifrol [BY] ^a Simpral [AF]	^a Pramipexole AN [EA] ^a Simipex 1 [RW]

pramipexole dihydrochloride monohydrate 250 microgram tablet, 100

9152Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	29.90	31.19	^a APO-Pramipexole [TX] ^a Sifrol [BY] ^a Simpral [AF]	^a Pramipexole AN [EA] ^a Simipex 0.25 [RW]

■ PRAMIPEXOLE

Caution Episodes of sudden onset of sleep without warning, during activity, have been reported with this drug.

Care should be taken when treating patients with advanced age and significant cognitive impairment with dopamine agonists.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Parkinson disease

pramipexole dihydrochloride monohydrate 2.25 mg modified release tablet, 30

5143Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	57.82	41.30	^a APO-Pramipexole ER [TX] ^a Sifrol ER [BY]	^a Pramipexole XR GP [AF] ^a SIMIPEX XR [RW]

pramipexole dihydrochloride monohydrate 375 microgram modified release tablet, 30

3418X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	19.78	21.07	^a APO-Pramipexole ER [TX] ^a Sifrol ER [BY]	^a Pramipexole XR GP [AF] ^a SIMIPEX XR [RW]

pramipexole dihydrochloride monohydrate 1.5 mg modified release tablet, 30

3420B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	42.56	41.30	^a APO-Pramipexole ER [TX] ^a Sifrol ER [BY]	^a Pramipexole XR GP [AF] ^a SIMIPEX XR [RW]

pramipexole dihydrochloride monohydrate 750 microgram modified release tablet, 30

3419Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	27.29	28.58	^a APO-Pramipexole ER [TX] ^a Sifrol ER [BY]	^a Pramipexole XR GP [AF] ^a SIMIPEX XR [RW]

pramipexole dihydrochloride monohydrate 3.75 mg modified release tablet, 30

5145T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	93.91	41.30	^a APO-Pramipexole ER [TX] ^a Sifrol ER [BY]	^a Pramipexole XR GP [AF] ^a SIMIPEX XR [RW]

pramipexole dihydrochloride monohydrate 4.5 mg modified release tablet, 30

3422D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	111.96	41.30	^a APO-Pramipexole ER [TX] ^a Sifrol ER [BY]	^a Pramipexole XR GP [AF] ^a SIMIPEX XR [RW]

pramipexole dihydrochloride monohydrate 3 mg modified release tablet, 30

3421C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	78.65	41.30	^a APO-Pramipexole ER [TX] ^a Sifrol ER [BY]	^a Pramipexole XR GP [AF] ^a SIMIPEX XR [RW]

■ PRAMIPEXOLE

Caution Episodes of sudden onset of sleep without warning, during activity, have been reported with this drug.

Care should be taken when treating patients with advanced age and significant cognitive impairment with dopamine agonists.

Note This drug is not PBS-subsidised for Restless Legs Syndrome secondary to other causes

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Restricted benefit

Primary severe restless legs syndrome

Clinical criteria:

- Patient must manifest all 4 diagnostic criteria for Restless Legs Syndrome, **AND**
- Patient must have a baseline International Restless Legs Syndrome Rating Scale (IRLSRS) score greater than or equal to 21 points prior to initiation of pramipexole.

The date and IRLSRS score must be documented in the patient's medical records at the time pramipexole treatment is initiated.

The diagnostic criteria for Restless Legs Syndrome are:

- (a) An urge to move the legs usually accompanied or caused by unpleasant sensations in the legs; and

- (b) The urge to move or unpleasant sensations begin or worsen during periods of rest or inactivity such as lying or sitting; and
- (c) The urge to move or unpleasant sensations are partially or totally relieved by movement, such as walking or stretching, at least as long as the activity continues; and
- (d) The urge to move or unpleasant sensations are worse in the evening or night than during the day or only occur during the evening or night.

pramipexole dihydrochloride monohydrate 125 microgram tablet, 30

9393J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	14.97	16.26	Sifrol [BY]

pramipexole dihydrochloride monohydrate 250 microgram tablet, 100

9394K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	29.90	31.19	Sifrol [BY]

▪ ROTIGOTINE**Restricted benefit**

Parkinson disease

Clinical criteria:

- The treatment must be as adjunctive therapy to a levodopa-decarboxylase inhibitor combination.

rotigotine 4 mg/24 hours patch, 28

2384L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	93.58	41.30	Neupro [UC]

rotigotine 6 mg/24 hours patch, 28

2410W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	104.57	41.30	Neupro [UC]

rotigotine 8 mg/24 hours patch, 28

11140H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	115.09	41.30	Neupro [UC]

▪ ROTIGOTINE**Restricted benefit**

Parkinson disease

Clinical criteria:

- The treatment must be as adjunctive therapy to a levodopa-decarboxylase inhibitor combination.

rotigotine 2 mg/24 hours patch, 28

2385M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	73.20	41.30	Neupro [UC]

Monoamine oxidase B inhibitors**▪ RASAGILINE****Note** No increase in the maximum quantity or number of units may be authorised.**Note** No increase in the maximum number of repeats may be authorised.**Restricted benefit**

Parkinson disease

rasagiline 1 mg tablet, 30

1952R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	86.62	41.30	^a Azilect [TB] ^a Rasagiline-Teva [EV]	^a Pharmacor Rasagiline [CR] ^a Rasalect [TI]

▪ SAFINAMIDE**Note** No increase in the maximum quantity or number of units may be authorised.**Note** No increase in the maximum number of repeats may be authorised.**Restricted benefit**

Parkinson disease

Clinical criteria:

- The treatment must be as adjunctive therapy to a levodopa-decarboxylase inhibitor combination.

safinamide 50 mg tablet, 30

11656L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	64.37	41.30	Xadago [CS]

safinamide 100 mg tablet, 30

11666B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	116.96	41.30	Xadago [CS]

SELEGILINE**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Late stage Parkinson disease

Clinical criteria:

- The treatment must be as adjunctive therapy to a levodopa-decarboxylase inhibitor combination.

selegiline hydrochloride 5 mg tablet, 100

1973W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	54.06	41.30	Eldepryl [AS]

*Other dopaminergic agents***ENTACAPONE****Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Parkinson disease

Clinical criteria:

- The treatment must be as adjunctive therapy to a levodopa-decarboxylase inhibitor combination, **AND**
- Patient must be experiencing fluctuations in motor function due to end-of-dose effect.

entacapone 200 mg tablet, 100

8367J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	4	..	*226.56	41.30	Comtan [SZ]

PSYCHOLEPTICS**ANTIPSYCHOTICS***Phenothiazines with aliphatic side-chain***CHLORPROMAZINE****Note Shared Care Model:**

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

chlorpromazine hydrochloride 50 mg/2 mL injection, 10 x 2 mL ampoules

1195X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	21.94	23.23	Largactil [SW]

chlorpromazine hydrochloride 10 mg tablet, 100

1196Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	15.35	16.64	Largactil [SW]

chlorpromazine hydrochloride 25 mg tablet, 100

1197B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	15.76	17.05	Largactil [SW]

chlorpromazine hydrochloride 5 mg/mL oral liquid, 100 mL

1201F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	16.61	17.90	Largactil [SW]

chlorpromazine hydrochloride 100 mg tablet, 100

1199D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	19.79	21.08	Largactil [SW]

Phenothiazines with piperidine structure

■ PERICIAZINE

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

periciazine 2.5 mg tablet, 100

3052P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	15.23	16.52	Neulactil [SW]

periciazine 10 mg tablet, 100

3053Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	18.22	19.51	Neulactil [SW]

Butyrophenone derivatives

■ HALOPERIDOL

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

haloperidol 500 microgram tablet, 100

2761H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	15.49	16.78	Serenace [AS]

haloperidol 2 mg/mL oral liquid, 100 mL

2763K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	22.54	23.83	Serenace [AS]

haloperidol 5 mg/mL injection, 10 x 1 mL ampoules

2768Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	23.22	24.51	Serenace [AS]

haloperidol 1.5 mg tablet, 100

2767P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	15.84	17.13	Serenace [AS]

haloperidol 5 mg tablet, 50

2770T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	15.70	16.99	Serenace [AS]

■ HALOPERIDOL DECANOATE

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

haloperidol (as decanoate) 150 mg/3 mL injection, 5 x 3 mL ampoules

2766N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	45.01	41.30	Haldol decanoate [JC]

haloperidol (as decanoate) 50 mg/mL injection, 5 x 1 mL ampoules

2765M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	28.47	29.76	Haldol decanoate [JC]

Indole derivatives

■ LURASIDONE

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**4246**

Schizophrenia

lurasidone hydrochloride 80 mg tablet, 30

10529E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	111.97	41.30	^a APO-Lurasidone [TX] ^a Latuda [SE]	^a Ardix Lurasidone [RX] ^a Lurasidone Lupin [GQ]

^a LURASIDONE SUN [RA]^a Pharmacor Lurasidone [CR]**lurasidone hydrochloride 40 mg tablet, 30**

10526B

Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
1	5	..	62.00	41.30	^a APO-Lurasidone [TX] ^a Latuda [SE] ^a LURASIDONE SUN [RA]	^a Ardix Lurasidone [RX] ^a Lurasidone Lupin [GQ] ^a Pharmacor Lurasidone [CR]

■ ZIPRASIDONE**Note Shared Care Model:**

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**4246**

Schizophrenia

Authority required (STREAMLINED)**5742**

Acute mania or mixed episodes

Clinical criteria:

- The condition must be associated with bipolar I disorder, **AND**
- The treatment must be as monotherapy, **AND**
- The treatment must be limited to up to 6 months per episode.

ziprasidone 60 mg capsule, 60

9072L

Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
1	5	..	147.24	41.30	^a APO-Ziprasidone [TX] ^a Ziprasidone GH [GQ]	^a Zeldox [UJ] ^a ZIPROX [RW]

ziprasidone 20 mg capsule, 60

9070J

Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
1	5	..	56.62	41.30	^a APO-Ziprasidone [TX] ^a Ziprasidone GH [GQ]	^a Zeldox [UJ] ^a ZIPROX [RW]

ziprasidone 40 mg capsule, 60

9071K

Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
1	5	..	101.37	41.30	^a APO-Ziprasidone [TX] ^a Ziprasidone GH [GQ]	^a Zeldox [UJ] ^a ZIPROX [RW]

ziprasidone 80 mg capsule, 60

9073M

Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
1	5	..	194.22	41.30	^a APO-Ziprasidone [TX] ^a Ziprasidone GH [GQ]	^a Zeldox [UJ] ^a ZIPROX [RW]

Thioxanthene derivatives**■ FLUPENTIXOL DECANOATE****Note Shared Care Model:**

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

flupentixol decanoate 100 mg/mL injection, 5 x 1 mL ampoules

2257T

Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
1	..	..	49.37	41.30	Fluanxol Concentrated Depot [LU]

flupentixol decanoate 20 mg/mL injection, 5 x 1 mL ampoules

2255Q

Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
1	..	..	24.27	25.56	Fluanxol Depot [LU]

■ ZUCLOPENTHIXOL DECANOATE**Note Shared Care Model:**

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

zuclopenthixol decanoate 200 mg/mL injection, 5 x 1 mL ampoules

8097E

Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
1	..	..	30.12	31.41	Clopixol Depot [LU]

Diazepines, oxazepines, thiazepines and oxepines

■ **ASENAPINE****Note Shared Care Model:**

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**4246**

Schizophrenia

Authority required (STREAMLINED)**5773**

Acute mania or mixed episodes

Clinical criteria:

- The condition must be associated with bipolar I disorder, **AND**
- The treatment must be limited to up to 6 months per episode.

Authority required (STREAMLINED)**5719**

Bipolar I disorder

Clinical criteria:

- The treatment must be maintenance therapy, **AND**
- The treatment must be as monotherapy.

asenapine 10 mg sublingual wafer, 60

5141N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	234.74	41.30	Saphris [OQ]

asenapine 5 mg sublingual wafer, 60

5140M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	143.63	41.30	Saphris [OQ]

■ **OLANZAPINE****Note Shared Care Model:**

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**5856**

Schizophrenia

Authority required (STREAMLINED)**5869**

Bipolar I disorder

Clinical criteria:

- The treatment must be maintenance therapy.

olanzapine 2.5 mg tablet, 28

8170B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	15.01	16.30	^a APO-Olanzapine [TX] ^a Olanzapine APOTEX [GX] ^a Olanzapine RBX [RA] ^a Ozin 2.5 [DO] ^a Zypine [AF]	^a Olanzapine AN [EA] ^a Olanzapine-DRLA [RZ] ^a Olanzapine Sandoz [SZ] ^a PRYZEX [RW]
			^B 2.79	17.80	16.30	^a Zyprexa [LY]	

olanzapine 5 mg tablet, 28

8185T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	17.53	18.82	^a APO-Olanzapine [TX] ^a Olanzapine APOTEX [GX] ^a Olanzapine RBX [RA] ^a Ozin 5 [DO] ^a Zypine [AF]	^a Olanzapine AN [EA] ^a Olanzapine-DRLA [RZ] ^a Olanzapine Sandoz [SZ] ^a PRYZEX [RW]
			^B 2.97	20.50	18.82	^a Zyprexa [LY]	

olanzapine 7.5 mg tablet, 28

8186W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	20.33	21.62	^a APO-Olanzapine [TX] ^a Olanzapine APOTEX [GX] ^a Olanzapine RBX [RA] ^a Ozin 7.5 [DO] ^a Zypine [AF]	^a Olanzapine AN [EA] ^a Olanzapine-DRLA [RZ] ^a Olanzapine Sandoz [SZ] ^a PRYZEX [RW]
			^B 3.00	23.33	21.62	^a Zyprexa [LY]	

olanzapine 10 mg tablet, 28

8187X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	23.09	24.38	^a APO-Olanzapine [TX] ^a Olanzapine APOTEX [GX] ^a Olanzapine RBX [RA] ^a Ozin 10 [DO] ^a Zypine [AF]	^a Olanzapine AN [EA] ^a Olanzapine-DRLA [RZ] ^a Olanzapine Sandoz [SZ] ^a PRYZEX [RW]
			^B 3.00	26.09	24.38	^a Zyprexa [LY]	

■ OLANZAPINE

Note Pharmaceutical benefits that have the form olanzapine tablet 5 mg (orally disintegrating) and pharmaceutical benefits that have the form olanzapine wafer 5 mg are equivalent for the purposes of substitution.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**5856**

Schizophrenia

Authority required (STREAMLINED)**5869**

Bipolar I disorder

Clinical criteria:

- The treatment must be maintenance therapy.

olanzapine 5 mg orally disintegrating tablet, 28

3381Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	17.53	18.82	^a APO-Olanzapine ODT [TX] ^a Olanzapine ODT generichealth 5 [GQ] ^a PRYZEX ODT [RW]	^a Olanzapine AN ODT [EA] ^a Olanzapine Sandoz ODT 5 [SZ]

olanzapine 5 mg wafer, 28

8433W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	17.53	18.82	^a Zypine ODT [AF]
			^B 2.97	20.50	18.82	^a Zyprexa Zydis [LY]

■ OLANZAPINE

Note Pharmaceutical benefits that have the form olanzapine tablet 10 mg (orally disintegrating) and pharmaceutical benefits that have the form olanzapine wafer 10 mg are equivalent for the purposes of substitution.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**5856**

Schizophrenia

Authority required (STREAMLINED)**5869**

Bipolar I disorder

Clinical criteria:

- The treatment must be maintenance therapy.

olanzapine 10 mg orally disintegrating tablet, 28

3382B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	23.09	24.38	^a APO-Olanzapine ODT [TX] ^a Olanzapine ODT generichealth 10 [GQ] ^a PRYZEX ODT [RW]	^a Olanzapine AN ODT [EA] ^a Olanzapine Sandoz ODT 10 [SZ]

olanzapine 10 mg wafer, 28

8434X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	23.09	24.38	^a Zypine ODT [AF]
			^B 3.00	26.09	24.38	^a Zyprexa Zydis [LY]

■ OLANZAPINE

Note Pharmaceutical benefits that have the form olanzapine tablet 15 mg (orally disintegrating) and pharmaceutical benefits that have the form olanzapine wafer 15 mg are equivalent for the purposes of substitution.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical

practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)

5856

Schizophrenia

Authority required (STREAMLINED)

5869

Bipolar I disorder

Clinical criteria:

- The treatment must be maintenance therapy.

olanzapine 15 mg wafer, 28

8952E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	28.63	29.92	^a Zypine ODT [AF]
			^B 3.00	31.63	29.92	^a Zyprexa Zydis [LY]

olanzapine 15 mg orally disintegrating tablet, 28

3384D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	28.63	29.92	^a APO-Olanzapine ODT [TX]	^a Olanzapine AN ODT [EA]
						^a Olanzapine Sandoz ODT 15 [SZ]	^a PRYZEX ODT [RW]

■ **OLANZAPINE**

Note Pharmaceutical benefits that have the form olanzapine tablet 20 mg (orally disintegrating) and pharmaceutical benefits that have the form olanzapine wafer 20 mg are equivalent for the purposes of substitution.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)

5856

Schizophrenia

Authority required (STREAMLINED)

5869

Bipolar I disorder

Clinical criteria:

- The treatment must be maintenance therapy.

olanzapine 20 mg wafer, 28

8953F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	34.17	35.46	^a Zypine ODT [AF]
			^B 3.00	37.17	35.46	^a Zyprexa Zydis [LY]

olanzapine 20 mg orally disintegrating tablet, 28

3385E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	34.17	35.46	^a APO-Olanzapine ODT [TX]	^a Olanzapine AN ODT [EA]
						^a Olanzapine Sandoz ODT 20 [SZ]	^a PRYZEX ODT [RW]

■ **OLANZAPINE**

Caution Monitor for post-injection syndrome for at least two hours after each injection.

Note Special Pricing Arrangements apply.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)

4304

Schizophrenia

olanzapine 405 mg modified release injection [1 vial] (&) inert substance diluent [3 mL vial], 1 pack

9303P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	505.12	41.30	Zyprexa Relprevv [LY]

olanzapine 300 mg modified release injection [1 vial] (&) inert substance diluent [3 mL vial], 1 pack

9295F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*817.58	41.30	Zyprexa Relprevv [LY]

olanzapine 210 mg modified release injection [1 vial] (&) inert substance diluent [3 mL vial], 1 pack

9294E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*505.12	41.30	Zyprexa Relprevv [LY]

■ QUETIAPINE**Note Shared Care Model:**

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**4246**

Schizophrenia

Authority required (STREAMLINED)**5611**

Acute mania

Clinical criteria:

- The condition must be associated with bipolar I disorder, **AND**
- The treatment must be as monotherapy, **AND**
- The treatment must be limited to up to 6 months per episode.

Authority required (STREAMLINED)**5639**

Bipolar I disorder

Clinical criteria:

- The treatment must be maintenance therapy.

quetiapine 300 mg tablet, 60

8580N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	45.05	41.30	^a APO-Quetiapine [TX] ^a Pharmacor Quetiapine 300 [CR] ^a Quetiapine Actavis 300 [ED] ^a Quetiapine APOTEX [GX] ^a Quetiapine GH 300 [GQ] ^a Quetiapine Sandoz [SZ] ^a Syquet [AF]	^a Kaptan [ER] ^a Quetia 300 [RW] ^a Quetiapine AN [EA] ^a Quetiapine-DRLA [RZ] ^a Quetiapine RBX [RA] ^a Quetiapine Sandoz Pharma [HX]
			^b 5.00	50.05	41.30	^a Seroquel [AP]	

quetiapine 200 mg modified release tablet, 60

9203J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	46.00	41.30	^a APO-Quetiapine XR [TX] ^a QUEPINE XR [RF] ^a Quetia XR [OW]	^a APX-Quetiapine XR [TY] ^a QUETIAPINE-AS XR [RW] ^a Tevatiapine XR [SZ]
			^b 2.15	48.15	41.30	^a Seroquel XR [AP]	

quetiapine 300 mg modified release tablet, 60

9204K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	55.66	41.30	^a APO-Quetiapine XR [TX] ^a QUEPINE XR [RF] ^a Quetia XR [OW]	^a APX-Quetiapine XR [TY] ^a QUETIAPINE-AS XR [RW] ^a Tevatiapine XR [SZ]
			^b 2.15	57.81	41.30	^a Seroquel XR [AP]	

quetiapine 400 mg modified release tablet, 60

9205L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	71.51	41.30	^a APO-Quetiapine XR [TX] ^a QUEPINE XR [RF] ^a Quetia XR [OW]	^a APX-Quetiapine XR [TY] ^a QUETIAPINE-AS XR [RW] ^a Tevatiapine XR [SZ]
			^b 2.15	73.66	41.30	^a Seroquel XR [AP]	

quetiapine 200 mg tablet, 60

8458E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	34.52	35.81	^a APO-Quetiapine [TX] ^a Pharmacor Quetiapine 200 [CR] ^a Quetiapine Actavis 200 [ED] ^a Quetiapine APOTEX [GX] ^a Quetiapine GH 200 [GQ] ^a Quetiapine Sandoz [SZ] ^a Syquet [AF]	^a Kaptan [ER] ^a Quetia 200 [RW] ^a Quetiapine AN [EA] ^a Quetiapine-DRLA [RZ] ^a Quetiapine RBX [RA] ^a Quetiapine Sandoz Pharma [HX]

			^B 5.00	39.52	35.81	^a Seroquel [AP]	
quetiapine 150 mg modified release tablet, 60							
5458G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	28.54	29.83	^a APX-Quetiapine XR [TY] ^a Quetia XR [OW]	^a QUEPINE XR [RW] ^a Tevatiapine XR [SZ]
			^B 10.75	39.29	29.83	^a Seroquel XR [AP]	
quetiapine 100 mg tablet, 90							
8457D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	28.54	29.83	^a APO-Quetiapine [TX] ^a Pharmacor Quetiapine 100 [CR] ^a Quetiapine Actavis 100 [ED] ^a Quetiapine APOTEX [GX] ^a Quetiapine GH 100 [GQ] ^a Quetiapine Sandoz [SZ] ^a Syquet [AF]	^a Kaptan [ER] ^a Quetia 100 [RW] ^a Quetiapine AN [EA] ^a Quetiapine-DRLA [RZ] ^a Quetiapine RBX [RA] ^a Quetiapine Sandoz Pharma [HX]
			^B 4.99	33.53	29.83	^a Seroquel [AP]	
quetiapine 50 mg modified release tablet, 60							
9202H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	23.70	24.99	^a APO-Quetiapine XR [TX] ^a QUEPINE XR [RF] ^a Quetia XR [OW]	^a APX-Quetiapine XR [TY] ^a QUETIAPINE-AS XR [RW] ^a Tevatiapine XR [SZ]
			^B 5.16	28.86	24.99	^a Seroquel XR [AP]	

■ QUETIAPINE

Note No increase in the maximum quantity or number of units may be authorised.

Note Authority applications for increased repeats up to a maximum of 5 may be authorised for patients requiring dose optimisation for this condition not adequately provided by other strengths of this drug.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**7916**

Schizophrenia

Authority required (STREAMLINED)**7927**

Acute mania

Clinical criteria:

- The condition must be associated with bipolar I disorder, **AND**
- The treatment must be as monotherapy.

Authority required (STREAMLINED)**7893**

Bipolar I disorder

Clinical criteria:

- The treatment must be maintenance therapy.

quetiapine 25 mg tablet, 60

8456C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	17.86	19.15	^a APO-Quetiapine [TX] ^a Pharmacor Quetiapine 25 [CR] ^a Quetiapine AN [EA] ^a Quetiapine-DRLA [RZ] ^a Quetiapine RBX [RA] ^a Quetiapine Sandoz Pharma [HX]	^a Kaptan [ER] ^a Quetia 25 [RW] ^a Quetiapine APOTEX [GX] ^a Quetiapine GH 25 [GQ] ^a Quetiapine Sandoz [SZ] ^a Syquet [AF]
			^B 5.00	22.86	19.15	^a Seroquel [AP]	

Benzamides**■ AMISULPRIDE****Note Shared Care Model:**

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**4246**

Schizophrenia

amisulpride 400 mg tablet, 60

8596K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	66.39	41.30	^a Amipride 400 [RW]	^a Amisulpride 400 Winthrop [WA]
						^a Amisulpride AN [EA]	^a Amisulpride Sandoz [SZ]
						^a Amisulpride Sandoz Pharma [HX]	^a APO-Amisulpride [TX]
						^a Solian 400 [SW]	^a Sulprix [AF]

amisulpride 100 mg/mL oral liquid, 60 mL

8736T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	2	5	..	*142.86	41.30	Solian Solution [SW]	

amisulpride 200 mg tablet, 60

8595J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	42.01	41.30	^a Amisulpride 200 Winthrop [WA]	^a Amisulpride AN [EA]
						^a Amisulpride Sandoz [SZ]	^a Amisulpride Sandoz Pharma [HX]
						^a APO-Amisulpride [TX]	^a Solian 200 [SW]
						^a Sulprix [AF]	

amisulpride 100 mg tablet, 30

8594H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	18.98	20.27	^a Amisulpride 100 Winthrop [WA]	^a Amisulpride AN [EA]
						^a Amisulpride Sandoz [SZ]	^a Amisulpride Sandoz Pharma [HX]
						^a APO-Amisulpride [TX]	^a Solian 100 [SW]
						^a Sulprix [AF]	

Other antipsychotics**■ ARIPIPRAZOLE****Note Shared Care Model:**

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**4246**

Schizophrenia

aripiprazole 300 mg modified release injection [1 vial] (&) inert substance diluent [2 mL vial], 1 pack

10224D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	306.42	41.30	Abilify Maintena [LU]	

aripiprazole 400 mg modified release injection [1 vial] (&) inert substance diluent [2 mL vial], 1 pack

10219W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	381.27	41.30	Abilify Maintena [LU]	

aripiprazole 30 mg tablet, 30

8720Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	126.49	41.30	^a Abilify [OS]	^a Abyraz [AF]
						^a APO-Aripiprazole [TX]	^a Aripiprazole AN [EA]
						^a Aripiprazole generichealth [HQ]	^a Aripiprazole GH [GQ]
						^a Aripiprazole Sandoz [SZ]	^a Tevaripiprazole [TB]

aripiprazole 10 mg tablet, 30

8717T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	66.07	41.30	^a Abilify [OS]	^a Abyraz [AF]
						^a APO-Aripiprazole [TX]	^a Aripiprazole AN [EA]
						^a Aripiprazole generichealth [HQ]	^a Aripiprazole GH [GQ]
						^a Aripiprazole Sandoz [SZ]	^a Tevaripiprazole [TB]

aripiprazole 15 mg tablet, 30

8718W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	88.72	41.30	^a Abilify [OS]	^a Abyraz [AF]
						^a APO-Aripiprazole [TX]	^a Aripiprazole AN [EA]
						^a Aripiprazole generichealth [HQ]	^a Aripiprazole GH [GQ]

^a Aripiprazole Sandoz [SZ]^a Tevaripiprazole [TB]**aripiprazole 20 mg tablet, 30**

8719X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	105.39	41.30	^a Abilify [OS] ^a APO-Aripiprazole [TX] ^a Aripiprazole generichealth [HQ] ^a Aripiprazole Sandoz [SZ]	^a Abyraz [AF] ^a Aripiprazole AN [EA] ^a Aripiprazole GH [GQ] ^a Tevaripiprazole [TB]

■ BREXPIPIRAZOLE**Note Shared Care Model:**

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**4246**

Schizophrenia

brexpiprazole 1 mg tablet, 30

11189X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	144.86	41.30	Rexulti [LU]

brexpiprazole 3 mg tablet, 30

11190Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	144.86	41.30	Rexulti [LU]

brexpiprazole 2 mg tablet, 30

11188W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	144.86	41.30	Rexulti [LU]

brexpiprazole 4 mg tablet, 30

11184P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	144.86	41.30	Rexulti [LU]

■ PALIPERIDONE**Note Shared Care Model:**

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**4246**

Schizophrenia

paliperidone 150 mg modified release injection, 1 syringe

5109X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	380.73	41.30	Invega Sustenna [JC]

paliperidone 100 mg modified release injection, 1 syringe

5107T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	380.73	41.30	Invega Sustenna [JC]

paliperidone 75 mg modified release injection, 1 syringe

5103N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	311.11	41.30	Invega Sustenna [JC]

paliperidone 6 mg modified release tablet, 28

9141D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	127.69	41.30	Invega [JC]

paliperidone 50 mg modified release injection, 1 syringe

5102M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	240.77	41.30	Invega Sustenna [JC]

paliperidone 3 mg modified release tablet, 28

9140C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	68.90	41.30	Invega [JC]

paliperidone 9 mg modified release tablet, 28

9142E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	188.01	41.30	Invega [JC]

paliperidone 25 mg modified release injection, 1 syringe

5100K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	123.91	41.30	Invega Sustenna [JC]

■ PALIPERIDONE**Note Shared Care Model:**

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note Patient dosage is to be determined as per the dose transition table in the Product Information based on the maintenance dose of paliperidone once monthly injection.

Note No increase in the maximum number of repeats may be authorised.

Note No increase in the maximum quantity or number of units may be authorised.

Authority required (STREAMLINED)**6832**

Schizophrenia

Clinical criteria:

- Patient must have previously received and be stabilised on PBS-subsidised paliperidone once-monthly injection for at least 4 consecutive months.

paliperidone 175 mg/0.875 mL modified release injection, 0.875 mL syringe

11085K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	708.28	41.30	Invega Trinza [JC]

paliperidone 525 mg/2.625 mL modified release injection, 2.625 mL syringe

11066K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	1106.58	41.30	Invega Trinza [JC]

paliperidone 263 mg/1.315 mL modified release injection, 1.315 mL syringe

11072R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	912.33	41.30	Invega Trinza [JC]

paliperidone 350 mg/1.75 mL modified release injection, 1.75 mL syringe

11094X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	1106.58	41.30	Invega Trinza [JC]

■ RISPERIDONE**Note Shared Care Model:**

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**4246**

Schizophrenia

Authority required (STREAMLINED)**5907**

Acute mania

Clinical criteria:

- The condition must be associated with bipolar I disorder, **AND**
- The treatment must be as adjunctive therapy to mood stabilisers, **AND**
- The treatment must be limited to up to 6 months per episode.

risperidone 4 mg tablet, 60

3172Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	41.72	41.30	^a APO-Risperidone [TX]	^a NOUMED RISPERIDONE [VO]
						^a Rispa [RW]	^a Risperdal [JC]
						^a Risperidone AMNEAL [EF]	^a Risperidone generichealth [GQ]
						^a Risperidone Sandoz [SZ]	^a Rispernia [ER]
						^a Rixadone [AF]	

risperidone 1 mg tablet, 60

3169T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	18.26	19.55	^a APO-Risperidone [TX]	^a NOUMED RISPERIDONE [VO]
						^a Ozidal [RA]	^a Rispa [RW]
						^a Risperdal [JC]	^a Risperidone AMNEAL [EF]
						^a Risperidone generichealth [GQ]	^a Risperidone Sandoz [SZ]
						^a Rispernia [ER]	^a Rixadone [AF]

risperidone 1 mg/mL oral liquid, 100 mL

8100H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	±1	5	..	113.55	41.30	^a Risperdal [JC]	^a Rixadone [AF]

risperidone 2 mg tablet, 60

3170W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	26.29	27.58	^a APO-Risperidone [TX]	^a NOUMED RISPERIDONE [VO]
						^a Ozidal [RA]	^a Rispa [RW]
						^a Risperdal [JC]	^a Risperidone AMNEAL [EF]
						^a Risperidone generichealth [GQ]	^a Risperidone Sandoz [SZ]
						^a Rispernia [ER]	^a Rixadone [AF]

risperidone 3 mg tablet, 60

3171X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	34.02	35.31	^a APO-Risperidone [TX]	^a NOUMED RISPERIDONE [VO]
						^a Ozidal [RA]	^a Rispa [RW]
						^a Risperdal [JC]	^a Risperidone AMNEAL [EF]
						^a Risperidone generichealth [GQ]	^a Risperidone Sandoz [SZ]
						^a Rispernia [ER]	^a Rixadone [AF]

■ RISPERIDONE**Note Shared Care Model:**

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**6898**

Severe behavioural disturbances

Clinical criteria:

- Patient must have autism spectrum disorder, **AND**
- The treatment must be under the supervision of a paediatrician or psychiatrist, **AND**
- The treatment must be in combination with non-pharmacological measures.

Population criteria:

- Patient must be under 18 years of age.

Behaviour disturbances are defined as severe aggression and injuries to self or others where non-pharmacological methods alone have been unsuccessful.

The diagnosis of autism spectrum disorder must be made based on the Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-V) or ICD-10 international classification of mental and behavioural disorders.

Authority required (STREAMLINED)**6899**

Severe behavioural disturbances

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have autism spectrum disorder, **AND**
- Patient must have been commenced on PBS-subsidised treatment with risperidone prior to turning 18 years of age, **AND**
- The treatment must be under the supervision of a paediatrician or psychiatrist, **AND**
- The treatment must be in combination with non-pharmacological measures.

Population criteria:

- Patient must be aged 18 years or older.

Behaviour disturbances are defined as severe aggression and injuries to self or others where non-pharmacological methods alone have been unsuccessful.

The diagnosis of autism spectrum disorder must be made based on the Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-V) or ICD-10 international classification of mental and behavioural disorders.

risperidone 1 mg tablet, 60

8789N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	2	..	18.26	19.55	^a APO-Risperidone [TX]	^a NOUMED RISPERIDONE [VO]
						^a Ozidal [RA]	^a Rispa [RW]
						^a Risperdal [JC]	^a Risperidone AMNEAL [EF]
						^a Risperidone generichealth [GQ]	^a Risperidone Sandoz [SZ]
						^a Rispernia [ER]	^a Rixadone [AF]

risperidone 1 mg/mL oral liquid, 100 mL

9293D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	±1	2	..	113.55	41.30	^a Risperdal [JC]	^a Rixadone [AF]

■ RISPERIDONE**Note Shared Care Model:**

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**6897**

Severe behavioural disturbances

Clinical criteria:

- Patient must have autism spectrum disorder, **AND**
- The treatment must be under the supervision of a paediatrician or psychiatrist, **AND**
- The treatment must be in combination with non-pharmacological measures.

Population criteria:

- Patient must be under 18 years of age.

Behaviour disturbances are defined as severe aggression and injuries to self or others where non-pharmacological methods alone have been unsuccessful.

The diagnosis of autism spectrum disorder must be made based on the Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-V) or ICD-10 international classification of mental and behavioural disorders.

Authority required (STREAMLINED)**6938**

Severe behavioural disturbances

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have autism spectrum disorder, **AND**
- Patient must have been commenced on PBS-subsidised treatment with risperidone prior to turning 18 years of age, **AND**
- The treatment must be under the supervision of a paediatrician or psychiatrist, **AND**
- The treatment must be in combination with non-pharmacological measures.

Population criteria:

- Patient must be aged 18 years or older.

Behaviour disturbances are defined as severe aggression and injuries to self or others where non-pharmacological methods alone have been unsuccessful.

The diagnosis of autism spectrum disorder must be made based on the Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-V) or ICD-10 international classification of mental and behavioural disorders.

risperidone 2 mg tablet, 60

9079W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	2	..	26.29	27.58	^a APO-Risperidone [TX]	^a NOUMED RISPERIDONE [VO]
						^a Ozidal [RA]	^a Rispa [RW]
						^a Risperdal [JC]	^a Risperidone AMNEAL [EF]
						^a Risperidone generichealth [GQ]	^a Risperidone Sandoz [SZ]
						^a Rispernia [ER]	^a Rixadone [AF]

■ RISPERIDONE**Note Shared Care Model:**

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**4246**

Schizophrenia

Authority required (STREAMLINED)**5912**

Bipolar I disorder

Clinical criteria:

- The condition must be refractory to treatment, **AND**
- The treatment must be in combination with lithium or sodium valproate, **AND**
- The treatment must be maintenance therapy.

risperidone 25 mg modified release injection [1 vial] (&) inert substance diluent [2 mL syringe], 1 pack

8780D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*272.34	41.30	Risperdal Consta [JC]

risperidone 50 mg modified release injection [1 vial] (&) inert substance diluent [2 mL syringe], 1 pack

8782F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*431.80	41.30	Risperdal Consta [JC]

risperidone 37.5 mg modified release injection [1 vial] (&) inert substance diluent [2 mL syringe], 1 pack

8781E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*352.50	41.30	Risperdal Consta [JC]

■ RISPERIDONE**Note Shared Care Model:**

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note For item codes 8869T and 1846E, pharmaceutical benefits that have the form tablet 0.5 mg are equivalent for the purposes of substitution.

Authority required (STREAMLINED)**5903**

Schizophrenia

risperidone 500 microgram tablet, 20

1846E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	3	5	..	*19.17	20.46	^a Risperdal [JC]

risperidone 500 microgram tablet, 60

8869T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	19.15	20.44	^a APO-Risperidone [TX]	^a NOUMED RISPERIDONE [VO]
						^a Ozidal [RA]	^a Rispa [RW]
						^a Risperidone AMNEAL [EF]	^a Risperidone Sandoz [SZ]
						^a Rispernia [ER]	^a Rixadone [AF]

■ RISPERIDONE**Note Shared Care Model:**

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note For items 8787L and 1842Y, pharmaceutical benefits that have the form tablet 0.5 mg are equivalent for the purposes of substitution.

Authority required (STREAMLINED)**6898**

Severe behavioural disturbances

Clinical criteria:

- Patient must have autism spectrum disorder, **AND**
- The treatment must be under the supervision of a paediatrician or psychiatrist, **AND**
- The treatment must be in combination with non-pharmacological measures.

Population criteria:

- Patient must be under 18 years of age.

Behaviour disturbances are defined as severe aggression and injuries to self or others where non-pharmacological methods alone have been unsuccessful.

The diagnosis of autism spectrum disorder must be made based on the Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-V) or ICD-10 international classification of mental and behavioural disorders.

Authority required (STREAMLINED)**6899**

Severe behavioural disturbances

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have autism spectrum disorder, **AND**
- Patient must have been commenced on PBS-subsidised treatment with risperidone prior to turning 18 years of age, **AND**
- The treatment must be under the supervision of a paediatrician or psychiatrist, **AND**

- The treatment must be in combination with non-pharmacological measures.

Population criteria:

- Patient must be aged 18 years or older.

Behaviour disturbances are defined as severe aggression and injuries to self or others where non-pharmacological methods alone have been unsuccessful.

The diagnosis of autism spectrum disorder must be made based on the Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-V) or ICD-10 international classification of mental and behavioural disorders.

risperidone 500 microgram tablet, 20

1842Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	3	2	..	*19.17	20.46	^a Risperdal [JC]

risperidone 500 microgram tablet, 60

8787L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	2	..	19.15	20.44	^a APO-Risperidone [TX]	^a NOUMED RISPERIDONE [VO]
						^a Ozidal [RA]	^a Rispa [RW]
						^a Risperidone AMNEAL [EF]	^a Risperidone Sandoz [SZ]
						^a Rispermia [ER]	^a Rixadone [AF]

■ RISPERIDONE

Caution In placebo controlled trials in elderly patients with dementia there was a significantly higher incidence of cerebrovascular adverse events, such as stroke (including fatalities) and transient ischaemic attacks, in patients treated with risperidone compared with patients treated with placebo.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)**10020**

Behavioural disturbances

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be characterised by psychotic symptoms and aggression, **AND**
- Patient must have dementia of the Alzheimer type, **AND**
- Patient must have failed to respond to non-pharmacological methods of treatment, **AND**
- Patient must not receive more than 12 weeks of treatment under this restriction.

A patient may only qualify for 12 weeks of PBS-subsidised treatment under this restriction once in a 12 month period.

risperidone 1 mg tablet, 60

11877D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	2	..	18.26	19.55	^a APO-Risperidone [TX]	^a NOUMED RISPERIDONE [VO]
						^a Ozidal [RA]	^a Rispa [RW]
						^a Risperdal [JC]	^a Risperidone AMNEAL [EF]
						^a Risperidone generichealth [GQ]	^a Risperidone Sandoz [SZ]
						^a Rispermia [ER]	^a Rixadone [AF]

risperidone 1 mg/mL oral liquid, 100 mL

11874Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	±1	2	..	113.55	41.30	^a Risperdal [JC]	^a Rixadone [AF]

■ RISPERIDONE

Caution In placebo controlled trials in elderly patients with dementia there was a significantly higher incidence of cerebrovascular adverse events, such as stroke (including fatalities) and transient ischaemic attacks, in patients treated with risperidone compared with patients treated with placebo.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Behavioural disturbances

Treatment Phase: Continuing treatment, trial of dose reduction or cessation of treatment

Clinical criteria:

- The condition must be characterised by psychotic symptoms and aggression, **AND**

- Patient must have dementia of the Alzheimer type, **AND**
- Patient must have responded to an initial course of treatment with this drug for this condition, **AND**
- Patient must have failed to respond to non-pharmacological methods of treatment, **AND**
- The treatment must be for dose tapering purposes as part of a trial of treatment reduction or cessation; OR
- Patient must have trialled a period of treatment reduction or cessation with this drug for this condition and experienced worsening or re-emergence of symptoms during this trial, and retrials are considered periodically, **AND**
- Patient must be optimised on non-pharmacological methods of treatment.

The patient's response to treatment and a trial of treatment reduction or cessation must be discussed formally with a psychiatrist or geriatrician or in a documented clinical review process involving a least one other medical practitioner, or be reviewed by a psychiatrist or geriatrician.

Response to treatment is defined as a significant reduction in symptoms of psychosis or aggression.

Patients must cease treatment if there is no improvement in symptoms of psychosis and aggression, or worsening of symptoms with therapy.

Patients must be monitored for adverse effects such as falls, drowsiness leading to reduced self-care, incontinence, reduced nutrition, reduced ability to communicate needs/wishes and take part in activities. Therapy must be ceased if harms of therapy outweigh benefits.

Trials of reduction or cessation of therapy should be considered periodically with the intention of maintaining symptom control through non-pharmacological measures wherever possible and/or lowest effective dose therapy.

Evidence of patient benefit from therapy, failure of non-pharmacological approaches to manage symptoms in the absence of therapy, and recurrence of symptoms following reduction or cessation of therapy, trialled on at least 1 occasion, must be documented in the patient's medical records.

risperidone 1 mg tablet, 60

11879F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	2	..	18.26	19.55	^a APO-Risperidone [TX]	^a NOUMED RISPERIDONE [VO]
						^a Ozidal [RA]	^a Rispa [RW]
						^a Risperdal [JC]	^a Risperidone AMNEAL [EF]
						^a Risperidone generichealth [GQ]	^a Risperidone Sandoz [SZ]
						^a Rispernia [ER]	^a Rixadone [AF]

risperidone 1 mg/mL oral liquid, 100 mL

11882J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	±1	2	..	113.55	41.30	^a Risperdal [JC]	^a Rixadone [AF]

■ RISPERIDONE

Caution In placebo controlled trials in elderly patients with dementia there was a significantly higher incidence of cerebrovascular adverse events, such as stroke (including fatalities) and transient ischaemic attacks, in patients treated with risperidone compared with patients treated with placebo.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note Pharmaceutical benefits that have the form pack size risperidone 500 microgram tablet, 20 and pharmaceutical benefits that have the form pack size risperidone 500 microgram tablet, 60 are equivalent for the purposes of substitution.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

10020

Behavioural disturbances

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be characterised by psychotic symptoms and aggression, **AND**
- Patient must have dementia of the Alzheimer type, **AND**
- Patient must have failed to respond to non-pharmacological methods of treatment, **AND**
- Patient must not receive more than 12 weeks of treatment under this restriction.

A patient may only qualify for 12 weeks of PBS-subsidised treatment under this restriction once in a 12 month period.

risperidone 500 microgram tablet, 20

11872W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	3	2	..	*19.17	20.46	^a Risperdal [JC]

risperidone 500 microgram tablet, 60

11869Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	2	..	19.15	20.44	^a APO-Risperidone [TX]	^a NOUMED RISPERIDONE [VO]
						^a Ozidal [RA]	^a Rispa [RW]
						^a Risperidone AMNEAL [EF]	^a Risperidone Sandoz [SZ]
						^a Rispernia [ER]	^a Rixadone [AF]

■ RISPERIDONE

Caution In placebo controlled trials in elderly patients with dementia there was a significantly higher incidence of cerebrovascular adverse events, such as stroke (including fatalities) and transient ischaemic attacks, in patients treated with risperidone compared with patients treated with placebo.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note Pharmaceutical benefits that have the form pack size risperidone 500 microgram tablet, 20 and pharmaceutical benefits that have the form pack size risperidone 500 microgram tablet, 60 are equivalent for the purposes of substitution.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Behavioural disturbances

Treatment Phase: Continuing treatment, trial of dose reduction or cessation of treatment

Clinical criteria:

- The condition must be characterised by psychotic symptoms and aggression, **AND**
- Patient must have dementia of the Alzheimer type, **AND**
- Patient must have responded to an initial course of treatment with this drug for this condition, **AND**
- Patient must have failed to respond to non-pharmacological methods of treatment, **AND**
- The treatment must be for dose tapering purposes as part of a trial of treatment reduction or cessation; OR
- Patient must have trialled a period of treatment reduction or cessation with this drug for this condition and experienced worsening or re-emergence of symptoms during this trial, and retreats are considered periodically, **AND**
- Patient must be optimised on non-pharmacological methods of treatment.

The patient's response to treatment and a trial of treatment reduction or cessation must be discussed formally with a psychiatrist or geriatrician or in a documented clinical review process involving a least one other medical practitioner, or be reviewed by a psychiatrist or geriatrician.

Response to treatment is defined as a significant reduction in symptoms of psychosis or aggression.

Patients must cease treatment if there is no improvement in symptoms of psychosis and aggression, or worsening of symptoms with therapy.

Patients must be monitored for adverse effects such as falls, drowsiness leading to reduced self-care, incontinence, reduced nutrition, reduced ability to communicate needs/wishes and take part in activities. Therapy must be ceased if harms of therapy outweigh benefits.

Trials of reduction or cessation of therapy should be considered periodically with the intention of maintaining symptom control through non-pharmacological measures wherever possible and/or lowest effective dose therapy.

Evidence of patient benefit from therapy, failure of non-pharmacological approaches to manage symptoms in the absence of therapy, and recurrence of symptoms following reduction or cessation of therapy, trialled on at least 1 occasion, must be documented in the patient's medical records.

risperidone 500 microgram tablet, 20

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
11873X	3	2	..	*19.17	20.46	^a Risperdal [JC]

risperidone 500 microgram tablet, 60

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
11881H	1	2	..	19.15	20.44	^a APO-Risperidone [TX]	^a NOUMED RISPERIDONE [VO]
						^a Ozidal [RA]	^a Rispa [RW]
						^a Risperidone AMNEAL [EF]	^a Risperidone Sandoz [SZ]
						^a Rispernia [ER]	^a Rixadone [AF]

ANXIOLYTICS

Benzodiazepine derivatives

■ ALPRAZOLAM

Note The panic disorder must not be attributable to some known organic factor.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Panic disorder

Clinical criteria:

- The treatment must be for use when other treatments have failed; OR
- The treatment must be for use when other treatments are inappropriate.

alprazolam 250 microgram tablet, 10

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
11205R	1	..	..	18.75	20.04	Kalma 0.25 [AF]

alprazolam 500 microgram tablet, 10

11187T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	19.69	20.98	^a Alprax 0.5 [AS]	^a Kalma 0.5 [AF]

alprazolam 1 mg tablet, 10

11186R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	20.67	21.96	^a Alprax 1 [AS]	^a Kalma 1 [AF]

■ DIAZEPAM**diazepam 5 mg tablet, 50**

5072Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
DP	1	..	..	13.72	15.01	^a Antenex 5 [AF] ^a NOUMED DIAZEPAM [VO]	^a APO-Diazepam [TX] ^a Valpam 5 [RW]
			^B 3.07	16.79	15.01	^a Valium [RO]	

diazepam 2 mg tablet, 50

5071X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
DP	1	..	..	13.66	14.95	^a APO-Diazepam [TX]	^a Valpam 2 [RW]
			^B 2.78	16.44	14.95	^a Antenex 2 [AF]	

■ DIAZEPAM**Authority required**

Chronic spasticity

Population criteria:

- Patient must be under 18 years of age.

diazepam 1 mg/mL oral liquid, 100 mL

2669L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	..	..	43.88	41.30	Diazepam Elixir [ON]

■ DIAZEPAM

Note Authorities for increased maximum quantities and/or repeats for the oral forms of diazepam will be granted only for

- (i) the treatment of disabling spasticity; or
- (ii) malignant neoplasia (late stage); or
- (iii) use by patients who are receiving long-term nursing care on account of age, infirmity or other condition in hospitals, nursing homes or residential facilities and who have been demonstrated, within the past six months, to be benzodiazepine dependent by an unsuccessful attempt at gradual withdrawal; or
- (iv) use by a patient who is receiving long-term nursing care and in respect of whom a Carer Allowance is payable as a disabled adult and who has been demonstrated, within the past six months, to be benzodiazepine dependent by an unsuccessful attempt at gradual withdrawal.

Up to six months' treatment (i.e. one month's treatment with five repeats) may be requested.

diazepam 5 mg tablet, 50

3162K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	13.72	15.01	^a Antenex 5 [AF] ^a NOUMED DIAZEPAM [VO]	^a APO-Diazepam [TX] ^a Valpam 5 [RW]
			^B 3.07	16.79	15.01	^a Valium [RO]	

diazepam 2 mg tablet, 50

3161J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	13.66	14.95	^a APO-Diazepam [TX]	^a Valpam 2 [RW]
			^B 2.78	16.44	14.95	^a Antenex 2 [AF]	

■ OXAZEPAM**oxazepam 15 mg tablet, 25**

5192G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	1	..	..	13.55	14.84	Serepax [AS]

oxazepam 30 mg tablet, 25

5193H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
DP	1	..	..	13.23	14.52	^a APO-Oxazepam [TX]	^a Murelax [RW]
			^B 2.20	15.43	14.52	^a Serepax [AS]	

■ OXAZEPAM

Note Authorities for increased maximum quantities and/or repeats will not be granted except as detailed under the 'Authority required' listing of oxazepam.

oxazepam 15 mg tablet, 25

3132W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	13.55	14.84	Serepax [AS]

oxazepam 30 mg tablet, 25

3133X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	13.23	14.52	^a APO-Oxazepam [TX]	^a Murelax [RW]
			^B 2.20	15.43	14.52	^a Serepax [AS]	

■ OXAZEPAM**Authority required**

Malignant neoplasia (late stage)

Authority required

Anxiety

Clinical criteria:

- Patient must be receiving this drug for the management of anxiety, **AND**
- Patient must be receiving long-term nursing care on account of age, infirmity or other condition in a hospital, nursing home or residential facility, **AND**
- Patient must have demonstrated, within the past 6 months, benzodiazepine dependence by an unsuccessful attempt at gradual withdrawal.

Authority required

Anxiety

Clinical criteria:

- Patient must be receiving this drug for the management of anxiety, **AND**
- Patient must be receiving long-term nursing care, **AND**
- Patient must be one in respect of whom a Carer Allowance is payable as a disabled adult, **AND**
- Patient must have demonstrated, within the past 6 months, benzodiazepine dependence by an unsuccessful attempt at gradual withdrawal.

oxazepam 15 mg tablet, 25

3134Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*15.08	16.37	Serepax [AS]

oxazepam 30 mg tablet, 25

3135B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	2	5	..	*14.44	15.73	^a APO-Oxazepam [TX]	^a Murelax [RW]
			^B 4.40	*18.84	15.73	^a Serepax [AS]	

HYPNOTICS AND SEDATIVES*Benzodiazepine derivatives***■ NITRAZEPAM****nitrazepam 5 mg tablet, 25**

5189D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	1	..	..	13.73	15.02	Mogadon [IL]

■ NITRAZEPAM

Note Authorities for increased maximum quantities and/or repeats will not be granted except as detailed under the 'Authority required' listing of nitrazepam.

nitrazepam 5 mg tablet, 25

2723H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	13.73	15.02	Mogadon [IL]

■ NITRAZEPAM**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required

Myoclonic epilepsy

Authority required

Malignant neoplasia (late stage)

Authority required

Insomnia

Clinical criteria:

- Patient must be receiving this drug for the management of insomnia, **AND**

- Patient must be receiving long-term nursing care on account of age, infirmity or other condition in a hospital, nursing home or residential facility, **AND**
- Patient must have demonstrated, within the past 6 months, benzodiazepine dependence by an unsuccessful attempt at gradual withdrawal.

Authority required

Insomnia

Clinical criteria:

- Patient must be receiving this drug for the management of insomnia, **AND**
- Patient must be receiving long-term nursing care, **AND**
- Patient must be one in respect of whom a Carer Allowance is payable as a disabled adult, **AND**
- Patient must have demonstrated, within the past 6 months, benzodiazepine dependence by an unsuccessful attempt at gradual withdrawal.

nitrazepam 5 mg tablet, 25

2732T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*15.44	16.73	Mogadon [IL]

■ **TEMAZEPAM****temazepam 10 mg tablet, 25**

5221T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
DP	1	..	..	13.23	14.52	^a APO-Temazepam [TX]	^a Temaze [AF]
						^a Temtabs [LN]	
			^B 3.35	16.58	14.52	^a Normison [AS]	

■ **TEMAZEPAM**

Note Authorities for increased maximum quantities and/or repeats will not be granted except as detailed under the 'Authority required' listing of temazepam.

temazepam 10 mg tablet, 25

2089Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	13.23	14.52	^a APO-Temazepam [TX]	^a Temaze [AF]
						^a Temtabs [LN]	
			^B 3.35	16.58	14.52	^a Normison [AS]	

■ **TEMAZEPAM****Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required

Malignant neoplasia (late stage)

Authority required

Insomnia

Clinical criteria:

- Patient must be receiving this drug for the management of insomnia, **AND**
- Patient must be receiving long-term nursing care on account of age, infirmity or other condition in a hospital, nursing home or residential facility, **AND**
- Patient must have demonstrated, within the past 6 months, benzodiazepine dependence by an unsuccessful attempt at gradual withdrawal.

Authority required

Insomnia

Clinical criteria:

- Patient must be receiving this drug for the management of insomnia, **AND**
- Patient must be receiving long-term nursing care, **AND**
- Patient must be one in respect of whom a Carer Allowance is payable as a disabled adult, **AND**
- Patient must have demonstrated, within the past 6 months, benzodiazepine dependence by an unsuccessful attempt at gradual withdrawal.

temazepam 10 mg tablet, 25

2088X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	2	5	..	*14.44	15.73	^a APO-Temazepam [TX]	^a Temaze [AF]
						^a Temtabs [LN]	
			^B 6.70	*21.14	15.73	^a Normison [AS]	

■ **PSYCHOANALEPTICS****ANTIDEPRESSANTS***Non-selective monoamine reuptake inhibitors*

■ AMITRIPTYLINE

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

amitriptyline hydrochloride 10 mg tablet, 50

2417F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	2	..	14.76	16.05	^a Amitriptyline Alphapharm 10 [AL]	^a Amitriptyline Lupin [GQ]
			^B 1.82	16.58	16.05	^a APO-Amitriptyline 10 [TX]	^a ENTRIP [RW]
						^a Endep 10 [AF]	

amitriptyline hydrochloride 50 mg tablet, 50

2429W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	2	..	15.31	16.60	^a Amitriptyline Alphapharm 50 [AL]	^a Amitriptyline Lupin [GQ]
			^B 1.82	17.13	16.60	^a APO-Amitriptyline 50 [TX]	^a ENTRIP [RW]
						^a Endep 50 [AF]	

amitriptyline hydrochloride 25 mg tablet, 50

2418G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	2	..	14.94	16.23	^a Amitriptyline Alphapharm 25 [AL]	^a Amitriptyline Lupin [GQ]
			^B 1.82	16.76	16.23	^a APO-Amitriptyline 25 [TX]	^a ENTRIP [RW]
						^a Endep 25 [AF]	

■ CLOMIPRAMINE

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Cataplexy

Clinical criteria:

- The condition must be associated with narcolepsy.

Restricted benefit

Obsessive-compulsive disorder

Restricted benefit

Phobic disorders

Population criteria:

- Patient must be an adult.

clomipramine hydrochloride 25 mg tablet, 50

1561E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	2	..	17.96	19.25	^a APO-Clomipramine [TX]	^a GenRx Clomipramine [GX]
			^B 4.41	22.37	19.25	^a Placil [AF]	
						^a Anafranil 25 [SZ]	

■ DOSULEPIN (DOTHIEPIN)

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

dosulepin (dothiepin) hydrochloride 25 mg capsule, 50

1357K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	14.82	16.11	^a Dosulepin Mylan [AL]
			^B 1.81	16.63	16.11	^a Dothep 25 [AF]

dosulepin (dothiepin) hydrochloride 75 mg tablet, 30

1358L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	14.82	16.11	^a Dosulepin Mylan [AL]
			^B 1.81	16.63	16.11	^a Dothep 75 [AF]

■ DOXEPIN

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

NERVOUS SYSTEM

doxepin 25 mg capsule, 50

1013H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	16.29	17.58	Deptran 25 [AF]

doxepin 50 mg tablet, 50

1012G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	15.45	16.74	Deptran 50 [AF]

doxepin 10 mg capsule, 50

1011F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	15.41	16.70	Deptran 10 [AF]

■ IMIPRAMINE

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

imipramine hydrochloride 10 mg tablet, 50

2420J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	16.40	17.69	Tofranil 10 [GH]

■ IMIPRAMINE

Note Pharmaceutical benefits that have the form imipramine hydrochloride 25 mg tablet in a pack size of 50 can be substituted for a pack size of 100 in the case of a shortage.

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

imipramine hydrochloride 25 mg tablet, 100

12113M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	0.5	2	..	*26.58	27.87	^a Imipramine (Leading) [QY]

imipramine hydrochloride 25 mg tablet, 50

2421K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	19.66	20.95	^a Tofranil 25 [GH]

■ NORTRIPTYLINE

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Major depression

Clinical criteria:

- The treatment must be for use when other anti-depressant therapy has failed.

Restricted benefit

Major depression

Clinical criteria:

- The treatment must be for use when other anti-depressant therapy is contraindicated.

nortriptyline 10 mg tablet, 50

2522R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	2	..	15.98	17.27	^a Allegron [RW]	^a NortriTABS 10 mg [GH]

nortriptyline 25 mg tablet, 50

2523T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	2	..	16.89	18.18	^a Allegron [RW]	^a NortriTABS 25 mg [GH]

Selective serotonin reuptake inhibitors

■ CITALOPRAM

Restricted benefit

Major depressive disorders

citalopram 20 mg tablet, 28

8220P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	13.64	14.93	^a APO-Citalopram [TX]	^a Celapram [AF]

NP				^a Citalopram Actavis [ED]	^a Citalopram AN [EA]
				^a Citalopram Sandoz [SZ]	^a NOUMED CITALOPRAM [VO]
				^a Talam [RW]	
				^a Cipramil [LU]	
	^B 10.37	24.01	14.93		

citalopram 40 mg tablet, 28

8703C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	14.47	15.76	^a APO-Citalopram [TX]	^a Celapram [AF]
						^a Citalopram Actavis [ED]	^a Citalopram AN [EA]
						^a Citalopram Sandoz [SZ]	^a NOUMED CITALOPRAM [VO]
						^a Talam [RW]	

citalopram 10 mg tablet, 28

8702B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	13.33	14.62	^a Celapram [AF]	^a Citalopram Actavis [EA]
						^a Citalopram AN [EF]	^a Talam [RW]

■ ESCITALOPRAM**Restricted benefit**

Major depressive disorders

escitalopram 20 mg tablet, 28

8701Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	14.40	15.69	^a APO-Escitalopram [TX]	^a Blooms the Chemist Escitalopram [IB]
						^a Cilopam-S [ER]	^a Escitalopram AN [EA]
						^a Escitalopram GH [HQ]	^a Escitalopram Sandoz [HX]
						^a Esipram [CF]	^a Lexam 20 [RW]
						^a LoxaLate [AF]	
				^B 11.80	26.20	15.69	^a Lexapro [LU]

escitalopram 10 mg tablet, 28

8700X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	14.38	15.67	^a APO-Escitalopram [TX]	^a Blooms the Chemist Escitalopram [IB]
						^a Cilopam-S [ER]	^a Escitalopram AN [EA]
						^a Escitalopram GH [HQ]	^a Escitalopram Sandoz [HX]
						^a Esipram [CF]	^a Lexam 10 [RW]
						^a LoxaLate [AF]	
				^B 11.46	25.84	15.67	^a Lexapro [LU]

■ ESCITALOPRAM**Restricted benefit**

Moderate to severe generalised anxiety disorder (GAD)

Clinical criteria:

- The condition must be defined by Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-V) criteria, **AND**
- Patient must not have responded to non-pharmacological therapy, **AND**
- Patient must be one for whom a GP Mental Health Care Plan, as described under items 2715 or 2717 of the Medicare Benefits Schedule, has been prepared.

Restricted benefit

Moderate to severe generalised anxiety disorder (GAD)

Clinical criteria:

- The condition must be defined by Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-V) criteria, **AND**
- Patient must not have responded to non-pharmacological therapy, **AND**
- Patient must have been assessed by a psychiatrist.

Restricted benefit

Moderate to severe social anxiety disorder (social phobia, SAD)

Clinical criteria:

- The condition must be defined by Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-V) criteria, **AND**
- Patient must not have responded to non-pharmacological therapy, **AND**
- Patient must be one for whom a GP Mental Health Care Plan, as described under items 2715 or 2717 of the Medicare Benefits Schedule, has been prepared.

Restricted benefit

Moderate to severe social anxiety disorder (social phobia, SAD)

Clinical criteria:

- The condition must be defined by Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-V) criteria, **AND**

- Patient must not have responded to non-pharmacological therapy, **AND**
- Patient must have been assessed by a psychiatrist.

escitalopram 20 mg tablet, 28

9433L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	14.40	15.69	^a Escitalopram GH [HQ]	^a Esipram [CF]
			^B 11.80	26.20	15.69	^a Lexapro [LU]	

escitalopram 10 mg tablet, 28

9432K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	14.38	15.67	^a Escitalopram GH [HQ]	^a Esipram [CF]
			^B 11.46	25.84	15.67	^a Lexapro [LU]	

■ ESCITALOPRAM**Restricted benefit**

Major depressive disorders

Restricted benefit

Moderate to severe generalised anxiety disorder (GAD)

Clinical criteria:

- The condition must be defined by Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-V) criteria, **AND**
- Patient must not have responded to non-pharmacological therapy, **AND**
- Patient must be one for whom a GP Mental Health Care Plan, as described under items 2715 or 2717 of the Medicare Benefits Schedule, has been prepared.

Restricted benefit

Moderate to severe generalised anxiety disorder (GAD)

Clinical criteria:

- The condition must be defined by Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-V) criteria, **AND**
- Patient must not have responded to non-pharmacological therapy, **AND**
- Patient must have been assessed by a psychiatrist.

Restricted benefit

Moderate to severe social anxiety disorder (social phobia, SAD)

Clinical criteria:

- The condition must be defined by Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-V) criteria, **AND**
- Patient must not have responded to non-pharmacological therapy, **AND**
- Patient must be one for whom a GP Mental Health Care Plan, as described under items 2715 or 2717 of the Medicare Benefits Schedule, has been prepared.

Restricted benefit

Moderate to severe social anxiety disorder (social phobia, SAD)

Clinical criteria:

- The condition must be defined by Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-V) criteria, **AND**
- Patient must not have responded to non-pharmacological therapy, **AND**
- Patient must have been assessed by a psychiatrist.

escitalopram 20 mg/mL oral liquid, 15 mL

10181W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	37.99	39.28	Lexapro [LU]

■ FLUOXETINE**Restricted benefit**

Major depressive disorders

Restricted benefit

Obsessive-compulsive disorder

fluoxetine 20 mg tablet, 28

8270G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	15.88	17.17	^a Lovan 20 Tab [AL]	^a Zactin Tablet [AF]

■ FLUOXETINE

Note Pharmaceutical benefits that have the form fluoxetine 20 mg capsule are equivalent for the purposes of substitution.

Restricted benefit

Major depressive disorders

Restricted benefit

Obsessive-compulsive disorder

fluoxetine 20 mg capsule, 28

1434L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	15.88	17.17	^a APO-Fluoxetine [TX]	^a Blooms the Chemist Fluoxetine [IB]
						^a BTC Fluoxetine [JB]	^a FLUOTEX [RF]
						^a Fluoxetine AN [EA]	^a Fluoxetine APOTEX [TY]
						^a Fluoxetine-GA [ED]	^a Fluoxetine generichealth [GQ]
						^a Fluoxetine Sandoz [SZ]	^a Lovan [AL]
						^a Zactin [AF]	
			^B 1.10	16.98	17.17	^a Prozac 20 [LY]	

fluoxetine 20 mg capsule, 100

12256C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	0.3	5	..	*24.01	25.30	^a Fluoxetine Capsule (USP) [DZ]

FLUVOXAMINE**Restricted benefit**

Major depressive disorders

Restricted benefit

Obsessive-compulsive disorder

fluvoxamine maleate 100 mg tablet, 30

8174F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	20.59	21.88	^a APO-Fluvoxamine [TX]	^a Faverin 100 [RW]
						^a Fluvoxamine AN [ED]	^a Fluvoxamine GA [EA]
						^a Movox 100 [AF]	
			^B 3.50	24.09	21.88	^a Luvox [GO]	

fluvoxamine maleate 50 mg tablet, 30

8512B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	17.69	18.98	^a APO-Fluvoxamine [TX]	^a Faverin 50 [RW]
						^a Fluvoxamine AN [ED]	^a Fluvoxamine GA [EA]
						^a Movox 50 [AL]	
			^B 3.49	21.18	18.98	^a Luvox [GO]	

PAROXETINE**Restricted benefit**

Major depressive disorders

Restricted benefit

Obsessive-compulsive disorder

Restricted benefit

Panic disorder

paroxetine 20 mg tablet, 30

2242B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	15.61	16.90	^a APO-Paroxetine [TX]	^a Extine 20 [RW]
						^a Paroxetine AN [EA]	^a Paroxetine GH [GQ]
						^a Paroxetine Sandoz [SZ]	^a Paxtine [AF]
						^a Roxet 20 [DO]	
			^B 2.41	18.02	16.90	^a Aropax [AS]	

SERTRALINE**Note** Pharmaceutical benefits that have the form sertraline 50 mg tablet are equivalent for the purposes of substitution.**Restricted benefit**

Major depressive disorders

sertraline 50 mg tablet, 30

12305P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	30.91	32.20	^a Sertraline tablets USP (Medsurge) [DZ]

sertraline 50 mg tablet, 30

2236Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	14.00	15.29	^a APO-Sertraline [TX]	^a Auro-Sertraline 50 [DO]
						^a Eleva 50 [AF]	^a NOUMED SERTRALINE [VO]
						^a Sertra 50 [RW]	^a Sertraline AN [EA]
						^a Sertraline generichealth [GQ]	^a Sertraline Sandoz [SZ]
						^a Setrona [RA]	
			^B 5.51	19.51	15.29	^a Zoloft [UJ]	

■ **SERTRALINE**

Note Pharmaceutical benefits that have the form sertraline 100 mg tablet are equivalent for the purposes of substitution.

Restricted benefit

Major depressive disorders

sertraline 100 mg tablet, 30

12317G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	30.91	32.20	^a Sertraline tablets USP (Medsurge) [DZ]

sertraline 100 mg tablet, 30

2237R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	14.00	15.29	^a APO-Sertraline [TX] ^a Eleva 100 [AF] ^a Sertra 100 [RW] ^a Sertraline generichealth [GQ] ^a Setrona [RA]	^a Auro-Sertraline 100 [DO] ^a NOUMED SERTRALINE [VO] ^a Sertraline AN [EA] ^a Sertraline Sandoz [SZ] ^a Zolofit [UJ]

■ **SERTRALINE**

Note Pharmaceutical benefits that have the form sertraline 100 mg tablet are equivalent for the purposes of substitution.

Restricted benefit

Obsessive-compulsive disorder

Restricted benefit

Panic disorder

Clinical criteria:

- The treatment must be for use when other treatments have failed; OR
- The treatment must be for use when other treatments are inappropriate.

sertraline 100 mg tablet, 30

12296E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	30.91	32.20	^a Sertraline tablets USP (Medsurge) [DZ]

sertraline 100 mg tablet, 30

8837D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	14.00	15.29	^a Auro-Sertraline 100 [DO] ^a NOUMED SERTRALINE [VO] ^a Zolofit [UJ]	^a Eleva 100 [AF] ^a Sertraline AN [EA]

■ **SERTRALINE**

Note Pharmaceutical benefits that have the form sertraline 50 mg tablet are equivalent for the purposes of substitution.

Restricted benefit

Obsessive-compulsive disorder

Restricted benefit

Panic disorder

Clinical criteria:

- The treatment must be for use when other treatments have failed; OR
- The treatment must be for use when other treatments are inappropriate.

sertraline 50 mg tablet, 30

12299H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	30.91	32.20	^a Sertraline tablets USP (Medsurge) [DZ]

sertraline 50 mg tablet, 30

8836C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	14.00	15.29	^a Auro-Sertraline 50 [DO] ^a NOUMED SERTRALINE [VO]	^a Eleva 50 [AF] ^a Sertraline AN [EA]
			^B 5.51	19.51	15.29	^a Zolofit [UJ]	

Monoamine oxidase inhibitors, non-selective■ **PHENELZINE**

Caution This drug is an irreversible monoamine oxidase inhibitor.

Note Pharmaceutical benefits that have the form phenelzine 15 mg tablet, 60 and pharmaceutical benefits that have the form phenelzine 15 mg tablet, 100 are equivalent for the purposes of substitution.

Restricted benefit

Depression

Clinical criteria:

- The treatment must be for when all other anti-depressant therapy has failed; OR
- The treatment must be for when all other anti-depressant therapy is inappropriate.

phenelzine 15 mg tablet, 60

12239E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	
	1.7	1	..	*175.18	41.30	^a Phenelzine sulfate USP (Generic Health) [GQ]	

phenelzine 15 mg tablet, 100

2856H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	
	1	1	..	81.19	41.30	^a Nardil [LM]	

■ TRANYLCPROMINE

Caution This drug is an irreversible monoamine oxidase inhibitor.

tranylcypromine 10 mg tablet, 50

2444P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	
	1	2	..	50.34	41.30	Parnate [GH]	

*Monoamine oxidase A inhibitors***■ MOCLOBEMIDE****Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Major depressive disorders

moclobemide 150 mg tablet, 60

1900B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	
NP	1	5	..	21.76	23.05	^a Amira 150 [AF]	^a APO-Moclobemide [TX]
						^a Clobemix [ED]	^a Moclobemide AN [EA]
						^a Moclobemide Sandoz [SZ]	^a Mohexal [HX]
						^b 3.00	^a Aurorix [GO]

moclobemide 300 mg tablet, 60

8003F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	
NP	1	5	..	28.29	29.58	^a Amira 300 [AF]	^a APO-Moclobemide [TX]
						^a Clobemix [ED]	^a Moclobemide AN [EA]
						^a Moclobemide Sandoz [SZ]	
						^b 3.00	^a Aurorix 300 mg [GO]

*Other antidepressants***■ DESVENLAFAXINE**

Note Pharmaceutical benefits that have the forms desvenlafaxine tablet (modified release) 100 mg, desvenlafaxine tablet (modified release) 100 mg (as benzoate) and desvenlafaxine tablet (extended release) 100 mg (as succinate) are equivalent for the purposes of substitution.

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Major depressive disorders

desvenlafaxine 100 mg modified release tablet, 28

10231L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	
NP	1	5	..	24.51	25.80	^a Desfax [AF]	^a DESVEN [RW]
						^a Desvenlafaxine Actavis [EA]	^a Desvenlafaxine Sandoz [SZ]

desvenlafaxine 100 mg modified release tablet, 28

10245F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	
NP	1	5	..	24.51	25.80	^a APO-Desvenlafaxine MR [TX]	^a Desvenlafaxine GH XR [GQ]

desvenlafaxine 100 mg modified release tablet, 28

9367B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	
NP	1	5	..	24.51	25.80	^a Pristiq [PF]	

■ DESVENLAFAXINE

Note Pharmaceutical benefits that have the forms desvenlafaxine tablet (modified release) 50 mg, desvenlafaxine tablet (modified release) 50 mg (as benzoate) and desvenlafaxine tablet (extended release) 50 mg (as succinate) are equivalent for the purposes of substitution.

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a

patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Major depressive disorders

desvenlafaxine 50 mg modified release tablet, 28

10234P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	22.19	23.48	^a APO-Desvenlafaxine MR [TX]	^a Desvenlafaxine GH XR [GQ]

desvenlafaxine 50 mg modified release tablet, 28

10241B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	22.19	23.48	^a Desfax [AF] ^a Desvenlafaxine Actavis [EA]	^a DESVEN [RW] ^a Desvenlafaxine Sandoz [SZ]

desvenlafaxine 50 mg modified release tablet, 28

9366Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	22.19	23.48	^a Pristiq [PF]

■ DULOXETINE**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Major depressive disorders

duloxetine 30 mg enteric capsule, 28

9155W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	16.29	17.58	^a Andeptra [EL] ^a Duloxecor [CR] ^a Duloxetine Sandoz [HX] ^a DYTREX 30 [RW]	^a APO-Duloxetine [TX] ^a Duloxetine AN [EA] ^a Duloxetine Sandoz 30 [SZ] ^a Tixel [AL]
			^B 7.88	24.17	17.58	^a Cymbalta [LY]	

duloxetine 60 mg enteric capsule, 28

9156X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	17.94	19.23	^a Andeptra [EL] ^a Duloxecor [CR] ^a Duloxetine Sandoz [HX] ^a DYTREX 60 [RW]	^a APO-Duloxetine [TX] ^a Duloxetine AN [EA] ^a Duloxetine Sandoz 60 [SZ] ^a Tixel [AL]
			^B 8.00	25.94	19.23	^a Cymbalta [LY]	

■ LITHIUM CARBONATE**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

lithium carbonate 250 mg tablet, 200

3059B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	24.12	25.41	Lithicarb [AS]

lithium carbonate 450 mg modified release tablet, 100

8290H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	2	..	*36.50	37.79	Quilonum SR [AS]

■ MIANSERIN

Caution Neutropenia and agranulocytosis are more frequent in the elderly, especially in the early months of therapy.

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Severe depression

mianserin hydrochloride 10 mg tablet, 50

1627P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	19.82	21.11	Lumin 10 [AF]

mianserin hydrochloride 20 mg tablet, 50

1628Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	28.47	29.76	Lumin 20 [AF]

■ MIRTAZAPINE**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Major depressive disorders

mirtazapine 45 mg orally disintegrating tablet, 30

8857E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	20.32	21.61	^a Milivin OD 45 [DO] ^a Mirtazapine AN ODT [EA]	^a MIRTANZA ODT [RF] ^a Mirtazapine Sandoz ODT 45 [SZ]

mirtazapine 15 mg orally disintegrating tablet, 30

8855C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	16.28	17.57	^a Milivin OD 15 [DO] ^a Mirtazapine AN ODT [EA]	^a MIRTANZA ODT [RF] ^a Mirtazapine Sandoz ODT 15 [SZ]

mirtazapine 45 mg tablet, 30

8883M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	18.02	19.31	^a APO-Mirtazapine [TX] ^a MIRTANZA [RF] ^a Mirtazapine GH [GQ] ^a Mirtazon [RW]	^a Axit 45 [AF] ^a Mirtazapine AN [EA] ^a Mirtazapine Sandoz [SZ]

mirtazapine 30 mg tablet, 30

8513C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	15.78	17.07	^a APO-Mirtazapine [TX] ^a MIRTANZA [RF] ^a Mirtazapine Sandoz [SZ]	^a Axit 30 [AF] ^a Mirtazapine AN [EA] ^a Mirtazon [RW]
			^B 4.84	20.62	17.07	^a Avanza [AL]	

mirtazapine 15 mg tablet, 30

9365X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	14.66	15.95	^a APO-Mirtazapine [TX] ^a MIRTANZA [RF]	^a Axit 15 [AF] ^a Mirtazapine AN [EA]

mirtazapine 30 mg orally disintegrating tablet, 30

8856D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	17.56	18.85	^a Milivin OD 30 [DO] ^a Mirtazapine AN ODT [EA]	^a MIRTANZA ODT [RF] ^a Mirtazapine Sandoz ODT 30 [SZ]

■ REBOXETINE**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Major depressive disorders

reboxetine 4 mg tablet, 60

8583R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	38.73	40.02	Edronax [PF]

■ VENLAFAXINE**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Major depressive disorders

venlafaxine 37.5 mg modified release capsule, 28

8868R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	14.65	15.94	^a Efexor-XR [UJ]	^a Elaxine SR 37.5 [ZP]

^a Venlafaxine AN SR [EA]**NP Venlafaxine 150 mg modified release capsule, 28**

8302Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	16.51	17.80	^a APO-Venlafaxine XR [TX] ^a Efexor-XR [UJ] ^a Enlafax-XR [AF] ^a Venlafaxine AN SR [EA]	^a Blooms the Chemist Venlafaxine XR [IB] ^a Elaxine SR 150 [ZP] ^a Sandoz Venlafaxine XR [HX] ^a Venlafaxine generichealth XR [GQ]

venlafaxine 75 mg modified release capsule, 28

8301X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	15.78	17.07	^a APO-Venlafaxine XR [TX] ^a Efexor-XR [UJ] ^a Enlafax-XR [AF] ^a Venlafaxine AN SR [EA]	^a Blooms the Chemist Venlafaxine XR [IB] ^a Elaxine SR 75 [ZP] ^a Sandoz Venlafaxine XR [HX] ^a Venlafaxine generichealth XR [GQ]

PSYCHOSTIMULANTS, AGENTS USED FOR ADHD AND NOOTROPICS*Centrally acting sympathomimetics***■ ARMODAFINIL**

Note This drug is not PBS-subsidised when used in combination with PBS-subsidised dexamfetamine sulfate or modafinil.

Authority required

Narcolepsy

Treatment Phase: Initial 1 - treatment of narcolepsy without cataplexy

Treatment criteria:

- Must be treated by a qualified sleep medicine practitioner or neurologist.

Clinical criteria:

- The treatment must be for use when therapy with dexamfetamine sulfate poses an unacceptable medical risk; OR
- The treatment must be for use when intolerance to dexamfetamine sulfate is of a severity to necessitate treatment withdrawal, **AND**
- Patient must have experienced excessive daytime sleepiness, recurrent naps or lapses into sleep occurring almost daily for at least 3 months, **AND**
- Patient must have a mean sleep latency less than or equal to 10 minutes on a Multiple Sleep Latency Test (MSLT); OR
- Patient must have an electroencephalographic (EEG) recording showing the pathologically rapid development of REM sleep, **AND**
- Patient must not have any medical or psychiatric disorder that could otherwise account for the hypersomnia.

The presence of any one of the following indicates treatment with dexamfetamine sulfate poses an unacceptable medical risk:

- a psychiatric disorder;
 - a cardiovascular disorder;
 - a history of substance abuse;
 - glaucoma;
 - any other absolute contraindication to dexamfetamine sulfate as specified in the TGA-approved Product Information.
- The MSLT must be preceded by nocturnal polysomnography. Sleep prior to the MSLT must be at least 6 hours in duration. The authority application must be made in writing and must include the following:
- a completed authority prescription form; and
 - a completed Narcolepsy Initial PBS authority application and Supporting information form; and
 - details of the contraindication or intolerance to dexamfetamine sulfate; and
 - either:
 - the result and date of the polysomnography test and Multiple Sleep Latency Test (MSLT) conducted by, or under the supervision of, a qualified sleep medicine practitioner; or
 - the result and date of the electroencephalograph (EEG), conducted by, or under the supervision of, a neurologist.

The polysomnography, MSLT or EEG test reports must be provided with the authority application.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos. Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Narcolepsy

Treatment Phase: Initial 2 - treatment of narcolepsy with cataplexy

Treatment criteria:

- Must be treated by a qualified sleep medicine practitioner or neurologist.

Clinical criteria:

- The treatment must be for use when therapy with dexamfetamine sulfate poses an unacceptable medical risk; OR
 - The treatment must be for use when intolerance to dexamfetamine sulfate is of a severity to necessitate treatment withdrawal, **AND**
 - Patient must have experienced excessive daytime sleepiness, recurrent naps or lapses into sleep occurring almost daily for at least 3 months, **AND**
 - Patient must have a definite history of cataplexy documented in their medical records for auditing purposes, **AND**
 - Patient must not have any medical or psychiatric disorder that could otherwise account for the hypersomnia.
- The presence of any one of the following indicates treatment with dexamfetamine sulfate poses an unacceptable medical risk:

- (a) a psychiatric disorder;
- (b) a cardiovascular disorder;
- (c) a history of substance abuse;
- (d) glaucoma;
- (e) any other absolute contraindication to dexamfetamine sulfate as specified in the TGA-approved Product Information.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required**Narcolepsy**

Treatment Phase: Continuing or change of treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition; OR
- Patient must have previously received PBS-subsidised treatment with modafinil for this condition.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

armodafinil 150 mg tablet, 30

10912H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	156.38	41.30	Nuvigil [TB]

armodafinil 50 mg tablet, 30

10922W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*106.86	41.30	Nuvigil [TB]

armodafinil 250 mg tablet, 30

10919Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	255.95	41.30	Nuvigil [TB]

■ ATOMOXETINE

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)**7876****Attention deficit hyperactivity disorder**

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a paediatrician or psychiatrist.

Clinical criteria:

- The condition must be or have been diagnosed according to the DSM-5 criteria, **AND**
- Patient must have a contraindication to dexamfetamine, methylphenidate or lisdexamfetamine as specified in TGA-approved product information; OR
- Patient must have a comorbid mood disorder that has developed or worsened as a result of dexamfetamine, methylphenidate or lisdexamfetamine treatment and is of a severity necessitating treatment withdrawal; OR
- Patient must be at an unacceptable medical risk of a severity necessitating permanent stimulant treatment withdrawal if given a stimulant treatment with another agent; OR
- Patient must have experienced adverse reactions of a severity necessitating permanent treatment withdrawal following treatment with dexamfetamine, methylphenidate and lisdexamfetamine (not simultaneously).

Population criteria:

- Patient must be or have been diagnosed between the ages of 6 and 18 years inclusive.

Authority required (STREAMLINED)**7890**

Attention deficit hyperactivity disorder
Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition.

atomoxetine 40 mg capsule, 28

9095Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	5	..	*89.78	41.30	^a APO-Atomoxetine [TX] ^a Atomoxetine Amneal [EA] ^a Strattera [LY]	^a ATOMERRA [RW] ^a Atomoxetine Sandoz [SZ]

atomoxetine 10 mg capsule, 28

9092M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	5	..	*89.78	41.30	^a APO-Atomoxetine [TX] ^a Atomoxetine Amneal [EA] ^a Strattera [LY]	^a ATOMERRA [RW] ^a Atomoxetine Sandoz [SZ]

atomoxetine 100 mg capsule, 28

9290Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	62.57	41.30	^a APO-Atomoxetine [TX] ^a Atomoxetine Amneal [EA] ^a Strattera [LY]	^a ATOMERRA [RW] ^a Atomoxetine Sandoz [SZ]

atomoxetine 80 mg capsule, 28

9289X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	62.57	41.30	^a APO-Atomoxetine [TX] ^a Atomoxetine Amneal [EA] ^a Strattera [LY]	^a ATOMERRA [RW] ^a Atomoxetine Sandoz [SZ]

atomoxetine 25 mg capsule, 28

9094P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	5	..	*89.78	41.30	^a APO-Atomoxetine [TX] ^a Atomoxetine Amneal [EA] ^a Strattera [LY]	^a ATOMERRA [RW] ^a Atomoxetine Sandoz [SZ]

atomoxetine 18 mg capsule, 28

9093N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	5	..	*89.78	41.30	^a APO-Atomoxetine [TX] ^a Atomoxetine Amneal [EA] ^a Strattera [LY]	^a ATOMERRA [RW] ^a Atomoxetine Sandoz [SZ]

atomoxetine 60 mg capsule, 28

9096R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	5	..	*89.78	41.30	^a APO-Atomoxetine [TX] ^a Atomoxetine Amneal [EA] ^a Strattera [LY]	^a ATOMERRA [RW] ^a Atomoxetine Sandoz [SZ]

■ DEXAMFETAMINE

Note Care must be taken to comply with the provisions of State/Territory law when prescribing this drug.

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required

Attention deficit hyperactivity disorder

Treatment must be in accordance with the law of the relevant State or Territory.

Authority required

Narcolepsy

dexamfetamine sulfate 5 mg tablet, 100

1165H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	23.22	24.51	Aspen Pharma Pty Ltd [AS]

■ LISDEXAMFETAMINE**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note In accordance with the Therapeutic Goods Administration (TGA)-approved Product Information, this PBS listing currently intends for once daily dosing only. Divided dosing is not intended (e.g. 20 mg in the mornings, 30 mg in the evenings). Where applications (either on the same day or on separate days) for multiple strengths are sought, repeats should only be sought for the listed target strength.

Note Care must be taken to comply with the provisions of State/Territory law when prescribing this drug.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Authority required

Attention deficit hyperactivity disorder

Clinical criteria:

- Patient must require continuous coverage over 12 hours, **AND**
- The treatment must not exceed a maximum daily dose of 70 mg with this drug.

Population criteria:

- Patient must be aged between the ages of 6 and 18 years inclusive; OR
- Patient must have had a diagnosis of ADHD prior to turning 18 years of age if PBS-subsidised treatment is continuing beyond 18 years of age; OR
- Patient must have a retrospective diagnosis of ADHD if PBS-subsidised treatment is commencing after turning 18 years of age; OR
- Patient must have had a retrospective diagnosis of ADHD if PBS-subsidised treatment is continuing in a patient who commenced PBS-subsidised treatment after turning 18 years of age.

A retrospective diagnosis of ADHD for the purposes of administering this restriction is:

(i) the presence of pre-existing childhood symptoms of ADHD (onset during the developmental period, typically early to mid-childhood); and

(ii) documentation in the patient's medical records that an in-depth clinical interview with, or, obtainment of evidence from, either a: (a) parent, (b) teacher, (c) sibling, (d) third party, has occurred and which supports point (i) above.

lisdexamfetamine dimesilate 70 mg capsule, 30

10492F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	98.64	41.30	Vyvanse [TK]

lisdexamfetamine dimesilate 30 mg capsule, 30

10486X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	98.64	41.30	Vyvanse [TK]

lisdexamfetamine dimesilate 20 mg capsule, 30

11884L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	98.64	41.30	Vyvanse [TK]

lisdexamfetamine dimesilate 60 mg capsule, 30

11897E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	98.64	41.30	Vyvanse [TK]

lisdexamfetamine dimesilate 40 mg capsule, 30

11898F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	98.64	41.30	Vyvanse [TK]

lisdexamfetamine dimesilate 50 mg capsule, 30

10474G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	98.64	41.30	Vyvanse [TK]

■ METHYLPHENIDATE

Note Care must be taken to comply with the provisions of State/Territory law when prescribing this drug.

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required

Attention deficit hyperactivity disorder

Treatment must be in accordance with the law of the relevant State or Territory.

methylphenidate hydrochloride 10 mg tablet, 100

8839F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	25.96	27.25	^a Artige [NM]
			^B 3.76	29.72	27.25	^a Ritalin 10 [NV]

■ METHYLPHENIDATE

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note In accordance with the Therapeutic Goods Administration (TGA)-approved Product Information, this PBS listing currently intends for once daily dosing only. Divided dosing is not intended (e.g. 18 mg in the mornings, 36 mg in the evenings).

Note Care must be taken to comply with the provisions of State/Territory law when prescribing this drug.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Attention deficit hyperactivity disorder

Population criteria:

- Patient must be or have been diagnosed between the ages of 6 and 18 years inclusive.

Clinical criteria:

- Patient must have demonstrated a response to immediate-release methylphenidate hydrochloride with no emergence of serious adverse events, **AND**
- Patient must require continuous coverage over 12 hours, **AND**
- The treatment must not exceed a maximum daily dose of 72 mg with this drug.

methylphenidate hydrochloride 54 mg modified release tablet, 30

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
2432B	1	5	..	71.93	41.30	Concerta [JC]

NP

methylphenidate hydrochloride 36 mg modified release tablet, 30

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
2388Q	1	5	..	62.78	41.30	Concerta [JC]

NP

methylphenidate hydrochloride 27 mg modified release tablet, 30

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
2172H	1	5	..	58.65	41.30	Concerta [JC]

NP

methylphenidate hydrochloride 18 mg modified release tablet, 30

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
2387P	1	5	..	54.51	41.30	Concerta [JC]

NP

■ **METHYLPHENIDATE**

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note In accordance with the Therapeutic Goods Administration (TGA)-approved Product Information, this PBS listing currently intends for once daily dosing only. Divided dosing is not intended (e.g. 20 mg in the mornings, 30 mg in the evenings).

Note Care must be taken to comply with the provisions of State/Territory law when prescribing this drug.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Authority required

Attention deficit hyperactivity disorder

Population criteria:

- Patient must be or have been diagnosed between the ages of 6 and 18 years inclusive.

Clinical criteria:

- Patient must have demonstrated a response to immediate-release methylphenidate hydrochloride with no emergence of serious adverse events, **AND**
- Patient must require continuous coverage over 8 hours, **AND**
- The treatment must not exceed a maximum daily dose of 80 mg with this drug.

methylphenidate hydrochloride 60 mg modified release capsule, 30

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
12116Q	1	5	..	71.67	41.30	Ritalin LA [NV]

NP

methylphenidate hydrochloride 20 mg modified release capsule, 30

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
2276T	1	5	..	47.76	41.30	Ritalin LA [NV]

NP

methylphenidate hydrochloride 30 mg modified release capsule, 30

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
2280B	1	5	..	55.22	41.30	Ritalin LA [NV]

NP

methylphenidate hydrochloride 40 mg modified release capsule, 30

2283E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	57.75	41.30	Ritalin LA [NV]

methylphenidate hydrochloride 10 mg modified release capsule, 30

3440C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	38.49	39.78	Ritalin LA [NV]

■ MODAFINIL

Note This drug is not PBS-subsidised when used in combination with PBS-subsidised dexamfetamine sulfate or armodafinil.

Authority required

Narcolepsy

Treatment Phase: Initial 1 - treatment of narcolepsy without cataplexy

Treatment criteria:

- Must be treated by a qualified sleep medicine practitioner or neurologist.

Clinical criteria:

- The treatment must be for use when therapy with dexamfetamine sulfate poses an unacceptable medical risk; OR
- The treatment must be for use when intolerance to dexamfetamine sulfate is of a severity to necessitate treatment withdrawal, **AND**
- Patient must have experienced excessive daytime sleepiness, recurrent naps or lapses into sleep occurring almost daily for at least 3 months, **AND**
- Patient must have a mean sleep latency less than or equal to 10 minutes on a Multiple Sleep Latency Test (MSLT); OR
- Patient must have an electroencephalographic (EEG) recording showing the pathologically rapid development of REM sleep, **AND**
- Patient must not have any medical or psychiatric disorder that could otherwise account for the hypersomnia.

The presence of any one of the following indicates treatment with dexamfetamine sulfate poses an unacceptable medical risk:

- a psychiatric disorder;
 - a cardiovascular disorder;
 - a history of substance abuse;
 - glaucoma;
 - any other absolute contraindication to dexamfetamine sulfate as specified in the TGA-approved Product Information.
- The MSLT must be preceded by nocturnal polysomnography. Sleep prior to the MSLT must be at least 6 hours in duration. The authority application must be made in writing and must include the following:
- a completed authority prescription form; and
 - a completed Narcolepsy Initial PBS authority application and Supporting information form; and
 - details of the contraindication or intolerance to dexamfetamine sulfate; and
 - either:
 - the result and date of the polysomnography test and Multiple Sleep Latency Test (MSLT) conducted by, or under the supervision of, a qualified sleep medicine practitioner; or
 - the result and date of the electroencephalograph (EEG), conducted by, or under the supervision of, a neurologist.
- The polysomnography, MSLT or EEG test reports must be provided with the authority application.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Narcolepsy

Treatment Phase: Initial 2 - treatment of narcolepsy with cataplexy

Treatment criteria:

- Must be treated by a qualified sleep medicine practitioner or neurologist.

Clinical criteria:

- The treatment must be for use when therapy with dexamfetamine sulfate poses an unacceptable medical risk; OR
- The treatment must be for use when intolerance to dexamfetamine sulfate is of a severity to necessitate treatment withdrawal, **AND**
- Patient must have experienced excessive daytime sleepiness, recurrent naps or lapses into sleep occurring almost daily for at least 3 months, **AND**
- Patient must have a definite history of cataplexy documented in their medical records for auditing purposes, **AND**
- Patient must not have any medical or psychiatric disorder that could otherwise account for the hypersomnia.

The presence of any one of the following indicates treatment with dexamfetamine sulfate poses an unacceptable medical risk:

- (a) a psychiatric disorder;
- (b) a cardiovascular disorder;
- (c) a history of substance abuse;
- (d) glaucoma;
- (e) any other absolute contraindication to dexamfetamine sulfate as specified in the TGA-approved Product Information.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Narcolepsy

Treatment Phase: Continuing or change of treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition; OR
- Patient must have previously received PBS-subsidised treatment with armodafinil for this condition.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

modafinil 100 mg tablet, 60

8816B	Max. Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	5	..	*206.82	41.30	^a APO-Modafinil [TX] ^a Modafinil AN [EA] ^a Modafinil Mylan [AF] ^a Modavigil [TB]	^a Modafin [RW] ^a Modafinil GH [GQ] ^a Modafinil Sandoz [SZ]

ANTI-DEMENTIA DRUGS

Anticholinesterases

▪ **DONEPEZIL**

Authority required

Mild to moderately severe Alzheimer disease

Treatment Phase: Initial 2

Clinical criteria:

- Patient must have a baseline Mini-Mental State Examination (MMSE) or Standardised Mini-Mental State Examination (SMMSE) score of 10 or more, **AND**
- The condition must be confirmed by, or in consultation with, a specialist/consultant physician (including a psychiatrist), **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

The authority application must include the result of the baseline MMSE or SMMSE. If this score is 25 - 30 points, the result of a baseline Alzheimer Disease Assessment Scale, cognitive sub-scale (ADAS-Cog) may also be specified.

Application through this treatment restriction must be made in writing.

Where a course of PBS-subsidised treatment with this drug with this strength was approved under the Initial 1 restriction, no more than 1 month's therapy and sufficient repeats to complete 6 months' initial treatment with this strength of this drug will be authorised under this restriction.

Where no prior approval has been issued before this application, up to a maximum of 1 month's therapy plus 5 repeats will be authorised.

Authority required

Mild to moderately severe Alzheimer disease

Treatment Phase: Initial 2

Clinical criteria:

- Patient must have a baseline Mini-Mental State Examination (MMSE) or Standardised Mini-Mental State Examination (SMMSE) score of 9 or less, **AND**
- The condition must be confirmed by, or in consultation with, a specialist/consultant physician (including a psychiatrist), **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

A patient who is unable to register a score of 10 or more for reasons other than their Alzheimer disease, as specified below.

Such patients will need to be assessed using the Clinicians Interview Based Impression of Severity (CIBIS) scale. The authority application must include the result of the baseline (S)MMSE and specify to which group(s) (see below) the patient belongs.

Patients who qualify under this criterion are from 1 or more of the following groups:

- (1) Unable to communicate adequately because of lack of competence in English, in people of non-English speaking background;
- (2) Limited education, as defined by less than 6 years of education, or who are illiterate or innumerate;
- (3) Aboriginal or Torres Strait Islanders who, by virtue of cultural factors, are unable to complete an (S)MMSE test;
- (4) Intellectual (developmental or acquired) disability, eg Down's syndrome;
- (5) Significant sensory impairment despite best correction, which precludes completion of an (S)MMSE test;

(6) Prominent dysphasia, out of proportion to other cognitive and functional impairment.

Application through this treatment restriction must be made in writing.

Where a course of PBS-subsidised treatment with this drug with this strength was approved under the Initial 1 restriction, no more than 1 month's therapy and sufficient repeats to complete 6 months' initial treatment with this strength of this drug will be authorised under this restriction.

Where no prior approval has been issued before this application, up to a maximum of 1 month's therapy plus 5 repeats will be authorised.

donepezil hydrochloride 10 mg tablet, 28

11924N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	22.08	23.37	^a APO-Donepezil [TX] ^a Aridon 10 [RW] ^a Donepezil AN [EA] ^a Donepezil GH [HQ]	^a Arazil [AF] ^a Aridon APN 10 [RF] ^a Donepezil-DRLA [RZ] ^a Donepezil Sandoz [SZ]
			^B 9.00	31.08	23.37	^a Aricept [PF]	

donepezil hydrochloride 5 mg tablet, 28

11922L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	22.08	23.37	^a APO-Donepezil [TX] ^a Aridon 5 [RW] ^a Donepezil AN [EA] ^a Donepezil GH [HQ]	^a Arazil [AF] ^a Aridon APN 5 [RF] ^a Donepezil-DRLA [RZ] ^a Donepezil Sandoz [SZ]
			^B 3.68	25.76	23.37	^a Aricept [PF]	

■ DONEPEZIL

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)

10108

Mild to moderately severe Alzheimer disease

Treatment Phase: Continuing

Clinical criteria:

- Patient must have received six months of sole PBS-subsidised initial therapy with this drug and has received a written authority approval, **AND**
- Patient must demonstrate a clinically meaningful response to the initial treatment, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Prior to continuing treatment, a comprehensive assessment must be undertaken and documented, involving the patient, the patient's family or carer and the treating physician to establish agreement that treatment is continuing to produce worthwhile benefit.

Treatment should cease if there is no agreement of benefit as there is always the possibility of harm from unnecessary use.

Re-assessments for a clinically meaningful response are to be undertaken and documented every six months.

Clinically meaningful response to treatment is demonstrated in the following areas:

Patient's quality of life including but not limited to level of independence and happiness;

Patient's cognitive function including but not limited to memory, recognition and interest in environment;

Patient's behavioural symptoms, including but not limited to hallucination, delusions, anxiety, marked agitation or associated aggressive behaviour.

donepezil hydrochloride 10 mg tablet, 28

2479L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	22.08	23.37	^a APO-Donepezil [TX] ^a Aridon 10 [RW] ^a Donepezil AN [EA] ^a Donepezil GH [HQ]	^a Arazil [AF] ^a Aridon APN 10 [RF] ^a Donepezil-DRLA [RZ] ^a Donepezil Sandoz [SZ]
			^B 9.00	31.08	23.37	^a Aricept [PF]	

donepezil hydrochloride 5 mg tablet, 28

2532G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	22.08	23.37	^a APO-Donepezil [TX] ^a Aridon 5 [RW] ^a Donepezil AN [EA] ^a Donepezil GH [HQ]	^a Arazil [AF] ^a Aridon APN 5 [RF] ^a Donepezil-DRLA [RZ] ^a Donepezil Sandoz [SZ]
			^B 3.68	25.76	23.37	^a Aricept [PF]	

■ DONEPEZIL

Authority required

Mild to moderately severe Alzheimer disease

Treatment Phase: Initial 1

Clinical criteria:

- Patient must have a baseline Mini-Mental State Examination (MMSE) or Standardised Mini-Mental State Examination (SMMSE) score of 10 or more, **AND**
- The condition must be confirmed by, or in consultation with, a specialist/consultant physician (including a psychiatrist), **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

The authority application must include the result of the baseline MMSE or SMMSE. If this score is 25 - 30 points, the result of a baseline Alzheimer Disease Assessment Scale, cognitive sub-scale (ADAS-Cog) may also be specified.

Up to a maximum of 2 months' initial therapy will be authorised for this drug, for this strength under this treatment restriction. This application must be followed by a written authority application for no more than 1 month's therapy and sufficient repeats to complete a maximum of up to 6 months' initial treatment with this drug with this strength.

Authority required

Mild to moderately severe Alzheimer disease

Treatment Phase: Initial 1

Clinical criteria:

- Patient must have a baseline Mini-Mental State Examination (MMSE) or Standardised Mini-Mental State Examination (SMMSE) score of 9 or less, **AND**
- The condition must be confirmed by, or in consultation with, a specialist/consultant physician (including a psychiatrist), **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

A patient who is unable to register a score of 10 or more for reasons other than their Alzheimer disease, as specified below. Such patients will need to be assessed using the Clinicians Interview Based Impression of Severity (CIBIS) scale. The authority application must include the result of the baseline (S)MMSE and specify to which group(s) (see below) the patient belongs.

Patients who qualify under this criterion are from 1 or more of the following groups:

- (1) Unable to communicate adequately because of lack of competence in English, in people of non-English speaking background;
- (2) Limited education, as defined by less than 6 years of education, or who are illiterate or innumerate;
- (3) Aboriginal or Torres Strait Islanders who, by virtue of cultural factors, are unable to complete an (S)MMSE test;
- (4) Intellectual (developmental or acquired) disability, eg Down's syndrome;
- (5) Significant sensory impairment despite best correction, which precludes completion of an (S)MMSE test;
- (6) Prominent dysphasia, out of proportion to other cognitive and functional impairment.

Up to a maximum of 2 months' initial therapy will be authorised for this drug, for this strength under this treatment restriction. This application must be followed by a written authority application for no more than 1 month's therapy and sufficient repeats to complete a maximum of up to 6 months' initial treatment with this drug with this strength.

donepezil hydrochloride 10 mg tablet, 28

8496E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	1	..	22.08	23.37	^a APO-Donepezil [TX] ^a Aridon 10 [RW] ^a Donepezil AN [EA] ^a Donepezil GH [HQ]	^a Arazil [AF] ^a Aridon APN 10 [RF] ^a Donepezil-DRLA [RZ] ^a Donepezil Sandoz [SZ]
			^B 9.00	31.08	23.37	^a Aricept [PF]	

donepezil hydrochloride 5 mg tablet, 28

8495D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	1	..	22.08	23.37	^a APO-Donepezil [TX] ^a Aridon 5 [RW] ^a Donepezil AN [EA] ^a Donepezil GH [HQ]	^a Arazil [AF] ^a Aridon APN 5 [RF] ^a Donepezil-DRLA [RZ] ^a Donepezil Sandoz [SZ]
			^B 3.68	25.76	23.37	^a Aricept [PF]	

■ GALANTAMINE

Authority required

Mild to moderately severe Alzheimer disease

Treatment Phase: Initial 2

Clinical criteria:

- Patient must have a baseline Mini-Mental State Examination (MMSE) or Standardised Mini-Mental State Examination (SMMSE) score of 10 or more, **AND**
- The condition must be confirmed by, or in consultation with, a specialist/consultant physician (including a psychiatrist), **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

The authority application must include the result of the baseline MMSE or SMMSE. If this score is 25 - 30 points, the result of a baseline Alzheimer Disease Assessment Scale, cognitive sub-scale (ADAS-Cog) may also be specified.

Application through this treatment restriction must be made in writing.

Where a course of PBS-subsidised treatment with this drug with this strength was approved under the Initial 1 restriction, no more than 1 month's therapy and sufficient repeats to complete 6 months' initial treatment with this strength of this drug will be authorised under this restriction.

Where no prior approval has been issued before this application, up to a maximum of 1 month's therapy plus 5 repeats will be authorised.

Authority required

Mild to moderately severe Alzheimer disease

Treatment Phase: Initial 2

Clinical criteria:

- Patient must have a baseline Mini-Mental State Examination (MMSE) or Standardised Mini-Mental State Examination (SMMSE) score of 9 or less, **AND**
- The condition must be confirmed by, or in consultation with, a specialist/consultant physician (including a psychiatrist), **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

A patient who is unable to register a score of 10 or more for reasons other than their Alzheimer disease, as specified below. Such patients will need to be assessed using the Clinicians Interview Based Impression of Severity (CIBIS) scale. The authority application must include the result of the baseline (S)MMSE and specify to which group(s) (see below) the patient belongs.

Patients who qualify under this criterion are from 1 or more of the following groups:

- (1) Unable to communicate adequately because of lack of competence in English, in people of non-English speaking background;
- (2) Limited education, as defined by less than 6 years of education, or who are illiterate or innumerate;
- (3) Aboriginal or Torres Strait Islanders who, by virtue of cultural factors, are unable to complete an (S)MMSE test;
- (4) Intellectual (developmental or acquired) disability, eg Down's syndrome;
- (5) Significant sensory impairment despite best correction, which precludes completion of an (S)MMSE test;
- (6) Prominent dysphasia, out of proportion to other cognitive and functional impairment.

Application through this treatment restriction must be made in writing.

Where a course of PBS-subsidised treatment with this drug with this strength was approved under the Initial 1 restriction, no more than 1 month's therapy and sufficient repeats to complete 6 months' initial treatment with this strength of this drug will be authorised under this restriction.

Where no prior approval has been issued before this application, up to a maximum of 1 month's therapy plus 5 repeats will be authorised.

galantamine 8 mg modified release capsule, 28

11917F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	33.90	35.19	^a APO-Galantamine MR [TX] ^a Galantyl [AF] ^a Reminyl [JC]	^a Galantamine AN SR [EA] ^a Gamine XR [RW]

galantamine 16 mg modified release capsule, 28

11918G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	38.78	40.07	^a APO-Galantamine MR [TX] ^a Galantyl [AF] ^a Reminyl [JC]	^a Galantamine AN SR [EA] ^a Gamine XR [RW]

galantamine 24 mg modified release capsule, 28

11899G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	43.91	41.30	^a APO-Galantamine MR [TX] ^a Galantyl [AF] ^a Reminyl [JC]	^a Galantamine AN SR [EA] ^a Gamine XR [RW]

■ GALANTAMINE**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**10108**

Mild to moderately severe Alzheimer disease

Treatment Phase: Continuing

Clinical criteria:

- Patient must have received six months of sole PBS-subsidised initial therapy with this drug and has received a written authority approval, **AND**
- Patient must demonstrate a clinically meaningful response to the initial treatment, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Prior to continuing treatment, a comprehensive assessment must be undertaken and documented, involving the patient, the patient's family or carer and the treating physician to establish agreement that treatment is continuing to produce worthwhile benefit.

Treatment should cease if there is no agreement of benefit as there is always the possibility of harm from unnecessary use.

Re-assessments for a clinically meaningful response are to be undertaken and documented every six months.

Clinically meaningful response to treatment is demonstrated in the following areas:

Patient's quality of life including but not limited to level of independence and happiness;

Patient's cognitive function including but not limited to memory, recognition and interest in environment;

Patient's behavioural symptoms, including but not limited to hallucination, delusions, anxiety, marked agitation or associated aggressive behaviour.

galantamine 8 mg modified release capsule, 28

2463P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	33.90	35.19	^a APO-Galantamine MR [TX] ^a Galantyl [AF] ^a Reminyl [JC]	^a Galantamine AN SR [EA] ^a Gamine XR [RW]

galantamine 16 mg modified release capsule, 28

2537M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	38.78	40.07	^a APO-Galantamine MR [TX] ^a Galantyl [AF] ^a Reminyl [JC]	^a Galantamine AN SR [EA] ^a Gamine XR [RW]

galantamine 24 mg modified release capsule, 28

2531F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	43.91	41.30	^a APO-Galantamine MR [TX] ^a Galantyl [AF] ^a Reminyl [JC]	^a Galantamine AN SR [EA] ^a Gamine XR [RW]

■ GALANTAMINE

Authority required

Mild to moderately severe Alzheimer disease

Treatment Phase: Initial 1

Clinical criteria:

- Patient must have a baseline Mini-Mental State Examination (MMSE) or Standardised Mini-Mental State Examination (SMMSE) score of 10 or more, **AND**
- The condition must be confirmed by, or in consultation with, a specialist/consultant physician (including a psychiatrist), **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

The authority application must include the result of the baseline MMSE or SMMSE. If this score is 25 - 30 points, the result of a baseline Alzheimer Disease Assessment Scale, cognitive sub-scale (ADAS-Cog) may also be specified.

Up to a maximum of 2 months' initial therapy will be authorised for this drug, for this strength under this treatment restriction. This application must be followed by a written authority application for no more than 1 month's therapy and sufficient repeats to complete a maximum of up to 6 months' initial treatment with this drug with this strength.

Authority required

Mild to moderately severe Alzheimer disease

Treatment Phase: Initial 1

Clinical criteria:

- Patient must have a baseline Mini-Mental State Examination (MMSE) or Standardised Mini-Mental State Examination (SMMSE) score of 9 or less, **AND**
- The condition must be confirmed by, or in consultation with, a specialist/consultant physician (including a psychiatrist), **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

A patient who is unable to register a score of 10 or more for reasons other than their Alzheimer disease, as specified below. Such patients will need to be assessed using the Clinicians Interview Based Impression of Severity (CIBIS) scale. The authority application must include the result of the baseline (S)MMSE and specify to which group(s) (see below) the patient belongs.

Patients who qualify under this criterion are from 1 or more of the following groups:

- (1) Unable to communicate adequately because of lack of competence in English, in people of non-English speaking background;
- (2) Limited education, as defined by less than 6 years of education, or who are illiterate or innumerate;
- (3) Aboriginal or Torres Strait Islanders who, by virtue of cultural factors, are unable to complete an (S)MMSE test;
- (4) Intellectual (developmental or acquired) disability, eg Down's syndrome;
- (5) Significant sensory impairment despite best correction, which precludes completion of an (S)MMSE test;
- (6) Prominent dysphasia, out of proportion to other cognitive and functional impairment.

Up to a maximum of 2 months' initial therapy will be authorised for this drug, for this strength under this treatment restriction. This application must be followed by a written authority application for no more than 1 month's therapy and sufficient repeats to complete a maximum of up to 6 months' initial treatment with this drug with this strength.

galantamine 8 mg modified release capsule, 28

8770N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	1	..	33.90	35.19	^a APO-Galantamine MR [TX] ^a Galantyl [AF] ^a Reminyl [JC]	^a Galantamine AN SR [EA] ^a Gamine XR [RW]

galantamine 16 mg modified release capsule, 28

8771P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	1	..	38.78	40.07	^a APO-Galantamine MR [TX] ^a Galantyl [AF]	^a Galantamine AN SR [EA] ^a Gamine XR [RW]

^a Reminyl [JC]**galantamine 24 mg modified release capsule, 28**

8772Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	1	..	43.91	41.30	^a APO-Galantamine MR [TX] ^a Galantyl [AF] ^a Reminyl [JC]	^a Galantamine AN SR [EA] ^a Gamine XR [RW]

■ RIVASTIGMINE**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**10108**

Mild to moderately severe Alzheimer disease

Treatment Phase: Continuing

Clinical criteria:

- Patient must have received six months of sole PBS-subsidised initial therapy with this drug and has received a written authority approval, **AND**
- Patient must demonstrate a clinically meaningful response to the initial treatment, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Prior to continuing treatment, a comprehensive assessment must be undertaken and documented, involving the patient, the patient's family or carer and the treating physician to establish agreement that treatment is continuing to produce worthwhile benefit.

Treatment should cease if there is no agreement of benefit as there is always the possibility of harm from unnecessary use.

Re-assessments for a clinically meaningful response are to be undertaken and documented every six months.

Clinically meaningful response to treatment is demonstrated in the following areas:

Patient's quality of life including but not limited to level of independence and happiness;

Patient's cognitive function including but not limited to memory, recognition and interest in environment;

Patient's behavioural symptoms, including but not limited to hallucination, delusions, anxiety, marked agitation or associated aggressive behaviour.

rivastigmine 13.3 mg/24 hours patch, 30

10538P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	81.52	41.30	Exelon Patch 15 [NV]

rivastigmine 9.5 mg/24 hours patch, 30

2551G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	81.52	41.30	Exelon Patch 10 [NV]

rivastigmine 1.5 mg capsule, 56

2475G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	89.24	41.30	Exelon [NV]

rivastigmine 3 mg capsule, 56

2493F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	89.24	41.30	Exelon [NV]

rivastigmine 6 mg capsule, 56

2526Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	89.24	41.30	Exelon [NV]

rivastigmine 4.6 mg/24 hours patch, 30

2477J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	81.52	41.30	Exelon Patch 5 [NV]

rivastigmine 4.5 mg capsule, 56

2494G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	89.24	41.30	Exelon [NV]

■ RIVASTIGMINE**Authority required**

Mild to moderately severe Alzheimer disease

Treatment Phase: Initial 1

Clinical criteria:

- Patient must have a baseline Mini-Mental State Examination (MMSE) or Standardised Mini-Mental State Examination (SMMSE) score of 10 or more, **AND**

- The condition must be confirmed by, or in consultation with, a specialist/consultant physician (including a psychiatrist), **AND**

- The treatment must be the sole PBS-subsidised therapy for this condition.

The authority application must include the result of the baseline MMSE or SMMSE. If this score is 25 - 30 points, the result of a baseline Alzheimer Disease Assessment Scale, cognitive sub-scale (ADAS-Cog) may also be specified.

Up to a maximum of 2 months' initial therapy will be authorised for this drug, for this strength under this treatment restriction. This application must be followed by a written authority application for no more than 1 month's therapy and sufficient repeats to complete a maximum of up to 6 months' initial treatment with this drug with this strength.

Authority required

Mild to moderately severe Alzheimer disease

Treatment Phase: Initial 1

Clinical criteria:

- Patient must have a baseline Mini-Mental State Examination (MMSE) or Standardised Mini-Mental State Examination (SMMSE) score of 9 or less, **AND**

- The condition must be confirmed by, or in consultation with, a specialist/consultant physician (including a psychiatrist), **AND**

- The treatment must be the sole PBS-subsidised therapy for this condition.

A patient who is unable to register a score of 10 or more for reasons other than their Alzheimer disease, as specified below. Such patients will need to be assessed using the Clinicians Interview Based Impression of Severity (CIBIS) scale. The authority application must include the result of the baseline (S)MMSE and specify to which group(s) (see below) the patient belongs.

Patients who qualify under this criterion are from 1 or more of the following groups:

- (1) Unable to communicate adequately because of lack of competence in English, in people of non-English speaking background;
- (2) Limited education, as defined by less than 6 years of education, or who are illiterate or innumerate;
- (3) Aboriginal or Torres Strait Islanders who, by virtue of cultural factors, are unable to complete an (S)MMSE test;
- (4) Intellectual (developmental or acquired) disability, eg Down's syndrome;
- (5) Significant sensory impairment despite best correction, which precludes completion of an (S)MMSE test;
- (6) Prominent dysphasia, out of proportion to other cognitive and functional impairment.

Up to a maximum of 2 months' initial therapy will be authorised for this drug, for this strength under this treatment restriction. This application must be followed by a written authority application for no more than 1 month's therapy and sufficient repeats to complete a maximum of up to 6 months' initial treatment with this drug with this strength.

rivastigmine 13.3 mg/24 hours patch, 30

10541T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	81.52	41.30	Exelon Patch 15 [NV]

rivastigmine 9.5 mg/24 hours patch, 30

9162F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	81.52	41.30	Exelon Patch 10 [NV]

rivastigmine 1.5 mg capsule, 56

8497F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	89.24	41.30	Exelon [NV]

rivastigmine 3 mg capsule, 56

8498G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	89.24	41.30	Exelon [NV]

rivastigmine 6 mg capsule, 56

8500J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	89.24	41.30	Exelon [NV]

rivastigmine 4.6 mg/24 hours patch, 30

9161E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	81.52	41.30	Exelon Patch 5 [NV]

rivastigmine 4.5 mg capsule, 56

8499H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	89.24	41.30	Exelon [NV]

▪ RIVASTIGMINE

Authority required

Mild to moderately severe Alzheimer disease

Treatment Phase: Initial 2

Clinical criteria:

- Patient must have a baseline Mini-Mental State Examination (MMSE) or Standardised Mini-Mental State Examination (SMMSE) score of 10 or more, **AND**

- The condition must be confirmed by, or in consultation with, a specialist/consultant physician (including a psychiatrist), **AND**
 - The treatment must be the sole PBS-subsidised therapy for this condition.
- The authority application must include the result of the baseline MMSE or SMMSE. If this score is 25 - 30 points, the result of a baseline Alzheimer Disease Assessment Scale, cognitive sub-scale (ADAS-Cog) may also be specified.

Application through this treatment restriction must be made in writing.

Where a course of PBS-subsidised treatment with this drug with this strength was approved under the Initial 1 restriction, no more than 1 month's therapy and sufficient repeats to complete 6 months' initial treatment with this strength of this drug will be authorised under this restriction.

Where no prior approval has been issued before this application, up to a maximum of 1 month's therapy plus 5 repeats will be authorised.

Authority required

Mild to moderately severe Alzheimer disease

Treatment Phase: Initial 2

Clinical criteria:

- Patient must have a baseline Mini-Mental State Examination (MMSE) or Standardised Mini-Mental State Examination (SMMSE) score of 9 or less, **AND**
 - The condition must be confirmed by, or in consultation with, a specialist/consultant physician (including a psychiatrist), **AND**
 - The treatment must be the sole PBS-subsidised therapy for this condition.
- A patient who is unable to register a score of 10 or more for reasons other than their Alzheimer disease, as specified below. Such patients will need to be assessed using the Clinicians Interview Based Impression of Severity (CIBIS) scale. The authority application must include the result of the baseline (S)MMSE and specify to which group(s) (see below) the patient belongs.

Patients who qualify under this criterion are from 1 or more of the following groups:

- (1) Unable to communicate adequately because of lack of competence in English, in people of non-English speaking background;
- (2) Limited education, as defined by less than 6 years of education, or who are illiterate or innumerate;
- (3) Aboriginal or Torres Strait Islanders who, by virtue of cultural factors, are unable to complete an (S)MMSE test;
- (4) Intellectual (developmental or acquired) disability, eg Down's syndrome;
- (5) Significant sensory impairment despite best correction, which precludes completion of an (S)MMSE test;
- (6) Prominent dysphasia, out of proportion to other cognitive and functional impairment.

Application through this treatment restriction must be made in writing.

Where a course of PBS-subsidised treatment with this drug with this strength was approved under the Initial 1 restriction, no more than 1 month's therapy and sufficient repeats to complete 6 months' initial treatment with this strength of this drug will be authorised under this restriction.

Where no prior approval has been issued before this application, up to a maximum of 1 month's therapy plus 5 repeats will be authorised.

rivastigmine 13.3 mg/24 hours patch, 30

11904M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	81.52	41.30	Exelon Patch 15 [NV]

rivastigmine 9.5 mg/24 hours patch, 30

11903L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	81.52	41.30	Exelon Patch 10 [NV]

rivastigmine 1.5 mg capsule, 56

11925P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	89.24	41.30	Exelon [NV]

rivastigmine 3 mg capsule, 56

11901J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	89.24	41.30	Exelon [NV]

rivastigmine 6 mg capsule, 56

11916E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	89.24	41.30	Exelon [NV]

rivastigmine 4.6 mg/24 hours patch, 30

11912Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	81.52	41.30	Exelon Patch 5 [NV]

rivastigmine 4.5 mg capsule, 56

11923M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	89.24	41.30	Exelon [NV]

Other anti-dementia drugs

■ MEMANTINE

Authority required

Moderately severe Alzheimer disease

Treatment Phase: Initial 2

Clinical criteria:

- Patient must have a baseline Mini-Mental State Examination (MMSE) or Standardised Mini-Mental State Examination (SMMSE) score of 10 to 14, **AND**
- The condition must be confirmed by, or in consultation with, a specialist/consultant physician (including a psychiatrist), **AND**

- The treatment must be the sole PBS-subsidised therapy for this condition.

The authority application must include the result of the baseline MMSE or SMMSE of 10 to 14.

Application through this treatment restriction must be made in writing.

Where a course of PBS-subsidised treatment with this drug with this strength was approved under the Initial 1 restriction, no more than 1 month's therapy and sufficient repeats to complete 6 months' initial treatment with this strength of this drug will be authorised under this restriction.

Where no prior approval has been issued before this application, up to a maximum of 1 month's therapy plus 5 repeats will be authorised.

Authority required

Moderately severe Alzheimer disease

Treatment Phase: Initial 2

Clinical criteria:

- Patient must have a baseline Mini-Mental State Examination (MMSE) or Standardised Mini-Mental State Examination (SMMSE) score of 9 or less, **AND**
- The condition must be confirmed by, or in consultation with, a specialist/consultant physician (including a psychiatrist), **AND**

- The treatment must be the sole PBS-subsidised therapy for this condition.

A patient who is unable to register a score of 10 to 14 for reasons other than their Alzheimer disease, as specified below.

Such patients will need to be assessed using the Clinicians Interview Based Impression of Severity (CIBIS) scale. The authority application must include the result of the baseline (S)MMSE and specify to which group(s) (see below) the patient belongs.

Patients who qualify under this criterion are from 1 or more of the following groups:

- (1) Unable to communicate adequately because of lack of competence in English, in people of non-English speaking background;
- (2) Limited education, as defined by less than 6 years of education, or who are illiterate or innumerate;
- (3) Aboriginal or Torres Strait Islanders who, by virtue of cultural factors, are unable to complete an (S)MMSE test;
- (4) Intellectual (developmental or acquired) disability, eg Down's syndrome;
- (5) Significant sensory impairment despite best correction, which precludes completion of an (S)MMSE test;
- (6) Prominent dysphasia, out of proportion to other cognitive and functional impairment.

Application through this treatment restriction must be made in writing.

Where a course of PBS-subsidised treatment with this drug with this strength was approved under the Initial 1 restriction, no more than 1 month's therapy and sufficient repeats to complete 6 months' initial treatment with this strength of this drug will be authorised under this restriction.

Where no prior approval has been issued before this application, up to a maximum of 1 month's therapy plus 5 repeats will be authorised.

memantine hydrochloride 10 mg tablet, 56

11902K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	42.22	41.30	^a APO-Memantine [TX] ^a Memantine generichealth [GQ]	^a Ebixa [LU] ^a Memanxa [RW]

memantine hydrochloride 20 mg tablet, 28

11905N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	42.22	41.30	^a APO-Memantine [TX] ^a Memantine generichealth [GQ]	^a Ebixa [LU]

■ MEMANTINE

Authority required

Moderately severe Alzheimer disease

Treatment Phase: Initial 1

Clinical criteria:

- Patient must have a baseline Mini-Mental State Examination (MMSE) or Standardised Mini-Mental State Examination (SMMSE) score of 10 to 14, **AND**
- The condition must be confirmed by, or in consultation with, a specialist/consultant physician (including a psychiatrist), **AND**

- The treatment must be the sole PBS-subsidised therapy for this condition.

The authority application must include the result of the baseline MMSE or SMMSE of 10 to 14.

Up to a maximum of 2 months' initial therapy will be authorised for this drug, for this strength under this treatment restriction. This application must be followed by a written authority application for no more than 1 month's therapy and sufficient repeats to complete a maximum of up to 6 months' initial treatment with this drug with this strength.

Authority required

Moderately severe Alzheimer disease

Treatment Phase: Initial 1

Clinical criteria:

- Patient must have a baseline Mini-Mental State Examination (MMSE) or Standardised Mini-Mental State Examination (SMMSE) score of 9 or less, **AND**
- The condition must be confirmed by, or in consultation with, a specialist/consultant physician (including a psychiatrist), **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

A patient who is unable to register a score of 10 to 14 for reasons other than their Alzheimer disease, as specified below.

Such patients will need to be assessed using the Clinicians Interview Based Impression of Severity (CIBIS) scale. The authority application must include the result of the baseline (S)MMSE and specify to which group(s) (see below) the patient belongs.

Patients who qualify under this criterion are from 1 or more of the following groups:

- (1) Unable to communicate adequately because of lack of competence in English, in people of non-English speaking background;
- (2) Limited education, as defined by less than 6 years of education, or who are illiterate or innumerate;
- (3) Aboriginal or Torres Strait Islanders who, by virtue of cultural factors, are unable to complete an (S)MMSE test;
- (4) Intellectual (developmental or acquired) disability, eg Down's syndrome;
- (5) Significant sensory impairment despite best correction, which precludes completion of an (S)MMSE test;
- (6) Prominent dysphasia, out of proportion to other cognitive and functional impairment.

Up to a maximum of 2 months' initial therapy will be authorised for this drug, for this strength under this treatment restriction. This application must be followed by a written authority application for no more than 1 month's therapy and sufficient repeats to complete a maximum of up to 6 months' initial treatment with this drug with this strength.

memantine hydrochloride 10 mg tablet, 56

1956Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	1	..	42.22	41.30	^a APO-Memantine [TX] ^a Memantine generichealth [GQ]	^a Ebixa [LU] ^a Memanxa [RW]

memantine hydrochloride 20 mg tablet, 28

9306T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	1	..	42.22	41.30	^a APO-Memantine [TX] ^a Memantine generichealth [GQ]	^a Ebixa [LU]

MEMANTINE

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)

10104

Moderately severe Alzheimer disease

Treatment Phase: Continuing

Clinical criteria:

- Patient must have received six months of sole PBS-subsidised initial therapy with this drug and has received a written authority approval, **AND**
- Patient must demonstrate a clinically meaningful response to the initial treatment, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Prior to continuing treatment, a comprehensive assessment must be undertaken and documented, involving the patient, the patient's family or carer and the treating physician to establish agreement that treatment is continuing to produce worthwhile benefit.

Treatment should cease if there is no agreement of benefit as there is always the possibility of harm from unnecessary use.

Re-assessments for a clinically meaningful response are to be undertaken and documented every six months.

Clinically meaningful response to treatment is demonstrated in the following areas:

Patient's quality of life including but not limited to level of independence and happiness;

Patient's cognitive function including but not limited to memory, recognition and interest in environment;

Patient's behavioural symptoms, including but not limited to hallucination, delusions, anxiety, marked agitation or associated aggressive behaviour.

memantine hydrochloride 10 mg tablet, 56

2492E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	42.22	41.30	^a APO-Memantine [TX] ^a Memantine generichealth [GQ]	^a Ebixa [LU] ^a Memanxa [RW]

NP

NERVOUS SYSTEM

memantine hydrochloride 20 mg tablet, 28

2513G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	42.22	41.30	^a APO-Memantine [TX] ^a Memantine generichealth [GQ]	^a Ebixa [LU]

OTHER NERVOUS SYSTEM DRUGS

PARASYMPATHOMIMETICS

Anticholinesterases

■ PYRIDOSTIGMINE

pyridostigmine bromide 180 mg modified release tablet, 50

2608G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*117.74	41.30	Mestinon Timespan [IL]

pyridostigmine bromide 60 mg tablet, 150

1959D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	59.94	41.30	Mestinon [IL]

pyridostigmine bromide 10 mg tablet, 50

2724J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*23.74	25.03	Mestinon [IL]

Choline esters

■ BETHANECHOL

bethanechol chloride 10 mg tablet, 100

1062X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	34.95	36.24	Uro-Carb [YN]

DRUGS USED IN ADDICTIVE DISORDERS

Drugs used in nicotine dependence

■ BUPROPION

Note Clinical review is recommended within 2 to 3 weeks of the original prescription being requested.

Note The period between commencing a course of bupropion hydrochloride and varenicline tartrate must be at least 6 months.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

6881

Nicotine dependence

Treatment Phase: Completion of a short-term (9 weeks) course of treatment

Clinical criteria:

- The treatment must be as an aid to achieving abstinence from smoking, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug during this current course of treatment, **AND**
- Patient must not receive more than 9 weeks of PBS-subsidised treatment with this drug per 12-month period.

Treatment criteria:

- Patient must be undergoing concurrent counselling for smoking cessation through a comprehensive support and counselling program.

bupropion hydrochloride 150 mg modified release tablet, 90

8710K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	169.05	41.30	Zyban [AS]

■ BUPROPION

Note Clinical review is recommended within 2 to 3 weeks of the original prescription being requested.

Note The period between commencing a course of bupropion hydrochloride and varenicline tartrate must be at least 6 months.

Note A patient may only qualify for PBS-subsidised treatment under this treatment phase restriction once during a short-term course of treatment.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

6882

Nicotine dependence

Treatment Phase: Commencement of a short-term (9 weeks) course of treatment

Clinical criteria:

- The treatment must be as an aid to achieving abstinence from smoking, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have indicated they are ready to cease smoking, **AND**
- Patient must not receive more than 9 weeks of PBS-subsidised treatment with this drug per 12-month period.

Treatment criteria:

- Patient must be undergoing concurrent counselling for smoking cessation through a comprehensive support and counselling program or is about to enter such a program at the time PBS-subsidised treatment is initiated.

Details of the support and counselling program must be documented in the patient's medical records at the time treatment is initiated.

bupropion hydrochloride 150 mg modified release tablet, 30

8465M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	63.46	41.30	Zyban [AS]

■ NICOTINE

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Restricted benefit

Nicotine dependence

Clinical criteria:

- The treatment must be as an aid to achieving abstinence from smoking, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have indicated they are ready to cease smoking, **AND**
- Patient must not receive more than 12 weeks of PBS-subsidised nicotine replacement therapy per 12-month period.

Treatment criteria:

- Patient must be undergoing concurrent counselling for smoking cessation through a comprehensive support and counselling program or is about to enter such a program at the time PBS-subsidised treatment is initiated.

Details of the support and counselling program must be documented in the patient's medical records at the time treatment is initiated.

nicotine 7 mg/24 hours patch, 28

5573H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	49.90	41.30	Nicotinell Step 3 [ON]

nicotine 14 mg/24 hours patch, 28

5572G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	49.90	41.30	Nicotinell Step 2 [ON]

nicotine 21 mg/24 hours patch, 28

3414Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	49.90	41.30	Nicotinell Step 1 [ON]

■ NICOTINE

Note Only 2 courses of PBS-subsidised nicotine replacement therapy may be prescribed per 12-month period. Benefit is improved if used in conjunction with a comprehensive support and counselling program.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Restricted benefit

Nicotine dependence

Population criteria:

- Patient must be an Aboriginal or a Torres Strait Islander person.

Clinical criteria:

- The treatment must be the sole PBS-subsidised therapy for this condition.

nicotine 21 mg/24 hours patch, 28

5571F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	49.90	41.30	Nicotinell Step 1 [ON]

■ NICOTINE

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Restricted benefit

Nicotine dependence

Population criteria:

- Patient must be an Aboriginal or a Torres Strait Islander person.

Clinical criteria:

- The treatment must be the sole PBS-subsidised therapy for this condition.

Note Only 2 courses of PBS-subsidised nicotine replacement therapy may be prescribed per 12-month period. Benefit is improved if used in conjunction with a comprehensive support and counselling program.

Restricted benefit

Nicotine dependence

Clinical criteria:

- The treatment must be as an aid to achieving abstinence from smoking, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have indicated they are ready to cease smoking, **AND**
- Patient must not receive more than 12 weeks of PBS-subsidised nicotine replacement therapy per 12-month period.

Treatment criteria:

- Patient must be undergoing concurrent counselling for smoking cessation through a comprehensive support and counselling program or is about to enter such a program at the time PBS-subsidised treatment is initiated.

Details of the support and counselling program must be documented in the patient's medical records at the time treatment is initiated.

nicotine 4 mg lozenge, 216

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
11619M	1	2	..	49.90	41.30	Nicotinell [ON]

nicotine 4 mg chewing gum, 216

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
11612E	1	2	..	49.90	41.30	Nicotinell [ON]

nicotine 2 mg lozenge, 216

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
11617K	1	2	..	49.90	41.30	Nicotinell [ON]

nicotine 21 mg/24 hours patch, 28

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
5465P	1	2	..	49.90	41.30	Nicabate P [GC]

nicotine 25 mg/16 hours patch, 28

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
10076H	1	2	..	49.90	41.30	nicorette 16hr Invisipatch [JT]

nicotine 2 mg chewing gum, 216

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
11618L	2	1	..	*68.84	41.30	Nicotinell [ON]

■ VARENICLINE

Note A course of treatment with this drug is 12 weeks or up to 24 weeks, if initial treatment of 12 weeks has been successful.

Note A patient may only qualify for PBS-subsidised treatment under this treatment phase restriction once during a short-term course of treatment.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

6885

Nicotine dependence

Treatment Phase: Completion of a short-term (24 weeks) course of treatment

Clinical criteria:

- The treatment must be as an aid to achieving abstinence from smoking, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug during this current course of treatment, **AND**
- Patient must have ceased smoking in the process of completing an initial 12-weeks or ceased smoking following an initial 12-weeks of PBS-subsidised treatment with this drug in the current course of treatment.

Treatment criteria:

- Patient must be undergoing concurrent counselling for smoking cessation through a comprehensive support and counselling program.

varenicline 1 mg tablet, 56

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
5469W	1	2	..	100.63	41.30	Champix [PF]

■ VARENICLINE

Note A course of treatment with this drug is 12 weeks or up to 24 weeks, if initial treatment of 12 weeks has been successful.

Note A patient may only qualify for PBS-subsidised treatment under this treatment phase restriction once during a short-term course of treatment.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

7483

Nicotine dependence

Treatment Phase: Continuation of a short-term (12 weeks or 24 weeks) course of treatment

Clinical criteria:

- The treatment must be as an aid to achieving abstinence from smoking, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have previously received treatment with this drug during this current course of treatment.

Treatment criteria:

- Patient must be undergoing concurrent counselling for smoking cessation through a comprehensive support and counselling program.

varenicline 1 mg tablet, 56

9129L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	..	..	*193.10	41.30	Champix [PF]

■ **VARENICLINE**

Note A course of treatment with this drug is 12 weeks or up to 24 weeks, if initial treatment of 12 weeks has been successful.

Note The period between commencing varenicline and bupropion or a new course of varenicline must be at least 6 months.

Note A patient may only qualify for PBS-subsidised treatment under this treatment phase restriction once during a short-term course of treatment.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

6871

Nicotine dependence

Treatment Phase: Commencement of a short-term (12 weeks or 24 weeks) course of treatment

Clinical criteria:

- The treatment must be as an aid to achieving abstinence from smoking, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have indicated they are ready to cease smoking, **AND**
- Patient must not receive more than 24 weeks of PBS-subsidised treatment with this drug per 12-month period.

Treatment criteria:

- Patient must be undergoing concurrent counselling for smoking cessation through a comprehensive support and counselling program or is about to enter such a program at the time PBS-subsidised treatment is initiated.

Details of the support and counselling program must be documented in the patient's medical records at the time treatment is initiated.

Clinical review is recommended within 2 to 3 weeks of the initial prescription being requested.

varenicline 500 microgram tablet [11] (&) varenicline 1 mg tablet [42], 53

9128K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	..	..	87.18	41.30	Champix [PF]

Drugs used in alcohol dependence

■ **ACAMPROSATE**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

5366

Alcohol dependence

Clinical criteria:

- The treatment must be part of a comprehensive treatment program with the goal of maintaining abstinence.

acamprosate calcium 333 mg enteric tablet, 180

8357W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	1	..	114.87	41.30	^a Acamprosate Mylan [AL] ^a Campral [AF]	^a APO-Acamprosate [TX]

■ **NALTREXONE**

Caution Naltrexone hydrochloride is contraindicated in patients receiving opioid drugs.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Alcohol dependence

Clinical criteria:

- The treatment must be part of a comprehensive treatment program with the goal of maintaining abstinence.

naltrexone hydrochloride 50 mg tablet, 30

8370M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	1	..	130.40	41.30	^a APO-Naltrexone [TX]	^a Naltrexone GH [GQ]

OTHER NERVOUS SYSTEM DRUGS*Other nervous system drugs***■ RILUZOLE****Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required

Amyotrophic lateral sclerosis

Treatment Phase: Initial treatment

Clinical criteria:

- The condition must be diagnosed by a neurologist, **AND**
- Patient must not have had the disease for more than 5 years, **AND**
- Patient must have at least 60 percent of predicted forced vital capacity within the 2 months before commencing therapy with this drug, **AND**
- Patient must be ambulatory; OR
- Patient must not be ambulatory, and must be able to either use upper limbs or to swallow, **AND**
- Patient must not have undergone a tracheostomy, **AND**
- Patient must not have experienced respiratory failure.

The date of diagnosis and the date and results of spirometry (in terms of percent of predicted forced vital capacity) must be supplied with the initial authority application.

Authority required

Amyotrophic lateral sclerosis

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition, **AND**
- Patient must be ambulatory; OR
- Patient must not be ambulatory, and must be able to either use upper limbs or to swallow, **AND**
- Patient must not have undergone a tracheostomy, **AND**
- Patient must not have experienced respiratory failure.

riluzole 50 mg/10 mL oral liquid, 300 mL

11662T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*244.40	41.30	Teglutik [CS]

riluzole 50 mg tablet, 56

8664B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	208.43	41.30	^a APO-Riluzole [TX] ^a Rilutek [SW]	^a Pharmacor Riluzole [CR] ^a Riluzole Sandoz [SZ]

■ TETRABENAZINE**Note Continuing Therapy Only:**

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**5340**

Hyperkinetic extrapyramidal disorders

tetrabenazine 25 mg tablet, 112

1330B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	296.57	41.30	iNova Pharmaceuticals (Australia) Pty Ltd [IL]

■ ANTIPARASITIC PRODUCTS, INSECTICIDES AND REPELLENTS**■ ANTIPROTOZOALS****AGENTS AGAINST AMOEBIASIS AND OTHER PROTOZOAL DISEASES***Other agents against amoebiasis and other protozoal diseases*

■ ATOVAQUONE

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)

5609

Mild to moderate *Pneumocystis carinii* pneumonia

Population criteria:

- Patient must be an adult, **AND**
- Patient must be intolerant of trimethoprim/sulfamethoxazole therapy.

atovaquone 750 mg/5 mL oral liquid, 210 mL

8300W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	..	..	981.04	41.30	Wellvone [AS]

ANTIMALARIALS

Biguanides

■ ATOVAQUONE + PROGUANIL

Note This drug is not PBS-subsidised for prophylaxis of malaria.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Confirmed or suspected *Plasmodium falciparum* malaria

Population criteria:

- Patient must be aged 3 years or older.

Clinical criteria:

- The treatment must be used where quinine containing regimens are inappropriate.

atovaquone 250 mg + proguanil hydrochloride 100 mg tablet, 12

9439T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	67.09	41.30	Malarone [GK]

Methanolquinolines

■ QUININE

Caution Severe thrombocytopenia has been reported with this drug.

Authority required (STREAMLINED)

5633

Malaria

quinine sulfate dihydrate 300 mg tablet, 50

1975Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	20.84	22.13	Quinate [RW]

Artemisinin and derivatives, combinations

■ ARTEMETHER + LUMEFANTRINE

Note This drug is not PBS-subsidised for prophylaxis of malaria.

Restricted benefit

Confirmed or suspected *Plasmodium falciparum* malaria

Clinical criteria:

- Patient must be unable to swallow a solid dosage form of artemether with lumefantrine.

artemether 20 mg + lumefantrine 120 mg dispersible tablet, 18

5296R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	100.48	41.30	Riamet 20mg/120mg Dispersible [SZ]

■ ARTEMETHER + LUMEFANTRINE

Note This drug is not PBS-subsidised for prophylaxis of malaria.

Restricted benefit

Confirmed or suspected *Plasmodium falciparum* malaria

artemether 20 mg + lumefantrine 120 mg tablet, 24

9498X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	82.35	41.30	Riamet [SZ]

■ **ANTHELMINTICS****ANTITREMATODALS***Quinoline derivatives and related substances*■ **PRAZIQUANTEL**Authority required (STREAMLINED)**5659**

Schistosomiasis

praziquantel 600 mg tablet, 8

9447F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	37.62	38.91	Biltricide [BN]

ANTINEMATODAL AGENTS*Benzimidazole derivatives*■ **ALBENDAZOLE**Authority required (STREAMLINED)**5607**

Hydatid disease

Clinical criteria:

- The treatment must be in conjunction with surgery; OR
- The treatment must be used when a surgical cure cannot be achieved; OR
- The treatment must be used when surgery cannot be used.

albendazole 400 mg tablet, 60

8459F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	169.18	41.30	Eskazole [AS]

■ **ALBENDAZOLE**Authority required (STREAMLINED)**5680**

Tapeworm infestation

albendazole 200 mg tablet, 6

8503M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	35.67	36.96	Zentel [AS]

■ **ALBENDAZOLE**Authority required (STREAMLINED)**5817**

Whipworm infestation

Population criteria:

- Patient must be an Aboriginal or a Torres Strait Islander person.

Authority required (STREAMLINED)**5712**

Strongyloidiasis

Authority required (STREAMLINED)**5797**

Hookworm infestation

albendazole 200 mg tablet, 6

9047E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	35.67	36.96	Zentel [AS]

Tetrahydropyrimidine derivatives■ **PYRANTEL****pyrantel 250 mg tablet, 6**

3048K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	23.58	24.87	Anthel 250 [AF]

pyrantel 125 mg tablet, 6

3047J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	17.80	19.09	Anthel 125 [AF]

Avermectines

■ IVERMECTIN

Authority required (STREAMLINED)

4319

Onchocerciasis

ivermectin 3 mg tablet, 4

8359Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	29.60	30.89	Stromectol [MK]

■ IVERMECTIN

Authority required (STREAMLINED)

4328

Strongyloidiasis

Authority required (STREAMLINED)

4565

Crusted (Norwegian) scabies

Clinical criteria:

- The condition must be established by clinical and/or parasitological examination, **AND**
- Patient must be undergoing topical therapy for this condition; OR
- Patient must have a contraindication to topical treatment.

Population criteria:

- Patient must weigh 15 kg or over, **AND**
- Patient must be 5 years of age or older.

Authority required (STREAMLINED)

4566

Human sarcoptic scabies

Clinical criteria:

- The condition must be established by clinical and/or parasitological examination, **AND**
- Patient must have completed and failed sequential treatment with topical permethrin and benzyl benzoate and finished the most recent course of topical therapy at least 4 weeks prior to initiating oral therapy; OR
- Patient must have a contraindication to topical treatment.

Population criteria:

- Patient must weigh 15 kg or over, **AND**
- Patient must be 5 years of age or older.

Note This drug is not PBS-subsidised for first line treatment of typical scabies.

ivermectin 3 mg tablet, 4

2868Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	2	..	*47.18	41.30	Stromectol [MK]

■ ECTOPARASITICIDES, INCL. SCABICIDES, INSECTICIDES AND REPELLENTS

ECTOPARASITICIDES, INCL. SCABICIDES

Pyrethrines, incl. synthetic compounds

■ PERMETHRIN

permethrin 5% cream, 30 g

3054R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	1	..	19.33	20.62	Lyclear [JT]

■ RESPIRATORY SYSTEM

■ NASAL PREPARATIONS

DECONGESTANTS AND OTHER NASAL PREPARATIONS FOR TOPICAL USE

Other nasal preparations

■ MUPIROCIN

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)

6647

Staphylococcus aureus infection

Clinical criteria:

- Patient must have nasal colonisation with the bacteria.

Population criteria:

- Patient must be an Aboriginal or a Torres Strait Islander person.

mupirocin 2% ointment, 3 g

9440W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	..	..	25.54	26.83	Bactroban [GK]

mupirocin 2% ointment, 5 g

11822F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	..	..	25.54	26.83	Mupirocin Nasal (Medsurge) [DZ]

DRUGS FOR OBSTRUCTIVE AIRWAY DISEASES**ADRENERGICS, INHALANTS***Selective beta-2-adrenoreceptor agonists***FORMOTEROL (EFORMOTEROL)**Restricted benefit

Asthma

Clinical criteria:

- Patient must experience frequent episodes of the condition, **AND**
- Patient must be currently receiving treatment with oral corticosteroids; OR
- Patient must be currently receiving treatment with optimal doses of inhaled corticosteroids.

formoterol (eformoterol) fumarate dihydrate 6 microgram/actuation powder for inhalation, 60 actuations

8239P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	26.13	27.42	Oxis Turbuhaler [AP]

formoterol (eformoterol) fumarate dihydrate 12 microgram powder for inhalation, 60 capsules

8136F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	33.86	35.15	Foradile [SZ]

formoterol (eformoterol) fumarate dihydrate 12 microgram/actuation powder for inhalation, 60 actuations

8240Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	33.22	34.51	Oxis Turbuhaler [AP]

INDACATEROL**Note** This drug is not PBS-subsidised for the treatment of asthma.**Note** The treatment must not be used in combination with an ICS/LABA, or LAMA/LABA**Note** A LAMA/LABA includes aclidinium/formoterol, glycopyrronium/indacaterol, tiotropium/olodaterol, or umeclidinium/vilanterol.**Note** An ICS/LABA includes budesonide/formoterol, fluticasone/salmeterol, or fluticasone/vilanterol**Note** Diagnosis of COPD should include measurement of airflow obstruction using spirometry, with confirmation of post-bronchodilator airflow obstruction.**Note** Adherence to current treatment and device (inhaler) technique should be reviewed at each clinical visit and before "stepping up" a patient's medication regimen.Restricted benefit

Chronic obstructive pulmonary disease (COPD)

indacaterol 150 microgram powder for inhalation, 30 capsules

5134F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	60.35	41.30	Onbrez [NV]

indacaterol 300 microgram powder for inhalation, 30 capsules

5137J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	60.35	41.30	Onbrez [NV]

SALBUTAMOL**Note** Pharmaceutical benefits that have the form salbutamol 100 microgram/actuation inhalation, 200 actuations are equivalent for the purposes of substitution.**salbutamol 100 microgram/actuation inhalation, 200 actuations**

12109H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	2	5	..	*21.68	22.97	^a Zempreon CFC-Free with dose counter [GC]	
			^B 5.00	*26.68	22.97	^a Asmol CFC-Free with dose counter [AF]	^a Ventolin CFC-Free with dose counter [GK]

salbutamol 100 microgram/actuation inhalation, 200 actuations

8288F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*20.40	21.69	^a Asmol CFC-free [AL]
			^B 3.66	*24.06	21.69	^a Ventolin CFC-free [GK]

■ SALBUTAMOL

Restricted benefit

Bronchospasm

Clinical criteria:

- Patient must be unable to achieve co-ordinated use of other metered dose inhalers containing this drug.

salbutamol 100 microgram/actuation breath activated inhalation, 200 actuations

8354Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*40.00	41.29	Airomir Autohaler [IL]

■ SALBUTAMOL

Note Pharmaceutical benefits that have a 30 x 2 pack size and a 20 x 3 pack size are equivalent for the purposes of substitution.

Restricted benefit

Asthma

Clinical criteria:

- Patient must be unable to use this drug delivered from an oral pressurised inhalation device via a spacer.

Restricted benefit

Chronic obstructive pulmonary disease (COPD)

Clinical criteria:

- Patient must be unable to use this drug delivered from an oral pressurised inhalation device via a spacer.

Note Diagnosis of COPD should include measurement of airflow obstruction using spirometry, with confirmation of post-bronchodilator airflow obstruction.

Note Adherence to current treatment and device (inhaler) technique should be reviewed at each clinical visit and before "stepping up" a patient's medication regimen.

salbutamol 5 mg/2.5 mL inhalation solution, 30 x 2.5 mL ampoules

2001H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	2	5	..	*20.44	21.73	^a APO-Salbutamol [TX]	^a Salbutamol Actavis [EA]
						^a Salbutamol AN [ED]	^a Salbutamol Cipla [LR]
			^B 0.50	*20.94	21.73	^a Asmol 5 uni-dose [AF]	

salbutamol 5 mg/2.5 mL inhalation solution, 20 x 2.5 mL ampoules

11095Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	3	5	^B 4.98	*25.44	21.75	^a Ventolin Nebules [GK]

salbutamol 2.5 mg/2.5 mL inhalation solution, 30 x 2.5 mL ampoules

2000G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	2	5	..	*20.02	21.31	^a APO-Salbutamol [TX]	^a Salbutamol Actavis [EA]
						^a Salbutamol AN [ED]	^a Salbutamol Cipla [LR]
			^B 0.50	*20.52	21.31	^a Asmol 2.5 uni-dose [AF]	

salbutamol 2.5 mg/2.5 mL inhalation solution, 20 x 2.5 mL ampoules

11130T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	3	5	^B 5.01	*25.05	21.33	^a Ventolin Nebules [GK]

■ SALMETEROL

Restricted benefit

Asthma

Clinical criteria:

- Patient must experience frequent episodes of the condition, **AND**
- Patient must be currently receiving treatment with oral corticosteroids; OR
- Patient must be currently receiving treatment with optimal doses of inhaled corticosteroids.

salmeterol 50 microgram/actuation powder for inhalation, 60 actuations

8141L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	33.86	35.15	Serevent Accuhaler [GK]

■ TERBUTALINE

Authority required (STREAMLINED)

9828

Bronchospasm

Clinical criteria:

- Patient must be unable to achieve co-ordinated use of a metered dose inhaler containing a short-acting beta-2 agonist; OR
- Patient must have developed a clinically important product-related adverse event during treatment with another short-acting beta-2 agonist.

Device (inhaler) technique should be reviewed at each clinical visit and before initiating treatment with this medicine.

terbutaline sulfate 500 microgram/actuation powder for inhalation, 100 actuations

2817G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*26.68	27.97	Bricanyl Turbuhaler [AP]

terbutaline sulfate 500 microgram/actuation powder for inhalation, 120 actuations

12267P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*26.68	27.97	Bricanyl Turbuhaler [AP]

Adrenergics in combination with corticosteroids or other drugs, excl. anticholinergics

■ BECLOMETASONE + FORMOTEROL (EFORMOTEROL)

Note This product is not indicated for the initiation of treatment in asthma

Note This drug is not PBS-subsidised for the treatment of chronic obstructive pulmonary disease (COPD).

Note The patient must not be on a concomitant single agent long-acting-beta-2-agonist (LABA)

Note A LABA includes olodaterol, indacaterol, salmeterol, formoterol or vilanterol.

Note Adherence to current treatment and device (inhaler) technique should be reviewed at each clinical visit and before "stepping up" a patient's medication regimen.

Note This product is not PBS-subsidised for use as 'maintenance and reliever' therapy.

Note This product is not PBS-subsidised for use as 'anti-inflammatory reliever' therapy for mild asthma.

Authority required (STREAMLINED)

11057

Asthma

Clinical criteria:

- Patient must have previously had frequent episodes of asthma while receiving treatment with oral corticosteroids or optimal doses of inhaled corticosteroids.

Population criteria:

- Patient must be aged 18 years or older.

beclometasone dipropionate 100 microgram/actuation + formoterol (eformoterol) fumarate dihydrate 6 microgram/actuation inhalation, 120 actuations

12183F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	43.94	41.30	Fostair [EU]

■ BUDESONIDE + FORMOTEROL (EFORMOTEROL)

Note This drug is not PBS-subsidised for the treatment of chronic obstructive pulmonary disease (COPD) or for allergen-induced or exercise-induced bronchoconstriction in the absence of asthma.

Note A LABA includes olodaterol, indacaterol, salmeterol, formoterol or vilanterol.

Authority required (STREAMLINED)

10482

Mild asthma

Clinical criteria:

- Patient must have asthma and require an anti-inflammatory reliever therapy, **AND**
- Patient must not be on a concomitant single agent long-acting-beta-2-agonist (LABA).

Population criteria:

- Patient must be aged 12 years or over.

Device (inhaler) technique should be reviewed at each clinical visit and before initiating treatment with this medicine.

budesonide 100 microgram/actuation + formoterol (eformoterol) fumarate dihydrate 3 microgram/actuation inhalation, 120 actuations

12042T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±2	2	..	*48.40	41.30	Symbicort Rapihaler 100/3 [AP]

■ BUDESONIDE + FORMOTEROL (EFORMOTEROL)

Note Pharmaceutical benefits that have the brand DuoResp Spiromax 200/6 powder for inhalation, 120 actuations and pharmaceutical benefits that have the brand Symbicort Turbuhaler 200/6 powder for inhalation, 120 actuations are equivalent for the purposes of substitution.

Note Patient must be aged 18 years or older.

Note This drug is not PBS-subsidised for the treatment of chronic obstructive pulmonary disease (COPD) or for allergen-induced or exercise-induced bronchoconstriction in the absence of asthma.

Note A LABA includes olodaterol, indacaterol, salmeterol, formoterol or vilanterol.

Authority required (STREAMLINED)

10464

Mild asthma

Clinical criteria:

- Patient must have asthma and require an anti-inflammatory reliever therapy, **AND**
- Patient must not be on a concomitant single agent long-acting-beta-2-agonist (LABA).

Device (inhaler) technique should be reviewed at each clinical visit and before initiating treatment with this medicine.

budesonide 200 microgram/actuation + formoterol (eformoterol) fumarate dihydrate 6 microgram/actuation powder for inhalation, 120 actuations

12029D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	2	..	43.94	41.30	^a DuoResp Spiromax [AF]

■ BUDESONIDE + FORMOTEROL (EFORMOTEROL)

Note Pharmaceutical benefits that have the brand DuoResp Spiromax 200/6 powder for inhalation, 120 actuations and pharmaceutical benefits that have the brand Symbicort Turbuhaler 200/6 powder for inhalation, 120 actuations are equivalent for the purposes of substitution.

Note Patient must be aged 12 years or over.

Note This drug is not PBS-subsidised for the treatment of chronic obstructive pulmonary disease (COPD) or for allergen-induced or exercise-induced bronchoconstriction in the absence of asthma.

Note A LABA includes olodaterol, indacaterol, salmeterol, formoterol or vilanterol.

Authority required (STREAMLINED)**10464**

Mild asthma

Clinical criteria:

- Patient must have asthma and require an anti-inflammatory reliever therapy, **AND**
- Patient must not be on a concomitant single agent long-acting-beta-2-agonist (LABA).
Device (inhaler) technique should be reviewed at each clinical visit and before initiating treatment with this medicine.

budesonide 200 microgram/actuation + formoterol (eformoterol) fumarate dihydrate 6 microgram/actuation powder for inhalation, 120 actuations

12041R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	2	..	43.94	41.30	^a Symbicort Turbuhaler 200/6 [AP]

■ BUDESONIDE + FORMOTEROL (EFORMOTEROL)

Note This product is not indicated for the initiation of treatment in asthma

Note This drug is not PBS-subsidised for the treatment of chronic obstructive pulmonary disease (COPD).

Note The patient must not be on a concomitant single agent long-acting-beta-2-agonist (LABA)

Note A LABA includes olodaterol, indacaterol, salmeterol, formoterol or vilanterol.

Note Adherence to current treatment and device (inhaler) technique should be reviewed at each clinical visit and before "stepping up" a patient's medication regimen.

Authority required (STREAMLINED)**4380**

Asthma

Clinical criteria:

- Patient must have previously had frequent episodes of asthma while receiving treatment with oral corticosteroids or optimal doses of inhaled corticosteroids; OR
- Patient must have experienced frequent asthma symptoms while receiving treatment with oral or inhaled corticosteroids and require single maintenance and reliever therapy; OR
- Patient must have experienced frequent asthma symptoms while receiving treatment with a combination of an inhaled corticosteroid and long acting beta-2 agonist and require single maintenance and reliever therapy.

Population criteria:

- Patient must be aged 12 years or over.

budesonide 100 microgram/actuation + formoterol (eformoterol) fumarate dihydrate 6 microgram/actuation powder for inhalation, 120 actuations

8796Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	48.60	41.30	Symbicort Turbuhaler 100/6 [AP]

■ BUDESONIDE + FORMOTEROL (EFORMOTEROL)

Note This product is not indicated for the initiation of treatment in asthma

Note This drug is not PBS-subsidised for the treatment of chronic obstructive pulmonary disease (COPD).

Note The patient must not be on a concomitant single agent long-acting-beta-2-agonist (LABA)

Note A LABA includes olodaterol, indacaterol, salmeterol, formoterol or vilanterol.

Note Adherence to current treatment and device (inhaler) technique should be reviewed at each clinical visit and before "stepping up" a patient's medication regimen.

Authority required (STREAMLINED)**4397**

Asthma

Clinical criteria:

- Patient must have previously had frequent episodes of asthma while receiving treatment with oral corticosteroids or optimal doses of inhaled corticosteroids; OR
- Patient must have experienced frequent asthma symptoms while receiving treatment with oral or inhaled corticosteroids and require single maintenance and reliever therapy; OR

- Patient must have experienced frequent asthma symptoms while receiving treatment with a combination of an inhaled corticosteroid and long acting beta-2 agonist.

Population criteria:

- Patient must be aged 12 years or over.

budesonide 100 microgram/actuation + formoterol (eformoterol) fumarate dihydrate 3 microgram/actuation inhalation, 120 actuations

10015D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*48.40	41.30	Symbicort Rapihaler 100/3 [AP]

budesonide 50 microgram/actuation + formoterol (eformoterol) fumarate dihydrate 3 microgram/actuation inhalation, 120 actuations

10024N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*46.78	41.30	Symbicort Rapihaler 50/3 [AP]

■ BUDESONIDE + FORMOTEROL (EFORMOTEROL)

Note Unlike Symbicort Turbuhaler 200/6, Symbicort Rapihaler 200/6 is not recommended nor PBS-subsidised for use as 'maintenance and reliever' therapy as the approved Product Information does not specify such use.

Note This product is not indicated for the initiation of treatment in asthma

Note The patient must not be on a concomitant single agent long-acting-beta-2-agonist (LABA)

Note A LABA includes olodaterol, indacaterol, salmeterol, formoterol or vilanterol.

Note Adherence to current treatment and device (inhaler) technique should be reviewed at each clinical visit and before "stepping up" a patient's medication regimen.

Authority required (STREAMLINED)**10538**

Asthma

Clinical criteria:

- Patient must have failed PBS-subsidised fluticasone propionate and salmeterol as a fixed dose combination for this condition.

Treatment criteria:

- Must be treated by a respiratory physician; OR
- Must be treated by a paediatrician.

budesonide 200 microgram/actuation + formoterol (eformoterol) fumarate dihydrate 6 microgram/actuation inhalation, 120 actuations

12082X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±2	5	..	*71.96	41.30	Symbicort Rapihaler 200/6 [AP]

■ BUDESONIDE + FORMOTEROL (EFORMOTEROL)

Note This product is not indicated for the initiation of treatment in asthma

Note This drug is not PBS-subsidised for the treatment of chronic obstructive pulmonary disease (COPD).

Note The patient must not be on a concomitant single agent long-acting-beta-2-agonist (LABA)

Note A LABA includes olodaterol, indacaterol, salmeterol, formoterol or vilanterol.

Note Adherence to current treatment and device (inhaler) technique should be reviewed at each clinical visit and before "stepping up" a patient's medication regimen.

Authority required (STREAMLINED)**10538**

Asthma

Clinical criteria:

- Patient must have failed PBS-subsidised fluticasone propionate and salmeterol as a fixed dose combination for this condition.

Treatment criteria:

- Must be treated by a respiratory physician; OR
- Must be treated by a paediatrician.

budesonide 200 microgram/actuation + formoterol (eformoterol) fumarate dihydrate 6 microgram/actuation powder for inhalation, 120 actuations

12093L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	±1	5	..	43.94	41.30	^a DuoResp Spiromax [AF]	^a Symbicort Turbuhaler 200/6 [AP]

budesonide 100 microgram/actuation + formoterol (eformoterol) fumarate dihydrate 3 microgram/actuation inhalation, 120 actuations

12089G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±2	5	..	*48.40	41.30	Symbicort Rapihaler 100/3 [AP]

budesonide 50 microgram/actuation + formoterol (eformoterol) fumarate dihydrate 3 microgram/actuation inhalation, 120 actuations

12100W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±2	5	..	*46.78	41.30	Symbicort Rapihaler 50/3 [AP]

budesonide 100 microgram/actuation + formoterol (eformoterol) fumarate dihydrate 6 microgram/actuation powder for inhalation, 120 actuations

12101X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	5	..	48.60	41.30	Symbicort Turbuhaler 100/6 [AP]

■ BUDESONIDE + FORMOTEROL (EFORMOTEROL)

Note Pharmaceutical benefits that have the brand DuoResp Spiromax 200/6 powder for inhalation, 120 actuations and pharmaceutical benefits that have the brand Symbicort Turbuhaler 200/6 powder for inhalation, 120 actuations are equivalent for the purposes of substitution.

Note Patient must be aged 18 years or older.

Note This product is not indicated for the initiation of treatment in asthma

Note This drug is not PBS-subsidised for the treatment of chronic obstructive pulmonary disease (COPD).

Note The patient must not be on a concomitant single agent long-acting-beta-2-agonist (LABA)

Note A LABA includes olodaterol, indacaterol, salmeterol, formoterol or vilanterol.

Note Adherence to current treatment and device (inhaler) technique should be reviewed at each clinical visit and before "stepping up" a patient's medication regimen.

Authority required (STREAMLINED)**7970**

Asthma

Clinical criteria:

- Patient must have previously had frequent episodes of asthma while receiving treatment with oral corticosteroids or optimal doses of inhaled corticosteroids; OR
- Patient must have experienced frequent asthma symptoms while receiving treatment with oral or inhaled corticosteroids and require single maintenance and reliever therapy; OR
- Patient must have experienced frequent asthma symptoms while receiving treatment with a combination of an inhaled corticosteroid and long acting beta-2 agonist and require single maintenance and reliever therapy.

budesonide 200 microgram/actuation + formoterol (eformoterol) fumarate dihydrate 6 microgram/actuation powder for inhalation, 120 actuations

11273H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	5	..	43.94	41.30	^a DuoResp Spiromax [AF]

NP

■ BUDESONIDE + FORMOTEROL (EFORMOTEROL)

Note Pharmaceutical benefits that have the brand DuoResp Spiromax 200/6 powder for inhalation, 120 actuations and pharmaceutical benefits that have the brand Symbicort Turbuhaler 200/6 powder for inhalation, 120 actuations are equivalent for the purposes of substitution.

Note Patient must be aged 12 years or over.

Note This product is not indicated for the initiation of treatment in asthma

Note This drug is not PBS-subsidised for the treatment of chronic obstructive pulmonary disease (COPD).

Note The patient must not be on a concomitant single agent long-acting-beta-2-agonist (LABA)

Note A LABA includes olodaterol, indacaterol, salmeterol, formoterol or vilanterol.

Note Adherence to current treatment and device (inhaler) technique should be reviewed at each clinical visit and before "stepping up" a patient's medication regimen.

Authority required (STREAMLINED)**7970**

Asthma

Clinical criteria:

- Patient must have previously had frequent episodes of asthma while receiving treatment with oral corticosteroids or optimal doses of inhaled corticosteroids; OR
- Patient must have experienced frequent asthma symptoms while receiving treatment with oral or inhaled corticosteroids and require single maintenance and reliever therapy; OR
- Patient must have experienced frequent asthma symptoms while receiving treatment with a combination of an inhaled corticosteroid and long acting beta-2 agonist and require single maintenance and reliever therapy.

budesonide 200 microgram/actuation + formoterol (eformoterol) fumarate dihydrate 6 microgram/actuation powder for inhalation, 120 actuations

8625Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	5	..	43.94	41.30	^a Symbicort Turbuhaler 200/6 [AP]

NP

■ BUDESONIDE + FORMOTEROL (EFORMOTEROL)**Authority required (STREAMLINED)****4404**

Asthma

Clinical criteria:

- Patient must have previously had frequent episodes of asthma while receiving treatment with oral corticosteroids or optimal doses of inhaled corticosteroids.

Population criteria:

- Patient must be aged 12 years or over.

Note Unlike Symbicort Turbuhaler 200/6, Symbicort Rapihaler 200/6 is not recommended nor PBS-subsidised for use as 'maintenance and reliever' therapy as the approved Product Information does not specify such use.

Note This product is not indicated for the initiation of treatment in asthma

Note The patient must not be on a concomitant single agent long-acting-beta-2-agonist (LABA)

Note A LABA includes olodaterol, indacaterol, salmeterol, formoterol or vilanterol.

Note Adherence to current treatment and device (inhaler) technique should be reviewed at each clinical visit and before "stepping up" a patient's medication regimen.

Authority required (STREAMLINED)**10121**

Chronic obstructive pulmonary disease (COPD)

Clinical criteria:

- Patient must have significant symptoms despite regular beta-2 agonist bronchodilator therapy, **AND**
- Patient must have experienced at least one severe COPD exacerbation, which required hospitalisation, or two or more moderate exacerbations in the previous 12 months.

Note This product is not indicated for the initiation of bronchodilator therapy in COPD.

Note The treatment must not be used in combination with LABA monotherapy or LAMA/LABA combination therapy.

Note A LAMA/LABA includes aclidinium/formoterol, glycopyrronium/indacaterol, tiotropium/olodaterol, or umeclidinium/vilanterol.

Note Diagnosis of COPD should include measurement of airflow obstruction using spirometry, with confirmation of post-bronchodilator airflow obstruction.

budesonide 200 microgram/actuation + formoterol (eformoterol) fumarate dihydrate 6 microgram/actuation inhalation, 120 actuations

10018G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*71.96	41.30	Symbicort Rapihaler 200/6 [AP]

■ BUDESONIDE + FORMOTEROL (EFORMOTEROL)

Note Pharmaceutical benefits that have the brand DuoResp Spiromax 400/12 powder for inhalation, 2 x 60 actuations and pharmaceutical benefits that have the brand Symbicort Turbuhaler 400/12 powder for inhalation, 2 x 60 actuations are equivalent for the purposes of substitution.

Authority required (STREAMLINED)**7979**

Asthma

Clinical criteria:

- Patient must have previously had frequent episodes of asthma while receiving treatment with oral corticosteroids or optimal doses of inhaled corticosteroids.

Note Patient must be aged 18 years or older.

Note Symbicort 400/12 is not recommended nor PBS-subsidised for use as 'maintenance and reliever' therapy.

Note This product is not indicated for the initiation of treatment in asthma

Note The patient must not be on a concomitant single agent long-acting-beta-2-agonist (LABA)

Note A LABA includes olodaterol, indacaterol, salmeterol, formoterol or vilanterol.

Note Adherence to current treatment and device (inhaler) technique should be reviewed at each clinical visit and before "stepping up" a patient's medication regimen.

Authority required (STREAMLINED)**10121**

Chronic obstructive pulmonary disease (COPD)

Clinical criteria:

- Patient must have significant symptoms despite regular beta-2 agonist bronchodilator therapy, **AND**
- Patient must have experienced at least one severe COPD exacerbation, which required hospitalisation, or two or more moderate exacerbations in the previous 12 months.

Note This product is not indicated for the initiation of bronchodilator therapy in COPD.

Note The treatment must not be used in combination with LABA monotherapy or LAMA/LABA combination therapy.

Note A LAMA/LABA includes aclidinium/formoterol, glycopyrronium/indacaterol, tiotropium/olodaterol, or umeclidinium/vilanterol.

Note Diagnosis of COPD should include measurement of airflow obstruction using spirometry, with confirmation of post-bronchodilator airflow obstruction.

budesonide 400 microgram/actuation + formoterol (eformoterol) fumarate dihydrate 12 microgram/actuation powder for inhalation, 2 x 60 actuations

11301T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	64.60	41.30	^a DuoResp Spiromax [AF]

budesonide 400 microgram/actuation + formoterol (eformoterol) fumarate dihydrate 12 microgram/actuation powder for inhalation, 2 x 60 actuations

8750M

Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
±1	5	..	64.60	41.30	^a Symbicort Turbuhaler 400/12 [AP]

NP

■ FLUTICASONE FUROATE + VILANTEROL**Note** This drug is not recommended nor PBS-subsidised for use as 'maintenance and reliever' therapy.**Note** This drug is not PBS-subsidised for the treatment of chronic obstructive pulmonary disease (COPD).**Note** This product is not indicated for the initiation of treatment in asthma**Note** The patient must not be on a concomitant single agent long-acting-beta-2-agonist (LABA)**Note** A LABA includes olodaterol, indacaterol, salmeterol, formoterol or vilanterol.**Note** Adherence to current treatment and device (inhaler) technique should be reviewed at each clinical visit and before "stepping up" a patient's medication regimen.**Authority required (STREAMLINED)****4731**

Asthma

Clinical criteria:

- Patient must have previously had frequent episodes of asthma while receiving treatment with oral corticosteroids or optimal doses of inhaled corticosteroids.

Population criteria:

- Patient must be aged 12 years or over.

fluticasone furoate 200 microgram/actuation + vilanterol 25 microgram/actuation powder for inhalation, 30 actuations

11129R

Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
±1	5	..	72.55	41.30	Breo Ellipta 200/25 [GK]

NP

■ FLUTICASONE FUROATE + VILANTEROL**Authority required (STREAMLINED)****4711**

Asthma

Clinical criteria:

- Patient must have previously had frequent episodes of asthma while receiving treatment with oral corticosteroids or optimal doses of inhaled corticosteroids.

Population criteria:

- Patient must be aged 12 years or over.

Note This drug is not recommended nor PBS-subsidised for use as 'maintenance and reliever' therapy.**Note** This product is not indicated for the initiation of treatment in asthma**Note** The patient must not be on a concomitant single agent long-acting-beta-2-agonist (LABA)**Note** A LABA includes olodaterol, indacaterol, salmeterol, formoterol or vilanterol.**Note** Adherence to current treatment and device (inhaler) technique should be reviewed at each clinical visit and before "stepping up" a patient's medication regimen.**Authority required (STREAMLINED)****10121**

Chronic obstructive pulmonary disease (COPD)

Clinical criteria:

- Patient must have significant symptoms despite regular beta-2 agonist bronchodilator therapy, **AND**
- Patient must have experienced at least one severe COPD exacerbation, which required hospitalisation, or two or more moderate exacerbations in the previous 12 months.

Note This product is not indicated for the initiation of bronchodilator therapy in COPD.**Note** The treatment must not be used in combination with LABA monotherapy or LAMA/LABA combination therapy.**Note** A LAMA/LABA includes aclidinium/formoterol, glycopyrronium/indacaterol, tiotropium/olodaterol, or umeclidinium/vilanterol.**Note** Diagnosis of COPD should include measurement of airflow obstruction using spirometry, with confirmation of post-bronchodilator airflow obstruction.**fluticasone furoate 100 microgram/actuation + vilanterol 25 microgram/actuation powder for inhalation, 30 actuations**

11124L

Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
±1	5	..	57.05	41.30	Breo Ellipta 100/25 [GK]

NP

■ FLUTICASONE PROPIONATE + FORMOTEROL (EFORMOTEROL)**Note** Flutiform is not recommended nor PBS-subsidised for use as 'maintenance and reliever' therapy.**Note** Flutiform is not indicated or PBS-subsidised for bronchodilator therapy in COPD.**Note** This product is not indicated for the initiation of treatment in asthma**Note** The patient must not be on a concomitant single agent long-acting-beta-2-agonist (LABA)**Note** A LABA includes olodaterol, indacaterol, salmeterol, formoterol or vilanterol.

Note Adherence to current treatment and device (inhaler) technique should be reviewed at each clinical visit and before "stepping up" a patient's medication regimen.

Authority required (STREAMLINED)

4395

Asthma

Clinical criteria:

- Patient must have previously had frequent episodes of asthma while receiving treatment with oral corticosteroids or optimal doses of inhaled corticosteroids.

Population criteria:

- Patient must be aged 12 years or over.

fluticasone propionate 50 microgram/actuation + formoterol (eformoterol) fumarate dihydrate 5 microgram/actuation inhalation, 120 actuations

2827T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	37.84	39.13	flutiform 50/5 [MF]

fluticasone propionate 250 microgram/actuation + formoterol (eformoterol) fumarate dihydrate 10 microgram/actuation inhalation, 120 actuations

10008R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	60.61	41.30	flutiform 250/10 [MF]

fluticasone propionate 125 microgram/actuation + formoterol (eformoterol) fumarate dihydrate 5 microgram/actuation inhalation, 120 actuations

10007Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	45.70	41.30	flutiform 125/5 [MF]

■ **FLUTICASONE PROPIONATE + SALMETEROL**

Note This product is not indicated for the initiation of treatment in asthma

Note This drug is not PBS-subsidised for the treatment of chronic obstructive pulmonary disease (COPD).

Note The patient must not be on a concomitant single agent long-acting-beta-2-agonist (LABA)

Note A LABA includes olodaterol, indacaterol, salmeterol, formoterol or vilanterol.

Note Adherence to current treatment and device (inhaler) technique should be reviewed at each clinical visit and before "stepping up" a patient's medication regimen.

Authority required (STREAMLINED)

4930

Asthma

Clinical criteria:

- Patient must have previously had frequent episodes of asthma while receiving treatment with oral corticosteroids or optimal doses of inhaled corticosteroids.

Population criteria:

- Patient must be aged 4 years or older.

fluticasone propionate 50 microgram/actuation + salmeterol 25 microgram/actuation inhalation, 120 actuations

8517G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	±1	5	..	42.49	41.30	^a PAVTIDE MDI 50/25 [GC]	^a Seretide MDI 50/25 [GK]

fluticasone propionate 250 microgram/actuation + salmeterol 50 microgram/actuation powder for inhalation, 60 actuations

8431R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	±1	5	..	49.18	41.30	^a PAVTIDE ACCUHALER 250/50 [GC]	^a Seretide Accuhaler 250/50 [GK]

fluticasone propionate 125 microgram/actuation + salmeterol 25 microgram/actuation inhalation, 120 actuations

8518H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	±1	5	..	49.18	41.30	^a Fluticasone + Salmeterol Cipla 125/25 [LR]	^a Pavtide [GC]
						^a SalplusF Inhaler 125/25 [SZ]	^a Seroflo 125/25 [YC]
			^b 4.00	53.18	41.30	^a Seretide MDI 125/25 [GK]	

fluticasone propionate 100 microgram/actuation + salmeterol 50 microgram/actuation powder for inhalation, 60 actuations

8430Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	±1	5	..	42.49	41.30	^a PAVTIDE ACCUHALER 100/50 [GC]	^a Seretide Accuhaler 100/50 [GK]

■ **FLUTICASONE PROPIONATE + SALMETEROL**

Authority required (STREAMLINED)

4930

Asthma

Clinical criteria:

- Patient must have previously had frequent episodes of asthma while receiving treatment with oral corticosteroids or optimal doses of inhaled corticosteroids.

Population criteria:

- Patient must be aged 4 years or older.

Note This product is not indicated for the initiation of treatment in asthma

Note The patient must not be on a concomitant single agent long-acting-beta-2-agonist (LABA)

Note A LABA includes olodaterol, indacaterol, salmeterol, formoterol or vilanterol.

Note Adherence to current treatment and device (inhaler) technique should be reviewed at each clinical visit and before "stepping up" a patient's medication regimen.

Authority required (STREAMLINED)**10121**

Chronic obstructive pulmonary disease (COPD)

Clinical criteria:

- Patient must have significant symptoms despite regular beta-2 agonist bronchodilator therapy, **AND**
- Patient must have experienced at least one severe COPD exacerbation, which required hospitalisation, or two or more moderate exacerbations in the previous 12 months.

Note This product is not indicated for the initiation of bronchodilator therapy in COPD.

Note The treatment must not be used in combination with LABA monotherapy or LAMA/LABA combination therapy.

Note A LAMA/LABA includes aclidinium/formoterol, glycopyrronium/indacaterol, tiotropium/olodaterol, or umeclidinium/vilanterol.

Note Diagnosis of COPD should include measurement of airflow obstruction using spirometry, with confirmation of post-bronchodilator airflow obstruction.

fluticasone propionate 250 microgram/actuation + salmeterol 25 microgram/actuation inhalation, 120 actuations

8519J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	±1	5	..	62.32	41.30	^a Fluticasone + Salmeterol Cipla 250/25 [LR]	^a Pavtide [GC]
						^a SalplusF Inhaler 250/25 [SZ]	^a SeroFlo 250/25 [YC]
			^B 4.00	66.32	41.30	^a Seretide MDI 250/25 [GK]	

fluticasone propionate 500 microgram/actuation + salmeterol 50 microgram/actuation powder for inhalation, 60 actuations

8432T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	±1	5	..	62.32	41.30	^a PAVTIDE ACCUHALER 500/50 [GC]	^a Seretide Accuhaler 500/50 [GK]

■ INDACATEROL + MOMETASONE

Note This product is not indicated for the initiation of treatment in asthma

Note This product is not PBS-subsidised for the treatment of chronic obstructive pulmonary disease (COPD).

Note The patient must not be on a concomitant single agent long-acting-beta-2-agonist (LABA)

Note A LABA includes olodaterol, indacaterol, salmeterol, formoterol or vilanterol.

Note Adherence to current treatment and device (inhaler) technique should be reviewed at each clinical visit and before "stepping up" a patient's medication regimen.

Authority required (STREAMLINED)**11360**

Asthma

Clinical criteria:

- Patient must have previously had frequent episodes of asthma while receiving treatment with oral corticosteroids or optimal doses of inhaled corticosteroids.

Population criteria:

- Patient must be aged 12 years or over.

indacaterol 125 microgram + mometasone furoate 62.5 microgram powder for inhalation, 30 capsules

12269R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	37.84	39.13	Aectura Breezhaler [NV]

indacaterol 125 microgram + mometasone furoate 260 microgram powder for inhalation, 30 capsules

12279G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	60.61	41.30	Aectura Breezhaler [NV]

indacaterol 125 microgram + mometasone furoate 127.5 microgram powder for inhalation, 30 capsules

12289T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	43.94	41.30	Aectura Breezhaler [NV]

Adrenergics in combination with anticholinergics incl. triple combinations with corticosteroids

■ ACLIDINIUM + FORMOTEROL (EFORMOTEROL)

Note This product is not PBS-subsidised for the treatment of asthma.

Note This product is not indicated for the initiation of bronchodilator therapy in COPD.

Note The treatment must not be used in combination with an ICS/LABA, LAMA, LABA, or SAMA

Note A LAMA includes tiotropium, glycopyrronium, aclidinium or umeclidinium.

Note A LABA includes olodaterol, indacaterol, salmeterol, formoterol or vilanterol.

Note A SAMA includes ipratropium

Note Diagnosis of COPD should include measurement of airflow obstruction using spirometry, with confirmation of post-bronchodilator airflow obstruction.

Note Adherence to current treatment and device (inhaler) technique should be reviewed at each clinical visit and before "stepping up" a patient's medication regimen.

Authority required (STREAMLINED)

7798

Chronic obstructive pulmonary disease (COPD)

Clinical criteria:

- Patient must have COPD symptoms that persist despite regular bronchodilator treatment with a long acting muscarinic antagonist (LAMA); OR
- Patient must have COPD symptoms that persist despite regular bronchodilator treatment with a long acting beta 2 agonist (LABA); OR
- Patient must have been stabilised on a combination of a LAMA and a LABA.

aclidinium 340 microgram/actuation + formoterol (eformoterol) fumarate dihydrate 12 microgram/actuation powder for inhalation, 60 actuations

10565C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	85.61	41.30	Brimica Genuair [FK]

■ FLUTICASONE FUROATE + UMECLIDINIUM + VILANTEROL

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note Formal assessment and correction of inhaler technique should be performed in accordance with the COPD-X Plan (available at <http://copdx.org.au/>); the assessment and adherence to correct technique should be documented in the patient's medical records.

Note Diagnosis of COPD should include measurement of airflow obstruction using spirometry, with confirmation of post-bronchodilator airflow obstruction.

Note This product is not PBS-subsidised for the treatment of asthma or the initiation of bronchodilator therapy in COPD.

Note The treatment must not be used in combination with an ICS/LABA, LABA/LAMA or LAMA, LABA or ICS monotherapy.

Note A LAMA includes tiotropium, glycopyrronium, aclidinium or umeclidinium.

Note A LABA includes olodaterol, indacaterol, salmeterol, formoterol or vilanterol.

Note An ICS includes fluticasone propionate, fluticasone furoate, budesonide, beclometasone or ciclesonide.

Authority required (STREAMLINED)

10167

Chronic obstructive pulmonary disease (COPD)

Clinical criteria:

- Patient must have experienced at least one severe COPD exacerbation, which required hospitalisation, or two or more moderate exacerbations in the previous 12 months, with significant symptoms despite regular bronchodilator therapy with a long acting muscarinic antagonist (LAMA) and a long acting beta-2 agonist (LABA) or an inhaled corticosteroid (ICS) and a LABA; OR
- Patient must have been stabilised on a combination of a LAMA, LABA and an ICS for this condition.

fluticasone furoate 100 microgram/actuation + umeclidinium 62.5 microgram/actuation + vilanterol 25 microgram/actuation powder for inhalation, 30 actuations

11379X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	94.57	41.30	Trelegy Ellipta 100/62.5/25 [GK]

■ INDACATEROL + GLYCOPYRRONIUM

Note This product is not PBS-subsidised for the treatment of asthma.

Note This product is not indicated for the initiation of bronchodilator therapy in COPD.

Note The treatment must not be used in combination with an ICS/LABA, LAMA, LABA, or SAMA

Note A LAMA includes tiotropium, glycopyrronium, aclidinium or umeclidinium.

Note A LABA includes olodaterol, indacaterol, salmeterol, formoterol or vilanterol.

Note A SAMA includes ipratropium

Note Diagnosis of COPD should include measurement of airflow obstruction using spirometry, with confirmation of post-bronchodilator airflow obstruction.

Note Adherence to current treatment and device (inhaler) technique should be reviewed at each clinical visit and before "stepping up" a patient's medication regimen.

Authority required (STREAMLINED)

7798

Chronic obstructive pulmonary disease (COPD)

Clinical criteria:

- Patient must have COPD symptoms that persist despite regular bronchodilator treatment with a long acting muscarinic antagonist (LAMA); OR
- Patient must have COPD symptoms that persist despite regular bronchodilator treatment with a long acting beta 2 agonist (LABA); OR
- Patient must have been stabilised on a combination of a LAMA and a LABA.

indacaterol 110 microgram + glycopyrronium 50 microgram powder for inhalation, 30 capsules

10156M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	89.42	41.30	ultibro breezhaler 110/50 [NV]

■ INDACATEROL + GLYCOPYRRONIUM + MOMETASONE

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 888 333.

Note Formal assessment and correction of inhaler technique should be performed in accordance with the National Asthma Council (NAC) Information Paper for Health Professionals on Inhaler Technique (available at www.nationalasthma.org.au); the assessment and adherence to correct technique should be documented in the patient's medical records. Patients can obtain support with inhaler technique through their local Asthma Foundation (1800 645 130).

Note This drug is not PBS-subsidised for the treatment of chronic obstructive pulmonary disease (COPD).

Note This product is not indicated for the initiation of treatment in asthma

Note The treatment must not be used in combination with an ICS/LABA, LABA/LAMA or LAMA, LABA or ICS monotherapy.

Note A LAMA includes tiotropium, glycopyrronium, aclidinium or umeclidinium.

Note A LABA includes olodaterol, indacaterol, salmeterol, formoterol or vilanterol.

Note An ICS includes fluticasone propionate, fluticasone furoate, budesonide, beclometasone or ciclesonide.

Authority required

Severe asthma

Clinical criteria:

- Patient must have experienced at least one severe exacerbation, which has required documented use of systemic corticosteroids, in the previous 12 months while receiving optimised asthma therapy, despite formal assessment of and adherence to correct inhaler technique, which has been documented.

Population criteria:

- Patient must be aged 18 years or older.

Optimised asthma therapy includes adherence to the maintenance combination of an inhaled corticosteroid (at least 800 micrograms budesonide per day or equivalent) and a long acting beta-2 agonist.

indacaterol 114 microgram + glycopyrronium 46 microgram + mometasone furoate 136 microgram powder for inhalation, 30 capsules

12295D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	91.60	41.30	Enerzair Breezhaler [NV]

indacaterol 114 microgram + glycopyrronium 46 microgram + mometasone furoate 68 microgram powder for inhalation, 30 capsules

12298G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	74.93	41.30	Enerzair Breezhaler [NV]

■ TIOTROPIUM + OLODATEROL

Note Ensure that the patient understands that medication cartridges in the refill only pack are for the re-usable inhaler device only, or that medication cartridges supplied in the pack containing the non-reusable inhaler device are for use with the non-reusable device only.

Note This product is not PBS-subsidised for the treatment of asthma.

Note This product is not indicated for the initiation of bronchodilator therapy in COPD.

Note The treatment must not be used in combination with an ICS/LABA, LABA, LABA, or SAMA

Note A LAMA includes tiotropium, glycopyrronium, aclidinium or umeclidinium.

Note A LABA includes olodaterol, indacaterol, salmeterol, formoterol or vilanterol.

Note A SAMA includes ipratropium

Note Diagnosis of COPD should include measurement of airflow obstruction using spirometry, with confirmation of post-bronchodilator airflow obstruction.

Note Adherence to current treatment and device (inhaler) technique should be reviewed at each clinical visit and before "stepping up" a patient's medication regimen.

Authority required (STREAMLINED)

7798

Chronic obstructive pulmonary disease (COPD)

Clinical criteria:

- Patient must have COPD symptoms that persist despite regular bronchodilator treatment with a long acting muscarinic antagonist (LAMA); OR
- Patient must have COPD symptoms that persist despite regular bronchodilator treatment with a long acting beta 2 agonist (LABA); OR
- Patient must have been stabilised on a combination of a LAMA and a LABA.

tiotropium 2.5 microgram/actuation + olodaterol 2.5 microgram/actuation inhalation solution, 60 actuations

10557P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	73.60	41.30	Spiolto Respimat [BY]

■ UMECLIDINIUM + VILANTEROL

Note This product is not PBS-subsidised for the treatment of asthma.

Note This product is not indicated for the initiation of bronchodilator therapy in COPD.

Note The treatment must not be used in combination with an ICS/LABA, LAMA, LABA, or SAMA

Note A LAMA includes tiotropium, glycopyrronium, aclidinium or umeclidinium.

Note A LABA includes olodaterol, indacaterol, salmeterol, formoterol or vilanterol.

Note A SAMA includes ipratropium

Note Diagnosis of COPD should include measurement of airflow obstruction using spirometry, with confirmation of post-bronchodilator airflow obstruction.

Note Adherence to current treatment and device (inhaler) technique should be reviewed at each clinical visit and before "stepping up" a patient's medication regimen.

Authority required (STREAMLINED)**7798**

Chronic obstructive pulmonary disease (COPD)

Clinical criteria:

- Patient must have COPD symptoms that persist despite regular bronchodilator treatment with a long acting muscarinic antagonist (LAMA); OR
- Patient must have COPD symptoms that persist despite regular bronchodilator treatment with a long acting beta 2 agonist (LABA); OR
- Patient must have been stabilised on a combination of a LAMA and a LABA.

umeclidinium 62.5 microgram/actuation + vilanterol 25 microgram/actuation powder for inhalation, 30 actuations

10188F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	89.41	41.30	Anoro Ellipta 62.5/25 [GK]

OTHER DRUGS FOR OBSTRUCTIVE AIRWAY DISEASES, INHALANTS*Glucocorticoids***■ BECLOMETASONE****beclometasone dipropionate 100 microgram/actuation inhalation, 200 actuations**

8407L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	31.13	32.42	Qvar 100 [IL]

beclometasone dipropionate 50 microgram/actuation inhalation, 200 actuations

8406K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	21.12	22.41	Qvar 50 [IL]

■ BECLOMETASONE**Restricted benefit**

Asthma

Clinical criteria:

- Patient must be unable to achieve co-ordinated use of other metered dose inhalers containing this drug.

beclometasone dipropionate 50 microgram/actuation breath activated inhalation, 200 actuations

8408M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	27.17	28.46	Qvar 50 Autohaler [IL]

beclometasone dipropionate 100 microgram/actuation breath activated inhalation, 200 actuations

8409N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	35.13	36.42	Qvar 100 Autohaler [IL]

■ BUDESONIDE**budesonide 200 microgram/actuation powder for inhalation, 200 actuations**

2071B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	29.43	30.72	Pulmicort Turbuhaler [AP]

budesonide 400 microgram/actuation powder for inhalation, 200 actuations

2072C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	40.39	41.30	Pulmicort Turbuhaler [AP]

budesonide 100 microgram/actuation powder for inhalation, 200 actuations

2070Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	23.98	25.27	Pulmicort Turbuhaler [AP]

■ BUDESONIDE**Authority required (STREAMLINED)****6340**

Severe chronic asthma

Clinical criteria:

- Patient must require long-term steroid therapy, **AND**
- Patient must not be able to use other forms of inhaled steroid therapy.

budesonide 500 microgram/2 mL inhalation solution, 30 x 2 mL ampoules

2065Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	34.23	35.52	Pulmicort Respules [AP]

budesonide 1 mg/2 mL inhalation solution, 30 x 2 mL ampoules

2066R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	42.95	41.30	Pulmicort Respules [AP]

■ CICLESONIDE**ciclesonide 80 microgram/actuation inhalation, 120 actuations**

8853Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	25.95	27.24	Alvesco 80 [EU]

ciclesonide 160 microgram/actuation inhalation, 120 actuations

8854B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	37.46	38.75	Alvesco 160 [EU]

■ FLUTICASONE FUROATE**fluticasone furoate 100 microgram/actuation powder for inhalation, 30 actuations**

11719T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	28.85	30.14	Arnuity Ellipta [GK]

fluticasone furoate 200 microgram/actuation powder for inhalation, 30 actuations

11729H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	42.99	41.30	Arnuity Ellipta [GK]

■ FLUTICASONE PROPIONATE**fluticasone propionate 250 microgram/actuation inhalation, 120 actuations**

8346G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	±1	1	..	42.99	41.30	^a Axotide [GC]	^a Fluticasone Cipla Inhaler [LR]
			^b 3.00	45.99	41.30	^a Flixotide [GK]	

fluticasone propionate 125 microgram/actuation inhalation, 120 actuations

8345F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	±1	5	..	28.85	30.14	^a Axotide [GC]	^a Fluticasone Cipla Inhaler [LR]
			^b 3.00	31.85	30.14	^a Flixotide [GK]	

fluticasone propionate 500 microgram/actuation powder for inhalation, 60 actuations

8149X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	1	..	42.99	41.30	Flixotide Accuhaler [GK]

fluticasone propionate 50 microgram/actuation inhalation, 120 actuations

8516F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	±1	5	..	19.43	20.72	^a Axotide Junior [GC]	^a Flixotide Junior [GK]

fluticasone propionate 250 microgram/actuation powder for inhalation, 60 actuations

8148W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	±1	5	..	28.85	30.14	^a Axotide Accuhaler [GC]	^a Flixotide Accuhaler [GK]

fluticasone propionate 100 microgram/actuation powder for inhalation, 60 actuations

8147T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	±1	5	..	19.43	20.72	^a Axotide Junior Accuhaler [GC]	^a Flixotide Junior Accuhaler [GK]

Anticholinergics

■ **ACLIDINIUM**

Note The treatment must not be used in combination with a LAMA/LABA or SAMA

Note A LAMA/LABA includes aclidinium/formoterol, glycopyrronium/indacaterol, tiotropium/olodaterol, or umeclidinium/vilanterol.

Note A SAMA includes ipratropium

Note Diagnosis of COPD should include measurement of airflow obstruction using spirometry, with confirmation of post-bronchodilator airflow obstruction.

Note Adherence to current treatment and device (inhaler) technique should be reviewed at each clinical visit and before "stepping up" a patient's medication regimen.

Restricted benefit

Chronic obstructive pulmonary disease (COPD)

aclidinium 322 microgram/actuation inhalation: powder for, 60 actuations

10124W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	60.35	41.30	Bretaris Genuair [FK]

■ **GLYCOPYRRONIUM**

Note The treatment must not be used in combination with a LAMA/LABA or SAMA

Note A LAMA/LABA includes aclidinium/formoterol, glycopyrronium/indacaterol, tiotropium/olodaterol, or umeclidinium/vilanterol.

Note A SAMA includes ipratropium

Note Diagnosis of COPD should include measurement of airflow obstruction using spirometry, with confirmation of post-bronchodilator airflow obstruction.

Note Adherence to current treatment and device (inhaler) technique should be reviewed at each clinical visit and before "stepping up" a patient's medication regimen.

Restricted benefit

Chronic obstructive pulmonary disease (COPD)

glycopyrronium 50 microgram powder for inhalation, 30 capsules

10059K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	60.35	41.30	seebri breezhaler [NV]

■ **IPRATROPIUM****ipratropium bromide monohydrate 21 microgram/actuation inhalation, 200 actuations**

8671J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*35.86	37.15	Atrovent [BY]

■ **IPRATROPIUM****Restricted benefit**

Asthma

Clinical criteria:

- Patient must be unable to use this drug delivered from an oral pressurised inhalation device via a spacer.

Restricted benefit

Chronic obstructive pulmonary disease (COPD)

Clinical criteria:

- Patient must be unable to use this drug delivered from an oral pressurised inhalation device via a spacer.

Note Diagnosis of COPD should include measurement of airflow obstruction using spirometry, with confirmation of post-bronchodilator airflow obstruction.

Note Adherence to current treatment and device (inhaler) technique should be reviewed at each clinical visit and before "stepping up" a patient's medication regimen.

ipratropium bromide 500 microgram/mL inhalation solution, 30 x 1 mL ampoules

8238N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	2	5	..	*31.20	32.49	^a Aeron 500 [AL]	^a APO-Ipratropium [TX]
						^a Ipratrin Adult [AF]	
			^b 0.50	*31.70	32.49	^a Atrovent Adult [BY]	

ipratropium bromide 250 microgram/mL inhalation solution, 30 x 1 mL ampoules

1542E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	2	5	..	*28.26	29.55	^a Aeron 250 [AL]	^a APO-Ipratropium [TX]
						^a Ipratrin [AF]	
			^b 0.50	*28.76	29.55	^a Atrovent [BY]	

■ **TIOTROPIUM**

Note Ensure that the patient understands that medication cartridges in the refill only pack are for the re-usable inhaler device only, or that medication cartridges supplied in the pack containing the non-reusable inhaler device are for use with the non-reusable device only.

Note Formal assessment and correction of inhaler technique should be performed in accordance with the National Asthma Council (NAC) Information Paper for Health Professionals on Inhaler Technique (available at www.humanservices.gov.au or

www.nationalasthma.org.au); the assessment and adherence to correct technique should be documented in the patient's medical records. Patients can obtain support with inhaler technique through their local Asthma Foundation (1800 645 130).

Restricted benefit

Severe asthma

Clinical criteria:

- Patient must have experienced at least one severe exacerbation, which has required documented use of systemic corticosteroids, in the previous 12 months while receiving optimised asthma therapy, despite formal assessment of and adherence to correct inhaler technique, which has been documented, **AND**
- The treatment must be used in combination with a maintenance combination of an inhaled corticosteroid (ICS) and a long acting beta-2 agonist (LABA) unless a LABA is contraindicated.

Population criteria:

- Patient must be aged 18 years or older.

Optimised asthma therapy includes adherence to the maintenance combination of an inhaled corticosteroid (at least 800 micrograms budesonide per day or equivalent) and a long acting beta-2 agonist.

tiotropium 2.5 microgram/actuation inhalation solution, 60 actuations

11043F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	43.01	41.30	Spiriva Respimat [BY]

■ TIOTROPIUM

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note Ensure that the patient understands that medication cartridges in the refill only pack are for the re-usable inhaler device only, or that medication cartridges supplied in the pack containing the non-reusable inhaler device are for use with the non-reusable device only.

Note Formal assessment and correction of inhaler technique should be performed in accordance with the National Asthma Council (NAC) Information Paper for Health Professionals on Inhaler Technique (available at www.humanservices.gov.au or www.nationalasthma.org.au); the assessment and adherence to correct technique should be documented in the patient's medical records. Patients can obtain support with inhaler technique through their local Asthma Foundation (1800 645 130).

Note Adherence to current treatment and device (inhaler) technique should be reviewed at each clinical visit and before "stepping up" a patient's medication regimen.

Authority required (STREAMLINED)

8606

Severe asthma

Treatment criteria:

- Must be treated by a respiratory physician, paediatric respiratory physician, clinical immunologist, allergist, paediatrician or general physician experienced in the management of patients with severe asthma; or in consultation with one of these specialists.

Clinical criteria:

- Patient must have failed to achieve adequate control with optimised asthma therapy, despite formal assessment of and adherence to correct inhaler technique, which has been documented, **AND**
- Patient must have experienced at least one severe exacerbation prior to receiving PBS-subsidised treatment with this drug for this condition, which has required documented use of systemic corticosteroids in the previous 12 months while receiving optimised asthma therapy; OR
- Patient must have experienced frequent episodes of moderate asthma exacerbations prior to receiving PBS-subsidised treatment with this drug for this condition, **AND**
- The treatment must be used in combination with a maintenance combination of an inhaled corticosteroid (ICS) and a long acting beta-2 agonist (LABA) unless a LABA is contraindicated.

Population criteria:

- Patient must be aged 6 to 17 years inclusive.

Optimised asthma therapy includes adherence to the maintenance combination of a medium to high dose ICS and a LABA. If LABA therapy is contraindicated, not tolerated or not effective, montelukast, cromoglycate or nedocromil may be used as an alternative

tiotropium 2.5 microgram/actuation inhalation solution, 60 actuations

11629C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	43.01	41.30	Spiriva Respimat [BY]

■ TIOTROPIUM

Note Ensure that the patient understands that medication cartridges in the refill only pack are for the re-usable inhaler device only, or that medication cartridges supplied in the pack containing the non-reusable inhaler device are for use with the non-reusable device only.

Note The treatment must not be used in combination with a LAMA/LABA or SAMA

Note A LAMA/LABA includes aclidinium/formoterol, glycopyrronium/indacaterol, tiotropium/olodaterol, or umeclidinium/vilanterol.

Note A SAMA includes ipratropium

Note Diagnosis of COPD should include measurement of airflow obstruction using spirometry, with confirmation of post-bronchodilator airflow obstruction.

RESPIRATORY SYSTEM

Note Adherence to current treatment and device (inhaler) technique should be reviewed at each clinical visit and before "stepping up" a patient's medication regimen.

Restricted benefit

Bronchospasm and dyspnoea associated with chronic obstructive pulmonary disease

Treatment Phase: Long-term maintenance treatment

tiotropium 2.5 microgram/actuation inhalation solution, 60 actuations

10509D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	43.01	41.30	Spiriva Respimat [BY]

■ TIOTROPIUM

Note Pharmaceutical benefits that have the form tiotropium 18 microgram powder for inhalation and pharmaceutical benefits that have the form tiotropium 13 microgram powder for inhalation are equivalent for the purposes of substitution.

Note The treatment must not be used in combination with a LAMA/LABA or SAMA

Note A LAMA/LABA includes acclidinium/formoterol, glycopyrronium/indacaterol, tiotropium/olodaterol, or umeclidinium/vilanterol.

Note A SAMA includes ipratropium

Note Diagnosis of COPD should include measurement of airflow obstruction using spirometry, with confirmation of post-bronchodilator airflow obstruction.

Note Adherence to current treatment and device (inhaler) technique should be reviewed at each clinical visit and before "stepping up" a patient's medication regimen.

Restricted benefit

Chronic obstructive pulmonary disease (COPD)

tiotropium 13 microgram powder for inhalation, 30 capsules

11892X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	43.01	41.30	^a Braltus [AF]

tiotropium 18 microgram powder for inhalation, 30 capsules

8626B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	43.01	41.30	^a Spiriva [BY]

■ UMECLIDINIUM

Note The treatment must not be used in combination with a LAMA/LABA or SAMA

Note A LAMA/LABA includes acclidinium/formoterol, glycopyrronium/indacaterol, tiotropium/olodaterol, or umeclidinium/vilanterol.

Note A SAMA includes ipratropium

Note Diagnosis of COPD should include measurement of airflow obstruction using spirometry, with confirmation of post-bronchodilator airflow obstruction.

Note Adherence to current treatment and device (inhaler) technique should be reviewed at each clinical visit and before "stepping up" a patient's medication regimen.

Restricted benefit

Chronic obstructive pulmonary disease (COPD)

umeclidinium 62.5 microgram/actuation inhalation: powder for, 30 actuations

10187E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	60.35	41.30	Incruse Ellipta [GK]

Antiallergic agents, excl. corticosteroids

■ CROMOGLYCAT

sodium cromoglycate 1 mg/actuation inhalation, 200 actuations

8767K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	35.57	36.86	Intal CFC-Free [SW]

sodium cromoglycate 5 mg/actuation inhalation, 112 actuations

8334P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	39.75	41.04	Intal Forte CFC-Free [SW]

■ NEDOCROMIL

nedocromil sodium 2 mg/actuation inhalation, 112 actuations

8365G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	35.64	36.93	Tilade CFC-Free [SW]

ADRENERGICS FOR SYSTEMIC USE

Alpha- and beta-adrenoreceptor agonists

■ ADRENALINE (EPINEPHRINE)

adrenaline (epinephrine) 1 in 1000 (1 mg/mL) injection, 5 x 1 mL ampoules

1016L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	1	..	21.68	22.97	Link Medical Products Pty Ltd [LM]

adrenaline (epinephrine) 1 in 1000 (1 mg/mL) injection, 5 x 1 mL ampoules

5004J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	1	..	..	21.68	22.97	Link Medical Products Pty Ltd [LM]

■ ADRENALINE (EPINEPHRINE)

Note The auto-injector should be provided in the framework of a comprehensive anaphylaxis prevention program and an emergency action plan including training in recognition of the symptoms of anaphylaxis and the use of the auto-injector device. (For further information see the Australasian Society of Clinical Immunology and Allergy website at www.allergy.org.au.)

Note Authority approvals will be limited to a maximum quantity of 2 auto-injectors at any one time.

Note No applications for repeats will be authorised.

Authority required

Acute allergic reaction with anaphylaxis

Treatment Phase: Initial sole PBS-subsidised supply for anticipated emergency treatment

Clinical criteria:

- Patient must have been assessed to be at significant risk of anaphylaxis by, or in consultation with a clinical immunologist; OR
- Patient must have been assessed to be at significant risk of anaphylaxis by, or in consultation with an allergist; OR
- Patient must have been assessed to be at significant risk of anaphylaxis by, or in consultation with a paediatrician; OR
- Patient must have been assessed to be at significant risk of anaphylaxis by, or in consultation with a respiratory physician.

The name of the specialist consulted must be provided at the time of application for initial supply.

Authority required

Acute allergic reaction with anaphylaxis

Treatment Phase: Initial sole PBS-subsidised supply for anticipated emergency treatment

Clinical criteria:

- Patient must have been discharged from hospital or an emergency department after treatment with adrenaline (epinephrine) for acute allergic reaction with anaphylaxis.

Authority required

Acute allergic reaction with anaphylaxis

Treatment Phase: Continuing sole PBS-subsidised supply for anticipated emergency treatment

Clinical criteria:

- Patient must have previously been issued with an authority prescription for this drug.

adrenaline (epinephrine) 150 microgram/0.3 mL injection, 0.3 mL pen device

8697R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	84.26	41.30	^a Adrenaline Jr Mylan [AF]	^a EpiPen Jr. [AL]

adrenaline (epinephrine) 300 microgram/0.3 mL injection, 0.3 mL pen device

8698T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	84.26	41.30	^a Adrenaline Mylan [AF]	^a EpiPen [AL]

Selective beta-2-adrenoreceptor agonists

■ SALBUTAMOL

salbutamol 2 mg/5 mL oral liquid, 150 mL

1103C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*25.74	27.03	Ventolin [GK]

■ TERBUTALINE

terbutaline sulfate 500 microgram/mL injection, 5 x 1 mL ampoules

1034K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	29.09	30.38	Bricanyl [AP]

OTHER SYSTEMIC DRUGS FOR OBSTRUCTIVE AIRWAY DISEASES

Xanthines

■ THEOPHYLLINE

Caution Because of variable effects of food on absorption of sustained release theophylline preparations, patients stabilised on one brand should not be changed to another without appropriate monitoring.

Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a

patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

theophylline 250 mg modified release tablet, 100

2634P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	16.96	18.25	Nuelin-SR 250 [IL]

theophylline 133.3 mg/25 mL oral liquid, 500 mL

2614N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	16.88	18.17	Nuelin [IL]

theophylline 200 mg modified release tablet, 100

8230E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	16.20	17.49	Nuelin-SR 200 [IL]

theophylline 300 mg modified release tablet, 100

8231F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	17.88	19.17	Nuelin-SR 300 [IL]

Leukotriene receptor antagonists**■ MONTELUKAST**

Note This drug is not PBS-subsidised for use in a child aged 2 to 5 years with moderate to severe asthma. It is not intended as an alternative for a child aged 2 to 5 years who requires a corticosteroid as a preventer medication.

Note This drug is not subsidised in a child aged 2 to 5 years for use in combination with other preventer medications. PBS subsidy for this drug will therefore cease for a child aged 2 to 5 years who requires a preventer medication in addition to this drug.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)**6666**

Asthma

Treatment Phase: First-line prevention

Population criteria:

- Patient must be aged 2 to 5 years inclusive.

Clinical criteria:

- The condition must be frequent intermittent; OR
- The condition must be mild persistent, **AND**
- The treatment must be the single preventer agent, **AND**
- The treatment must be an alternative to sodium cromoglycate; OR
- The treatment must be an alternative to nedocromil sodium.

montelukast 4 mg chewable tablet, 28

8627C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	19.17	20.46	^a APO-Montelukast [TX] ^a MONTELAIR 4 [RF] ^a Montelukast APOTEX [GX] ^a Montelukast Lupin [HQ] ^a Montelukast Sandoz 4 [SZ]	^a Auro-Montelukast Tabs 4 [DO] ^a Montelukast AN [EA] ^a Montelukast GH [GQ] ^a Montelukast Mylan [AF] ^a Respikast 4 [RW]

■ MONTELUKAST

Note This drug is not PBS-subsidised for use in a patient aged 15 years or older, or for use in addition to a long-acting beta-agonist in any age group, or for use as a single second line preventer, as an alternative to corticosteroids, in a child aged 6 to 14 years with moderate to severe asthma.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)**6674**

Asthma

Treatment Phase: First-line prevention

Clinical criteria:

- The condition must be frequent intermittent; OR
- The condition must be mild persistent, **AND**
- The treatment must be the single preventer agent, **AND**
- The treatment must be an alternative to sodium cromoglycate; OR
- The treatment must be an alternative to nedocromil sodium.

Population criteria:

- Patient must be aged 6 to 14 years inclusive.

Authority required (STREAMLINED)

7781

Asthma

Treatment Phase: Prevention of condition

Clinical criteria:

- The condition must be exercise-induced, **AND**
- The treatment must be as an alternative to adding salmeterol xinafoate; OR
- The treatment must be an alternative to adding formoterol fumarate, **AND**
- The condition must be otherwise well controlled while receiving optimal dose inhaled corticosteroid, **AND**
- Patient must require short-acting beta-2 agonist 3 or more times per week for prevention or relief of residual exercise-related symptoms.

Population criteria:

- Patient must be aged 6 to 14 years inclusive.

montelukast 5 mg chewable tablet, 28

8628D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	18.75	20.04	^a APO-Montelukast [TX] ^a MONTELAIR 5 [RF] ^a Montelukast APOTEX [GX] ^a Montelukast Mylan [AF] ^a Respikast 5 [RW]	^a Auro-Montelukast Tabs 5 [DO] ^a Montelukast AN [EA] ^a Montelukast Lupin [HQ] ^a Montelukast Sandoz 5 [SZ]

■ ANTIHISTAMINES FOR SYSTEMIC USE
ANTIHISTAMINES FOR SYSTEMIC USE
Phenothiazine derivatives
■ PROMETHAZINE
promethazine hydrochloride 50 mg/2 mL injection, 5 x 2 mL ampoules

1948M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	..	..	*35.60	36.89	DBL Promethazine Hydrochloride [PF]

■ SENSORY ORGANS
■ OPHTHALMOLOGICALS
ANTIINFECTIVES
Antibiotics
■ AZITHROMYCIN
Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Restricted benefit

Trachoma

azithromycin 500 mg tablet, 2

8336R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	2	..	16.41	17.70	^a APO-Azithromycin [TX] ^a Azithromycin Sandoz [SZ] ^a Zithromax [PF]	^a Azithromycin Mylan [AF] ^a ZITHRO [RW]

azithromycin 200 mg/5 mL powder for oral liquid, 15 mL

8201P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	‡1	..	..	#27.36	29.02	Zithromax [PF]

■ CHLORAMPHENICOL
Restricted benefit

For treatment of a patient identifying as Aboriginal or Torres Strait Islander

chloramphenicol 0.5% eye drops, 10 mL

11112W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
OP NP MW	‡1	2	..	16.13	17.42	Chlorsig [AS]

■ GENTAMICIN
Restricted benefit

Perioperative use in ophthalmic surgery

Restricted benefit

Suspected Pseudomonas eye infection

gentamicin 0.3% eye drops, 5 mL

5566Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
OP	±1	2	..	22.34	23.63	Genoptic [AG]

■ GENTAMICIN

Restricted benefit

Invasive ocular infection

Restricted benefit

Perioperative use in ophthalmic surgery

Restricted benefit

Suspected Pseudomonal eye infection

gentamicin 0.3% eye drops, 5 mL

1441W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	2	..	22.34	23.63	Genoptic [AG]

■ TOBRAMYCIN

Restricted benefit

Perioperative use in ophthalmic surgery

Restricted benefit

Suspected Pseudomonal eye infection

tobramycin 0.3% eye ointment, 3.5 g

5570E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
OP	±1	..	..	25.20	26.49	Tobrex [NV]

tobramycin 0.3% eye drops, 5 mL

5569D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
OP	±1	2	..	22.64	23.93	Tobrex [NV]

■ TOBRAMYCIN

Restricted benefit

Invasive ocular infection

Restricted benefit

Perioperative use in ophthalmic surgery

Restricted benefit

Suspected Pseudomonal eye infection

tobramycin 0.3% eye ointment, 3.5 g

2329N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	25.20	26.49	Tobrex [NV]

tobramycin 0.3% eye drops, 5 mL

2328M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	2	..	22.64	23.93	Tobrex [NV]

Antivirals

■ ACICLOVIR

Restricted benefit

Herpes simplex keratitis

aciclovir 3% eye ointment, 4.5 g

11654J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
OP	±1	..	..	65.24	41.30	Acivision [DZ]

■ ACICLOVIR

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Herpes simplex keratitis

aciclovir 3% eye ointment, 4.5 g

11652G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	..	..	65.24	41.30	Acivision [DZ]

Fluoroquinolones

■ CIPROFLOXACIN

Authority required

Bacterial keratitis

Treatment criteria:

- Must be treated by an ophthalmologist or in consultation with an ophthalmologist.

ciprofloxacin 0.3% eye drops, 5 mL

1217C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*31.20	32.49	^a CiloQuin [NM]
			^B 4.36	*35.56	32.49	^a Ciloxan [NV]

■ CIPROFLOXACIN

Authority required

Bacterial keratitis

Treatment criteria:

- Must be treated by an ophthalmologist or in consultation with an ophthalmologist.

ciprofloxacin 0.3% eye drops, 5 mL

5564W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
OP	2	..	..	*31.20	32.49	^a CiloQuin [NM]
			^B 4.36	*35.56	32.49	^a Ciloxan [NV]

■ OFLOXACIN

Authority required

Bacterial keratitis

Treatment criteria:

- Must be treated by an ophthalmologist or in consultation with an ophthalmologist.

ofloxacin 0.3% eye drops, 5 mL

5567B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
OP	2	..	..	*32.42	33.71	Ocuflax [AG]

■ OFLOXACIN

Authority required

Bacterial keratitis

Treatment criteria:

- Must be treated by an ophthalmologist or in consultation with an ophthalmologist.

ofloxacin 0.3% eye drops, 5 mL

8383F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*32.42	33.71	Ocuflax [AG]

ANTIINFLAMMATORY AGENTS

Corticosteroids, plain

■ DEXAMETHASONE

dexamethasone 0.1% eye drops, 5 mL

1288T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	2	..	15.82	17.11	Maxidex [NV]

■ DEXAMETHASONE

Authority required

Non-infectious posterior segment uveitis

Treatment criteria:

- Must be treated by an ophthalmologist or in consultation with an ophthalmologist.

Clinical criteria:

- Patient must have documented visual impairment defined as a best corrected visual acuity score of approximate Snellen equivalent 6/12 or worse in the eye proposed for treatment, secondary to vitreous haze or macular oedema, **AND**
- Patient must have unilateral, asymmetric or bilateral flare-up where systemic treatment or further intensification of systemic treatment is not clinically indicated.

dexamethasone 700 microgram implant, 1

11317P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	1360.00	41.30	Ozurdex [AG]

■ DEXAMETHASONE

Note No applications for increased maximum quantities will be authorised.

Note No applications for increased repeats will be authorised.

dexamethasone 0.1% eye drops, 5 mL

5565X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
OP	‡1	..	..	15.82	17.11	Maxidex [NV]

■ DEXAMETHASONE

Note Special Pricing Arrangements apply.

Note No increase in the maximum number of repeats may be authorised.

Note No increase in the maximum quantity or number of units may be authorised for applications for treatment of one eye.

Note Where both eyes are affected by the condition, a quantity of 2 units can be requested on the same authority prescription form.

Authority required

Diabetic macular oedema (DMO)

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by an ophthalmologist or by an accredited ophthalmology registrar in consultation with an ophthalmologist.

Clinical criteria:

- Patient must have visual impairment due to diabetic macular oedema, **AND**
- Patient must have documented visual impairment defined as a best corrected visual acuity score between 78 and 39 letters based on the early treatment diabetic retinopathy study chart administered at a distance of 4 metres (approximate Snellen equivalent 20/32 to 20/160), in the eye proposed for treatment, **AND**
- The condition must be diagnosed by optical coherence tomography; OR
- The condition must be diagnosed by fluorescein angiography, **AND**
- Patient must have had a cataract removed in the treated eye; OR
- Patient must be scheduled for cataract surgery in the treated eye, **AND**
- Patient must have a contraindication to vascular endothelial growth factor (VEGF) inhibitors; OR
- Patient must be unsuitable for treatment with VEGF inhibitors; OR
- Patient must have failed prior treatment with VEGF inhibitors, **AND**
- The treatment must be as monotherapy; OR
- The treatment must be in combination with laser photocoagulation, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Authority approval for initial treatment of each eye must be sought.

The first authority application for each eye must be made in writing.

A written application must include:

- a) a completed authority prescription form;
- b) a completed Diabetic Macular Oedema (DMO) - PBS Supporting Information Form; and
- c) a copy of the optical coherence tomography or fluorescein angiogram report.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Diabetic macular oedema (DMO)

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by an ophthalmologist or by an accredited ophthalmology registrar in consultation with an ophthalmologist.

Clinical criteria:

- Patient must have had a cataract removed in the treated eye; OR
- Patient must be scheduled for cataract surgery in the treated eye, **AND**
- Patient must have previously been issued with an authority prescription for this drug for the same eye, **AND**
- The treatment must be as monotherapy; OR
- The treatment must be in combination with laser photocoagulation, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

dexamethasone 700 microgram implant, 1

10943Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	1360.00	41.30	Ozurdex [AG]

▪ DEXAMETHASONE

Note Special Pricing Arrangements apply.

Note No increase in the maximum number of repeats may be authorised.

Note No increase in the maximum quantity or number of units may be authorised for applications for treatment of one eye.

Note Where both eyes are affected by the condition, a quantity of 2 units can be requested on the same authority prescription form.

Authority required

Branch retinal vein occlusion with macular oedema

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by an ophthalmologist or by an accredited ophthalmology registrar in consultation with an ophthalmologist.

Clinical criteria:

- Patient must have visual impairment due to macular oedema secondary to branched retinal vein occlusion (BRVO), **AND**
- Patient must have documented visual impairment defined as a best corrected visual acuity score between 73 and 20 letters based on the early treatment diabetic retinopathy study chart administered at a distance of 4 metres (approximate Snellen equivalent 20/40 to 20/400), in the eye proposed for treatment, **AND**
- The condition must be diagnosed by optical coherence tomography; OR
- The condition must be diagnosed by fluorescein angiography, **AND**
- Patient must have a contraindication to vascular endothelial growth factor (VEGF) inhibitors; OR
- Patient must have failed prior treatment with VEGF inhibitors, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Authority approval for initial treatment of each eye must be sought.

The first authority application for each eye must be made in writing.

A written application must include:

- a) a completed authority prescription form;
- b) a completed Retinal Vein Occlusion Initial PBS authority application Supporting information form; and
- c) a copy of the optical coherence tomography or fluorescein angiogram report.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Branch retinal vein occlusion with macular oedema

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by an ophthalmologist or by an accredited ophthalmology registrar in consultation with an ophthalmologist.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition for the same eye, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Central retinal vein occlusion with macular oedema

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by an ophthalmologist or by an accredited ophthalmology registrar in consultation with an ophthalmologist.

Clinical criteria:

- Patient must have visual impairment due to macular oedema secondary to central retinal vein occlusion (CRVO), **AND**
- Patient must have documented visual impairment defined as a best corrected visual acuity score between 73 and 24 letters based on the early treatment diabetic retinopathy study chart administered at a distance of 4 metres (approximate Snellen equivalent 20/40 to 20/320), in the eye proposed for treatment, **AND**
- The condition must be diagnosed by optical coherence tomography; OR

- The condition must be diagnosed by fluorescein angiography, **AND**
- Patient must have a contraindication to vascular endothelial growth factor (VEGF) inhibitors; OR
- Patient must have failed prior treatment with VEGF inhibitors, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Authority approval for initial treatment of each eye must be sought.

The first authority application for each eye must be made in writing.

A written application must include:

- a) a completed authority prescription form;
- b) a completed Retinal Vein Occlusion Initial PBS authority application Supporting information form; and
- c) a copy of the optical coherence tomography or fluorescein angiogram report.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Central retinal vein occlusion with macular oedema

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by an ophthalmologist or by an accredited ophthalmology registrar in consultation with an ophthalmologist.

Clinical criteria:

- Patient must have previously received PBS-subsidised treatment with this drug for this condition for the same eye, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

dexamethasone 700 microgram implant, 1

11469P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	1	..	1360.00	41.30	Ozurdex [AG]

FLUOROMETHOLONE

fluorometholone 0.1% eye drops, 5 mL

1204J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	15.18	16.47	FML Liquifilm [AG]

FLUOROMETHOLONE

Note No applications for increased maximum quantities will be authorised.

Note No applications for increased repeats will be authorised.

fluorometholone 0.1% eye drops, 5 mL

5513E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
OP	±1	..	..	15.18	16.47	FML Liquifilm [AG]

FLUOROMETHOLONE ACETATE

fluorometholone acetate 0.1% eye drops, 5 mL

1438Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	2	..	15.18	16.47	Flarex [NV]

FLUOROMETHOLONE ACETATE

Note No applications for increased maximum quantities will be authorised.

Note No applications for increased repeats will be authorised.

fluorometholone acetate 0.1% eye drops, 5 mL

5533F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
OP	±1	..	..	15.18	16.47	Flarex [NV]

■ HYDROCORTISONE ACETATE

hydrocortisone acetate 1% eye ointment, 5 g

2441L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	..	..	17.50	18.79	Hycor [AS]

■ HYDROCORTISONE ACETATE

Note No applications for increased maximum quantities will be authorised.

Note No applications for increased repeats will be authorised.

hydrocortisone acetate 1% eye ointment, 5 g

5516H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
OP	±1	..	..	17.50	18.79	Hycor [AS]

Corticosteroids and mydriatics in combination

■ PREDNISOLONE ACETATE + PHENYLEPHRINE

Restricted benefit

Severe eye inflammation

Clinical criteria:

- Patient must have had a cataract removed in the treated eye; OR
- Patient must be scheduled for cataract surgery in the treated eye.

Population criteria:

- Patient must identify as Aboriginal or Torres Strait Islander.

prednisolone acetate 1% + phenylephrine hydrochloride 0.12% eye drops, 10 mL

11908R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	28.89	30.18	Prednefrin Forte [AG]

■ PREDNISOLONE ACETATE + PHENYLEPHRINE

Restricted benefit

Corneal grafts

Restricted benefit

Uveitis

prednisolone acetate 1% + phenylephrine hydrochloride 0.12% eye drops, 10 mL

3112T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	2	..	28.89	30.18	Prednefrin Forte [AG]

■ PREDNISOLONE ACETATE + PHENYLEPHRINE

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Restricted benefit

Uveitis

prednisolone acetate 1% + phenylephrine hydrochloride 0.12% eye drops, 10 mL

5568C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
OP	±1	..	..	28.89	30.18	Prednefrin Forte [AG]

ANTIGLAUCOMA PREPARATIONS AND MIOTICS

Sympathomimetics in glaucoma therapy

■ APRACLOMIDINE

Restricted benefit

Intra-ocular pressure

Clinical criteria:

- The treatment must be for short-term reduction of intra-ocular pressure, **AND**
- Patient must already be on maximally tolerated anti-glaucoma therapy.

apraclonidine 0.5% eye drops, 10 mL

8083K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	2	..	37.08	38.37	lopidine 0.5% [NV]

■ BRIMONIDINE

brimonidine tartrate 0.15% eye drops, 5 mL

5298W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	5	..	23.95	25.24	Alphagan P 1.5 [AG]

brimonidine tartrate 0.2% eye drops, 5 mL

8351M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	5	..	23.95	25.24	^a Enidin [PE]
			^B 1.42	25.37	25.24	^a Alphagan [AG]

■ BRIMONIDINE

Note For prescribing in accordance with Optometry Board of Australia guidelines.

brimonidine tartrate 0.15% eye drops, 5 mL

5563T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	5	..	23.95	25.24	Alphagan P 1.5 [AG]

brimonidine tartrate 0.2% eye drops, 5 mL

5534G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	5	..	23.95	25.24	^a Enidin [PE]
			^B 1.42	25.37	25.24	^a Alphagan [AG]

■ BRIMONIDINE + TIMOLOL

Restricted benefit

Elevated intra-ocular pressure

Clinical criteria:

- The condition must have been inadequately controlled with monotherapy, **AND**
- Patient must have open-angle glaucoma; OR
- Patient must have ocular hypertension.

brimonidine tartrate 0.2% + timolol 0.5% eye drops, 5 mL

8826M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	5	..	29.07	30.36	Combigan [AG]

■ BRIMONIDINE + TIMOLOL

Note For prescribing in accordance with Optometry Board of Australia guidelines.

Restricted benefit

Elevated intra-ocular pressure

Clinical criteria:

- The condition must have been inadequately controlled with monotherapy, **AND**
- Patient must have open-angle glaucoma; OR
- Patient must have ocular hypertension.

brimonidine tartrate 0.2% + timolol 0.5% eye drops, 5 mL

5535H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	5	..	29.07	30.36	Combigan [AG]

Parasympathomimetics

■ PILOCARPINE

pilocarpine hydrochloride 1% eye drops, 15 mL

2595N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	5	..	17.37	18.66	Isopto Carpine [NV]

pilocarpine hydrochloride 2% eye drops, 15 mL

2596P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	5	..	18.42	19.71	Isopto Carpine [NV]

pilocarpine hydrochloride 4% eye drops, 15 mL

2598R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	5	..	20.90	22.19	Isopto Carpine [NV]

■ PILOCARPINE

Note For prescribing in accordance with Optometry Board of Australia guidelines.

pilocarpine hydrochloride 1% eye drops, 15 mL

5536J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	5	..	17.37	18.66	Isopto Carpine [NV]

pilocarpine hydrochloride 2% eye drops, 15 mL

5537K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	5	..	18.42	19.71	Isopto Carpine [NV]

pilocarpine hydrochloride 4% eye drops, 15 mL

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
5538L	±1	5	..	20.90	22.19	Isopto Carpine [NV]

OP

Carbonic anhydrase inhibitors
■ ACETAZOLAMIDE
Note Continuing Therapy Only:

For prescribing by nurse practitioners as continuing therapy only, where the treatment of, and prescribing of medicine for, a patient has been initiated by a medical practitioner. Further information can be found in the Explanatory Notes for Nurse Practitioners.

acetazolamide 250 mg tablet, 100

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
1004W	1	3	..	25.55	26.84	Diamox [RW]

NP

■ BRINZOLAMIDE
brinzolamide 1% eye drops, 5 mL

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8483L	±1	5	..	26.24	27.53	^a BrinzoQuin [NM]
			^B 4.78	31.02	27.53	^a Azopt [NV]

■ BRINZOLAMIDE

Note For prescribing in accordance with Optometry Board of Australia guidelines.

brinzolamide 1% eye drops, 5 mL

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
5540N	±1	5	..	26.24	27.53	^a BrinzoQuin [NM]
			^B 4.78	31.02	27.53	^a Azopt [NV]

OP

■ BRINZOLAMIDE + BRIMONIDINE
Restricted benefit

Elevated intra-ocular pressure

Clinical criteria:

- The condition must have been inadequately controlled with monotherapy, **AND**
- Patient must have open-angle glaucoma; OR
- Patient must have ocular hypertension.

brinzolamide 1% + brimonidine tartrate 0.2% eye drops, 5 mL

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
10536M	±1	5	..	28.58	29.87	Simbrinza 1%/0.2% [NV]

■ BRINZOLAMIDE + BRIMONIDINE

Note For prescribing in accordance with Optometry Board of Australia guidelines.

Restricted benefit

Elevated intra-ocular pressure

Clinical criteria:

- The condition must have been inadequately controlled with monotherapy, **AND**
- Patient must have open-angle glaucoma; OR
- Patient must have ocular hypertension.

brinzolamide 1% + brimonidine tartrate 0.2% eye drops, 5 mL

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
10547D	±1	5	..	28.58	29.87	Simbrinza 1%/0.2% [NV]

OP

■ BRINZOLAMIDE + TIMOLOL
Restricted benefit

Elevated intra-ocular pressure

Clinical criteria:

- The condition must have been inadequately controlled with monotherapy, **AND**
- Patient must have open-angle glaucoma; OR
- Patient must have ocular hypertension.

brinzolamide 1% + timolol 0.5% eye drops, 5 mL

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
3438Y	±1	5	..	29.81	31.10	Azarga [NV]

■ BRINZOLAMIDE + TIMOLOL

Note For prescribing in accordance with Optometry Board of Australia guidelines.

Restricted benefit

Elevated intra-ocular pressure

Clinical criteria:

- The condition must have been inadequately controlled with monotherapy, **AND**
- Patient must have open-angle glaucoma; OR
- Patient must have ocular hypertension.

brinzolamide 1% + timolol 0.5% eye drops, 5 mL

5562R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
OP	±1	5	..	29.81	31.10	Azarga [NV]

■ DORZOLAMIDE

dorzolamide 2% eye drops, 5 mL

8488R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	±1	5	..	19.87	21.16	^a APO-Dorzolamide [TX] ^a Trusopt [MF]	^a Trusamide [AF]

■ DORZOLAMIDE

Note For prescribing in accordance with Optometry Board of Australia guidelines.

dorzolamide 2% eye drops, 5 mL

5541P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
OP	±1	5	..	19.87	21.16	^a APO-Dorzolamide [TX] ^a Trusopt [MF]	^a Trusamide [AF]

■ DORZOLAMIDE + TIMOLOL

Restricted benefit

Elevated intra-ocular pressure

Clinical criteria:

- The condition must have been inadequately controlled with monotherapy, **AND**
- Patient must have open-angle glaucoma; OR
- Patient must have ocular hypertension.

dorzolamide 2% + timolol 0.5% eye drops, 5 mL

8567X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	±1	5	..	22.61	23.90	^a APO-Dorzolamide/Timolol 20/5 [TX]	^a Cosdor [AF]
			^B 1.00	23.61	23.90	^a Cosopt [MF]	

■ DORZOLAMIDE + TIMOLOL

Note For prescribing in accordance with Optometry Board of Australia guidelines.

Restricted benefit

Elevated intra-ocular pressure

Clinical criteria:

- The condition must have been inadequately controlled with monotherapy, **AND**
- Patient must have open-angle glaucoma; OR
- Patient must have ocular hypertension.

dorzolamide 2% + timolol 0.5% eye drops, 5 mL

5542Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
OP	±1	5	..	22.61	23.90	^a APO-Dorzolamide/Timolol 20/5 [TX]	^a Cosdor [AF]
			^B 1.00	23.61	23.90	^a Cosopt [MF]	

Beta blocking agents

■ BETAXOLOL

betaxolol 0.5% eye drops, 5 mL

2825Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	5	..	19.28	20.57	^a BetoQuin [NM]
			^B 4.76	24.04	20.57	^a Betoptic [NV]

■ BETAXOLOL

Note For prescribing in accordance with Optometry Board of Australia guidelines.

betaxolol 0.5% eye drops, 5 mL

5544T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
OP	±1	5	..	19.28	20.57	^a BetoQuin [NM]
			^B 4.76	24.04	20.57	^a Betoptic [NV]

■ TIMOLOL
timolol 0.5% eye drops, 2.5 mL

1926J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	5	..	17.81	19.10	Timoptol XE [MF]

timolol 0.5% eye drops, 5 mL

1279H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	5	..	17.75	19.04	Timoptol [MF]

■ TIMOLOL

Note For prescribing in accordance with Optometry Board of Australia guidelines.

timolol 0.5% eye drops, 2.5 mL

5550D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
OP	±1	5	..	17.81	19.10	Timoptol XE [MF]

timolol 0.5% eye drops, 5 mL

5548B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
OP	±1	5	..	17.75	19.04	Timoptol [MF]

Prostaglandin analogues
■ BIMATOPROST
bimatoprost 0.03% eye drops, 3 mL

8620Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	±1	5	..	36.94	38.23	^a APO-Bimatoprost [TX]	^a Bimatoprost Sandoz [SZ]
						^a Bimtop [AF]	^a Lumigan [AG]

bimatoprost 0.03% eye drops, 30 x 0.4 mL unit doses

10046R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	5	..	32.94	34.23	Lumigan PF [AG]

■ BIMATOPROST

Note For prescribing in accordance with Optometry Board of Australia guidelines.

bimatoprost 0.03% eye drops, 3 mL

5551E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
OP	±1	5	..	36.94	38.23	^a APO-Bimatoprost [TX]	^a Bimatoprost Sandoz [SZ]
						^a Bimtop [AF]	^a Lumigan [AG]

bimatoprost 0.03% eye drops, 30 x 0.4 mL unit doses

10053D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
OP	±1	5	..	32.94	34.23	Lumigan PF [AG]

■ BIMATOPROST + TIMOLOL
Restricted benefit

Elevated intra-ocular pressure

Clinical criteria:

- The condition must have been inadequately controlled with monotherapy, **AND**
- Patient must have open-angle glaucoma; OR
- Patient must have ocular hypertension.

bimatoprost 0.03% + timolol 0.5% eye drops, 30 x 0.4 mL unit doses

10107Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	5	..	37.73	39.02	GANfort PF 0.3/5 [AG]

■ BIMATOPROST + TIMOLOL
Restricted benefit

Elevated intra-ocular pressure

Clinical criteria:

- The condition must have been inadequately controlled with monotherapy, **AND**
- Patient must have open-angle glaucoma; OR
- Patient must have ocular hypertension.

bimatoprost 0.03% + timolol 0.5% eye drops, 3 mL

9464D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	5	..	41.39	41.30	Ganfort 0.3/5 [AG]

■ BIMATOPROST + TIMOLOL

Note For prescribing in accordance with Optometry Board of Australia guidelines.

Restricted benefit

Elevated intra-ocular pressure

Clinical criteria:

- The condition must have been inadequately controlled with monotherapy, **AND**
- Patient must have open-angle glaucoma; OR
- Patient must have ocular hypertension.

bimatoprost 0.03% + timolol 0.5% eye drops, 3 mL

5558M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	5	..	41.39	41.30	Ganfort 0.3/5 [AG]

bimatoprost 0.03% + timolol 0.5% eye drops, 30 x 0.4 mL unit doses

10108B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	5	..	37.73	39.02	GANfort PF 0.3/5 [AG]

■ LATANOPROST

latanoprost 0.005% eye drops, 2.5 mL

8243W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	±1	5	..	17.85	19.14	^a APO-Latanoprost [TX] ^a Latanoprost Sandoz [SZ] ^a Xalatan [AS]	^a Latanoprost Actavis [EA] ^a Xalaprost [AF]

■ LATANOPROST

Note For prescribing in accordance with Optometry Board of Australia guidelines.

latanoprost 0.005% eye drops, 2.5 mL

5552F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	±1	5	..	17.85	19.14	^a APO-Latanoprost [TX] ^a Latanoprost Sandoz [SZ] ^a Xalatan [AS]	^a Latanoprost Actavis [EA] ^a Xalaprost [AF]

■ LATANOPROST + TIMOLOL

Restricted benefit

Elevated intra-ocular pressure

Clinical criteria:

- The condition must have been inadequately controlled with monotherapy, **AND**
- Patient must have open-angle glaucoma; OR
- Patient must have ocular hypertension.

latanoprost 0.005% + timolol 0.5% eye drops, 2.5 mL

8895E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	±1	5	..	22.30	23.59	^a APO-Latanoprost/Timolol 0.05/5 [TX] ^a Xalacom [AS]	^a Latanoprost/Timolol Sandoz 50/5 [SZ] ^a Xalamol 50/5 [AF]

■ LATANOPROST + TIMOLOL

Note For prescribing in accordance with Optometry Board of Australia guidelines.

Restricted benefit

Elevated intra-ocular pressure

Clinical criteria:

- The condition must have been inadequately controlled with monotherapy, **AND**
- Patient must have open-angle glaucoma; OR
- Patient must have ocular hypertension.

latanoprost 0.005% + timolol 0.5% eye drops, 2.5 mL

5553G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	±1	5	..	22.30	23.59	^a APO-Latanoprost/Timolol 0.05/5 [TX] ^a Xalacom [AS]	^a Latanoprost/Timolol Sandoz 50/5 [SZ] ^a Xalamol 50/5 [AF]

■ TAFLUPROST

tafluprost 0.0015% eye drops, 30 x 0.3 mL unit doses

2755B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	5	..	34.65	35.94	Saflutan [MF]

■ TAFLUPROST

Note For prescribing in accordance with Optometry Board of Australia guidelines.

tafluprost 0.0015% eye drops, 30 x 0.3 mL unit doses

2748P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	5	..	34.65	35.94	Saflutan [MF]

OP

■ TRAVOPROST

travoprost 0.004% eye drops, 2.5 mL

8597L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	5	..	37.38	38.67	Travatan [NV]

■ TRAVOPROST

Note For prescribing in accordance with Optometry Board of Australia guidelines.

travoprost 0.004% eye drops, 2.5 mL

5554H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	5	..	37.38	38.67	Travatan [NV]

OP

■ TRAVOPROST + TIMOLOL

Restricted benefit

Elevated intra-ocular pressure

Clinical criteria:

- The condition must have been inadequately controlled with monotherapy, **AND**
- Patient must have open-angle glaucoma; OR
- Patient must have ocular hypertension.

travoprost 0.004% + timolol 0.5% eye drops, 2.5 mL

9057Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	5	..	41.84	41.30	Duotrav [NV]

■ TRAVOPROST + TIMOLOL

Note For prescribing in accordance with Optometry Board of Australia guidelines.

Restricted benefit

Elevated intra-ocular pressure

Clinical criteria:

- The condition must have been inadequately controlled with monotherapy, **AND**
- Patient must have open-angle glaucoma; OR
- Patient must have ocular hypertension.

travoprost 0.004% + timolol 0.5% eye drops, 2.5 mL

5555J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	5	..	41.84	41.30	Duotrav [NV]

OP

MYDRIATICS AND CYCLOPLEGICS

Anticholinergics

■ ATROPINE SULFATE

atropine sulfate monohydrate 1% eye drops, 15 mL

1093M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	2	..	22.87	24.16	Atropt [AS]

NP

OCULAR VASCULAR DISORDER AGENTS

Antineovascularisation agents

■ AFLIBERCEPT

Note Special Pricing Arrangements apply.

Note No increase in the maximum number of repeats may be authorised.

Note No increase in the maximum quantity or number of units may be authorised for applications for treatment of one eye.

Note Where both eyes are affected by the condition, a quantity of 2 units can be requested on the same authority prescription form.

Note Pharmaceutical benefits that have the form aflibercept 0.09mL injection syringe and pharmaceutical benefits that have the form aflibercept 0.1mL injection vial are equivalent for the purposes of substitution.

Authority required

Diabetic macular oedema (DMO)

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by an ophthalmologist or by an accredited ophthalmology registrar in consultation with an ophthalmologist.

Clinical criteria:

- Patient must have visual impairment due to diabetic macular oedema, **AND**
- Patient must have documented visual impairment defined as a best corrected visual acuity score between 78 and 39 letters based on the early treatment diabetic retinopathy study chart administered at a distance of 4 metres (approximate Snellen equivalent 20/32 to 20/160), in the eye proposed for treatment, **AND**
- The condition must be diagnosed by optical coherence tomography; OR
- The condition must be diagnosed by fluorescein angiography, **AND**
- The treatment must be as monotherapy; OR
- The treatment must be in combination with laser photocoagulation, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Authority approval for initial treatment of each eye must be sought.

The first authority application for each eye must be made in writing.

A written application must include:

- a completed authority prescription form;
- a completed Diabetic Macular Oedema (DMO) - PBS Supporting Information Form; and
- a copy of the optical coherence tomography or fluorescein angiogram report.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Diabetic macular oedema (DMO)

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by an ophthalmologist or by an accredited ophthalmology registrar in consultation with an ophthalmologist.

Clinical criteria:

- Patient must have previously been issued with an authority prescription for this drug for the same eye, **AND**
- The treatment must be as monotherapy; OR
- The treatment must be in combination with laser photocoagulation, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

aflibercept 4 mg/0.1 mL injection, 0.1 mL vial

10505X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1042.89	41.30	^a Eylea [BN]

aflibercept 3.6 mg/0.09 mL injection, 0.09 mL syringe

12153P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1042.89	41.30	^a Eylea [BN]

■ AFLIBERCEPT

Note Special Pricing Arrangements apply.

Note No increase in the maximum number of repeats may be authorised.

Note No increase in the maximum quantity or number of units may be authorised for applications for treatment of one eye.

Note Where both eyes are affected by the condition, a quantity of 2 units can be requested on the same authority prescription form.

Note Pharmaceutical benefits that have the form aflibercept 0.09mL injection syringe and pharmaceutical benefits that have the form aflibercept 0.1mL injection vial are equivalent for the purposes of substitution.

Authority required

Subfoveal choroidal neovascularisation (CNV)

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by an ophthalmologist or by an accredited ophthalmology registrar in consultation with an ophthalmologist.

Clinical criteria:

- The condition must be due to pathologic myopia (PM), **AND**
- The condition must be diagnosed by optical coherence tomography; OR
- The condition must be diagnosed by fluorescein angiography, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Authority approval for initial treatment of each eye must be sought.

The first authority application for each eye must be made in writing.

A written application must include:

- a) a completed authority prescription form;
- b) a completed Subfoveal Choroidal Neovascularisation (CNV) - PBS Supporting Information Form; and
- c) a copy of the optical coherence tomography or fluorescein angiogram report.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos.

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Subfoveal choroidal neovascularisation (CNV)

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by an ophthalmologist or by an accredited ophthalmology registrar in consultation with an ophthalmologist.

Clinical criteria:

- The condition must be due to pathologic myopia (PM), **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition for the same eye.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

aflibercept 4 mg/0.1 mL injection, 0.1 mL vial

12131L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	1042.89	41.30	^a Eylea [BN]

aflibercept 3.6 mg/0.09 mL injection, 0.09 mL syringe

12141B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	1042.89	41.30	^a Eylea [BN]

▪ **AFLIBERCEPT**

Note Special Pricing Arrangements apply.

Note No increase in the maximum quantity or number of units may be authorised for applications for treatment of one eye.

Note Where both eyes are affected by the condition, a quantity of 2 units can be requested on the same authority prescription form.

Note Pharmaceutical benefits that have the form aflibercept 0.09mL injection syringe and pharmaceutical benefits that have the form aflibercept 0.1mL injection vial are equivalent for the purposes of substitution.

Authority required

Subfoveal choroidal neovascularisation (CNV)

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by an ophthalmologist or by an accredited ophthalmology registrar in consultation with an ophthalmologist.

Clinical criteria:

- The condition must be due to age-related macular degeneration (AMD), **AND**
- The condition must be diagnosed by optical coherence tomography; OR
- The condition must be diagnosed by fluorescein angiography, **AND**

- The treatment must be the sole PBS-subsidised therapy for this condition. Authority approval for initial treatment of each eye must be sought. The first authority application for each eye must be made in writing. A written application must include:

- a) a completed authority prescription form;
- b) a completed Subfoveal Choroidal Neovascularisation (CNV) - PBS Supporting Information Form; and
- c) a copy of the optical coherence tomography or fluorescein angiogram report.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Subfoveal choroidal neovascularisation (CNV)

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by an ophthalmologist or by an accredited ophthalmology registrar in consultation with an ophthalmologist.

Clinical criteria:

- The condition must be due to age-related macular degeneration (AMD), **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have previously been granted an authority prescription for the same eye.

Note No increase in the maximum number of repeats may be authorised.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

afibercept 4 mg/0.1 mL injection, 0.1 mL vial

2168D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	1042.89	41.30	^a Eylea [BN]

afibercept 3.6 mg/0.09 mL injection, 0.09 mL syringe

12152N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	1042.89	41.30	^a Eylea [BN]

▪ AFLIBERCEPT

Note Special Pricing Arrangements apply.

Note No increase in the maximum number of repeats may be authorised.

Note No increase in the maximum quantity or number of units may be authorised for applications for treatment of one eye.

Note Where both eyes are affected by the condition, a quantity of 2 units can be requested on the same authority prescription form.

Note Pharmaceutical benefits that have the form afibercept 0.09mL injection syringe and pharmaceutical benefits that have the form afibercept 0.1mL injection vial are equivalent for the purposes of substitution.

Authority required

Branch retinal vein occlusion with macular oedema

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by an ophthalmologist or by an accredited ophthalmology registrar in consultation with an ophthalmologist.

Clinical criteria:

- Patient must have visual impairment due to macular oedema secondary to branched retinal vein occlusion (BRVO), **AND**
- Patient must have documented visual impairment defined as a best corrected visual acuity score between 73 and 20 letters based on the early treatment diabetic retinopathy study chart administered at a distance of 4 metres (approximate Snellen equivalent 20/40 to 20/400), in the eye proposed for treatment, **AND**
- The condition must be diagnosed by optical coherence tomography; OR
- The condition must be diagnosed by fluorescein angiography, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Authority approval for initial treatment of each eye must be sought.

The first authority application for each eye must be made in writing.

A written application must include:

- a) a completed authority prescription form;

- b) a completed Retinal Vein Occlusion Initial PBS authority application Supporting information form; and
- c) a copy of the optical coherence tomography or fluorescein angiogram report.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Branch retinal vein occlusion with macular oedema
Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by an ophthalmologist or by an accredited ophthalmology registrar in consultation with an ophthalmologist.

Clinical criteria:

- Patient must have previously been issued with an authority prescription for this drug for the same eye, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Central retinal vein occlusion with macular oedema
Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by an ophthalmologist or by an accredited ophthalmology registrar in consultation with an ophthalmologist.

Clinical criteria:

- Patient must have visual impairment due to macular oedema secondary to central retinal vein occlusion (CRVO), **AND**
- Patient must have documented visual impairment defined as a best corrected visual acuity score between 73 and 24 letters based on the early treatment diabetic retinopathy study chart administered at a distance of 4 metres (approximate Snellen equivalent 20/40 to 20/320), in the eye proposed for treatment, **AND**
- The condition must be diagnosed by optical coherence tomography; OR
- The condition must be diagnosed by fluorescein angiography, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Authority approval for initial treatment of each eye must be sought.

The first authority application for each eye must be made in writing.

A written application must include:

- a) a completed authority prescription form;
- b) a completed Retinal Vein Occlusion Initial PBS authority application Supporting information form; and
- c) a copy of the optical coherence tomography or fluorescein angiogram report.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Central retinal vein occlusion with macular oedema
Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by an ophthalmologist or by an accredited ophthalmology registrar in consultation with an ophthalmologist.

Clinical criteria:

- Patient must have previously been issued with an authority prescription for this drug for the same eye, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

aflibercept 4 mg/0.1 mL injection, 0.1 mL vial

11991D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	1042.89	41.30	^a Eylea [BN]

aflibercept 3.6 mg/0.09 mL injection, 0.09 mL syringe

12132M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	1042.89	41.30	^a Eylea [BN]

■ **RANIBIZUMAB**

Note Special Pricing Arrangements apply.

Note No increase in the maximum number of repeats may be authorised.

Note No increase in the maximum quantity or number of units may be authorised for applications for treatment of one eye.

Note Where both eyes are affected by the condition, a quantity of 2 units can be requested on the same authority prescription form.

Note Pharmaceutical benefits that have the form ranibizumab 0.165 mL injection syringe and pharmaceutical benefits that have the form ranibizumab 0.23 mL injection vial are equivalent for the purposes of substitution.

Authority required

Subfoveal choroidal neovascularisation (CNV)

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by an ophthalmologist or by an accredited ophthalmology registrar in consultation with an ophthalmologist.

Clinical criteria:

- The condition must be due to age-related macular degeneration (AMD), **AND**
- The condition must be diagnosed by optical coherence tomography; OR
- The condition must be diagnosed by fluorescein angiography, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Authority approval for initial treatment of each eye must be sought.

The first authority application for each eye must be made in writing.

A written application must include:

- a) a completed authority prescription form;
- b) a completed Subfoveal Choroidal Neovascularisation (CNV) - PBS Supporting Information Form; and
- c) a copy of the optical coherence tomography or fluorescein angiogram report.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos. Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Subfoveal choroidal neovascularisation (CNV)

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by an ophthalmologist or by an accredited ophthalmology registrar in consultation with an ophthalmologist.

Clinical criteria:

- The condition must be due to age-related macular degeneration (AMD), **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have previously been granted an authority prescription for the same eye.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

ranibizumab 1.65 mg/0.165 mL injection, 0.165 mL syringe

10138N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	1042.89	41.30	^a Lucentis [NV]

ranibizumab 2.3 mg/0.23 mL injection, 0.23 mL vial

1382R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	1042.89	41.30	^a Lucentis [NV]

▪ RANIBIZUMAB

Note Special Pricing Arrangements apply.

Note No increase in the maximum number of repeats may be authorised.

Note No increase in the maximum quantity or number of units may be authorised for applications for treatment of one eye.

Note Where both eyes are affected by the condition, a quantity of 2 units can be requested on the same authority prescription form.

Note Pharmaceutical benefits that have the form ranibizumab 0.165 mL injection syringe and pharmaceutical benefits that have the form ranibizumab 0.23 mL injection vial are equivalent for the purposes of substitution.

Authority required

Diabetic macular oedema (DMO)

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by an ophthalmologist or by an accredited ophthalmology registrar in consultation with an ophthalmologist.

Clinical criteria:

- Patient must have visual impairment due to diabetic macular oedema, **AND**
- Patient must have documented visual impairment defined as a best corrected visual acuity score between 78 and 39 letters based on the early treatment diabetic retinopathy study chart administered at a distance of 4 metres (approximate Snellen equivalent 20/32 to 20/160), in the eye proposed for treatment, **AND**
- The condition must be diagnosed by optical coherence tomography; OR
- The condition must be diagnosed by fluorescein angiography, **AND**
- The treatment must be as monotherapy; OR
- The treatment must be in combination with laser photocoagulation, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Authority approval for initial treatment of each eye must be sought.

The first authority application for each eye must be made in writing.

A written application must include:

- a completed authority prescription form;
- a completed Diabetic Macular Oedema (DMO) - PBS Supporting Information Form; and
- a copy of the optical coherence tomography or fluorescein angiogram report.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia

Complex Drugs

Reply Paid 9826

HOBART TAS 7001

Authority required

Diabetic macular oedema (DMO)

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by an ophthalmologist or by an accredited ophthalmology registrar in consultation with an ophthalmologist.

Clinical criteria:

- Patient must have previously been issued with an authority prescription for this drug for the same eye, **AND**
- The treatment must be as monotherapy; OR
- The treatment must be in combination with laser photocoagulation, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

ranibizumab 1.65 mg/0.165 mL injection, 0.165 mL syringe

10374B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1042.89	41.30	^a Lucentis [NV]

ranibizumab 2.3 mg/0.23 mL injection, 0.23 mL vial

10373Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	1042.89	41.30	^a Lucentis [NV]

▪ RANIBIZUMAB

Note Special Pricing Arrangements apply.

Note No increase in the maximum quantity or number of units may be authorised for applications for treatment of one eye.

Note Where both eyes are affected by the condition, a quantity of 2 units can be requested on the same authority prescription form.

Note Pharmaceutical benefits that have the form ranibizumab 0.165 mL injection syringe and pharmaceutical benefits that have the form ranibizumab 0.23 mL injection vial are equivalent for the purposes of substitution.

Authority required

Subfoveal choroidal neovascularisation (CNV)

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by an ophthalmologist or by an accredited ophthalmology registrar in consultation with an ophthalmologist.

Clinical criteria:

- The condition must be due to pathologic myopia (PM), **AND**
- The condition must be diagnosed by optical coherence tomography; OR
- The condition must be diagnosed by fluorescein angiography, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Authority approval for initial treatment of each eye must be sought.

The first authority application for each eye must be made in writing.

A written application must include:

- a) a completed authority prescription form;
- b) a completed Subfoveal Choroidal Neovascularisation (CNV) - PBS Supporting Information Form; and
- c) a copy of the optical coherence tomography or fluorescein angiogram report.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au. Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos. Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Subfoveal choroidal neovascularisation (CNV)

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by an ophthalmologist or by an accredited ophthalmology registrar in consultation with an ophthalmologist.

Clinical criteria:

- The condition must be due to pathologic myopia (PM), **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition for the same eye.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Subfoveal choroidal neovascularisation (CNV)

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by an ophthalmologist or by an accredited ophthalmology registrar in consultation with an ophthalmologist.

Clinical criteria:

- The condition must not be due to pathologic myopia, **AND**
- The condition must not be due to age-related macular degeneration, **AND**
- The condition must be diagnosed by optical coherence tomography; OR
- The condition must be diagnosed by fluorescein angiography, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Authority approval for initial treatment of each eye must be sought.

The first authority application for each eye must be made in writing.

A written application must include:

- a) a completed authority prescription form;
- b) a completed Subfoveal Choroidal Neovascularisation (CNV) - PBS Supporting Information Form; and
- c) a copy of the optical coherence tomography or fluorescein angiogram report.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday). Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au

Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Subfoveal choroidal neovascularisation (CNV)

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by an ophthalmologist or by an accredited ophthalmology registrar in consultation with an ophthalmologist.

Clinical criteria:

- The condition must not be due to pathologic myopia, **AND**
- The condition must not be due to age-related macular degeneration, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition, **AND**
- Patient must have previously received PBS-subsidised treatment with this drug for this condition for the same eye.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

ranibizumab 1.65 mg/0.165 mL injection, 0.165 mL syringe

11480F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	1042.89	41.30	^a Lucentis [NV]

ranibizumab 2.3 mg/0.23 mL injection, 0.23 mL vial

11471R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	1042.89	41.30	^a Lucentis [NV]

▪ **RANIBIZUMAB**

Note Special Pricing Arrangements apply.

Note No increase in the maximum number of repeats may be authorised.

Note No increase in the maximum quantity or number of units may be authorised for applications for treatment of one eye.

Note Where both eyes are affected by the condition, a quantity of 2 units can be requested on the same authority prescription form.

Note Pharmaceutical benefits that have the form ranibizumab 0.165 mL injection syringe and pharmaceutical benefits that have the form ranibizumab 0.23 mL injection vial are equivalent for the purposes of substitution.

Authority required

Branch retinal vein occlusion with macular oedema

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by an ophthalmologist or by an accredited ophthalmology registrar in consultation with an ophthalmologist.

Clinical criteria:

- Patient must have visual impairment due to macular oedema secondary to branched retinal vein occlusion (BRVO), **AND**
- Patient must have documented visual impairment defined as a best corrected visual acuity score between 73 and 20 letters based on the early treatment diabetic retinopathy study chart administered at a distance of 4 metres (approximate Snellen equivalent 20/40 to 20/400), in the eye proposed for treatment, **AND**
- The condition must be diagnosed by optical coherence tomography; OR
- The condition must be diagnosed by fluorescein angiography, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Authority approval for initial treatment of each eye must be sought.

The first authority application for each eye must be made in writing.

A written application must include:

- a) a completed authority prescription form;
- b) a completed Retinal Vein Occlusion Initial PBS authority application Supporting information form; and
- c) a copy of the optical coherence tomography or fluorescein angiogram report.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos
Or mailed to:
Services Australia
Complex Drugs

Reply Paid 9826
HOBART TAS 7001

Authority required

Branch retinal vein occlusion with macular oedema
Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by an ophthalmologist or by an accredited ophthalmology registrar in consultation with an ophthalmologist.

Clinical criteria:

- Patient must have previously been issued with an authority prescription for this drug for the same eye, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Central retinal vein occlusion with macular oedema
Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by an ophthalmologist or by an accredited ophthalmology registrar in consultation with an ophthalmologist.

Clinical criteria:

- Patient must have visual impairment due to macular oedema secondary to central retinal vein occlusion (CRVO), **AND**
- Patient must have documented visual impairment defined as a best corrected visual acuity score between 73 and 24 letters based on the early treatment diabetic retinopathy study chart administered at a distance of 4 metres (approximate Snellen equivalent 20/40 to 20/320), in the eye proposed for treatment, **AND**
- The condition must be diagnosed by optical coherence tomography; OR
- The condition must be diagnosed by fluorescein angiography, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Authority approval for initial treatment of each eye must be sought.

The first authority application for each eye must be made in writing.

A written application must include:

- a) a completed authority prescription form;
- b) a completed Retinal Vein Occlusion Initial PBS authority application Supporting information form; and
- c) a copy of the optical coherence tomography or fluorescein angiogram report.

Note Any queries concerning the arrangements to prescribe may be directed to Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).
Prescribing information (including Authority Application forms and other relevant documentation as applicable) is available on the Services Australia website at www.servicesaustralia.gov.au
Applications for authority to prescribe should be submitted online using the form upload facility in Health Professional Online Services (HPOS) at www.servicesaustralia.gov.au/hpos

Or mailed to:

Services Australia
Complex Drugs
Reply Paid 9826
HOBART TAS 7001

Authority required

Central retinal vein occlusion with macular oedema
Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by an ophthalmologist or by an accredited ophthalmology registrar in consultation with an ophthalmologist.

Clinical criteria:

- Patient must have previously been issued with an authority prescription for this drug for the same eye, **AND**
- The treatment must be the sole PBS-subsidised therapy for this condition.

Note Applications for authorisation under this restriction may be made in real time using the Online PBS Authorities system (see www.servicesaustralia.gov.au/HPOS) or by telephone by contacting Services Australia on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

ranibizumab 1.65 mg/0.165 mL injection, 0.165 mL syringe

11975G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	1042.89	41.30	^a Lucentis [NV]

ranibizumab 2.3 mg/0.23 mL injection, 0.23 mL vial

11981N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	1042.89	41.30	^a Lucentis [NV]

OTHER OPHTHALMOLOGICALS

Other ophthalmologicals

■ CARBOMER-974P

Authority required (STREAMLINED)

6172

Severe dry eye syndrome

Clinical criteria:

- Patient must be sensitive to preservatives in multi-dose eye drops.

carbomer-974P 0.3% eye gel, 30 x 500 mg unit doses

5502N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	
OP	3	5	..	*32.97	34.26	Poly Gel [AQ]	

carbomer-974P 0.3% eye gel, 30 x 500 mg unit doses

8514D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	
NP	3	5	..	*32.97	34.26	Poly Gel [AQ]	

■ CARBOMER-980

Restricted benefit

Severe dry eye syndrome, including Sjogren's syndrome

carbomer-980 0.2% eye gel, 10 g

5503P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
OP	±1	5	..	15.05	16.34	^a Optifresh eye gel [PP]	^a PAA [UL]
			^B 3.64	18.69	16.34	^a Viscotears [UO]	

carbomer-980 0.2% eye gel, 10 g

8384G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	±1	5	..	15.05	16.34	^a Optifresh eye gel [PP]	^a PAA [UL]
			^B 3.64	18.69	16.34	^a Viscotears [UO]	

■ CARBOMER-980

Authority required (STREAMLINED)

6172

Severe dry eye syndrome

Clinical criteria:

- Patient must be sensitive to preservatives in multi-dose eye drops.

carbomer-980 0.2% eye drops, 30 x 0.6 mL unit doses

5504Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	
OP	3	5	..	*37.83	39.12	Viscotears Gel PF [UO]	

carbomer-980 0.2% eye drops, 30 x 0.6 mL unit doses

8578L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	
NP	3	5	..	*37.83	39.12	Viscotears Gel PF [UO]	

■ CARBOMER-980

Note No applications for increased maximum quantities will be authorised.

Note No applications for repeats will be authorised.

Restricted benefit

Severe dry eye syndrome, including Sjogren's syndrome

Clinical criteria:

- Patient must be receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.

carbomer-980 0.2% eye gel, 10 g

9210R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	±1	11	..	15.05	16.34	^a Optifresh eye gel [PP]	^a PAA [UL]
			^B 3.64	18.69	16.34	^a Viscotears [UO]	

■ CARMELLOSE SODIUM

Restricted benefit

Severe dry eye syndrome, including Sjogren's syndrome

carmellose sodium 1% eye drops, 15 mL

5508X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	
OP	±1	5	..	15.17	16.46	Refresh Liquigel [AG]	

carmellose sodium 0.5% eye drops, 15 mL

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
5507W	±1	5	..	15.17	16.46	Refresh Tears Plus [AG]

■ **CARMELLOSE SODIUM**

Restricted benefit

Severe dry eye syndrome, including Sjogren's syndrome

carmellose sodium 1% eye drops, 15 mL

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8593G	±1	5	..	15.17	16.46	Refresh Liquigel [AG]

carmellose sodium 0.5% eye drops, 15 mL

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8548X	±1	5	..	15.17	16.46	Refresh Tears Plus [AG]

■ **CARMELLOSE SODIUM**

Authority required (STREAMLINED)

6172

Severe dry eye syndrome

Clinical criteria:

- Patient must be sensitive to preservatives in multi-dose eye drops.

carmellose sodium 0.5% eye drops, 30 x 0.4 mL unit doses

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
2338C	3	5	..	*29.61	30.90	^a Optifresh Tears [PP]
			^B 6.21	*35.82	30.90	^a Cellufresh [AG]

carmellose sodium 0.5% eye drops, 30 x 0.4 mL unit doses

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
5506T	3	5	..	*29.61	30.90	^a Optifresh Tears [PP]
			^B 6.21	*35.82	30.90	^a Cellufresh [AG]

carmellose sodium 1% eye drops, 30 x 0.4 mL unit doses

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
2324H	3	5	..	*29.61	30.90	^a Optifresh Plus [PP]
			^B 6.21	*35.82	30.90	^a Celluvisc [AG]

carmellose sodium 1% eye drops, 30 x 0.4 mL unit doses

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
5505R	3	5	..	*29.61	30.90	^a Optifresh Plus [PP]
			^B 6.21	*35.82	30.90	^a Celluvisc [AG]

■ **CARMELLOSE SODIUM**

Note The in-use shelf life of Evolve carmellose 0.5% and Evolve hypromellose 0.3% is 3 months from the date of opening.

Authority required (STREAMLINED)

6172

Severe dry eye syndrome

Clinical criteria:

- Patient must be sensitive to preservatives in multi-dose eye drops.

carmellose sodium 0.5% eye drops, 10 mL

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
11852T	±1	5	..	30.30	31.59	Evolve Carmellose [CX]

carmellose sodium 0.5% eye drops, 10 mL

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
11853W	±1	5	..	30.30	31.59	Evolve Carmellose [CX]

■ **CARMELLOSE SODIUM**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Restricted benefit

Severe dry eye syndrome, including Sjogren's syndrome

Clinical criteria:

- Patient must be receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.

carmellose sodium 1% eye drops, 15 mL

9212W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	11	..	15.17	16.46	Refresh Liquigel [AG]

carmellose sodium 0.5% eye drops, 15 mL

9211T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	11	..	15.17	16.46	Refresh Tears Plus [AG]

■ CARMELLOSE SODIUM + GLYCEROL
Restricted benefit

Severe dry eye syndrome, including Sjogren's syndrome

carmellose sodium 0.5% + glycerol 0.9% eye drops, 15 mL

5556K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	3	..	15.17	16.46	Optive [AG]

carmellose sodium 0.5% + glycerol 0.9% eye drops, 15 mL

9355J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	3	..	15.17	16.46	Optive [AG]

■ CARMELLOSE SODIUM + GLYCEROL
Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Restricted benefit

Severe dry eye syndrome, including Sjogren's syndrome

Clinical criteria:

- Patient must be receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.

carmellose sodium 0.5% + glycerol 0.9% eye drops, 15 mL

9356K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	7	..	15.17	16.46	Optive [AG]

■ DEXTRAN-70 + HYPROMELLOSE
Restricted benefit

Severe dry eye syndrome, including Sjogren's syndrome

dextran-70 0.1% + hypromellose 0.3% eye drops, 15 mL

1509K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	5	..	15.72	17.01	^a Poly-Tears [IQ]
			^B 3.84	19.56	17.01	^a Tears Naturale [AQ]

■ DEXTRAN-70 + HYPROMELLOSE
Restricted benefit

Severe dry eye syndrome, including Sjogren's syndrome

dextran-70 0.1% + hypromellose 0.3% eye drops, 15 mL

5520M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	5	..	15.72	17.01	^a Poly-Tears [IQ]
			^B 3.84	19.56	17.01	^a Tears Naturale [AQ]

■ DEXTRAN-70 + HYPROMELLOSE
Authority required (STREAMLINED)
6172

Severe dry eye syndrome

Clinical criteria:

- Patient must be sensitive to preservatives in multi-dose eye drops.

dextran-70 0.1% + hypromellose 0.3% eye drops, 28 x 0.4 mL unit doses

5521N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	5	..	*36.93	38.22	Bion Tears [AQ]

dextran-70 0.1% + hypromellose 0.3% eye drops, 28 x 0.4 mL unit doses

8299T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	5	..	*36.93	38.22	Bion Tears [AQ]

■ DEXTRAN-70 + HYPROMELLOSE
Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Restricted benefit

Severe dry eye syndrome, including Sjogren's syndrome

Clinical criteria:

- Patient must be receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.

dextran-70 0.1% + hypromellose 0.3% eye drops, 15 mL

9216C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	11	..	15.72	17.01	^a Poly-Tears [IQ]
			^B 3.84	19.56	17.01	^a Tears Naturale [AQ]

■ **HYALURONATE SODIUM**

Note The in-use shelf life of Hylo-Fresh and Hylo-Forte is 6 months from the date of opening.

Authority required (STREAMLINED)

4105

Severe dry eye syndrome

Clinical criteria:

- Patient must be sensitive to preservatives in multi-dose eye drops.

hyaluronate sodium 0.2% eye drops, 10 mL

2171G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
OP	±1	5	..	34.49	35.78	Hylo-Forte [AE]

hyaluronate sodium 0.2% eye drops, 10 mL

2253N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	34.49	35.78	Hylo-Forte [AE]

hyaluronate sodium 0.1% eye drops, 10 mL

2181T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	34.49	35.78	Hylo-Fresh [AE]

hyaluronate sodium 0.1% eye drops, 10 mL

2184Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
OP	±1	5	..	34.49	35.78	Hylo-Fresh [AE]

■ **HYPROMELLOSE**

Restricted benefit

Severe dry eye syndrome, including Sjogren's syndrome

hypromellose 0.5% eye drops, 15 mL

2956N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	15.55	16.84	Methopt [AF]

hypromellose 0.3% w/w eye drops, 10 mL

11625W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	14.51	15.80	^a In a Wink Moisturising [IQ]
			^B 4.09	18.60	15.80	^a Genteal [AQ]

■ **HYPROMELLOSE**

Restricted benefit

Severe dry eye syndrome, including Sjogren's syndrome

hypromellose 0.5% eye drops, 15 mL

5517J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
OP	±1	5	..	15.55	16.84	Methopt [AF]

hypromellose 0.3% w/w eye drops, 10 mL

11634H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
OP	±1	5	..	14.51	15.80	^a In a Wink Moisturising [IQ]
			^B 4.09	18.60	15.80	^a Genteal [AQ]

■ **HYPROMELLOSE**

Note The in-use shelf life of Evolve carmellose 0.5% and Evolve hypromellose 0.3% is 3 months from the date of opening.

Authority required (STREAMLINED)

6172

Severe dry eye syndrome

Clinical criteria:

- Patient must be sensitive to preservatives in multi-dose eye drops.

hypromellose 0.3% w/v eye drops, 10 mL

11842G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	30.30	31.59	Evolve Hypromellose [CX]

hypromellose 0.3% w/v eye drops, 10 mL

11849P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
OP	±1	5	..	30.30	31.59	Evolve Hypromellose [CX]

■ HYPROMELLOSE

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Restricted benefit

Severe dry eye syndrome, including Sjogren's syndrome

Clinical criteria:

- Patient must be receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.

hypromellose 0.5% eye drops, 15 mL

9214Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	11	..	15.55	16.84	Methopt [AF]

hypromellose 0.3% w/w eye drops, 10 mL

11643T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	11	..	14.51	15.80	^a In a Wink Moisturising [IQ]
			^B 4.09	18.60	15.80	^a Genteal [AQ]

■ HYPROMELLOSE + CARBOMER-980

Restricted benefit

Severe dry eye syndrome, including Sjogren's syndrome

hypromellose 0.3% + carbomer-980 0.2% eye gel, 10 g

5519L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
OP	±1	5	..	15.55	16.84	^a HPMC PAA [IQ]
			^B 4.62	20.17	16.84	^a Genteal gel [AQ]

■ HYPROMELLOSE + CARBOMER-980

Restricted benefit

Severe dry eye syndrome, including Sjogren's syndrome

hypromellose 0.3% + carbomer-980 0.2% eye gel, 10 g

8564R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	15.55	16.84	^a HPMC PAA [IQ]
			^B 4.62	20.17	16.84	^a Genteal gel [AQ]

■ HYPROMELLOSE + CARBOMER-980

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Restricted benefit

Severe dry eye syndrome, including Sjogren's syndrome

Clinical criteria:

- Patient must be receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.

hypromellose 0.3% + carbomer-980 0.2% eye gel, 10 g

9215B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	11	..	15.55	16.84	^a HPMC PAA [IQ]
			^B 4.62	20.17	16.84	^a Genteal gel [AQ]

■ OCRIPLASMIN

Note No increase in the maximum number of repeats may be authorised.

Note Special Pricing Arrangements apply.

Note Where authority approval for treatment for both eyes simultaneously is being sought, a maximum quantity of 2 vials may be requested.

Authority required

Vitreomacular traction syndrome

Treatment criteria:

- Must be treated by an ophthalmologist.

Clinical criteria:

- Patient must have visual impairment due to vitreomacular traction (VMT) without a full thickness macular hole (FTMH); OR
- Patient must have visual impairment due to vitreomacular traction (VMT) with a full thickness macular hole (FTMH) of a diameter of less than or equal to 400 micrometres, **AND**
- Patient must have documented visual impairment defined as a best corrected visual acuity score of approximate Snellen equivalent 20/25 or worse in the eye proposed for treatment, **AND**
- The condition must be diagnosed by optical coherence tomography, **AND**
- The condition must have a vitreomacular adhesion diameter less than or equal to 1500 micrometres, **AND**
- Patient must not have an epiretinal membrane attached to the vitreomacular traction, **AND**
- The condition must be previously untreated in the eye proposed for treatment, **AND**
- Patient must not have received prior vitrectomy in the eye proposed for treatment, **AND**
- Patient must be symptomatic.

The prescriber must state which eye(s) is being treated at the time of application.

ocriplasmin 375 microgram/0.3 mL intraocular injection, 0.3 mL vial

11798Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	4161.16	41.30	Jetrea RTU [IJ]

▪ **PARAFFIN**

paraffin 1 g/g eye ointment, 2 x 3.5 g

1750D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	‡1	5	..	24.35	25.64	Poly Visc [IQ]
			^B 1.84	26.19	25.64	^a Ircal [PE]
						^a Refresh Night Time [AG]

paraffin 1 g/g eye ointment, 2 x 3.5 g

5522P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
OP	‡1	5	..	24.35	25.64	Poly Visc [IQ]
			^B 1.84	26.19	25.64	^a Ircal [PE]
						^a Refresh Night Time [AG]

paraffin 1 g/g eye ointment, 3.5 g

1754H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*24.90	26.19	Poly Visc [IQ]

paraffin 1 g/g eye ointment, 3.5 g

5523Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
OP	2	5	..	*24.90	26.19	Poly Visc [IQ]

▪ **PARAFFIN**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Restricted benefit

For use in patients who are receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.

paraffin 1 g/g eye ointment, 2 x 3.5 g

9218E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	‡1	11	..	24.35	25.64	Poly Visc [IQ]
			^B 1.84	26.19	25.64	^a Ircal [PE]
						^a Refresh Night Time [AG]

▪ **PARAFFIN**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Restricted benefit

For use in patients who are receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.

paraffin 1 g/g eye ointment, 3.5 g

9217D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	11	..	*24.90	26.19	Poly Visc [IQ]

▪ **PERFLUOROHEXYLOCTANE**

Note The in-use shelf life of Novatears is 6 months from the date of opening.

Authority required (STREAMLINED)

6172

Severe dry eye syndrome

Clinical criteria:

- Patient must be sensitive to preservatives in multi-dose eye drops.

perfluorohexyloctane 100% eye drops, 3 mL

11439C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
OP	1	5	..	33.52	34.81	Novatears [AE]

perfluorohexyloctane 100% eye drops, 3 mL

11446K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	33.52	34.81	Novatears [AE]

■ POLYETHYLENE GLYCOL-400 + PROPYLENE GLYCOL
Restricted benefit

Severe dry eye syndrome, including Sjogren's syndrome

polyethylene glycol-400 0.4% + propylene glycol 0.3% eye drops, 15 mL

5524R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
OP	±1	5	..	15.18	16.47	Systane [AQ]

■ POLYETHYLENE GLYCOL-400 + PROPYLENE GLYCOL
Restricted benefit

Severe dry eye syndrome, including Sjogren's syndrome

polyethylene glycol-400 0.4% + propylene glycol 0.3% eye drops, 15 mL

8676P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	15.18	16.47	Systane [AQ]

■ POLYETHYLENE GLYCOL-400 + PROPYLENE GLYCOL
Authority required (STREAMLINED)
6172

Severe dry eye syndrome

Clinical criteria:

- Patient must be sensitive to preservatives in multi-dose eye drops.

polyethylene glycol-400 0.4% + propylene glycol 0.3% eye drops, 28 x 0.8 mL unit doses

5532E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
OP	2	5	..	*31.56	32.85	Systane [AQ]

polyethylene glycol-400 0.4% + propylene glycol 0.3% eye drops, 28 x 0.8 mL unit doses

9170P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*31.56	32.85	Systane [AQ]

■ POLYETHYLENE GLYCOL-400 + PROPYLENE GLYCOL
Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Restricted benefit

Severe dry eye syndrome, including Sjogren's syndrome

Clinical criteria:

- Patient must be receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.

polyethylene glycol-400 0.4% + propylene glycol 0.3% eye drops, 15 mL

9219F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	11	..	15.18	16.47	Systane [AQ]

■ POLYVINYL ALCOHOL
Restricted benefit

Severe dry eye syndrome, including Sjogren's syndrome

polyvinyl alcohol 1.4% eye drops, 15 mL

2682E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	15.55	16.84	^a PVA Tears [PE]
			^b 1.29	16.84	16.84	^a Liquifilm Tears [AG]

■ POLYVINYL ALCOHOL
Restricted benefit

Severe dry eye syndrome, including Sjogren's syndrome

polyvinyl alcohol 1.4% eye drops, 15 mL

5526W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
OP	±1	5	..	15.55	16.84	^a PVA Tears [PE]
			^B 1.29	16.84	16.84	^a Liquifilm Tears [AG]

■ POLYVINYL ALCOHOL

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Restricted benefit

Severe dry eye syndrome, including Sjogren's syndrome

Clinical criteria:

- Patient must be receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.

polyvinyl alcohol 1.4% eye drops, 15 mL

9220G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	11	..	15.55	16.84	^a PVA Tears [PE]
			^B 1.29	16.84	16.84	^a Liquifilm Tears [AG]

■ RETINOL PALMITATE + PARAFFIN

Note The in-use shelf life of VitA-POS is 6 months from the date of opening.

retinol palmitate 0.0138% + paraffin eye ointment, 5 g

2167C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
OP	2	5	..	*24.90	26.19	VitA-POS [AE]

retinol palmitate 0.0138% + paraffin eye ointment, 5 g

2222Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*24.90	26.19	VitA-POS [AE]

■ RETINOL PALMITATE + PARAFFIN

Note The in-use shelf life of VitA-POS is 6 months from the date of opening.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Restricted benefit

For use in patients who are receiving treatment under a GP Management Plan or Team Care Arrangements where Medicare benefits were or are payable for the preparation of the Plan or coordination of the Arrangements.

retinol palmitate 0.0138% + paraffin eye ointment, 5 g

2202X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	11	..	*24.90	26.19	VitA-POS [AE]

■ SOY LECITHIN + TOCOPHEROL + VITAMIN A

Authority required (STREAMLINED)

6172

Severe dry eye syndrome

Clinical criteria:

- Patient must be sensitive to preservatives in multi-dose eye drops.

soy lecithin 1% + tocopherol 0.002% + vitamin A palmitate 0.025% spray, 100 actuations

5545W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
OP	2	5	..	*34.06	35.35	tearsagain [RB]

soy lecithin 1% + tocopherol 0.002% + vitamin A palmitate 0.025% spray, 100 actuations

9448G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*34.06	35.35	tearsagain [RB]

■ OTOLOGICALS

ANTIINFECTIVES

Antiinfectives

■ CIPROFLOXACIN

Authority required

Chronic suppurative otitis media

Population criteria:

- Patient must be an Aboriginal or a Torres Strait Islander person, **AND**
- Patient must be aged 1 month or older.

Authority required

Chronic suppurative otitis media

Population criteria:

- Patient must be less than 18 years of age.

Clinical criteria:

- Patient must have perforation of the tympanic membrane.

Authority required

Chronic suppurative otitis media

Population criteria:

- Patient must be less than 18 years of age.

Clinical criteria:

- Patient must have a grommet in situ.

ciprofloxacin 0.3% ear drops, 5 mL

2480M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	1	..	37.18	38.47	Ciloxan [NV]

CORTICOSTEROIDS AND ANTIINFECTIVES IN COMBINATION*Corticosteroids and antiinfectives in combination***■ FRAMYCETIN SULFATE + GRAMICIDIN + DEXAMETHASONE****framycetin sulfate 0.5% + gramicidin 0.005% + dexamethasone 0.05% ear drops, 8 mL**

2781J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	2	..	15.28	16.57	^a Otodex [AV]
			^B 2.24	17.52	16.57	^a Sofradex [SW]

■ TRIAMCINOLONE + NEOMYCIN + GRAMICIDIN + NYSTATIN**triamcinolone acetate 0.1% + neomycin 0.25% + gramicidin 0.025% + nystatin 100 000 units/g ointment, 5 g**

2974M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	2	..	13.85	15.14	^a Otocomb Otic [LN]
			^B 1.58	15.43	15.14	^a Kenacomb Otic [AS]

triamcinolone acetate 0.09% + neomycin 0.225% + gramicidin 0.0225% + nystatin 90 000 units/mL ear drops, 7.5 mL

2971J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	2	..	16.21	17.50	^a Otocomb Otic [LN]
			^B 1.58	17.79	17.50	^a Kenacomb Otic [AS]

■ OPHTHALMOLOGICAL AND OTOLOGICAL PREPARATIONS**ANTIINFECTIVES***Antiinfectives***■ FRAMYCETIN SULFATE****framycetin sulfate 0.5% eye/ear drops, 8 mL**

1440T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP MW	±1	2	..	15.28	16.57	Soframycin [SW]

framycetin sulfate 0.5% eye/ear drops, 8 mL

5557L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
OP	±1	2	..	15.28	16.57	Soframycin [SW]

■ VARIOUS**■ ALLERGENS****ALLERGENS***Allergen extracts***■ HONEY BEE VENOM****honey bee venom 550 microgram injection [1 vial] (&) inert substance diluent [9 mL vial] (&) inert substance diluent [3 x 1.8 mL vials], 1 pack**

2886X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	232.31	41.30	Albey Bee Venom [DE]

honey bee venom 550 microgram injection [1 vial] (&) inert substance diluent [9 mL vial], 1 pack

10621B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	296.46	41.30	Hymenoptera Honey Bee Venom [DE]

■ PAPER WASP VENOM

Note Paper wasp venom is not European wasp venom

paper wasp venom 550 microgram injection [1 vial] (&) inert substance diluent [9 mL vial] (&) inert substance diluent [3 x 1.8 mL vials], 1 pack

2918N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	232.31	41.30	Albey Paper Wasp Venom [DE]

paper wasp venom 550 microgram injection [1 vial] (&) inert substance diluent [9 mL vial], 1 pack

10620Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	346.72	41.30	Hymenoptera Paper Wasp Venom [DE]

■ YELLOW JACKET VENOM**yellow jacket venom 550 microgram injection [1 vial] (&) inert substance diluent [9 mL vial], 1 pack**

10622C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	401.45	41.30	Hymenoptera Yellow Jacket Venom [DE]

yellow jacket venom 550 microgram injection [1 vial] (&) inert substance diluent [9 mL vial] (&) inert substance diluent [3 x 1.8 mL vials], 1 pack

2883R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	232.31	41.30	Albey Yellow Jacket Venom [DE]

■ ALL OTHER THERAPEUTIC PRODUCTS**ALL OTHER THERAPEUTIC PRODUCTS***Antidotes***■ NALOXONE****naloxone hydrochloride 1 mg/mL injection, 2 mL syringe**

11077B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	1	..	..	35.99	37.28	Prenoxad [FF]

naloxone hydrochloride 1 mg/mL injection, 2 mL syringe

11078C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	35.99	37.28	Prenoxad [FF]

naloxone hydrochloride 400 microgram/mL injection, 5 x 1 mL ampoules

10783M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	..	..	39.19	40.48	^a Junalox [JO]	^a Naloxone Hydrochloride (DBL) [PF]
						^a Naloxone Juno [JU]	^a NALOXONE SXP [XC]

naloxone hydrochloride 400 microgram/mL injection, 5 x 1 mL ampoules

10787R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
DP	1	..	..	39.19	40.48	^a Junalox [JO]	^a Naloxone Hydrochloride (DBL) [PF]
						^a Naloxone Juno [JU]	^a NALOXONE SXP [XC]

naloxone 1.8 mg/actuation nasal spray, 2 x 1 actuation

11816X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
DP	±1	..	..	48.96	41.30	Nyxoid [MF]

naloxone 1.8 mg/actuation nasal spray, 2 x 1 actuation

11817Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	..	..	48.96	41.30	Nyxoid [MF]

*Drugs for treatment of hyperkalemia and hyperphosphatemia***■ LANTHANUM**

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**5491**

Hyperphosphataemia

Treatment Phase: Maintenance following initiation and stabilisation

Clinical criteria:

- The condition must not be adequately controlled by calcium, **AND**
- Patient must have a serum phosphate of greater than 1.6 mmol per L at the commencement of therapy; OR
- The condition must be where a serum calcium times phosphate product is greater than 4 at the commencement of therapy, **AND**
- The treatment must not be used in combination with any other non-calcium phosphate binding agents.

Treatment criteria:

- Patient must be undergoing dialysis for chronic kidney disease.

lanthanum 750 mg chewable tablet, 6 x 15

9404Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	388.58	41.30	Fosrenol [TK]

lanthanum 500 mg chewable tablet, 2 x 45

9403X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	259.69	41.30	Fosrenol [TK]

lanthanum 1 g chewable tablet, 6 x 15

9405B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	436.56	41.30	Fosrenol [TK]

SEVELAMER**Note Shared Care Model:**

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note Pharmaceutical benefits that have the forms sevelamer hydrochloride 800 mg and sevelamer carbonate 800 mg tablet are equivalent for the purposes of substitution

Authority required (STREAMLINED)**5491**

Hyperphosphataemia

Treatment Phase: Maintenance following initiation and stabilisation

Clinical criteria:

- The condition must not be adequately controlled by calcium, **AND**
- Patient must have a serum phosphate of greater than 1.6 mmol per L at the commencement of therapy; OR
- The condition must be where a serum calcium times phosphate product is greater than 4 at the commencement of therapy, **AND**
- The treatment must not be used in combination with any other non-calcium phosphate binding agents.

Treatment criteria:

- Patient must be undergoing dialysis for chronic kidney disease.

sevelamer carbonate 800 mg tablet, 180

11856B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	5	..	207.12	41.30	^a Sevelamer Apotex [TX]	^a Sevelamer Lupin [GQ]

sevelamer hydrochloride 800 mg tablet, 180

2142R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	207.12	41.30	^a Renagel [GZ]

SUCROFERRIC OXYHYDROXIDE**Note Shared Care Model:**

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required (STREAMLINED)**5491**

Hyperphosphataemia

Treatment Phase: Maintenance following initiation and stabilisation

Clinical criteria:

- The condition must not be adequately controlled by calcium, **AND**
- Patient must have a serum phosphate of greater than 1.6 mmol per L at the commencement of therapy; OR
- The condition must be where a serum calcium times phosphate product is greater than 4 at the commencement of therapy, **AND**

- The treatment must not be used in combination with any other non-calcium phosphate binding agents.

Treatment criteria:

- Patient must be undergoing dialysis for chronic kidney disease.

sucroferric oxyhydroxide 2.5 g (iron 500 mg) chewable tablet, 90

10250L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	411.06	41.30	Velphoro [VL]

Detoxifying agents for antineoplastic treatment▪ **FOLINIC ACID****folinic acid 100 mg/10 mL injection, 10 x 10 mL ampoules**

1704Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	59.11	41.30	Leucovorin Calcium (Pfizer Australia Pty Ltd) [PF]

folinic acid 300 mg/30 mL injection, 30 mL vial

9041W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	4	1	..	*57.30	41.30	Leucovorin Calcium (Hospira Pty Limited) [PF]

▪ **FOLINIC ACID**

Note For item codes 8740B and 1610R, pharmaceutical benefits that have the form injection equivalent to 50 mg folinic acid in 5 mL are equivalent for the purposes of substitution.

folinic acid 50 mg/5 mL injection, 10 x 5 mL ampoules

1610R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	53.85	41.30	Leucovorin Calcium (Pfizer Australia Pty Ltd) [PF]

▪ **FOLINIC ACID****Restricted benefit**

Megaloblastic anaemias

Clinical criteria:

- The condition must be a result of folic acid deficiency from the use of folic acid antagonists.

folinic acid 15 mg tablet, 10

2308L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	93.74	41.30	Leucovorin Calcium (Hospira Pty Limited) [PF]

▪ **MESNA****Restricted benefit**

Urothelial toxicity

Treatment Phase: Prophylaxis or reduction of toxicity

Clinical criteria:

- The treatment must be adjunctive therapy to ifosfamide or high dose cyclophosphamide.

mesna 400 mg/4 mL injection, 15 x 4 mL ampoules

8078E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	83.54	41.30	Uromitexan [BX]

mesna 1 g/10 mL injection, 15 x 10 mL ampoules

8079F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	177.08	41.30	Uromitexan [BX]

Drugs for treatment of hypercalcaemia▪ **PHOSPHORUS****Authority required (STREAMLINED)**

5089

Hypophosphataemic rickets

Authority required (STREAMLINED)

5114

Vitamin D-resistant rickets

Authority required (STREAMLINED)

5095

Familial hypophosphataemia

Authority required (STREAMLINED)

5123

Hypercalcaemia

phosphorus 500 mg effervescent tablet, 100

2946C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	67.55	41.30	PHOSPHATE PHEBRA [FG]

Other therapeutic products**■ POLYLACTIC ACID**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Authority applications may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Severe facial lipoatrophy

Treatment Phase: Initial PBS-subsidised treatment

Clinical criteria:

- The treatment must be for facial administration only, **AND**
- The condition must be caused by therapy for HIV infection.

Accreditation following completion of injection administration training with Galderma is required to prescribe poly-L-lactic acid under the PBS. Patients must be referred from the HIV physician to the accredited injector.

polylactic acid 150 mg injection, 1 vial

9475Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	4	..	*385.92	41.30	Sculptra [GA]

■ POLYLACTIC ACID

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Maintenance treatment is limited to one re-treatment (maximum 2 vials) every 2 years.

Note Authority applications may be made by telephone to the Department of Human Services on 1800 700 270 (hours of operation 8 a.m. to 5 p.m. EST Monday to Friday).

Authority required

Severe facial lipoatrophy

Treatment Phase: Maintenance PBS-subsidised treatment

Clinical criteria:

- The treatment must be for facial administration only, **AND**
- The condition must be caused by therapy for HIV infection.

Accreditation following completion of injection administration training with Galderma is required to prescribe poly-L-lactic acid under the PBS. Patients must be referred from the HIV physician to the accredited injector.

polylactic acid 150 mg injection, 1 vial

9476R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*385.92	41.30	Sculptra [GA]

■ DIAGNOSTIC AGENTS**URINE TESTS****■ GLUCOSE AND KETONE INDICATOR URINE****Restricted benefit**

For treatment of a patient identifying as Aboriginal or Torres Strait Islander

glucose and ketone indicator urine diagnostic strip, 50

3107M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	2	..	*19.80	21.09	Keto-Diastix [DX]

■ GLUCOSE INDICATOR URINE**Restricted benefit**

For treatment of a patient identifying as Aboriginal or Torres Strait Islander

glucose indicator urine diagnostic strip, 50

3104J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	2	..	*21.48	22.77	Diastix [DX]

■ GENERAL NUTRIENTS**OTHER NUTRIENTS****■ MEDIUM CHAIN TRIGLYCERIDES**

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)**6147**

Ketogenic diet

Clinical criteria:

- Patient must have intractable seizures requiring treatment with a ketogenic diet; OR
- Patient must have a glucose transport protein defect; OR
- Patient must have pyruvate dehydrogenase deficiency.

Authority required (STREAMLINED)**6191**

Dietary management of conditions requiring a source of medium chain triglycerides

Clinical criteria:

- Patient must have chylous ascites; OR
- Patient must have chylothorax; OR
- Patient must have hyperlipoproteinaemia type 1; OR
- Patient must have long chain fatty acid oxidation disorders; OR
- Patient must have fat malabsorption due to liver disease; OR
- Patient must have fat malabsorption due to short gut syndrome; OR
- Patient must have fat malabsorption due to cystic fibrosis; OR
- Patient must have fat malabsorption due to gastrointestinal disorders.

medium chain triglycerides oral liquid, 15 x 225 mL bottles

11444H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*274.64	41.30	Betaquik [VF]

■ MEDIUM CHAIN TRIGLYCERIDES**Note** No increase in the maximum quantity or number of units may be authorised.**Note** No increase in the maximum number of repeats may be authorised.**Authority required (STREAMLINED)****6181**

Chylous ascites

Authority required (STREAMLINED)**6134**

Chylothorax

Authority required (STREAMLINED)**6164**

Fat malabsorption

Clinical criteria:

- The condition must be due to liver disease; OR
- The condition must be due to short gut syndrome; OR
- The condition must be due to cystic fibrosis; OR
- The condition must be due to gastrointestinal disorders.

Authority required (STREAMLINED)**6203**

Hyperlipoproteinaemia type 1

Authority required (STREAMLINED)**6155**

Intractable childhood epilepsy

Clinical criteria:

- Patient must require a ketogenic diet.

Authority required (STREAMLINED)**6135**

Cerebrospinal fluid glucose transporter defect

Clinical criteria:

- Patient must require a ketogenic diet.

Authority required (STREAMLINED)**6146**

Long chain fatty acid oxidation disorders

medium chain triglycerides oral oil, 500 mL

3128P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*45.70	41.30	MCT Oil [SB]

medium chain triglycerides oral liquid, 250 mL bottle

9327X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	8	5	..	*169.18	41.30	Liquigen [SB]

■ PROTEIN FORMULA WITH CARBOHYDRATE, FAT, VITAMINS AND MINERALS**Note** No increase in the maximum number of repeats may be authorised.

Note Not indicated for the treatment of intractable childhood epilepsy or cerebrospinal fluid glucose transporter defect requiring a ketogenic diet.

Restricted benefit

Dietary management of conditions requiring a source of medium chain triglycerides

Clinical criteria:

- Patient must have fat malabsorption due to liver disease; OR
- Patient must have fat malabsorption due to short gut syndrome; OR
- Patient must have fat malabsorption due to cystic fibrosis; OR
- Patient must have fat malabsorption due to gastrointestinal disorders.

Population criteria:

- Patient must be aged from 1 to 10 years inclusive.

protein formula with carbohydrate, fat, vitamins and minerals oral liquid, 12 x 500 mL bottles

11939J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	7	5	..	*1582.25	41.30	Nutrini Peptisorb Energy [NU]

Fat/carbohydrates/proteins/minerals/vitamins, combinations

■ AMINO ACID SYNTHETIC FORMULA

Note Authorities for increased maximum quantities, up to a maximum of 52, may be authorised.

Authority required

Eosinophilic oesophagitis

Treatment Phase: Initial treatment for up to 3 months

Treatment criteria:

- Must be treated by a clinical immunologist, suitably qualified allergist or gastroenterologist.

Clinical criteria:

- Patient must require an amino acid based formula as a component of a dietary elimination program.

Population criteria:

- Patient must be 18 years of age or less.

Treatment with oral steroids should not be commenced during the period of initial treatment.

Eosinophilic oesophagitis is demonstrated by the following criteria:

- Chronic symptoms of reflux that persisted despite a 2-month trial of a proton pump inhibitor or chronic dysphagia; and
- A lack of demonstrable anatomic abnormality with the exception of stricture, which can be attributable to eosinophilic oesophagitis; and
- Eosinophilic infiltration of the oesophagus, demonstrated by oesophageal biopsy specimens obtained by endoscopy and where the most densely involved oesophageal biopsy had 20 or more eosinophils in any single 400 x high powered field, along with normal antral and duodenal biopsies.

The date of birth of the patient must be included in the authority application.

Authority required

Eosinophilic oesophagitis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a clinical immunologist, suitably qualified allergist or gastroenterologist.

Clinical criteria:

- Patient must have responded to an initial course of PBS-subsidised treatment.

Population criteria:

- Patient must be 18 years of age or less.

Response to initial treatment is demonstrated by oesophageal biopsy specimens obtained by endoscopy, where the most densely involved oesophageal biopsy had 5 or less eosinophils in any single 400 x high powered field, along with normal antral and duodenal biopsies. The response criteria will not be deemed to have been met if oral steroids were commenced during initial treatment.

amino acid synthetic formula powder for oral liquid, 400 g

1521C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	12	5	..	*437.82	41.30	Neocate Junior Vanilla [SB]

amino acid synthetic formula powder for oral liquid, 400 g

2250K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	12	5	..	*437.82	41.30	EleCare [AB]

■ AMINO ACID SYNTHETIC FORMULA

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Cows' milk protein enteropathy

Treatment Phase: Initial treatment for up to 6 months

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist, or in consultation with a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist.

Clinical criteria:

- The condition must not be isolated infant colic or reflux, **AND**
- Patient must be intolerant to both soy protein and protein hydrolysate formulae, as demonstrated when the child has failed to respond to a strict cows' milk protein free and strict soy protein free diet with a protein hydrolysate (with or without medium chain triglycerides) as the principal formula.

Population criteria:

- Patient must be up to the age of 24 months.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Severe cows' milk protein enteropathy with failure to thrive

Treatment Phase: Initial treatment for up to 6 months

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist, or in consultation with a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist.

Clinical criteria:

- The condition must not be isolated infant colic or reflux.

Population criteria:

- Patient must be up to the age of 24 months.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Combined intolerance to cows' milk protein, soy protein and protein hydrolysate formulae

Treatment Phase: Initial treatment for up to 6 months

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist.

Clinical criteria:

- The condition must not be isolated infant colic or reflux.

Population criteria:

- Patient must be older than 24 months of age.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Proven combined immunoglobulin E (IgE) mediated allergy to cows' milk protein and soy protein

Treatment Phase: Initial treatment for up to 6 months

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist, or in consultation with a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist.

Clinical criteria:

- Patient must have failed a trial of protein hydrolysate formulae (with or without medium chain triglycerides).

Population criteria:

- Patient must be up to the age of 24 months.

The name of the specialist and the date of birth of the patient must be included in the authority application.

amino acid synthetic formula powder for oral liquid, 400 g

1180D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	8	5	..	*294.22	41.30	Neocate Junior Vanilla [SB]

amino acid synthetic formula powder for oral liquid, 400 g

8574G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	8	5	..	*294.22	41.30	EleCare [AB]

■ AMINO ACID SYNTHETIC FORMULA

Note Authorities for increased maximum quantities, up to a maximum of 20, may be authorised.

Authority required

Cows' milk anaphylaxis

Treatment criteria:

- Must be treated by a specialist allergist or clinical immunologist, or in consultation with a specialist allergist or clinical immunologist.

Population criteria:

- Patient must be up to the age of 24 months.

Anaphylaxis is defined as a severe and/or potentially life threatening allergic reaction.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Cows' milk protein enteropathy

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist, or have an appointment to be assessed by one of these specialists.

Clinical criteria:

- The condition must not be isolated infant colic or reflux, **AND**
- Patient must be intolerant to both soy protein and protein hydrolysate formulae, as demonstrated when the child has failed to respond to a strict cows' milk protein free and strict soy protein free diet with a protein hydrolysate (with or without medium chain triglycerides) as the principal formula.

Population criteria:

- Patient must be up to the age of 24 months.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Severe cows' milk protein enteropathy with failure to thrive

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist, or have been assessed at least once or have an appointment to be assessed by one of these specialists.

Clinical criteria:

- The condition must not be isolated infant colic or reflux, **AND**
- Patient must have had failure to thrive prior to commencement with initial treatment.

Population criteria:

- Patient must be up to the age of 24 months.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Combined intolerance to cows' milk protein, soy protein and protein hydrolysate formulae

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist at intervals not greater than 12 months.

Clinical criteria:

- The condition must not be isolated infant colic or reflux.

Population criteria:

- Patient must be older than 24 months of age.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Proven combined immunoglobulin E (IgE) mediated allergy to cows' milk protein and soy protein

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist.

Clinical criteria:

- Patient must have failed a trial of protein hydrolysate formulae (with or without medium chain triglycerides) prior to commencement with initial treatment.

Population criteria:

- Patient must be up to the age of 24 months.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Severe intestinal malabsorption including short bowel syndrome

Clinical criteria:

- Patient must have failed to respond to protein hydrolysate formulae; OR
- Patient must have been receiving parenteral nutrition.

amino acid synthetic formula powder for oral liquid, 400 g

1192R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	8	5	..	*294.22	41.30	Neocate Junior Vanilla [SB]

amino acid synthetic formula powder for oral liquid, 400 g

8575H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	8	5	..	*294.22	41.30	EleCare [AB]

■ AMINO ACID SYNTHETIC FORMULA SUPPLEMENTED WITH LONG CHAIN POLYUNSATURATED FATTY ACIDS

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Cows' milk protein enteropathy

Treatment Phase: Initial treatment for up to 6 months

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist, or in consultation with a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist.

Clinical criteria:

- The condition must not be isolated infant colic or reflux, **AND**
- Patient must be intolerant to both soy protein and protein hydrolysate formulae, as demonstrated when the child has failed to respond to a strict cows' milk protein free and strict soy protein free diet with a protein hydrolysate (with or without medium chain triglycerides) as the principal formula.

Population criteria:

- Patient must be up to the age of 24 months.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Severe cows' milk protein enteropathy with failure to thrive

Treatment Phase: Initial treatment for up to 6 months

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist, or in consultation with a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist.

Clinical criteria:

- The condition must not be isolated infant colic or reflux.

Population criteria:

- Patient must be up to the age of 24 months.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Combined intolerance to cows' milk protein, soy protein and protein hydrolysate formulae

Treatment Phase: Initial treatment for up to 6 months

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist.

Clinical criteria:

- The condition must not be isolated infant colic or reflux.

Population criteria:

- Patient must be older than 24 months of age.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Proven combined immunoglobulin E (IgE) mediated allergy to cows' milk protein and soy protein

Treatment Phase: Initial treatment for up to 6 months

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist, or in consultation with a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist.

Clinical criteria:

- Patient must have failed a trial of protein hydrolysate formulae (with or without medium chain triglycerides).

Population criteria:

- Patient must be up to the age of 24 months.

The name of the specialist and the date of birth of the patient must be included in the authority application.

amino acid synthetic formula supplemented with long chain polyunsaturated fatty acids powder for oral liquid, 400 g

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
2246F	8	5	..	*315.42	41.30	Neocate LCP [SB]

NP

amino acid synthetic formula supplemented with long chain polyunsaturated fatty acids powder for oral liquid, 400 g

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
9339M	8	5	..	*315.42	41.30	EleCare LCP [AB]

NP

■ AMINO ACID SYNTHETIC FORMULA SUPPLEMENTED WITH LONG CHAIN POLYUNSATURATED FATTY ACIDS

Note Authorities for increased maximum quantities, up to a maximum of 20, may be authorised.

Authority required

Cows' milk anaphylaxis

Treatment criteria:

- Must be treated by a specialist allergist or clinical immunologist, or in consultation with a specialist allergist or clinical immunologist.

Population criteria:

- Patient must be up to the age of 24 months.

Anaphylaxis is defined as a severe and/or potentially life threatening allergic reaction.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Cows' milk protein enteropathy

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist, or have an appointment to be assessed by one of these specialists.

Clinical criteria:

- The condition must not be isolated infant colic or reflux, **AND**
- Patient must be intolerant to both soy protein and protein hydrolysate formulae, as demonstrated when the child has failed to respond to a strict cows' milk protein free and strict soy protein free diet with a protein hydrolysate (with or without medium chain triglycerides) as the principal formula.

Population criteria:

- Patient must be up to the age of 24 months.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Severe cows' milk protein enteropathy with failure to thrive

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist, or have been assessed at least once or have an appointment to be assessed by one of these specialists.

Clinical criteria:

- The condition must not be isolated infant colic or reflux, **AND**
- Patient must have had failure to thrive prior to commencement with initial treatment.

Population criteria:

- Patient must be up to the age of 24 months.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Combined intolerance to cows' milk protein, soy protein and protein hydrolysate formulae

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist at intervals not greater than 12 months.

Clinical criteria:

- The condition must not be isolated infant colic or reflux.

Population criteria:

- Patient must be older than 24 months of age.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Proven combined immunoglobulin E (IgE) mediated allergy to cows' milk protein and soy protein

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist.

Clinical criteria:

- Patient must have failed a trial of protein hydrolysate formulae (with or without medium chain triglycerides) prior to commencement with initial treatment.

Population criteria:

- Patient must be up to the age of 24 months.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Severe intestinal malabsorption including short bowel syndrome

Clinical criteria:

- Patient must have failed to respond to protein hydrolysate formulae; OR
- Patient must have been receiving parenteral nutrition.

amino acid synthetic formula supplemented with long chain polyunsaturated fatty acids powder for oral liquid, 400 g

2560R

NP

Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8	5	..	*315.42	41.30	Neocate LCP [SB]

amino acid synthetic formula supplemented with long chain polyunsaturated fatty acids powder for oral liquid, 400 g

9340N

NP

Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8	5	..	*315.42	41.30	EleCare LCP [AB]

■ AMINO ACID SYNTHETIC FORMULA SUPPLEMENTED WITH LONG CHAIN POLYUNSATURATED FATTY ACIDS AND MEDIUM CHAIN TRIGLYCERIDES

Note Authorities for increased maximum quantities, up to a maximum of 52, may be authorised.

Authority required

Eosinophilic oesophagitis

Treatment Phase: Initial treatment for up to 3 months

Treatment criteria:

- Must be treated by a clinical immunologist, suitably qualified allergist or gastroenterologist.

Clinical criteria:

- Patient must require an amino acid based formula as a component of a dietary elimination program.

Population criteria:

- Patient must be 18 years of age or less.

Treatment with oral steroids should not be commenced during the period of initial treatment.

Eosinophilic oesophagitis is demonstrated by the following criteria:

- (i) Chronic symptoms of reflux that persisted despite a 2-month trial of a proton pump inhibitor or chronic dysphagia; and
- (ii) A lack of demonstrable anatomic abnormality with the exception of stricture, which can be attributable to eosinophilic oesophagitis; and
- (iii) Eosinophilic infiltration of the oesophagus, demonstrated by oesophageal biopsy specimens obtained by endoscopy and where the most densely involved oesophageal biopsy had 20 or more eosinophils in any single 400 x high powered field, along with normal antral and duodenal biopsies.

The date of birth of the patient must be included in the authority application.

Authority required

Eosinophilic oesophagitis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a clinical immunologist, suitably qualified allergist or gastroenterologist.

Clinical criteria:

- Patient must have responded to an initial course of PBS-subsidised treatment.

Population criteria:

- Patient must be 18 years of age or less.

Response to initial treatment is demonstrated by oesophageal biopsy specimens obtained by endoscopy, where the most densely involved oesophageal biopsy had 5 or less eosinophils in any single 400 x high powered field, along with normal antral and duodenal biopsies. The response criteria will not be deemed to have been met if oral steroids were commenced during initial treatment.

amino acid synthetic formula supplemented with long chain polyunsaturated fatty acids and medium chain triglycerides powder for oral liquid, 400 g

1545H

Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
12	5	..	*469.62	41.30	Neocate Gold [SB]

NP

■ AMINO ACID SYNTHETIC FORMULA SUPPLEMENTED WITH LONG CHAIN POLYUNSATURATED FATTY ACIDS AND MEDIUM CHAIN TRIGLYCERIDES

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Cows' milk protein enteropathy

Treatment Phase: Initial treatment for up to 6 months

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist, or in consultation with a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist.

Clinical criteria:

- The condition must not be isolated infant colic or reflux, **AND**
- Patient must be intolerant to both soy protein and protein hydrolysate formulae, as demonstrated when the child has failed to respond to a strict cows' milk protein free and strict soy protein free diet with a protein hydrolysate (with or without medium chain triglycerides) as the principal formula.

Population criteria:

- Patient must be up to the age of 24 months.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Severe cows' milk protein enteropathy with failure to thrive

Treatment Phase: Initial treatment for up to 6 months

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist, or in consultation with a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist.

Clinical criteria:

- The condition must not be isolated infant colic or reflux.

Population criteria:

- Patient must be up to the age of 24 months.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Combined intolerance to cows' milk protein, soy protein and protein hydrolysate formulae

Treatment Phase: Initial treatment for up to 6 months

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist.

Clinical criteria:

- The condition must not be isolated infant colic or reflux.

Population criteria:

- Patient must be older than 24 months of age.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Proven combined immunoglobulin E (IgE) mediated allergy to cows' milk protein and soy protein

Treatment Phase: Initial treatment for up to 6 months

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist, or in consultation with a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist.

Clinical criteria:

- Patient must have failed a trial of protein hydrolysate formulae (with or without medium chain triglycerides).

Population criteria:

- Patient must be up to the age of 24 months.

The name of the specialist and the date of birth of the patient must be included in the authority application.

amino acid synthetic formula supplemented with long chain polyunsaturated fatty acids and medium chain triglycerides powder for oral liquid, 400 g

5466Q

Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8	5	..	*315.42	41.30	Neocate Gold [SB]

NP

■ AMINO ACID SYNTHETIC FORMULA SUPPLEMENTED WITH LONG CHAIN POLYUNSATURATED FATTY ACIDS AND MEDIUM CHAIN TRIGLYCERIDES

Note Authorities for increased maximum quantities, up to a maximum of 20, may be authorised.

Authority required

Cows' milk anaphylaxis

Treatment criteria:

- Must be treated by a specialist allergist or clinical immunologist, or in consultation with a specialist allergist or clinical immunologist.

Population criteria:

- Patient must be up to the age of 24 months.

Anaphylaxis is defined as a severe and/or potentially life threatening allergic reaction.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Cows' milk protein enteropathy

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist, or have an appointment to be assessed by one of these specialists.

Clinical criteria:

- The condition must not be isolated infant colic or reflux, **AND**
- Patient must be intolerant to both soy protein and protein hydrolysate formulae, as demonstrated when the child has failed to respond to a strict cows' milk protein free and strict soy protein free diet with a protein hydrolysate (with or without medium chain triglycerides) as the principal formula.

Population criteria:

- Patient must be up to the age of 24 months.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Severe cows' milk protein enteropathy with failure to thrive

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist, or have been assessed at least once or have an appointment to be assessed by one of these specialists.

Clinical criteria:

- The condition must not be isolated infant colic or reflux, **AND**
- Patient must have had failure to thrive prior to commencement with initial treatment.

Population criteria:

- Patient must be up to the age of 24 months.
- The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Combined intolerance to cows' milk protein, soy protein and protein hydrolysate formulae

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist at intervals not greater than 12 months.

Clinical criteria:

- The condition must not be isolated infant colic or reflux.

Population criteria:

- Patient must be older than 24 months of age.
- The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Proven combined immunoglobulin E (IgE) mediated allergy to cows' milk protein and soy protein

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist.

Clinical criteria:

- Patient must have failed a trial of protein hydrolysate formulae (with or without medium chain triglycerides) prior to commencement with initial treatment.

Population criteria:

- Patient must be up to the age of 24 months.
- The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Severe intestinal malabsorption including short bowel syndrome

Clinical criteria:

- Patient must have failed to respond to protein hydrolysate formulae; OR
- Patient must have been receiving parenteral nutrition.

amino acid synthetic formula supplemented with long chain polyunsaturated fatty acids and medium chain triglycerides powder for oral liquid, 400 g

5467R

Max. Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8	5	..	*315.42	41.30	Neocate Gold [SB]

NP

■ AMINO ACID SYNTHETIC FORMULA SUPPLEMENTED WITH LONG CHAIN POLYUNSATURATED FATTY ACIDS AND MEDIUM CHAIN TRIGLYCERIDES

Note Authorities for increased maximum quantities, up to a maximum of 20, may be authorised.

Authority required

Cows' milk protein enteropathy

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist, or have an appointment to be assessed by one of these specialists.

Clinical criteria:

- The condition must not be isolated infant colic or reflux, **AND**
- Patient must be intolerant to both soy protein and protein hydrolysate formulae, as demonstrated when the child has failed to respond to a strict cows' milk protein free and strict soy protein free diet with a protein hydrolysate (with or without medium chain triglycerides) as the principal formula.

Population criteria:

- Patient must be up to the age of 24 months.
- The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Severe cows' milk protein enteropathy with failure to thrive

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist, or have been assessed at least once or have an appointment to be assessed by one of these specialists.

Clinical criteria:

- The condition must not be isolated infant colic or reflux, **AND**
- Patient must have had failure to thrive prior to commencement with initial treatment.

Population criteria:

- Patient must be up to the age of 24 months.
- The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Combined intolerance to cows' milk protein, soy protein and protein hydrolysate formulae

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist at intervals not greater than 12 months.

Clinical criteria:

- The condition must not be isolated infant colic or reflux.

Population criteria:

- Patient must be older than 24 months of age.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Cows' milk anaphylaxis

Treatment criteria:

- Must be treated by a specialist allergist or clinical immunologist, or in consultation with a specialist allergist or clinical immunologist.

Population criteria:

- Patient must be up to the age of 24 months.

Anaphylaxis is defined as a severe and/or potentially life threatening allergic reaction.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Proven combined immunoglobulin E (IgE) mediated allergy to cows' milk protein and soy protein

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist.

Clinical criteria:

- Patient must have failed a trial of protein hydrolysate formulae (with or without medium chain triglycerides) prior to commencement with initial treatment.

Population criteria:

- Patient must be up to the age of 24 months.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Severe intestinal malabsorption including short bowel syndrome

Clinical criteria:

- Patient must have failed to respond to protein hydrolysate formulae; OR
- Patient must have been receiving parenteral nutrition.

Authority required

Eosinophilic oesophagitis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a clinical immunologist, suitably qualified allergist or gastroenterologist.

Clinical criteria:

- Patient must have responded to an initial course of PBS-subsidised treatment.

Population criteria:

- Patient must be 18 years of age or less.

Response to initial treatment is demonstrated by oesophageal biopsy specimens obtained by endoscopy, where the most densely involved oesophageal biopsy had 5 or less eosinophils in any single 400 x high powered field, along with normal antral and duodenal biopsies. The response criteria will not be deemed to have been met if oral steroids were commenced during initial treatment.

amino acid synthetic formula supplemented with long chain polyunsaturated fatty acids and medium chain triglycerides powder for oral liquid, 400 g

2900P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	8	5	..	*315.42	41.30	Alfamino [NT]

■ AMINO ACID SYNTHETIC FORMULA SUPPLEMENTED WITH LONG CHAIN POLYUNSATURATED FATTY ACIDS AND MEDIUM CHAIN TRIGLYCERIDES

Authority required

Cows' milk protein enteropathy

Treatment Phase: Initial treatment for up to 6 months

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist, or in consultation with a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist.

Clinical criteria:

- The condition must not be isolated infant colic or reflux, **AND**
- Patient must be intolerant to both soy protein and protein hydrolysate formulae, as demonstrated when the child has failed to respond to a strict cows' milk protein free and strict soy protein free diet with a protein hydrolysate (with or without medium chain triglycerides) as the principal formula.

Population criteria:

- Patient must be up to the age of 24 months.
- The name of the specialist and the date of birth of the patient must be included in the authority application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Severe cows' milk protein enteropathy with failure to thrive

Treatment Phase: Initial treatment for up to 6 months

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist, or in consultation with a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist.

Clinical criteria:

- The condition must not be isolated infant colic or reflux.

Population criteria:

- Patient must be up to the age of 24 months.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Combined intolerance to cows' milk protein, soy protein and protein hydrolysate formulae

Treatment Phase: Initial treatment for up to 6 months

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist.

Clinical criteria:

- The condition must not be isolated infant colic or reflux.

Population criteria:

- Patient must be older than 24 months of age.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Proven combined immunoglobulin E (IgE) mediated allergy to cows' milk protein and soy protein

Treatment Phase: Initial treatment for up to 6 months

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist, or in consultation with a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist.

Clinical criteria:

- Patient must have failed a trial of protein hydrolysate formulae (with or without medium chain triglycerides).

Population criteria:

- Patient must be up to the age of 24 months.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Eosinophilic oesophagitis

Treatment Phase: Initial treatment for up to 3 months

Treatment criteria:

- Must be treated by a clinical immunologist, suitably qualified allergist or gastroenterologist.

Clinical criteria:

- Patient must require an amino acid based formula as a component of a dietary elimination program.

Population criteria:

- Patient must be 18 years of age or less.

Treatment with oral steroids should not be commenced during the period of initial treatment.

Eosinophilic oesophagitis is demonstrated by the following criteria:

- Chronic symptoms of reflux that persisted despite a 2-month trial of a proton pump inhibitor or chronic dysphagia; and
- A lack of demonstrable anatomic abnormality with the exception of stricture, which can be attributable to eosinophilic oesophagitis; and
- Eosinophilic infiltration of the oesophagus, demonstrated by oesophageal biopsy specimens obtained by endoscopy and where the most densely involved oesophageal biopsy had 20 or more eosinophils in any single 400 x high powered field, along with normal antral and duodenal biopsies.

The date of birth of the patient must be included in the authority application.

Note Authorities for increased maximum quantities, up to a maximum of 20, may be authorised.

amino acid synthetic formula supplemented with long chain polyunsaturated fatty acids and medium chain triglycerides powder for oral liquid, 400 g

2928D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	8	5	..	*315.42	41.30	Alfamino [NT]

■ PROTEIN HYDROLYSATE FORMULA WITH MEDIUM CHAIN TRIGLYCERIDES

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required (STREAMLINED)**6174**

Cows' milk protein enteropathy and intolerance to soy protein

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist, specialist paediatrician or specialist paediatric gastroenterologist and hepatologist, or in consultation with a specialist allergist, clinical immunologist, specialist paediatrician or specialist paediatric gastroenterologist and hepatologist.

Clinical criteria:

- The condition must not be isolated infant colic or reflux, **AND**
- Patient must have failed to respond to a strict soy-based cows' milk protein free diet.

Population criteria:

- Patient must be up to the age of 24 months.

Authority required (STREAMLINED)**6193**

Cows' milk protein enteropathy and intolerance to soy protein

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist, specialist paediatrician or specialist paediatric gastroenterologist and hepatologist, or in consultation with a specialist allergist, clinical immunologist, specialist paediatrician or specialist paediatric gastroenterologist and hepatologist.

Clinical criteria:

- The condition must not be isolated infant colic or reflux, **AND**
- Patient must have demonstrated a clinical improvement with the protein hydrolysate formula with medium chain triglycerides.

Population criteria:

- Patient must be up to the age of 24 months.

Authority required (STREAMLINED)**6204**

Cows' milk protein enteropathy and intolerance to soy protein

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist, specialist paediatrician or specialist paediatric gastroenterologist and hepatologist.

Clinical criteria:

- The condition must not be isolated infant colic or reflux, **AND**
- Patient must have failed to respond to a strict soy-based cows' milk protein free diet.

Population criteria:

- Patient must be older than 24 months of age.
- The name of the specialist must be documented in the patient's medical records

Authority required (STREAMLINED)**6137**

Proven combined immunoglobulin E (IgE) mediated allergy to cows' milk protein and soy protein

Treatment Phase: Initial treatment for up to 6 months

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist, or in consultation with a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist.

Population criteria:

- Patient must be up to the age of 24 months.
- The name of the specialist must be documented in the patient's medical records

Authority required (STREAMLINED)**6182**

Proven combined immunoglobulin E (IgE) mediated allergy to cows' milk protein and soy protein

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist.

Population criteria:

- Patient must be up to the age of 24 months.
- The name of the specialist must be documented in the patient's medical records

Authority required (STREAMLINED)**6194**

Biliary atresia

Authority required (STREAMLINED)**6157**

Chronic liver failure with fat malabsorption

Authority required (STREAMLINED)**6205**

Chylous ascites

Authority required (STREAMLINED)**6195**

Cystic fibrosis

Authority required (STREAMLINED)**6158**

Enterokinase deficiency

Authority required (STREAMLINED)**6166**

Proven fat malabsorption

Authority required (STREAMLINED)**6148**

Severe diarrhoea of greater than 2 weeks duration

Population criteria:

- Patient must be aged less than 4 months.

Authority required (STREAMLINED)**6138**

Severe intestinal malabsorption including short bowel syndrome

protein hydrolysate formula with medium chain triglycerides powder for oral liquid, 450 g

8259Q

Max. Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8	5	..	*88.38	41.30	Aptamil Gold+ Pepti-Junior [NU]

NP

■ PROTEIN HYDROLYSATE FORMULA WITH MEDIUM CHAIN TRIGLYCERIDES**Note** No increase in the maximum quantity or number of units may be authorised.**Note** No increase in the maximum number of repeats may be authorised.**Authority required (STREAMLINED)****6174**

Cows' milk protein enteropathy and intolerance to soy protein

Treatment Phase: Initial treatment

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist, specialist paediatrician or specialist paediatric gastroenterologist and hepatologist, or in consultation with a specialist allergist, clinical immunologist, specialist paediatrician or specialist paediatric gastroenterologist and hepatologist.

Clinical criteria:

- The condition must not be isolated infant colic or reflux, **AND**
- Patient must have failed to respond to a strict soy-based cows' milk protein free diet.

Population criteria:

- Patient must be up to the age of 24 months.

Authority required (STREAMLINED)**6193**

Cows' milk protein enteropathy and intolerance to soy protein

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist, specialist paediatrician or specialist paediatric gastroenterologist and hepatologist, or in consultation with a specialist allergist, clinical immunologist, specialist paediatrician or specialist paediatric gastroenterologist and hepatologist.

Clinical criteria:

- The condition must not be isolated infant colic or reflux, **AND**
- Patient must have demonstrated a clinical improvement with the protein hydrolysate formula with medium chain triglycerides.

Population criteria:

- Patient must be up to the age of 24 months.

Authority required (STREAMLINED)**6204**

Cows' milk protein enteropathy and intolerance to soy protein

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist, specialist paediatrician or specialist paediatric gastroenterologist and hepatologist.

Clinical criteria:

- The condition must not be isolated infant colic or reflux, **AND**
- Patient must have failed to respond to a strict soy-based cows' milk protein free diet.

Population criteria:

- Patient must be older than 24 months of age.

The name of the specialist must be documented in the patient's medical records

Authority required (STREAMLINED)**6137**

Proven combined immunoglobulin E (IgE) mediated allergy to cows' milk protein and soy protein

Treatment Phase: Initial treatment for up to 6 months

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist, or in consultation with a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist.

Population criteria:

- Patient must be up to the age of 24 months.

The name of the specialist must be documented in the patient's medical records

Authority required (STREAMLINED)**6182**

Proven combined immunoglobulin E (IgE) mediated allergy to cows' milk protein and soy protein

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist.

Population criteria:

- Patient must be up to the age of 24 months.

The name of the specialist must be documented in the patient's medical records

Authority required (STREAMLINED)**6194**

Biliary atresia

Authority required (STREAMLINED)**6157**

Chronic liver failure with fat malabsorption

Authority required (STREAMLINED)**6205**

Chylous ascites

Authority required (STREAMLINED)**6195**

Cystic fibrosis

Authority required (STREAMLINED)**6158**

Enterokinase deficiency

Authority required (STREAMLINED)**6166**

Proven fat malabsorption

Authority required (STREAMLINED)**6148**

Severe diarrhoea of greater than 2 weeks duration

Population criteria:

- Patient must be aged less than 4 months.

Authority required (STREAMLINED)**6138**

Severe intestinal malabsorption including short bowel syndrome

Authority required (STREAMLINED)**6206**

Chylothorax

protein hydrolysate formula with medium chain triglycerides powder for oral liquid, 400 g

2676W

NP

Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8	5	..	*135.34	41.30	Alfaré [NT]

■ TRIGLYCERIDES MEDIUM CHAIN FORMULA

Note No increase in the maximum number of repeats may be authorised.

Note No increase in the maximum quantity or number of units may be authorised.

Note Not indicated for the treatment of intractable childhood epilepsy or cerebrospinal fluid glucose transporter defect requiring a ketogenic diet.

Restricted benefit

Dietary management of conditions requiring a source of medium chain triglycerides

Clinical criteria:

- Patient must have fat malabsorption due to liver disease; OR
- Patient must have fat malabsorption due to short gut syndrome; OR
- Patient must have fat malabsorption due to cystic fibrosis; OR
- Patient must have fat malabsorption due to gastrointestinal disorders.

triglycerides medium chain formula powder for oral liquid, 400 g

10152H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
(NP)	8	5	..	*402.94	41.30	Monogen [SB]

triglycerides medium chain formula powder for oral liquid, 400 g

10155L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
(NP)	8	5	..	*363.82	41.30	Lipistart [VF]

triglycerides medium chain formula oral liquid, 8 x 500 mL pouches

10375C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
(NP)	8	5	..	*717.98	41.30	Nutrini Peptisorb [SB]

▪ TRIGLYCERIDES MEDIUM CHAIN FORMULA

Note No increase in the maximum number of repeats may be authorised.

Note Authorities for increased maximum quantities, up to a maximum of 20, may be authorised.

Note Not indicated for the treatment of intractable childhood epilepsy or cerebrospinal fluid glucose transporter defect requiring a ketogenic diet.

Restricted benefit

Dietary management of conditions requiring a source of medium chain triglycerides

Clinical criteria:

- Patient must have fat malabsorption due to liver disease; OR
- Patient must have fat malabsorption due to short gut syndrome; OR
- Patient must have fat malabsorption due to cystic fibrosis; OR
- Patient must have fat malabsorption due to gastrointestinal disorders.

triglycerides medium chain formula powder for oral liquid, 400 g

10154K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
(NP)	8	5	..	*337.18	41.30	Peptamen Junior [NT]

▪ TRIGLYCERIDES MEDIUM CHAIN FORMULA

Note No increase in the maximum number of repeats may be authorised.

Note No increase in the maximum quantity or number of units may be authorised.

Note Not indicated for the treatment of intractable childhood epilepsy or cerebrospinal fluid glucose transporter defect requiring a ketogenic diet.

Restricted benefit

Hyperlipoproteinaemia type 1

Restricted benefit

Long chain fatty acid oxidation disorders

Restricted benefit

Chylous ascites

Restricted benefit

Chyllothorax

triglycerides medium chain formula powder for oral liquid, 400 g

1938B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
(NP)	8	5	..	*363.82	41.30	Lipistart [VF]

triglycerides medium chain formula powder for oral liquid, 400 g

8478F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
(NP)	8	5	..	*402.94	41.30	Monogen [SB]

Carbohydrates**▪ MODIFIED LONG CHAIN AMYLOPECTIN****Restricted benefit**

Glycogen storage disease

modified long chain amylopectin powder for oral liquid, 30 x 60 g sachets

9386B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
(NP)	4	5	..	*650.90	41.30	Glycosade [VF]

Amino acids/carbohydrates/minerals/vitamins, combinations

■ AMINO ACID FORMULA WITH FAT, CARBOHYDRATE, VITAMINS, MINERALS, TRACE ELEMENTS AND MEDIUM CHAIN TRIGLYCERIDES

Authority required

Cows' milk protein enteropathy

Treatment Phase: Initial treatment for up to 6 months

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist, or in consultation with a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist.

Clinical criteria:

- The condition must not be isolated infant colic or reflux, **AND**
- Patient must be intolerant to both soy protein and protein hydrolysate formulae, as demonstrated when the child has failed to respond to a strict cows' milk protein free and strict soy protein free diet with a protein hydrolysate (with or without medium chain triglycerides) as the principal formula.

Population criteria:

- Patient must be up to the age of 24 months.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Severe cows' milk protein enteropathy with failure to thrive

Treatment Phase: Initial treatment for up to 6 months

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist, or in consultation with a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist.

Clinical criteria:

- The condition must not be isolated infant colic or reflux.

Population criteria:

- Patient must be up to the age of 24 months.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Combined intolerance to cows' milk protein, soy protein and protein hydrolysate formulae

Treatment Phase: Initial treatment for up to 6 months

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist.

Clinical criteria:

- The condition must not be isolated infant colic or reflux.

Population criteria:

- Patient must be older than 24 months of age.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Proven combined immunoglobulin E (IgE) mediated allergy to cows' milk protein and soy protein

Treatment Phase: Initial treatment for up to 6 months

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist, or in consultation with a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist.

Clinical criteria:

- Patient must have failed a trial of protein hydrolysate formulae (with or without medium chain triglycerides).

Population criteria:

- Patient must be up to the age of 24 months.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Eosinophilic oesophagitis

Treatment Phase: Initial treatment for up to 3 months

Treatment criteria:

- Must be treated by a clinical immunologist, suitably qualified allergist or gastroenterologist.

Clinical criteria:

- Patient must require an amino acid based formula as a component of a dietary elimination program.

Population criteria:

- Patient must be 18 years of age or less.
- Treatment with oral steroids should not be commenced during the period of initial treatment.
- Eosinophilic oesophagitis is demonstrated by the following criteria:

- (i) Chronic symptoms of reflux that persisted despite a 2-month trial of a proton pump inhibitor or chronic dysphagia; and
- (ii) A lack of demonstrable anatomic abnormality with the exception of stricture, which can be attributable to eosinophilic oesophagitis; and
- (iii) Eosinophilic infiltration of the oesophagus, demonstrated by oesophageal biopsy specimens obtained by endoscopy and where the most densely involved oesophageal biopsy had 20 or more eosinophils in any single 400 x high powered field, along with normal antral and duodenal biopsies.

The date of birth of the patient must be included in the authority application.

Note Authorities for increased maximum quantities, up to a maximum of 20, may be authorised.

amino acid formula with fat, carbohydrate, vitamins, minerals, trace elements and medium chain triglycerides powder for oral liquid, 400 g

10522T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	8	5	..	*342.94	41.30	Alfamino Junior [NT]

amino acid formula with fat, carbohydrate, vitamins, minerals, trace elements and medium chain triglycerides powder for oral liquid, 400 g

11161K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	8	5	..	*342.94	41.30	Neocate Junior [SB]

■ AMINO ACID FORMULA WITH FAT, CARBOHYDRATE, VITAMINS, MINERALS, TRACE ELEMENTS AND MEDIUM CHAIN TRIGLYCERIDES

Note Authorities for increased maximum quantities, up to a maximum of 20, may be authorised.

Authority required

Cows' milk protein enteropathy

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist, or have an appointment to be assessed by one of these specialists.

Clinical criteria:

- The condition must not be isolated infant colic or reflux, **AND**
- Patient must be intolerant to both soy protein and protein hydrolysate formulae, as demonstrated when the child has failed to respond to a strict cows' milk protein free and strict soy protein free diet with a protein hydrolysate (with or without medium chain triglycerides) as the principal formula.

Population criteria:

- Patient must be up to the age of 24 months.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Severe cows' milk protein enteropathy with failure to thrive

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist, or have been assessed at least once or have an appointment to be assessed by one of these specialists.

Clinical criteria:

- The condition must not be isolated infant colic or reflux, **AND**
- Patient must have had failure to thrive prior to commencement with initial treatment.

Population criteria:

- Patient must be up to the age of 24 months.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Combined intolerance to cows' milk protein, soy protein and protein hydrolysate formulae

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist at intervals not greater than 12 months.

Clinical criteria:

- The condition must not be isolated infant colic or reflux.

Population criteria:

- Patient must be older than 24 months of age.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Proven combined immunoglobulin E (IgE) mediated allergy to cows' milk protein and soy protein

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist.

Clinical criteria:

- Patient must have failed a trial of protein hydrolysate formulae (with or without medium chain triglycerides) prior to commencement with initial treatment.

Population criteria:

- Patient must be up to the age of 24 months.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Cows' milk anaphylaxis

Treatment criteria:

- Must be treated by a specialist allergist or clinical immunologist, or in consultation with a specialist allergist or clinical immunologist.

Population criteria:

- Patient must be up to the age of 24 months.

Anaphylaxis is defined as a severe and/or potentially life threatening allergic reaction.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Severe intestinal malabsorption including short bowel syndrome

Clinical criteria:

- Patient must have failed to respond to protein hydrolysate formulae; OR
- Patient must have been receiving parenteral nutrition.

Authority required

Eosinophilic oesophagitis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a clinical immunologist, suitably qualified allergist or gastroenterologist.

Clinical criteria:

- Patient must have responded to an initial course of PBS-subsidised treatment.

Population criteria:

- Patient must be 18 years of age or less.

Response to initial treatment is demonstrated by oesophageal biopsy specimens obtained by endoscopy, where the most densely involved oesophageal biopsy had 5 or less eosinophils in any single 400 x high powered field, along with normal antral and duodenal biopsies. The response criteria will not be deemed to have been met if oral steroids were commenced during initial treatment.

amino acid formula with fat, carbohydrate, vitamins, minerals, trace elements and medium chain triglycerides powder for oral liquid, 400 g

10527C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	8	5	..	*342.94	41.30	Alfamino Junior [NT]

amino acid formula with fat, carbohydrate, vitamins, minerals, trace elements and medium chain triglycerides powder for oral liquid, 400 g

11183N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	8	5	..	*342.94	41.30	Neocate Junior [SB]

Milk substitutes

▪ MILK POWDER SYNTHETIC LOW CALCIUM

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Restricted benefit

Hypercalcaemia

Population criteria:

- Patient must be under the age of 4 years.

milk powder synthetic low calcium powder for oral liquid, 400 g

3092R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	8	5	..	*363.18	41.30	Locasol [SB]

Other combinations of nutrients

▪ AMINO ACID FORMULA SUPPLEMENTED WITH PREBIOTICS, PROBIOTICS AND LONG CHAIN POLYUNSATURATED FATTY ACIDS

Note Authorities for increased maximum quantities, up to a maximum of 52, may be authorised.

Authority required

Eosinophilic oesophagitis

Treatment Phase: Initial treatment for up to 3 months

Treatment criteria:

- Must be treated by a clinical immunologist, suitably qualified allergist or gastroenterologist.

Clinical criteria:

- Patient must require an amino acid based formula as a component of a dietary elimination program.

Population criteria:

- Patient must be 18 years of age or less.

Treatment with oral steroids should not be commenced during the period of initial treatment.

Eosinophilic oesophagitis is demonstrated by the following criteria:

- (i) Chronic symptoms of reflux that persisted despite a 2-month trial of a proton pump inhibitor or chronic dysphagia; and
- (ii) A lack of demonstrable anatomic abnormality with the exception of stricture, which can be attributable to eosinophilic oesophagitis; and
- (iii) Eosinophilic infiltration of the oesophagus, demonstrated by oesophageal biopsy specimens obtained by endoscopy and where the most densely involved oesophageal biopsy had 20 or more eosinophils in any single 400 x high powered field, along with normal antral and duodenal biopsies.

The date of birth of the patient must be included in the authority application.

Authority required

Eosinophilic oesophagitis

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a clinical immunologist, suitably qualified allergist or gastroenterologist.

Clinical criteria:

- Patient must have responded to an initial course of PBS-subsidised treatment.

Population criteria:

- Patient must be 18 years of age or less.

Response to initial treatment is demonstrated by oesophageal biopsy specimens obtained by endoscopy, where the most densely involved oesophageal biopsy had 5 or less eosinophils in any single 400 x high powered field, along with normal antral and duodenal biopsies. The response criteria will not be deemed to have been met if oral steroids were commenced during initial treatment.

amino acid formula supplemented with prebiotics, probiotics and long chain polyunsaturated fatty acids powder for oral liquid, 400 g

11343B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	12	5	..	*500.82	41.30	Neocate Syneo [SB]

■ AMINO ACID FORMULA SUPPLEMENTED WITH PREBIOTICS, PROBIOTICS AND LONG CHAIN POLYUNSATURATED FATTY ACIDS

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Cows' milk protein enteropathy

Treatment Phase: Initial treatment for up to 6 months

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist, or in consultation with a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist.

Clinical criteria:

- The condition must not be isolated infant colic or reflux, **AND**
- Patient must be intolerant to both soy protein and protein hydrolysate formulae, as demonstrated when the child has failed to respond to a strict cows' milk protein free and strict soy protein free diet with a protein hydrolysate (with or without medium chain triglycerides) as the principal formula.

Population criteria:

- Patient must be up to the age of 24 months.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Severe cows' milk protein enteropathy with failure to thrive

Treatment Phase: Initial treatment for up to 6 months

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist, or in consultation with a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist.

Clinical criteria:

- The condition must not be isolated infant colic or reflux.

Population criteria:

- Patient must be up to the age of 24 months.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Combined intolerance to cows' milk protein, soy protein and protein hydrolysate formulae

Treatment Phase: Initial treatment for up to 6 months

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist.

Clinical criteria:

- The condition must not be isolated infant colic or reflux.

Population criteria:

- Patient must be older than 24 months of age.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Proven combined immunoglobulin E (IgE) mediated allergy to cows' milk protein and soy protein

Treatment Phase: Initial treatment for up to 6 months

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist, or in consultation with a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist.

Clinical criteria:

- Patient must have failed a trial of protein hydrolysate formulae (with or without medium chain triglycerides).

Population criteria:

- Patient must be up to the age of 24 months.

The name of the specialist and the date of birth of the patient must be included in the authority application.

amino acid formula supplemented with prebiotics, probiotics and long chain polyunsaturated fatty acids powder for oral liquid, 400 g

11331J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	8	5	..	*336.22	41.30	Neocate Syneo [SB]

■ AMINO ACID FORMULA SUPPLEMENTED WITH PREBIOTICS, PROBIOTICS AND LONG CHAIN POLYUNSATURATED FATTY ACIDS

Note Authorities for increased maximum quantities, up to a maximum of 20, may be authorised.

Authority required

Cows' milk anaphylaxis

Treatment criteria:

- Must be treated by a specialist allergist or clinical immunologist, or in consultation with a specialist allergist or clinical immunologist.

Population criteria:

- Patient must be up to the age of 24 months.

Anaphylaxis is defined as a severe and/or potentially life threatening allergic reaction.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Cows' milk protein enteropathy

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist, or have an appointment to be assessed by one of these specialists.

Clinical criteria:

- The condition must not be isolated infant colic or reflux, **AND**
- Patient must be intolerant to both soy protein and protein hydrolysate formulae, as demonstrated when the child has failed to respond to a strict cows' milk protein free and strict soy protein free diet with a protein hydrolysate (with or without medium chain triglycerides) as the principal formula.

Population criteria:

- Patient must be up to the age of 24 months.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Severe cows' milk protein enteropathy with failure to thrive

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist, or have been assessed at least once or have an appointment to be assessed by one of these specialists.

Clinical criteria:

- The condition must not be isolated infant colic or reflux, **AND**
- Patient must have had failure to thrive prior to commencement with initial treatment.

Population criteria:

- Patient must be up to the age of 24 months.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Combined intolerance to cows' milk protein, soy protein and protein hydrolysate formulae

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist at intervals not greater than 12 months.

Clinical criteria:

- The condition must not be isolated infant colic or reflux.

Population criteria:

- Patient must be older than 24 months of age.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Proven combined immunoglobulin E (IgE) mediated allergy to cows' milk protein and soy protein

Treatment Phase: Continuing treatment

Treatment criteria:

- Must be treated by a specialist allergist, clinical immunologist or specialist paediatric gastroenterologist and hepatologist.

Clinical criteria:

- Patient must have failed a trial of protein hydrolysate formulae (with or without medium chain triglycerides) prior to commencement with initial treatment.

Population criteria:

- Patient must be up to the age of 24 months.

The name of the specialist and the date of birth of the patient must be included in the authority application.

Authority required

Severe intestinal malabsorption including short bowel syndrome

Clinical criteria:

- Patient must have failed to respond to protein hydrolysate formulae; OR
- Patient must have been receiving parenteral nutrition.

Note A risk/benefit analysis prior to treatment, and continuous patient monitoring from a health care professional is required for the use of this product, for this indication.

amino acid formula supplemented with prebiotics, probiotics and long chain polyunsaturated fatty acids powder for oral liquid, 400 g

11340W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
(NP)	8	5	..	*336.22	41.30	Neocate Syneo [SB]

AMINO ACID FORMULA WITH CARBOHYDRATE WITHOUT PHENYLALANINE

Note This product does not contain any vitamins, minerals or trace elements and is not intended as a sole source of nutrition

Restricted benefit

Phenylketonuria

amino acid formula with carbohydrate without phenylalanine modified release tablet, 6 x 77

12072J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
(NP)	4	5	..	*1321.66	41.30	PKU Easy Tablet [OH]

AMINO ACID FORMULA WITH CARBOHYDRATE, VITAMINS, MINERALS AND TRACE ELEMENTS WITHOUT PHENYLALANINE

Restricted benefit

Phenylketonuria

amino acid formula with carbohydrate, vitamins, minerals and trace elements without phenylalanine powder for oral liquid, 30 x 20 g sachets

10806R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
(NP)	4	5	..	*1003.30	41.30	PKU Go [OH]

AMINO ACID FORMULA WITH FAT, CARBOHYDRATE WITHOUT METHIONINE

Note This product does not contain any vitamins, minerals or trace elements and is not intended as a sole source of nutrition

Restricted benefit

Pyridoxine non-responsive homocystinuria

amino acid formula with fat, carbohydrate without methionine modified release tablet, 6 x 77

12006X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
(NP)	5	5	..	*2961.34	41.30	HCU Easy Tablet [OH]

AMINO ACID FORMULA WITH FAT, CARBOHYDRATE WITHOUT PHENYLALANINE

Restricted benefit

Phenylketonuria

amino acid formula with fat, carbohydrate without phenylalanine tablet: modified release, 4 x 110 g

10683G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
(NP)	7	5	..	*1849.23	41.30	PKU Easy Microtabs [OH]

■ AMINO ACID FORMULA WITH FAT, CARBOHYDRATE WITHOUT PHENYLALANINE AND TYROSINE

Note This product does not contain any vitamins, minerals or trace elements and is not intended as a sole source of nutrition

Restricted benefit

Tyrosinaemia

amino acid formula with fat, carbohydrate without phenylalanine and tyrosine modified release tablet, 6 x 77

12015J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	5	..	*2401.30	41.30	TYR Easy Tablet [OH]

■ AMINO ACID FORMULA WITH FAT, CARBOHYDRATE WITHOUT VALINE, LEUCINE AND ISOLEUCINE

Note This product does not contain any vitamins, minerals or trace elements and is not intended as a sole source of nutrition

Restricted benefit

Maple syrup urine disease

amino acid formula with fat, carbohydrate without valine, leucine and isoleucine modified release tablet, 6 x 77

12014H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	5	5	..	*2961.34	41.30	MSUD Easy Tablet [OH]

■ AMINO ACID FORMULA WITH FAT, CARBOHYDRATE, VITAMINS, MINERALS AND TRACE ELEMENTS WITHOUT METHIONINE AND SUPPLEMENTED WITH DOCOSAHEXAENOIC ACID

Restricted benefit

Pyridoxine non-responsive homocystinuria

amino acid formula with fat, carbohydrate, vitamins, minerals and trace elements without methionine and supplemented with docosahexaenoic acid oral liquid, 36 x 125 mL bottles

3417W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	5	..	*2180.34	41.30	HCU Anamix junior LQ [SB]

■ AMINO ACID FORMULA WITH FAT, CARBOHYDRATE, VITAMINS, MINERALS AND TRACE ELEMENTS WITHOUT PHENYLALANINE

Restricted benefit

Phenylketonuria

amino acid formula with fat, carbohydrate, vitamins, minerals and trace elements without phenylalanine oral liquid: powder for, 30 x 34 g bottles

10632N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	5	5	..	*1826.24	41.30	PKU Easy Shake & Go [OH]

■ AMINO ACID FORMULA WITH FAT, CARBOHYDRATE, VITAMINS, MINERALS AND TRACE ELEMENTS WITHOUT PHENYLALANINE AND TYROSINE, AND SUPPLEMENTED WITH DOCOSAHEXAENOIC ACID

Restricted benefit

Tyrosinaemia

amino acid formula with fat, carbohydrate, vitamins, minerals and trace elements without phenylalanine and tyrosine, and supplemented with docosahexaenoic acid oral liquid, 36 x 125 mL bottles

9330C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	5	..	*2180.34	41.30	TYR Anamix junior LQ [SB]

■ AMINO ACID FORMULA WITH FAT, CARBOHYDRATE, VITAMINS, MINERALS AND TRACE ELEMENTS, WITHOUT PHENYLALANINE AND TYROSINE

Restricted benefit

Tyrosinaemia

amino acid formula with fat, carbohydrate, vitamins, minerals and trace elements, without phenylalanine and tyrosine powder for oral liquid, 30 x 34 g bottles

10934L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	5	..	*2965.58	41.30	TYR Easy Shake & Go [OH]

■ AMINO ACID FORMULA WITH VITAMINS AND MINERALS WITHOUT LYSINE AND LOW IN TRYPTOPHAN

Restricted benefit

Proven glutaric aciduria type 1

amino acid formula with vitamins and minerals without lysine and low in tryptophan powder for oral liquid, 30 x 18 g sachets

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
10715Y	8	5	..	*1830.70	41.30	GA1 Anamix Junior [NU]

■ AMINO ACID FORMULA WITH VITAMINS AND MINERALS WITHOUT LYSINE AND LOW IN TRYPTOPHAN**Restricted benefit**

Proven glutaric aciduria type 1

amino acid formula with vitamins and minerals without lysine and low in tryptophan powder for oral liquid, 400 g

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
2650L	8	5	..	*665.58	41.30	GA1 Anamix infant [SB]

amino acid formula with vitamins and minerals without lysine and low in tryptophan powder for oral liquid, 500 g

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
10466W	9	5	..	*2668.77	41.30	XLYS, LOW TRY Maxamum [SB]

■ AMINO ACID FORMULA WITH VITAMINS AND MINERALS WITHOUT LYSINE AND LOW IN TRYPTOPHAN**Restricted benefit**

Proven glutaric aciduria type 1

Restricted benefit

Pyridoxine dependent epilepsy

Clinical criteria:

- Patient must be managed on a low lysine diet for pyridoxine dependent epilepsy, **AND**
- The condition must be treated by or in consultation with a metabolic physician.

amino acid formula with vitamins and minerals without lysine and low in tryptophan powder for oral liquid, 30 x 24 g sachets

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
9438R	4	5	..	*1830.66	41.30	GA gel [VF]

amino acid formula with vitamins and minerals without lysine and low in tryptophan powder for oral liquid, 30 x 25 g sachets

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
5484P	4	5	..	*2733.06	41.30	GA express 15 [VF]

■ AMINO ACID FORMULA WITH VITAMINS AND MINERALS WITHOUT METHIONINE**Restricted benefit**

Pyridoxine non-responsive homocystinuria

amino acid formula with vitamins and minerals without methionine oral liquid, 30 x 87 mL sachets

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
2639X	4	5	..	*1742.34	41.30	HCU cooler 10 [VF]

amino acid formula with vitamins and minerals without methionine powder for oral liquid, 30 x 24 g sachets

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8677Q	4	5	..	*1742.34	41.30	HCU gel [VF]

amino acid formula with vitamins and minerals without methionine powder for oral liquid, 30 x 25 g sachets

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8744F	4	5	..	*2558.94	41.30	HCU express 15 [VF]

amino acid formula with vitamins and minerals without methionine powder for oral liquid, 500 g

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8416Y	8	5	..	*2239.26	41.30	XMET Maxamum [SB]

amino acid formula with vitamins and minerals without methionine oral liquid, 30 x 130 mL pouches

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
9133Q	4	5	..	*2558.94	41.30	HCU cooler 15 [VF]

amino acid formula with vitamins and minerals without methionine oral liquid, 30 x 125 mL pouches

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
1548L	3	5	..	*2558.91	41.30	HCU Lophlex LQ 20 [SB]

amino acid formula with vitamins and minerals without methionine oral liquid, 30 x 174 mL sachets

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
2640Y	4	5	..	*3358.18	41.30	HCU cooler 20 [VF]

NP

■ AMINO ACID FORMULA WITH VITAMINS AND MINERALS WITHOUT METHIONINE**Restricted benefit**

Pyridoxine non-responsive homocystinuria

amino acid formula with vitamins and minerals without methionine oral liquid: powder for, 30 x 36 g sachets

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
10693T	4	5	..	*1742.34	41.30	HCU Anamix Junior [NU]

NP

amino acid formula with vitamins and minerals without methionine powder for oral liquid, 30 x 29 g sachets

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
12273Y	4	5	..	*3358.18	41.30	HCU Lophlex [SB]

NP

■ AMINO ACID FORMULA WITH VITAMINS AND MINERALS WITHOUT METHIONINE**Restricted benefit**

Pyridoxine non-responsive homocystinuria

Population criteria:

- Patient must be an infant or a very young child.

amino acid formula with vitamins and minerals without methionine powder for oral liquid, 400 g

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8417B	8	5	..	*632.62	41.30	HCU Anamix infant [SB]

NP

■ AMINO ACID FORMULA WITH VITAMINS AND MINERALS WITHOUT METHIONINE, THREONINE AND VALINE AND LOW IN ISOLEUCINE**Restricted benefit**

Methylmalonic acidaemia

Restricted benefit

Propionic acidaemia

amino acid formula with vitamins and minerals without methionine, threonine and valine and low in isoleucine powder for oral liquid, 400 g

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8058D	8	5	..	*632.62	41.30	MMA/PA Anamix infant [SB]

NP

amino acid formula with vitamins and minerals without methionine, threonine and valine and low in isoleucine powder for oral liquid, 30 x 24 g sachets

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
3444G	4	5	..	*1742.34	41.30	MMA/PA gel [VF]

NP

amino acid formula with vitamins and minerals without methionine, threonine and valine and low in isoleucine powder for oral liquid, 500 g

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8061G	8	5	..	*2239.26	41.30	XMTVI Maxamum [SB]

NP

amino acid formula with vitamins and minerals without methionine, threonine and valine and low in isoleucine powder for oral liquid, 30 x 25 g sachets

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
3443F	4	5	..	*2558.94	41.30	MMA/PA express 15 [VF]

NP

■ AMINO ACID FORMULA WITH VITAMINS AND MINERALS WITHOUT METHIONINE, THREONINE AND VALINE AND LOW IN ISOLEUCINE**Restricted benefit**

Methylmalonic acidaemia

Restricted benefit

Propionic acidaemia

amino acid formula with vitamins and minerals without methionine, threonine and valine and low in isoleucine powder for oral liquid, 30 x 18 g sachets

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
10730R	8	5	..	*1742.30	41.30	MMA/PA Anamix Junior [NU]

NP

■ AMINO ACID FORMULA WITH VITAMINS AND MINERALS WITHOUT PHENYLALANINE**Restricted benefit**

Phenylketonuria

amino acid formula with vitamins and minerals without phenylalanine oral liquid, 30 x 174 mL cans

2474F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	5	..	*1690.86	41.30	PKU Cooler 20 [VF]

amino acid formula with vitamins and minerals without phenylalanine oral liquid, 30 x 130 mL pouches

10410X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	5	..	*1268.66	41.30	PKU Air 15 [VF]

amino acid formula with vitamins and minerals without phenylalanine powder for oral liquid, 30 x 24 g sachets

8555G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	5	..	*867.10	41.30	PKU gel [VF]

amino acid formula with vitamins and minerals without phenylalanine oral liquid, 30 x 174 mL pouches

10411Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	5	..	*1690.86	41.30	PKU Air 20 [VF]

amino acid formula with vitamins and minerals without phenylalanine powder for oral liquid, 500 g

2739E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	8	5	..	*1108.06	41.30	XP Maxamum [SB]

amino acid formula with vitamins and minerals without phenylalanine oral liquid, 60 x 62.5 mL pouches

9397N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*867.68	41.30	PKU Lophlex LQ 10 [SB]

amino acid formula with vitamins and minerals without phenylalanine powder for oral liquid, 30 x 34 g sachets

1909L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	5	..	*1690.86	41.30	PKU express 20 [VF]

amino acid formula with vitamins and minerals without phenylalanine oral liquid, 36 x 125 mL bottles

9396M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	5	..	*1040.30	41.30	PKU Anamix Junior LQ [SB]

amino acid formula with vitamins and minerals without phenylalanine oral semi-solid, 36 x 109 g tubs

2806Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	3	5	..	*1519.20	41.30	PKU Lophlex Sensation 20 [SB]

amino acid formula with vitamins and minerals without phenylalanine oral liquid, 18 x 250 mL

8746H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	5	5	..	*1075.94	41.30	Easiphen [SB]

amino acid formula with vitamins and minerals without phenylalanine oral liquid, 30 x 87 mL cans

2382J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	5	..	*848.18	41.30	PKU Cooler 10 [VF]

amino acid formula with vitamins and minerals without phenylalanine oral liquid, 30 x 130 mL cans

8846N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	5	..	*1268.66	41.30	PKU Cooler 15 [VF]

amino acid formula with vitamins and minerals without phenylalanine oral liquid, 30 x 85 g sachets

5483N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	5	..	*867.10	41.30	PKU squeezie [VF]

amino acid formula with vitamins and minerals without phenylalanine powder for oral liquid, 30 x 28 g sachets

12136R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	5	..	*1671.54	41.30	PKU Lophlex [SB]

amino acid formula with vitamins and minerals without phenylalanine powder for oral liquid, 30 x 25 g sachets

8591E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	5	..	*1269.34	41.30	PKU express 15 [VF]

amino acid formula with vitamins and minerals without phenylalanine oral liquid, 30 x 125 mL pouches

9021T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	5	..	*1671.54	41.30	PKU Lophlex LQ 20 [SB]

■ AMINO ACID FORMULA WITH VITAMINS AND MINERALS WITHOUT PHENYLALANINE

Note Changes in vitamin D levels and amino acid composition have occurred with a recent formulation change.

Restricted benefit

Phenylketonuria

amino acid formula with vitamins and minerals without phenylalanine oral liquid: powder for, 30 x 36 g sachets

10258X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	5	..	*868.06	41.30	PKU Anamix Junior [SB]

■ AMINO ACID FORMULA WITH VITAMINS AND MINERALS WITHOUT PHENYLALANINE AND TYROSINERestricted benefit

Tyrosinaemia

amino acid formula with vitamins and minerals without phenylalanine and tyrosine powder for oral liquid, 500 g

3078B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	8	5	..	*2239.26	41.30	XPhen, Tyr Maxamum [SB]

amino acid formula with vitamins and minerals without phenylalanine and tyrosine powder for oral liquid, 400 g

8445L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	8	5	..	*632.62	41.30	TYR Anamix infant [SB]

amino acid formula with vitamins and minerals without phenylalanine and tyrosine oral liquid, 30 x 87 mL pouches

2674R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	5	..	*1742.34	41.30	TYR cooler 10 [VF]

amino acid formula with vitamins and minerals without phenylalanine and tyrosine powder for oral liquid, 30 x 34 g sachets

11151X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	5	..	*3358.18	41.30	TYR express 20 [VF]

amino acid formula with vitamins and minerals without phenylalanine and tyrosine powder for oral liquid, 30 x 25 g sachets

8667E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	5	..	*2558.94	41.30	TYR express 15 [VF]

amino acid formula with vitamins and minerals without phenylalanine and tyrosine powder for oral liquid, 30 x 24 g sachets

8631G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	5	..	*1742.34	41.30	TYR gel [VF]

amino acid formula with vitamins and minerals without phenylalanine and tyrosine oral liquid, 30 x 130 mL pouches

9132P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	5	..	*2558.94	41.30	TYR cooler 15 [VF]

amino acid formula with vitamins and minerals without phenylalanine and tyrosine powder for oral liquid, 30 x 28 g sachets

12274B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	5	..	*3358.18	41.30	TYR Lophlex [SB]

amino acid formula with vitamins and minerals without phenylalanine and tyrosine oral liquid, 30 x 174 mL pouches

2701E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	5	..	*3358.18	41.30	TYR cooler 20 [VF]

amino acid formula with vitamins and minerals without phenylalanine and tyrosine oral liquid, 30 x 125 mL pouches

1547K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	3	5	..	*2558.91	41.30	TYR Lophlex LQ 20 [SB]

■ AMINO ACID FORMULA WITH VITAMINS AND MINERALS WITHOUT PHENYLALANINE AND TYROSINE

Note Changes in vitamin D levels and amino acid composition have occurred with a recent formulation change.

Restricted benefit

Tyrosinaemia

amino acid formula with vitamins and minerals without phenylalanine and tyrosine oral liquid: powder for, 30 x 36 g sachets

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
10260B	4	5	..	*1742.34	41.30	TYR Anamix Junior [SB]

NP

AMINO ACID FORMULA WITH VITAMINS AND MINERALS WITHOUT VALINE, LEUCINE AND ISOLEUCINE**Restricted benefit**

Maple syrup urine disease

amino acid formula with vitamins and minerals without valine, leucine and isoleucine powder for oral liquid, 500 g

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8057C	8	5	..	*2239.26	41.30	MSUD Maxamum [SB]

NP

amino acid formula with vitamins and minerals without valine, leucine and isoleucine powder for oral liquid, 500 g

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8310J	4	5	..	*2212.58	41.30	MSUD AID III [SB]

NP

amino acid formula with vitamins and minerals without valine, leucine and isoleucine powder for oral liquid, 30 x 24 g sachets

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8592F	4	5	..	*1742.34	41.30	MSUD gel [VF]

NP

amino acid formula with vitamins and minerals without valine, leucine and isoleucine oral liquid, 30 x 125 mL pouches

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
1546J	3	5	..	*2558.91	41.30	MSUD Lophlex LQ 20 [SB]

NP

amino acid formula with vitamins and minerals without valine, leucine and isoleucine powder for oral liquid, 30 x 25 g sachets

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8632H	4	5	..	*2558.94	41.30	MSUD express 15 [VF]

NP

amino acid formula with vitamins and minerals without valine, leucine and isoleucine oral liquid, 30 x 174 mL pouches

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
2654Q	4	5	..	*3358.18	41.30	MSUD cooler 20 [VF]

NP

amino acid formula with vitamins and minerals without valine, leucine and isoleucine powder for oral liquid, 30 x 34 g sachets

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
1914R	4	5	..	*3367.74	41.30	MSUD express 20 [VF]

NP

amino acid formula with vitamins and minerals without valine, leucine and isoleucine powder for oral liquid, 400 g

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
2380G	8	5	..	*632.62	41.30	MSUD Anamix infant [SB]

NP

amino acid formula with vitamins and minerals without valine, leucine and isoleucine oral liquid, 30 x 87 mL pouches

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
2651M	4	5	..	*1742.34	41.30	MSUD cooler 10 [VF]

NP

amino acid formula with vitamins and minerals without valine, leucine and isoleucine powder for oral liquid, 30 x 28 g sachets

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
12285N	4	5	..	*3358.18	41.30	MSUD Lophlex [SB]

NP

amino acid formula with vitamins and minerals without valine, leucine and isoleucine oral liquid, 30 x 130 mL pouches

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
2375B	4	5	..	*2558.94	41.30	MSUD cooler 15 [VF]

NP

■ AMINO ACID FORMULA WITH VITAMINS AND MINERALS WITHOUT VALINE, LEUCINE AND ISOLEUCINE

Note Changes in vitamin D levels and amino acid composition have occurred with a recent formulation change.

Restricted benefit

Maple syrup urine disease

amino acid formula with vitamins and minerals without valine, leucine and isoleucine oral liquid: powder for, 30 x 36 g sachets

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
10259Y	4	5	..	*1742.34	41.30	MSUD Anamix Junior [SB]

NP

■ AMINO ACID FORMULA WITH VITAMINS AND MINERALS WITHOUT VALINE, LEUCINE AND ISOLEUCINE WITH FAT, CARBOHYDRATE AND TRACE ELEMENTS AND SUPPLEMENTED WITH DOCOSAHEXAENOIC ACID

Restricted benefit

Maple syrup urine disease

amino acid formula with vitamins and minerals without valine, leucine and isoleucine with fat, carbohydrate and trace elements and supplemented with docosahexaenoic acid oral liquid, 36 x 125 mL bottles

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
9499Y	4	5	..	*2180.34	41.30	MSUD Anamix Junior LQ [SB]

NP

■ AMINO ACID FORMULA WITH VITAMINS AND MINERALS, LOW PHENYLALANINE AND SUPPLEMENTED WITH DOCOSAHEXAENOIC ACID AND ARACHIDONIC ACID

Restricted benefit

Phenylketonuria

amino acid formula with vitamins and minerals, low phenylalanine and supplemented with docosahexaenoic acid and arachidonic acid powder for oral liquid, 30 x 25 g sachets

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
11836Y	4	5	..	*1003.26	41.30	PKU Explore 10 [VF]

NP

amino acid formula with vitamins and minerals, low phenylalanine and supplemented with docosahexaenoic acid and arachidonic acid powder for oral liquid, 30 x 12.5 g sachets

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
11185Q	8	5	..	*1003.26	41.30	PKU Anamix First Spoon [SB]

NP

amino acid formula with vitamins and minerals, low phenylalanine and supplemented with docosahexaenoic acid and arachidonic acid powder for oral liquid, 30 x 12.5 g sachets

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
11859E	8	5	..	*1003.26	41.30	PKU Explore 5 [VF]

NP

■ AMINO ACID FORMULA WITH VITAMINS, MINERALS AND LONG CHAIN POLYUNSATURATED FATTY ACIDS WITHOUT PHENYLALANINE

Restricted benefit

Phenylketonuria

amino acid formula with vitamins, minerals and long chain polyunsaturated fatty acids without phenylalanine powder for oral liquid, 400 g

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
8479G	8	5	..	*578.70	41.30	PKU Anamix infant [SB]

NP

■ AMINO ACID FORMULA WITH VITAMINS, MINERALS AND LONG CHAIN POLYUNSATURATED FATTY ACIDS WITHOUT PHENYLALANINE

Note The level of iron in this product is below the recommended daily intake (RDI) for infants and should be supplemented by other sources where appropriate.

Restricted benefit

Phenylketonuria

amino acid formula with vitamins, minerals and long chain polyunsaturated fatty acids without phenylalanine powder for oral liquid, 400 g

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
11653H	8	5	..	*615.18	41.30	PKU Start [VF]

NP

■ AMINO ACID FORMULA WITHOUT PHENYLALANINE

Restricted benefit

Phenylketonuria

amino acid formula without phenylalanine 1 g tablet, 75

8678R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	24	5	..	*1169.34	41.30	Phlexy-10 [SB]

■ AMINO ACID FORMULA WITHOUT VALINE, LEUCINE AND ISOLEUCINE**Restricted benefit**

Maple syrup urine disease

amino acid formula without valine, leucine and isoleucine containing 5 g of protein equivalent powder for oral liquid, 30 x 6 g sachets

10161T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	12	5	..	*2965.50	41.30	MSUD amino5 [VF]

■ ARACHIDONIC ACID AND DOCOSAHEXAENOIC ACID WITH CARBOHYDRATE**Restricted benefit**

Peroxisomal biogenesis disorders

arachidonic acid and docosahexaenoic acid with carbohydrate containing 200 mg arachidonic acid and 100 mg docosahexaenoic acid oral liquid: powder for, 30 x 4 g sachets

10036F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	5	..	*352.50	41.30	keyomega [VF]

■ ARGININE WITH CARBOHYDRATE**Note** Arginine with carbohydrate is not indicated for the treatment of arginase deficiency and other inborn errors of protein metabolism.**Restricted benefit**

Urea cycle disorders

arginine with carbohydrate containing 5 g arginine oral liquid: powder for, 30 x 7.6 g sachets

10093F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	5	..	*880.14	41.30	Arginine 5000 [VF]

arginine with carbohydrate containing 2 g arginine oral liquid: powder for, 30 x 4 g sachets

5482M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	5	..	*666.90	41.30	Arginine 2000 [VF]

arginine with carbohydrate containing 500 mg arginine oral liquid: powder for, 30 x 4 g sachets

9437Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	5	..	*446.94	41.30	Arginine 500 [VF]

■ CARBOHYDRATE, FAT, VITAMINS, MINERALS AND TRACE ELEMENTS**Restricted benefit**

Proven inborn errors of protein metabolism

Clinical criteria:

- Patient must be unable to meet their energy requirements with permitted food and formulae.

carbohydrate, fat, vitamins, minerals and trace elements powder for oral liquid, 400 g

8369L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	8	5	..	*257.58	41.30	Energivit [SB]

■ CARBOHYDRATES, FAT, VITAMINS, MINERALS, TRACE ELEMENTS AND SUPPLEMENTED WITH ARACHIDONIC ACID AND DOCOSAHEXAENOIC ACID**Restricted benefit**

Proven inborn errors of protein metabolism

Clinical criteria:

- Patient must be unable to meet their energy requirements with permitted food and formulae.

carbohydrates, fat, vitamins, minerals, trace elements and supplemented with arachidonic acid and docosahexaenoic acid providing 200 kilocalories powder for oral liquid, 30 x 43 g sachets

10039J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	5	..	*453.66	41.30	basecal 200 [VF]

■ CITRULLINE**Note** Citrulline is not indicated for the treatment of arginase deficiency and other inborn errors of protein metabolism**Restricted benefit**

Urea cycle disorders

Clinical criteria:

- The treatment must be for preventing low plasma arginine levels; OR
- The treatment must be for preventing low citrulline levels.

citrulline 1 g tablet, 300

10736C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	5	..	1143.50	41.30	Citrulline Easy [OH]

■ CITRULLINE WITH CARBOHYDRATE

Note Citrulline with carbohydrate is not indicated for the treatment of arginase deficiency and other inborn errors of protein metabolism.

Restricted benefit

Urea cycle disorders

Clinical criteria:

- The treatment must be for preventing low plasma arginine levels; OR
- The treatment must be for preventing low citrulline levels.

citrulline with carbohydrate containing 1 g citrulline powder for oral liquid, 30 x 4 g sachets

5481L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	5	..	*446.94	41.30	Citrulline 1000 [VF]

■ DOCOSAHEXAENOIC ACID WITH CARBOHYDRATE**Restricted benefit**

Peroxisomal biogenesis disorders

docosahexaenoic acid with carbohydrate containing 200 mg docosahexaenoic acid oral liquid: powder for, 30 x 4g sachets

10040K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	5	..	*352.50	41.30	docomega [VF]

■ ESSENTIAL AMINO ACIDS FORMULA**Restricted benefit**

Gyrate atrophy of the choroid and retina

Restricted benefit

Urea cycle disorders

essential amino acids formula powder for oral liquid, 2 x 200 g

9329B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	6	5	..	*1031.82	41.30	Essential Amino Acid Mix [SB]

■ ESSENTIAL AMINO ACIDS FORMULA WITH MINERALS AND VITAMIN C**Restricted benefit**

Gyrate atrophy of the choroid and retina

Restricted benefit

Urea cycle disorders

essential amino acids formula with minerals and vitamin C powder for oral liquid, 400 g

2027Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	5	5	..	*521.84	41.30	Dialamine [SB]

■ ESSENTIAL AMINO ACIDS FORMULA WITH VITAMINS AND MINERALS**Restricted benefit**

Gyrate atrophy of the choroid and retina

Restricted benefit

Urea cycle disorders

essential amino acids formula with vitamins and minerals powder for oral liquid, 50 x 12.5 g sachets

9385Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	5	..	*1304.62	41.30	EAA Supplement [VF]

■ GLYCINE WITH CARBOHYDRATE**Restricted benefit**

Isovaleric acidemia

glycine with carbohydrate containing 500 mg glycine powder for oral liquid, 30 x 4 g sachets

10195N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	5	..	*495.78	41.30	Glycine500 [VF]

■ GLYCOMACROPEPTIDE AND ESSENTIAL AMINO ACID FORMULA WITH VITAMINS, MINERALS, AND LOW IN TYROSINE AND PHENYLALANINE

Restricted benefit

Tyrosinaemia

glycomacropeptide and essential amino acid formula with vitamins, minerals, and low in tyrosine and phenylalanine powder for oral liquid, 30 x 35 g sachets

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
11832R	4	5	..	*4097.18	41.30	TYR Sphere20 [VF]

glycomacropeptide and essential amino acid formula with vitamins, minerals, and low in tyrosine and phenylalanine powder for oral liquid, 30 x 31 g sachets

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
12175T	4	5	..	*3358.18	41.30	Tylactin Build 20 [QH]

■ GLYCOMACROPEPTIDE AND ESSENTIAL AMINO ACIDS WITH VITAMINS AND MINERALS

Restricted benefit

Tyrosinaemia

glycomacropeptide and essential amino acids with vitamins and minerals containing 15 g of protein equivalent oral liquid, 30 x 250 mL cartons

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
10528D	4	5	..	*3113.18	41.30	Tylactin RTD [QH]

glycomacropeptide and essential amino acids with vitamins and minerals containing 15 g of protein equivalent bar, 14 x 81 g

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
11290F	8	5	..	*2916.38	41.30	Tylactin Complete [QH]

■ GLYCOMACROPEPTIDE AND ESSENTIAL AMINO ACIDS WITH VITAMINS AND MINERALS

Restricted benefit

Phenylketonuria

glycomacropeptide and essential amino acids with vitamins and minerals powder for oral liquid, 60 x 20 g sachets

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
11084J	5	5	..	*1316.44	41.30	PKU Restore [QH]

glycomacropeptide and essential amino acids with vitamins and minerals powder for oral liquid, 30 x 51 g sachets

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
10992M	4	5	..	*2083.98	41.30	PKU Bettermilk Lite [QH]

glycomacropeptide and essential amino acids with vitamins and minerals powder for oral liquid, 30 x 49 g sachets

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
10652P	4	5	..	*1578.94	41.30	Camino Pro Bettermilk [QH]

glycomacropeptide and essential amino acids with vitamins and minerals bar, 7 x 81 g

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
2644E	14	5	..	*1292.94	41.30	Camino Pro Complete [QH]

■ GLYCOMACROPEPTIDE AND ESSENTIAL AMINO ACIDS WITH VITAMINS AND MINERALS

Note This product contains higher vitamin A levels than other PBS-listed glycomacropeptide products.

Restricted benefit

Phenylketonuria

glycomacropeptide and essential amino acids with vitamins and minerals containing 15 g of protein equivalent oral liquid, 30 x 250 mL cartons

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
11640P	4	5	..	*1578.90	41.30	PKU Glytactin RTD 15 Lite [QH]

glycomacropeptide and essential amino acids with vitamins and minerals containing 15 g protein oral liquid, 30 x 250 mL cartons

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
10332T	4	5	..	*1578.90	41.30	PKU Glytactin RTD 15 [QH]

■ GLYCOMACROPEPTIDE AND ESSENTIAL AMINO ACIDS WITH VITAMINS AND MINERALS

Note This product is low in folic acid, choline and methionine and is not intended as a sole source of nutrition.

Restricted benefit

Phenylketonuria

glycomacropeptide and essential amino acids with vitamins and minerals containing 10 g of protein equivalent powder for oral liquid, 60 x 16 g sachets

11287C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
(NP)	2	5	..	*1073.90	41.30	PKU Build 10 [QH]

glycomacropeptide and essential amino acids with vitamins and minerals containing 20 g of protein equivalent powder for oral liquid, 30 x 32 g sachets

11279P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
(NP)	4	5	..	*2083.98	41.30	PKU Build 20 [QH]

■ GLYCOMACROPEPTIDE FORMULA WITH DOCOSAHEXAENOIC ACID AND LOW PHENYLALANINE

Restricted benefit

Phenylketonuria

glycomacropeptide formula with docosahexaenoic acid and low phenylalanine powder for oral liquid, 30 x 35 g sachets

11071Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
(NP)	4	5	..	*2083.98	41.30	PKU Sphere20 [VF]

glycomacropeptide formula with docosahexaenoic acid and low phenylalanine powder for oral liquid, 30 x 27 g sachets

11245W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
(NP)	4	5	..	*1578.94	41.30	PKU Sphere15 [VF]

■ GLYCOMACROPEPTIDE FORMULA WITH DOCOSAHEXAENOIC ACID AND LOW PHENYLALANINE

Note This product contains higher vitamin A levels than other PBS-listed glycomacropeptide products.

Restricted benefit

Phenylketonuria

glycomacropeptide formula with docosahexaenoic acid and low phenylalanine oral liquid, 18 x 250 mL cartons

11844J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
(NP)	10	5	..	*1578.84	41.30	PKU GMPro LQ [SB]

■ HIGH FAT FORMULA WITH VITAMINS, MINERALS AND TRACE ELEMENTS AND LOW IN PROTEIN AND CARBOHYDRATE

Note Authorities for increased maximum quantities, up to a maximum of 11, may be authorised.

Restricted benefit

Ketogenic diet

Clinical criteria:

- Patient must have intractable seizures requiring treatment with a ketogenic diet; OR
- Patient must have a glucose transport protein defect; OR
- Patient must have pyruvate dehydrogenase deficiency.

Keyo should only be used under strict supervision of a dietitian, together with a metabolic physician and/or neurologist.

high fat formula with vitamins, minerals and trace elements and low in protein and carbohydrate oral semi-solid, 48 x 100 g tubs

11108P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
(NP)	3	5	..	*793.38	41.30	Keyo [VF]

■ HIGH FAT FORMULA WITH VITAMINS, MINERALS AND TRACE ELEMENTS AND LOW IN PROTEIN AND CARBOHYDRATE

Note Authorities for increased maximum quantities, up to a maximum of 11, may be authorised.

Restricted benefit

Ketogenic diet

Clinical criteria:

- Patient must have intractable seizures requiring treatment with a ketogenic diet; OR
- Patient must have a glucose transport protein defect; OR
- Patient must have pyruvate dehydrogenase deficiency.

KetoCal 4:1 should only be used under strict supervision of a dietitian, together with a metabolic physician and/or neurologist.

high fat formula with vitamins, minerals and trace elements and low in protein and carbohydrate (4:1 ratio long chain fat to carbohydrate plus protein) oral liquid, 32 x 200 mL cartons

10185C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	5	5	..	*938.59	41.30	KetoCal 4:1 LQ [SB]

■ HIGH FAT FORMULA WITH VITAMINS, MINERALS AND TRACE ELEMENTS AND LOW IN PROTEIN AND CARBOHYDRATE

Note Authorities for increased maximum quantities, up to a maximum of 48, may be authorised.

Restricted benefit

Ketogenic diet

Clinical criteria:

- Patient must have intractable seizures requiring treatment with a ketogenic diet; OR
- Patient must have a glucose transport protein defect; OR
- Patient must have pyruvate dehydrogenase deficiency.

KetoCal 3:1 should only be used under strict supervision of a dietitian, together with a metabolic physician and/or neurologist.

high fat formula with vitamins, minerals and trace elements and low in protein and carbohydrate (3:1 ratio long chain fat to carbohydrate plus protein) powder for oral liquid, 300 g

2652N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	24	5	..	*983.82	41.30	KetoCal 3:1 [SB]

■ HIGH FAT FORMULA WITH VITAMINS, MINERALS AND TRACE ELEMENTS AND LOW IN PROTEIN AND CARBOHYDRATE

Note Authorities for increased maximum quantities, up to a maximum of 48, may be authorised.

Restricted benefit

Ketogenic diet

Clinical criteria:

- Patient must have intractable seizures requiring treatment with a ketogenic diet; OR
- Patient must have a glucose transport protein defect; OR
- Patient must have pyruvate dehydrogenase deficiency.

KetoCal 4:1 should only be used under strict supervision of a dietitian, together with a metabolic physician and/or neurologist.

high fat formula with vitamins, minerals and trace elements and low in protein and carbohydrate (4:1 ratio long chain fat to carbohydrate plus protein) powder for oral liquid, 300 g

9446E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	24	5	..	*983.82	41.30	KetoCal 4:1 [SB]

■ ISOLEUCINE WITH CARBOHYDRATE

Restricted benefit

Maple syrup urine disease

isoleucine with carbohydrate containing 50 mg isoleucine oral liquid: powder for, 30 x 4 g sachets

9134R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	5	..	*446.94	41.30	Isoleucine 50 [VF]

isoleucine with carbohydrate containing 1 g isoleucine oral liquid: powder for, 30 x 4 g sachets

9436P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	5	..	*490.90	41.30	Isoleucine 1000 [VF]

■ MILK PROTEIN AND FAT FORMULA WITH VITAMINS AND MINERALS CARBOHYDRATE FREE

Restricted benefit

Ketogenic diet

Clinical criteria:

- Patient must have intractable seizures requiring treatment with a ketogenic diet; OR
- Patient must have a glucose transport protein defect; OR
- Patient must have pyruvate dehydrogenase deficiency; OR
- Patient must be an infant or young child with glucose-galactose intolerance and multiple monosaccharide intolerance.

milk protein and fat formula with vitamins and minerals carbohydrate free powder for oral liquid, 225 g

8630F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	24	5	..	*533.34	41.30	Carbohydrate Free Mixture [SB]

■ PHENYLALANINE WITH CARBOHYDRATE

Restricted benefit

Tyrosinaemia

phenylalanine with carbohydrate containing 50 mg phenylalanine oral liquid: powder for, 30 x 4 g sachets

9384X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	5	..	*446.94	41.30	Phenylalanine 50 [VF]

■ PROTEIN FORMULA WITH AMINO ACIDS, CARBOHYDRATES, VITAMINS AND MINERALS WITHOUT PHENYLALANINE, AND SUPPLEMENTED WITH DOCOSAHEXAENOIC ACID**Restricted benefit**

Phenylketonuria

protein formula with amino acids, carbohydrates, vitamins and minerals without phenylalanine, and supplemented with docosahexaenoic acid oral liquid, 30 x 130 mL pouches

10658Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	5	5	..	*1825.24	41.30	PKU Easy [OH]

■ PROTEIN FORMULA WITH VITAMINS AND MINERALS, AND LOW IN POTASSIUM, PHOSPHORUS, CALCIUM, CHLORIDE AND VITAMIN A**Authority required (STREAMLINED)****11070**

Chronic renal failure

Population criteria:

- Patient must be a child aged 3 years or older.

Clinical criteria:

- Patient must require treatment with a low protein and a low phosphorus diet; OR
- Patient must require treatment with a low protein, low phosphorus and low potassium diet.

protein formula with vitamins and minerals, and low in potassium, phosphorus, calcium, chloride and vitamin A oral liquid, 24 x 125 mL bottles

12191P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	8	5	..	*1328.06	41.30	Renastep [VF]

■ SOY PROTEIN AND FAT FORMULA WITH VITAMINS AND MINERALS CARBOHYDRATE FREE**Restricted benefit**

Ketogenic diet

Clinical criteria:

- Patient must have intractable seizures requiring treatment with a ketogenic diet; OR
- Patient must have a glucose transport protein defect; OR
- Patient must have pyruvate dehydrogenase deficiency; OR
- Patient must be an infant or young child with glucose-galactose intolerance and multiple monosaccharide intolerance.

soy protein and fat formula with vitamins and minerals carbohydrate free oral liquid, 384 mL can

8577K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	120	5	..	*551.34	41.30	RCF [AB]

■ TRIGLYCERIDES LONG CHAIN WITH GLUCOSE POLYMER**Restricted benefit**

Proven inborn errors of protein metabolism

Clinical criteria:

- Patient must be unable to meet their energy requirements with permitted food and formulae.

triglycerides long chain with glucose polymer oral liquid, 18 x 250 mL cartons

9308X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	6	5	..	*290.16	41.30	ProZero [VF]

triglycerides long chain with glucose polymer oral liquid, 6 x 1 L cartons

9309Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	5	..	*258.02	41.30	ProZero [VF]

triglycerides long chain with glucose polymer oral liquid, 27 x 200 mL cartons

10189G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	5	..	*158.34	41.30	Sno-Pro [SB]

■ TRIGLYCERIDES MEDIUM CHAIN AND LONG CHAIN WITH GLUCOSE POLYMER**Restricted benefit**

Proven inborn errors of protein metabolism

Clinical criteria:

- Patient must be unable to meet their energy requirements with permitted food and formulae.

triglycerides medium chain and long chain with glucose polymer powder for oral liquid, 400 g

3136C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	8	5	..	*238.30	41.30	Duocal [SB]

■ TRIGLYCERIDES MEDIUM CHAIN FORMULA

Note No increase in the maximum quantity or number of units may be authorised.

Note No increase in the maximum number of repeats may be authorised.

Note Not indicated for the treatment of intractable childhood epilepsy or cerebrospinal fluid glucose transporter defect requiring a ketogenic diet.

Authority required (STREAMLINED)**6165**

Chylous ascites

Authority required (STREAMLINED)**6192**

Chylothorax

Authority required (STREAMLINED)**6173**

Fat malabsorption

Clinical criteria:

- The condition must be due to liver disease; OR
- The condition must be due to short gut syndrome; OR
- The condition must be due to cystic fibrosis; OR
- The condition must be due to gastrointestinal disorders.

Authority required (STREAMLINED)**6156**

Hyperlipoproteinaemia type 1

Authority required (STREAMLINED)**6136**

Long chain fatty acid oxidation disorders

triglycerides medium chain formula powder for oral liquid, 30 x 16 g sachets

9383W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	5	..	*202.54	41.30	MCT Pro-Cal [VF]

■ TYROSINE WITH CARBOHYDRATE

Note This formulation is suitable for patients aged 3 and older.

Restricted benefit

Phenylketonuria

tyrosine with carbohydrate containing 1 g tyrosine oral liquid: powder for, 30 x 4 g sachets

9165J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	5	..	*446.94	41.30	Tyrosine 1000 [VF]

■ VALINE WITH CARBOHYDRATE**Restricted benefit**

Maple syrup urine disease

valine with carbohydrate containing 1 g valine oral liquid: powder for, 30 x 4 g sachets

9434M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	5	..	*490.90	41.30	Valine 1000 [VF]

valine with carbohydrate containing 50 mg valine oral liquid: powder for, 30 x 4 g sachets

9135T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	5	..	*446.94	41.30	Valine 50 [VF]

■ VITAMINS, MINERALS AND TRACE ELEMENTS

Note Phlexy-Vits must only be used under strict supervision of a dietician and a paediatrician.

Restricted benefit

Dietary management of conditions requiring a highly restrictive therapeutic diet

Clinical criteria:

- Patient must have insufficient vitamin and mineral intake due to a specific diagnosis requiring a highly restrictive therapeutic diet, **AND**
- Patient must be unable to adequately meet vitamin, mineral and trace element needs with other proprietary vitamin and mineral preparations.

Population criteria:

- Patient must be aged 3 years or older.

vitamins, minerals and trace elements powder for oral liquid, 30 x 7 g sachets

11200L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	280.83	41.30	Phlexy-Vits [SB]

■ VITAMINS, MINERALS AND TRACE ELEMENTS WITH CARBOHYDRATE

Note FruitiVits must only be used under strict supervision of a dietitian and a paediatrician.

Restricted benefit

Dietary management of conditions requiring a highly restrictive therapeutic diet

Clinical criteria:

- Patient must have insufficient vitamin and mineral intake due to a specific diagnosis requiring a highly restrictive therapeutic diet, **AND**
- Patient must be unable to adequately meet vitamin, mineral and trace element needs with other proprietary vitamin and mineral preparations.

Population criteria:

- Patient must be aged 3 years or older.

vitamins, minerals and trace elements with carbohydrate powder for oral liquid, 30 x 6 g sachets

10149E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	5	..	253.44	41.30	FruitiVits [VF]

■ VITAMINS, MINERALS AND TRACE ELEMENTS WITH CARBOHYDRATE

Note Paediatric Seravit must only be used under strict supervision of a dietitian and a paediatrician.

Restricted benefit

Dietary management of conditions requiring a highly restrictive therapeutic diet

Clinical criteria:

- Patient must have insufficient vitamin and mineral intake due to a specific diagnosis requiring a highly restrictive therapeutic diet, **AND**
- Patient must be unable to adequately meet vitamin, mineral and trace element needs with other proprietary vitamin and mineral preparations.

Population criteria:

- Patient must be an infant or a child.

vitamins, minerals and trace elements with carbohydrate powder for oral liquid, 200 g

9328Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	6	5	..	*335.58	41.30	Paediatric Seravit [SB]

■ WHEY PROTEIN FORMULA SUPPLEMENTED WITH AMINO ACIDS, LONG CHAIN POLYUNSATURATED FATTY ACIDS, VITAMINS AND MINERALS, LOW IN PROTEIN, PHOSPHATE, POTASSIUM AND LACTOSE**Authority required (STREAMLINED)**

6190

Chronic renal failure

Population criteria:

- Patient must be an infant or a young child.

Clinical criteria:

- Patient must require treatment with a low protein and a low phosphorus diet; OR
- Patient must require treatment with a low protein, low phosphorus and low potassium diet.

whey protein formula supplemented with amino acids, long chain polyunsaturated fatty acids, vitamins and minerals, low in protein, phosphate, potassium and lactose powder for oral liquid, 6 x 400 g cans

2870C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	5	..	*1363.10	41.30	Renastart [VF]

■ WHEY PROTEIN FORMULA SUPPLEMENTED WITH AMINO ACIDS, VITAMINS AND MINERALS, AND LOW IN PROTEIN, PHOSPHATE, POTASSIUM AND LACTOSE**Authority required (STREAMLINED)**

6190

Chronic renal failure

Population criteria:

- Patient must be an infant or a young child.

Clinical criteria:

- Patient must require treatment with a low protein and a low phosphorus diet; OR
- Patient must require treatment with a low protein, low phosphorus and low potassium diet.

whey protein formula supplemented with amino acids, vitamins and minerals, and low in protein, phosphate, potassium and lactose powder for oral liquid, 400 g

8587Y

NP

Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
16	5	..	*873.02	41.30	Kindergen [SB]

Palliative Care

ALIMENTARY TRACT AND METABOLISM.....	994
STOMATOLOGICAL PREPARATIONS	994
STOMATOLOGICAL PREPARATIONS.....	994
DRUGS FOR FUNCTIONAL GASTROINTESTINAL DISORDERS	994
BELLADONNA AND DERIVATIVES, PLAIN	994
PROPULSIVES.....	994
DRUGS FOR CONSTIPATION	994
DRUGS FOR CONSTIPATION.....	994
MUSCULO-SKELETAL SYSTEM.....	996
ANTIINFLAMMATORY AND ANTIRHEUMATIC PRODUCTS.....	996
ANTIINFLAMMATORY AND ANTIRHEUMATIC PRODUCTS, NON- STEROIDS.....	996
NERVOUS SYSTEM.....	997
ANALGESICS.....	997
OPIOIDS	997
OTHER ANALGESICS AND ANTIPYRETICS	1003
ANTIEPILEPTICS.....	1003
ANTIEPILEPTICS	1003
PSYCHOLEPTICS.....	1004
ANXIOLYTICS	1004
HYPNOTICS AND SEDATIVES	1004

ALIMENTARY TRACT AND METABOLISM

STOMATOLOGICAL PREPARATIONS

STOMATOLOGICAL PREPARATIONS

Other agents for local oral treatment

BENZYDAMINE

Authority required (STREAMLINED)

6197

Painful mouth

Clinical criteria:

- Patient must be receiving palliative care.

benzydamine hydrochloride 0.15% mouthwash, 500 mL

5385K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	3	..	23.21	24.50	Diffiam [IL]

DRUGS FOR FUNCTIONAL GASTROINTESTINAL DISORDERS

BELLADONNA AND DERIVATIVES, PLAIN

Belladonna alkaloids, semisynthetic, quaternary ammonium compounds

HYOSCINE BUTYLBROMIDE

Authority required (STREAMLINED)

6207

For use in patients receiving palliative care

hyoscine butylbromide 20 mg/mL injection, 5 x 1 mL ampoules

5317W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	6	3	..	*76.68	41.30	^a Buscopan [VZ]	^a HYOSCINE BUTYLBROMIDE MEDSURGE [DZ]
						^a HYOSCINE BUTYLBROMIDE SXP [XC]	

PROPULSIVES

Propulsives

METOCLOPRAMIDE

Authority required (STREAMLINED)

6084

Nausea or gastric stasis

Clinical criteria:

- Patient must be receiving palliative care.

metoclopramide hydrochloride 10 mg/2 mL injection, 10 x 2 mL ampoules

10762K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	4	..	..	*34.86	36.15	Maxolon [IL]

DRUGS FOR CONSTIPATION

DRUGS FOR CONSTIPATION

Contact laxatives

BISACODYL

Restricted benefit

Constipation

Clinical criteria:

- Patient must be receiving palliative care.

bisacodyl 10 mg suppository, 12

5304E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	3	3	..	*22.38	23.67	Petrus Bisacodyl Suppositories [PP]

bisacodyl 10 mg suppository, 10

5303D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	3	3	..	*24.66	25.95	^a Petrus Bisacodyl Suppositories [PP]
			^B 1.29	*25.95	25.95	^a Dulcolax [VZ]

bisacodyl 5 mg enteric tablet, 200

5301B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	3	..	18.01	19.30	Lax-Tab [AE]

*Bulk-forming laxatives***■ RHAMNUS FRANGULA + STERCULIA**Restricted benefit

Constipation

Clinical criteria:

- Patient must be receiving palliative care.

rhamnus frangula 80 mg/g + sterculia 620 mg/g granules, 500 g

5322D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	3	..	26.12	27.41	Normacol Plus [NE]

*Osmotically acting laxatives***■ MACROGOL-3350**Authority required (STREAMLINED)**6170**

Constipation

Clinical criteria:

- Patient must be receiving palliative care.

macrogol-3350 1 g/g powder for oral liquid, 510 g

5426N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	3	..	*27.16	28.45	OsmoLax [KY]

■ MACROGOL-3350 + SODIUM CHLORIDE + BICARBONATE + POTASSIUM CHLORIDEAuthority required (STREAMLINED)**6171**

Constipation

Clinical criteria:

- Patient must be receiving palliative care.

macrogol-3350 13.125 g/25 mL + sodium chloride 350.7 mg/25 mL + sodium bicarbonate 178.5 mg/25 mL + potassium chloride 46.6 mg/25 mL oral liquid, 500 mL

10127B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	3	..	*22.10	23.39	Movicol Liquid [NE]

macrogol-3350 13.125 g + sodium chloride 350.7 mg + sodium bicarbonate 178.5 mg + potassium chloride 46.6 mg powder for oral liquid, 30 sachets

5389P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	2	3	..	*27.16	28.45	^a APOHEALTH Macrogol with Electrolytes [GX] ^a Chemists' Own Macrogol with Electrolytes [RW] ^a lax-sachets [AE] ^a Molaxole [GO]	^a APO-MACROGOL plus ELECTROLYTES [TX] ^a LaxaCon [EA] ^a Macrovic [RF] ^a Movicol [NE]

*Enemas***■ BISACODYL**Restricted benefit

Constipation

Clinical criteria:

- Patient must be receiving palliative care.

bisacodyl 10 mg/5 mL enema, 25 x 5 mL

5302C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	3	..	39.43	40.72	Bisalax [AS]

■ CITRIC ACID + LAURYL SULFOACETATE SODIUM + SORBITOLRestricted benefit

Constipation

Clinical criteria:

- Patient must be receiving palliative care.

MUSCULO-SKELETAL SYSTEM

sodium citrate dihydrate 450 mg/5 mL + lauryl sulfoacetate sodium 45 mg/5 mL + sorbitol 3.125 g/5 mL enema, 12 x 5 mL

5331N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	3	..	*30.28	31.57	Micolette [AE]

Peripheral opioid receptor antagonists

■ METHYLNALTREXONE

Authority required (STREAMLINED)

6180

Opioid-induced constipation

Clinical criteria:

- The treatment must be in combination with oral laxatives, **AND**
- Patient must be receiving palliative care, **AND**
- Patient must have failed to respond to laxatives.

methylnaltrexone bromide 12 mg/0.6 mL injection, 7 x 0.6 mL vials

5424L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	243.55	41.30	Relistor [LM]

methylnaltrexone bromide 12 mg/0.6 mL injection, 0.6 mL vial

5423K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	7	..	..	*243.57	41.30	Relistor [LM]

■ MUSCULO-SKELETAL SYSTEM

■ ANTIINFLAMMATORY AND ANTIRHEUMATIC PRODUCTS

ANTIINFLAMMATORY AND ANTIRHEUMATIC PRODUCTS, NON-STERIODS

Acetic acid derivatives and related substances

■ DICLOFENAC

Restricted benefit

Severe pain

Clinical criteria:

- Patient must be receiving palliative care.

diclofenac sodium 100 mg suppository, 20

5363G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	3	..	*28.44	29.73	Voltaren 100 [NV]

diclofenac sodium 50 mg enteric tablet, 50

5362F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	3	..	14.34	15.63	^a APO-Diclofenac [TX]	^a Clonac 50 [RW]
						^a Diclofenac Amneal [ED]	^a Diclofenac AN [EA]
						^a Diclofenac Sandoz [SZ]	^a Fenac [AF]
			^B 3.21	17.55	15.63	^a Voltaren 50 [NV]	

diclofenac sodium 25 mg enteric tablet, 50

5361E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	2	3	..	*15.18	16.47	^a APO-Diclofenac [TX]	^a Clonac 25 [RW]
						^a Diclofenac Amneal [ED]	^a Diclofenac AN [EA]
						^a Diclofenac Sandoz [SZ]	^a Fenac 25 [AF]
			^B 3.22	*18.40	16.47	^a Voltaren 25 [NV]	

■ INDOMETACIN

Restricted benefit

Severe pain

Clinical criteria:

- Patient must be receiving palliative care.

indometacin 25 mg capsule, 50

5377B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	3	..	*17.64	18.93	^a Arthrexin [AF]
			^B 4.02	*21.66	18.93	^a Indocid [AS]

indometacin 100 mg suppository, 20

5378C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	3	..	*26.00	27.29	Indocid [AS]

Propionic acid derivatives

■ IBUPROFEN

Restricted benefit

Severe pain

Clinical criteria:

- Patient must be receiving palliative care.

ibuprofen 400 mg tablet, 30

5368M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	3	3	..	*16.95	18.24	^a APO-Ibuprofen 400 [TX]
			^B 7.47	*24.42	18.24	^a Brufen [GO]

■ NAPROXEN

Restricted benefit

Severe pain

Treatment criteria:

- Patient must be undergoing palliative care.

Clinical criteria:

- Patient must be unable to take a solid dose form of a non-steroidal anti-inflammatory agent.

naproxen 125 mg/5 mL oral liquid, 474 mL

5397C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	±1	3	..	122.71	41.30	Phebra Naproxen Suspension [FF]

■ NAPROXEN

Restricted benefit

Severe pain

Clinical criteria:

- Patient must be receiving palliative care.

naproxen 500 mg tablet, 50

5346J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	3	..	17.42	18.71	Naprosyn [IX]

naproxen 750 mg modified release tablet, 28

5347K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	3	..	17.01	18.30	^a Proxen SR 750 [IY]
			^B 0.99	18.00	18.30	^a Naprosyn SR750 [IX]

naproxen 1 g modified release tablet, 28

5348L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	3	..	18.58	19.87	^a Proxen SR 1000 [IY]
			^B 1.12	19.70	19.87	^a Naprosyn SR1000 [IX]

naproxen 250 mg tablet, 50

5345H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	3	..	*21.52	22.81	Naprosyn [IX]

■ NAPROXEN

Note Naproxen sodium 550 mg is approximately equivalent to 500 mg of naproxen acid.

Restricted benefit

Severe pain

Clinical criteria:

- Patient must be receiving palliative care.

naproxen sodium 550 mg tablet, 50

5353R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	3	..	17.56	18.85	^a Crysanal [IY]
			^B 1.87	19.43	18.85	^a Anaprox 550 [IX]

■ NERVOUS SYSTEM

■ ANALGESICS

OPIOIDS

Natural opium alkaloids

■ MORPHINE

Caution The risk of drug dependence is high.

Note Telephone approvals are limited to 1 month's therapy.

Authority required

Chronic severe disabling pain

Clinical criteria:

- Patient must be receiving palliative care, **AND**
- The condition must be unresponsive to non-opioid analgesics.

morphine sulfate pentahydrate 200 mg modified release tablet, 28

5391R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	119.42	41.30	MS Contin [MF]

■ MORPHINE
Caution The risk of drug dependence is high.

Note Telephone approvals are limited to 1 month's therapy.

Authority required

Severe disabling pain

Clinical criteria:

- Patient must be receiving palliative care, **AND**
- The condition must be unresponsive to non-opioid analgesics.

morphine sulfate pentahydrate 10 mg tablet, 20

5393W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	21.42	22.71	Sevredol [MF]

morphine sulfate pentahydrate 20 mg tablet, 20

5394X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	2	..	22.19	23.48	Sevredol [MF]

■ MORPHINE
Caution The risk of drug dependence is high.

Morphine sulfate pentahydrate 10 and 20 mg modified release capsules must not be co-prescribed with immediate release oral morphine, when it has been prescribed for the reduction of chronic breathlessness.

Note Treatment should be initiated by a specialist knowledgeable in the use of potent opioids for the management of chronic breathlessness.

Note Applications for an increased maximum quantity to provide for 1 month's supply of this drug will be authorised.

Note Where consultation with a palliative care specialist or service has occurred, applications for increased repeats for up to 3 months' supply may be authorised.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Restricted benefit

Chronic Breathlessness

Clinical criteria:

- Patient must be receiving palliative care.

morphine sulfate pentahydrate 10 mg modified release capsule, 28

11760Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	26.30	27.59	Kapanol [YN]

morphine sulfate pentahydrate 20 mg modified release capsule, 28

11761B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	30.63	31.92	Kapanol [YN]

Phenylpiperidine derivatives
■ FENTANYL
Caution The risk of drug dependence is high.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Breakthrough pain

Treatment Phase: Initial treatment for dose titration

Clinical criteria:

- Patient must have cancer, **AND**
- Patient must have pain directly attributable to cancer, **AND**

- Patient must be assessed as receiving adequate management of their persistent pain with opioids, **AND**
- Patient must have previously experienced inadequate pain relief following adequate doses of short acting opioids for the treatment of breakthrough pain; OR
- The treatment must be used as short acting opioids are considered clinically inappropriate; OR
- Patient must have previously experienced adverse effects following the use of short acting opioids for breakthrough pain.

Treatment criteria:

- Patient must be undergoing palliative care.

fentanyl 200 microgram sublingual tablet, 10

10600X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	..	..	*165.36	41.30	Abstral [FK]

fentanyl 800 microgram lozenge, 9

5404K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	99.07	41.30	Actiq [TB]

fentanyl 800 microgram sublingual tablet, 10

10612M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	89.93	41.30	Abstral [FK]

fentanyl 1200 microgram lozenge, 9

5405L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	99.07	41.30	Actiq [TB]

fentanyl 400 microgram sublingual tablet, 10

10603C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	89.93	41.30	Abstral [FK]

fentanyl 600 microgram lozenge, 9

5403J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	99.07	41.30	Actiq [TB]

fentanyl 200 microgram lozenge, 9

5401G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	99.07	41.30	Actiq [TB]

fentanyl 1600 microgram lozenge, 9

5406M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	99.07	41.30	Actiq [TB]

fentanyl 400 microgram lozenge, 9

5402H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	99.07	41.30	Actiq [TB]

fentanyl 600 microgram sublingual tablet, 10

10604D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	89.93	41.30	Abstral [FK]

fentanyl 300 microgram sublingual tablet, 10

10606F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	..	..	89.93	41.30	Abstral [FK]

fentanyl 100 microgram sublingual tablet, 10

10601Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	..	..	*165.36	41.30	Abstral [FK]

■ FENTANYL

Caution The risk of drug dependence is high.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Breakthrough pain

Treatment Phase: Initial treatment for dose titration

Clinical criteria:

- Patient must have cancer, **AND**

- Patient must have pain directly attributable to cancer, **AND**
- Patient must be assessed as receiving adequate management of their persistent pain with opioids, **AND**
- Patient must have previously experienced inadequate pain relief following adequate doses of short acting opioids for the treatment of breakthrough pain; OR
- The treatment must be used as short acting opioids are considered clinically inappropriate; OR
- Patient must have previously experienced adverse effects following the use of short acting opioids for breakthrough pain.

Treatment criteria:

- Patient must be undergoing palliative care.

fentanyl 800 microgram orally disintegrating tablet, 4

10723J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	..	..	*75.32	41.30	Fentora [TB]

fentanyl 100 microgram orally disintegrating tablet, 4

10729Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	..	..	*75.32	41.30	Fentora [TB]

fentanyl 600 microgram orally disintegrating tablet, 4

10722H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	..	..	*75.32	41.30	Fentora [TB]

fentanyl 400 microgram orally disintegrating tablet, 4

10739F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	..	..	*75.32	41.30	Fentora [TB]

fentanyl 200 microgram orally disintegrating tablet, 4

10697B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	..	..	*75.32	41.30	Fentora [TB]

■ FENTANYL

Caution The risk of drug dependence is high.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note For first continuing supply, applications for increased repeats for up to 3 months' supply may be authorised.

Note Where consultation with a palliative care specialist or service has occurred, applications for increased repeats for up to 3 months' supply may be authorised.

Note Telephone approvals are limited to 1 months' therapy.

Authority required

Breakthrough pain

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have cancer, **AND**
- Patient must have pain directly attributable to cancer, **AND**
- Patient must be assessed as receiving adequate management of their persistent pain with opioids, **AND**
- Patient must have previously experienced inadequate pain relief following adequate doses of short acting opioids for the treatment of breakthrough pain; OR
- The treatment must be used as short acting opioids are considered clinically inappropriate; OR
- Patient must have previously experienced adverse effects following the use of short acting opioids for breakthrough pain.

Treatment criteria:

- Patient must be undergoing palliative care.

fentanyl 1200 microgram lozenge, 30

5411T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	..	..	*587.60	41.30	Actiq [TB]

fentanyl 400 microgram lozenge, 30

5408P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	..	..	*587.60	41.30	Actiq [TB]

fentanyl 300 microgram sublingual tablet, 30

10610K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	..	..	*472.44	41.30	Abstral [FK]

fentanyl 100 microgram sublingual tablet, 30

10602B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	..	..	*472.44	41.30	Abstral [FK]

fentanyl 600 microgram lozenge, 30

5409Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	..	..	*587.60	41.30	Actiq [TB]

fentanyl 800 microgram sublingual tablet, 30

10611L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	..	..	*472.44	41.30	Abstral [FK]

fentanyl 600 microgram sublingual tablet, 30

10613N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	..	..	*472.44	41.30	Abstral [FK]

fentanyl 200 microgram lozenge, 30

5407N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	..	..	*587.60	41.30	Actiq [TB]

fentanyl 800 microgram lozenge, 30

5410R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	..	..	*587.60	41.30	Actiq [TB]

fentanyl 200 microgram sublingual tablet, 30

10607G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	..	..	*472.44	41.30	Abstral [FK]

fentanyl 400 microgram sublingual tablet, 30

10608H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	..	..	*472.44	41.30	Abstral [FK]

fentanyl 1600 microgram lozenge, 30

5412W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	..	..	*587.60	41.30	Actiq [TB]

■ FENTANYL

Caution The risk of drug dependence is high.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Note For first continuing supply, applications for increased repeats for up to 3 months' supply may be authorised.

Note Where consultation with a palliative care specialist or service has occurred, applications for increased repeats for up to 3 months' supply may be authorised.

Note Telephone approvals are limited to 1 months' therapy.

Authority required

Breakthrough pain

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must have cancer, **AND**
- Patient must have pain directly attributable to cancer, **AND**
- Patient must be assessed as receiving adequate management of their persistent pain with opioids, **AND**
- Patient must have previously experienced inadequate pain relief following adequate doses of short acting opioids for the treatment of breakthrough pain; OR
- The treatment must be used as short acting opioids are considered clinically inappropriate; OR
- Patient must have previously experienced adverse effects following the use of short acting opioids for breakthrough pain.

Treatment criteria:

- Patient must be undergoing palliative care.

fentanyl 200 microgram orally disintegrating tablet, 28

10698C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	..	..	*441.74	41.30	Fentora [TB]

fentanyl 800 microgram orally disintegrating tablet, 28

10738E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	..	..	*441.74	41.30	Fentora [TB]

fentanyl 400 microgram orally disintegrating tablet, 28

10737D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	..	..	*441.74	41.30	Fentora [TB]

fentanyl 600 microgram orally disintegrating tablet, 28

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
10713W	2	..	..	*441.74	41.30	Fentora [TB]

fentanyl 100 microgram orally disintegrating tablet, 28

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
10684H	2	..	..	*441.74	41.30	Fentora [TB]

Diphenylpropylamine derivatives
■ METHADONE

Caution The risk of drug dependence is high.

Note Where consultation with a palliative care specialist or service has occurred, applications for increased repeats for up to 3 months' supply may be authorised.

Note Telephone approvals are limited to 1 month's therapy.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required

Chronic severe disabling pain

Treatment Phase: Initial treatment, for up to 3 months

Clinical criteria:

- Patient must be receiving palliative care, **AND**
- The condition must be unresponsive to non-opioid analgesics.

methadone hydrochloride 5 mg/mL oral liquid, 200 mL

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
5399E	1	2	..	25.32	26.61	Aspen Methadone Syrup [AS]

■ METHADONE

Caution The risk of drug dependence is high.

Note Where consultation with a palliative care specialist or service has occurred, applications for increased repeats for up to 3 months' supply may be authorised.

Note Telephone approvals are limited to 1 month's therapy.

Note Shared Care Model:

For prescribing by nurse practitioners where care of a patient is shared between a nurse practitioner and medical practitioner in a formalised arrangement with an agreed management plan. Further information can be found in the Explanatory Notes for Nurse Practitioners.

Authority required

Chronic severe disabling pain

Treatment Phase: Continuing treatment

Clinical criteria:

- Patient must be receiving palliative care, **AND**
- The condition must be unresponsive to non-opioid analgesics.

methadone hydrochloride 5 mg/mL oral liquid, 200 mL

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
5400F	1	..	..	25.32	26.61	Aspen Methadone Syrup [AS]

Oripavine derivatives
■ BUPRENORPHINE

Caution The risk of drug dependence is high.

Note Telephone approvals are limited to 1 month's therapy.

Authority required

Chronic severe disabling pain

Clinical criteria:

- Patient must be receiving palliative care, **AND**
- The condition must be unresponsive to non-opioid analgesics.

buprenorphine 25 microgram/hour patch, 2

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
10964C	2	2	..	*95.86	41.30	^a Bupredermal [TX] ^a Norspan [MF]	^a Burprenorphine Sandoz [SZ]

buprenorphine 30 microgram/hour patch, 2

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
10949G	2	2	..	*107.16	41.30	^a Bupredermal [TX]	^a Buprenorphine Sandoz [SZ]

^a Norspan [MF]

NP	buprenorphine 5 microgram/hour patch, 2						
	10957Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP		2	2	..	*41.20	41.30	^a Bupredermal [TX] ^a Norspan [MF]
buprenorphine 10 microgram/hour patch, 2							
NP	10948F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
		2	2	..	*60.74	41.30	^a Bupredermal [TX] ^a Norspan [MF]
buprenorphine 20 microgram/hour patch, 2							
NP	10970J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
		2	2	..	*84.58	41.30	^a Bupredermal [TX] ^a Norspan [MF]
buprenorphine 40 microgram/hour patch, 2							
NP	10959T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
		2	2	..	*130.38	41.30	^a Bupredermal [TX] ^a Norspan [MF]
buprenorphine 15 microgram/hour patch, 2							
NP	10953L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
		2	2	..	*72.64	41.30	^a Bupredermal [TX] ^a Norspan [MF]

OTHER ANALGESICS AND ANTIPYRETICS

Anilides

■ PARACETAMOL

Restricted benefit

Analgesia or fever

Clinical criteria:

- Patient must be receiving palliative care, **AND**
- Patient must be intolerant to alternative therapy.

paracetamol 500 mg suppository, 10

NP	12210P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
		10	3	..	*97.04	41.30	Panadol [GC]

■ PARACETAMOL

Note Pharmaceutical benefits that have the form paracetamol 665 mg tablet: modified release, 96 and pharmaceutical benefits that have the form paracetamol 665 mg tablet: modified release, 192 are equivalent for the purposes of substitution.

Restricted benefit

Analgesia or fever

Clinical criteria:

- Patient must be receiving palliative care, **AND**
- Patient must be intolerant to alternative therapy.

paracetamol 665 mg modified release tablet, 192

NP	10796F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
		1	3	..	19.48	20.77	^a Osteomol 665 Paracetamol [CR]

paracetamol 665 mg modified release tablet, 96

NP	5343F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
		2	3	..	*19.48	20.77	^a APOHEALTH Osteo Relief Paracetamol 665 mg [TX] ^a Osteomol 665 Paracetamol [CR]

■ ANTIEPILEPTICS

ANTIEPILEPTICS

Benzodiazepine derivatives

■ CLONAZEPAM

Note No increase in the maximum number of repeats may be authorised.

Authority required

Myoclonus

Clinical criteria:

- The treatment must be for prophylaxis or prevention of the indication, **AND**

- Patient must be receiving palliative care.

clonazepam 2 mg tablet, 100

5338Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	3	..	22.73	24.02	Paxam 2 [AF]

clonazepam 2.5 mg/mL (0.1 mg/drop) oral liquid, 10 mL

5339B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	3	..	*19.52	20.81	Rivotril [RO]

CLONAZEPAM

Note Pharmaceutical benefits that have form pack size clonazepam 500 microgram tablet, 100 and clonazepam 500 microgram tablet, 50 are equivalent for the purposes of substitution.

Note No increase in the maximum number of repeats may be authorised.

Authority required

Myoclonus

Clinical criteria:

- The treatment must be for prophylaxis or prevention of the indication, **AND**
- Patient must be receiving palliative care.

clonazepam 500 microgram tablet, 100

5337X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	1	3	..	17.72	19.01	^a Paxam 0.5 [AF]

clonazepam 500 microgram tablet, 50

11520H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	3	^B 1.84	*19.56	19.01	^a Rivotril [RO]

PSYCHOLEPTICS**ANXIOLYTICS***Benzodiazepine derivatives***DIAZEPAM**

Note No increase in the maximum number of repeats may be authorised.

Authority required

Anxiety

Clinical criteria:

- Patient must be receiving palliative care.

diazepam 5 mg tablet, 50

5356X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	3	..	13.72	15.01	^a Antenex 5 [AF]	^a APO-Diazepam [TX]
			^B 3.07	16.79	15.01	^a NOUMED DIAZEPAM [VO]	^a Valpam 5 [RW]
						^a Valium [RO]	

diazepam 2 mg tablet, 50

5355W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	1	3	..	13.66	14.95	^a APO-Diazepam [TX]	^a Valpam 2 [RW]
			^B 2.78	16.44	14.95	^a Antenex 2 [AF]	

OXAZEPAM

Note No increase in the maximum number of repeats may be authorised.

Authority required

Anxiety

Clinical criteria:

- Patient must be receiving palliative care.

oxazepam 15 mg tablet, 25

5371Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	3	..	*15.08	16.37	Serepax [AS]

oxazepam 30 mg tablet, 25

5372R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	2	3	..	*14.44	15.73	^a APO-Oxazepam [TX]	^a Murelax [RW]
			^B 4.40	*18.84	15.73	^a Serepax [AS]	

HYPNOTICS AND SEDATIVES*Benzodiazepine derivatives*

■ NITRAZEPAM

Note No increase in the maximum number of repeats may be authorised.

Authority required

Insomnia

Clinical criteria:

- Patient must be receiving palliative care.

nitrazepam 5 mg tablet, 25

5359C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
NP	2	3	..	*15.44	16.73	Mogadon [IL]

■ TEMAZEPAM

Note No increase in the maximum number of repeats may be authorised.

Authority required

Insomnia

Clinical criteria:

- Patient must be receiving palliative care.

temazepam 10 mg tablet, 25

5375X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
NP	2	3	..	*14.44	15.73	^a APO-Temazepam [TX]	^a Temaze [AF]
						^a Temtabs [LN]	
			^B 6.70	*21.14	15.73	^a Normison [AS]	

Repatriation Pharmaceutical Benefits Scheme

BENEFICIARIES' ENTITLEMENT CARDS AND ELIGIBILITY FOR REPATRIATION PHARMACEUTICAL BENEFITS

Gold card

This card is issued to those veterans of Australia's defence force, their widows/widowers and dependants entitled to treatment for all medical conditions.

White card

A White Card is issued to Australian veterans or mariners under the Veterans' Entitlements Act 1986 with:

- an accepted war or service-caused injury or disease;
- malignant cancer (neoplasia) whether war-caused or not;
- pulmonary tuberculosis whether war-caused or not;
- post-traumatic stress disorder whether war-caused or not; or
- anxiety and/or depression whether war-caused or not.

Orange card

Orange Repatriation pharmaceutical benefits cards are issued to Commonwealth and allied veterans and mariners who:

- have qualifying service from World War I or II and
- are aged 70 or over and
- have been resident in Australia for 10 years or more.



For more information go to the Department of Veterans' Affairs website:
<http://www.dva.gov.au>

RPBS Explanatory Notes

Introduction

The Australian Repatriation System

- The Australian Repatriation system is based primarily on the principle of compensation to veterans and eligible dependants for injury or death related to war service. In certain cases, treatment is also provided for accepted injuries or conditions that are not service-related or have occurred as a result of other than war service.
- Through the *Veterans' Entitlements Act 1986* the Department of Veterans' Affairs provides programs of compensation, income support and treatment for eligible veterans and their dependants. One of the defined benefits for eligible veterans is the Repatriation Pharmaceutical Benefits Scheme. This range of medications and dressings is more comprehensive than is available through the Pharmaceutical Benefits Scheme.

RPBS prescribing provisions

- Unless otherwise stated, Repatriation Pharmaceutical Benefits Scheme (RPBS) prescriptions must conform with the requirements of Pharmaceutical Benefits Scheme (PBS) prescriptions, as detailed in Section 1 – Explanatory Notes in the *Schedule of Pharmaceutical Benefits* book. The prescriber shall ensure that a prescription contains the following details:
 - the category of benefit, i.e., RPBS, by placing a cross in the relevant box;
 - the patient's full name and address;
 - the prescription date;
 - the DVA file number of the patient as evidence of entitlement;
 - in the case of authority prescriptions, the Authority approval number or the four digit streamlined authority code;
 - the item, form, strength, quantity and directions;
 - the number of repeats, if applicable;
 - indicate when brand substitution is not permitted; and
 - the name, signature, the prescriber number and address of the prescriber.

Prior Approval Arrangements

- The prior approval of the Department is required to prescribe the following:
 - 'Authority required' items (excluding 'Authority required (STREAMLINED)' items) listed in either the PBS or RPBS Schedule;
 - increased quantities and/or repeats of items listed in either the PBS or RPBS Schedule;
 - items listed under section 100 of the *National Health Act 1953*; and
 - other items not listed in either Schedule (non-Schedule items).
- The above items are to be prescribed on the common PBS/RPBS authority prescription form in accordance with the directions stated in the Explanatory Notes in the *Schedule of Pharmaceutical Benefits* (See also information regarding dental prescribing and prescribing by optometrists under the RPBS in these Notes.)
- All Authority required prescriptions and requests for non-Schedule items must receive prior approval from the Department. This can be achieved by either:
 - using the Department's national free call number 1800 552 580; or
 - by mailing the written authority prescription to the Veterans' Affairs Pharmaceutical Advisory Centre (VAPAC) at the reply paid address shown at the end of these RPBS Explanatory Notes.

Prior approval is not required from DVA to prescribe an Authority required (STREAMLINED) item (except where increased quantities and/or repeats are required). Instead the authority prescription form must include a four digit streamlined authority code.

- Some requests for prior approval (including some non-Schedule items) need to be referred by VAPAC to the Repatriation Pharmaceutical Reference Committee for consideration. In such cases a VAPAC pharmacist will advise the prescriber to submit a request in writing that provides the following information:
 - A current clinical report on the patient's condition (such as age, co-morbidities, renal, liver failure) and clinical reports including pathology, biochemistry, diagnostic and other investigations if appropriate.
 - Details of past and current therapy for the condition. Include details of PBS, RPBS and non-Schedule items utilised, and the results of those therapies.
 - Details of the proposed treatment regimen. Include intended dose and duration of treatment and objective measures of response.
 - When the proposed use of the item is outside the TGA-approved indications for use in Australia, provide copies of articles from peer reviewed publications supporting the proposed treatment.
 - Signed, informed patient consent where the item is to be used for a non-TGA-approved indication.
 - For items without Australian marketing approval, a copy of the TGA Special Access Scheme approval to prescribe the drug.
- Requests for prior approval to prescribe a non-Schedule (PBS or RPBS) item that is of the same therapeutic class (ATC level 3) as an item that is listed on the Schedule, will not be approved unless unequivocal clinical evidence is presented to demonstrate that the requested item is essential for effective treatment of the nominated patient.
- A pharmacist should not supply an item prescribed on an RPBS Authority Prescription Form unless the form has been approved and stamped by VAPAC, or has been endorsed by the prescriber with a telephone Authority approval number provided by VAPAC. Medicare Australia will not accept RPBS Authority prescriptions that have not been approved by the Department of Veterans' Affairs for payment.

Palliative Care Drugs

- The following medications may be available, or made available in increased quantities or doses under prior approval arrangements for use only in the palliative care of terminal disease:
 - clonazepam
 - cyclizine

- dexamethasone
- disodium pamidronate
- fentanyl
- glycopyrrolate
- hyoscine butylbromide
- hyoscine hydrobromide
- ketamine
- midazolam
- octreotide
- For further information telephone VAPAC on 1800 552 580.

Dental Prescribing

- Under Department of Veterans' Affairs arrangements, financial responsibility for pharmaceutical benefits prescribed by a Local Dental Officer (LDO) is limited to treatment to which holders of the following cards are entitled: Where possible the LDO shall prescribe in accordance with the provisions governing dental prescribing under the Pharmaceutical Benefits Scheme (PBS).
 - a Gold Repatriation Health Card – For All Conditions; or
 - a White Repatriation Health Card – For Specific Conditions; or
 - an Orange Repatriation Pharmaceutical Benefits Card.
- Prescriptions for PBS Dental Schedule items for Gold, White and Orange Card holders are to be dispensed at the PBS concessional rate. Claims for payment by the dispensing pharmacist are to be included with other Repatriation prescriptions. The card holder is required to meet the cost of any applicable brand premium.
- When a non-PBS Dental Schedule item is prescribed for an eligible card holder, the LDO's private prescription form should be used. The dispensing pharmacist may charge the patient the full cost of the prescription. The patient may claim a refund for the full cost of a non-Schedule item from the Department if an itemised receipt (not a cash register receipt) and a copy of the prescription are provided.

Prescribing by optometrists

- Optometrists approved as 'PBS prescribers' may write RPBS prescriptions as outlined in Section 1 for medicines listed in Section 2 of the PBS Schedule as pharmaceutical benefits for optometrical use.
- Medicines in the optometrist list include non-Authority and Authority required items. Procedures for obtaining VAPAC approval to prescribe 'Authority required' optometrist items or increased quantities and/or repeats of optometrist items under the RPBS are the same as indicated under prior approval arrangements above.
- The list of medicines for prescribing by optometrists under the RPBS is the same as applies under the PBS. There are no optometrist listings in the RPBS Schedule for prescribing for veterans only. There is no provision for optometrist prescribers to request approval to prescribe items that are not included in the PBS optometrist list (non-Schedule items).
- Optometrist PBS/RPBS prescription forms are for use for prescribing non-Authority or Authority required optometrist items under the RPBS with one item per form only.

Provisions governing pricing and payment for RPBS benefits

Introduction

- Unless otherwise stated, the pricing and payment principles and arrangements for approved pharmacists supplying pharmaceutical benefits under the RPBS will be the same as those arrangements applying under the PBS.
- Where a pharmaceutical benefit that is not listed on the PBS or RPBS Schedule is dispensed on an RPBS Authority prescription, a pharmacist will price the benefit and enter the serial number, prescription identifying number and price on the sticker or stamp imprint affixed to the prescription.

Pricing of Schedule Items

- Items supplied under the RPBS from the PBS Schedule, both ready-prepared and extemporaneously-prepared, will be paid on the same basis as benefits supplied under the PBS. Items supplied under the RPBS from the Repatriation Schedule, including wound dressings, will be paid on the basis of the price as given in the Repatriation Pharmaceutical Benefits section (Section 1 – RPBS Schedule, Drugs, Medicines and Dressings) of the *Schedule of Pharmaceutical Benefits*.

Pricing of Non-Schedule Ready Prepared Items

- Non-Schedule ready-prepared items are to be priced on the basis of the invoiced, GST-exclusive wholesale price to pharmacists plus the appropriate PBS mark-up and the PBS dispensing fee. Where the item price to pharmacists is greater than \$100.00, a copy of the invoice pertaining to the supply of that item is to be submitted together with the appropriate copy of the authority prescription as part of the claim for payment.

Pricing of Non-Schedule Extemporaneously Prepared Items

- When an ingredient drug is not listed in the PBS Drug Tariff, the recovery price will be based on the invoiced wholesale price to pharmacists, increased by a mark-up of 100%, calculated in accordance with the directions contained in the pricing instructions for pricing of PBS extemporaneously-prepared benefits in this Schedule. The price paid by the pharmacist for the commercial pack from which the ingredient is used shall be endorsed on the prescription form.

Miscellaneous Pricing Rules

- The price to pharmacists used as the basis of pricing will be the invoiced, GST-exclusive price from the wholesaler.
- If multiple quantities of a manufacturer's original pack are supplied, the PBS mark-up is applied to the price to pharmacist of each pack and then totalled. The PBS dispensing fee, and the PBS dangerous drug fee if applicable, are then added to the total of the marked-up prices.

- When the quantity prescribed corresponds with the quantity of a manufacturer's original pack, in no circumstances will the price payable for one pack exceed that payable for multiples or combinations of packs to supply the quantity prescribed.
- The list of ingredient drugs and prices included in the PBS Drug Tariff are common to both the PBS and RPBS. Certain restrictions apply regarding the prescribing and dispensing of some of these ingredient drugs as pharmaceutical benefits, e.g., use as additive only.
- For items prescribed generically, including non-Schedule and wound dressings, the pharmacist should indicate on the prescription the quantity and brand supplied. If prescriptions are not endorsed, the Department will pay the lowest priced acceptable product available.

General

Packaging Material, Postage or Freight

- Payment to a pharmacist for the costs of packaging materials, postage or freight required to supply a pharmaceutical benefit is to be paid by the patient, who may then claim reimbursement from the Department through the provision of a pharmacist's itemised receipt.

Payment for Items Supplied at Short Intervals

- For all items dispensed at specific short intervals of time, the Department will pay a separate PBS dispensing fee for each occasion that the drug is supplied and which is acknowledged on receipt by the patient or agent.
- The price payable on the items supplied will be based on the individual dose quantity supplied. Where applicable, a PBS dangerous drug fee and a minimum container charge will be payable for each supply.

Receipts for Patient Charges

- Where a charge is paid by a patient in any of the circumstances of paragraphs 13 or 24, the pharmacist is required to provide a printed receipt to the patient with the details of the items or services provided, the amount paid, date of supply and the patient's name and address. The patient may apply for reimbursement from the Department.

Special Patient Contributions

- The Special Patient Contribution for items listed as Special Pharmaceutical Benefits in the PBS Schedule is not payable by veterans entitled to pharmaceutical benefits under the RPBS. Eligible veterans receiving Special Pharmaceutical Benefits under the RPBS are required to pay only the concessional patient contribution and any applicable brand premium. If a Safety Net Entitlement card is held, the veteran should receive a Special Pharmaceutical Benefit free of charge, subject to any brand premium applicable. Medicare Australia will reimburse the dispensing pharmacist the total dispensed price, less the concessional patient contribution and/or brand premium if applicable.

Therapeutic Group Premiums — Authority Processing

- Items attracting a therapeutic group premium are dual listed. Dispensing pharmacists are therefore required to select the appropriate code for those items that are dual listed as authority and non-authority items, in order to correctly charge the patient and claim from Medicare Australia. Those authority prescriptions that grant exemption from a therapeutic group premium will have the letters 'TPX' at the beginning of the telephone Authority approval number, or, in the case of a written approval, will be stamped with the words "This prescription does not attract a therapeutic group premium".

Contact the Department of Veterans' Affairs

Authority Prescription Applications

Applications for authority to prescribe under the Repatriation Pharmaceutical Benefits Scheme (RPBS) should be sent to the Veterans' Affairs Pharmaceutical Advisory Centre (VAPAC) using the free postal service:

REPLY PAID 9998

VAPAC (Veterans' Affairs Pharmaceutical Advisory Centre)

Department of Veterans' Affairs

GPO Box 9998

BRISBANE QLD 4001

For RPBS enquiries and telephone approvals 24 hours a day the Freecall number is: 1800 552 580

Departmental pharmacists answer applications for prior approval for non-Schedule items and Authority application calls.

WOUND ASSESSMENT AND DRESSING IDENTIFICATION

It is essential to define the aetiology of the wound before selecting a dressing. Recommendations are based on wound type, colour of wound base, depth of wound, and amount of exudate.

This wound chart adheres to the MOIST WOUND concept of healing and wound dressings are described below as ABSORBING or MOISTURE DONATING.

Most wound healing products are designed to remain in situ for several days, with the exception of those for infected wounds which should be changed daily. The quantities and repeats listed in the Repatriation Schedule are considered to be adequate to manage the treatment of a wound for two weeks to one month, when an assessment of the wound's healing process should be undertaken.

DRESSINGS

Pink Epithelialising Wound

Aim: To protect and promote epithelialisation. Epithelialising wounds normally are superficial and only produce a light exudate.

(A) Covering	- Film; - Film Island	- Gauze—Paraffin; - Non-adherent
(B) Absorbing	- Foam (Light Exudate); - Hydroactive (Superficial Wound—Light Exudate)	Hydrocolloid (Superficial Wound—Light Exudate)

Red Granulating Wound

Aims: (1) to protect the granulating tissue; (2) to encourage epithelialisation; (3) to absorb excess exudate.

LIGHT EXUDATE:	Superficial	Cavity
(A) Absorbing	- Foam (Light Exudate); - Hydroactive (Superficial Wound—Light Exudate); - Hydrocolloid (Superficial Wound—Light Exudate)	Hydrocolloid (Cavity Wound)
(B) Moisture donating	- Hydrogel—Amorphous; - Hydrogel—Sheet	Hydrogel—Amorphous
HIGH EXUDATE:	Superficial	Cavity
(A) Absorbing	- Alginate (Superficial Wound); - Foam—Heavy Exudate; - Hydroactive (Superficial Wound—Moderate Exudate); - Hydrocolloid (Superficial Wound—Moderate/High Exudate)	- Alginate (Cavity Wound); - Foam—Moderate Exudate (see “cavity conforming” product); - Hydroactive (Cavity Wound); - Hydrocolloid (Cavity Wound)
(B) Moisture donating	NOT APPROPRIATE	

Yellow Sloughy Wound

Aims: (1) to remove slough; (2) to encourage granulation; (3) to absorb excess exudate.

LIGHT EXUDATE:	Superficial	Cavity
(A) Absorbing	- Cadexomer Iodine; - Foam—Light Exudate; - Foam with Charcoal; - Hydroactive (Superficial Wound—Moderate Exudate); - Hydrocolloid (Superficial Wound—Moderate Exudate)	- Cadexomer Iodine; - Hydrocolloid (Cavity Wound)
(B) Moisture Donating	- Hydrogel—Amorphous; - Hydrogel—Sheet	Hydrogel—Amorphous
HIGH EXUDATE:	Superficial	Cavity
(A) Absorbing	- Alginate (Superficial Wound); - Cadexomer Iodine; - Foam—Heavy Exudate; - Hydroactive (Superficial Wound—Moderate/High Exudate); - Hydrocolloid (Superficial Wound—Moderate/High Exudate)	- Alginate (Cavity Wound); - Cadexomer Iodine; - Hydrocolloid (Cavity Wound)
(B) Moisture donating	NOT APPROPRIATE	

Black Necrotic Wound

Aims: To remove eschar by — (1) sharp debridement, e.g., scissor/scalpel and/or (2) rehydration and autolytic debridement. (These wounds usually produce a LIGHT EXUDATE.)

DRY / LIGHT EXUDATE:	Superficial	Cavity
(A) Absorbing	• Hydroactive (Superficial Wound—Light Exudate); • Hydrocolloid (Superficial Wound—Light/Moderate Exudate)	• Hydrocolloid (Cavity Wound)
(B) Moisture donating	• Hydrogel—Amorphous; • Hydrogel—Sheet	• Hydrogel—Amorphous; • Hydrogel—Sheet

Infected Wounds

Aims: (1) to clear the infection with systemic antibiotics; (2) to absorb excess exudate; (3) to remove slough if present; (4) to decrease bacterial burden - by applying a Silver dressing or Cadexomer Iodine dressing.

Malodorous Wounds

Aims: (1) to clear infection if present; (2) to remove slough if present; (3) to clear colonising odour-producing bacteria in slough — by applying metronidazole gel, a Silver dressing or a Cadexomer Iodine dressing; (4) to absorb excess exudate.

Products: Activated Charcoal; Alginate with Charcoal; Foam with Charcoal; Silver dressing; Cadexomer Iodine dressing.

Minor Skin Trauma

Aims: (1) to stop bleeding; (2) to prevent infection; (3) to minimise the surface defect; (4) to promote epithelialisation.

Ordering Products

Ordering Coloplast Products

Coloplast dressings are available via a range of distributors. However, Coloplast's principal agreement to ensure correct RPBS Price to Pharmacy and ready supply has been secured with Independence Australia on 1300 788 855 and BrightSky on 1300 290 400. Please note that Coloplast is unable to guarantee ready supply or rebate for price differences on purchases outside these distributors.

Ordering Hartmann Products

Hartmann wound dressings are available through HARTMANN and Independence Australia only. If you would like to order Hartmann Wound Care products, please call HARTMANN customer service on 1800 805 839 or Independence Australia on 1300 788 855.

Ordering Molnlycke Healthcare Products

Molnlycke Healthcare products are distributed through leading pharmacy distributors. To best ensure product availability at RPBS agreed prices, special arrangements have been made with API and Independence Australia Health Solutions (IAHS). IAHS orders can be placed on: Tel: 1300 788 855; or Email customerservice@independenceaustralia.com. Molnlycke Healthcare are not able to ensure product availability or pricing on listed products beyond these two suppliers.

Ordering Smith & Nephew Products

Smith & Nephew products are distributed via the three major wholesalers, API, SIGMA & Symbion. To best ensure product availability at RPBS agreed prices, please order from one of these suppliers. In the event that your preferred wholesaler cannot supply, please contact Smith & Nephew Customer Service on 13 13 60. Smith & Nephew cannot ensure RPBS agreed pricing from distributors other than those aforementioned.

ALIMENTARY TRACT AND METABOLISM.....	1016
STOMATOLOGICAL PREPARATIONS	1016
STOMATOLOGICAL PREPARATIONS.....	1016
DRUGS FOR FUNCTIONAL GASTROINTESTINAL DISORDERS.....	1016
DRUGS FOR FUNCTIONAL GASTROINTESTINAL DISORDERS	1016
BELLADONNA AND DERIVATIVES, PLAIN	1016
DRUGS FOR CONSTIPATION	1016
DRUGS FOR CONSTIPATION.....	1016
ANTIDIARRHEALS, INTESTINAL ANTIINFLAMMATORY/ANTIINFECTIVE AGENTS	1018
ELECTROLYTES WITH CARBOHYDRATES	1018
ANTIPROPULSIVES.....	1018
ANTIOBESITY PREPARATIONS, EXCL. DIET PRODUCTS	1018
ANTIOBESITY PREPARATIONS, EXCL. DIET PRODUCTS	1018
VITAMINS.....	1019
VITAMIN B1, PLAIN AND IN COMBINATION WITH VITAMIN B6 AND B12.....	1019
VITAMIN B-COMPLEX, INCL. COMBINATIONS	1019
MINERAL SUPPLEMENTS	1019
CALCIUM.....	1019
OTHER MINERAL SUPPLEMENTS.....	1019
BLOOD AND BLOOD FORMING ORGANS	1020
ANTITHROMBOTIC AGENTS.....	1020
ANTITHROMBOTIC AGENTS	1020
ANTIANEMIC PREPARATIONS	1020
IRON PREPARATIONS	1020
VITAMIN B12 AND FOLIC ACID	1021
BLOOD SUBSTITUTES AND PERFUSION SOLUTIONS.....	1021
IRRIGATING SOLUTIONS	1021
CARDIOVASCULAR SYSTEM	1021
VASOPROTECTIVES	1021
AGENTS FOR TREATMENT OF HEMORRHOIDS AND ANAL FISSURES FOR TOPICAL USE.....	1021
DERMATOLOGICALS	1022
ANTIFUNGALS FOR DERMATOLOGICAL USE	1022
ANTIFUNGALS FOR TOPICAL USE.....	1022
ANTIFUNGALS FOR SYSTEMIC USE.....	1022
EMOLLIENTS AND PROTECTIVES	1023
EMOLLIENTS AND PROTECTIVES	1023
PROTECTIVES AGAINST UV-RADIATION	1023
ANTIPRURITICS, INCL. ANTIHISTAMINES, ANESTHETICS, ETC.	1024
ANTIPRURITICS, INCL. ANTIHISTAMINES, ANESTHETICS, ETC.....	1024
ANTIPSORIATICS.....	1024
ANTIPSORIATICS FOR TOPICAL USE.....	1024
ANTIBIOTICS AND CHEMOTHERAPEUTICS FOR DERMATOLOGICAL USE	1024
ANTIBIOTICS FOR TOPICAL USE	1024
CORTICOSTEROIDS, DERMATOLOGICAL PREPARATIONS	1024

CORTICOSTEROIDS, PLAIN	1024
ANTISEPTICS AND DISINFECTANTS	1025
ANTISEPTICS AND DISINFECTANTS	1025
OTHER DERMATOLOGICAL PREPARATIONS	1025
OTHER DERMATOLOGICAL PREPARATIONS	1025
GENITO URINARY SYSTEM AND SEX HORMONES	1027
GYNECOLOGICAL ANTIINFECTIVES AND ANTISEPTICS	1027
ANTIINFECTIVES AND ANTISEPTICS, EXCL. COMBINATIONS WITH CORTICOSTEROIDS	1027
OTHER GYNECOLOGICALS	1027
OTHER GYNECOLOGICALS	1027
UROLOGICALS	1027
UROLOGICALS	1027
DRUGS USED IN BENIGN PROSTATIC HYPERTROPHY	1030
ANTIINFECTIVES FOR SYSTEMIC USE	1031
ANTIBACTERIALS FOR SYSTEMIC USE	1031
MACROLIDES, LINCOSAMIDES AND STREPTOGRAMINS	1031
ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS	1031
ANTINEOPLASTIC AGENTS	1031
ANTIMETABOLITES	1031
IMMUNOSUPPRESSANTS	1031
IMMUNOSUPPRESSANTS	1031
MUSCULO-SKELETAL SYSTEM	1032
TOPICAL PRODUCTS FOR JOINT AND MUSCULAR PAIN	1032
TOPICAL PRODUCTS FOR JOINT AND MUSCULAR PAIN	1032
DRUGS FOR TREATMENT OF BONE DISEASES	1032
DRUGS AFFECTING BONE STRUCTURE AND MINERALIZATION	1032
NERVOUS SYSTEM	1033
ANALGESICS	1033
OPIOIDS	1033
OTHER ANALGESICS AND ANTIPYRETICS	1033
PSYCHOLEPTICS	1035
ANXIOLYTICS	1035
HYPNOTICS AND SEDATIVES	1035
OTHER NERVOUS SYSTEM DRUGS	1036
DRUGS USED IN ADDICTIVE DISORDERS	1036
ANTIPARASITIC PRODUCTS, INSECTICIDES AND REPELLENTS	1036
ANTHELMINTICS	1036
ANTINEMATODAL AGENTS	1036
RESPIRATORY SYSTEM	1037
NASAL PREPARATIONS	1037

DECONGESTANTS AND OTHER NASAL PREPARATIONS FOR TOPICAL USE	1037
NASAL DECONGESTANTS FOR SYSTEMIC USE.....	1037
ANTIHISTAMINES FOR SYSTEMIC USE	1037
ANTIHISTAMINES FOR SYSTEMIC USE.....	1037
SENSORY ORGANS	1038
OTOLOGICALS	1038
OTHER OTOLOGICALS.....	1038
VARIOUS	1038
ALL OTHER THERAPEUTIC PRODUCTS	1038
ALL OTHER THERAPEUTIC PRODUCTS.....	1038
GENERAL NUTRIENTS	1039
OTHER NUTRIENTS	1039
ALL OTHER NON-THERAPEUTIC PRODUCTS	1039
ALL OTHER NON-THERAPEUTIC PRODUCTS	1039

ALIMENTARY TRACT AND METABOLISM

STOMATOLOGICAL PREPARATIONS

STOMATOLOGICAL PREPARATIONS

Antifungives and antiseptics for local oral treatment

CHLORHEXIDINE

chlorhexidine gluconate 0.2% mouthwash, 250 mL

4161B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	‡1	..	..	16.86	6.60	Plaqacide [OB]

chlorhexidine gluconate 0.2% mouthwash, 300 mL

4204G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	‡1	..	..	19.72	6.60	Savacol Mouth and Throat Rinse [OM]

NYSTATIN

nystatin 100 000 units/mL oral liquid, 24 mL

10854G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	‡1	1	..	18.50	6.60	^a Pharmacy Action Nystatin Oral Drops [GQ]	^a Trust Nystatin Oral Drops [CR]
			..	19.74	6.60	^a Mycostatin Oral Drops [LN]	

DRUGS FOR FUNCTIONAL GASTROINTESTINAL DISORDERS

DRUGS FOR FUNCTIONAL GASTROINTESTINAL DISORDERS

Synthetic anticholinergics, esters with tertiary amino group

MEBEVERINE

mebeverine hydrochloride 135 mg tablet, 90

4328T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	29.84	6.60	^a APO-Mebeverine [TX]	^a Colese [AF]
			..	34.34	6.60	^a Colofac [GO]	

BELLADONNA AND DERIVATIVES, PLAIN

Belladonna alkaloids, semisynthetic, quaternary ammonium compounds

HYOSCINE BUTYLBROMIDE

hyoscine butylbromide 20 mg/mL injection, 5 x 1 mL ampoules

4279F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	22.80	6.60	^a Buscopan [VZ]	^a HYOSCINE BUTYLBROMIDE SXP [XC]

DRUGS FOR CONSTIPATION

DRUGS FOR CONSTIPATION

Softeners, emollients

DOCUSATE

docusate sodium 50 mg tablet, 100

4200C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	18.88	6.60	Coloxyl 50 [AS]

Contact laxatives

BISACODYL

bisacodyl 10 mg suppository, 12

10580W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	4	..	*22.38	6.60	Petrus Bisacodyl Suppositories [PP]

bisacodyl 10 mg suppository, 10

10578R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	5	..	*24.66	6.60	^a Petrus Bisacodyl Suppositories [PP]
			..	*25.95	6.60	^a Dulcolax [VZ]

▪ **DOCUSATE + SENNOSIDE B****docosate sodium 50 mg + sennoside B 8 mg tablet, 100**

4028B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	18.97	6.60	Soflax [EA]

docosate sodium 50 mg + sennoside B 8 mg tablet, 90

10177P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	2	..	17.86	6.60	^a Pharmacy Action Laxative with Senna [GQ]	
			..	17.90	6.60	^a Chemists' Own Laxative with Senna [RW]	^a Colaxsen [AS]
						^a Coloxyl with Senna [LN]	^a Co-Senna [PP]
						^a Trust Coloxease [CR]	

▪ **SENNOSIDE B****sennoside B 7.5 mg tablet, 100**

4455L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	17.42	6.60	^a Senna-Gen [PP]
			..	18.49	6.60	^a Senokot [RC]

Bulk-forming laxatives▪ **DRY PSYLLIUM HUSK****dry psyllium husk 3.5 g powder for oral liquid, 30 sachets**

4285M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	1	..	21.78	6.60	Fybogel [RC]

▪ **PSYLLIUM HUSK POWDER****psyllium husk powder 3.4 g/7 g powder for oral liquid, 336 g**

4422R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	1	..	22.11	6.60	Fibre Health Natural Granular [PP]
			..	25.28	6.60	Metamucil Natural Granular [PY]

psyllium husk powder 3.4 g/5.9 g powder for oral liquid, 283 g

4419N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	1	..	25.28	6.60	Metamucil Orange Smooth [PY]

▪ **RHAMNUS FRANGULA + STERCULIA****rhamnus frangula 80 mg/g + sterculia 620 mg/g granules, 500 g**

4558X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	1	..	26.12	6.60	Normacol Plus [NE]

Enemas▪ **CITRIC ACID + LAURYL SULFOACETATE SODIUM + SORBITOL****sodium citrate dihydrate 450 mg/5 mL + lauryl sulfoacetate sodium 45 mg/5 mL + sorbitol 3.125 g/5 mL enema, 4 x 5 mL**

4462W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	17.02	6.60	Micolette [AE]

Other drugs for constipation▪ **GLYCEROL****glycerol 2.8 g suppository, 12**

4246L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	5	..	*24.75	6.60	Petrus Pharmaceuticals Pty Ltd [PP]

glycerol 700 mg suppository, 12

10586E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	5	..	*23.91	6.60	Petrus Pharmaceuticals Pty Ltd [PP]

glycerol 1.4 g suppository, 12

10596Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	5	..	*24.27	6.60	Petrus Pharmaceuticals Pty Ltd [PP]

■ ANTIDIARRHEALS, INTESTINAL ANTIINFLAMMATORY/ANTIINFECTIVE AGENTS

ELECTROLYTES WITH CARBOHYDRATES

Oral rehydration salt formulations

■ SODIUM CHLORIDE + POTASSIUM CHLORIDE + GLUCOSE MONOHYDRATE + CITRIC ACID

sodium chloride 470 mg + potassium chloride 300 mg (potassium 4 mmol) + glucose monohydrate 3.56 g + sodium acid citrate 530 mg powder for oral liquid, 10 x 4.9 g sachets

10574M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	‡1	..	..	16.83	6.60	restore O.R.S. [EA]

ANTIPROPULSIVES

Antipropulsives

■ LOPERAMIDE

loperamide hydrochloride 2 mg capsule, 12

10592L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	13.51	6.60	Gastrex [CR]

loperamide hydrochloride 2 mg capsule, 20

11135C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	13.82	6.60	^a Gastrex [CR]	^a Pharmacy Action Diarrhoea Relief [GQ]

■ ANTI OBESITY PREPARATIONS, EXCL. DIET PRODUCTS

ANTI OBESITY PREPARATIONS, EXCL. DIET PRODUCTS

Peripherally acting antiobesity products

■ ORLISTAT

Note The patient should be ideally enrolled in an exercise program and be receiving supplemental vitamins.

Authority required

Obesity

Treatment Phase: Initial treatment

Clinical criteria:

- Patient must have a Body Mass Index (BMI) greater than or equal to 35 with no known co-morbidities; OR
- Patient must have a BMI greater than or equal to 30 with 1 or more of the following co-morbidities;(i) diabetes;(ii) ischaemic heart disease;(iii) psychiatric conditions;(iv) hypertension, **AND**
- Patient must be receiving, or enrolled to receive, professional dietetic and weight management advice (where this is available), **AND**
- The treatment must not exceed 12 months in total from initial application, **AND**
- Patient must not receive more than 1 continuous treatment in a lifetime.

The prescriber must provide the patient's initial body weight and BMI at the time of application.

Authority required

Obesity

Treatment Phase: Continuing treatment (3 to 6 months following commencement)

Clinical criteria:

- Patient must have previously been issued with an authority prescription for this drug, **AND**
- Patient must have reduced their initial body weight by 2.5 kg or 2.5% (whichever is the lesser) during the period 3 to 6 months following commencement of treatment with this drug, **AND**
- The treatment must not exceed 12 months in total from initial application, **AND**
- Patient must not receive more than 1 continuous treatment in a lifetime, **AND**
- Patient must be receiving, or enrolled to receive, professional dietetic and weight management advice (where this is available).

Authority required

Obesity

Treatment Phase: Continuing treatment (6 to 12 months following commencement)

Clinical criteria:

- Patient must have previously been issued with an authority prescription for this drug, **AND**
- Patient must have reduced their initial body weight by 5 kg or 5% (whichever is the lesser) during the period 6 to 12 months following commencement of treatment with this drug, **AND**
- The treatment must not exceed 12 months in total from initial application, **AND**
- Patient must not receive more than 1 continuous treatment in a lifetime, **AND**
- Patient must be receiving, or enrolled to receive, professional dietetic and weight management advice (where this is available).

orlistat 120 mg capsule, 84

4570M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	134.68	6.60	Xenical [PB]

■ **VITAMINS****VITAMIN B1, PLAIN AND IN COMBINATION WITH VITAMIN B6 AND B12***Vitamin B1, plain*■ **THIAMINE****thiamine hydrochloride 100 mg tablet, 100**

4043T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	16.05	6.60	Betavit [PP]

VITAMIN B-COMPLEX, INCL. COMBINATIONS*Vitamin B-complex, plain*■ **LYSINE + THIAMINE + PYRIDOXINE + CYANOCOBALAMIN + FERRIC PYROPHOSPHATE****lysine hydrochloride 300 mg/10 mL + thiamine hydrochloride 10 mg/10 mL + pyridoxine hydrochloride 5 mg/10 mL + cyanocobalamin 25 microgram/10 mL + iron (as ferric pyrophosphate) 10 mg/10 mL oral liquid, 200 mL**

4493L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	2	..	18.04	6.60	Accomin Adult Tonic [PF]

■ **MINERAL SUPPLEMENTS****CALCIUM***Calcium*■ **CALCIUM****Restricted benefit**

Hyperphosphataemia

Clinical criteria:

- The condition must be associated with chronic renal failure.

calcium carbonate 1.5 g (calcium 600 mg) tablet, 120

4142B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	1	..	*23.16	6.60	^a CAL-600 [PP]	^a Cal-care 600 mg [CR]

calcium carbonate 1.25 g (calcium 500 mg) chewable tablet, 120

11845K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	1	..	*27.88	6.60	Cal-500 [PP]

■ **CALCIUM****Restricted benefit**

Hypocalcaemia

Restricted benefit

Osteoporosis

Restricted benefit

Proven calcium malabsorption

calcium carbonate 1.5 g (calcium 600 mg) tablet, 120

4082W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	1	..	17.59	6.60	^a CAL-600 [PP]	^a Cal-care 600 mg [CR]

calcium carbonate 1.25 g (calcium 500 mg) chewable tablet, 120

11862H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	19.95	6.60	Cal-500 [PP]

OTHER MINERAL SUPPLEMENTS*Magnesium*■ **MAGNESIUM****Restricted benefit**

Hypomagnesaemia

The condition must be documented in the patient's medical records.

BLOOD AND BLOOD FORMING ORGANS

magnesium 37.4 mg tablet, 50

4321K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	18.04	6.60	Amcal Mag-A [IG] Pharmacy Care Magnesium [SI]	Mag-Sup [PP]
			..	18.61	6.60	Magmin [BB]	

■ BLOOD AND BLOOD FORMING ORGANS

■ ANTITHROMBOTIC AGENTS

ANTITHROMBOTIC AGENTS

Platelet aggregation inhibitors excl. heparin

■ ASPIRIN

aspirin 100 mg tablet, 112

10590J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	14.14	6.60	Spren 100 [OW]

aspirin 100 mg tablet, 90

4076M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	20.06	6.60	Cardiprin 100 [RC]

■ ASPIRIN

Note The enteric coated preparations are for patients with a significant risk of gastrointestinal bleeding.

aspirin 100 mg enteric tablet, 84

4077N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	1	..	18.36	6.60	Cardasa [AF] ^a Cartia [AS]	^a Pharmacy Action Low Dose Aspirin [GQ]
						^a Trust Aspirin EC 100 [CR]	

aspirin 100 mg enteric capsule, 84

4078P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	19.15	6.60	Astrix [YN]

■ CLOPIDOGREL

Note Pharmaceutical benefits that have the forms clopidogrel tablet 75 mg (as besilate) and clopidogrel tablet 75 mg (as hydrogen sulfate) are equivalent for the purposes of substitution.

Authority required

For use in patients pre- and post-angioplasty

clopidogrel 75 mg tablet, 28

10169F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	3	..	16.23	6.60	^a BTC Clopidogrel [JB] ^a Clopidogrel GH [GQ]	^a Clopidogrel APOTEX [GX] ^a Plidogrel [RF]

clopidogrel 75 mg tablet, 28

4179Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	3	..	16.23	6.60	^a APO-Clopidogrel [TX] ^a Clopidogrel AN [EA] ^a Iscover [AV] ^a Plavacor 75 [CR]	^a Blooms the Chemist Clopidogrel [IB] ^a Clopidogrel Sandoz Pharma [HX] ^a Piax [AF]

■ ANTIANEMIC PREPARATIONS

IRON PREPARATIONS

Iron bivalent, oral preparations

■ FERROUS FUMARATE

ferrous fumarate 200 mg (iron 65.7 mg) tablet, 60

10594N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	18.69	6.60	Ferro-tab [AE]

Iron in combination with folic acid

■ FERROUS FUMARATE + FOLIC ACID

ferrous fumarate 310 mg (iron 100 mg) + folic acid 350 microgram tablet, 60

10579T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	19.71	6.60	Ferro-f-tab [AE]

VITAMIN B12 AND FOLIC ACID

Vitamin B12 (cyanocobalamin and analogues)

■ HYDROXOCOBALAMIN

Note One injection of hydroxocobalamin 1 mg every three months provides appropriate maintenance therapy in vitamin B₁₂ deficiencies.

Note Pharmaceutical benefits that have the form hydroxocobalamin injection 1 mg (as acetate) in 1 mL and pharmaceutical benefits that have the form hydroxocobalamin injection 1 mg (as chloride) in 1 mL are equivalent for the purposes of substitution.

Restricted benefit

Pernicious anaemia

Restricted benefit

Proven vitamin B12 deficiencies other than pernicious anaemia

Restricted benefit

Anaemias associated with vitamin B12 deficiency

Clinical criteria:

- Patient must have had a gastrectomy, **AND**
- The treatment must be for prophylaxis.

hydroxocobalamin 1 mg/mL injection, 3 x 1 mL ampoules

10577Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	16.40	6.60	^a Vita-B12 [GH]

hydroxocobalamin 1 mg/mL injection, 3 x 1 mL ampoules

10587F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	16.40	6.60	^a Neo-B12 [PF]

Folic acid and derivatives

■ FOLIC ACID

folic acid 500 microgram tablet, 100

10584C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	..	..	*16.22	6.60	^a Foltabs 500 [PP]	^a Megafol 0.5 [AF]

■ FOLIC ACID

Note The 5 mg strength tablet should be used in malabsorption states only.

folic acid 5 mg tablet, 100

10573L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	1	..	*18.30	6.60	Megafol 5 [AF]

■ BLOOD SUBSTITUTES AND PERFUSION SOLUTIONS

IRRIGATING SOLUTIONS

Salt solutions

■ SODIUM CHLORIDE

sodium chloride 0.9% (9 g/L) solution, 1 L bottle

4461T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	2	..	15.85	6.60	Baxter Healthcare Pty Ltd [BX]

sodium chloride 0.9% (4.5 g/500 mL) solution, 500 mL bottle

4460R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	2	..	15.59	6.60	Baxter Healthcare Pty Ltd [BX]

■ CARDIOVASCULAR SYSTEM

■ VASOPROTECTIVES

AGENTS FOR TREATMENT OF HEMORRHOIDS AND ANAL FISSURES FOR TOPICAL USE

Other agents for treatment of hemorrhoids and anal fissures for topical use

DERMATOLOGICALS

■ ZINC OXIDE + PERU BALSAM + BENZYL BENZOATE

zinc oxide 300 mg + peru balsam 50 mg + benzyl benzoate 33 mg suppository, 12

4040P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	1	..	18.05	6.60	Anusol [JT]

zinc oxide 10.75% + peru balsam 1.88% + benzyl benzoate 1.25% ointment, 50 g

4039N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	1	..	18.99	6.60	Anusol [JT]

■ DERMATOLOGICALS

■ ANTIFUNGALS FOR DERMATOLOGICAL USE

ANTIFUNGALS FOR TOPICAL USE

Imidazole and triazole derivatives

■ CLOTRIMAZOLE

clotrimazole 1% cream, 20 g

4004R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	1	..	14.07	6.60	^a Pharmacy Action Anti-Fungal Cream [GQ]
			..	14.38	6.60	^a Clonea [AF]

Other antifungals for topical use

■ AMOROLFINE

Restricted benefit

Onychomycosis

amorolfine 5% solution, 5 mL

4010C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	1	..	48.58	6.60	^a Myconail [AE]
			..	68.66	6.60	^a Pharmacy Action Anti-Fungal Nail Treatment [GQ]
			..	84.83	6.60	^a Aporyl [TX]
			..	93.58	6.60	^a Loceryl [GA]

■ TERBINAFINE

Restricted benefit

Tinea pedis

terbinafine hydrochloride 1% cream, 15 g

4473K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	±1	1	..	25.47	6.60	^a Lamisil [GK]	^a Pharmacy Action Pharmisil [GQ]
						^a Trust Terbinafine Cream [CR]	

terbinafine 1% gel, 15 g

4463X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	26.74	6.60	Lamisil DermGel [GK]

■ TOLNAFTATE

tolnaftate 0.07% spray, 100 g

4481W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	19.56	6.60	Tinaderm [BN]

ANTIFUNGALS FOR SYSTEMIC USE

Antifungals for systemic use

■ TERBINAFINE

Authority required

Onychomycosis

Clinical criteria:

- The condition must be due to dermatophyte infection proven by microscopy and confirmed by an Approved Pathology Provider; OR
- The condition must be due to dermatophyte infection proven by culture and confirmed by an Approved Pathology Provider.

terbinafine 250 mg tablet, 42

4011D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	1	..	30.93	6.60	^a APO-Terbinafine [TX]	^a Lamisil (Novartis Pharmaceuticals Australia Pty Limited) [NV]
						^a Tamsil [RW]	^a Terbinafine Sandoz [SZ]
						^a Tinasil [AF]	

EMOLLIENTS AND PROTECTIVES**EMOLLIENTS AND PROTECTIVES***Soft paraffin and fat products***WOOL ALCOHOLS****wool alcohols 6% ointment, 100 g**

4041Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	‡1	1	..	18.77	6.60	Eucerin [BE]

*Carbamide products***UREA****urea 10% cream, 100 g**

4042R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	‡1	2	..	17.10	6.60	Aquacare H.P. [AG]
			..	17.30	6.60	Urederm [KY]

*Other emollients and protectives***GELATIN + PECTIN + CARMELLOSE SODIUM****gelatin 16.7% + pectin 16.7% + carmellose sodium 16.7% paste, 5 g**

4518T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	‡1	..	..	16.82	6.60	Orabase [AS]

GLYCEROL + WHITE SOFT PARAFFIN**glycerol 5% + white soft paraffin 5% lotion, 1 L**

11712K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	‡1	1	..	31.22	6.60	QV Skin Lotion [EO]

SKIN EMOLLIENT**SKIN EMOLLIENT Bath oil 500 mL, 1**

4122Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	‡1	2	..	21.52	6.60	Alpha Keri Bath Oil [MT]
			..	23.62	6.60	QV Bath Oil [EO]
			..	23.70	6.60	Hamilton Skin Therapy Oil [KY]

SKIN EMOLLIENT Lotion 500 mL, 1

4107E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	‡1	2	..	21.52	6.60	Alpha Keri Lotion [MT]

PROTECTIVES AGAINST UV-RADIATION*Protectives against UV-radiation for topical use***BEMOTRIZINOL + OCTOCRYLENE + DIETHYLAMINO HYDROXYBENZOYL HEXYL BENZOATE + TITANIUM DIOXIDE****bemotrizinol 1% + octocrylene 2% + diethylamino hydroxybenzoyl hexyl benzoate 3.5% + titanium dioxide 2% lotion, 125 mL**

11387H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	‡1	2	..	22.13	6.60	Sunsense Ultra SPF 50+ [EO]

SUNSCREENS**SUNSCREENS Cream 75 g, 1**

4307Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	‡1	2	..	22.13	6.60	Sunsense Sensitive SPF 50+ [EO]

DERMATOLOGICALS

SUNSCREENS Lotion (non-alcoholic) 125 mL, 1

4546G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	2	..	20.33	6.60	Aquasun Lotion SPF 18 [PF]

■ ANTIPRURITICS, INCL. ANTIHISTAMINES, ANESTHETICS, ETC.

ANTIPRURITICS, INCL. ANTIHISTAMINES, ANESTHETICS, ETC.

Anesthetics for topical use

■ LIDOCAINE (LIGNOCAINE)

lidocaine (lignocaine) hydrochloride 2% oral liquid, 200 mL

4308R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	104.14	6.60	Xylocaine Viscous [AS]

Other antipruritics

■ TAR + TROLAMINE LAURIL SULFATE

Note For patients who have failed to respond to simple moisturising agents.

tar 2.3% + trolamine lauril sulfate 6% solution, 500 mL

4408B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	2	..	26.37	6.60	Pinetarsol [EO]

■ ANTIPSORIATICS

ANTIPSORIATICS FOR TOPICAL USE

Tars

■ COAL TAR SOLUTION + PHENOL + PRECIPITATED SULFUR

coal tar solution 5% + phenol 0.5% + precipitated sulfur 0.5% gel, 30 g

4505D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	2	..	20.37	6.60	Egopsoryl-TA [EO]

■ ANTIBIOTICS AND CHEMOTHERAPEUTICS FOR DERMATOLOGICAL USE

ANTIBIOTICS FOR TOPICAL USE

Other antibiotics for topical use

■ MUPIROCIN

mupirocin 2% ointment, 15 g

4350Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	20.64	6.60	^a Medicianz Mupirocin Ointment [DZ]
			..	21.44	6.60	^a APO-Mupirocin [TX]
			..	24.51	6.60	^a Bactroban [GK]

■ MUPIROCIN

Restricted benefit

Secondarily infected traumatic skin lesions

mupirocin 2% cream, 15 g

4348W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	24.51	6.60	Bactroban [GK]

■ CORTICOSTEROIDS, DERMATOLOGICAL PREPARATIONS

CORTICOSTEROIDS, PLAIN

Corticosteroids, weak (group I)

■ HYDROCORTISONE ACETATE

hydrocortisone acetate 1% cream, 30 g

11710H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	±1	1	..	17.24	6.60	^a Pharmacy Action Hydrocortisone Cream 1% [GQ]	^a Trust HydroCortic Cream [CR]

■ HYDROCORTISONE ACETATE

Restricted benefit

Corticosteroid-responsive dermatoses

hydrocortisone acetate 1% ointment, 30 g

10831C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	1	..	17.24	6.60	Cortic-DS 1% [AS]

Corticosteroids, potent (group III)■ **BETAMETHASONE VALERATE****betamethasone (as valerate) 0.1% ointment, 30 g**

4132L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	2	..	25.94	6.60	Betnovate [AS]

betamethasone (as valerate) 0.1% cream, 30 g

4131K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	2	..	25.94	6.60	Betnovate [AS]

■ **MOMETASONE****Note** Application to large areas of skin for longer than four weeks is not recommended.**mometasone furoate 0.1% ointment, 50 g**

4343N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	±1	..	..	35.55	6.60	^a Elocon [AL]	^a Momasone [AS]

mometasone furoate 0.1% cream, 50 g

4342M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	35.55	6.60	Elocon [AL]

■ **ANTISEPTICS AND DISINFECTANTS****ANTISEPTICS AND DISINFECTANTS***Iodine products*■ **POVIDONE-IODINE****povidone-iodine 10% solution, 100 mL**

4411E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	25.66	6.60	Betadine Antiseptic Liquid [SW]

■ **OTHER DERMATOLOGICAL PREPARATIONS****OTHER DERMATOLOGICAL PREPARATIONS***Medicated shampoos*■ **SALICYLIC ACID + BENZALKONIUM CHLORIDE + ALCOHOL + COAL TAR SOLUTION + POLYOXYETHYLENE ETHERS****SALICYLIC ACID with COAL TAR SOLUTION Scalp cleanser 20 mg-50 mg per mL (2%-5%), 200 mL, 1**

4560B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	2	..	24.16	6.60	Ionil-T [GA]

■ **SELENIUM SULFIDE****selenium sulfide 2.5% shampoo, 125 mL**

4452H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	18.73	6.60	Selsun [DQ]

■ **TAR + COAL TAR SOLUTION + SALICYLIC ACID****tar 1% + coal tar solution 1% + salicylic acid 2% solution, 250 mL**

4447C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	2	..	22.82	6.60	Sebitar [EO]

Wart and anti-corn preparations■ **SALICYLIC ACID + LACTIC ACID****salicylic acid 16.7% + lactic acid 15% solution, 15 mL**

11959K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	22.22	6.60	Duofilm Solution [GK]

Other dermatologicals

DERMATOLOGICALS

■ DICLOFENAC

Note Maximum quantity of four tubes (original + 3 repeats) in 12 months.

Authority required

Solar (actinic) keratosis

Treatment Phase: Management

Clinical criteria:

- Patient must require topical drug therapy as field treatment for clinically visible and subclinical lesions where other standard treatments are inappropriate.

diclofenac sodium 3% gel, 25 g

4046Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	3	..	59.08	6.60	Solaraze 3% Gel [FK]

■ GLYCEROL

glycerol 15% solution, 1 kg

11708F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	1	..	30.73	6.60	QV Gentle Wash [EO]

■ ICHTHAMMOL

Note For patients who have failed to respond to simple moisturising agents.

ichthammol 1% cream, 50 g

4281H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	2	..	22.18	6.60	Egoderm Cream [EO]

■ ICHTHAMMOL + ZINC OXIDE

Note For patients who have failed to respond to simple moisturising agents.

ichthammol 1% + zinc oxide 15% ointment, 50 g

4280G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	2	..	22.18	6.60	Egoderm Ointment [EO]

■ IMIQUIMOD

Authority required

Superficial basal cell carcinoma

Treatment Phase: Primary treatment

Clinical criteria:

- The condition must be confirmed by a histological diagnosis, **AND**
- The condition must be one where other standard treatments are inappropriate, **AND**
- The condition must require topical drug therapy.

imiquimod 5% cream, 12 x 250 mg sachets

4559Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	1	..	87.55	6.60	^a Aldiq [AF]	^a APO-Imiquimod [TX]
			..	89.83	6.60	^a Aldara [IL]	

■ IMIQUIMOD

Note Pharmaceutical benefits that have the form imiquimod single use sachets and pharmaceutical benefits that have the form imiquimod multi-use pump are equivalent for the purposes of substitution.

Authority required

Solar keratosis

Clinical criteria:

- Patient must require topical drug therapy on the face and scalp as field treatment for clinically visible and subclinical lesions where other standard treatments are inappropriate.

imiquimod 5% cream, 12 x 250 mg sachets

4134N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	1	..	87.55	6.60	^a Aldiq [AF]	^a APO-Imiquimod [TX]
			..	89.83	6.60	^a Aldara [IL]	

imiquimod 5% cream, 2 x 2 g

10106X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	92.10	6.60	^a Aldara Pump [IL]

■ LIGHT LIQUID PARAFFIN + COCOAMPHODIACETATE DISODIUM

light liquid paraffin 3.5% + cocoamphodiacetate disodium 3% lotion, 500 mL

4549K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	2	..	24.47	6.60	Hamilton Skin Therapy Wash [KY]

■ PANTHENOL

Note To be used in conjunction with the scalp cleanser salicylic acid with coal tar solution and pine tar (code 4447C).

panthenol conditioner, 200 g

4510J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	2	..	18.83	6.60	SebiRinse [EO]

■ ZINC OXIDE + MAIZE STARCH + PURIFIED TALC + CHLORPHENESIN

zinc oxide 25% + maize starch 55.85% + purified talc 18.07% + chlorphenesin 1% powder, 100 g

4497Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	1	..	17.15	6.60	Z.S.C. [RW]

■ GENITO URINARY SYSTEM AND SEX HORMONES

■ GYNECOLOGICAL ANTIINFECTIVES AND ANTISEPTICS

ANTIINFECTIVES AND ANTISEPTICS, EXCL. COMBINATIONS WITH CORTICOSTEROIDS

Antibiotics

■ NYSTATIN

nystatin 20 000 units/g vaginal cream, 75 g

4013F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	1	..	18.43	6.60	Nilstat [AS]

Imidazole derivatives

■ CLOTRIMAZOLE

clotrimazole 1% vaginal cream, 35 g

4016J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	±1	..	..	18.34	6.60	^a Clonea 6 Day Cream [AF]	^a Pharmacy Action FemCream [GQ]
			..	19.55	6.60	^a Trust Fem V 6 Day Cream [CR]	
						^a APO-Clotrimazole 6 Day Cream [TX]	

clotrimazole 2% vaginal cream, 20 g

4017K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	±1	..	..	19.55	6.60	^a APO-Clotrimazole 3 Day Cream [TX]	^a Clonea 3 Day Cream [AF]
						^a Pharmacy Action FemCream 3 Day Cream [GQ]	^a Trust Fem V 3 Day Cream [CR]

■ OTHER GYNECOLOGICALS

OTHER GYNECOLOGICALS

■ ACETIC ACID + HYDROXYQUINOLINE + RICINOLEIC ACID

acetic acid 0.94% + oxyquinoline sulfate 0.025% + ricinoleic acid 0.75% vaginal gel, 100 g

4434J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	35.05	6.60	Aci-Jel [CU]

■ UROLOGICALS

UROLOGICALS

Drugs used in erectile dysfunction

■ ALPROSTADIL

Authority required

Erectile dysfunction

Clinical criteria:

- The condition must be vasculogenic; OR
- The condition must be psychogenic; OR

GENITO URINARY SYSTEM AND SEX HORMONES

- The condition must be neurogenic, **AND**
- Patient must have a specific accepted war-caused or service-related disability.

Population criteria:

- Patient must be male.

Authorisation will not be given for any additional prescriptions within 6 months or for any increased quantities or repeats.

alprostadil 20 microgram injection [1 chamber] (&) inert substance diluent [0.5 mL chamber], 2 dual chamber syringes

4580C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	3	..	*134.91	6.60	Caverject Impulse [PF]

alprostadil 10 microgram injection [1 chamber] (&) inert substance diluent [0.5 mL chamber], 2 dual chamber syringes

4579B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	3	..	*106.59	6.60	Caverject Impulse [PF]

■ AVANAFIL

Authority required

Erectile dysfunction

Clinical criteria:

- The condition must be vasculogenic; OR
- The condition must be psychogenic; OR
- The condition must be neurogenic, **AND**
- Patient must have a specific accepted war-caused or service-related disability.

Population criteria:

- Patient must be male.

Authorisation will not be given for any additional prescriptions within 6 months or for any increased quantities or repeats.

avanafil 50 mg tablet, 4

11837B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	50.73	6.60	Spedra [FK]

avanafil 200 mg tablet, 4

11860F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	55.03	6.60	Spedra [FK]

avanafil 100 mg tablet, 4

11861G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	52.88	6.60	Spedra [FK]

■ SILDENAFIL

Authority required

Erectile dysfunction

Clinical criteria:

- The condition must be vasculogenic; OR
- The condition must be psychogenic; OR
- The condition must be neurogenic, **AND**
- Patient must have a specific accepted war-caused or service-related disability.

Population criteria:

- Patient must be male.

Authorisation will not be given for any additional prescriptions within 6 months or for any increased quantities or repeats.

sildenafil 25 mg tablet, 4

4584G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	25.09	6.60	^a Viagra [UJ]	
			..	55.97	6.60	^a Sildenafil Actavis [EA]	^a Sildenafil APOTEX [GX]
						^a Sildenafil Sandoz [SZ]	^a Vasafil 25 [RW]
						^a Vedafile [AF]	
			..	55.98	6.60	^a APO-Sildenafil [TX]	

sildenafil 100 mg tablet, 4

4586J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	32.02	6.60	^a Viagra [UJ]	
			..	72.54	6.60	^a APO-Sildenafil [TX]	^a NOUMED Sildenafil [VO]
						^a Sildenafil Actavis [EA]	^a Sildenafil APOTEX [GX]
						^a Sildenafil generichealth [GQ]	^a Sildenafil Sandoz [SZ]
						^a Vasafil 100 [RW]	^a Vedafile [AF]

sildenafil 50 mg tablet, 4

4585H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	28.66	6.60	^a Viagra [UJ]	
			..	67.97	6.60	^a APO-Sildenafil [TX]	^a NOUMED Sildenafil [VO]
						^a Sildenafil Actavis [EA]	^a Sildenafil APOTEX [GX]
						^a Sildenafil Sandoz [SZ]	^a Vasafil 50 [RW]
						^a Vedafile [AF]	

■ TADALAFIL**Authority required**

Erectile dysfunction

Clinical criteria:

- The condition must be vasculogenic; OR
- The condition must be psychogenic; OR
- The condition must be neurogenic, **AND**
- Patient must have a specific accepted war-caused or service-related disability.

Population criteria:

- Patient must be male.

Authorisation will not be given for any additional prescriptions within 6 months or for any increased quantities or repeats.

tadalafil 10 mg tablet, 4

4596X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	104.78	6.60	^a Cialis [LY]	^a Cipla Tadalafil [LR]
						^a Tadalafil Sandoz [SZ]	

tadalafil 20 mg tablet, 4

4597Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	104.78	6.60	^a Cialis [LY]	^a Cipla Tadalafil [LR]
						^a Tadalacord [CR]	^a Tadalafil Sandoz [SZ]

■ VARDENAFIL**Authority required**

Erectile dysfunction

Clinical criteria:

- The condition must be vasculogenic; OR
- The condition must be psychogenic; OR
- The condition must be neurogenic, **AND**
- Patient must have a specific accepted war-caused or service-related disability.

Population criteria:

- Patient must be male.

Authorisation will not be given for any additional prescriptions within 6 months or for any increased quantities or repeats.

varafenafil 20 mg tablet, 4

4302K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	82.11	6.60	Levitra [BN]

varafenafil 10 mg tablet, 4

4290T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	72.36	6.60	Levitra [BN]

Other urologicals**■ BICARBONATE + CITRIC ACID + TARTARIC ACID**

Note Pharmaceutical benefits that have the forms sodium bicarbonate 1.76 g + sodium citrate 630 mg + citric acid 720 mg + tartaric acid 890 mg powder for oral liquid and sodium bicarbonate 1.76 g + citric acid 720 mg + sodium citrate 630 mg + tartaric acid 890 mg effervescent granules are equivalent for the purposes of substitution.

Restricted benefit

Urinary symptoms

Clinical criteria:

- The treatment must be for when antibiotic or other therapy alone is inappropriate.

sodium bicarbonate 1.76 g + sodium citrate 630 mg + citric acid 720 mg + tartaric acid 890 mg powder for oral liquid, 28 x 4 g sachets

4049D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	±1	4	..	18.22	6.60	^a Urafol [EA]	^a Ural Sachets [AS]

sodium bicarbonate 1.76 g + citric acid 720 mg + sodium citrate 630 mg + tartaric acid 890 mg effervescent granules, 28 x 4 g sachets

12179B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	4	..	18.22	6.60	^a Trust Cystitis Relief [CR]

DRUGS USED IN BENIGN PROSTATIC HYPERTROPHY

Alpha-adrenoreceptor antagonists

■ ALFUZOSIN

Authority required

Benign prostatic hyperplasia

Clinical criteria:

- Patient must have lower urinary tract symptoms.

alfuzosin hydrochloride 10 mg modified release tablet, 30

4277D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	63.78	6.60	Xatral SR [SW]

■ DUTASTERIDE + TAMSULOSIN

Authority required

Benign prostatic hyperplasia

Clinical criteria:

- Patient must have lower urinary tract symptoms.

dutasteride 500 microgram + tamsulosin hydrochloride 400 microgram modified release capsule, 30

10102Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	35.19	6.60	Duodart 500ug/400ug [GK]

■ SILODOSIN

Authority required

Benign prostatic hyperplasia

Clinical criteria:

- Patient must have lower urinary tract symptoms.

silodosin 4 mg capsule, 30

12079R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	27.78	6.60	Urorec [YN]

silodosin 8 mg capsule, 30

12077P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	43.56	6.60	Urorec [YN]

■ TAMSULOSIN

Authority required

Benign prostatic hyperplasia

Clinical criteria:

- Patient must have lower urinary tract symptoms.

tamsulosin hydrochloride 400 microgram modified release tablet, 30

4070F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	63.78	6.60	^a Flomaxtra [LS]	^a Tamsulosin Sandoz SR [SZ]

Testosterone-5-alpha reductase inhibitors

■ DUTASTERIDE

Authority required

Benign prostatic hyperplasia

Clinical criteria:

- Patient must have lower urinary tract symptoms.

dutasteride 500 microgram capsule, 30

10095H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	28.69	6.60	^a APO-Dutasteride [TX]
			..	35.69	6.60	^a Avodart [GK]

■ FINASTERIDE

Authority required

Benign prostatic hyperplasia

Clinical criteria:

- Patient must have lower urinary tract symptoms.

finasteride 5 mg tablet, 30

4233T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	76.81	6.60	^a Auro-Finasteride [DO]	^a Finasteride AN [EA]
						^a Finasteride GH 5 [GQ]	^a Finide [AL]

..	94.66	6.60	^a Finnacar [RW]	
..	99.01	6.60	^a APO-Finasteride [TX]	^a Finasta [SZ]
			^a Finasteride-GA 5 [GN]	^a Pharmacor Finasteride 5 [CR]
			^a Proscar [OQ]	

finasteride 5 mg tablet, 28

4303L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	89.15	6.60	Finpro [RZ]

■ **ANTIINFECTIVES FOR SYSTEMIC USE**

■ **ANTIBACTERIALS FOR SYSTEMIC USE**

MACROLIDES, LINCOSAMIDES AND STREPTOGRAMINS

Macrolides

■ **AZITHROMYCIN**

Restricted benefit

Upper and lower respiratory tract infections

azithromycin 500 mg tablet, 3

4115N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	33.84	6.60	Zedd 500 [RW]	
						^a APO-Azithromycin [TX]	^a Azithromycin-GA [EA]
						^a Azithromycin Mylan [AF]	^a Azithromycin Sandoz [SZ]
						^a ZITHRO [RF]	^a Zithromax [PF]

■ **ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS**

■ **ANTINEOPLASTIC AGENTS**

ANTIMETABOLITES

Pyrimidine analogues

■ **FLUOROURACIL**

fluorouracil 5% cream, 20 g

4222F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	±1	..	..	66.21	6.60	^a APOC-5FU [TX]	^a Efudix [IL]

■ **IMMUNOSUPPRESSANTS**

IMMUNOSUPPRESSANTS

Tumor necrosis factor alpha (TNF-alpha) inhibitors

■ **INFLIXIMAB**

Note Any queries concerning the arrangements to prescribe infliximab may be directed to the Veterans' Affairs Pharmaceutical Advisory Centre (VAPAC) on 1800 552 580.

Written applications for authority to prescribe infliximab should be forwarded to:

Reply Paid 9998

Veterans' Affairs Pharmaceutical Advisory Centre (VAPAC)

Department of Veterans' Affairs

GPO Box 9998

BRISBANE QLD 4001

Authority required

Initial treatment, in combination with methotrexate, of specific accepted war-caused or service-related disability of refractory rheumatoid arthritis. Initial treatment may be prescribed by rheumatologists or consultant physicians for the reduction of signs and symptoms and prevention of structural joint damage in adult patients with active rheumatoid arthritis who satisfy all of the following criteria:

(1) (a) Proven raised erythrocyte sedimentation rate (ESR) and/or C-reactive protein (CRP); and

(1) (b) Proven erosive rheumatoid arthritis without end-stage disease;

(2) Failure of an adequate trial of methotrexate and 2 other disease modifying anti-rheumatic drugs (such as sulfasalazine, hydroxychloroquine, leflunomide or cyclosporin) — unless these drugs were contraindicated or intolerance had developed;

(3) No history of active tuberculosis requiring treatment in the last 3 years;

(4) No history of opportunistic infection in the last 2 months;

(5) Female patients of child-bearing age are not pregnant, not breast-feeding, and are using an effective form of contraception.

Applications for authorisation must be in writing and must include:

(1) a completed authority prescription form; and

(2) a completed Infliximab (Remicade) RPBS Authority Application - Supporting Information form (contact the VAPAC on 1800 552 580 for a copy of the form)

Authority required

Continuing treatment, in combination with methotrexate, of specific accepted war-caused or service-related disability of refractory rheumatoid arthritis. Continuing treatment may be prescribed by rheumatologists or consultant physicians, following initial therapy of 3 doses, in patients who satisfy the following criteria:

- (1) There is improvement in ESR and/or CRP; and
 - (2) An ACR20 (American College of Rheumatology) response is achieved by 14 weeks after the commencement of therapy.
- Applications for authorisation must be in writing and must include:

- (1) a completed authority prescription form; and
- (2) a completed Infliximab (Remicade) RPBS Authority Application - Supporting Information form (contact the VAPAC on 1800 552 580 for a copy of the form)

infliximab 100 mg injection, 1 vial

4284L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	355.14	6.60	Remicade [JC]

MUSCULO-SKELETAL SYSTEM

TOPICAL PRODUCTS FOR JOINT AND MUSCULAR PAIN

TOPICAL PRODUCTS FOR JOINT AND MUSCULAR PAIN

Preparations with salicylic acid derivatives

METHYL SALICYLATE + MENTHOL + CAMPHOR + EUCALYPTUS OIL + PINE OIL PUMILIO + TURPENTINE OIL + PEPPERMINT OIL + CAJUPUT OIL + CAPSICUM EXTRACT

methyl salicylate 20% + menthol 5% + camphor 3.5% + eucalyptus oil 3% + pine oil pumilio 1% + turpentine oil 1% + peppermint oil 0.5% + cajuput oil 0.5% + capsicum extract 0.15% cream, 100 g

11707E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	1	..	18.63	6.60	Goanna Heat Cream [GQ]

DRUGS FOR TREATMENT OF BONE DISEASES

DRUGS AFFECTING BONE STRUCTURE AND MINERALIZATION

Bisphosphonates

RISEDRONATE

Authority required

Preservation of bone mineral density

Clinical criteria:

- Patient must be on long-term glucocorticoid therapy, **AND**
- Patient must be undergoing continuous treatment with a dose equal to or greater than 7.5 mg of prednisone or equivalent per day, **AND**
- Patient must be osteopenic (bone mineral density t-score of less than -1.0).

Prescribers need to demonstrate that the patient has been on continuous therapy for 3 months or more.

risedronate sodium 35 mg enteric tablet, 4

2191H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	36.03	6.60	Actonel EC [TT]

risedronate sodium 35 mg tablet, 4

4444X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	36.03	6.60	^a Acris Once-a-Week [AF] ^a Risedronate AN [EA] ^a Risedronate Sandoz [SZ]	^a APO-Risedronate [TX] ^a Risedronate-GA [GN] ^a Risedro once a week [RW]

risedronate sodium 5 mg tablet, 28

4443W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	5	..	36.03	6.60	Actonel [TT]

Bisphosphonates, combinations

ALENDRONATE + COLECALCIFEROL

Authority required

Preservation of bone mineral density

Clinical criteria:

- Patient must be on long-term glucocorticoid therapy, **AND**
- Patient must be undergoing continuous treatment with a dose equal to or greater than 7.5 mg of prednisone or equivalent per day, **AND**
- Patient must be osteopenic (bone mineral density t-score of less than -1.0).

Prescribers need to demonstrate that the patient has been on continuous therapy for 3 months or more.

alendronate 70 mg + colecalciferol 140 microgram (5600 units) tablet, 4

2224C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	18.87	6.60	^a ALENDRONATE PLUS D3 70mg/140ug APOTEX [GX]	^a Alendronate plus D3-DRLA [RZ]
			..	22.87	6.60	^a FonatPlus [AF]	
						^a Fosamax Plus 70 mg/140 mcg [MQ]	

alendronate 70 mg + colecalciferol 70 microgram (2800 units) tablet, 4

2194L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	18.87	6.60	^a ALENDRONATE PLUS D3 70mg/70ug APOTEX [GX]	^a Alendronate plus D3-DRLA [RZ]
						^a FonatPlus [AF]	
			..	22.87	6.60	^a Fosamax Plus [MQ]	

■ RISEDRONATE (&) CALCIUM CARBONATE**Authority required**

Preservation of bone mineral density

Clinical criteria:

- Patient must be on long-term glucocorticoid therapy, **AND**
 - Patient must be undergoing continuous treatment with a dose equal to or greater than 7.5 mg of prednisone or equivalent per day, **AND**
 - Patient must be osteopenic (bone mineral density t-score of less than -1.0).
- Prescribers need to demonstrate that the patient has been on continuous therapy for 3 months or more.

risedronate sodium 35 mg tablet [4] (&) calcium (as carbonate) 500 mg tablet [24], 28

4059P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	5	..	36.03	6.60	Acris Combi [AF]

■ NERVOUS SYSTEM**■ ANALGESICS****OPIOIDS***Natural opium alkaloids***■ MORPHINE****Caution** The risk of drug dependence is high.**Note** Authorities for increased maximum quantities and/or repeats will be granted only for:

- chronic severe disabling pain associated with proven malignant neoplasia; or
- chronic severe disabling pain where treatment has been initiated by a specialist with appropriate expertise in pain management.

Restricted benefit

Chronic severe disabling pain

Clinical criteria:

- The condition must be unresponsive to non-opioid analgesics.

morphine sulfate pentahydrate 200 mg modified release tablet, 28

4349X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	119.42	6.60	MS Contin [MF]

*Opioids in combination with non-opioid analgesics***■ ASPIRIN + CODEINE****aspirin 300 mg + codeine phosphate hemihydrate 8 mg dispersible tablet, 40**

4286N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	18.77	6.60	Aspalgin 40 [GO]

■ PARACETAMOL + CODEINE**paracetamol 500 mg + codeine phosphate hemihydrate 8 mg tablet, 40**

4275B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	2	..	15.88	6.60	Panamax Co. 40 [SW]

OTHER ANALGESICS AND ANTIPYRETICS*Anilides*

■ PARACETAMOL

paracetamol 500 mg tablet, 100

10582Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	1	..	13.69	6.60	^a APO-Paracetamol [TX] ^a Mendelev Paracetamol [HX] ^a Paracetamol (Sandoz) [SZ] ^a Parapane [AF]	^a Febridol [EA] ^a Panamax [SW] ^a Paralgin [OW] ^a PHARMACY CARE PARACETAMOL [SI]

paracetamol 240 mg/5 mL oral liquid, 200 mL

10599W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	±1	2	..	15.87	6.60	^a Panamax 240 Elixir [SW]	^a Trust for Kids Paracetamol 6 to 12 years [CR]

■ PARACETAMOL

Restricted benefit

Persistent pain

Clinical criteria:

- The condition must be associated with osteoarthritis.

paracetamol 665 mg modified release tablet, 96

10598T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	2	5	..	*19.48	6.60	^a Osteomol 665 Paracetamol [CR]	^a Pharmacy Action Paracetamol Osteo 665 [GQ]

■ PARACETAMOL

Restricted benefit

Chronic arthropathies

paracetamol 500 mg tablet, 100

10585D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	3	4	..	*17.04	6.60	^a APO-Paracetamol [TX] ^a Mendelev Paracetamol [HX] ^a Paracetamol (Sandoz) [SZ] ^a Parapane [AF]	^a Febridol [EA] ^a Panamax [SW] ^a Paralgin [OW] ^a PHARMACY CARE PARACETAMOL [SI]

Other analgesics and antipyretics

■ GABAPENTIN

Authority required

Refractory neuropathic pain

Clinical criteria:

- The condition must be unable to be controlled by other drugs.

gabapentin 800 mg tablet, 100

4595W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	50.99	6.60	^a APO-Gabapentin [TX] ^a Gabapentin APOTEX [TY] ^a Gantin [ED] ^a Pharmacor Gabapentin 800 [CR]	^a Gabapentin AN [EA] ^a Gabapentin Aspen 800 [RW] ^a Neurontin [UJ]

gabapentin 600 mg tablet, 100

4594T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	41.24	6.60	^a APO-Gabapentin [TX] ^a Gabapentin APOTEX [TY] ^a Neurontin [UJ]	^a Gabapentin AN [EA] ^a Gabapentin Aspen 600 [RW] ^a Pharmacor Gabapentin 600 [CR]

gabapentin 400 mg capsule, 100

4593R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	30.43	6.60	^a APO-Gabapentin [TX] ^a Gabapentin APOTEX [TY] ^a Gabapentin generichealth [HQ] ^a Neurontin [UJ]	^a Gabacor [CR] ^a Gabapentin Aspen 400 [RW] ^a Gantin [EA] ^a Nupentin 400 [AF]

gabapentin 300 mg capsule, 100

4592Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	25.52	6.60	^a APO-Gabapentin [TX] ^a Gabapentin APOTEX [TY] ^a Gabapentin generichealth [HQ] ^a Neurontin [UJ]	^a Gabacor [CR] ^a Gabapentin Aspen 300 [RW] ^a Gantin [EA] ^a Nupentin 300 [AF]

gabapentin 100 mg capsule, 100

4591P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	5	..	16.19	6.60	^a APO-Gabapentin [TX] ^a Gabapentin APOTEX [TY] ^a Neurontin [UJ]	^a Gabacor [CR] ^a Gabapentin Aspen 100 [RW] ^a Nupentin 100 [AF]

■ **PSYCHOLEPTICS****ANXIOLYTICS***Benzodiazepine derivatives*■ **BROMAZEPAM**

Note This drug should not be used as the first line of treatment.

Note Other PBS-listed benzodiazepines should have been adequately tried and found to be ineffective or inappropriate.

Note Authorities for increased quantities and/or repeats may be granted to patients with terminal disease, and other patients who have been shown to be dependent on this item by an unsuccessful attempt at gradual withdrawal.

Authority required

Terminal disease

Clinical criteria:

- The treatment must be for the short-term, **AND**
- Patient must be receiving palliative care.

Authority required

Refractory phobic or anxiety states

Clinical criteria:

- The treatment must be for the short-term.

bromazepam 6 mg tablet, 30

4151L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*37.82	6.60	Lexotan [PB]

bromazepam 3 mg tablet, 30

4150K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*32.08	6.60	Lexotan [PB]

Azaspirodecanedione derivatives■ **BUSPIRONE****Authority required**

Anxiety

Clinical criteria:

- The treatment must be for the short-term.

buspirone hydrochloride 5 mg tablet, 50

4144D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	39.47	6.60	Buspar [AS]

buspirone hydrochloride 10 mg tablet, 50

4145E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	55.94	6.60	Buspar [AS]

HYPNOTICS AND SEDATIVES*Benzodiazepine derivatives*■ **FLUNITRAZEPAM**

Note This drug should not be used as the first line of treatment.

Note Other PBS-listed benzodiazepines should have been adequately tried and found to be ineffective or inappropriate.

Note Authorities for increased quantities and/or repeats may be granted to patients with terminal disease, and other patients who have been shown to be dependent on this item by an unsuccessful attempt at gradual withdrawal.

Authority required

Terminal disease

Clinical criteria:

- The treatment must be for the short-term, **AND**
- Patient must be receiving palliative care.

Authority required

Refractory phobic or anxiety states

Clinical criteria:

- The treatment must be for the short-term.

ANTIPARASITIC PRODUCTS, INSECTICIDES AND REPELLENTS

flunitrazepam 1 mg tablet, 30

4216X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	21.88	6.60	Hypnodorm [AF]

Benzodiazepine related drugs

■ ZOPICLONE

Restricted benefit

Insomnia

Clinical criteria:

- The treatment must be for the short-term.

zopiclone 7.5 mg tablet, 30

4522B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	25.36	6.60	^a APO-Zopiclone [TX]	^a Imoclone [RW]
						^a Imrest [AF]	^a Pharmacor Zopiclone [CR]
						^a Zopiclone GH [GQ]	
			..	28.10	6.60	^a Imovane [SW]	

■ OTHER NERVOUS SYSTEM DRUGS

DRUGS USED IN ADDICTIVE DISORDERS

Drugs used in nicotine dependence

■ NICOTINE

Note Studies have shown that successful therapy with this drug is enhanced by patient participation in a support and counselling program.

Authority required

Nicotine dependence

Clinical criteria:

- Patient must have indicated they are ready to cease smoking, **AND**
- Patient must have entered a comprehensive support and counselling program.

nicotine 7 mg/24 hours patch, 7

4571N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*52.50	6.60	QuitX [AF]

nicotine 21 mg/24 hours patch, 7

4573Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	2	..	*58.62	6.60	QuitX [AF]
			..	*68.68	6.60	Nicabate CQ 21 [GC]

nicotine 14 mg/24 hours patch, 7

4572P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*55.66	6.60	QuitX [AF]
			..	*68.68	6.60	Nicabate CQ 14 [GC]

■ ANTIPARASITIC PRODUCTS, INSECTICIDES AND REPELLENTS

■ ANTHELMINTICS

ANTINEMATODAL AGENTS

Benzimidazole derivatives

■ MEBENDAZOLE

Note Pharmaceutical benefits that have the forms mebendazole 100 mg tablet and mebendazole 100 mg chewable tablet are equivalent for the purposes of substitution.

mebendazole 100 mg tablet, 6

4325P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	18.22	6.60	^a Pharmacy Action Worm Treatment [GQ]

mebendazole 100 mg chewable tablet, 6

12194T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	18.22	6.60	^a Trust Deworm [CR]

RESPIRATORY SYSTEM

NASAL PREPARATIONS

DECONGESTANTS AND OTHER NASAL PREPARATIONS FOR TOPICAL USE

Sympathomimetics, plain

OXYMETAZOLINE

oxymetazoline hydrochloride 0.05% nasal spray, 20 mL

11711J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	±1	..	..	21.01	6.60	^a Pharmacy Action Nasal Decongestant [GQ]	^a Trust Decongestant Nasal Spray [CR]

oxymetazoline hydrochloride 0.05% nasal spray, 15 mL

4378K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	21.35	6.60	Drixine [BN]

oxymetazoline hydrochloride 0.05% nasal spray, 18 mL

4379L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	21.01	6.60	Logicin Rapid Relief [AS]

Antiallergic agents, excl. corticosteroids

CROMOGLYCATE

sodium cromoglycate 2% nasal spray, 26 mL

4468E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	5	..	26.36	6.60	Rynacrom [SW]

Corticosteroids

BUDESONIDE

Restricted benefit

Severe intractable rhinitis

budesonide 64 microgram/actuation nasal spray, 120 actuations

4092J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	42.61	6.60	Budamax Aqueous [JT]

Other nasal preparations

IPRATROPIUM

Restricted benefit

Severe intractable rhinorrhoea

Clinical criteria:

- The condition must be associated with perennial rhinitis, **AND**
- The condition must be unresponsive to insufflated nasal steroids.

ipratropium bromide monohydrate 44 microgram/actuation nasal spray, 180 actuations

4090G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	5	..	33.56	6.60	Atrovent Nasal Forte [VZ]

ipratropium bromide monohydrate 22 microgram/actuation nasal spray, 180 actuations

4089F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	5	..	27.41	6.60	Atrovent Nasal Aqueous [VZ]

NASAL DECONGESTANTS FOR SYSTEMIC USE

Sympathomimetics

PSEUDOEPHEDRINE

pseudoephedrine hydrochloride 60 mg tablet, 12

4029C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	15.55	6.60	^a Pharmacy Action Sinus & Nasal Decongestant Relief [GQ]	^a Trust Sinus & Nasal Decongestant [CR]
			..	16.15	6.60	^a Logicin Sinus [AS]	

ANTI-HISTAMINES FOR SYSTEMIC USE

ANTI-HISTAMINES FOR SYSTEMIC USE

Piperazine derivatives

SENSORY ORGANS

■ CETIRIZINE

cetirizine hydrochloride 10 mg tablet, 30

4175R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	28.99	6.60	^a Pharmacy Action Cetrelief [GQ]	^a Trust Cetirizine [CR]
			..	32.22	6.60	^a Alzene [AF]	

Other antihistamines for systemic use

■ FEXOFENADINE

fexofenadine hydrochloride 60 mg tablet, 20

4237B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	3	..	..	*56.10	6.60	Telfast [SW]	

fexofenadine hydrochloride 120 mg tablet, 30

4238C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	28.81	6.60	^a Pharmacy Action Fexorelief 120 [GQ]	^a Trust Fexit 120 [CR]
			..	32.02	6.60	^a Xergic [AF]	
			..	48.23	6.60	^a Telfast 120 [SW]	

■ LORATADINE

loratadine 10 mg tablet, 30

4313B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	1	..	..	31.42	6.60	^a Pharmacy Action Lorastyne [GQ]	^a Trust Loratadine [CR]
						^a Trust Loratadine Antihistamine [RM]	
			..	35.12	6.60	^a Allereze [AF]	
			..	44.75	6.60	^a Lorano [SZ]	
			..	47.02	6.60	^a Claratyne [BN]	

■ SENSORY ORGANS

■ OTOLOGICALS

OTHER OTOLOGICALS

Indifferent preparations

■ CARBAMIDE PEROXIDE

carbamide peroxide 6.5% ear drops, 12 mL

4176T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	±1	..	..	21.50	6.60	Ear Clear for Ear Wax Removal [KY]	

■ DOCUSATE

docusate sodium 0.5% ear drops, 10 mL

4199B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	±1	..	..	19.02	6.60	Waxsol [GO]	

■ ORTHO-DICHLOROBENZENE + PARA-DICHLOROBENZENE + CHLOROBUTANOL + ARACHIS OIL

ortho-dichlorobenzene 14% + para-dichlorobenzene 2% + chlorobutanol hemihydrate 5% + arachis oil 57% ear drops, 10 mL

4180B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	±1	..	..	18.68	6.60	Cerumol [UN]	

■ VARIOUS

■ ALL OTHER THERAPEUTIC PRODUCTS

ALL OTHER THERAPEUTIC PRODUCTS

Drugs for treatment of hyperkalemia and hyperphosphatemia

■ SODIUM POLYSTYRENE SULFONATE

sodium polystyrene sulfonate 999.3 mg/g powder, 454 g

4470G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer	Brand Name and Manufacturer
	±1	2	..	70.84	6.60	Resonium-A [SW]	

GENERAL NUTRIENTS

OTHER NUTRIENTS

Other combinations of nutrients

PROTEIN FORMULA WITH ARGININE, VITAMIN C AND E

Restricted benefit

Stage 2 and above pressure injury

Clinical criteria:

- The treatment must be for special medical purposes to support healing of pressure injuries.

protein formula with arginine, vitamin C and E powder for oral liquid, 14 x 9.2 g sachets

10850C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	4	5	..	*161.90	6.60	Arginaid [NT]

PROTEIN FORMULA WITH ARGININE, VITAMIN C, E AND ZINC

Restricted benefit

Stage 2 and above pressure injury

Clinical criteria:

- The treatment must be for special medical purposes to support healing of pressure injuries.

protein formula with arginine, vitamin C, E and zinc oral liquid, 27 x 237 mL cartons

10841N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*269.16	6.60	Arginaid Extra [NT]

protein formula with arginine, vitamin C, E and zinc oral liquid, 24 x 200 mL bottles

11401C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	5	..	*205.90	6.60	Cubitan [SB]

ALL OTHER NON-THERAPEUTIC PRODUCTS

ALL OTHER NON-THERAPEUTIC PRODUCTS

LUBRICATING AGENT

lubricating agent gel, 100 g

4306P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	15.43	6.60	Lubri-Gel [PP]

Other non-therapeutic auxiliary products

BANDAGE ABSORBENT WOOL

bandage absorbent wool 10 cm x 3 m bandage, 1

4653X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	24.11	6.60	Surepress 650948 [CC]

BANDAGE CALICO

bandage calico large triangular bandage, 1

4717G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	18.03	6.60	Handy 36361414 [BV]

BANDAGE COMPRESSION

bandage compression 10 cm x 3.5 m soft bandage [1] (&) bandage compression 10 cm x 6 m short stretch bandage [1], 1 pack

11714M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	38.61	6.60	Rosidal TCS 26484 [LC]

BANDAGE COMPRESSION

Note Treatment of varices and oedema associated with venous disease and lymphoedema; contraindicated in arterial disease.

bandage compression 10 cm x 3 m high stretch bandage, 1

4748X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	5	..	..	*72.49	6.60	Surepress 650947 [CC]
			..	*152.69	6.60	Tensopress 71723-00 [BV]

bandage compression 8 cm x 2.6 m short stretch bandage, 1

4654Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	5	..	..	*77.84	6.60	Comprilan 01027-00 [BV]

■ **BANDAGE COMPRESSION**

Note Treatment of varices and oedema associated with venous disease and lymphoedema; contraindicated in arterial disease.

Note Smith & Nephew products are distributed via the three major wholesalers, API, Sigma & Symbion. To best ensure product availability at RPBS agreed prices, please order from one of these suppliers. In the event that your preferred wholesaler cannot supply, please contact Smith & Nephew Customer Service on 13 13 60. Smith & Nephew cannot ensure RPBS pricing from distributors other than those aforementioned.

bandage compression four layer bandage, 1

4598B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	5	..	..	*186.09	6.60	Profore Lite 66050415 [SN]

bandage compression four layer bandage, 1

4658E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	5	..	..	*278.14	6.60	Profore 66050016 [SN]

■ **BANDAGE COMPRESSION**

Note Treatment of varices and oedema associated with venous disease and lymphoedema; contraindicated in arterial disease.

Note Molnlycke Health Care products are distributed through leading pharmacy distributors. To best ensure product availability at the RPBS agreed prices, special arrangements have been made with API and Independence Australia Health Solutions (IAHS). IAHS orders can be placed on: Tel: 1300 788 855; or Email: customerservice@independenceaustralia.com. Molnlycke Health Care is not able to ensure product availability or pricing on listed products beyond these two suppliers.

bandage compression 10 cm x 3.5 m high stretch bandage, 1

4657D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	5	..	..	*77.64	6.60	Setopress 3505 [MH]

■ **BANDAGE COMPRESSION**

Note Treatment of varices and oedema associated with venous disease and lymphoedema; contraindicated in arterial disease.

Note Bandage can be left in situ for up to 7 days as per manufacturer's instructions.

Restricted benefit

Venous ulcer

Treatment Phase: Initial treatment

Restricted benefit

Venous ulcer

Treatment Phase: Continuing treatment

bandage compression two layer bandage, 1

4050E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	43.78	6.60	Coban 2 [MM]

■ **BANDAGE RETENTION COHESIVE HEAVY****bandage retention cohesive heavy 15 cm x 1.3 m bandage, 1**

4814J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*30.94	6.60	Peg 7425 [MM]

bandage retention cohesive heavy 10 cm x 1.3 m bandage, 1

4813H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*24.76	6.60	Peg 7423 [MM]

bandage retention cohesive heavy 5 cm x 1.3 m bandage, 1

4811F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*18.62	6.60	Peg 7420 [MM]

bandage retention cohesive heavy 7.5 cm x 1.3 m bandage, 1

4812G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*21.48	6.60	Peg 7422 [MM]

bandage retention cohesive heavy 10 cm x 2 m bandage, 1

4660G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*23.28	6.60	Coban 1584 [MM]

■ **BANDAGE RETENTION COHESIVE LIGHT****bandage retention cohesive light 2.5 cm x 2 m bandage, 2**

4718H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	18.16	6.60	Handygauze Cohesive 8631 [BV]

bandage retention cohesive light 6 cm x 2 m bandage, 1

4719J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*20.54	6.60	Handygauze Cohesive 8633 [BV]

bandage retention cohesive light 10 cm x 2 m bandage, 1

4662J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*34.40	6.60	Handygauze Cohesive 8635 [BV]

▪ BANDAGE RETENTION COTTON CREPE**bandage retention cotton crepe 7.5 cm x 2.3 m bandage, 1**

4728W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*25.76	6.60	Telfa 8253F [KE]
			..	*28.80	6.60	Tensocrepe 36307501 [BV]

bandage retention cotton crepe 10 cm x 2.3 m bandage, 1

4729X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*28.50	6.60	Telfa 8254F [KE]
			..	*33.94	6.60	Tensocrepe 36301001 [BV]

bandage retention cotton crepe 5 cm x 2.3 m bandage, 1

4727T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*21.58	6.60	Telfa 8252F [KE]
			..	*24.24	6.60	Tensocrepe 36300501 [BV]

▪ BANDAGE TUBULAR**bandage tubular size D (25 cm to 43 cm) straight bandage, 1**

4664L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	19.82	6.60	Elastoplast 2226 [BE]

bandage tubular size C (15 cm to 25 cm) straight bandage, 1

4663K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	19.82	6.60	Elastoplast 2225 [BE]

bandage tubular size E (35 cm to 45 cm) straight bandage, 1

4665M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	19.82	6.60	Elastoplast 2227 [BE]

▪ BANDAGE TUBULAR

Note Molnlycke Health Care products are distributed through leading pharmacy distributors. To best ensure product availability at the RPBS agreed prices, special arrangements have been made with API and Independence Australia Health Solutions (IAHS). IAHS orders can be placed on: Tel: 1300 788 855; or Email: customerservice@independenceaustralia.com. Molnlycke Health Care is not able to ensure product availability or pricing on listed products beyond these two suppliers.

bandage tubular 8.75 cm x 1 m bandage, 1

4858Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	19.46	6.60	Tubigrip E 1547 [MH]

bandage tubular 6.25 cm x 1 m bandage, 1

4855M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	19.46	6.60	Tubigrip B 1520 [MH]

bandage tubular 6.75 cm x 1 m bandage, 1

4856N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	19.46	6.60	Tubigrip C 1545 [MH]

bandage tubular 7.5 cm x 1 m bandage, 1

4857P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	19.46	6.60	Tubigrip D 1546 [MH]

bandage tubular 10 cm x 1 m bandage, 1

4859R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	19.46	6.60	Tubigrip F 1548 [MH]

■ BANDAGE TUBULAR FINGER

BANDAGE-TUBULAR (FINGER) Complete pack including applicator, 1

4798M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	21.89	6.60	Tubegauz 0501633 [SS]

■ BANDAGE TUBULAR LIGHT WEIGHT

Note Molnlycke Health Care products are distributed through leading pharmacy distributors. To best ensure product availability at the RPBS agreed prices, special arrangements have been made with API and Independence Australia Health Solutions (IAHS). IAHS orders can be placed on: Tel: 1300 788 855; or Email: customerservice@independenceaustralia.com.

Molnlycke Health Care is not able to ensure product availability or pricing on listed products beyond these two suppliers.

bandage tubular light weight 10 m large limb size bandage, 1

4673Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	30.90	6.60	Tubifast 2438 [MH]

bandage tubular light weight 10 m small limb size bandage, 1

4671W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	26.23	6.60	Tubifast 2434 [MH]

bandage tubular light weight 10 m medium limb size bandage, 1

4672X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	29.67	6.60	Tubifast 2436 [MH]

■ BANDAGE TUBULAR LONG STOCKING

Note Molnlycke Health Care products are distributed through leading pharmacy distributors. To best ensure product availability at the RPBS agreed prices, special arrangements have been made with API and Independence Australia Health Solutions (IAHS). IAHS orders can be placed on: Tel: 1300 788 855; or Email: customerservice@independenceaustralia.com.

Molnlycke Health Care is not able to ensure product availability or pricing on listed products beyond these two suppliers.

bandage tubular long stocking small size bandage, 1

4674B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*38.72	6.60	Tubigrip 1482 [MH]

bandage tubular long stocking large size bandage, 1

4799N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*38.72	6.60	Tubigrip 1484 [MH]

bandage tubular long stocking medium size bandage, 1

4797L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*38.72	6.60	Tubigrip 1483 [MH]

bandage tubular long stocking XX/large size bandage, 1

4675C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*38.74	6.60	Tubigrip 1486 [MH]

■ BANDAGE TUBULAR SHORT STOCKING

Note Molnlycke Health Care products are distributed through leading pharmacy distributors. To best ensure product availability at the RPBS agreed prices, special arrangements have been made with API and Independence Australia Health Solutions (IAHS). IAHS orders can be placed on: Tel: 1300 788 855; or Email: customerservice@independenceaustralia.com.

Molnlycke Health Care is not able to ensure product availability or pricing on listed products beyond these two suppliers.

bandage tubular short stocking large D/E size bandage, 1

4816L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*28.50	6.60	Tubigrip 1481 [MH]

bandage tubular short stocking medium C/D size bandage, 1

4815K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*28.50	6.60	Tubigrip 1480 [MH]

bandage tubular short stocking small B/C size bandage, 1

4661H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*28.50	6.60	Tubigrip 1479 [MH]

■ BANDAGE ZINC PASTE

Note Used as an adjunct in the management of leg ulceration and associated eczema and skin conditions.

Note Molnlycke Health Care products are distributed through leading pharmacy distributors. To best ensure product availability at the RPBS agreed prices, special arrangements have been made with API and Independence Australia Health Solutions

(IAHS). IAHS orders can be placed on: Tel: 1300 788 855; or Email: customerservice@independenceaustralia.com.
Molnlycke Health Care is not able to ensure product availability or pricing on listed products beyond these two suppliers.

bandage zinc paste 7.5 cm x 6 m bandage, 1

4669R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	3	..	*32.22	6.60	Steripaste 3610 [MH]

▪ BANDAGE ZINC PASTE

Note Used as an adjunct in the management of leg ulceration and associated eczema and skin conditions.

Note Smith & Nephew products are distributed via the three major wholesalers, API, Sigma & Symbion. To best ensure product availability at RPBS agreed prices, please order from one of these suppliers. In the event that your preferred wholesaler cannot supply, please contact Smith & Nephew Customer Service on 13 13 60. Smith & Nephew cannot ensure RPBS pricing from distributors other than those aforementioned.

bandage zinc paste 7.5 cm x 6 m bandage, 1

4750B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	3	..	*92.42	6.60	Viscopaste 4948 [SN]

bandage zinc paste 80 cm (stockings) bandage, 4

4760M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	3	..	105.90	6.60	ZipZoc 66000747 [SN]

▪ BETAINE + POLYAMINOPROPYL BIGUANIDE

betaine 0.1% + polyaminopropyl biguanide 0.1% solution, 6 x 40 mL ampoules

2525X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	29.63	6.60	Prontosan Wound Irrigation Solution [BR]

▪ CADEXOMER-IODINE

Note Suitable for yellow sloughy infected and malodorous wounds.

Note Smith & Nephew products are distributed via the three major wholesalers, API, Sigma and Symbion. To best ensure product availability at RPBS agreed prices, please order from one of these suppliers. In the event that your preferred wholesaler cannot supply, please contact Smith & Nephew Customer Service on 13 13 60. Smith & Nephew cannot ensure RPBS agreed pricing from distributors other than those aforementioned.

cadexomer-iodine 8 cm x 6 cm dressing, 3 x 10 g sheet

4936T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	2	..	180.90	6.60	Iodosorb 66051340 [SN]

cadexomer-iodine 50% ointment, 2 x 20 g

4933P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	2	..	130.92	6.60	Iodosorb Ointment 66051230 [SN]

cadexomer-iodine 6 cm x 4 cm dressing, 5 x 5 g sheet

4935R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	2	..	124.85	6.60	Iodosorb 66051330 [SN]

cadexomer-iodine 50% ointment, 4 x 10 g

4932N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	2	..	132.18	6.60	Iodosorb Ointment 66051240 [SN]

cadexomer-iodine 3 g powder, 7 sachets

4931M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	2	..	83.12	6.60	Iodosorb Powder 66051070 [SN]

cadexomer-iodine 10 cm x 8 cm dressing, 2 x 17 g sheet

4937W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	190.68	6.60	Iodosorb 66051360 [SN]

▪ DRESSING ACTIVATED CHARCOAL MALODOROUS WOUND

dressing activated charcoal malodorous wound 10 cm x 10 cm dressing, 10

4742N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	77.98	6.60	CarboFLEX 403202 [CC]

dressings activated charcoal malodorous wound 15 cm x 20 cm dressing, 5

4743P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	‡1	..	..	87.88	6.60	CarboFLEX 403204 [CC]

dressings activated charcoal malodorous wound 10.5 cm x 10.5 cm dressing, 1

4681J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	10	..	..	*97.94	6.60	Actisorb Plus MAP105 [KI]

■ DRESSING ALGINATE CAVITY WOUND

Note This dressing should be used only on moderately to heavily exuding wounds and should remain in place until saturated or for a maximum of 3 days.

dressings alginate cavity wound 2 g rope, 5 x 2 g

1905G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*110.96	6.60	Kaltostat 168117 [CC]

dressings alginate cavity wound 2 g rope, 1

4832H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	10	..	..	*105.44	6.60	Sorbsan 1411 [UM]

■ DRESSING ALGINATE CAVITY WOUND

Note This dressing should be used only on moderately to heavily exuding wounds and should remain in place until saturated or for a maximum of 3 days.

Note Coloplast dressings are available via a range of distributors. However, Coloplast's principal agreement to ensure correct RPBS Price to Pharmacy and ready supply has been secured with Independence Australia on 1300 788 855 and BrightSky on 1300 290 400. Please note that Coloplast is unable to guarantee ready supply or rebate for price differences on purchases outside these distributors.

dressings alginate cavity wound 2 g (40 cm) rope, 6 x 2 g

4682K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*132.34	6.60	Comfeel SeaSorb Filler 3740 [CT]

■ DRESSING ALGINATE SUPERFICIAL WOUND

Note This dressing should be used only on moderately to heavily exuding wounds and should remain in place until saturated or for a maximum of 3 days.

dressings alginate superficial wound 7.5 cm x 12 cm dressing, 10

4683L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	‡1	1	..	88.98	6.60	Kaltostat 168212 [CC]

■ DRESSING ALGINATE SUPERFICIAL WOUND

Note This dressing should be used only on moderately to heavily exuding wounds and should remain in place until saturated or for a maximum of 3 days.

Note Smith & Nephew products are distributed via the three major wholesalers, API, Sigma and Symbion. To best ensure product availability at RPBS agreed prices, please order from one of these suppliers. In the event that your preferred wholesaler cannot supply, please contact Smith & Nephew Customer Service on 13 13 60. Smith & Nephew cannot ensure RPBS agreed pricing from distributors other than those aforementioned.

dressings alginate superficial wound 5 cm x 5 cm dressing, 10

4699H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	‡1	1	..	50.50	6.60	Kaltostat 168210 [CC]
			..	64.94	6.60	Algisite M 66000519 [SN]

dressings alginate superficial wound 15 cm x 20 cm dressing, 10

4691X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	‡1	1	..	298.40	6.60	Algisite M 66000521 [SN]

dressings alginate superficial wound 10 cm x 10 cm dressing, 10

4700J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	‡1	1	..	122.16	6.60	Algisite M 66000520 [SN]

■ DRESSING ALGINATE SUPERFICIAL WOUND

Note This dressing should be used only on moderately to heavily exuding wounds and should remain in place until saturated or for a maximum of 3 days.

Note Coloplast dressings are available via a range of distributors. However, Coloplast's principal agreement to ensure correct RPBS Price to Pharmacy and ready supply has been secured with Independence Australia on 1300 788 855 and BrightSky on 1300 290 400. Please note that Coloplast is unable to guarantee ready supply or rebate for price differences on purchases outside these distributors.

dressings alginate superficial wound 5 cm x 5 cm dressing, 1

4684M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	10	1	..	*48.04	6.60	Comfeel SeaSorb Dressing 3705 [CT]

dressings alginate superficial wound 10 cm x 10 cm dressing, 1

4831G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	10	1	..	*82.94	6.60	Sorbsan 1410 [UM]
			..	*88.14	6.60	Comfeel SeaSorb Dressing 3710 [CT]

■ DRESSING ALGINATE WITH MANUKA HONEY

Note Suitable for yellow sloughy infected and malodorous wounds.

dressings alginate with manuka honey 10 cm x 10 cm dressing, 5

10849B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	4	..	64.49	6.60	Algivon Plus CR4225 [DJ]

dressings alginate with manuka honey 2.5 cm x 20 cm ribbon, 5

10857K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	4	..	115.97	6.60	Algivon Plus Ribbon & Probe CR4231 [DJ]

■ DRESSING FILM**dressings film 15 cm x 20 cm dressing, 1**

4688R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	6	..	..	*33.06	6.60	Tegaderm Transparent 1628 [MM]

dressings film 10 cm x 12 cm dressing, 4

4687Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	23.52	6.60	Nexcare Tegaderm Transparent H1626 [MM]

dressings film 6 cm x 7 cm dressing, 8

4686P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	20.04	6.60	Nexcare Tegaderm Transparent H1624 [MM]

■ DRESSING FILM

Note Smith & Nephew products are distributed via the three major wholesalers, API, Sigma and Symbion. To best ensure product availability at RPBS agreed prices, please order from one of these suppliers. In the event that your preferred wholesaler cannot supply, please contact Smith & Nephew Customer Service on 13 13 60. Smith & Nephew cannot ensure RPBS agreed pricing from distributors other than those aforementioned.

dressings film 10 cm x 12 cm dressing, 10

4893M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	41.35	6.60	Op-Site Flexigrid 4629 [SN]

■ DRESSING FILM ISLAND**dressings film island 9 cm x 10 cm dressing, 1**

4690W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	10	..	..	*30.54	6.60	Tegaderm Transparent Island 3586 [MM]

dressings film island 5 cm x 7 cm dressing, 1

4689T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	10	..	..	*20.54	6.60	Tegaderm Transparent Island 3582 [MM]

■ DRESSING FILM ISLAND

Note Smith & Nephew products are distributed via the three major wholesalers, API, Sigma and Symbion. To best ensure product availability at RPBS agreed prices, please order from one of these suppliers. In the event that your preferred wholesaler cannot supply, please contact Smith & Nephew Customer Service on 13 13 60. Smith & Nephew cannot ensure RPBS agreed pricing from distributors other than those aforementioned.

dressings film island 8 cm x 10 cm dressing, 5

4899W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*50.52	6.60	Cutifilm Plus 36361371 [SN]

dressing film island 5 cm x 7.2 cm dressing, 5

4898T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*33.80	6.60	Cutifilm Plus 36361370 [SN]

▪ DRESSING FOAM HEAVY EXUDATE

Note This dressing should remain in place until saturated or up to a maximum of 7 days. Allow a minimum of 2 cm to 3 cm in excess of the wound size of the dressing around the wound.

Note Smith & Nephew products are distributed via the three major wholesalers, API, Sigma and Symbion. To best ensure product availability at RPBS agreed prices, please order from one of these suppliers. In the event that your preferred wholesaler cannot supply, please contact Smith & Nephew Customer Service on 13 13 60. Smith & Nephew cannot ensure RPBS agreed pricing from distributors other than those aforementioned.

dressing foam heavy exudate 10 cm x 10 cm dressing, 10

4795J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	1	..	154.34	6.60	Allevyn 66007637 [SN]

▪ DRESSING FOAM MODERATE EXUDATE

Note Smith & Nephew products are distributed via the three major wholesalers, API, Sigma and Symbion. To best ensure product availability at RPBS agreed prices, please order from one of these suppliers. In the event that your preferred wholesaler cannot supply, please contact Smith & Nephew Customer Service on 13 13 60. Smith & Nephew cannot ensure RPBS agreed pricing from distributors other than those aforementioned.

dressing foam moderate exudate cavity conforming foam, 20 g sachet

4694C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	1	..	99.85	6.60	Cavicare 4563 [SN]

▪ DRESSING FOAM MODERATE EXUDATE

Note This dressing should remain in place until saturated or up to a maximum of 7 days. Allow a minimum of 2 cm to 3 cm in excess of the wound size of the dressing around the wound.

Note Smith & Nephew products are distributed via the three major wholesalers, API, Sigma and Symbion. To best ensure product availability at RPBS agreed prices, please order from one of these suppliers. In the event that your preferred wholesaler cannot supply, please contact Smith & Nephew Customer Service on 13 13 60. Smith & Nephew cannot ensure RPBS agreed pricing from distributors other than those aforementioned.

dressing foam moderate exudate 12.5 cm x 12.5 cm dressing, 10

4590N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	153.22	6.60	Allevyn Adhesive 66000044 [SN]

▪ DRESSING FOAM WITH SILICONE

Note Smith & Nephew products are distributed via the three major wholesalers, API, Sigma and Symbion. To best ensure product availability at RPBS agreed prices, please order from one of these suppliers. In the event that your preferred wholesaler cannot supply, please contact Smith & Nephew Customer Service on 13 13 60. Smith & Nephew cannot ensure RPBS agreed pricing from distributors other than those aforementioned.

dressing foam with silicone 12.9 cm x 12.9 cm dressing, 10

10029W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	91.36	6.60	Allevyn Life 66801068 [SN]

dressing foam with silicone 10.3 cm x 10.3 cm dressing, 10

10017F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	64.90	6.60	Allevyn Life 66801067 [SN]

dressing foam with silicone 16 cm x 16 cm dressing, 10

11393P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	110.77	6.60	Allevyn Life Non-Bordered 66801749 [SN]

dressing foam with silicone 21 cm x 21 cm dressing, 10

10021K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	253.55	6.60	Allevyn Life 66801070 [SN]

dressing foam with silicone 15.4 cm x 15.4 cm dressing, 10

10023M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	125.74	6.60	Allevyn Life 66801069 [SN]

dressing foam with silicone 10.5 cm x 10.5 cm dressing, 10

11384E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	64.90	6.60	Allevyn Life Non-Bordered 66801748 [SN]

▪ DRESSING FOAM WITH SILICONE AND SILVER

Note Molnlycke Health Care products are distributed through leading pharmacy distributors. To best ensure product availability at the RPBS agreed prices, special arrangements have been made with API and Independence Australia Health Solutions (IAHS). IAHS orders can be placed on: Tel: 1300 788 855; or Email: customerservice@independenceaustralia.com. Molnlycke Health Care is not able to ensure product availability or pricing on listed products beyond these two suppliers.

Authority required

Wounds

Clinical criteria:

- Patient must have a wound where there is evidence of critical colonisation; OR
- Patient must have a well-assessed chronic wound that has not responded to conventional dressings.

dressing foam with silicone and silver 10 cm x 10 cm dressing, 5

2439J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	105.55	6.60	Mepilex Ag [MH]

dressing foam with silicone and silver 10 cm x 10 cm dressing, 5

2470B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	112.63	6.60	Mepilex Border Ag [MH]

▪ DRESSING FOAM WITH SILICONE HEAVY EXUDATE

dressing foam with silicone heavy exudate 22 cm x 23 cm dressing, 6

12195W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	167.55	6.60	Mepilex Border Heel 282750 [MH]

dressing foam with silicone heavy exudate 16 cm x 20 cm dressing, 5

12216Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	84.11	6.60	Mepilex Border Sacrum 282050 [MH]

dressing foam with silicone heavy exudate 7.5 cm x 7.5 cm dressing, 10

12184G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	52.99	6.60	Mepilex Border Flex 595211 [MH]

dressing foam with silicone heavy exudate 22 cm x 25 cm dressing, 5

12207L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	127.82	6.60	Mepilex Border Sacrum 282450 [MH]

dressing foam with silicone heavy exudate 10 cm x 10 cm dressing, 10

12206K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	71.05	6.60	Mepilex Border Flex 595311 [MH]

dressing foam with silicone heavy exudate 15 cm x 20 cm dressing, 10

12185H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	177.95	6.60	Mepilex Border Flex 595611 [MH]

▪ DRESSING FOAM WITH SILICONE HEAVY EXUDATE

Note Smith & Nephew products are distributed via the three major wholesalers, API, Sigma and Symbion. To best ensure product availability at RPBS agreed prices, please order from one of these suppliers. In the event that your preferred wholesaler cannot supply, please contact Smith & Nephew Customer Service on 13 13 60. Smith & Nephew cannot ensure RPBS agreed pricing from distributors other than those aforementioned.

dressing foam with silicone heavy exudate 7.5 cm x 7.5 cm dressing, 10

4207K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	61.45	6.60	Allevyn Gentle Border 66800269 [SN]

dressing foam with silicone heavy exudate 10 cm x 10 cm dressing, 10

4196W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	86.16	6.60	Allevyn Gentle 66800248 [SN]

dressing foam with silicone heavy exudate 10 cm x 10 cm dressing, 10

4230P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	86.16	6.60	Allevyn Gentle Border 66800270 [SN]

■ DRESSING FOAM WITH SILICONE HEAVY EXUDATE

Note Molnlycke Healthcare products are distributed through leading pharmacy distributors. To best ensure product availability at RPBS agreed prices, special arrangements have been made with API and Independence Australia Health Solutions (IAHS). IAHS orders can be placed on: Tel: 1300 788 855; or Email customerservice@independenceaustralia.com. Molnlycke Healthcare are not able to ensure product availability or pricing on listed products beyond these two suppliers.

■ dressing foam with silicone heavy exudate 10 cm x 10 cm dressing, 5

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
4643J	±1	..	..	43.78	6.60	Mepilex Border 295300 [MH]

■ dressing foam with silicone heavy exudate 7.5 cm x 7.5 cm dressing, 5

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
4642H	±1	..	..	33.19	6.60	Mepilex Border 295200 [MH]

■ DRESSING FOAM WITH SILICONE LIGHT EXUDATE

Note Molnlycke Healthcare products are distributed through leading pharmacy distributors. To best ensure product availability at RPBS agreed prices, special arrangements have been made with API and Independence Australia Health Solutions (IAHS). IAHS orders can be placed on: Tel: 1300 788 855; or Email customerservice@independenceaustralia.com. Molnlycke Healthcare are not able to ensure product availability or pricing on listed products beyond these two suppliers.

■ dressing foam with silicone light exudate 10 cm x 10 cm dressing, 5

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
4645L	±1	..	..	39.66	6.60	Mepilex Lite 284100 [MH]

■ dressing foam with silicone light exudate 6 cm x 8.5 cm dressing, 5

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
4644K	±1	..	..	30.84	6.60	Mepilex Lite 284000 [MH]

■ DRESSING FOAM WITH SILICONE MODERATE EXUDATE

Note Molnlycke Healthcare products are distributed through leading pharmacy distributors. To best ensure product availability at RPBS agreed prices, special arrangements have been made with API and Independence Australia Health Solutions (IAHS). IAHS orders can be placed on: Tel: 1300 788 855; or Email customerservice@independenceaustralia.com. Molnlycke Healthcare are not able to ensure product availability or pricing on listed products beyond these two suppliers.

■ dressing foam with silicone moderate exudate 10 cm x 10 cm dressing, 5

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
4626L	±1	..	..	43.78	6.60	Mepilex 294100 [MH]

■ DRESSING FOAM WITH SILVER

Note Smith & Nephew products are distributed via the three major wholesalers, API, Sigma and Symbion. To best ensure product availability at RPBS agreed prices, please order from one of these suppliers. In the event that your preferred wholesaler cannot supply, please contact Smith & Nephew Customer Service on 13 13 60. Smith & Nephew cannot ensure RPBS agreed pricing from distributors other than those aforementioned.

■ Authority required

Wounds

■ Clinical criteria:

- Patient must have a wound where there is evidence of critical colonisation; OR
- Patient must have a well-assessed chronic wound that has not responded to conventional dressings.

■ dressing foam with silver 7.5 cm x 7.5 cm dressing, 10

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
4252T	±1	..	..	157.66	6.60	Allevyn Ag Adhesive 66800073 [SN]

■ dressing foam with silver 7.5 cm x 7.5 cm dressing, 10

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
4263J	±1	..	..	157.66	6.60	Allevyn Ag Gentle Border 66800460 [SN]

■ dressing foam with silver 10 cm x 10 cm dressing, 10

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
4255Y	±1	..	..	234.67	6.60	Allevyn Ag Adhesive 66800075 [SN]

■ dressing foam with silver 10 cm x 10 cm dressing, 10

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
4259E	±1	..	..	239.19	6.60	Allevyn Ag Non-Adhesive 66800086 [SN]

dressings foam with silver 10 cm x 10 cm dressing, 10

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
4266M	‡1	..	..	234.67	6.60	Allevyn Ag Gentle Border 66800461 [SN]

dressings foam with silver 12.5 cm x 12.5 cm dressing, 10

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
4258D	‡1	..	..	291.61	6.60	Allevyn Ag Adhesive 66800078 [SN]

dressings foam with silver 12.5 cm x 12.5 cm dressing, 10

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
4270R	‡1	..	..	291.61	6.60	Allevyn Ag Gentle Border 66800462 [SN]

■ DRESSING GAUZE**dressings gauze eye pad, 12 pads**

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
4768Y	‡1	..	..	17.61	6.60	Curity 4112 [KE]

■ DRESSING GAUZE ABSORBENT**dressings gauze absorbent 5 cm x 5 cm pad, 100**

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
4707R	‡1	..	..	19.62	6.60	Handy 71117-05 [BV]

dressings gauze absorbent 10 cm x 10 cm pad, 100

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
4708T	‡1	..	..	33.35	6.60	Handy 71117-06 [BV]

■ DRESSING GAUZE PARAFFIN

Note Smith & Nephew products are distributed via the three major wholesalers, API, Sigma and Symbion. To best ensure product availability at RPBS agreed prices, please order from one of these suppliers. In the event that your preferred wholesaler cannot supply, please contact Smith & Nephew Customer Service on 13 13 60. Smith & Nephew cannot ensure RPBS agreed pricing from distributors other than those aforementioned.

dressings gauze paraffin 10 cm x 10 cm dressing, 10

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
4759L	‡1	..	..	27.43	6.60	Jelonet 7404 [SN]

■ DRESSING GAUZE PARAFFIN WITH CHLORHEXIDINE ACETATE

Note Smith & Nephew products are distributed via the three major wholesalers, API, Sigma and Symbion. To best ensure product availability at RPBS agreed prices, please order from one of these suppliers. In the event that your preferred wholesaler cannot supply, please contact Smith & Nephew Customer Service on 13 13 60. Smith & Nephew cannot ensure RPBS agreed pricing from distributors other than those aforementioned.

dressings gauze paraffin with chlorhexidine acetate 10 cm x 10 cm dressing, 10

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
4845B	‡1	2	..	35.01	6.60	Bactigras 7457 [SN]

■ DRESSING GELLING FIBRE**dressings gelling fibre 5 cm x 5 cm dressing, 10**

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
12187K	‡1	..	..	40.73	6.60	Exufiber 709900 [MH]

dressings gelling fibre 10 cm x 10 cm dressing, 10

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
12181D	‡1	..	..	75.13	6.60	Exufiber 709901 [MH]

dressings gelling fibre 15 cm x 15 cm dressing, 10

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
12202F	‡1	..	..	158.41	6.60	Exufiber 709903 [MH]

dressings gelling fibre 2 cm x 45 cm dressing, 5

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
12182E	‡1	..	..	52.02	6.60	Exufiber 709909 [MH]

dressings gelling fibre 1 cm x 45 cm dressing, 5

12213T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	‡1	..	..	52.02	6.60	Exufiber 709908 [MH]

▪ DRESSING GELLING FIBRE

Note Smith & Nephew products are distributed via the three major wholesalers, API, Sigma and Symbion. To best ensure product availability at RPBS agreed prices, please order from one of these suppliers. In the event that your preferred wholesaler cannot supply, please contact Smith & Nephew Customer Service on 13 13 60. Smith & Nephew cannot ensure RPBS agreed pricing from distributors other than those aforementioned.

dressings gelling fibre 10 cm x 10 cm dressing, 10

2486W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	‡1	1	..	110.94	6.60	Durafiber 66800560 [SN]

dressings gelling fibre 15 cm x 15 cm dressing, 5

2445Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	1	..	*231.28	6.60	Durafiber 66800561 [SN]

dressings gelling fibre 2 cm x 45 cm rope, 5

2462N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	‡1	1	..	95.64	6.60	Durafiber 66800563 [SN]

▪ DRESSING HYDROACTIVE DEBRIDEMENT

Note Hartmann wound dressings are available through HARTMANN and Independence Australia only. If you would like to order Hartmann Wound Care products, please call HARTMANN customer service on 1800 805 839 or Independence Australia on 1300 788 855.

dressings hydroactive debridement 7.5 cm x 7.5 cm dressing, 10

4950M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	‡1	..	..	110.95	6.60	TenderWet 24 Active 609213 [HR]

dressings hydroactive debridement 5.5 cm dressing, 10

4948K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	‡1	..	..	85.21	6.60	TenderWet Active Cavity 609272 [HR]

dressings hydroactive debridement 4 cm dressing, 10

4949L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	‡1	..	..	83.35	6.60	TenderWet 24 Active 609210 [HR]

▪ DRESSING HYDROACTIVE SUPERFICIAL WOUND HIGH EXUDATE SEMI-PERMEABLE ABSORBENT FOAM**dressings hydroactive superficial wound high exudate semi-permeable absorbent foam 18 cm x 18 cm dressing: island, 5 dressings**

4696E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	‡1	..	..	130.55	6.60	Tielle MTL103 [KI]

dressings hydroactive superficial wound high exudate semi-permeable absorbent foam 11 cm x 11 cm dressing: island, 10 dressings

4695D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	‡1	..	..	107.31	6.60	Tielle MTL101E [KI]

▪ DRESSING HYDROACTIVE SUPERFICIAL WOUND HIGH EXUDATE SEMI-PERMEABLE ABSORBENT FOAM

Note Coloplast dressings are available via a range of distributors. However, Coloplast's principal agreement to ensure correct RPBS Price to Pharmacy and ready supply has been secured with Independence Australia on 1300 788 855 and BrightSky on 1300 290 400. Please note that Coloplast is unable to guarantee ready supply or rebate for price differences on purchases outside these distributors.

dressings hydroactive superficial wound high exudate semi-permeable absorbent foam 18 cm x 18 cm waterproof pad, 5

4930L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	‡1	2	..	91.47	6.60	Biatain Adhesive 3423 [CT]

dressings hydroactive superficial wound high exudate semi-permeable absorbent foam 12 cm x 12 cm waterproof pad, 10

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
4929K	‡1	1	..	94.32	6.60	Biatain Adhesive 3420 [CT]

dressings hydroactive superficial wound high exudate semi-permeable absorbent foam 15 cm x 15 cm waterproof pad, 5

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
4928J	‡1	2	..	84.77	6.60	Biatain Non-adhesive 3413 [CT]

dressings hydroactive superficial wound high exudate semi-permeable absorbent foam 10 cm x 10 cm waterproof pad, 10

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
4927H	‡1	1	..	86.12	6.60	Biatain Non-adhesive 3410 [CT]

▪ DRESSING HYDROACTIVE SUPERFICIAL WOUND LIGHT EXUDATE

Note Smith & Nephew products are distributed via the three major wholesalers, API, Sigma and Symbion. To best ensure product availability at RPBS agreed prices, please order from one of these suppliers. In the event that your preferred wholesaler cannot supply, please contact Smith & Nephew Customer Service on 13 13 60. Smith & Nephew cannot ensure RPBS agreed pricing from distributors other than those aforementioned.

dressings hydroactive superficial wound light exudate 5 cm x 6 cm dressing, 10

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
4905E	‡1	1	..	73.59	6.60	Allevyn Thin 66047576 [SN]

dressings hydroactive superficial wound light exudate 10 cm x 10 cm dressing, 5

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
4906F	2	1	..	*131.92	6.60	Allevyn Thin 66047578 [SN]

▪ DRESSING HYDROACTIVE SUPERFICIAL WOUND MODERATE EXUDATE

Note Smith & Nephew products are distributed via the three major wholesalers, API, Sigma and Symbion. To best ensure product availability at RPBS agreed prices, please order from one of these suppliers. In the event that your preferred wholesaler cannot supply, please contact Smith & Nephew Customer Service on 13 13 60. Smith & Nephew cannot ensure RPBS agreed pricing from distributors other than those aforementioned.

dressings hydroactive superficial wound moderate exudate 10 cm x 10 cm dressing, 5

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
4886E	2	1	..	*99.04	6.60	Cutinova Hydro 66047443 [SN]

dressings hydroactive superficial wound moderate exudate 5 cm x 6 cm dressing, 10

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
4885D	‡1	1	..	60.50	6.60	Cutinova Hydro 66047441 [SN]

▪ DRESSING HYDROCOLLOID CAVITY WOUND

Note This dressing should remain in place until saturated or strike through occurs for a maximum of 7 days.

dressings hydrocolloid cavity wound paste, 30 g

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
4896Q	10	..	..	*139.44	6.60	DuoDERM Paste 187930 [CC]

▪ DRESSING HYDROCOLLOID SUPERFICIAL WOUND LIGHT EXUDATE

Note This dressing should be applied to a thickness of 3 mm to 5 mm. It should be covered with a hydrocolloid dressing and may be left in place for up to 7 days.

dressings hydrocolloid superficial wound light exudate 10 cm x 10 cm dressing, 10

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
4907G	‡1	1	..	71.38	6.60	DuoDERM Extra Thin 187955 [CC]

▪ DRESSING HYDROCOLLOID SUPERFICIAL WOUND LIGHT EXUDATE

Note This dressing should be applied to a thickness of 3 mm to 5 mm. It should be covered with a hydrocolloid dressing and may be left in place for up to 7 days.

Note Coloplast dressings are available via a range of distributors. However, Coloplast's principal agreement to ensure correct RPBS Price to Pharmacy and ready supply has been secured with Independence Australia on 1300 788 855 and BrightSky on 1300 290 400. Please note that Coloplast is unable to guarantee ready supply or rebate for price differences on purchases outside these distributors.

dressing hydrocolloid superficial wound light exudate 5 cm x 7 cm dressing, 10

4888G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	1	..	42.82	6.60	Comfeel Plus Transparent 3530 [CT]

dressing hydrocolloid superficial wound light exudate 9 cm x 14 cm dressing, 10

4889H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	1	..	83.02	6.60	Comfeel Plus Transparent 3536 [CT]

dressing hydrocolloid superficial wound light exudate 10 cm x 10 cm dressing, 10

4924E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	1	..	69.62	6.60	Comfeel Plus Transparent 3533 [CT]

▪ DRESSING HYDROCOLLOID SUPERFICIAL WOUND LIGHT EXUDATE

Note This dressing should be applied to a thickness of 3 mm to 5 mm. It should be covered with a hydrocolloid dressing and may be left in place for up to 7 days.

Note Hartmann wound dressings are available through HARTMANN and Independence Australia only. If you would like to order Hartmann Wound Care products, please call HARTMANN customer service on 1800 805 839 or Independence Australia on 1300 788 855.

dressing hydrocolloid superficial wound light exudate 10 cm x 10 cm dressing, 10

4947J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	1	..	49.25	6.60	Hydrocoll Thin 900758 [HR]

▪ DRESSING HYDROCOLLOID SUPERFICIAL WOUND MODERATE EXUDATE

Note This dressing should remain in place until saturated or strike through occurs for a maximum of 7 days.

dressing hydrocolloid superficial wound moderate exudate 20 cm x 20 cm dressing, 5

4920Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	1	..	*214.82	6.60	DuoDERM CGF 187662 [CC]

dressing hydrocolloid superficial wound moderate exudate 10 cm x 10 cm dressing, 5

4897R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	1	..	*80.18	6.60	DuoDERM CGF 187660 [CC]

▪ DRESSING HYDROCOLLOID SUPERFICIAL WOUND MODERATE EXUDATE

Note This dressing should remain in place until saturated or strike through occurs for a maximum of 7 days.

Note Smith & Nephew products are distributed via the three major wholesalers, API, Sigma and Symbion. To best ensure product availability at RPBS agreed prices, please order from one of these suppliers. In the event that your preferred wholesaler cannot supply, please contact Smith & Nephew Customer Service on 13 13 60. Smith & Nephew cannot ensure RPBS agreed pricing from distributors other than those aforementioned.

dressing hydrocolloid superficial wound moderate exudate 10 cm x 10 cm dressing, 10

4921B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	1	..	100.08	6.60	Replicare Ultra 66000434 [SN]

▪ DRESSING HYDROCOLLOID SUPERFICIAL WOUND MODERATE EXUDATE

Note This dressing should remain in place until saturated or strike through occurs for a maximum of 7 days.

Note Hartmann wound dressings are available through HARTMANN and Independence Australia only. If you would like to order Hartmann Wound Care products, please call HARTMANN customer service on 1800 805 839 or Independence Australia on 1300 788 855.

dressing hydrocolloid superficial wound moderate exudate 10 cm x 10 cm dressing, 10

4945G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	1	..	49.25	6.60	Hydrocoll 900744 [HR]

dressing hydrocolloid superficial wound moderate exudate 15 cm x 15 cm dressing, 10

4946H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	1	..	87.92	6.60	Hydrocoll 900936 [HR]

▪ DRESSING HYDROCOLLOID SUPERFICIAL WOUND MODERATE EXUDATE

Note This dressing should remain in place until saturated or strike through occurs for a maximum of 7 days.

Note Coloplast dressings are available via a range of distributors. However, Coloplast's principal agreement to ensure correct RPBS Price to Pharmacy and ready supply has been secured with Independence Australia on 1300 788 855 and BrightSky on 1300 290 400. Please note that Coloplast is unable to guarantee ready supply or rebate for price differences on purchases outside these distributors.

DRESSING-HYDROCOLLOID (SUPERFICIAL WOUND-MODERATE EXUDATE) Dressings with alginate 10 cm x 10 cm, 10, 1

4923D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	‡1	1	..	80.73	6.60	Comfeel Plus Ulcer Dressing 3110 [CT]

dressing hydrocolloid superficial wound moderate exudate 10cm (round) dressing, 1

4679G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	5	..	..	*60.39	6.60	Comfeel Plus Pressure Relieving 3353 [CT]

dressing hydrocolloid superficial wound moderate exudate 7cm (butterfly shape) dressing, 1

4678F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	5	..	..	*56.29	6.60	Comfeel Plus Pressure Relieving 3350 [CT]

▪ DRESSING HYDROFIBRE ALTERNATE TO ALGINATES**dressing hydrofibre alternate to alginates 15 cm x 15 cm dressing, 5**

2803M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	1	..	*200.52	6.60	Aquacel Extra 420673 [CC]

dressing hydrofibre alternate to alginates 10 cm x 10 cm dressing, 10

10837J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	‡1	..	..	121.59	6.60	Aquacel Foam Non-Adhesive [CC]

dressing hydrofibre alternate to alginates 10 cm x 10 cm dressing, 10

2797F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	‡1	1	..	97.98	6.60	Aquacel Extra 420672 [CC]

dressing hydrofibre alternate to alginates 2 g (30 cm) rope, 5 x 2 g

4698G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	‡1	1	..	82.28	6.60	Aquacel 403770 [CC]

dressing hydrofibre alternate to alginates 12.5 cm x 12.5 cm dressing, 10

10832D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	‡1	..	..	117.94	6.60	Aquacel Foam Adhesive [CC]

▪ DRESSING HYDROFIBRE WITH SILVER**Authority required**

Wounds

Clinical criteria:

- Patient must have a wound where there is evidence of critical colonisation; OR
- Patient must have a well-assessed chronic wound that has not responded to conventional dressings.

dressing hydrofibre with silver 15 cm x 15 cm dressing, 5

10098L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	‡1	1	..	274.44	6.60	Aquacel Ag 403710 [CC]

dressing hydrofibre with silver 10 cm x 10 cm dressing, 10

10097K	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	‡1	1	..	256.54	6.60	Aquacel Ag 403708 [CC]

dressing hydrofibre with silver 2 cm x 45 cm rope, 5

10105W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	‡1	1	..	217.04	6.60	Aquacel Ag 403771 [CC]

▪ DRESSING HYDROGEL**dressing hydrogel 10 cm x 10 cm dressing, 5**

11709G	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	‡1	..	..	30.57	6.60	Suprasorb G 33631 [LC]

dressing hydrogel 10 cm x 10 cm dressing, 20

2471C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	185.39	6.60	Sorbact Absorption Dressing S98222 [YB]

dressing hydrogel 7.5 cm x 15 cm dressing, 10

11395R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	‡1	..	..	114.27	6.60	Sorbact Gel Dressing S98137 [YB]

▪ DRESSING HYDROGEL AMORPHOUS

Note This dressing should be applied to a thickness of 3 mm to 5 mm and remain in situ in infected wounds for 24 hours and in clean wounds for up to 3 days. It should be covered with a secondary dressing such as foam or film. It should not be covered with gauze or combine.

dressing hydrogel amorphous gel, 3 x 30 g

4913N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	1	..	*94.47	6.60	DuoDERM Gel H7987 [CC]

dressing hydrogel amorphous gel, 50 g

4914P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	3	..	*35.25	6.60	Solugel 10336 [JJ]

▪ DRESSING HYDROGEL AMORPHOUS

Note This dressing should be applied to a thickness of 3 mm to 5 mm and remain in situ in infected wounds for 24 hours and in clean wounds for up to 3 days. It should be covered with a secondary dressing such as foam or film. It should not be covered with gauze or combine.

Note Coloplast dressings are available via a range of distributors. However, Coloplast's principal agreement to ensure correct RPBS Price to Pharmacy and ready supply has been secured with Independence Australia on 1300 788 855 and BrightSky on 1300 290 400. Please note that Coloplast is unable to guarantee ready supply or rebate for price differences on purchases outside these distributors.

dressing hydrogel amorphous gel, 10 x 15 g

4912M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	‡1	1	..	64.80	6.60	DuoDERM Gel 187990 [CC]
			..	71.72	6.60	Comfeel Purilon Gel 3900 [CT]

▪ DRESSING HYDROGEL AMORPHOUS

Note This dressing should be applied to a thickness of 3 mm to 5 mm and remain in situ in infected wounds for 24 hours and in clean wounds for up to 3 days. It should be covered with a secondary dressing such as foam or film. It should not be covered with gauze or combine.

Note Smith & Nephew products are distributed via the three major wholesalers, API, Sigma and Symbion. To best ensure product availability at RPBS agreed prices, please order from one of these suppliers. In the event that your preferred wholesaler cannot supply, please contact Smith & Nephew Customer Service on 13 13 60. Smith & Nephew cannot ensure RPBS agreed pricing from distributors other than those aforementioned.

dressing hydrogel amorphous gel, 25 g

4894N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	4	3	..	*78.58	6.60	Intrasite Gel 7313 [SN]

dressing hydrogel amorphous gel, 50 g

4599C	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	3	3	..	*38.37	6.60	SoloSite Gel 36361338 [SN]

▪ DRESSING HYDROGEL FOAM**dressing hydrogel foam 10 cm x 10 cm dressing, 10**

2533H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	125.57	6.60	Sorbact Foam Dressing S98310 [YB]

▪ DRESSING HYDROGEL RIBBON**dressing hydrogel ribbon 5 cm x 200 cm dressing, 10**

2529D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	185.39	6.60	Sorbact Ribbon Gauze S98120 [YB]

dressing hydrogel ribbon 1 cm x 50 cm dressing, 20

2512F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	1	..	..	187.65	6.60	Sorbact Ribbon Gauze S98118 [YB]

▪ DRESSING HYDROGEL SHEET

Note This dressing should be applied to a thickness of 3 mm to 5 mm and remain in situ in infected wounds for 24 hours and in clean wounds for up to 3 days. It should be covered with a secondary dressing such as foam or film. It should not be covered with gauze or combine.

dressing hydrogel sheet 9.5 cm x 10.2 cm dressing, 5

4911L	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*81.82	6.60	Nu-Gel 2497 [KI]

▪ DRESSING HYDROGEL SHEET

Note This dressing should be applied to a thickness of 3 mm to 5 mm and remain in situ in infected wounds for 24 hours and in clean wounds for up to 3 days. It should be covered with a secondary dressing such as foam or film. It should not be covered with gauze or combine.

Note Hartmann wound dressings are available through HARTMANN and Independence Australia only. If you would like to order Hartmann Wound Care products, please call HARTMANN customer service on 1800 805 839 or Independence Australia on 1300 788 855.

dressing hydrogel sheet 10 cm x 10 cm dressing, 5

4806Y	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*54.38	6.60	Hydrosorb 900854 [HR]

▪ DRESSING HYDROPHOBIC**dressing hydrophobic 10 cm x 10 cm dressing, 10**

11392N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	198.94	6.60	Sorbact Foam Gentle Border 98532 [YB]

dressing hydrophobic 10 cm x 10 cm dressing, 10

11402D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	89.97	6.60	Sorbact Superabsorbent 98501 [YB]

dressing hydrophobic 15 cm x 15 cm dressing, 10

11394Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	311.84	6.60	Sorbact Foam Gentle Border 98533 [YB]

dressing hydrophobic 20 cm x 20 cm dressing, 10

11403E	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	244.10	6.60	Sorbact Superabsorbent 98503 [YB]

dressing hydrophobic 15 cm x 15 cm foam dressing, 10

11404F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	204.59	6.60	Sorbact Foam Dressing S98315 [YB]

▪ DRESSING NON ADHERENT

Note Molnlycke Healthcare products are distributed through leading pharmacy distributors. To best ensure product availability at RPBS agreed prices, special arrangements have been made with API and Independence Australia Health Solutions (IAHS). IAHS orders can be placed on: Tel: 1300 788 855; or Email customerservice@independenceaustralia.com. Molnlycke Healthcare are not able to ensure product availability or pricing on listed products beyond these two suppliers.

DRESSING SELF ADHESIVE NON-ADHERENT DRY ABSORBENT Dressings, non-woven, with silicone 7.5 cm x 10 cm, 10, 1

4244J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	104.02	6.60	Mepitel 290710 [MH]

DRESSING SELF ADHESIVE NON-ADHERENT DRY ABSORBENT Dressings, non-woven, with silicone 5 cm x 7.5 cm, 10, 1

4243H	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	64.02	6.60	Mepitel 290510 [MH]

▪ DRESSING NON ADHERENT

Note Hartmann wound dressings are available through HARTMANN and Independence Australia only. If you would like to order Hartmann Wound Care products, please call HARTMANN customer service on 1800 805 839 or Independence Australia on 1300 788 855.

dressing non adherent 7.5 cm x 10 cm dressing, 10

4944F	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	19.69	6.60	Atrauman 499513 [HR]

▪ DRESSING NON ADHERENT

Note Smith & Nephew products are distributed via the three major wholesalers, API, Sigma and Symbion. To best ensure product availability at RPBS agreed prices, please order from one of these suppliers. In the event that your preferred wholesaler

cannot supply, please contact Smith & Nephew Customer Service on 13 13 60. Smith & Nephew cannot ensure RPBS agreed pricing from distributors other than those aforementioned.

dressings non adherent 10 cm x 10 cm dressing, 10

4861W	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	41.27	6.60	Melolin 66974933 [SN]

dressings non adherent 5 cm x 5 cm dressing, 5

4819P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*20.98	6.60	Cutilin Non-Stick Wound Pad 36361374 [SN]

dressings non adherent 5 cm x 5 cm dressing, 5

4860T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*22.48	6.60	Melolin 36361357 [SN]

dressings non adherent 10 cm x 10 cm dressing, 5

4862X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	2	..	..	*30.22	6.60	Cutilin Non-Stick Wound Pad 36361375 [SN]

▪ DRESSING NON-ADHERENT ABSORBENT

dressings non-adherent absorbent 22 cm x 32 cm hydroactive dressing, 10

11718R	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	49.65	6.60	Vliwasorb Pro 32644 [LC]

dressings non-adherent absorbent 12.5 cm x 12.5 cm hydroactive dressing, 10

11717Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	26.21	6.60	Vliwasorb Pro 32641 [LC]

dressings non-adherent absorbent 22 cm x 22 cm hydroactive dressing, 10

11715N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	37.07	6.60	Vliwasorb Pro 32643 [LC]

▪ DRESSING NON-ADHERENT WITH SILICONE

dressings non-adherent with silicone 5 cm x 7.5 cm dressing, 10

12208M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	91.69	6.60	Mepitel One 289100 [MH]

dressings non-adherent with silicone 10 cm x 18 cm dressing, 10

12196X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	308.67	6.60	Mepitel One 289500 [MH]

▪ DRESSING TULLE NON-ADHERENT PRIMARY WOUND CONTACT LAYER PARAFFIN

dressings tulle non-adherent primary wound contact layer paraffin 7.6 cm x 7.6 cm dressing, 1

4909J	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	10	1	..	*20.14	6.60	Adaptic 2012 [KI]

▪ DRESSING WITH SILVER

Note Coloplast dressings are available via a range of distributors. However, Coloplast's principal agreement to ensure correct RPBS Price to Pharmacy and ready supply has been secured with Independence Australia on 1300 788 855 and BrightSky on 1300 290 400. Please note that Coloplast is unable to guarantee ready supply or rebate for price differences on purchases outside these distributors.

Authority required

Wounds

Clinical criteria:

- Patient must have a wound where there is evidence of critical colonisation; OR
- Patient must have a well-assessed chronic wound that has not responded to conventional dressings.

dressings with silver 12.5 cm x 12.5 cm hydroactive dressing, 5

4647N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	183.48	6.60	Biatain Ag 9632 [CT]

dressings with silver 10 cm x 10 cm hydroactive dressing, 5

4646M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	168.77	6.60	Biatain Ag 9622 [CT]

■ DRESSING WITH SILVER

Note Hartmann wound dressings are available through HARTMANN and Independence Australia only. If you would like to order Hartmann Wound Care products, please call HARTMANN customer service on 1800 805 839 or Independence Australia on 1300 788 855.

Authority required

Wounds

Clinical criteria:

- Patient must have a wound where there is evidence of critical colonisation; OR
- Patient must have a well-assessed chronic wound that has not responded to conventional dressings.

dressings with silver 10 cm x 10 cm tulle dressing, 3

4648P	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	44.88	6.60	Atrauman Ag 499572 [HR]

■ GAUZE AND COTTON TISSUE COMBINE ROLL

gauze and cotton tissue combine roll 10 cm x 10 m wrapped pack roll, 1

4761N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	21.40	6.60	JJ 12010 [JJ]

gauze and cotton tissue combine roll 9 cm x 10 m wrapped pack roll, 1

4767X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	16.79	6.60	BSN 2902165 [BV]

■ PAD WOUND DEBRIDEMENT

Note If the wound has not healed during this period, further use is to be discontinued after initial pack, no repeats. Where wounds remain unresponsive to standard treatment, patient should be referred on to a specialist.

pad wound debridement pad, 5

11391M	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	92.39	6.60	Debrisoft Lolly [LC]

pad wound debridement 10 cm x 10 cm pad, 5

11383D	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	92.39	6.60	Debrisoft [LC]

■ POVIDONE-IODINE

povidone-iodine 9.5 cm x 9.5 cm dressing, 25

10847X	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	2	..	78.57	6.60	Inadine [KI]

■ SODIUM CHLORIDE + HYPOCHLOROUS ACID + SODIUM HYPOCHLORITE

sodium chloride 0.022% + hypochlorous acid 0.004% + sodium hypochlorite 0.004% irrigation solution, 250 mL

11134B	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	3	..	31.37	6.60	Microdacyn [TF]

■ TAPE NON WOVEN RETENTION POLYACRYLATE

tape non woven retention polyacrylate 2.5 cm x 9.1 m tape, 1 roll

4915Q	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	17.65	6.60	Medipore 2961 [MM]

■ TAPE NON WOVEN RETENTION POLYACRYLATE

Note Molnlycke Healthcare products are distributed through leading pharmacy distributors. To best ensure product availability at RPBS agreed prices, special arrangements have been made with API and Independence Australia Health Solutions (IAHS). IAHS orders can be placed on: Tel: 1300 788 855; or Email customerservice@independenceaustralia.com. Molnlycke Healthcare are not able to ensure product availability or pricing on listed products beyond these two suppliers.

tape non woven retention polyacrylate 2.5 cm x 10 m tape, 1 roll

4917T	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	16.17	6.60	Mefix 310250 [MH]

■ TAPE PLASTER ADHESIVE ELASTIC

tape plaster adhesive elastic 2.5 cm x 2.5 m tape, 1 roll

4780N	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
	±1	..	..	18.55	6.60	Leukoplast 01071-00 [BV]

tape plaster adhesive elastic 7.5 cm x 2.5 m tape, 1 roll

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
4782Q	±1	..	..	28.17	6.60	Leukoplast 01073-00 [BV]

tape plaster adhesive elastic 5 cm x 2.5 m tape, 1 roll

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
4781P	±1	..	..	24.49	6.60	Leukoplast 01072-00 [BV]

■ TAPE PLASTER ADHESIVE HYPOALLERGENIC**tape plaster adhesive hypoallergenic 2.5 cm x 5 m tape, 1 roll**

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
4787Y	±1	..	..	18.90	6.60	Leukosilk 1022 [BV]

tape plaster adhesive hypoallergenic 2.5 cm x 5 m tape, 1 roll

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
4794H	±1	..	..	18.37	6.60	Leukopor 2472 [BV]

tape plaster adhesive hypoallergenic 5 cm x 5 m tape, 1 roll

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
4788B	±1	..	..	21.40	6.60	Leukoflex 1124 [BV]

tape plaster adhesive hypoallergenic 5 cm x 5 m tape, 1 roll

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
4789C	±1	..	..	22.72	6.60	Leukosilk 1024 [BV]

tape plaster adhesive hypoallergenic 5 cm x 5 m tape, 1 roll

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
4790D	±1	..	..	21.89	6.60	Leukopor 2474 [BV]

tape plaster adhesive hypoallergenic 1.9 cm x 7.3 m dispenser tape, 1 roll

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
4849F	±1	..	..	16.18	6.60	Nexcare Gentle Paper First Aid Tape 789 [MM]

tape plaster adhesive hypoallergenic 1.9 cm x 5.4 m dispenser tape, 1 roll

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
4848E	±1	..	..	16.18	6.60	Nexcare Durable Cloth First Aid Tape 799 [MM]

tape plaster adhesive hypoallergenic 1.25 cm x 5 m tape, 1 roll

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
4783R	±1	..	..	16.09	6.60	Leukopor 2471 [BV]

tape plaster adhesive hypoallergenic 1.25 cm x 5 m tape, 1 roll

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
4785W	±1	..	..	16.36	6.60	Leukosilk 1021 [BV]

■ TAPE PLASTER ADHESIVE WITH SILICONE

Note Molnlycke Healthcare products are distributed through leading pharmacy distributors. To best ensure product availability at RPBS agreed prices, special arrangements have been made with API and Independence Australia Health Solutions (IAHS). IAHS orders can be placed on: Tel: 1300 788 855; or Email customerservice@independenceaustralia.com. Molnlycke Healthcare are not able to ensure product availability or pricing on listed products beyond these two suppliers.

tape plaster adhesive with silicone 2 cm x 3 m tape, 1 roll

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
4239D	±1	..	..	25.02	6.60	Mepitac 298300 [MH]

tape plaster adhesive with silicone 4 cm x 1.5 m tape, 1 roll

	Max.Qty Packs	No. of Rpts	Premium \$	DPMQ \$	MRVSN \$	Brand Name and Manufacturer
4240E	±1	..	..	25.02	6.60	Mepitac 298400 [MH]

Extemporaneously Prepared Benefits

Drug Tariff

Drug	Standard	Recovery Prices			
		0.1 g/mL \$	1 g/mL \$	10 g/mL \$	100 g/mL \$
Acacia Mucilage (by weight)	APF 15	0.01	0.11	0.87	7.77
Acacia, powdered	BP	0.02	0.19	1.55	13.82
Acetic Acid (33 per cent)	BP	0.01	0.06	0.47	4.20
Acetic Acid (6 per cent)	BP	0.01	0.02	0.16	1.47
Acetone (use as additive only)	BP	0.02	0.19	1.52	13.48
Alum	BP	0.01	0.08	0.60	5.32
Aluminium Acetate Solution	BP	0.02	0.17	1.38	12.27
Anise Water Concentrated 1 in 40	BP	0.01	0.08	0.60	5.33
Aqueous Cream (for use only as a base combined with active ingredients)	APF	0.01	0.05	0.39	3.44
Ascorbic Acid (for use only as an ingredient of ferrous sulfate mixtures)	BP	0.39	3.14	25.11	223.22
Aspirin	BP	0.19	1.51	12.09	107.43
Belladonna Tincture	BP	0.20	1.59	12.74	113.26
Benzocaine	BP	0.14	1.14	9.11	80.98
Benzoic Acid	BP	0.07	0.56	4.50	39.98
Benzoic Acid Compound Ointment	APF	0.02	0.19	1.48	13.12
Benzoic Acid Solution	BP	0.02	0.14	1.15	10.24
Benzoin Compound Tincture	BP	0.07	0.55	4.39	38.98
Boric Acid (use as additive only)	BP	0.02	0.17	1.38	12.30
Boric Acid, Olive Oil and Zinc Oxide Ointment	QHF	0.02	0.15	1.22	10.83
Calcium Hydroxide	BP	0.12	0.94	7.54	67.02
Calcium Hydroxide Solution	BP	0.01	0.03	0.20	1.80
Castor Oil (use as additive only)	BP	0.02	0.16	1.24	11.03
Cetomacrogol Aqueous Cream (for use only as a base combined with active ingredients)	APF	0.01	0.05	0.42	3.70
Cetrimide Aqueous Cream (for use only as a base combined with active ingredients)	APF	0.02	0.17	1.37	12.21
Chlorhexidine Acetate (use as additive only)	BP	0.63	5.03	40.21	357.42
Chlorhexidine Aqueous Cream (for use only as a base combined with active ingredients)	APF	0.03	0.24	1.95	17.36
Chloroform (use as additive only)	BP	0.10	0.80	6.37	56.63
Chloroform Spirit	BP	0.01	0.09	0.74	6.62
Chloroform Water Concentrated 1 in 40	APF 15	0.02	0.12	0.93	8.29
Citric Acid Monohydrate	BP	0.04	0.30	2.40	21.30
Coal Tar	BP	0.32	2.54	20.30	180.45
Coal Tar Solution	BP	0.02	0.19	1.55	13.78
Cocaine Hydrochloride	BP	5.81	46.47	371.74	3304.35
Coconut Oil	BP	0.01	0.05	0.36	3.17
Codeine Linctus	APF	0.02	0.13	1.04	9.25
Codeine Phosphate (may only be prescribed in linctuses, mixtures or mixtures for children)	BP	2.88	23.07	184.56	1640.50
Collodion Flexible	BP	0.20	1.57	12.54	111.51
Dithranol	BP	5.03	40.20	321.56	2858.28
Emulsifying Ointment (for use only as a base combined with active ingredients)	BP	0.02	0.16	1.26	11.16
Ephedrine Hydrochloride (may only be prescribed in nasal instillations)	BP	2.61	20.90	167.21	1486.33
Ethanol (90 per cent) (use as additive only)	BP	0.01	0.04	0.31	2.77
Ethanol (96 per cent) (use as additive only)	BP	0.01	0.06	0.44	3.90
Ether Solvent (use as additive only)	BP	0.29	2.35	18.83	167.36
Eucalyptus Oil (use as additive only)	BP	0.03	0.21	1.68	14.96
Ferrous Sulfate	BP	0.05	0.40	3.19	28.38

Drug	Standard	Recovery Prices			
		0.1 g/mL \$	1 g/mL \$	10 g/mL \$	100 g/mL \$
Formaldehyde Solution	BP	0.07	0.53	4.26	37.86
Gentian Alkaline Mixture	APF	0.01	0.09	0.68	6.00
Glycerol	BP	0.02	0.13	1.04	9.26
Honey Purified (use as additive only)	BP 1993	0.01	0.03	0.27	2.44
Hydroxybenzoate Compound Solution	APF	0.09	0.68	5.42	48.17
Iodine	BP	0.34	2.72	21.79	193.72
Iodine Alcoholic Solution	BP	0.04	0.32	2.59	22.99
Iodine Aqueous Oral Solution	BP	0.04	0.33	2.62	23.27
Kaolin Mixture	BPC 1968	0.03	0.25	1.99	17.68
Kaolin and Opium Mixture	APF 14	0.01	0.10	0.82	7.27
Lactic Acid	BP	0.38	3.02	24.13	214.48
Lavender Spike Oil	BPC 1968	0.13	1.04	8.32	73.91
Liquorice Liquid Extract	BP	0.03	0.23	1.84	16.37
Magnesium Carbonate Light	BP	0.05	0.37	2.92	25.99
Magnesium Sulfate (may only be prescribed for other than oral use)	BP	0.01	0.04	0.29	2.60
Magnesium Trisilicate	BP	0.06	0.44	3.52	31.25
Menthol, Racemic or Levomenthol	BP	0.28	2.24	17.94	159.50
Methyl Hydroxybenzoate	BP	0.46	3.71	29.71	264.11
Methyl Hydroxybenzoate Solution	APF	0.05	0.38	3.05	27.09
Methylated Industrial Spirit (use as additive only)	BP	0.01	0.02	0.12	1.04
Olive Oil (use as additive only)	BP	0.02	0.13	1.05	9.32
Paraffin Hard	BP	0.06	0.49	3.91	34.76
Paraffin Light Liquid	BP	0.02	0.17	1.39	12.32
Paraffin Liquid (use as additive only)	BP	0.01	0.08	0.60	5.34
Paraffin Soft White	BP	0.01	0.06	0.49	4.33
Paraffin Soft Yellow	BP	0.01	0.05	0.42	3.75
Peppermint Oil (use as additive only)	BP	0.08	0.66	5.28	46.92
Peppermint Water Concentrated 1 in 40 (use as additive only)	APF 16	0.04	0.31	2.48	22.05
Phenobarbitone Sodium (may only be prescribed for the treatment of epilepsy)	BP	9.37	74.92	599.36	5327.61
Phenol Liquefied (not available for ear drops)	BP	0.26	2.08	16.60	147.57
Podophyllum Resin	BP	4.55	36.41	291.31	2589.44
Potassium Citrate	BP	0.03	0.22	1.79	15.92
Potassium Iodide	BP	0.47	3.79	30.28	269.13
Potassium Permanganate	BP	0.06	0.50	4.01	35.65
Propyl Hydroxybenzoate	BP	0.42	3.36	26.91	239.22
Propylene Glycol	BP	0.02	0.12	0.96	8.52
Red Syrup	APF 15	0.02	0.13	1.04	9.23
Resorcinol	BP	0.45	3.57	28.59	254.15
Salicylic Acid	BP	0.04	0.33	2.64	23.49
Salicylic Acid Ointment	APF	0.02	0.17	1.38	12.23
Salicylic Acid Ointment	BP	0.02	0.17	1.38	12.23
Simple Ointment (white) (for use only as a base combined with active ingredients)	BP	0.02	0.14	1.11	9.88
Simple Ointment (yellow) (for use only as a base combined with active ingredients)	BP	0.02	0.14	1.11	9.88
Sodium Bicarbonate	BP	0.03	0.21	1.65	14.65
Sodium Chloride	BP	0.03	0.20	1.57	13.94
Sodium Chloride Solution	BP	0.01	0.02	0.12	1.04
Sodium Citrate	BP	0.07	0.56	4.45	39.54
Sodium Thiosulfate (use as additive only)	BP	0.05	0.40	3.20	28.46
Starch	BP	0.02	0.12	0.95	8.48
Sulfur Ointment (for use only as a base combined with active ingredients)	BP 1980	0.02	0.16	1.24	10.99
Sulfur Precipitated	BP 1980	0.03	0.26	2.09	18.57
Syrup	BP	0.01	0.06	0.47	4.16
Talc Purified, sterilised	BP	0.07	0.52	4.13	36.71
Thymol	BP	0.56	4.44	35.51	315.66
Thymol Compound Mouth Wash	APF 15	0.02	0.13	1.06	9.40
Tragacanth Compound Powder	BP 1980	0.07	0.57	4.58	40.73
Tragacanth Mucilage	APF 13	0.01	0.09	0.72	6.36
Tragacanth Mucilage	BPC 1973	0.01	0.08	0.64	6.36
Tragacanth, powdered	BP	0.45	3.59	28.75	255.53

Drug	Standard	Recovery Prices			
		0.1 g/mL \$	1 g/mL \$	10 g/mL \$	100 g/mL \$
Trichloroacetic Acid	BP 1980	0.40	3.17	25.32	225.05
Triethanolamine	BP	0.16	1.28	10.27	91.32
Water For Injections, sterilised (b) (extemporaneously prepared eye drops and eye lotions)	BP	0.00	0.00	11.60	11.60
Water Purified	BP	0.01	0.01	0.10	0.86
Wool Alcohols Ointment (white) (for use only as a base combined with active ingredients)	BP	0.03	0.21	1.64	14.55
Wool Alcohols Ointment (yellow) (for use only as a base combined with active ingredients)	BP	0.02	0.19	1.48	13.18
Wool Fat	BP	0.03	0.21	1.68	14.92
Wool Fat Hydrous	BP	0.04	0.30	2.42	21.53
Zinc Compound Paste	BP	0.05	0.40	3.17	28.17
Zinc Cream (for use only as a base combined with active ingredients)	BP	0.01	0.09	0.73	6.52
Zinc Oxide	BP	0.03	0.23	1.85	16.46
Zinc Sulfate	BP	0.06	0.46	3.67	32.61
Zinc and Salicylic Acid Paste	BP	0.05	0.37	2.99	26.56

Container Prices

Type	Container	Price \$
Dispensing Bottles	25mL	1.74
Dispensing Bottles	50mL	1.39
Dispensing Bottles	100mL	0.89
Dispensing Bottles	200mL	1.18
Dispensing Bottles	500mL	1.28
Poison Bottles	25mL	0.90
Poison Bottles	50mL	0.81
Poison Bottles	100mL	0.90
Poison Bottles	200mL	1.12
Poison Bottles	500mL	2.06
Dropper Containers (Glass)	15mL	1.13
Dropper Containers (Polythene)	15mL	0.98
Screw Cap Jars	25g	0.99
Screw Cap Jars	50g	1.19
Screw Cap Jars	100g	1.24
Screw Cap Jars	200g	0.77
Screw Cap Jars	500g	2.19

Standard Formula Preparations

Code	Item	Reference	DPMQ \$	MRVSN \$
	Creams			
7502W	(Maximum Quantity 100 g and 1 Repeat) Salicylic Acid and Sulfur Aqueous	APF	15.50	17.16
	Dusting Powders			
7458M	(Maximum Quantity 100 g and 1 Repeat) Zinc, Starch and Talc	APF 15 & BPC 1973	38.22	39.88
	Ear Drops			
7643G	(Maximum Quantity 15 ml and 2 Repeats) Aluminium Acetate	BP	12.83	14.49
7642F	Aluminium Acetate	APF	12.08	13.74
7314Y	Sodium Bicarbonate	APF & BP	11.60	13.26
7313X	Spirit	APF	11.06	12.72
	Inhalations			
7484X	(Maximum Quantity 50 ml and 1 Repeat) Benzoin and Menthol	APF	34.76	36.42
7308P	Menthol	APF	14.39	16.05
7310R	Menthol and Eucalyptus	BP1980	15.59	17.25
	Linctuses containing Codeine Phosphate			
7530H	(Maximum Quantity 100 ml and 0 Repeat) Codeine	APF	19.92	21.58
	Lotions			
7709R	(Maximum Quantity 200 ml and 2 Repeats) Aluminium Acetate Aqueous	APF	13.91	15.57
	Mixtures, Other			
7604F	(Maximum Quantity 200 ml and 4 Repeats) Gentian Alkaline	APF	22.96	24.62
7348R	Kaolin	BPC 1968	46.33	41.30
7301G	Kaolin and Opium	APF 14	25.50	27.16
7342K	Magnesium Trisilicate	BPC 1968	22.27	23.93
7343L	Magnesium Trisilicate and Belladonna	BPC 1968	34.85	36.51
	Mouth Washes			
7457L	(Maximum Quantity 200 ml and 1 Repeat) Thymol Compound	APF 15	29.69	31.35
	Ointments, Waxes			
7914M	(Maximum Quantity 100 g and 1 Repeat) Benzoic Acid Compound	APF & BP	24.14	25.80
7902X	Boric Acid, Olive Oil and Zinc Oxide	QHF	21.85	23.51
7926E	Salicylic Acid	APF	23.25	24.91
7928G	Salicylic Acid (extemporaneous formula)	BP	23.25	24.91
	Paints			
7567G	(Maximum Quantity 25 ml and 1 Repeat) Podophyllin Compound	APF 16 & BP	158.19	41.30
7568H	Salicylic Acid	APF	42.87	41.30
	Pastes, Other			
7558T	(Maximum Quantity 100 g and 1 Repeat) Zinc	APF & BP	39.19	40.85
	Powders for Internal Use			
7545D	(Maximum Quantity 100 g and 2 Repeats) Magnesium Trisilicate	BP	41.80	41.30

Codes, Maximum Quantities, and Number of Repeats for Extemporaneously Prepared Benefits

Code	Preparation	Maximum Quantity	Number of Repeats
13Q	Creams	100 g	1
48M	Dusting Powders	100 g	1
15T	Ear Drops	15 ml	2
19B	Eye Drops containing Cocaine Hydrochloride	15 ml	..
22E	Eye Drops, Other	15 ml	5
23F	Eye Lotions	200 ml	2
29M	Inhalations	50 ml	1
64J	Linctuses containing Codeine Phosphate	100 ml	..
34T	Linctuses, Other	100 ml	2
39C	Lotions	200 ml	2
65K	Mixtures containing Codeine Phosphate	200 ml	..
66L	Mixtures for Children containing Codeine Phosphate	100 ml	..
41E	Mixtures for Children, Other	100 ml	4
40D	Mixtures, Other	200 ml	4
30N	Mouth Washes	200 ml	1
42F	Nasal Instillations	15 ml	2
43G	Ointments, Waxes	100 g	1
44H	Paints	25 ml	1
63H	Pastes containing Cocaine Hydrochloride	25 g	..
45J	Pastes, Other	100 g	1
49N	Powders for Internal Use	100 g	2
52R	Solutions	200 ml	2

Index of Manufacturers' Code

Code	Manufacturer	Code	Manufacturer
AB	Abbott Australasia Pty Ltd	GZ	sanofi-aventis Australia Pty Ltd
AE	AFT Pharmaceuticals (AU) Pty Ltd	HB	Besins Healthcare Australia Pty Ltd
AF	Alphapharm Pty Ltd	HN	Horizon Hospital Healthcare Pty Ltd
AG	Allergan Australia Pty Limited	HQ	Generic Health Pty Ltd
AL	Alphapharm Pty Ltd	HR	Paul Hartmann Pty Ltd
AN	Amgen Australia Pty Limited	HX	Sandoz Pty Ltd
AP	AstraZeneca Pty Ltd	IB	Apotex Pty Ltd
AQ	Alcon Laboratories (Australia) Pty Ltd	IG	Sigma Company Limited
AS	Aspen Pharmacare Australia Pty Limited	IJ	I-Care Pharma Distributors Pty Ltd
AV	sanofi-aventis Australia Pty Ltd	IL	iNova Pharmaceuticals (Australia) Pty Limited
BB	Blackmores Limited	IM	iNova Pharmaceuticals (Australia) Pty Limited
BD	Biogen Australia Pty Ltd	IO	BioMarin Pharmaceutical Australia Pty Ltd
BE	Beiersdorf Australia Ltd	IQ	Alcon Laboratories (Australia) Pty Ltd
BG	Sandoz Pty Ltd	IR	Indivior Pty Ltd
BN	Bayer Australia Ltd	IS	Ipsen Pty Ltd
BQ	Bristol-Myers Squibb Australia Pty Ltd	IX	Clinect Pty Ltd
BR	B. Braun Australia Pty Ltd	IY	Clinect Pty Ltd
BV	BSN medical (Aust.) Pty Ltd	JB	Apotex Pty Ltd
BX	Baxter Healthcare Pty Limited	JC	Janssen-Cilag Pty Ltd
BY	Boehringer Ingelheim Pty Ltd	JJ	Johnson & Johnson Medical Pty Ltd
BZ	Boucher & Muir Pty Ltd	JO	Juno Pharmaceuticals Pty Ltd
CC	ConvaTec A Division of Bristol-Myers Squibb Australia Pty Ltd	JT	Johnson & Johnson Pacific Pty Limited
CF	CNS Pharma Pty Ltd	JU	Juno Pharmaceuticals Pty Ltd
CH	Apotex Pty Ltd	JX	Juno Pharmaceuticals Pty Ltd
CJ	Celgene Pty Limited	KE	Kendall Australasia Pty Ltd
CR	Pharmacor Pty Limited	KI	KCI Medical Australia Pty Ltd
CS	Seqirus (Australia) Pty Ltd	KP	Eli Lilly Australia Pty Ltd
CT	Coloplast Pty Ltd	KY	Key Pharmaceuticals Pty Ltd
CU	Care Pharmaceuticals Pty Limited	LC	Lohmann & Rauscher Pty Ltd
CX	Contact Lens Centre Australia Limited	LI	Luminarie Pty Ltd
DE	Stallergenes Australia Pty Ltd	LL	Astellas Pharma Australia Pty Ltd
DJ	De Fries Industries Pty Ltd	LM	Link Medical Products Pty Ltd
DO	Fair-Med Healthcare (Australia) Pty Ltd	LN	Aspen Pharmacare Australia Pty Limited
DQ	Church & Dwight (Australia) Pty Ltd	LO	Leo Pharma Pty Ltd
DV	Medical Developments International Limited	LQ	Astellas Pharma Australia Pty Ltd
DX	Ascensia Diabetes Care Australia Pty Ltd	LR	Cipla Australia Pty Ltd
DZ	Medsurge Healthcare Pty Ltd	LS	Astellas Pharma Australia Pty Ltd
EA	Amneal Pharmaceuticals Pty Ltd	LU	Lundbeck Australia Pty Ltd
ED	Amneal Pharmaceuticals Pty Ltd	LV	Luminarie Pty Ltd
EF	Amneal Pharmaceuticals Pty Ltd	LX	Lawley Pharmaceuticals Pty Ltd
EI	Eisai Australia Pty Ltd	LY	Eli Lilly Australia Pty Ltd
EJ	Encapsulate Pharma Pty Ltd	MF	Mundipharma Pty Limited
EL	Eli Lilly Australia Pty Ltd	MH	Molnlycke Health Care Pty Ltd
EO	Ego Pharmaceuticals Proprietary Limited	MK	Merck Sharp & Dohme (Australia) Pty Ltd
ER	Eris Pharmaceuticals (Australia) Pty Ltd	MM	3M Pharmaceuticals Australia Pty Ltd
EU	Chiesi Australia Pty Ltd	MQ	Alphapharm Pty Ltd
EV	Teva Pharma Australia Pty Limited	MT	Mentholatum Australasia Pty Ltd
EW	Celltrion Healthcare Australia Pty Ltd	MW	Biomed Aust Pty Limited
FB	Pierre Fabre Australia Pty Ltd	NE	Norgine Pty Limited
FD	Dr Falk Pharma Australia Pty Ltd	NF	Novo Nordisk Pharmaceuticals Pty Limited
FF	Phebra Pty Ltd	NI	Novo Nordisk Pharmaceuticals Pty Limited
FG	Phebra Pty Ltd	NM	Novartis Pharmaceuticals Australia Pty Limited
FI	Boehringer Ingelheim Pty Ltd	NO	Novo Nordisk Pharmaceuticals Pty Limited
FK	A.Menarini Australia Pty Limited	NQ	Takeda Pharmaceuticals Australia Pty Ltd
FO	For Benefit Medicines Pty Ltd	NT	Nestle Australia Ltd
FP	Ferring Pharmaceuticals Pty Limited	NU	Nutricia Australia Pty Limited
FX	Gedeon Richter Australia Pty Ltd	NV	Novartis Pharmaceuticals Australia Pty Limited
FZ	Pfizer Australia Pty Ltd	OB	Oral B Laboratories Pty Ltd
GA	Galderma Australia Pty Ltd	OC	Accord Healthcare Pty Ltd
GC	GlaxoSmithKline Australia Pty Ltd	OD	Accord Healthcare Pty Ltd
GH	Amdipharm Mercury (Australia) Pty Limited	OE	Omegapharm Pty Ltd
GI	Gilead Sciences Pty Limited	OH	Orpharma Pty Ltd
GK	GlaxoSmithKline Australia Pty Ltd	OM	Colgate Oral Care
GN	Actavis Pty Ltd	ON	Orion Laboratories Pty Ltd
GO	Mylan Health Pty Ltd	OQ	Organon Pharma Pty Ltd
GQ	Generic Health Pty Ltd	OS	Otsuka Australia Pharmaceutical Pty Ltd
GT	Mylan Health Pty Ltd	OU	Oraderm Pharmaceuticals Pty Ltd
GX	Apotex Pty Ltd	OV	Organon Pharma Pty Ltd
		OW	Arrow Pharma Pty Ltd

Code	Manufacturer
PB	Pharmaco (Australia) Limited
PE	Allergan Australia Pty Limited
PF	Pfizer Australia Pty Ltd
PK	Fresenius Kabi Australia Pty Limited
PP	Petrus Pharmaceuticals Pty Ltd
PY	Procter & Gamble Pharmaceuticals Australia Pty Ltd
QH	Cortex Health Pty Ltd
QY	Pro Pharmaceuticals Group Pty Ltd
RA	Sun Pharma ANZ Pty Ltd
RB	Bio Revive Pty Ltd
RC	Reckitt Benckiser (Australia) Pty Limited
RF	Arrow Pharma Pty Ltd
RI	Dr Reddy's Laboratories (Australia) Pty Ltd
RJ	Recordati Rare Diseases Australia Pty. Ltd.
RM	Pharmacor Pty Limited
RN	Sun Pharma ANZ Pty Ltd
RO	Roche Products Pty Ltd
RW	Arrow Pharma Pty Ltd
RX	Servier Laboratories (Aust.) Pty Ltd
RZ	Dr Reddy's Laboratories (Australia) Pty Ltd
SA	SciGen (Australia) Pty Limited
SB	Nutricia Australia Pty Limited
SE	Servier Laboratories (Aust.) Pty Ltd
SG	Merck Healthcare Pty Ltd
SI	Sigma Company Limited
SN	Smith & Nephew Pty Limited
SS	SSL Australia Pty Ltd
SW	sanofi-aventis Australia Pty Ltd
SY	Bayer Australia Ltd
SZ	Sandoz Pty Ltd
TB	Teva Pharma Australia Pty Limited
TD	STADA Pharmaceuticals Australia Pty Limited
TF	Te Arai BioFarma Limited
TI	Teva Pharma Australia Pty Limited
TK	Takeda Pharmaceuticals Australia Pty Ltd
TM	Technipro PulmoMed Pty Ltd
TQ	Terumo BCT Australia Pty Limited
TT	Theramex Australia Pty Ltd
TW	Apotex Pty Ltd
TX	Apotex Pty Ltd
TY	Apotex Pty Ltd
UC	UCB Australia Proprietary Limited
UJ	Upjohn Australia Pty Ltd
UL	Bausch & Lomb (Australia) Pty Ltd
UM	Unomedical Pty Ltd
UN	Unilever Australia Limited
UO	Bausch & Lomb (Australia) Pty Ltd
UR	Camurus Pty Ltd
VE	AbbVie Pty Ltd
VF	VitaFlo Australia Pty Limited
VI	ViiV Healthcare Pty Ltd
VL	Vifor Pharma Pty Limited
VO	Avallon Pharmaceuticals Pty Limited
VR	Vertex Pharmaceuticals (Australia) Pty Ltd
VZ	Sanofi-aventis Healthcare Pty Ltd
WA	sanofi-aventis Australia Pty Ltd
WC	Wockhardt Bio Pty Ltd
XA	Pharmaxis Ltd
XC	Southern Cross Pharma Pty Ltd
XH	MS Health Pty Ltd
XI	Alexion Pharmaceuticals Australasia Pty Ltd
XM	The Medicines Company (Australia) Pty Limited
XT	Arrotex Pharmaceuticals Pty Ltd
YB	Bayport Brands Pty Ltd
YC	Cipla Australia Pty Ltd
YN	Mayne Pharma International Pty Ltd
YT	Mayne Products Pty Ltd
ZA	AstraZeneca Pty Ltd
ZE	Seekwell Pty Ltd

Code	Manufacturer
ZP	Medis Pharma Pty Ltd
ZX	Zenex Pharmaceuticals Pty Ltd

Generic/Proprietary Index

ABATACEPT	350, 353	ADALIMUMAB398, 400, 402, 404, 405, 409, 411, 413, 415, 418, 420, 423, 428, 433, 436, 439, 442, 446, 452, 457, 462	
ABEMACICLIB.....	302	ADAPALENE + BENZOYL PEROXIDE.....	185
Abilify Maintena(LU)	857	Adaptic 2012(KI).....	1056
Abilify(OS).....	857, 858	Addos XR 30(RW).....	137
ABIRATERONE	346	Addos XR 60(RW).....	137
Abisart 150(AL).....	145	Adefin 10(AF).....	137
Abisart 300(AL).....	145	Adefin 20(AF).....	137
Abisart 75(AL).....	145	Adefin XL 30 (AF).....	137
Abisart HCTZ 150/12.5(AL)	148	Adefin XL 60 (AF).....	137
Abisart HCTZ 300/12.5(AL)	148	Adenuric(FK).....	768
Abisart HCTZ 300/25(AL)	148	Adesan HCT 16/12.5(AF).....	147
Abstral(FK).....	999, 1000, 1001	Adesan HCT 32/12.5(AF).....	147
Abyraz (AF).....	857, 858	Adesan HCT 32/25(AF).....	147
ACALABRUTINIB	314	Adesan(AF).....	144, 145
ACAMPROSATE	897	ADRENALINE (EPINEPHRINE)	
Acamprosate Mylan(AL)	897	.CARDIOVASCULAR SYSTEM.....	125
ACARBOSE.....	83	.Prescriber Bag	19
Acarbose Mylan(AF).....	83	.RESPIRATORY SYSTEM	919
Accomin Adult Tonic(PF)	1019	Adrenaline Jr Mylan(AF)	
Accupril(PF).....	140	.CARDIOVASCULAR SYSTEM.....	126
Accuretic 10/12.5mg(PF).....	143	.RESPIRATORY SYSTEM	919
Accuretic 20/12.5mg(PF).....	143	Adrenaline Mylan(AF)	
ACETAZOLAMIDE	929	.CARDIOVASCULAR SYSTEM.....	126
Acefec(AL).....	138, 139	.RESPIRATORY SYSTEM	919
ACETIC ACID + HYDROXYQUINOLINE + RICINOLEIC ACID.....	1027	ADT Booster(CS).....	19, 253
ACICLOVIR		ADVAGRAF XL(LQ).....	755
.ANTIINFECTIVES FOR SYSTEMIC USE	246	Advantan (Fatty)(LO).....	180, 181, 182
.SENSORY ORGANS	922	Advantan(LO).....	180, 181, 182
Aciclovir AN(ED).....	246	Aeron 250(AL).....	916
Aciclovir APOTEX (TY)	246	Aeron 500(AL).....	916
Aciclovir APOTEX(TY).....	246	AFATINIB	288, 289
Aciclovir GH(GQ).....	246	Afinitor(NV).....	307, 308, 309, 310
Aciclovir Sandoz (HX).....	246	AFLIBERCEPT	933, 934, 935, 936
Aciclovir Sandoz(HX).....	246	Aggrastat(AS).....	112
Aci-Jel(CU).....	1027	Airomir Autohaler(IL)	903
Acimax Tablets(AL)	37, 38, 39	Akynzeo(MF).....	46
ACITRETIN.....	175	ALBENDAZOLE	900
AcVision(DZ).....	922	Albey Bee Venom(DE).....	951
Aclasta(HX).....	771	Albey Paper Wasp Venom(DE).....	952
ACLIDINIUM.....	916	Albey Yellow Jacket Venom(DE).....	952
ACLIDINIUM + FORMOTEROL (EFORMOTEROL).....	911	Aldactone(PF).....	131, 132
Aclor 125(MQ).....	226	Aldara Pump(IL).....	190, 1026
Aclor 250(MQ).....	226, 227	Aldara(IL).....	190, 1026
Acpio 15(RF).....	84	Aldiq(AF).....	190, 1026
Acpio 30(RF).....	84	Aldomet(AS).....	128
Acpio 45(RF).....	84	Alecensa(RO).....	297
ACQUIN (RF).....	140	ALECTINIB	296
Acquin Aspen 10 (RW)	140	Alendro Once Weekly(RW)	769
Acquin Aspen 20(RW)	140	Alendrobell 70mg(GQ).....	769
Acquin Aspen 5(RW)	140	ALENDRONATE	768
ACQUIN(RF).....	140	ALENDRONATE + COLECALCIFEROL	772, 1032
Acris Combi(AF).....	774, 1033	ALENDRONATE + COLECALCIFEROL (&) CALCIUM CARBONATE	773
Acris Once-a-Month(AF).....	770	ALENDRONATE PLUS D3 70mg/140ug APOTEX(GX) 772, 1033	
Acris Once-a-Week(AF).....	770, 1032	ALENDRONATE PLUS D3 70mg/70ug APOTEX(GX) 773, 1033	
Actaze (RW)	84	Alendronate Plus D3 Sandoz(SZ)	772, 773
Actemra ACTPen(RO) ...697, 699, 700, 702, 705, 706, 708, 713, 716, 720, 725, 730		Alendronate plus D3-DRLA (RZ)	772, 773, 1033
Actemra Subcutaneous Injection(RO) ...697, 699, 700, 702, 704, 706, 708, 713, 716, 720, 725, 730		Alendronate Sandoz (SZ)	769
Actiq(TB).....	999, 1000, 1001	Alfamino Junior(NT).....	972, 973
Actisorb Plus MAP105(KI)	1044	Alfamino(NT)	965, 967
Actonel EC(TT).....	770, 1032	Alfaré(NT).....	969
Actonel Once-a-Month (TT)	770	ALFUZOSIN	1030
Actonel(TT).....	770, 771, 1032	Algisite M 66000519(SN).....	1044
Actos(TK).....	84	Algisite M 66000520(SN).....	1044
Actrapid Penfill 3 mL(NO)	62	Algisite M 66000521(SN).....	1044
Actrapid(NO).....	61		

<i>Algivon Plus CR4225(DJ)</i>	1045
<i>Algivon Plus Ribbon & Probe CR4231(DJ)</i>	1045
<i>Alkeran(AS)</i>	253
<i>Allegron(RW)</i>	870
<i>Allereze(AF)</i>	1038
<i>Allevyn 66007637(SN)</i>	1046
<i>Allevyn Adhesive 66000044(SN)</i>	1046
<i>Allevyn Ag Adhesive 66800073(SN)</i>	1048
<i>Allevyn Ag Adhesive 66800075(SN)</i>	1048
<i>Allevyn Ag Adhesive 66800078(SN)</i>	1049
<i>Allevyn Ag Gentle Border 66800460(SN)</i>	1048
<i>Allevyn Ag Gentle Border 66800461(SN)</i>	1049
<i>Allevyn Ag Gentle Border 66800462(SN)</i>	1049
<i>Allevyn Ag Non-Adhesive 66800086(SN)</i>	1048
<i>Allevyn Gentle 66800248(SN)</i>	1047
<i>Allevyn Gentle Border 66800269(SN)</i>	1047
<i>Allevyn Gentle Border 66800270(SN)</i>	1047
<i>Allevyn Life 66801067(SN)</i>	1046
<i>Allevyn Life 66801068(SN)</i>	1046
<i>Allevyn Life 66801069(SN)</i>	1046
<i>Allevyn Life 66801070(SN)</i>	1046
<i>Allevyn Life Non-Bordered 66801748(SN)</i>	1046
<i>Allevyn Life Non-Bordered 66801749(SN)</i>	1046
<i>Allevyn Thin 66047576(SN)</i>	1051
<i>Allevyn Thin 66047578(SN)</i>	1051
<i>Allmercap(LM)</i>	255
<i>ALLOPURINOL</i>	768
<i>Allopurinol APOTEX(GX)</i>	768
<i>Allopurinol Sandoz (SZ)</i>	768
<i>Allosig(RF)</i>	768
<i>ALOGLIPTIN</i>	84
<i>ALOGLIPTIN + METFORMIN</i>	65
<i>Aloxi(MF)</i>	48
<i>Alpha Keri Bath Oil(MT)</i>	1023
<i>Alpha Keri Lotion(MT)</i>	1023
<i>AlphaClav Duo Forte(AF)</i>	221, 222, 223
<i>AlphaClav Duo(AF)</i>	222, 223
<i>Alphagan P 1.5(AG)</i>	927, 928
<i>Alphagan(AG)</i>	928
<i>Alphamox 125(AF)</i>	215
<i>Alphamox 250(AF)</i>	215, 216, 217
<i>Alphamox 500(AF)</i>	215, 216, 217
<i>Alphapress 25(AF)</i>	130
<i>Alphapress 50(AF)</i>	129
<i>Alprax 0.5(AS)</i>	866
<i>Alprax 1(AS)</i>	866
<i>ALPRAZOLAM</i>	865
<i>Alprim(AF)</i>	229
<i>ALPROSTADIL</i>	1027
<i>Alunbrig(TK)</i>	297
<i>Alvesco 160(EU)</i>	915
<i>Alvesco 80(EU)</i>	915
<i>Alzene(AF)</i>	1038
<i>AMANTADINE</i>	845
<i>Amaryl(SW)</i>	65
<i>Amcal Mag-A(IG)</i>	1020
<i>AMCLAVOX DUO 500/125 (RW)</i>	222, 223
<i>AMCLAVOX DUO FORTE 875/125 (RW)</i>	221, 222, 223
<i>Amdipharm Mercury (Australia) Pty Limited(GH)</i>	
CARDIOVASCULAR SYSTEM	133
GENITO URINARY SYSTEM AND SEX HORMONES	
.....	203
<i>AMILORIDE + HYDROCHLOROTHIAZIDE</i>	132
<i>AMILOXYN (RF)</i>	215, 216, 217
<i>AMINO ACID FORMULA SUPPLEMENTED WITH</i>	
PREBIOTICS, PROBIOTICS AND LONG CHAIN	
POLYUNSATURATED FATTY ACIDS.....	973, 974, 975
<i>AMINO ACID FORMULA WITH CARBOHYDRATE</i>	
WITHOUT PHENYLALANINE	976

<i>AMINO ACID FORMULA WITH CARBOHYDRATE,</i>	
VITAMINS, MINERALS AND TRACE ELEMENTS	
WITHOUT PHENYLALANINE	976
<i>AMINO ACID FORMULA WITH FAT, CARBOHYDRATE</i>	
WITHOUT METHIONINE.....	976
<i>AMINO ACID FORMULA WITH FAT, CARBOHYDRATE</i>	
WITHOUT PHENYLALANINE	976
<i>AMINO ACID FORMULA WITH FAT, CARBOHYDRATE</i>	
WITHOUT PHENYLALANINE AND TYROSINE.....	977
<i>AMINO ACID FORMULA WITH FAT, CARBOHYDRATE</i>	
WITHOUT VALINE, LEUCINE AND ISOLEUCINE.....	977
<i>AMINO ACID FORMULA WITH FAT, CARBOHYDRATE,</i>	
VITAMINS, MINERALS AND TRACE ELEMENTS	
WITHOUT METHIONINE AND SUPPLEMENTED WITH	
DOCOSAHEXAENOIC ACID.....	977
<i>AMINO ACID FORMULA WITH FAT, CARBOHYDRATE,</i>	
VITAMINS, MINERALS AND TRACE ELEMENTS	
WITHOUT PHENYLALANINE	977
<i>AMINO ACID FORMULA WITH FAT, CARBOHYDRATE,</i>	
VITAMINS, MINERALS AND TRACE ELEMENTS	
WITHOUT PHENYLALANINE AND TYROSINE, AND	
SUPPLEMENTED WITH DOCOSAHEXAENOIC ACID	
.....	977
<i>AMINO ACID FORMULA WITH FAT, CARBOHYDRATE,</i>	
VITAMINS, MINERALS AND TRACE ELEMENTS,	
WITHOUT PHENYLALANINE AND TYROSINE.....	977
<i>AMINO ACID FORMULA WITH FAT, CARBOHYDRATE,</i>	
VITAMINS, MINERALS, TRACE ELEMENTS AND	
MEDIUM CHAIN TRIGLYCERIDES	971, 972
<i>AMINO ACID FORMULA WITH VITAMINS AND</i>	
MINERALS WITHOUT LYSINE AND LOW IN	
TRYPTOPHAN	977, 978
<i>AMINO ACID FORMULA WITH VITAMINS AND</i>	
MINERALS WITHOUT METHIONINE	978, 979
<i>AMINO ACID FORMULA WITH VITAMINS AND</i>	
MINERALS WITHOUT METHIONINE, THREONINE	
AND VALINE AND LOW IN ISOLEUCINE.....	979
<i>AMINO ACID FORMULA WITH VITAMINS AND</i>	
MINERALS WITHOUT PHENYLALANINE	979, 980
<i>AMINO ACID FORMULA WITH VITAMINS AND</i>	
MINERALS WITHOUT PHENYLALANINE AND	
TYROSINE	981
<i>AMINO ACID FORMULA WITH VITAMINS AND</i>	
MINERALS WITHOUT VALINE, LEUCINE AND	
ISOLEUCINE	982, 983
<i>AMINO ACID FORMULA WITH VITAMINS AND</i>	
MINERALS WITHOUT VALINE, LEUCINE AND	
ISOLEUCINE WITH FAT, CARBOHYDRATE AND	
TRACE ELEMENTS AND SUPPLEMENTED WITH	
DOCOSAHEXAENOIC ACID.....	983
<i>AMINO ACID FORMULA WITH VITAMINS AND</i>	
MINERALS, LOW PHENYLALANINE AND	
SUPPLEMENTED WITH DOCOSAHEXAENOIC ACID	
AND ARACHIDONIC ACID.....	983
<i>AMINO ACID FORMULA WITH VITAMINS, MINERALS</i>	
AND LONG CHAIN POLYUNSATURATED FATTY	
ACIDS WITHOUT PHENYLALANINE.....	983
<i>AMINO ACID FORMULA WITHOUT PHENYLALANINE</i>	
<i>AMINO ACID FORMULA WITHOUT VALINE, LEUCINE</i>	
AND ISOLEUCINE.....	984
<i>AMINO ACID SYNTHETIC FORMULA</i>	957, 958
<i>AMINO ACID SYNTHETIC FORMULA SUPPLEMENTED</i>	
WITH LONG CHAIN POLYUNSATURATED FATTY	
ACIDS.....	959, 960
<i>AMINO ACID SYNTHETIC FORMULA SUPPLEMENTED</i>	
WITH LONG CHAIN POLYUNSATURATED FATTY	
ACIDS AND MEDIUM CHAIN TRIGLYCERIDES	962,
963, 964, 965	
<i>AMIODARONE</i>	124
<i>Amiodarone Sandoz(SZ)</i>	125

Amipride 400(RW)	857
Amira 150(AF)	875
Amira 300(AF)	875
AMISULPRIDE	856
Amisulpride 100 Winthrop(WA)	857
Amisulpride 200 Winthrop(WA)	857
Amisulpride 400 Winthrop (WA)	857
Amisulpride AN (EA)	857
Amisulpride AN(EA)	857
Amisulpride Sandoz (SZ)	857
Amisulpride Sandoz Pharma (HX)	857
Amisulpride Sandoz Pharma(HX)	857
Amisulpride Sandoz(SZ)	857
AMITRIPTYLINE	869
Amitriptyline Alphapharm 10(AL)	869
Amitriptyline Alphapharm 25(AL)	869
Amitriptyline Alphapharm 50(AL)	869
Amitriptyline Lupin (GQ)	869
Amlo 10(RW)	136
Amlo 5(RW)	136
AMLODIPINE	136
AMLODIPINE + ATORVASTATIN	172
AMLODIPINE + VALSARTAN	150
AMLODIPINE + VALSARTAN + HYDROCHLOROTHIAZIDE	151
Amlodipine Amneal (EF)	136
Amlodipine AN(EA)	136
Amlodipine APOTEX (GX)	136
Amlodipine GH(GQ)	136
Amlodipine Sandoz (SZ)	136
AMOROLFINE	1022
AMOXICILLIN	215, 216, 217
AMOXICILLIN + CLAVULANIC ACID	221, 222
AMOXICLAV AMNEAL 500/125(ED)	222, 223
AMOXICLAV AMNEAL 875/125(ED)	221, 222, 223
Amoxil Forte(AS)	215
Amoxil(AS)	215, 216, 217
Amoxycillin AN(EA)	215, 216, 217
Amoxycillin generichealth 500 (GQ)	215, 216, 217
Amoxycillin Sandoz (SZ)	215, 216, 217
Amoxycillin Sandoz(BG)	216
Amoxycillin Sandoz(SZ)	215, 216, 217
Amoxycillin/Clavulanic Acid 500/125 APOTEX (TY)	222, 223
Amoxyclav AN 500/125(EA)	222, 223
Amoxyclav AN 875/125 (EA)	221, 222, 223
AmoxyClav generichealth 875/125(HQ)	221, 222, 223
AMPHOTERICIN B	29
AMPICILLIN	217
Ampicyn(AF)	217
Anafranil 25(SZ)	869
Anamorph(RW)	784, 785, 788
Anandron(SW)	345
Anaprox 550(IX)	765, 997
Anastrol(AS)	345
ANASTROZOLE	345
Anastrozole FBM (FO)	345
Anastrozole GH(GQ)	345
Anastrozole Sandoz (SZ)	345
Andepra(EL)	876
Andriol Testocaps(MK)	196
Androcur(BN)	
.ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS	344
.GENITO URINARY SYSTEM AND SEX HORMONES	201
Androcur-100(BN)	
.ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS	344

.GENITO URINARY SYSTEM AND SEX HORMONES	201
Androderm(TB)	195
AndroForte 5(LX)	195
Angiomax(XM)	112
Anoro Ellipta 62.5/25(GK)	914
Anpec 80(AF)	137
Antenex 2(AF)	866, 1004
Antenex 5(AF)	866, 1004
ANTERONE 100(RW)	
.ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS	344
.GENITO URINARY SYSTEM AND SEX HORMONES	201
ANTERONE 50(RW)	
.ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS	344
.GENITO URINARY SYSTEM AND SEX HORMONES	201
Anthel 125(AF)	900
Anthel 250(AF)	900
Antroquoril(OV)	178
Anusol(JT)	1022
Apidra SoloStar(SW)	61
Apidra(AV)	61
Apidra(SW)	61
APIXABAN	114, 115, 116
APO- Hydroxychloroquine(TX)	766, 767
APO- Paracetamol/Codeine 500/30(TX)	815, 816, 817
APO-Acamprosate (TX)	897
APO-Aciclovir (TX)	246
APO-Aciclovir(TX)	246
APO-Alendronate (TX)	769
APO-Amiodarone (TX)	125
APO-Amisulpride (TX)	857
APO-Amisulpride(TX)	857
APO-Amitriptyline 10(TX)	869
APO-Amitriptyline 25(TX)	869
APO-Amitriptyline 50(TX)	869
APO-Amlodipine(TX)	136
APO-Amoxycillin (TX)	215, 216, 217
APO-Amoxycillin and Clavulanic Acid (TX)	221, 222, 223
APO-Amoxycillin(TX)	215, 216, 217
APO-Amoxycillin/ Clavulanic Acid 500/125 (TX)	222, 223
APO-Anastrozole(TX)	345
APO-Aripiprazole(TX)	857, 858
APO-Atenolol(TX)	134
APO-Atomoxetine(TX)	880
APO-Atorvastatin(TX)	153, 154
APO-Azathioprine(TX)	755
APO-Azithromycin(TX)	
.Repatriation Pharmaceutical Benefits Scheme	1031
APO-Azithromycin(TX)	
.ANTIINFECTIVES FOR SYSTEMIC USE	230
.SENSORY ORGANS	921
APO-Baclofen(TX)	767
APO-Bicalutamide(TX)	344
APO-Bimatoprost(TX)	931
APO-Bisoprolol(TX)	134
APOC-5FU(TX)	1031
APO-Cabergoline(TX)	192
APO-Calcitriol(TX)	
.ALIMENTARY TRACT AND METABOLISM	102
.MUSCULO-SKELETAL SYSTEM	774
APO-Candesartan (TX)	144, 145
APO-Candesartan HCTZ 16/12.5 (TX)	147
APO-Candesartan HCTZ 32/12.5 (TX)	147
APO-Candesartan HCTZ 32/25 (TX)	147
APO-Carvedilol(TX)	136
APO-Cefaclor (TX)	226, 227

APO-Cefaclor CD(TX)	226	APO-Latanoprost(TX)	932
APO-Celecoxib(TX)	766	APO-Latanoprost/Timolol 0.05/5(TX)	932
APO-Cephalexin(TX)	223, 224, 225	APO-Letrozole(TX)	346
APO-Ciprofloxacin(TX)	235, 236	APO-Levetiracetam(TX)	837, 838
APO-Citalopram(TX)	870, 871	APO-Levodopa/Carbidopa(TX)	843
APO-Clindamycin(TX)	232, 233	APO-Lisinopril(TX)	139
APO-Clomipramine(TX)	869	APO-Lurasidone(TX)	850, 851
APO-Clonidine(TX)	128	APO-MACROGOL plus ELECTROLYTES (TX)	53, 995
APO-Clopidogrel(TX)	109, 110, 1020	APO-Mebeverine(TX)	1016
APO-Clopidogrel/Aspirin 75/100(TX)	111	APO-Meloxicam(TX)	761, 762
APO-Clotrimazole 3 Day Cream(TX)	1027	APO-Memantine(TX)	892, 893, 894
APO-Clotrimazole 6 Day Cream(TX)	1027	APO-Metformin 1000(TX)	64
APO-Cyproterone (TX)	201, 344	APO-Metformin 500(TX)	63
APO-Desvenlafaxine MR(TX)	875, 876	APO-Metformin 850(TX)	63
APO-Diazepam (TX)	866, 1004	APO-Metformin XR 1000(TX)	64
APO-Diazepam(TX)	866, 1004	APO-Metformin XR 500(TX)	64
APO-Diclofenac(TX)	760, 996	APO-Metoclopramide(TX)	44
APO-Donepezil(TX)	885, 886	APO-Metoprolol(TX)	135
APO-Dorzolamide(TX)	930	Apomine Intermittent(PF)	845
APO-Dorzolamide/Timolol 20/5(TX)	930	Apomine Solution for Infusion(PF)	845
APO-Doxycycline(TX)	213, 214	APO-Mirtazapine(TX)	877
APO-Duloxetine (TX)	876	APO-Moclobemide (TX)	875
APO-Dutasteride(TX)	203, 1030	APO-Modafinil(TX)	884
APO-Enalapril (TX)	138, 139	APO-Montelukast(TX)	920, 921
APO-Eplerenone(TX)	131	APOMORPHINE	845
APO-Escitalopram(TX)	871	APO-Moxonidine(TX)	129
APO-Esomeprazole(TY)	30, 31, 32, 33, 34	APO-Mupirocin(TX)	1024
APO-Exemestane(TX)	346	APO-Mycophenolate(TX)	370
APO-Ezetimibe(TX)	167	APO-Naltrexone(TX)	898
APO-Ezetimibe/Simvastatin 10/10(TX)	172	APO-Nebivolol(TX)	135
APO-Ezetimibe/Simvastatin 10/20(TX)	172	APO-Nicorandil(TX)	127
APO-Ezetimibe/Simvastatin 10/40(TX)	171	APO-Nifedipine XR(TX)	137
APO-Ezetimibe/Simvastatin 10/80(TX)	171	APO-Nitrofurantoin(TX)	238
APO-Famciclovir(TX)	247, 248	APO-Norfloxacin(TX)	236
APO-Fenofibrate(TX)	158, 159	APO-Olanzapine ODT(TX)	853, 854
APO-Fentanyl(TX)	806, 807, 809, 810, 812	APO-Olanzapine(TX)	852, 853
APO-Finasteride(TX)	1031	APO-Olmesartan(TX)	146
APO-Flecainide(TX)	124	APO-Olmesartan/Amlodipine/HCTZ 20/5/12.5(TX)	152
APO-Flucloxacillin(TX)	220, 221	APO-Olmesartan/Amlodipine/HCTZ 40/10/12.5(TX)	152
APO-Fluconazole(TX)	240	APO-Olmesartan/Amlodipine/HCTZ 40/10/25(TX)	152
APO-Fluoxetine(TX)	873	APO-Olmesartan/Amlodipine/HCTZ 40/5/12.5 tablet(TX)	152
APO-Fluvoxamine(TX)	873	APO-Olmesartan/Amlodipine/HCTZ 40/5/25 tablet(TX)	152
APO-Fosinopril HCTZ 20/12.5(TX)	142	APO-Olmesartan/HCTZ 20/12.5(TX)	148
APO-Fosinopril(TX)	139	APO-Olmesartan/HCTZ 40/12.5(TX)	148
APO-Frusemide(TX)	130, 131	APO-Olmesartan/HCTZ 40/25(TX)	148
APO-Gabapentin(TX)		APO-Omeprazole(TX)	36, 37, 38, 39
Repatriation Pharmaceutical Benefits Scheme	1034, 1035	APO-Ondansetron ODT(TX)	47, 48
APO-Gabapentin(TX)		APO-Ondansetron(TX)	46
NERVOUS SYSTEM	834	APO-Oxazepam(TX)	866, 867, 1004
APO-Galantamine MR(TX)	887, 888, 889	APO-Pantoprazole(TX)	39, 40, 41
APO-Gliclazide MR(TX)	64	APO-Paracetamol(TX)	823, 1034
APO-Gliclazide(TX)	64	APO-Paroxetine(TX)	873
APO-Glimepiride(TX)	65	APO-Perindopril Arginine(TX)	139, 140
APOHEALTH Macrogol with Electrolytes(GX)	53, 995	APO-Perindopril Arginine/Amlodipine 10/10(TX)	144
APOHEALTH Osteo Relief Paracetamol 665 mg(TX)	824, 1003	APO-Perindopril Arginine/Amlodipine 10/5(TX)	143
APO-Ibuprofen 400(TX)	763, 997	APO-Perindopril Arginine/Amlodipine 5/10(TX)	143
APO-Imiquimod (TX)	190, 1026	APO-Perindopril Arginine/Amlodipine 5/5(TX)	143
APO-Indapamide SR(TX)	130	APO-Perindopril(TX)	139, 140
APO-Ipratropium (TX)	916	APO-PINDOL(DZ)	133
APO-Irbesartan (TX)	145	APO-Piroxicam(TX)	762, 763
APO-Irbesartan HCTZ (TX)	148	APO-Pramipexole ER(TX)	847
APO-Isosorbide Mononitrate(TX)	126	APO-Pramipexole(TX)	846, 847
APO-Isotretinoin(TX)	186	APO-Pravastatin(TX)	154, 155
APO-Itraconazole(TX)	242	APO-Prazosin(TX)	129
APO-Ivabradine(TX)	127	APO-Primidone(LM)	828
APO-Lamotrigine(TX)	836, 837	APO-Prochlorperazine(TX)	50
APO-Lansoprazole ODT(TX)	34, 35, 36	APO-Propranolol(TX)	133
APO-Lansoprazole(TX)	34, 35, 36	APO-Quetiapine XR(TX)	855, 856
		APO-Quetiapine(TX)	855, 856

APO-Quinapril (TX).....	140	Ardix Lurasidone (RX)	850, 851
APO-Quinapril(TX).....	140	Arginaid Extra(NT).....	1039
APO-Rabeprazole(TX).....	41, 42, 43	Arginaid(NT).....	1039
APO-Raloxifene(TX).....	776	Arginine 2000(VF).....	984
APO-Ramipril(TX).....	140, 141	Arginine 500(VF).....	984
APO-Ranitidine(TX).....	30	Arginine 5000(VF).....	984
APO-Riluzole(TX).....	898	ARGININE WITH CARBOHYDRATE	984
APO-Risedronate (TX).....	770, 1032	Arianna 1 (AF).....	345
APO-Risedronate(TX).....	770	Aricept(PF).....	885, 886
APO-Risperidone(TX).....	859, 860, 861, 862, 863, 864, 865	Aridon 10(RW).....	885, 886
APO-Rizatriptan(TX).....	826	Aridon 5(RW).....	885, 886
APO-Rosuvastatin(TX).....	155, 156, 157	Aridon APN 10 (RF).....	885, 886
APO-Roxithromycin(TX).....	231, 232	Aridon APN 5 (RF).....	885, 886
Aporyl(TX).....	1022	Arimidex(AP).....	345
APO-Salbutamol(TX).....	22, 903	ARIPIRAZOLE	857
APO-Sertraline(TX).....	873, 874	Aripiprazole AN (EA)	857, 858
APO-Sildenafil(TX).....	1028, 1029	Aripiprazole generichealth(HQ).....	857, 858
APO-Simvastatin(TX).....	157, 158	Aripiprazole GH (GQ).....	857, 858
APO-Sotalol(TX).....	125	Aripiprazole Sandoz(SZ).....	857, 858
APO-Sumatriptan(TX).....	826, 827	Aristocort 0.02%(AS).....	176
APO-Telmisartan HCTZ 40/12.5(TX).....	149	Arixtra(AS).....	120
APO-Telmisartan HCTZ 80/12.5(TX).....	149	ARMODAFINIL	878
APO-Telmisartan HCTZ 80/25(TX).....	149	Arnuiity Ellipta(GK).....	915
APO-Telmisartan(TX).....	146	Aromasin(PF).....	346
APO-Temazepam(TX).....	868, 1005	Aropax(AS).....	873
APO-Temozolomide(TX).....	254, 255	Arrow Pharma Pty Ltd(RW).....	245
APO-Terbinafine(TX).....	174, 1023	Artane(RW).....	842
APO-TERIFLUNOMIDE(TX).....	373	ARTEMETHER + LUMEFANTRINE.....	899
APOTEX-Pioglitazone (TX).....	84	Arthrexin(AF).....	761, 996
APO-Topiramate(TX).....	840, 841	Artige(NM).....	881
APO-Tramadol SR(TX).....	822	Asacol(EU).....	57
APO-Tramadol(TX).....	819, 821	Asartan HCT 16/12.5(DO).....	147
APO-Tranexamic Acid(TX).....	121	Asartan HCT 32/12.5(DO).....	147
APO-Ursodeoxycholic acid(TX).....	51	Asartan HCT 32/25(DO).....	147
APO-Valaciclovir(TX).....	249	ASENAPINE.....	852
APO-Venlafaxine XR(TX).....	878	Asmol 2.5 uni-dose(AF).....	22, 903
APO-Ziprasidone(TX).....	851	Asmol 5 uni-dose(AF).....	22, 903
APO-Zolmitriptan(TX).....	827	Asmol CFC-Free with dose counter(AF).....	22, 902
APO-Zonisamide(TX).....	841, 842	Asmol CFC-free(AL).....	22, 902
APO-Zopiclone(TX).....	1036	Aspalgin 40(GO).....	1033
APRACLOINIDINE	927	Aspecillin VK(AF).....	218, 219
APREMILAST	358	Aspen Methadone Syrup(AS) Palliative Care.....	1002
APREPITANT	48	Aspen Pharma Pty Ltd(AS) NERVOUS SYSTEM	778, 779, 880
Aprepitant APOTEX(TX).....	49	ASPIRIN	109, 1020
Aptamil Gold+ Pepti-Junior(NU).....	968	ASPIRIN + CODEINE.....	1033
APX-Clopidogrel/Aspirin 75/100 (TY).....	111	Astrix(YN).....	1020
APX-Gabapentin (GX).....	834	Atacand Plus 16/12.5(AP).....	147
APX-Meloxicam (TY).....	761, 762	Atacand Plus 32/12.5(AP).....	147
APX-Metformin (TY).....	63, 64	Atacand Plus 32/25(AP).....	147
APX-Paracetamol/Codeine (TY).....	815, 816, 817	Atacand(AP).....	144, 145
APX-Quetiapine XR (TY).....	855, 856	Ataris 10(AF).....	369
APX-Quetiapine XR(TY).....	856	Ataris 20(AF).....	369, 370
Aquacare H.P.(AG).....	1023	Aectura Breezhaler(NV).....	911
Aquacel 403770(CC).....	1053	ATENOLOL	134
Aquacel Ag 403708(CC).....	1053	Atenolol Amneal (EF).....	134
Aquacel Ag 403710(CC).....	1053	Atenolol GH (GQ).....	134
Aquacel Ag 403771(CC).....	1053	Atenolol Sandoz(SZ).....	134
Aquacel Extra 420672(CC).....	1053	Atenolol-AFT(AE).....	134
Aquacel Extra 420673(CC).....	1053	Atenolol-GA(ED).....	134
Aquacel Foam Adhesive(CC).....	1053	ATOMERRA (RW).....	880
Aquacel Foam Non-Adhesive(CC).....	1053	ATOMOXETINE	879
Aquasun Lotion SPF 18(PF).....	1024	Atomoxetine Amneal(EA).....	880
Arabloc(AV).....	369, 370	Atomoxetine Sandoz (SZ).....	880
ARACHIDONIC ACID AND DOCOSAHEXAENOIC ACID WITH CARBOHYDRATE	984	Atorvachol (RF).....	153, 154
Aratac 100(AF).....	125	ATORVASTATIN	153
Aratac 200(AF).....	125	Atorvastatin Amneal(EF).....	153, 154
Arava (SW).....	369, 370	Atorvastatin GH (GQ).....	153, 154
Arazil (AF).....	885, 886	Atorvastatin SZ(HX).....	153, 154
ARDIX GLICLAZIDE 60mg MR(RX).....	64		

ATOVAQUONE	899	Azathioprine Sandoz(SZ)	755
ATOVAQUONE + PROGUANIL	899	Azilect(TB)	848
Atozet(AF)	170	AZITHROMYCIN	
Atrauman 499513(HR)	1055	ANTIINFECTIVES FOR SYSTEMIC USE	230
Atrauman Ag 499572(HR)	1057	Repatriation Pharmaceutical Benefits Scheme	1031
ATROPINE SULFATE		SENSORY ORGANS	921
ALIMENTARY TRACT AND METABOLISM	43	Azithromycin Mylan (AF)	230, 921
Prescriber Bag	19	Azithromycin Mylan(AF)	
SENSORY ORGANS	933	Repatriation Pharmaceutical Benefits Scheme	1031
Atropt(AS)	933	Azithromycin Sandoz (SZ)	
Atrovent Adult(BY)	916	Repatriation Pharmaceutical Benefits Scheme	1031
Atrovent Nasal Aqueous(VZ)	1037	Azithromycin Sandoz(SZ)	
Atrovent Nasal Forte(VZ)	1037	ANTIINFECTIVES FOR SYSTEMIC USE	230
Atrovent(BY)	916	SENSORY ORGANS	921
Aubagio (GZ)	373	Azithromycin-GA (EA)	1031
Augmentin Duo 400(AS)	222	Azopt(NV)	929
Augmentin Duo forte(AS)	221, 222, 223	BACLOFEN	767
Augmentin Duo(AS)	222, 223	Bactigras 7457(SN)	1049
AURANOFIN	767	Bactroban(GK)	
Auro-Amlodipine 10 (DO)	136	Repatriation Pharmaceutical Benefits Scheme	1024
Auro-Amlodipine 5 (DO)	136	Bactroban(GK)	
Auro-Famciclovir 125 (DO)	247	RESPIRATORY SYSTEM	902
Auro-Famciclovir 250 (DO)	247	BALSALAZIDE	57
Auro-Famciclovir 500 (DO)	248	BANDAGE ABSORBENT WOOL	1039
Auro-Finasteride(DO)	1030	BANDAGE CALICO	1039
Auro-Lisinopril 10 (DO)	139	BANDAGE COMPRESSION	1039, 1040
Auro-Lisinopril 20 (DO)	139	BANDAGE RETENTION COHESIVE HEAVY	1040
Auro-Lisinopril 5 (DO)	139	BANDAGE RETENTION COHESIVE LIGHT	1040
Auro-Montelukast Tabs 4 (DO)	920	BANDAGE RETENTION COTTON CREPE	1041
Auro-Montelukast Tabs 5 (DO)	921	BANDAGE TUBULAR	1041
Aurorix 300 mg(GO)	875	BANDAGE TUBULAR FINGER	1042
Aurorix(GO)	875	BANDAGE TUBULAR LIGHT WEIGHT	1042
Auro-Sertraline 100 (DO)	874	BANDAGE TUBULAR LONG STOCKING	1042
Auro-Sertraline 100(DO)	874	BANDAGE TUBULAR SHORT STOCKING	1042
Auro-Sertraline 50 (DO)	873	BANDAGE ZINC PASTE	1042, 1043
Auro-Sertraline 50(DO)	874	Barbloc 5(AF)	133
Ausfam 20(RW)	29	BARICITINIB	359, 362
Ausfam 40(RW)	29	basecal 200(VF)	984
Ausgem(RW)	159	Baxter Healthcare Pty Ltd(BX)	1021
Ausran (RW)	30	BECLOMETASONE	914
Austrapen(AL)	217	BECLOMETASONE + FORMOTEROL (EFORMOTEROL)	904
AVANAFIL	1028	Bemfola(FX)	199, 200
Avanza(AL)	877	BEMOTRIZINOL + OCTOCRYLENE + DIETHYLAMINO	
Avapro HCT 150/12.5(AV)	148	HYDROXYBENZOYL HEXYL BENZOATE + TITANIUM	
Avapro HCT 300/12.5(AV)	148	DIOXIDE	1023
Avapro HCT 300/25(AV)	148	BenPen(CS)	19, 217, 218
Avapro(AV)	145	BENZATHINE BENZYL PENICILLIN	19, 217
Avodart(GK)	203, 1030	BENZATROPINE	19, 842
Avonex(BD)	348	Benzatropine Injection(FF)	19, 842
AVSARTAN HCT 150/12.5(RF)	148	Benztrop(FF)	842
AVSARTAN HCT 300/12.5(RF)	148	BENZYDAMINE	29, 994
AVSARTAN HCT 300/25(RF)	148	BENZYL PENICILLIN	19, 217
AVSARTAN(RF)	145	Beprol 10 (DO)	134
Axit 15 (AF)	877	Beprol 2.5 (DO)	134
Axit 30 (AF)	877	Beprol 5 (DO)	134
Axit 45 (AF)	877	Betadine Antiseptic Liquid(SW)	1025
AXITINIB	313	Betaferon(BN)	349
Axotide Accuhaler(GC)	915	BETAINE	103
Axotide Junior Accuhaler(GC)	915	BETAINE + POLYAMINOPROPYL BIGUANIDE	1043
Axotide Junior(GC)	915	Betaloc(AP)	135
Axotide(GC)	915	BETAMETHASONE ACETATE + BETAMETHASONE	
Aylide 1 (AF)	65	SODIUM PHOSPHATE	206
Aylide 2 (AF)	65	BETAMETHASONE DIPROPIONATE	176, 177, 178
Aylide 3 (AF)	65	BETAMETHASONE VALERATE	178, 179, 1025
Aylide 4 (AF)	65	Betaquik(VF)	956
Azapin (RW)	755	Betavit(PP)	102, 1019
Azarga(NV)	929, 930	BETAXOLOL	930
AZATHIOPRINE	755	BETHANECHOL	894
Azathioprine AN(EA)	755	Betnovate 1/2(AS)	178, 179, 180
Azathioprine GH (GQ)	755		

<i>Betnovate 1/5(AS)</i>	178	<i>Breo Ellipta 200/25(GK)</i>	909
<i>Betnovate(AS)</i>	1025	<i>Bretaris Genuair(FK)</i>	916
<i>Betoptic(NV)</i>	930, 931	<i>Brevinor(PF)</i>	193
<i>BetoQuin(NM)</i>	930, 931	<i>Brevinor-1(PF)</i>	193
<i>Bi ELIGARD CP(MF)</i>	342	<i>BREXPIRAZOLE</i>	858
<i>Biatain Adhesive 3420(CT)</i>	1051	<i>Bricanyl Turbuhaler(AP)</i>	904
<i>Biatain Adhesive 3423(CT)</i>	1050	<i>Bricanyl(AP)</i>	919
<i>Biatain Ag 9622(CT)</i>	1056	<i>BRIGATINIB</i>	297
<i>Biatain Ag 9632(CT)</i>	1056	<i>Brilinta(AP)</i>	111
<i>Biatain Non-adhesive 3410(CT)</i>	1051	<i>Brimica Genuair(FK)</i>	912
<i>Biatain Non-adhesive 3413(CT)</i>	1051	<i>BRIMONIDINE</i>	927, 928
<i>Biassig (AV)</i>	231, 232	<i>BRIMONIDINE + TIMOLOL</i>	928
<i>Bicalox (ER)</i>	344	<i>BRINZOLAMIDE</i>	929
<i>BICALUTAMIDE</i>	344	<i>BRINZOLAMIDE + BRIMONIDINE</i>	929
<i>BICARBONATE</i>	202	<i>BRINZOLAMIDE + TIMOLOL</i>	929, 930
<i>BICARBONATE + CITRIC ACID + TARTARIC ACID</i> ..	1029	<i>BrinzoQuin(NM)</i>	929
<i>Bicard 10(RW)</i>	134	<i>BRIVARACETAM</i>	832, 833
<i>Bicard 2.5(RW)</i>	134	<i>Briviact(UC)</i>	832, 833, 834
<i>Bicard 5(RW)</i>	134	<i>BROMAZEPAM</i>	1035
<i>Bicillin L-A(PF)</i>	19, 217	<i>BROMOCRIPTINE</i>	
<i>Bicor(AL)</i>	134	.GENITO URINARY SYSTEM AND SEX HORMONES	191
<i>Biltricide(BN)</i>	900	.NERVOUS SYSTEM	846
<i>BIMATOPROST</i>	931	<i>Brufen(GO)</i>	763, 997
<i>BIMATOPROST + TIMOLOL</i>	931, 932	<i>BSN 2902165(BV)</i>	1057
<i>Bimatoprost Sandoz (SZ)</i>	931	<i>BTC Amlodipine (JB)</i>	136
<i>Bimtop(AF)</i>	931	<i>BTC Clopidogrel(JB)</i>	109, 110, 1020
<i>BINIMETINIB</i>	300	<i>BTC Fluoxetine(JB)</i>	873
<i>Bion Tears(AQ)</i>	945	<i>BTC Lercanidipine(JB)</i>	137
BISACODYL		<i>BTC Perindopril(JB)</i>	139, 140
.ALIMENTARY TRACT AND METABOLISM	51, 53	<i>BTC PREGABALIN(JB)</i>	824
.Palliative Care	994, 995	<i>BTC Rosuvastatin(JB)</i>	155, 156, 157
.Repatriation Pharmaceutical Benefits Scheme	1016	<i>Budamax Aqueous(JT)</i>	1037
<i>Bisalax(AS)</i>	54, 995	<i>Budenofalk(FD)</i>	56
<i>BISOPROLOL</i>	134	BUDESONIDE	
<i>Bisoprolol AN (EA)</i>	134	.ALIMENTARY TRACT AND METABOLISM	56
<i>Bisoprolol generichealth(GQ)</i>	134	.Repatriation Pharmaceutical Benefits Scheme	1037
<i>Bisoprolol Sandoz (SZ)</i>	134	.RESPIRATORY SYSTEM	914, 915
<i>Bispro 10(AF)</i>	134	BUDESONIDE + FORMOTEROL (EFORMOTEROL) ..	904,
<i>Bispro 2.5(AF)</i>	134	905, 906, 907, 908	
<i>Bispro 5(AF)</i>	134	<i>Bupredermal(TX)</i>	814, 815, 1002, 1003
BIVALIRUDIN	112	BUPRENORPHINE	
<i>Bivalirudin APOTEX (TX)</i>	112	.NERVOUS SYSTEM	813
<i>Blooms the Chemist Amlodipine(IB)</i>	136	.Palliative Care	1002
<i>Blooms the Chemist Atorvastatin (IB)</i>	153, 154	<i>Buprenorphine Sandoz (SZ)</i>	814, 815, 1002, 1003
<i>Blooms the Chemist Candesartan (IB)</i>	144	BUPROPION	894
<i>Blooms the Chemist Candesartan HCTZ 16/12.5 (IB)</i> ..	147	<i>Burprenorphine Sandoz (SZ)</i>	814, 1002
<i>Blooms the Chemist Candesartan HCTZ 32/12.5 (IB)</i> ..	147	<i>Buscopan(VZ)</i>	20, 994, 1016
<i>Blooms the Chemist Candesartan HCTZ 32/25 (IB)</i>	147	<i>Buspar(AS)</i>	1035
<i>Blooms the Chemist Candesartan(IB)</i>	144, 145	BUSPIRONE	1035
<i>Blooms the Chemist Celecoxib (IB)</i>	766	BUSULFAN	253
<i>Blooms the Chemist Clopidogrel (IB)</i>	109, 110, 1020	<i>Bydureon(AP)</i>	94
<i>Blooms the Chemist Escitalopram (IB)</i>	871	<i>Byetta 10 microgram(AP)</i>	95
<i>Blooms The Chemist Ezetimibe (IB)</i>	167	<i>Byetta 5 microgram(AP)</i>	95
<i>Blooms the Chemist Fenofibrate (IB)</i>	158, 159	<i>Cabaser(PF)</i>	846
<i>Blooms the Chemist Fluoxetine (IB)</i>	873	CABERGOLINE	
<i>Blooms the Chemist Irbesartan (IB)</i>	145	.GENITO URINARY SYSTEM AND SEX HORMONES	192
<i>Blooms the Chemist Irbesartan HCTZ 150/12.5 (IB)</i>	148	.NERVOUS SYSTEM	846
<i>Blooms the Chemist Irbesartan HCTZ 300/12.5 (IB)</i>	148	<i>Cabometyx(IS)</i>	318
<i>Blooms the Chemist Irbesartan HCTZ 300/25 (IB)</i>	148	CABOZANTINIB	317, 318
<i>Blooms the Chemist Metformin XR 1000 (IB)</i>	64	CADEXOMER-IODINE	1043
<i>Blooms the Chemist Metformin XR 500 (IB)</i>	64	<i>Cadivast 10/10(AF)</i>	172
<i>Blooms the Chemist Perindopril (IB)</i>	139, 140	<i>Cadivast 10/20(AF)</i>	172
<i>Blooms the Chemist Rosuvastatin (IB)</i>	155, 156, 157	<i>Cadivast 10/40(AF)</i>	172
<i>Blooms the Chemist Venlafaxine XR (IB)</i>	878	<i>Cadivast 10/80(AF)</i>	172
<i>Bondronat(IX)</i>	769	<i>Cadivast 5/10(AF)</i>	172
<i>Braffovi(FB)</i>	295	<i>Cadivast 5/20(AF)</i>	172
<i>Braltus(AF)</i>	918	<i>Cadivast 5/40(AF)</i>	172
<i>Brenzys(RF)</i>	502, 503, 505, 507, 512, 514, 517, 519, 520,	<i>Cadivast 5/80(AF)</i>	173
522, 525, 532, 536, 539, 544, 549, 554, 562			
<i>Breo Ellipta 100/25(GK)</i>	909		

<i>Caduet 10/10(AS)</i>	172	<i>CarboFLEX 403202(CC)</i>	1043
<i>Caduet 10/20(AS)</i>	172	<i>CarboFLEX 403204(CC)</i>	1044
<i>Caduet 10/40(AS)</i>	172	<i>Carbohydrate Free Mixture(SB)</i>	988
<i>Caduet 10/80(AS)</i>	172	CARBOHYDRATE, FAT, VITAMINS, MINERALS AND	
<i>Caduet 5/40(AS)</i>	172	TRACE ELEMENTS	984
<i>Caduet 5/80(AS)</i>	173	CARBOHYDRATES, FAT, VITAMINS, MINERALS, TRACE	
<i>Cal-500(PP)</i>	103, 1019	ELEMENTS AND SUPPLEMENTED WITH	
<i>CAL-600(PP)</i>	1019	ARACHIDONIC ACID AND DOCOSAHEXAENOIC ACID	
<i>Cal-care 600 mg (CR)</i>	1019		984
CALCIPOTRIOL + BETAMETHASONE DIPROPIONATE		CARBOMER-974P	943
.....	175	CARBOMER-980	943
<i>Calcipotriol/Betamethasone Sandoz 50/500(SZ)</i>	175	<i>Cardasa(AF)</i>	1020
<i>Calciprox (ER)</i>	102, 774	<i>Cardiprin 100(RC)</i>	1020
<i>Calci-Tab 600(AE)</i>	103	<i>Cardizem CD(SW)</i>	138
CALCITONIN SALMON (SALCATONIN)	212	<i>Cardizem(SW)</i>	138
CALCITRIOL		<i>Cardol (AF)</i>	125
..ALIMENTARY TRACT AND METABOLISM	102	<i>Carlevent(TX)</i>	844
..MUSCULO-SKELETAL SYSTEM	774	CARMELLOSE SODIUM	943, 944
<i>Calcitriol AN(EA)</i>		CARMELLOSE SODIUM + GLYCEROL	945
..ALIMENTARY TRACT AND METABOLISM	102	CARMUSTINE	253
..MUSCULO-SKELETAL SYSTEM	774	<i>Cartia(AS)</i>	1020
CALCIUM	103, 1019	CARVEDILOL	135
<i>Calindamin (RW)</i>	232, 233	<i>Carvedilol AN (EA)</i>	136
CALITROL (ED)		<i>Carvedilol Sandoz(SZ)</i>	136
..ALIMENTARY TRACT AND METABOLISM	102	<i>Carvidol (RF)</i>	136
..MUSCULO-SKELETAL SYSTEM	774	<i>Carvidol(RF)</i>	136
<i>Calquence(AP)</i>	315	<i>Catapres 100 (IX)</i>	128
<i>Calutex(AS)</i>	344	<i>Catapres(IX)</i>	128
<i>Camino Pro Bettermilk(QH)</i>	986	<i>Caverject Impulse(PF)</i>	1028
<i>Camino Pro Complete(QH)</i>	986	<i>Cavicare 4563(SN)</i>	1046
<i>Campral(AF)</i>	897	<i>Cavstat (AF)</i>	155, 156, 157
CANDESAN (RF)	144, 145	<i>Ceclor CD(AL)</i>	226
CANDESAN COMBI 16/12.5(RF)	147	<i>Ceclor(AL)</i>	226, 227
CANDESAN COMBI 32/12.5(RF)	147	CEFACLOR	226
CANDESAN COMBI 32/25(RF)	147	<i>Cefaclor GH (GQ)</i>	226
CANDESAN(RF)	144	CEFALEXIN	223, 224, 225
CANDESARTAN	144	<i>Cefalexin Sandoz (SZ)</i>	223, 224, 225
CANDESARTAN + HYDROCHLOROTHIAZIDE	147	<i>Cefalexin Sandoz(SZ)</i>	223
<i>Candesartan AN (EA)</i>	144	CEFAZOLIN	225, 226
<i>Candesartan AN(EA)</i>	144, 145	<i>Cefazolin-AFT(AE)</i>	225, 226
<i>Candesartan Aspen 16(RW)</i>	145	CEFEPIME	229
<i>Candesartan Aspen 32 (RW)</i>	144	<i>Cefepime Alphapharm (AF)</i>	229
<i>Candesartan Aspen 4 (RW)</i>	145	<i>Cefepime Kabi(PK)</i>	229
<i>Candesartan Aspen 8 (RW)</i>	144	<i>Cefepime-AFT(AE)</i>	229
<i>Candesartan Combi Aspen 16/12.5 (RW)</i>	147	CEFOTAXIME	227
<i>Candesartan Combi Aspen 32/12.5 (RW)</i>	147	CEFTRIAXONE	228
<i>Candesartan Combi Aspen 32/25 (RW)</i>	147	<i>Ceftriaxone Alphapharm(AF)</i>	228, 229
<i>Candesartan GH (GQ)</i>	145	<i>Ceftriaxone-AFT(AE)</i>	228, 229
<i>Candesartan GH(GQ)</i>	144	CEFUROXIME	227
<i>Candesartan HCT GH 16/12.5(GQ)</i>	147	<i>Celapram (AF)</i>	870, 871
<i>Candesartan HCT GH 32/12.5(GQ)</i>	147	<i>Celapram(AF)</i>	871
<i>Candesartan HCT GH 32/25(GQ)</i>	147	<i>Celaxib(AF)</i>	766
<i>Candesartan HCTZ AN 16/12.5(EA)</i>	147	<i>Celebrex (UJ)</i>	766
<i>Candesartan HCTZ AN 32/12.5(EA)</i>	147	CELECOXIB	765
<i>Candesartan HCTZ AN 32/25(EA)</i>	147	<i>Celecoxib AN(EA)</i>	766
<i>Candesartan Sandoz (SZ)</i>	144	<i>Celecoxib APOTEX (TY)</i>	766
<i>Candesartan Sandoz(SZ)</i>	144, 145	<i>Celecoxib BTC(JB)</i>	766
<i>Candesartan/HCT Sandoz (SZ)</i>	147	<i>Celecoxib GH (GQ)</i>	766
CAPECITABINE	256	<i>Celecoxib Sandoz(SZ)</i>	766
<i>Capecitabine Alphapharm(AF)</i>	256	<i>Celestone Chronodose(OQ)</i>	206
<i>Capecitabine Sandoz(SZ)</i>	256	<i>Celestone-M(OQ)</i>	178
<i>Capecitabine-DRLA (RZ)</i>	256	<i>Celexi (RW)</i>	766
<i>Capecitabine-DRLA(RZ)</i>	256	<i>CellCept (RO)</i>	370
<i>Capoten(RW)</i>	138	<i>CellCept(RO)</i>	370
CAPTOPRIL	138	<i>Cellufresh(AG)</i>	944
<i>Captopril Sandoz(SZ)</i>	138	<i>Celluvisc(AG)</i>	944
CARBAMAZEPINE	830	<i>Cephalex 500(CR)</i>	223, 224, 225
<i>Carbamazepine Sandoz(SZ)</i>	830	<i>Cephalexin AN (EA)</i>	223, 224, 225
CARBAMIDE PEROXIDE	1038	<i>Cephalexin AN(EA)</i>	223, 224, 225
CARBIMAZOLE	210	<i>Cephalexin generichealth(GQ)</i>	223, 224, 225

<i>Cephazolin Alphapharm(AF)</i>	225, 226
<i>Ceptolate(AF)</i>	370
<i>CERITINIB</i>	297
<i>Certican(NV)</i>	368
<i>CERTOLIZUMAB PEGOL</i>	469, 471, 473, 475, 478, 479, 484, 487, 489, 493, 496
<i>Cerumol(UN)</i>	1038
<i>CETIRIZINE</i>	1038
<i>C-Flox 250 (AL)</i>	236
<i>C-Flox 500 (AL)</i>	235
<i>C-Flox 750 (AL)</i>	235
<i>Champix(PF)</i>	896, 897
<i>Chem mart Meloxicam (CH)</i>	762
<i>Chem mart Metformin(CH)</i>	64
<i>Chemists' Own Laxative with Senna(RW)</i>	1017
<i>Chemists' Own Macrogol with Electrolytes(RW)</i>	53, 995
<i>CHLORAMBUCIL</i>	253
<i>CHLORAMPHENICOL</i>	921
<i>CHLORHEXIDINE</i>	1016
<i>CHLORPROMAZINE</i>	19, 849
<i>Chlorsig(AS)</i>	921
<i>CHLORTALIDONE</i>	130
<i>Chlorvescent(AS)</i>	103
<i>Cholstat 10(AF)</i>	155
<i>Cholstat 20 (AF)</i>	155
<i>Cholstat 40 (AF)</i>	154, 155
<i>Cialis(LY)</i>	1029
<i>CICLESONIDE</i>	915
<i>CICLOSPORIN</i>	754
<i>Cifran(RA)</i>	235
<i>Cilamox (AL)</i>	215, 216, 217
<i>Cilamox(AL)</i>	216, 217
<i>Cilicaine V(AF)</i>	218
<i>Cilicaine VK(AF)</i>	218, 219
<i>Cilicaine(AF)</i>	19, 219
<i>Cilopam-S(ER)</i>	871
<i>CiloQuin(NM)</i>	923
<i>Ciloxan(NV)</i>	
.SENSORY ORGANS	923, 951
<i>Cimzia(UC)</i>	469, 471, 473, 475, 478, 479, 484, 486, 489, 493, 496, 500
<i>CINACALCET</i>	212
<i>Cinacalcet Mylan(AF)</i>	212
<i>Cipla Gefitinib(LR)</i>	291
<i>Cipla Icatibant(LR)</i>	123
<i>CIPLA IMATINIB ADULT(LR)</i> 264, 265, 266, 267, 268, 269, 270, 272, 273, 275, 276, 277, 278, 279, 280	
<i>CIPLA MELOXICAM 15(LR)</i>	762
<i>CIPLA MELOXICAM 7.5(LR)</i>	761
<i>Cipla Pregabalin (LR)</i>	824
<i>Cipla Tadalafil (LR)</i>	1029
<i>Cipramil(LU)</i>	871
<i>CIPROFLOXACIN</i>	
.ANTIINFECTIVES FOR SYSTEMIC USE	234, 235
.SENSORY ORGANS	923, 950
<i>Ciprofloxacin AN (EA)</i>	235
<i>Ciprofloxacin AN(EA)</i>	235
<i>Ciprofloxacin GH(HQ)</i>	235
<i>Ciprofloxacin Sandoz (SZ)</i>	235
<i>Ciprofloxacin Sandoz(SZ)</i>	236
<i>Ciprol 250 (RW)</i>	236
<i>Ciprol 500(RW)</i>	235
<i>Ciprol 750(RW)</i>	235
<i>CITALOPRAM</i>	870
<i>Citalopram Actavis (EA)</i>	871
<i>Citalopram Actavis(ED)</i>	871
<i>Citalopram AN (EA)</i>	871
<i>Citalopram AN(EF)</i>	871
<i>Citalopram Sandoz(SZ)</i>	871

<i>CITRIC ACID + LAURYL SULFOACETATE SODIUM + SORBITOL</i>	54, 995, 1017
<i>CITRULLINE</i>	984
<i>Citrulline 1000(VF)</i>	985
<i>Citrulline Easy(OH)</i>	985
<i>CITRULLINE WITH CARBOHYDRATE</i>	985
<i>CLADRIBINE</i>	
.ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS	367
<i>Clarac(ED)</i>	230
<i>Claratyne(BN)</i>	1038
<i>Clarithro 250 (RW)</i>	230
<i>CLARITHROMYCIN</i>	230, 231
<i>Clarithromycin AN(EA)</i>	230
<i>Clarithromycin Sandoz (SZ)</i>	230
<i>Clexane Safety-Lock(AV)</i>	108, 109
<i>Climara 100(BN)</i>	197
<i>Climara 25(BN)</i>	197
<i>Climara 50(BN)</i>	197
<i>Climara 75(BN)</i>	197
<i>CLINDAMYCIN</i>	232
<i>Clindamycin BNM(BZ)</i>	232, 233
<i>Clindamycin LU (LV)</i>	232, 233
<i>Clindamycin(AF)</i>	232, 233
<i>Clobemix(ED)</i>	875
<i>CLOBETASOL</i>	185
<i>Clobex(GA)</i>	185
<i>Clofen 10 (AF)</i>	767
<i>Clofen 25 (AF)</i>	767
<i>Clomid(SW)</i>	201
<i>CLOMIFENE</i>	201
<i>CLOMIPRAMINE</i>	869
<i>Clonac 25 (RW)</i>	760, 996
<i>Clonac 50 (RW)</i>	760, 996
<i>CLONAZEPAM</i>	19, 829, 1003, 1004
<i>Clonea 3 Day Cream (AF)</i>	1027
<i>Clonea 6 Day Cream(AF)</i>	1027
<i>Clonea(AF)</i>	1022
<i>CLONIDINE</i>	128
<i>CLOPIDOGREL</i>	109, 1020
<i>CLOPIDOGREL + ASPIRIN</i>	110
<i>Clopidogrel AN(EA)</i>	109, 110, 1020
<i>Clopidogrel APOTEX (GX)</i>	109, 110, 1020
<i>Clopidogrel GH (GQ)</i>	109, 110
<i>Clopidogrel GH(GQ)</i>	1020
<i>Clopidogrel Sandoz (SZ)</i>	109, 110
<i>Clopidogrel Sandoz Pharma (HX)</i>	1020
<i>Clopidogrel Sandoz Pharma(HX)</i>	109, 110
<i>Clopidogrel Winthrop (WA)</i>	109, 110
<i>Clopidogrel Winthrop plus aspirin(WA)</i>	111
<i>Clopidogrel/Aspirin Actavis 75/100(EA)</i>	111
<i>CLOPIDOGREL/ASPIRIN AN 75/100 (ED)</i>	111
<i>Clopidogrel-GA(EA)</i>	109, 110
<i>Clopixol Depot(LU)</i>	851
<i>CLOTTRIMAZOLE</i>	
.Repatriation Pharmaceutical Benefits Scheme	1022, 1027
<i>Clovix 75(RW)</i>	109, 110
<i>COAL TAR SOLUTION + PHENOL + PRECIPITATED SULFUR</i>	1024
<i>Cobal-B12(JU)</i>	122
<i>Coban 1584(MM)</i>	1040
<i>Coban 2(MM)</i>	1040
<i>COBIMETINIB</i>	300, 301
<i>Codalgin Forte(AF)</i>	815, 816, 817
<i>Codapane Forte 500/30 (AL)</i>	815, 816, 817
<i>CODEINE</i>	778
<i>Co-Diovan 160/12.5(NV)</i>	149
<i>Co-Diovan 160/25(NV)</i>	149
<i>Co-Diovan 320/12.5(NV)</i>	150

Co-Diovan 320/25(NV)	150	Crosuva 20(RW).....	155, 156
Co-Diovan 80/12.5(NV)	149	Crosuva 40(RW).....	156, 157
Colaxsen (AS).....	1017	Crosuva 5(RW).....	155, 156
Colazide(PK).....	57	Crysanal(IY).....	765, 997
COLCHICINE.....	768	Cubitan(SB).....	1039
Colese (AF).....	1016	Curam Duo 500/125(SZ)	222, 223
COLESTYRAMINE	159	Curam Duo Forte 875/125(SZ).....	221, 222, 223
Colgout(AS).....	768	Curam Duo(SZ)	222
Colifoam(GO).....	56	Curam(SZ).....	222
Colofac(GO).....	1016	Curity 4112(KE).....	1049
Coloxyl 50(AS).....	1016	Cutifilm Plus 36361370(SN)	1046
Coloxyl with Senna(LN)	1017	Cutifilm Plus 36361371(SN)	1045
Combigan(AG).....	928	Cutilin Non-Stick Wound Pad 36361374(SN)	1056
Comfarol Forte(SZ).....	815, 816, 817	Cutilin Non-Stick Wound Pad 36361375(SN)	1056
Comfeel Plus Pressure Relieving 3350(CT)	1053	Cutinova Hydro 66047441(SN)	1051
Comfeel Plus Pressure Relieving 3353(CT)	1053	Cutinova Hydro 66047443(SN)	1051
Comfeel Plus Transparent 3530(CT)	1052	Cyclonex(ZX).....	253
Comfeel Plus Transparent 3533(CT)	1052	CYCLOPHOSPHAMIDE.....	253
Comfeel Plus Transparent 3536(CT)	1052	Cyclosporin Sandoz(SZ).....	754
Comfeel Plus Ulcer Dressing 3110(CT)	1053	Cyklokapron (PF)	121
Comfeel Purilon Gel 3900(CT).....	1054	Cymbalta(LY)	876
Comfeel SeaSorb Dressing 3705(CT)	1045	Cyprone 100(AF)	
Comfeel SeaSorb Dressing 3710(CT)	1045	.ANTINEOPLASTIC AND IMMUNOMODULATING	
Comfeel SeaSorb Filler 3740(CT).....	1044	AGENTS	344
Comprilan 01027-00(BV)	1039	.GENITO URINARY SYSTEM AND SEX HORMONES	
Comtan(SZ).....	849		201
Concerta(JC)	882	Cyprone 50(AL)	
Copaxone(TB)	350	.ANTINEOPLASTIC AND IMMUNOMODULATING	
CoPlavix (SW)	111	AGENTS	344
Coralan (SE)	127	.GENITO URINARY SYSTEM AND SEX HORMONES	
Corbeton 40(AF).....	133		201
Cordarone X 100 (SW)	125	Cyprostat-100 (SY)	
Cordarone X 200 (SW)	125	.ANTINEOPLASTIC AND IMMUNOMODULATING	
Cordilox 180 SR(GT)	137	AGENTS	344
Cordilox SR(GT)	138	.GENITO URINARY SYSTEM AND SEX HORMONES	
Cortate(AS).....	207		201
Cortic-DS 1%(AS).....	1025	CYPROTERONE	
Cortic-DS 1%(LN).....	176	.ANTINEOPLASTIC AND IMMUNOMODULATING	
CORTISONE	207	AGENTS	344
Cortival 1/2(LN).....	178, 179	.GENITO URINARY SYSTEM AND SEX HORMONES	
Cortival 1/5(LN).....	178		201
Cosamide 50 (AF).....	344	Cyproterone AN (EA)	
Cosdor (AF)	930	.ANTINEOPLASTIC AND IMMUNOMODULATING	
Co-Senna (PP)	1017	AGENTS	344
Cosentyx(NV) 650, 652, 654, 655, 657, 658, 661, 664, 667,		.GENITO URINARY SYSTEM AND SEX HORMONES	
671, 674, 678, 684			201
Cosopt(MF).....	930	Cyproterone AN(EA)	
Cosudex(AP)	344	.ANTINEOPLASTIC AND IMMUNOMODULATING	
Cotellic(RO).....	301	AGENTS	344
Coumadin(GO)	107	.GENITO URINARY SYSTEM AND SEX HORMONES	
Coveram 10/10(SE).....	144		201
Coveram 10/5(SE).....	143	Cyproterone Sandoz (HX)	
Coveram 5/10(SE).....	143	.ANTINEOPLASTIC AND IMMUNOMODULATING	
Coveram 5/5(SE).....	143	AGENTS	344
Coversyl 10mg(SE).....	140	.GENITO URINARY SYSTEM AND SEX HORMONES	
Coversyl 2.5mg(SE).....	139		201
Coversyl 5mg(SE).....	140	Cyproterone Sandoz(HX)	
Coversyl Plus 5mg/1.25mg(SE).....	142	.ANTINEOPLASTIC AND IMMUNOMODULATING	
Coversyl Plus LD 2.5mg/0.625mg(SE)	142	AGENTS	344
Cozavan(AF).....	146	.GENITO URINARY SYSTEM AND SEX HORMONES	
Creon 10,000(GO).....	60		201
Creon 25,000(GO).....	60	Cystadane(RJ)	104
Creon 40,000(GO).....	60	DABIGATRAN	112, 113
Creon Micro(GO).....	60	DABRAFENIB	293, 294
Crestor(AP).....	155, 156, 157	Daivobet (LO).....	175
CRIZOTINIB	298	Daivobet 50/500 gel(LO)	175
CROMOGLYCATE		Daktarin Tincture(JT).....	173
.Repatiation Pharmaceutical Benefits Scheme	1037	Daktarin(JT).....	173
.RESPIRATORY SYSTEM.....	918	Dalacin C (PF).....	232, 233
Crosuva 10(RW).....	156, 157	DALTEPARIN SODIUM.....	107

<i>Dantrium</i> (PF)	767, 768	<i>Diapride 3</i> (RW).....	65
DANTROLENE	767	<i>Diapride 4</i> (RW).....	65
<i>Daonil</i> (SW)	64	<i>Diasp SR</i> (RW).....	111
DAPAGLIFLOZIN	96, 97	<i>Diastix</i> (DX)	955
DAPAGLIFLOZIN + METFORMIN	66	DIAZEPAM	866, 1004
<i>Dapa-Tabs</i> (AF)	130	<i>Diazepam Elixir</i> (ON).....	866
DAPSONE		<i>Dibenyline</i> (GH).....	
.ANTIINFECTIVES FOR SYSTEMIC USE	245	.CARDIOVASCULAR SYSTEM.....	133
.DERMATOLOGICALS	190	.GENITO URINARY SYSTEM AND SEX HORMONES	202
DASATINIB	257, 259, 261		202
<i>DBL Cefotaxime</i> (PF).....	227, 228	<i>Dicarz</i> (AF)	136
<i>DBL Heparin Sodium</i> (PF)	109	DICLOFENAC	
<i>DBL Methotrexate</i> (PF)	255	.MUSCULO-SKELETAL SYSTEM	760
<i>DBL Morphine Sulfate Pentahydrate</i> (PF) 21, 784, 785, 787, 791		.Palliative Care.....	996
<i>DBL Promethazine Hydrochloride</i> (PF)	21	.Repatriation Pharmaceutical Benefits Scheme	1026
.Prescriber Bag	21	<i>Diclofenac Amneal</i> (ED)	760, 996
<i>DBL Promethazine Hydrochloride</i> (PF)		<i>Diclofenac AN</i> (EA)	760, 996
.ALIMENTARY TRACT AND METABOLISM	50	<i>Diclofenac Sandoz</i> (SZ).....	760, 996
<i>DBL Promethazine Hydrochloride</i> (PF)		DICLOXACILLIN	219
.RESPIRATORY SYSTEM.....	921	<i>Dicloxacillin Mylan 250</i> (AL)	219
<i>DBL Tobramycin</i> (PF)	234	<i>Dicloxacillin Mylan 500</i> (AL)	219
<i>Debrisoft Lolly</i> (LC)	1057	<i>Difflam</i> (IL).....	29, 994
<i>Debrisoft</i> (LC)	1057	<i>Diffucan</i> (PF)	240
DEGARELIX	347	<i>Diflucan</i> (PF)	240, 241
DENOSUMAB.....	774, 775	DIGOXIN	123
<i>Denpax</i> (AF)	805, 807, 808, 810, 812	<i>Dilantin Infatabs</i> (UJ)	828
<i>Densate 70</i> (DO).....	769	<i>Dilantin Sodium</i> (UJ).....	828
<i>Depo-Medrol</i> (PF)	208	<i>Dilantin</i> (UJ).....	828
<i>Depo-Nisolone</i> (FZ)	208	<i>Dilart HCT 160/12.5</i> (AF).....	149
<i>Depo-Provera</i> (PF)	194	<i>Dilart HCT 160/25</i> (AF).....	149
<i>Depo-Ralovera</i> (FZ)	194	<i>Dilart HCT 320/12.5</i> (AF).....	150
<i>Deptran 10</i> (AF)	870	<i>Dilart HCT 320/25</i> (AF).....	150
<i>Deptran 25</i> (AF)	870	<i>Dilart HCT 80/12.5</i> (AF).....	149
<i>Deptran 50</i> (AF)	870	<i>Dilart</i> (AF).....	146
<i>Deralin 10</i> (AF)	133	<i>Dilatrend 12.5</i> (PB).....	136
<i>Deralin 160</i> (AF)	133	<i>Dilatrend 25</i> (PB).....	136
<i>Deralin 40</i> (AF)	133	<i>Dilatrend 6.25</i> (PB).....	136
<i>Dermatane</i> (ER).....	186	<i>Dilaudid</i> (MF).....	780, 781, 782
<i>Dermatane</i> (ER).....	186	<i>Dilaudid-HP</i> (MF).....	782
<i>Desfax</i> (AF).....	875, 876	DILTIAZEM	138
DESMOPRESSIN	204, 205	<i>Diltiazem AN</i> (EA).....	138
<i>Desmopressin Acetate</i> (Medsurge)(DZ)	204, 205	<i>Diltiazem Sandoz CD</i> (SZ)	138
DESVEN (RW).....	875, 876	DIMETHYL FUMARATE.....	755, 756
DESVENLAFAXINE.....	875	<i>Dimirel</i> (AV)	65
<i>Desvenlafaxine Actavis</i> (EA).....	875, 876	<i>Diovan</i> (NV).....	146
<i>Desvenlafaxine GH XR</i> (GQ).....	875, 876	<i>Dipentum</i> (IX)	59
<i>Desvenlafaxine Sandoz</i> (SZ)	875, 876	DIPHENOXYLATE + ATROPINE SULFATE	56
DEXAMETHASONE		<i>Diphereline</i> (IS)	342
.SENSORY ORGANS	923, 924, 925	DIPHThERIA + TETANUS VACCINE	19, 253
.SYSTEMIC HORMONAL PREPARATIONS, EXCL. SEX HORMONES AND INSULINS	207	<i>Diprosone</i> (OQ)	176, 177, 178
DEXAMFETAMINE	880	DIPYRIDAMOLE + ASPIRIN	111
<i>Dexmethsone</i> (AS)	207	DISOPYRAMIDE	124
DEXTRAN-70 + HYPROMELLOSE	945	<i>Distaph 250</i> (AF)	219
<i>Diabex 1000</i> (AL).....	64	<i>Distaph 500</i> (AF)	219
<i>Diabex 850</i> (AL).....	64	<i>Dithiazide</i> (FF).....	130
<i>Diabex XR 1000</i> (AL).....	64	<i>Ditropan</i> (SW).....	202
<i>Diabex XR 500</i> (AL).....	64	<i>Dizole 100</i> (AF)	240
<i>Diabex</i> (AL).....	63	<i>Dizole 200</i> (AF)	240
<i>Diacomit</i> (EU)	839, 840	<i>Dizole 50</i> (AF)	240
<i>Diaformin 1000</i> (AF)	64	<i>docomega</i> (VF).....	985
<i>Diaformin 850</i> (AF)	64	DOCOSAHEXAENOIC ACID WITH CARBOHYDRATE	985
<i>Diaformin XR 1000</i> (AF).....	64	DOCUSATE	
<i>Diaformin</i> (AF)	63	.Repatriation Pharmaceutical Benefits Scheme	1016, 1038
<i>Dialamine</i> (SB).....	985	DOCUSATE + SENNOSIDE B	1017
<i>Diamicron 60mg MR</i> (SE)	64	<i>Dolapril 0.5</i> (RW).....	141
<i>Diamox</i> (RW)	929	<i>Dolapril 1</i> (RW).....	141
<i>Diapride 1</i> (RW)	65	<i>Dolapril 2</i> (RW).....	141
<i>Diapride 2</i> (RW)	65	<i>Dolapril 4</i> (RW).....	141
		DOMPERIDONE	44

DONEPEZIL	884, 885	DRESSING NON-ADHERENT ABSORBENT	1056
<i>Donepezil AN(EA)</i>	885, 886	DRESSING NON-ADHERENT WITH SILICONE	1056
<i>Donepezil GH(HQ)</i>	885, 886	DRESSING TULLE NON-ADHERENT PRIMARY WOUND	
<i>Donepezil Sandoz (SZ)</i>	885, 886	CONTACT LAYER PARAFFIN	1056
<i>Donepezil-DRLA (RZ)</i>	885, 886	DRESSING WITH SILVER	1056, 1057
<i>Doryx(YN)</i>	213, 214	<i>Drixine(BN)</i>	1037
DORZOLAMIDE	930	DRY PSYLLIUM HUSK	1017
DORZOLAMIDE + TIMOLOL	930	DULAGLUTIDE	92
<i>Dostinex (PF)</i>	192	<i>Dulcolax(VZ)</i>	52, 994, 1016
DOSULEPIN (DOTHIEPIN)	869	<i>Duloxecor(CR)</i>	876
<i>Dosulepin Mylan(AL)</i>	869	DULOXETINE	876
<i>Dothep 25(AF)</i>	869	<i>Duloxetine AN (EA)</i>	876
<i>Dothep 75(AF)</i>	869	<i>Duloxetine Sandoz 30 (SZ)</i>	876
<i>Doubluts(GC)</i>	203	<i>Duloxetine Sandoz 60 (SZ)</i>	876
DOXEPIN	869	<i>Duloxetine Sandoz(HX)</i>	876
<i>Doxsig(RW)</i>	213, 214	<i>Duocal(SB)</i>	990
DOXYCYCLINE	213, 214	<i>DuoCover(AV)</i>	111
<i>Doxycycline AN (EA)</i>	213, 214	<i>Duodart 500ug/400ug(GK)</i>	203, 1030
<i>Doxycycline Sandoz (HX)</i>	213, 214	<i>DuoDERM CGF 187660(CC)</i>	1052
<i>Doxycycline Sandoz(HX)</i>	213	<i>DuoDERM CGF 187662(CC)</i>	1052
<i>Doxylin 100(AF)</i>	213, 214	<i>DuoDERM Extra Thin 187955(CC)</i>	1051
<i>Doxylin 50(AF)</i>	214	<i>DuoDERM Gel 187990(CC)</i>	1054
<i>D-Penamamine(AL)</i>	767	<i>DuoDERM Gel H7987(CC)</i>	1054
DRESSING ACTIVATED CHARCOAL MALODOROUS		<i>DuoDERM Paste 187930(CC)</i>	1051
WOUND	1043	<i>Duodopa(VE)</i>	844
DRESSING ALGINATE CAVITY WOUND	1044	<i>Duofilm Solution(GK)</i>	1025
DRESSING ALGINATE SUPERFICIAL WOUND	1044	<i>DuoPlidogrel (GZ)</i>	111
DRESSING ALGINATE WITH MANUKA HONEY	1045	<i>DuoResp Spiromax(AF)</i>	905, 906, 907, 908
DRESSING FILM	1045	<i>Duotrav(NV)</i>	933
DRESSING FILM ISLAND	1045	DUPILUMAB	186
DRESSING FOAM HEAVY EXUDATE	1046	<i>Dupixent(SW)</i>	189
DRESSING FOAM MODERATE EXUDATE	1046	<i>Durafiber 66800560(SN)</i>	1050
DRESSING FOAM WITH SILICONE	1046	<i>Durafiber 66800561(SN)</i>	1050
DRESSING FOAM WITH SILICONE AND SILVER	1047	<i>Durafiber 66800563(SN)</i>	1050
DRESSING FOAM WITH SILICONE HEAVY EXUDATE		<i>Duride (AF)</i>	126
.....	1047, 1048	<i>Durogesic 100 (JC)</i>	812
DRESSING FOAM WITH SILICONE LIGHT EXUDATE		<i>Durogesic 12 (JC)</i>	806
.....	1048	<i>Durogesic 25 (JC)</i>	807
DRESSING FOAM WITH SILICONE MODERATE		<i>Durogesic 50 (JC)</i>	809
EXUDATE	1048	<i>Durogesic 75 (JC)</i>	810
DRESSING FOAM WITH SILVER	1048	DUTASTERIDE	203, 1030
DRESSING GAUZE	1049	DUTASTERIDE + TAMSULOSIN	203, 1030
DRESSING GAUZE ABSORBENT	1049	<i>Dutran 100(EA)</i>	812
DRESSING GAUZE PARAFFIN	1049	<i>Dutran 12(EA)</i>	805
DRESSING GAUZE PARAFFIN WITH CHLORHEXIDINE		<i>Dutran 25(EA)</i>	807
ACETATE	1049	<i>Dutran 50(EA)</i>	809
DRESSING GELLING FIBRE	1049, 1050	<i>Dutran 75(EA)</i>	810
DRESSING HYDROACTIVE DEBRIDEMENT	1050	<i>DYTREX 30(RW)</i>	876
DRESSING HYDROACTIVE SUPERFICIAL WOUND		<i>DYTREX 60(RW)</i>	876
HIGH EXUDATE SEMI-PERMEABLE ABSORBENT		<i>EAA Supplement(VF)</i>	985
FOAM	1050	<i>Ear Clear for Ear Wax Removal(KY)</i>	1038
DRESSING HYDROACTIVE SUPERFICIAL WOUND		<i>Easiphen(SB)</i>	980
LIGHT EXUDATE	1051	<i>Ebixa (LU)</i>	892, 893, 894
DRESSING HYDROACTIVE SUPERFICIAL WOUND		<i>Edecrin(FK)</i>	131
MODERATE EXUDATE	1051	<i>Edronax(PF)</i>	877
DRESSING HYDROCOLLOID CAVITY WOUND	1051	<i>Efexor-XR(UJ)</i>	877, 878
DRESSING HYDROCOLLOID SUPERFICIAL WOUND		<i>Efudix (IL)</i>	1031
LIGHT EXUDATE	1051, 1052	<i>Egoderm Cream(EO)</i>	1026
DRESSING HYDROCOLLOID SUPERFICIAL WOUND		<i>Egoderm Ointment(EO)</i>	1026
MODERATE EXUDATE	1052	<i>Egopsoryl-TA(EO)</i>	1024
DRESSING HYDROFIBRE ALTERNATE TO ALGINATES		<i>Elastoplast 2225(BE)</i>	1041
.....	1053	<i>Elastoplast 2226(BE)</i>	1041
DRESSING HYDROFIBRE WITH SILVER	1053	<i>Elastoplast 2227(BE)</i>	1041
DRESSING HYDROGEL	1053	<i>Elaxine SR 150 (ZP)</i>	878
DRESSING HYDROGEL AMORPHOUS	1054	<i>Elaxine SR 37.5 (ZP)</i>	877
DRESSING HYDROGEL FOAM	1054	<i>Elaxine SR 75 (ZP)</i>	878
DRESSING HYDROGEL RIBBON	1054	<i>Eldepryl(AS)</i>	849
DRESSING HYDROGEL SHEET	1054, 1055	<i>Eleanor 150/30 ED(EA)</i>	193
DRESSING HYDROPHOBIC	1055	<i>EleCare LCP(AB)</i>	960, 961
DRESSING NON ADHERENT	1055	<i>EleCare(AB)</i>	957, 958, 959

ELETRIPTAN.....	825	Eryc(YN).....	231
Eleuphrat(OV).....	176, 177, 178	ERYTHROMYCIN.....	231
Eleva 100 (AF).....	874	ERYTHROMYCIN ETHYLSUCCINATE.....	231
Eleva 100(AF).....	874	Esbriet(RO).....	758, 759, 760
Eleva 50 (AF).....	874	ESCITALOPRAM.....	871, 872
Eleva 50(AF).....	873	Escitalopram AN (EA).....	871
Elidel(GO).....	190	Escitalopram GH(HQ).....	871, 872
Eligard 1 month(MF).....	341	Escitalopram Sandoz (HX).....	871
Eligard 3 month(MF).....	341	Esipram (CF).....	872
Eligard 4 month(MF).....	341	Esipram(CF).....	871
Eligard 6 month(MF).....	341	Eskazole(AS).....	900
Eliquis(BQ).....	114, 115, 116	ESOMEPRAZOLE.....	30, 31, 32, 33
Elocon Alcohol Free(AL).....	182, 183, 184	ESOMEPRAZOLE (&) CLARITHROMYCIN (&)	
Elocon(AL).....	182, 183, 184, 185, 1025	AMOXICILLIN.....	43
Emend IV(MK).....	50	Esomeprazole ACTAVIS(EA).....	30, 31, 32, 33, 34
EMEXLON (RW).....	44	Esomeprazole Apotex (TX).....	30, 31, 32, 33, 34
EMPAGLIFLOZIN.....	98, 99	Esomeprazole GH(GQ).....	30, 31, 32, 33, 34
EMPAGLIFLOZIN + LINAGLIPTIN.....	68, 69	Esomeprazole GxP (AF).....	30, 31, 32, 33, 34
EMPAGLIFLOZIN + METFORMIN.....	69, 70	Esomeprazole Mylan(AL).....	30, 31, 32, 33, 34
E-Mycin 200(AF).....	231	Esomeprazole RBX (RA).....	30, 31, 32, 33, 34
E-Mycin 400(AF).....	231	ESOMEPRAZOLE SANDOZ Hp7(SZ).....	43
ENALAPRIL.....	138	Esomeprazole Sandoz(SZ).....	30, 31, 32, 33, 34
ENALAPRIL + HYDROCHLOROTHIAZIDE.....	142	Esomeprazole SZ (HX).....	30, 31, 32, 33, 34
Enalapril Actavis(ED).....	138, 139	Esopreze(BG).....	30, 31, 32, 33, 34
Enalapril generichealth (GQ).....	138, 139	ESPLER(RW).....	131
Enalapril Sandoz(SZ).....	138, 139	Essential Amino Acid Mix(SB).....	985
Enalapril/HCT Sandoz(SZ).....	142	ESSENTIAL AMINO ACIDS FORMULA.....	985
Enbrel (PF)	514, 517, 519, 520, 522, 525, 532, 536, 539, 544, 549, 554, 562	ESSENTIAL AMINO ACIDS FORMULA WITH MINERALS	
Enbrel(PF)	509, 514, 517, 519, 522, 525, 529, 532, 536, 539, 544, 549, 554, 562, 573	AND VITAMIN C.....	985
ENCORAFENIB.....	295	ESSENTIAL AMINO ACIDS FORMULA WITH VITAMINS	
Endep 10(AF).....	869	AND MINERALS.....	985
Endep 25(AF).....	869	Estalis continuous 50/140(SZ).....	198
Endep 50(AF).....	869	Estalis continuous 50/250(SZ).....	198
Endone(AF).....	795, 796, 799	Estalis sequi 50/140(SZ).....	199
Energivit(SB).....	984	Estalis sequi 50/250(SZ).....	199
Enerzair Breezhaler(NV).....	913	Estraderm MX 100(JU).....	197
Enidin(PE).....	928	Estraderm MX 25(JU).....	196
Enlax-XR(AF).....	878	Estraderm MX 50(JU).....	197
ENOXAPARIN SODIUM.....	108	ESTRADIOL.....	196
Enstilar(LO).....	175	ESTRADIOL (&) ESTRADIOL + DYDROGESTERONE.....	198
ENTACAPONE.....	849	ESTRADIOL + DYDROGESTERONE.....	198
ENTRECTINIB.....	318	ESTRADIOL + NORETHISTERONE ACETATE.....	198
Entresto(NV).....	152, 153	Estradot 100(SZ).....	197
ENTRIP (RW).....	869	Estradot 25(SZ).....	196
ENZALUTAMIDE.....	344	Estradot 37.5(SZ).....	197
Epclusa(GI).....	252	Estradot 50(SZ).....	197
Epiduo(GA).....	185	Estradot 75(SZ).....	197
Epilim EC(SW).....	832	ESTRIOL.....	197
Epilim Liquid(SW).....	832	ETACRYNIC ACID.....	131
Epilim Syrup(SW).....	832	ETANERCEPT500, 503, 505, 507, 509, 512, 514, 517, 520, 522, 525, 529, 532, 536, 539, 544, 549, 554, 562	
Epilim(SW).....	832	ETHOSUXIMIDE.....	828
EpiPen (AL).....	126, 919	ETONOGESTREL.....	193
EpiPen Jr. (AL).....	126, 919	ETOPOSIDE.....	257
Epiramax 100 (RW).....	840	Eucerin(BE).....	1023
Epiramax 200 (RW).....	840	Eutroxsig(LN).....	210
Epiramax 25 (RW).....	841	Evelyn 150/30 ED (GQ).....	193
Epiramax 50 (RW).....	841	Evenity(AN).....	777
EPLERENONE.....	131	EVEROLIMUS	
Eplerenone AN (EA).....	131	ANTINEOPLASTIC AND IMMUNOMODULATING	
EPROSARTAN.....	145	AGENTS.....	307, 308, 309, 310, 368
EPROSARTAN + HYDROCHLOROTHIAZIDE.....	147	Everolimus Sandoz (SZ).....	308, 309
EPTIFIBATIDE.....	111	Evifyne (EL).....	776
Erivedge(RO).....	335	Evista(LY).....	776
ERLOTINIB.....	289, 290	EVOLOCUMAB.....	159, 160, 162
Erlotinib Sandoz(SZ).....	290	Evolve Carmellose(CX).....	944
ERTUGLIFLOZIN.....	100, 101	Evolve Hypromellose(CX).....	947
ERTUGLIFLOZIN + METFORMIN.....	71, 72	Exelon Patch 10(NV).....	889, 890, 891
ERTUGLIFLOZIN + SITAGLIPTIN.....	73	Exelon Patch 15(NV).....	889, 890, 891
		Exelon Patch 5(NV).....	889, 890, 891

Exelon(NV)	889, 890, 891
EXEMESTANE	345, 346
Exemestane GH (GQ)	346
Exemestane Sandoz(SZ)	346
EXENATIDE	93, 94
Exforge 10/160(NV)	150
Exforge 10/320(NV)	150
Exforge 5/160(NV)	150
Exforge 5/320(NV)	150
Exforge 5/80(NV)	150
Exforge HCT 10/160/12.5(NV)	151
Exforge HCT 10/160/25(NV)	151
Exforge HCT 10/320/25(NV)	152
Exforge HCT 5/160/12.5(NV)	151
Exforge HCT 5/160/25(NV)	151
Extine 20 (RW)	873
Exufiber 709900(MH)	1049
Exufiber 709901(MH)	1049
Exufiber 709903(MH)	1049
Exufiber 709908(MH)	1050
Exufiber 709909(MH)	1049
Eylea(BN)	934, 935, 936, 938
Ezalo Composite Pack 10mg+10mg(AF)	168
Ezalo Composite Pack 10mg+20mg(AF)	168
Ezalo Composite Pack 10mg+40mg(AF)	168
Ezalo Composite Pack 10mg+5mg(AF)	169
EZEMICHOL(RW)	166, 167
EZETIMIBE	166
EZETIMIBE (&) ROSUVASTATIN	167, 168
EZETIMIBE + ATORVASTATIN	169, 170
EZETIMIBE + SIMVASTATIN	170, 171
Ezetimibe GH (GQ)	167
Ezetimibe Sandoz(SZ)	167
Ezetimibe/Atorvastatin GH 10/10 (GQ)	170
Ezetimibe/Atorvastatin GH 10/20 (GQ)	170
Ezetimibe/Atorvastatin GH 10/40 (GQ)	170
Ezetimibe/Atorvastatin GH 10/80 (GQ)	170
EZETIMIBE/SIMVASTATIN SANDOZ (SZ)	171, 172
EZETORIN(RW)	171, 172
Ezetrol(AL)	166, 167
Ezovir (AF)	247
Ezovir(AF)	247, 248
EzSimva GH 10/10 (GQ)	172
EzSimva GH 10/20 (GQ)	172
EzSimva GH 10/40 (GQ)	171
EzSimva GH 10/80 (GQ)	171
FAMCICLOVIR	247, 248
Famciclovir AN (EA)	247, 248
Famciclovir AN(EA)	247, 248
Famciclovir generichealth 250 (GQ)	247
Famciclovir generichealth 500 (GQ)	248
Famciclovir-GA (ED)	247
Famciclovir-GA(ED)	247, 248
FAMOTIDINE	29
Famotidine AN (EA)	29
Famvir (IX)	247, 248
Famvir(IX)	247, 248
Fareston(AS)	344
Faverin 100 (RW)	873
Faverin 50 (RW)	873
Favic 125(RW)	247
Favic 250 (RW)	247
Favic 500 (RW)	248
Favic 500(RW)	248
Febriadol (EA)	823, 1034
FEBUXOSTAT	768
Feldene(PF)	762, 763
Feldene-D(PF)	762
Felodil XR 10(RW)	137
Felodil XR 5(RW)	137

FELODIPINE	137
Felodur ER 10 mg (TX)	137
Felodur ER 2.5 mg(TX)	137
Felodur ER 5 mg (TX)	137
Femara 2.5 mg (NV)	346
Femme-Tab ED 20/100(AE)	193
Femme-Tab ED 30/150(AE)	193
Femolet(AF)	346
Femoston 1/10(GO)	198
Femoston 2/10(GO)	199
Femoston-Conti(GO)	198
Fenac (AF)	760, 996
Fenac 25 (AF)	760, 996
Fendex ER (AF)	137
Fendex ER(AF)	137
Fenocol(YC)	158, 159
FENOFIBRATE	158
Fenofibrate Cipla (LR)	158, 159
Fenofibrate Mylan (AF)	158, 159
Fenofibrate Mylan(AF)	158, 159
FENOFIBRATE RBX (RA)	158, 159
FENOFIBRATE RBX(RA)	158, 159
Fenofibrate Sandoz(SZ)	158, 159
Fenpatch 100 (ZP)	812
Fenpatch 12 (ZP)	805
Fenpatch 25 (ZP)	807
Fenpatch 50 (ZP)	809
Fenpatch 75 (ZP)	810
FENTANYL 804, 806, 807, 809, 810, 998, 999, 1000, 1001	
Fentanyl Sandoz(SZ)	806, 807, 809, 810, 812
Fentora(TB)	1000, 1001, 1002
Fera (AS)	346
Ferinject(VL)	121
FERRIC CARBOXYMALTOSE	121
FERRIC DERISOMALTOSE	121
Ferro-f-tab(AE)	122, 1021
Ferro-Liquid(AE)	121
Ferrosig(SI)	121, 122
Ferro-tab(AE)	121, 1020
FERROUS FUMARATE	121, 1020
FERROUS FUMARATE + FOLIC ACID	122, 1021
FERROUS SULFATE	121
FEXOFENADINE	1038
Fiasp FlexTouch(NO)	61
Fiasp(NO)	61
Fibre Health Natural Granular(PP)	1017
Fibsol 10(RW)	139
Fibsol 20(RW)	139
Fibsol 5(RW)	139
Finasta (SZ)	1031
FINASTERIDE	1030
Finasteride AN (EA)	1030
Finasteride GH 5(GQ)	1030
Finasteride-GA 5(GN)	1031
FINGOLIMOD	368
Finide (AL)	1030
Finnacar(RW)	1031
Finpro(RZ)	1031
Firmagon 120mg(FP)	347
Firmagon 80mg(FP)	347
Fixta 60 (DO)	776
Flagyl S(SW)	237, 238
Flagyl(SW)	237, 238
Flamazine(SN)	176
Flarex(NV)	926
FLECAINIDE	124
Flecainide Sandoz (SZ)	124
Flecatab(AF)	124
Flixotide Accuhaler (GK)	915
Flixotide Accuhaler(GK)	915

<i>Flixotide Junior (GK)</i>	915	MUSCULO-SKELETAL SYSTEM	777
<i>Flixotide Junior Accuhaler (GK)</i>	915	SYSTEMIC HORMONAL PREPARATIONS, EXCL. SEX	
<i>Flixotide (GK)</i>	915	HORMONES AND INSULINS	212
<i>Flomaxtra (LS)</i>	1030	<i>Forxiga (AP)</i>	97, 98
<i>Flopen (AL)</i>	220, 221	<i>Fosamax Plus 70 mg/140 mcg (MQ)</i>	772, 1033
<i>Florinef (AS)</i>	206	<i>Fosamax Plus (MQ)</i>	773, 1033
<i>Fluanxol Concentrated Depot (LU)</i>	851	FOSAPREPITANT	49
<i>Fluanxol Depot (LU)</i>	851	<i>Fosetic 20/12.5 (ZP)</i>	142
<i>Flubiclox (JU)</i>	220	FOSINOPRIL	139
<i>Flucil (AS)</i>	220	FOSINOPRIL + HYDROCHLOROTHIAZIDE	142
<i>Flucil (LN)</i>	220	<i>Fosinopril/HCT Actavis 20/12.5 (EA)</i>	142
FLUCLOXACILLIN	219, 220, 221	<i>Fosipril 10 (RW)</i>	139
FLUCONAZOLE	238, 239, 240	<i>Fosipril 20 (RW)</i>	139
<i>Fluconazole Alphapharm (AF)</i>	239	<i>Fosrenol (TK)</i>	953
<i>Fluconazole APOTEX (GX)</i>	240	<i>Fostair (EU)</i>	904
<i>Fluconazole Sandoz (SZ)</i>	240	<i>Fragmin (PF)</i>	107, 108
<i>Fluconazole Sandoz (SZ)</i>	240	<i>Frakas (RW)</i>	214
<i>Fludara (GZ)</i>	255	FRAMYCETIN SULFATE	951
FLUDARABINE	255	FRAMYCETIN SULFATE + GRAMICIDIN +	
FLUDROCORTISONE ACETATE	206	DEXAMETHASONE	951
FLUNITRAZEPAM	1035	<i>Fruiti Vits (VF)</i>	991
FLUOROMETHOLONE	926	<i>Frusax (ER)</i>	130
FLUOROMETHOLONE ACETATE	926	<i>Frusemix (TY)</i>	130
FLUOROURACIL	1031	<i>Frusemix-M (TY)</i>	131
<i>FLUOTEX (RF)</i>	873	<i>Frusemix-M (TY)</i>	20, 131
FLUOXETINE	872	<i>Fucidin (LO)</i>	237
<i>Fluoxetine AN (EA)</i>	873	FULVESTRANT	343
<i>Fluoxetine APOTEX (TY)</i>	873	<i>Fulvestrant Sandoz (SZ)</i>	343
<i>Fluoxetine Capsule (USP) (DZ)</i>	873	<i>Fungilin (AS)</i>	29
<i>Fluoxetine generichealth (GQ)</i>	873	FUROSEMIDE (FRUSEMIDE)	20, 130, 131
<i>Fluoxetine Sandoz (SZ)</i>	873	FUROSEMIDE AN (EA)	130
<i>Fluoxetine-GA (ED)</i>	873	FUROSEMIDE AN (EA)	131
FLUPENTIXOL DECANOATE	851	FUSIDATE	237
FLUTAMIDE	345	<i>Fybogel (RC)</i>	1017
<i>Flutamin (AF)</i>	345	<i>Fycompa (EI)</i>	838, 839
<i>Fluticasone + Salmeterol Cipla 125/25 (LR)</i>	910	GA express 15 (VF)	978
<i>Fluticasone + Salmeterol Cipla 250/25 (LR)</i>	911	GA gel (VF)	978
<i>Fluticasone Cipla Inhaler (LR)</i>	915	GA1 Anamix infant (SB)	978
FLUTICASONE FUROATE	915	GA1 Anamix Junior (NU)	978
FLUTICASONE FUROATE + UMECLIDINIUM +		<i>Gabacor (CR)</i>	1034, 1035
VILANTEROL	912	<i>Gabacor (CR)</i>	
FLUTICASONE FUROATE + VILANTEROL	909	NERVOUS SYSTEM	834
FLUTICASONE PROPIONATE	915	GABAPENTIN	
FLUTICASONE PROPIONATE + FORMOTEROL		NERVOUS SYSTEM	834
(EFORMOTEROL)	909	Repatriation Pharmaceutical Benefits Scheme	1034
FLUTICASONE PROPIONATE + SALMETEROL	910	<i>Gabapentin AN (EA)</i>	834, 1034
<i>flutiform 125/5 (MF)</i>	910	<i>Gabapentin AN (EA)</i>	
<i>flutiform 250/10 (MF)</i>	910	NERVOUS SYSTEM	834
<i>flutiform 50/5 (MF)</i>	910	<i>Gabapentin AN (EA)</i>	
FLUVASTATIN	154	NERVOUS SYSTEM	834
FLUVOXAMINE	873	<i>Gabapentin APOTEX (TY)</i>	
<i>Fluvoxamine AN (ED)</i>	873	NERVOUS SYSTEM	834
<i>Fluvoxamine GA (EA)</i>	873	<i>Gabapentin APOTEX (TY)</i>	
<i>Fluzole 200 (RW)</i>	240	Repatriation Pharmaceutical Benefits Scheme	1034, 1035
<i>FML Liquifilm (AG)</i>	926	<i>Gabapentin APOTEX (TY)</i>	
FOLIC ACID	122, 123, 1021	NERVOUS SYSTEM	834
FOLINIC ACID	954	<i>Gabapentin Aspen 100 (RW)</i>	
FOLLITROPIN ALFA	199	Repatriation Pharmaceutical Benefits Scheme	1035
FOLLITROPIN BETA	200	<i>Gabapentin Aspen 100 (RW)</i>	
<i>Foltabs 500 (PP)</i>	123, 1021	NERVOUS SYSTEM	834
<i>Fonat (AL)</i>	769	<i>Gabapentin Aspen 300 (RW)</i>	
<i>Fonat Plus (AF)</i>	772, 773	Repatriation Pharmaceutical Benefits Scheme	1034
<i>Fonat Plus (AF)</i>	1033	<i>Gabapentin Aspen 300 (RW)</i>	
FONDAPARINUX	120	NERVOUS SYSTEM	834
<i>Foradile (SZ)</i>	902	<i>Gabapentin Aspen 400 (RW)</i>	
<i>Formet 1000 (RW)</i>	64	Repatriation Pharmaceutical Benefits Scheme	1034
FORMET 500 (RF)	63	<i>Gabapentin Aspen 400 (RW)</i>	
FORMET 850 (RF)	64	NERVOUS SYSTEM	834
FORMOTEROL (EFORMOTEROL)	902	<i>Gabapentin Aspen 600 (RW)</i>	
<i>Forteo (LY)</i>			

.Repatriation Pharmaceutical Benefits Scheme	1034	GLIBENCLAMIDE	64
<i>Gabapentin Aspen 600 (RW)</i>		GLICLAZIDE	64
.NERVOUS SYSTEM	834	GLIMEPIRIDE	64
<i>Gabapentin Aspen 800 (RW)</i>		<i>Glimepiride AN(EA)</i>	65
.Repatriation Pharmaceutical Benefits Scheme	1034	<i>Glimepiride APOTEX (GX)</i>	65
<i>Gabapentin Aspen 800 (RW)</i>		<i>Glimepiride Sandoz(SZ)</i>	65
.NERVOUS SYSTEM	834	GLIPIZIDE	65
<i>Gabapentin generichealth(HQ)</i>		<i>Glivec (AF)</i> 264, 265, 266, 267, 268, 269, 270, 272, 273,	
.Repatriation Pharmaceutical Benefits Scheme	1034	275, 276, 277, 278, 279, 280	
<i>Gabapentin generichealth(HQ)</i>		<i>Glivec(AF)</i>	263, 264, 265, 266, 269, 270, 271
.NERVOUS SYSTEM	834	<i>Glucagen Hypokit(NO)</i>	20, 211
<i>Gabapentin Sandoz (SZ)</i>		GLUCAGON HYDROCHLORIDE	20, 211
.NERVOUS SYSTEM	834	<i>Glucobay 100(BN)</i>	83
<i>Gabitril(TB)</i>	831	<i>Glucobete 1000(DO)</i>	64
GALANTAMINE	886, 887, 888	<i>Glucobete 500(DO)</i>	63
<i>Galantamine AN SR (EA)</i>	887, 888, 889	<i>Glucobete 850 (DO)</i>	64
<i>Galantyl(AF)</i>	887, 888, 889	GLUCOSE AND KETONE INDICATOR URINE	955
<i>Galvumet 50/1000(NV)</i>	83	GLUCOSE INDICATOR URINE	955
<i>Galvumet 50/500(NV)</i>	83	<i>Glyade (AF)</i>	64
<i>Galvumet 50/850(NV)</i>	83	<i>Glyade MR (AF)</i>	64
<i>Galvus(NV)</i>	92	GLYBOSAY (RW)	83
<i>Gamine XR (RW)</i>	887, 888, 889	GLYCEROL	
<i>Ganfort 0.3/5(AG)</i>	932	.Repatriation Pharmaceutical Benefits Scheme	1017,
<i>GANfort PF 0.3/5(AG)</i>	931, 932	1026	
<i>Gantin (EA)</i>		GLYCEROL + WHITE SOFT PARAFFIN	1023
.Repatriation Pharmaceutical Benefits Scheme	1034	GLYCERYL TRINITRATE	20, 126
<i>Gantin(ED)</i>	1034	GLYCINE WITH CARBOHYDRATE	985
GAPENTIN (RF)		<i>Glycine500(VF)</i>	985
.NERVOUS SYSTEM	834	GLYCOMACROPEPTIDE AND ESSENTIAL AMINO ACID	
GAPENTIN(RF)	834	FORMULA WITH VITAMINS, MINERALS, AND LOW IN	
<i>Gastrex(CR)</i>	56, 1018	TYROSINE AND PHENYLALANINE	986
<i>Gastro-Stop (AS)</i>	56	GLYCOMACROPEPTIDE AND ESSENTIAL AMINO	
GAUZE AND COTTON TISSUE COMBINE ROLL	1057	ACIDS WITH VITAMINS AND MINERALS	986, 987
GEFITINIB	291	GLYCOMACROPEPTIDE FORMULA WITH	
GELATIN + PECTIN + CARMELLOSE SODIUM	1023	DOCOSAHEXAENOIC ACID AND LOW	
GEMFIBROZIL	159	PHENYLALANINE	987
<i>Genoptic(AG)</i>	922	GLYCOPYRRONIUM	916
<i>Genox 20(AF)</i>	343, 344	<i>Glycosade(VF)</i>	970
<i>GenRx Aciclovir (GX)</i>	246	<i>Glyxambi(BY)</i>	68, 69
<i>GenRx Aciclovir(GX)</i>	246	<i>Goanna Heat Cream(GQ)</i>	1032
<i>GenRx Celecoxib(GX)</i>	766	GOLIMUMAB 573, 575, 577, 580, 581, 585, 587, 589, 594,	
<i>GenRx Clomipramine (GX)</i>	869	598, 602	
<i>GenRx Cyproterone Acetate (GX)</i>		<i>Gonal-f Pen(SG)</i>	199
.ANTINEOPLASTIC AND IMMUNOMODULATING		<i>Gopten(GO)</i>	141
AGENTS	344	GOSERELIN	340
.GENITO URINARY SYSTEM AND SEX HORMONES		GOSERELIN (&) BICALUTAMIDE	340
.....	201	GRANISETRON	44, 45
<i>GenRx Cyproterone Acetate(GX)</i>		<i>Granisetron Kabi (PK)</i>	44
.ANTINEOPLASTIC AND IMMUNOMODULATING		<i>Granisetron-AFT(AE)</i>	44
AGENTS	344	GRISEOFULVIN	174
.GENITO URINARY SYSTEM AND SEX HORMONES		<i>Grisovin 500(AS)</i>	174
.....	201	<i>Grisovin(AS)</i>	174
<i>GenRx Famotidine(GX)</i>	29	GUANFACINE	128
<i>GenRx Indapamide (GX)</i>	130	GUSELKUMAB	607
<i>GenRx Isosorbide Mononitrate(GX)</i>	126	<i>Gynotril(ER)</i>	346
<i>GenRx Norfloxacin (GX)</i>	236	<i>Haldol decanoate(JC)</i>	850
<i>GenRx Perindopril/ Indapamide 4/1.25(GX)</i>	142	HALOPERIDOL	19, 850
<i>GenRx Tamoxifen (GX)</i>	343	HALOPERIDOL DECANOATE	850
GENTAMICIN		<i>Hamilton Skin Therapy Oil(KY)</i>	1023
.ANTIINFECTIVES FOR SYSTEMIC USE	233	<i>Hamilton Skin Therapy Wash(KY)</i>	1027
.SENSORY ORGANS	921, 922	<i>Handy 36361414(BV)</i>	1039
<i>Genteal gel(AQ)</i>	947	<i>Handy 71117-05(BV)</i>	1049
<i>Genteal(AQ)</i>	946, 947	<i>Handy 71117-06(BV)</i>	1049
<i>Gilenya(NV)</i>	368, 369	<i>Handygauze Cohesive 8631(BV)</i>	1040
<i>Gilmot(CR)</i> 264, 265, 266, 267, 268, 269, 270, 272, 273,		<i>Handygauze Cohesive 8633(BV)</i>	1041
275, 276, 277, 278, 279, 280		<i>Handygauze Cohesive 8635(BV)</i>	1041
<i>Giotrif(BY)</i>	289	<i>Harvoni(GI)</i>	251
GLATIRAMER ACETATE	349	<i>HCU Anamix infant(SB)</i>	979
GLECAPREVIR + PIBRENTASVIR	250	<i>HCU Anamix junior LQ(SB)</i>	977
<i>Gliadel(EI)</i>	253	<i>HCU Anamix Junior(NU)</i>	979

<i>HCU cooler 10(VF)</i>	978	<i>HYOSCINE BUTYLBROMIDE SXP(XC)</i>	20, 994
<i>HCU cooler 15(VF)</i>	978	<i>Hypnodorm(AF)</i>	1036
<i>HCU cooler 20(VF)</i>	979	<i>HYPROMELLOSE</i>	946, 947
<i>HCU Easy Tablet(OH)</i>	976	<i>HYPROMELLOSE + CARBOMER-980</i>	947
<i>HCU express 15(VF)</i>	978	<i>Hysone 20(AF)</i>	207
<i>HCU gel(VF)</i>	978	<i>Hysone 4(AF)</i>	207
<i>HCU Lophlex LQ 20(SB)</i>	978	<i>IBANDRONATE</i>	769
<i>HCU Lophlex(SB)</i>	979	<i>Ibavyr(IX)</i>	251, 252
<i>HEPARIN</i>	109	<i>Ibilex 125 (AF)</i>	223
<i>Hequinel (RW)</i>	766, 767	<i>Ibilex 250 (AF)</i>	223, 224, 225
<i>Herceptin SC(RO)</i>	331	<i>Ibilex 500 (AF)</i>	223, 224, 225
HIGH FAT FORMULA WITH VITAMINS, MINERALS AND TRACE ELEMENTS AND LOW IN PROTEIN AND CARBOHYDRATE	987, 988	<i>Ibrance(PF)</i>	304
<i>Hiprex(IL)</i>	238	<i>IBRUTINIB</i>	315
HONEY BEE VENOM	951	<i>IBUPROFEN</i>	763, 997
<i>HPMC PAA(IQ)</i>	947	<i>ICATIBANT</i>	123
<i>Humalog KwikPen(KP)</i>	61	<i>ICHTHAMMOL</i>	1026
<i>Humalog Mix25 KwikPen(KP)</i>	63	<i>ICHTHAMMOL + ZINC OXIDE</i>	1026
<i>Humalog Mix25(LY)</i>	63	<i>Iclusig(TK)</i>	285, 286, 288
<i>Humalog Mix50 KwikPen(KP)</i>	62	<i>Idaprex 2 (SZ)</i>	139
<i>Humalog Mix50(LY)</i>	63	<i>Idaprex 4 (SZ)</i>	140
<i>Humalog U200 Kwikpen(LY)</i>	61	<i>Idaprex 8 (SZ)</i>	140
<i>Humalog(LY)</i>	61	<i>Idaprex Combi 4/1.25 (SZ)</i>	142
HUMAN CHORIONIC GONADOTROPHIN	200	<i>IDELALISIB</i>	316, 317
<i>Humira(VE)</i> 400, 402, 404, 405, 409, 411, 413, 415, 418, 420, 423, 428, 433, 435, 436, 438, 442, 446, 451, 452, 456, 462, 469		<i>Ikorel (SW)</i>	127
<i>Humulin 30/70(LY)</i>	62	<i>Ikotab(AF)</i>	127
<i>Humulin NPH(LY)</i>	62	<i>Ilumya(RA)</i>	687, 694
<i>Humulin R (LY)</i>	61, 62	IMATINIB263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 275, 276, 277, 278, 279, 280	
HYALURONATE SODIUM	946	<i>Imatinib GH (GQ)</i> ... 264, 265, 266, 267, 268, 269, 270, 272, 273, 275, 276, 277, 278, 279, 280	
<i>Hycor(AS)</i>	927	IMATINIB RBX (RA)	265, 266, 269, 270
HYDRALAZINE	129	IMATINIB RBX(RA) 264, 265, 266, 267, 268, 269, 270, 272, 273, 275, 276, 277, 278, 279, 280	
<i>Hydrea(BQ)</i>	337	<i>Imatinib-APOTEX (TX)</i> .. 264, 265, 266, 267, 268, 269, 270, 272, 273, 275, 276, 277, 278, 279, 280	
HYDROCHLOROTHIAZIDE	130	IMATINIB-DRLA(RZ) 264, 265, 266, 267, 268, 269, 270, 272, 273, 275, 276, 277, 278, 279, 280	
<i>Hydrocoll 900744(HR)</i>	1052	<i>Imatinib-Teva (SZ)</i> . 264, 265, 266, 267, 268, 269, 270, 272, 273, 275, 276, 277, 278, 279, 280	
<i>Hydrocoll 900936(HR)</i>	1052	<i>Imazan (ER)</i>	755
<i>Hydrocoll Thin 900758(HR)</i>	1052	<i>Imbruvica(JC)</i>	315, 316
HYDROCORTISONE	207	<i>Imdur 120 mg(IX)</i>	126
HYDROCORTISONE ACETATE		<i>Imdur Durule(IX)</i>	127
.ALIMENTARY TRACT AND METABOLISM	56	<i>Imigran FDT(AS)</i>	826, 827
.DERMATOLOGICALS	176	<i>Imigran(AS)</i>	827
.Repatriation Pharmaceutical Benefits Scheme	1024	<i>Imigran(LN)</i>	827
.SENSORY ORGANS	927	IMIPRAMINE	870
<i>Hydrocortisone Mylan 20(AL)</i>	207	<i>Imipramine (Leading)(QY)</i>	870
<i>Hydrocortisone Mylan 4(AL)</i>	207	IMIQUIMOD	190, 1026
HYDROCORTISONE SODIUM SUCCINATE	20, 207	<i>Imoclone (RW)</i>	1036
HYDROMORPHONE	780, 781, 782	<i>Imovane(SW)</i>	1036
<i>HYDROMORPHONE JUNO (JU)</i>	782	<i>Implanon NXT(OQ)</i>	193
<i>HYDROMORPHONE JUNO-HP (JU)</i>	782	<i>Imrest(AF)</i>	1036
<i>Hydrosorb 900854(HR)</i>	1055	<i>Imuran (AS)</i>	755
<i>Hydroxo-B12(AS)</i>	122	<i>Imuran(AS)</i>	755
HYDROXOCOBALAMIN	122, 1021	<i>In a Wink Moisturising(IQ)</i>	946, 947
HYDROXYCARBAMIDE (HYDROXYUREA)	337	<i>Inadine(KI)</i>	1057
HYDROXYCARBAMIDE MEDSURGE (DZ)	337	<i>Incruse Ellipta(GK)</i>	918
HYDROXYCHLOROQUINE	766	INDACATEROL	902
<i>Hydroxychloroquine AN(EA)</i>	766, 767	INDACATEROL + GLYCOPYRRONIUM	912
<i>Hydroxychloroquine GH (GQ)</i>	766, 767	INDACATEROL + GLYCOPYRRONIUM + MOMETASONE	
HYDROXYETHYL STARCH 130/0.4 + SODIUM CHLORIDE	123	 913	
<i>Hygroton 25(GH)</i>	130	INDACATEROL + MOMETASONE	911
<i>Hylo-Forte(AE)</i>	946	INDAPAMIDE	130
<i>Hylo-Fresh(AE)</i>	946	INDAPAMIDE AN SR (EA)	130
<i>Hymenoptera Honey Bee Venom(DE)</i>	952	<i>Indapamide AN(EA)</i>	130
<i>Hymenoptera Paper Wasp Venom(DE)</i>	952	<i>Inderal(AP)</i>	133
<i>Hymenoptera Yellow Jacket Venom(DE)</i>	952	<i>Indocid(AS)</i>	761, 996
HYOSCINE BUTYLBROMIDE	20, 994, 1016	INDOMETACIN	761, 996
HYOSCINE BUTYLBROMIDE MEDSURGE (DZ) ...	20, 994	<i>Indosyl Combi 4/1.25(RW)</i>	142
HYOSCINE BUTYLBROMIDE SXP (XC)	1016		

<i>Indosyl Mono 2(RW)</i>	139	<i>Isoleucine 1000(VF)</i>	988
<i>Indosyl Mono 4(RW)</i>	140	<i>Isoleucine 50(VF)</i>	988
<i>Indosyl Mono 8(RW)</i>	140	ISOLEUCINE WITH CARBOHYDRATE	988
INFLIXIMAB	1031	<i>Isomoni(SZ)</i>	126
<i>Inlyta(PF)</i>	313, 314	ISONIAZID	245
<i>iNova Pharmaceuticals (Australia) Pty Ltd(IL)</i>	898	<i>Isoptin 180 SR(GO)</i>	137
<i>Inpler (AF)</i>	131	<i>Isoptin SR(GO)</i>	138
<i>Insig (RW)</i>	130	<i>Isoptin(GO)</i>	137
<i>Inspra(UJ)</i>	131	<i>Isopto Carpine(NV)</i>	928, 929
INSULIN ASPART	61	<i>Isordil Sublingual(RW)</i>	126
INSULIN ASPART + INSULIN ASPART PROTAMINE	62	<i>Isosorbide AN (EA)</i>	126
INSULIN DEGLUDEC + INSULIN ASPART	62	ISOSORBIDE DINITRATE	126
INSULIN DETEMIR	63	ISOSORBIDE MONONITRATE	126
INSULIN GLARGINE	63	ISOTRETINOIN	185, 186
INSULIN GLULISINE	61	<i>Isotretinoin AN(EA)</i>	186
INSULIN ISOPHANE HUMAN	62	<i>Isotretinoin GX (SZ)</i>	186
INSULIN ISOPHANE HUMAN + INSULIN NEUTRAL		<i>Isotretinoin Lupin(GQ)</i>	186
HUMAN	62	<i>Isotretinoin SCP 20 (CR)</i>	186
INSULIN LISPRO	61	<i>Itacap (AF)</i>	242
INSULIN LISPRO + INSULIN LISPRO PROTAMINE	62	ITRACONAZOLE	241
INSULIN NEUTRAL HUMAN	61	<i>ITRANOX(RW)</i>	242
<i>Intal CFC-Free(SW)</i>	918	IVABRADINE	127
<i>Intal Forte CFC-Free(SW)</i>	918	IVERMECTIN	901
<i>Integrilin(MK)</i>	111	IXEKIZUMAB	615, 617, 621, 625, 632
INTERFERON ALFA-2A	347	<i>Jakavi(NV)</i>	312, 313
INTERFERON BETA-1A	348	<i>Janumet XR(MK)</i>	79, 81
INTERFERON BETA-1B	348	<i>Janumet(MK)</i>	79, 81
<i>Intrasite Gel 7313(SN)</i>	1054	<i>Januvia(MK)</i>	89, 91
<i>Intuniv(TK)</i>	129	<i>Jardiamet 12.5 mg/1000 mg(BY)</i>	70, 71
<i>Invega Sustenna(JC)</i>	858, 859	<i>Jardiamet 12.5 mg/500 mg(BY)</i>	70, 71
<i>Invega Trinza(JC)</i>	859	<i>Jardiamet 5 mg/1000 mg(BY)</i>	70, 71
<i>Invega(JC)</i>	858, 859	<i>Jardiamet 5 mg/500 mg(BY)</i>	70, 71
<i>Iodosorb 66051330(SN)</i>	1043	<i>Jardiance(BY)</i>	99, 100
<i>Iodosorb 66051340(SN)</i>	1043	<i>Jelonet 7404(SN)</i>	1049
<i>Iodosorb 66051360(SN)</i>	1043	<i>Jetrea RTU(IJ)</i>	948
<i>Iodosorb Ointment 66051230(SN)</i>	1043	<i>Jinarc(OS)</i>	132, 133
<i>Iodosorb Ointment 66051240(SN)</i>	1043	<i>JJ 12010(JJ)</i>	1057
<i>Iodosorb Powder 66051070(SN)</i>	1043	<i>Junalox(JO)</i>	21, 952
<i>Ionil-T(GA)</i>	1025	<i>Jurnista(JC)</i>	784
<i>Iopidine 0.5%(NV)</i>	927	<i>Kalixocin(AF)</i>	230
<i>I-Pantoprazole (CR)</i>	39, 40, 41	<i>Kalma 0.25(AF)</i>	865
<i>Ipratrin Adult(AF)</i>	916	<i>Kalma 0.5 (AF)</i>	866
<i>Ipratrin(AF)</i>	916	<i>Kalma 1 (AF)</i>	866
IPRATROPIUM		<i>Kaltostat 168117(CC)</i>	1044
.Repatriation Pharmaceutical Benefits Scheme	1037	<i>Kaltostat 168210(CC)</i>	1044
.RESPIRATORY SYSTEM	916	<i>Kaltostat 168212(CC)</i>	1044
<i>Iptam (AL)</i>	826, 827	<i>Kapanol(YN)</i>	794, 795, 998
IRBESARTAN	145	<i>Kaptan (ER)</i>	855, 856
IRBESARTAN + HYDROCHLOROTHIAZIDE	147	<i>Karlor CD(MQ)</i>	226
<i>Irbesartan Actavis 150(ED)</i>	145	<i>Karvea (SW)</i>	145
<i>Irbesartan Actavis 300(ED)</i>	145	<i>Karvezide 150/12.5 (SW)</i>	148
<i>Irbesartan Actavis 75(ED)</i>	145	<i>Karvezide 300/12.5 (SW)</i>	148
<i>Irbesartan AMNEAL (EF)</i>	145	<i>Karvezide 300/25 (SW)</i>	148
<i>Irbesartan AN(EA)</i>	145	<i>Keflex(AS)</i>	223, 224, 225
<i>Irbesartan GH (GQ)</i>	145	<i>Keflor CD (AF)</i>	226
<i>Irbesartan HCT Actavis 150/12.5(ED)</i>	148	<i>Keflor(AF)</i>	226, 227
<i>Irbesartan HCT Actavis 300/12.5(ED)</i>	148	<i>Kenacomb Otic(AS)</i>	951
<i>Irbesartan HCT Actavis 300/25(ED)</i>	148	<i>Kenacort-A10(AS)</i>	209
<i>Irbesartan HCT GH 150/12.5 (GQ)</i>	148	<i>Keppra (UC)</i>	837, 838
<i>Irbesartan HCT GH 300/12.5 (GQ)</i>	148	<i>Kerron 1000(DO)</i>	837
<i>Irbesartan HCT GH 300/25 (GQ)</i>	148	<i>Kerron 250(DO)</i>	837
<i>Irbesartan HCTZ AMNEAL (EF)</i>	148	<i>Kerron 500(DO)</i>	837
<i>Irbesartan Sandoz(SZ)</i>	145	<i>Kerron(DO)</i>	838
<i>Irbesartan/HCT Sandoz(SZ)</i>	148	<i>KetoCal 3</i>	
<i>Ircal(PE)</i>	948	1(SB)	988
<i>Iressa (AP)</i>	291	<i>KetoCal 4</i>	
IRON POLYMALTOSE	121	1 LQ(SB)	988
IRON SUCROSE	122	1(SB)	988
<i>Iscover(AV)</i>	109, 110, 1020	KETOCONAZOLE	173
ISOBIDE MR (RF)	126	<i>Keto-Diastix(DX)</i>	955

KETOPROFEN	763	Leflunomide Sandoz(SZ)	370
Kevtam 1000 (AF)	837	Lenest 30 ED (AF)	193
Kevtam 250 (AF)	837	Lengout(LN)	768
Kevtam 500 (AF)	837	LENVATINIB	320
Keyo(VF)	987	Lenvima(EI)	320, 321
keyomega(VF)	984	Lercadip (EA)	137
Kindergen(SB)	992	Lercan(RW)	137
Kinson (AF)	843	LERCANIDIPINE	137
Kisqali(NV)	305, 306, 307	LERCANIDIPINE + ENALAPRIL	143
Klacid(GO)	230, 231	Lercanidipine APOTEX (GX)	137
Kombiglyze XR 2.5/1000(AP)	77, 79	Lescol XL(NV)	154
Kombiglyze XR 5/1000(AP)	77, 79	LETOZOLE	346
Kombiglyze XR 5/500(AP)	77, 79	Letrozole APOTEX (GX)	346
Konakion MM(PB)	21	Letrozole FBM(FO)	346
Kosteo(RW)		Letrozole GH (HQ)	346
ALIMENTARY TRACT AND METABOLISM	102	Letrozole Sandoz(SZ)	346
MUSCULO-SKELETAL SYSTEM	774	Leucovorin Calcium (Hospira Pty Limited)(PF)	954
KSART HCT 150/12.5(RW)	148	Leucovorin Calcium (Pfizer Australia Pty Ltd)(PF)	954
KSART HCT 300/12.5(RW)	148	Leukeran(AS)	253
KSART HCT 300/25(RW)	148	Leukoflex 1124(BV)	1058
Kuvan(IO)	104, 105, 106	Leukoplast 01071-00(BV)	1057
Kyleena(SY)	191	Leukoplast 01072-00(BV)	1058
Kytril(IX)	44, 45	Leukoplast 01073-00(BV)	1058
L.C.E. Sandoz (HX)	844	Leukopor 2471(BV)	1058
LABETALOL	136	Leukopor 2472(BV)	1058
LACOSAMIDE	834, 835	Leukopor 2474(BV)	1058
Lamictal(AS)	836, 837	Leukosilk 1021(BV)	1058
Lamisil (Novartis Pharmaceuticals Australia Pty Limited)		Leukosilk 1022(BV)	1058
(NV)	174, 1023	Leukosilk 1024(BV)	1058
Lamisil DermGel(GK)	1022	LEUPRORELIN	341
Lamisil(GK)	174, 1022	LEUPRORELIN (&) BICALUTAMIDE	342
LAMITAN (RF)	836, 837	Levactam(ER)	837
LAMOTRIGINE	836	Levecetam 1000 (RZ)	837
Lamotrigine AN(EA)	836, 837	Levecetam 250 (RZ)	837
Lamotrigine Aspen 100 (RW)	836	Levecetam 500 (RZ)	837
Lamotrigine Aspen 200 (RW)	836	Levemir FlexPen(NF)	63
Lamotrigine Aspen 25 (RW)	836	Levemir Penfill(NO)	63
Lamotrigine Aspen 5 (RW)	836	LEVETIRACETAM	837
Lamotrigine Aspen 50 (RW)	837	Levetiracetam AN(EA)	837
Lamotrigine GH(GQ)	836, 837	Levetiracetam GH (GQ)	837
Lamotrigine Sandoz (SZ)	836, 837	Levetiracetam Mylan(AL)	837
Lanoxin(AS)	124	Levetiracetam SZ (SZ)	837
Lanoxin-PG(AS)	124	Levetiracetam-AFT (AE)	838
LANSOPRAZOLE	34, 35	Levi 1000(RW)	837
Lansoprazole ODT GH (GQ)	34, 35, 36	Levi 250(RW)	837
LANTHANUM	952	Levi 500(RW)	837
Lanvis(AS)	255	Levitra(BN)	1029
Lanzopran (RA)	34, 35, 36	Levlen ED(SY)	193
LAPATINIB	310	LEVODOPA + BENSERAZIDE	842
Largactil(SW)	19, 849	LEVODOPA + CARBIDOPA	843, 844
Lasix(SW)	20, 130, 131	LEVODOPA + CARBIDOPA + ENTACAPONE	844
Lasix-M(SW)	20, 131	LEVONORGESTREL	
LATANOPROST	932	GENITO URINARY SYSTEM AND SEX HORMONES	191, 193
LATANOPROST + TIMOLOL	932	LEVONORGESTREL + ETHINYLESTRADIOL	
Latanoprost Actavis (EA)	932	GENITO URINARY SYSTEM AND SEX HORMONES	193
Latanoprost Sandoz(SZ)	932	LEVOTHYROXINE	209
Latanoprost/Timolol Sandoz 50/5 (SZ)	932	Lexam 10 (RW)	871
Latuda(SE)	850, 851	Lexam 20 (RW)	871
LaxaCon (EA)	53, 995	Lexapro(LU)	871, 872
lax-sachets(AE)	53, 995	Lexotan(PB)	1035
Lax-Tab(AE)	51, 995	LIDOCAINE (LIGNOCAINE)	
Lecteva(TB)	844	CARDIOVASCULAR SYSTEM	124
LEDIPASVIR + SOFOSBUVIR	250, 251	Prescriber Bag	20
LEFLUNOMIDE	369	Repatriation Pharmaceutical Benefits Scheme	1024
Leflunomide AN (EA)	369, 370	LIGHT LIQUID PARAFFIN + COCOAMPHODIACETATE	
Leflunomide APOTEX (GX)	369	DISODIUM	1027
Leflunomide APOTEX(GX)	370	LINAGLIPTIN	85
Leflunomide generichealth (HQ)	370	LINAGLIPTIN + METFORMIN	74
Leflunomide generichealth(HQ)	369		
Leflunomide Sandoz (SZ)	369		

<i>Lincocin</i> (PF).....	233	<i>Lovir</i> (EA).....	246
LINCOMYCIN.....	233	<i>LoxaLate</i> (AF).....	871
LINCOMYCIN SXP(XC).....	233	<i>Loxip 500</i> (DO).....	235
<i>Link Medical Products Pty Ltd</i> (LM)		<i>Loxip 750</i> (DO).....	235
.Prescriber Bag.....	19	<i>Lozanoc</i> (YN).....	242
<i>Link Medical Products Pty Ltd</i> (LM)		<i>LPV</i> (IL).....	218, 219
.CARDIOVASCULAR SYSTEM.....	125	LUBRICATING AGENT.....	1039
<i>Link Medical Products Pty Ltd</i> (LM)		<i>Lubri-Gel</i> (PP).....	1039
.CARDIOVASCULAR SYSTEM.....	125	<i>Lucentis</i> (NV).....	938, 939, 941, 942
<i>Link Medical Products Pty Ltd</i> (LM)		<i>Lucrin Depot 3 Month PDS</i> (VE).....	341
.DERMATOLOGICALS.....	190	<i>Lucrin Depot 4 Month PDS</i> (VE).....	341
<i>Link Medical Products Pty Ltd</i> (LM)		<i>Lucrin Depot 6-Month</i> (VE).....	341
.DERMATOLOGICALS.....	190	<i>Lucrin Depot 7.5mg PDS</i> (VE).....	341
<i>Link Medical Products Pty Ltd</i> (LM)		<i>Lucrin Depot Paediatric 30 mg PDS</i> (VE).....	341, 342
.ANTIINFECTIVES FOR SYSTEMIC USE.....	245	<i>Lumigan</i> (AG).....	931
<i>Link Medical Products Pty Ltd</i> (LM)		<i>Lumigan PF</i> (AG).....	931
.ANTIINFECTIVES FOR SYSTEMIC USE.....	245	<i>Lumin 10</i> (AF).....	876
<i>Link Medical Products Pty Ltd</i> (LM)		<i>Lumin 20</i> (AF).....	877
.RESPIRATORY SYSTEM.....	919	<i>Lunava 10</i> (ZP).....	370
<i>Link Medical Products Pty Ltd</i> (LM)		<i>Lunava 20</i> (ZP).....	370
.RESPIRATORY SYSTEM.....	919	LURASIDONE.....	850
<i>Lioresal 10</i> (NV).....	767	<i>Lurasidone Lupin</i> (GQ).....	850, 851
<i>Lioresal 25</i> (NV).....	767	LURASIDONE SUN(RA).....	851
LIOTHYRONINE.....	210	<i>Luvox</i> (GO).....	873
<i>Lipex 10</i> (AL).....	157, 158	<i>Lyclear</i> (JT).....	901
<i>Lipex 20</i> (AL).....	157, 158	<i>Lynparza</i> (AP).....	336, 337
<i>Lipex 40</i> (AL).....	157, 158	LYPRALIN(RW).....	824
<i>Lipidil</i> (GO).....	158, 159	<i>Lyrice</i> (UJ).....	824
<i>Lipigem</i> (AF).....	159	LYSINE + THIAMINE + PYRIDOXINE +	
<i>Lipistart</i> (VF).....	970	CYANOCOBALAMIN + FERRIC PYROPHOSPHATE.....	1019
<i>Lipitor</i> (AS).....	153, 154	<i>Lyralon</i> (AF).....	824
<i>Lipostat 10</i> (RF).....	155	<i>Mabthera SC</i> (RO).....	329, 330
<i>Lipostat 20</i> (RF).....	155	<i>Macroclatin</i> (PF).....	238
<i>Lipostat 40</i> (RF).....	154, 155	MACROGOL-3350.....	52, 995
<i>Lipostat 80</i> (RF).....	154, 155	MACROGOL-3350 + SODIUM CHLORIDE +	
<i>Liquifilm Tears</i> (AG).....	949, 950	BICARBONATE + POTASSIUM CHLORIDE.....	53, 995
<i>Liquigen</i> (SB).....	956	<i>Macrovic</i> (RF).....	53, 995
LISDEXAMFETAMINE.....	880	<i>Madopar 125</i> (RO).....	842, 843
LISINOPRIL.....	139	<i>Madopar 62.5</i> (RO).....	843
<i>Lisinopril AN</i> (EA).....	139	<i>Madopar HBS</i> (RO).....	842
<i>Lisinopril generichealth</i> (GQ).....	139	<i>Madopar Rapid 125</i> (RO).....	843
<i>Lisinopril Sandoz</i> (SZ).....	139	<i>Madopar Rapid 62.5</i> (RO).....	842
<i>Lithicarb</i> (AS).....	876	<i>Madopar</i> (RO).....	842, 843
LITHIUM CARBONATE.....	876	<i>MagMin</i> (PBS)(BB).....	103
<i>Locasol</i> (SB).....	973	<i>Magmin</i> (BB).....	1020
<i>Loceryl</i> (GA).....	1022	MAGNESIUM.....	103, 1019
<i>Lofenoxal</i> (IL).....	56	<i>Mag-Sup</i> (PP).....	103, 1020
<i>Logem</i> (AL).....	836, 837	<i>Malarone</i> (GK).....	899
<i>Logicin Rapid Relief</i> (AS).....	1037	<i>Malean</i> (RW).....	138, 139
<i>Logicin Sinus</i> (AS).....	1037	<i>Marevan</i> (GT).....	107
<i>Logynon ED</i> (SY).....	193	<i>Marevan</i> (GT).....	107
<i>Lomotil</i> (IM).....	56	<i>Mavenclad</i> (SG).....	367
<i>Loniten</i> (PF).....	130	<i>Maviret</i> (VE).....	250
<i>Lonsurf 15/6.14</i> (SE).....	256, 257	<i>Maxalt</i> (AL).....	826
<i>Lonsurf 20/8.19</i> (SE).....	256, 257	<i>Maxamox</i> (SZ).....	215, 216
LOPERAMIDE.....	56, 1018	<i>Maxidex</i> (NV).....	923, 924
<i>Lopresor 100</i> (NV).....	135	<i>Maxolon</i> (IL).....	21, 44, 994
<i>Lopresor 50</i> (NV).....	135	<i>Maxor</i> (AF).....	36, 37, 38
<i>Lorano</i> (SZ).....	1038	<i>Maxor EC Tabs</i> (AF).....	37, 38, 39
LORATADINE.....	1038	<i>Mayne Pharma Doxycycline</i> (YT).....	213, 214
LORLATINIB.....	299	<i>Mayne Pharma Erythromycin</i> (YT).....	231
<i>Lorstat 10</i> (AF).....	153, 154	<i>Mayne Pharma Oxycodone IR</i> (YN).....	795, 796, 799
<i>Lorstat 20</i> (AF).....	153, 154	<i>Mayzent</i> (NV).....	372
<i>Lorstat 40</i> (AF).....	153, 154	<i>MCT Oil</i> (SB).....	956
<i>Lorstat 80</i> (AF).....	153, 154	<i>MCT Pro-Cal</i> (VF).....	990
<i>Lorviqua</i> (PF).....	300	MEBENDAZOLE.....	1036
LOSARTAN.....	146	MEBEVERINE.....	1016
<i>Losac Tablets</i> (AP).....	36, 37, 38, 39	<i>Medicianz Mupirocin Ointment</i> (DZ).....	1024
<i>Lovan</i> (AL).....	873	<i>Medipore 2961</i> (MM).....	1057
<i>Lovan 20 Tab</i> (AL).....	872		

MEDIUM CHAIN TRIGLYCERIDES	955, 956	<i>Metformin Sandoz (SZ)</i>	63, 64
MEDROXYPROGESTERONE		<i>Metformin XR 500 APOTEX (GX)</i>	64
ANTINEOPLASTIC AND IMMUNOMODULATING		METHADONE	
AGENTS	339	NERVOUS SYSTEM	812
GENITO URINARY SYSTEM AND SEX HORMONES		Palliative Care	1002
.....	194, 197, 198	METHENAMINE HIPPURATE	238
<i>MEDSURGE HYDROMORPHONE 2 mg/1 mL(DZ)</i>	782	<i>Methoblastin PFS(PF)</i>	757, 758
<i>MEDSURGE HYDROMORPHONE HP 10 mg/1 mL(DZ)</i>		<i>Methoblastin(PF)</i>	
.....	782	ANTINEOPLASTIC AND IMMUNOMODULATING	
MEFENAMIC ACID	765	AGENTS	255, 756
<i>Mefix 310250(MH)</i>	1057	<i>Methopt(AF)</i>	946, 947
<i>Megafof 0.5 (AF)</i>	123, 1021	METHOTREXATE	
<i>Megafof 5(AF)</i>	123, 1021	ANTINEOPLASTIC AND IMMUNOMODULATING	
<i>Mekinist(NV)</i>	301, 302	AGENTS	255, 756, 757, 758
<i>Mektovi(FB)</i>	300	<i>Methotrexate Accord(OD)</i>	255
<i>Melizide(AF)</i>	65	METHOXYFLURANE	20
MELOBIC (RF)	761, 762	METHYL SALICYLATE + MENTHOL + CAMPHOR +	
MELOBIC(RF)	762	EUCALYPTUS OIL + PINE OIL PUMILIO +	
<i>Melolin 36361357(SN)</i>	1056	TURPENTINE OIL + PEPPERMINT OIL + CAJUPUT	
<i>Melolin 66974933(SN)</i>	1056	OIL + CAPSICUM EXTRACT	1032
<i>Meloxiauro 15(DO)</i>	762	METHYLDOPA	128
<i>Meloxiauro 7.5(DO)</i>	761	METHYLNALTREXONE	996
<i>Meloxibell (GQ)</i>	761, 762	METHYLPHENIDATE	881, 882
MELOXICAM	761, 762	<i>Methylpred(AL)</i>	208
<i>Meloxicam AN(EA)</i>	761, 762	METHYLPREDNISOLONE	
<i>Meloxicam Sandoz (SZ)</i>	762	DERMATOLOGICALS	180, 181, 182
<i>Meloxicam Sandoz(SZ)</i>	761, 762	SYSTEMIC HORMONAL PREPARATIONS, EXCL. SEX	
<i>Meloxicam-GA (ED)</i>	761, 762	HORMONES AND INSULINS	208
MELPHALAN	253	<i>Methylprednisolone Alphapharm (AF)</i>	208
MEMANTINE	892, 893	METOCLOPRAMIDE	21, 44, 994
<i>Memantine generichealth(GQ)</i>	892, 893, 894	<i>Metoclopramide AN(EA)</i>	44
<i>Memanza (RW)</i>	892, 893	<i>Metoprolol AN(EA)</i>	135
<i>Mendelev Paracetamol(HX)</i>	823, 1034	<i>Metoprolol Sandoz (SZ)</i>	135
<i>Mepilex 294100(MH)</i>	1048	METOPROLOL SUCCINATE	134
<i>Mepilex Ag(MH)</i>	1047	METOPROLOL TARTRATE	135
<i>Mepilex Border 295200(MH)</i>	1048	<i>Metrogyl 200(AF)</i>	237
<i>Mepilex Border 295300(MH)</i>	1048	<i>Metrogyl 400(AF)</i>	238
<i>Mepilex Border Ag(MH)</i>	1047	<i>Metrol 100(RW)</i>	135
<i>Mepilex Border Flex 595211(MH)</i>	1047	<i>Metrol 50(RW)</i>	135
<i>Mepilex Border Flex 595311(MH)</i>	1047	<i>Metrol-XL 190(RW)</i>	135
<i>Mepilex Border Flex 595611(MH)</i>	1047	<i>Metrol-XL 23.75(RW)</i>	134
<i>Mepilex Border Heel 282750(MH)</i>	1047	<i>Metrol-XL 47.5(RW)</i>	135
<i>Mepilex Border Sacrum 282050(MH)</i>	1047	<i>Metrol-XL 95(RW)</i>	134
<i>Mepilex Border Sacrum 282450(MH)</i>	1047	METRONIDAZOLE	237, 238
<i>Mepilex Lite 284000(MH)</i>	1048	<i>Metronide 200 (AV)</i>	237
<i>Mepilex Lite 284100(MH)</i>	1048	<i>Metronide 400 (AV)</i>	238
<i>Mepitac 298300(MH)</i>	1058	<i>Mezavant(TK)</i>	57
<i>Mepitac 298400(MH)</i>	1058	<i>Miacalcic 100(EU)</i>	212
<i>Mepitel 290510(MH)</i>	1055	MIANSERIN	876
<i>Mepitel 290710(MH)</i>	1055	<i>Micardis Plus 40/12.5 mg(BY)</i>	149
<i>Mepitel One 289100(MH)</i>	1056	<i>Micardis Plus 80/12.5 mg(BY)</i>	149
<i>Mepitel One 289500(MH)</i>	1056	<i>Micardis Plus 80/25 mg(BY)</i>	149
MERCAPTOPURINE	255	<i>Micardis(BY)</i>	146
MERCAPTOPURINE-LINK(LM)	255	<i>Micolette(AE)</i>	54, 996, 1017
MESALAZINE	57, 58	MICONAZOLE	173
<i>Mesasal(GO)</i>	58	<i>Microdacyn(TF)</i>	1057
MESNA	954	<i>Microgynon 50 ED(BN)</i>	193
<i>Mestinon Timespan(IL)</i>	894	<i>Microlut 28(BN)</i>	193
<i>Mestinon(IL)</i>	894	<i>Micronelle 30 ED(TX)</i>	193
<i>Metalyse(BY)</i>	112	MIDAZOLAM	21
<i>Metamucil Natural Granular(PY)</i>	1017	MIFEPRISTONE (&) MISOPROSTOL	202
<i>Metamucil Orange Smooth(PY)</i>	1017	<i>Milivin OD 15(DO)</i>	877
METEX XR (RF)	64	<i>Milivin OD 30(DO)</i>	877
METEX XR 1000(RW)	64	<i>Milivin OD 45(DO)</i>	877
<i>Metex XR(RW)</i>	64	MILK POWDER SYNTHETIC LOW CALCIUM	973
METFORMIN	63, 64	MILK PROTEIN AND FAT FORMULA WITH VITAMINS	
<i>Metformin (Medsurge)(DZ)</i>	64	AND MINERALS CARBOHYDRATE FREE	988
<i>Metformin AN (EA)</i>	63, 64	<i>Minax 100 (AF)</i>	135
<i>Metformin AN(EA)</i>	64	<i>Minax 50 (AF)</i>	135
<i>Metformin GH(HQ)</i>	63, 64	<i>Minax XL (AF)</i>	134, 135

Minidiab (PF)	65	Monodur 60 mg(IY).....	127
Minipress (PF)	129	Monofeme 28(FZ).....	193
Minirin Melt(FP)	205	Monofer(PF)	121
Minirin Nasal Spray(FP).....	204, 205	Monogen(SB)	970
Minirin(FP)	204	MONTELAIR 4(RF)	920
Minitran 10(IL).....	126	MONTELAIR 5(RF)	921
Minitran 15(IL).....	126	MONTELUKAST	920
Minitran 5(IL).....	126	Montelukast AN (EA)	920, 921
MINOCYCLINE	214	Montelukast APOTEX(GX)	920, 921
Minomycin-50(AS)	215	Montelukast GH (GQ).....	920
MINOXIDIL	130	Montelukast Lupin (HQ).....	921
Mirena(BN)	191	Montelukast Lupin(HQ).....	920
MIRTANZA ODT (RF).....	877	Montelukast Mylan (AF).....	920
MIRTANZA(RF)	877	Montelukast Mylan(AF).....	921
MIRTAZAPINE.....	877	Montelukast Sandoz 4(SZ)	920
Mirtazapine AN (EA)	877	Montelukast Sandoz 5 (SZ)	921
Mirtazapine AN ODT(EA).....	877	MORPHINE	21, 784, 785, 787, 788, 790, 791, 792, 795, 997, 998, 1033
Mirtazapine GH(GQ)	877	Morphine Juno(JU).....	21, 784, 785, 787, 791
Mirtazapine Sandoz (SZ)	877	Morphine MR AN (EA).....	794
Mirtazapine Sandoz ODT 15 (SZ).....	877	MORPHINE MR APOTEX(TX)	794
Mirtazapine Sandoz ODT 30 (SZ).....	877	Morphine MR Mylan (AF)	794
Mirtazapine Sandoz ODT 45 (SZ).....	877	MORPHINE SULFATE 10 mg/1 mL MEDSURGE (DZ) 785, 791	
Mirtazapine Sandoz(SZ)	877	MORPHINE SULFATE 15 mg/1 mL MEDSURGE (DZ) ..21, 784, 787	
Mirtazon (RW).....	877	MORPHINE SULFATE 30 mg/1 mL MEDSURGE (DZ) ..21, 784, 787	
Mirtazon(RW).....	877	Motilium(JC)	44
Mistrom (ER).....	135	Movalis 15 (RW).....	762
Mixtard 30/70 InnoLet(NI).....	62	Movalis 15(RW).....	762
Mixtard 30/70 Penfill 3 mL (NO).....	62	Movalis 7.5 (RW).....	761
Mixtard 50/50 Penfill 3 mL(NO).....	62	Movalis 7.5(RW).....	762
Mizart (AF)	146	Movapo Pen(TD)	845
Mizart HCT 40/12.5 mg (AF).....	149	Movapo PFS(TD).....	845
Mizart HCT 80/12.5 mg (AF).....	149	Movapo(TD)	845
Mizart HCT 80/25 mg (AF).....	149	Movicol (NE).....	53, 995
MMA/PA Anamix infant(SB).....	979	Movicol Liquid(NE)	53, 995
MMA/PA Anamix Junior(NU)	979	Movox 100(AF).....	873
MMA/PA express 15(VF)	979	Movox 50(AL).....	873
MMA/PA gel(VF).....	979	Moxicam (AF).....	762
Mobic(BY)	761, 762	Moxicam 15(AF)	762
Mobilis 10 (AF).....	762	Moxicam 7.5(AF)	761
Mobilis 20 (AF).....	763	MOXONIDINE	129
Mobilis D-10(AF).....	763	Moxonidine GH (GQ).....	129
Mobilis D-20(AF).....	762	Moxonidine GX(SZ).....	129
MOCLOBEMIDE	875	Moxonidine MYL (AF).....	129
Moclobemide AN (EA)	875	Moxotens(RF).....	129
Moclobemide Sandoz(SZ)	875	MS Contin Suspension 100 mg(MF).....	794
Modafin (RW).....	884	MS Contin Suspension 20 mg(MF).....	794
MODAFINIL	883	MS Contin Suspension 200 mg(MF).....	792, 795
Modafinil AN(EA)	884	MS Contin Suspension 30 mg(MF).....	794
Modafinil GH (GQ).....	884	MS Contin Suspension 60 mg(MF).....	795
Modafinil Mylan(AF).....	884	MS Contin(MF).....	792, 794, 795, 998, 1033
Modafinil Sandoz (SZ)	884	MS Mono(MF)	794
Modavigil(TB).....	884	MS-2 Step(XH)	202
MODIFIED LONG CHAIN AMYLOPECTIN	970	MSUD AID III(SB).....	982
Moduretic(AS)	132	MSUD amino5(VF)	984
Mogadon(IL)		MSUD Anamix infant(SB)	982
. Palliative Care	1005	MSUD Anamix Junior LQ(SB)	983
Mogadon(IL)		MSUD Anamix Junior(SB)	983
. NERVOUS SYSTEM.....	830, 867, 868	MSUD cooler 10(VF)	982
Mohexal (HX).....	875	MSUD cooler 15(VF)	982
Molaxole(GO)	53, 995	MSUD cooler 20(VF)	982
Momasone (AS)	1025	MSUD Easy Tablet(OH)	977
Momasone(AS).....	182, 183, 184, 185	MSUD express 15(VF)	982
MOMETASONE.....	182, 183, 184, 1025	MSUD express 20(VF)	982
Momex SR 10(RW).....	794	MSUD gel(VF)	982
Momex SR 100(RW).....	794	MSUD Lophlex LQ 20(SB)	982
Momex SR 30(RW).....	794	MSUD Lophlex(SB)	982
Momex SR 60(RW).....	794		
Monace 10(AF).....	139		
Monace 20(AF).....	139		
Monodur 120 mg(IY).....	126		

MSUD Maxamum(SB)	982
MUPIROCIN	
Repatriation Pharmaceutical Benefits Scheme	1024
RESPIRATORY SYSTEM	901
Mupirocin Nasal (Medsurge)(DZ)	902
Murelax (RW)	866, 867, 1004
MYCOBACTERIUM BOVIS (BACILLUS CALMETTE AND GUERIN (BCG)) TICE STRAIN	350
MycoCept (RF)	370
Myconail(AE)	1022
MYCOPHENOLATE	370
Mycophenolate AN(EA)	370
Mycophenolate APOTEX (GX)	370
Mycophenolate GH(GQ)	370
Mycophenolate Sandoz (SZ)	370
Mycophenolate Sandoz(SZ)	370
Mycostatin Oral Drops(LN)	1016
Myfortic(NV)	370
Myleran(AS)	253
Mysoline(LM)	828
NAFARELIN	205
NALOXONE	21, 952
Naloxone Hydrochloride (DBL) (PF)	21, 952
Naloxone Juno(JU)	21, 952
NALOXONE SXP (XC)	21, 952
NALTREXONE	897
Naltrexone GH (GQ)	898
Naprosyn SR1000(IX)	764, 765, 997
Naprosyn SR750(IX)	764, 765, 997
Naprosyn(IX)	764, 765, 997
NAPROXEN	764, 765, 997
Naramig(AS)	825, 826
NARATRIPTAN	825
NARCAN(FE)	21
Nardil(LM)	875
Natrilix SR(SE)	130
Natrilix(SE)	130
Navelbine(FB)	257
Nebilet (FK)	135
NEBIVOLOL	135
Nebivolol Sandoz(SZ)	135
NEDOCROMIL	918
Neo-B12 (PF)	122
Neo-B12(PF)	1021
Neocate Gold(SB)	962, 963, 964
Neocate Junior Vanilla(SB)	957, 958, 959
Neocate Junior(SB)	972, 973
Neocate LCP(SB)	960, 961
Neocate Syneo(SB)	974, 975, 976
Neo-Mercazole(GH)	210
Neoral 10(NV)	754
Neoral 100 (NV)	754
Neoral 25 (NV)	754
Neoral 50 (NV)	754
Neoral(NV)	754
Neotigason(TB)	175
Nepiten (AF)	135
Nesina Met 12.5/1000(TK)	66
Nesina Met 12.5/500(TK)	66
Nesina Met 12.5/850(TK)	66
Nesina(TK)	85
NETUPITANT + PALONOSETRON	45
Neulactil(SW)	850
Neupro(UC)	848
Neuroccord (CR)	824
Neurontin (UJ)	834, 1034
Neurontin(UJ)	
Repatriation Pharmaceutical Benefits Scheme	1034, 1035
Neurontin(UJ)	

NERVOUS SYSTEM	834
Nexavar(BN)	325, 326
Nexazole (RW)	30, 31, 32, 33, 34
Nexcare Durable Cloth First Aid Tape 799(MM)	1058
Nexcare Gentle Paper First Aid Tape 789(MM)	1058
Nexcare Tegaderm Transparent H1624(MM)	1045
Nexcare Tegaderm Transparent H1626(MM)	1045
Nexium Hp7(AP)	43
Nexium(AP)	30, 31, 32, 33, 34
Nexole(RF)	30, 31, 32, 33, 34
Nicabate CQ 14(GC)	1036
Nicabate CQ 21(GC)	1036
Nicabate P(GC)	896
NICORANDIL	127
nicorette 16hr Invisipatch(JT)	896
NICOTINE	895, 1036
Nicotinell Step 1(ON)	895
Nicotinell Step 2(ON)	895
Nicotinell Step 3(ON)	895
Nicotinell(ON)	896
Nidem(RW)	64
NIFEDIPINE	137
NILOTINIB	280, 282
Nilstat(AS)	
Repatriation Pharmaceutical Benefits Scheme	1027
Nilstat(AS)	
ALIMENTARY TRACT AND METABOLISM	54, 55
NILUTAMIDE	345
NINTEDANIB	321
NITRAZEPAM	
NERVOUS SYSTEM	829, 867
Palliative Care	1005
NITROFURANTOIN	238
Nitrolingual Pumpspray(SW)	20, 126
Nizac(RF)	29
NIZATIDINE	29
Nizoral 2% Cream(JT)	173
Nizoral 2%(JT)	173
Nolvadex-D (AP)	344
Nolvadex-D(AP)	343
Nordette 28(PF)	193
Nordip(AF)	136
NORETHISTERONE	
GENITO URINARY SYSTEM AND SEX HORMONES	194, 198
NORETHISTERONE + ETHINYLESTRADIOL	193
NORETHISTERONE + MESTRANOL	193
NORETHISTERONE ACETATE + ESTRADIOL (&)	
ESTRADIOL	199
NORFLOXACIN	236
Noriday 28 Day(PF)	194
Norimin 28 Day(FZ)	193
Norimin-1 28 Day(FZ)	193
Norinyl-1/28(PF)	193
Normacol Plus(NE)	52, 995, 1017
Normison(AS)	868, 1005
Norprolac(FP)	192
Norspan(MF)	814, 815, 1002, 1003
NORTRIPTYLINE	870
NortriTABS 10 mg (GH)	870
NortriTABS 25 mg (GH)	870
Norvapine (ED)	136
Norvasc(AS)	136
Noten (AF)	134
NOUMED ALLOPURINOL (VO)	768
NOUMED AMOXICILLIN (VO)	216, 217
NOUMED AMOXICILLIN(VO)	215
NOUMED ATORVASTATIN(VO)	153, 154
NOUMED CEFALEXIN(VO)	223, 224, 225
NOUMED CIPROFLOXACIN(VO)	235

NOUMED CITALOPRAM (VO)	871	Olanzapine Sandoz ODT 20(SZ)	854
NOUMED DIAZEPAM(VO)	866, 1004	Olanzapine Sandoz ODT 5 (SZ)	853
NOUMED ESOMEPRAZOLE (VO)	30, 31, 32, 33, 34	Olanzapine-DRLA (RZ)	852, 853
NOUMED LAMOTRIGINE (VO)	836, 837	OLAPARIB	335, 336, 337
NOUMED LANSOPRAZOLE(VO)	34, 35, 36	Olmekar HCT 20/5/12.5 (RF)	152
NOUMED LEVETIRACETAM (VO)	837	Olmekar HCT 40/10/12.5 (RF)	152
NOUMED PANTOPRAZOLE (VO)	39	Olmekar HCT 40/10/25 (RF)	152
NOUMED PANTOPRAZOLE(VO)	39, 40, 41	Olmekar HCT 40/5/12.5 (RF)	152
NOUMED PERINDOPRIL (VO)	139, 140	Olmekar HCT 40/5/25 (RF)	152
NOUMED PIOGLITAZONE(VO)	84	OLMEKAR(RW)	150, 151
NOUMED RISPERIDONE (VO)	859, 860, 861, 862, 863, 864, 865	OLMERTAN (RW)	146
Noumed Rosuvastatin (VO)	155, 156, 157	OLMERTAN COMBI 20/12.5 (RW)	148
NOUMED SERTRALINE (VO)	873, 874	OLMERTAN COMBI 40/12.5 (RW)	148
NOUMED SERTRALINE(VO)	874	OLMERTAN COMBI 40/25 (RW)	148
NOUMED Sildenafil (VO)	1028, 1029	OLMESARTAN	146
NOUMED TERBINAFINE(VO)	174	Olmesartan - MYL (AF)	146
Novacodone(HX)	801	OLMESARTAN + AMLODIPINE + HYDROCHLOROTHIAZIDE	152
Novasone (AF)	182, 183, 184, 185	Olmesartan AN(EA)	146
Novatears(AE)	949	Olmesartan HCT - MYL 20/12.5 (AF)	148
Novatin (TX)	175	Olmesartan HCT - MYL 40/12.5 (AF)	148
NovoMix 30 FlexPen(NF)	62	Olmesartan HCT - MYL 40/25 (AF)	148
NovoMix 30 Penfill 3 mL(NO)	62	Olmesartan HCT AN 20/12.5(EA)	148
NovoRapid FlexPen(NF)	61	Olmesartan HCT AN 40/12.5(EA)	148
NovoRapid Penfill 3 mL(NO)	61	Olmesartan HCT AN 40/25(EA)	148
NovoRapid(NO)	61	OLMESARTAN MEDOXOMIL + AMLODIPINE	150
Noxicid Caps (AL)	30, 31, 32, 33, 34	OLMESARTAN MEDOXOMIL + HYDROCHLOROTHIAZIDE	148
Nuelin(IL)	920	Olmesartan Sandoz(SZ)	146
Nuelin-SR 200(IL)	920	Olmesartan/Amlodipine - MYL 20/5(AF)	151
Nuelin-SR 250(IL)	920	Olmesartan/Amlodipine - MYL 40/10(AF)	150
Nuelin-SR 300(IL)	920	Olmesartan/Amlodipine - MYL 40/5(AF)	150
Nufloxib(AF)	236	Olmesartan/Amlodipine 20/5 APOTEX (TX)	151
Nu-Gel 2497(KI)	1055	Olmesartan/Amlodipine 40/10 APOTEX (TX)	150
Nupentin 100 (AF)		Olmesartan/Amlodipine 40/5 APOTEX (TX)	150
.Repatriation Pharmaceutical Benefits Scheme	1035	Olmesartan/HCT Sandoz(SZ)	148
Nupentin 100 (AF)		Olmetec Plus(AL)	148
.NERVOUS SYSTEM	834	Olmetec(AL)	146
Nupentin 300 (AF)		OLSALAZINE	59
.Repatriation Pharmaceutical Benefits Scheme	1034	Olumiant(LY)	362, 367
Nupentin 300(AF)		Omegapharm Pty Ltd (OE)	229
.NERVOUS SYSTEM	834	Omepral (ZA)	37, 38, 39
Nupentin 400 (AF)		OMEPRAZOLE	36, 37, 38
.Repatriation Pharmaceutical Benefits Scheme	1034	Omeprazole AN(EA)	37, 38, 39
Nupentin 400(AF)		Omeprazole generichealth (GQ)	37, 38, 39
.NERVOUS SYSTEM	834	Omeprazole Sandoz(HX)	36, 37, 38
Nutrini Peptisorb Energy(NU)	957	Omeprazole Sandoz(SZ)	37, 38, 39
Nutrini Peptisorb(SB)	970	Onbrez(NV)	902
Nuvigil(TB)	879	OncoTICE(MK)	350
NYSTATIN		ONDANSETRON	46, 47
.ALIMENTARY TRACT AND METABOLISM	54	Ondansetron AN (EA)	46
.Repatriation Pharmaceutical Benefits Scheme	1016, 1027	Ondansetron AN ODT (EA)	47, 48
Nyxoid(MF)	952	Ondansetron APOTEX(GX)	46
O.R.S.(AF)	55	Ondansetron Mylan ODT(AF)	47, 48
OCRIPLASMIN	947	Ondansetron Mylan Tablets(AF)	46
Ocuflox(AG)	923	Ondansetron ODT GH(GQ)	47, 48
Odaplix SR(AF)	130	Ondansetron ODT-DRLA (RZ)	47, 48
Odomzo(RA)	334	Ondansetron SZ (HX)	46
Ofev(BY)	322	Ondansetron SZ ODT (HX)	47, 48
OFLOXACIN	923	Ondansetron-DRLA (RZ)	46
OLANZAPINE	852, 853, 854	Onglyza(AP)	87, 89
Olanzapine AN (EA)	852, 853	Op-Site Flexigrid 4629(SN)	1045
Olanzapine AN ODT (EA)	853, 854	Optifresh eye gel(PP)	943
Olanzapine APOTEX(GX)	852, 853	Optifresh Plus(PP)	944
Olanzapine ODT generichealth 10(GQ)	853	Optifresh Tears(PP)	944
Olanzapine ODT generichealth 5(GQ)	853	Optisulin SoloStar(WA)	63
Olanzapine RBX(RA)	852, 853	Optisulin(GZ)	63
Olanzapine Sandoz (SZ)	852, 853	Optive(AG)	945
Olanzapine Sandoz ODT 10 (SZ)	853	Orabase(AS)	1023
Olanzapine Sandoz ODT 15(SZ)	854	Oratane (RF)	186

Oratane(OU)	186
Oratane(RF)	186
Ordine 10(MF)	785, 788
Ordine 2(MF)	785, 788
Ordine 5(MF)	785, 788
Orencia ClickJect(BQ)	353, 358
Orencia(BQ)	353, 358
ORLISTAT	1018
Oroxine(AS)	210
ORTHO-DICHLOROBENZENE + PARA-DICHLOROBENZENE + CHLOROBUTANOL + ARACHIS OIL	1038
Orudis SR 200(SW)	764
Oruvail SR(AV)	763, 764
OSIMERTINIB	291, 292
OsmoLax(KY)	53, 995
Ospolot(FF)	840
Osteomol 665 Paracetamol (CR)	824, 1003
Osteomol 665 Paracetamol(CR)	824, 1003, 1034
Osteovan (SZ)	771
Otezla(AN)	359
Otocomb Otic(LN)	951
Otodex(AV)	951
Ovestin Ovula(AS)	197
Ovestin(AS)	197
OXAZEPAM	866, 867, 1004
OXCARBAZEPINE	831
Oxis Turbuhaler(AP)	902
OXPRENOLOL	133
OXYBUTYNIN	202
Oxybutynin Sandoz (SZ)	202
OXYCODONE	795, 796, 797, 798, 799, 801
OXYCODONE + NALOXONE	802
Oxycodone BNM(BZ)	796, 797, 799
Oxycodone Mylan (AL)	795, 796, 799
Oxycodone Sandoz (SZ)	801
OxyContin(MF)	801, 802
OXYMETAZOLINE	1037
OxyNorm (MF)	796, 797, 799
OxyNorm Liquid 1mg/mL(MF)	796, 799
OxyNorm(MF)	795, 796
Oxytrol(TT)	202
OZANIMOD	370
Ozempic(NO)	96
Ozidal(RA)	860, 861, 862, 863, 864, 865
Ozin 10(DO)	853
Ozin 2.5(DO)	852
Ozin 5(DO)	852
Ozin 7.5(DO)	852
Ozmep(ZP)	37, 38, 39
Ozole (RA)	240
Ozole(RA)	240
Ozpan (RA)	39, 40, 41
Ozpan(RA)	39
Ozurdex(AG)	923, 925, 926
PAA (UL)	943
Pacrolim(AF)	755
PAD WOUND DEBRIDEMENT	1057
Paediatric Seravit(SB)	991
PALBOCICLIB	303
Palexia SR(CS)	818, 819
PALIPERIDONE	858, 859
PALONOSETRON	48
PAMIDRONATE DISODIUM	769
Pamisol(PF)	769, 770
Panadeine Forte(SW)	815, 816, 817
Panadol(GC)	1003
Panafcort(AS)	209
Panafcortelone(AS)	208
Panamax (SW)	823, 1034

Panamax 240 Elixir(SW)	823, 1034
Panamax Co. 40(SW)	1033
Panamax(SW)	823
PANCREATIC EXTRACT	60
PANCRELIPASE	60
PANTHENOL	1027
Panthron (ER)	39
Panthron(ER)	39, 40, 41
PANTOPRAZOLE	39, 40
Pantoprazole Actavis (ED)	39, 40, 41
Pantoprazole AN(EA)	39, 40, 41
Pantoprazole APOTEX (TY)	39, 40, 41
Pantoprazole generichealth(HQ)	39, 40, 41
Pantoprazole Sandoz (SZ)	39, 40, 41
Panzylrat 25000(TM)	60, 61
PAPER WASP VENOM	952
PARACETAMOL	823, 824, 1003, 1034
Paracetamol (Sandoz)(SZ)	823, 1034
PARACETAMOL + CODEINE	815, 816, 1033
Paracetamol/Codeine GH 500/30 (GQ)	815, 816, 817
PARAFFIN	948
Paralgin (OW)	823, 1034
Parapane(AF)	823, 824, 1034
Parbezol (RW)	41, 42, 43
Pariet(JC)	41, 42, 43
Parlodel(SZ)	
.GENITO URINARY SYSTEM AND SEX HORMONES	191, 192
.NERVOUS SYSTEM	846
Parnate(GH)	875
PAROXETINE	873
Paroxetine AN(EA)	873
Paroxetine GH (GQ)	873
Paroxetine Sandoz(SZ)	873
Pavtide (GC)	910, 911
PAVTIDE ACCUHALER 100/50(GC)	910
PAVTIDE ACCUHALER 250/50(GC)	910
PAVTIDE ACCUHALER 500/50(GC)	911
PAVTIDE MDI 50/25(GC)	910
Paxam 0.5(AF)	829, 1004
Paxam 2(AF)	829, 1004
Paxtine (AF)	873
PAZOPANIB	322, 323, 324
Peg 7420(MM)	1040
Peg 7422(MM)	1040
Peg 7423(MM)	1040
Peg 7425(MM)	1040
Pegasys(RO)	349
PEGINTERFERON ALFA-2A	349
PEGINTERFERON BETA-1A	349
Pemzo (RW)	36, 37, 38
PENICILLAMINE	767
Pentasa(FP)	57, 58
Penthrox(DV)	20
Peptamen Junior(NT)	970
PERAMPANEL	838, 839
PERFLUOROHEXYLOCTANE	948
PERHEXILINE	127
PERICIAZINE	850
Perindo Combi 4/1.25(AF)	142
Perindo(AF)	139, 140
PERINDOPRIL	139, 140
PERINDOPRIL + AMLODIPINE	143
PERINDOPRIL + INDAPAMIDE	142
Perindopril Actavis 2 (EA)	139
Perindopril Actavis 4 (ED)	140
Perindopril Actavis 8 (ED)	140
Perindopril AN(EF)	139, 140
Perindopril and Indapamide AN 4/1.25 (EF)	142
Perindopril APOTEX (TY)	139, 140

<i>Perindopril Combi Actavis 4/1.25(ED)</i>	142	<i>Pharmacor Quetiapine 300(CR)</i>	855
<i>Perindopril generichealth(GQ)</i>	140	<i>Pharmacor Rasagiline (CR)</i>	848
<i>Perindopril/ Indapamide GH 4/1.25 (GQ)</i>	142	<i>Pharmacor Riluzole (CR)</i>	898
<i>PERMETHRIN</i>	901	<i>Pharmacor Rosuvastatin 10(CR)</i>	156, 157
<i>Petrus Bisacodyl Suppositories(PP)</i>	52, 994, 1016	<i>Pharmacor Rosuvastatin 20(CR)</i>	156
<i>Petrus Pharmaceuticals Pty Ltd(PP)</i>	1017	<i>Pharmacor Rosuvastatin 40(CR)</i>	156, 157
<i>Pexsig(AS)</i>	127	<i>Pharmacor Rosuvastatin 5(CR)</i>	155, 156
<i>Pfizer Australia Pty Ltd(PF)</i>		<i>Pharmacor Sumatriptan 50(CR)</i>	826
<i>.Prescriber Bag</i>	19, 20, 21	<i>Pharmacor Tacrolimus 0.5 (CR)</i>	755
<i>Pfizer Australia Pty Ltd(PF)</i>		<i>Pharmacor Tacrolimus 1 (CR)</i>	755
<i>.ALIMENTARY TRACT AND METABOLISM</i>	43	<i>Pharmacor Tacrolimus 5 (CR)</i>	755
<i>Pfizer Australia Pty Ltd(PF)</i>		<i>Pharmacor Telmisartan 40(CR)</i>	146
<i>.ALIMENTARY TRACT AND METABOLISM</i>	44	<i>Pharmacor Telmisartan 80(CR)</i>	146
<i>Pfizer Australia Pty Ltd(PF)</i>		<i>Pharmacor Teriflunomide(CR)</i>	373
<i>.BLOOD AND BLOOD FORMING ORGANS</i>	109	<i>Pharmacor Zopiclone (CR)</i>	1036
<i>Pfizer Australia Pty Ltd(PF)</i>		<i>Pharmacy Action Anti-Fungal Cream(GQ)</i>	1022
<i>.ANTIINFECTIVES FOR SYSTEMIC USE</i>	233	<i>Pharmacy Action Anti-Fungal Nail Treatment(GQ)</i>	1022
<i>Pfizer Australia Pty Ltd(PF)</i>		<i>Pharmacy Action Cetrelief(GQ)</i>	1038
<i>.ANTIINFECTIVES FOR SYSTEMIC USE</i>	234	<i>Pharmacy Action Diarrhoea Relief (GQ)</i>	1018
<i>Pharmacor Amlodipine(CR)</i>	136	<i>Pharmacy Action FemCream (GQ)</i>	1027
<i>Pharmacor Atorvastatin (CR)</i>	153, 154	<i>Pharmacy Action FemCream 3 Day Cream(GQ)</i>	1027
<i>Pharmacor Cefuroxime(CR)</i>	227	<i>Pharmacy Action Fexorelief 120(GQ)</i>	1038
<i>Pharmacor Cinacalcet (CR)</i>	212	<i>Pharmacy Action Hydrocortisone Cream 1%(GQ)</i>	1024
<i>Pharmacor Cyproterone 100 (CR)</i>		<i>Pharmacy Action Laxative with Senna(GQ)</i>	1017
<i>.ANTINEOPLASTIC AND IMMUNOMODULATING</i>		<i>Pharmacy Action Lorastyne(GQ)</i>	1038
<i>AGENTS</i>	344	<i>Pharmacy Action Low Dose Aspirin (GQ)</i>	1020
<i>.GENITO URINARY SYSTEM AND SEX HORMONES</i>		<i>Pharmacy Action Nasal Decongestant(GQ)</i>	1037
.....	201	<i>Pharmacy Action Nystatin Oral Drops(GQ)</i>	1016
<i>Pharmacor Cyproterone 50(CR)</i>		<i>Pharmacy Action Paracetamol Osteo 665 (GQ)</i>	1034
<i>.ANTINEOPLASTIC AND IMMUNOMODULATING</i>		<i>Pharmacy Action Pharmisil (GQ)</i>	1022
<i>AGENTS</i>	344	<i>Pharmacy Action Sinus & Nasal Decongestant Relief(GQ)</i>	
<i>.GENITO URINARY SYSTEM AND SEX HORMONES</i>			1037
.....	201	<i>Pharmacy Action Worm Treatment(GQ)</i>	1036
<i>Pharmacor Ezetimibe 10 (CR)</i>	166, 167	<i>Pharmacy Care Magnesium(SI)</i>	1020
<i>Pharmacor Ezetimibe Simvastatin 10/10(CR)</i>	172	<i>PHARMACY CARE PARACETAMOL (SI)</i>	823, 824, 1034
<i>Pharmacor Ezetimibe Simvastatin 10/20(CR)</i>	172	<i>Phebra Naproxen Suspension(FF)</i>	764, 997
<i>Pharmacor Ezetimibe Simvastatin 10/40(CR)</i>	171	<i>Pheburane(OH)</i>	106
<i>Pharmacor Ezetimibe Simvastatin 10/80(CR)</i>	171	<i>PHENELZINE</i>	874
<i>Pharmacor Finasteride 5 (CR)</i>	1031	<i>Phenelzine sulfate USP (Generic Health)(GQ)</i>	875
<i>Pharmacor Gabapentin 600 (CR)</i>		<i>Phenobarb(RW)</i>	827
<i>.Repatriation Pharmaceutical Benefits Scheme</i>	1034	<i>PHENOBARBITAL (PHENOBARBITONE)</i>	827
<i>Pharmacor Gabapentin 600(CR)</i>		<i>Phenobarbitone Injection (Aspen Pharmacare Australia Pty Ltd)(AS)</i>	828
<i>.NERVOUS SYSTEM</i>	834	<i>PHENOXYBENZAMINE</i>	
<i>Pharmacor Gabapentin 800(CR)</i>		<i>.CARDIOVASCULAR SYSTEM</i>	133
<i>.Repatriation Pharmaceutical Benefits Scheme</i>	1034	<i>.GENITO URINARY SYSTEM AND SEX HORMONES</i>	
<i>Pharmacor Gabapentin 800(CR)</i>			202
<i>.NERVOUS SYSTEM</i>	834	<i>PHENOXYMETHYLPENICILLIN</i>	218, 219
<i>Pharmacor Gliclazide MR (CR)</i>	64	<i>Phenoxymethylpenicillin-AFT(AE)</i>	218
<i>Pharmacor Gliclazide MR(CR)</i>	64	<i>Phenylalanine 50(VF)</i>	989
<i>Pharmacor Letrozole 2.5 (CR)</i>	346	<i>PHENYLALANINE WITH CARBOHYDRATE</i>	988
<i>Pharmacor Lurasidone (CR)</i>	851	<i>PHENYTOIN</i>	828
<i>Pharmacor Meloxicam 15 (CR)</i>	762	<i>Phlexy-10(SB)</i>	984
<i>Pharmacor Meloxicam 7.5 (CR)</i>	761	<i>Phlexy-Vits(SB)</i>	991
<i>Pharmacor Metformin XR (CR)</i>	64	<i>PHOSPHATE PHEBRA(FG)</i>	955
<i>Pharmacor Metformin XR(CR)</i>	64	<i>PHOSPHORUS</i>	954
<i>Pharmacor Mycophenolate 250 (CR)</i>	370	<i>Physeptone(AS)</i>	813
<i>Pharmacor Mycophenolate 500(CR)</i>	370	<i>Physiotens (GO)</i>	129
<i>Pharmacor Olmesartan 20 (CR)</i>	146	<i>PHYTOMENADIONE</i>	21
<i>Pharmacor Olmesartan 40 (CR)</i>	146	<i>Piax (AF)</i>	109, 110, 1020
<i>Pharmacor Olmesartan Amlodipine 20/5 (CR)</i>	151	<i>Piax Plus Aspirin(AF)</i>	111
<i>Pharmacor Olmesartan Amlodipine 40/10 (CR)</i>	150	<i>PILOCARPINE</i>	928
<i>Pharmacor Olmesartan Amlodipine 40/5 (CR)</i>	150	<i>PIMECROLIMUS</i>	189
<i>Pharmacor Olmesartan HCTZ 20/12.5 (CR)</i>	148	<i>PINDOLOL</i>	133
<i>Pharmacor Olmesartan HCTZ 40/12.5 (CR)</i>	148	<i>Pinetarsol(EO)</i>	1024
<i>Pharmacor Olmesartan HCTZ 40/25 (CR)</i>	148	<i>PIOGLITAZONE</i>	83
<i>Pharmacor Omeprazole 20(CR)</i>	36, 37, 38	<i>Pioglitazone AN (EA)</i>	84
<i>Pharmacor Posaconazole(CR)</i>	242	<i>Pioglitazone AN(EA)</i>	84
<i>Pharmacor Quetiapine 100(CR)</i>	856	<i>Pioglitazone Sandoz(SZ)</i>	84
<i>Pharmacor Quetiapine 200(CR)</i>	855	<i>PIRFENIDONE</i>	758, 759
<i>Pharmacor Quetiapine 25(CR)</i>	856		

<i>Pirmella 1/35(DZ)</i>	193	<i>PRAVASTATIN</i>	154, 155
<i>PIROXICAM</i>	762	<i>Pravastatin AN (EA)</i>	154, 155
<i>PIZOTIFEN</i>	827	<i>Pravastatin AN(EA)</i>	154, 155
<i>PKU Air 15(VF)</i>	980	<i>Pravastatin generichealth (GQ)</i>	154, 155
<i>PKU Air 20(VF)</i>	980	<i>Pravastatin generichealth(GQ)</i>	155
<i>PKU Anamix First Spoon(SB)</i>	983	<i>Pravastatin Sandoz (SZ)</i>	155
<i>PKU Anamix infant(SB)</i>	983	<i>Pravastatin Sandoz(SZ)</i>	154, 155
<i>PKU Anamix Junior LQ(SB)</i>	980	<i>PRAZQUANTEL</i>	900
<i>PKU Anamix Junior(SB)</i>	981	<i>PRazosin</i>	129
<i>PKU Bettermilk Lite(QH)</i>	986	<i>PredMix(LN)</i>	208
<i>PKU Build 10(QH)</i>	987	<i>Prednefrin Forte(AG)</i>	927
<i>PKU Build 20(QH)</i>	987	<i>PREDNISOLONE</i>	208
<i>PKU Cooler 10(VF)</i>	980	<i>PREDNISOLONE ACETATE + PHENYLEPHRINE</i>	927
<i>PKU Cooler 15(VF)</i>	980	<i>PREDNISOLONE SODIUM PHOSPHATE</i>	
<i>PKU Cooler 20(VF)</i>	980	ALIMENTARY TRACT AND METABOLISM	56
<i>PKU Easy Microtabs(OH)</i>	976	SYSTEMIC HORMONAL PREPARATIONS, EXCL. SEX	
<i>PKU Easy Shake & Go(OH)</i>	977	HORMONES AND INSULINS	208
<i>PKU Easy Tablet(OH)</i>	976	<i>PREDNISONE</i>	209
<i>PKU Easy(OH)</i>	989	<i>Predsol(AS)</i>	56, 57
<i>PKU Explore 10(VF)</i>	983	<i>Predsolone(LN)</i>	208
<i>PKU Explore 5(VF)</i>	983	<i>Predsone(LN)</i>	209
<i>PKU express 15(VF)</i>	980	<i>PREGABALIN</i>	824
<i>PKU express 20(VF)</i>	980	<i>Pregabalin AMNEAL(EA)</i>	824
<i>PKU gel(VF)</i>	980	<i>Pregabalin APOTEX (GX)</i>	824
<i>PKU Glytactin RTD 15 Lite(QH)</i>	986	<i>Pregabalin GH (GQ)</i>	824
<i>PKU Glytactin RTD 15(QH)</i>	986	<i>Pregabalin Sandoz(SZ)</i>	824
<i>PKU GMPPro LQ(SB)</i>	987	<i>PREGABALIN-DRLA(RZ)</i>	824
<i>PKU Go(OH)</i>	976	<i>Pregabalin-Teva (TB)</i>	824
<i>PKU Lophlex LQ 10(SB)</i>	980	<i>Pregnyl(OQ)</i>	201
<i>PKU Lophlex LQ 20(SB)</i>	980	<i>Prenoxad(FF)</i>	952
<i>PKU Lophlex Sensation 20(SB)</i>	980	<i>Presolol 100(AF)</i>	136
<i>PKU Lophlex(SB)</i>	980	<i>PREXUM 10 (RW)</i>	140
<i>PKU Restore(QH)</i>	986	<i>PREXUM 2.5 (RW)</i>	139
<i>PKU Sphere15(VF)</i>	987	<i>PREXUM 5 (RW)</i>	140
<i>PKU Sphere20(VF)</i>	987	<i>Prexum Combi 5/1.25(RW)</i>	142
<i>PKU squeeze(VF)</i>	980	<i>PREXUM Combi LD 2.5/0.625(RW)</i>	142
<i>PKU Start(VF)</i>	983	<i>Prilace (RF)</i>	140, 141
<i>Placil(AF)</i>	869	<i>Prilace(RF)</i>	141
<i>Plaquacide(OB)</i>	1016	<i>PRIMIDONE</i>	828
<i>Plaquenil(SW)</i>	766, 767	<i>Primolut N(BN)</i>	198
<i>Plavacor 75(CR)</i>	110, 1020	<i>Pristiq(PF)</i>	875, 876
<i>Plegriby(BD)</i>	349	<i>Pritor/Amlodipine(FI)</i>	151
<i>Plendil ER(GX)</i>	137	<i>Pro-Banthine(RW)</i>	202
<i>Plidogrel (RF)</i>	109, 110, 1020	<i>PROBENECID</i>	768
<i>Poly Gel(AQ)</i>	943	<i>Probitor (SZ)</i>	36, 37, 38
<i>Poly Visc(IQ)</i>	948	<i>PROCAINE BENZYL PENICILLIN (PROCAINE</i>	
<i>POLYETHYLENE GLYCOL-400 + PROPYLENE GLYCOL</i>		<i>PENICILLIN)</i>	19, 219
.....	949	<i>ProCalm (RW)</i>	50
<i>POLYLACTIC ACID</i>	955	<i>PROCHLORPERAZINE</i>	21, 50
<i>Poly-Tears(IQ)</i>	945, 946	<i>Prochlorperazine AN(EA)</i>	50
<i>POLYVINYL ALCOHOL</i>	949, 950	<i>Prochlorperazine GH (GQ)</i>	50
<i>PONATINIB</i>	284, 285, 286	<i>Pro-Cid(FF)</i>	768
<i>Ponstan(PF)</i>	765	<i>Prodeine Forte(AV)</i>	815, 816, 817
<i>POSACONAZOLE</i>	242	<i>Profore 66050016(SN)</i>	1040
<i>Posaconazole ARX (XT)</i>	242	<i>Profore Lite 66050415(SN)</i>	1040
<i>POSACONAZOLE DR.REDDY'S(RZ)</i>	242	<i>Progout 100(AF)</i>	768
<i>Posaconazole Juno (JU)</i>	242	<i>Progout 300(AF)</i>	768
<i>Posaconazole Sandoz(SZ)</i>	242	<i>Prograf(LL)</i>	755
<i>POTASSIUM CHLORIDE</i>	103	<i>Progynova(BN)</i>	196
<i>POTASSIUM CHLORIDE + POTASSIUM BICARBONATE</i>		<i>Proladone(FF)</i>	796, 798
+ <i>POTASSIUM CARBONATE</i>	103	<i>Prolia(AN)</i>	775
<i>POVIDONE-IODINE</i>		<i>PROMETHAZINE</i>	
<i>.Repatriation Pharmaceutical Benefits Scheme</i>	1025,	ALIMENTARY TRACT AND METABOLISM	50
1057		<i>.Prescriber Bag</i>	21
<i>Pradaxa(BY)</i>	113, 114	RESPIRATORY SYSTEM	921
<i>Pramin (AF)</i>	44	<i>Prontosan Wound Irrigation Solution(BR)</i>	1043
<i>PRAMIPEXOLE</i>	846, 847	<i>PROPANTHELINE</i>	202
<i>Pramipexole AN (EA)</i>	846, 847	<i>PROPRANOLOL</i>	133
<i>Pramipexole XR GP (AF)</i>	847	<i>PROPYLTHIOURACIL</i>	210
<i>Pravachol(RW)</i>	154, 155	<i>Proscar(OQ)</i>	1031

<i>Protaphane</i> (NO)	62	<i>Quetiapine Sandoz Pharma</i> (HX)	855, 856
<i>Protaphane InnoLet</i> (NI)	62	<i>Quetiapine Sandoz Pharma</i> (HX)	856
<i>Protaphane Penfill</i> 3 mL (NO).....	62	<i>Quetiapine Sandoz</i> (SZ)	855, 856
PROTEIN FORMULA WITH AMINO ACIDS, CARBOHYDRATES, VITAMINS AND MINERALS WITHOUT PHENYLALANINE, AND SUPPLEMENTED WITH DOCOSAHEXAENOIC ACID	989	QUETIAPINE-AS XR (RW)	855, 856
PROTEIN FORMULA WITH ARGININE, VITAMIN C AND E	1039	<i>Quetiapine-DRLA</i> (RZ)	855, 856
PROTEIN FORMULA WITH ARGININE, VITAMIN C, E AND ZINC	1039	<i>Quetiapine-DRLA</i> (RZ)	856
PROTEIN FORMULA WITH CARBOHYDRATE, FAT, VITAMINS AND MINERALS	956	<i>Quilonum SR</i> (AS)	876
PROTEIN FORMULA WITH VITAMINS AND MINERALS, AND LOW IN POTASSIUM, PHOSPHORUS, CALCIUM, CHLORIDE AND VITAMIN A	989	QUINAGOLIDE	192
PROTEIN HYDROLYSATE FORMULA WITH MEDIUM CHAIN TRIGLYCERIDES	967, 968	QUINAPRIL	140
<i>Provera</i> (PF) .ANTINEOPLASTIC AND IMMUNOMODULATING AGENTS	339, 340	QUINAPRIL + HYDROCHLOROTHIAZIDE.....	143
.GENITO URINARY SYSTEM AND SEX HORMONES	198	<i>Quinapril generichealth</i> (GQ).....	140
<i>Proxen SR 1000</i> (IY).....	764, 765, 997	<i>Quinate</i> (RW)	899
<i>Proxen SR 750</i> (IY).....	764, 765, 997	QUININE	899
<i>Prozac</i> 20(LY)	873	<i>QuitX</i> (AF)	1036
<i>ProZero</i> (VF).....	989	<i>QV Bath Oil</i> (EO).....	1023
<i>PRYZEX</i> (RW)	852, 853	<i>QV Gentle Wash</i> (EO)	1026
<i>PRYZEX ODT</i> (RW)	854	<i>QV Skin Lotion</i> (EO)	1023
<i>PRYZEX ODT</i> (RW)	853	<i>Qvar 100 Autohaler</i> (IL)	914
PSEUDOEPHEDRINE	1037	<i>Qvar 100</i> (IL)	914
PSYLLIUM HUSK POWDER	1017	<i>Qvar 50 Autohaler</i> (IL)	914
<i>PTU</i> (FF).....	210	<i>Qvar 50</i> (IL)	914
<i>Pulmicort Respules</i> (AP).....	915	RABEPRAZOLE	41, 42
<i>Pulmicort Turbuhaler</i> (AP)	914, 915	<i>Rabeprazole AN</i> (EA)	41, 42, 43
<i>Puregon 300 IU/0.36 mL</i> (OQ).....	200	<i>Rabeprazole Mylan</i> (AF).....	41, 42, 43
<i>Puregon 600 IU/0.72 mL</i> (OQ).....	200	<i>Rabeprazole Sandoz</i> (SZ)	41
<i>Puregon 900 IU/1.08 mL</i> (OQ).....	200	<i>Rabeprazole Sandoz</i> (SZ)	42, 43
<i>Purinethol</i> (AS)	255	<i>Rabeprazole SUN</i> (RN)	42, 43
<i>PVA Tears</i> (PE)	949, 950	<i>Ralovera</i> (FZ)	198
<i>Pyralin EN</i> (FZ)	59	RALOVISTA(RF)	776
PYRANTEL	900	RALOXIFENE	776
PYRIDOSTIGMINE	894	<i>Raloxifene AMNEAL</i> (ED)	776
<i>Qpril 10</i> (AF)	140	<i>Raloxifene AN</i> (EA)	776
<i>Qpril 20</i> (AF)	140	<i>Raloxifene GH</i> (GQ)	776
<i>Qpril 5</i> (AF)	140	<i>Ramace 1.25 mg</i> (AV)	141
<i>Qtern 5/10</i> (AP).....	77	<i>Ramace 10 mg</i> (AV)	140
QUEPINE XR (RW)	856	<i>Ramace 2.5 mg</i> (AV)	141
QUEPINE XR(RF)	855, 856	<i>Ramace 5 mg</i> (AV).....	141
<i>Questran Lite</i> (GO)	159	RAMIPRIL	140, 141
<i>Quetia 100</i> (RW).....	856	RAMIPRIL + FELODIPINE	144
<i>Quetia 200</i> (RW).....	855	<i>Ramipril AN</i> (EA)	140, 141
<i>Quetia 25</i> (RW).....	856	<i>Ramipril Sandoz</i> (SZ)	140, 141
<i>Quetia 300</i> (RW).....	855	<i>Ramipril Winthrop</i> (WA).....	140, 141
<i>Quetia XR</i> (OW).....	855, 856	<i>Rani 2</i> (AF).....	30
QUETIAPINE	855, 856	RANIBIZUMAB	938, 939, 941
<i>Quetiapine Actavis 100</i> (ED).....	856	RANITIDINE	30
<i>Quetiapine Actavis 200</i> (ED).....	855	<i>Ranitidine AN</i> (EA)	30
<i>Quetiapine Actavis 300</i> (ED).....	855	<i>Ransim</i> (RA)	157, 158
<i>Quetiapine AN</i> (EA)	855, 856	<i>Rapamune</i> (PF).....	372
<i>Quetiapine AN</i> (EA)	856	<i>Rapilysin 10 U</i> (TB)	112
<i>Quetiapine APOTEX</i> (GX)	856	RASAGILINE	848
<i>Quetiapine APOTEX</i> (GX)	855, 856	<i>Rasagiline-Teva</i> (EV)	848
<i>Quetiapine GH 100</i> (GQ)	856	<i>Rasalect</i> (TI).....	848
<i>Quetiapine GH 200</i> (GQ)	855	<i>RBX Topiramate</i> (RA).....	840, 841
<i>Quetiapine GH 25</i> (GQ)	856	RCF(AB)	989
<i>Quetiapine GH 300</i> (GQ)	855	<i>Reandron 1000</i> (BN)	196
<i>Quetiapine RBX</i> (RA).....	855, 856	<i>Reaptan 10/10</i> (RW).....	144
<i>Quetiapine RBX</i> (RA).....	856	<i>Reaptan 10/5</i> (RW).....	143
<i>Quetiapine Sandoz</i> (SZ)	856	<i>Reaptan 5/10</i> (RW).....	143
		<i>Reaptan 5/5</i> (RW).....	143
		<i>Rebif 44</i> (SG)	348
		REBOXETINE	877
		<i>ReddyMax Plus D-Cal</i> (RZ)	773
		<i>Redipred</i> (AS).....	208
		<i>Reedos 100</i> (DO)	837
		<i>Reedos 200</i> (DO)	836
		<i>Reedos 25</i> (DO)	836
		<i>Reedos 50</i> (DO)	837
		<i>Refresh Liquigel</i> (AG)	943, 944, 945
		<i>Refresh Night Time</i> (AG)	948
		<i>Refresh Tears Plus</i> (AG)	944, 945

<i>Relistor(LM)</i>	996	<i>Rixadone (AF)</i>	860, 861, 862, 863, 864, 865
<i>Relpax(UJ)</i>	825	<i>Rixadone(AF)</i>	859
<i>Remicade(JC)</i>	1032	<i>RIXALT (RF)</i>	826
<i>Reminyl(JC)</i>	887, 888, 889	<i>RIZATRIPTAN</i>	826
<i>Renagel(GZ)</i>	953	<i>Rizatriptan AN ODT(EA)</i>	826
<i>Renastart(VF)</i>	991	<i>Rizatriptan ODT APOTEX(GX)</i>	826
<i>Renastep(VF)</i>	989	<i>Rizatriptan ODT GH (GQ)</i>	826
<i>Renitec 20(AF)</i>	139	<i>Rizatriptan Wafers-10mg (AF)</i>	826
<i>Renitec Plus 20/6(AF)</i>	142	<i>Rizatriptan-AU (DZ)</i>	826
<i>Renitec(AF)</i>	138	<i>Roaccutane(RO)</i>	186
<i>Repatha(AN)</i>	160, 161, 162, 166	<i>Rocaltrol(IX)</i>	
<i>Replicare Ultra 66000434(SN)</i>	1052	ALIMENTARY TRACT AND METABOLISM	102
<i>Resonium-A(SW)</i>	1038	MUSCULO-SKELETAL SYSTEM	774
<i>Respikast 4 (RW)</i>	920	<i>Rocta 10(RW)</i>	186
<i>Respikast 5(RW)</i>	921	<i>Rocta 20 (RW)</i>	186
<i>Resprim Forte(AF)</i>	229, 230	<i>Roferon-A(RO)</i>	347
<i>restore O.R.S. (EA)</i>	55	<i>ROMOSOZUMAB</i>	776
<i>restore O.R.S.(EA)</i>	55, 1018	<i>Rosidal TCS 26484(LC)</i>	1039
<i>RETEPLASE</i>	112	<i>Rostor 10 (DO)</i>	156, 157
<i>RETINOL PALMITATE + PARAFFIN</i>	950	<i>Rostor 20 (DO)</i>	156
<i>Rexulti(LU)</i>	858	<i>Rostor 40 (DO)</i>	156, 157
<i>RHAMNUS FRANGULA + STERCULIA</i>	52, 995, 1017	<i>Rostor 5 (DO)</i>	155, 156
<i>Riamet 20mg/120mg Dispersible(SZ)</i>	899	<i>ROSUVASTATIN</i>	155, 156
<i>Riamet(SZ)</i>	899	<i>Rosuvastatin AMNEAL(EF)</i>	155, 156, 157
<i>RIBAVIRIN</i>	251	<i>Rosuvastatin APOTEX (GX)</i>	155, 156, 157
<i>RIBOCICLIB</i>	304, 305, 306	<i>Rosuvastatin generichealth(HQ)</i>	155, 156, 157
<i>Ridaura(BZ)</i>	767	<i>Rosuvastatin Lupin (GQ)</i>	155, 156, 157
<i>Ridaura(GH)</i>	767	<i>Rosuvastatin RBX(RA)</i>	155, 156, 157
<i>Rifadin(SW)</i>	245	<i>Rosuvastatin Sandoz (SZ)</i>	155, 156, 157
<i>RIFAMPICIN</i>		<i>Rosuzet Composite Pack(AL)</i>	168, 169
ANTIINFECTIVES FOR SYSTEMIC USE	244, 245	<i>ROTIGOTINE</i>	848
<i>RIFAXIMIN</i>	55	<i>Roxar 150(RW)</i>	231, 232
<i>Rilutek(SW)</i>	898	<i>Roxar 300(RW)</i>	231, 232
<i>RILUZOLE</i>	898	<i>Roxet 20(DO)</i>	873
<i>Riluzole Sandoz (SZ)</i>	898	<i>Roximycin (AF)</i>	231, 232
<i>Rimycin 150(AF)</i>		<i>Roxin (RW)</i>	236
ANTIINFECTIVES FOR SYSTEMIC USE	244, 245, 246	<i>ROXITHROMYCIN</i>	231, 232
<i>Rimycin 300(AF)</i>		<i>Roxithromycin AN(EA)</i>	231, 232
ANTIINFECTIVES FOR SYSTEMIC USE	244, 245, 246	<i>Roxithromycin Sandoz(SZ)</i>	231, 232
<i>Rinvoq(VE)</i>	393, 398	<i>Roxithromycin-GA (ED)</i>	231, 232
<i>RISANKIZUMAB</i>	638, 642	<i>Rozlytrek(RO)</i>	320
<i>Risedro once a week (RW)</i>	1032	<i>Rulide D(SW)</i>	232
<i>Risedro once a week(RW)</i>	770	<i>Rulide(SW)</i>	231, 232
<i>RISEDRONATE</i>	770, 1032	<i>RUXOLITINIB</i>	311, 312
<i>RISEDRONATE (&) CALCIUM CARBONATE</i>	773, 1033	<i>Rynacrom(SW)</i>	1037
<i>Risedronate AN(EA)</i>	770, 1032	<i>Rythmodan(SW)</i>	124
<i>Risedronate Sandoz (SZ)</i>	770	<i>Ryzodeg Flextouch(NO)</i>	62
<i>Risedronate Sandoz(SZ)</i>	1032	<i>Ryzodeg Penfill(NO)</i>	62
<i>Risedronate-GA (GN)</i>	1032	<i>Sabril(SW)</i>	832
<i>Rispa (RW)</i>	860, 861, 862, 863, 864, 865	<i>SACUBITRIL + VALSARTAN</i>	152
<i>Rispa(RW)</i>	859	<i>SAFINAMIDE</i>	848
<i>Risperdal (JC)</i>	859	<i>Saflutan(MF)</i>	933
<i>Risperdal Consta(JC)</i>	862	<i>Salazopyrin(PF)</i>	59
<i>Risperdal(JC)</i>	860, 861, 862, 863, 864, 865	<i>Salazopyrin-EN(PF)</i>	59
<i>RISPERIDONE</i>	859, 860, 861, 862, 863, 864, 865	<i>SALBUTAMOL</i>	
<i>Risperidone AMNEAL (EF)</i>	860, 861, 863, 864	Prescriber Bag	22
<i>Risperidone AMNEAL(EF)</i>	859, 862, 863, 864, 865	RESPIRATORY SYSTEM	902, 903, 919
<i>Risperidone generichealth (GQ)</i>	859	<i>Salbutamol Actavis (EA)</i>	22, 903
<i>Risperidone generichealth(GQ)</i>	860, 861, 863, 864	<i>Salbutamol AN(ED)</i>	22, 903
<i>Risperidone Sandoz (SZ)</i>	860, 861, 862, 863, 864, 865	<i>Salbutamol Cipla (LR)</i>	22, 903
<i>Risperidone Sandoz(SZ)</i>	859	<i>SALICYLIC ACID + BENZALKONIUM CHLORIDE +</i>	
<i>Rispernia (ER)</i>	859	ALCOHOL + COAL TAR SOLUTION +	
<i>Rispernia(ER)</i>	860, 861, 862, 863, 864, 865	POLYOXYETHYLENE ETHERS	1025
<i>Ritalin 10(NV)</i>	881	<i>SALICYLIC ACID + LACTIC ACID</i>	1025
<i>Ritalin LA(NV)</i>	882, 883	<i>SALMETEROL</i>	903
<i>Rithmik 200(RW)</i>	125	<i>Salofalk(FD)</i>	57, 58, 59
<i>RITUXIMAB</i>	329, 330	<i>SalplusF Inhaler 125/25(SZ)</i>	910
<i>RIVAROXYABAN</i>	116, 117, 118, 119	<i>SalplusF Inhaler 250/25(SZ)</i>	911
<i>RIVASTIGMINE</i>	889, 890	<i>Salpraz(AF)</i>	39, 40, 41
<i>Rivotril(RO)</i>	19, 829, 1004	<i>Sandomigran 0.5(AE)</i>	827

Sandoz Lamotrigine (HX)..... 836, 837
Sandoz Metformin (HX)..... 64
Sandoz Venlafaxine XR (HX)..... 878
Sandrena (AS)..... 197
Saphris (OQ)..... 852
SAPROPTERIN..... 104, 105
Savacol Mouth and Throat Rinse (OM)..... 1016
SAXAGLIPTIN..... 87
SAXAGLIPTIN + DAPAGLIFLOZIN..... 76, 77
SAXAGLIPTIN + METFORMIN..... 77
Sculptra (GA)..... 955
SebiRinse (EO)..... 1027
Sebitar (EO)..... 1025
SECUKINUMAB ... 649, 650, 652, 655, 657, 658, 661, 664, 667, 671, 674, 678
seebri breezhaler (NV)..... 916
Segluromet 2.5/1000 (MK)..... 72, 73
Segluromet 2.5/500 (MK)..... 72, 73
Segluromet 7.5/1000 (MK)..... 72, 73
Segluromet 7.5/500 (MK)..... 72, 73
SELEGILINE..... 849
SELENIUM SULFIDE..... 1025
Selsun (DQ)..... 1025
SEMAGLUTIDE..... 95
Semglee (AF)..... 63
Senna-Gen (PP)..... 1017
SENNOSIDE B..... 1017
Senokot (RC)..... 1017
Seprtin Forte (RW)..... 229, 230
Seprtin (RW)..... 230
Serenace (AS)..... 19, 850
Serepax (AS)..... 866, 867, 1004
Seretide Accuhaler 100/50 (GK)..... 910
Seretide Accuhaler 250/50 (GK)..... 910
Seretide Accuhaler 500/50 (GK)..... 911
Seretide MDI 125/25 (GK)..... 910
Seretide MDI 250/25 (GK)..... 911
Seretide MDI 50/25 (GK)..... 910
Serevent Accuhaler (GK)..... 903
Seroflo 125/25 (YC)..... 910
Seroflo 250/25 (YC)..... 911
Seroquel XR (AP)..... 855, 856
Seroquel (AP)..... 855, 856
Sertra 100 (RW)..... 874
Sertra 50 (RW)..... 873
SERTRALINE..... 873, 874
Sertraline AN (EA)..... 873, 874
Sertraline generichealth (GQ)..... 873, 874
Sertraline Sandoz (SZ)..... 873, 874
Sertraline tablets USP (Medsurge) (DZ)..... 873, 874
Setopress 3505 (MH)..... 1040
Setrona (RA)..... 873, 874
SEVELAMER..... 953
Sevelamer Apotex (TX)..... 953
Sevelamer Lupin (GQ)..... 953
Sevikar 20/5 (AL)..... 151
Sevikar 40/10 (AL)..... 150
Sevikar 40/5 (AL)..... 150
Sevikar HCT 20/5/12.5 (AF)..... 152
Sevikar HCT 40/10/12.5 (AF)..... 152
Sevikar HCT 40/10/25 (AF)..... 152
Sevikar HCT 40/5/12.5 (AF)..... 152
Sevikar HCT 40/5/25 (AF)..... 152
Sevredol (MF)..... 790, 998
Shilova 500 (DO)..... 249
Sical (AF)
 ALIMENTARY TRACT AND METABOLISM 102
 MUSCULO-SKELETAL SYSTEM 774
Sifrol ER (BY)..... 847
Sifrol (BY)..... 846, 847, 848

Sigmacort (AS)..... 176
Sigmaxin (LN)..... 124
Sigmaxin-PG (LN)..... 124
SILDENAFIL
 Repatriation Pharmaceutical Benefits Scheme 1028
Sildenafil Actavis (EA)..... 1028, 1029
Sildenafil APOTEX (GX)..... 1028, 1029
Sildenafil generichealth (GQ)..... 1028
Sildenafil Sandoz (SZ)..... 1028
Sildenafil Sandoz (SZ)..... 1028, 1029
SILODOSIN..... 1030
SILVER SULFADIAZINE..... 175
Simbrinza 1%/0.2% (NV)..... 929
Simipex 0.125 (RW)..... 846
Simipex 0.25 (RW)..... 847
Simipex 1 (RW)..... 846
SIMIPEX XR (RW)..... 847
Simponi (JC)... 575, 577, 579, 580, 581, 585, 587, 589, 594, 598, 602, 607
Simpral (AF)..... 846, 847
Simvar 10 (RW)..... 157, 158
Simvar 20 (RW)..... 157, 158
Simvar 40 (RW)..... 157, 158
Simvar 80 (RW)..... 157, 158
SIMVASTATIN..... 157
Simvastatin AN (EA)..... 157, 158
Simvastatin AN (EA)..... 157, 158
Simvastatin generichealth (GQ)..... 157, 158
Simvastatin generichealth (GQ)..... 157, 158
Simvastatin Sandoz (SZ)..... 157, 158
Simvastatin Sandoz (SZ)..... 157, 158
SINADOPA 100/25 (RW)..... 843
SINADOPA 250/25 (RW)..... 843
Sinemet 100/25 (AL)..... 843
Sinemet CR Prolonged-Release Tablets (OQ)..... 843
Sinemet CR (AL)..... 843
Sinemet (AL)..... 843
SIPONIMOD..... 371
SIROLIMUS..... 372
SITAGLIPTIN..... 89
SITAGLIPTIN + METFORMIN..... 79, 80
SKIN EMOLLIENT..... 1023
Skyrizi (VE)..... 642, 649
Sno-Pro (SB)..... 989
Sodibic (AS)..... 202
SODIUM CHLORIDE..... 1021
SODIUM CHLORIDE + HYPOCHLOROUS ACID + SODIUM HYPOCHLORITE..... 1057
SODIUM CHLORIDE + POTASSIUM CHLORIDE + GLUCOSE MONOHYDRATE + CITRIC ACID... 55, 1018
SODIUM PHENYL BUTYRATE..... 106
SODIUM POLYSTYRENE SULFONATE..... 1038
Sodium Valproate Sandoz (SZ)..... 831, 832
Soflax (EA)..... 1017
SOFOSBUVIR + VELPATASVIR..... 252
SOFOSBUVIR + VELPATASVIR + VOXILAPREVIR..... 252
Sofradex (SW)..... 951
Soframycin (SW)..... 951
Solaraze 3% Gel (FK)..... 1026
Solavert (RF)..... 125
Solian 100 (SW)..... 857
Solian 200 (SW)..... 857
Solian 400 (SW)..... 857
Solian Solution (SW)..... 857
Solone (IL)..... 208
SoloSite Gel 36361338 (SN)..... 1054
Solu-Cortef (PF)..... 20, 207, 208
Solugel 10336 (JJ)..... 1054
Solu-Medrol (PF)..... 208
Somac (NQ)..... 39, 40, 41

Somac(NQ).....	39, 40, 41	SUNSCREENS	1023
Sone (IL).....	209	Sunsense Sensitive SPF 50+(EO)	1023
SONIDEGIB.....	332	Sunsense Ultra SPF 50+(EO)	1023
SORAFENIB.....	324, 325	Suprasorb G 33631(LC)	1053
Sorbact Absorption Dressing S98222(YB).....	1053	Surepress 650947(CC).....	1039
Sorbact Foam Dressing S98310(YB).....	1054	Surepress 650948(CC).....	1039
Sorbact Foam Dressing S98315(YB).....	1055	Sutent(PF)	326, 327, 328, 329
Sorbact Foam Gentle Border 98532(YB).....	1055	Symbicort Rapihale 100/3(AP)	904, 906
Sorbact Foam Gentle Border 98533(YB).....	1055	Symbicort Rapihale 200/6(AP)	906, 908
Sorbact Gel Dressing S98137(YB).....	1054	Symbicort Rapihale 50/3(AP)	906, 907
Sorbact Ribbon Gauze S98118(YB).....	1054	Symbicort Turbuhaler 100/6(AP)	905, 907
Sorbact Ribbon Gauze S98120(YB).....	1054	Symbicort Turbuhaler 200/6 (AP)	906
Sorbact Superabsorbent 98501(YB).....	1055	Symbicort Turbuhaler 200/6(AP)	905, 907
Sorbact Superabsorbent 98503(YB).....	1055	Symbicort Turbuhaler 400/12(AP)	909
Sorbsan 1410(UM)	1045	Symmetrel 100(NV).....	845
Sorbsan 1411(UM)	1044	Synacthen Depot 1 mg/1 mL(IX)	203
Sotacor(RW).....	125	Synarel(PF)	206
SOTALOL	125	Syquet (AF)	856
Sotalol Sandoz (SZ).....	125	Syquet(AF)	855, 856
SOY LECITHIN + TOCOPHEROL + VITAMIN A.....	950	Systane(AQ).....	949
SOY PROTEIN AND FAT FORMULA WITH VITAMINS		Tacidine (AF).....	29
AND MINERALS CARBOHYDRATE FREE	989	Tacrogaf (RW).....	755
Sozol(RW).....	39, 40, 41	TACROLIMUS.....	755
Span-K(AS).....	103	TACROLIMUS APOTEX(TX)	755
Spedra(FK)	1028	Tacrolimus Sandoz (SZ)	755
Spiolto Respimat(BY)	914	Tacrolimus Sandoz(SZ).....	755
Spiractin 100(AF).....	132	Tadalaccord(CR)	1029
Spiractin 25(AF).....	131	TADALAFIL	
Spiriva Respimat(BY)	917, 918	Repatriation Pharmaceutical Benefits Scheme	1029
Spiriva(BY).....	918	Tadalafil Sandoz (SZ).....	1029
SPIRONOLACTONE	131	Tadalafil Sandoz(SZ).....	1029
Spirolactone Mylan 100 (AL).....	132	Tafinlar(NV)	294, 295
Spirolactone Mylan 25 (AL).....	131	TAFLUPROST.....	933
Sporanox (JC).....	242	Tagrisso(AP)	291, 292, 293
Spren 100(OW).....	109, 1020	Talam (RW)	871
Sprycel(BQ)	259, 261, 263	Talam(RW)	871
Stalevo 100/25/200mg (SZ)	844	Taltz(LY).....	617, 621, 625, 632, 638
Stalevo 125/31.25/200mg (SZ)	844	Tamate (AF)	840, 841
Stalevo 150/37.5/200mg (SZ)	844	Tambacor (IL).....	124
Stalevo 200/50/200mg (SZ)	844	Tambacor(IL).....	124
Stalevo 50/12.5/200mg (SZ).....	844	Tamosin(AS)	343
Stalevo 75/18.75/200mg (SZ).....	844	TAMOXIFEN	343
Staphylex 250(AF)	220, 221	Tamoxifen Sandoz (SZ).....	343
Staphylex 500(AF)	220, 221	Tamsil (RW)	174
Steglatro 15(MK).....	101, 102	Tamsil(RW).....	1023
Steglatro 5(MK).....	101, 102	TAMSULOSIN	1030
Steglujan 15/100(MK)	73, 74	Tamsulosin Sandoz SR (SZ)	1030
Steglujan 5/100(MK)	73, 74	TAPE NON WOVEN RETENTION POLYACRYLATE.....	1057
Stelara(JC).....	733, 736, 742, 747, 754	TAPE PLASTER ADHESIVE ELASTIC.....	1057
Stelax 10 (RW)	767	TAPE PLASTER ADHESIVE HYPOALLERGENIC	1058
Stelax 25 (RW)	767	TAPE PLASTER ADHESIVE WITH SILICONE.....	1058
Stemetil(SW).....	21, 50	TAPENTADOL	817
Stemzine(AV).....	50	TAR + COAL TAR SOLUTION + SALICYLIC ACID	1025
Steripaste 3610(MH).....	1043	TAR + TROLAMINE LAURIL SULFATE.....	1024
STIRIPENTOL	839	Targin 10/5mg(MF).....	804
Strattera(LY)	880	Targin 15/7.5mg(MF).....	804
Stromectol(MK).....	901	Targin 2.5/1.25 mg(MF).....	804
SUCROFERRIC OXYHYDROXIDE.....	953	Targin 20/10mg(MF).....	804
SULFASALAZINE.....	59	Targin 30/15 mg(MF).....	804
Sulprix (AF).....	857	Targin 40/20mg(MF).....	804
Sulprix(AF).....	857	Targin 5/2.5mg(MF).....	804
SULTHIAME	840	Targin 60/30(MF).....	804
Sumatran (OW).....	826	Targin 80/40(MF).....	804
Sumatran(OW).....	827	Tarka 2/180(GO)	144
SUMATRIPTAN.....	826	Tarka 4/240(GO)	144
Sumatriptan AN(EA)	827	Tasigna(NV)	282, 284
Sumatriptan generichealth (GQ).....	827	Tazac(RW).....	29
Sumatriptan Sandoz (SZ)	827	Tears Naturale(AQ)	945, 946
Sumatriptan Sandoz(SZ)	827	tearsagain(RB).....	950
SUNITINIB	326, 327, 328	Tecfidera(BD).....	756

<i>Tegaderm Transparent 1628(MM)</i>	1045	<i>Terbinafine Sandoz(SZ)</i>	174
<i>Tegaderm Transparent Island 3582(MM)</i>	1045	<i>Terbinafine-DRLA (RZ)</i>	174
<i>Tegaderm Transparent Island 3586(MM)</i>	1045	TERBUTALINE.....	
<i>Teglutik(CS)</i>	898	RESPIRATORY SYSTEM.....	903, 919
<i>Tegretol 100(NV)</i>	830	<i>TERIFLAGIO (RW)</i>	373
<i>Tegretol 200(NV)</i>	830	TERIFLUNOMIDE.....	372
<i>Tegretol CR 200(NV)</i>	830	<i>Teriflunomide Dr.Reddy's(RZ)</i>	373
<i>Tegretol CR 400(NV)</i>	830, 831	<i>Teriflunomide GH (GQ)</i>	373
<i>Tegretol Liquid(NV)</i>	830	<i>Teriflunomide Sandoz(SZ)</i>	373
<i>Telfa 8252F(KE)</i>	1041	<i>Terimide (AF)</i>	373
<i>Telfa 8253F(KE)</i>	1041	TERIPARATIDE.....	
<i>Telfa 8254F(KE)</i>	1041	MUSCULO-SKELETAL SYSTEM.....	777
<i>Telfast 120(SW)</i>	1038	SYSTEMIC HORMONAL PREPARATIONS, EXCL. SEX	
<i>Telfast(SW)</i>	1038	HORMONES AND INSULINS.....	211
TELMISARTAN.....	146	<i>Terry White Chemists Meloxicam(TW)</i>	762
TELMISARTAN + AMLODIPINE.....	151	<i>Terry White Chemists Metformin(TW)</i>	64
TELMISARTAN + HYDROCHLOROTHIAZIDE.....	149	<i>Tertroxin(AS)</i>	210
<i>Telmisartan AN (EA)</i>	146	<i>Testavan(FP)</i>	195
<i>Telmisartan GH (GQ)</i>	146	<i>Testogel(HB)</i>	195
<i>Telmisartan HCT GH 40/12.5(GQ)</i>	149	TESTOSTERONE.....	194
<i>Telmisartan HCT GH 80/12.5(GQ)</i>	149	TESTOSTERONE UNDECANOATE.....	195
<i>Telmisartan HCT GH 80/25(GQ)</i>	149	TETRABENAZINE.....	898
<i>Telmisartan HCTZ AN 40/12.5(EA)</i>	149	TETRACOSACTIDE (TETRACOSACTRIN).....	203
<i>Telmisartan HCTZ AN 80/12.5(EA)</i>	149	<i>Tevaripirazole (TB)</i>	857, 858
<i>Telmisartan HCTZ AN 80/25(EA)</i>	149	<i>Tevatiapine XR (SZ)</i>	855, 856
<i>Telmisartan Sandoz(SZ)</i>	146	<i>Teveten Plus 600/12.5(GO)</i>	147
<i>Telmisartan/HCT Sandoz (SZ)</i>	149	<i>Teveten(GO)</i>	145
<i>Telmisartan-DRLA(RZ)</i>	146	THEOPHYLLINE.....	919
<i>Teltartan (RW)</i>	146	THIAMINE.....	102, 1019
<i>Teltartan HCT 40/12.5 (RW)</i>	149	<i>Thioprine 50 (AF)</i>	755
<i>Teltartan HCT 80/12.5 (RW)</i>	149	<i>Thyrogen(GZ)</i>	204
<i>Teltartan HCT 80/25 (RW)</i>	149	THYROTROPIN ALFA.....	203
<i>Temaze (AF)</i>	868, 1005	TIAGABINE.....	831
TEMAZEPAM.....	868, 1005	TICAGRELOR.....	111
<i>Temizole 100 (AL)</i>	254	<i>Tielle MTL101E(KI)</i>	1050
<i>Temizole 140 (AL)</i>	254	<i>Tielle MTL103(KI)</i>	1050
<i>Temizole 20 (AL)</i>	254	<i>Tilade CFC-Free(SW)</i>	918
<i>Temizole 250 (AL)</i>	254	TILDRAKIZUMAB.....	684, 687
<i>Temizole 5 (AL)</i>	254, 255	TIMOLOL.....	931
<i>Temodal (MK)</i>	254	<i>Timoptol XE(MF)</i>	931
<i>Temodal(MK)</i>	254	<i>Timoptol(MF)</i>	931
<i>Temolide(JU)</i>	254	<i>Tinaderm(BN)</i>	1022
TEMOZOLOMIDE.....	254	<i>Tinasil (AF)</i>	174
<i>Temozolomide Alphapharm (AF)</i>	254	<i>Tinasil(AF)</i>	1023
<i>Temozolomide Alphapharm(AF)</i>	254, 255	TIOGUANINE.....	255
<i>Temozolomide Amneal(JO)</i>	254	TITROPIUM.....	916, 917, 918
<i>Temozolomide Juno (JX)</i>	254, 255	TITROPIUM + OLODATEROL.....	913
<i>Temozolomide Juno(JX)</i>	254	TIROFIBAN.....	111
<i>Temtabs(LN)</i>	868, 1005	<i>Tirofiban AC (JO)</i>	112
<i>Tenaxil SR (RW)</i>	130	<i>Tirofiban Juno(JU)</i>	112
<i>TenderWet 24 Active 609210(HR)</i>	1050	<i>Tixol (AL)</i>	876
<i>TenderWet 24 Active 609213(HR)</i>	1050	<i>TOBI podhaler(GO)</i>	234
<i>TenderWet Active Cavity 609272(HR)</i>	1050	<i>Tobi(GO)</i>	233
TENECTEPLASE.....	112	<i>Tobra-Day(FF)</i>	233
TENOFOVIR DISOPROXIL + EMTRICITABINE.....	252	TOBRAMYCIN.....	
<i>Tenofovir Disoproxil Emtricitabine Mylan 300/200(AF)</i> ..	252	ANTIINFECTIVES FOR SYSTEMIC USE.....	233, 234
<i>Tenofovir EMT GH(GQ)</i>	253	SENSORY ORGANS.....	922
<i>Tenofovir/Emtricitabine 300/200 APOTEX(TX)</i>	253	<i>Tobramycin Mylan (AF)</i>	234
<i>Tenolten 50(DO)</i>	134	TOBRAMYCIN SUN (RA).....	233
<i>Tenormin(AP)</i>	134	<i>Tobramycin WKT(LI)</i>	233
<i>Tensig (RW)</i>	134	TOBRAMYCIN WOCKHARDT (WC).....	233
<i>Tensocrepe 36300501(BV)</i>	1041	<i>Tobrex(NV)</i>	922
<i>Tensocrepe 36301001(BV)</i>	1041	TOCILIZUMAB694, 697, 699, 700, 702, 705, 707, 708, 713,	
<i>Tensocrepe 36307501(BV)</i>	1041	716, 720, 725.....	
<i>Tensopress 71723-00(BV)</i>	1039	TOFACITINIB.....	373, 376, 379, 384
TERBINAFINE.....		<i>Tofranil 10(GH)</i>	870
DERMATOLOGICALS.....	173, 174	<i>Tofranil 25(GH)</i>	870
Repatiation Pharmaceutical Benefits Scheme.....	1022	TOLNAFTATE.....	1022
<i>Terbinafine AN(EA)</i>	174	TOLVAPTAN.....	132
<i>Terbinafine Sandoz (SZ)</i>	1023	<i>Topamax Sprinkle(JC)</i>	840, 841

<i>Topamax</i> (JC).....	840, 841	TRIMETHOPRIM + SULFAMETHOXAZOLE	229, 230
TOPIRAMATE	840, 841	<i>Trimethoprim Mylan</i> (AL)	229
<i>Topiramate AN</i> (EA)	840, 841	<i>Triphasil 28</i> (PF)	193
<i>Topiramate Sandoz</i> (SZ).....	840, 841	<i>Triprim</i> (RW).....	229
<i>Toprol-XL 190</i> (AP).....	135	TRIPTORELIN.....	342
<i>Toprol-XL 47.5</i> (AP).....	135	<i>Triquilar ED</i> (BN)	193
<i>Toprol-XL 95</i> (AP).....	135	<i>Tritace</i> (SW)	140
TOREMIFENE	344	<i>Tritace 1.25 mg</i> (SW)	141
<i>Torvastat 10</i> (RW)	153, 154	<i>Tritace 10 mg</i> (SW)	140
<i>Torvastat 20</i> (RW)	153, 154	<i>Tritace 2.5 mg</i> (SW)	141
<i>Torvastat 40</i> (RW)	153, 154	<i>Tritace 5 mg</i> (SW)	141
<i>Torvastat 80</i> (RW)	153, 154	TROPISETRON	48
<i>Toujeo Solostar</i> (SW)	63	<i>Tropisetron-AFT</i> (AE)	48
<i>Trajenta</i> (BY)	85, 87	<i>Trovas</i> (RA)	153, 154
<i>Trajentamet</i> (BY)	74, 76	<i>Trulicity</i> (LY)	93
TRAMADOL.....	22, 819, 820, 821	<i>Trusamide</i> (AF)	930
<i>Tramadol ACT</i> (JO)	22, 819, 821	<i>Trusopt</i> (MF).....	930
<i>Tramadol AMNEAL</i> (EF).....	819, 821	<i>Trust Aspirin EC 100</i> (CR).....	1020
<i>Tramadol AN</i> (JU).....	22, 819, 821	<i>Trust Cetirizine</i> (CR).....	1038
<i>Tramadol AN SR</i> (EA)	822	<i>Trust Coloxease</i> (CR).....	1017
<i>Tramadol AN</i> (EA)	819, 821	<i>Trust Cystitis Relief</i> (CR).....	1029
<i>Tramadol Sandoz</i> (SZ).....	819, 821	<i>Trust Decongestant Nasal Spray</i> (CR).....	1037
<i>Tramadol Sandoz SR</i> (SZ).....	822	<i>Trust Deworm</i> (CR)	1036
<i>Tramadol Sandoz</i> (SZ).....	22, 819, 821	<i>Trust Fem V 3 Day Cream</i> (CR).....	1027
<i>Tramadol SR generichealth</i> (GQ)	822	<i>Trust Fem V 6 Day Cream</i> (CR)	1027
<i>Tramal 100</i> (CS).....	22, 819, 821	<i>Trust Fexit 120</i> (CR).....	1038
<i>Tramal SR 100</i> (CS)	822	<i>Trust for Kids Paracetamol 6 to 12 years</i> (CR).....	1034
<i>Tramal SR 150</i> (CS)	823	<i>Trust HydroCortic Cream</i> (CR).....	1024
<i>Tramal SR 200</i> (CS)	822	<i>Trust Loratadine</i> (CR).....	1038
<i>Tramal SR 50</i> (CS)	822	<i>Trust Loratadine Antihistamine</i> (RM).....	1038
<i>Tramal</i> (CS).....	819, 820, 821	<i>Trust Nystatin Oral Drops</i> (CR)	1016
<i>Tramedo SR</i> (AL).....	822	<i>Trust Sinus & Nasal Decongestant</i> (CR)	1037
<i>Tramedo</i> (AF)	819, 821	<i>Trust Terbinafine Cream</i> (CR)	1022
TRAMETINIB.....	301, 302	<i>Tryzan Caps 1.25</i> (AF)	141
<i>Tranalpha</i> (AF).....	141	<i>Tryzan Caps 10</i> (AF)	140
<i>Trandate</i> (AS)	136	<i>Tryzan Caps 2.5</i> (AF)	141
TRANDOLAPRIL	141	<i>Tryzan Caps 5</i> (AF)	141
TRANDOLAPRIL + VERAPAMIL.....	144	<i>Tryzan Tabs 1.25</i> (AF).....	141
TRANEXAMIC ACID	121	<i>Tryzan Tabs 10</i> (AF).....	140
<i>Transiderm-Nitro 25</i> (SZ).....	126	<i>Tryzan Tabs 2.5</i> (AF).....	141
<i>Transiderm-Nitro 50</i> (SZ).....	126	<i>Tryzan Tabs 5</i> (AF).....	141
TRANLYCYPROMINE	875	<i>Tubegauz 0501633</i> (SS)	1042
TRASTUZUMAB.....	330, 331	<i>Tubifast 2434</i> (MH).....	1042
<i>Travatan</i> (NV)	933	<i>Tubifast 2436</i> (MH).....	1042
TRAVOPROST	933	<i>Tubifast 2438</i> (MH).....	1042
TRAVOPROST + TIMOLOL	933	<i>Tubigrip 1479</i> (MH).....	1042
<i>Trelegy Elipta 100/62.5/25</i> (GK).....	912	<i>Tubigrip 1480</i> (MH).....	1042
<i>Tremfya</i> (JC).....	615	<i>Tubigrip 1481</i> (MH).....	1042
<i>Trexject</i> (LM).....	757, 758	<i>Tubigrip 1482</i> (MH).....	1042
TRIAMCINOLONE		<i>Tubigrip 1483</i> (MH).....	1042
DERMATOLOGICALS	176	<i>Tubigrip 1484</i> (MH).....	1042
SYSTEMIC HORMONAL PREPARATIONS, EXCL. SEX		<i>Tubigrip 1486</i> (MH).....	1042
HORMONES AND INSULINS	209	<i>Tubigrip B 1520</i> (MH).....	1041
TRIAMCINOLONE + NEOMYCIN + GRAMICIDIN +		<i>Tubigrip C 1545</i> (MH)	1041
NYSTATIN	951	<i>Tubigrip D 1546</i> (MH)	1041
<i>Triasyn 2.5/2.5</i> (SW).....	144	<i>Tubigrip E 1547</i> (MH)	1041
<i>Triasyn 5.0/5.0</i> (SW).....	144	<i>Tubigrip F 1548</i> (MH)	1041
<i>Tricortone</i> (LN).....	176	<i>Twynsta</i> (BY).....	151
<i>Trifeme 28</i> (FZ).....	193	<i>Tykerb</i> (NV)	310, 311
TRIFLURIDINE + TIPIRACIL	256	<i>Tylactin Build 20</i> (QH)	986
TRIGLYCERIDES LONG CHAIN WITH GLUCOSE		<i>Tylactin Complete</i> (QH).....	986
POLYMER.....	989	<i>Tylactin RTD</i> (QH)	986
TRIGLYCERIDES MEDIUM CHAIN AND LONG CHAIN		<i>TYR Anamix infant</i> (SB)	981
WITH GLUCOSE POLYMER.....	989	<i>TYR Anamix junior LQ</i> (SB).....	977
TRIGLYCERIDES MEDIUM CHAIN FORMULA		<i>TYR Anamix Junior</i> (SB)	982
VARIOUS	969, 970, 990	<i>TYR cooler 10</i> (VF).....	981
TRIHENXYPHENIDYL (BENZHEXOL)	842	<i>TYR cooler 15</i> (VF).....	981
<i>Trihexyphenidyl hydrochloride USP (Medsurge)</i> (DZ)	842	<i>TYR cooler 20</i> (VF).....	981
<i>Trileptal</i> (NV).....	831	<i>TYR Easy Shake & Go</i> (OH)	977
TRIMETHOPRIM	229	<i>TYR Easy Tablet</i> (OH).....	977

TYR express 15(VF)	981	Valtrex(RW)	249, 250
TYR express 20(VF)	981	Vancocin(AS)	55
TYR gel(VF)	981	VANCOMYCIN	
TYR Lophlex LQ 20(SB)	981	ALIMENTARY TRACT AND METABOLISM	55
TYR Lophlex(SB)	981	ANTIINFECTIVES FOR SYSTEMIC USE	236, 237
TYR Sphere20(VF)	986	Vancomycin Alphapharm(AF)	236, 237
Tyrosine 1000(VF)	990	VARDENAFIL	1029
TYROSINE WITH CARBOHYDRATE	990	VARENICLINE	896, 897
ultibro breezhaler 110/50(NV)	913	Vasafil 100(RW)	1028
UMECLIDINIUM	918	Vasafil 25 (RW)	1028
UMECLIDINIUM + VILANTEROL	914	Vasafil 50 (RW)	1029
UPADACITINIB	389, 393	Vasocardol (AV)	138
Uracol(EA)	1029	Vasocardol CD (AV)	138
Ural Sachets (AS)	1029	Vedafil (AF)	1028
Uramet (AS)	238	Vedafil(AF)	1028, 1029
UREA	1023	Vedilol 12.5(RW)	136
Urederm(KY)	1023	Vedilol 25(RW)	136
Uremide(AF)	130	Vedilol 3.125 (RW)	136
Urex(RW)	130	Vedilol 6.25(RW)	136
Urex-Forte(RW)	130	Velphoro(VL)	954
Urex-M (RW)	20, 131	VEMURAFENIB	295, 296
Uro-Carb(YN)	894	Venclexta(VE)	338, 339
Uromitexan(BX)	954	VENETOCLAX	337, 338, 339
Urorec(YN)	1030	VENLAFAXINE	877
URSODEOXYCHOLIC ACID	50	Venlafaxine AN SR(EA)	878
Ursodox GH (GQ)	51	Venlafaxine generichealth XR (GQ)	878
Ursofalk(FD)	51	Venofer(VL)	122
Ursosan (BZ)	51	Ventolin CFC-Free with dose counter (GK)	22, 902
USTEKINUMAB	730, 733, 736, 742, 747	Ventolin CFC-free(GK)	22, 902
Vaclovir (AF)	249	Ventolin Nebules(GK)	22, 903
Vaclovir(AF)	249	Ventolin(GK)	919
Vagifem Low(NO)	196	Vepesid(LM)	257
VALACICLOVIR	248, 249	VERAPAMIL	137
Valaciclovir AN (EA)	249	Verzenio(LY)	303
Valaciclovir AN(EA)	249	Vexazone (AF)	84
Valaciclovir APOTEX (GX)	249	Vfend(PF)	243, 244
Valaciclovir APOTEX(GX)	249	Viagra(UJ)	1028, 1029
Valaciclovir generichealth (GQ)	249	VIGABATRIN	832
Valaciclovir generichealth(GQ)	249	VILDAGLIPTIN	91
Valaciclovir RBX (RA)	249	VILDAGLIPTIN + METFORMIN	81
Valaciclovir RBX(RA)	249	Vimpat(UC)	835, 836
Valaciclovir Sandoz (SZ)	249	VINORELBINE	257
Valaciclovir Sandoz(SZ)	249	Viscopaste 4948(SN)	1043
Valaciclovir SZ(HX)	249	Viscotears Gel PF(UO)	943
Valacor 500 (CR)	249	Viscotears(UO)	943
Valine 1000(VF)	990	VISMODEGIB	334
Valine 50(VF)	990	Vita-B12 (GH)	122
VALINE WITH CARBOHYDRATE	990	Vita-B12(GH)	1021
Valium(RO)	866, 1004	VITAMINS, MINERALS AND TRACE ELEMENTS	990
Valpam 2 (RW)	866, 1004	VITAMINS, MINERALS AND TRACE ELEMENTS WITH	
Valpam 5 (RW)	866, 1004	CARBOHYDRATE	991
Valprease 200 (RW)	832	VitA-POS(AE)	950
Valprease 500 (RW)	831	Vliwasorb Pro 32641(LC)	1056
Valpro EC 200 (AF)	832	Vliwasorb Pro 32643(LC)	1056
Valpro EC 500 (AF)	832	Vliwasorb Pro 32644(LC)	1056
VALPROATE	831	Volirop 12.5 (DO)	136
Valproate Winthrop EC 200(WA)	832	Volirop 25 (DO)	136
Valproate Winthrop EC 500(WA)	832	Volirop 3.125(DO)	136
VALSARTAN	146	Volirop 6.25 (DO)	136
VALSARTAN + HYDROCHLOROTHIAZIDE	149	Voltaren 100(NV)	760, 996
Valsartan/Amlodipine Novartis 160/10(NM)	150	Voltaren 25(NV)	760, 996
Valsartan/Amlodipine Novartis 160/5(NM)	150	Voltaren 50(NV)	760, 996
Valsartan/Amlodipine Novartis 320/10(NM)	150	Voluvon 6%(PK)	123
Valsartan/Amlodipine Novartis 320/5(NM)	150	VORICONAZOLE	242, 243, 244
Valsartan/Amlodipine Novartis 80/5(NM)	150	Voriconazole APO (GX)	243
Valsartan/Amlodipine/HCT Novartis 160/10/12.5(NM)	151	Voriconazole APOTEX(TX)	243
Valsartan/Amlodipine/HCT Novartis 160/10/25(NM)	151	Voriconazole Sandoz (SZ)	243
Valsartan/Amlodipine/HCT Novartis 160/5/12.5(NM)	151	VORINOSTAT	331, 332
Valsartan/Amlodipine/HCT Novartis 160/5/25(NM)	151	Vosevi(GI)	252
Valsartan/Amlodipine/HCT Novartis 320/10/25(NM)	151	Votrient(NV)	322, 323, 324

Vttack(AF).....	243	Zilfojim ODT 8(DO).....	47
Vytorin (AL).....	171, 172	Zimstat (AF).....	157, 158
Vyvanse(TK).....	881	Zimstat(AF).....	157, 158
Vzole (RW).....	243	ZINC OXIDE + MAIZE STARCH + PURIFIED TALC +	
WARFARIN.....	107	CHLORPHENESIN.....	1027
Waxsol(GO).....	1038	ZINC OXIDE + PERU BALSAM + BENZYL BENZOATE	
Wellvone(AS).....	899		1022
WHEY PROTEIN FORMULA SUPPLEMENTED WITH		Zinnat (AS).....	227
AMINO ACIDS, LONG CHAIN POLYUNSATURATED		Zinnat(AS).....	227
FATTY ACIDS, VITAMINS AND MINERALS, LOW IN		Zinopril 10(AL).....	139
PROTEIN, PHOSPHATE, POTASSIUM AND LACTOSE		Zinopril 20(AL).....	139
.....	991	Zinopril 5(AL).....	139
WHEY PROTEIN FORMULA SUPPLEMENTED WITH		ZIPRASIDONE.....	851
AMINO ACIDS, VITAMINS AND MINERALS, AND LOW		Ziprasidone GH(GQ).....	851
IN PROTEIN, PHOSPHATE, POTASSIUM AND		ZIPROX (RW).....	851
LACTOSE.....	991	ZipZoc 66000747(SN).....	1043
WOOL ALCOHOLS.....	1023	Zircol 10(AL).....	137
Xadago(CS).....	848, 849	Zircol 20(AL).....	137
Xalacom(AS).....	932	ZITHRO (RW)	
Xalamol 50/5 (AF).....	932	.ANTIINFECTIVES FOR SYSTEMIC USE.....	230
Xalaprost (AF).....	932	.SENSORY ORGANS.....	921
Xalatan(AS).....	932	ZITHRO(RF)	
Xalkori(PF).....	298, 299	.Repatriation Pharmaceutical Benefits Scheme.....	1031
Xarelto(BN).....	117, 118, 119, 120	Zithromax (PF)	
Xatral SR(SW).....	1030	.Repatriation Pharmaceutical Benefits Scheme.....	1031
Xelabine (AL).....	256	Zithromax(PF)	
Xeljanz(PF).....	376, 379, 384, 389	.ANTIINFECTIVES FOR SYSTEMIC USE.....	230
Xenical(PB).....	1019	.SENSORY ORGANS.....	921
Xergic(AF).....	1038	Zocor (MQ).....	157, 158
Xgeva(AN).....	775	Zofran syrup 50 mL(AS).....	46, 47
Xifaxan(NE).....	55	Zofran Zydys(AS).....	47, 48
Xigduo XR 10/1000(AP).....	66, 68	Zofran(AS).....	46
Xigduo XR 10/500(AP).....	66, 68	ZolaCos CP 10.8/50(28)(AP).....	340
Xigduo XR 5/1000(AP).....	66, 68	ZolaCos CP 10.8/50(84)(AP).....	341
XLYS, LOW TRY Maxamum(SB).....	978	ZolaCos CP 3.6/50(AP).....	341
XMET Maxamum(SB).....	978	Zoladex 10.8 Implant(AP).....	340
XMTVI Maxamum(SB).....	979	Zoladex Implant(AP).....	340
XP Maxamum(SB).....	980	Zoledasta(TX).....	771
XPhen, Tyr Maxamum(SB).....	981	ZOLEDRONIC ACID.....	771
Xtandi(LL).....	345	Zoledronic Acid SUN (RA).....	771
Xylocaine Viscous(AS).....	1024	Zolinza(MK).....	332
Xylocard 500(AS).....	124	ZOLMITRIPTAN.....	827
YELLOW JACKET VENOM.....	952	Zoloft (UJ).....	874
Z.S.C.(RW).....	1027	Zoloft(UJ).....	873, 874
Zabep(AL).....	42, 43	Zoltrip (RW).....	827
Zactin Tablet (AF).....	872	Zomig(AP).....	827
Zactin(AF).....	873	Zonegran (EI).....	841, 842
Zan-Extra 10/10(GO).....	143	ZONISAMIDE.....	841
Zan-Extra 10/20(GO).....	143	ZOPICLONE.....	1036
Zanidip(GO).....	137	Zopiclone GH(GQ).....	1036
Zantac Syrup(AS).....	30	Zopral (AF).....	34, 35, 36
Zantac(AS).....	30	Zopral ODT(AF).....	34, 35, 36
Zarontin(IX).....	828, 829	Zopral(AF).....	34
Zatamil(EO).....	182, 183, 184, 185	Zoton FasTabs(PF).....	34, 35, 36
Zedd 500(RW).....	1031	Zotren 4 (RF).....	46
Zeklen 10/10 mg(AF).....	172	Zotren 8 (RF).....	46
Zeklen 10/20 mg(AF).....	172	Zotren ODT (RF).....	47
Zeklen 10/40 mg(AF).....	171	Zotren ODT(RF).....	47, 48
Zeklen 10/80 mg(AF).....	171	ZUCLOPENTHIXOL DECANOATE.....	851
Zelboraf(RO).....	296	Zumenon(GO).....	196
Zeldox (UJ).....	851	Zyban(AS).....	894, 895
Zelitrex (RF).....	249	Zydelig(GI).....	317
Zelitrex(RF).....	249, 250	Zydol (RW).....	819, 821
Zempreon CFC-Free with dose counter(GC).....	22, 902	Zydol SR 100 (RW).....	822
Zentel(AS).....	900	Zydol SR 150 (RW).....	822
Zeposia(CJ).....	371	Zydol SR 200 (RW).....	822
Zestril(AP).....	139	Zykadia(NV).....	298
ZETIN(RW).....	175	Zyloprim(RW).....	768
Zient 10mg(AF).....	166, 167	Zypine ODT(AF).....	853, 854
Zilfojim ODT 4(DO).....	47	Zypine(AF).....	852, 853

Zyprexa Relprevv(LY) 854, 855
Zyprexa Zydys(LY) 853, 854

Zyprexa(LY).....852, 853
Zytiga(JC).....346, 347